


NOVEL PYRIDINE DERIVATIVES

The present invention relates to organic compounds useful for therapy and/or prophylaxis in a mammal, and in particular to compounds that are preferential inhibitors of the cysteine protease cathepsin, in particular of the cysteine protease cathepsin S or L.

5 The invention relates in particular to a compound of formula (I)



wherein

A^1 is -S- or -S(O)₂-;

A^2 is nitrogen or -(CH)-;

10 R^1 is halogen or haloalkyl;

R^2 is hydrogen or halogen;

R^3 is hydrogen, halogen, haloalkyl, pyrazolyl, [1,2,3]-triazolyl or [1,2,4]-triazolyl;

R^4 and R^6 are independently selected from hydrogen, alkyl, haloalkyl and halophenyl; and

R^5 is hydrogen, halogen, haloalkyl, alkoxy, haloalkoxy, alkylpyridinyl, halopyridinyl or alkylpyrazolyl;

5 or a pharmaceutically acceptable salt thereof.

The compounds of the invention are preferential inhibitors of the cysteine protease Cathepsin (Cat), in particular Cathepsin S or Cathepsin L and are therefore useful to treat metabolic diseases like diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction of cardiovascular events in chronic kidney disease,
10 glomerulonephritis, age related macular degeneration, diabetic nephropathy and diabetic retinopathy. In addition, immune mediated diseases like rheumatoid arthritis, crohn's disease, multiple sclerosis, sjorgen syndrome, lupus erythematosus, neuropathic pain, diabetes type I, asthma and allergy and skin related immune disease are suitable diseases to be treated with a cathepsin S inhibitor.

15 Objects of the present invention are the compounds of formula (I) and their aforementioned salts per se and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts, the use of the said compounds and salts for the prophylaxis and/or therapy of illnesses,
20 especially in the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction of cardiovascular events in chronic kidney disease and diabetic nephropathy, and the use of the said compounds and salts for the production of medicaments for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction
25 of cardiovascular events in chronic kidney disease and diabetic nephropathy.

Mammalian cathepsins are cysteine-type proteases involved in key steps of biological and pathological events. Cathepsins are considered tractable drug targets as it is feasible to inhibit enzymatic activity with small molecules and are therefore of interest to the pharmaceutical industry (Bromme, D. (2001), 'Papain-like cysteine proteases', Curr
30 Protoc Protein Sci, Chapter 21, Unit 21 2; Roberts, R. (2005), 'Lysosomal cysteine proteases: structure, function and inhibition of cathepsins', Drug News Perspect, 18 (10), 605-14).

Cathepsin S is prominently expressed in antigen presenting cells like macrophages and dendritic cells and smooth muscle cells (Hsing, L. C. and Rudensky, A. Y. (2005),

'The lysosomal cysteine proteases in MHC class II antigen presentation', *Immunol Rev*, 207, 229-41; Rudensky, A. and Beers, C. (2006), 'Lysosomal cysteine proteases and antigen presentation', *Ernst Schering Res Found Workshop*, (56), 81-95). While Cathepsin S is only weakly expressed in normal arterial tissue, strong upregulation is seen in
5 atherosclerotic arteries (Liu, J., et al. (2006), 'Increased serum cathepsin S in patients with atherosclerosis and diabetes', *Atherosclerosis*, 186 (2), 411-9; Sukhova, G. K., et al. (1998), 'Expression of the elastolytic cathepsins S and K in human atheroma and regulation of their production in smooth muscle cells', *J Clin Invest*, 102 (3), 576-83).

Preclinical data suggest that the function of Cathepsin S is critical for atherosclerosis
10 as Cathepsin S deficient mice have a reduced atherosclerosis-phenotype when tested in appropriate mouse models. In LDL-Rec deficient mice reduced lipid accumulation, elastin-fibre breakdown and chronic arterial inflammation is reported. In APO E deficient mice a significant reduction of acute plaque rupture events was reported. When chronic renal disease is introduced into CatS/In APO-E deficient mice a strong reduction of accelerated
15 calcification is seen on top of the anti atherosclerotic activity in arteries and heart valves Aikawa, E., et al. (2009), 'Arterial and aortic valve calcification abolished by elastolytic cathepsin S deficiency in chronic renal disease', *Circulation*, 119 (13), 1785-94; de Nooijer, R., et al. (2009), 'Leukocyte cathepsin S is a potent regulator of both cell and matrix turnover in advanced atherosclerosis', *Arterioscler Thromb Vasc Biol*, 29 (2), 188-
20 94; Rodgers, K. J., et al. (2006), 'Destabilizing role of cathepsin S in murine atherosclerotic plaques', *Arterioscler Thromb Vasc Biol*, 26 (4), 851-6; Sukhova et al. (2003), 'Deficiency of cathepsin S reduces atherosclerosis in LDL receptor-deficient mice', *J Clin Invest*, 111 (6), 897-906). This suggests a potential inhibitor of Cathepsin S would stabilise atherosclerotic plaque by reducing extracellular matrix breakdown, by reducing
25 the proinflammatory state and by reducing accelerated calcification and subsequently its clinical manifestations.

These phenotypes described in atherosclerosis models are in agreement with known cellular functions of Cathepsin S. Firstly, Cathepsin S is involved in the degradation of extracellular matrix that stabilises the plaque. In particular, Cathepsin S has potent
30 elastolytic activity and can exert this at neutral pH, a feature that distinguishes Cathepsin S from all other Cathepsins. Secondly, Cathepsin S is the major protease involved in antigen processing, in particular cleavage of the invariant chain in antigen presenting cells, resulting in reduced contribution of Tcells to the chronic inflammation of the atherosclerotic tissue. Elevated inflammation results in further oxidative and proteolytic
35 tissue damage and subsequently plaque destabilisation (Cheng, X. W., et al. (2004), 'Increased expression of elastolytic cysteine proteases, cathepsins S and K, in the neointima of balloon-injured rat carotid arteries', *Am J Pathol*, 164 (1), 243-51; Driessen,

C., et al. (1999), 'Cathepsin S controls the trafficking and maturation of MHC class II molecules in dendritic cells', *J Cell Biol*, 147 (4), 775-90; Rudensky, A. and Beers, C. (2006), 'Lysosomal cysteine proteases and antigen presentation', *Ernst Schering Res Found Workshop*, (56), 81-95).

5 The anti-inflammatory and anti-elastinolytic properties of a Cat S inhibitor make it also a prominent target for chronic obstructive pulmonary disease (Williams, A. S., et al. (2009), 'Role of cathepsin S in ozone-induced airway hyperresponsiveness and inflammation', *Pulm Pharmacol Ther*, 22 (1), 27-32). Furthermore due to its extracellular functions in matrix degradation, inhibition of cathepsin S will impact neointima formation
10 and angiogenesis (Burns-Kurtis, C. L., et al. (2004), 'Cathepsin S expression is up-regulated following balloon angioplasty in the hypercholesterolemic rabbit', *Cardiovasc Res*, 62 (3), 610-20; Cheng, X. W., et al. (2004), 'Increased expression of elastolytic cysteine proteases, cathepsins S and K, in the neointima of balloon-injured rat carotid arteries', *Am J Pathol*, 164 (1), 243-51; Shi, G. P., et al. (2003), 'Deficiency of the cysteine
15 protease cathepsin S impairs microvessel growth', *Circ Res*, 92 (5), 493-500; Wang, B., et al. (2006), 'Cathepsin S controls angiogenesis and tumor growth via matrix-derived angiogenic factors', *J Biol Chem*, 281 (9), 6020-9). An inhibitor of Cathepsin S might therefore be useful in several different disease settings.

 Cathepsin S plays also a role in the reduction of tumor growth and tumor cell
20 invasion as described by Roberta E. Burden in *Clin Cancer Res* 2009;15(19). In addition, nephrectomized Cathepsin S knock out mice showed a significant reduction of arterial calcification when compared to nephrectomized wild type mice. This indicates that inhibition of Cathepsin S may have a beneficial effect on the reduction of cardiovascular events in chronic kidney disease patients (Elena Aikawa, *Circulation*, 2009, 1785-1794).

25 Cathepsin L shows a broader expression profile than cathepsin S and there are also data which suggest a role of cathepsin L in atherosclerosis, e.g. LDLrec & Cat L deficient mice show a reduced atherosclerotic phenotype (Kitamoto, S., et al. (2007), 'Cathepsin L deficiency reduces diet-induced atherosclerosis in low-density lipoprotein receptor-knockout mice', *Circulation*, 115 (15), 2065-75). In addition, Cat L was suggested to be
30 involved in metabolic syndrome as it controls adipogenesis and peripheral glucose tolerance. In renal disease Cathepsin L is described to regulate podocyte function by proteolytically processing dynamin and thereby proteinuria (Sever, S., et al. (2007), 'Proteolytic processing of dynamin by cytoplasmic cathepsin L is a mechanism for proteinuric kidney disease', *J Clin Invest*, 117 (8), 2095-104).

35 Tissue remodelling, extracellular matrix degradation, the generation of active neuropeptides and roles in antigen presentation in thymic epithelial cells are cellular

- activities described for Cathepsin L (Funkelstein et al. (2008), (a) Major role of cathepsin L for producing the peptide hormones ACTH, β -Endorphin, and α -MSH, illustrated by protease gene knockout and expression, Journal of Biological Chemistry, 283(51), 35652-35659; (b) Cathepsin L participates in the production of neuropeptide Y in secretory
- 5 vesicles, demonstrated by protease gene knockout and expression, Journal of Neurochemistry, 106(1), 384-391, Rudensky and Beers 2006).

In the present description the term “alkyl”, alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, in particular a straight or branched-chain alkyl group with 1 to 6 carbon atoms and particularly a straight

10 or branched-chain alkyl group with 1 to 4 carbon atoms. Examples of straight-chain and branched C1-C8 alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, in particular methyl, ethyl, propyl, isopropyl, isobutyl and tert.-butyl, more particularly methyl.

- 15 The term “alkoxy”, alone or in combination, signifies a group of the formula alkyl-O- in which the term “alkyl” has the previously given significance, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec. butoxy and tert.butoxy, in particular methoxy.

The term “oxy”, alone or in combination, signifies the -O- group.

- 20 The term “halogen” or “halo”, alone or in combination, signifies fluorine, chlorine, bromine or iodine.

The terms “haloalkyl” and “haloalkoxy”, alone or in combination, denote an alkyl group and an alkoxy group substituted with at least one halogen, in particular substituted with one to five halogens, particularly one to three halogens. A particular “haloalkyl” is

25 trifluoromethyl. Particular haloalkoxy are trifluoroethoxy and trifluoropropoxy.

- The term “pharmaceutically acceptable salts” refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, in
- 30 particular, hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein. In addition these salts may be prepared from addition of an inorganic base or an organic base to the

free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and
5 basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polyamine resins. The compound of formula (I) can also be present in the form of zwitterions. Particular pharmaceutically acceptable salts of compound of formula (I) are the salts of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid and
10 methanesulfonic acid.

If one of the starting materials or compounds of formula (I) contains one or more functional groups which are not stable or are reactive under the reaction conditions of one or more reaction steps, appropriate protecting groups (as described e.g. in "Protective Groups in Organic Chemistry" by T. W. Greene and P. G. M. Wuts, 3rd Ed., 1999, Wiley,
15 New York) can be introduced before the critical step applying methods well known in the art. Such protecting groups can be removed at a later stage of the synthesis using standard methods described in the literature. Examples of protecting groups are tert-butoxycarbonyl (Boc), 9-fluorenylmethyl carbamate (Fmoc), 2-trimethylsilylethyl carbamate (Teoc), carbobenzyloxy (Cbz) and p-methoxybenzyloxycarbonyl (Moz).

20 The compound of formula (I) can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.

The term "asymmetric carbon atom" means a carbon atom with four different
25 substituents. According to the Cahn-Ingold-Prelog Convention an asymmetric carbon atom can be of the "R" or "S" configuration.

The invention relates in particular to the following:

A compound of formula (I) wherein A^1 is $-S(O)_2-$;

A compound of formula (I) wherein A^2 is $-(CH)-$;

30 A compound of formula (I) wherein R^1 is chloro, bromo, iodo or trifluoromethyl;

A compound of formula (I) wherein R^2 is halogen;

A compound of formula (I) wherein R^2 is chloro or fluoro;

A compound of formula (I) wherein R³ is hydrogen, halogen or haloalkyl;

A compound of formula (I) wherein R³ is halogen;

A compound of formula (I) wherein R³ is chloro;

5 A compound of formula (I) wherein R⁴ and R⁶ are independently selected from hydrogen and haloalkyl;

A compound of formula (I) wherein R⁴ and R⁶ are independently selected from hydrogen and trifluoromethyl;

A compound of formula (I) wherein R⁵ is hydrogen, alkoxy, haloalkoxy, halogen, alkylpyridinyl or alkylpyrazolyl; and

10 A compound of formula (I) wherein R⁵ is hydrogen, methoxy, trifluoroethoxy, fluoro, trifluoropropoxy, bromo, methylpyridinyl or methylpyrazolyl.

The invention further relates to a compound of formula (I) selected from:

(2S,4R)-4-(2-Chloro-4-methoxy-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

15 (2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

20 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

25 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 5 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-fluorophenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 10 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 15 (2S,4S)-4-(2-Chloro-4-fluoro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-5-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 25 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-phenylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-((S)-1,1,1-trifluoropropan-2-yloxy)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

(2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-(2-methylpyridin-4-yl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

(2S,4R)-4-[2-Chloro-4-(2-chloro-pyridin-4-yl)-benzenesulfonyl]-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

5 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

10 (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-
15 amide;

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
20

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

25 (2S,4R)-4-(2-Fluoro-benzenesulfonyl)-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-fluoro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
30

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- 5 ((2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide; and

- 10 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide.

The invention relates in particular to a compound of formula (I) selected from:

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- 15 (2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- 20 (2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-fluorophenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

- 25 (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-((S)-1,1,1-trifluoropropan-2-yloxy)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

(2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-(2-methylpyridin-4-yl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;

- 5 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 10

(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 15

(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20

- (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide; and
- 25

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide.

The compound of formula (I) can be prepared using procedures known in the art.

- 30 The compound of formula (I) can also be prepared using the following procedures.

The following abbreviations are used in the present specification.

- AcOEt: Ethyl acetate;
- ACN: Acetonitrile;
- boc: *tert*-Butyloxycarbonyl;
- BOP: Benzotriazolyl-N-oxy-tris(dimethylamino)-phosphonium hexafluorophosphate;
- 5 BOP-Cl: Bis-(2-oxo-3-oxazolidinyl)-phosphinic acid chloride;
- Cbz: Carbobenzyloxy;
- CDI: 1,1'-Carbonyldiimidazole;
- DCM : Dichloromethane;
- DIEA: Diisopropyl ethyl amine;
- 10 DMAP: 4-Dimethylaminopyridine;
- DMF: N,N-Dimethylformamide;
- EDCI: N-(3-Dimethylaminopropyl)-N'-ethyl-carbodiimide hydrochloride;
- EtOAc: Ethyl acetate;
- Fmoc: 9-Fluorenylmethyloxycarbonyl;
- 15 h: hour;
- HATU: O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate;
- HOBT: 1-Hydroxybenzotriazole;
- Hunig's Base: Ethyl-diisopropyl-amine;
- KHMDS: Potassium bis(trimethylsilyl)amide;
- 20 LDA: Lithiumdiisopropylamide;
- LHMDS: Lithium bis(trimethylsilyl)amide;
- mCPBA or MCPBA: meta-Chloroperoxybenzoic acid;
- MeOH: Methanol;

Mes-Cl: Mesityl chloride;

min: minute;

Moz: Methoxybenzyl carbonyl;

Na₂SO₄: Sodium sulfate;

5 Nos-Cl: 3-Nitrobenzenesulfonyl chloride;

Pd₂(dba)₃: Tris(dibenzylideneacetone)dipalladium;

PyBOP: Benzotriazol-1-yl-oxytripyrrolidinephosphonium hexafluorophosphate;

TBTU: O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate;


Teoc: Trimethylsilyl ethoxycarbonyl;

10 THF: Tetrahydrofurane;

TFA: Trifluoroacetic acid; and

Tos-Cl: Toluene-4-sulfonyl chloride.


Scheme 1



15 R¹-R² are as defined above; Base 1 is e.g. NaOtBu, KOtBu, NaH, LiHMDS, KHMDS or LDA; Base 2 is e.g. LiOH, NaOH or KOH.

A pyridine derivative such as **1** is treated with cyclopropanecarbonitrile **2** in the presence of a base (Base 1 as defined above) to yield the pyridine derivative **3**. Compound **3** is treated with a base (Base 2 as defined above) to yield the final carboxylic acid
 20 derivative **4** as free acid or as a salt thereof.


Scheme 2



LG is a leaving group such as triflate, mesylate, tosylate, brosylate or nosylate; A^2 and R^3 - R^6 are as defined above; R is e.g. Methyl, Ethyl, iPropyl or Benzyl.

- A Boc-protected proline derivative **5** is reacted with a phenylthiol derivative in the presence of a base such as triethyl amine, DIEA, 2,6-lutidine, etc. to yield the thioether derivative **6**. Oxidation of **6** with a peroxide reagent such as H_2O_2 , oxone, mCPBA yields the sulfone derivative **7**. Saponification of the ester to the acid with a base such as LiOH, NaOH or KOH yields the corresponding carboxylic acid **8** or salts thereof. Amide coupling is accomplished by reaction of **8** with 1-aminocarbonitrile derivative and a coupling reagent, such as EDCI, CDI, BOP-Cl, TBTU, HATU, PyBOP or BOP, in the presence of a base, such as DIEA, triethyl amine or lutidine, to yield amide **9**. Finally, the Boc-protecting group is removed by treating compound **9** with an acid such as TFA, HCl in an organic solvent (e.g. AcOEt, dioxane) or formic acid to yield amine **10**.


Scheme 3



A^2 and R^1 - R^6 are as defined above.

Carboxylic acid **4** is reacted with amine **10** in the presence of one of the amide coupling reagents, such as EDCI, CDI, BOP-Cl, TBTU, HATU, PyBOP or BOP, in the presence of a base such as DIEA, triethyl amine or 2,6-lutidine, to yield amide **11**.


Scheme 4



R^1 - R^4 and R^6 are as defined above; LG is a leaving group such as Cl, Br, I; R^5 is phenyl, substituted phenyl, heterocyclyl or substituted heterocyclyl as defined above; R is H or methyl, or both R together with the boron atom to which they are attached form 2,4,4,5,5-pentamethyl-[1,3,2]dioxaborolane.

Compound **12** is reacted with a boronic acid or ester derivative **13** in the presence of a base such as Na_2CO_3 , K_2CO_3 , Cs_2CO_3 , KOtBu , K_3PO_4 , and a catalyst known in the art for performing Suzuki reactions such as e.g. $\text{Pd}(\text{PPh}_3)_4$, $\text{Pd}_2(\text{dba})_3$ or a Pd-source with a phosphine ligand, to yield the biaryl derivative **11**.

Scheme 5




LG is a leaving group such as triflate, mesylate, tosylate, brosylate or nosylate; A² and R¹-R⁶ are as defined above; X is F, Cl, Br, I or X = O-LG.

- 5 The aminolactone **14** or a corresponding salt thereof such as hydrochloride, hydrobromide, phosphate, hydrogenphosphate, sulfate, hydrogensulfate, methansulfonate etc. is reacted with carboxylic acid **4** in the presence of an amide coupling reagent, such as EDCI, CDI, BOP-Cl, TBTU, HATU, PyBOP or BOP, in the presence of a base such as DIEA, triethyl amine, 2,6-lutidine, or alternatively, in the presence of an acid halogenide

such as phosgene, triphosgene, oxalylchloride or thionylchloride, to yield amide **15**.

Opening of the lactone **15** by an amine is performed in the presence of an appropriate base such as sodium 2-ethylhexanoate, TEA, DIEA, DMAP, 2,6-lutidine or pyridine to yield the alcohol **16**. Compound **16** is treated with X-LG in the presence of a base such as TEA, DIEA, DMAP, 2,6-lutidine or pyridine to yield the intermediate **17** which is subsequently reacted with thiols to yield the thioether **18**. Oxidation of thioether **18** to the sulfone **19** is achieved by the reaction of **18** with oxidizing reagents such as H₂O₂, oxone, MCPBA.

Scheme 6




- 10 **R³** is a leaving group such as F, Cl, or S(O)₂-Me; **X** is either N or CH; Base is an inorganic base such as Na₂CO₃, K₂CO₃, Cs₂CO₃ or an organic base such as DIEA, triethylamine or 2,6-lutidine.

- 15 Compound **20** is dissolved in an appropriate solvent such as DMF, DMA or THF, a base as defined above and the nitrogen containing 5-membered heterocycle is added to the reaction mixture. The mixture is initially stirred at room temperature and subsequently heated to an elevated temperature from 30 - 100 °C until the reaction is completed.


The invention also relates to a process for the preparation of a compound of formula (I) as defined above, comprising one of the following steps:

- (a) The reaction of a compound of formula (A)




in the presence of acid, wherein A^1 , A^2 and R^1 to R^6 are as defined above and wherein PG is an amine protecting group;

(b) The reaction of a compound of formula (B1)




5 with a compound of formula (B2)



in the presence of a base and an amide coupling agent and a base, wherein A^1 , A^2 and R^1 to R^6 are as defined above;


(c) The reaction of a compound of formula (C)



10 in the presence of $R^5B(OR)_2$, a base and a Suzuki catalyst, wherein wherein A^1 , A^2 and R^1 to R^4 and R^6 are as defined above, LG is a leaving group, R^5 is alkylpyridinyl, halopyridinyl or alkylpyrazolyl and R is hydrogen or methyl, or both

R, together with the boron atom to which they are attached, form 2,4,4,5,5-pentamethyl-[1,3,2]dioxaborolane; or

(d) The reaction of a compound of formula (D)



5 in the presence of an oxidizing agent, wherein wherein A¹ and R¹ to R⁶ are as defined above.

In step (a), the acid is for example TFA, HCl or formic acid.

In step (a), the amine protecting group is for example boc, Fmoc, Cbz, Teoc, benzyl or Moz.

10 In step (b), the amide coupling agent is for example EDCI, CDI, BOP-Cl, TBTU, HATU, PyBOP or BOP.

In step (b), the base is for example DIEA, triethyl amine or 2,6-lutidine.

In step (c), the leaving group is for example Cl, Br or I.

In step (c), the base is for example Na₂CO₃, K₂CO₃, Cs₂CO₃, KOtBu or K₃PO₄.

15 In step (c), the Suzuki catalyst is for example Pd(PPh₃)₄, Pd₂(dba)₃ or a Pd-source with a phosphine ligand.

In step (d), the oxidizing agent is for example H₂O₂, oxone or MCPBA.

A compound of formula (I), when manufactured according to the above process is also an object of the invention.

20 The compounds of formula (I) and their pharmaceutically acceptable salts can be used as medicaments (e.g. in the form of pharmaceutical preparations). The

pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also be effected parentally, such as
5 intramuscularly or intravenously (e.g. in the form of injection solutions).

The compounds of formula (I) and their pharmaceutically acceptable salts can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatin capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such
10 adjuvants for tablets, dragées and hard gelatin capsules.

Suitable adjuvants for soft gelatin capsules are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols.

Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar and glucose.

15 Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils.

Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid and liquid polyols.

Moreover, the pharmaceutical preparations can contain preservatives, solubilizers,
20 viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

The invention thus also relates in particular to the following:

A compound of formula (I) for use as therapeutically active substance;

25 A pharmaceutical composition comprising a compound of formula (I) and a therapeutically inert carrier;

The use of a compound of formula (I) for the preparation of a medicament for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction of cardiovascular events in chronic kidney
30 disease, diabetic nephropathy, diabetic retinopathy or age related macular degeneration;


A compound of formula (I) for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction of cardiovascular events in chronic kidney disease, diabetic nephropathy, diabetic rethinopathy or age related macular degeneration; and

- 5 A method for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease, cancer, reduction of cardiovascular events in chronic kidney disease, diabetic nephropathy, diabetic rethinopathy or age related macular degeneration, which method comprises administering an effective amount of a compound of formula (I).

- 10 The invention will be illustrated by the following examples which have no limiting character.


Examples**Example 1**

(2S,4R)-4-(2-Chloro-4-methoxy-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



5

a) 1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonitrile




To a solution of 5-chloro-2-fluoropyridine (2 g, 1.53 ml, 15.2 mmol, Eq: 1.00) and cyclopropanecarbonitrile (1.02 g, 1.15 ml, 15.2 mmol, Eq: 1.00) in toluene (20.0 ml) was added dropwise over 5 min. KHMDS 0.5 M in toluene (30.4 ml, 15.2 mmol, Eq: 1.00) at 0°C. The solution turned brown. After 45 min, the reaction mixture was allowed to warm up to 22 °C and stirred for 2.5 h. Saturated aqueous NH₄Cl solution (50 ml) was then added and the aqueous phase was extracted with AcOEt (3 x 60 ml). The combined organic phases were dried over Na₂SO₄, filtered and concentrated under reduced pressure.

The crude material was purified by flash chromatography (silica gel, 70 g, 0% to 20% EtOAc in heptane) to yield the title compound as a white solid (840 mg; 31 %). m/z = 179.0373 [M+H]⁺.

10

15

b) 1-(5-Chloro-pyridin-2-yl)-cyclopropanecarboxylic acid




Compound 1a) (600 mg, 3.36 mmol, Eq: 1.00) was dissolved in 1 % aqueous KOH solution (18 ml, 207 mg, 3.7 mmol, Eq: 1.1) . The reaction mixture was stirred 17 h at

20


100 °C. The crude reaction mixture was concentrated in vacuo and was acidified to pH 4. The crude material was purified by preparative HPLC to yield the title compound as a white solid (339 mg; 51 %). $m/z = 198.1$ $[M+H]^+$.

c) (2*S*,4*R*)-4-(2-Chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide HCl-salt



CAS 1252640-17-7 (600 mg, 1.24 mmol, Eq: 1.00) was dissolved in HCl/dioxane (1.55 ml, 6.2 mmol, Eq: 5.00) and stirred at 22 °C for 4 h. The crude reaction mixture was concentrated in vacuo to yield a white solid (309 mg; 65 %) which was used without further purification. $m/z = 384.2$ $[M+H]^+$.


d) (2*S*,4*R*)-4-(2-Chloro-4-methoxy-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 1b) (61.8 mg, 313 μ mol, Eq: 1.20) was dissolved in DMF (2 ml). HATU (198 mg, 521 μ mol, Eq: 2.00), DIEA (67.3 mg, 91.0 μ l, 521 μ mol, Eq: 2.00) and example 1c) (100 mg, 261 μ mol, Eq: 1.00) were added to the solution and stirred at 22 °C for 15 h. The crude material was purified by preparative HPLC to yield the title compound as a white solid (106 mg; 72 %). $m/z = 563.2$ $[M+H]^+$.

Example 2


20 (2*S*,4*R*)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 2 was prepared in analogy to example 1 starting from CAS 1252638-10-0 to yield the title compound as a white solid (106 mg; 72 %). $m/z = 533.2$ $[M+H]^+$.

Example 3

- 5 **(2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**




- 10 Example 3 was prepared in analogy to example 1 starting from CAS 1252634-04-0 to yield the title compound as a white solid (50 mg; 36 %). $m/z = 631.1$ $[M+H]^+$.

Example 4

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

- 26 -




Example 4 was prepared in analogy to the methods described in example 1 starting from 5-chloro-2,3-difluoropyridine and example 1c) to yield the title compound as a white solid (45 mg; 30 %). $m/z = 581.1$ $[M+H]^+$.

5

Example 5

(2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 5 was prepared in analogy to the methods described in example 1 starting from 5-chloro-2,3-difluoropyridine and CAS 1252638-10-0 to yield the title compound as a white solid (99 mg; 64 %). $m/z = 551.1$ $[M+H]^+$.


10

Example 6

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

15

- 27 -




Example 6 was prepared in analogy to the methods described in example 1 starting from 5-chloro-2,3-difluoropyridine and CAS 1252634-04-0 to yield the title compound as a white solid (114 mg; 79 %). $m/z = 649.2$ $[M+H]^+$.

5

Example 7

(2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 7 was prepared in analogy to the methods described in example 1 starting from 5-bromo-2,3-difluoropyridine and example 1c) to yield the title compound as a white solid (14 mg; 17 %). $m/z = 627.0$ $[M+H]^+$.

Example 8

(2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

- 28 -




Example 8 was prepared in analogy to the methods described in example 1 starting from 5-bromo-2,3-difluoropyridine and CAS 1252638-10-0 to yield the title compound as a white foam (48 mg; 57 %). $m/z = 597.0$ $[M+H]^+$.


5

Example 9

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




10 *a) (1S,4S)-5-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-2-oxa-5-aza-bicyclo[2.2.1]heptan-3-one*



To a milky suspension of 5-chloro-3-fluoro-pyridine-2-carboxylic acid prepared in analogy of example 1b) (670 mg, 3.11 mmol, Eq: 1.00) in toluene (6 ml) at 25 °C was

added DMF (11.4 mg, 12.0 μ l, 155 μ mol, Eq: 0.05). The mixture was cooled down to 0 °C, then a solution of oxalyl chloride (434 mg, 299 μ l, 3.42 mmol, Eq: 1.10) in toluene (2.00 ml) was dropped in within 10 min. The reaction mixture was stirred at 0 °C for 30 min, then without cooling for 3 h. At 0 °C, (1*S*,4*S*)-2-oxa-5-aza-bicyclo[2.2.1]heptan-3-one
 5 methanesulfonate (CAS 769167-53-5) (650 mg, 3.11 mmol, Eq: 1.00) and THF (4.00 ml) were added to the reaction mixture, followed by TEA (1.18 g, 1.62 ml, 11.7 mmol, Eq: 3.75), dropped within 10 min (exothermic). The mixture was stirred at 22 °C for 16 h. The reaction mixture was poured into 20% aqueous citric acid solution (25 ml) and extracted with EtOAc (3 x 20 ml). The organic layers were dried over Na₂SO₄ and concentrated in
 10 vacuo. The crude material was purified by flash chromatography (silica gel, 40 g, 0% to 50% EtOAc in heptane) to yield the title compound as an orange oil (850 mg; 88 %).
 m/z = 311.1 [M+H]⁺.


*b) (2*S*,4*S*)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-hydroxy-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide*



15


A mixture of example 9a) (850 mg, 2.74 mmol, Eq: 1.00), 1-aminocyclopropane-carbonitrile hydrochloride (422 mg, 3.56 mmol, Eq: 1.30), sodium 2-ethylhexanoate (705 mg, 4.24 mmol, Eq: 1.55) in water (3 ml) and THF (2.00 ml) was stirred at 55 °C for 18 h. To the reaction mixture were added hydrochloric acid (189 mg, 157 μ l, 1.91 mmol, Eq: 0.70) and sodium chloride (1.36 g, 1.36 ml, 23.3 mmol, Eq: 8.50). The mixture was stirred
 20 for 15 min, then poured into AcOEt (25 ml) and extracted. The aqueous layer was back-extracted with AcOEt (3 x 20 ml). The organic layers were dried over Na₂SO₄ and concentrated in vacuo. The crude material was purified by flash chromatography (silica gel, 40 g, 0% to 90% EtOAc in heptane) to yield the title compound as white foam (560 mg;
 25 52 %). m/z = 393.0 [M+H]⁺.

*c) 3-Nitro-benzenesulfonic acid (3*S*,5*S*)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-5-(1-cyano-cyclopropylcarbamoyl)-pyrrolidin-3-yl ester*



Example 9b) (560 mg, 1.43 mmol, Eq: 1.00) was dissolved in DCM (15 ml) and 3-nitrobenzene-1-sulfonyl chloride (335 mg, 1.51 mmol, Eq: 1.06) was added. The mixture was cooled down to 0 °C and TEA (433 mg, 596 µl, 4.28 mmol, Eq: 3.00) was slowly and
 5 carefully added with a syringe. The icebath was removed and the reaction mixture was stirred at 25 °C for 18 h. The reaction mixture was extracted with aqueous 10 % Na₂CO₃ and 0.1 N aqueous HCl solutions. The organic layers were dried over Na₂SO₄, filtered and evaporated. The crude material was purified by flash chromatography (silica gel, 40 g, 0% to 85% EtOAc in heptane) to yield the title compound as off-white solid (510 mg; 62 %).
 10 m/z = 578.0 [M+H]⁺.


d) (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 9c) (70 mg, 121 µmol, Eq: 1.00) and 4'-fluorobiphenyl-3-thiol (27.2 mg, 133
 15 µmol, Eq: 1.10) were dissolved in propionitrile (1 ml). TEA (30.6 mg, 42.2 µl, 303 µmol, Eq: 2.50) was added and the reaction mixture was stirred at 90 °C for 3 h. The reaction mixture was poured into 0.1 M aqueous HCl solution (10 ml) and extracted with EtOAc (3 x 10 ml). The organic layers were combined, dried over Na₂SO₄ and concentrated in vacuo. The crude material was purified by flash chromatography (silica gel, 10 g, 0 % to 66 %
 20 EtOAc in heptane) to yield the title compound as off-white oil (46 mg; 68%). m/z = 579.1 [M+H]⁺.

Example 10


(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



- 5 Example 10 was prepared in analogy to the methods described in example 9 starting from 3-chloropyridine-2-thiol and example 9c) to yield the title compound as a light yellow oil (50 mg; 79 %). $m/z = 522.0$ $[M+H]^+$.

Example 11

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-fluorophenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide




Example 11 was prepared in analogy to the methods described in example 1 starting from 5-chloro-2,3-difluoropyridine and CAS 1252633-65-0 to yield the title compound as a white solid (46 mg; 30 %). $m/z = 569.0632$ $[M+H]^+$.

15

Example 12

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

- 32 -




Example 12 was prepared in analogy to the methods described in example 9 starting from pyridine-2-thiol and example 9c) to yield the title compound as a colorless oil (4 mg; 7 %). $m/z = 486.1 [M+H]^+$.

5

Example 13

(2S,4R)-1-[1-(5-Chloro-3-fluoropyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluorobiphenyl-3-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 9 (41 mg, 70.8 μmol , Eq: 1.00) was dissolved in DCM (1 ml) and mCPBA (25.7 mg, 149 μmol , Eq: 2.10) was added. The reaction mixture was stirred for 3 h at 22 °C. The reaction mixture was poured into 10% aqueous Na_2CO_3 (5 ml) solution and extracted with DCM (3 x 5 ml). The organic layers were dried over Na_2SO_4 , filtered and concentrated in vacuo to yield the title compound as a white solid (42 mg; 97 %). $m/z = 611.0 [M+H]^+$.

Example 14

(2S,4R)-1-[1-(5-Chloro-3-fluoropyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloropyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

- 33 -




Example 14 was prepared in analogy to the methods described in example 13 starting from example 10 to yield the title compound as a white solid (44 mg; 99 %). $m/z = 552.1$ $[M+H]^+$.

5

Example 15

(2S,4S)-4-(2-Chloro-4-fluorobenzenesulfonyl)-1-[1-(5-chloro-3-fluoropyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 15 was obtained as a by-product during the synthesis of example 11 as light yellow solid (36 mg; 21 %).

10

Example 16

(2S,4R)-1-[1-(5-Chloro-3-fluoropyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-5-trifluoromethylpyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 16 was prepared in analogy to the methods described in example 9 starting from 3-chloro-5-(trifluoromethyl)pyridine-2-thiol and example 9c) to yield the title compound as a light yellow oil (4 mg; 5 %). $m/z = 585.9$ $[M+H]^+$.

5

Example 17

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 17 was prepared in analogy to the methods described in example 9 starting from 5-chloropyridine-2-thiol and example 9c) to yield the title compound as an off-white solid (30 mg; 83 %). $m/z = 522.0$ $[M+H]^+$.

10

Example 18

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

- 35 -




Example 18 was prepared in analogy to the methods described in example 13 starting from example 10 to yield the title compound as a off-white solid (3 mg; 94 %). $m/z = 518.1$ $[M+H]^+$.

5

Example 19

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 19 was prepared in analogy to the methods described in example 9 starting from 6-methylpyridine-2-thiol and example 9c) to yield the title compound as an off-white solid (17 mg; 49 %). $m/z = 500.1$ $[M+H]^+$.

Example 20

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

15

- 36 -




Example 20 was prepared in analogy to the methods described in example 9 starting from 5-(trifluoromethyl)pyridine-2-thiol and example 9c) to yield the title compound as a light yellow solid (32 mg; 84 %). $m/z = 554.1$ $[M+H]^+$.

5

Example 21

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridin-2-ylsulfanylmethyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




10 Example 21 was prepared in analogy to the methods described in example 9 starting from 3-(trifluoromethyl)pyridine-2-thiol and example 9c) to yield the title compound as a yellow solid (33 mg; 86 %). $m/z = 554.1$ $[M+H]^+$.

Example 22

15 **(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**

- 37 -




Example 22 was prepared in analogy to the methods described in example 13 starting from example 17 to yield the title compound as a white foam (22 mg; 80 %). $m/z = 552.1$ $[M+H]^+$.

5

Example 23

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 23 was prepared in analogy to the methods described in example 13 starting from example 19 to yield the title compound as a white foam (13 mg; 82 %). $m/z = 532.0$ $[M+H]^+$.

10

Example 24

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

15




Example 24 was prepared in analogy to the methods described in example 13 starting from example 20 to yield the title compound as a white solid (29 mg; 98 %). $m/z = 585.9$ $[M+H]^+$.

5

Example 25


(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



10 Example 25 was prepared in analogy to the methods described in example 13 starting from example 21 to yield the title compound as a white solid (16 mg; 52 %). $m/z = 585.9$ $[M+H]^+$.

Example 26

15 **(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-phenylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**




Example 26 was prepared in analogy to the methods described in example 9 starting from 2-chloro-5-(trifluoromethyl)benzenethiol and example 9c) to yield the title compound as a white solid (10 mg; 25 %). $m/z = 586.9$ $[M+H]^+$.

5

Example 27

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-((S)-1,1,1-trifluoropropan-2-yloxy)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide




- 10 Example 11 (27 mg, 47.4 μmol , Eq: 1.00) was dissolved in DMF (1 ml). Cs_2CO_3 (23.2 mg, 71.1 μmol , Eq: 1.50) and (S)-1,1,1-trifluoropropan-2-ol (5.95 mg, 52.2 μmol , Eq: 1.10) were added to the solution and stirred at 40 °C for 4 h. The crude material was purified by preparative HPLC to yield the title compound as a white solid (17 mg; 54 %). $m/z = 663.2$ $[M+H]^+$.


15

Example 28


(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide



a) (2*S*,4*R*)-4-(2-Chloro-5-trifluoromethyl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



- 5 Example 28a) was prepared in analogy to the methods described for CAS 1252638-10-0 (see Haap et al.; US20100267722 and Hardegger et al.; Angewandte Chemie, International Edition, 50(1), 314-318, S314/1-S314/145; 2011) starting from 2-chloro-5-trifluoromethyl-benzenethiol to yield the title compound as a light yellow solid (125 mg; 74 %) $m/z = 421.9 [M+H]^+$.
- 10 b) (2*S*,4*R*)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide




Example 28b) was prepared in analogy to example 1 starting from example 28a) and 5-chloro-2,3-difluoropyridine to yield the title compound as a white solid (37 mg; 50 %).

- 15 $m/z = 619.1 [M+H]^+$.

Example 29

(2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




5

Example 29 was prepared in analogy to example 1 starting from example 28a) and example 1b) to yield the title compound as a white solid (37 mg; 50 %). $m/z = 601.1$ $[M+H]^+$.

Example 30


10 **(2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**



15 Example 30 was prepared in analogy to example 28 starting from 4-bromo-2-chlorobenzenethiol to yield the title compound as a white foam (860 mg; 79 %) $m/z = 631.0$ $[M+H]^+$.

Example 31

(2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-(2-methylpyridin-4-yl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide




- 5 Example 30 (100 mg, 159 μ mol, Eq: 1.00) was dissolved in 1,2-dimethoxyethane (2 ml). 2-Methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (45.2 mg, 206 μ mol, Eq: 1.30), triphenylphosphine (8.32 mg, 31.7 μ mol, Eq: 0.20), 2 M aqueous Na_2CO_3 solution (500 μ l) and $\text{Pd}(\text{OAc})_2$ (3.56 mg, 15.9 μ mol, Eq: 0.10) were added and stirred at 45 $^\circ\text{C}$ for 4 h. The reaction mixture was poured into 0.1 M aqueous HCl solution (10 ml) and
- 10 extracted with DCM (3 x 10 ml). The organic layers were dried over Na_2SO_4 and concentrated in vacuo. The crude material was purified by preparative HPLC to yield the title compound as an off-white foam (61 mg; 60 %). $m/z = 642.1$ $[\text{M}+\text{H}]^+$.

Example 32

- (2S,4R)-4-[2-Chloro-4-(2-chloro-pyridin-4-yl)-benzenesulfonyl]-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**
- 15

- 43 -




Example 32 was prepared in analogy to example 31 starting from example 30 and 2-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine to yield the title compound as a white solid (27 mg; 26 %) $m/z = 664.1$ $[M+H]^+$.

5

Example 33


(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



10 Example 33 was prepared in analogy to example 31 starting from example 30 and 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole to yield the title compound as a white solid (25 mg; 25 %). $m/z = 631.1$ $[M+H]^+$.

Example 34


(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



- 5 Example 34 was prepared in analogy to example 31 starting from example 30 and 2-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine to yield the title compound as a white foam (47 mg; 38 %) $m/z = 642.2$ $[M+H]^+$.


Example 35

- 10 **(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide**




a) (2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-2-(1-cyano-cyclopropylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester

- 45 -



Example 35a) was prepared in analogy to the methods described for CAS 1252631-66-5 (see Haap et al.; US20100267722) starting from (2*S*,4*R*)-1,2-pyrrolidinedicarboxylic acid 4-hydroxy- 1-(1,1-dimethylethyl) ester and 4-bromo-2-chloro-benzenethiol to yield the
5 title compound as a white solid (3.2 g; 58 %). $m/z = 434.1$ $[M+H-Boc]^+$.


b) (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-2-(1-cyano-cyclopropylcarbamoyl)-pyrrolidine-1-carboxylic acid tert-butyl ester



Example 35a) (700 mg, 1.31 mmol, Eq: 1.00) was dissolved in 1,2-dimethoxyethane (8
10 ml). 2-Methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (345 mg, 1.58 mmol, Eq: 1.20), triphenylphosphine (68.9 mg, 263 μ mol, Eq: 0.20), 2 M aqueous Na_2CO_3 solution (2 ml) and palladium (II) acetate (29.5 mg, 131 μ mol, Eq: 0.10) were added and stirred at 22 °C for 24 h. After that, the reaction mixture was stirred at 50 °C for 24 h. Then, 2-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (57.6 mg, 263
15 μ mol, Eq: 0.2) was added to the reaction mixture which was then stirred at 60 °C for 6 h. The reaction mixture was poured into 0.1 M aqueous HCl solution (50 ml) and extracted with DCM (3 x 20 ml). The organic layers were dried over Na_2SO_4 and concentrated in vacuo. The crude material was purified by flash chromatography (silica gel, 40 g, 0% to 100% EtOAc in heptane) to yield the title compound as a light yellow oil (200 mg; 28%).
20 $m/z = 545.3$ $[M+H]^+$.

c) (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide


- 46 -



Example 35b) (200 mg, 367 μmol , Eq: 1.00) was dissolved in formic acid (2.4 g, 2 ml, 52.1 mmol, Eq: 142) and stirred at 22 °C for 15 h. The reaction mixture was adjusted carefully with icecold aqueous 10 % Na_2CO_3 -solution to pH 8 and extracted with CH_2Cl_2 .


- 5 The water layer was washed totally 3 times with $\text{CH}_2\text{Cl}_2/\text{THF}$ (1:1; 30 ml), the combined organic layers were dried over Na_2SO_4 , filtered and evaporated to yield the title compound as a white foam (142 mg; 87 %). $m/z = 445.2$ $[\text{M}+\text{H}]^+$.

d) 1-(3-Fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarboxylic acid



- 10 Example 35 d) was prepared in analogy to the methods described for examples 1a) and b) to yield the title compound as a light brown solid (50 mg; 41 %) $m/z = 250.0$ $[\text{M}+\text{H}]^+$.


e) (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



The title compound was prepared in analogy to example 1d) starting from example 35c) (45 mg) and 35d) (30 mg) to yield an off-white solid (34 mg 50 %) $m/z = 676.3 [M+H]^+$.

Example 36


(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 36 was prepared in analogy to example 35 starting from example 35c) and 1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarboxylic acid, which was prepared in analogy to examples 1a) and 1b), to yield the title compound as a off-white solid (35 mg; 47 %) $m/z = 734.2 [M+H]^+$.

Example 37

(2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




Example 37 was prepared in analogy to example 35 starting from example 35c) and 1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarboxylic acid, which was prepared in analogy to examples 1a) and 1b), to yield the title compound as an off-white solid (26 mg; 39 %) $m/z = 660.2$ $[M+H]^+$.


5

Example 38

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




- 10 a) (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



- 15 Example 38a) was prepared in analogy to example 35c) starting from examples 35a) and 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole to yield the title compound as a light brown foam (173 mg; 95 %). $m/z = 434.2$ $[M+H]^+$.


b) (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



Example 38b) was prepared in analogy to example 35 starting from example 38a) and 1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarboxylic acid, which was prepared in analogy to examples 1a) and 1b), to yield the title compound as a off-white solid (40 mg; 48 %) $m/z = 723.1$ $[M+H]^+$.

Example 39

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



10


Example 39 was prepared in analogy to example 38b) starting from example 38a) and 35d) to yield the title compound as a off-white solid (16 mg; 60 %). $m/z = 665.1$ $[M+H]^+$.

Example 40

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

15

- 50 -




Example 40 was prepared in analogy to example 35 starting from example 38a) and 1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarboxylic acid, which was prepared in analogy to examples 1a) and 1b), to yield the title compound as an off-white solid (29 mg; 39 %)

5 $m/z = 649.2 [M+H]^+$.

Example 41

(2S,4R)-4-(2-Fluoro-benzenesulfonyl)-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




10 Example 41 was prepared in analogy to example 8 starting from 2-fluorobenzenethiol and 1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarboxylic acid to yield the title compound as a white foam (328 mg; 68 %). $m/z = 627.3 [M+H]^+$.

Example 42

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-fluoro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide

15

- 51 -




Example 42 was prepared in analogy to example 41 starting from 2-fluorobenzenethiol and 1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarboxylic acid to yield the title compound as a white foam (212 mg; 67 %). $m/z = 535.4$ $[M+H]^+$.

5

Example 43

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




To a 5 mL tube were added example 41 (60 mg, 95.8 μ mol, Eq: 1.00), 1H-pyrazole (9.78 mg, 144 μ mol, Eq: 1.50), Cs_2CO_3 (37.4 mg, 115 μ mol, Eq: 1.20) and DMF (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was again 1H-pyrazole (9.78 mg, 144 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to yield the title compound as a white foam (8 mg; 12.4%). $m/z = 675.0647$ $[M+H]^+$.

15

Example 44


(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



To a 5 mL tube were added example 41 (60 mg, 95.8 μ mol, Eq: 1.00), 1H-1,2,3-triazole (9.92 mg, 8.32 μ l, 144 μ mol, Eq: 1.50), Cs₂CO₃ (37.4 mg, 115 μ mol, Eq: 1.20) and DMA (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was
 5 again 1H-1,2,3-triazole (9.92 mg, 8.32 μ l, 144 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to yield the title compound as a white foam (25 mg; 23%; purity 50-80%). m/z = 676.0638 [M+H]⁺.

Example 45


(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-
 10 [1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-
 cyclopropyl)-amide



To a 5 mL tube were added example 41 (60 mg, 95.8 μ mol, Eq: 1.00), 1H-1,2,4-triazole (9.92 mg, 144 μ mol, Eq: 1.50), 1H-1,2,4-triazole (9.92 mg, 144 μ mol, Eq: 1.50) and DMA
 15 (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was again 1H-1,2,4-triazole (9.92 mg, 144 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to the title compound as a white foam (15 mg; 18.8%; purity 50-80%). m/z = 676.0629 [M+H]⁺.

Example 46

((2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




To a 5 mL tube were added example 42 (50 mg, 93.5 μ mol, Eq: 1.00), 1H-pyrazole (9.54 mg, 140 μ mol, Eq: 1.50), Cs₂CO₃ (36.5 mg, 112 μ mol, Eq: 1.20) and DMA (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was again 1H-pyrazole (9.54 mg, 140 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to yield the title compound as a white foam (10 mg; 15%; purity 80%). m/z = 583.1319 [M+H]⁺.

10

Example 47

((2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide




To a 5 mL tube were added example 42 (50 mg, 93.5 μ mol, Eq: 1.00), 1H-1,2,3-triazole (9.68 mg, 8.12 μ l, 140 μ mol, Eq: 1.50), Cs₂CO₃ (36.5 mg, 112 μ mol, Eq: 1.20) and DMA (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was again 1H-1,2,3-triazole (9.68 mg, 8.12 μ l, 140 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to yield the title compound as a white foam (10 mg; 11%; purity 60%). m/z = 584.1272 [M+H]⁺.

20

Example 48

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide



5

To a 5 mL tube were added example 42 (50 mg, 93.5 μ mol, Eq: 1.00), 1H-1,2,4-triazole (9.68 mg, 140 μ mol, Eq: 1.50), Cs_2CO_3 (36.5 mg, 112 μ mol, Eq: 1.20) and DMA (1 ml). The reaction mixture was stirred for 24 h at 22 °C. To the reaction mixture was again 1H-1,2,4-triazole (9.68 mg, 140 μ mol, Eq: 1.50) added and stirred for 24 h at 50 °C. The crude material was purified by preparative HPLC to yield the title compound as a white foam (25 mg; 35%; purity 77%). $m/z = 584.1281$ $[\text{M}+\text{H}]^+$.

10

Exemple 49**Cathepsin enzyme inhibition assay**

Enzyme activity is measured by observing the increase in fluorescence intensity caused by cleavage of a peptide substrate containing a fluorophore whose emission is quenched in the intact peptide.

15

Assay buffer: 100 mM potassium phosphate pH 6.5, EDTA-Na 5 mM, Triton X-100 0.001%, DTT 5 mM.

Enzymes (all at 1 nM): human and mouse Cathepsin S, Cat K, Cat B, Cat L.

Substrate (20 μ M): Z-Val-Val-Arg-AMC, except for Cat K which uses Z-Leu-Arg-AMC (both from Bachem).

20

Z = Benzyloxycarbonyl.

AMC = 7-Amino-4-Methyl-Coumarin.

DTT = dithiothreitol.

Final volume: 100 μ L.

Excitation 360 nm, Emission 465 nm.

Enzyme is added to the substance dilutions in 96-well microtitre plates and the reaction is
 5 started with substrate. Fluorescence emission is measured over 20 minutes, during which
 time a linear increase is observed in the absence of inhibitor. IC₅₀ are calculated by
 standard methods.

Inhibition of human Cat S, mouse Cat S, human Cat K, human Cat B, human Cat L and
 mouse Cat L have been measured separately. The results obtained for human Cat S and L
 10 for representative compounds of the invention are expressed in the following table in μ M.

Example	IC50 h S	IC50 h L	Example	IC50 h S	IC50 h L
1	0.000606	0.4097	25	0.003213	13.1975
2	0.000492	0.3402	26	0.003466	0.0524
3	0.000825	0.429	27	0.001596	0.09
4	0.000393	0.0119	28	0.000543	0.0568
5	0.000398	0.0129	29	0.000618	0.3618
6	0.000675	0.0445	30	0.000474	0.0726
7	0.000734	0.0255	31	0.000505	0.0616
8	0.000438	0.0192	32	0.000686	0.1121
9	0.011785	0.4165	33	0.000375	0.0287
10	0.96365	15.8722	34	0.000322	0.0642

11	0.000602	0.0898	35	0.000768	0.0589
12	0.5941	8.2265	36	0.000382	0.0098
13	0.00077	0.2474	37	0.000779	0.0722
14	0.002844	2.117	38	0.000678	0.0032
15	0.011075	1.679	39	0.000696	0.0436
16	1.031	>25	40	0.000703	0.1111
17	0.2635	5.648	41	0.0006	0.003
18	0.00424	3.4345	42	0.0007	0.031
19	0.12715	4.6095	43	0.016985	0.0906
20	0.3926	5.0635	44	0.002372	1.5585
21	0.27455	10.011	45	0.3272	28.383
22	0.002996	1.9695	46	0.1587	17.445
23	0.002796	5.635	47	0.004268	35.2
24	0.006244	2.694	48	0.058535	21.1025

The compounds of the invention are preferential inhibitors of Cathepsin-S and L over Cathepsin-K and B.

The compounds according to the invention have, in the foregoing assay, an IC₅₀ at Cat S and/or L which is between 0.00001 and 100 µM. Particular compounds of the invention
5 have an IC₅₀ at Cat S and/or L between 0.00001 and 50 µM and mor particularly between

0.00001 and 20 μM . The particular compounds of the invention have an IC_{50} in at least one of the foregoing assay below 0.09 μM .

Example A

A compound of formula (I) can be used in a manner known per se as the active ingredient
5 for the production of tablets of the following composition:

	<u>Per tablet</u>
Active ingredient	200 mg
Microcrystalline cellulose	155 mg
Corn starch	25 mg
10 Talc	25 mg
Hydroxypropylmethylcellulose	<u>20 mg</u>
	425 mg


Example B

15 A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of capsules of the following composition:

	<u>Per capsule</u>
Active ingredient	100.0 mg
Corn starch	20.0 mg
20 Lactose	95.0 mg
Talc	4.5 mg
Magnesium stearate	<u>0.5 mg</u>
	220.0 mg

Claims

1. A compound of formula (I)



wherein

- 5 A^1 is -S- or -S(O)₂-;
- A^2 is nitrogen or -(CH)-;
- R^1 is halogen or haloalkyl;
- R^2 is hydrogen or halogen;
- R^3 is hydrogen, halogen, haloalkyl, pyrazolyl, [1,2,3]-triazolyl or [1,2,4]-triazolyl;
- 10 R^4 and R^6 are independently selected from hydrogen, alkyl, haloalkyl and halophenyl; and
- R^5 is hydrogen, halogen, haloalkyl, alkoxy, haloalkoxy, alkylpyridinyl, halopyridinyl or alkylpyrazolyl;
- or a pharmaceutically acceptable salt thereof.
- 15 2. A compound according to claim 1, wherein A^1 is -S(O)₂-.
3. A compound according to claim 1 or 2, wherein A^2 is -(CH)-.
4. A compound according to any one of claims 1 to 3, wherein R^1 is chloro, bromo, iodo or trifluoromethyl.
5. A compound according to any one of claims 1 to 4, wherein R^2 is halogen.

6. A compound according to any one of claims 1 to 5, wherein R² is chloro or fluoro.
7. A compound according to any one of claims 1 to 6, wherein R³ is halogen.
8. A compound according to any one of claims 1 to 7, wherein R³ is chloro.
9. A compound according to any one of claims 1 to 8, wherein R⁴ and R⁶ are
5 independently selected from hydrogen and haloalkyl.
10. A compound according to any one of claims 1 to 9, wherein R⁴ and R⁶ are independently selected from hydrogen and trifluoromethyl.
11. A compound according to any one of claims 1 to 10, wherein R⁵ is hydrogen, alkoxy, haloalkoxy, halogen, alkylpyridinyl or alkylpyrazolyl.
- 10 12. A compound according to any one of claims 1 to 11, wherein R⁵ is hydrogen, methoxy, trifluoroethoxy, fluoro, trifluoropropoxy, bromo, methylpyridinyl or mehtylpyrazolyl.
13. A compound according to any one of claims 1 to 12 selected from

(2S,4R)-4-(2-Chloro-4-methoxy-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-
15 cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-pyridin-2-yl)-
cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-
trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-
20 cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-
methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-
amide;

(2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-
25 cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-
(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-
cyclopropyl)-amide;

- (2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 5 (2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 10 (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-fluorophenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(4'-fluoro-biphenyl-3-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 15 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4S)-4-(2-Chloro-4-fluoro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-chloro-5-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 25 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 5 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridin-2-ylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-chloro-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 10 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(6-methyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(5-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 15 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(3-trifluoromethyl-pyridine-2-sulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-phenylsulfanyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20 (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-((S)-1,1,1-trifluoropropan-2-yloxy)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-
- 25 carboxamide;
- (2S,4R)-1-[1-(5-Chloro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-5-trifluoromethyl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 30 (2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-(2-methylpyridin-4-yl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- 5 (2S,4R)-4-[2-Chloro-4-(2-chloro-pyridin-4-yl)-benzenesulfonyl]-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 10 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 15 (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20 (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 25 (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 30 (2S,4R)-4-(2-Fluoro-benzenesulfonyl)-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-fluoro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

5 (2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(3-Fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

10

((2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-pyrazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,3]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide; and

15

(2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-[1,2,4]triazol-1-yl-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide.

14. A compound according to any one of claims 1 to 13 selected from

20 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-4-(2-Chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

25 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2,2,2-trifluoro-ethoxy)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

(2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-4-methoxy-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

30

- (2S,4R)-1-[1-(5-Bromo-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-(2-chloro-benzenesulfonyl)-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-fluorophenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- 5 (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-((S)-1,1,1-trifluoropropan-2-yloxy)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-carboxamide;
- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-5-(trifluoromethyl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-
- 10 carboxamide;
- (2S,4R)-4-(4-Bromo-2-chloro-benzenesulfonyl)-1-[1-(5-chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-1-(1-(5-chloro-3-fluoropyridin-2-yl)cyclopropanecarbonyl)-4-(2-chloro-4-(2-methylpyridin-4-yl)phenylsulfonyl)-N-(1-cyanocyclopropyl)pyrrolidine-2-
- 15 carboxamide;
- (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 20 (2S,4R)-1-[1-(5-Chloro-3-fluoro-pyridin-2-yl)-cyclopropanecarbonyl]-4-[2-chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid
- 25 (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- (2S,4R)-4-[2-Chloro-4-(2-methyl-pyridin-3-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;
- 30


(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-iodo-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide;

5 (2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3-fluoro-5-trifluoromethyl-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide; and

(2S,4R)-4-[2-Chloro-4-(1-methyl-1H-pyrazol-4-yl)-benzenesulfonyl]-1-[1-(3,5-dichloro-pyridin-2-yl)-cyclopropanecarbonyl]-pyrrolidine-2-carboxylic acid (1-cyano-cyclopropyl)-amide.


10 15. A process for the preparation of a compound of formula (I) as defined in any one of claims 1 to 14, comprising one of the following steps:

(a) The reaction of a compound of formula (A)




15 in the presence of acid, wherein A¹, A² and R¹ to R⁶ are as defined in any one of claims 1 to 12 and wherein PG is an amine protecting group;

(b) The reaction of a compound of formula (B1)




with a compound of formula (B2)

- 66 -



in the presence of a base and an amide coupling agent and a base, wherein A^1 , A^2 and R^1 to R^6 are as defined in any one of claims 1 to 12;

(c) The reaction of a compound of formula (C)




5

in the presence of $R^5B(OR)_2$, a base and a Suzuki catalyst, wherein wherein A^1 , A^2 and R^1 to R^4 and R^6 are as defined in any one of claims 1 to 12, LG is a leaving group, R^5 is alkylpyridinyl, halopyridinyl or alkylpyrazolyl and R is hydrogen or methyl, or both R, together with the boron atom to which they are attached, form 2,4,4,5,5-pentamethyl-[1,3,2]dioxaborolane; or

10

(d) The reaction of a compound of formula (D)



in the presence of an oxidizing agent, wherein wherein A^1 and R^1 to R^6 are as defined in any one of claims 1 to 12.

16. A compound according to any one of claims 1 to 14, when manufactured according to a process of claim 15.
- 5 17. A compound according to any one of claims 1 to 14 for use as a therapeutically active substance.
18. A pharmaceutical composition comprising a compound in accordance with any one of claims 1 to 14 and a therapeutically inert carrier.
19. The use of a compound according to any one of claims 1 to 14 for the preparation of
10 a medicament for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease or diabetic nephropathy.
20. A compound according to any one of claims 1 to 14 for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease or diabetic nephropathy.
- 15 21. A method for the treatment or prophylaxis of diabetes, atherosclerosis, abdominal aortic aneurysm, peripheral arterial disease or diabetic nephropathy, which method comprises administering an effective amount of a compound as defined in any one of claims 1 to 14.
22. The invention as hereinbefore described.