

TSH receptor antagonizing tetrahydroquinoline compounds

The invention relates to a compound having TSH receptor antagonist activity, in particular a tetrahydroquinoline derivative and the use thereof for the treatment and prevention of TSH receptor responsive disorders and to a pharmaceutical composition and a kit containing the same.

Thyrotropin or thyroid-stimulating hormone (TSH) plays an important role in the regulation of metabolism and development. TSH is released from the anterior pituitary under the influence of thyrotropin-releasing hormone. It targets the thyroid gland to stimulate the release of the thyroid hormones triiodothyronine and thyroxine and thyroid growth. The actions of TSH are mediated by a specific G protein-coupled receptor which couples preferentially to G_s proteins leading to activation of adenylyl cyclase. This signal transduction pathway is predominantly responsible for the production of thyroid hormones and proliferation of the thyrocytes (Krohn et al. (2005) *Endocrine Rev.* **26**, 504-524).

The TSH receptor on the thyroid is also directly involved in the pathogenesis and pathophysiology of Graves' disease. Graves' disease is characterized by hyperstimulation of the thyroid as a result of circulating TSH receptor-stimulating immunoglobulins (TSI), which persistently activate the receptor (Gerding et al. (2000) *Clin. Endocrinol.* **52**, 267-271). TSI may also directly participate in the pathogenesis and pathophysiology of Graves' ophthalmopathy and Graves'-associated pretibial dermopathy as TSH receptors are present in orbital tissue and affected skin regions of these patients, respectively (Gerding et al. (2000) *Clin. Endocrinol.* **52**, 267-271; Daumerie et al. (2002) *Eur. J. Endocrinol.* **146**, 35-38). In addition, the TSH receptor plays a crucial role in toxic and non-toxic nodular goitre. In (toxic) nodular goitre, the thyroid gland contains autonomously functioning thyroid nodules that secrete excess thyroid hormone as a result of mutations in the TSH receptors which render them constitutively active with significantly increased cAMP levels (Krohn et al. (2005) *Endocrine Rev.* **26**, 504-524).

Blocking the TSH receptor or inhibiting the signaling which is induced after TSH-, or TSI-mediated receptor stimulation, or as a result of constitutively active mutations of the TSH receptor will inhibit thyroid hormone secretion and thyrocyte proliferation. Low molecular weight TSH receptor antagonists could therefore be used to treat or prevent hyperthyroidism, Graves' disease, nodular goiter, Graves' ophthalmopathy and Graves'-associated pretibial dermopathy. In addition, low molecular weight TSH receptor antagonists could be used to prevent stimulation of growth of residues or metastases of thyroid cancer, which is thought to be promoted by (over)stimulation of the TSH receptor. In a more general sense, TSH receptor antagonists could be used to prevent or treat all of those ailments in which (over)activation of the TSH receptor plays a role.

Tetrahydroquinoline derivative compounds are described in WO2003/004028, WO2004/056779 and WO2004/056780. These compounds can be used to regulate fertility.


FSH receptor modulators reported in the literature have high specificity towards the FSH receptor. Yanofsky *et al* (2006, J. Biol. Chem. **281**, 13226-13233) and Pelletier *et al* (2005, Bioorg. & Med. Chem. **13**, 5986-5995) demonstrated that low nanomolar potent LMW FSH receptor agonists with a thiazolidinone scaffold are neither TSH receptor antagonists nor agonists. Also (bis)sulfonic acid, (bis)benzamides, have been identified as FSH receptor antagonists, but they show no or little ability to inhibit TSH receptor activity (Wrobel *et al.*, 2002, Biorg. Med. Chem. **10**, 639-656). Another series of low micromolar FSH receptor antagonists (diazonaphthylsulfonic acid derivatives) did not show affinity towards the TSH receptor (Arey *et al.*, 2002, Endocrinology **143**, 3822-3829).

In addition, mutagenesis studies on the FSH receptor and the TSH receptor have demonstrated that activation of the TSH receptor is distinct from that of the FSH receptor indicating that FSH receptor antagonists will not necessarily inhibit TSH receptor as well (Schulz *et al.*, 1999, Mol Endocrinol **13**, 181-190). This differential activation is underscored by the knowledge that the TSH receptor, in sharp contrast to the FSH receptor, also couples efficiently to phospholipase C via Gq proteins, a

pathway which is required for thyroid hormone synthesis (Kero *et al*, 2007, J.Clin.Invest. **117**, 2399-2407) and also activated by TSI in Graves'disease.

The present invention describes tetrahydroquinoline compounds that inhibit TSH receptor activation. The compounds of the invention can be used as (partial) antagonists of the TSH receptor.

It has now been found, that the following class of tetrahydroquinoline compounds of formula I or pharmaceutically acceptable salts thereof, have TSH receptor antagonistic activity:



Formula I

wherein

R¹ is H or methyl;

X is a bond, O, NH or N((1-4C)alkyl);

R² is (1-4C)alkyl, R⁵(1-4C)alkyl or R⁹(2-4C)alkyl,

if X is NH, R² in addition is R⁸(1-2C)alkoxycarbonyl; or,

if X is a bond, R² in addition is H, halogen or (2-5C)heteroaryl or phenyl, both optionally substituted with one or more substituents selected from (1-3C)alkyl, (1-3C)alkoxy or halogen;

Y is a bond, O or NH, whereas

if Y is a bond, R³ is phenyl, optionally one or more substituents selected from halogen, hydroxy, methoxy, phenoxy, phenyl, (1-4C)alkyl, nitro, amino or (di)[(1-4C)alkyl]amino or

R^3 is R^6 (1-6C)alkyl, R^6 (3-6C)cycloalkyl, R^7 oxy(1-6C)alkyl, or 2-pyridyl or
 R^3 is a 5-membered heteroaryl, optionally substituted with one or more substituents
selected from (1-3C)alkoxy, (1-3C)alkyl or halogen;
if Y is O, R^3 is R^6 (1-6C)alkyl, R^7 oxy(2-6C)alkyl or (3-7C)cycloalkyl; and
if Y is NH, R^3 is R^6 (1-6C)alkyl, R^7 oxy(2-6C)alkyl;
 R^4 is H, (di)[(1-3C)alkyl]amino or (1-3C)alkoxy;
 R^5 is CN or pyridyl;
 R^6 is H, or (3-5C)cycloalkyl, (2-5C)heteroaryl or phenyl, the latter three groups
optionally substituted with one or more substituents selected from halogen, (1-
4C)alkoxy or (1-4C)alkyl, the latter two optionally substituted with one or more
halogen;
 R^7 is (2-5C)heteroaryl or phenyl, both optionally substituted with one or more
substituents selected from halogen, (1-4C)alkoxy or (1-4C)alkyl;
 R^8 is H or (2-5C)heteroaryl or phenyl, both optionally substituted with one or more
substituents selected from (1-3C)alkyl, (1-3C)alkoxy or halogen; and
 R^9 is (di)[(1-4C)alkyl]amino or (2-6C)heterocycloalkyl.

In the above Formula 1, the two R^1 groups are always the same and are either H or methyl.

In particular, the compounds of the present invention show antagonistic TSH receptor activity.

Thus, the TSH receptor antagonistic compounds of the present invention may be used to treat a mammal, including a human, with disorders responsive to TSH receptor mediated pathways. They can also be used to prevent such TSH receptor mediated disorders. The TSH receptor antagonistic compounds of the present invention also fully inhibit TSH receptor-mediated phospholipase C activity with equal potency.

The following terms are intended to have the indicated meanings denoted below as used in the specification and claims.

The term (1-3C)alkyl as used in the definition means a branched or unbranched alkyl group having 1-3 carbon atoms, being methyl, ethyl, propyl or isopropyl.

The term (1-4C)alkyl as used in the definition means a branched or unbranched alkyl group having 1-4 carbon atoms, being methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and *tert*-butyl.

The term (2-4C)alkyl as used in the definition means a branched or unbranched alkyl group having 2-4 carbon atoms, being, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl and *tert*-butyl.

The term (1-6C)alkyl as used in the definition means a branched or unbranched alkyl group having 1-6 carbon atoms for example methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl and n-hexyl. (1-5C)Alkyl groups are preferred.

The term (2-6C)alkyl as used in the definition means a branched or unbranched alkyl group having 2-6 carbon atoms such as ethyl, propyl, isopropyl, butyl, pentyl or hexyl.

The term (3-6C)cycloalkyl means a cycloalkyl group having 3-6 carbon atoms, being cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term (3-5C)cycloalkyl means a cycloalkyl group having 3-5 carbon atoms, being cyclopropyl, cyclobutyl and cyclopentyl.

The term (3-7C)cycloalkyl means a mono or bicycloalkyl group having 3-7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and norbornyl.

The term (2-5C)heteroaryl means an aromatic group having 2-5 carbon atoms and at least including one heteroatom selected from N, O and/or S. More than one heteroatom, including different heteroatoms may be included if feasible. Preferred heteroaryl groups are thienyl, furyl, pyridyl, isoxazolyl, pyrimidyl, pyrrolyl. The (2-5C)heteroaryl group may be attached via a carbon atom or a heteroatom, if feasible.

The term 5-membered heteroaryl means an aromatic group having 2-4 carbon atoms and at least including one heteroatom selected from N, O and/or S. More than one heteroatom, including different heteroatoms may be included if feasible. Preferred heteroaryl groups are thienyl, furyl, imidazolyl, isoxazolyl, pirazolyl, pyrrolyl, oxadiazolyl, oxazolyl or thiazolyl. The 5-membered heteroaryl group may be attached via a carbon atom or a heteroatom, if feasible.

The term (1-3C)alkoxy means an alkoxy group having 1-3 carbon atoms, the alkyl moiety having the same meaning as previously defined.

The term (1-4C)alkoxy means an alkoxy group having 1-4 carbon atoms, the alkyl moiety having the same meaning as previously defined. (1-2C)Alkoxy groups are preferred.

The term (1-2)alkoxycarbonyl means carbonyl group with an attached alkoxy group having 1-2 carbon atoms.

The term (2-6C)heterocycloalkyl means a heterocycloalkyl group, having 2-6 carbon atoms and at least including one heteroatom selected from N, O and/or S, which may be attached via a heteroatom, if feasible, or a C atom. More than one heteroatom, including different heteroatoms may be included if feasible. The preferred number of heteroatoms is 1-2. The preferred number of C-atoms is 3-6. Preferred groups are morpholinyl, homomorpholinyl, piperidinyl and homopiperidinyl. Most preferred is morpholinyl.

The term oxy(1-6C)alkyl means an oxyalkyl group having 1-6 carbon atoms, the alkyl moiety having the same meaning as previously defined.

The term oxy(2-6C)alkyl means an oxyalkyl group having 2-6 carbon atoms, the alkyl moiety having the same meaning as previously defined.

The term (di)[(1-4C)alkyl]amino as used herein means an amino group, monosubstituted or disubstituted with alkyl groups, each of which contains 1-4 carbon atoms and has the same meaning as previously defined.

The term (di)[(1-3C)]alkylamino as used herein means an amino group, monosubstituted or disubstituted with alkyl groups, each of which contains 1-3 carbon atoms and has the same meaning as previously defined.

The term halogen means fluorine, chlorine, bromine or iodine.

The term pharmaceutically acceptable salt represents those salts which are, within the scope of medical judgement, suitable for use in contact for the tissues of humans and/or animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art.

The compounds of the present invention possess at least one chiral carbon atom and may therefore be obtained as pure enantiomers, or as a mixture of enantiomers, or as a mixture of diastereomers.

The compounds of the invention may form hydrates or solvates. It is known to those of skill in the art that charged compounds form hydrated species when lyophilized with water, or form solvated species when concentrated in a solution with an appropriate organic solvent. The compounds of this invention include the hydrates or solvates of the compounds listed.

Another aspect of the present invention relates to the compounds and the use of compounds of Formula I wherein Y is a bond, O or NH, whereas

if Y is a bond, R³ is phenyl, optionally mono/di substituted with halogen, hydroxy, methoxy, phenoxy, phenyl, (1-4C)alkyl, nitro, amino or (di)[(1-4C)alkyl]amino or R³ is R⁶(1-6C)alkyl, R⁶(3-6C)cycloalkyl, R⁷oxy(1-6C)alkyl, or 2-pyridyl or

R³ is a 5-membered heteroaryl, optionally substituted with (1-3C)alkoxy, (1-3C)alkyl or halogen;

if Y is O, R³ is R⁶(1-6C)alkyl or R⁷oxy(2-6C)alkyl; and

if Y is NH, R³ is R⁶(2-6C)alkyl, R⁷oxy(2-6C)alkyl;

wherein Y is a bond, O or NH, whereas if Y is O, R³ is R⁶(1-6C)alkyl or R⁷oxy(2-6C)alkyl and wherein if Y is NH and R³ is R⁶(2-6C)alkyl or R⁷oxy(2-6C)alkyl.

Another aspect of the present invention relates to the compounds and the use of compounds of Formula I wherein X is a bond or O.

Yet another aspect of the invention relates to the compounds and the use of compounds of Formula I wherein R² is H and X is a bond.

Another aspect of the invention relates to the compounds and the use of compounds according to Formula I wherein R² is R⁵(1-4C)alkyl and X=O.

The present invention also provides for the compounds and the use of compounds according to Formula I wherein R⁴ is H, or (di)[(1-3C)alkyl]amino.

The invention also provides the compounds and the use of compounds according to Formula I wherein X is a bond, O or NH.

The invention also relates to compounds and use of compounds according to Formula I which are enantiomers showing the (+) optical rotation.

Yet another aspect of the invention concerns the compounds and use of compounds wherein all specific definitions of the groups R¹ through R⁹ and X and Y as defined here above are combined in the compound of formula I.

Suitable methods for the preparation of the compounds of the invention are outlined below.

If R¹=Me, the compounds of general formula I can be prepared starting with the well-documented Skraup reaction. Performing this reaction on N-*tert*-butoxycarbonyl (N-Boc) protected 1,4-phenylenediamine **1** gives 1,2-dihydroquinoline derivative **2**. The above mentioned reaction is typically conducted at elevated temperature in acetone or mesityl oxide in the presence of iodine or protic acid such as hydrochloric acid, *p*-toluenesulfonic acid or aqueous hydrogen iodide. Alternatively, compound **2** can be prepared by reacting compound **1** with acetone in the presence of MgSO₄, 4-*tert*-butylcatechol and iodine (L.G. Hamann, R.I. Higuchi, L. Zhi, J.P. Edwards and X.-N. Wang, J. (1998) Med. Chem. **41**, 623-639). In yet another procedure, the reaction can be performed in acetone using lanthanide triflates (e.g. scandium triflate) as catalysts. In this case, the reaction can be run at room temperature or at elevated temperatures using conventional heating or microwave irradiation (M. E. Theoclitou and L. A. Robinson (2002) Tetrahedron Lett. **43**:3907-3910).

Subsequent 1-*N*-acetylation of compound **2** can be carried out using standard conditions. Compound **2** can be acylated in a solvent such as tetrahydrofuran, with acetyl chloride in the presence of a base such as pyridine to give 1-*N*-acetyl-4-methyl-1,2-dihydroquinoline **3**.


Cleavage of the Boc protective group under conditions well known to those skilled in the art affords 6-amino-1,2-dihydroquinoline **4**. This reaction is typically conducted in dichloromethane in the presence of trifluoroacetic acid.

Introduction of the requisite substituted phenyl group at position 4 of the dihydroquinoline scaffold can be accomplished via Friedel-Crafts alkylation of benzene, anisole or bromobenzene with compound **4** to yield compounds of general formula **5**.

This reaction is conducted at ambient or elevated temperatures in benzene, anisole or bromobenzene under catalysis of a Lewis acid (e.g. AlCl_3).

Subsequent 6-*N*-functionalisation of the compounds of general formula **5** can be carried out using standard conditions to give the compounds of general formula **6**, wherein R3 is as previously defined. For example, compounds of formula **5** may react in a solvent such as dichloromethane, tetrahydrofuran or toluene with an acyl halide (R3-C(O)-hal , where $\text{hal}=\text{Cl}$, Br or I) in the presence of a base such as *N,N*-diisopropylethylamine or pyridine. Alternatively, acylation can be accomplished by reaction with an appropriate carboxylic acid ($\text{R3-CO}_2\text{H}$) in the presence of a coupling reagent such as *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) or *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and a tertiary base, e.g. *N,N*-diisopropylethylamine, in a solvent such as *N,N*-dimethylformamide or dichloromethane at ambient or elevated temperature.

Furthermore, compounds of general formula **5** can be converted to carbamates **6b** or ureas **6c** via the isocyanate **7**. In a typical reaction, compound **5** is converted into the isocyanate **7** in a solvent such as ethyl acetate with trichloromethyl chloroformate in the presence of activated carbon at elevated temperature. The isocyanate **7** can be converted with the appropriate alcohols R3-OH into carbamates **6b** (wherein R3 is as previously defined) or with the appropriate amines R3-NH_2 into ureas **6c** (wherein R3 is as previously defined) in a solvent such as tetrahydrofuran or dichloromethane in the presence of a base such as triethyl amine or *N,N*-diisopropylethylamine at ambient or elevated temperature. Alternatively, compounds of general formula **5** can be converted in a solvent such as tetrahydrofuran, dichloromethane or *N,N*-dimethylformamide with appropriate chloroformates R3-O-C(O)-Cl or isocyanates R3-N=C=O into carbamates **6b** or ureas **6c** respectively, wherein R3 is as previously defined, at ambient or elevated temperature.




If $R_1=H$, the compounds of general formula I can be prepared starting with a Wittig reaction. Performing this reaction with a ketone of general formula **8** with (diethoxyphosphoryl)-acetic acid ethyl ester yields the α,β -unsaturated esters of general formula **9** which can be converted to the carboxylic acids of general formula **10** by sodium hydroxide (2N) in ethanol at room temperature.

Subsequent acylation of aniline with acids of general formula **10** can be carried out using standard conditions to give compounds of general formula **11**. Acylation can be accomplished in the presence of a coupling reagent such as *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) or *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) and a tertiary base, e.g. *N,N*-diisopropylethylamine, in a solvent such as *N,N*-dimethylformamide or dichloromethane at ambient or elevated temperature.

The ring closure of compounds of general formula **11** can be performed with concentrated H_2SO_4 at room temperature yielding compounds of the general formula **12** which can be subsequently reduced using $BH_3-S(CH_3)_2$ in toluene at elevated temperature to yield tetrahydroquinolines of general formula **13**.

Compounds of general formula **13** can be acetylated using standard conditions. In a typical experiment, compounds of general formula **13** are reacted in a solvent such as dichloromethane or tetrahydrofuran with acetyl chloride in the presence of a base such as triethylamine or pyridine to give compounds of general formula **14**.

Introduction of the nitro-group at position 6 of the tetrahydroquinolines **14** can be accomplished by using a mixture of nitric acid and acetic anhydride in dichloromethane as the solvent at room temperature. The obtained compounds of general formula **15** can be converted to the aniline derivatives **16**, by a reduction using zinc in the presence of acetic acid and tetrahydrofuran as the solvent at 0°C.



Subsequent 6-*N*-acylation of the compounds of general formula **16**, to give the compounds of general formula **17a**, wherein R₃ is as previously defined can be carried out using standard conditions, as described for the acylation of compounds of the general formula **5**. Similarly, carbamates **17b**, wherein R₃ is as previously defined, and ureas **17c**, wherein R₃ is as previously defined, can be prepared, starting from compounds of general formula **16**, as described for carbamates **6b** and ureas **6c**.

Furthermore, compounds of general formula **20** and **23** can be prepared as described for compounds of general formula **30** and **33** respectively, starting from compounds of general formula **17**.


Compounds of general formula **17** and **6** (if R = Br) can be converted via a Suzuki reaction with boronic acids or boronic esters into compounds of general formula **24**. In a typical experiment, a bromide of general formula **17** or **6** can react with a boronic acid or boronic ester in a solvent such as dimethoxyethane or dioxane using cesium fluoride or potassium phosphate as a base and a palladium catalyst, such as palladium tetrakis(triphenylphosphine) or palladium dichloroditriphenylphosphine, in the presence of triphenylphosphine or triphenylarsine at elevated temperature. Furthermore, compounds of general formula **17** and **6** (if R = Br) can be converted via Buchwald chemistry into the anilines of general formula **25**. Compounds of general formula **17** or **6** may react in dimethoxyethane with benzophenone imine in the presence of sodium *tert*-butoxide as a base, tris(dibenzylideneacetone)dipalladium as a catalyst and 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl at elevated temperature followed by a hydrolysis with aqueous hydrochloric acid in tetrahydrofuran at ambient temperature to give the aniline derivatives of general formula **25**.

Alkylation of compounds of general formula **25**, yielding compounds of general formula **27a** wherein R₂ is as previously defined, can be carried out using the appropriate alkyl halides (R₂-hal, wherein hal = Cl, Br or I) in a solvent such as ethanol, tetrahydrofuran or *N,N*-dimethylformamide with a base such as triethyl amine, potassium carbonate, cesium carbonate or sodium bicarbonate in the presence of sodium iodide or potassium iodide as a catalyst. Furthermore, compounds of general formula **25** can be converted to carbamates of general formula **27b** via the isocyanates of general formula **26**. In a typical reaction, compounds of general formula **25** are converted into the isocyanates of general formula **26** in a solvent such as ethyl acetate with trichloromethyl chloroformate in the presence of activated carbon at elevated temperature. The isocyanates of general formula **26** can be converted into carbamates of general formula **27b**, wherein R₂ is as previously defined, in a solvent such as tetrahydrofuran or dichloromethane with the appropriate alcohol in the presence of a base such as triethyl amine or *N,N*-diisopropylethylamine. Alternatively, compounds of general formula **25** can be converted into carbamates of general formula **27b**, wherein R₂ is as previously defined, in a solvent such as tetrahydrofurane, dichloromethane or *N,N*-dimethylformamide with an appropriate chloroformate at ambient or elevated temperature.

Introduction of the nitro-group onto compounds of general formula **27** can be accomplished via a nitration using a mixture of (fuming) nitric acid and glacial acid in dichloromethane or concentrated sulfuric acid at ambient temperature or elevated temperature to yield compounds of general formula **28**. The aniline derivatives **29** can be obtained via a reduction of compounds of general formula **28** with zinc in tetrahydrofuran in the presence of glacial acetic acid at ambient or elevated temperature. The primary amine function in compounds of general formula **29** can be converted into alkylated anilines of general formula **30** via a reductive alkylation with aliphatic aldehydes in methanol using sodium cyanoborohydride in the presence of glacial acetic acid.

Furthermore, compounds of general formula **29** can be converted into alcohols of general formula **32** via the diazonium salts of general formula **31** by using methods well known to those skilled in the art. In a typical experiment, compounds of general formula **29** react in a solvent such as water with NaNO_2 and H_2SO_4 followed by addition of urea, $\text{Cu}(\text{NO}_3)_2$ and Cu_2O at 0°C or ambient temperature to yield the alcohols of general formula **32**.

Alkylation of compounds of general formula **32** can be carried out using alkyl halides in a solvent such as ethanol, tetrahydrofuran or *N,N*-dimethylformamide with a base such as triethyl amine, potassium carbonate, cesium carbonate or sodium bicarbonate in the presence of sodium iodide or potassium iodide as a catalyst to yield compounds of general formula **33**.




A procedure to obtain compounds of general formula **39** starts with the conversion of compounds of general formula **5** or **16** with 9-fluorenylmethyl chloroformate (Fmoc-Cl) in dichloromethane in the presence of triethyl amine at 0°C to give compounds of general formula **34**.

Subsequent introduction of the nitro-group can be accomplished via a nitration using a mixture of fuming nitric acid and glacial acid in dichloromethane at ambient temperature to yield compounds of general formula **35**. The aniline derivatives **36** can be obtained via a reduction of compounds **35** with zinc in tetrahydrofuran in the presence of glacial acetic acid at ambient temperature. The primary amine function in compounds **36** can be converted into alkylated anilines of general formula **37** via a reductive alkylation with aliphatic aldehydes in methanol using sodium cyanoborohydride in the presence of glacial acetic acid. The Fmoc-group removal can be accomplished with piperidine in dichloromethane to give the aniline derivatives of general formula **38**.


Subsequent 6-*N*-acylation of the compounds of general formula **38** can be carried out using standard conditions, as described for the acylation of compounds of general

formula **5**, to give the compounds of general formula **39a**, wherein R3 is as previously defined. Similarly, carbamates **39b**, wherein R3 is as previously defined, and ureas **39c**, wherein R3 is as previously defined, can be prepared, starting from compounds of general formula **38**, as described for carbamates **6b** and ureas **6c**.



Alternatively, compounds of general formula **39** can be prepared starting from compounds of general formula **35**. The Fmoc-group removal can be accomplished with piperidine in dichloromethane to give the aniline derivatives of general formula **40**.


Subsequent 6-N-functionalization of compounds of general formula **40** can be carried out using standard conditions, as described for the preparation of compounds of general formula **6**, to give compounds of general formula **41**, wherein R3 is as previously defined. The aniline derivatives **42** can be obtained via a reduction of compounds **41** with zinc in tetrahydrofuran in the presence of glacial acetic acid at ambient temperature. The primary amine function in compounds **42** can be converted into alkylated anilines of general formula **39** via a reductive alkylation with aldehydes in methanol using sodium cyanoborohydride in the presence of glacial acetic acid.




A procedure to obtain compounds of general formula **51** and **52** starts with the reaction of compound **43** with 9-fluorenylmethyl chloroformate (Fmoc-Cl) in tetrahydrofuran in the presence of pyridine at 0°C to give compound **44**. The aniline derivative **45** can be obtained via a reduction of compound **44** with zinc in tetrahydrofuran in the presence of glacial acetic acid at ambient temperature. Dihydroquinoline derivative **46** can be prepared via the previously mentioned Skraup reaction followed by *N*-acylation using acetyl chloride in pyridine/dichloromethane (1/1) as the solvent in the presence of *N,N*-dimethyl aniline to yield compound **47**. The Fmoc-group removal can be accomplished with piperidine in dichloromethane to give the aniline derivative **48**. Modification of the 6-amino group to amides **49a**, carbamates **49b** and ureas **49c**, can be performed using conditions as mentioned for the preparation of compounds of general formula **6**.

Introduction of the phenyl group at position 4 of the dihydroquinoline scaffold can be accomplished via Friedel-Crafts alkylation of benzene with the compounds of general formula **49**. This reaction is conducted at ambient or elevated temperatures in benzene, under catalysis of a Lewis acid (e.g. AlCl₃). Under these conditions demethylation takes place to yield compounds of general formula **50**. Compounds of general formula **52** can be obtained if the reaction is carried out in anisole.

Alcohols of general formula **50** can be alkylated using alkyl halides in a solvent such as acetonitrile, tetrahydrofuran or *N,N*-dimethylformamide in the presence of a base such as potassium carbonate, potassium *tert*-butoxide or sodium hydride to yield the compounds of general formula **51**.



A procedure to obtain compounds of general formula **54** and **57** starts with the demethylation of compounds with general formula **6** or **17** where R=OMe. If R=OMe, demethylation can be effected in a solvent such as dichloromethane in the presence of boron tribromide (BBr₃) at ambient or elevated temperature to yield alcohols of general formula **53**. Alkylation of compounds of general formula **53** can be carried out using alkyl halides (R₂-hal) in a solvent such as ethanol, tetrahydrofuran or *N,N*-dimethylformamide with a base such as triethyl amine, potassium carbonate, cesium carbonate or sodium bicarbonate in the presence of sodium iodide or potassium iodide as a catalyst to yield compounds of general formula **54**. Introduction of the nitro-group onto compounds of general formula **54** to yield compounds of general formula **55**, followed by nitro group reduction to compounds of general formula **56** and amino alkylation to yield compounds of general formula **57** can be accomplished using conditions as described for the conversion of compounds of general formula **27** into compounds of general formula **30**.



Additionally suitable methods for the preparation of some compounds of the invention are described in WO2004/056779 and WO2004/056780.

Some of the compounds of the invention, which can be in the form of a free base, may be isolated from the reaction mixture in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salts may also be obtained by treating the free base of formula I with an organic or inorganic acid such as hydrogen chloride, hydrogen bromide, hydrogen iodide, sulfuric acid, phosphoric acid, acetic acid, propionic acid, glycolic acid, maleic acid, malonic acid, methanesulphonic acid, fumaric acid, succinic acid, tartaric acid, citric acid, benzoic acid, and ascorbic acid.

Methods for obtaining the pure enantiomers are well known in the art, *e.g.* crystallization of salts which are obtained from optically active acids and the racemic mixture, or chromatography using chiral columns. For diastereomers, straight phase or reversed phase columns may be used. Optical rotation can easily be measured *e.g.* with a polarimeter.

The compounds of the present invention inhibit the TSH receptor. The skilled artisan will recognize that desirable IC_{50} values are dependent on the compound tested. For example, a compound with an IC_{50} value of less than $1E-5$ M is generally considered a candidate for drug selection. In general, the IC_{50} values are preferably lower than $1E-7$ M. However, a compound which has a larger IC_{50} value but is more selective for the

particular receptor may be even a better candidate. The compounds of the present invention might also possess FSH receptor activity but they do not have a FSH receptor selectivity over the TSH receptor. Generally it can be stated that the compounds of the present invention have high TSH receptor activity and are at least equipotent on FSH and TSH receptors.

Methods to determine receptor binding, as well as *in vitro* and *in vivo* assays to determine biological activity of thyrotropin or the TSH receptor are well known. In the *in vitro* assays, the expressed TSH receptor is incubated with the test compound and receptor binding or stimulation/inhibition of a functional response is measured. *Ex vivo* human or animal thyrocytes/thyroid slices or thyroid cell lines may be used (Fuhrer et al. (2003) Endocrinology **144**, 4018-4030). Alternatively, cDNA encoding the TSH receptor may heterologously be expressed in suitable host cells, e.g. Chinese Hamster Ovary cells, but other cell lines, preferably of mammalian origin, are also suitable (Fuhrer et al. (2003) Endocrinology **144**, 4018-4030). Methods to construct cells expressing recombinant TSHR are well known in the art.

Techniques for site-directed DNA mutagenesis, DNA ligation, PCR and construction of suitable expression systems are all well known in the art. Portions, or all, of the DNA encoding the desired receptor can be constructed synthetically using standard solid phase techniques, preferably to include restriction sites for ease of ligation. Suitable control elements for transcription and translation of the inserted coding sequence can be ligated to the DNA coding sequence. As is well known, expression systems are available, which are compatible with a wide variety of hosts, including bacteria and eukaryotic hosts such as yeast, insect cells, avian, mammalian cells and the like.

Cells expressing the TSH receptor are incubated with the test compound to observe binding, stimulation or inhibition of a functional response. Alternatively, isolated cell membranes containing the expressed receptor may be used to determine binding of test compound. For measurement of binding, radioactively labeled or fluorescently labeled compounds may be used. Also competitive binding assays can be performed. Another biochemical assay involves the screening for TSH receptor-mediated cAMP accumulation. Such an assay involves incubation of the TSH receptor-expressing cells

with the test compound for a sufficient period of time and measurement of cAMP, often in the presence of a cAMP phosphodiesterase inhibitor to block cAMP degradation. The amount of cAMP can be decreased or increased, depending on the inhibitory or stimulating effect of the test compound upon binding to the receptor. Screening for TSH receptor antagonists involves incubation of the TSH receptor expressing cells with a concentration range of test compound in the presence of a fixed, submaximally effective TSH concentration (i.e., a TSH concentration that induces approximately 80% of the maximal stimulation of cAMP accumulation in the absence of compound). Instead of TSH, serum from Graves' disease patients, or (partially) purified TSI can be used to stimulate the TSH receptor (Gerding et al. (2000) Clin. Endocrinol. **52**, 267-271). Alternatively, constitutively active TSH receptors can be heterologously expressed and the antagonistic effect of test compounds on the active receptor may result in reduction in the increased basal cAMP levels (Fuhrer et al. (2003) Endocrinology **144**, 4018-4030). From the concentration-effect curves, the IC₅₀ value and the percentage of inhibition of TSH receptor-induced cAMP accumulation can be determined for each of the test compounds.

In addition to a direct measurement of cAMP, cell lines can be used, which in addition to transfection with receptor cDNA are also transfected with a second cDNA encoding a reporter gene, which expression is dependent on the intracellular level of cAMP. Such reporter genes might be cAMP-inducible or might be constructed in such a way that they are connected to novel cAMP responsive elements. In general, reporter gene expression might be controlled by any response element reacting to changing levels of cAMP. Suitable reporter genes are e.g. the genes encoding firefly luciferase, beta-galactosidase, alkaline phosphatase or beta-lactamase. The principles of such transactivation assays are well known in the art and are described, e.g. in Evans et al. (1999) J.Clin. Endocrinol. Metab. **84**, 374-377. As reference compound human (recombinant) TSH or bovine TSH can be used.

The present invention also relates to a pharmaceutical composition comprising a tetrahydroquinoline derivative or pharmaceutically acceptable salts thereof having the general formula I in admixture with pharmaceutically acceptable auxiliaries and

optionally other therapeutic agents. The auxiliaries must be “acceptable” in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

Compositions include e.g. those suitable for oral, ocular, sublingual, subcutaneous, intravenous, intramuscular, local, or rectal administration, and the like, all in unit dosage forms for administration.

For oral administration, the active ingredient may be presented as discrete units, such as tablets, capsules, powders, granulates, solutions, suspensions, and the like.

For parenteral administration, the pharmaceutical composition of the invention may be presented in unit-dose or multi-dose containers, e.g. injection liquids in predetermined amounts, for example in sealed vials and ampoules, and may also be stored in a freeze dried (lyophilized) condition requiring only the addition of sterile liquid carrier, e.g. water, prior to use.

For an ophthalmic formulation, the active ingredient may be presented as a solution, suspension, ointment, or gel for application on the conjunctiva or cornea, or for retrobulbar injection.

Mixed with such pharmaceutically acceptable auxiliaries, e.g. as described in the standard reference, Gennaro, A.R. et al., Remington: *The Science and Practice of Pharmacy* (20th Edition., Lippincott Williams & Wilkins, 2000, see especially Part 5: Pharmaceutical Manufacturing), the active agent may be compressed into solid dosage units, such as pills, tablets, or be processed into capsules or suppositories. By means of pharmaceutically acceptable liquids the active agent can be applied as a fluid composition, e.g. as an injection preparation, in the form of a solution, suspension, emulsion, or as a spray, e.g. a nasal spray.

For making solid dosage units, the use of conventional additives such as fillers, colorants, polymeric binders and the like is contemplated. In general any pharmaceutically acceptable additive which does not interfere with the function of the active compounds can be used. Suitable carriers with which the active agent of the invention can be administered as solid compositions include lactose, starch, cellulose derivatives and the like, or mixtures thereof, used in suitable amounts. For parenteral administration, aqueous suspensions, isotonic saline solutions and sterile injectable

solutions may be used, containing pharmaceutically acceptable dispersing agents and/or wetting agents, such as propylene glycol or butylene glycol.

The invention further includes a pharmaceutical composition, as hereinbefore described, in combination with packaging material suitable for said composition, said packaging material including instructions for the use of the composition for the use as herein described.

Thus, the present invention also provides a kit to treat disorders responsive to TSH receptor mediated pathways.

The kits comprise: A) a pharmaceutical composition comprising a compound of Formula I or a pharmaceutically acceptable salt thereof; and B) instructions describing a method of using the pharmaceutical composition to treat disorders responsive to TSH receptor mediated pathways.

In another aspect the kit can be used to prevent disorders responsive to TSH receptor mediated pathways.

A "kit" as used in the instant application includes a container for containing the pharmaceutical compositions and may also include divided containers such as a divided bottle or a divided foil packet. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material

The tetrahydroquinoline derivatives of the invention can also be administered in the form of implantable pharmaceutical devices, consisting of a core of active material, encased by a release rate-regulating membrane. Such implants are to be applied subcutaneously or locally, and will release the active ingredient at an approximately constant rate over relatively large periods of time, for instance from weeks to years. Methods for the preparation of implantable pharmaceutical devices as such are known in the art, for example as described in European Patent 0,303,306 (AKZO Nobel N.V.).

The exact dose and regimen of administration of the active ingredient, or a pharmaceutical composition thereof, will necessarily be dependent upon the therapeutic effect to be achieved, and may vary with the particular compound, the route of

administration, and the age and condition of the individual subject to whom the medicament is to be administered.

In general parenteral administration requires lower dosages than other methods of administration which are more dependent upon absorption. However, a dosage for humans preferably contains 0.0001-25 mg per kg body weight. The desired dose may be presented as one dose or as multiple subdoses administered at appropriate intervals throughout the day.

A further aspect of the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I as described hereinabove for the manufacture of a medicament to be used for the treatment of disorders responsive to TSH receptor mediated pathways. Thus, patients in need thereof can be administered with suitable amounts of the compounds according to the invention.

In another aspect the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the manufacture of a medicament to be used for the treatment of patients with a need to inhibit the actions of TSH, TSI or constitutive activity of TSH receptors.

In another aspect the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the treatment of hyperthyroidism.

In yet another aspect the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the treatment of Graves' disease.

In still another aspect the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the treatment of Graves' ophthalmopathy.

Another aspect of the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the treatment of Graves' associated pretibial dermopathy.

Still another aspect the invention resides in the use of a tetrahydroquinoline derivative compound having the general formula I for the treatment of nodular goitre.

The compounds according to the invention can also be used for the treatment of thyroid cancer.

In another aspect the invention relates to the use of a tetrahydroquinoline derivative compound having the general formula I for the prevention of the here above identified disorders.

The invention is illustrated by the following examples.

Examples

General comments

The following abbreviations are used in the examples: DMA = *N,N*-dimethylaniline, DIPEA = *N,N*-diisopropylethylamine; TFA = trifluoroacetic acid, DtBAD = di-*tert*-butyl azodicarboxylate; TBTU = *O*-Benzotriazole-1-yl-*N,N,N',N'*-tetramethyluronium tetrafluoroborate; HATU = *O*-(7-azabenzotriazole-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate; Fmoc = 9-fluorenylmethoxycarbonyl; Fmoc-Cl = 9-fluorenylmethoxycarbonylchloride; DMF = *N,N*-dimethylformamide; Boc = *tert*-butoxycarbonyl; THF = tetrahydrofuran; DMAP = dimethyl-pyridin-4-yl-amine; HOAc = acetic acid; Et₃N = triethyl amine; EtOAc = ethyl acetate; DCM = dichloro methane; MeOH = methanol; MeI = methyl iodide; DME = 1,2-dimethoxy ethane.

The names of the final products described in the examples are generated using the Beilstein Autonom program (version: 4.01.304).

Unless stated otherwise, all final products of the examples below are lyophilized from water/1,4-dioxane mixtures or water/acetonitrile mixtures. If the compound was prepared as a HCl- or TFA salt, the respective acids were added in appropriate amounts to the solvent mixture before lyophilization.

Example 1

Hexanoic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

(a) (2,2,4-Trimethyl-1,2-dihydro-quinolin-6-yl)-carbamic acid *tert*-butyl ester

To a stirred solution of N-boc-1,4-phenylene diamine (13.33 g, 63.9 mmol) in mesityl oxide (40 ml, 367 mmol), under an N₂ atmosphere, was added iodine (3.47 g, 12.8 mmol) portionwise over 5 minutes. The reaction was stirred at 100 °C for 3 hours, then

cooled to ambient temperature and reduced to yield the title compound as a crude brown oil (18.4 g, 100%). The product was used without further purification.

(b) 1-(1-Acetyl-2,2,4-trimethyl-1,2-dihydro-quinolin-6-yl)-carbamic acid *tert*-butyl ester

A stirred solution of the compound described in example 1 (a) (16.4 g, 57 mmol) in THF (200 ml) and pyridine (5.07 ml, 63 mmol), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). Acetylchloride (4.46 ml, 63 mmol) in THF (50 ml) was then added dropwise over a period of 15 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 3 hours. The reaction was then quenched with water. The organics were extracted with EtOAc, washed with water, dried (Na₂SO₄), filtered and reduced to an oil. The oil was purified by flash chromatography (heptane/EtOAc 8:2) to yield the title compound (11.95 g, 57%).

(c) 1-(6-Amino-2,2,4-trimethyl-2*H*-quinolin-1-yl)-ethanone

A stirred solution of the compound described in example 1 (b) (11.95 g, 36 mmol) in DCM (250 ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). Trifluoroacetic acid (27.7 ml, 360 mmol) was then added dropwise over a period of 15 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 1 hour. The reaction was then quenched by the addition of sodium carbonate (s) under vigorous stirring and water was added. The organics were extracted with EtOAc, washed with water, dried (Na₂SO₄), filtered and reduced to an oil. The oil was purified by flash chromatography (heptane/EtOAc 6:4) to yield the title compound (2.67 g, 32%).

(d) 1-(6-Amino-2,2,4-trimethyl-4-phenyl-3,4-dihydro-2*H*-quinolin-1-yl)-ethanone

A stirred solution of the compound described in example 1 (c) (100 mg, 494 µmol) in dry benzene (2ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). Aluminium trichloride (198 mg, 1.48 mmol) was then added portionwise over 15 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 16 hours. The reaction was then quenched with aqueous ammonium chloride. The organics were extracted with EtOAc, washed with water, dried (Na₂SO₄), filtered and reduced to yield the title compound as a crude brown foam (124 mg, 81%). Product foam was used without further purification.

(e) Hexanoic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

A stirred solution of the compound described in example 1 (d) (880 mg, 2.85 mmol) and triethylamine (477 μ L, 3.42 mmol) in dry DCM (7.5 ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). A solution of hexanoyl chloride (479 μ L, 3.42 mmol) in DCM (2.5 ml) was then added drop wise over 15 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 1 hour. The reaction was then quenched with 1N aqueous hydrochloric acid. The organics were extracted with DCM, washed with NaHCO₃ in H₂O solution, then water, then dried (PE filter) and reduced to yield an oil. The oil was purified by flash chromatography then the product crystallized from EtOAc to yield the title compound as a white solid (871 mg, 75%). Data : (m/z) = 407.3 (M+H)⁺.

Example 2

1-Methyl-1H-pyrrole-2-carboxylic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

DIPEA (70.6 μ L, 405 μ mol), HATU (46.2 μ L, 122 μ mol) and N-methylpyrrole-2-carboxylic acid (15.2 mg, 122 μ mol) in dry DCM (0.75ml) were stirred together, under an N₂ atmosphere, for 15 minutes. The reaction mixture was then cooled to 0°C (ice/water bath) then a solution of the compound described in example 1(d) (25 mg, 81.1 μ mol) in dry DCM (0.25 ml) was added dropwise over 3 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 1 hour. The reaction was then quenched with 1N aqueous hydrochloric acid. The organics were extracted with DCM, washed with NaHCO₃ in H₂O solution, then water, then dried (PE filter) and reduced to yield an oil. The oil was purified by flash chromatography to yield the title compound as a white solid (15 mg, 46%). Data : (m/z) = 416.3 (M+H)⁺.

Example 3

N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-5-bromo-2-methylamino-benzamide

This compound was prepared, in an analogous manner as described in Example 2, from the compound described in example 1 (d) (264 mg, 78%) Data : (m/z) = 521 (M+H)⁺.

Example 4

Pyridine-2-carboxylic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

This compound was prepared, in an analogous manner as described in Example 2, from the compound described in example 1 (d) (320 mg, 46%) Data : (m/z) = 414 (M+H)⁺.

Example 5

N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3,4-dimethyl-benzamide

This compound was prepared, in an analogous manner as described in Example 1, from the compound described in example 1 (d), to yield the title compound (320 mg, 46%) Data : (m/z) = 414 (M+H)⁺.

Example 6

(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 2-phenoxy-ethyl ester

To a stirred solution of the compound described in example 1 (d) (124 mg, 402 μ mol) and a catalytic amount of activated charcoal in dry EtOAc (4.6 ml), under an N₂ atmosphere, was added trichloromethyl chloroformate (97 μ l, 804 μ mol). The reaction was stirred at reflux for 2 hours, then cooled to ambient temperature and filtered over dikalite and concentrated under reduced pressure to yield the isocyanate as a crude oil (134 mg, 100%).

The isocyanate (48.8 mg, 146 μ mol) in THF (1 ml) was then added to a solution of 2-phenoxyethanol (182 μ l, 1.46 mmol) and TEA (211 μ l, 1.46 mmol) in THF (2 ml), under an N₂ atmosphere. The reaction was stirred for 15 hours at ambient temperature, then quenched with water, extracted with EtOAc, washed with brine, dried (Na₂SO₄), filtered and reduced to an oil. The oil was purified by flash chromatography (heptane/EtOAc) to yield the title compound (51 mg, 74%). Data : (m/z) = 473.5 (M+H)⁺.

Example 7

(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid butyl ester

A stirred solution of the compound described in example 1 (d) (20 mg, 65 μ mol) in DCM (2 ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). N-

butylchloroformate (16.7 μ l, 130 μ mol) was added and the cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 16 hours. The reaction was then quenched with water, extracted with EtOAc, washed with brine, dried (Na_2SO_4), filtered and reduced to an oil. The oil was purified by flash chromatography (heptane/EtOAc) to yield the title compound (10 mg, 37%) Data : $(m/z) = 409.3 (M+H)^+$..

Example 8

1-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-cyclopentyl-urea

A stirred solution of the compound described in example 1 (d) (25 mg, 81.1 μ mol) and triethylamine (16.9 μ L, 122 μ mol) in dry DCM (0.75 ml), under an N_2 atmosphere, was cooled to 0°C (ice/water bath). A solution of cyclopentyl isocyanate (11.0 mg, 97.3 μ mol) in DCM (0.25 ml) was then added dropwise over 3 minutes. The cooling was removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 1 hour. The reaction was quenched with 1N aqueous hydrochloric acid. The organics were extracted with DCM, washed with NaHCO_3 in H_2O solution and water, then dried (PE filter) and reduced to yield an oil. The oil was purified by flash chromatography to yield the title compound as a white solid (3 mg, 9%). Data : $(m/z) = 420.5 (M+H)^+$.

Example 9

Hexanoic acid [1-acetyl-4-(4-bromo-phenyl)-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

(a) (E)-3-(4-Bromo-phenyl)-but-2-enoic acid ethyl ester

A solution of potassium tert-butoxide (10.45g, 90.43mmol) in THF (250ml) was stirred under nitrogen atmosphere. A solution of triethyl phosphonoacetate (18.1ml, 90.43mmol) in THF (100ml) was then added and the reaction mixture was stirred for 30 minutes at room temperature. A solution of 1-(4-Bromo-phenyl)-ethanone (6g, 30.14mmol) in THF (100ml) was then added dropwise over 5 minutes. The reaction mixture was heated to reflux. After 3 hours the reaction mixture was cooled to ambient temperature and THF was removed under reduced pressure. The reaction mixture was dissolved in H_2O , extracted with EtOAc, washed with H_2O and brine, dried (Na_2SO_4)

and concentrated under reduced pressure. The crude product was purified by column chromatography on silica (Hept : EtOAc 8:2) to afford the title compound (8.1g, 99%).

(b) (E)-3-(4-Bromo-phenyl)-but-2-enoic acid

To a stirred solution of the compound described in example 9 (a) (8.16g, 30.3mmol) in ethanol (50ml), 2N aqueous sodium hydroxide (50ml) was added. The reaction mixture was stirred over night at room temperature. The reaction mixture was cooled and quenched with 2N aqueous hydrochloric acid. The organics were extracted into EtOAc, washed with H₂O and brine, dried (Na₂SO₄) and concentrated under reduced pressure to yield the title compound as a white solid (7.42g, 99%). Product used without further purification.

(c) (E)-3-(4-Bromo-phenyl)-but-2-enoic acid phenylamide

To a stirred solution of the compound described in example 9 (b) (2g, 8.296mmol) in CH₂Cl₂ (40ml), under N₂ atmosphere, DIPEA (2.89ml, 16.59mmol) and o-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluroniumhexafluorophosphate (3.79g, 9.96mmol) were added. The reaction mixture was stirred for 30 minutes at room temperature and a solution of aniline (910μl, 9.96mmol) in CH₂Cl₂ (30ml) was added dropwise. The reaction mixture was stirred for 3.5 hours and then quenched with 3% aqueous citric acid (100ml), extracted with CH₂Cl₂, washed with H₂O, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica (Hept : EtOAc 8:2) to afford the title compound (2.48g, 95%).

(d) 4-(4-Bromo-phenyl)-4-methyl-3,4-dihydro-1H-quinolin-2-one

The compound described in example 9 (c) (2.28g, 7.2mmol) was dissolved in concentrated sulfuric acid (20ml) and stirred for 45 minutes. The reaction mixture was poured into an ice/water mixture and the organics were extracted into CH₂Cl₂, washed with aqueous NaHCO₃ and H₂O, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica (Hept : EtOAc 8:2) to afford the title compound as a yellow solid (1.9g, 83%).

(e) 4-(4-Bromo-phenyl)-4-methyl-1,2,3,4-tetrahydro-quinoline

To a solution of the compound described in example 9 (d) (9.09g, 28.75mmol) in toluene (475ml), under N₂ atmosphere, a 2M solution of borane-methyl sulfide complex in toluene (37.4ml, 74.74mmol) was added dropwise over 20 minutes. The reaction

mixture was heated to reflux for 2 hours, then cooled to ambient temperature, quenched with H₂O and stirred for 50 minutes at room temperature. The organics were extracted into CH₂Cl₂, washed with H₂O, dried (Na₂SO₄) and concentrated under reduced pressure to afford the title compound as an oil (8.6g, 99%). The product was used without further purification.

(f) 1-[4-(4-Bromo-phenyl)-4-methyl-3,4-dihydro-2H-quinolin-1-yl]-ethanone

To a stirred solution of the compound described in example 9 (e) (8.69g, 28.75mmol) in CH₂Cl₂ (240ml), under N₂ atmosphere, pyridine (4.64ml, 57.5mmol) and acetyl chloride (3.08ml, 43.13mmol) were added and stirred at room temperature for 3 hours. The reaction mixture was then poured into ice water. The organics were extracted into CH₂Cl₂, washed with H₂O, dried (Na₂SO₄) and concentrated under reduced pressure to afford the title compound as an oil (9.8g, 99%). The product was used without further purification.

(g) 1-[4-(4-Bromo-phenyl)-4-methyl-6-nitro-3,4-dihydro-2H-quinolin-1-yl]-ethanone

To a stirred solution of the compound described in example 9 (f) (9.8g, 28.47mmol) in CH₂Cl₂ (210ml), under N₂ atmosphere, acetic anhydride (267μl, 2.85mmol) was added. A solution of fuming nitric acid (7.17ml, 0.17mol) in CH₂Cl₂ (10ml) was then added dropwise over 15 minutes and the reaction mixture was stirred overnight at room temperature. The reaction mixture was then poured into ice water. The organics were extracted into CH₂Cl₂, washed with aqueous NaHCO₃ and H₂O, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica (Hept : EtOAc 9:1) to afford the title compound (7.2g, 65%).

(h) 1-[6-Amino-4-(4-bromo-phenyl)-4-methyl-3,4-dihydro-2H-quinolin-1-yl]-ethanone

A stirred solution of the compound described in example 9 (g) (7.3g, 18.8 mmol) and acetic acid (10.73ml, 187.6mmol) in THF (690ml), under N₂ atmosphere, was cooled to 0°C (ice/water bath). Zinc (dust) (38.1g, 0.56mol) was then added portion wise over 15 minutes. The reaction mixture was stirred for 30 minutes at 0°C. The cooling was then removed and the reaction mixture was allowed to warm slowly to ambient temperature and stirred for 16 hours. The solids were filtered off and the filtrate was deluded with EtOAc and water. The organic layer was washed with aqueous NaHCO₃ and H₂O, dried (Na₂SO₄) and concentrated under reduced pressure to afford the title compound as an orange oil (6.67g, 99%). The product was used without further purification.

(i) Hexanoic acid [1-acetyl-4-(4-bromo-phenyl)-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

A stirred solution of the compound described in example 9 (h) (7.54g, 21.0 mmol) and triethylamine (3.52ml, 25.2mmol) in CH₂Cl₂ (750ml), under N₂ atmosphere, was cooled to 0°C (ice/water bath). N-hexanoyl chloride (3.54ml, 25.2mmol) was then added dropwise over 5 minutes and the reaction mixture was stirred for 1 hour at 0°C. Then the cooling was removed and the reaction mixture was allowed to stir overnight at room temperature. The reaction was then quenched with aqueous NaHCO₃. The organics were extracted into CH₂Cl₂, washed with H₂O, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica (Hept : EtOAc 1:1) to afford the title compound as a yellow foam (5.66g, 59%). Data : (m/z) = 457.3/459.3 (M+H)⁺.

Example 10

N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-2-phenoxy-acetamide

(a) 1-(6-Amino-4-methyl-4-phenyl-3,4-dihydro-2H-quinolin-1-yl)-ethanone

This compound has been prepared in an analogous manner as described for example 9 (h).

(b) N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-2-phenoxy-acetamide

To a solution of phenoxyacetic acid (19.5 mg, 0.13 mmol) in DCM (1ml), TBTU (52 mg, 0.16 mmol) and DIPEA (26μl, 0.15 mmol) were added and the reaction mixture was stirred at room temperature. After 10 minutes a solution of the compound described in example 10 (a) (30 mg, 0.11 mmol) in DCM (1ml) was added. The reaction mixture was stirred overnight at room temperature. The reaction was quenched with a saturated NaHCO₃ solution in H₂O and extracted with DCM. The organic layer was dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified with column chromatography on silica to afford the title compound (36 mg, 81 %). Data: (m/z) = 415.5 (M+H)⁺.

Example 11

N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide

This compound was prepared, in an analogous manner as described for example 10, from the compound described in example 10 (a), to afford the title compound (15mg, 32 %). Data: (m/z) = 447.3 (M+H)⁺.

Example 12

N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-thiophen-2-yl-propionamide

This compound was prepared, in an analogous manner as described in Example 11, from the compound described in example 10 (a), to afford the title compound (21mg, 47%) Data : (m/z) = 419 (M+H)⁺.

Example 13

1-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-benzyl)-urea

This compound was prepared, in an analogous manner as described in example 19 (d), from the compound described in example 10 (a), to afford the title compound (55mg, 87%). Data : (m/z) = 448 (M+H)⁺.

Example 14

Hexanoic acid {1-acetyl-4-[4-(3-fluoro-pyridin-4-yl)-phenyl]-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl}-amide

A stirred solution of the compound described in Example 9 (50mg, 0.11mmol), 3-fluoropyridine-4-boronic acid (42mg, 0.30mmol), potassium phosphate tribasic heptahydrate (44mg, 0.13mmol), bis(triphenylphosphine)palladium(II) chloride (4.6mg, 6.6μmol), triphenylarsine (2.3mg, 7.6μmol) and H₂O (0.5ml) in dioxane (3ml), under N₂ atmosphere, was heated to reflux for 5 hours. The reaction mixture was then cooled to ambient temperature, quenched with aqueous NaHCO₃ and diluted with EtOAc. The organic layer was washed with H₂O and brine, dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica, followed by preparative HPLC. Freeze-drying afforded the title compound (10mg, 19%). Data : (m/z) = 474.5 (M+H)⁺.

Example 15

[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid 3-chloro-benzyl ester

(a) Hexanoic acid [1-acetyl-4-(4-amino-phenyl)-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

To a stirred solution of the compound described in Example 9 (5.08g, 11.1mmol), tris(dibenzylideneacetone)dipalladium(0)chloroform adduct (850mg, 0.82mmol), 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (850mg, 1.37mmol) and sodium tert-butoxide (2.13g, 22.2mmol) in DME (200ml), under N₂ atmosphere, benzophenone imine (2.8ml, 16.8mmol) was added and heated to 80°C for 16 hours. The reaction mixture was then cooled to ambient temperature, diluted with EtOAc and solids were filtered off. The filtrate was concentrated under reduced pressure and purified by column chromatography on silica (Hept : EtOAc 7:3) to yield an orange solid. To a stirred solution of this orange solid in THF (21ml), under N₂ atmosphere, 2N aqueous hydrochloric acid was added. After 3 hours the reaction mixture was quenched with EtOAc, made basic with 2N aqueous sodium hydroxide, washed with H₂O and brine, dried (Na₂SO₄) and concentrated under reduced pressure to afford the title compound as an orange oil (3.95g, 90%). The product was used without further purification.

(b) 4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid 3-chloro-benzyl ester

This compound was prepared, in an analogous manner as described in Example 6, from the compound described in example 15 (a), to afford the title compound (16mg, 54%).
Data : (m/z) = 562 (M+H)⁺.

Example 16

[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid methyl ester

This compound was prepared, in an analogous manner as described in Example 6, from the compound described in example 15 (a), to afford the title compound (10mg, 41%).
Data : (m/z) = 452 (M+H)⁺.

Example 17

Cyclopentanecarboxylic acid (1-acetyl-7-dimethylamino-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

(a) (1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

A stirred solution of the compound described in example 1 (d) (2.39 g, 7.75 mmol) and pyridine (660 µL, 8.14 mmol) in dry DCM (15 ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). A solution of 9-fluorenylmethyl chloroformate (2.11 g, 8.14 mmol) in dry DCM (10 ml) was then added dropwise over 15 minutes. After 30

minutes the reaction was quenched with H₂O. The organics were extracted and reduced to yield an oil. The oil was purified by flash chromatography. The product was crystallised from MeOH/DCM then filtered off to yield the title compound as a yellow solid (3.94 g, 96%).

(b) (1-Acetyl-2,2,4-trimethyl-7-nitro-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

To a stirred solution of the compound described in example 17 (a) (3.94 g, 7.42 mmol) in glacial acetic acid (20 ml) and dry DCM (10 ml), under an N₂ atmosphere, was charged with concentrated nitric acid (310 µL, 7.42 mmol), dropwise over 3 minutes. Stirred at ambient temperature for 30 minutes then charged with concentrated nitric acid (100 µL, 2.36 mmol). Stirred for 15 minutes at ambient temperature then the reaction was then quenched with H₂O. The organics were extracted. MeOH (2 ml) was added to the organics and the DCM was removed by rotary evaporation. The product was crystallised from MeOH/DCM then filtered off to yield the title compound as a yellow solid (2.62 g, 61%).

(c) (1-Acetyl-7-amino-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

A stirred solution of the compound described in example 17 (b) (100 mg, 173 µmol) and glacial acetic acid (100 µL, 1.73 mmol) in dry THF (3 ml), under an N₂ atmosphere, was cooled to 0°C (ice/water bath). Zinc dust (227 mg, 3.47 mmol) was added portionwise over 10 minutes. The cooling was removed and the reaction was allowed to warm to ambient temperature and stirred for 2 hours. Fresh glacial acetic acid (100 µL, 1.73 mmol) and zinc dust (227 mg, 3.47 mmol) were added. After 10 minutes the reaction was filtered through decalite. DCM was added. The organics were washed with NaHCO₃, brine and dried (MgSO₄), filtered and reduced to yield the title compound as an orange oil (95 mg, 100%).

(d) (1-Acetyl-7-dimethylamino-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

To a stirred solution of the compound described in example 17 (c) (103 mg, 189 µmol) in methanol (2ml) and glacial acetic acid (135 µL, 2.36 mmol) was added sodium cyanoborohydride (18.8 mg, 302 µmol) then a 37% solution of formaldehyde in water (14 µL, 189 µmol). After 2 hours the reaction was quenched with NaHCO₃ and the organics

were extracted into EtOAc. The organics were then washed with brine, dried (MgSO₄), filtered and reduced to yield the title compound as a red solid (106 mg, 100%).

(e) 1-(6-Amino-7-dimethylamino-2,2,4-trimethyl-4-phenyl-3,4-dihydro-2H-quinolin-1-yl)-ethanone

To a stirred solution of the compound described in example 17 (d) (380 mg, 662 μ mol) in dry DCM (5ml) was added piperidine (1ml). After 2 minutes the reaction mixture was reduced to an oil by rotary evaporation. The oil was purified by flash chromatography to yield the title compound as a pink solid (232 mg, 100%).

Cyclopentanecarboxylic acid (1-acetyl-7-dimethylamino-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

To a stirred solution of HATU (224 mg, 590 μ mol) and DIPEA (340 μ L, 1.96 mmol) in DCM (5ml) was added cyclopentane carbonyl chloride (128 μ L, 1.18 mmol). Stirred under an N₂ atmosphere for 15 minutes then a solution of the compound described in example 17 (e) (138 mg, 393 μ mol) in DCM (5ml) was added dropwise. After 18 hours the reaction was quenched with 0.5N hydrochloric acid. The organics were washed with NaHCO₃, H₂O then dried (PE-filter) and reduced to yield an oil. The oil was purified by flash chromatography to yield the title compound as a white solid (147 mg, 85%). Data : (m/z) = 448.5 (M+H)⁺.

Example 18

N-(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-phenyl)-propionamide

This compound was prepared, in an analogous manner as described in Example 17, from the compound described in example 10a to afford the title compound (14mg, 46%)
Data : (m/z) = 490 (M+H)⁺.

Example 19

1-(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(2-methoxy-benzyl)-urea

(a) (1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

This compound was prepared, in an analogous manner as described in example 17 (a), from the compound described in example 10 (a) to afford the title compound (2.2g, 82%)

(b) 1-(1-Acetyl-4-methyl-7-nitro-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

This compound was prepared, in an analogous manner as described in example 17(b), from the compound described in example 19a to afford the title compound (1.1g, 58%)

(c) 1-(6-Amino-4-methyl-7-nitro-4-phenyl-3,4-dihydro-2H-quinolin-1-yl)-ethanone

To a stirred solution of the compound described in example 19 (b) (112mg, 0.2mmol) in CH₂Cl₂ (2 ml), piperidine (0.2ml, 2mmol) was added. The reaction was stirred overnight at room temperature. The reaction mixture was poured into H₂O and extracted into CH₂Cl₂. The organics were washed with H₂O and brine, dried (Na₂SO₄) and concentrated under reduced pressure. The product was purified by column chromatography on silica to give the title compound (50mg, 76%) Data : (m/z) = 326 (M+H)⁺.

(d) 1-(1-Acetyl-4-methyl-7-nitro-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(2-methoxy-benzyl)-urea

To a stirred solution of the compound described in example 19 (c) (20mg, 61.5μmol) in EtOAc (3ml), a cat. amount of activated carbon and diphosgene (15μl, 123μmol) were added. The reaction was heated under reflux for 4 hours then filtered over decalite and concentrated under reduced pressure. The crude isocyanate was dissolved in CH₂Cl₂ (1ml) and added to a stirred solution of 2-methoxybenzylamine (16μl, 123μmol) and Et₃N (17μl, 123μmol) in CH₂Cl₂ (2 ml). The reaction was stirred for 3 days at room temperature. The reaction mixture was quenched with H₂O and acidified with 2N HCl. The product was extracted into CH₂Cl₂, dried and purified by column chromatography on silica to give the title compound (23mg, 77%). Data : (m/z) = 489 (M+H)⁺.

(e) 1-(1-Acetyl-7-amino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(2-methoxy-benzyl)-urea

To a stirred solution of the compound described in example 19 (d) (23mg, 47μmol) in dry THF (2ml), HOAc (27μl, 470μmol) and zinc (62mg, 940μmol) were added. The reaction was stirred at room temperature for 3 hours. The reaction mixture was filtered over decalite and poured into H₂O. The product was extracted into EtOAc. The organics were washed with aqueous NaHCO₃ and brine, dried (Na₂SO₄) and concentrated under reduced pressure to give the title compound (22mg, 100%). Data : (m/z) = 459 (M+H)⁺.

(f) 1-(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(2-methoxy-benzyl)-urea

To a stirred solution of the crude compound described in example 19 (e) (22mg, 47 μ mol), in MeOH (3ml) and HOAc (30 μ l, 588 μ mol), formaldehyde (37% in H₂O, (8 μ l, 94 μ mol)) and NaCNBH₃ (6mg, 94 μ mol) were added. The reaction was stirred overnight at room temperature. The reaction mixture was poured into H₂O and extracted into EtOAc. The organics were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure. The product was purified by column chromatography on silica to give the title compound (15.7mg, 69%) Data : (m/z) = 487(M+H)⁺.

Example 20

(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid thiophen-2-ylmethyl ester

This compound was prepared, in an analogous manner as described in Example 19, from the compound described in example 19 (c) using 2-thiophenemethanol in the first step to afford the title compound (12mg, 56%) Data : (m/z) = 464(M+H)⁺.

Example 21

Biphenyl-4-carboxylic acid (1-acetyl-7-methoxy-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

(a) (2-Methoxy-4-nitro-phenyl)-carbamic acid 9H-fluoren-9-ylmethyl ester

2-methoxy-4-nitroaniline (3g, 17.8 mmol) was dissolved in THF (60 ml) and pyridine (1.6 ml, 19.6 mmol) was added. The reaction mixture was cooled to 0 °C and Fmoc-Cl (5.07 g, 19.6 mmol) was added in small portions. The reaction was allowed to come to ambient temperature and held for 5 hours. The reaction mixture was concentrated under reduced pressure. The crude product was purified by recrystallisation from CH₂Cl₂ and MeOH to afford the title compound (6.08 g, 88 %). Data: (m/z) = 391 (M+H)⁺.

(b) (4-Amino-2-methoxy-phenyl)-carbamic acid 9H-fluoren-9-ylmethyl ester

The compound described in example 21(a) (64.8 g, 0.17 mol) was dissolved in THF (1.5 l) and acetic acid (95 ml, 1.7 mol) was added. The reaction mixture was cooled to 0 °C and zinc (217.1 g, 3.4 mol) was added in small portions. The reaction was allowed to come to ambient temperature and held for 0.5 hour. The reaction mixture was filtered over decalite and the filtrate was washed with a saturated NaHCO₃ solution in H₂O (2x) and brine (1x), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by recrystallisation from CH₂Cl₂ and MeOH to afford the title compound (55.6 g, 93 %). Data: (m/z) = 361 (M+H)⁺.

(c) (7-Methoxy-2,2,4-trimethyl-1,2-dihydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

The compound described in example 21(b) (4.45 g, 12.4 mmol), I₂ (157 mg, 0.62 mmol), MgSO₄ (7.4 g, 62 mmol) and catechol (61 mg, 0.37 mmol) were dissolved/suspended in acetone (350 ml). The reaction mixture was heated to reflux and held for 5h. The reaction was cooled to ambient temperature and filtered over decalite. The filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography on silica to afford the title compound (4.24 g, 78 %). Data: (m/z) = 441 (M+H)⁺.

(d) (1-Acetyl-7-methoxy-2,2,4-trimethyl-1,2-dihydro-quinolin-6-yl)-carbamic acid 9H-fluoren-9-ylmethyl ester

The compound described in example 21 (c) (4.24 g, 9.5 mmol) was dissolved in CH₂Cl₂ (25 ml). First pyridine (25 ml) and a catalytic amount of DMAP were added, and then a solution of acetylchloride (2.0 ml, 28.5 mmol) in CH₂Cl₂ (20 ml) was added dropwise. The reaction mixture was held at ambient temperature for 15 minutes. The reaction mixture was diluted with CH₂Cl₂ and extracted with H₂O (3x), 0.1 M HCl in H₂O (3x), 0.5 M HCl in H₂O (1x), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica to afford the title compound (3.91 g, 85 %). Data: (m/z) = 483 (M+H)⁺.

(e) 1-(6-Amino-7-methoxy-2,2,4-trimethyl-2H-quinolin-1-yl)-ethanone

The compound described in example 21 (d) (236 mg, 0.49 mmol), was dissolved in CH₂Cl₂ (4 ml). Piperidine (485 µl, 4.9 mmol) was added and the reaction mixture was held at ambient temperature for 1 hour. The reaction mixture was concentrated under reduced pressure and coevaporated with toluene (2x). The crude product was purified by column chromatography on silica to afford the title compound (90 mg, 71 %). Data: (m/z) = 261 (M+H)⁺.

(f) Biphenyl-4-carboxylic acid (1-acetyl-7-methoxy-2,2,4-trimethyl-1,2-dihydro-quinolin-6-yl)-amide

The compound described in example 21 (e) (2.22 g, 8.5 mmol) was dissolved in toluene (48 ml) and pyridine (2ml). 4-biphenylcarbonylchloride (2.21 g, 10.2 mmol) was added and the reaction was held at ambient temperature for 3 hours. An extra equivalent of 4-biphenylcarbonylchloride was added and the reaction was held at ambient temperature for 1 hour. The reaction mixture was concentrated under reduced pressure and

coevaporated with toluene (2x). The residue was taken up in EtOAc and extracted with a saturated NaHCO₃ solution in H₂O, H₂O and a 1 M HCl solution in H₂O, dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by recrystallisation from CH₂Cl₂ and MeOH to afford the title compound (3.1 g, 82 %). Data: (m/z) = 441 (M+H)⁺.

(g) Biphenyl-4-carboxylic acid (1-acetyl-7-hydroxy-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

The compound described in example 21 (f) (3.1 g, 7.05 mmol) was dissolved in benzene (100 ml). AlCl₃ (5.6 g, 42.3 mmol) was added and the reaction was held at ambient temperature for 20 hours. The reaction was quenched with H₂O and the reaction mixture was brought to pH 8 by addition of a 2 M NaOH solution in H₂O (63 ml) and extracted. The organics were washed with H₂O and brine, dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by recrystallisation from acetonitril to afford the title compound (195 mg, 5%). Data: (m/z) = 505 (M+H)⁺.

Biphenyl-4-carboxylic acid (1-acetyl-7-methoxy-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

The compound described in example 21 (g) (600 mg, 1.2 mmol) was dissolved in acetonitril (50 ml). K₂CO₃ (821 mg, 5.9 mmol) was added and the reaction mixture was heated to 45 °C and held for 15 minutes. MeI (84 µl, 1.3 mmol) was added and the reaction mixture was held at 45 °C for 3 hours. An extra 0.2 equivalents of MeI were added and the reaction was completed in 0.5 hour. The reaction mixture was concentrated under reduced pressure, taken up in EtOAc and washed with H₂O and brine, dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by recrystallisation from heptane/EtOAc to afford the title compound (351 mg, 56 %). Data: (m/z) = 519 (M+H)⁺.

Example 22

Furan-2-carboxylic acid [1-acetyl-7-methoxy-4-(4-methoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

(a) Furan-2-carboxylic acid (1-acetyl-7-methoxy-2,2,4-trimethyl-1,2-dihydro-quinolin-6-yl)-amide

The compound described in example 21 (e) (406 mg, 1.56 mmol) was dissolved in CH₂-Cl₂ (5 ml). 2-furancarbonylchloride (170 µl, 1.72 mmol) and DIPEA (815 µl, 4.68

mmol) were added and the reaction mixture was held at ambient temperature for 15 hours. The reaction was quenched with H₂O and extracted. The organics were concentrated under reduced pressure and the crude product was purified by column chromatography on silica to afford the title compound (370 mg, 67 %). Data: (m/z) = 355 (M+H)⁺.

Furan-2-carboxylic acid [1-acetyl-7-methoxy-4-(4-methoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

The compound described in example 22 (a) (260 mg, 0.73 mmol) and AlCl₃ (catalytic amount) were dissolved in anisol (5ml) and held at ambient temperature for 15 hours. The reaction mixture was extracted with H₂O and EtOAc. The crude product was purified by column chromatography on silica to afford the title compound (301 mg, 89 %). Data: (m/z) = 463 (M+H)⁺.

Example 23

Biphenyl-4-carboxylic acid [1-acetyl-4-(4-cyanomethoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

(a) Biphenyl-4-carboxylic acid [1-acetyl-4-(4-methoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

This compound was prepared, in an analogous manner as described in Example 22, from the compound described in example 1c to afford the title compound (466 mg, 60 %). Data: (m/z) = 519 (M+H)⁺.

(b) Biphenyl-4-carboxylic acid [1-acetyl-4-(4-hydroxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

The compound described in example 23 (a) (466 mg, 0.9 mmol) was dissolved in CH₂-Cl₂ (7 ml) and the reaction mixture was cooled to 0 °C. BBr₃ (680 mg, 2.7 mmol) was added and the reaction was allowed to come to ambient temperature and held for 3 hours. The reaction mixture was cooled to 0 °C and slowly was added a 1 M NaOH solution in H₂O and EtOAc. This mixture was acidified and extracted. The organics were dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica to afford the title compound (125 mg, 28 %). Data: (m/z) = 505 (M+H)⁺.

Biphenyl-4-carboxylic acid [1-acetyl-4-(4-cyanomethoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide

The compound described in example 23 (b) (118 mg, 0.2 mmol) was dissolved in DMF (5 ml). CsCO_3 (325 mg, 0.84 mmol) and (2-chloro-ethyl)-diethyl-amine hydrochloride (43.3 mg, 0.25 mmol) were added and the reaction mixture was held at ambient temperature for 15 hours. The reaction was quenched with water and extracted with EtOAc. The organics were dried (MgSO_4) and concentrated under reduced pressure. The crude product was purified by preparative HPLC to afford the title compound (61 mg, 51 %). Data: $(m/z) = 596 (M+H)^+$.

Example 24

Biphenyl-4-carboxylic acid {1-acetyl-2,2,4-trimethyl-4-[4-(pyridin-4-ylmethoxy)-phenyl]-1,2,3,4-tetrahydro-quinolin-6-yl}-amide

This compound was prepared, in an analogous manner as described in Example 23, from the compound described in example 23 (b) to afford the title compound (320 mg, 46 %). Data: $(m/z) = 414 (M+H)^+$.

Example 25

[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid furan-2-ylmethyl ester

This compound was prepared, in an analogous manner as described in Example 6, from the compound described in example 15 (a), to afford the title compound (13mg, 47%). Data : $(m/z) = 518 (M+H)^+$.

Example 26

N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide

This compound was prepared, in an analogous manner as described in Example 2, from the compound described in example 1(d) (12 mg, 4%) Data : $(m/z) = 475 (M+H)$.

Example 27

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-methyl-butyl ester

To a stirred solution of the compound described in example 10 (a) (124 mg, 402 μmol) and a catalytic amount of activated charcoal in dry EtOAc (4.6 ml), under an N_2 atmosphere, was added trichloromethyl chloroformate (97 μl , 804 μmol). The reaction was stirred at reflux for 2 hours, then cooled to ambient temperature and filtered over

dikalite and concentrated under reduced pressure to yield a crude oil (134 mg, 100%). This oil (20 mg, 0.065mmol) was dissolved in THF (1 ml) and added to a solution of 3-methyl-1-butanol (70.8 μ l, 0.65 mmol) and TEA (94 μ l, 0.65mmol) in THF (2 ml), under a N₂ atmosphere. The reaction was stirred overnight at ambient temperature, then quenched with water, extracted with EtOAc, washed with brine, dried (Na₂SO₄), filtered and reduced to an oil. The oil was purified by flash chromatography (heptane/EtOAc) to yield the title compound (10.9 mg, 42.5%). Data : (m/z) = 395 (M+H).

Example 28

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 1-methyl-cyclopropylmethyl ester

This compound has been prepared in an analogous manner as described for example 27, using 1-methyl cyclopropane methanol (19.1 mg, 68%). Data : (m/z) = 393 (M+H).

Example 29

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid cyclobutylmethyl ester

This compound has been prepared in an analogous manner as described for example 27, using cyclobutane methanol (11 mg, 40%). Data : (m/z) = 393 (M+H).

Example 30

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid (R)-1,2-dimethyl-propyl ester

This compound has been prepared in an analogous manner as described for example 27, using (R)-(-)-3-methyl-2-butanol (6 mg, 2%). Data : (m/z) = 395 (M+H).

Example 31

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid (1R,2S,4S)-bicyclo[2.2.1]hept-2-yl ester

This compound has been prepared in an analogous manner as described for example 27, using endo-norborneol (8 mg, 27%). Data : (m/z) = 419 (M+H).

Example 32

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4,4a,8a-hexahydro-quinolin-6-yl)-carbamic acid cyclopentyl ester

This compound has been prepared in an analogous manner as described for example 27, using pentenol (46%). Data : (m/z) = 393 (M+H).

Example 33

N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-trifluoromethyl-phenyl)-propionamide

This compound was prepared, in an analogous manner as described for example 10, from the compound described in example 10 (a), to afford the title compound (35 %).

Data: (m/z) = 481 (M+H)⁺.

Example 34

(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-chloro-4-fluoro-benzyl ester

This compound has been prepared in an analogous manner as described for example 27, using (3-chloro-4-fluoro-phenyl)-methanol (73.3%). Data : (m/z) = 467 (M+H).

Example 35

5-Bromo-thiophene-2-carboxylic acid (1-acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide

This compound has been prepared in an analogous manner as described for example 10, using 5-bromo-thiophene-2-carboxylic acid (66%). Data : (m/z) = 470 (M+H).

Example 36

(+)-1-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-benzyl)-urea

This compound has been obtained starting from example 13, via chiral HPLC. Column AD-H (5 μ) 25x0.46 cm. Eluens : heptane/iso-propyl alcohol 90/10. Retention time :

26.8 min. $[\alpha]_D^{20} = + 243^\circ$ (ethyl alcohol, 5 mg/ml)

Example 37

(+)-N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide

This compound has been obtained starting from example 26, via chiral HPLC. Column AD-H (5 μ) 25x0.46 cm. Eluens : heptane/iso-propyl alcohol 80/20. Retention time : 7.0

min. $[\alpha]_D^{20} = + 349^\circ$ (ethyl alcohol, 5 mg/ml)

Example 38

(+)-N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-trifluoromethyl-phenyl)-propionamide

This compound has been obtained starting from example 33, via chiral HPLC. Column AD-H (5 μ) 25x0.46 cm. Eluens : heptane/iso-propyl alcohol 90/10. Retention time : 26.8 min. $[\alpha]_D^{20} = + 229^\circ$ (ethyl alcohol, 5 mg/ml)

Example 39

(+)-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4,4a,8a-hexahydro-quinolin-6-yl)-carbamic acid cyclopentyl ester

This compound has been obtained starting from example 32, via chiral HPLC. Column OD-H (5 μ) 25x0.46 cm. Eluens : heptane/ethyl alcohol 85/15. Retention time : 6.6 min.

$[\alpha]_D^{20} = + 278^\circ$ (ethyl alcohol, 5 mg/ml)

Example 40

(+)-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-chloro-4-fluoro-benzyl ester

This compound has been obtained starting from example 34, via chiral HPLC. Column OD (10 μ) 25x0.46 cm. Eluens : heptane/ethyl alcohol 80/20. Retention time : 8.1 min.

$[\alpha]_D^{20} = + 236^\circ$ (ethyl alcohol, 5 mg/ml)

Example 41

Antagonist activity of compounds against TSH at the human TSH receptor expressed in CHO cells

Antagonist activity of the compounds against TSH at the human TSH receptor was tested in Chinese Hamster Ovary (CHO) cells stably transfected with a plasmid encoding the human TSH receptor and a second plasmid with a cAMP responsive element (CRE) / promotor directing the expression of a firefly luciferase reporter gene. Binding of bovine TSH to the Gs-coupled TSH receptor will result in an increase of cAMP, which in turn will induce an increased transactivation of the luciferase reporter. Luciferase activity was quantified using a luminescence counter. The cells were incubated with the test compounds (concentration between 0.316 nM and 10 μ M)

together with 18 nM bovine TSH (which, at this concentration in the absence of test compound, induced 80% of the maximal luciferase stimulation). The IC_{50} (concentration of test compound causing half-maximal (50 %) inhibition of the maximally attainable inhibition of the luciferase stimulation by the compound) and efficacy of the compounds were determined using the software program XLfit (Excel version 4.1, ID Business Solutions Limited). The compounds described in the preceding examples all have an IC_{50} of less than 10^{-6} M. The examples 12, 13, 23, 18, 1, 4, 20, 7, 6, 11, 26, 27, 28, 29, 30, 33, 31, 32, 34, 35, 36, 37, 38, 39 and 40 show IC_{50} lower than $1E-7$ M.

Example 42

Antagonist activity of compounds against human TSI at the human TSH receptor expressed in CHO cells

Antagonist activity of three test compounds against human TSI at the human TSH receptor was tested in Chinese Hamster Ovary (CHO) cells (cultured 16 h in the absence of serum prior to the assay) stably expressing the human TSH receptor and a CRE-driven firefly luciferase reporter gene. TSI was partially purified from the serum of a GD patient by filtration over a 0.45 μ m filter, protein G-Sepharose column chromatography, dialysis against phosphate-buffered saline and subsequent concentrating on a 10K Amicon filter. It was confirmed that the TSI preparation does not activate luciferase activity in control CHO cells lacking the human TSHR. The cAMP phosphodiesterase inhibitor rolipram was included in the assay medium (10 μ M) to augment TSHR-induced CRE-luciferase synthesis, which was quantified using a luminescence counter. The cells were incubated with compound A, B or C (0.316 nM - 10 μ M) together with 3.16 mg/ml TSI (or bovine TSH at a equieffective concentration of 18 nM). Compound A is hexanoic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide (see example 1). Compound B is an enantiomer of (1-Acetyl-4-methyl-4-phenyl-1,2,3,4,4a,8a-hexahydro-quinolin-6-yl) -carbamic acid cyclopentyl ester (see example 39). Compound C is an enantiomer of (1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-chloro-4-fluorobenzyl ester (see example 40). All 3 compounds are full antagonists and show IC_{50} of less than 10^{-6} M (Table I).

Table I


	IC50 (TSI)	IC50 (bTSH)
Compound A	88 nM	368 nM
Compound B	12 nM	48 nM
Compound C	47 nM	192 nM

Example 43Antagonist activity of compounds at constitutively active human TSH receptors expressed in CHO cells

Antagonist activity of two compounds was tested in Chinese Hamster Ovary (CHO) cells transiently transfected with a plasmid encoding one of the 5 most prevalent, constitutively active human TSHR mutants (Thr632Ile, Ala623Val, Ile568Thr, Asp619Gly and Asp633Glu) (van Sande et al. (1995) J. Clin. Endocrinol. Metabol. **80**, 2577-2585), which have been identified in autonomously functioning thyroid nodules causing hyperthyroidism. The cells were incubated with 10^{-6} M of either the compound of example 35 or example 1 in the presence of 10 μ M rolipram. Activity of the TSH receptor mutants was quantified using a Packard cAMP Alphascreen assay. Both compounds inhibited constitutive activity of all 5 receptor mutants by >80% at a concentration of 10^{-6} M.

Claims

1. Use of the compounds according to general formula I



Formula I

or a pharmaceutically acceptable salt thereof, wherein

R¹ is H or methyl;

X is a bond, O, NH or N((1-4C)alkyl);

R² is (1-4C)alkyl, R⁵(1-4C)alkyl or R⁹(2-4C)alkyl,

if X is NH, R² in addition is R⁸(1-2C)alkoxycarbonyl; or,

if X is a bond, R² in addition is H, halogen or (2-5C)heteroaryl or phenyl, both optionally substituted with one or more substituents selected from (1-3C)alkyl, (1-3C)alkoxy or halogen;

Y is a bond, O or NH, whereas

if Y is a bond, R³ is phenyl, optionally substituted with one or more substituents selected from halogen, hydroxy, methoxy, phenoxy, phenyl, (1-4C)alkyl, nitro, amino or (di)[(1-4C)alkyl]amino or

R³ is R⁶(1-6C)alkyl, R⁶(3-6C)cycloalkyl, R⁷oxy(1-6C)alkyl, or 2-pyridyl or

R³ is a 5-membered heteroaryl, optionally substituted with one or more substituents selected from (1-3C)alkoxy, (1-3C)alkyl or halogen;

if Y is O, R³ is R⁶(1-6C)alkyl, R⁷oxy(2-6C)alkyl or (3-7C)cycloalkyl; and

if Y is NH, R³ is R⁶(1-6C)alkyl, R⁷oxy(2-6C)alkyl;

R⁴ is H, (di)[(1-3C)alkyl]amino or (1-3C)alkoxy;

R⁵ is CN or pyridyl;

R⁶ is H, or (3-5C)cycloalkyl, (2-5C)heteroaryl or phenyl, the latter three groups

optionally substituted with one or more substituents selected from halogen, (1-4C)alkoxy or (1-4C)alkyl, the latter groups both optionally substituted with one or more halogen;

R⁷ is (2-5C)heteroaryl or phenyl, both optionally substituted with one or more substituents selected from halogen, (1-4C)alkoxy or (1-4C)alkyl;

R⁸ is H or (2-5C)heteroaryl or phenyl, both optionally substituted with one or more substituents selected from (1-3C)alkyl, (1-3C)alkoxy or halogen; and

R⁹ is (di)[(1-4C)alkyl]amino or (2-6C)heterocycloalkyl

for the manufacture of a medicament for the treatment or prevention of disorders responsive to TSH receptor mediated pathways.

2. The use according to claim 1 wherein X in Formula I is a bond or O.
3. The use according to claim 2 wherein X is a bond and R² in Formula I is H.
4. The use according to claims 1 and 2 wherein R² in Formula I is R⁵(1-4C)alkyl and X is O.
5. The use according to claims 1-4 wherein R⁴ in Formula I is H, or (di)[(1-3C)alkyl]amino.
6. The use according to claims 1-5 wherein the compound of Formula I has the (+) optical rotation.
7. A compound according to Formula I as described in claims 1-5 which compound has the (+) optical rotation.
8. A compound selected from the group of
 Hexanoic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
 1-Methyl-1*H*-pyrrole-2-carboxylic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-5-bromo-2-methylamino-benzamide,
 Pyridine-2-carboxylic acid (1-acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3,4-dimethyl-benzamide,
 (1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic

acid 2-phenoxy-ethyl ester,
(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid butyl ester,
1-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-cyclopentyl-urea,
Hexanoic acid [1-acetyl-4-(4-bromo-phenyl)-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide,
N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-2-phenoxy-acetamide,
N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide,
N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-thiophen-2-yl-propionamide,
1-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-benzyl)-urea,
Hexanoic acid {1-acetyl-4-[4-(3-fluoro-pyridin-4-yl)-phenyl]-4-methyl-1,2,3,4-tetrahydro-quinolin-6-yl}-amide,
[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid 3-chloro-benzyl ester,
[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid methyl ester,
Cyclopentanecarboxylic acid (1-acetyl-7-dimethylamino-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
N-(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-phenyl)-propionamide,
1-(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(2-methoxy-benzyl)-urea,
(1-Acetyl-7-dimethylamino-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid thiophen-2-ylmethyl ester,
Biphenyl-4-carboxylic acid (1-acetyl-7-methoxy-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
Furan-2-carboxylic acid [1-acetyl-7-methoxy-4-(4-methoxy-phenyl)-2,2,4-

trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide,
Biphenyl-4-carboxylic acid [1-acetyl-4-(4-cyanomethoxy-phenyl)-2,2,4-trimethyl-1,2,3,4-tetrahydro-quinolin-6-yl]-amide,
Biphenyl-4-carboxylic acid {1-acetyl-2,2,4-trimethyl-4-[4-(pyridin-4-ylmethoxy)-phenyl]-1,2,3,4-tetrahydro-quinolin-6-yl}-amide,
[4-(1-Acetyl-6-hexanoylamino-4-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-phenyl]-carbamic acid furan-2-ylmethyl ester,
N-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-methyl-butyl ester,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 1-methyl-cyclopropylmethyl ester,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid cyclobutylmethyl ester,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid (R)-1,2-dimethyl-propyl ester,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid (1R,2S,4S)-bicyclo[2.2.1]hept-2-yl ester,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4,4a,8a-hexahydro-quinolin-6-yl)-carbamic acid cyclopentyl ester,
N-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-trifluoromethyl-phenyl)-propionamide,
(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid 3-chloro-4-fluoro-benzyl ester,
5-Bromo-thiophene-2-carboxylic acid (1-acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-amide,
(+)-1-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(4-chloro-benzyl)-urea,
(+)-*N*-(1-Acetyl-2,2,4-trimethyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-chloro-phenyl)-propionamide,
(+)-*N*-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-3-(3-

trifluoromethyl-phenyl)-propionamide,
(+)-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4,4a,8a-hexahydro-quinolin-6-yl)-
carbamic acid cyclopentyl ester or
(+)-(1-Acetyl-4-methyl-4-phenyl-1,2,3,4-tetrahydro-quinolin-6-yl)-carbamic acid
3-chloro-4-fluoro-benzyl ester.

9. A pharmaceutical composition comprising the compound as defined in claims 1-8 and pharmaceutically acceptable auxiliaries for the treatment or prevention of disorders responsive to the actions of TSH, TSI or constitutive activity of TSH receptor.
10. The use of claims 1-6 wherein the disease, condition, or disorder is selected from the group of hyperthyroidism, Graves' disease, Graves' ophthalmopathy, Graves' associated pretibial dermopathy, nodular goitre and thyroid cancer.
11. A kit for use in treating or preventing disorders responsive to TSH receptor mediated pathways, the kit comprising: A) a pharmaceutical composition comprising a compound of Formula I as defined in claims 1-8 or a pharmaceutically acceptable salt thereof; and B) instructions describing a method of using the pharmaceutical composition to treat or prevent disorders responsive to TSH receptor mediated pathways.