

GENETICALLY MODIFIED MAJOR HISTOCOMPATIBILITY COMPLEX MICE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit of priority to U.S. Provisional Patent Application Nos. 61/552,582 and 61/552,587, both filed October 28, 2011, and U.S. Provisional Patent Application No. 61/700,908, filed September 14, 2012, all of which are hereby incorporated by reference herein in their entireties.

FIELD OF THE INVENTION

[0002] Present invention relates to a genetically modified non-human animal, e.g., a rodent (e.g., a mouse or a rat), that expresses a human or humanized Major Histocompatibility Complex (MHC) class I molecule. The invention also relates to a genetically modified non-human animal, e.g., a mouse or a rat, that expresses a human or humanized MHC I protein (e.g., MHC I α chain) and/or a human or humanized β 2 microglobulin; as well as embryos, tissues, and cells expressing the same. The invention further provides methods for making a genetically modified non-human animal that expresses human or humanized MHC class I protein (e.g., MHC I α chain) and/or β 2 microglobulin. Also provided are methods for identifying and evaluating peptides in the context of a humanized cellular immune system *in vitro* or in a genetically modified non-human animal, and methods of modifying an MHC I and/or a β 2 microglobulin locus of a non-human animal, e.g., a mouse or a rat, to express a human or humanized MHC I and/or β 2 microglobulin.

BACKGROUND OF THE INVENTION

[0003] In the adaptive immune response, foreign antigens are recognized by receptor molecules on B lymphocytes (e.g., immunoglobulins) and T lymphocytes (e.g., T cell

receptor or TCR). These foreign antigens are presented on the surface of cells as peptide fragments by specialized proteins, generically referred to as major histocompatibility complex (MHC) molecules. MHC molecules are encoded by multiple loci that are found as a linked cluster of genes that spans about 4 Mb. In mice, the MHC genes are found on chromosome 17, and for historical reasons are referred to as the histocompatibility 2 (H-2) genes. In humans, the genes are found on chromosome 6 and are called human leukocyte antigen (HLA) genes. The loci in mice and humans are polygenic; they include three highly polymorphic classes of MHC genes (class I, II and III) that exhibit similar organization in human and murine genomes (see FIG. 2 and FIG. 3, respectively).

[0004] MHC loci exhibit the highest polymorphism in the genome; some genes are represented by >300 alleles (e.g., human HLA-DR β and human HLA-B). All class I and II MHC genes can present peptide fragments, but each gene expresses a protein with different binding characteristics, reflecting polymorphisms and allelic variants. Any given individual has a unique range of peptide fragments that can be presented on the cell surface to B and T cells in the course of an immune response.

[0005] Both humans and mice have class I MHC genes (see FIG. 2 and FIG. 3). In humans, the classical class I genes are termed HLA-A, HLA-B and HLA-C, whereas in mice they are H-2K, H-2D and H-2L. Class I molecules consist of two chains: a polymorphic α -chain (sometimes referred to as heavy chain) and a smaller chain called β 2-microglobulin (also known as light chain), which is generally not polymorphic (FIG. 1). These two chains form a non-covalent heterodimer on the cell surface. The α -chain contains three domains (α 1, α 2 and α 3). Exon 1 of the α -chain gene encodes the leader sequence, exons 2 and 3 encode the α 1 and α 2 domains, exon 4 encodes the α 3 domain, exon 5 encodes the transmembrane domain, and exons 6 and 7 encode the cytoplasmic tail. The α -chain forms a peptide-binding cleft involving the α 1 and α 2 domains (which resemble Ig-like domains) followed by the α 3 domain, which is similar to β 2-microglobulin.

[0006] β 2 microglobulin is a non-glycosylated 12 kDa protein; one of its functions is to stabilize the MHC class I α -chain. Unlike the α -chain, the β 2 microglobulin does not span the membrane. The human β 2 microglobulin locus is on chromosome 15, while the mouse locus is on chromosome 2. β 2 microglobulin gene consists of 4 exons and 3 introns. Circulating forms of β 2 microglobulin are present in the serum, urine, and other body fluids; thus, the non-covalently MHC I-associated β 2 microglobulin can be exchanged with circulating β 2 microglobulin under physiological conditions.

[0007] Class I MHC molecules are expressed on all nucleated cells, including tumor cells. They are expressed specifically on T and B lymphocytes, macrophages, dendritic cells and neutrophils, among other cells, and function to display peptide fragments (typically 8-10 amino acids in length) on the surface to CD8+ cytotoxic T lymphocytes (CTLs). CTLs are specialized to kill any cell that bears an MHC I-bound peptide recognized by its own membrane-bound TCR. When a cell displays peptides derived from cellular proteins not normally present (e.g., of viral, tumor, or other non-self origin), such peptides are recognized by CTLs, which become activated and kill the cell displaying the peptide.

[0008] Typically, presentation of normal (i.e., self) proteins in the context of MHC I molecules does not elicit CTL activation due to the tolerance mechanisms. However, in some diseases (e.g., cancer, autoimmune diseases) peptides derived from self-proteins become a target of the cellular component of the immune system, which results in destruction of cells presenting such peptides. Although there has been advancement in recognizing some self-derived antigens that elicit cellular immune response (e.g., antigens associated with various cancers), in order to improve identification of peptides recognized by human CTLs through MHC class I molecules there remains a need for both *in vivo* and *in vitro* systems that mimic aspects of the human cellular immune system. Systems that mimic the human cellular immune system can be used in identifying disease-associated antigens in order to develop human therapeutics, e.g., vaccines and other biologics. Systems for assessing antigen recognition in the context of the human immune system can assist in identifying therapeutically useful CTL populations (e.g., useful for studying and combatting human disease). Such systems can also assist in enhancing the activity of human CTL populations to more effectively combat infections and foreign antigen-bearing entities. Thus, there is a need for biological systems (e.g., genetically engineered animals) that can generate an immune system that displays components that mimic the function of human immune system.

SUMMARY OF THE INVENTION

[0009] A biological system for generating or identifying peptides that associate with human MHC class I proteins and chimeras thereof, and bind to CD8+ T cells, is provided. Non-human animals comprising non-human cells that express human or humanized molecules that function in the cellular immune response are provided. Humanized rodent loci that encode human or human or humanized MHC I and $\beta 2$ microglobulin proteins are also provided. Humanized rodent cells that express human or humanized MHC and $\beta 2$ microglobulin molecules are also provided. *In vivo* and *in vitro* systems are provided that comprise humanized rodent cells, wherein the rodent cells express one or more human or humanized immune system molecules.

[0010] Provided herein is a non-human animal, e.g., a rodent (e.g., a mouse or a rat), comprising in its genome a nucleotide sequence encoding a chimeric human/non-human (e.g., human/rodent, e.g., human/mouse or human/rat) MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human MHC I polypeptide. Specifically, provided herein is a non-human animal comprising at an endogenous MHC I locus a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human MHC I polypeptide, and wherein the animal expresses the chimeric human/non-human MHC I polypeptide. In one aspect, the animal does not express an extracellular domain of an endogenous non-human MHC I polypeptide from an endogenous non-human MHC I locus. In one aspect of the invention, the non-human animal (e.g., a rodent, e.g., a mouse or a rat) comprises two copies of the MHC I locus comprising a nucleotide sequence encoding chimeric human/non-human (e.g., human/rodent, e.g., human/mouse or human/rat) MHC I polypeptide. In another aspect of the invention, the animal comprises one copy of the MHC I locus comprising a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide. Thus, the animal may be homozygous or heterozygous for the MHC I locus comprising a nucleotide sequence encoding chimeric human/non-human MHC I polypeptide. In various embodiments, the nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide is comprised in the germline of the non-human animal (e.g., rodent, e.g., rat or mouse).

[0011] In one aspect, the nucleotide sequence encoding the chimeric human/non-human MHC I is operably linked to endogenous non-human regulatory elements, e.g., promoter, enhancer, silencer, etc. In one embodiment, a human portion of the chimeric polypeptide comprises a human leader sequence. In an additional embodiment, the human portion of the chimeric polypeptide comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of the human MHC I polypeptide. The human MHC I polypeptide may be selected from a group consisting of HLA-A, HLA-B, and HLA-C. In one embodiment, the human MHC I polypeptide is an HLA-A2 polypeptide, e.g., an HLA-A2.1 polypeptide.

[0012] In one aspect, the genetically engineered non-human animal is a rodent. In one embodiment, the rodent is a mouse. Thus, in one embodiment, the endogenous non-human locus is a mouse locus, e.g., a mouse H-2K, H-2D or H-2L locus. In one embodiment, the non-human portion of the chimeric human/non-human MHC I polypeptide comprises transmembrane and cytoplasmic domains of the endogenous non-human MHC I polypeptide. Thus, in an embodiment wherein the non-human animal is a mouse, the endogenous non-human MHC I locus may be an H-2K locus (e.g., H-2Kb locus) and the endogenous non-human MHC I polypeptide may be an H-2K polypeptide; therefore, the

chimeric human/non-human MHC I polypeptide may comprise transmembrane and cytoplasmic domains of H-2K polypeptide. In another embodiment wherein the non-human animal is a mouse, the endogenous non-human MHC I locus may be an H-2D locus and the endogenous non-human MHC I polypeptide may be an H-2D polypeptide; therefore, the chimeric human/non-human MHC I polypeptide may comprise transmembrane and cytoplasmic domains of H-2D polypeptide. Similarly, in another embodiment, the endogenous non-MHC I locus may be an H-2L locus and the endogenous non-human MHC I polypeptide may be an H-2L polypeptide; therefore, the chimeric human/non-human MHC I polypeptide may comprise transmembrane and cytoplasmic domains of H-2L polypeptide.

[0013] Also provided herein is a mouse comprising at an endogenous H-2K locus a nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human HLA-A (e.g., HLA-A2) polypeptide and a mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K polypeptide, and wherein the mouse expresses the chimeric human/mouse MHC I polypeptide. In some embodiments, the mouse does not express an extracellular domain of the mouse H-2K polypeptide from an endogenous H-2K locus. In one aspect, the nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide is operably linked to endogenous mouse regulatory elements. The human portion of the chimeric polypeptide may comprise a human leader sequence. It may also comprise $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of the human MHC I polypeptide. The human MHC I polypeptide may be HLA-A polypeptide, e.g., HLA-A2.1 polypeptide. In one aspect, the mouse H-2K locus is an H-2Kb locus.

[0014] Another aspect of the invention relates to a non-human animal, e.g., a rodent (e.g., a mouse or a rat), comprising in its genome a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. Thus, provided herein is a non-human animal comprising at an endogenous non-human $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide, wherein the animal expresses the human or humanized $\beta 2$ microglobulin polypeptide. In one aspect, the animal does not express a functional endogenous non-human $\beta 2$ microglobulin polypeptide from an endogenous non-human $\beta 2$ microglobulin locus. In one aspect, the animal comprises two copies of the $\beta 2$ microglobulin locus encoding the human or humanized $\beta 2$ microglobulin polypeptide; in another embodiment, the animal comprises one copy of the $\beta 2$ microglobulin locus encoding the human or humanized $\beta 2$ microglobulin polypeptide. Thus, the animal may be homozygous or heterozygous for the $\beta 2$ microglobulin locus encoding the human or humanized $\beta 2$ microglobulin polypeptide. In various embodiments, the nucleotide sequence

encoding the human or humanized $\beta 2$ microglobulin polypeptide is comprised in the germline of the non-human animal (e.g., rodent, e.g., rat or mouse). In one embodiment, a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide comprises a nucleotide sequence encoding a polypeptide comprising a human $\beta 2$ microglobulin amino acid sequence. In one embodiment, the polypeptide is capable of binding to an MHC I protein.

[0015] In some embodiments, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide is operably linked to endogenous non-human $\beta 2$ microglobulin regulatory elements. In one aspect, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide comprises a nucleotide sequence set forth in exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. In another aspect, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide comprises nucleotide sequences set forth in exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. In a further aspect, the nucleotide sequence also comprises a nucleotide sequence set forth in exon 1 of a non-human $\beta 2$ microglobulin gene. In some embodiments, the non-human animal is a rodent (e.g., mouse or a rat); thus, the non-human $\beta 2$ microglobulin locus is a rodent (e.g., a mouse or a rat) $\beta 2$ microglobulin locus.

[0016] Also provided is a mouse comprising at an endogenous $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide, wherein the mouse expresses the human or humanized $\beta 2$ microglobulin polypeptide. In some embodiments, the mouse does not express a functional endogenous mouse $\beta 2$ microglobulin from an endogenous $\beta 2$ microglobulin locus. The nucleotide sequence may be linked to endogenous mouse regulatory elements. In one aspect, the nucleotide sequence comprises a nucleotide sequence set forth in exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. Alternatively, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide may comprise nucleotide sequences set forth in exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. The nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide may further comprise a nucleotide sequence of exon 1 of a mouse $\beta 2$ microglobulin gene. In one embodiment, a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide comprises a nucleotide sequence encoding a polypeptide comprising a human $\beta 2$ microglobulin amino acid sequence. In one embodiment, the polypeptide is capable of binding to an MHC I protein.

[0017] The invention further provides a non-human animal (e.g., a rodent, e.g., a mouse or a rat) comprising in its genome a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide and a nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide. In one embodiment, the invention provides a non-human animal comprising in its genome a first nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human MHC I polypeptide; and a second nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide, wherein the first nucleotide sequence is located at an endogenous non-human MHC I locus, and the second nucleotide sequence is located at an endogenous non-human β 2 microglobulin locus, and wherein the animal expresses the chimeric human/non-human MHC I polypeptide and the human or humanized β 2 microglobulin polypeptide. In one aspect, the animal is a mouse. Thus, the endogenous MHC I locus may be selected from a group consisting of H-2K, H-2D, and H-2L locus. In one embodiment, the endogenous mouse locus is an H-2K locus (e.g., H-2Kb locus). In one embodiment, the human MHC I polypeptide is selected from the group consisting of HLA-A, HLA-B, and HLA-C polypeptide. In one aspect, the human MHC I polypeptide is HLA-A, e.g., HLA-A2 (e.g., HLA-A2.1). In various embodiments, the first and the second nucleotide sequences are comprised in the germline of the non-human animal (e.g., rodent, e.g., mouse or rat).

[0018] Therefore, the invention provides a mouse comprising in its genome a first nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human HLA-A (e.g., HLA-A2) and a mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K; and a second nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide, wherein the first nucleotide sequence is located at an endogenous H-2K locus and the second nucleotide sequence is located at an endogenous mouse β 2 microglobulin locus, and wherein the mouse expresses the chimeric human/mouse MHC I polypeptide and the human or humanized β 2 microglobulin polypeptide. In one embodiment, the non-human animal (e.g., the mouse) comprising both the chimeric MHC I polypeptide and human or humanized β 2 microglobulin polypeptide does not express an extracellular domain of an endogenous non-human MHC I polypeptide (e.g., the mouse H-2K polypeptide) and/or a functional endogenous non-human (e.g., the mouse) β 2 microglobulin polypeptides from their respective endogenous loci. In one aspect, the animal (e.g., the mouse) comprises two copies of each of the first and the second nucleotide sequence. In another aspect, the animal (e.g., the mouse) comprises one copy of the first

and one copy of the second nucleotide sequences. Thus, the animal may be homozygous or heterozygous for both the first and the second nucleotide sequences.

[0019] In one aspect, the first nucleotide sequence is operably linked to endogenous non-human (e.g., mouse) MHC I regulatory elements, and the second nucleotide sequence is operably linked to endogenous non-human (e.g., mouse) $\beta 2$ microglobulin elements. The human portion of the chimeric polypeptide may comprise $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the human MHC I polypeptide. The second nucleotide sequence may comprise a nucleotide sequence set forth in exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. Alternatively, the second nucleotide sequence may comprise nucleotide sequences set forth in exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. In one aspect, the mouse comprising both the chimeric MHC I polypeptide and human or humanized $\beta 2$ microglobulin polypeptide may be such that the expression of human or humanized $\beta 2$ microglobulin increases the expression of the chimeric human/mouse MHC I polypeptide as compared to the expression of the chimeric human/mouse MHC I polypeptide in the absence of expression of human or humanized $\beta 2$ microglobulin polypeptide.

[0020] Also provided are methods of making genetically engineered non-human animals (e.g., rodents, e.g., mice or rats) described herein. Thus, in one embodiment, provided is a method of modifying an MHC I locus of a rodent (e.g., a mouse or a rat) to express a chimeric human/rodent (e.g., human/mouse or human/rat) MHC I polypeptide, wherein the method comprises replacing at the endogenous MHC I locus a nucleotide sequence encoding an extracellular domain of a rodent MHC I polypeptide with a nucleotide sequence encoding an extracellular domain of a human MHC I polypeptide. In another embodiment, provided is a method of modifying a $\beta 2$ microglobulin locus of a rodent (e.g., a mouse or a rat) to express a human or humanized $\beta 2$ microglobulin polypeptide, wherein the method comprises replacing at the endogenous rodent (e.g., mouse or rat) $\beta 2$ microglobulin locus a nucleotide sequence encoding a rodent (e.g., a mouse or a rat) $\beta 2$ microglobulin polypeptide with a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In such methods, the replacement may be made in a single ES cell, and the single ES cell may be introduced into a rodent (e.g., a mouse or a rat) to make an embryo. The resultant rodent (e.g., a mouse or a rat) can be bred to generate a double humanized animal.

[0021] Thus, the invention also provides a method of making double humanized animals, e.g., rodents (e.g., mice or rats). In one embodiment, provided is a method of making a genetically modified mouse comprising (a) modifying an MHC I locus of a first mouse to express a chimeric human/mouse MHC I polypeptide comprising replacing at the endogenous mouse MHC I locus a nucleotide sequence encoding an extracellular domain of

a mouse MHC I polypeptide with a nucleotide sequence encoding an extracellular domain of a human MHC I polypeptide, (b) modifying a $\beta 2$ microglobulin locus of a second mouse to express a human or humanized $\beta 2$ microglobulin polypeptide comprising replacing at the endogenous mouse $\beta 2$ microglobulin locus a nucleotide sequence encoding a mouse $\beta 2$ microglobulin polypeptide with a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide; and (c) breeding the first and the second mouse to generate a genetically modified mouse comprising in its genome a first nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide and a second nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide, wherein the genetically modified mouse expresses the chimeric human/mouse MHC I polypeptide and the human or humanized $\beta 2$ microglobulin polypeptide. In some embodiments, the MHC I locus is selected from H-2K, H-2D, and H-2L; in some embodiments, the human MHC I polypeptide is selected from HLA-A, HLA-B, and HLA-C. In one embodiment, the MHC I locus is an H-2K locus, the human MHC I polypeptide is HLA-A (e.g., HLA-A2), and the mouse expresses a chimeric HLA-A/H-2K polypeptide (e.g., HLA-A2/H-2K polypeptide). In one aspect, the chimeric HLA-A2/H-2K polypeptide comprises an extracellular domain of the HLA-A2 polypeptide and cytoplasmic and transmembrane domains of H-2K polypeptide. In one aspect, the second nucleotide sequence comprises nucleotide sequences set forth in exons 2, 3, and 4 (e.g., exon 2 to exon 4) of a human $\beta 2$ microglobulin gene, and a nucleotide sequence set forth in exon 1 of a mouse $\beta 2$ microglobulin gene.

[0022] Also provided herein are cells, e.g., isolated antigen-presenting cells, derived from the non-human animals (e.g., rodents, e.g., mice or rats) described herein. Tissues and embryos derived from the non-human animals described herein are also provided.

[0023] In yet another embodiment, the invention provides methods for identification of antigens or antigen epitopes that elicit immune response, methods for evaluating a vaccine candidate, methods for identification of high affinity T cells to human pathogens or cancer antigens.

[0024] Any of the embodiments and aspects described herein can be used in conjunction with one another, unless otherwise indicated or apparent from the context. Other embodiments will become apparent to those skilled in the art from a review of the ensuing detailed description. The following detailed description includes exemplary representations of various embodiments of the invention, which are not restrictive of the invention as claimed. The accompanying figures constitute a part of this specification and, together with the description, serve only to illustrate embodiments and not to limit the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] FIG. 1 is a schematic drawing of the four domains of a class I MHC molecule: α -chain containing the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains and the non-covalently associated fourth domain, $\beta 2$ -microglobulin ($\beta 2m$). The gray circle represents a peptide bound in the peptide-binding cleft.

[0026] FIG. 2 is a schematic representation (not to scale) of the relative genomic structure of the human HLA, showing class I, II and III genes.

[0027] FIG. 3 is a schematic representation (not to scale) of the relative genomic structure of the mouse MHC, showing class I, II and III genes.

[0028] FIG. 4 illustrates a viral vector construct containing a cDNA encoding a chimeric HLA-A/H-2K polypeptide with an IRES-GFP reporter (A); and histograms comparing expression of human HLA-A2 in MG87 cells transduced with HLA-A2 (dashed line), HLA-A2/H-2K (dotted line), or no transduction (solid line) either alone (left) or co-transduced with humanized $\beta 2$ microglobulin (right) (B). Data from horizontal gates presented graphically in (B) is illustrated as percent of cells expressing the construct in the table in (C).

[0029] FIG. 5 is a schematic diagram (not to scale) of the targeting strategy used for making a chimeric H-2K locus that expresses an extracellular region of a human HLA-A2 protein. Mouse sequences are represented in black and human sequences are represented in white. L=leader, UTR=untranslated region, TM=transmembrane domain, CYT=cytoplasmic domain, HYG=hygromycin.

[0030] FIG. 6A demonstrates expression (% total cells) of HLA-A2 (left) and H-2K (right) in cells isolated from either a wild-type (WT) mouse or a heterozygous mouse carrying the chimeric HLA-A2/H-2K locus (HLA-A/H-2K HET).

[0031] FIG. 6B is a dot plot of *in vivo* expression of the chimeric HLA-A2/H-2K protein in a heterozygous mouse harboring a chimeric HLA-A2/H-2K locus.

[0032] FIG. 7 shows a targeting strategy (not to scale) for humanization of a $\beta 2$ microglobulin gene at a mouse $\beta 2$ microglobulin locus. Mouse sequences are in black and human sequences are in white. NEO=neomycin.

[0033] FIG. 8 shows a representative dot plot of HLA class I and human $\beta 2$ microglobulin expression on cells isolated from the blood of wild-type (WT) mice, mice heterozygous for chimeric HLA-A2/H-2K, and mice heterozygous for chimeric HLA-A2/H-2K and heterozygous for humanized $\beta 2$ microglobulin (double heterozygous; class I/ $\beta 2m$ HET).

[0034] FIG. 9 shows a representative histogram of human HLA class I expression (X axis) on cells isolated from the blood of wild-type (WT), chimeric HLA-A2/H-2K heterozygous (class I HET), and chimeric HLA-A2/H2K/humanized β 2 microglobulin double heterozygous (class I/ β 2m HET) mice.

[0035] FIG. 10 shows the results of IFN γ Elispot assays for human T cells exposed to antigen-presenting cells (APCs) from wild-type mice (WT APCs) or mice heterozygous for both chimeric HLA-A2/H-2K and humanized β 2 microglobulin (double HET APCs) in the presence of flu (left) or EBV (right) peptides. Statistical analysis was performed using one way ANOVA with a Tukey's Multiple Comparison Post Test.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0036] The present invention provides genetically modified non-human animals (e.g., mice, rats, rabbits, etc.) that express human or humanized MHC I and/or β 2 microglobulin polypeptides; embryos, cells, and tissues comprising the same; methods of making the same; as well as methods of using the same. Unless defined otherwise, all terms and phrases used herein include the meanings that the terms and phrases have attained in the art, unless the contrary is clearly indicated or clearly apparent from the context in which the term or phrase is used.

[0037] The term "conservative," when used to describe a conservative amino acid substitution, includes substitution of an amino acid residue by another amino acid residue having a side chain R group with similar chemical properties (e.g., charge or hydrophobicity). Conservative amino acid substitutions may be achieved by modifying a nucleotide sequence so as to introduce a nucleotide change that will encode the conservative substitution. In general, a conservative amino acid substitution will not substantially change the functional properties of interest of a protein, for example, the ability of MHC I to present a peptide of interest. Examples of groups of amino acids that have side chains with similar chemical properties include aliphatic side chains such as glycine, alanine, valine, leucine, and isoleucine; aliphatic-hydroxyl side chains such as serine and threonine; amide-containing side chains such as asparagine and glutamine; aromatic side chains such as phenylalanine, tyrosine, and tryptophan; basic side chains such as lysine, arginine, and histidine; acidic side chains such as aspartic acid and glutamic acid; and, sulfur-containing side chains such as cysteine and methionine. Conservative amino acids substitution groups include, for example, valine/leucine/isoleucine, phenylalanine/tyrosine, lysine/arginine, alanine/valine, glutamate/aspartate, and asparagine/glutamine. In some embodiments, a conservative amino acid substitution can be a substitution of any native residue in a protein with alanine,

as used in, for example, alanine scanning mutagenesis. In some embodiments, a conservative substitution is made that has a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet et al. ((1992) Exhaustive Matching of the Entire Protein Sequence Database, Science 256:1443-45), hereby incorporated by reference. In some embodiments, the substitution is a moderately conservative substitution wherein the substitution has a nonnegative value in the PAM250 log-likelihood matrix.

[0038] Thus, also encompassed by the invention is a genetically modified non-human animal whose genome comprises a nucleotide sequence encoding a human or humanized MHC I polypeptide and/or $\beta 2$ microglobulin polypeptide, wherein the polypeptide(s) comprises conservative amino acid substitutions of the amino acid sequence(s) described herein.

[0039] One skilled in the art would understand that in addition to the nucleic acid residues encoding a human or humanized MHC I polypeptide and/or $\beta 2$ microglobulin described herein, due to the degeneracy of the genetic code, other nucleic acids may encode the polypeptide(s) of the invention. Therefore, in addition to a genetically modified non-human animal that comprises in its genome a nucleotide sequence encoding MHC I and/or $\beta 2$ microglobulin polypeptide(s) with conservative amino acid substitutions, a non-human animal whose genome comprises a nucleotide sequence(s) that differs from that described herein due to the degeneracy of the genetic code is also provided.

[0040] The term "identity" when used in connection with sequence includes identity as determined by a number of different algorithms known in the art that can be used to measure nucleotide and/or amino acid sequence identity. In some embodiments described herein, identities are determined using a ClustalW v. 1.83 (slow) alignment employing an open gap penalty of 10.0, an extend gap penalty of 0.1, and using a Gonnet similarity matrix (MacVector™ 10.0.2, MacVector Inc., 2008). The length of the sequences compared with respect to identity of sequences will depend upon the particular sequences. In various embodiments, identity is determined by comparing the sequence of a mature protein from its N-terminal to its C-terminal. In various embodiments when comparing a chimeric human/non-human sequence to a human sequence, the human portion of the chimeric human/non-human sequence (but not the non-human portion) is used in making a comparison for the purpose of ascertaining a level of identity between a human sequence and a human portion of a chimeric human/non-human sequence (e.g., comparing a human ectodomain of a chimeric human/mouse protein to a human ectodomain of a human protein).

[0041] The terms "homology" or "homologous" in reference to sequences, e.g., nucleotide or amino acid sequences, means two sequences which, upon optimal alignment and comparison, are identical in at least about 75% of nucleotides or amino acids, at least about 80% of nucleotides or amino acids, at least about 90-95% nucleotides or amino acids, e.g., greater than 97% nucleotides or amino acids. One skilled in the art would understand that, for optimal gene targeting, the targeting construct should contain arms homologous to endogenous DNA sequences (i.e., "homology arms"); thus, homologous recombination can occur between the targeting construct and the targeted endogenous sequence.

[0042] The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. As such, a nucleic acid sequence encoding a protein may be operably linked to regulatory sequences (e.g., promoter, enhancer, silencer sequence, etc.) so as to retain proper transcriptional regulation. In addition, various portions of the chimeric or humanized protein of the invention may be operably linked to retain proper folding, processing, targeting, expression, and other functional properties of the protein in the cell. Unless stated otherwise, various domains of the chimeric or humanized proteins of the invention are operably linked to each other.

[0043] The term "MHC I complex" or the like, as used herein, includes the complex between the MHC I α chain polypeptide and the β 2-microglobulin polypeptide. The term "MHC I polypeptide" or the like, as used herein, includes the MHC I α chain polypeptide alone. Typically, the terms "human MHC" and "HLA" can be used interchangeably.

[0044] The term "replacement" in reference to gene replacement refers to placing exogenous genetic material at an endogenous genetic locus, thereby replacing all or a portion of the endogenous gene with an orthologous or homologous nucleic acid sequence. As demonstrated in the Examples below, nucleic acid sequences of endogenous loci encoding portions of mouse MHC I and β 2 microglobulin polypeptides were replaced by nucleotide sequences encoding portions of human MHC I and β 2 microglobulin polypeptides, respectively.

[0045] "Functional" as used herein, e.g., in reference to a functional polypeptide, refers to a polypeptide that retains at least one biological activity normally associated with the native protein. For example, in some embodiments of the invention, a replacement at an endogenous locus (e.g., replacement at an endogenous non-human MHC I and/or β 2 microglobulin locus) results in a locus that fails to express a functional endogenous polypeptide.

[0046] Several aspects described herein below for the genetically modified MHC I non-human animals, e.g., animal type; animal strains; cell types; screening, detection and other methods; methods of use; etc., will be applicable to the genetically engineered $\beta 2$ microglobulin and MHC I/ $\beta 2$ microglobulin animals.

Genetically Modified MHC I Animals

[0047] In various embodiments, the invention generally provides genetically modified non-human animals that comprise in their genome a nucleotide sequence encoding a human or humanized MHC I polypeptide; thus, the animals express a human or humanized MHC I polypeptide.

[0048] MHC genes are categorized into three classes: class I, class II, and class III, all of which are encoded either on human chromosome 6 or mouse chromosome 17. A schematic of the relative organization of the human and mouse MHC classes is presented in FIGs. 2 and 3, respectively. The MHC genes are among the most polymorphic genes of the mouse and human genomes. MHC polymorphisms are presumed to be important in providing evolutionary advantage; changes in sequence can result in differences in peptide binding that allow for better presentation of pathogens to cytotoxic T cells.

[0049] MHC class I protein comprises an extracellular domain (which comprises three domains: α_1 , α_2 , and α_3), a transmembrane domain, and a cytoplasmic tail. The α_1 and α_2 domains form the peptide-binding cleft, while the α_3 interacts with $\beta 2$ -microglobulin.

[0050] In addition to its interaction with $\beta 2$ -microglobulin, the α_3 domain interacts with the TCR co-receptor CD8, facilitating antigen-specific activation. Although binding of MHC class I to CD8 is about 100-fold weaker than binding of TCR to MHC class I, CD8 binding enhances the affinity of TCR binding. Wooldridge et al. (2010) MHC Class I Molecules with Superenhanced CD8 Binding Properties Bypass the Requirement for Cognate TCR Recognition and Nonspecifically Activate CTLs, J. Immunol. 184:3357-3366. Interestingly, increasing MHC class I binding to CD8 abrogated antigen specificity in CTL activation. *Id.*

[0051] CD8 binding to MHC class I molecules is species-specific; the mouse homolog of CD8, Lyt-2, was shown to bind H-2D^d molecules at the α_3 domain, but it did not bind HLA-A molecules. Connolly et al. (1988) The Lyt-2 Molecule Recognizes Residues in the Class I α_3 Domain in Allogeneic Cytotoxic T Cell Responses, J. Exp. Med. 168:325-341. Differential binding was presumably due to CDR-like determinants (CDR1- and CDR2-like) on CD8 that was not conserved between humans and mice. Sanders et al. (1991) Mutations in CD8 that Affect Interactions with HLA Class I and Monoclonal Anti-CD8 Antibodies, J. Exp. Med. 174:371-379; Vitiello et al. (1991) Analysis of the HLA-restricted Influenza-specific

Cytotoxic T Lymphocyte Response in Transgenic Mice Carrying a Chimeric Human-Mouse Class I Major Histocompatibility Complex, J. Exp. Med. 173:1007-1015; and, Gao et al. (1997) Crystal structure of the complex between human CD8 $\alpha\alpha$ and HLA-A2, Nature 387:630-634. It has been reported that CD8 binds HLA-A2 in a conserved region of the α_3 domain (at position 223-229). A single substitution (V245A) in HLA-A reduced binding of CD8 to HLA-A, with a concomitant large reduction in T cell-mediated lysis. Salter et al. (1989), Polymorphism in the α_3 domain of HLA-A molecules affects binding to CD8, Nature 338:345-348. In general, polymorphism in the α_3 domain of HLA-A molecules also affected binding to CD8. *Id.* In mice, amino acid substitution at residue 227 in H-2D^d affected the binding of mouse Lyt-2 to H-2D^d, and cells transfected with a mutant H-2D^d were not lysed by CD8+ T cells. Potter et al. (1989) Substitution at residue 227 of H-2 class I molecules abrogates recognition by CD8-dependent, but not CD8-independent, cytotoxic T lymphocytes, Nature 337:73-75.

[0052] Therefore, due to species specificity of interaction between the MHC class I α_3 domain and CD8, an MHC I complex comprising a replacement of an H-2K α_3 domain with a human HLA-A2 α_3 domain was nonfunctional in a mouse (*i.e., in vivo*) in the absence of a human CD8. In animals transgenic for HLA-A2, substitution of human α_3 domain for the mouse α_3 domain resulted in restoration of T cell response. Irwin et al. (1989) Species-restricted interactions between CD8 and the α_3 domain of class I influence the magnitude of the xenogeneic response, J. Exp. Med. 170:1091-1101; Vitiello et al. (1991), *supra*.

[0053] The transmembrane domain and cytoplasmic tail of mouse MHC class I proteins also have important functions. One function of MHC I transmembrane domain is to facilitate modulation by HLA-A2 of homotypic cell adhesion (to enhance or inhibit adhesion), presumably as the result of cross-linking (or ligation) of surface MHC molecules. Wagner et al. (1994) Ligation of MHC Class I and Class II Molecules Can Lead to Heterologous Desensitization of Signal Transduction Pathways That Regulate Homotypic Adhesion in Human Lymphocytes, J. Immunol. 152:5275-5287. Cell adhesion can be affected by mAbs that bind at diverse epitopes of the HLA-A2 molecule, suggesting that there are multiple sites on HLA-A2 implicated in modulating homotypic cell adhesion; depending on the epitope bound, the affect can be to enhance or to inhibit HLA-A2-dependent adhesion. *Id.*

[0054] The cytoplasmic tail, encoded by exons 6 and 7 of the MHC I gene, is reportedly necessary for proper expression on the cell surface and for LIR1-mediated inhibition of NK cell cytotoxicity. Gruda et al. (2007) Intracellular Cysteine Residues in the Tail of MHC Class I Proteins Are Crucial for Extracellular Recognition by Leukocyte Ig-Like Receptor 1, J. Immunol. 179:3655-3661. A cytoplasmic tail is required for multimerization of at least some

MHC I molecules through formation of disulfide bonds on its cysteine residues, and thus may play a role in clustering and in recognition by NK cells. Lynch *et al.* (2009) Novel MHC Class I Structures on Exosomes, *J. Immunol.* 183:1884-1891.

[0055] The cytoplasmic domain of HLA-A2 contains a constitutively phosphorylated serine residue and a phosphorylatable tyrosine, although—in Jurkat cells—mutant HLA-A2 molecules lacking a cytoplasmic domain appear normal with respect to expression, cytoskeletal association, aggregation, and endocytic internalization. Gur *et al.* (1997) Structural Analysis of Class I MHC Molecules: The Cytoplasmic Domain Is Not Required for Cytoskeletal Association, Aggregation, and Internalization, *Mol. Immunol.* 34(2):125-132. Truncated HLA-A2 molecules lacking the cytoplasmic domain are apparently normally expressed and associate with β 2 microglobulin. *Id.*

[0056] However, several studies have demonstrated that the cytoplasmic tail is critical in intracellular trafficking, dendritic cell (DC)-mediated antigen presentation, and CTL priming. A tyrosine residue encoded by exon 6 was shown to be required for MHC I trafficking through endosomal compartments, presentation of exogenous antigens, and CTL priming; while deletion of exon 7 caused enhancement of anti-viral CTL responses. Lizee *et al.* (2003) Control of Dendritic Cross-Presentation by the Major Histocompatibility Complex Class I Cytoplasmic Domain, *Nature Immunol.* 4:1065-73; Basha *et al.* (2008) MHC Class I Endosomal and Lysosomal Trafficking Coincides with Exogenous Antigen Loading in Dendritic Cells, *PLoS ONE* 3: e3247; and Rodriguez-Cruz *et al.* (2011) Natural Splice Variant of MHC Class I Cytoplasmic Tail Enhances Dendritic Cell-Induced CD8⁺ T-Cell Responses and Boosts Anti-Tumor Immunity, *PLoS ONE* 6:e22939.

[0057] In various embodiments, the invention provides a genetically modified non-human animal (e.g., mouse, rat, rabbit, etc.) that comprises in its genome a nucleotide sequence encoding a human or humanized MHC class I polypeptide. The non-human animal may comprise in its genome a nucleotide sequence that encodes an MHC I polypeptide that is partially human and partially non-human, e.g., a non-human animal that expresses a chimeric human/non-human MHC I polypeptide. In one aspect, the non-human animal only expresses the human or humanized MHC I polypeptide, e.g., chimeric human/non-human MHC I polypeptide, and does not express an endogenous non-human MHC I protein from an endogenous MHC I locus.

[0058] In one embodiment, the chimeric human/non-human MHC I polypeptide comprises in its human portion a peptide binding domain of a human MHC I polypeptide. In one aspect, the human portion of the chimeric polypeptide comprises an extracellular domain of a human MHC I. In this embodiment, the human portion of the chimeric

polypeptide comprises an extracellular domain of an α chain of a human MHC I. In one embodiment, the human portion of the chimeric polypeptide comprises $\alpha 1$ and $\alpha 2$ domains of a human MHC I. In another embodiment, the human portion of the chimeric polypeptide comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a human MHC I.

[0059] The human or humanized MHC I polypeptide may be derived from a functional human HLA molecule encoded by any of HLA-A, HLA-B, HLA-C, HLA-E, HLA-F, or HLA-G loci. A list of commonly used HLA antigens is described in Shankarkumar et al. ((2004) The Human Leukocyte Antigen (HLA) System, Int. J. Hum. Genet. 4(2):91-103), incorporated herein by reference. Shankarkumar et al. also present a brief explanation of HLA nomenclature used in the art. Additional information regarding HLA nomenclature and various HLA alleles can be found in Holdsworth et al. (2009) The HLA dictionary 2008: a summary of HLA-A, -B, -C, -DRB1/3/4/5, and DQB1 alleles and their association with serologically defined HLA-A, -B, -C, -DR, and -DQ antigens, Tissue Antigens 73:95-170, and a recent update by Marsh et al. (2010) Nomenclature for factors of the HLA system, 2010, Tissue Antigens 75:291-455, both incorporated herein by reference. Thus, the human or humanized MHC I polypeptide may be derived from any functional human HLA class I molecules described therein.

[0060] In one specific aspect, the human or humanized MHC I polypeptide is derived from human HLA-A. In a specific embodiment, the HLA-A polypeptide is an HLA-A2 polypeptide (e.g., and HLA-A2.1 polypeptide). In one embodiment, the HLA-A polypeptide is a polypeptide encoded by an HLA-A*0201 allele, e.g., HLA-A*02:01:01:01 allele. The HLA-A*0201 allele is commonly used amongst the North American population. Although the present Examples describe this particular HLA sequence, any suitable HLA-A sequence is encompassed herein, e.g., polymorphic variants of HLA-A2 exhibited in human population, sequences with one or more conservative or non-conservative amino acid modifications, nucleic acid sequences differing from the sequence described herein due to the degeneracy of genetic code, etc.

[0061] In one aspect, a non-human animal that expresses a human HLA-A2 sequence is provided, wherein the human HLA-A2 sequence comprises one or more conservative or non-conservative modifications.

[0062] In one aspect, a non-human animal that expresses a human HLA-A2 sequence is provided, wherein the human HLA-A2 sequence is at least about 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a human HLA-A2 sequence. In a specific embodiment, the human HLA-A2 sequence is at least about 90%, 95%, 96%, 97%, 98%, or 99% identical to the human HLA-A2 sequence described in the Examples. In one embodiment, the human

HLA-A2 sequence comprises one or more conservative substitutions. In one embodiment, the human HLA-A2 sequence comprises one or more non-conservative substitutions.

[0063] In another specific aspect, the human or humanized MHC I polypeptide is derived from human MHC I selected from HLA-B and HLA-C. In one aspect, the human or humanized MHC I is derived from HLA-B, e.g., HLA-B27.

[0064] In one aspect, the non-human portion of the chimeric human/non-human MHC I polypeptide comprises transmembrane and/or cytoplasmic domains of the non-human MHC I polypeptide. In one embodiment, the non-human animal is a mouse, and the non-human MHC I polypeptide is selected from H-2K, H-2D, and H-2L. In one embodiment, the non-human MHC I polypeptide is H-2K, e.g., H-2Kb. Although specific H-2K sequences are described in the Examples, any suitable H-2K sequences, e.g., polymorphic variants, conservative/non-conservative amino acid substitutions, etc., are encompassed herein.

[0065] The non-human animal described herein may comprise in its genome a nucleotide sequence encoding a human or humanized MHC I polypeptide, e.g., chimeric human/non-human MHC I polypeptide, wherein the nucleotide sequence encoding such polypeptide is located at an endogenous non-human MHC I locus (e.g., H-2K locus). In one aspect, this results in a replacement of an endogenous MHC I gene or a portion thereof with a nucleotide sequence encoding a human or humanized MHC I polypeptide, e.g., a chimeric gene encoding a chimeric human/non-human MHC I polypeptide described herein. In one embodiment, the replacement comprises a replacement of an endogenous nucleotide sequence encoding a non-human MHC I peptide binding domain or a non-human MHC I extracellular domain with a human nucleotide sequence (e.g., HLA-A2 nucleotide sequence) encoding the same. In this embodiment, the replacement does not comprise a replacement of an MHC I sequence encoding transmembrane and/or cytoplasmic domains of a non-human MHC I polypeptide (e.g., H-2K polypeptide). Thus, the non-human animal contains chimeric human/non-human nucleotide sequence at an endogenous non-human MHC I locus, and expresses chimeric human/non-human MHC polypeptide from the endogenous non-human MHC I locus.

[0066] A chimeric human/non-human polypeptide may be such that it comprises a human or a non-human leader (signal) sequence. In one embodiment, the chimeric polypeptide comprises a non-human leader sequence of an endogenous MHC I protein. In another embodiment, the chimeric polypeptide comprises a leader sequence of a human MHC I protein, e.g., HLA-A2 protein (for instance, HLA-A2.1 leader sequence). Thus, the nucleotide sequence encoding the chimeric MHC I polypeptide may be operably linked to a nucleotide sequence encoding a human MHC I leader sequence.

[0067] A chimeric human/non-human MHC I polypeptide may comprise in its human portion a complete or substantially complete extracellular domain of a human MHC I polypeptide. Thus, the human portion may comprise at least 80%, preferably at least 85%, more preferably at least 90%, e.g., 95% or more of the amino acids encoding an extracellular domain of a human MHC I polypeptide (e.g., HLA-A2 polypeptide). In one example, substantially complete extracellular domain of the human MHC I polypeptide lacks a human MHC I leader sequence. In another example, the chimeric human/non-human MHC I polypeptide comprises a human MHC I leader sequence.

[0068] Moreover, the chimeric MHC I polypeptide may be expressed under the control of endogenous non-human regulatory elements, e.g., rodent MHC I regulatory animals. Such arrangement will facilitate proper expression of the chimeric MHC I polypeptide in the non-human animal, e.g., during immune response in the non-human animal.

[0069] The genetically modified non-human animal may be selected from a group consisting of a mouse, rat, rabbit, pig, bovine (e.g., cow, bull, buffalo), deer, sheep, goat, chicken, cat, dog, ferret, primate (e.g., marmoset, rhesus monkey). For the non-human animals where suitable genetically modifiable ES cells are not readily available, other methods are employed to make a non-human animal comprising the genetic modification. Such methods include, e.g., modifying a non-ES cell genome (e.g., a fibroblast or an induced pluripotent cell) and employing nuclear transfer to transfer the modified genome to a suitable cell, e.g., an oocyte, and gestating the modified cell (e.g., the modified oocyte) in a non-human animal under suitable conditions to form an embryo.

[0070] In one aspect, the non-human animal is a mammal. In one aspect, the non-human animal is a small mammal, e.g., of the superfamily Dipodoidea or Muroidea. In one embodiment, the genetically modified animal is a rodent. In one embodiment, the rodent is selected from a mouse, a rat, and a hamster. In one embodiment, the rodent is selected from the superfamily Muroidea. In one embodiment, the genetically modified animal is from a family selected from Calomyscidae (e.g., mouse-like hamsters), Cricetidae (e.g., hamster, New World rats and mice, voles), Muridae (true mice and rats, gerbils, spiny mice, crested rats), Nesomyidae (climbing mice, rock mice, with-tailed rats, Malagasy rats and mice), Platacanthomyidae (e.g., spiny dormice), and Spalacidae (e.g., mole rates, bamboo rats, and zokors). In a specific embodiment, the genetically modified rodent is selected from a true mouse or rat (family Muridae), a gerbil, a spiny mouse, and a crested rat. In one embodiment, the genetically modified mouse is from a member of the family Muridae. In one embodiment, the animal is a rodent. In a specific embodiment, the rodent is selected from a mouse and a rat. In one embodiment, the non-human animal is a mouse.

[0071] In a specific embodiment, the non-human animal is a rodent that is a mouse of a C57BL strain selected from C57BL/A, C57BL/An, C57BL/GrFa, C57BL/KaLwN, C57BL/6, C57BL/6J, C57BL/6ByJ, C57BL/6NJ, C57BL/10, C57BL/10ScSn, C57BL/10Cr, and C57BL/Ola. In another embodiment, the mouse is a 129 strain selected from the group consisting of a strain that is 129P1, 129P2, 129P3, 129X1, 129S1 (e.g., 129S1/SV, 129S1/SvIm), 129S2, 129S4, 129S5, 129S9/SvEvH, 129S6 (129/SvEvTac), 129S7, 129S8, 129T1, 129T2 (see, e.g., Festing *et al.* (1999) Revised nomenclature for strain 129 mice, Mammalian Genome 10:836, see also, Auerbach et al (2000) Establishment and Chimera Analysis of 129/SvEv- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines). In a specific embodiment, the genetically modified mouse is a mix of an aforementioned 129 strain and an aforementioned C57BL/6 strain. In another specific embodiment, the mouse is a mix of aforementioned 129 strains, or a mix of aforementioned BL/6 strains. In a specific embodiment, the 129 strain of the mix is a 129S6 (129/SvEvTac) strain. In another embodiment, the mouse is a BALB strain, e.g., BALB/c strain. In yet another embodiment, the mouse is a mix of a BALB strain and another aforementioned strain.

[0072] In one embodiment, the non-human animal is a rat. In one embodiment, the rat is selected from a Wistar rat, an LEA strain, a Sprague Dawley strain, a Fischer strain, F344, F6, and Dark Agouti. In one embodiment, the rat strain is a mix of two or more strains selected from the group consisting of Wistar, LEA, Sprague Dawley, Fischer, F344, F6, and Dark Agouti.

[0073] Thus, in one embodiment, the invention relates to a genetically modified mouse that comprises in its genome a nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises a peptide binding domain or an extracellular domain of a human MHC I (e.g., human HLA-A, e.g., human HLA-A2, e.g., human HLA-A2.1). In some embodiments, the mouse does not express a peptide binding or an extracellular domain of an endogenous mouse polypeptide from its endogenous mouse locus. The peptide binding domain of the human MHC I may comprise $\alpha 1$ and $\alpha 2$ domains. Alternatively, the peptide binding domain of the human MHC I may comprise $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains. In one aspect, the extracellular domain of the human MHC I comprises an extracellular domain of a human MHC I α chain. In one embodiment, the endogenous mouse locus is an H-2K (e.g., H-2Kb) locus, and the mouse portion of the chimeric polypeptide comprises transmembrane and cytoplasmic domains of a mouse H-2K (e.g., H-2Kb) polypeptide.

[0074] Thus, in one embodiment, a genetically modified mouse is provided, wherein the mouse comprises at an endogenous H-2K (e.g., H-2Kb) locus a nucleotide sequence

encoding a chimeric human/mouse MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human HLA-A2 (e.g., HLA-A2.1) polypeptide and a mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K (e.g., H-2Kb) polypeptide. In one aspect, the mouse does not express an extracellular domain of the mouse H-2K (e.g., H-2Kb) polypeptide from an endogenous MHC I locus. In one embodiment, the mouse expresses a chimeric HLA-A2/H-2K (e.g., a chimeric HLA-A2.1/H-2Kb) polypeptide from an endogenous H-2K (e.g., H-2Kb) locus. In various embodiments, expression of the chimeric gene is under control of endogenous mouse MHC class I regulatory elements. In some aspects, the mouse comprises two copies of the chimeric MHC I locus containing a nucleotide sequence encoding a chimeric HLA-A2/H-2K polypeptide; while in other aspects, the mouse comprises one copy of the chimeric MHC I locus containing a nucleotide sequence encoding a chimeric HLA-A2/H-2K polypeptide. Thus, the mouse may be homozygous or heterozygous for the nucleotide sequence encoding the chimeric HLA-A2/H-2K polypeptide.

[0075] In some embodiments described herein, a mouse is provided that comprises a chimeric MHC I locus located at an endogenous mouse H-2K locus. The chimeric locus comprises a nucleotide sequence that encodes an extracellular domain of a human HLA-A2 protein, e.g., $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a human HLA-A2 gene. The chimeric locus lacks a nucleotide sequence that encodes an extracellular domain of a mouse H-2K protein (e.g., $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of the mouse H-2K). In one aspect, the chimeric locus lacks a nucleotide sequence that encodes a leader peptide, $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a mouse H-2K; and comprises a leader peptide, $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a human HLA-A2, and transmembrane and cytoplasmic domains of a mouse H-2K. The various domains of the chimeric locus are operably linked to each other such that the chimeric locus expresses a functional chimeric human/mouse MHC I protein.

[0076] In various embodiments, a non-human animal, e.g., a rodent (e.g., a mouse or a rat), that expresses a functional chimeric MHC I protein from a chimeric MHC I locus as described herein displays the chimeric protein on a cell surface. In one embodiment, the non-human animal expressed the chimeric MHC I protein on a cell surface in a cellular distribution that is the same as observed in a human. In one aspect, the cell displays a peptide fragment (an antigen fragment) bound to an extracellular portion (e.g., human HLA-A2 extracellular portion) of the chimeric MHC I protein. In an embodiment, the extracellular portion of such chimeric protein interacts with other proteins on the surface of said cell, e.g., $\beta 2$ -microglobulin.

[0077] In various embodiments, a cell displaying the chimeric MHC I protein, e.g., HLA-A2/H-2K protein, is a nucleated cell. In various aspects, the cell is an antigen-presenting cell (APC). Although most cells in the body can present an antigen in the context of MHC I, some nonlimiting examples of antigen presenting cells include macrophages, dendritic cells, and B cells. Other antigen presenting cells, including professional and nonprofessional APCs, are known in the art, and are encompassed herein. In some embodiments, the cell displaying the chimeric MHC I protein is a tumor cell, and a peptide fragment presented by the chimeric protein is derived from a tumor. In other embodiments, the peptide fragment presented by the chimeric MHC I protein is derived from a pathogen, e.g., a bacterium or a virus.

[0078] The chimeric MHC I protein described herein may interact with other proteins on the surface of the same cell or a second cell. In some embodiments, the chimeric MHC I protein interacts with endogenous non-human proteins on the surface of said cell. The chimeric MHC I protein may also interact with human or humanized proteins on the surface of the same cell or a second cell.

[0079] On the same cell, HLA class I molecules may functionally interact with both non-human (e.g., rodent, e.g., mouse or rat) and human β 2-microglobulin. Thus, in one embodiment, the chimeric MHC I protein, e.g., HLA-A2/H-2K protein, interacts with a mouse β 2-microglobulin. Although interaction between some human HLA class I molecules and mouse β 2-microglobulin is possible, it nevertheless may be greatly reduced in comparison to interaction between human HLA class I and human β 2-microglobulin. Thus, in the absence of human β 2-microglobulin, expression of human MHC I on the cell surface may be reduced. Perarnau et al. (1988) Human β 2-microglobulin Specifically Enhances Cell-Surface Expression of HLA Class I Molecules in Transfected Murine Cells, *J. Immunol.* 141:1383-89. Other HLA molecules, e.g., HLA-B27, do not interact with mouse β 2-microglobulin; see, e.g., Tishon et al. (2000) Transgenic Mice Expressing Human HLA and CD8 Molecules Generate HLA-Restricted Measles Virus Cytotoxic T Lymphocytes of the Same Specificity as Humans with Natural Measles Virus Infection, *Virology* 275:286-293, which reports that HLA-B27 function in transgenic mice requires both human β 2-microglobulin and human CD8. Therefore, in another embodiment, the chimeric MHC I protein interacts with a human or humanized β 2-microglobulin. In some such embodiments, as described herein below, the non-human animal, e.g., a rodent (e.g., a mouse or a rat), comprises in its genome a human or humanized β 2-microglobulin gene, and the animal expresses a functional human or humanized β 2-microglobulin polypeptide; therefore, the chimeric MHC I protein interacts with a human or humanized β 2-microglobulin polypeptide.

[0080] In various aspects, the chimeric protein (e.g., HLA-A2/H-2K protein) also interacts with proteins on the surface of a second cell (through its extracellular portion). The second cell may be a cell of a non-human, e.g., a mouse, or a human origin. The second cell may be derived from the same non-human animal or the same non-human animal specie as the cell expressing the chimeric MHC I polypeptide. Nonlimiting examples of proteins with which the extracellular portion of a chimeric protein (e.g., HLA-A2/H-2K) may interact include T cell receptor (TCR) and its co-receptor CD8. Thus, a second cell may be a T cell. In addition, the extracellular portion of the chimeric MHC I protein may bind a protein on the surface of Natural Killer (NK) cells, e.g., killer immunoglobulin receptors (KIRs) on the surface of NK cells.

[0081] A T cell or NK cell may bind a complex formed between the chimeric MHC I polypeptide and its displayed peptide fragment. Such binding may result in T cell activation or inhibition of NK-mediated cell killing, respectively. One hypothesis is that NK cells have evolved to kill either infected or tumor cells that have evaded T cell mediated cytotoxicity by downregulating their MHC I complex. However, when MHC I complex is expressed on cell surface, NK cell receptors recognize it, and NK-mediated cell killing is inhibited. Thus, in some aspects, when an NK cell binds a complex formed between the chimeric MHC I polypeptide (e.g., HLA-A2/H-2K polypeptide) and a displayed peptide fragment on the surface of infected or tumor cell, the NK-mediated cell killing is inhibited.

[0082] In one example, the chimeric MHC I polypeptide described herein, e.g., a chimeric HLA-A2/H-2K polypeptide, interacts with CD8 protein on the surface of a second cell. In one embodiment, the chimeric HLA-A2/H-2K polypeptide interacts with endogenous rodent (e.g., mouse or rat) CD8 protein on the surface of a second cell. In one embodiment, the second cell is a T cell. In another embodiment, the second cell is engineered to express CD8. In certain aspects, the chimeric HLA-A2/H-2K polypeptide interacts with a human CD8 on the surface of the second cell (e.g., a human cell or a rodent cell). In some such embodiments, the non-human animal, e.g., a mouse or a rat, comprises a human CD8 transgene, and the mouse or the rat expresses a functional human CD8 protein.

[0083] The chimeric MHC I polypeptide described herein may also interact with a non-human (e.g., a mouse or a rat) TCR, a human TCR, or a humanized TCR on a second cell. The chimeric MHC I polypeptide may interact with an endogenous TCR (e.g., mouse or rat TCR) on the surface of a second cell. The chimeric MHC I polypeptide may also interact with a human or humanized TCR expressed on the surface of a second cell, wherein the cell is derived from the same animal or animal specie (e.g., mouse or rat) as the cell expressing the chimeric MHC I polypeptide. The chimeric MHC I polypeptide may interact with a human TCR expressed on the surface of a human cell.

[0084] In addition to genetically engineered non-human animals, a non-human embryo (e.g., a rodent embryo, e.g., mouse or a rat embryo) is also provided, wherein the embryo comprises a donor ES cell that is derived from a non-human animal (e.g., a rodent, e.g., a mouse or a rat) as described herein. In one aspect, the embryo comprises an ES donor cell that comprises the chimeric MHC I gene, and host embryo cells.

[0085] Also provided is a tissue, wherein the tissue is derived from a non-human animal (e.g., a mouse or a rat) as described herein, and expresses the chimeric MHC I polypeptide (e.g., HLA-A2/H-2K polypeptide).

[0086] In addition, a non-human cell isolated from a non-human animal as described herein is provided. In one embodiment, the cell is an ES cell. In one embodiment, the cell is an antigen-presenting cell, e.g., dendritic cell, macrophage, B cell. In one embodiment, the cell is an immune cell. In one embodiment, the immune cell is a lymphocyte.

[0087] Also provided is a non-human cell comprising a chromosome or fragment thereof of a non-human animal as described herein. In one embodiment, the non-human cell comprises a nucleus of a non-human animal as described herein. In one embodiment, the non-human cell comprises the chromosome or fragment thereof as the result of a nuclear transfer.

[0088] In one aspect, a non-human induced pluripotent cell comprising gene encoding a chimeric MHC I polypeptide (e.g., HLA-A2/H-2K polypeptide) as described herein is provided. In one embodiment, the induced pluripotent cell is derived from a non-human animal as described herein.

[0089] In one aspect, a hybridoma or quadroma is provided, derived from a cell of a non-human animal as described herein. In one embodiment, the non-human animal is a mouse or rat.

[0090] Also provided is a method for making a genetically engineered non-human animal (e.g., a genetically engineered rodent, e.g., a mouse or a rat) described herein. The method for making a genetically engineered non-human animal results in the animal whose genome comprises a nucleotide sequence encoding a chimeric MHC I polypeptide. In one embodiment, the method results in a genetically engineered mouse, whose genome comprises at an endogenous MHC I locus, e.g., H-2K locus, a nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide, wherein a human portion of the chimeric MHC I polypeptide comprises an extracellular domain of a human HLA-A2 and a mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K. In some embodiments, the method utilizes a targeting construct made using VELOCIGENE® technology, introducing the construct into ES cells, and introducing targeted ES cell clones

into a mouse embryo using VELOCIMOUSE® technology, as described in the Examples. In one embodiment, the ES cells are a mix of 129 and C57BL/6 mouse strains; in another embodiment, the ES cells are a mix of BALB/c and 129 mouse strains.

[0091] Thus, a nucleotide construct used for generating genetically engineered non-human animals described herein is also provided. In one aspect, the nucleotide construct comprises: 5' and 3' non-human homology arms, a human DNA fragment comprising human HLA-A gene sequences, and a selection cassette flanked by recombination sites. In one embodiment, the human DNA fragment is a genomic fragment that comprises both introns and exons of a human HLA-A gene. In one embodiment, the non-human homology arms are homologous to a non-human MHC class I locus (e.g., a mouse H-2K locus).

[0092] In one embodiment, the genomic fragment comprises a human HLA-A leader, an $\alpha 1$ domain, an $\alpha 2$ domain and an $\alpha 3$ domain coding sequence. In one embodiment, the human DNA fragment comprises, from 5' to 3': an HLA-A leader sequence, an HLA-A leader/ $\alpha 1$ intron, an HLA-A $\alpha 1$ exon, an HLA-A $\alpha 1$ - $\alpha 2$ intron, an HLA-A $\alpha 2$ exon, an HLA-A $\alpha 2$ - $\alpha 3$ intron, and an HLA-A $\alpha 3$ exon.

[0093] A selection cassette is a nucleotide sequence inserted into a targeting construct to facilitate selection of cells (e.g., ES cells) that have integrated the construct of interest. A number of suitable selection cassettes are known in the art. Commonly, a selection cassette enables positive selection in the presence of a particular antibiotic (e.g., Neo, Hyg, Pur, CM, Spec, etc.). In addition, a selection cassette may be flanked by recombination sites, which allow deletion of the selection cassette upon treatment with recombinase enzymes. Commonly used recombination sites are *loxP* and *Frt*, recognized by Cre and Flp enzymes, respectively, but others are known in the art.

[0094] In one embodiment, the selection cassette is located at the 5' end the human DNA fragment. In another embodiment, the selection cassette is located at the 3' end of the human DNA fragment. In another embodiment, the selection cassette is located within the human DNA fragment. In another embodiment, the selection cassette is located within an intron of the human DNA fragment. In another embodiment, the selection cassette is located within the $\alpha 2$ - $\alpha 3$ intron.

[0095] In one embodiment, the 5' and 3' non-human homology arms comprise genomic sequence at 5' and 3' locations of an endogenous non-human (e.g., murine) MHC class I gene locus, respectively (e.g., 5' of the first leader sequence and 3' of the $\alpha 3$ exon of the non-human MHC I gene). In one embodiment, the endogenous MHC class I locus is selected from mouse H-2K, H-2D and H-2L. In a specific embodiment, the endogenous MHC class I locus is mouse H-2K.

[0096] In one aspect, a nucleotide construct is provided, comprising, from 5' to 3': a 5' homology arm containing mouse genomic sequence 5' of the endogenous mouse H-2K locus, a first human DNA fragment comprising a first genomic sequence of an HLA-A gene, a 5' recombination sequence site (e.g., *loxP*), a selection cassette, a 3' recombination sequence site (e.g., *loxP*), a second human DNA fragment comprising a second genomic sequence of an HLA-A gene and a 3' homology arm containing mouse genomic sequence 3' of an endogenous H-2K $\alpha 3$ exon. In one embodiment, the nucleotide construct comprises, from 5' to 3': a 5' homology arm containing mouse genomic sequence 5' of the endogenous mouse H-2K locus, a human genomic sequence including an HLA-A leader, an HLA-A leader/ $\alpha 1$ intron sequence, an HLA-A $\alpha 1$ exon, an HLA-A $\alpha 1$ - $\alpha 2$ intron, an HLA-A $\alpha 2$ exon, a first 5' portion of an $\alpha 2$ - $\alpha 3$ intron, a selection cassette flanked by recombination sites, a second 3' portion of an $\alpha 2$ - $\alpha 3$ intron, an HLA-A $\alpha 3$ exon, and a 3' homology arm containing non-mouse genomic sequence 3' of the endogenous mouse H-2K $\alpha 3$ exon. In one embodiment, a 5' homology arm sequence is set forth in SEQ ID NO:1, and a 3' homology arm sequence is set forth in SEQ ID NO:2.

[0097] Upon completion of gene targeting, ES cells or genetically modified non-human animals are screened to confirm successful incorporation of exogenous nucleotide sequence of interest or expression of exogenous polypeptide. Numerous techniques are known to those skilled in the art, and include (but are not limited to) Southern blotting, long PCR, quantitative PCT (e.g., real-time PCR using TAQMAN®), fluorescence *in situ* hybridization, Northern blotting, flow cytometry, Western analysis, immunocytochemistry, immunohistochemistry, etc. In one example, non-human animals (e.g., mice) bearing the genetic modification of interest can be identified by screening for loss of mouse allele and/or gain of human allele using a modification of allele assay described in Valenzuela *et al.* (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech.* 21(6):652-659. Other assays that identify a specific nucleotide or amino acid sequence in the genetically modified animals are known to those skilled in the art.

[0098] The disclosure also provides a method of modifying an MHC I locus of a non-human animal to express a chimeric human/non-human MHC I polypeptide described herein. In one embodiment, the invention provides a method of modifying an MHC I locus of a mouse to express a chimeric human/mouse MHC I polypeptide wherein the method comprises replacing at an endogenous MHC I locus a nucleotide sequence encoding a peptide binding domain of a mouse MHC polypeptide with a nucleotide sequence encoding a peptide binding domain of a human MHC I polypeptide. In some aspects, a nucleotide sequence of an extracellular domain of a mouse MHC I is replaced by a nucleotide

sequence of an extracellular domain of a human MHC I. The mouse may fail to express the peptide binding or the extracellular domain of the mouse MHC I from an endogenous MHC I locus. In some embodiments, a nucleotide sequence of an extracellular domain of a mouse H-2K is replaced by a nucleotide sequence of an extracellular domain of a human HLA-A2, such that the modified mouse MHC I locus expresses a chimeric HLA-A2/H-2K polypeptide.

[0099] In one aspect, a method for making a chimeric human HLA class I/non-human MHC class I molecule is provided, comprising expressing in a single cell a chimeric HLA-A/H-2K protein from a nucleotide construct, wherein the nucleotide construct comprises a cDNA sequence that encodes an $\alpha 1$, $\alpha 2$, and $\alpha 3$ domain of an HLA-A protein and a transmembrane and cytoplasmic domain of a non-human H-2K protein, e.g., mouse H-2K protein. In one embodiment, the nucleotide construct is a viral vector; in a specific embodiment, the viral vector is a lentiviral vector. In one embodiment, the cell is selected from a CHO, COS, 293, HeLa, and a retinal cell expressing a viral nucleic acid sequence (e.g., a PERC.6™ cell).

[00100] In one aspect, a cell that expresses a chimeric human HLA Class I/non-human MHC I protein (e.g., HLA-A/H-2K protein) is provided. In one embodiment, the cell comprises an expression vector comprising a chimeric MHC class I gene, wherein the chimeric MHC class I gene comprises a sequence of a human HLA-A gene fused in operable linkage with a sequence of a non-human H-2K gene, e.g., mouse H-2K gene. In one embodiment, the sequence of the human HLA-A gene comprises the exons that encode $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of an HLA-A protein. In one embodiment, the sequence of the non-human H-2K gene comprises the exons that encode transmembrane and cytoplasmic domains of an H-2K protein. In one embodiment, the cell is selected from CHO, COS, 293, HeLa, and a retinal cell expressing a viral nucleic acid sequence (e.g., a PERC.6™ cell).

[00101] A chimeric MHC class I molecule made by a non-human animal as described herein is also provided, wherein the chimeric MHC class I molecule comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains from a human HLA-A protein and transmembrane and cytoplasmic domains from a non-human, e.g., mouse, H-2K protein. The chimeric MHC I polypeptide described herein may be detected by anti-HLA-A antibodies. Thus, a cell displaying chimeric human/non-human MHC I polypeptide may be detected and/or selected using anti-HLA-A antibody. In some instances, the chimeric MHC I polypeptide described herein may be detected by an anti-HLA-A2 antibody.

[00102] Although the following Examples describe a genetically engineered animal whose genome comprises a replacement of a nucleotide sequence encoding an extracellular domain of mouse H-2K polypeptide with the sequence encoding an extracellular domain of a

human HLA-A at the endogenous mouse H-2K locus, one skilled in the art would understand that a similar strategy may be used to replace other mouse MHC I loci (H-2D, H-2L, etc.) with their corresponding human HLA loci (HLA-B, HLA-C, etc.). Thus, a non-human animal comprising in its genome a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide wherein a human portion of the polypeptide is derived from another HLA class I protein is also provided. The replacement of multiple MHC I loci is also contemplated.

Genetically Modified β 2 Microglobulin Animals

[00103] The invention generally provides genetically modified non-human animals that comprise in their genome a nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide; thus, the animals express a human or humanized β 2 microglobulin polypeptide.

[00104] β 2 microglobulin or the light chain of the MHC class I complex (also abbreviated " β 2M") is a small (12 kDa) non-glycosylated protein, that functions primarily to stabilize the MHC I α chain. The human β 2 microglobulin gene encodes a protein of 119 amino acids, with 20 N-terminal amino acids encoding a leader sequence. The mature protein comprises 99 amino acids. The gene contains 4 exons, with the first exon containing the 5' untranslated region, the entire leader sequence and the first two amino acids of the mature polypeptide; the second exon encoding the majority of the mature protein; the third exon encoding the last four amino acids of the mature protein and a stop codon; and the fourth exon containing the 3' non-translated region. Gussow et al. (1987) The β 2-Microglobulin Gene. Primary Structure and Definition of the Transcriptional Unit, J. Immunol. 139:3131-38. β 2 microglobulin is non-covalently associated with MHC I. Unbound β 2 microglobulin is found in body fluids, such as plasma, and is carried to the kidney for excretion. Kidney dysfunction causes accumulation of β 2 microglobulin, which can be pathogenic (e.g., Dialysis Related Amyloidosis); the accumulated protein forms filamentous fibrils resembling amyloid plaques in joints and connective tissues.

[00105] In addition to Dialysis Related Amyloidosis, β 2 microglobulin has been implicated in a number of other disorders. Elevated levels of β 2 microglobulin were detected in lymphocytic malignancies, e.g., non-Hodgkin's lymphoma and multiple myeloma. See, e.g., Shi et al. (2009) β 2 Microglobulin: Emerging as a Promising Cancer Therapeutic Target, Drug Discovery Today 14:25-30. Some other malignancies with elevated levels of β 2 microglobulin include breast cancer, prostate cancer, lung cancer, renal cancer, gastrointestinal and nasopharyngeal cancers. Overexpression of β 2 microglobulin has been

suggested to have tumor growth promoting effects. *Id.* It has also been recently shown that β 2 microglobulin drives epithelial to mesenchymal transition, promoting cancer bone and soft tissue metastasis in breast, prostate, lung and renal cancers. Jossion et al. (2011) β 2 microglobulin Induces Epithelial to Mesenchymal Transition and Confers Cancer Lethality and Bone Metastasis in Human Cancer Cells. *Cancer Res.* 71(7): 1-11. β 2 microglobulin interacts with a non-classical MHC I member, hemochromatosis (HFE) protein, and with the transferrin receptor, and modulates iron homeostasis. *Id.* Involvement of β 2 microglobulin in other hallmarks of malignancy (self-renewal, angiogenesis enhancement, resistance to treatment) is widely documented in the art.

[00106] Mice deficient in β 2 microglobulin have been reported. See, Koller et al. (1990) Normal development of mice deficient in β 2m, MHC class I proteins, and CD8+ T cells, *Science* 248: 1227-1230. As reported in Koller et al., these mice appeared healthy, however, MHC class I expression was not detected. Further, most T cell populations appeared normal in some tissues, while a marked decrease of CD8+ T cells was observed in others. This purported lack of MHC I expression disagrees with previous results obtained by Allen et al. ((1986) β 2 microglobulin Is Not Required for Cell Surface Expression of the Murine Class I Histocompatibility Antigen H-2D^b or of a Truncated H-2D^b, *Proc. Natl. Acad. Sci. USA* 83:7447-7451). Allen et al. reported that β 2 microglobulin was not absolutely required for cell surface expression of all MHC I complexes, because cells lacking β 2 microglobulin were able to express H-2D^b. However, the function of H-2D^b in these cells was presumably compromised, and conformation of H-2D^b was different from the native protein, which explains the inability of Koller and colleagues to detect this protein using antibodies against native H-2D^b. However, cells lacking β 2 microglobulin can reportedly present endogenous antigen to CD8+ T cells (including exogenous CD8+ T cells from normal mice), and β 2 microglobulin is reportedly not required in order to develop high levels of H-2^d MHC class I-restricted CD8+ CTLs in response to antigen challenge in mice, although it is required in order to sustain an effective immune response. Quinn et al. (1997) Virus-Specific, CD8+ Major Histocompatibility Complex Class I-Restricted Cytotoxic T Lymphocytes in Lymphocytic Choriomeningitis Virus-Infected β 2-Microglobulin-Deficient Mice, *J. Virol.* 71:8392-8396. It is of note that the ability to generate high levels of such T cells in the absence of β 2 microglobulin is reportedly limited to an H-2^d MHC class I-restricted response. β 2 microglobulin deficient mice have been reported to have a host of dramatic characteristics, such as, for example, an increased susceptibility to some parasitic diseases, an increased susceptibility to hepatitis infections, a deficiency in iron metabolism, and an impaired breeding phenotype. Cooper et al. (2007) An impaired breeding phenotype

in mice with a genetic deletion of Beta-2 microglobulin and diminished MHC class I expression: Role in reproductive fitness, Biol. Reprod. 77:274-279.

[00107] Mice that express human $\beta 2$ microglobulin as well as human HLA class I molecules (i.e., HLA-B7) on a randomly inserted transgene have been reported. Chamberlain et al. (1988) Tissue-specific and cell surface expression of human major histocompatibility complex class I heavy (HLA-B7) and light ($\beta 2$ -microglobulin) chain genes in transgenic mice, Proc. Natl. Acad. Sci. USA 85:7690-7694. The expression of human HLA class I was consistent with that of endogenous class I with a marked decrease in the liver. *Id.* The expression of human $\beta 2$ microglobulin was also consistent with the endogenous $\beta 2$ microglobulin, while expression of the human HLA class I molecule was increased 10- to 17-fold in double transgenic mice. *Id.* However, the authors did not attempt a replacement of a mouse endogenous $\beta 2$ microglobulin locus with a human $\beta 2$ microglobulin locus.

[00108] Therefore, disclosed herein is a genetically engineered non-human animal (e.g., a rodent, e.g., a mouse or a rat) whose genome comprises a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In one aspect, the animal does not express an endogenous non-human $\beta 2$ microglobulin from an endogenous non-human $\beta 2$ microglobulin locus. In some embodiments, the nucleotide sequence encodes a $\beta 2$ microglobulin polypeptide that is partially human and partially non-human, e.g., it contains some amino acids that correspond to human and some amino acids that correspond to non-human $\beta 2$ microglobulin. In one aspect, the non-human animal does not express an endogenous non-human $\beta 2$ microglobulin polypeptide from an endogenous non-human locus, and only expresses the human or humanized $\beta 2$ microglobulin polypeptide. In one example, the non-human animal does not express a complete endogenous non-human $\beta 2$ microglobulin polypeptide but only expresses a portion of a non-human endogenous $\beta 2$ microglobulin polypeptide from an endogenous $\beta 2$ microglobulin locus. Thus, in various embodiments, the animal does not express a functional non-human $\beta 2$ microglobulin polypeptide from an endogenous non-human $\beta 2$ microglobulin locus. In a specific aspect, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin is located at an endogenous non-human $\beta 2$ microglobulin locus. In one aspect, the animal comprises two copies of $\beta 2$ microglobulin locus comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In another aspect, the animal comprises one copy of $\beta 2$ microglobulin locus comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. Thus, the animal may be homozygous or heterozygous for $\beta 2$ microglobulin locus comprising a nucleotide sequence that encodes a

human or humanized $\beta 2$ microglobulin polypeptide. The nucleotide sequence of the human or humanized $\beta 2$ microglobulin may be derived from a collection of $\beta 2$ microglobulin sequences that are naturally found in human populations. In various embodiments, the genetically engineered non-human animal of the invention comprises in its germline a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin. In one embodiment, a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide comprises a nucleotide sequence encoding a polypeptide comprising a human $\beta 2$ microglobulin amino acid sequence. In one embodiment, the polypeptide is capable of binding to an MHC I protein.

[00109] The nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide may comprise nucleic acid residues corresponding to the entire human $\beta 2$ microglobulin gene. Alternatively, the nucleotide sequence may comprise nucleic acid residues encoding amino acid sequence set forth in amino acids 21-119 of a human $\beta 2$ microglobulin protein (i.e., amino acid residues corresponding to the mature human $\beta 2$ microglobulin). In an alternative embodiment, the nucleotide sequence may comprise nucleic acid residues encoding amino acid sequence set forth in amino acids 23-115 of a human $\beta 2$ microglobulin protein, for example, amino acid sequence set forth in amino acids 23-119 of a human $\beta 2$ microglobulin protein. The nucleic and amino acid sequences of human $\beta 2$ microglobulin are described in Gussow et al., *supra*, incorporated herein by reference.

[00110] Thus, the human or humanized $\beta 2$ microglobulin polypeptide may comprise amino acid sequence set forth in amino acids 23-115 of a human $\beta 2$ microglobulin polypeptide, e.g., amino acid sequence set forth in amino acids 23-119 of a human $\beta 2$ microglobulin polypeptide, e.g., amino acid sequence set forth in amino acids 21-119 of a human $\beta 2$ microglobulin polypeptide. Alternatively, the human $\beta 2$ microglobulin may comprise amino acids 1-119 of a human $\beta 2$ microglobulin polypeptide.

[00111] In some embodiments, the nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin comprises a nucleotide sequence set forth in exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. Alternatively, the nucleotide sequence comprises nucleotide sequences set forth in exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. In this embodiment, the nucleotide sequences set forth in exons 2, 3, and 4 are operably linked to allow for normal transcription and translation of the gene. Thus, in one embodiment, the human sequence comprises a nucleotide sequence corresponding to exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. In a specific embodiment, the human sequence comprises a

nucleotide sequence corresponding to exon 2 to about 267 bp after exon 4 of a human $\beta 2$ microglobulin gene. In a specific embodiment, the human sequence comprises about 2.8 kb of a human $\beta 2$ microglobulin gene.

[00112] Thus, the human or humanized $\beta 2$ microglobulin polypeptide may be encoded by a nucleotide sequence comprising nucleotide sequence set forth in exon 2 to exon 4 of a human $\beta 2$ microglobulin, e.g., nucleotide sequence corresponding to exon 2 to exon 4 of a human $\beta 2$ microglobulin gene. Alternatively, the polypeptide may be encoded by a nucleotide sequence comprising nucleotide sequences set forth in exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. In a specific embodiment, the human or humanized $\beta 2$ microglobulin polypeptide is encoded by a nucleotide sequence corresponding to exon 2 to about 267 bp after exon 4 of a human $\beta 2$ microglobulin gene. In another specific embodiment, the human or humanized polypeptide is encoded by a nucleotide sequence comprising about 2.8 kb of a human $\beta 2$ microglobulin gene. As exon 4 of the $\beta 2$ microglobulin gene contains the 5' untranslated region, the human or humanized polypeptide may be encoded by a nucleotide sequence comprising exons 2 and 3 of the $\beta 2$ microglobulin gene.

[00113] It would be understood by those of ordinary skill in the art that although specific nucleic acid and amino acid sequences to generate genetically engineered animals are described in the present examples, sequences of one or more conservative or non-conservative amino acid substitutions, or sequences differing from those described herein due to the degeneracy of the genetic code, are also provided.

[00114] Therefore, a non-human animal that expresses a human $\beta 2$ microglobulin sequence is provided, wherein the $\beta 2$ microglobulin sequence is at least about 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a human $\beta 2$ microglobulin sequence. In a specific embodiment, the $\beta 2$ microglobulin sequence is at least about 90%, 95%, 96%, 97%, 98%, or 99% identical to the human $\beta 2$ microglobulin sequence described in the Examples. In one embodiment, the human $\beta 2$ microglobulin sequence comprises one or more conservative substitutions. In one embodiment, the human $\beta 2$ microglobulin sequence comprises one or more non-conservative substitutions.

[00115] In addition, provided are non-human animals wherein the nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin protein also comprises a nucleotide sequence set forth in exon 1 of a non-human $\beta 2$ microglobulin gene. Thus, in a specific embodiment, the non-human animal comprises in its genome a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin wherein the nucleotide sequence

comprises exon 1 of a non-human $\beta 2$ microglobulin and exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. Thus, the human or humanized $\beta 2$ microglobulin polypeptide is encoded by exon 1 of a non-human $\beta 2$ microglobulin gene and exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene (e.g., exons 2 and 3 of a human $\beta 2$ microglobulin gene).

[00116] Similarly to a non-human animal comprising a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, the non-human animal comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin may be selected from a group consisting of a mouse, rat, rabbit, pig, bovine (e.g., cow, bull, buffalo), deer, sheep, goat, chicken, cat, dog, ferret, primate (e.g., marmoset, rhesus monkey). In some embodiments of the invention, the non-human animal is a mammal. In a specific embodiment, the non-human animal is a murine, e.g., a rodent (e.g., a mouse or a rat). In one embodiment, the animal is a mouse.

[00117] Thus, in some aspects, a genetically engineered mouse is provided, wherein the mouse comprises a nucleotide sequence encoding a human or a humanized $\beta 2$ microglobulin polypeptide as described herein. A genetically engineered mouse is provided, wherein the mouse comprises at its endogenous $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide (e.g., a human or substantially human $\beta 2$ microglobulin polypeptide). In some embodiments, the mouse does not express an endogenous $\beta 2$ microglobulin polypeptide (e.g., a functional endogenous $\beta 2$ microglobulin polypeptide) from an endogenous $\beta 2$ microglobulin locus. In some embodiments, the genetically engineered mouse comprises a nucleotide sequence comprising exon 1 of a mouse $\beta 2$ microglobulin gene and exons 2, 3, and 4 of a human $\beta 2$ microglobulin gene. In some embodiments, the mouse expresses the human or humanized $\beta 2$ microglobulin polypeptide.

[00118] In one aspect, a modified non-human $\beta 2$ microglobulin locus is provided that comprises a heterologous $\beta 2$ microglobulin sequence. In one embodiment, the heterologous $\beta 2$ microglobulin sequence is a human or a humanized sequence.

[00119] In one embodiment, the modified locus is a rodent locus. In a specific embodiment, the rodent locus is selected from a mouse or rat locus. In one embodiment, the non-human locus is modified with at least one human $\beta 2$ microglobulin coding sequence.

[00120] In one embodiment, the heterologous $\beta 2$ microglobulin sequence is operably linked to endogenous regulatory elements, e.g., endogenous promoter and/or expression control sequence. In a specific embodiment, the heterologous $\beta 2$ microglobulin sequence is

a human sequence and the human sequence is operably linked to an endogenous promoter and/or expression control sequence.

[00121] In one aspect, a modified non-human $\beta 2$ microglobulin locus is provided that comprises a human sequence operably linked to an endogenous promoter and/or expression control sequence.

[00122] In various aspects, the human or humanized $\beta 2$ microglobulin expressed by a genetically modified non-human animal, or cells, embryos, or tissues derived from a non-human animal, preserves all the functional aspects of the endogenous and/or human $\beta 2$ microglobulin. For example, it is preferred that the human or humanized $\beta 2$ microglobulin binds the α chain of MHC I polypeptide (e.g., endogenous non-human or human MHC I polypeptide). The human or humanized $\beta 2$ microglobulin polypeptide may bind, recruit or otherwise associate with any other molecules, e.g., receptor, anchor or signaling molecules that associate with endogenous non-human and/or human $\beta 2$ microglobulin (e.g., HFE, etc.).

[00123] In addition to genetically modified animals (e.g., rodents, e.g., mice or rats), also provided is a tissue or cell, wherein the tissue or cell is derived from a non-human animal as described herein, and comprises a heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence, i.e., nucleotide and/or amino acid sequence. In one embodiment, the heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence is a human or humanized $\beta 2$ microglobulin gene or human or humanized $\beta 2$ microglobulin sequence. Preferably, the cell is a nucleated cell. The cell may be any cell known to express MHC I complex, e.g., an antigen presenting cell. The human or humanized $\beta 2$ microglobulin polypeptide expressed by said cell may interact with endogenous non-human MHC I (e.g., rodent MHC I), to form a functional MHC I complex. The resultant MHC I complex may be capable of interacting with a T cell, e.g., a cytotoxic T cell. Thus, also provided is an *in vitro* complex of a cell from a non-human animal as described herein and a T cell.

[00124] Also provided are non-human cells that comprise human or humanized $\beta 2$ microglobulin gene or sequence, and an additional human or humanized sequence, e.g., chimeric MHC I polypeptide presently disclosed. In such an instance, the human or humanized $\beta 2$ microglobulin polypeptide may interact with, e.g., a chimeric human/non-human MHC I polypeptide, and a functional MHC I complex may be formed. In some aspects, such complex is capable of interacting with a TCR on a T cell, e.g., a human or a non-human T cell. Thus, also provided in an *in vitro* complex of a cell from a non-human animal as described herein and a human or a non-human T cell.

[00125] Another aspect of the disclosure is a rodent embryo (e.g., a mouse or a rat embryo) comprising a heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence as described herein. In one embodiment, the embryo comprises an ES donor cell that comprises the heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence, and host embryo cells. The heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence is a human or humanized $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence.

[00126] This invention also encompasses a non-human cell comprising a chromosome or fragment thereof of a non-human animal as described herein (e.g., wherein the chromosome or fragment thereof comprises a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide). The non-human cell may comprise a nucleus of a non-human animal as described herein. In one embodiment, the non-human cell comprises the chromosome or fragment thereof as the result of a nuclear transfer.

[00127] In one aspect, a non-human induced pluripotent cell comprising a heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence is provided. In one embodiment, the induced pluripotent cell is derived from a non-human animal as described herein. In one embodiment, the heterologous $\beta 2$ microglobulin gene or $\beta 2$ microglobulin sequence is a human or humanized gene or sequence.

[00128] Also provided is a hybridoma or quadroma, derived from a cell of a non-human animal as described herein. In one embodiment, the non-human animal is a mouse or rat.

[00129] The disclosure also provides methods for making a genetically engineered non-human animal (e.g., a genetically engineered rodent, e.g., a mouse or a rat) described herein. The methods result in an animal whose genome comprises a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In one aspect, the methods result in a genetically engineered mouse, whose genome comprises at an endogenous $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In some instances, the mouse does not express a functional mouse $\beta 2$ microglobulin from an endogenous mouse $\beta 2$ microglobulin locus. In some aspects, the methods utilize a targeting construct made using VELOCIGENE® technology, introducing the construct into ES cells, and introducing targeted ES cell clones into a mouse embryo using VELOCIMOUSE® technology, as described in the Examples. In one embodiment, the ES cells are mix of 129 and C57BL/6 mouse strains; in another embodiment, the ES cells are a mix of BALB/c and 129 mouse strains.

[00130] Also provided is a nucleotide construct used for generating genetically engineered non-human animals. The nucleotide construct may comprise: 5' and 3' non-

human homology arms, a human DNA fragment comprising human $\beta 2$ microglobulin sequences, and a selection cassette flanked by recombination sites. In one embodiment, the human DNA fragment is a genomic fragment that comprises both introns and exons of a human $\beta 2$ microglobulin gene. In one embodiment, the non-human homology arms are homologous to a non-human $\beta 2$ microglobulin locus. The genomic fragment may comprise exons 2, 3, and 4 of the human $\beta 2$ microglobulin gene. In one instance, the genomic fragment comprises, from 5' to 3': exon 2, intron, exon 3, intron, and exon 4, all of human $\beta 2$ microglobulin sequence. The selection cassette may be located anywhere in the construct outside the $\beta 2$ microglobulin coding region, e.g., it may be located 3' of exon 4 of the human $\beta 2$ microglobulin. The 5' and 3' non-human homology arms may comprise genomic sequence 5' and 3' of endogenous non-human $\beta 2$ microglobulin gene, respectively. In another embodiment, the 5' and 3' non-human homology arms comprise genomic sequence 5' of exon 2 and 3' of exon 4 of endogenous non-human gene, respectively.

[00131] Another aspect of the invention relates to a method of modifying a $\beta 2$ microglobulin locus of a non-human animal (e.g., a rodent, e.g., a mouse or a rat) to express a human or humanized $\beta 2$ microglobulin polypeptide described herein. One method of modifying a $\beta 2$ microglobulin locus of a mouse to express a human or humanized $\beta 2$ microglobulin polypeptide comprises replacing at an endogenous $\beta 2$ microglobulin locus a nucleotide sequence encoding a mouse $\beta 2$ microglobulin with a nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide. In one embodiment of such method, the mouse does not express a functional $\beta 2$ microglobulin polypeptide from an endogenous mouse $\beta 2$ microglobulin locus. In some specific embodiments, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide comprises nucleotide sequence set forth in exons 2 to 4 of the human $\beta 2$ microglobulin gene. In other embodiments, the nucleotide sequence encoding the human or humanized $\beta 2$ microglobulin polypeptide comprises nucleotide sequences set forth in exons 2, 3, and 4 of the human $\beta 2$ microglobulin gene.

Genetically Modified MHC I / $\beta 2$ Microglobulin Animals

[00132] In various embodiments, the invention generally provides genetically modified non-human animals that comprise in their genome nucleotide sequences encoding both human or humanized MHC I and $\beta 2$ microglobulin polypeptides; thus, the animals express both human or humanized MHC I and $\beta 2$ microglobulin polypeptides.

[00133] Functional differences arise in the use of mixed human/non-human system components. HLA class I binds $\beta 2$ microglobulin tighter than mouse class I. Bernabeu

(1984) β 2-microglobulin from serum associates with MHC class I antigens on the surface of cultured cells, *Nature* 308:642-645. Attempts to abrogate functional differences are reflected in the construction of particular humanized MHC mice. H-2 class I and class II knockout mice (in a mouse β 2 microglobulin KO background) that express a randomly integrated human HLA-A2.1/HLA-DR1 chimeric transgene having an α 1 and α 2 of human HLA-A2.1, and α 3 of mouse H-2D^b, attached at its N-terminal via a linker to the C-terminus of human β 2-microglobulin have been developed. See, e.g., Pajot et al. (2004) A mouse model of human adaptive immune functions: *HLA-A2.1/HLA-DR1-transgenic H-2 class I-/class II-knockout mice*, *Eur. J. Immunol.* 34:3060-3069. These mice reportedly generate antigen-specific antibody and CTL responses against hepatitis B virus, whereas mice merely transgenic for HLA-A2.1 or H-2 class I/class II knockout mice do not. The deficiency of mice that are merely transgenic for the genes presumably stems from the ability of such mice to employ endogenous class I and/or class II genes to circumvent any transgene, an option not available to MHC knockout mice. However, the mice may express at least H-2D^b, presumably due to breedings into the mouse β 2 microglobulin knockout mouse background (see, Pajot et al., *supra*; which apparently comprised an intact endogenous class I and class II locus).

[00134] Cell surface expression of the chimeric fusion with human β 2 microglobulin is reportedly lower than endogenous MHC expression, but survivability/rate of NK killing is not reported, nor is the rate of NK self-killing. Pajot et al., *supra*. Some improvement in CD8+ T cell numbers was observed over MHC class I-deficient β 2-microglobulin knockout mice (2-3% of total splenocytes, vs. 0.6-1% in the β 2 KO mice). However, T cell variable region usage exhibited altered profiles for BV 5.1, BV 5.2, and BV 11 gene segments. Both CD8+ and CD4+ T cell responses were reportedly restricted to the appropriate hepatitis B antigen used to immunize the mice, although at least two mice killed cells bearing either of the antigens, where the mice were immunized with only one antigen, which might be due to a lack of NK cell inhibition or lack of NK cell selectivity.

[00135] As mentioned above, mice transgenic for both human MHC I and human β 2 microglobulin comprise a nucleotide sequence encoding a chimeric MHC I/ β 2 microglobulin protein, wherein the MHC I and β 2 microglobulin portions are contained within a single polypeptide chain, resulting in MHC I α chain and β 2 microglobulin being covalently linked to each other and thereby tethered at the cell surface. A mouse which comprises in its genome two independent nucleotide sequences, one encoding a human or humanized MHC I polypeptide and the other encoding a human or humanized β 2 microglobulin polypeptide is provided. The mouse provided herein would express an MHC I complex that more closely

resembles an MHC I complex present in nature, wherein MHC I α chain and β 2 microglobulin are provided on two separate polypeptide chains with β 2 microglobulin non-covalently associating with the MHC I α chain.

[00136] Thus, the present disclosure provides a non-human animal comprising in its genome: a first nucleotide sequence encoding a human or humanized MHC I polypeptide, and a second nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide. In one aspect, provided is a non-human animal comprising in its genome: (a) a first nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein the human portion of the chimeric polypeptide comprises a peptide binding domain or an extracellular domain of a human MHC I (e.g., HLA-A, HLA-B, or HLA-C; e.g., HLA-A2), and (b) a second nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide.

[00137] The first nucleotide sequence may be located at an endogenous non-human MHC I locus such that the animal comprises in its genome a replacement at the MHC I locus of all or a portion of endogenous MHC I gene (e.g., a portion encoding a peptide binding domain or an extracellular domain) with the corresponding human MHC I sequence. Thus, the animal may comprise at an endogenous MHC I locus a nucleotide sequence encoding an extracellular domain of a human MHC I (e.g., HLA-A, HLA-B, or HLA-C; e.g., HLA-A2) and transmembrane and cytoplasmic domains of endogenous non-human MHC I (e.g., H-2K, H-2D, etc., e.g., H-2Kb). In one aspect, the animal is a mouse, and the first nucleotide sequence comprises a nucleotide sequence encoding an extracellular domain of a human HLA-A2 (e.g., HLA-A2.1) and transmembrane and cytoplasmic domains of a mouse H-2K (e.g., H-2Kb).

[00138] The second nucleotide sequence may be located at an endogenous non-human β 2 microglobulin locus such that the animal comprises in its genome a replacement at the β 2 microglobulin locus of all or a portion of endogenous β 2 microglobulin gene with the corresponding human β 2 microglobulin sequence. The second nucleotide sequence may comprise a nucleotide sequence set forth in exon 2 to exon 4 of a human β 2 microglobulin gene. Alternatively, the second nucleotide sequence may comprise nucleotide sequences set forth in exons 2, 3, and 4 of a human β 2 microglobulin gene. In this embodiment, nucleotide sequences are operably linked to each other. The second nucleotide sequence may further comprise the sequence of exon 1 of a non-human β 2 microglobulin gene.

[00139] In one aspect, the animal does not express a functional MHC I from an endogenous non-human MHC I locus (e.g., does not express either a peptide binding domain or an extracellular domain of the endogenous MHC I), and the animal does not

express a functional $\beta 2$ microglobulin polypeptide from an endogenous non-human $\beta 2$ microglobulin locus. In some aspects, the animal is homozygous for both an MHC I locus comprising a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide and a $\beta 2$ microglobulin locus comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin. In other aspects, the animal is heterozygous for both an MHC I locus comprising a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide and a $\beta 2$ microglobulin locus comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin.

[00140] Preferably, the first and the second nucleotide sequences are operably linked to endogenous expression control elements (e.g., promoters, enhancers, silencers, etc.).

[00141] Various other embodiments of the first and second nucleotide sequences (and the polypeptides they encode) encompassed herein may be readily understood from the embodiments described throughout the specification, e.g., those described in the sections related to genetically engineered MHC I animals and genetically engineered $\beta 2$ microglobulin animals.

[00142] In one aspect, the disclosure provides a mouse comprising in its genome (a) a first nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide (specifically, HLA-A2/H-2K polypeptide), wherein the human portion of the chimeric polypeptide comprises an extracellular domain of a human HLA-A2 and the mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K, and (b) a second nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide (e.g., wherein the nucleotide sequence comprises a nucleotide sequence set forth in exon 2 to exon 4 of the human $\beta 2$ microglobulin gene or nucleotide sequences set forth in exon 2, 3, and 4 of the human $\beta 2$ microglobulin gene), wherein the first nucleotide sequence is located at an endogenous H-2K locus, and the second sequence is located at an endogenous $\beta 2$ microglobulin locus. In one embodiment, the mouse does not express functional H-2K and mouse $\beta 2$ microglobulin polypeptides from their respective endogenous loci. In one embodiment, the mouse expresses both the chimeric human/mouse MHC I polypeptide and the human or humanized $\beta 2$ microglobulin polypeptide.

[00143] As shown in the following Examples, animals genetically engineered to co-express both the human or humanized MHC I and $\beta 2$ microglobulin displayed increased expression of chimeric MHC class I on cell surface in comparison to animals humanized for MHC I alone. In some embodiments, co-expression of human or humanized MHC I and $\beta 2$ microglobulin increases cell surface expression of human or humanized MHC I by more than

about 10%, e.g., more than about 20%, e.g., about 50% or more, e.g., about 70%, over the expression of human or humanized MHC I in the absence of human or humanized $\beta 2$ microglobulin.

[00144] The disclosure also provides a method of making genetically engineered non-human animals (e.g., rodents, e.g., rats or mice) whose genome comprises a first and a second nucleotide sequence as described herein. The method generally comprises generating a first genetically engineered non-human animal whose genome comprises a first nucleotide sequence described herein (i.e., a human or humanized MHC I sequence), generating a second genetically engineered non-human animal whose genome comprises a second nucleotide sequence described herein (i.e., a human or humanized $\beta 2$ microglobulin sequence), and breeding the first and the second animal to obtain progeny whose genome contains both nucleotide sequences. In one embodiment, the first and the second animal are heterozygous for the first and the second nucleotide sequence, respectively. In one embodiment, the first and the second animal are homozygous for the first and the second nucleotide sequence, respectively. In one embodiment, the first and second animals are generated through replacement of endogenous non-human loci with the first and the second nucleotide sequences, respectively. In one aspect, the first and the second animals are generated through utilization of constructs generated via VELOCIGENE® technology, and introducing targeted ES cell clones bearing such constructs into an embryo (e.g., a rodent embryo, e.g., a mouse or a rat embryo) via the VELOCIMOUSE® method.

Use of Genetically Modified Animals

[00145] In various embodiments, the genetically modified non-human animals described herein make APCs with human or humanized MHC I and/or $\beta 2$ microglobulin on the cell surface and, as a result, present peptides derived from cytosolic proteins as epitopes for CTLs in a human-like manner, because substantially all of the components of the complex are human or humanized. The genetically modified non-human animals of the invention can be used to study the function of a human immune system in the humanized animal; for identification of antigens and antigen epitopes that elicit immune response (e.g., T cell epitopes, e.g., unique human cancer epitopes), e.g., for use in vaccine development; for identification of high affinity T cells to human pathogens or cancer antigens (i.e., T cells that bind to antigen in the context of human MHC I complex with high avidity), e.g., for use in adoptive T cell therapy; for evaluation of vaccine candidates and other vaccine strategies; for studying human autoimmunity; for studying human infectious diseases; and otherwise for devising better therapeutic strategies based on human MHC expression.

[00146] The MHC I complex binds peptides and presents them on cell surface. Once presented on the surface in the context of such a complex, the peptides are recognizable by T cells. For example, when the peptide is derived from a pathogen or other antigen of interest (e.g., a tumor antigen), T cell recognition can result in T cell activation, macrophage killing of cells bearing the presented peptide sequence, and B cell activation of antibodies that bind the presented sequence.

[00147] T cells interact with cells expressing MHC I complex through the peptide-bound MHC class I ectodomain and the T cell's CD8 ectodomain. CD8+ T cells that encounter APC's that have suitable antigens bound to the MHC class I molecule will become cytotoxic T cells. Thus, antigens that in the context of MHC class I bind with high avidity to a T cell receptor are potentially important in the development of treatments for human pathologies. However, presentation of antigens in the context of mouse MHC I is only somewhat relevant to human disease, since human and mouse MHC complexes recognize antigens differently, e.g., a mouse MHC I may not recognize the same antigens or may present different epitopes than a human MHC I. Thus, the most relevant data for human pathologies is obtained through studying the presentation of antigen epitopes by human MHC I.

[00148] Thus, in various embodiments, the genetically engineered animals of the present invention are useful, among other things, for evaluating the capacity of an antigen to initiate an immune response in a human, and for generating a diversity of antigens and identifying a specific antigen that may be used in human vaccine development.

[00149] In one aspect, a method for determining antigenicity in a human of a peptide sequence is provided, comprising exposing a genetically modified non-human animal as described herein to a molecule comprising the peptide sequence, allowing the non-human animal to mount an immune response, and detecting in the non-human animal a cell that binds a sequence of the peptide presented by a chimeric human/non-human MHC I, or a humanized MHC I complex (comprising a chimeric human/non-human MHC I and a human or humanized $\beta 2$ microglobulin) as described herein.

[00150] In one aspect, a method for determining whether a peptide will provoke a cellular immune response in a human is provided, comprising exposing a genetically modified non-human animal as described herein to the peptide, allowing the non-human animal to mount an immune response, and detecting in the non-human animal a cell that binds a sequence of the peptide by a chimeric human/non-human MHC class I molecule as described herein. In one embodiment, the non-human animal following exposure comprises an MHC class I-restricted CD8+ cytotoxic T lymphocyte (CTL) that binds the peptide. In one embodiment, the CTL kills a cell bearing the peptide.

[00151] In one aspect, a method for identifying a human CTL epitope is provided, comprising exposing a non-human animal as described herein to an antigen comprising a putative CTL epitope, allowing the non-human animal to mount an immune response, isolating from the non-human animal an MHC class I-restricted CD8+ CTL that binds the epitope, and identifying the epitope bound by the MHC class I-restricted CD8+ CTL.

[00152] In one aspect, a method is provided for identifying an HLA class I-restricted peptide whose presentation by a human cell and binding by a human lymphocyte (e.g., human T cell) will result in cytotoxicity of the peptide-bearing cell, comprising exposing a non-human animal (or MHC class I-expressing cell thereof) as described herein to a molecule comprising a peptide of interest, isolating a cell of the non-human animal that expresses a chimeric human/non-human class I molecule that binds the peptide of interest, exposing the cell to a human lymphocyte that is capable of conducting HLA class I-restricted cytotoxicity, and measuring peptide-induced cytotoxicity.

[00153] In one aspect, a method is provided for identifying an antigen that generates a cytotoxic T cell response in a human, comprising exposing a putative antigen to a mouse as described herein, allowing the mouse to generate an immune response, and identifying the antigen bound by the HLA-A-restricted molecule.

[00154] In one embodiment, the antigen comprises a bacterial or viral surface or envelope protein. In one embodiment, the antigen comprises an antigen on the surface of a human tumor cell. In one embodiment, the antigen comprises a putative vaccine for use in a human. In one embodiment, the antigen comprises a human epitope that generates antibodies in a human. In another embodiment, the antigen comprises a human epitope that generates high affinity CTLs that target the epitope/MHC I complex.

[00155] In one aspect, a method is provided for determining whether a putative antigen contains an epitope that upon exposure to a human immune system will generate an HLA-A-restricted immune response (e.g., HLA-A2-restricted response), comprising exposing a mouse as described herein to the putative antigen and measuring an antigen-specific HLA-A-restricted (e.g., HLA-A2-restricted) immune response in the mouse.

[00156] In one embodiment, the putative antigen is selected from a biopharmaceutical or fragment thereof, a non-self protein, a surface antigen of a non-self cell, a surface antigen of a tumor cell, a surface antigen of a bacterial or yeast or fungal cell, a surface antigen or envelope protein of a virus.

[00157] In addition, the genetically engineered non-human animals described herein may be useful for identification of T cell receptors, e.g., high-avidity T cell receptors, that recognize an antigen of interest, e.g., a tumor or another disease antigen. The method may

comprise: exposing the non-human animal described herein to an antigen, allowing the non-human animal to mount an immune response to the antigen, isolating from the non-human animal a T cell comprising a T cell receptor that binds the antigen presented by a human or humanized MHC I, and determining the sequence of said T cell receptor.

[00158] In one aspect, a method for identifying a T cell receptor variable domain having high affinity for a human tumor antigen is provided, comprising exposing a mouse comprising humanized MHC I $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains (e.g., HLA-A2 $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains) to a human tumor antigen; allowing the mouse to generate an immune response; and, isolating from the mouse a nucleic acid sequence encoding a T cell receptor variable domain, wherein the T cell receptor variable domain binds the human tumor antigen with a K_D of no higher than about 1 nanomolar.

[00159] In one embodiment, the mouse further comprises a replacement at the endogenous mouse T cell receptor variable region gene locus with a plurality of unrearranged human T cell receptor variable region gene segments, wherein the unrearranged human T cell receptor variable region gene segments recombine to encode a chimeric human-mouse T cell receptor gene comprising a human variable region and a mouse constant region. In one embodiment, the mouse comprises a human CD8 transgene, and the mouse expresses a functional human CD8 protein.

[00160] T cell receptors having high avidity to tumor antigens are useful in cell-based therapeutics. T cell populations with high avidity to human tumor antigens have been prepared by exposing human T cells to HLA-A2 that has been mutated to minimize CD8 binding to the $\alpha 3$ subunit, in order to select only those T cells with extremely high avidity to the tumor antigen (i.e., T cell clones that recognize the antigen in spite of the inability of CD8 to bind $\alpha 3$). See, Pittet et al. (2003) $\alpha 3$ Domain Mutants of Peptide/MHC Class I Multimers Allow the Selective Isolation of High Avidity Tumor-Reactive CD8 T Cells, *J. Immunol.* 171:1844-1849. The non-human animals, and cells of the non-human animals, are useful for identifying peptides that will form a complex with human HLA class I that will bind with high avidity to a T cell receptor, or activate a lymphocyte bearing a T cell receptor.

[00161] Antigen/HLA class I binding to a T cell, or activation of a T cell, can be measured by any suitable method known in the art. Peptide-specific APC-T cell binding and activation are measurable. For example, T cell engagement of antigen-presenting cells that express HLA-A2 reportedly causes PIP2 to accumulate at the immunosynapse, whereas cross-linking MHC class I molecules does not. See, Fooksman et al. (2009) Cutting Edge: Phosphatidylinositol 4,5-Bisphosphate Concentration at the APC Side of the Immunological Synapse Is Required for Effector T Cell Function, *J. Immunol.* 182:5179-5182.

[00162] Functional consequences of the interaction of a lymphocyte bearing a TCR, and a class I-expressing APC, are also measurable and include cell killing by the lymphocyte. For example, contact points on the $\alpha 2$ subunit of HLA-A2 by CD8+ CTLs reportedly generate a signal for Fas-independent killing. HLA-A2-expressing Jurkat cells apoptose when contacted (by antibodies) at epitopes on the HLA-A2 molecule known (from crystallographic studies) to contact CD8, without any apparent reliance on the cytoplasmic domain. See, Pettersen et al. (1998) The TCR-Binding Region of the HLA Class I $\alpha 2$ Domain Signals Rapid Fas-Independent Cell Death: A Direct Pathway for T Cell-Mediated Killing of Target Cells? J. Immunol. 160:4343-4352. It has been postulated that the rapid killing induced by HLA-A2 $\alpha 2$ contact with a CD8 of a CD8+ CTL may primarily be due to this Fas-independent HLA-A2-mediated pathway (*id.*), as distinguished from TCR-independent $\alpha 3$ domain-mediated killing—which by itself can induce apoptosis (see, Woodle et al. (1997) Anti-human class I MHC antibodies induce apoptosis by a pathway that is distinct from the Fas antigen-mediated pathway, J. Immunol. 158:2156-2164).

[00163] The consequence of interaction between a T cell and an APC displaying a peptide in the context of MHC I can also be measured by a T cell proliferation assay. Alternatively, it can be determined by measuring cytokine release commonly associated with activation of immune response. In one embodiment, IFN γ ELISPOT can be used to monitor and quantify CD8+ T cell activation.

[00164] As described herein, CD8+ T cell activation can be hampered in the genetically modified non-human animals described herein due to species-specific binding of CD8 to MHC I. For embodiments where a species-specific CD8 interaction is desired, a cell of a genetically modified animal as described herein (e.g., a rodent, e.g., a mouse or a rat) is exposed (e.g., *in vitro*) to a human cell, e.g., a human CD8-bearing cell, e.g., a human T cell. In one embodiment, an MHC class I-expressing cell of a mouse as described herein is exposed *in vitro* to a T cell that comprises a human CD8 and a T cell receptor. In a specific embodiment, the T cell is a human T cell. In one embodiment, the MHC class I-expressing cell of the mouse comprises a peptide bound to a chimeric human/mouse MHC I or a humanized MHC I complex (which includes human $\beta 2$ microglobulin), the T cell is a human T cell, and the ability of the T cell to bind the peptide-displaying mouse cell is determined. In one embodiment, activation of the human T cell by the peptide-displaying mouse cell is determined. In one embodiment, an *in vitro* method for measuring activation of a human T cell by the peptide-displaying cell is provided, comprising exposing a mouse or a mouse cell as described herein to an antigen of interest, exposing a cell from said mouse or said mouse cell (presumably bearing a peptide derived from the antigen in complex with human or humanized MHC I) to a human T cell, and measuring activation of the human T cell. In one

embodiment, the method is used to identify a T cell epitope of a human pathogen or a human neoplasm. In one embodiment, the method is used to identify an epitope for a vaccine.

[00165] In one embodiment, a method is provided for determining T cell activation by a putative human therapeutic, comprising exposing a genetically modified animal as described herein to a putative human therapeutic (or e.g., exposing a human or humanized MHC I-expressing cell of such an animal to a peptide sequence of the putative therapeutic), exposing a cell of the genetically modified animal that displays a human or humanized MHC I/peptide complex to a T cell comprising a human T cell receptor and a CD8 capable of binding the cell of the genetically modified animal, and measuring activation of the human T cell that is induced by the peptide-displaying cell of the genetically modified animal.

[00166] In various embodiments, a complex formed between a human or humanized MHC class I-expressing cell from an animal as described herein is made with a T cell that comprises a human CD8 sequence, e.g., a human T cell, or a T cell of a non-human animal that comprises a transgene that encodes human CD8. Mice transgenic for human CD8 are known in the art. Tishon et al. (2000) Transgenic Mice Expressing Human HLA and CD8 Molecules Generate HLA-Restricted Measles Virus Cytotoxic T Lymphocytes of the Same Specificity as Humans with Natural Measles Virus Infection, *Virology* 275(2):286-293; also, LaFace et al. (1995) Human CD8 Transgene Regulation of HLA Recognition by Murine T Cells, *J. Exp. Med.* 182:1315-1325.

[00167] In addition to the ability to identify antigens and antigen epitopes from human pathogens or neoplasms, the genetically modified animals of the invention can be used to identify autoantigens of relevance to human autoimmune diseases, e.g., type I diabetes, multiple sclerosis, etc. For example, Takaki et al. ((2006) HLA-A*0201-Restricted T Cells from Humanized NOD Mice Recognize Autoantigens of Potential Clinical Relevance to Type 1 Diabetes, *J. Immunol.* 176:3257-65) describe the utility of NOD mice bearing HLA/β2 microglobulin monochain in identifying type 1 diabetes autoantigens. Also, the genetically modified animals of the invention can be used to study various aspects of human autoimmune disease. As some polymorphic alleles of human MHC I are known to be associated with development of certain diseases, e.g., autoimmune diseases (e.g., Graves' disease, myasthenia gravis, psoriasis, etc.; see Bakker et al. (2006) A high-resolution HLA and SNP haplotype map for disease association studies in the extended human MHC, *Nature Genetics* 38:1166-72 and Supplementary Information and International MHC and Autoimmunity Genetics Network (2009) Mapping of multiple susceptibility variants within the MHC region for 7 immune-mediated diseases, *Proc. Natl. Acad. Sci. USA* 106:18680-85, both incorporated herein by reference), a genetically modified animal of the invention

comprising a humanized MHC I locus including such an allele may be useful as an autoimmune disease model. In one embodiment, the disease allele is HLA-B27, and the disease is ankylosing spondylitis or reactive arthritis; thus, in one embodiment, the animal used for the study of these diseases comprises a human or humanized HLA-B27.

[00168] Other aspects of cellular immunity that involve MHC I complexes are known in the art; therefore, genetically engineered non-human animals described herein can be used to study these aspects of immune biology. For instance, binding of TCR to MHC class I is modulated *in vivo* by additional factors. Leukocyte immunoglobulin-like receptor subfamily B member (LILRB1, or LIR-1) is expressed on MHC Class I-restricted CTLs and down-regulates T cell stimulation by binding a specific determinant on the $\alpha 3$ subunit of MHC class I molecules on APCs. Structural studies show that the binding site for LIR-1 and CD8 overlap, suggesting that inhibitory LIR-1 competes with stimulatory CD8 for binding with MHC class I molecules. Willcox et al. (2003) Crystal structure of HLA-A2 bound to LIR-1, a host and viral major histocompatibility complex receptor, *Nature Immunology* 4(9):913-919; also, Shirioishi et al. (2003) Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G, *Proc. Natl. Acad. Sci. USA* 100(15):8856-8861. LIR-1 transduces inhibitory signals through its (intracellular) immunoreceptor tyrosine-based inhibitory motif (ITIM). In NK cells, studies have shown that KIRs (inhibitory killer cell Ig-like receptors) lacking ITIMs (normally incapable of inhibition) can inhibit in the presence of LIR-1 (presumably through the LIR-1 ITIM) bound to the $\alpha 3$ domain of an MHC class I molecule (see, Kirwin et al. (2005) Killer Cell Ig-Like Receptor-Dependent Signaling by Ig-Like Transcript 2 (ILT2/CD85j/LILRB1/LIR-1) *J. Immunol.* 175:5006-5015), suggesting cooperation between LIR-1 bound to MHC class I and KIRs and thus a role for HLA $\alpha 3$ domain binding in modulating NK cell inhibition.

[00169] As described above, MHC molecules interact with cells that do not express a TCR. Among these cells are NK cells. NK cells are cytotoxic lymphocytes (distinguished from CTLs, or cytotoxic T lymphocytes) that play a central role in the cellular immune response, and in particular innate immunity. NK cells are the first line of defense against invading microorganisms, viruses, and other non-self (e.g., tumor) entities. NK cells are activated or inhibited through surface receptors, and they express CD8 but do not express TCRs. NK cells can interact with cells that express MHC class I, but interaction is through the CD8-binding $\alpha 3$ domain rather than the TCR-binding, peptide-bearing α_1 and α_2 domains. A primary function of NK cells is to destroy cells that lack sufficient MHC class I surface protein.

[00170] Cross-linking MHC class I molecules on the surface of human natural killer (NK) cells results in intracellular tyrosine phosphorylation, migration of the MHC class I molecule from the immunosynapse, and down-regulation of tumor cell killing. Rubio et al. (2004) Cross-linking of MHC class I molecules on human NK cells inhibits NK cell function, segregates MHC I from the NK cell synapse, and induces intracellular phosphotyrosines, J. Leukocyte Biol. 76:116-124.

[00171] Another function of MHC class I in NK cells is apparently to prevent self-killing. NK cells bear both activating receptor 2B4 and the 2B4 ligand CD48; MHC class I appears to bind 2B4 and prevent its activation by CD48. Betser-Cohen (2010) The Association of MHC Class I Proteins with the 2B4 Receptor Inhibits Self-Killing of Human NK Cells, J. Immunol. 184:2761-2768.

[00172] Thus, the genetically engineered non-human animals described herein can be used to study these non-TCR or non-CTL mediated processes and to design approaches for their modulation.

EXAMPLES

[00173] The invention will be further illustrated by the following nonlimiting examples. These Examples are set forth to aid in the understanding of the invention but are not intended to, and should not be construed to, limit its scope in any way. The Examples do not include detailed descriptions of conventional methods that would be well known to those of ordinary skill in the art (molecular cloning techniques, etc.). Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is indicated in Celsius, and pressure is at or near atmospheric.

Example 1. Construction and Characterization of Genetically Modified HLA-A2 Mice

Example 1.1: Expression of HLA-A2/H-2K in MG87 Cells.

[00174] A viral construct containing a chimeric HLA-A2/H-2K gene sequence (FIG. 4A) was made using standard molecular cloning techniques known to a skilled artisan in order to analyze chimeric human/mouse MHC I expression in transfected cells.

[00175] Briefly, a chimeric human HLA-A/mouse H-2K viral construct was made using the exon sequences encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the α chain and cloning them in frame with the mouse coding sequences for the transmembrane and cytoplasmic domains from the H-2K gene (FIG. 4A, pMIG-HLA-A2/H2K). As illustrated in FIG. 4, the construct contained an IRES-GFP reporter sequence, which allowed for determining if the construct was able to express in cells upon transfection.

[00176] Viruses containing the chimeric construct described above were made and propagated in human embryonic kidney 293 (293T) cells. 293T cells were plated on 10 cm dishes and allowed to grow to 95% confluency. A DNA transfection mixture was prepared with 25 µg of pMIG-HLA-A2/H2K, pMIG-human HLA-A2, or pMIG-humanized β2 microglobulin, and 5 µg of pMDG (envelope plasmid), 15 µg of pCL-Eco (packaging construct without packaging signal Ψ), 1 mL of Opti-MEM (Invitrogen). Added to this 1 mL DNA mixture was 80 µL of Lipofectamine-2000 (Invitrogen) in 1 mL of Opti-MEM, which was previously mixed together and allowed to incubate at room temperature for 5 minutes. The Lipofectamine/DNA mixture was allowed to incubate for an additional 20 minutes at room temperature, and then was added to 10 cm dishes, and the plates were incubated at 37°C. Media from the cells was collected after 24 hours and a fresh 10 mL of R10 (RPMI 1640 + 10% FBS) media was added to the cells. This media exchange was repeated twice. After a total of four days, the collected media was pooled, centrifuged and passed through a sterile filter to remove cellular debris.

[00177] The propagated viruses made above were used to transduce MG87 (mouse fibroblast) cells. MG87 cells from a single T-75 flask were washed once with PBS. 3 mL of 0.25% Trypsin + EDTA was added to the cells and allowed to incubate at room temperature for three minutes. 7 mL of D10 (high glucose DMEM; 10% Fetal Bovine Serum) was added to the cells/trypsin mixture and transferred to a 15 mL tube to centrifuge at 1300 rpm for five minutes. After centrifuging the cells, the media was aspirated and the cells resuspended in 5 mL D10. Cells were counted and $\sim 3.0 \times 10^5$ cells were placed per well in a 6-well plate. pMIG-human HLA-A2 or pMIG-HLA-A2/H-2K either alone or with pMIG-humanized β2 microglobulin virus were added to the wells, with non-transduced cells as a control. Cells were incubated at 37°C with 5% CO₂ for 2 days. Cells were prepared for FACS analysis (using anti-HLA-A2 antibody, clone BB7.2) for HLA-A2 expression with or without β2 microglobulin.

[00178] The graphs (FIG. 4B), as well as the table summarizing the data obtained from the graphs (FIG. 4C) demonstrate that co-transduction with humanized β2 microglobulin increases the expression of human HLA-A2 or chimeric human/non-human HLA-A2/H-2K, as demonstrated by the shift of curves to the right.

Example 1.2. Engineering a Chimeric HLA-A2/H-2K Locus.

[00179] The mouse H-2K gene was humanized in a single step by construction of a unique targeting vector from human and mouse bacterial artificial chromosome (BAC) DNA using VELOCIGENE® technology (see, e.g., US Pat. No. 6,586,251 and Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution

expression analysis. *Nat. Biotech.* 21(6): 652-659). DNA from mouse BAC clone RP23-173k21 (Invitrogen) was modified by homologous recombination to replace the genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the mouse H-2K gene with human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ subunits of the human HLA-A gene (FIG. 5).

[00180] Briefly, the genomic sequence encoding the mouse the $\alpha 1$, $\alpha 2$ and $\alpha 3$ subunits of the H-2K gene is replaced with the human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the human HLA-A*0201 gene in a single targeting event using a targeting vector comprising a hygromycin cassette flanked by *loxP* sites with a 5' mouse homology arm containing sequence 5' of the mouse H-2K locus including the 5' untranslated region (UTR; 5' homology arm is set forth in SEQ ID NO: 1) and a 3' mouse homology arm containing genomic sequence 3' of the mouse H-2K $\alpha 3$ coding sequence (3' homology arm is set forth in SEQ ID NO: 2).

[00181] The final construct for targeting the endogenous H-2K gene locus from 5' to 3' included (1) a 5' homology arm containing ~200 bp of mouse genomic sequence 5' of the endogenous H-2K gene including the 5'UTR, (2) ~1339 bp of human genomic sequence including the HLA-A*0201 leader sequence, the HLA-A*0201 leader/ $\alpha 1$ intron, the HLA-A*0201 $\alpha 1$ exon, the HLA-A*0201 $\alpha 1$ - $\alpha 2$ intron, the HLA-A*0201 $\alpha 2$ exon, ~316 bp of the 5' end of the $\alpha 2$ - $\alpha 3$ intron, (3) a 5' *loxP* site, (4) a hygromycin cassette, (5) a 3' *loxP* site, (6) ~580 bp of human genomic sequence including ~304 bp of the 3' end of the $\alpha 2$ - $\alpha 3$ intron, the HLA-A*0201 $\alpha 3$ exon, and (7) a 3' homology arm containing ~200 bp of mouse genomic sequence including the intron between the mouse H-2K $\alpha 3$ and transmembrane coding sequences (see FIG. 5 for schematic representation of the H-2K targeting vector). The sequence of 149 nucleotides at the junction of the mouse/human sequences at the 5' of the targeting vector is set forth in SEQ ID NO: 3, and the sequence of 159 nucleotides at the junction of the human/mouse sequences at the 3' of the targeting vector is set forth in SEQ ID NO:4. Homologous recombination with this targeting vector created a modified mouse H-2K locus containing human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the HLA-A*0201 gene operably linked to the endogenous mouse H-2K transmembrane and cytoplasmic domain coding sequences which, upon translation, leads to the formation of a chimeric human/mouse MHC class I protein.

[00182] The targeted BAC DNA was used to electroporate mouse F1H4 ES cells to create modified ES cells for generating mice that express a chimeric MHC class I protein on the surface of nucleated cells (e.g., T and B lymphocytes, macrophages, neutrophils). ES cells containing an insertion of human HLA sequences were identified by a quantitative TAQMAN™ assay. Specific primer sets and probes were designed for detecting insertion of

human HLA sequences and associated selection cassettes (gain of allele, GOA) and loss of endogenous mouse sequences (loss of allele, LOA). Table 1 identifies the names and locations detected for each of the probes used in the quantitative PCR assays.

Table 1: Probes Used For Genotyping

Probe	Assay	Region Detected by Probe	Sequence	SEQ ID NO
HYG	GOA	Hygromycin cassette	ACGAGCGGGT TCGGCCCATTC	5
1665H1	GOA	Human HLA-A2 α 2- α 3 intron	AGTCCTTCAG CCTCCTCA GGTCAGG	6
1665H2	GOA	Human HLA-A2 α 2 exon	TACCACCAGT ACGCCTACGA CGGCA	7
5112H2	GOA	Human HLA-A2 α 2- α 3 intron	ATCCTGTACC AGAGAGTG	8

[00183] The selection cassette may be removed by methods known by the skilled artisan. For example, ES cells bearing the chimeric human/mouse MHC class I locus may be transfected with a construct that expresses Cre in order to remove the “/oxed” hygromycin cassette introduced by the insertion of the targeting construct containing human HLA-A*0201 gene sequences (See FIG. 5). The hygromycin cassette may optionally be removed by breeding to mice that express Cre recombinase. Optionally, the hygromycin cassette is retained in the mice.

[00184] Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, e.g., US Pat. No. 7,294,754 and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses Nature Biotech. 25(1):91-99). VELOCIMICE® (F0 mice fully derived from the donor ES cell) independently bearing a chimeric MHC class I gene were identified by genotyping using a modification of allele assay (Valenzuela *et al.*, *supra*) that detects the presence of the unique human HLA-A*0201 gene sequences.

Example 1.3. In Vivo Expression of Chimeric HLA-A/H-2K in Genetically Modified Mice.

[00185] A heterozygous mouse carrying a genetically modified H-2K locus as described in Example 1.2 was analyzed for expression of the chimeric HLA-A/H-2K protein in the cells of the animal.

[00186] Blood was obtained separately from a wild-type and a HLA-A/H-2K chimeric heterozygote (A2/H2K) mouse. Cells were stained for human HLA-A2 with a phycoerythrin-conjugated (PE) anti-HLA-A antibody, and exposed to an allophycocyanin-conjugated anti-H-2K^b antibody for one hour at 4°C. Cells were analyzed for expression by flow cytometry using antibodies specific for HLA-A and H-2K^b. FIG. 6A shows the expression of H-2K^b and HLA-A2 in the wild-type and chimeric heterozygote, with chimeric heterozygote expressing both proteins. FIG. 6B shows expression of both the H-2K^b and the chimeric HLA-A2/H2K in the heterozygous mouse.

Example 2: Construction and Characterization of Genetically Modified β 2 Microglobulin Mice

Example 2.1: Engineering a Humanized β 2 Microglobulin Locus

[00187] The mouse β 2 microglobulin (β 2m) gene was humanized in a single step by construction of a unique targeting vector from human and mouse bacterial artificial chromosome (BAC) DNA using VELOCIGENE® technology (see, e.g., US Pat. No. 6,586,251 and Valenzuela et al., *supra*).

[00188] Briefly, a targeting vector was generated by bacterial homologous recombination containing mouse β 2m upstream and downstream homology arms from BAC clone 89C24 from the RPCI-23 library (Invitrogen). The mouse homology arms were engineered to flank a 2.8 kb human β 2m DNA fragment extending from exon 2 to about 267 nucleotides downstream of non-coding exon 4 (FIG. 7). A drug selection cassette (neomycin) flanked by recombinase recognition sites (e.g., *loxP* sites) was engineered into the targeting vector to allow for subsequent selection. The final targeting vector was linearized and electroporated into a F1H4 mouse ES cell line (Valenzuela et al., *supra*).

[00189] Targeted ES cell clones with drug cassette removed (by introduction of Cre recombinase) were introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, e.g., US Pat. No. 7,294,754 and Poueymirou et al., *supra*). VELOCIMICE® (F0 mice fully derived from the donor ES cell) bearing the humanized β 2m gene were identified by screening for loss of mouse allele and gain of human allele using a modification of allele assay (Valenzuela et al., *supra*).

Example 2.2: Characterization of Humanized β 2 Microglobulin Mice

[00190] Mice heterozygous for a humanized β 2 microglobulin (β 2m) gene were evaluated for expression using flow cytometry (FIGs 8. and 9).

[00191] Briefly, blood was isolated from groups (n=4 per group) of wild type, humanized β 2m, humanized MHC (i.e., human HLA) class I, and double humanized β 2m and MHC

class I mice using techniques known in art. The blood from each of the mice in each group was treated with ACK lysis buffer (Lonza Walkersville) to eliminate red blood cells. Remaining cells were stained using fluorochrome conjugated anti-CD3 (17A2), anti-CD19 (1D3), anti-CD11b (M1/70), anti-human HLA class I, and anti-human β 2 microglobulin (2M2) antibodies. Flow cytometry was performed using BD-FACSCANTO™ (BD Biosciences).

[00192] Expression of human HLA class I was detected on cells from single humanized and double humanized animals, while expression of β 2 microglobulin was only detected on cells from double humanized mice (FIG. 8). Co-expression of human β 2m and human HLA class I resulted in an increase of detectable amount of human HLA class I on the cell surface compared to human HLA class I expression in the absence of human β 2m (FIG.9; mean fluorescent intensity of 2370 versus 1387).

Example 3. Immune Response to Flu and Epstein-Barr Virus (EBV) Peptides Presented by APCs from Genetically Modified Mice Expressing HLA-A2/H-2K and Humanized β 2 Microglobulin.

[00193] PBMCs from several human donors were screened for both HLA-A2 expression and their ability to mount a response to flu and EBV peptides. A single donor was selected for subsequent experiments.

[00194] Human T cells are isolated from PBMCs of the selected donor using negative selection. Splenic non-T cells were isolated from a mouse heterozygous for a chimeric HLA-A2/H-2K and heterozygous for a humanized β 2-microglobulin gene, and a wild-type mouse. About 50,000 splenic non-T cells from the mice were added to an Elispot plate coated with anti-human IFN γ antibody. Flu peptide (10 micromolar) or a pool of EBV peptides (5 micromolar each) was added. Poly IC was added at 25 micrograms/well, and the wells were incubated for three hours at 37°C at 5% CO₂. Human T cells (50,000) and anti-human CD28 were added to the splenic non T cells and the peptides, and the wells were incubated for 40 hours at 37°C at 5% CO₂, after which an IFN γ Elispot assay was performed.

[00195] As shown in FIG. 10, human T cells were able to mount a response to flu and EBV peptides when presented by mouse APCs that expressed the chimeric HLA-A2/H-2K and humanized β 2 microglobulin on their surface.

Equivalents

[00196] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

[00197] Entire contents of all non-patent documents, patent applications and patents cited throughout this application are incorporated by reference herein in their entirety.

CLAIMS

1. A nucleic acid comprising a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein the nucleotide sequence comprises a first nucleic acid sequence that encodes $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of a human HLA class I polypeptide operably linked to a second nucleic acid sequence that encodes transmembrane and cytoplasmic domains of the non-human MHC I polypeptide.
2. The nucleic acid of claim 1, wherein the non-human MHC I polypeptide is a rodent MHC I polypeptide, and wherein the second nucleic acid sequence encodes transmembrane and cytoplasmic domains of the rodent MHC I polypeptide.
3. The nucleic acid of claim 1 or claim 2, wherein the second nucleic acid sequence encodes transmembrane and cytoplasmic domains of the mouse MHC I polypeptide.
4. The nucleic acid of claim 3, wherein the mouse MHC I polypeptide is selected from the group consisting of H-2K, H-2D and H-2L.
5. The nucleic acid of any of the preceding claims, wherein the human HLA class I polypeptide is selected from the group consisting of HLA-A, HLA-B, and HLA-C.
6. The nucleic acid of any of the preceding claims, wherein the first nucleic acid sequence encodes $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of a human HLA-A polypeptide and the second nucleic acid sequence encodes transmembrane and cytoplasmic domains of a mouse H-2K polypeptide.
7. The nucleic acid of claim 6, wherein the HLA-A polypeptide is a HLA-A2 polypeptide.
8. The nucleic acid of claim 7, wherein the HLA-A2 polypeptide is a HLA-A2.1 polypeptide.
9. The nucleic acid of any one of claims 6 to 8, wherein the H-2K polypeptide is a H-2Kb polypeptide.
10. The nucleic acid of any one of the preceding claims, further comprising a regulatory element and a leader sequence operably linked to the nucleotide sequence.

11. The nucleic acid of claim 10, wherein the regulatory element is a non-human regulatory element and the leader sequence is selected from a human HLA class I polypeptide leader sequence, a non-human MHC I polypeptide leader sequence, and a combination thereof.
12. The nucleic acid of claim 11, wherein the regulatory element is a rodent regulatory element.
13. The nucleic acid of claim 12, wherein the regulatory element is a mouse regulatory element.
14. An isolated cell comprising the nucleotide sequence of any of the preceding claims.
15. The isolated cell of claim 14, wherein the cell expresses the chimeric human/non-human MHC I polypeptide on its surface.
16. The isolated cell of claim 14 or claim 15, further comprising a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide operably linked to endogenous non-human $\beta 2$ microglobulin regulatory elements.
17. The isolated cell of any one of claims 14 to 16, wherein the cell is an embryonic stem cell.
18. A chimeric human/non-human animal MHC I polypeptide, wherein a human portion of the chimeric polypeptide comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a human HLA class I polypeptide and a non-human portion of the chimeric polypeptide comprises transmembrane and cytoplasmic domains of a non-human animal MHC I polypeptide.
19. The chimeric human/non-human animal MHC I polypeptide of claim 18, wherein the chimeric human/non-human animal MHC I polypeptide is non-covalently bound to a human or humanized $\beta 2$ microglobulin.
20. The chimeric human/non-human animal MHC I polypeptide of claim 18 or 19, wherein the non-human portion of the chimeric human/non-human animal MHC I polypeptide comprises transmembrane and cytoplasmic domains of a rodent MHC I polypeptide.

21. The chimeric human/non-human animal MHC I polypeptide of any of claims 18 - 20, wherein the non-human portion of the chimeric human/non-human animal MHC I polypeptide comprises transmembrane and cytoplasmic domains of a mouse MHC I polypeptide.
22. The chimeric human/non-human animal MHC I polypeptide of claim 21, wherein the mouse MHC I polypeptide is selected from the group consisting of H-2K, H-2D and H-2L.
23. The chimeric human/non-human animal MHC I polypeptide of any one of claims 18 to 22, wherein the human HLA class I polypeptide is selected from the group consisting of HLA-A, HLA-B, and HLA-C.
24. The chimeric human/non-human animal MHC I polypeptide of any one of claims 18 to 23, wherein the human HLA class I polypeptide is a human HLA-A polypeptide and the non-human animal MHC I polypeptide is a mouse H-2K polypeptide.
25. The chimeric human/non-human animal MHC I polypeptide of claim 24, wherein the HLA-A polypeptide is a HLA-A2 polypeptide.
26. The chimeric human/non-human animal MHC I polypeptide of claim 25, wherein the HLA-A2 polypeptide is a HLA-A2.1 polypeptide.
27. The chimeric human/non-human animal MHC I polypeptide of any one of claims 24 – 26, wherein the H-2K polypeptide is a H-2Kb polypeptide.