

Description**Field of the Invention**

5 The present invention is directed to a non-fat cream cheese product and to a method for manufacture thereof. Generally, the method includes the steps of combining a cultured skim milk retentate, dry curd cottage cheese and non-fat dry milk with an emulsifying salt, and a gum system in a series of steps to produce a non-fat cream cheese product that resembles a reduced fat or full fat cream cheese in texture, taste and mouthfeel.

Background of the Invention

10 Cream cheese is a soft, mild acid-coagulated uncured cheese made of cream from a mixture of cream and milk. Cream cheese is stored under refrigeration conditions and the body of cream cheese is smooth and butter-like. The texture and body of cream cheese at refrigeration temperatures is such that the cream cheese can be sliced and spread. In making cream cheese, sweet whole milk and/or skim milk and sweet cream are blended in pre-selected proportions to form a cream cheese mix. The cream cheese mix normally has a butterfat content of from about 10% to about 14%. After processing, the finished cream cheese has a butterfat content of from about 33% to about 35% by weight.

20 The cream cheese mix is pasteurized and homogenized after which it is cooled, usually to a temperature between 62° F. and 92° F. (between 16.7°C and 33.3°C) and is then inoculated with a lactic acid culture. Rennet may be used to aid the coagulation of the mix. The mix is held at the inoculation temperature until it has ripened and a coagulum is formed. The acidity of the coagulum is from about 0.6% to about 0.9% (calculated as percent equivalent lactic acid).

25 After the desired acidity is obtained, the curd is separated from the whey and is thereafter packaged. One well known process for making cream cheese and separating cream cheese curd from whey includes a mechanical separation of the curd. This process is disclosed in U.S. Patent No. 2,387,176 to Link. In accordance with the method of the Link patent, after the mix is ripened to form a coagulum, the coagulum is heated to an elevated temperature to break the viscosity of the mix. Thereafter, the heated mix is centrifuged at the elevated temperature to separate the curd from the whey.

30 There has been considerable technical efforts directed to providing a cream cheese type product which has the texture, smoothness and organoleptic properties of cream cheese, but with reduced levels of fat. With increasing consumer awareness, the focal point is on reducing fat and calorie consumption. Low fat, low calorie foods which look and taste similar to their full fat, higher calorie counterparts are eagerly sought by the consumer. Researchers in the food industry have concentrated on developing food products which are nutritious and palatable, containing substantially reduced levels of high calorie fat containing ingredients. This is particularly true in the dairy industry where low calorie, low-fat products such as skim milk, yogurt and reduced fat ice cream have been successfully marketed.

40 The high fat levels in some dairy products, such as cream cheese which has a fat content of at least about 33%, have been thought to be necessary to maintain a desirable creamy mouthfeel and to avoid the grainy texture associated with prior attempts at producing low fat cream cheese products.

Many efforts have been made to develop imitation cream cheese products which contain reduced fat levels. Examples of such efforts are disclosed in U.S. Patent 2,160,159 Lundstedt, et al and U.S. Patent 3,929,892 to Hynes, et al. However, the fat content of the cream cheese products produced by the methods of these patents still exceeds about 10% fat. It would be desirable to reduce the fat content well below 10%. In particular, it would be highly desirable to provide a non-fat cream cheese product.

50 More recently, methods have been developed for making very low butterfat content imitation cream cheese products having low calorie contents which are intended for diet conscious consumers. U.S. Patent No. 4,244,983 to Baker and U.S. Patent No. 4,379,175 to Baker disclose imitation cream cheese products and a method for their manufacture having butterfat content of less than about 5%, preferably less than about 2% and which have about 60 calories per serving. However, as admitted by the inventor of these patents, in U.S. Patent No. 4,724,152 to Baker, that, while the very low butterfat content of these products is desirable, the products do not closely duplicate the creamy and full-bodied consistency of full fat cream cheese.

55 U.S. Patent No. 4,724,152 to Baker describes a method for making a low fat cream cheese product. The method includes the steps of admixing milk, a fat-containing carrier and non-fat dry milk solids to form a dressing mixture. The dressing mixture is pasteurized and homogenized and is thereafter heated to a temperature in the range of from about 145°F to about 195°F (from about 62.8°C to about 90.6°C),

preferably 165°F to 190°F (73.9°C to 87.8°C), with constant agitation. A stabilizer is admixed into the mixture. Thereafter, while maintaining the temperature of the mixture in a range of 150°F to 175°F (65.6°C to 79.4°C), soft unripened cultured cheese curd, such as cottage or baker's cheese curd is added to the stabilizer-containing dressing mix. The admixture is pumped to a homogenizer where it is homogenized under conventional elevated pressure conditions of 500 to 5000 psig (35.5 to 345.5 bar), preferably 1500 to 3000 psig (104.4 to 207.7 bar). Following homogenization, the imitation cream cheese product, at temperatures of at least 40°F (at least 4.4°C), is packaged into convenient packages.

While the '152 Baker patent provides a cream cheese product with reduced fat, i.e., a fat level in the range of 2 to 9% by weight, the Baker patent does not provide a non-fat cream cheese product.

Accordingly, it is a principal object of the present invention to provide a non-fat cream cheese type product, having the appearance, taste, consistency and texture of fat-containing cream cheese.

It is another object to provide a method for making an imitation cream cheese product which has substantially no fat and which is adapted to large scale commercial operations.

These and other objects of the invention will become more apparent from the following detailed description.

Summary of the Invention

In the method of the present invention for making a non-fat cream cheese product, a fermented skim milk retentate is introduced into a mixing tank provided with agitation means. The skim milk retentate is agitated and transferred through a recirculating loop as dry cottage cheese curd and an emulsifying salt are added to the mixing tank. Agitation and recirculation are continued for a period of time sufficient to provide a dispersion of the dry cottage cheese curd and emulsifying salt homogeneously throughout the retentate. Heating of the dispersion is then started by injecting steam into the dispersion as it passes through the recirculating loop. While the temperature of the dispersion is within the range of from about 70°F to about 150°F (about 21.1°C to about 65.6°C), non-fat dry milk is added to the dispersion. Agitation and heating is continued until the dispersion temperature is within the range of from about 140°F to about 165°F (about 60°C to about 73.9°C). Xanthan gum, as the first gum of a stabilizing system is then added to the mixing tank. Agitation and heating is continued until the dispersion temperature is in the range of from about 165°F to about 190°F (about 73.9°C to about 87.8°C). The steam is shut off and salt, sugar and carrageenan, as the second gum of a stabilizer system, are added to the mixing tank. Agitation is continued for a period of from about 1 to about 4 minutes to provide a non-fat cream cheese mix. The cream cheese mix is then homogenized to provide a non-fat cream cheese product.

Detailed Description of the Invention

Generally, in accordance with the invention, a protein base is provided from three separate concentrated skim milk sources, these being skim milk retentate, dry cottage cheese curd and non-fat dry milk solids. The skim milk retentate preferably has a non-fat milk solids content of from about 20 to about 30%. All percentages used herein are by weight based on the weight of the non-fat cream cheese product, unless otherwise indicated. The skim milk retentate is produced by ultrafiltration treatment in accordance with well known procedures. Dry cultured cottage cheese curd is a well known dairy product, which is a drained, uncreamed skim milk cheese curd made by inoculating skim milk with culture and allowing the product to coagulate. Dry cottage cheese curd may also be made by a direct acidification method. Generally, dry cottage cheese curd comprises about 17% to 26% by weight non-fat milk solids and 74% to 83% by weight water. When used herein, the term "dry cottage cheese curd" refers to a soft cheese curd such as cottage cheese or baker's cheese curd product having the above solids and moisture characteristics. Non-fat dry milk solids are produced by drying skim milk and usually have about 4% moisture.

The skim milk used to provide the dry cottage cheese curd may be subjected to well known heat treatment conditions to co-precipitate the serum proteins with that of skim milk. Such conditions are taught in U.S. Patent No. 3,039,879 to Vakaleris.

Prior to use in the method of the present invention, the skim milk retentate produced by ultrafiltration is fermented to reduce the pH of the skim milk retentate. Fermentation with a lactic starter culture is a well known cheese making step. Preferably, fermentation of the skim milk retentate proceeds to a level such that a coagulum does not form, i.e., the fermentation process is stopped while the pH is above the isoelectric point and preferably the pH is at a level of from about 4.8 to 5.2. In this connection, the use of dry cottage cheese curd provides a bland flavor component which reduces the level of flavor produced by fermenting the retentate. Accordingly, when the dry curd cottage cheese is present in the non-fat cream cheese

product at a level of from about 20% to about 50%, the pH of the retentate is preferably on the low side of the pH range, i.e., from about 4.8 to about 5.0 to provide more flavor. When dry curd cottage cheese is not used or is present at a level up to about 20%, the pH of the retentate is preferably on the higher side of the pH range, i.e., from about 5.0 to about 5.2.

5 The final fluid system of the present invention is a complex network of protein, both casein and serum protein and a particular gum system. The various components require introduction and mixing in a particular sequence to provide the highly desirable appearance, taste, consistency and texture of cream cheese resembling that of a fat-containing cream cheese, but containing substantially no fat.

10 The fermented skim milk retentate is transferred to a mixing tank provided with agitation means. The mixer is a type which imparts high shear and is preferably a blender type mixer, such as is manufactured by Bredo, inc. and sold under the trade name Bredo™. A Bredo mixer has a circular disk mounted on a rotating shaft. The circular disk is provided with a plurality of upstanding blades. The mixing action in this type of mixer is similar to that provided by a household blender. The fermented skim milk is recirculated through a recirculating loop including a conduit by means of a pump as it is being agitated. The conduit
15 exits from the bottom of the mixing tank and extends to an outlet in the top of the mixing tank.

After the fermented retentate is added to the mixer, agitation and recirculation is started and the dry curd cottage cheese, if used, and an emulsifying salt are added to the retentate. Agitation and recirculation are continued until a homogeneous dispersion of the dry curd cottage cheese and emulsifying salts are obtained prior to the start of heating by steam injection. Agitation and recirculation for a period of from
20 about 1 to about 5 minutes has been found to be sufficient. While not wishing to be bound by any theory, it is believed that the emulsifying salt serves two functions. It is believed that the emulsifying salts assists in processing the dry curd cottage cheese to break down the curd protein structure and provide a smooth homogeneous dispersion. It is also believed that the emulsifying salt serves to bind the calcium which is present in the retentate. If free calcium is present in the dispersion in any substantial amount when steam
25 heating is started, a stringy pasta filata type curd is formed at least locally at the point of steam injection. Such stringy curd is highly undesirable.

Any of the emulsifying salts commonly used in the manufacture of process cheese can be used. Such emulsifying salts include monosodium phosphate, disodium phosphate, dipotassium phosphate, trisodium phosphate, sodium metaphosphate (sodium hexametaphosphate), sodium acid pyrophosphate, sodium
30 tripolyphosphate, tetrasodium pyrophosphate, potassium citrate, calcium citrate, sodium aluminum phosphate, sodium citrate, sodium tartrate and sodium potassium tartrate. Since there is not any substantial level of fat present in the non-fat cream cheese product of the invention, the emulsifying salt is not present for the same purpose as it is used for in the manufacture of process cheese, i.e., to emulsify the fat and protein. Instead, the emulsifying salt is used as a curd processing agent and calcium binding agent as
35 indicated hereinabove.

After a homogeneous dispersion is attained, the dispersion is heated by means of steam injection. A suitable type of steam injection device is described in U.S. Patent No. 4,112,131 to Bosy, et al. It has been determined that heating by steam injection is required to effect heating of the various components providing the non-fat cream cheese product of the present invention within the time constraints required after addition
40 of the components.

In general, the remaining components which provide the non-fat cream cheese product of the present invention are added to the dispersion at particular temperatures while the dispersion is being heated to a temperature in the range of from about 165°F to 190°F (about 73.9°C to about 87.8°C) within a period of less than about 45 minutes.

45 Non-fat dry milk is added to the dispersion when the temperature of the dispersion is within the range of from about 70°F to about 150°F (about 21.1°C to about 65.6°C), preferably while the temperature is within the range of from about 140°F to about 150°F (about 60°C to about 65.6°C).

The xanthan gum, which is the first gum of the stabilizer system of the invention, is added to the dispersion after the temperature of the dispersion has increased to within the range of from about 140°F to about 165°F (about 60°C to about 63.9°C), preferably after the temperature has been increased to within the range of from about 155°F to about 165°F (about 68.3°C to about 73.9°C). After the dispersion has been heated to a temperature within the range of from about 165°F to about 190°F (about 73.9°C to about 87.8°C), preferably within the range of from about 180°F to about 190°F (about 82.2°C to about 87.8°C), the steam injection is stopped and salt, sugar and carrageenan, which is the second gum in the stabilizer
55 system of the present invention, are added to the dispersion. A preservative may also be added if increased shelf life of the non-fat cream cheese is desired. Suitable preservatives include sorbic acid, benzoic acid, potassium sorbate, sodium benzoate and calcium propionate and mixtures thereof. The preservative, if used, is preferably added at a level of from about 0.05% to about 0.2% and is added at the same time as

the salt, sugar and carrageenan.

The use of both xanthan gum and carrageenan as a stabilizer system at particular levels and ratio is important to provide the structural and organoleptic properties of the invention. The xanthan gum is used at a level of from about 0.2% to about 0.4%. Carrageenan is used at a level of from about 0.05% to about 0.3%. The ratio of the level of use of the xanthan gum to the level of use of carrageenan is preferably in the range of from about 1.5:1 to about 4:1 by weight. If the level of xanthan gum is too high, the product produced is too firm. If the level of xanthan gum is too low, the product has the consistency of sour cream. If the level of carrageenan is too low, the product has undesirable cut properties. If the level of carrageenan is too high, the product has a gummy consistency.

After the last components of the non-fat cream cheese product have been added to the dispersion while the dispersion is at a temperature in the range of about 165°F to about 190°F (about 73.9°C to about 87.8°C), steam injection is stopped. Agitation is continued for an additional period of from about 1 to about 5 minutes, and the mixture is transferred through a conventional homogenizer, such as a Gaulin homogenizer and is homogenized at about 5,000 psig (about 345.5 bar), single stage to provide a non-fat cream cheese product.

After homogenization, the non-fat cream cheese product has an aerated texture. The non-fat cream cheese product is then subjected to a deaeration step in a vacuum deaerator prior to packaging to provide a firm non-aerated product.

While the cream cheese product of the present invention is characterized as being a non-fat product, from a practical standpoint, it is impossible to remove all butterfat from milk in ordinary commercial cream separation process. Usually, a few tenths of a percent of butterfat remains in the skim milk after separating cream from the milk in the most efficient separators, i.e., skim milk usually has from about 0.05% to about 0.5% of fat. When the skim milk is further concentrated, such as by providing a skim milk retentate or dry curd cottage cheese, the butterfat content is increased in proportion to the degree of concentration. Accordingly, the non-fat cream cheese product of the present invention has from about 0.05% to about 1.5% butterfat in the final cream cheese product.

The non-fat cream cheese product of the present invention has from about 40% to about 80% of skim milk retentate. The skim milk retentate has from about 20% to about 30% solids. Dry curd cottage cheese is present in the non-fat cream cheese product at a level of from more than 0% to about 50%, preferably from about 20% to about 45%. Dry curd cottage cheese generally has a solids content of from about 20% to about 23%. The emulsifying salt is present in the non-fat cream cheese product at a level of from about 0.8% to about 1.2%. Non-fat dry milk is present in the non-fat cream cheese product at a level of from about 2% to about 4%. Non-fat dry milk generally has a solids content of from about 95% to about 97%. Salt is present at a level of from about 0.4% to about 0.8%, and sugar is present at a level of from about 0.7% to about 1.1%. Additionally, vitamins and spices may optionally be present at minor levels in the non-fat cream cheese product.

The non-fat cream cheese product of the invention has from about 23% to about 26% of total solids and from about 14% to about 18% of protein.

The following example further illustrates various features of the present invention.

Example 1

A non-fat cream cheese product is prepared in accordance with the method of the invention containing the following ingredients at the indicated levels:

Table 1

<u>Ingredient</u>	<u>Percentage As Is</u>	<u>Percent Solids</u>
UF concentrated skim milk	48.5	11.9
Dry curd cottage cheese	32.3	7.0
Water/Steam	13.2	0
Non-fat dry milk	2.7	2.6
Sodium Tripolyphosphate	0.9	0.9
Salt	0.55	0.5
Sugar	0.82	0.8
Titanium Dioxide	0.46	0.5
Xanthan Gum	0.27	0.3
Kappa Carrageenan (GP911)	0.09	0.9
Potassium Sorbate	0.06	0.06
Calcium Propionate	0.05	0.05
Starter Culture	0.025	0
Vitamin A Palmitate	0.016	0
Oleoresin Paprika	0.001	0
Total Solids		24.70

The skim milk retentate was retained in a holding tank at a temperature of 72°F (22.2°C). The retentate was inoculated with a culture of *S. lactis* and was fermented for a time of 16 hours until a pH of 5.0 was reached. 969.5 pounds (439.75 kg) of the fermented skim milk retentate were then transferred to a Bredo mixer having a capacity of 300 gallons (1135.6 litres). The mixer was turned on and 646.7 pounds (293.34 kg) of dry curd cottage cheese at a temperature of 40°F (4.4°C) was added to the mixer. 18.24 pounds (8.27 kg) of tripolyphosphate was added to the mixer and agitation and recirculation through a recirculation loop was continued for three minutes. At this time, steam at a pressure of 45 psig (4.1 bar) was injected into the recirculating loop. After reaching a temperature of 145°F (62.8°C), 54.76 pounds (24.84 kg) of non-fat dry milk was added. After reaching a temperature of 160°F (71.1°C), 5.48 pounds (2.49 kg) of xanthan gum was added. After reaching a temperature of 187°F (86.1°C), steam injection was stopped and 10.96 pounds (4.97 kg) of salt, 16.42 pounds (7.44 kg) of sugar, 9.14 pounds (4.15 kg) of titanium dioxide, 1.16 pounds (0.53 kg) of potassium sorbate, 0.92 pounds (0.42 kg) of calcium propionate, 1.8 pounds (0.82 kg) of carrageenan, 0.32 pounds (0.15 kg) of Vitamin A palmitate and 0.02 pounds (0.01 kg) of oleoresin paprika was added. Agitation of the mixture was continued for a total of 3 minutes.

The non-fat cream cheese mix resulting from the sequence of heating and addition steps described hereinabove, was then homogenized in a Gaulin homogenizer at a pressure of 5,000 psig (345.5 bar). After homogenization, the non-fat cream cheese product was transferred through a vacuum deaerator and was filled into suitable size packages. The resulting non-fat cream cheese product had the characteristics, mouthfeel and organoleptic properties of a cream cheese product that contains a substantial level of fat.

Claims

1. A method for making a non-fat cream cheese product comprising
 - introducing a cultured skim milk retentate into a mixing tank provided with agitation means,
 - adding dry cottage cheese curd and an emulsifying salt to said mixing tank and agitating said retentate for a first predetermined time period to provide a dispersion,
 - agitating said dispersion and recirculating said dispersion through a recirculating loop while injecting steam into said dispersion,
 - adding non-fat dry milk to said mixing tank at a first predetermined temperature,
 - adding xanthan gum as the first gum of a gum system to said mixing tank at a second predetermined temperature,
 - heating said dispersion to a third predetermined temperature and stopping said steam injection,
 - adding salt, sugar and carrageenan as the second gum of said gum system, to said mixing tank,
 - continuing said agitation for a second predetermined time period, and

**Table 1**

Ingredient	Percentage As Is	Percent Solids
UF concentrated skim milk	48.5	11.9
Dry curd cottage cheese	32.3	7.0
Water/Steam	13.2	0
Non-fat dry milk	2.7	2.6
Sodium Tripolyphosphate	0.9	0.9
Salt	0.55	0.5
Sugar	0.82	0.8
Titanium Dioxide	0.46	0.5
Xanthan Gum	0.27	0.3
Kappa Carrageenan (GP911)	0.09	0.9
Potassium Sorbate	0.06	0.06
Calcium Propionate	0.05	0.05
Starter Culture	0.025	0
Vitamin A Palmitate	0.016	0
Oleoresin Paprika	0.001	0
Total Solids		24.70

The skim milk retentate was retained in a holding tank at a temperature of 72°F (22.2°C). The retentate was inoculated with a culture of *S. lactis* and was fermented for a time of 16 hours until a pH of 5.0 was reached. 969.5 pounds (439.75 kg) of the fermented skim milk retentate were then transferred to a Bredo mixer having a capacity of 300 gallons (1135.6 litres). The mixer was turned on and 646.7 pounds (293.34 kg) of dry curd cottage cheese at a temperature of 40°F (4.4°C) was added to the mixer. 18.24 pounds (8.27 kg) of tripolyphosphate was added to the mixer and agitation and recirculation through a recirculation loop was continued for three minutes. At this time, steam at a pressure of 45 psig (4.1 bar) was injected into the recirculating loop. After reaching a temperature of 145°F (62.8°C), 54.76 pounds (24.84 kg) of non-fat dry milk was added. After reaching a temperature of 160°F (71.1°C), 5.48 pounds (2.49 kg) of xanthan gum was added. After reaching a temperature of 187°F (86.1°C), steam injection was stopped and 10.96 pounds (4.97 kg) of salt, 16.42 pounds (7.44 kg) of sugar, 9.14 pounds (4.15 kg) of titanium dioxide, 1.16 pounds (0.53 kg) of potassium sorbate, 0.92 pounds (0.42 kg) of calcium propionate, 1.8 pounds (0.82 kg) of carrageenan, 0.32 pounds (0.15 kg) of Vitamin A palmitate and 0.02 pounds (0.01 kg) of oleoresin paprika was added. Agitation of the mixture was continued for a total of 3 minutes.

The non-fat cream cheese mix resulting from the sequence of heating and addition steps described hereinabove, was then homogenized in a Gaulin homogenizer at a pressure of 5,000 psig (345.5 bar). After homogenization, the non-fat cream cheese product was transferred through a vacuum deaerator and was filled into suitable size packages. The resulting non-fat cream cheese product had the characteristics, mouthfeel and organoleptic properties of a cream cheese product that contains a substantial level of fat.

Claims

1. A method for making a non-fat cream cheese product comprising
 - introducing a cultured skim milk retentate into a mixing tank provided with agitation means,
 - adding dry cottage cheese curd and an emulsifying salt to said mixing tank and agitating said retentate for a first predetermined time period to provide a dispersion,
 - agitating said dispersion and recirculating said dispersion through a recirculating loop while injecting steam into said dispersion,
 - adding non-fat dry milk to said mixing tank at a first predetermined temperature,
 - adding xanthan gum as the first gum of a gum system to said mixing tank at a second predetermined temperature,
 - heating said dispersion to a third predetermined temperature and stopping said steam injection,
 - adding salt, sugar and carrageenan as the second gum of said gum system, to said mixing tank,
 - continuing said agitation for a second predetermined time period, and



homogenizing said non-fat cream cheese mix to provide a non-fat cream cheese product.

2. A method in accordance with Claim 1, wherein said first predetermined time period is from about 1 to about 5 minutes.
3. A method in accordance with Claim 1 or Claim 2, wherein said second predetermined time period is from about 1 to about 5 minutes.
4. A method in accordance with any one of Claims 1 to 3, wherein said first predetermined temperature is from about 70 °F to about 150 °F (about 21.1 °C to about 65.6 °C).
5. A method in accordance with any one of Claims 1 to 4, wherein said second predetermined temperature is from about 140 °F to about 165 °F (about 60 °C to about 73.9 °C).
6. A method in accordance with any one of Claims 1 to 5, wherein said third predetermined temperature is from about 165 °F to about 190 °F (about 73.9 °C to about 87.8 °C).
7. A method in accordance with any one of Claims 1 to 6, wherein said xanthan gum is added at a level of from about 0.2% to about 0.4%, said carrageenan is added at a level of from about 0.05% to about 0.3% and the ratio of the level of use of the xanthan gum to the level of use of carrageenan is in the range of from about 1.5:1 to about 4:1 by weight.
8. A method in accordance with any one of claims 1 to 7, wherein said skim milk retentate is present at a level of from about 40% to about 80%, said dry curd cottage cheese is present at a level of from more than 0% to about 50%, said emulsifying salt is present at a level of from about 0.8% to about 1.2%, said non-fat dry milk is present at a level of from about 2% to about 4%, said salt is present at a level of from about 0.4% to about 0.8% and said sugar is present at a level of from about 0.7% to about 1.1%.
9. A method in accordance with any one of Claims 1 to 8, wherein said non-fat cream cheese product has from about 23% to about 26% of total solids and from about 14% to about 18% of protein and has less than about 1.5% butterfat.
10. A method in accordance with any one of Claims 1 to 9, wherein a preservative is added while said dispersion is at a temperature of from about 180 °F to about 190 °F (about 82.2 °C to about 87.8 °C).
11. A method in accordance with Claim 10 wherein said preservative is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate, sodium benzoate and calcium propionate and mixtures thereof.

Patentansprüche

1. Verfahren zur Herstellung eines fettfreien Rahmfrischkäseprodukts umfassend
 - Einbringen eines Retentats von saurer Magermilch in einen Mischbehälter, der mit Rührmittel ausgestattet ist,
 - Zugabe von trockenem Hüttenkäsebruch und einem emulgierenden Salz in diesen Mischbehälter und Rühren des Retentats über einen vorbestimmten Zeitraum unter Bereitstellung einer Dispersion;
 - Rühren der Dispersion und Pumpen der Dispersion durch eine Umwälzschleife, während Dampf in die Dispersion gespritzt wird;
 - bei einer ersten vorbestimmter Temperatur Zugabe von fettfreier Trockenmilch in den Mischbehälter;
 - Zugabe von Xanthangummi als ersten Gummi eines Gummisystems in den Mischbehälter bei einer zweiten vorbestimmten Temperatur;
 - Erhitzen der Dispersion auf eine dritte vorbestimmten Temperatur und Stoppen der Dampfeinspritzung;
 - Zugabe von Salz, Zucker und Carrageenan als zweiten Gummi des Gummisystems in den Mischbehälter;



- Fortsetzen des Rührens über einen zweiten vorbestimmten Zeitraum, und
 - Homogenisieren der fettfreien Rahmfrischkäsemischung unter Bereitstellung eines fettfreien Rahmfrischkäseprodukts.
- 5 2. Verfahren nach Anspruch 1, in dem der erste vorbestimmte Zeitraum etwa 1 bis etwa 5 Minuten ist.
3. Verfahren nach Anspruch 1 oder 2, in dem der zweite vorbestimmte Zeitraum etwa 1 bis etwa 5 Minuten ist.
- 10 4. Verfahren nach einem der Ansprüche 1 bis 3, in dem die erste vorbestimmte Temperatur zwischen etwa 70 °F und etwa 150 °F (etwa 21,1 °C und etwa 65,6 °C) liegt.
5. Verfahren nach einem der Ansprüche 1 bis 4, in dem die zweite vorbestimmte Temperatur zwischen etwa 140 °F und etwa 165 °C (etwa 60 °C und etwa 73,9 °C) liegt.
- 15 6. Verfahren nach einem der Ansprüche 1 bis 5, in dem die dritte vorbestimmte Temperatur zwischen etwa 165 °F und etwa 190 °F (etwa 73,9 °C und etwa 87,8 °C) liegt.
- 20 7. Verfahren nach einem der Ansprüche 1 bis 6, in dem der Xanthangummi zu einem Level von etwa 0,2 % bis etwa 0,4 %, das Carrageenan zu einem Level von etwa 0,05 % bis etwa 0,3 % zugegeben wird, und das Verhältnis des Verwendungslevel von Xanthangummi zu dem Verwendungslevel von Carrageenan im Bereich von etwa 1,5:1 bis etwa 4:1 liegt (Gewichtsangaben).
- 25 8. Verfahren nach einem der Ansprüche 1 bis 7, in dem das Magermilch-Retentat mit einem Level von etwa 40 % bis etwa 80 % vorliegt; der trockene Hüttenkäsebruch mit einem Level von mehr als 0 % bis etwa 50 % vorliegt; das emulgierende Salz mit einem Level von etwa 0,8 % bis etwa 1,2 % vorliegt; die fettfreie Trockenmilch mit einem Level von etwa 2 % bis etwa 4 % vorliegt; das Salz mit einem Level von etwa 0,4 % bis etwa 0,8 % vorliegt; und der Zucker mit einem Level von etwa 0,7 % bis etwa 1,1 % vorliegt.
- 30 9. Verfahren nach einem der Ansprüche 1 bis 8, in dem das fettfreie Rahmfrischkäseprodukt etwa 23 % bis etwa 26 % Gesamtfestsubstanz und etwa 14 % bis etwa 18 % Protein und weniger als etwa 1,5 % Butterfett enthält.
- 35 10. Verfahren nach einem der Ansprüche 1 bis 9, in dem ein Konservierungsmittel zugesetzt wird, während die Dispersion eine Temperatur zwischen etwa 180 °F und etwa 190 °F (etwa 82,2 °C und etwa 87,8 °C) hat.
- 40 11. Verfahren nach Anspruch 1, in dem das Konservierungsmittel aus der aus Sorbinsäure, Benzoesäure, Kaliumsorbat, Natriumbenzoat, Kalziumpropionat und Mischungen derselben bestehenden Gruppe ausgewählt ist.

Revendications

- 45 1. Procédé de fabrication d'un fromage frais à teneur nulle en matière grasse comprenant les étapes suivantes :
- introduire un rétentat de lait écrémé acidifié dans un bac de mélange muni d'un moyen agitateur, ajouter du caillé de cottage cheese sec et un sel émulsifiant dans ledit bac de mélange et agiter ledit rétentat pendant un premier laps de temps prédéterminé pour provoquer une dispersion,
- 50 agiter ladite dispersion et faire recirculer ladite dispersion dans une boucle de recirculation tout en injectant de la vapeur dans ladite dispersion,
- ajouter du lait en poudre à teneur nulle en matière grasse dans ledit bac de mélange à une première température prédéterminée,
- ajouter de la gomme de xanthane comme première gomme d'un système de gomme dans ledit
- 55 bac de mélange à une deuxième température prédéterminée,
- chauffer ladite dispersion à une troisième température prédéterminée et arrêter ladite injection de vapeur,
- ajouter du sel, du sucre et du carrageen comme deuxième gomme dudit système de gomme,



dans ledit bac de mélange,
poursuivre ladite agitation pendant un second laps de temps prédéterminé, et
homogénéiser ledit mélange de fromage frais à teneur nulle en matière grasse pour obtenir un
fromage frais à teneur nulle en matière grasse.

- 5
2. Procédé selon la revendication 1, ou ledit premier laps de temps prédéterminé est d'environ 1 à environ 5 minutes.
- 10
3. Procédé selon la revendication 1 ou la revendication 2, où ledit second laps de temps prédéterminé est d'environ 1 à environ 5 minutes.
4. Procédé selon l'une quelconque des revendications 1 à 3, où ladite première température prédéterminée est comprise entre environ 70 °F et environ 150 °F (entre environ 21,1 °C et environ 65,6 °C).
- 15
5. Procédé selon l'une quelconque des revendications 1 à 4, où ladite deuxième température prédéterminée est comprise entre environ 140 °F et environ 165 °F (entre environ 60 °C et environ 73,9 °C).
6. Procédé selon l'une quelconque des revendications 1 à 5, où ladite troisième température prédéterminée est comprise entre environ 165 °F et environ 190 °F (entre environ 73,9 °C et environ 87,8 °C).
- 20
7. Procédé selon l'une quelconque des revendications 1 à 6, où on ajoute ladite gomme de xanthane dans des proportions allant environ de 0,2 % à environ 0,4 %, on ajoute ledit carragheen dans des proportions allant environ de 0,05 % à environ 0,3 % et le rapport du niveau d'utilisation de la gomme de xanthane avec le niveau d'utilisation du carragheen est compris entre environ 1,5:1 et environ 4:1 en poids.
- 25
8. Procédé selon l'une quelconque des revendications 1 à 7, où ledit rétentat de lait écrémé est présent dans des proportions allant environ de 40 % à environ 80 %, ledit caillé de cottage cheese sec est présent dans des proportions allant environ de plus de 0 % à environ 50 %, ledit sel émulsifiant est présent dans des proportions allant environ de 0,8 % à environ 1,2 %, ledit lait en poudre à teneur nulle en matière grasse est présent dans des proportions allant environ de 2 % à environ 4 %, ledit sel est présent dans des proportions allant environ de 0,4 % à environ 0,8 % et ledit sucre est présent dans des proportions allant environ de 0,7 % à environ 1,1 %.
- 30
9. Procédé selon l'une quelconque des revendications 1 à 8, où ledit fromage frais à teneur nulle en matière grasse a environ de 23 % à environ 26 % d'extrait sec total et environ de 14 % à environ 18 % de protéines et a moins de 1,5 % environ de matière grasse.
- 35
10. Procédé selon l'une quelconque des revendications 1 à 9, où l'on ajoute un conservateur pendant que ladite dispersion est à une température comprise entre environ 180 °F à environ 190 °F (entre environ 82,2 °C à environ 87,8 °C).
- 40
11. Procédé selon la revendication 10, où ledit conservateur est sélectionné parmi un groupe constitué d'acide sorbique, d'acide benzoïque, de sorbat de potassium, de benzoate de sodium et de propionate de calcium et d'un mélange de ceux-ci.
- 45
- 50
- 55