

5

PHARMACEUTICAL COMPOSITION FOR USE IN THE TREATMENT OR
PREVENTION OF OSTEOARTICULAR DISEASES

10 Field of the invention

[0001] The present invention is in the pharmaceutical field and is related to a new pharmaceutical composition for use in the treatment or the prevention of the acute and/or chronic osteoarticular diseases.

15 Background of the invention

[0002] Osteoarthritis, the most common form of arthritis, is a disease characterised by a slow degenerative processes in the articular cartilage, subchondral bone associated with marginal osteophyte formation, and low grade inflammation. Osteoarthritis is believed to affect 15% of the population in its chronic form. Of those, one-quarter are severely disabled. Most cases of osteoarthritis have no known cause and are referred to as primary osteoarthritis. When the cause of the osteoarthritis is known, the condition is referred to as secondary osteoarthritis. Secondary osteoarthritis is caused by another disease or condition. Conditions that can lead to secondary osteoarthritis include repeated trauma or surgery to the joint structures, abnormal joints at birth (congenital abnormalities), gout, diabetes, and other hormone disorders. Other forms of arthritis are systemic illnesses, such as rheumatoid arthritis and systemic lupus erythematosus (SLE).

[0003] Osteoarthritis involves mainly the hips, knees, spine, and the interphalangeal joints. The most common symptom of osteoarthritis is pain in the affected joint(s) after repetitive use. Joint pain is usually worse
5 later in the day. There can be swelling, warmth, and creaking of the affected joints. Pain and stiffness of the joints can also occur after long periods of inactivity. In severe osteoarthritis, complete loss of cartilage cushion causes friction between bones, causing pain at rest or pain
10 with limited motion.

[0004] Osteoarthritis is characterized by a slow degradation of cartilage over several years. In normal cartilage, a delicate balance exists between matrix synthesis and degradation; in osteoarthritis, however,
15 cartilage degradation exceeds synthesis. The balance between synthesis and degradation is affected by age and is regulated by several factors produced by the synovium and chondrocytes, including cytokines, growth factors, aggrecanases, and matrix metalloproteinases. In addition to
20 water, the extracellular matrix is composed of proteoglycans, made up of glycosaminoglycans attached to a backbone made of hyaluronic acid, entrapped within a collagenous framework or fibrillary matrix. A significant proteoglycan in articular cartilage is aggrecan, which
25 binds to hyaluronic acid and helps provide the compressibility and elasticity of cartilage. Aggrecan is cleaved by aggrecanases, leading to its degradation and to subsequent erosion of cartilage. The loss of aggrecan from the cartilage matrix is one of the first pathophysiological
30 changes observed in OA.

[0005] Cytokines produced by the synovium and chondrocytes, especially IL-1 β and Tumor Necrosis Factor alpha (TNF- α), are also key players in the degradation of cartilage. IL-1 β is spontaneously released from

cartilage of osteoarthritis but not normal cartilage. Both IL-1beta and TNF-alpha stimulate their own production and the production of other cytokines (e.g., IL-8, IL-6, and leukotriene inhibitory factor), proteases, and prostaglandin E₂ (PGE₂). Synthesis of the inflammatory cytokines IL-1beta and TNF-alpha and expression of their receptors are enhanced in osteoarthritis. Both cytokines have been shown to potently induce degradation of cartilage *in vitro*. Other proinflammatory cytokines overexpressed in osteoarthritis include IL-6, IL-8, IL-11, and IL-17, as well as leukotriene inhibitory factor.

[0006] The extracellular matrix (ECM) composing the cartilage is degraded by locally produced matrix metalloproteinases. Proinflammatory cytokines, including IL-1beta, TNF-alpha, IL-17, and IL-18, increase synthesis of matrix metalloproteinases, decrease matrix metalloproteinase enzyme inhibitors, and decrease extracellular matrix synthesis.

[0007] In an attempt to reverse the breakdown of the extracellular matrix, chondrocytes increase synthesis of matrix components including proteoglycans. Even though this activity increases, a net loss of proteoglycans in the upper cartilage layer is seen. Elevated anti-inflammatory cytokines found in the synovial fluid of osteoarthritis include IL-4, IL-10, and IL-13. Their role is to reduce production of IL-1beta, TNF-alpha, and matrix metalloproteinases, and inhibit prostaglandin release. Local production of growth and differentiation factors such as insulin-like growth factor 1, transforming growth factors, fibroblastic growth factors, and bone morphogenetic proteins also stimulate matrix synthesis.

State of the Art

[0008] Currently available pharmacological therapies target palliation of pain and include analgesics (i.e. acetaminophen, cyclooxygenase-2-specific inhibitors, non-selective nonsteroidal anti-inflammatory drugs, tramadol, 5 opioids). However, the clinical presentation of osteoarthritis is usually monoarticular or oligoarticular with fluctuations in intensity and localisation over time. It is therefore logical to consider local therapeutic 10 modalities in order to avoid untoward systemic effects. Several compounds have been used intra-articularly (e.g., glucocorticoids, hyaluronic acid) or topically (e.g., capsaicin, methylsalicylate). However, the benefit of intra-articular glucocorticoides lasts only a few days 15 (Barron, M. C., 2007, J. Am. Osteopath., 107, ES21-27).

[0009] Polysaccharides form a class of materials whose recognition of the potential utility is growing. Apart from their biological activity, one of the most important properties of polysaccharides in general is their 20 ability to form hydrogels. Hydrogel formation can occur by a number of mechanisms and is strongly influenced by the types of monosaccharide involved, as well as the presence and nature of substituent groups. Polysaccharide gel formation is generally of two types: hydrogen bonded and 25 ionic. Hydrogen-bonded gels are typical of molecules such as agarose (thermal gellation) and chitosan (pH-dependent gellation), whereas ionically bonded gels are characteristic of alginates and carrageenans.

[0010] Proteoglycans are one of the major 30 macromolecules found in articular cartilage. These molecules consist of a core protein and covalently attached glycosaminoglycans (GAG) chains. The GAGs are long, unbranched heteropolysaccharides, consisting of repeated

disaccharide units, with the general structure: Muronic acid amino sugar. The cartilage-specific GAGs include chondroitin 4-sulfate (glucuronic acid and N-acetyl-galactosamine with an SO₄ on the 4-carbon position),
5 chondroitin 6-sulfate (glucuronic acid and N-acetyl-galactosamine with an SO₄ on the 6-carbon position) and keratan sulfate (galactose and N-acetyl-glucosamine with an SO₄ on the 6-carbon position).

[0011] These molecules are capable of forming
10 hydrogel complexes with oppositely charged ionic polymers, particularly the cationic polysaccharide chitosan. This interaction may form the basis of a new materials approach to cartilage tissue engineering. The other important cartilage GAG is hyaluronic acid (glucuronic acid and N-
15 acetyl-glucosamine). This molecule is one of the major components in synovial fluid. Hyaluronic acid molecules are also present in cartilage matrix as the backbone structure in proteoglycan aggregates. In general, hyaluronic acid plays a major role as an organizer of the extracellular
20 matrix. Purified hyaluronic acid is employed as a structural biomaterial because of its high molecular weight and gel forming ability. The properties of the molecule may be broadly altered by chemical modification. For example, partial esterification of the carboxyl groups reduces the
25 water solubility of the polymer and increases its viscosity. Extensive esterification generates materials that form water-insoluble films or swellable gels. Ethyl and benzyl esterified hyaluronate membranes have excellent healing responses and biodegradability properties. The
30 fully esterified membranes have in vivo lifetimes of several months, whereas the partially esterified forms have been shown to degrade within a few weeks.

[0012] Hyaluronic acid is responsible for the viscoelastic quality of synovial fluid that acts as both a

lubricant and shock absorber. In synovial fluid, hyaluronic acid coats the surface of the articular cartilage and shares space deeper in the cartilage among collagen fibrils and sulfated proteoglycans. It protects the cartilage and blocks the loss of proteoglycans from the cartilage matrix into the synovial space, maintaining the normal cartilage matrix. In synovial fluid from knee joints in osteoarthritis, concentrations of hyaluronic acid, glycosaminoglycans, and keratan sulfate are lower than in synovial fluid from normal knee joints. Additionally, experiments using rabbit synovial cells showed that the proinflammatory cytokines IL-1 and TNF-alpha stimulate the expression of hyaluronic acid synthetase, which may contribute to the fragmentation of hyaluronic acid under inflammatory conditions. Exogenous hyaluronic acid may facilitate the production of newly synthesized hyaluronic acid. Hyaluronic acid and derivatives have been used as therapeutic aids in the treatment of osteoarthritis as a means of improving lubrication of articulating surfaces and thus reducing joint pain. Several in vitro culture studies have also demonstrated that hyaluronic acid has a beneficial effect by inhibiting chondrocytic chondrolysis mediated by fibronectin fragment. Hyaluronic acid has also been shown to have anti-inflammatory effects, as well as inhibitory effects on prostaglandin synthesis, and proteoglycan release and degradation.

[0013] Chitosan, a partially de-acetylated derivative of chitin, found in arthropod exoskeletons is another proteoglycan, a linear polysaccharide consisting of beta(1-4) linked D-glucosamine residues with a variable number of randomly located N-acetyl-glucosamine groups. It thus shares some characteristics with various GAGs and hyaluronic acid present in articular cartilage, Depending on the source and preparation procedure, chitosan's average

molecular weight may range from about 50 to about 1000 kDa. Commercially available preparations have degrees of deacetylation ranging from about 50 to about 90%. Chitosan is a semi-crystalline polymer and the degree of crystallinity is a function of the degree of deacetylation. Crystallinity is maximum for both chitin (i.e. 0% deacetylated) and fully deacetylated (i.e. 100%) chitosan. Minimum crystallinity is achieved at intermediate degrees of deacetylation.

10 [0014] Much of the potential of chitosan as a biomaterial stems from its cationic nature and high charge density. The charge density allows chitosan to form insoluble ionic complexes or complex coacervates with a wide variety of water-soluble anionic polymers.

15 [0015] In fact, chitosan oligosaccharides have been shown to have a stimulatory effect on macrophages, and the effect has been linked to the acetylated residues. Furthermore, both chitosan and its parent molecule, chitin, have been shown to exert chemoattractive effects on
20 neutrophils in vitro and in vivo. In vivo, chitosan is degraded by enzymatic hydrolysis. The mechanical properties of chitosan scaffolds are mainly dependent on the pore sizes and pore orientations.

[0016] Hyaluronic acid, a glycosaminoglycan, is
25 widely used for the treatment of osteoarthritis of the knee. A survey of 2 general practices in the United Kingdom showed that about 15% of patients with osteoarthritis received intra-articular treatment with hyaluronic acid preparations. Because of its viscoelastic quality, it may
30 replace synovial fluid. Furthermore, it may reduce the perception of pain. Beneficial molecular and cellular effects have also been reported. Hyaluronic acid is frequently applied by intra-articular injection, but the evidence concerning its clinical relevance is conflicting.

State-of-the-art systematic reviews and meta-analyses were published recently, and their authors concluded that intra-articular hyaluronic acid, at best, has a small effect, a clinically meaningful effect meaning an improvement of at least 15 mm on the visual analog scale of pain (Bellamy et al, 2006; Cochrane Database Syst Rev. 2006 Apr 19;(2):CD005321). These data form the basis for the use of intra-articular administration of hyaluronic acid in patients with osteoarthritis. The benefits are sometimes noticed only one year after injections and, in some experiments, injections must be performed three to five-times a week (Barron, M. C., 2007, J. Am. Osteopath., 107, ES21-27).

[0017] Alpha-2-adrenergic receptor ligands especially agonists are drugs commonly used in medical practice as antihypertensive substance and in clinical anesthesiology as component of general and locoregional anesthesia and analgesia. They produce anxiolysis, analgesia, sedation, anesthetic sparing effects and peri-operative hemodynamic stabilizing effects. Negative neurotoxicity studies allow their use (mainly clonidine) by systemic and perimedullar routes and for peripheral nerve blocks. Among the clinically available alpha-2-adrenoreceptors agonists, clonidine remains widely used: the substance is devoid of neurotoxicity and displays less side effects (i.e. hypotension and sedation) than the more potent and also more alpha-2-adrenergic receptor selective agonist, dexmedetomidine. Clonidine, a potent alpha-2-adrenergic receptors partial agonist, was used primarily for the treatment of hypertension. This drug stimulates alpha-2-adrenergic receptors in the vasomotor centers, causing a reduction of sympathetic outflow from the central nervous system. Both cardiac output and peripheral resistance are reduced resulting in a decrease in blood

pressure. Higher concentrations cause a vasoconstriction by activation of postsynaptic receptors in vascular smooth muscle. However, the significant advantages of the drug are counter balanced by side effects including dryness of the mouth, sedation and dizziness. Furthermore, other activities of these compounds such as anti-inflammatory effect have never been reported by oral administration.

[0018] Besides the well-known analgesic effects of spinally administered alpha-2-adrenergic (alpha-2-adrenoceptors) agonists, their peripheral use has been commonly reported in acute pain conditions. For peri-operative analgesic techniques, clonidine is added to local anaesthetic in peritroncular nerve blocks to enhance potency and duration of analgesia. ZOUHER A et al.; Paediatr Anaesth. 2005 Nov;15(11):964-70; Luiz-Cleber P. et al. ; Anesth Analg. 2005 Sep;101(3):807-11, Murphy DR "A non-surgical approach to low back pain" Med. Health R. I., 2000 April; 83(4): 104-7). Further, intra-articular injection of clonidine and its adjunction to local anaesthetic solution for intravenous regional anaesthesia have also displayed anti-nociceptive effect S. Armand et al; Br J Anaesth. 1998 Aug;81(2):126-34; Gentili M, et al. Pain 1996; 64: 593-596; Reuben S, et al. Anesthesiol. 1999; 91: 654-658. However, the effect is here designed not to last for days or weeks.

[0019] Alpha-2-adrenoceptors agonists are known to block the tissue content increase of pro-inflammatory cytokines, such as TNF-alpha and IL-1beta and increase the tissue content of anti-inflammatory cytokine TGFbeta. This has been shown in inflammatory neuropathic pain model by partial ligation of sciatic nerve, by applying locally alpha 2 adrenergic receptor agonists by peripheral nerve block.

Summary of the invention

[0019a] In the first aspect, there is provided a pharmaceutical composition, comprising

- possibly an adequate pharmaceutical carrier or diluent
- a glycosaminoglycan being hyaluronic acid or its derivative
- an effective amount of an anti-inflammatory compound selected from the group consisting of an anti-TNF agent, a steroidal compound, a disease modifying antirheumatic drug, an anticytokine agent or a mixture thereof and
- an effective amount of mesenchymal stem cells with the proviso that the stem cells are not human embryonic stem cells.

[0019b] In the second aspect, there is provided a kit of parts comprising

- a vial with the composition according to any of the preceding claims,
- a device for delivering the said composition to an inflamed joint of a mammal subject and having
- reservoir means for storing the said composition,
- piston means movable along the longitudinal axis of the reservoir for dispensing the said pharmaceutical composition,
- and an hollow needle mounted on said reservoir means for delivering the said pharmaceutical composition to the peripheral nerve of the mammal subject.

[0019c] In the third aspect, there is provided the use of the pharmaceutical composition as defined above in the preparation of a medicament for the treatment and/or the prevention of acute and/or chronic osteoarticular diseases and/or symptoms by intra-articular injection.

[0020] The invention is related to a new (intra-articular) pharmaceutical composition for use in the treatment and/or the prevention of acute or chronic
5 osteoarticular diseases and acute or chronic osteoarticular symptoms - especially osteoarthritis, this composition comprising

- possibly an adequate pharmaceutical carrier or diluent
- a polysaccharide and/or a glycosaminoglycan,
10 preferably a glycosaminoglycan (including proteoglycan),
- stem cells (differentiated or not), but with the proviso that these stem cells are not human embryonic stem cells, having potential anti-inflammatory
15 properties and
- a sufficient amount of an anti-inflammatory agent (compound).

[0021] In the pharmaceutical composition of the invention the polysaccharide or the glycosaminoglycan
20 (including the proteoglycan) are present as a film or a matrix, preferably in the form of a paste or a gel, more preferably an hydrogel with a sufficient amount of an aqueous solvent.

[0022] Preferably, the anti-inflammatory agent
25 (compound) is selected from the group consisting of a
steroidal (prednisolone, dexamethasone, betamethasone,
triamcinolone...), a non-steroidal anti-inflammatory
compound (ibuprofen, diclofenac, naproxen, cox-2 inhibitors
etc...), a disease modifying antirheumatic drug (DMARD such
30 as methotrexate, leflunomide, etc...), an alpha 2
adrenergic receptor agonist, an anti-CD20 agent, an anti-

cytokine agent (anti-IL1, anti-IL6, anti-IL17), an anti-TNF agent (infliximab, etanercept, adalimumab, rituximab, etc) or a mixture thereof.

[0023] Advantageously, the anti-inflammatory
5 compound is a compound interacting with the alpha-2-adrenergic receptor, preferably an alpha-2-adrenergic receptor agonist.

[0024] The alpha-2-adrenergic receptor agonist is selected from the group consisting of clonidine, p-aminoclonidine, tiamenidine, 5-bromo-6-(2 imidazolidine-2-ylamino) quinoxaline, dexmedetomidine, detomidine, medetomidine, alphas-methyldopa, oxymetazoline, brimonidine, tizanidine, mivazerol, lofexidine, xylazine, guanfacine, guanclofine, guanoxabenz, or a derivative or structural
10 analogue thereof, alpha-methylnorephrine, azepexole, indoramin, 6-allyl-2-amino-5,6,7,8-tetrahydro4H-thiazolo [4,5-d]azepine diHCl or a compound identified in the table 1.

[0025] In the composition according to the invention
20 the polysaccharide-based hydrogel and the glycosaminoglycan are not covalently linked or are covalently linked.

[0026] Preferably, the glycosaminoglycan is selected from the group consisting of hyaluronic acid with low (< 900 kDa) or high (> 900 kDa) molecular mass.

25 [0027] In the composition of the invention, the hyaluronic acid and the alpha 2 adrenergic receptor agonist are not covalently linked or are covalently linked.

[0028] In the composition according to the invention the glycosaminoglycan is selected from the group consisting
30 of proteoglycan, chondroitin sulfate, keratin sulphate, hyaluronic acid, (including their derivative), chitosan, a chitosan or chitin derivative, or a mixture thereof.

[0029] Advantageously, the non human embryonic stem cells of the composition are selected from the group

consisting of (cells obtained from) bone marrow or adipose tissue or bone tissue or joint tissue, a mix of hematopoietic and mesenchymal stromal cells, (expanded or not), mesenchymal stromal cells, osteoblastic cells, 5 chondrocytic cells or a combination thereof.

[0030] The inventors have discovered that the pharmaceutical composition according to the invention is suitable for a treatment or prevention of acute and/or chronic osteoarticular diseases and associated symptoms 10 (especially osteoarticular pain, mobility or function), of inflammatory origin, such as osteoarthritis, degenerative arthristis, degenerative arthritis, gonarthrosis, coxarthrosis, and other inflammatory general conditions in which joints are involved, such as autoimmune diseases, 15 especially rheumatoid arthritis and systemic lupus erythematosus (SLE) spondyloarthropathies, polymyalgia rheumatica, ankylosing spondylitis, Reiter's Syndrome, psoriatic arthropathy, enteropathic arthritis (related to inflammatory bowel disease, such as haemorrhagic colitis and Crohn's disease), neuropathic arthropathy, acute 20 rheumatic fever, gout, chondrocalcinosis, calcium hydroxyapatite crystal deposition disease, Lyme disease and all other degenerative joint diseases.

[0031] The composition of the invention is also 25 suitable for obtaining an efficient proliferation of stem cells therefore, for improving regenerative or damaged tissues. This effect upon regeneration of treated tissues is advantageously induced by the synergic effect of the active compounds present in the composition of the 30 invention.

[0032] Therefore, the composition according to the invention is characterized by an anti-inflammatory activity, a tissue or cell regenerative effect, and an advantageous efficient treatment and/or prevention of acute

and/or chronic osteoarticular diseases and/or symptoms. These effects could be combined in the composition according to the invention.

[0033] According to a preferred embodiment of the present invention the composition presents a preferred formulation, preferably being an injectable solution for delivering an efficient amount (therapeutic dose) of the active compound present in the composition to a mammal subject including a human patient, especially in the knees, the hips and the spine of the subject.

[0034] Preferably, this injectable solution comprises from about 0.1 mg to about 100 mg/kg preferably from about 1 mg to about 10 mg/kg of body weight of this polysaccharide and/or glycosaminoglycan, preferably in the form of a polysaccharide-based hydrogel and may comprise from about 0.1 mg to about 100 mg/kg of patient body weight preferably from about 0.1 mg to about 0.8 mg/kg of patient body weight of an anti-inflammatory compound, being preferably a compound activating the alpha-2-adrenergic receptor, preferably an alpha-2-adrenergic receptor agonist.

[0035] The injectable solution is adequate for intra-articular administration (percutaneous injection) in a joint of a mammal subject, preferably of a human patient.

[0036] This formulation is also adequate for local administration (percutaneous injection in/or in the vicinity of an inflamed joint of a mammal subject, preferably of a human patient), local administration injection that does concern the epidermis, the muscle or any deep organs.

[0037] Another aspect of the present invention is related to the use of the pharmaceutical composition for the manufacture of a medicament in the treatment and/or the

prevention of these mentioned diseases and/or symptoms (pain) induced by these diseases.

[0038] The present invention is related to the surprising discovery that this intra-articular
5 administration to a mammal subject, particularly a human patient of this pharmaceutical composition results in an improvement of symptoms associated with osteoarticular diseases, such as a relief of osteoarticular pain, an improvement of joint mobility and/or function, a decrease
10 in articular accumulation of inflammatory liquid, induced by the above mentioned diseases or pathologies.

[0039] The present invention is also related to the surprising discovery that the intra-articular
15 administration to a mammal subject, particularly a human patient of the pharmaceutical composition according to the invention results in both a reduction of inflammation, and of joint degeneration, resulting in disease improvement and joint regenerative effects, combined with the above effects in the improvement of symptoms associated with
20 osteoarticular diseases. These effects are obtained by the synergic combination of the elements present in the composition.

[0040] The present invention is also related to the surprising discovery that the intra-articular
25 administration to a mammal subject, particularly a human patient of the pharmaceutical composition according to the invention results also in a shortening of the time to onset of the therapeutic activity of the first component, an increase in the duration of action of the first component.

30 [0041] The composition and method of the present invention also allows a decrease of the number of required administrations to obtain a desired efficacy and allows for a faster onset of action.

[0042] The invention is also directed to a method for a treatment and/or a prevention (prophylaxis) of a condition selected from the group consisting of the above mentioned diseases or symptoms through an administration to
5 a mammal subject, preferably a human patient of a sufficient amount of the pharmaceutical composition of the invention, through an intra-articular administration, particularly in the knees, the hips and the spine or any joint (e.g., interphalangeal) of the mammal subject;
10 preferably the human patient.

[0043] A last aspect of the present invention is related to a kit of parts comprising one or more vials with the elements (carrier/diluent, polysaccharide, glycosaminoglycan, anti-inflammatory agent (or compound),
15 stem cells or a combination thereof) of the composition of the invention and a device for delivering these elements (simultaneously or successively) or this composition to an inflamed joint of a mammal subject, preferably a human patient suffering from the above mentioned osteoarticular
20 diseases or symptoms (in particular pain) and having

- reservoir means for storing this pharmaceutical composition,
- piston means movable along the longitudinal axis of the reservoir for dispensing this pharmaceutical
25 composition and,
- a hollow needle mounted on said reservoir means for delivering this pharmaceutical composition to the peripheral nerve of the mammal subject.

[0044] These and other objects and features of the
30 invention will become more fully apparent when the following detailed description of the invention is read in conjunction with the accompanying example.

Detailed description of the invention

[0045] The term "intra-articular" as used herein refers to a percutaneous injection of a joint with the pharmaceutical composition of the invention.

5 [0046] The term "local administration" as used herein refers to a percutaneous injection in or in the vicinity of an inflamed joint. Local administration injection thus concerns the epidermis, the dermis, the muscle, or any deep organ.

10 [0047] The main advantages to local administration are to selectively restrict the analgesic effect to the injured areas. Furthermore, the local administration allows for high local concentration levels with little or no systemic release. The local administration and the intra-
15 articular injection of hyaluronic acid is a recognised treatment of the above mentioned conditions.

[0048] The local administration and the intra-articular injection of the anti-inflammatory compound and stem cells are easily realizable and provide long lasting
20 effect in combination with proteoglycans. In consequence, the problems related to placement of an invasive drug delivery system and the problems of bothersome side-effects, due to systemic release, can be strongly minimized. Health-related quality of life, patient
25 satisfaction and economic assessment might be improved with such a treatment, especially in chronic conditions.

[0049] Preferably, the administered proteoglycan is selected from the group consisting of a polysaccharide-based hydrogel or a glycosaminoglycan, for example
30 hyaluronic acid or a salt thereof or an ester of hyaluronic acid with an alcohol of the aliphatic, heterocyclic or cycloaliphatic series, or a sulphated form of hyaluronic acid or combination of agents containing hyaluronic acid. Suitable dosages of a polysaccharide-based hydrogel or a

glycosaminoglycan or a derivative thereof will typically be from about 0.1 mg to about 100 mg /kg body weight per day or from about 0.5 mg to about 10 mg/kg body weight per day more preferably from about 2 mg to about 8 mg by body weight per day.

[0050] Advantageously, the administrated stem cells are from bone marrow, adipose tissue, bone tissue or joint tissue, composed of a mix of hematopoietic and mesenchymal stromal cells, of mesenchymal stromal cells - expanded or not, of osteoblastic cells or chondrocytic cells or any combination thereof.

[0051] According to a first embodiment, the administrated stem cells are osteoblastic cells, or preferably a cell population selected from the group consisting of osteoprogenitors, pre-osteoblasts and osteoblasts.

[0052] According to the second embodiment, the administrated stem cells are chondrocytic cells, or even preferably a cells population selected from the group consisting of chondroprogenitors, pre-chondrocytes and chondrocytes.

[0053] In a preferred embodiment, the alpha 2 adrenergic receptor agonist may be clonidine, p-aminoclonidine, tiamenidine, 5-bromo-6-(2 imidazolidine-2-ylamino) quinoxaline, dexmedetomidine, detomidine, medetomidine, alphas-methyldopa, oxymetazonline, brimonidine, tizanidine, mivazerol, lofexidine, xylazine, guanabenz, guanfacine, guanclofine, guanoxabenz, or a derivative or structural analogue thereof, alpha-methylnorephrine, azepevole, indoramin, 6-allyl-2-amino-5,6,7,8-tetrahydro4H-thiazolo [4,5-d]azepine diHCl analogs thereof or a compound selected from the table 1 and analogs thereof.

Table 1:**Classification of the alpha-2-adrenergic receptor agonists****guanidines**

- 5 agmatine
betanidine
biguanides
cimetidine
creatine
10 gabexate
guanethidine
guanfacine
guanidine
impromidine
15 iodo-3 benzylguanidine
methylguanidine
mitoguazone
nitrosoguanidines
pinacidil
20 robenidine
sulfaguanidine
zanamivir

25 imidazoles

- 4- (3-butoxy-4-méthoxybenzyl)imidazolidin-2-one
acide urocanique
amino-imidazole carboxamide
antazoline
30 biotine
bis(4-methyl-1-homo piperazinylthiocarbonyl)disulfide
carbimazole
cimetidine
clotrimazole
35 creatinine
dacarbazine
dexmedetomidine
econazole
enoximone
40 ethymizol
etomidate
fadrozole
fluspirilène

histamine
histidinol
idazoxan

5 imidazolidines

imidazolines
clonidine
tolazoline
impromidine

10 levamisole
losartane
medetomidine
miconazole
naphazoline

15 niridazole
nitroimidazoles
ondansetron
oxymetazoline
phentolamine

20 tetramisole
thiamazole
trimetaphan

25 Derivatives of clonidine

2,6-dimethylclonidine
4-azidoclonidine
4-carboxyclonidine-methyl 3,5-dichlorotyrosine
4-hydroxyc lonidine

30 4-iodoclonidine
alinidine
apraclonidine
chlorethylclonidine
clonidine 4-isothiocyanate
35 clonidine 4-methylisothiocyanate
clonidine receptor
clonidine-displacing substance
hydroxyphenacetyl aminoclonidine
N,N'-dimethylclonidine

40

[0054] The active compounds used in accordance with the invention are known. Pharmaceutical preparations containing hyaluronic acid are commercially available as are clonidine and other alpha 2 adrenergic receptor agonists. The compounds can be manufactured in a known manner essentially in accordance with the processes described in the prior art.

[0055] The term "clonidine" as used herein refers to N-(2,6-dichlorophenyl)-4,5-dihydro-1H-imidazol-2-amine and includes the pharmaceutically acceptable salts thereof, e.g., salts with inorganic acids, such as hydrohalic acids, or with organic acids, for example lower aliphatic monocarboxylic or dicarboxylic acids such as acetic acid, fumaric acid or tartaric acid or aromatic carboxylic acids such as salicylic acid are also suitable.

[0056] Clonidine is employed in a therapeutically effective amount. The present invention also encompasses the use of alpha 2 adrenergic receptor agonist for the manufacture of injectable formulations for a delivery to a joint or a close region thereof, a therapeutic dose of said agonist by intra-articular injection, wherein said solution comprises from about 3 µg to about 1500 µg of said alpha 2 adrenergic receptor agonist. Preferably, said formulation comprises from about 30 µg to about 500µg of said alpha 2 adrenergic receptor agonist. More preferably said solution comprises from 50 µg to 350 µg of said alpha 2 adrenergic receptor agonist. In a preferred embodiment, the alpha 2 adrenergic receptor agonist is clonidine. The actual concentration of clonidine may vary, depending on the nature and degree of the pain syndromes being treated and whether the drug is being administered for therapeutic or prophylactic purposes.

[0057] It is hypothesized that based on their anti-inflammatory properties, inhibition of pro-inflammatory

cytokines TNF-alpha and IL1-beta and the increase of anti-inflammatory cytokines such as TGFbeta, alpha 2 receptor agonists in injectable administration will have applications in osteoarticular inflammatory conditions and
5 in diseases where there are osteoarticular inflammatory conditions, such as those above mentioned.

[0058] According to the method and the formulation of the invention, an injection of the combination of hyaluronic acid and an alpha 2 adrenergic receptor agonist
10 form induces a long lasting pain relief, both in human case and in an animal model of osteoarticular pain. The method is safe, devoid of major drug's side effects and allows for acute as well as chronic treatment and or prophylaxis without the use of too invasive technique.

15 [0059] Preservatives may be incorporated in an amount effective for inhibiting growth of microbes, such as bacteria, yeast and molds, in the composition. Any conventional preservative against microbial contamination of the product can be employed so long as it is
20 pharmaceutically acceptable and is unreactive with clonidine. Preferred preservatives are antimicrobial aromatic alcohols, such as benzyl alcohol, phenoxyethanol, phenethyl alcohol, and the like, and esters of parahydroxybenzoic acid commonly referred to as paraben
25 compounds, such as methyl, ethyl, propyl, and butyl esters of parahydroxybenzoic acid and the like and mixtures thereof, but are not limited thereto. Particularly preferred are benzyl alcohol and phenoxyethanol.

[0060] Optionally, anaesthetic agent, such as
30 lidocaine and the like, can be included. For administration according to the invention the active quantities of the compounds that alleviate neuropathic pain can be contained

together with customary pharmaceutical excipients and/or additives in solid or liquid pharmaceutical formulations.

[0061] Liquid preparations such as solutions, suspensions or emulsions of the active ingredients, prepared by dissolving or dispersing the compounds, can contain the usual diluents such as water, oil and/or suspending aids such as polyethylene glycols and optional pharmaceutical adjuvants, in a carrier, such as, for example, aqueous saline, aqueous dextrose, glycerol, or ethanol, to form a solution or suspension and such like. Further additives such as preservatives, flavouring agents and such like may also be added.

Example

[0062] The effects of the combination of alpha 2 adrenoreceptor agonists (e.g., clonidine), of a glycosaminoglycan (e.g., hyaluronic acid) and of stem cells were tested in an inflammation model with stimulated Peripheral Blood Mononuclear Cells.

PBMCs (Peripheral blood mononuclear cells) from heparinized venous blood of healthy volunteers were isolated by Ficoll-gradient centrifugation. MNC (Mononuclear cells) were washed three times in PBS and resuspended in RPMI-1640 medium supplemented with 100 U/mL penicillin, 100µg/mL streptomycin, and 10% heat-inactivated FBS (Fetal Bovine Serum). Cells were seeded at 100.000 cells in 96-well plate in a total volume of 200µL/well.

PBMC, at 100.000 cells/200µL, were plated in 96-well microtiter plates and stimulated or not with 10µg/mL phytohemagglutinin (PHA). Effects of hyaluronic acid "HA"

(200µg/mL) and p-aminoclonidine "A2A" (5µM) were tested on PBMC.

MSC (Mesenchymal stromal cells) were derived from 20 to
5 60mL of heparinized bone marrow (BM) obtained from the
iliac crest of healthy volunteers. BM was mixed with
phosphate-buffered saline (PBS, 2v:v) and layered on
density gradient Ficoll solution. After centrifugation,
mononuclear cells were harvested from the interface and
10 washed twice in PBS. Cells were resuspended in standard MEM
medium supplemented with 15% FBS. The cells were plated at
2x10⁶ cells/75cm² flasks and maintained in a 37°C
humidified atmosphere containing 5% CO₂. Cells were allowed
to attach for 24 hours prior an initial medium change.
15 Cells were detached at day 10 using trypsin- EDTA solution
for 1-5 min at 37°C. Then, cells were irradiated (10
minutes at 2200Ci) before seeding in 96-well for
proliferative assays at 12.500 cells/well to obtain a ratio
1:8 (MSC:PBMC).

20

Proliferation Assay

The culture was incubated with 1µCi/mL 3H-thymidine for 24
hours before end of culture to measure the proliferation.
Then, cells were washed twice with ice-cold PBS and twice
25 with ice-cold 5% trichloroacetic acid (TCA). Finally, cells
were lysed with 0.1N NaOH-0.1% Triton-X100. Supernatants
were collected and analyzed on a beta-counter in presence
of scintillation liquid. Results were done in cpm ("count
per minute" = disintegration number of radioactive element
30 observed in series of successive counts of one minute).

IL-1β (beta) detection (By ELISA). Levels of IL-1β from
PBMC or culture supernatants were measured by Quantikine
ELISA kit (R&D Systems Inc, Mineapolis, USA). The minimum

detectable concentration of IL-1 β was estimated at 1.0 pg/mL.

The release of IL-1beta by PBMC and the proliferation of
 5 PBMC, induced by PHA stimulation, were set as controls, and compared with the measures for PBMC treated with HA and/or A2A.

The inhibition of PBMC release of IL-1beta and the
 10 inhibition of PBMC proliferation, induced by LPS stimulation of TH1 cells, were also measured and provided similar trends as for PHA stimulated cells.

15 Effects of HA alone

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
PBMC	ND*	769
PBMC + PHA	2185	16069
PBMC + HA	2095	364
PBMC + PHA + HA	6394	9087

*not determined

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC	ND*	5
PBMC + PHA	100	100
PBMC + HA	96	2.2
PBMC + PHA + HA	293	57

*:not determined

20

IL-1beta secretion induced by PHA stimulation shows a significant increase over basal condition (where IL-1beta levels were undetectable) and was set at 100. Proliferation of PMBC cells also significantly increased, by a factor of

about 20. Surprisingly, a similar cytokine increase (over basal condition) was observed with addition of HA alone, while PBMC proliferation was not affected. When stimulated by PHA, the addition of HA synergistically increases IL-1beta secretion, 3 times over the stimulated conditions, but was able to reduce PBMC proliferation by a factor 2.

Effects of A2A alone

10

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
PBMC	ND*	769
PBMC + A2A	ND*	786
PBMC + PHA	2185	16069
PBMC + PHA + A2A	ND*	16212

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC	ND*	5
PBMC + A2A	ND*	4.9
PBMC + PHA	100	100
PBMC + PHA + A2A	ND*	101

*:not determined

15 A2A had no effect on basal (unstimulated) conditions. Interestingly, when stimulated by PHA, the addition of A2A totally suppressed the increase in IL-1beta secretion due to PHA stimulation, to undetectable levels, but was not able to inhibit PBMC proliferation.

20

Effects of Combination of A2A & HA

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
PBMC + PHA	2185	16069
PBMC + PHA + HA	6394	9087
PBMC + PHA + HA + A2A	3053	5445

*:not determined

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC + PHA	100	100
PBMC + PHA + HA	292	57
PBMC + PHA + HA + A2A	139	34

*:not determined

- 5 Interestingly, the combination HA/A2A demonstrates a potent anti-inflammatory effect over the PHA/HA condition. Indeed, the addition of A2A was able to significantly revert (by over 50%) the increase in IL-1 β secretion due to PHA/HA stimulation, bringing its level to the one of the PHA alone condition. Surprisingly, A2A was further able to inhibit PBMC proliferation, by another 35%.

Effects of MSC addition

- Testing the effects of MSC on both PBMC-proliferation and IL-1 β secretion, a stimulatory effect was observed over basal (unstimulated) conditions. Surprisingly, the proliferative effect was maintained in stimulated conditions (e.g., presence of PHA), but instead of a pro-inflammatory effect on IL-1 β secretion, a potent inhibitory effect was observed (- 62%).

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
PBMC	ND*	769
PBMC + MSC	229	1612
PBMC + PHA	2185	16069
PBMC + PHA + MSC	841	26192

*:not determined

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC	ND*	100
PBMC + MSC	+++	209
PBMC + PHA	100	100
PBMC + PHA + MSC	38	163

*:not determined

In PHA stimulated conditions and in presence of hyaluronic acid, MSCs were modestly (about 10%) anti-inflammatory and anti-proliferative.

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
PBMC	ND*	769
PBMC + PHA	2185	16069
PBMC + PHA + HA	6394	9087
PBMC + PHA + HA + MSC	5769	8673

*:not determined

10

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC	ND*	5
PBMC + PHA	100	100
PBMC + PHA + HA	292	57
PBMC + PHA + HA + MSC	264	54

*:not determined.

Finally in PHA stimulated conditions and in presence of A2A, MSCs had no influence.

Conditions tested	IL-1 β detection (pg/mL)	Proliferation (cpm)
-------------------	--------------------------------	---------------------

PBMC	ND*	769
PBMC + PHA	2185	16069
PBMC + PHA + A2A	ND	16212
PBMC + PHA + A2A + MSC	314	17061

*:not determined

Conditions tested	IL-1 β detection (%)	Proliferation (%)
PBMC	ND*	5
PBMC + PHA	100	100
PBMC + PHA + A2A	ND	100
PBMC + PHA + A2A + MSC	14	106

*:not determined

5

During inflammation, lower molecular weight fragments of hyaluronan are known to be inflammatory and immune-stimulatory agents by inducing the secretion of cytokines such as IL-6 and monocyte chemoattractant protein (MCP-1).

- 10 HA has also the property to enhance the adhesion of lymphocytes and monocytes to the extracellular matrix (Yamawaki et al., 2009). The present experiments show that hyaluronic acid has a potent stimulatory effect on cytokine (e.g., IL-1 β) production without PBMC proliferation.

15

- It has been demonstrated that clonidine alters the Th1/Th2 cytokine production (Xu et al., 2007; Cook-Mills et al., 1998). The present experiments show that the addition of alpha 2 adrenoreceptor agonists totally suppresses the
- 20 increase in cytokine (e.g., IL-1 β) secretion due to PHA stimulation, but is not able to inhibit PBMC proliferation.

- Surprisingly, the combination HA/A2A demonstrates a potent anti-inflammatory effect both on cytokine production and
- 25 inflammatory cell proliferation. Indeed, the addition of

A2A to HA was able to significantly inhibit the increase in IL-1beta secretion and was further able to inhibit PBMC proliferation.

- 5 An in vitro property of MSCs is their ability to reduce the proliferation of lymphocytes of various types while enjoying immune privilege in vitro and in vivo (Bocelli-Tyndall et al., 2007). However, it was shown that the number of MSCs and the incubation time were key factors in
- 10 MSCs' lymphocyte regulation: inhibition of lymphocyte proliferation was dose-dependent on MSC concentrations (LeBlanc in 2003).

- Interestingly, these results show that MSC have a moderate
- 15 stimulatory effect on unstimulated PBMC by increasing both IL-1beta secretion and increase PBMC proliferation. On the contrary, MSC displayed a potent anti-inflammatory effect (-62%) on IL-1beta secretion but increased PBMC proliferation. The inventors have hypothesized that MSCs
- 20 did not induce proliferation of PBMC, but were able to increase their survival rate. This is illustrated by the attachment of PBMCs on the surface of MSCs (see enclosed figure 1). This seems to be confirmed by the increase in viability observed with PHA stimulated PBMCs in presence of
- 25 MSCs; a viability increased by over 20%.

Regenerative effects

MSCs were culture in chondrogenic medium for 7, 14 and 21 days and 4 conditions were tested

- 30 - control
- A2A: 5µMol aminoclonidine
 - HA: 200µg/ml of hyaluronic acid
 - HA + A2A: 200µg/ml of hyaluronic acid and 5µMol aminoclonidine

At day 14 and 21, pellets were fixed in 10% buffered formalin for 6 hours, and washed twice in PBS. Pellets were then embedded in OCT (Sakura Finetek, Belgium). Sections
5 were cut at a thickness of 5 μ m and were stained with Alcian blue, before counterstained with Nuclear fast red (Klinipath, Belgium)

Alcian blue is one of the most widely used cationic dyes
10 for the demonstration of glycosaminoglycans (GAGs). GAGs will be stained blue, while Nuclear fast red will counterstain cells nuclei in pink or red, and cytoplasm in pale pink. In brief, the procedure is 1) Stain in alcian blue solution for 30 minutes ; 3) Wash in running tap water
15 for 2 minutes ; 4) Rinse in distilled water; 5) Counterstain in nuclear fast red solution for 5 minutes; 6) Wash in running tap water for 1 minute; 7) Dehydrate through 95% alcohol, and 2 changes of absolute alcohol, 3 minutes each; 8) clear in Ultraclear, a xylene substitute
20 (Klinipath, Belgium); 9) Mount with Ultrakit mounting medium (Klinipaht, Belgium). Toluidine blue stains proteoglycan in red-purple (metachromatic staining) and the nucleus in blue (orthochromatic staining). Metachromasia, tissue elements staining a different color from the dye solution, is due to
25 the pH, dye concentration and temperature of the basic dye. Blue or violet dyes will show a red color shift, and red dyes will show a yellow color shift with metachromatic tissue elements. Toluidine Blue is an often used to stain proteoglycans and glycosaminoglycans in tissues such as
30 cartilage.

Results. Although pellets from different conditions were of similar size (mean of 0.7mm; range from 0.9 to 0.4mm), Alcian Blue staining reveals that cellularity of pellets

was significantly affected by the changes in conditions. For example, the addition of hyaluronic acid strongly reduced the cellularity of the pellet. In parallel, matrix (i.e., mucopolysaccharides and glycosaminoglycans) content was almost absent in control and A2A alone conditions as illustrated in Fig 2 by the absence of blue staining while in the hyaluronic acid and hyaluronic acid/clonidine conditions, the slides were uniformly blue stained, demonstrating an important matrix production. The combined condition HA/A2A appears to further increase the matrix production over HA alone condition.

Fig 2: Alcian blue staining at low magnification power (Top) and high magnification power (Bottom) for the different culture conditions: control, A2A (clonidine alone), HA (hyaluronic acid alone) and HA/A2A (combination of hyaluronic acid and clonidine)

Interestingly, cell nucleus staining demonstrate that the addition of clonidine (HA/A2A) to hyaluronic acid (HA) increases the cellularity of pellet.

In conclusion, the combination of hyaluronic acid, an anti-inflammatory compound such as an alpha2 adrenoreceptor agonist and mesenchymal stromal cell displays surprisingly synergistic anti-inflammatory and regenerative effects:

- The presence of mesenchymal stromal cells are required for the *de novo* production of cartilage matrix;
- This production is significantly increased (and accelerated) by the addition of hyaluronic acid, but the addition of hyaluronic acid has potent pro-inflammatory effects;
- These hyaluronic acid potent pro-inflammatory effects are totally neutralized by the addition of alpha2-adrenergic agonists;

- The addition of alpha2-adrenergic agonist agents supports the regenerative effects of the combination of mesenchymal stromal cells and hyaluronic acid by increasing the cellularity of the cell preparation and the production of matrix.

References

- Bocelli-Tyndall et al. (Rheumatology; 46:403-408 (2007)).
- Bondeson et al. (Arthritis Res Ther; 8(6):R187 (2006)).
- Cook-Mills et al., 1995. (Immunology. 85 544-549 (1995)).
- 5** Le Blanc et al., (Scandinavian Journal of Immunology, 57, 11-20 (2003)).
- Levesque et al., (The Journal of Immunology, 166: 188-196 (2001)).
- Mchugh et al. (Clin Exp Immunol. 99:160-167 (1995)).
- 10** Pelletier et al. (Arthritis Rheum. 44(6):1237-47 (2001)).
- Yamawaki H et al. (Glycobiology, 19, 83-92 (2009)).

CLAIMS

1. A pharmaceutical composition, comprising
 - possibly an adequate pharmaceutical carrier or diluent
 - a glycosaminoglycan being hyaluronic acid or its derivative
 - an effective amount of an anti-inflammatory compound selected from the group consisting of an anti-TNF agent, a steroidal compound, a disease modifying antirheumatic drug, an anticytokine agent or a mixture thereof and
 - an effective amount of mesenchymal stem cells with the proviso that the stem cells are not human embryonic stem cells.
2. The pharmaceutical composition according to the claim 1 for use in the treatment and/or the prevention of acute and/or chronic osteoarticular diseases and/or symptoms by intra-articular injection.

3. The pharmaceutical composition according to any of the preceding claims, which comprises from 0.1 to 100 mg/kg of body weight of glycosaminoglycan.

4. The pharmaceutical composition according to any of the preceding claims, which comprises from 0.1 to 100 mg/kg of body weight of anti-inflammatory compound.

5. The pharmaceutical composition according to any of the preceding claims, wherein the mesenchymal stem cells are human adult mesenchymal stem cells or non human embryonic mesenchymal stem cells.

6. The pharmaceutical composition according to any of the preceding claims which is an injectable solution.

7. The pharmaceutical composition of claim 6, wherein the injection of the injectable solution is done twice a week.

8. The pharmaceutical composition of claim 6, wherein the injection of the injectable solution is done once a week.

9. The pharmaceutical composition of claim 6, wherein the injection of the injectable solution is done once in two or three weeks.

10. The pharmaceutical composition according to any of the preceding claims, wherein the acute or chronic osteoarticular symptom is selected from the group consisting of pain, mobility and/or function.

11. The pharmaceutical composition according to any of the preceding claims, wherein the osteoarticular disease is selected from the group consisting of osteoarthritis, degenerative arthritis, gonarthrosis, coxarthrosis, and other inflammatory general conditions or symptoms in which joints are involved, such as systemic lupus erythematosus (SLE), spondyloarthropathies, polymyalgia rheumatica, ankylosing spondylitis, Reiter's Syndrome, psoriatic arthropathy, enteropathic arthritis (related to inflammatory bowel disease such as haemorrhagic colitis and Crohn's disease) rheumatoid arthritis, neuropathic arthropathy, acute rheumatic fever, gout, chondrocalcinosis, calcium hydroxyapatite crystal deposition disease, and Lyme disease.

12. A kit of parts comprising

- a vial with the composition according to any of the preceding claims,
- a device for delivering the said composition to an inflamed joint of a mammal subject and having
- reservoir means for storing the said composition,
- piston means movable along the longitudinal axis of the reservoir for dispensing the said pharmaceutical composition,
- and an hollow needle mounted on said reservoir means for delivering the said pharmaceutical composition to the peripheral nerve of the mammal subject.

13. Use of the pharmaceutical composition according to any of claims 1 to 11 in the preparation of a medicament for the treatment and/or the prevention of acute and/or chronic osteoarticular diseases and/or symptoms by intra-articular injection.