CLAIMS

[Claim 1]

A urination disorder-alleviating agent comprising acotiamide or a salt thereof as an active ingredient, wherein the acotiamide or a salt thereof is orally administered in an amount of 200 to 300 mg per administration for a total of 600 to 900 mg per day to an adult.

[Claim 2]

The urination disorder-alleviating agent according to claim 1, wherein an administration subject is a patient selected from the group consisting of patients with underactive bladder and patients with underactive bladder accompanied by overactive bladder.

[Claim 3]

The urination disorder-alleviating agent according to claim 1, wherein the number of oral administrations of acotiamide or a salt thereof per day is 2 to 3.

[Claim 4]

The urination disorder-alleviating agent according to claim 1, wherein an administration subject is a patient with underactive bladder.

[Claim 5]

Use of acotiamide or a salt thereof for producing a urination disorder-alleviating agent, wherein the

acotiamide or a salt thereof is orally administered in an amount of 200 to 300 mg per administration for a total of 600 to 900 mg per day to an adult.

[Claim 6]

The use according to claim 5, wherein an administration subject is a patient selected from the group consisting of patients with underactive bladder and patients with underactive bladder accompanied by overactive bladder.

[Claim 7]

The use according to claim 5, wherein the number of oral administrations of acotiamide or a salt thereof per day is 2 to 3.

[Claim 8]

The use according to claim 5, wherein an administration subject is a patient with underactive bladder.

[Claim 9]

Acotiamide or a salt thereof for use in alleviating urination disorder, wherein the acotiamide or a salt thereof is orally administered in an amount of 200 to 300 mg per administration for a total of 600 to 900 mg per day to an adult.

[Claim 10]

The acotiamide or a salt thereof according to claim

9, wherein an administration subject is a patient selected from the group consisting of patients with underactive bladder and patients with underactive bladder accompanied by overactive bladder.

[Claim 11]

The acotiamide or a salt thereof according to claim 9 or 10, wherein the number of oral administrations per day is 2 to 3.

[Claim 12]

The acotiamide or a salt thereof according to any one of claims 9 to 11, wherein an administration subject is a patient with underactive bladder.