

SINGAPORE PATENTS ACT (CHAPTER 221)
PATENTS RULES

PATENTS FORM 1/8



**Request for the Grant of a Patent / Statement of
Inventorship and Right to Grant of Patent**

Pre-requisites:

- a. Among other requirements, a Date of Filing will be issued only if a description of the invention is filed or if a complete and valid statement is made under Part 9 of this form.

Estimated Time:

This form may take approximately 15 - 20 minutes to complete.

General:

- a. * denotes mandatory field.

PART 1 Claiming the Filing Date of an Earlier Singapore Application

This application is

☒

A divisional of an earlier Singapore application.

Application No.

11201809928V

Filing Date
(DD/MM/YYYY)

25 May 2017

☐

An application filed in response to an order by the Registrar after determination of a question regarding the entitlement of an earlier Singapore application.

Application No.

Filing Date
(DD/MM/YYYY)

I choose not to rely on the auto-populated information, changes will be made to the auto-populated information as below.

Yes ☐ No ☒

PART 2 Reference

Applicant/Agent
Reference

WT/JAN/ksh/C.20221702 (4861/SG/2)

PART 3 Title of Invention*

Title of Invention*

Modified Antimicrobial Peptide Derived From An Arginine-rich Domain

PART 4 Details of Applicant*

Note: If there is insufficient space, please use the continuation sheet CS 1.

Applicants

Type of Applicant

☐

Corporate

☒

Individual

UEN/ Company Code

Name

SHIH, Chiaho

Address Type

☐

Singapore

☒

Foreign

	Address	128 Academia Rd. Sec. 2 Nangung, Institute of Biomedical Science, Academia Sinica, Taipei City, 11529, Taiwan
	Is SME	☐ Yes ☐ No
	Citizenship	United States of America
	Residency Country	
	Sole Proprietor or Partners' Names	

PART 5 Declaration of Priority under Section 17(2)

Declaration of Priority(s) under Section 17(2)	Application No.	Country	Filing Date
	62/342,415	U.S.A.	27/05/2016

PART 6 Inventor*

A. Are all applicants named above also inventors?	Yes ☐ No ☒
B. If yes, are there any other inventors?	Yes ☐ No ☐

PART 7 Details of Inventor

Note: If there is insufficient space, please use the continuation sheet CS 5.

Inventor	Name	CHEN, Heng-Li
	Company UEN	
	Care Of	
	Address Type	☐ Singapore ☒ Foreign
	Address	128 Academia Sinica Rd., Sec. 2 Nangung, Institute of Biomedical Sciences, Academia Sinica, Taipei City, 11529, Taiwan
	Nationality	
	Country of Residency	Taiwan, Republic of China
	The above inventor was resident of Singapore at any time during the period of invention ☐	

PART 8 Derivation of Right

The Applicants Derived the Right from the Inventor to be Granted the Said Application* (Note: Only one checkbox may be crossed)	☐ By Contract of Employment
	☒ By Assignment of Invention

☐ Others (Please specify below)	
PART 9 Micro-Organism Deposited under the Budapest Treaty	
☐	The invention requires the use of a micro-organism and for disclosure purposes, a culture of the micro-organism has been deposited for with an international depository authority under the Budapest Treaty.
PART 10 Description Incorporated by Reference to an Earlier Relevant Application	
☐	No description is filed with this request. Instead the description of the invention for which the patent is sought is incorporated in the application by reference to, and is completely contained in, the following earlier relevant application, as filed: Country <input type="text"/> Application No. <input type="text"/> Filing Date (DD/MM/YYYY) <input type="text"/>
PART 11 Contact Details*	
<i>Note:</i> a. This part need not be filled up if the person's address in Part 4 is to be used as an address for service in Singapore. b. The email address and telephone number provided are for contact purposes in case of query. c. The official correspondence will be sent electronically via IP2SG.	
Agent UEN/Company Code	<input type="text" value="200601697N"/>
Agent Name	<input type="text" value="AMICA LAW LLC"/>
Representative or C/O Name	<input type="text"/>
	Address for Service in Singapore <input type="text"/>
	Block/ House No. <input type="text" value="77"/>
	Street Name <input type="text" value="Robinson Road"/>
	Level - Unit <input type="text" value="#22-01"/>
	Building Name <input type="text" value="Robinson 77"/>
	Postal Code <input type="text" value="068896"/>
Contact Person	<input type="text" value="Winnie Tham"/>
Direct Telephone No.	<input type="text" value="63036217"/>
Email Address	<input type="text" value="winnie.tham@amicalaw.com"/>
PART 12 Checklist*	

Checklist A	The application consists of the following no. of sheets	
Description	<input type="text" value="20"/>	sheet(s)
Claim	<input type="text" value="5"/>	sheet(s)
Drawing	<input type="text" value="9"/>	sheet(s)
Abstract	<input type="text" value="1"/>	sheet(s)
Sequence Listing	<input type="text" value="13"/>	sheet(s)
<i>(Note: Please indicate on the abstract the figure of the drawing, if any, which should accompany the abstract when published)</i>		
Total number of sheets	<input type="text" value="48"/>	sheet(s)
PART 13 Declaration*		
Declaration	<p><u>Statement</u></p> <p>I/ We believe that the person(s) stated in Part 7 is/ are the inventor(s) of the invention in respect of which the patent application is made.</p> <p>I/ We also consent to the publication of the details contained herein to each of the inventors as stated in Part 7.</p> <p><u>By Person Filing the Application</u></p> <p>I do hereby declare that the information furnished above is true to the best of my knowledge. I understand that I may be liable for criminal prosecution for providing any false information in this application.</p> <p><u>By Agent</u></p> <p>I do hereby declare that:</p> <ol style="list-style-type: none"> I have been duly authorised to act as an agent, for the purposes of this application, on behalf of the person(s) filing this form. The information furnished above on behalf of the person(s) filing this form is true to the best of the person(s)' knowledge. I understand that I may be liable for criminal prosecution for providing any false information in this application. 	
Name	<input type="text" value="Winnie Tham"/>	
Signature		
Date (DD/MM/YYYY)	<input type="text" value="27 Mar 2022"/>	
No. of Extra Sheets Attached to this Form	<input type="text" value="6"/>	sheet(s)

GST Information

The tax invoice will be issued to:

SHIH, Chiaho

Name:

SHIH, Chiaho

Application Number:

INTELLECTUAL PROPERTY OFFICE OF SINGAPORE

CS1  *1013*	CONTINUATION SHEET 1
PART 1	
Continuation Sheet to Part Application Number <i>(if any)</i>	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; width: 100px; text-align: center;">1</div> <div>of Form</div> <div style="border: 1px solid black; width: 100px; text-align: center;">1/8</div> </div> <div style="border: 1px solid black; height: 20px; margin-top: 5px;"></div>
PART 2 Details of Person/ Requestor/ Applicant/ Proprietor/ Registered Owner/ Holder Licensor/ Licensee/ Grantor/ Opponent	
UEN/ Company Code <i>(if applicable)</i> Name	<div style="border: 1px solid black; height: 20px; margin-bottom: 5px;"></div> <div style="border: 1px solid black; padding: 5px; min-height: 40px;"> ACADEMIA SINICA </div>
Address	Singapore Address ☐ This Singapore address is to be used as the address for service for the purposes of this application. Block/ House No. <div style="border: 1px solid black; width: 100px; height: 20px;"></div> Street Name <div style="border: 1px solid black; width: 300px; height: 25px;"></div> Level - Unit <div style="display: flex; align-items: center;"> <div style="border: 1px solid black; width: 40px; height: 20px;"></div> <div style="margin: 0 5px;">-</div> <div style="border: 1px solid black; width: 60px; height: 20px;"></div> </div> Building Name <div style="border: 1px solid black; width: 300px; height: 25px;"></div> Postal Code <div style="border: 1px solid black; width: 100px; height: 20px;"></div>
	Foreign Address Line 1 <div style="border: 1px solid black; padding: 2px 5px;">Office of Public Affairs (Technology Transfer)</div> Line 2 <div style="border: 1px solid black; padding: 2px 5px;">(05) 128, Sec. 2, Academia Sinica Rd.</div> Line 3 <div style="border: 1px solid black; padding: 2px 5px;">Nan-Kang, Taipei, Taiwan</div>
Nationality or Country of Incorporation* State of Incorporation <i>(mandatory for USA corporations)</i>	<div style="border: 1px solid black; padding: 5px; margin-bottom: 5px;">Taiwan, Republic of China</div> <div style="border: 1px solid black; height: 20px;"></div>

Country of Residency
(mandatory for individuals)

**Sole Proprietor or
Partners' Name**
*(if sole proprietorship or
partnership)*

Company Profile
*(for Singapore incorporated
businesses only. Please cross
if applicable)*

☐

Small and Medium Enterprise¹

¹ Small and medium-sized enterprises in Singapore will be defined as businesses with annual sales turnover of not more than \$100 million or employing no more than 200 staff.

INTELLECTUAL PROPERTY OFFICE OF SINGAPORE

CS5  *1017*	CONTINUATION SHEET 5
PART 1	
Continuation Sheet to Part Application Number <i>(if any)</i>	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; width: 100px; height: 25px; text-align: center;">2</div> <div>of Form</div> <div style="border: 1px solid black; width: 100px; height: 25px; text-align: center;">1/8</div> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <div style="border: 1px solid black; width: 100%; height: 25px;"></div> </div>
PART 2 Details of Inventor*	
Name	<div style="border: 1px solid black; padding: 5px; min-height: 40px;">SHIH, Chiaho</div>
Address	<div> Singapore Address ☐ This Singapore address is to be used as the address for service for the purposes of this form. </div> <div style="margin-top: 10px;"> Block/ House No. <div style="border: 1px solid black; width: 100px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Street Name <div style="border: 1px solid black; width: 300px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Level - Unit <div style="border: 1px solid black; width: 50px; height: 25px;"></div> - <div style="border: 1px solid black; width: 100px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Building Name <div style="border: 1px solid black; width: 300px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Postal Code <div style="border: 1px solid black; width: 100px; height: 25px;"></div> </div>
	<div>Foreign Address</div> <div style="margin-top: 10px;"> Line 1 <div style="border: 1px solid black; width: 300px; height: 25px; padding: 2px;">128 Academia Rd. Sec 2 Nangung,</div> </div> <div style="margin-top: 10px;"> Line 2 <div style="border: 1px solid black; width: 300px; height: 25px; padding: 2px;">Institute of Biomedical Science Academia Sinica</div> </div> <div style="margin-top: 10px;"> Line 3 <div style="border: 1px solid black; width: 300px; height: 25px; padding: 2px;">Taipei City, 11529, Taiwan</div> </div>
Nationality* State of Residency <i>(mandatory for USA)</i> Country of Permanent Residence* <i>(mandatory for individuals)</i>	<div style="border: 1px solid black; padding: 5px; min-height: 20px;">United States of America</div> <div style="border: 1px solid black; width: 200px; height: 25px; margin-top: 10px;"></div> <div style="border: 1px solid black; padding: 5px; min-height: 20px; margin-top: 10px;">Taiwan, Republic of China</div>

The Above Inventor was
Resident in Singapore at
Any Time during the
Period of Invention?

☐

Yes

☒

No

INTELLECTUAL PROPERTY OFFICE OF SINGAPORE

CS5  *1017*	CONTINUATION SHEET 5
PART 1	
Continuation Sheet to Part Application Number <i>(if any)</i>	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; padding: 2px 10px; text-align: center;">3</div> <div>of Form</div> <div style="border: 1px solid black; padding: 2px 10px; text-align: center;">1/8</div> </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> <div style="border: 1px solid black; width: 100%; height: 20px;"></div> </div>
PART 2 Details of Inventor*	
Name	<div style="border: 1px solid black; padding: 5px; min-height: 40px;"> SU, Pei-Yi </div>
Address	<div> Singapore Address ☐ This Singapore address is to be used as the address for service for the purposes of this form. </div> <div style="margin-top: 10px;"> Block/ House No. <div style="border: 1px solid black; display: inline-block; width: 60px; height: 20px;"></div> </div> <div style="margin-top: 10px;"> Street Name <div style="border: 1px solid black; display: inline-block; width: 350px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Level - Unit <div style="border: 1px solid black; display: inline-block; width: 60px; height: 20px;"></div> - <div style="border: 1px solid black; display: inline-block; width: 100px; height: 20px;"></div> </div> <div style="margin-top: 10px;"> Building Name <div style="border: 1px solid black; display: inline-block; width: 350px; height: 25px;"></div> </div> <div style="margin-top: 10px;"> Postal Code <div style="border: 1px solid black; display: inline-block; width: 80px; height: 20px;"></div> </div>

The Above Inventor was
Resident in Singapore at
Any Time during the
Period of Invention?

☐

Yes

☒

No

TITLE OF INVENTION: MODIFIED ANTIMICROBIAL PEPTIDE DERIVED FROM AN ARGININE-RICH DOMAIN

BACKGROUND

5 Antibiotics have been used for the treatment of bacterial infection for more than 60 years. Recently, the increasing number of antibiotics-resistant bacteria has become a major threat to public health. The development of new antibiotics for clinical treatment is an urgent need. Antimicrobial peptides (AMPs) from various species can serve as a defense weapon of the host against pathogenic microbes. Because they can kill bacteria and fungi via different
10 mode of actions, they have been considered as potential candidates to overcome the problem of antibiotic-resistance.

SUMMARY

In one aspect, described herein is an antimicrobial peptide. The peptide contains 2 to 20 variable domains, each variable domain is a sequence of 2 to 20 consecutive basic amino
15 acids, wherein (a) the variable domains are separated from each other by a variable linker, (b) the variable linker can have 1 to 20 any amino acids other than two or more consecutive basic amino acids, and (c) the peptide has no more than 100 amino acids. In one embodiment, peptide has at least 3 or 4 variable domains. The peptide can have a C-terminal cysteine. In some embodiments, at least one of the basic amino acids in the variable domains is an
20 arginine. For example, all of the basic amino acids in each variable domain in the peptide can be arginine residues. Alternatively or in addition, at least one variable domain in the antimicrobial peptide has a lysine. In one embodiment, at least one variable domain has a histidine. The peptide can have a cyclic structure.

At least one of the basic amino acids in the peptide can be a chemically-modified
25 amino acid. In one embodiment, the chemically-modified amino acid is a D-amino acid, e.g., D-arginine. The variable domains and the variable linkers can be derived from the arginine-rich domain of a hepadnavirus core protein (HBcARD). In one embodiment, the HBcARD contains a sequence from residue 147 to the C-terminal residue of a hepadnavirus core protein. Each variable domain in the peptide can have three or four arginine residues and
30 each variable linker in the peptide can have 2 to 4 amino acids. The peptide can exhibit a

broad spectrum antimicrobial activity against a gram-positive bacterium, gram-negative bacterium, fungus, parasite, or virus.

In one embodiment, the antimicrobial peptide contains a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)$ (SEQ ID NO: 1), wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , and X_8 , individually, is any amino acid or absent,

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)$ (SEQ ID NO:2), wherein each of X_1 , X_3 , and X_6 , individually, is a variable domain, and each of X_2 , X_4 , and X_5 , individually, is any amino acid or absent, and

(iii) $(X_1)X_2PX_3P(X_4)X_5QX_6P(X_7)$ (SEQ ID NO: 3), wherein each of X_1 , X_4 , and X_7 , individually, is a variable domain, and each of X_2 , X_3 , X_5 , and X_6 , individually, is any amino acid or absent. Each variable domain is a sequence of 2 to 20 consecutive basic amino acids.

The peptide can have a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}C$ (SEQ ID NO: 4), and

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}Q$ (SEQ ID NO: 5), wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , X_8 , and X_{10} , individually, is any amino acid or absent.

In one embodiment, the antimicrobial peptide contains a sequence selected from the group consisting of:

(i) TVVRRRGRSPRRRTSPRRRRSQSPRRRRSQSRESQC (SEQ ID NO: 6), in which at least one of the arginine residues is D-arginine,

(ii) RRRGRSPRRRTSPRRRRSQSPRRRRSC (SEQ ID NO: 7),

(iii) RRRGRSPRRRTSPRRRRSQSPRRRRSQ (SEQ ID NO: 8),

(iv) RRRGRPRRRPPRRRRQPRRRRC (SEQ ID NO: 9),

(v) RRRGRSPRRRTSPRRRRRC (SEQ ID NO: 10),

(vi) RRRGRPRRRPPRRRRRC (SEQ ID NO: 11),

(vii) RRRTSPRRRRSQSPRRRRRC (SEQ ID NO: 12), and

(viii) RRRPPRRRRQPRRRRC (SEQ ID NO: 13).

For example, the antimicrobial peptide can contain the sequence of RRRGRSPRRRTSPRRRRSQSPRRRRSC (SEQ ID NO: 7), in which each of the arginine residues in the sequence is L-arginine. Alternatively, the peptide can have the sequence of

RRRGRSPRRRTSPRRRRSQSPRRRRSC (SEQ ID NO: 7), in which at least one of the arginine residues in the sequence is D-arginine.

In one embodiment, the antimicrobial peptide contains the sequence of rRrGRSPrRrTPSPrRrRSQSPrRrRSC (SEQ ID NO: 7), in which R is L-arginine and r is D-arginine. Alternatively, the peptide can contain the sequence of RrRGRSPRrRTPSPRrRrSQSPRrRrRSC (SEQ ID NO: 7), in which R is L-arginine and r is D-arginine.

The antimicrobial peptide can include the sequence of RRRGRPPRRPPrRRRRQPrRRRC (SEQ ID NO: 9), in which at least one of the arginine residues (e.g., 20%, 30%, 40%, or 50% of the arginine residues) in the sequence is D-arginine. For example, the sequence can be rRrGRPrRrPPrRrRQPrRrRC (SEQ ID NO: 9), in which R is L-arginine and r is D-arginine.

In one embodiment, the antimicrobial peptide further contains a non-HBcARD peptide (e.g., an affinity tag, a signal sequence, a ligand, or another antimicrobial peptide or fragment thereof). The non-HBcARD peptide can be a poly-histidine or an analog thereof. In one embodiment, the peptide has a sequence selected from the group consisting of:

- (i) RRRGRSPRRRTSPRRRRSQSPRRRRSHHHHHH (SEQ ID NO: 14),
- (ii) HHHHHHRRRGRSPRRRTSPRRRRSQSPRRRRS (SEQ ID NO: 15),
- (iii) RRRGRPPRRPPrRRRRQPrRRRH HH HH HH (SEQ ID NO: 16), and
- (iv) HHHHHHRRRGRPPRRPPrRRRRQPrRRR (SEQ ID NO: 17).

In another aspect, described herein is an antimicrobial peptide conjugate, which contains the antimicrobial peptide disclosed herein and a non-peptide moiety.

In yet another aspect, a pharmaceutical composition is described herein. The composition includes the antimicrobial peptide or the antimicrobial conjugate, and a pharmaceutically acceptable carrier.

Also contemplated herein is a method of treating an infection in a subject in need thereof. The method includes administering to the subject an antimicrobial peptide, an antimicrobial peptide conjugate, or a pharmaceutical composition described herein.

The details of one or more embodiments are set forth in the accompanying drawing and the description below. Other features, objects, and advantages of the embodiments will be apparent from the description and drawing, and from the claims.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 includes two sets of sequence alignments of HBcARD domains. (A) HBcARD sequences are highly conserved among human (SEQ ID NO: 6), wooly monkey (SEQ ID NO: 18), ground squirrel (SEQ ID NO: 19), woodchuck (SEQ ID NO: 20), and bat (SEQ ID NO: 21). (B) There are also four positive charge clusters separated in the HBc C-terminus of duck (SEQ ID NO: 22), heron (SEQ ID NO: 23), parrot (SEQ ID NO: 24), Ross's goose (SEQ ID NO: 25) and snow goose hepatitis B virus (SEQ ID NO: 26).

FIG. 2 is a set of graphs showing comparison of serum resistance between L- and D-HBcARD peptides. Peptides (L-HBcARD and D-HBcARD) were incubated with MBC buffer (10 mM sodium phosphate and 50 mM sodium chloride, pH 7.2) containing 5% fetal bovine serum (A), 5% mouse serum (B) or human serum (male and female) (C) at 37°C for 3 hours. The amounts of peptides were determined using SDS PAGE electrophoresis and green angel staining. (D) Peptide D-HBcARD exhibited 10,000-fold higher potency than that of L-HBcARD in MBC assay. Peptides (L-HBcARD and D-HBcARD) were incubated with *S. aureus* ATCC19636 with or without 5% mouse serum at 37°C for 3 hours. The antimicrobial activity was determined by colony formation assay. ***P<0.0001.

FIG. 3 is a graph showing comparison of hemolytic effect between L- and D-HBcARD peptides. Human red blood cells were incubated with different concentrations of the peptides. Hemolysis is presented as the percentage of Triton X-100-induced hemolysis. ***P<0.0001.

FIG. 4 is a set of graphs showing comparison of *in vivo* protection activities between L- and D-form HBcARD peptides in a mouse sepsis model infected with *S. aureus*. (A) Three week-old ICR mice were inoculated with *S. aureus* (4×10^6 CFU/mouse) and i.p. injected with PBS, L-HBcARD, or D-HBcARD at 2 hours post-inoculation. Each group contained ten mice. (B) and (C) Three week-old ICR mice were immunized with L- and D-HBcARD peptides (5 mg/kg), respectively, at day 0, 3 and 6. On day 14, the mice were inoculated with *S. aureus* (4×10^6 CFU/mouse) and i.p. injected with PBS, L-HBcARD or D-HBcARD (10 mg/kg) at 1 hour post-inoculation, respectively. Another group of mice were treated with PBS in parallel as a control. Each group of animals contained five mice. *P<0.05; ***P<0.0001; ns, no significance.

FIG. 5 is a graph showing *in vivo* protection efficacies of various modified HBcARD peptides at different doses in an ICR mouse sepsis model.

FIG. 6 is a graph showing *in vivo* protection efficacies of 150-177C and 150-177Q peptides in a BALB/c mouse lung infection model. Colistin-resistant *A. baumannii* was inoculated via intra-tracheal route.

FIG. 7 is a set of graphs showing lower *in vivo* toxicity of HBcARD peptide D-150-177C in comparison with polymyxin B. At day 0, male ICR mice (5 mice/group) were ip. injected with different doses of D-150-177C peptide (20-80 mg/kg) and polymyxin B (50 mg/kg), respectively. (A) The survival rates of all groups were monitored for 7 days. (B) Serum samples collected from mice treated with D-150-177C and polymyxin B were determined for alanine aminotransferase activity (ALT) at day 1. The dash line represents the mean of ALT value (45 U/L) of ICR mice (Charles River Laboratories).

FIG. 8 is a graph showing *in vivo* protection efficacies of all-D-arginine and partial-D-arginine substituted 150-177C peptides in a mouse sepsis model. Two hours after the inoculation, the mice (n=10/group) were treated with 5 mg/kg of various peptides D-, DL-, LD-150-177C, and PBS, respectively. Peptides D- and DL-150-177C protected all *S. aureus*-infected mice from death, while peptide LD-150-177C protected only 80% of mice.

FIG. 9 is a graph showing that deletion of serine and threonine residues from peptide DL-150-177C improved the *in vivo* protection efficacies at different doses in a mouse sepsis model.

DETAILED DESCRIPTION

It was unexpectedly discovered that certain modified peptides derived from an arginine-rich domain of HBV core protein exhibited broad spectrum antimicrobial activities.

Described herein is an antimicrobial peptide. It contains at least two (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) variable domains. The variable domains are in tandem and each separated from another by a variable linker. The antimicrobial peptide can have a maximum length of 100 amino acids (e.g., less than 10, 10, 14, 15, 20, 21, 22, 25, 28, 30, 35, 37, 40, 45, 47, 50, 55, 57, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acids). In one embodiment, the peptide has a C-terminal cysteine.

Each of the variable domains, individually, is a sequence of at least two (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) consecutive basic amino acids, e.g., arginine, histidine, and lysine. Each variable domain can contain a sequence of identical basic amino acids or different basic amino acids. For example, the antimicrobial peptides

can contain 2 to 20 identical or different variable domains, each being $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}X_{14}X_{15}X_{16}X_{17}X_{18}X_{19}X_{20}$ (SEQ ID NO: 27), wherein each of X_1 - X_{20} , individually, is an arginine, histidine, or lysine (natural or chemically modified) and any of X_3 - X_{20} can be present or absent. For example, each variable domain
 5 can have 2 to 10 basic amino acids.

In one embodiment, at least one variable domain in the peptide consists solely of arginine residues. In another embodiment, all of the variable domains in the peptide contain only arginine residues. Alternatively, the peptide can contain at least one variable domain that has one or more histidine or lysine residues.

10 Each variable linker has at least one amino acid (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20) and can contain any amino acid. It cannot have two or more consecutive basic amino acids.

The term “amino acid,” refers to any of the 20 standard amino acids (i.e., alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan,
 15 tyrosine, and valine). The term can also refer to a non-standard, non-proteinogenic, or chemically-modified amino acid, or an amino acid analog. An amino acid can be in the L- or D-form stereoisomer.

The term “basic amino acid,” refers to arginine, lysine, or histidine, the L- or D-form thereof, or an analog thereof.
 20

Non-standard amino acids include selenocysteine, pyrrolysine, N-formylmethionine, non-proteinogenic amino acids, amino acid analogs, and chemically-modified amino acids. A chemically-modified amino acid or amino acid analog typically has a different side chain from its naturally-occurring counterpart. Amino acid analogs and methods of incorporating
 25 them into a polypeptide are known in the art. See, e.g., Nguyen et al., *Biochemica et Biophysica Acta* 1808 (2011), 2297-2303; Knappe, *Antimicrobial Agents and Chemotherapy* 54(9) 2010, 4003-4005; US7879979; US5972940; US8835162; and US20080199964. Amino acid analogs are also commercially available. Amino acid analogs can be incorporated in the antimicrobial peptide to improve its stability, bioavailability,
 30 pharmacokinetics, tissue distribution, safety, tolerability, and/or efficacy.

Any of the antimicrobial peptides described herein can contain one or more residues that are not one of the twenty standard amino acids. In particular, one or more (e.g., 1, 2, 3,

4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 2%, 3%, 5%, 10%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%) of the basic amino acids in the variable domains or in the entire antimicrobial peptide can be a D-form of natural arginine, lysine, or histidine, or an analog of natural arginine, lysine, or histidine.

The antimicrobial peptide can be derived from the arginine-rich domain of a hepadnavirus core protein (HBcARD). HBcARD refers to a highly conserved arginine-rich C-terminal region of a core protein (HBc). See, Fig. 1. The HBc can be a mammalian HBc or an avian HBc. A mammalian HBc can be human HBc, woolly monkey HBc, ground squirrel HBc, woodchuck HBc, and bat HBc. An avian HBc can be a duck, heron, parrot, Ross's goose, or snow goose HBc. The HBc can be from a hepadnavirus of any genotype.

For example, the antimicrobial peptide can include a fragment of the HBcARD or a variant thereof (e.g., containing one or more amino acid substitutions, deletions, or insertions). The variable domains and the linkers can be derived from an HBcARD. For example, the sequence between any two arginine repeats in an HBcARD or a variant thereof can be used as a linker.

In one embodiment, the antimicrobial peptide contains a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)$ (SEQ ID NO: 1), wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , and X_8 , individually, is any amino acid or absent,

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)$ (SEQ ID NO:2), wherein each of X_1 , X_3 , and X_6 , individually, is a variable domain, and each of X_2 , X_4 , and X_5 , individually, is any amino acid or absent, and

(iii) $(X_1)X_2PX_3P(X_4)X_5QX_6P(X_7)$ (SEQ ID NO: 3), wherein each of X_1 , X_4 , and X_7 , individually, is a variable domain, and each of X_2 , X_3 , X_5 , and X_6 , individually, is any amino acid or absent. Each variable domain is a sequence of 2 to 20 consecutive basic amino acids.

The peptide can have a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}C$ (SEQ ID NO: 4), and

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}Q$ (SEQ ID NO: 5),

wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , X_8 , and X_{10} , individually, is any amino acid or absent.

The antimicrobial peptide can be a fusion or chimeric peptide that further contains a non-HBcARD peptide. A non-HBcARD peptide is not derived from any HBcARD and does not contain at least two variable domains connected by a linker as described above. A non-HBcARD peptide can be a peptide derived from another source (e.g., from a protein other than an HBcARD), an engineered peptide (e.g., another antimicrobial peptide), an affinity tag (e.g., a FLAG, poly-His, Myc, HA, CBP, HBH, or V5 tag), a signal sequence (e.g., a leader sequence or a localization signal), or a ligand (e.g., a receptor ligand).

For example, the antimicrobial peptide can have up to 100 amino acids and contain a sequence selected from the group consisting of:

(i) TVVRRRGRSPRRRTTPSPRRRRSQSPRRRRSQSRESQC (SEQ ID NO: 6), in which at least one of the arginine residues is D-arginine,

(ii) RRRGRSPRRRTTPSPRRRRSQSPRRRRSC (SEQ ID NO: 7),

(iii) RRRGRSPRRRTTPSPRRRRSQSPRRRRSQ (SEQ ID NO: 8),

(iv) RRRGRPRRRPPRRRRQPRRRRC (SEQ ID NO: 9),

(v) RRRGRSPRRRTTPSPRRRRRC (SEQ ID NO: 10),

(vi) RRRGRPRRRPPRRRRRC (SEQ ID NO: 11),

(vii) RRRTTPSPRRRRSQSPRRRRRC (SEQ ID NO: 12),

(viii) RRRPPRRRRQPRRRRC (SEQ ID NO: 13).

(ix) RRRGRSPRRRTTPSPRRRRSQSPRRRRSHHHHHH (SEQ ID NO: 14),

(x) HHHHHHRRRGRSPRRRTTPSPRRRRSQSPRRRRS (SEQ ID NO: 15),

(xi) RRRGRPRRRPPRRRRQPRRRHHHHHH (SEQ ID NO: 16), and

(xii) HHHHHHRRRGRPRRRPPRRRRQPRRRR (SEQ ID NO: 17).

As described herein, the antimicrobial peptide can have one or more modified amino acids. For example, in any of the above-described sequences or consensus sequences, one or more of the basic amino acids in the variable domains or in the entire antimicrobial peptide can be a D-form of natural arginine, lysine, or histidine, or an analog of natural arginine, lysine, or histidine.

Any of the above-described sequences or consensus sequences can be at the N- or C-terminus of the antimicrobial peptide.

The antimicrobial peptide described herein can also be conjugated to a non-peptide moiety at the N- or C-terminus to form a peptide conjugate. A non-peptide moiety can be a polymer (e.g., a polyethylene glycol polymer), oligosaccharide, lipid, glycolipid, solid

support (e.g., a bead or nanoparticle), small molecule drug, biotin, nucleic acid molecule, antibody, vitamin, carrier protein (e.g., KLH, BSA, or OVA), or detectable label (e.g., fluorescent, radioactive, or enzymatic label). The peptide conjugate also exhibits an antimicrobial activity. Methods of generating peptide conjugates are known in the art.

5 The antimicrobial peptide or peptide conjugate described herein can be mixed with a pharmaceutically acceptable carrier to form a pharmaceutical composition.

10 The composition can be formulated with a pharmaceutically acceptable carrier such as a phosphate buffered saline, a bicarbonate solution, and/or an adjuvant. Suitable pharmaceutical carriers and diluents, as well as pharmaceutical necessities for their use, are known in the art. This composition may be prepared as an injectable, liquid solution, emulsion, or another suitable formulation.

15 An effective amount of the composition described above may be administered by intranasal inhalation, topical application, or parenteral routes, e.g., intravenous injection, subcutaneous injection or intramuscular injection. Alternatively, other modes of administration including suppositories and oral formulations may be desirable. For suppositories, binders and carriers may include, for example, polyalkylene glycols or triglycerides. Oral formulations may include normally employed excipients such as pharmaceutical grades of saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders.

20 The above-described composition can be administered to a subject (e.g., a human, another mammal, or a laboratory animal) to treat a microbial infection or to inhibit growth of a microbe in the subject. The composition can be used to treat an infection caused by a Gram-positive or Gram-negative bacteria, fungus, parasite, or virus, e.g., *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Shigella dysenteriae*, *Escherichia coli*, *Staphylococcus aureus*, *Acinetobacter baumannii*, *Clostridium Difficile*, *Candida*, *Aspergillus*, *Blastomyces*, *Cryptococcus neoformans*, *Cryptococcus gattii*, *Coccidioides*, *Histoplasma*, *Pneumocystis jirovecii*, ringworm, *Sporothrix*, *Exserohilum*, or *Cladosporium*.

30 The specific disclosure below is to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. Without further elaboration, it is believed that one skilled in the art can, based on the description herein,

utilize the present disclosure to its fullest extent. All publications cited herein are hereby incorporated by reference in their entirety.

Modified antimicrobial peptides derived from human HBcARD

A novel antimicrobial peptide from human hepatitis B virus (HBV) core protein (HBc) arginine-rich domain (ARD) was previously identified. See Chen et al. (2013), PLoS Pathog 9:e1003425 doi:10.1371/journal.ppat.1003425, and Chen et al., (2016), Appl Microbiol Biotechnol. 100(21): 9125-9132. This HBcARD peptide showed a broad spectrum antimicrobial activity against Gram-negative and Gram-positive bacteria. In our mouse sepsis model, injection of HBcARD peptide at 2 hour postinoculation can protect 40% of *S. aureus*-infected mice from death. See Chen et al. (2013).

The HBcARD peptides are highly conserved in the hepadnaviruses of different species. To improve the antimicrobial efficacy of HBcARD, we compared the sequences and antimicrobial activities of various HBcARD peptides of mammalian, rodent, and avian hepadnaviruses. HBcARD peptide of human hepadnaviruses displayed the strongest antimicrobial activity.

To further improve the potency of our lead compound HBcARD, we tested the antimicrobial activity of each of several HBcARD derivatives, including peptides modified by truncation and D-arginine substitution for L-arginine. See Table 1.

Table 1. Sequences of various HBcARD peptides tested for their bactericidal activity

HBcARD peptide	amino acid sequence
L-147-183	TVVRRRGRSPRRRTTPSPRRRRSQSPRRRRSQSRESQC (SEQ ID NO: 6)
D-147-183	TVVrrrGrSPrrrTPSPrrrrSQSPrrrrSQSrESQC (SEQ ID NO: 6)
150-177C	RRRGRSPRRRTTPSPRRRRSQSPRRRRSC (SEQ ID NO: 7)
150-177Q	RRRGRSPRRRTTPSPRRRRSQSPRRRRSQ (SEQ ID NO: 8)
150-171C	RRRGRSPRRRTTPSPRRRRSC (SEQ ID NO: 10)
157-177C	RRRTTPSPRRRRSQSPRRRRSC (SEQ ID NO: 12)
164-177C	RRRRSQSPRRRRSC (SEQ ID NO: 28)
D-150-177C	rrrGrSPrrrTPSPrrrrSQSPrrrrSC (SEQ ID NO: 7)

DL-150-177C	rRrGRSPRrTPSPRrRSQSPRrRSC (SEQ ID NO: 7)
LD-150-177C	RrRGRSPRrRTPSPRrRrSQSPRrRrSC (SEQ ID NO: 7)
DL-dST-150-177C	rRrGRPrRrPPrRrRQPrRrRC (SEQ ID NO: 9)

R: L-arginine; r: D-arginine

Comparison of sequences and antimicrobial activities of human and non-human HBcARD peptides

By multiple sequence alignment, we compared the sequences of HBcARD peptides of various hepadnaviruses, including human (AAP31571.1), wooly monkey (AAO74859.1), ground squirrel (AAB08031.1), woodchuck (AAA46761.1), bat (AGT17576.1), duck (AAO49490.1), heron (AAA45737.1), parrot (AFY97786.1), Ross's goose (AAR89928.1) and snow goose (AAD22001.1). See Fig. 1. Similar to the HBcARD peptide of human hepatitis B virus, HBcARD peptides of other mammalian hepadnaviruses (wooly monkey, ground squirrel, woodchuck and bat) contained four clustering arginine-rich domains. While the sequence homology of HBcARD peptides between avian and mammalian hepadnaviruses was low, they both contained four highly positive-charged domains. To compare their respective antimicrobial activities, we determined the minimal bactericidal concentrations of four HBcARD peptides derived from bat, woodchuck, duck and heron hepadnaviruses. The results indicated that peptides derived from bat and woodchuck hepadnaviruses exhibited potent antimicrobial activity comparable to that from human hepatitis B virus. See Table 2. In contrast, peptides derived from avian hepadnaviruses displayed lower antimicrobial activity. Because HBcARD peptides from human HBV is shorter in length (147-183; 37 amino acids), we focused on this peptide in our subsequent modification and optimization experiments.

Table 2. Antimicrobial activity of human, bat, woodchuck, duck and heron HBcARD peptides against Gram-positive and Gram-negative bacteria

Bacterial strain ^a	Minimal bactericidal concentration (mg/L)				
	Human HBcARD	Bat HBcARD	Woodchuck HBcARD	Duck HBcARD	Heron HBcARD
<i>P. aeruginosa</i>	9.2	10	9.8	>70.6	>69.3

ATCC 9027					
<i>K. pneumoniae</i> ATCC 13884	9.2	5	9.8	>70.6	>69.3
<i>E. coli</i> ATCC 25922	18.4	20	19.6	ND	ND
<i>A. baumannii</i> ATCC 17978	2.3	2.5	4.9	70.6	69.3
<i>S. aureus</i> ATCC 19636	18.4	20	19.6	ND	ND

^a ATCC: American Type Culture Collection; ND: not detectable

Comparison of antimicrobial activity of L- and D-HBcARD peptides

We used PeptideCutter to predict the potential protease cleavage sites on the HBcARD 147-183 peptide (37-mer). The results showed that more than 70% of the protease cleavage sites were mapped to the arginine residues of HBcARD peptide (data not shown). To improve the peptide stability in the serum, we synthesized a modified peptide, D-HBcARD 147-183 peptide, by D-amino acid replacement, in which all arginine residues were replaced with its D-isomer.

To investigate whether D-arginine substitution will improve the antimicrobial activity, L- and D-HBcARD peptides (37-mer) were tested side-by-side against a wide variety of bacteria, including *P. aeruginosa*, *K. pneumoniae*, *A. baumannii*, *E. coli*, *S. dysenteriae* and *S. aureus*. Relative to the L-HBcARD, the D-HBcARD peptide displayed similar antimicrobial activity against *A. baumannii* (MBC=2.3-4.6 mg/L), and better antimicrobial activity against *P. aeruginosa* (MBC=4.6 mg/L), *K. pneumoniae* (MBC=4.6 mg/L), and *S. aureus* (MBC=4.6-9.2 mg/L). However, for reasons that remain unclear, the antimicrobial activity of D-HBcARD peptide against *E. coli* and *S. dysenteriae* was decreased from 18.4 to 73.6 mg/L. See Table 3.

Table 3. Comparison of minimal bactericidal concentration (MBC) between wild type HBcARD 147-183 peptide L-HBcARD and D-HBcARD

Bacterial strain ^a	Minimal bactericidal concentration (mg/L)
-------------------------------	---

	L-HBcARD (37-mer)	D-HBcARD (37-mer)
Gram-negative		
<i>P. aeruginosa</i> ATCC 9027	9.2	4.6
<i>P. aeruginosa</i> ATCC 27853	9.2-18.4	4.6
<i>K. pneumoniae</i> ATCC 13884	9.2	4.6
<i>E. coli</i> ATCC 25922	18.4	73.6
<i>A. baumannii</i> ATCC 17978	2.3	2.3
<i>A. baumannii</i> ATCC 17978CR ^b	2.3	2.3
<i>A. baumannii</i> ATCC 19606	2.3	2.3
<i>A. baumannii</i> ATCC 19606CR ^b	2.3	2.3
<i>A. baumannii</i> TCGH 45530 ^b	2.3	2.3
<i>A. baumannii</i> TCGH 46709 ^b	4.6	4.6
<i>S. dysenteriae</i> Xen27 ^c	18.4	73.6
Gram-positive		
<i>S. aureus</i> ATCC 19636	18.4	9.2
<i>S. aureus</i> ATCC 25923	18.4	4.6
<i>S. aureus</i> ATCC 29213	18.4	~4.6

^a ATCC: American Type Culture Collection

^b Colistin-resistant *A. baumannii* (Laboratory induced and Clinical isolates from Tzu Chi General Hospital (Chen et al. PLoS Pathog. 2013;9(6):e1003425.)

^c Caliper Life Sciences, Inc.

5 Comparison of serum resistance between L-HBcARD and D-HBcARD

To compare the stability of the 37-mer L- and D-HBcARD peptides against serum protease degradation, we performed a protease resistance assay. After incubation with 5% bovine serum at 37°C for 3 hours, more than 80% of L-HBcARD peptide was no longer detectable by staining with Green Angel on SDS-PAGE. In contrast, the signal intensity of the D-HBcARD peptide was not reduced under the same condition. See Fig. 2, A. Similar results were obtained, when L-HBcARD was incubated with 5% mouse or human (male and female) sera. See Fig. 2, B and C. These results suggested that D-HBcARD was more resistant than L-HBcARD to protease digestion of fetal bovine, mouse and human serums.

Furthermore, we determined the *in vitro* antimicrobial activity of L- and D-HBcARD peptides in the presence of serum. The results showed that both L-HBcARD and D-HBcARD peptides at 18.4 mg/L concentration were able to kill *S. aureus* in the MBC assay. See Fig. 2, D. Upon the addition of 5% mouse serum, there was a significant difference in the antimicrobial activities between L- and D-HBcARD peptides ($P < 0.0001$). While the treatment with L-HBcARD peptide in the presence of 5% mouse serum produced 10^7 bacterial colonies, treatment with D-HBcARD peptide produced around 10^3 colonies. Therefore, our D-HBcARD peptide exhibited higher antimicrobial activity than that of the L-HBcARD peptide in the presence of 5% mouse serum. This observation suggested that D-HBcARD was more stable and resistant to proteolytic degradation.

Hemolytic activity of D-HBcARD peptide

In the hemolysis assay, human RBCs were incubated with serially-diluted doses (0 to 460 mg/L) of D-HBcARD, L-HBcARD, or melittin, at 37°C for one hour. As shown in Fig. 3, melittin caused 100% hemolysis at the concentration of 8.9 mg/L. In contrast, no hemolytic activity of both D- and L-HBcARD peptides was detected up to 460 mg/L ($P < 0.0001$).

Comparison of *in vivo* protection efficacies between L- and D-form HBcARD peptides in a mouse sepsis model infected with *S. aureus*

To investigate the *in vivo* protection efficacy of the 37-mer L- and D-HBcARD peptides, we performed a mouse sepsis model infected with *S. aureus*. ICR mice were first i.p. inoculated with *S. aureus* (4×10^6 CFU/mouse), followed by one single i.p. injection of L-HBcARD, D-HBcARD and PBS at two hours post-inoculation, respectively. All mice ($n=10$) administered with PBS died at day 1. See Fig. 4, A. At 7 days post-inoculation, 40% of mice treated with L-HBcARD (10 mg/kg) survived. See Fig. 4, A. In contrast, D-HBcARD peptide was able to achieve a much higher survival rate (60% for 5 mg/kg dose, and 100% for 10 mg/kg dose). See Fig. 4, A. These results indicated that the *in vivo* efficacy of HBcARD can be improved by the D-arginine replacement strategy ($P < 0.0001$).

Comparison of the immunogenicity between L- and D-form HBcARD peptides in a mouse sepsis model infected with *S. aureus*

To investigate whether repeated treatments with the 37-mer HBcARD peptides may

compromise the efficacy by inducing neutralization antibody *in vivo*, we immunized 3-week old mice three times with L- and D-HBcARD peptides, respectively, prior to bacterial infection. Two weeks after the first immunization, immunized mice were then inoculated with *S. aureus*, followed by i.p. injection with either L- or D-HBcARD peptides 1 hour post-inoculation. Consistent with our previous report (Chen et al., 2013), L-HBcARD still protected all mice from death, irrespective of the prior immunizations with or without L-HBcARD peptide. See Fig. 4, B. Control mice immunized with PBS and treated with PBS showed a mortality near 80% within 24 hours after bacterial challenge ($P < 0.05$). Seven days post-inoculation, all mice treated with L- and D-HBcARD (10 mg/kg) survived bacterial challenge, irrespective of the prior immunization with or without D-HBcARD. See Fig. 4, C. Therefore, prior immunizations with either D-form or L-form peptides induced no neutralization activity against the *in vivo* antimicrobial activity of subsequent treatments with HBcARD peptides.

Significance of terminal cysteine

We tested the antimicrobial activity of additional HBcARD derivatives. See Table 1. The antimicrobial activities of these peptides were determined by minimal bactericidal concentration (MBC). Unlike the parental peptide HBcARD 147-183 (37-mer) (see Chen et al., 2013), the derivative peptide HBcARD 150-177Q (28-mer) showed no detectable bactericidal activity against *S. aureus*. See Table 4. To mimic the parental peptide HBcARD 147-183 with a cysteine at the carboxyl terminus, we designed another 28-mer derivative HBcARD 150-177C by replacing the terminal Q (glutamine) residue of HBcARD 150-177Q with a C (cysteine) residue. See Table 1. Interestingly, this Q-to-C substitution effectively rescued the bactericidal activity against *S. aureus*. See Table 4.

Table 4. Minimal bactericidal concentrations (MBC) of modified HBcARD peptides against various Gram-negative and Gram-positive bacteria

Bacteria strains	MBC (uM)				
	150-177C	150-177Q	150-171C	157-177C	164-177C
Gram-negative					
<i>P. aeruginosa</i> ATCC9027	2	—	—	—	—

<i>P. aeruginosa</i>					
ATCC27853	2	—	—	—	—
<i>E. coli</i> ATCC25922	2	—	—	—	—
<i>A. baumannii</i> ATCC17978	0.5	—	—	—	—
<i>A. baumannii</i> ATCC19606	0.25	—	—	—	—
<i>A. baumannii</i> ATCC45530	0.5	—	—	—	—
<i>A. baumannii</i> ATCC46709	1	—	—	—	—
Gram-positive					
<i>S. aureus</i> ATCC19636	2	ND	ND	ND	ND

—, not determined; ND, no detectable antimicrobial activity

Significance of length or arginine content

Given the fact that the terminal cysteine appears to be important for the bactericidal activity, we asked if the total length of the HBcARD 150-177C peptide (28-mer) can be further reduced. We compared the potencies against *S. aureus* among peptides 150-171C (20-mer), 157-177C (21-mer), and 164-177C (14-mer). As shown in Table 4, none of these peptides showed detectable activity. The results here indicated that shortened peptides with a reduced number of arginines exhibited no antimicrobial activity against *S. aureus*. See Table 4. On the other hand, we have shown that shorter peptides containing 3 variable domains are sufficient to kill other bacteria (data not shown).

The 28-mer peptide D-150-177C improved protection efficacy in a sepsis mouse model

We compared the *in vivo* protection efficacy of HBcARD 150-177C peptides (28-mer) containing either D-form or L-form arginines. See Table 1 and Fig. 5. ICR mice (n=10/group) were ip. inoculated with *S. aureus* ATCC19636 (4×10^6 /mouse), followed by treatment with modified peptides at two hours post-inoculation. All PBS control mice died on day 1, and only 20% of the mice treated with control peptide (150-177Q) survived. Administration of L-150-177C peptide at the dose of 5 and 10 mg/kg protected 30% and 70% mice from death, respectively. See Fig. 5. When treated with 5 mg/kg of D-150-177C, all mice (100%) survived.

The 28-mer peptide D-150-177C protected mice from lung infection with colistin-resistant *A. baumannii*

BALB/c mice (n=8/group) were intra-tracheally inoculated with colistin -resistant *A. baumannii* TCGH 46709 (3.4×10^8 cfu/mouse). These lung-infected mice were ip. treated with colistin (5 mg/kg/day) or D-150-177C (5 and 10 mg/kg/day), respectively. All mice treated with colistin died at 60 hours post-inoculation with drug-resistant *A. baumannii*. See Fig. 6. In contrast, there was a dose-dependent protection effect from D-150-177C. A significant difference was observed when the mice were treated with 10 mg/kg/day of D-150-177C peptide ($p < 0.05$). See Fig. 6.

Comparison of *in vivo* toxicity between peptide D-150-177C and polymyxin B

Using the sepsis mouse model, we examined the survival rates of ICR mice intra-peritoneally (i.p.) injected with the 28-mer D-150-177C peptide (20 – 80 mg/kg body weight) and polymyxin B (a colistin-related compound) (50 mg/kg body weight). See Fig. 7, A. All mice treated with polymyxin B (50 mg/kg) died at day one, as expected from the notorious nephrotoxicity of colistin and polymyxin B to kidney. See Fig. 7, A. In contrast, no acute toxicity was observed in the mice treated with 20 and 40 mg/kg D-150-177C. See Fig. 7, A. When the doses were increased to 60 and 80 mg/kg, survival rates of mice were decreased to 80% and 40%, respectively. See, Fig. 7, A. Liver injury can be detected by the serum ALT level. Mice treated with D-150-177C in the dose range of 20 to 40 mg/kg showed a higher ALT level than polymyxin B, but the p value is insignificant. See Fig. 7, B. The ALT levels of mice treated with 60 mg/kg were significantly higher than those treated at 20 mg/kg polymyxin B ($p < 0.01$). See Fig. 7, B.

Comparison of *in vivo* protection efficacies among various modified 150-177C peptides

The 28-mer peptide D-150-177C contains a total of 14 L-arginines substituted with 14 D-arginines. See Table 1 and Fig. 6. D-arginine is far more expensive than L-arginine. To reduce the cost of peptide synthesis, we compared the *in vivo* protection efficacies between peptides containing complete or partial D-arginine substitution. See Table 1 and Fig. 8. Peptide DL-150-177C is only partially D-arginine substituted, and exhibited very similar protection efficacy to all-D-arginine substituted peptide D-150-177C (100% substitution). See Fig. 8. In contrast, the protection efficacy of peptide LD-150-177C (also partially

substituted) appeared to be less than DL-150-177C. See Fig. 8.

Serine and threonine are dispensable

We engineered a new 22-mer peptide, DL-dST-150-177C (see Table 1), with two major modifications: (1) partial-D-arginine substitution by using the backbone of peptide DL-150-177C; and (2) deletion of 5 serine and one threonine residues from the parental 28-mer peptide DL-150-177C. This 22-mer peptide DL-dST-150-177C showed remarkably improved protection efficacy compared to the 28-mers DL-150-177C and L-150-177C. See Fig. 9.

Bacterial isolates

The antimicrobial activities of HBcARD peptides were tested using a number of bacterial strains, including *Pseudomonas aeruginosa* Migula strains (ATCC 27853, ampicillin-resistant and ATCC 9027, ampicillin-resistant), *Klebsiella pneumoniae* strain (ATCC 13884), *Shigella dysenteriae* Xen27 (Caliper Co.), *Escherichia coli* strain (ATCC 25922), *Staphylococcus aureus* subsp. strains (ATCC 25923, methicillin-resistant ATCC 29213, methicillin-resistant and ATCC 19636, methicillin-resistant) and *Acinetobacter baumannii* strains (ATCC 17978, ATCC 17978 CR, ATCC 19606, ATCC 19606 CR, TCGH 45530 and TCGH 46709). Clinical isolates TCGH 45530 and TCGH 46709 were obtained from Tzu-Chi Buddhist General Hospital (TCGH) in Taiwan, and were identified using the Vitek system (Biomérieux Vitek, Inc., Hazelwood, MO, USA). See Chang et al. (2012), J Microbiol Immunol Infect 45:37-42 doi:10.1016/j.jmii.2011.09.019.

Antimicrobial assay

L- and D-HBcARD peptides were purchased from Yao-Hong Biotechnology Inc. (Taipei, Taiwan). Antimicrobial activity was determined as described. See, Chen et al., 2013. Briefly, bacteria were grown in MH broth (Difco) to mid-logarithmic phase at 37°C, and were diluted to 10⁶ CFU (colony formation unit)/ml in phosphate buffer (10 mM sodium phosphate and 50 mM sodium chloride, pH 7.2). Peptides were serially diluted in the same buffer. Fifty microliters (μl) of bacteria were mixed with fifty μl of peptides at varying concentrations, followed by incubation at 37°C for 3 hours without shaking. At the end of incubation, bacteria were placed on Mueller-Hinton broth agar plates, and allowed to grow at

37°C overnight for measurement of minimal bactericidal concentration (MBC). The lowest peptide concentration that displayed no bacterial growth (zero colony) was defined as MBC. All peptides were tested in triplicate.

Stability to proteases

L- and D-HBcARD peptides (0.5 nmol) were mixed with MBC buffer in the presence or absence of 5% serum collected from bovine, mouse and human origins. After incubation at 37°C for 3 hours, the amounts of peptides surviving the protease digestion were determined by SDS-PAGE electrophoresis and staining with Green Angel. The images were quantified using image J software and the intensities were normalized with the no serum control. To investigate the effect of serum on the antimicrobial activity of HBcARD peptides, L- and D-HBcARD peptides at their respective MBC concentrations were incubated with *S. aureus* ATCC19636 strain (10^6 CFU/ml) at 37°C for three hours. Bacteria were plated on MH agar and the antimicrobial activity was determined by colony formation.

Hemolytic activity

The hemolytic activities of peptides were determined by hemolysis against human red blood cells (hRBCs). Human blood was obtained in EDTA-containing tube and was centrifuged at 450 g for 10 min. The pellet was washed three times with PBS buffer, and a solution of 10% hRBCs was prepared. hRBCs solution was mixed with serial dilutions of peptides in PBS buffer, and the reaction mixtures were incubated for 1 h at 37°C. After centrifugation at 450 g for 10 min, the percentage of hemolysis was determined by measuring the absorbance at the wavelength of 405 nm of the supernatant. Blank and 100% hemolysis were determined in PBS buffer and in the presence of 1% Triton X-100, respectively.

***In vivo* animal studies**

Three-week old male ICR mice (19 to 21 g) were purchased from BioLASCO (Taiwan). To test the *in vivo* protection efficacy of the HBcARD peptides, all mice were inoculated intraperitoneally with *S. aureus* ATCC 19636 (4×10^6 CFU/mouse). HBc147-183 (L- or D-HBcARD) or the PBS control was administered intraperitoneally at 2 hours after bacterial inoculation, respectively. Each group contained 10 mice. Mortality was monitored daily for 7 days following the bacterial inoculation.

To investigate the potential immunogenicity of HBcARD peptides, we determined the *in vivo* antimicrobial activity of HBcARD peptide in peptide-immunized mice. Briefly, three week-old male mice were immunized three times with 0.2 ml of L- and D-HBcARD peptides (5 mg/kg) at day 0, 3, 6, respectively. Immunized mice were inoculated with *S. aureus* ATCC 19636 (4×10^6 CFU/mouse) at day 14, and were administered with 0.2 ml of PBS, or L- and D-HBcARD peptide (10 mg/kg) one-hour postinoculation. Another group of mice received the identical protocol with PBS as a control. Each group contained 5 mice. Mortality was monitored daily for 7 days following the bacterial inoculation.

Statistical analysis

Statistical analysis was performed using Graphpad software. The results were shown in mean $\pm$ SD, and the difference between individual groups was analyzed by student t test. Survival curves were plotted by Kaplan-Meier method and analyzed by log-rank test.

OTHER EMBODIMENTS

All of the features disclosed in this specification may be combined in any combination. Each feature disclosed in this specification may be replaced by an alternative feature serving the same, equivalent, or similar purpose. Thus, unless explicitly stated otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

From the above description, one skilled in the art can easily ascertain the essential characteristics of the described embodiments, and without departing from the spirit and scope thereof, can make various changes, derivatives, and modifications of the embodiments to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

CLAIMS

1. An antimicrobial peptide, the peptide comprising 2 to 20 variable domains, each variable domain is a sequence of 2 to 20 consecutive basic amino acids, wherein (a) the
5 variable domains are separated from each other by a variable linker, (b) the variable linker can have 1 to 20 any amino acids other than two or more consecutive basic amino acids, and (c) the peptide has no more than 100 amino acids.

2. The antimicrobial peptide of claim 1, wherein the peptide has at least 3
10 variable domains.

3. The antimicrobial peptide of claim 1, wherein the peptide has a C-terminal cysteine.

4. The antimicrobial peptide of claim 1, wherein at least one of the basic amino
15 acids in the variable domains is an arginine.

5. The antimicrobial peptide of claim 4, wherein all of the basic amino acids in each variable domain are arginine residues.
20

6. The antimicrobial peptide of claim 1, wherein at least one of the basic amino acids is a chemically-modified amino acid.

7. The antimicrobial peptide of claim 6, wherein the chemically-modified amino
25 acid is a D-amino acid.

8. The antimicrobial peptide of claim 7, wherein the D-amino acid is D-arginine.

9. The antimicrobial peptide of any of claims 1-8, wherein the variable domains
30 and the variable linkers are derived from the arginine-rich domain of a hepadnavirus core protein (HBcARD).

10. The antimicrobial peptide of claim 9, wherein the HBcARD contains a sequence from residue 147 to the C-terminal residue of a hepadnavirus core protein.

11. The antimicrobial peptide of claim 10, wherein each variable domain has three or four arginine residues and each variable linker has 2 to 4 amino acids.

12. The peptide of claim 11, wherein the peptide contains a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)$, wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , and X_8 , individually, is any amino acid or absent,

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)$, wherein each of X_1 , X_3 , and X_6 , individually, is a variable domain, and each of X_2 , X_4 , and X_5 , individually, is any amino acid or absent, and

(iii) $(X_1)X_2PX_3P(X_4)X_5QX_6P(X_7)$, wherein each of X_1 , X_4 , and X_7 , individually, is a variable domain, and each of X_2 , X_3 , X_5 , and X_6 , individually, is any amino acid or absent,

13. The antimicrobial peptide of claim 12, wherein the peptide has a C-terminal cysteine.

14. The antimicrobial peptide of claim 12, wherein the peptide has a consensus sequence selected from the group consisting of:

(i) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}C$, and

(ii) $(X_1)GRX_2P(X_3)X_4PX_5P(X_6)X_7QX_8P(X_9)X_{10}Q$,

wherein each of X_1 , X_3 , X_6 , and X_9 , individually, is a variable domain, and each of X_2 , X_4 , X_5 , X_7 , X_8 , and X_{10} , individually, is any amino acid or absent.

15. The antimicrobial peptide of claim 12, wherein the peptide contains a sequence selected from the group consisting of:

(i) TVVRRRGRSPRRRTSPRRRRRSQSPRRRRRSQSRESQC, in which at least one of the arginine residues is D-arginine,

(ii) RRRGRSPRRRTSPRRRRRSQSPRRRRSC,

(iii) RRRGRSPRRRTSPRRRRRSQSPRRRRSQ,

- (iv) RRRGRPRRRPPRRRRQPRRRRC,
- (v) RRRGRSPRRRTSPRRRRRC,
- (vi) RRRGRPRRRPPRRRRRC,
- (vii) RRRTSPRRRRSQSPRRRRRC, or
- (viii) RRRPPRRRRQPRRRRC.

16. The antimicrobial peptide of claim 15, wherein the peptide contains the sequence of RRRGRSPRRRTSPRRRRSQSPRRRRSC, in which each of the arginine residues in the sequence is L-arginine.

17. The antimicrobial peptide of claim 15, wherein the peptide contains the sequence of RRRGRSPRRRTSPRRRRSQSPRRRRSC, in which at least one of the arginine residues in the sequence is D-arginine.

18. The antimicrobial peptide of claim 17, wherein each of the arginine residues in the sequence is D-arginine.

19. The antimicrobial peptide of claim 17, wherein the sequence is rRrGRSPrRrTPSPrRrRSQSPrRrRSC, in which R is L-arginine and r is D-arginine.

20. The antimicrobial peptide of claim 17, wherein the sequence is RrRGRSPRrRTPSPRrRrSQSPRrRrSC, in which R is L-arginine and r is D-arginine.

21. The antimicrobial peptide of claim 15, wherein the peptide contains the sequence of RRRGRPRRRPPRRRRQPRRRRC, in which at least one of the arginine residues in the sequence is D-arginine.

22. The antimicrobial peptide of claim 15, wherein the sequence is rRrGRPrRrPPrRrRQPrRrRC, in which R is L-arginine and r is D-arginine.

23. The antimicrobial peptide of any of claims 15-22, wherein the sequence is at the C-terminus of the peptide.

24. The antimicrobial peptide of any of claims 1-23, further comprising a non-HBcARD peptide.

25. The antimicrobial peptide of claim 24, wherein the non-HBcARD peptide is an affinity tag, a signal sequence, a ligand, or another antimicrobial peptide or fragment thereof.

26. The antimicrobial peptide of claim 25, wherein the non-HBcARD peptide is a poly-histidine or an analog thereof.

27. The antimicrobial peptide of claim 26, wherein the peptide has a sequence selected from the group consisting of:

(i) RRRGRSPRRRTSPRRRRSQSPRRRRSHHHHHH,

(ii) HHHHHHRRRGRSPRRRTSPRRRRSQSPRRRRS,

(iii) RRRGRPRRRPPRRRRQPRRRHHHHHH, and

(iv) HHHHHHRRRGRPRRRPPRRRRQPRRRR

28. The antimicrobial peptide of any of claims 1-4, wherein at least one variable domain has a lysine.

29. The antimicrobial peptide of any of claims 1-4, wherein at least one variable domain has a histidine.

30. The peptide of any of claims 1-29, wherein the peptide has a cyclic structure.

31. The antimicrobial peptide of any of claims 1-30, wherein the peptide exhibits a broad spectrum antimicrobial activity against a gram-positive bacteria, gram-negative bacteria, fungus, parasite, or virus.

32. An antimicrobial peptide conjugate, the conjugate comprising the peptide of any of claims 1-31 and a non-peptide moiety.

33. A pharmaceutical composition comprising the antimicrobial peptide of any of claims 1-31 and a pharmaceutically acceptable carrier.

34. A pharmaceutical composition comprising the antimicrobial peptide conjugate
5 of claim 32 and a pharmaceutically acceptable carrier.

35. A method of treating a microbial infection, the method comprising administering the composition of claim 33 or 34 to a subject in need thereof.

**TITLE OF INVENTION: MODIFIED ANTIMICROBIAL PEPTIDE
DERIVED FROM AN ARGININE-RICH DOMAIN**

ABSTRACT

5 An antimicrobial peptide, the peptide comprising 2 to 20 variable domains, each variable domain is a sequence of 2 to 20 consecutive basic amino acids, wherein (a) the variable domains are separated from each other by a variable linker, (b) the variable linker can have 1 to 20 any amino acids other than two or more consecutive basic amino acids, and (c) the peptide has no more than 100 amino acids.

10

FIG. 1

SEQUENCE LISTING

<110> Academia Sinica
 <120> Modified Antimicrobial Peptide Derived From An Arginine-Rich Domain
 <130> 218048-0276PCT
 <150> US 62/342,415
 <151> 2016-05-27
 <160> 28
 <170> PatentIn version 3.5
 <210> 1
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Synthetic polypeptide
 <220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine, and histidine
 <220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> Xaa can be any amino acid or absent
 <220>
 <221> MISC_FEATURE
 <222> (6)..(6)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine, and histidine
 <220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa can be any amino acid or absent
 <220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> Xaa can be any amino acid or absent
 <220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> Xaa can be 2-20 amino acids each selected from arginine, lysine, and histidine
 <220>
 <221> MISC_FEATURE
 <222> (12)..(12)
 <223> Xaa can be any amino acid or absent
 <220>
 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (16)..(16)
 <223> Xaa can be 2-20 amino acids each selected from arginine, lysine,
 and histidine

<400> 1

Xaa Gly Arg Xaa Pro Xaa Xaa Pro Xaa Pro Xaa Xaa Gln Xaa Pro Xaa
 1 5 10 15

<210> 2
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (6)..(6)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<400> 2

Xaa Gly Arg Xaa Pro Xaa Xaa Pro Xaa Pro Xaa
 1 5 10

<210> 3
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (2)..(2)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (6)..(6)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<400> 3

Xaa	Xaa	Pro	Xaa	Pro	Xaa	Xaa	Gln	Xaa	Pro	Xaa
1				5					10	

<210> 4
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (6)..(6)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine, and histidine

<220>
 <221> MISC_FEATURE
 <222> (12)..(12)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (16)..(16)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine, and histidine

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> Xaa can be any amino acid or absent

<400> 4

Xaa Gly Arg Xaa Pro Xaa Xaa Pro Xaa Pro Xaa Xaa Gln Xaa Pro Xaa
 1 5 10 15

Xaa Cys

<210> 5
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<220>
 <221> MISC_FEATURE
 <222> (1)..(1)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine, and histidine

<220>
 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> Xaa can be any amino acid or absent

<220>

<221> MISC_FEATURE
 <222> (6)..(6)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (12)..(12)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (14)..(14)
 <223> Xaa can be any amino acid or absent

<220>
 <221> MISC_FEATURE
 <222> (16)..(16)
 <223> Xaa can be 2 to 20 amino acids selected from arginine, lysine,
 and histidine

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> Xaa can be any amino acid or absent

<400> 5

Xaa Gly Arg Xaa Pro Xaa Xaa Pro Xaa Pro Xaa Xaa Gln Xaa Pro Xaa
 1 5 10 15

Xaa Gln

<210> 6
 <211> 37
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<400> 6

Thr Val Val Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser
 1 5 10 15

Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser
 20 25 30

Arg Glu Ser Gln Cys
35

<210> 7
<211> 28
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 7

Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro Arg Arg
1 5 10 15

Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Cys
20 25

<210> 8
<211> 28
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 8

Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro Arg Arg
1 5 10 15

Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln
20 25

<210> 9
<211> 22
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 9

Arg Arg Arg Gly Arg Pro Arg Arg Arg Pro Pro Arg Arg Arg Arg Gln
1 5 10 15

Pro Arg Arg Arg Arg Cys
20

<210> 10
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 10

Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro Arg Arg
1 5 10 15

Arg Arg Cys

<210> 11
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 11

Arg Arg Arg Gly Arg Pro Arg Arg Arg Pro Pro Arg Arg Arg Arg Cys
1 5 10 15

<210> 12
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 12

Arg Arg Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg
1 5 10 15

Arg Arg Arg Cys
20

<210> 13
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 13

Arg Arg Arg Pro Pro Arg Arg Arg Arg Gln Pro Arg Arg Arg Arg Cys
1 5 10 15

<210> 14
<211> 33
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<400> 14

Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg Thr Pro Ser Pro Arg Arg
1 5 10 15

Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser His His His His His

His

<210> 15
 <211> 33
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<400> 15

His His His His His His Arg Arg Arg Gly Arg Ser Pro Arg Arg Arg
 1 5 10 15

Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg
 20 25 30

Ser

<210> 16
 <211> 27
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<400> 16

Arg Arg Arg Gly Arg Pro Arg Arg Arg Pro Pro Arg Arg Arg Arg Gln
 1 5 10 15

Pro Arg Arg Arg Arg His His His His His His
 20 25

<210> 17
 <211> 27
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic polypeptide

<400> 17

His His His His His His Arg Arg Arg Gly Arg Pro Arg Arg Arg Pro
 1 5 10 15

Pro Arg Arg Arg Arg Gln Pro Arg Arg Arg Arg
 20 25

<210> 18
 <211> 36
 <212> PRT
 <213> Woolly monkey HepB virus

<400> 18

Thr Val Val Arg Arg Arg Arg Pro Ser Gly Arg Arg Thr Pro Ser Pro
1 5 10 15

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro
20 25 30

Ala Ser Ser Cys
35

<210> 19

<211> 41

<212> PRT

<213> Ground squirrel HepB virus

<400> 19

Thr Val Ile Arg Arg Arg Gly Ser Ala Arg Val Val Arg Ser Pro Arg
1 5 10 15

Arg Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg
20 25 30

Arg Pro Gln Ser Pro Ala Ser Asn Cys
35 40

<210> 20

<211> 42

<212> PRT

<213> Woodchuck HepB virus

<400> 20

Thr Val Ile Arg Arg Arg Gly Gly Ala Arg Ala Ser Arg Ser Pro Arg
1 5 10 15

Arg Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg
20 25 30

Arg Arg Ser Gln Ser Pro Ser Ala Asn Cys
35 40

<210> 21

<211> 43

<212> PRT

<213> Bat HepB virus

<400> 21

Thr Ile Val Arg Arg Arg Gly Gly Ser Arg Ala Thr Arg Ser Pro Arg
1 5 10 15

Arg Arg Thr Pro Ser Pro Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg
20 25 30

Arg Arg Ser Gln Ser Pro Ala Ser Ser Asn Cys

<210> 22
 <211> 36
 <212> PRT
 <213> Duck HepB virus

<400> 22

Arg Lys Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr
 1 5 10 15

Val Val Tyr Gly Arg Arg Arg Ser Lys Ser Arg Glu Arg Arg Ala Pro
 20 25 30

Thr Pro Gln Arg
 35

<210> 23
 <211> 36
 <212> PRT
 <213> Heron HepB virus

<400> 23

Arg Lys Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr
 1 5 10 15

Val Val Tyr Gly Arg Arg Arg Ser Lys Ser Arg Gly Arg Arg Ser Ser
 20 25 30

Pro Ser Gln Arg
 35

<210> 24
 <211> 36
 <212> PRT
 <213> Parrot HepB virus

<400> 24

Arg Lys Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr
 1 5 10 15

Val Val Tyr Gly Arg Arg Arg Ser Lys Ser Arg Glu Arg Ser Ser Ser
 20 25 30

Ser Pro Gln Arg
 35

<210> 25
 <211> 36
 <212> PRT
 <213> Ross' Goose HepB virus

<400> 25

Arg Lys Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr
 1 5 10 15

Val Val Tyr Gly Arg Arg Arg Ser Lys Ser Arg Glu Arg Arg Ala Pro
20 25 30

Thr Pro Gln Arg
35

<210> 26
<211> 36
<212> PRT
<213> Snow goose HepB virus

<400> 26

Arg Lys Pro Arg Gly Leu Glu Pro Arg Arg Arg Lys Val Lys Thr Thr
1 5 10 15

Val Val Tyr Gly Arg Arg Arg Ser Lys Ser Arg Glu Arg Arg Ala Ser
20 25 30

Ser Pro Gln Arg
35

<210> 27
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic polypeptide

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa can be arginine, lysine, or histidine

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> Xaa can be arginine, lysine, or histidine

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>

```

<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> Xaa can be arginine, lysine, or histidine or absent

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> Xaa can be arginine, lysine, or histidine or absent

```

<400> 27

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15

Xaa Xaa Xaa Xaa
20

<210> 28

<211> 14

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic polypeptide

<400> 28

Arg Arg Arg Arg Ser Gln Ser Pro Arg Arg Arg Arg Ser Cys
1 5 10

FIG. 1

	I	II	III	IV
a				
Human	<u>TVVRRRG-</u>	-----RSPRRR	TPSPRRRRSQSP	<u>RRRRSQS-RE</u> SQC
Woolly monkey	<u>TVVRRR--</u>	-----RPSGRR	TPSPRRRRSQSP	<u>RRRRSQS-PA</u> SSC
Ground squirrel	<u>TVIRRRGS</u>	ARVVRSPRRR	TPSPRRRRSQSP	<u>RRR-PQS-PA</u> SNC
Woodchuck	<u>TVIRRRGG</u>	ARASRSPRRR	TPSPRRRRSQSP	<u>RRRRSQS-PS</u> ANC
Bat	<u>TIVRRRRGG</u>	SRATRSPPRR	TPSPRRRRSQSP	<u>RRRRSQSPAS</u> SNC
b				
Duck	<u>RKPRGL</u>	<u>EPRRRKVKTT</u>	VVYGRRRSKS	<u>RERRAPTPQR</u>
Heron	<u>RKPRGL</u>	<u>EPRRRKVKTT</u>	VVYGRRRSKS	<u>RGRRSSPSQR</u>
Parrot	<u>RKPRGL</u>	<u>EPRRRKVKTT</u>	VVYGRRRSKS	<u>RERSSSSPQR</u>
Ross's goose	<u>RKPRGL</u>	<u>EPRRRKVKTT</u>	VVYGRRRSKS	<u>RERRAPTPQR</u>
Snow goose	<u>RKPRGL</u>	<u>EPRRRKVKTT</u>	VVYGRRRSKS	<u>RERRASSPQR</u>

FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6





FIG. 7

A



B




FIG. 8




FIG. 9

