

GENETIC ALTERATION IN ANTISENSE LONG NON-CODING RNA AS A
BIOMARKER FOR SUSCEPTIBILITY TO DISEASE TREATMENT**CROSS-REFERENCE TO RELATED APPLICATIONS**

- 5 [0001] This application claims the benefit of priority of US provisional application No. 62/345,081, filed 03 June 2016, the contents of it being hereby incorporated by reference in its entirety for all purposes.

FIELD OF THE INVENTION

- 10 [0002] The present invention relates generally to the field of molecular biology. In particular, the present invention relates to the use of biomarkers for determining treatment modalities for various diseases, such as cancer.

BACKGROUND OF THE INVENTION

- 15 [0003] Pathway-directed therapeutics, for example those targeting the epidermal growth factor receptor (EGFR)-related pathways, have established clinical activity in various types, of cancer, for example, head and neck squamous cell cancers (HNSCC). It is noted that site-specific differences in treatment outcomes have been observed.

- [0004] Although there have been significant improvements in multimodality approaches
20 to the treatment of cancer, these so far appear to have a full remission rate of only 50%, that is to say only 50% of the cancers are cured. To date, the standard of care provides limited options for the treatment of recurrent, metastatic disease, for example with platinum-based chemotherapy which are known to confer a median overall survival of 6 to 9 months. Despite evidence that a large subset of, for example, HNSCC cancers are dependent on EGFR-
25 signalling, so far only moderate success has been achieved with known treatments based on monoclonal antibodies and/or tyrosine kinase inhibitors (TKI). For example, in the metastatic setting, cetuximab monotherapy is associated with response rates of 13%, while efficacy of epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors across several phase II trials are more heterogeneous, with response rates ranging from 1.8–20 %.

- 30 [0005] Thus, there is a need for biomarkers, genetic, protein or otherwise, that are capable of predicting a subjects susceptibility to a specific drug treatment.

SUMMARY

[0006] In one aspect, the present invention refers to a method of predicting susceptibility of a subject suffering from cancer to a treatment with an anti-cancer drug, wherein the method comprises detecting the presence or absence of a genetic alteration in long non-coding RNA (lncRNA) that resides in an antisense strand of an oncogene, wherein the genetic alteration alters or disrupts expression of the oncogene; wherein in case the genetic alteration is present, the subject is predicted to show improved susceptibility to the treatment with the anti-cancer drug compared to a subject not carrying the mutation.

[0007] In another aspect, the present invention refers to a method of predicting the susceptibility of a subject suffering from cancer related to EGFR to a treatment with an EGFR inhibitor, the method comprising determining whether either one or two or all of the following is given: i) the subject has a silent G>A mutation (genetic alteration) at Q787Q position in exon 20 of EGFR (nucleotide 2361; as shown in NCBI sequence ID: NM_005228.4 and SEQ ID NO: 27); ii) the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to subject that does not respond to treatment with an EGFR inhibitor; iii) the subject has higher EGFR isoform D/isoform A ratio compared to subject that does not respond to treatment with an EGFR inhibitor; wherein in case one or two or all of i) to iii) is given the subject is predicted to show improved susceptibility to a treatment with an EGFR inhibitor compared to a subject not showing any of i) to iii).

[0008] In yet another aspect, the present invention refers to a method of treating cancer in a subject suffering from said cancer, wherein the method comprises detecting the presence or absence of a mutation in a long non-coding RNA on an antisense strand of an oncogene related to the cancer, wherein the genetic alteration alters or disrupts expression of the oncogene; administering an anti-cancer drug for the cancer type the subject suffers from in case the presence of the genetic alteration is confirmed.

[0009] In a further aspect, the present invention refers to a method of treating a subject suffering from cancer, comprising administering to the subject an effective amount of a therapeutic agent affecting expression of a non-coding RNA an oncogene, wherein the genetic alteration alters or disrupts expression of the oncogene; and administering to the subject an effective amount of an anti-cancer drug specific for the cancer related to the oncogene.

[0010] In one aspect, the present invention refers to a method of treating a subject suffering from an EGFR-related cancer, comprising administering to the subject an effective amount of a therapeutic agent affecting EGFR-AS1 lncRNA expression, or affecting EGFR-AS1 expression, or increasing the amount of EGFR isoform D and/or decreasing EGFR isoform A or a combination thereof; and administering to the subject an effective amount of a tyrosine kinase inhibitor used to treat the EGFR-related cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] The invention will be better understood with reference to the detailed description when considered in conjunction with the non-limiting examples and the accompanying drawings, in which:

[0012] **FIG. 1** is a collection of **FIG. 1A – FIG. 1E** showing the correlation of a point mutation in a lncRNA of EGFR, referred to herein as Q787Q, with sensitivity of primary HNSCC cell lines to EGFR tyrosine kinase inhibitors. **FIG. 1A** is a set of line graphs showing the result of a cell proliferation assays and the effect of a treatment with gefitinib, erlotinib or afatinib on the inhibition of cell proliferation. Error bars indicate one standard deviation. **FIG. 1B** is a set of DNA sequence trace chromatograms of Sanger sequencing, showing the disclosed AA genotype in cell lines with increased sensitivity to EGFR tyrosine kinase inhibitors (NCC-HN19 and NCC-HN64). **FIG. 1C** is a set of images of western blot results showing EGFR pathway activation with and without tyrosine kinase inhibitors treatment in HNSCC cell lines with the indicated Q787Q genotype. Each cell line is compared in terms of rate of phosphorylation and also in terms of the effect of a treatment of gefitinib on the rate of phosphorylation, that is the presence of the p-version of the protein, versus the unphosphorylated version of the protein. For example, the first two rows compare the level of phosphorylation between EGFR and p-EGFR (that is the phosphorylated version of EGFR). As gefitinib is a known tyrosine kinase inhibitor, it is expected that the rate of phosphorylation is anti-proportional to the amount of gefitinib administered. That is to say the rate of phosphorylation will decrease with an increase in the amount of gefitinib administered. GAPDH acts as a loading control. **FIG. 1D** is a graph of IC₅₀ values showing sensitivity to gefitinib in correctly targeted G/A^{AAV} clones (CL16, CL19 and CL63) compared to G/G^{AAV} negative controls (clones CL12, CL76 and CL77). This shows that the presence of the G/A mutation results in a stark difference in IC₅₀ values for gefitinib. In terms

of IC_{50} values for drugs, it is noted that the lower the IC_{50} value, the more effective a drug is considered to be. Error bars indicate one standard deviation and p-value as indicated based on student's t-test. **FIG. 1E** is a set of images of western blot results showing inhibition of EGFR pathway activation with 0.1 μ M gefitinib in G/A^{AAV} clones CL16, CL63, CL19 compared to G/G^{AAV} negative control (clone CL12).

[0013] **FIG. 2** is a collection of **FIG. 2A** – **FIG. 2E** showing the implication of the long non-coding RNA EGFR-AS1 as the mechanism for EGFR tyrosine kinase inhibitors sensitivity. **FIG. 2A** is a schematic showing the genomic position of the EGFR-AS1 lncRNA relative to exon 20 of the EGFR gene, and position of the Q787Q SNP, with arrows indicating position of the siRNA designed to knockdown this EGFR-AS1 lncRNA. **FIG. 2B** is a set of column graphs showing expression levels of the EGFR-AS1 lncRNA transcripts measured by real-time RT-PCR, in lines with AA (NCC-HN19 and NCC-HN64) and G/G genotypes (NCC-HN1 and NCC-HN43), and the isogenic NCC-HN1 clones with correct targeting (G/A^{AAV}: CL16, CL63, and CL19) and negative controls (G/G^{AAV}: CL12, CL76, and CL77). Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 2C** is a set of line graphs showing trend of EGFR-AS1 levels (measured by real-time RT-PCR) after Actinomycin D treatment in lines with AA (NCC-HN19 and NCC-HN64) and G/G genotypes (NCC-HN1 and NCC-HN43), and the isogenic NCC-HN1 clones with correct targeting (G/A^{AAV}: CL16, and CL19) and negative controls (G/G^{AAV}: CL12 and CL77). Error bars indicate one standard deviation. **FIG. 2D** is a graph showing IC_{50} values of gefitinib in cell lines with knockdown of EGFR-AS1 (siAS1) compared to non-targeting controls (NT). As stated previously, it is noted that the lower the IC_{50} value, the more effective a drug is considered to be. Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 2E** is a line graph showing tumour growth levels in AS1-high PDXs HN124 and HN159 after treatment with AS1-targeting LNA and non-targeting control. Error bars indicate one standard deviation.

[0014] **FIG. 3** is a collection of **FIG. 3A** – **FIG. 3F** showing effect of EGFR-AS1 long non-coding RNA on EGFR isoforms and sensitivity to tyrosine kinase inhibitors (TKI). **FIG. 3A** is a bar graph showing ratio of EGFR isoform D to A transcripts measured by real-time RT-PCR in the different lines and targeted NCC-HN1 (G/A^{AAV}) clones, with Q787Q genotype as indicated. Error bars indicate one standard deviation. Asterisks denote

significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 3B** is a bar graph showing ratio of EGFR isoform D to A transcripts measured by real-time RT-PCR after knockdown of EGFR-AS1 (siAS1) compared to non-targeted controls (NT). Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 3C** is a set of graphs showing IC_{50} values of the different HNSCC cell lines and NCC-HN1 clones (G/A^{AAV} and G/G^{AAV}; genotypes as indicated) after successful isoform D knockdown (shIsoD) compared to non-targeted controls (NT) treated with gefitinib. Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 3D** is a set of images showing western blot results of the EGFR pathway activation after treatment with gefitinib after isoform D knockdown (shIsoD), compared to non-targeting controls. GAPDH is used here as a loading control. **FIG. 3E** is a graph showing IC_{50} values of the different HNSCC cell lines and NCC-HN1 clones (G/A^{AAV} and G/G^{AAV}; genotypes as indicated) after successful isoform D knockdown (shIsoD), with additional knockdown of EGFR-AS1 (siAS1). The clones being referred to herein are CL16 and CL19 (G/A^{AAV}) and negative controls CL12 and CL77 (G/G^{AAV}). Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$). **FIG. 3F** is a set of graphs showing IC_{50} values for the AA cell lines (NCC-HN19 and NCC-HN64) and G/A HN1 clones treated with gefitinib in RPMI with serum (FCS), serum-free (SF), or serum-free but with added EGF (EGF). Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$).

[0015] **FIG. 4** is a collection of **FIG. 4A** – **FIG. 4D** showing *in vivo* correlation of Q787Q genotype with EGFR-AS1 and EGFR isoform transcript and response to gefitinib treatment. **FIG. 4A** is a set of photographs showing the result of RNA-in situ hybridization using RNAscope showing levels of AS1, EGFR isoform A and D in two tumour specimen which have a A/A and G/G genotypes. PPIB is the positive control and DAPB is the negative control. **FIG. 4B** is a table showing correlation between Q787Q genotype, relative transcript levels of AS1 and isoform D/A ratio (by real-time RT-PCR), along with the formalin-fixed paraffin embedded (FFPE) tissue score (RNA-scope scoring) and IC_{50} levels for a panel of patient derived tumour tissue and cell lines, thereby summarising the information provided in previous figures. **FIG. 4C** is a set of photographs and a line graph showing patient-derived xenograft models for HN137 primary (pri) and metastatic (met) tumours treated with control

or gefitinib. The arrow on the line graph indicates when treatment was initiated. **FIG. 4D** is a set of clinical and CT-scan images showing response of patient HN137 to gefitinib treatment. Arrows indicate location of lung metastasis that subsequently responded to treatment.

[0016] **FIG. 5** is a set of column graphs showing the relative change in EGFR copy number in various cell lines, based on real-time PCR, with the gefitinib IC₅₀ values as indicated. This graph indicates that there is no association between EGFR copy number and the IC₅₀ values. Error bars indicate one standard deviation.

[0017] **FIG. 6** is a schematic showing the targeting construct and strategy designed to alter the Q787Q genotype from G to A. The binding location of the primers used for PCR screening and location of EGFR-AS1 in this construct are indicated.

[0018] **FIG. 7** is a set of DNA gel images showing the PCR screening results showing correct targeting in clones CL16, CL63 and CL19 with random integration in CL12, CL76 and CL77. Random integration means that the clones CL12, CL76 and CL77 do not have the desired effect of a 'knock-in' for the G>A switch, but instead have randomly integrated into the cell line genome, thereby producing a negative control.

[0019] **FIG. 8** is a bar graph showing the relative fold change in levels of EGFR-AS1, as measured by real-time RT-PCR after knockdown (siAS1), compared to non-targeting controls (NT) in cell lines as indicated. The LNA control is a non-targeting LNA. Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- p<0.05, **- p<0.01, ***- p<0.001).

[0020] **FIG. 9** is a bar graph showing real-time RT-PCR result showing relative fold change in the transcript levels of AS1 and isoform D/A ratio after *in vivo* targeting of EGFR-AS1 in HN124 and HN159 patient-derived xenografts. Error bars indicate one standard deviation. Asterisks denote significance by student t-test (*- p<0.05, **- p<0.01, ***- p<0.001).

[0021] **FIG. 10** is an alternative splice diagram of the EGFR gene showing the four commonly described isoforms (A, B, C, and D) and the relevant exons.

[0022] **FIG. 11** is a bar graph showing the relative fold change in levels of the EGFR isoform A and D transcripts, as measured by real-time RT-PCR after knockdown of isoform D with targeted shRNA (shIsoD, also referred to as shEGFR4) compared to non-targeting controls (NT) in cell lines and isogenic NCC-HN1 clones. Error bars indicate one standard

deviation. Asterisks denote significance by student t-test (*- $p < 0.05$, **- $p < 0.01$, ***- $p < 0.001$).

[0023] **FIG. 12** is a table showing clinico-pathologic characteristics of patients from which cell lines were derived.

- 5 [0024] **FIG. 13** is a table showing mutational status of EGFR in the cell lines NCC-HN1, NCC-HN43, NCC-HN19, NCC-HN64, NCC-HN26, and NCC-HN73.

DEFINITION OF TERMS

- [0025] The term “genotype switching” refers to a genetic recombination event (also
10 termed a “knock-in”) where a nucleotide at a given position is switched with another nucleotide, thereby changing the genotype of the subject from one genotype to another. For example, in the experiments has shown herein, the nucleotide residue for position Q787Q at position 2361 on the EGFR gene (as shown in NCBI sequence ID: NM_005228.4) has been switched from a G to an A.

- 15 [0026] The term “tyrosine kinase” refers to an enzyme that can transfer a phosphate group from adenosine triphosphate (ATP) to a target protein in a cell. It functions as an “on” or “off” switch in many cellular functions. Tyrosine kinases are a subclass of protein kinase, of which the tyrosine kinase is named as such because it transfers the phosphate group from ATP to a tyrosine residue within the target protein. There are two known families of tyrosine
20 kinase, namely receptor tyrosine kinase (RTK) and non-receptor or cytoplasmic tyrosine kinase, whereby receptor tyrosine kinases comprise a transmembrane domain and one or more extracellular ligand-binding domains. Cytoplasmic tyrosine kinases do not possess such a transmembrane domain or any extracellular ligand-binding domains.

- [0027] The term “oncogene” refers to a gene that has the potential to cause cancer. An
25 oncogene can also refer to a dominant mutant allele of a cellular gene (a proto-oncogene) that disrupts cell growth and division and is capable of transforming a normal cell into a cancerous cell. In tumour cells, oncogenes are often mutated or expressed at high levels. Proto-oncogenes typically encode proteins involved in, but not limited to, positive control of the cell division cycle, such as, for example, growth factor receptors, signal transduction
30 proteins and transcription factors. Mutations in these genes tend to relax control mechanisms and accelerate cell division, leading to the cell proliferation that is characteristic of cancer. Some oncogenic mutations cause inhibition of programmed cell death (apoptosis), so that

cancerous cells are less likely to be destroyed by the host's immune system. Most normal cells will undergo apoptosis when critical functions are altered. Instead, activated oncogenes can cause those cells designated for apoptosis to proliferate and survive. Some oncogenes can require an additional step, for example, such as mutations in another gene, or environmental factors, such as viral infection, to cause cancer.

[0028] The term "RNA", that is "ribonucleic acid" refers to an organic molecule consisting of a long chain of nucleotides in which the sugar is ribose (or variations thereof) and the bases are adenine, cytosine, guanine, and uracil. There are various types of RNA, for example, but not limited to, messenger RNA (mRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), transfer-messenger RNA (tmRNA), small nuclear RNA (snRNA), antisense RNA (asRNA) and Piwi-interacting RNA (piRNA). All types of RNA are either coding or non-coding, that is the RNA either results in expression of a protein (for example, messenger RNA) or it does not (for example, transfer RNA). A particular example of non-coding RNA is long, non-coding RNA (lncRNA), which refers to non-coding RNA transcripts of 200 nucleotides or longer in length (at least 200 nucleotides). LncRNA are transcripts that range from 200 nucleotides to 100 000 nucleotides (or 200 bases to 100 kb), and distributed are throughout the genome. In some example, the lncRNAs are between 200 nucleotides to 100000 nucleotides, between 200 to 400 nucleotides, between 300 to 700 nucleotides, between 500 to 1000 nucleotides, between 900 to 1300 nucleotides, between 1200 to 2500 nucleotides, between 2400 to 3600 nucleotides, between 3500 to 4800 nucleotides, between 4500 to 10 000 nucleotides, between 9000 and 50000 nucleotides, between 50000 to 75000 nucleotides, between 70000 to 100000 nucleotides in length. In another example, the lncRNAs are at least 200 nucleotides, at least 250 nucleotides, at least 280 nucleotides, at least 320 nucleotides, at least 480 nucleotides, at least 520 nucleotides, at least 550 nucleotides, at least 640 nucleotides, at least 760 nucleotides, at least 830 nucleotides, at least 950 nucleotides, at least 1100 nucleotides, at least 1250 nucleotides, at least 1400 nucleotides, at least 1600 nucleotides, at least 1800 nucleotides, at least 2100 nucleotides, at least 2800 nucleotides, at least 5500 nucleotides, at least 10500 nucleotides, at least 25000 nucleotides, at least 35000 nucleotides, at least 48000 nucleotides, at least 55000 nucleotides, at least 68000 nucleotides, at least 80000 nucleotides in length. Although having little or no known protein-coding capability, they have diverse functions including transcriptional regulation, epigenetic modulation through chromatin modification, and post-transcriptional

regulation. Several of these functions relate to direct (homology-based) binding to DNA, pre-mRNA and mature mRNA. However, it is believed that the three-dimensional structural conformation of lncRNAs plays an important role in extending their wide ranging repertoire and also influences stability of these molecules themselves.

- 5 [0029] The term “RNAi” refers to RNA interference, a process in which RNA molecules inhibit gene function. This interference is based on the ability of double-stranded RNA to interfere with, or suppress, the expression of a gene with a corresponding base sequence. For example, two types of small ribonucleic acid (RNA) molecules – microRNA (miRNA) and small interfering RNA (siRNA) – are important to RNA interference. RNA molecules (or
- 10 RNAs) are the direct products of genes, and these small RNAs can bind, for example, to other specific messenger RNA (mRNA) molecules, thereby either increase or decrease their activity, for example by preventing an mRNA from producing a protein. RNA interference plays an important role in development and in defending cells against parasitic nucleotide sequences from, for example, viruses and transposons.
- 15 [0030] The term “sense strand”, also known as a coding strand, refers to a segment within double-stranded DNA that runs from 5′ to 3′, and which is complementary to the antisense strand of DNA, which runs from 3′ to 5′. The sense strand is the strand of DNA that has the same sequence as the mRNA, which takes the antisense strand as its template during transcription, and eventually (albeit typically, not always) undergoes translation into a
- 20 protein. The antisense strand is thus responsible for the RNA that is later translated to protein, while the sense strand possesses a nearly identical makeup to that of the mRNA. It is noted that for each segment of double-stranded DNA (dsDNA), there will possibly be two sets of sense and antisense, depending on which direction one reads the DNA, since the naming of sense and antisense is relative to perspective. It is ultimately the gene product, or mRNA, that
- 25 dictates which strand of one segment of dsDNA is called sense or antisense. However, it is noted that, for example in prokaryotes, overlapping genes on opposite strands means the sense for one mRNA can be the antisense for another mRNA. In the context of the present invention, the antisense strand obtained from the DNA refers to the RNA segment running in 3′ to 5′ direction.
- 30 [0031] The term “mutation” or “mutated” or “genetic alteration” refers to a natural or artificial modification, or genetic alteration of the genome or part of a nucleic acid sequence of any biological organism, virus or extra-chromosomal genetic element. This mutation can

be induced artificially using, but not limited to, chemicals and radiation, but can also occur spontaneously during nucleic acid replication in cell division. Mutations may or may not produce discernible changes in the observable characteristics (phenotype) of an organism. There are various types of mutations known, which can either be small-scale mutations or
5 large-scale mutations. Examples of small-scale mutations are, but are not limited to, substitution mutations, silent mutations, missense mutations, nonsense mutations, insertions, and deletions. Examples of large-scale mutations are, but are not limited to, amplifications, deletions, chromosomal translocations, interstitial deletions, chromosomal inversions and mutations that result in a loss of heterozygosity. Mutations can also be grouped by their effect
10 on the function of the resulting product. These include, but are not limited to, loss-of-function (inactivating) mutations, gain-of-function (activating) mutations, dominant-negative (antimorphic) mutations, lethal mutations and back or reverse mutations. Point mutations, for example, also known as single base modification, are a type of mutation that causes a single nucleotide base substitution, insertion, or deletion of the genetic material, DNA or RNA. The
15 term “frame-shift mutation” indicates the addition or deletion of a base pair.

[0032] For example, silent mutations are mutations in DNA that do not significantly alter the phenotype of the organism in which they occur. Silent mutations can occur in non-coding regions (outside of genes or within introns), or they may occur within exons. When they occur within exons, they either do not result in a change to the amino acid sequence of a
20 protein (also known as a synonymous substitution), or they result in the insertion of an alternative amino acid with similar properties to that of the original amino acid. In either case, there is no significant change in the resulting phenotype. The phrase silent mutation is often used interchangeably with the phrase synonymous mutation. However, synonymous mutations only occur within exons, and are not always silent mutations. Synonymous
25 mutations are mutations that can affect transcription, splicing, mRNA transport, and translation, any of which could alter phenotype, rendering the synonymous mutation non-silent.

[0033] The term “polymorphism” refers to the existence of two or more distinctly different forms (morphs) within, for example an animal species. In genetics, a (genetic)
30 polymorphism is used to describe essentially inter-individual, functionally silent differences in DNA sequence that make each human genome unique. In other words, a genetic polymorphism is the occurrence, in the same population, of multiple discrete allelic states, of

which at least two have high frequency. Conventionally, the high frequency is defined as being of 1% or more of the population in question. One example of a genetic polymorphism is a single nucleotide polymorphism (SNP), which is a variation in a single nucleotide that occurs at a specific position in the genome, where each variation is present to some appreciable degree within a population (for example, more than 1% of said population).

[0034] The term “susceptibility” refers to the propensity of something, for example a disease, to be likely affected by something else, for example, a treatment for said disease. This effect can be either positive or negative, depending on what is being referenced. For example, if a disease is sensitive to a particular treatment, then the susceptibility of said disease to a particular treatment is a positive effect. It can then be said that the disease is susceptible (or sensitive) to the treatment. On the other hand, if a disease is not susceptible to a given treatment, the disease is then considered to be unresponsive or resistant to said treatment.

[0035] As defined above, the term “predicting susceptibility” refers to the propensity of something, for example a disease, to be likely affected by something else, for example, a treatment for said disease. In other words, to predict susceptibility of a cancer to a particular treatment is to determine whether the cancer would react to a treatment with a certain medicament or anti-cancer drug. It is of note that the term “determining susceptibility” is not synonymous with, for example, “making a prognosis”. The former term only looks at the possible reaction of a disease to a specific drug or therapy, while the latter describes the likelihood of the patient to survive the disease or disease progression as a whole. While, in some cases, it may be possible to correlate the effect of one term on the other, that is to say that a disease reacting well to a given treatment (that is, the disease is susceptible to the treatment) may increase the likelihood of said patient receiving a positive prognosis in regards to the overall disease progression, this is not to be taken as a rule. As a person skilled in the art would appreciate, a positive prognosis depends on many factors patient-specific factors in addition to the disease’s susceptibility for treatment, for example, overall wellbeing of the patient prior to treatment, metabolism, diet, aggressiveness of the (primary) disease, secondary diseases and/or infections and the like. The term “expression” refers to either gene expression, that is the transcription of DNA into messenger RNA (mRNA) by the RNA polymerase, or protein expression, which is the translation of mRNA into a (functional) protein. An expression may be considered up-regulated (or over-expression) or down-

regulated (suppression, low or decreased expression, also termed under-expression), depending on whether an increase or decrease in expression is present, usually compared to a wild-type or a disease-free subject.

[0036] The term “isoform” or “protein isoform” refers to the different forms of a protein encoded from one and the same gene. These proteins are different in both structure and composition, whereby these differences are regulated by alternative splicing of mRNA. This alternative splicing has been shown to have a large impact in proteome diversity. The specificity of produced proteins is derived by protein structure/function, development stage and even the cell type. Isoform formation becomes more complicated when a protein has multiple subunits and each subunit has multiple isoforms.

[0037] The term “alternative splicing” refers to a regulated process during gene expression that results in a single gene coding for multiple proteins. In this process, particular exons (that is, parts of the genetic code that become part of the mature RNA) of a gene may be included within or excluded from the final, processed messenger RNA (mRNA) produced from that gene. The excluded sequences are termed introns, from intragenic region, that is a region inside a gene. The term intron and exon refers to both the DNA sequence within a gene and the corresponding sequence in RNA transcripts. Consequently, the proteins translated from alternatively spliced mRNAs will contain differences in their amino acid sequence and, often, in their biological functions.

[0038] The term “therapeutic agent” refers to a chemical compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient. For example, an anti-diabetic agent is considered a therapeutic agent, in the sense that it is administered to treat, for example, diabetes in a patient.

[0039] The term “locked nucleic acid”, “LNA” or “inaccessible RNA”, refers to a modified RNA nucleotide, in which the ribose moiety of an LNA nucleotide is modified with an extra bridge connecting the 2' oxygen and 4' carbon. The bridge “locks” the ribose in the 3'-endo (North) conformation, which is the same confirmation often found in the A-form duplexes. The, usually synthetic, locked nucleic acid nucleotides can be mixed with DNA or RNA residues in the oligonucleotide, whenever desired, and hybridize with DNA or RNA according to Watson-Crick base-pairing rules. The locked ribose conformation enhances base stacking and backbone pre-organization. This significantly increases the hybridization properties (melting temperature) of oligonucleotides.

[0040] The term “EGFR inhibitor” refers to compounds that are capable of inhibiting or blocking the activity of epidermal growth factor receptors. Various compounds and drugs are not limited to a single effect and can therefore be considered to be EGFR inhibitors, even if they are structurally different. That is to say, the inhibition of EGFR is the combining characteristic of these compounds.

[0041] The term “EGFR-AS1” refers to a 2.8 kb sequence that corresponds to intron and exon 20 (FIG. 2A) expressed by the EGFR gene.

[0042] As used herein, the term “haematological malignancy” or “haematological malignancies” refers to usually malignant neoplasms or cancers which are derived from blood-forming tissue, such as the bone marrow, or in the cells of the immune system. These cancers are also known in the art as blood cancers, or liquid cancers. Haematological malignancies may derive from either of the two major blood cell lineages: myeloid and lymphoid cell lines. The myeloid cell line normally produces granulocytes, erythrocytes, thrombocytes, macrophages and mast cells; the lymphoid cell line produces B, T, NK (natural killer cells) and plasma cells. Lymphomas, lymphocytic leukaemia, and myeloma are derived from the lymphoid line, while acute and chronic myelogenous leukaemia, myelodysplastic syndromes and myeloproliferative diseases are considered to be myeloid in origin. Examples of hematologic cancer are, but are not limited to, leukaemia, lymphoma, and multiple myeloma. Types of leukaemia are, but are not limited to, acute lymphoblastic leukaemia (ALL), acute myelogenous leukaemia (AML), chronic lymphocytic leukaemia (CLL), chronic myelogenous leukaemia (CML), and acute monocytic leukaemia (AMoL). Types of lymphomas are, but are not limited to, Hodgkin's lymphomas, which includes all four subtypes of Hodgkin's lymphomas; and all subtypes of Non-Hodgkin's lymphomas.

DETAILED DESCRIPTION OF THE PRESENT INVENTION

[0043] The success of targeted therapeutics is predicated based on high-precision predictive biomarkers. For example, cohort-based sequencing studies have failed to demonstrate activating mutations in exons 18 to 21 of epidermal growth factor receptor (EGFR), or demonstrate the presence of known predictors of (treatment) response, for example, EGFR amplifications, thereby highlighting a subset of tumours that remain EGFR-driven through non-genomic mechanisms. For example, in a completed phase II trial that examined the impact of induction gefitinib prior to chemo-radiotherapy for unresectable oral

squamous cell cancer, two patients who responded dramatically to gefitinib were observed. In both cases, the performed targeted EGFR-sequencing did not reveal “activating” mutations, thereby showing no apparent correlation between, for example, the treatment response and EGFR amplification, thereby underlining the need for biomarkers, whereby the presence or absence of these biomarkers serves as an indicator of how well a subject will respond to a certain treatment, or if a patient will respond to the intended treatment at all. Thus, in one example, the present invention refers to a method of predicting susceptibility of a subject suffering from cancer to a treatment with an anti-cancer drug.

[0044] The basis for determining subject susceptibility to treatment with a particular drug can be, for example, the presence or the absence of, for example genetic alternations or mutations within the genome, within genetic transcripts, such as RNA or within proteins expressed by genes. The genetic mutation or alteration in question may or may not result in a change in for example protein or RNA sequence, depending on the type of mutation in question. For example, different from an activating mutation, a silent mutation will not result in any discernible change in metabolism or expression product, as the mutation is silent. A point mutation may or may not be silent, depending on the exact mutation at hand and the result of said mutation. For example, if a point mutation results in the change of one amino acid for another, wherein the new amino acid results in a different tertiary structure of the resulting protein, then such a mutation is not considered to be a silent mutation. However, if the point mutation results in a different RNA sequence, which still results in the same amino acid at that point of the protein (due to the redundancy in the genetic code, that is the fact that multiple codons encode for the same amino acid), the point mutation will be considered a silent mutation.

[0045] The present application also discloses a method, which comprises detecting the presence or absence of a genetic alteration in an antisense strand of a long non-coding RNA (lncRNA). This long, non-coding RNA (lncRNA) sequence can reside in an antisense strand of an oncogene. In another example, the long non-coding RNA (lncRNA) sequence resides in the coding strand of an oncogene. Thus, in one example, a method is disclosed herein, which comprises detecting the presence or absence of a genetic alteration in an antisense strand of a long non-coding RNA (lncRNA). In another example, the genetic alteration alters or disrupts expression of the oncogene. In another example, disclosed herein is a method of predicting susceptibility of a subject suffering from cancer to a treatment with an anti-cancer drug,

wherein the method comprises detecting the presence or absence of a mutation in an antisense strand of a non-coding RNA of an oncogene, wherein the mutation alters or disrupts expression of the oncogene. In another example, the long non-coding RNA (lncRNA) sequence resides in the coding strand of an oncogene. In yet another example, if the genetic alteration or mutation is present, the subject is predicted to show improved susceptibility to the treatment with the anti-cancer drug compared to a subject not carrying the mutation. In one example, there is disclosed a method of predicting susceptibility of a subject suffering from cancer to a treatment with an anti-cancer drug, wherein the method comprises detecting the presence or absence of a genetic alteration in a long non-coding RNA (lncRNA) that resides in an antisense strand of an oncogene, wherein the genetic alteration alters or disrupts expression of the oncogene; wherein in case the genetic alteration is present, the subject is predicted to show improved susceptibility to the treatment with the anti-cancer drug compared to a subject not carrying the genetic alteration.

[0046] As used herein, the term “alters” refers to a change in a characteristic, usually in comparison to the same characteristic in a different state. A difference in, for example, expression level of a known gene can be considered to be an alteration of the gene expression of said gene. This is usually given in comparison to the disease-free (or healthy) state of the gene. Such a difference in expression, gene protein or otherwise can be given in absolute or in relative terms. For example, gene Z is expressed at a level of 50 in a disease-free state, given in absolute terms. In a diseases state, gene Z is expressed at a level of 25. In relative terms, the gene expression of gene Z would be 0.5 in the diseased state relative to the disease-free state (also termed to be a down-regulation of the expression of gene Z). In another example, gene Z is expressed at a level of 50 in a disease-free state, given in absolute terms. In a diseases state, gene Z is expressed at a level of 100. In relative terms, the gene expression of gene Z would be 2 in the diseased state relative to the disease-free state (also termed to be up-regulation of the expression of gene Z). In both of these examples, the gene expression of Z is altered.

[0047] As used herein, the term “disrupt” refers to the interruption, interference or termination of a process. For example, the presence of a mutation within a gene sequence can result in the disruption of the translation process, thereby usually resulting in either the truncation of the resulting protein (protein is only partially expressed, for example through

the introduction of a premature stop codon via the mutation) or the complete absence of the protein (no protein is expressed).

[0048] In another example, there is described a method of treating cancer in a subject suffering from said cancer, wherein the method comprises detecting the presence or absence of a mutation or a genetic alteration in a non-coding RNA on an antisense strand of an oncogene related to the cancer, wherein the mutation alters or disrupts expression of the oncogene; administering an anti-cancer drug for the cancer type the subject suffers from in case the presence of the mutation is confirmed. In another example, there is disclosed use of an anti-cancer drug in the manufacture of a medicament for treating cancer in a subject suffering from said cancer, wherein the medicament is to be administered to the subject when the presence or absence of a mutation or a genetic alteration in a non-coding RNA on an antisense strand of an oncogene related to the cancer is detected, wherein the mutation alters, or disrupts expression of the oncogene.

[0049] There are various targets known to function as oncogenetic switches in a cell. These target genes are, but are not limited to, regulatory genes within any given pathway involved in cell growth and regulation of cell proliferation and/or apoptosis. As phosphorylation is one of the best known examples of cellular switches used for regulating cellular pathways, genes that express kinases (enzymes responsible for the transfer of phosphor groups from one protein to another protein) are prominent targets in the development of anti-cancer drugs and regimens. Other categories of oncogenes include, but are not limited to, receptor tyrosine kinases, tyrosine kinases, cytoplasmic tyrosine kinases, cytoplasmic serine/threonine kinases, cytoplasmic serine/threonine kinases regulatory subunits, cyclin-dependent kinases, regulatory GTPases and transcription factors. Thus, in one example, the oncogene is a gene that results in, but is not limited to, the expression of a receptor tyrosine kinase, a protein target of a receptor tyrosine kinase or in a cytoplasmic tyrosine kinase. In another example, the oncogene is a gene that results in the expression of a receptor tyrosine kinase. In another example, the oncogene that results in the expression of a receptor tyrosine kinase is, but is not limited to, EGFR, IGF1R, PIK3CD, PIK3R3, PIK3CD, ERBB4, FGFR2, FGFR3, FGFR4, c-Kit, PDGFRA, PDGFRB, PIK3CA, PIK3Ra, PIK3R2, PIK3R3, ERBB2, ERBB3 and INSR. In one example, the oncogene is a gene that results in the expression of a cytoplasmic tyrosine kinase. In another example, the oncogene that results

in the expression of a cytoplasmic tyrosine kinase is, but is not limited to, mTOR, MAP3K1 (MEKK) MAPK8 (JNK) and BRAF.

[0050] In another example, an oncogene includes, but is not limited to, epidermal growth factor receptor (EGFR), platelet-derived growth factor receptor (PDGFR), vascular endothelial growth factor receptor (VEGFR), PIK3CB, PIK3R3, PIK3CD, ERBB4, BRAF, 5 FGFR2, FGFR3, FGFR4, c-Kit, MAPK3K1 (MEKK), MAPK8 (JNK), PDGFRA, PDGFRB, PIK3CA, PIK3R1, PIK3R2, ERBB3, INSR, abl, af4/hrx, akt-2, alk, alk/npm, aml1, aml1/mtg8, axl, bcl-2, bcl-3, bcl-6, bcr/abl, c-myc, dbl, dek/can, E2A/pbx1, enl/hrx, erg/TLS, erbB, erbB-2/ HER2/neu, ets-1, ewsfli1, fms, fos, fps, gli, gsp, hox11, hTERT, hst, IGF1R, 10 IL-3, int-2, jun, kit, kmt2b, kmt2c, kmt2d, KS3, K-sam, Lbc, lck, imo1, lmo2, L-myc, lyl-1, lyt-10, lyt-10/C-alpha-1, mas, mdm2, mll, mos, mtg8/aml1, myb, MYH11/CBFB, n-myc, ost, pax-5, pbx1/EA2, pim-1, PRAD-1, raf, rar/pml, ras, rasH, rasK, rasN, rel/nrg, ret, rhom1, rhom2, ros, ski, SRC, sis, tal1, tal2 (SCL), tan-1, tiam1, TSC2, and trk. In another example, the oncogene can include, but is not limited to, of EGFR, IGF1R, mTOR, PIK3CB, PIK3R3, 15 PIK3CD, ERBB4, BRAF, FGFR2, FGFR3, FGFR4, c-Kit, MAPK3K1 (MEKK), MAPK8 (JNK), PDGFRA, PDGFRB, PIK3CA, PIK3R1, PIK3R2, ERBB2, ERBB3 and INSR. In yet another example, the oncogene can be, but is not limited to, EGFR, IGF1R, mTOR, PIK3CB, PIK3R3, PIK3CD, and ERBB4. In a further example, the oncogene is EGFR.

[0051] As defined above, various genes become oncogenes, in some examples, due to a 20 mutation within their genetic sequence, thereby resulting in deviations from the usual function of said gene. In terms of gene targeting, there are multiple levels which can be targeted using available technology. That is to say, influencing gene expression and/or the resulting protein expression can be done on different levels. For example, one can influence the expression of a certain gene by silencing said gene using, for example, siRNA. Thus, the 25 present disclosure describes methods of detecting differences in gene expression, based on which a susceptibility of a subject to a particular treatment is inferred.

[0052] In one example, this inference is performed based on the presence or absence of a mutation within the oncogene, or in any expression products of the oncogene. In one example, the presence of a mutation has an effect on the functionality of the resulting protein. 30 In another example, the presence of the mutation does not have an effect on the functionality of the resulting protein.

[0053] Determination of the differences in gene expression can be performed using methods known to those skilled in the art. For example, gene sequencing can be used to ascertain if a mutation is present on a nucleic acid level. In another example, the comparison of gene expression can be done on a RNA level, that is for example, by ascertaining and comparing the levels of an RNA transcript of one or more target genes. In another example, the comparison of expression levels of a gene is made on the protein level. This can be performed, for example, by comparing the level of a protein expressed by a target gene in a diseased subject and comparing the level of the same protein in a disease-free subject.

[0054] In one example, the presence or absence of the mutation is determined in the RNA transcript of the gene. In another example, the presence or absence of the mutation is determined on the antisense strand of the RNA. In yet another example, the presence or absence of the mutation is determined in a non-coding region of the antisense strand of the RNA. In another example, the presence or absence of the mutation is determined on the antisense strand of a long non-coding RNA region (lncRNA) of the oncogene in question.

The antisense strand of a long non-coding RNA region (lncRNA) of an oncogene can be, but is not limited to, EGFR-AS1 for EGFR gene; IRAIN for IGF1R gene, MTOR-AS1 for MTOR gene; GAPDHP39 and RPL23AP40 for PIK3CB gene; LOC101929626 for PIK3R3 gene; PIK3CD-AS1, PIK3CD-AS2 and RPL26P7 for PIK3CD gene and RNA5SP119 for ERBB4 gene. In another example, the mutation is determined in exon 20 of EGFR-AS1 of EGFR. In yet another example, the mutation is a silent mutation at position Q787Q of exon 20 in exon 20 of EGFR-AS1 of EGFR. In another example, the silent mutation is a G>A mutation at position Q787Q of exon 20 in exon 20 of EGFR-AS1 of EGFR.

[0055] Mutations on a genetic level, for example in the mRNA or in the gene itself, can result in the expressed protein or RNA being different from that which is usually expressed in the majority of the population. This difference can be seen in various ways, for example when a genetic mutation results in the over- or under-expression of the resulting (functional) protein or RNA transcript. In one example, the genetic mutation results in an over- or under-expression of the resulting RNA transcript in a subject. In a further example, the genetic mutation results in the expression of a truncated RNA transcript or no expression of an RNA transcript. In another example, the genetic mutation results in a non-functional protein being expressed. In yet another example, the genetic mutation results in a truncated protein being expressed. In another example, the genetic mutation results in a different isoform of the

- protein being expressed. In yet another example, the genetic mutation results in a change in ratio of various proteins, for example, the increased expression of a normally under-expressed isoform. In one example, a mutation in the antisense strand of the long non-coding RNA of EGFR (EGFR-AS1) results in the increased expression of EGFR isoform D. In another
- 5 example, a mutation in the antisense strand of the long non-coding RNA of EGFR (EGFR-AS1) results in an increased EGFR isoform D to EGFR isoform A ratio (EGFR isoform D/A). In yet another example, there is disclosed a method of treating a subject suffering from an EGFR-related cancer, comprising administering to the subject an effective amount of a therapeutic agent affecting EGFR-AS1 lncRNA expression, or affecting EGFR-AS1
- 10 expression, or increasing the amount of EGFR isoform D and/or decreasing EGFR isoform A or a combination thereof; and administering to the subject an effective amount of a tyrosine kinase inhibitor used to treat the EGFR-related cancer. In another example, there is disclosed the use of a tyrosine kinase inhibitor in the manufacture of a medicament for treating a subject suffering from an EGFR-related cancer.
- 15 [0056] In another example, there is described a method of predicting the susceptibility of a subject suffering from cancer related to EGFR to a treatment with an EGFR inhibitor, the method comprising determining whether either one or two or all of the following is given: i) the subject has a silent G>A mutation at Q787Q position in exon 20 of EGFR; ii) the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to subject that does
- 20 not respond to treatment with an EGFR inhibitor; iii) the subject has higher EGFR isoform D/isoform A ratio compared to subject that does not respond to treatment with an EGFR inhibitor; wherein in case one or two or all of i) to iii) is given the subject is predicted to show improved susceptibility to a treatment with an EGFR inhibitor compared to a subject not showing any of i) to iii).
- 25 [0057] A genetic alteration or mutation within a genome can result in a genotype that is either homozygous or heterozygous for said mutation. As used herein, the terms “homozygous” and “heterozygous” refer to the degree of similarity between the alleles of a certain characteristic, or trait, of an organism. This is based on the fact that most eukaryotes are diploid, that means they have two matching sets of chromosomes. Both sets of
- 30 chromosomes have the same loci on each of them. Thus, an organism that has a homozygous genotype is describing an organism (or cell) in which the alleles at a given locus are identical. On the other hand, if an organism is described as being heterozygous for a certain allele, this

means that in the same locus, one chromosome shows one genotype (for example the nucleotide A), while the other chromosome shows a different genotype (for example, the nucleotide T) in the same locus. The question of homo- or heterozygosity can also be determined based on the presence or absence of genetic alterations or mutations in the
5 respective RNA or RNA transcripts from each allele. Thus, in one example, if the RNA transcripts from alleles show the same mutation in the same locus, then the organism is considered to be homozygous for that mutation. In a preferred example, the mutation or genetic alteration is homozygous. In yet another example, the mutation or genetic alteration is heterozygous.

10 [0058] It is also possible to determine the presence or absence of a genetic alteration or mutation on, for example, the corresponding antisense RNA strand or in long non-coding RNA (lncRNA). The present examples disclose a mutation or genetic alteration at a specific position within the EGFR-AS1 sequence. Thus, in one example, a defined locus within an antisense strand of a long non-coding RNA sequence is homozygous for a mutation or
15 genetic alteration. In another example, the genetic alteration or mutation in long non-coding RNA (lncRNA) that resides in an antisense strand of an oncogene is homozygous. In one example, the mutation or genetic alteration is present in the EGFR-AS1 sequence. In another example, the mutation or genetic alteration is present in the Q787Q locus of the EGFR-AS1 sequence. In one example, this locus is homozygous for a mutation or genetic alteration. This
20 means that at this location, the mutation or genetic alteration (and as a result the genotype) on both alleles are the same. In another example, this locus is heterozygous for the mutation or genetic alteration. In a further preferred example, the homozygous mutation or genetic alteration is AA. In a further example, the heterozygous mutation or genetic alteration is GA.

[0059] The basis of comparison for determining the presence or absence of a mutation
25 and/or the effect of such a mutation on the expression of the resulting protein or RNA is the comparison with a subject who is either healthy, that is disease-free, or a subject who has the same disease as the subject in question, but who is known not to respond to the treatment being assessed for the subject in question. For example, subject A has head and neck cancer. A mutation is found within the oncogene closely related to the cancer, for example EGFR.
30 Subject B also has head and neck cancer and was treated with anti-cancer drug X, which was not effective in the treatment of the cancer. Subject C also has head and neck cancer and was successfully treated with anti-cancer drug X. Genetic comparison of the oncogene closely

related to the cancers in subjects B and C (in this example, EGFR) show that the oncogene of subject B does not have a mutation in a relevant, pre-determined region of the oncogene, while subject C does present a mutation in the relevant, pre-determined region of the oncogene. Thus, subject A, having a mutation present in the same region as subject C, is shown to be susceptible to treatment with anti-cancer drug X. In another example, the determination of possible susceptibility to treatment with a given anti-cancer drug can be done based on the comparison of, for example, protein or RNA levels, or in the case of multiple proteins or RNA transcripts, and/or ratios of the proteins concerned. In one example, it is determined whether the subject has a higher EGFR isoform D/isoform A ratio compared to a subject that does not respond to treatment with the anti-cancer drug. In one example, it is determined whether the subject has a lower RNA transcript level compared to a subject that does not respond to treatment with the anti-cancer drug. In another example, it is determined whether the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to a subject that does not respond to treatment with the anti-cancer drug. In yet another example, the method as described herein further comprises measuring of either one or both of the following for predicting the susceptibility: i) whether the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to a subject that does not respond to treatment with the anti-cancer drug; ii) whether the subject has higher EGFR isoform D/isoform A ratio compared to a subject that does not respond to treatment with the anti-cancer drug, wherein the anti-cancer drug is a tyrosine kinase inhibitor or an EGFR inhibitor.

[0060] Thus, in one example an increase in the amount of EGFR isoform D in a patient is indicative of an increased sensitivity to treatment with, for example, a tyrosine kinase inhibitor for EGFR-related cancers, or treatments used for other gene specific cancers. In another example, a decrease in the amount of EGFR isoform A in a patient is indicative of an increased sensitivity to treatment with a tyrosine kinase inhibitor. In other words, a high ratio of EGFR isoform D to EGFR isoform A is considered to be indicative an increased sensitivity to treatment with, for example, a tyrosine kinase inhibitor for EGFR-related cancers, or treatments used for other gene specific cancers. Conversely, a low ratio of EGFR isoform D to EGFR isoform A is considered to be indicative of a possible resistance to treatment with, for example, a tyrosine kinase inhibitor for EGFR-related cancers, or treatments used for other gene specific cancers.

- [0061] It is known that certain oncogenes are more closely associated with some types of cancer than others. For example, the gene HER2 is a known oncogene most closely associated, but not only associated, with certain subtypes of breast cancer. Thus, in the present disclosure, the absence or presence of a mutation in an oncogene is understood to have an effect on the subject's susceptibility to a treatment for, for example, a cancer related to said oncogene. In one example, the cancer related to EGFR, PIK3CB, PIK3R3 and PIK3CD is, but is not limited to, non-small cell lung carcinoma, head and neck cancer, colorectal carcinoma, breast cancer, brain malignancies including glioblastomas, haematological malignancies, prostate cancer, bladder cancer, renal cell carcinoma, pancreas cancer, cervical cancer, oesophageal cancer, gastric cancer and ovarian cancer. In another example, the cancer related to EGFR is head and neck cancer, or lung cancer. In yet another example, the head and neck cancer can be head and neck squamous cell cancer (HNSCC) or oral squamous cell cancer (OSCC). In yet another example, the lung cancer is non-small cell lung cancer (NSCLC). In one example, the cancer related to PIKCD is haematological malignancies. In one example, the cancer related to ERBB4 is breast cancer. In one example, the cancer related to EGFR is, but is not limited to, non-small cell lung cancer (NSCLC), head and neck cancer, colorectal carcinoma, breast cancer, brain malignancies including glioblastoma, prostate cancer, bladder cancer, renal cell carcinoma, pancreas cancer, cervical cancer, oesophageal cancer, gastric cancer and ovarian cancer.
- [0062] Just as there are known genes closely related to particular types of cancer, there are also treatments, for example anti-cancer treatments that are known to work better with certain cancer types compared to other treatments for the same type of cancer. Without being bound by theory, it is thought that such differences in treatment susceptibility are the result of the anti-cancer treatment pinpointing, for example, a defective cellular or signalling pathway. For example, a cancer related to the HER2 gene, wherein the HER2 gene is defective or results in a defective product (for example, RNA or protein RNA or protein), said cancer related to the HER2 gene can be more susceptible to treatment with for example trastuzumab, compared to a treatment with other anti-cancer drugs. Therefore, in one example, the anti-cancer drug is, but is not limited to gefitinib, erlotinib and afatinib for the treatment of cancer related to EGFR; OSI-906 (linsitinib) for the treatment of cancer related to IGF1R; everolimus (also known as RAD001) and sirolimus for the treatment of cancer related to mTOR; BKM120 (buparlisib) and BYL719 (alpelisib) for the treatment of cancer related to PIK3CB and

- PIK3R3; idelalisib for the treatment of cancer related to PIK3CD and dacomatinib and lapatinib for the treatment of cancer related to ERBB4, or combinations thereof. In one example, the anti-cancer drug used for treating EGFR-related cancers is, but is not limited to, gefitinib, erlotinib, afatinib or combinations thereof. In another example, the anti-cancer drug used for treating mTOR-related cancers is, but is not limited to, everolimus (RAD001), sirolimus, or combinations thereof. In another example, the anti-cancer drug used for treating IGF1R-related cancers is, but is not limited to, linsitinib. In another example, the anti-cancer drug used for treating PIK3CB and PIK3R3-related cancers is, but is not limited to, BKM120 (buparlisib), BYL719 (alpelisib) or combinations thereof. In another example, the anti-cancer drug used for treating PIK3CD-related cancers is, but is not limited to, idelalisib. In another example, the anti-cancer drug used for treating ERBB4-related cancers is, but is not limited to, dacomatinib, lapatinib, or combinations thereof. In one example, the anti-cancer drug is a tyrosine kinase inhibitor. In another example, the tyrosine kinase inhibitor is an EGFR inhibitor. In yet another example, the tyrosine kinase inhibitor is, but is not limited to, gefitinib, erlotinib, erlotinib HCl, lapatinib, dacomitinib, TAE684, afatinib, dasatinib, saracatinib, veratinib, AEE788, WZ4002, icotinib, osimertinib, BI1482694, ASP8273, EGF816, AZD3759, cetuximab, necitumumab, panitumumab, nimotuzumab and combinations thereof. In a further example, the tyrosine kinase inhibitor is, but is not limited to, gefitinib, erlotinib, lapatinib and combinations thereof.
- [0063] It is understood that some drugs/therapeutic agents, some of which are disclosed herein, can be used to treat most cancer types, albeit with differing efficacies. While, for example, gefitinib is not usually used in the treatment of, for example, head and neck squamous cell cancer (HNSCC), it is shown that according to the present invention, head and neck squamous cell cancer (HNSCC) can indeed be treated with gefitinib.
- [0064] Some approaches are based on multi-pronged approach for treating a disease, while other may be based on a single-pronged approach, that is single treatment regimen, for a particular disease. In one example, the disease is treated using a single treatment. In another example, the disease is treated using at least two, at least three, at least four or more treatments. These treatments may be given subsequently, simultaneously or in combinations thereof. In one example, the disease is treated using at least two, at least three, at least four or more drugs.

[0065] Many types of anti-cancer drugs or treatments are available based on the specific type of cancer to be treated. For example, breast cancer can be treated with, but not limited to, any one of the following anti-cancer drugs, or with combinations thereof: everolimus (RAD001), tamoxifen, toremifene, trastuzumab, fulvestrant, anastrozole, exemestane, lapatinib, letrozole, pertuzumab, ado-trastuzumab emtansine, and palbociclib. A known mutation in an oncogene can therefore result in a more effective anti-cancer being chosen, as opposed to any of the above listed drugs being chosen. For example, if a mutation is detected in the HER2 gene in a breast cancer sample obtained from a subject, then the selection of trastuzumab for treating said cancer would be made, as trastuzumab is known for its high success rate in the treatment of HER2-mutation positive breast cancers.

[0066] The following provides various types of cancer and those anti-cancer drugs used to treat them. For example, adenocarcinomas of the stomach or gastro-oesophageal junction can be treated with, but not limited to, trastuzumab, ramucirumab or combinations thereof. Basal cell carcinomas can be treated with, but not limited to vismodegib, sonidegib or combinations thereof. Bladder cancer can be treated with, for example but not limited to, atezolizumab, everolimus, sirolimus and combinations thereof. Brain cancer can be treated with, but not limited to, bevacizumab, everolimus or combinations thereof. Breast cancer can be treated with, but not limited to, everolimus, tamoxifen, toremifene, trastuzumab, fulvestrant, anastrozole, exemestane, linsitinib, lapatinib, letrozole, pertuzumab, ado-trastuzumab emtansine, palbociclib, alpelisib and combinations thereof. Cervical cancer can be treated with, for example, but not limited to, bevacizumab. Colorectal cancer can be treated with, but not limited to, cetuximab, panitumumab, bevacizumab, ziv-aflibercept, regorafenib, ramucirumab and combinations thereof. Dermatofibrosarcoma protuberans can be treated with, for example, but not limited to, imatinib mesylate. Endocrine/neuroendocrine tumours can be treated with, for example, but not limited to, lanreotide acetate. Head and neck cancers can be treated with, but are not limited to, cetuximab, gefitinib, erlotinib, afatinib, alpelisib, linsitinib, buparlisib, idelalisib, dacomitinib, lapatinib and combinations thereof. Gastrointestinal stromal tumours can be treated with, but are not limited to, imatinib mesylate, sunitinib, regorafenib, and combinations thereof. Giant cell tumour of the bone can be treated with, for example, but not limited to, denosumab. Kaposi sarcoma can be treated with, for example, but not limited to, alitretinoin. Kidney cancer can be treated with, but not limited to bevacizumab, sorafenib, sunitinib, pazopanib, temsirolimus, everolimus, axitinib,

- nivolumab, cabozantinib, lenvatinib mesylate and combinations thereof. Leukaemia can be treated with, for example, but not limited to, tretinoin, imatinib mesylate, dasatinib, nilotinib, bosutinib, rituximab, alemtuzumab, ofatumumab, obinutuzumab, ibrutinib, idelalisib, blinatumomab, venetoclax and combinations thereof. Liver cancer can be treated with, for example, but not limited to, sorafenib. Lung cancer can be treated with, for example, but not limited to, bevacizumab, crizotinib, erlotinib, gefitinib, afatinib dimaleate, ceritinib, ramucirumab, nivolumab, pembrolizumab, osimertinib, necitumumab, alectinib and combinations thereof. Lymphomas can be treated with, for example, but not limited to, ibritumomab tiuxetan, denileukin diftitox, brentuximab vedotin, rituximab, vorinostat, romidepsin, bexarotene, bortezomib, pralatrexate, ibrutinib, siltuximab, idelalisib, belinostat, obinutuzumab, nivolumab and combinations thereof. Melanomas can be treated with, for example, but not limited to, ipilimumab, vemurafenib, trametinib, dabrafenib, pembrolizumab, nivolumab, cobimetinib, everolimus, sirolimus and combinations thereof. Multiple myelomas can be treated with, for example, but not limited to, bortezomib, carfilzomib, panobinostat, daratumumab, ixazomib citrate, elotuzumab and combinations thereof. Myelodysplastic/myeloproliferative disorders can be treated with, for example, but not limited to, imatinib mesylate, ruxolitinib phosphate and combinations thereof. Neuroblastomas can be treated with, for example, but not limited to, dinutuximab. Ovarian epithelial/fallopian tube/primary peritoneal cancers can be treated with, for example, but not limited to, bevacizumab, olaparib and combinations thereof. Pancreatic cancer can be treated with, for example, but not limited to, erlotinib, everolimus, sunitinib and combinations thereof. Prostate cancer can be treated with, for example, but not limited to, cabazitaxel, enzalutamide, abiraterone acetate, radium 223 dichloride, linsitinib and combinations thereof. Soft tissue sarcoma can be treated with, for example, but not limited to, pazopanib. Systemic mastocytosis can be treated with, for example, but not limited to, imatinib mesylate. Thyroid cancer can be treated with, for example, but not limited to, cabozantinib, vandetanib, sorafenib, lenvatinib mesylate and combinations thereof. The listing of a particular anti-cancer drug in the categories above does not preclude its use for treating other types of cancer.
- [0067] Specific methods disclosed herein further comprise the administration of an additional therapeutic agent (i.e., a therapeutic agent other than the anti-cancer treatment disclosed herein). In certain examples, the anti-cancer treatments can be used in combination

with at least one other therapeutic agent. Therapeutic agents include, but are not limited to antibiotics, anti-emetic agents, antidepressants, anti-fungal agents, anti-inflammatory agents, antiviral agents, other anticancer agents, immunomodulatory agents, expression modulating agents, alpha-interferons, gene silencing agents, agents capable of suppressing expression of
5 RNA transcripts or proteins, agents capable of affecting RNA or protein expression, β -interferons, alkylating agents, hormones, or cytokines. In one example, the method encompasses the administration of an additional therapeutic agent that demonstrates gene silencing activity. In another example, the therapeutic agent is capable of RNA interference. In yet another example, the therapeutic agent is selected from, but not limited to, antisense
10 oligonucleotides, short hairpin RNA (shRNA), small interfering RNA (siRNA), double stranded RNA (dsRNA), microRNA (miRNA), locked nucleic acids (LNAs) ribozymes, histone modification, RNA-directed DNA methylation, paramutations or combinations thereof. In one example, the therapeutic agent affects RNA expression. In yet another example, a method of treating a subject suffering from cancer is disclosed, comprising
15 administering to the subject an effective amount of a therapeutic agent affecting expression of a non-coding RNA an oncogene, wherein the mutation alters or disrupts expression of the oncogene; and administering to the subject an effective amount of an anti-cancer drug specific for the cancer related to the oncogene. In another example, the therapeutic agent affects EGFR-AS1 expression. In a further example, the therapeutic agent affecting EGFR-
20 AS1 expression comprises, but is not limited to, miRNA, shRNA, locked nucleic acids (LNAs), for example locked RNA or DNA, or siRNA. In yet another example, the therapeutic agent is one or more locked nucleic acids (LNAs). In another example, the therapeutic agent suppressing EGFR-AS1 expression comprises EGFR-AS1 targeting locked nucleic acids (LNA). In another example, the therapeutic agent increasing EGFR isoform D
25 or decreasing EGFR isoform A comprises an agent modulating the alternative splicing of EGFR.

[0068] The drugs and treatments as disclosed herein and the other therapeutics agent can act additively or, synergistically. In one example, an anti-cancer drug is administered concurrently with the administration of another therapeutic agent, which can be part of the
30 same composition or in a different composition. In another example, the anti-cancer agent is administered prior to or subsequent to administration of another therapeutic agent. In a separate example, an anti-cancer drug is administered to a subject who has not previously

undergone or is not currently undergoing treatment with another therapeutic agent. In one example, the methods of the invention comprise the administration of one or more anti-cancer treatments without an additional therapeutic agent. In another example, the methods of the invention comprise the administration of one or more anti-cancer treatments with at least one
5 or more additional therapeutic agents.

[0069] The methods and the treatments disclosed herein may be performed or carried out simultaneously, separately, one after the other, or in combination with other treatments. In one example, these treatments are, but are not limited to, radiation therapies, chemotherapy, surgery, and combinations thereof. In case, for example, multiple treatments are
10 implemented, these treatments can be performed or carried out, for example, one after the other, with a time interval between each treatment step, wherein, for example, the time interval between the first and the second treatment step is at least 1 to 24 hours, or at least 1, at least 4, at least 6, at least 8, at least 12, at least , or at least 1 or 2 or 3 or 4 or 5 or 6 or 7 days, or at least one week.

15 ***Differential sensitivity to EGFR tyrosine kinase inhibitor is mediated by a silent polymorphism in EGFR exon 20 in patient derived oral squamous cell carcinoma cell lines***

[0070] Six patient-derived cell lines established in the laboratory were tested for sensitivity to EGFR inhibitors using gefitinib, erlotinib and afatinib (FIG. 1A). As shown,
20 majority of cell lines were insensitive to EGFR inhibition, except for NCCWHN19 and NCCWHN64, each with gefitinib IC₅₀ values within the therapeutic range (0.07 and 0.26 μ M respectively). Targeted re-sequencing did not identify sensitizing EGFR mutations, nor was any correlation between drug sensitivity and EGFR copy number demonstrated (FIG. 5). Instead, the two sensitive lines were homozygous for the same synonymous SNP identified in
25 a phase 2 trial (rs10251977, 2361 G>A, Q787Q; SEQ ID NO: 27) with the A/A genotype, while the insensitive cell lines were either homozygous wild type (G/G) or heterozygous (G/A) (FIG. 1B). In line with the observed phenotype, Western blots performed on the gefitinib-sensitive cell lines NCC-HN19 and NCC-HN64 (A/A genotype) showed a significant and consistent reduction in EGFR, AKT, ERK and S6 phosphorylation, after
30 treatment with therapeutic doses of gefitinib (FIG. 1C). In contrast, cell lines with the G/G genotype (NCC-HN1 and NCC-HN43) required much higher drug doses to show effect on phosphor-AKT and phosphor-S6 levels, with modest, if any effect on phosphor-ERK.

Single nucleotide targeting reverse EGFR tyrosine kinase inhibitor resistant phenotype in isogenic cell lines

[0071] The Horizon AAV targeting system was utilized to genetically knock-in the single nucleotide alteration and convert a resistant line (NCC-HN1) to a sensitive line in an isogenic cell line system (FIG. 6 and 7). Sanger sequencing of the expressed EGFR cDNA confirmed that the A genotype was expressed in the successfully targeted clones (G/A^{AAV}:NCC-HN1 CL16, 63 and 19 respectively), compared to negative controls where the vector has integrated randomly (G/G^{AAV}:NCC-HN1 CL12, 76 and 77 respectively) (data not shown). Drug treatment of the G/A^{AAV} clones showed an increased sensitivity to gefitinib compared to negative controls. IC₅₀ values of G/A^{AAV} clones ranged from 0.1-0.3 μ M, compared to negative controls (G/G^{AAV}) ranged from 6.4-7.2 μ M (IC₅₀ for NCC-HN1 parental cell line ranged from 7-11 μ M; FIG. 1D), with consistent modulation of downstream pathways (FIG. 1E).

15

EGFR-AS1 long non-coding RNA drives in vitro and in vivo EGFR addiction

[0072] *In silico* analyses showed no potential miRNA targeting sites that could affect EGFR mRNA transcription or translation (data not shown). However, it was found that this described SNP was within the transcribed portion of the EGFR-AS1 lncRNA (FIG. 2A). Real-time RT-PCR showed that transcript levels of EGFR-AS1 lncRNA were significantly higher in the resistant lines with the G/G genotype (NCC-HN1 and NCC-HN43), compared to the sensitive A/A genotype (NCC-HN19 and NCC-HN64; FIG. 2B). Similar findings were seen in the NCC-HN1 isogenic clones: lower EGFR-AS1 transcript levels in G/A^{AAV} compared to control G/G^{AAV} clones. Using Actinomycin D to block transcription, it was next demonstrated that the EGFR-AS1 lncRNA transcript was more stable in lines with the G/G genotype compared to the A/A genotype, which was recapitulated in the isogenic NCC-HN1 clones with genotype switch (FIG. 2C). Finally, knockdown of EGFR-AS1 (FIG. 9) was shown to be sufficient to significantly increase sensitivity of G/G cell lines to gefitinib, with reduction in mean IC₅₀ values from 8.9 to 2.3 μ M for NCC-HN1 and from 9.6 to 2.8 μ M for NCC-HN43 (FIG. 2D). To test if EGFR-AS1 was a bona fide driver, it was determined whether tumours with G/A or G/G genotype were dependent on the lncRNA levels by *in vivo* knock-down in a patient-derived xenograft (PDX) system. A panel of locked nucleic acid

30

(LNA) against EGFR-AS1 was designed and the most effective candidate selected through *in vitro* screens for effective knock-down (data not shown). *In vivo* grade version of this AS1-targeting LNA and control non-targeting LNA were subsequently injected (weekly dose of 5mg/kg) into the tail veins of NOD-scid-gamma (NSG) mice harbouring patient-derived xenografts of a tumour (HN124) with G/A Q787Q genotype and high AS1 levels (FIG. 4B). After one week of LNA-only treatment, mice were started on daily gefitinib doses (25 mg/kg) for the rest of the experiment. Successful knock-down of AS1 levels was seen in patient-derived xenografts one week after treatment with AS1-targeting LNA compared to non-targeting controls (FIG. 8). AS1-knockdown *in vivo* was sufficient to cause tumour regression even before gefitinib was initiated, but this regression was sustained after treatment, compared to controls where neither control LNA nor gefitinib showed any effect (FIG. 2E). The same experiment was repeated in a different patient-derived xenograft (HN159-G/G genotype, high AS1 levels), this time omitting the gefitinib treatment. Again, it was shown that mice treated with AS1-targeting LNA (with successful knockdown) resulted in sustained tumour regression compared to controls, demonstrating tumour addiction to the lncRNA.

EGFR tyrosine kinase inhibitor sensitivity mediated through differential expression of EGFR isoforms

[0073] Given that the lncRNA had no effect on transcript, protein stability, or results in aberrant splicing of EGFR (data not shown), it was proceeded to examine the effect on expression levels of the four known EGFR isoforms (A-D) (FIG. 10). Real-time PCR showed that there was a higher ratio of isoform D:A transcript levels in both lines with the A/A genotype (NCC-HN19 and NCC-HN64), compared to the lines with the G/G genotype (NCC-HN1 and NCC-HN43; FIG. 3A). These results were recapitulated in the NCC-HN1 genotype switched G/A^{AAV} clones. Moreover, targeted knock-down of the EGFR-AS1 lncRNA was sufficient to increase isoform D:A ratio in all lines examined, although this was more dramatic in the G/G genotype (NCC-HN1 and NCC-HN43) (FIG. 3B). The same effect of increase in isoform D:A ratio was also seen *in vivo* with LNA-mediated knockdown of EGFR-AS1 in the two patient-derived xenografts (HN124 and HN159; FIG. 8). Next, it was determined whether isoform D expression was necessary for EGFR sensitivity. Using the unique sequence of exon 16B, isoform D specific targeting shRNAs were designed. Stable

transfectants in NCC-HN19, NCC-HN64, NCC-HN1 and NCC-HN43 showed specific targeting of isoform D but not isoform A (FIG. 11).

[0074] In these models, previously gefitinib-sensitive NCC-HN19 and NCC-HN64 (A/A) were rendered more resistant (FIG. 3C), with no significant effect on the G/G genotype lines (NCC-HN1 and NCC-HN43). Similarly, targeting isoform D in the genotype switched G/A^{AAV} clones showed reduced sensitivity, while having no effect on negative controls. Relative to non-targeting controls, reduced expression of isoform D also attenuated the impact of gefitinib on downstream pathway modulation in the HNSCC cell lines with the AA-genotype (FIG. 3D). Importantly, the effect of EGFR-AS1 knock-down promoting gefitinib sensitivity was abrogated by concurrent isoform D knockdown in the cell lines with the G/G genotype (FIG. 3E). In order to determine whether sensitivity to gefitinib was ligand-dependent, cell lines were cultured in normal media with serum, serum-free media and media supplemented with epidermal growth factor (EGF), and treated with gefitinib. IC₅₀ values in cell lines with the A/A genotype and genotype switched G/A^{AAV} clones were significantly lower in serum- and EGF-enriched media, compared to serum-free conditions (FIG. 3F). This data confirms that the effect of the EGFR-AS1 lncRNA is mediated through alteration of the EGFR isoforms A and D, and this effect is likely ligand dependent.

EGFR-AS1 levels and isoform D:A ratio determine gefitinib response

[0075] To evaluate potential biomarkers, semi-quantitative assessment of AS1(1+ or 0) (FIG. 4A) and relative EGFR isoform A and D levels in primary tumour tissue using RNA-in situ hybridization using RNAscope technology and real-time PCR, respectively, was performed. In A/A genotype, AS1 was not detected with increased isoform D/A ratio. Conversely, in either G/A or G/G genotypes, AS1 levels were high (1+), with low isoform D/A ratios. Where available, IC₅₀ values for patient-derived tumour lines were consistent with both genotype statuses, AS1 levels and isoform D/A ratios (FIG. 4B). From this cohort, one patient presented with recurrent, metastatic disease, and hence amenable to a co-clinical trial under IMPACT-SG. This patient (HN137) was initially diagnosed with T4N2BM0 oral squamous cell cancer and underwent surgery followed by adjuvant chemo-radiation therapy. He subsequently developed dermal and lung metastases six months after adjuvant therapy. Sequencing confirmed the A/A genotype for Q787Q, and real-time PCR showed low EGFR-AS1 levels and high EGFR isoform D:A ratio in his original primary and metastatic tumour

compared to controls (FIG. 4B). Both original and metastatic tumours had been engrafted as patient derived xenografts (PDXs), and these were expanded and a clinical trial using oral gefitinib (50 mg/kg by oral gavage) was conducted in NSG mice. Both primary and metastatic patient derived xenograft tumours showed significant tumour regression when
5 treated with gefitinib, compared to untreated controls (FIG. 4C). The patient was concurrently initiated on a 250 mg daily dose of gefitinib; clinical and radiological assessment showed significant regression of dermal and lung metastases after six weeks of gefitinib monotherapy (FIG. 4D).

[0076] Despite the availability of FDA-approved EGFR targeting therapies in squamous
10 cell carcinoma of head and neck and lung origin, no robust predictive biomarkers have been established. Reported here is a mechanism of EGFR addiction mediated by an lncRNA EGFR-AS1. It is also further demonstrated how a synonymous single nucleotide variant can mediate EGFR tyrosine kinase inhibitor sensitivity through modulation of the lncRNA EGFR-AS1, with consequent increase in isoform D:A ratios, resulting in ligand-driven
15 EGFR-dependency. This is supported by three patients with EGFR wild type oral squamous cell cancer, who were exceptional responders to gefitinib, all of whom were A/A genotype. To date, majority of predictive biomarkers in cancer are genomic based and largely confined to non-synonymous mutations that alter protein structure and function. Without being bound by theory, this report describes how a “silent” SNP can impact on lncRNA levels, with
20 dramatic effect on vulnerability to EGFR inhibitors. Further depicted is a possible roadmap for EGFR activation, defining a comprehensive biomarker suite to precisely predict which patients might respond to EGFR tyrosine kinase inhibitors.

[0077] Several recent studies have also identified lncRNAs critical to tumorigenesis, and suggest that these may be targeted with therapeutic intent. Without being bound by theory,
25 the data shown herein points to a critical role to an exon 20 SNP that results in a single nucleotide variant within the lncRNA, that influences the stability of EGFR-AS1 – possibly due to changes in higher order structure. One consequence is a cis-mediated effect on alternative splicing of EGFR, with increased isoform D:A ratio. Certainly, the localization of EGFR-AS1 transcripts in the nucleus lends geographic support to this effect (seen by RNA *in situ* hybridisation; FIG. 4A), where it can affect transcription and/or pre-mRNA processing.

[0078] Manipulating the genotype from AA to GA, or reducing the levels of AS-1 lncRNA was sufficient to increase sensitivity to EGFR tyrosine kinase inhibitors in a

consistent and predictable manner, and this effect could be abrogated by knockdown of EGFR isoform D, which is likely the effector of this phenotype.

[0079] Naturally occurring EGFR isoforms have been poorly studied to date, with a few studies suggesting that alternate isoforms are secreted in plasma (as secreted EGFR or sEGFR) and may be prognostic in lung and cervical cancers. Analysis of sEGFR levels in non-small cell lung cancer patients treated with erlotinib supported the notion that higher levels of sEGFR was predictive of tyrosine kinase inhibitor response, although no distinction was made between the particular isoform measured. The relative increase in isoform D that lacks the tyrosine kinase domain seen here, over full length isoform A, as well as the requirement for ligand-driven EGFR activation, suggests a mechanism involving the extra-cellular domain region, for example, receptor dimerization. While the lack of specific antibodies for each EGFR isoform imposes limitations on spatial localisation and quantification, analysis of mRNA transcripts is a viable alternative as shown here using real-time RT-PCR or RNAscope. Notwithstanding, the precise mechanism through which EGFR-AS1 augments alternative splicing to result in high isoform D:A ratios, remains unclear.

[0080] There are several important implications to the data disclosed herein. Notably, the prevalence of this SNP (homozygous A/A genotype) is 4% in the Asian context (as examined herein) but 30% in Caucasian populations (<http://exac.broadinstitute.org>), highlighting important geographical disparities of genomic biomarkers. Furthermore, the biomarker roadmap for EGFR pathway activation – starting with genotype, levels of EGFR-AS1 lncRNA, relative EGFR isoform expression, and ligand expression – has been established, thereby underscoring the need to establish multi-dimensional biomarker suites for high precision patient enrichment strategies. Also, the ability for “silent” synonymous alterations to influence EGFR addiction opens new avenues to discover novel biomarkers of existing targeted therapies, with enormous potential for immediate re-purposing and clinical impact.

[0081] Finally, the lncRNA EGFR-AS1 has been demonstrated as a bona fide driver of EGFR addiction, and represents a novel therapeutic target in a subset of squamous cell cancers, potentially through RNAi based strategies that have entered clinical testing.

[0082] The invention illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "comprising", "including", "containing", etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed

herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the inventions embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention.

[0083] The invention has been described broadly and generically herein. Each of the narrower species and sub-generic groupings falling within the generic disclosure also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

[0084] Other embodiments are within the following claims and non-limiting examples. In addition, where features or aspects of the invention are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group.

[0085] Throughout this disclosure, certain embodiments may be disclosed in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the disclosed ranges. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

EXPERIMENTAL SECTION

Tumour specimens and primary cell cultures

[0086] Patient-derived tumours were collected and processed as previously described (Leong HS, et al. Stem Cells Translational Medicine 2014; 3:1055-65). All lines were cultured in RPMI media (Sigma Aldrich, St. Louis, MO) supplemented with 10% fetal bovine

serum (FBS) (Hyclone, Thermo Scientific Inc., Fremont, CA) and 1% Penicillin-Streptomycin (Life Technologies, Amsterdam, Netherlands) in a humidified atmosphere of 5% CO₂ at 37 °C. Details of cell line origin and clinico-pathologic characteristics of the patient they were derived from are outlined in FIG. 12. Cell line identity was authenticated by
5 targeted sequencing of the line and primary tumour, to determine identical SNPs, with significant mutations shown in FIG. 13.

Cell proliferation assay and determination of IC₅₀ values

[0087] Cells were seeded in 100 µl complete growth medium at a density of 2,000 - 4,000
10 cells/well in 96-well tissue culture plates. After serial dilutions, 100µl of complete growth medium containing Gefitinib or Erlotinib (BioVision, Milpitas, CA) were added to cells. DMSO was used as controls. Plates were incubated for 48 hours after which cell viability was assessed using CellTitre-Glo® Luminescent Assay according to the manufacturer s protocol (Promega, Madison, WI).

15

Inhibition of RNA replication

[0088] Cells were starved in serum free media for 24 hours. A final concentration of 5 µg/ml of Actinomycin D (Sigma Aldrich, St. Louis, MO), dissolved in dimethyl sulfoxide, was added with fresh serum free media and cells were harvested at 0 hours, 1 hours, 2 hours,
20 and 4 hours intervals.

Mutational Analysis

[0089] Total genomic DNA was extracted from primary cell lines using QIAamp DNA Mini kit according to the manufacturer s instructions (Qiagen, Valencia, CA). 10ng of DNA
25 was used to run the Ion AmpliSeq Colon and Lung Cancer Panel v1 (Life Technologies, Amsterdam, Netherlands) on the Ion Torrent platform (Life Technologies, Amsterdam, Netherlands). The Panel consists of primer pairs targeting hotspots in 22 known genes implicated in colon and lung cancer. The 22 genes are KRAS, EGFR, BRAF, PIK3CA, AKT1, ERBB2, PTEN, NRAS, STK11, MAP2K1, ALK, DDR2, CTNNB1, MET, TP53,
30 SMAD4, FBX7, FGFR3, NOTCH1, ERBB4, FGFR1, FGFR2.

rAAV production

[0090] In primary patient-derived cell lines, targeted knock-in of the Q787Z genotype in for example NCC-HN1 cell line was performed using the Horizon AAV system (Horizon Discovery Group, Cambridge, UK). 1×10^6 HEK293 cells were seeded in 100 mm dishes 24 hours prior to transfection in DMEM (Sigma Aldrich, St. Louis, MO) with 10% fetal bovine serum, 1% Penicillin-Streptomycin in a 5% CO₂ incubator. The pAAV EGFR(G or A) transfer vector (Horizon Discovery Group, Cambridge, UK) was co-transfected with pAAV-RC and pHelper plasmids (10 µg of each) from the AAV Helper-Free System (Stratagene, La Jolla, CA) using Lipofectamine 2000 (Life Technologies, Amsterdam, Netherlands) according to manufacturer's instruction. The virus was harvested 48 hours after transfection in accordance with the manufacturer's protocol. In brief, the media was aspirated from the flask together with HEK293 cells, and subjected to three freeze and thaw cycles. The lysate was clarified by centrifugation at 13,000 rpm at 4°C to remove cell debris, and the supernatant containing rAAV was divided into aliquots and frozen at -80°C for subsequent use.

Viral infection and screening for recombinants

[0091] Cells were grown in 100 mm dishes until 70-80% confluence. After removing existing media, 200 µl of rAAV lysate and 4 ml of Opti-MEM media (Life Technologies, Amsterdam, Netherlands) were added to the cells. The virus was allowed to infect cells at 37°C for 4 hours after which the media was replaced with 5 ml fresh media. Infected cells were allowed to grow for 48 hours and then harvested by trypsinization, and distributed into twelve 96-well plates with media containing Geneticin (Life Technologies, Amsterdam, Netherlands). Plates were incubated at 37°C for 2 to 3 weeks prior to harvest and screening for recombinant events. To verify successful integration of the Horizon vector into the EGFR gene at the expected site, primers flanking part of the Neomycin open reading frame that is present in the vector and Intron 20 of the EGFR were designed as presented in FIG. 6. PCR using genomic DNA was performed as previously described (Zhao Y, et al. Oncogene 2008; 27:1-8).

Lentivirus shRNA knockdown

[0092] In primary patient-derived cell lines, targeted knockdowns of the Q787Z genotype in cell line, for example NCC-HN1, were performed using shRNA lentiviral system (Sigma Aldrich, St. Louis, MO). Cells were seeded with density of 1×10^5 in a 6-well tissue culture plate 24 hours prior infection. The media was removed and the cells were infected with 1 MOI (multiplicity of infection) of pLKO. 1-EGFR Isoform D-puro Lentiviral particles (Sigma Aldrich, St. Louis, MO) and 4ml of Opti-MEM media were added to the cells. Lentival pLKO.1-puro empty vector control transduction particles were used as controls. The virus was allowed to infect cells at 37°C for 4 hours and replaced with 5 ml fresh media. Infected cells were allowed to grow for 48 hours and then selected with Puromycin-containing media (Life Technologies, Amsterdam, Netherlands).

DNA, RNA and Protein analysis

[0093] Genomic DNA extraction and PCR/sequencing were performed as previously described (Zhao Y, et al. Oncogene 2008; 27:1-8). Real-time PCR of genomic DNA and cDNA was performed using iTaq Universal SYBR Green Supermix (Bio-Rad Laboratories, Hercules, CA) and a CFX96 Real-time PCR system (Bio-Rad Laboratories, Hercules, CA) according to the manufacturer s instructions. Reactions were carried out in triplicate with Actin served as the normalizing control. Primer sequences used are presented in Table 1 and Sanger sequencing was used to confirm identity of all transcripts. SDS-PAGE Western blotting was performed as previously described (Leong HS, et al. Stem Cells Translational Medicine 2014; 3: 1055-65 and Zhao Y, et al. Oncogene 2008; 27:1-8). Antibodies used were as follow: rabbit polyclonal antibodies to EGFR, phosphor-EGFR (Tyr 1068), AKT, phosphor-AKT (Ser 473), Erk 1/2, phosphor-Erk 1/2 (Thr 202/204), S6, phosphor-S6 (Ser 235/236) (Cell Signalling Technology, Danvers, MA) and GAPDH (Cell Signalling Technology, Danvers, MA). Total RNA extraction and reverse transcription was performed as previously described (Zhao Y, et al. Oncogene 2008; 27:1-8).

RNA in situ hybridization (ISH) using RNAscope

[0094] *In situ* hybridization for EGFR AS1 was performed using the RNAscope 2.0 FFPE Assay according to the manufacturer s protocols (Advanced Cell Diagnostics, Hayward, DA). 5µm patient-derived formalin fixed paraffin embedded (FFPE) sections were sent to

Singapore Health Services Pte. Ltd. to perform the RNA in situ hybridization. Positive staining was observed as brown, punctate dots. A positive control probe, Homo sapiens peptidylprolyl isomerase B (PPIB), and a negative control probe for the bacterial gene DapB (DAPB) were performed for each tissue sample.

5

Patient-derived xenograft (PDX) and treatment studies

[0095] Primary and metastatic tumour tissue were placed in DMEM/F-12 medium and cut into small pieces (diameter 0.8 mm to 1.5 mm) using dissection scissors. These were mixed with 20% Matrigel (Corning, Tewksbury, MA) and subcutaneously injected into the flanks of
10 male NOD-scid gamma (NSG) mouse (NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ; 005557; Jackson Laboratory, Bar Harbor, ME). When tumour size reached 1.5 cm³, tumours were harvested, processed and re-injected for expansion (passage 1 or P1). This process was repeated to expand the tumour tissue. For the co-clinical trial, the P4 passage was used. Gefitinib (Iressa) (AstraZeneca, London, UK) was prepared by dissolving a 250 mg clinical grade tablet in
15 sterile water containing 0.05% Tween-80 (Sigma Aldrich, St. Louis, MO) to a concentration of 20 mg/ml. When tumours reached 0.5 cm³ in size (24 days for HN137 met and 12 days for HN137 pri), mice were started on treatment with 100mg/kg Gefitinib or controls with the same amount of 0.05% tween-80 in water. Tumour size was measured with a calliper once every 3 days. 12 days post treatment with Gefitinib (n=5 mice), when the tumours size of
20 treated group reached 2.0cm³, the mice were euthanized and analysed.

***In vivo* Locked nucleic acid (LNA) treatment**

[0096] *In vivo* EGFR-AS1 knockdown was performed using custom locked nucleic acid (LNA) oligonucleotide (Exiqon, Vedbaek, Denmark). Treatments with custom LNA
25 oligonucleotides were performed according to the manufacturer's protocol. The custom sequence of the LNA (15-mer RNA/LNA oligonucleotide) against EGFR-AS1 (LNA-AS1) used was ATCGCAAAGGTAATC (SEQ ID NO: 25). Negative control LNA 15-mer oligonucleotide (LNA-i-miR-NC) sequence was AACACGTCTATACGC (SEQ ID NO: 26). Both oligonucleotides contain phosphorothioate backbone modifications and were purified by
30 HPLC followed by Na⁺ salt exchange and lyophilisation for *in vivo* use. Selected patient-derived xenografts were passaged, and when tumour growth was deemed to reach a size of 0.5 cm³, LNA treatment was initiated. LNA treatment was delivered as weekly tail vein

injections of 5 mg/kg, and after the first week of treatment, one mice from the experimental and control arms were sacrificed to confirm knock-down. Treatment was stopped when the tumours in the experimental arm were unmeasurable or when tumours in the control arms exceeded 2.0 cm³.

5

Study Conduct

[0097] The study and protocols for tissue collection for cell line/ patient-derived xenograft propagation have been approved by the Singhealth Centralized Institutional Review Board (CIRB 2007/441/B). The co-clinical trial in a patient with oral squamous cell cancer (OSCC) and the corresponding patient-derived xenograft implanted to a NSG mouse was initiated under the IMPACT-SG (Individualized Molecular Profiling for Allocation of Clinical Trials) (CIRB 2011/441/B).

10

Tables

15 [0098] Table 1: Table of sequences

SEQ ID NO	Target/ description	Primer	Sequence 5'-3'	Amplicon size (bp), if applicable
1	EGFR Exon 18	Forward	CTGAGGTGACCCTTGTCTCTGTGTTCTT	186
2		Reverse	AGAGGCCTGTGCCAGGGACCTTA	
3	EGFR Exon 19	Forward	TCACTGGGCAGCATGTGGCA	241
4		Reverse	CAGCTGCCAGACATGAGAAA	
5	EGFR Exon 20	Forward	CCATGCGAAGCCACACTGA	248
6		Reverse	CGTATCTCCCTTCCCTGATTACC	
7	EGFR Exon 21	Forward	AGCAGGGTCTTCTCTGTTTCA	200
8		Reverse	TGACCTAAAGCCACCTCCTT	
9	EGFR VIII	Forward	GGGCTCTGGAGGAAAAGAAA	952
10		Reverse	ATTCCGTTACACACTTTGCGGC	
11	Neo ORF	Forward	CTGGCTGCTATTGGGCGAAG	2131
12		Reverse	TGCTGGAGTAAAAGGGGCTG	

13	EGFR Copy	Forward	CCAAGGCACGAGTAACAAGCT	119
14		Reverse	GCACATAGGTAATTTCCAAA	
15	Actin (genomic)	Forward	GTCCCCTTCCCTCCTCAGAT	130
16		Reverse	CGGACTCGTCATACTCCTGC	
17	EGFR Isoform A	Forward	ACTCTGAGTGCATACAGTGC	261
18		Reverse	TCGTTGGACAGCCTTCAAGAC	
19	EGFR Isoform D	Forward	ACTCTGAGTGCATACAGTGC	478
20		Reverse	TGAAGGCATGAGGCTCAGTG	
21	Actin (mRNA)	Forward	ATGTTTGAGACCTTCACACC	198
22		Reverse	AGGTAGTCAGTCAGGTCCCGG	
23	EGFR AS1	Forward	TCCAGGTGAAGACGCATGAA	78
24		Reverse	GTCTTTTGCAGGCACAGCTT	
25	LNA against EGFR-AS1 (LNA- AS1)	n.a.	ATCGCAAAGGTAATC	n.a.
26	Negative control (LNA-i- miR-NC)	n.a.	AACACGTCTATACGC	n.a.
27	EGFR as shown in NCBI sequence ID: NM_00522 8.4	n.a.	GTCCGGGCAGCCCCCGGCGCAGCGCGGCC GCAGCAGCCTCCGCCCCCGCACGGTGTG AGCGCCCGACGCGGCCGAGGCGGCCGGA GTCCCGAGCTAGCCCCGGCGGCCCGCCGC GCCCAGACCGGACGACAGGCCACCTCGTC GGCGTCCGCCCCGAGTCCCCGCCTCGCCGC CAACGCCACAACCACCGCGCACGGCCCCC TGACTCCGTCCAGTATTGATCGGGAGAGC CGGAGCGAGCTCTTCGGGGAGCAGCGATG CGACCCTCCGGGACGGCCGGGGCAGCGCT CCTGGCGCTGCTGGCTGCGCTCTGCCCCGC GAGTCGGGCTCTGGAGGAAAAGAAAGTTT GCCAAGGCACGAGTAACAAGCTCACGCAG TTGGGCACTTTTGAAGATCATTTTCTCAGC CTCCAGAGGATGTTCAATAACTGTGAGGT GGTCCTTGGAATTTGGAAATTACCTATGT GCAGAGGAATTATGATCTTTCCTTCTTAA GACCATCCAGGAGGTGGCTGGTTATGTCC TCATTGCCCTCAACACAGTGGAGCGAATT	n.a.

CCTTTGGAAAACCTGCAGATCATCAGAGG
AAATATGTACTACGAAAATTCCTATGC
CTTAGCAGTCTTATCTAACTATGATGCAAA
TAAAACCGGACTGAAGGAGCTGCCCATGA
GAAATTTACAGGAAATCCTGCATGGCGCC
GTGCGGTTTCAGCAACAACCCTGCCCTGTG
CAACGTGGAGAGCATCCAGTGGCGGGACA
TAGTCAGCAGTGACTTTCTCAGCAACATGT
CGATGGACTTCCAGAACCACCTGGGCAGC
TGCCAAAAGTGTGATCCAAGCTGTCCCAA
TGGGAGCTGCTGGGGTGCAGGAGAGGAG
AACTGCCAGAACTGACCAAAATCATCTG
TGCCAGCAGTGCTCCGGGCGCTGCCGTG
GCAAGTCCCCCAGTGACTGCTGCCACAAC
CAGTGTGCTGCAGGCTGCACAGGCCCCCG
GGAGAGCGACTGCCTGGTCTGCCGCAAAT
TCCGAGACGAAGCCACGTGCAAGGACACC
TGCCCCCACTCATGCTCTACAACCCACC
ACGTACCAGATGGATGTGAACCCCGAGGG
CAAATACAGCTTTGGTGCCACCTGCGTGA
AGAAGTGTCCCCGTAATTATGTGGTGACA
GATCACGGCTCGTGCGTCCGAGCCTGTGG
GGCCGACAGCTATGAGATGGAGGAAGAC
GGCGTCCGCAAGTGTAAGAAGTGCGAAGG
GCCTTGCCGCAAAGTGTGTAACGGAATAG
GTATTGGTGAATTTAAAGACTCACTCTCCA
TAAATGCTACGAATATTAAACACTTCAAA
AACTGCACCTCCATCAGTGCGGATCTCCA
CATCCTGCCGGTGGCATTAGGGGTGACT
CCTTACACATACTCCTCCTCTGGATCCAC
AGGAACTGGATATTCTGAAAACCGTAAAG
GAAATCACAGGGTTTTTGCTGATTCAGGCT
TGGCCTGAAAACAGGACGGACCTCCATGC
CTTTGAGAACCTAGAAATCATACGCGGCA
GGACCAAGCAACATGGTCAGTTTTCTCTTG
CAGTCGTCAGCCTGAACATAACATCCTTG
GGATTACGCTCCCTCAAGGAGATAAGTGA
TGGAGATGTGATAATTTACAGGAAACAAAA
ATTTGTGCTATGCAAATACAATAAACTGG
AAAAAACTGTTTGGGACCTCCGGTCAGAA
AACCAAAATTATAAGCAACAGAGGTGAAA
ACAGCTGCAAGGCCACAGGCCAGGTCTGC
CATGCCTTGTGCTCCCCGAGGGCTGCTGG
GGCCCGGAGCCCAGGGACTGCGTCTCTTG
CCGGAATGTCAGCCGAGGCAGGGAATGCG
TGGACAAGTGCAACCTTCTGGAGGGTGAG
CCAAGGGAGTTTGTGGAGAACTCTGAGTG
CATACAGTGCCACCCAGAGTGCCTGCCTC
AGGCCATGAACATCACCTGCACAGGACGG
GGACCAGACAACTGTATCCAGTGTGCCCA
CTACATTGACGGCCCCCACTGCGTCAAGA
CCTGCCCGGCAGGAGTCATGGGAGAAAAC

AACACCCTGGTCTGGAAGTACGCAGACGC
CGGCCATGTGTGCCACCTGTGCCATCCAA
ACTGCACCTACGGATGCACTGGGCCAGGT
CTTGAAGGCTGTCCAACGAATGGGCCTAA
GATCCCGTCCATCGCCACTGGGATGGTGG
GGGCCCTCCTCTTGCTGCTGGTGGTGGCC
TGGGGATCGGCCTCTTCATGCGAAGGCGC
CACATCGTTCGGAAGCGCACGCTGCGGAG
GCTGCTGCAGGAGAGGGAGCTTGTGGAGC
CTCTTACACCCAGTGGAGAAGCTCCCAAC
CAAGCTCTCTTGAGGATCTTGAAGGAAAC
TGAATTCAAAAAGATCAAAGTGCTGGGT
CCGGTGCGTTCGGCACGGTGTATAAGGGA
CTCTGGATCCCAGAAGGTGAGAAAAGTTAA
AATTCCCGTCGCTATCAAGGAATTAAGAG
AAGCAACATCTCCGAAAGCCAACAAGGAA
ATCCTCGATGAAGCCTACGTGATGGCCAG
CGTGGACAACCCCCACGTGTGCCGCCTGC
TGGGCATCTGCCTCACCTCCACCGTGCAGC
TCATCACGCAGCTCATGCCCTTCGGCTGCC
TCCTGGACTATGTCCGGGAACACAAAAGAC
AATATTGGCTCCCAGTACCTGCTCAACTGG
TGTGTGCAGATCGCAAAGGGCATGAACTA
CTTGGAGGACCGTCGCTTGGTGCACCGCG
ACCTGGCAGCCAGGAACGTACTGGTGAAA
ACACCGCAGCATGTCAAGATCACAGATTT
TGGGCTGGCCAAACTGCTGGGTGCGGAAG
AGAAAGAATACCATGCAGAAGGAGGCAA
AGTGCCTATCAAGTGGATGGCATTGGAAT
CAATTTTACACAGAATCTATACCCACCAG
AGTGATGTCTGGAGCTACGGGGTGACTGT
TTGGGAGTTGATGACCTTTGGATCCAAGC
CATATGACGGAATCCCTGCCAGCGAGATC
TCCTCCATCCTGGAGAAAGGAGAACGCCT
CCCTCAGCCACCCATATGTACCATCGATGT
CTACATGATCATGGTCAAGTGCTGGATGA
TAGACGCAGATAGTCGCCCAAAGTTCCGT
GAGTTGATCATCGAATTCTCCAAAATGGC
CCGAGACCCCCAGCGCTACCTTGTCAATTCA
GGGGGATGAAAGAATGCATTTGCCAAGTC
CTACAGACTCCAATTCTACCGTGCCCTGA
TGGATGAAGAAGACATGGACGACGTGGTG
GATGCCGACGAGTACCTCATCCCACAGCA
GGGCTTCTTCAGCAGCCCCTCCACGTACG
GACTCCCCTCCTGAGCTCTCTGAGTGCAAC
CAGCAACAATTCCACCGTGGCTTGCATTG
ATAGAAATGGGCTGCAAAGCTGTCCCATC
AAGGAAGACAGCTTCTTGCAGCGATACAG
CTCAGACCCACAGGCGCCTTGACTGAGG
ACAGCATAGACGACACCTTCCTCCCAGTG
CCTGAATACATAAACCAGTCCGTTCCCAA
AAGGCCCGCTGGCTCTGTGCAGAATCCTG

TCTATCACAATCAGCCTCTGAACCCCGCGC
CCAGCAGAGACCCACACTACCAGGACCCC
CACAGCACTGCAGTGGGCAACCCCGAGTA
TCTCAAACTGTCCAGCCACCTGTGTCAA
CAGCACATTCGACAGCCCTGCCACTGGG
CCCAGAAAGGCAGCCACCAAATTAGCCTG
GACAACCCTGACTACCAGCAGGACTTCTT
TCCCAAGGAAGCCAAGCCAAATGGCATCT
TTAAGGGCTCCACAGCTGAAAATGCAGAA
TACCTAAGGGTTCGCGCCACAAAGCAGTGA
ATTTATTGGAGCATGACCACGGAGGATAG
TATGAGCCCTAAAAATCCAGACTCTTTTCG
ATACCCAGGACCAAGCCACAGCAGGTCCT
CCATCCCAACAGCCATGCCCGCATTAGCT
CTTAGACCCACAGACTGGTTTTGCAACGTT
TACACCGACTAGCCAGGAAGTACTTCCAC
CTCGGGCACATTTTGGGAAGTTGCATTCTT
TTGTCTTCAAACCTGTGAAGCATTTACAGAA
ACGCATCCAGCAAGAATATTGTCCCTTTG
AGCAGAAATTTATCTTTCAAAGAGGTATA
TTTGAIAAAAAAAAAAAGTATATGTGAGG
ATTTTTATTGATTGGGGATCTTGGAGTTTT
TCATTGTCTGCTATTGATTTTTACTTCAATG
GGCTCTTCCAACAAGGAAGAAGCTTGCTG
GTAGCACTTGCTACCCTGAGTTCATCCAGG
CCCAACTGTGAGCAAGGAGCACAAAGCCAC
AAGTCTTCCAGAGGATGCTTGATTCCAGT
GGTTCTGCTTCAAGGCTTCCACTGCAAAAC
ACTAAAGATCCAAGAAGGCCTTCATGGCC
CCAGCAGGCCGGATCGGTACTGTATCAAG
TCATGGCAGGTACAGTAGGATAAGCCACT
CTGTCCCTTCCTGGGCAAAGAAGAAACGG
AGGGGATGGAATTCTTCCTTAGACTTACTT
TTGTAAAAATGTCCCCACGGTACTTACTCC
CCACTGATGGACCAGTGGTTTCCAGTCAT
GAGCGTTAGACTGACTTGTTTGTCTTCCAT
TCCATTGTTTTGAACTCAGTATGCTGCCC
CTGTCTTGCTGTCATGAAATCAGCAAGAG
AGGATGACACATCAAATAATAACTCGGAT
TCCAGCCCACATTGGATTCATCAGCATTTG
GACCAATAGCCACAGCTGAGAATGTGGA
ATACCTAAGGATAGCACCGCTTTTGTCTC
GCAAAACGTATCTCCTAATTTGAGGCTC
AGATGAAATGCATCAGGTCCTTTGGGGCA
TAGATCAGAAGACTACAAAAATGAAGCTG
CTCTGAAATCTCCTTTAGCCATCACCCCAA
CCCCCAAAATTAGTTTGTGTTACTTATGG
AAGATAGTTTTCTCCTTTTACTTCACTTCA
AAAGCTTTTTACTCAAAGAGTATATGTTCC
CTCCAGGTCAGCTGCCCCCAAACCCCTCC
TTACGCTTTGTACACAAAAAGTGTCTCTG
CCTTGAGTCATCTATTCAAGCACTTACAGC

TCTGGCCACAACAGGGCATTTTACAGGTG
CGAATGACAGTAGCATTATGAGTAGTGTG
GAATTCAGGTAGTAAATATGAACTAGGG
TTTGAAATTGATAATGCTTTCACAACATTT
GCAGATGTTTTAGAAGGAAAAAAGTTCCT
TCCTAAAATAATTTCTCTACAATTGGAAGA
TTGGAAGATTCAGCTAGTTAGGAGCCAC
CTTTTTTCCTAATCTGTGTGTGCCCTGTAA
CCTGACTGGTTAACAGCAGTCCTTTGTAAA
CAGTGTTTTAACTCTCCTAGTCAATATCC
ACCCCATCCAATTTATCAAGGAAGAAATG
GTTCAGAAAAATATTTTCAGCCTACAGTTAT
GTTCAGTCACACACACATACAAAATGTTC
CTTTTGCTTTTAAAGTAATTTTGTACTCCC
AGATCAGTCAGAGCCCCTACAGCATTGTT
AAGAAAGTATTTGATTTTTGTCTCAATGAA
AATAAACTATATTCATTTCCACTCTATTA
TGCTCTCAAATACCCCTAAGCATCTATACT
AGCCTGGTATGGGTATGAAAGATACAAAG
ATAAATAAAACATAGTCCCTGATTCTAAG
AAATTCACAATTTAGCAAAGGAAATGGAC
TCATAGATGCTAACCTTAAACAACGTGA
CAAATGCCAGACAGGACCCATCAGCCAGG
CACTGTGAGAGCACAGAGCAGGGAGGTTG
GGTCCTGCCTGAGGAGACCTGGAAGGGAG
GCCTCACAGGAGGATGACCAGGTCTCAGT
CAGCGGGGAGGTGGAAAGTGCAGGTGCAT
CAGGGGCACCCTGACCGAGGAAACAGCTG
CCAGAGGCCTCCACTGCTAAAGTCCACAT
AAGGCTGAGGTCAGTCACCCTAAACAACC
TGCTCCCTCTAAGCCAGGGGATGAGCTTG
GAGCATCCCACAAGTTCCTAAAAGTTGC
AGCCCCCAGGGGGATTTTGAGCTATCATC
TCTGCACATGCTTAGTGAGAAGACTACAC
AACATTTCTAAGAATCTGAGATTTTATATT
GTCAGTTAACCCTTTTATTATTTCATTAC
CTCAGGACATGCAGAAATATTTTCAGTCAG
AACTGGGAAACAGAAGGACCTACATTCTG
CTGTCACTTATGTGTCAAGAAGCAGATGA
TCGATGAGGCAGGTCAGTTGTAAGTGAGT
CACATTGTAGCATTAAATTCTAGTATTTTT
GTAGTTTGAAACAGTAACTTAATAAAAAGA
GCAAAAGCTAAAAAAAAAAAAAAAAAAAA

Claims

1. A method of predicting susceptibility of a subject suffering from cancer to a treatment with an anti-cancer drug, wherein the method comprises detecting the presence or absence of a genetic alteration in a long non-coding RNA (lncRNA) that resides in an antisense strand of an oncogene, wherein the genetic alteration alters or disrupts expression of the oncogene; wherein in case the genetic alteration is present, the subject is predicted to show improved susceptibility to the treatment with the anti-cancer drug compared to a subject not carrying the genetic alteration.
2. The method of claim 1, wherein the genetic alteration is homozygous.
3. The method of claim 1, wherein the oncogene is selected from the group consisting of EGFR encoding epidermal growth factor receptor, IGF1R encoding insulin growth factor 1 receptor, mTOR encoding mechanistic target of rapamycin, PIK3CB encoding phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit beta isoform, PIK3CD encoding phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform, PIK3R3 encoding phosphatidylinositol 3-kinase regulatory subunit gamma and ERBB4 encoding receptor tyrosine-protein kinase erbB-4.
4. The method of any one of the preceding claims, wherein the antisense strand of the non-coding RNA of the oncogene is selected from the group consisting of EGFR-AS1 for EGFR gene; IRAIN for IGF1R gene, MTOR-AS1 for mTOR gene; GAPDHP39 and RPL23AP40 for PIK3CB gene; LOC101929626 for PIK3R3 gene; PIK3CD-AS1, PIK3CD-AS2 and RPL26P7 for PIK3CD gene and RNA5SP119 for ERBB4 gene.
5. The method of claim 3 to 4, wherein the cancer related to EGFR is head and neck cancer or lung cancer.
6. The method of claim 5, wherein the head and neck cancer is head and neck squamous cell carcinoma (HNSCC) or oral squamous cell carcinoma (OSCC).
7. The method of claim 5, wherein the lung cancer is non-small cell lung cancer (NSCLC).

8. The method of any one of claims 3 to 4, wherein the cancer related to PIK3CD is haematological malignancy.
- 5 9. The method of any one of claims 3 to 4, wherein the cancer related to ERBB4 is breast cancer.
10. The method of claim 1, wherein the genetic alteration is a silent G>A mutation (genetic alteration) at Q787Q position in exon 20 of the oncogene EGFR.
- 10 11. The method of claim 10, wherein the genetic alteration results in a homozygous mutation (AA).
12. The method of claim 10, further comprising measuring of either one or both of the
15 following for predicting the susceptibility: i) whether the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to a subject that does not respond to treatment with the anti-cancer drug; ii) whether the subject has higher EGFR isoform D/isoform A ratio compared to a subject that does not respond to treatment with the anti-cancer drug, wherein the anti-cancer drug is a tyrosine kinase
20 inhibitor or an EGFR inhibitor.
13. The method of claim 10 or 12, wherein the anti-cancer drug, is selected from the group consisting of gefitinib, erlotinib and afatinib.
- 25 14. A method of predicting the susceptibility of a subject suffering from cancer related to EGFR to a treatment with an EGFR inhibitor, the method comprising determining whether either one or two or all of the following is given:
- 30 i) the subject has a silent G>A mutation (genetic alteration) at Q787Q position in exon 20 of EGFR (nucleotide 2361);
- ii) the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to subject that does not respond to treatment with an EGFR inhibitor;
- iii) the subject has higher EGFR isoform D/isoform A ratio compared to subject that does not respond to treatment with an EGFR inhibitor;

wherein in case one or two or all of i) to iii) is given the subject is predicted to show improved susceptibility to a treatment with an EGFR inhibitor compared to a subject not showing any of i) to iii).

- 5 15. A method of treating cancer in a subject suffering from said cancer, wherein the method comprises:
- a) detecting the presence or absence of a mutation in a long non-coding RNA on an antisense strand of an oncogene related to the cancer, wherein the genetic alteration alters or disrupts expression of the oncogene;
- 10 b) administering an anti-cancer drug for the cancer type the subject suffers from in case the presence of the genetic alteration is confirmed.
16. The method of claim 15, wherein the oncogene is selected from the group consisting of EGFR encoding epidermal growth factor receptor, IGF1R encoding insulin growth factor 1 receptor, mTOR encoding mechanistic target of rapamycin, PIK3CB encoding phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit beta isoform, PIK3CD encoding phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform, PIK3R3 encoding phosphatidylinositol 3-kinase regulatory subunit gamma and ERBB4 encoding receptor tyrosine-protein kinase erbB-4.
- 15 20
17. The method of claim 15 or 16, wherein the non-coding RNA on an antisense strand of an oncogene is selected from the group consisting of EGFR-AS1 for EGFR gene; IRAIN for IGF1R gene, MTOR-AS1 for MTOR gene; GAPDHP39 and RPL23AP40 for PIK3CB gene; LOC101929626 for PIK3R3 gene; PIK3CD-AS1, PIK3CD-AS2 and RPL26P7 for PIK3CD gene and RNA5SP119 for ERBB4 gene.
- 25
18. The method of claim 16 to 17, wherein the cancer related to EGFR is head and neck cancer or lung cancer.
- 30 19. The method of claim 18, wherein the head and neck cancer is head and neck squamous cell carcinoma (HNSCC) or oral squamous cell carcinoma (OSCC).

20. The method of claim 18, wherein the lung cancer is non-small cell lung cancer (NSCLC).
21. The method of any one of claims 15 to 17, wherein the cancer related to PIK3CD is haematological malignancy.
22. The method of any one of claims 15 to 17, wherein the cancer related to ERBB4 is breast cancer.
23. The method of any one of claims 15 to 17, wherein the cancer is an EGFR related cancer and the mutation (genetic alteration) is a silent G>A mutation at Q787Q position in exon 20 of EGFR.
24. The method of claim 15 or 23, wherein the method further comprises determining if the subject has lower EGFR-AS1 or EGFR-AS1 lncRNA expression level compared to subject that does not respond to treatment with a tyrosine kinase inhibitor.
25. The method of claim 15 or 23 or 24, wherein the method further comprises determining if the subject has higher EGFR isoform D/isoform A ratio compared to subject that does not respond to treatment with an EGFR inhibitor.
26. The method of any of the preceding claims, wherein the anti-cancer drug is selected from the group consisting of gefitinib, erlotinib and afatinib for the treatment of cancer related to EGFR; OSI-902(Linsitinib) for the treatment of cancer related to IGF1R; everolimus (RAD001) and sirolimus for the treatment of cancer related to mTOR; BKM120(Buparlisib) and BYL719(Alpelisib) for the treatment of cancer related to PIK3CB and PIK3R3; Idelalisib for the treatment of cancer related to PIK3CD and dacomatinib and lapatinib for the treatment of cancer related to ERBB4.
27. The method any of one of claims 15 to 26, wherein the oncogene related to cancer is EGFR; wherein the mutation in a) of claim 15 is a silent G>A mutation at Q787Q position in exon 20 of EGFR; and wherein the anti-cancer drug is gefitinib.

28. A method of treating a subject suffering from cancer, comprising:
- a) administering to the subject an effective amount of a therapeutic agent affecting expression of a non-coding RNA an oncogene, wherein the genetic alteration alters or disrupts expression of the oncogene; and
 - 5 b) administering to the subject an effective amount of an anti-cancer drug specific for the cancer related to the oncogene.
29. The method of claim 28, wherein a) and b) are carried out simultaneously or b) is carried out after a).
- 10 30. The method of claim 29, wherein when b) is carried out after a), the time interval between a) and b) is at least 1 to 24 hours, or at least 1 or 2 or 3 or 4 or 5 or 6 or 7 days, or at least one week.
- 15 31. The method of claim 28, wherein the therapeutic agent is selected from the group consisting of miRNAs (microRNAs), siRNA (small interfering RNA), locked nucleic acids (LNAs) and shRNA (short hairpin RNA).
32. A method of treating a subject suffering from an EGFR-related cancer, comprising:
- 20 a) administering to the subject an effective amount of a therapeutic agent affecting EGFR-AS1 lncRNA expression, or affecting EGFR-AS1 expression, or increasing the amount of EGFR isoform D and/or decreasing EGFR isoform A or a combination thereof; and
 - b) administering to the subject an effective amount of a tyrosine kinase inhibitor used to
 - 25 treat the EGFR-related cancer.
33. The method of claim 32, wherein a) and b) are carried out simultaneously or b) is carried out after a).
- 30 34. The method of claim 33, wherein when b) is carried out after a), the time interval between a) and b) is at least 1 to 24 hours, or at least 1 or 2 or 3 or 4 or 5 or 6 or 7 days, or at least one week.

35. The method of any one of claims 32 to 34, wherein the therapeutic agent affecting EGFR-AS1 expression comprises miRNA, shRNA, LNAs, siRNA or combinations thereof.
- 5 36. The method of any one of claims 32 to 34, wherein the therapeutic agent suppressing EGFR-AS1 comprises EGFR-AS1 targeting LNA.
- 10 37. The method of any one of claims 32 to 34, wherein the therapeutic agent increasing EGFR isoform D or decreasing EGFR isoform A comprises an agent modulating the alternative splicing of EGFR.
38. The method of any one of claims 32 to 37, wherein the tyrosine kinase inhibitor is an EGFR inhibitor.
- 15 39. The method of claim 38, wherein the tyrosine kinase inhibitor is selected from the group consisting of gefitinib, erlotinib, erlotinib HCl, lapatinib, dacomitinib, TAE684, afatinib, dasatinib, saracatinib, veratinib, AEE788, WZ4002, icotinib, osimertinib, BI1482694, ASP8273, EGF816, AZD3759, cetuximab, necitumumab, 20 pannitumumab, nimotuzumab and combinations thereof.
40. The method of claim 38, wherein the tyrosine kinase inhibitor is selected from the group consisting of gefitinib, erlotinib and lapatinib.