X is phen-1,4-ylene or pyridin-2,5-ylene, which are unsubstituted or carry 1, 2 or 3 radicals selected independently from each other from hydroxy, halogen, C_1-C_6-alkyl, C_1-C_6-haloalkyl, C_1-C_6-alkoxy, C_1-C_6-haloalkoxy and C_3-C_8-cycloalkyl;

L^3 is selected from -CH=CH-, -C≡C-, -CR^bR^c-CH(OH)-, -CR^bR^c-C(O)-, -CR^bR^c-O-, -CR^bR^c-NR^d-, -CR^bR^c-$S(O)_m$-, -CH(OH)-, -C(O)-, -C(O)-NR^d-, -O-, -NR^d-, -NR^d-C(O)-, -NR^dC(O)-O-, -NR^d-C(O)-NR^e-, -NR^d-$S(O)_n$-, -$S(O)_p$- and -$S(O)_q$-NR^d-, wherein m, n and p are 0, 1 or 2 and q is 1, or 2, and wherein

R^b and R^c are independently from each other selected from H, C_1-C_6-alkyl, C_3-C_8-cycloalkyl and wherein two radicals R^b and R^c bound to the same carbon atom together with said carbon atom may form a 3- to 8-membered ring, wherein said ring may contain 1 or 2 heteroatoms selected from O, N and S as ring member and wherein the ring members of said ring may optionally be independently substituted by hydroxy, halogen, C_1-C_6-alkyl, C_1-C_6-haloalkyl, C_1-C_6-alkoxy, C_1-C_6-haloalkoxy and C_3-C_8-cycloalkyl, and wherein

R^d and R^e independently from each other are H or C_1-C_6-alkyl;

Y is selected from ~~H, C_1-C_6-alkyl,~~ C_3-C_8-cycloalkyl, C_3-C_8-cycloalkyl-C_1-C_6-alkyl, C_3-C_8-cycloalkyl-C_2-C_6-alkenyl, phenyl, phenyl-C_1-C_6-alkyl, phenyl-C_2-C_6-alkenyl, naphthyl, naphthyl-C_1-C_6-alkyl, naphthyl-C_2-C_6-alkenyl, heterocyclyl, heterocyclyl-C_1-C_6-alkyl and heterocyclyl-C_2-C_6-alkenyl, wherein the heterocyclyl moieties are selected from 5- to 7-membered heterocyclic radicals and 5- to10-membered, bicyclic heterocyclic radicals, containing one, two or three heteroatoms, selected from O, N and S as ring members, and wherein

the C_1-C_6-alkyl and C_2-C_6-alkenyl moieties in the aforementioned radicals Y are unsubstituted or carry at least one substituent selected from hydroxy,