

DESCRIPTION

UREA DERIVATIVE OR PHARMACOLOGICALLY ACCEPTABLE SALT THEREOF

Technical Field

[0001]

5 The present invention relates to a urea derivative or a pharmacologically acceptable salt thereof that are useful as pharmaceuticals and have a formyl peptide receptor like 1 (hereinafter may be abbreviated as FPRL1) agonist effect, to a pharmaceutical composition containing the urea derivative or the
10 pharmacologically acceptable salt thereof, and to a pharmaceutical use thereof.

Background Art

[0002]

 FPRL1 (formyl peptide receptor like 1, also known as Lipoxin
15 A4 Receptor, ALXR, and FPR2) is a G protein-coupled receptor cloned as a subtype of N-formyl peptide receptors (FPRs) by Murphy et al. (Non-Patent Literature 1). The FPRL1 was discovered as a receptor that mediates calcium mobilization in response to high concentration of fMLF (formyl methionine leucyl phenylalanine peptide).

20 [0003]

 Expression of FPRL1 has been found in neutrophils, monocytes, T-lymphocytes, dendritic cells, etc. (Non-Patent Literature 2), but the role of FPRL1 in a living body is complicated and has therefore not been elucidated sufficiently (Non-Patent Literature 3).
25 However, in a paw edema model and an arthritis model using FPRL1

deficient mice, it has been recognized that the reactions become worse (Non-Patent Literature 4). Therefore, it is considered that FPRL1 contributes to the resolution of the inflammation.

[0004]

5 Endogenous lipid mediators such as Lipoxin A4 (LXA4) and Resolvin D1 (RvD1) and peptides such as WKYMVm have been reported as agonists that bind to FPRL1 (Non-Patent Literatures 5 and 6).

[0005]

Such FPRL1 agonists can reduce neutrophil chemotaxis in vitro
10 (Non-Patent Literatures 7 and 8). Although neutrophils perform host defense, they cause vascular injury, result in an increase in vascular permeability and edema, followed by release of chemotactic factors, and thereby contribute to inflammation (Non-Patent Literature 9). Therefore, it is considered that the FPRL1 agonists exhibit an
15 anti-inflammatory effect.

For example, it has been confirmed that peptide agonists exhibit an inhibitory effect on intestinal inflammation (Non-Patent Literature 10), an inhibitory effect on airway inflammation (Non-Patent Literature 11), an inhibitory effect on septicemia
20 (Non-Patent Literature 12), and an inhibitory effect on a cancer model (Non-Patent Literature 13). It has also been recognized that QuinC1, a non-peptide low-molecular weight compound, inhibits bleomycin-induced lung inflammation (Non-Patent Literature 14).

[0006]

25 Therefore, FPRL1 can be considered as a target of various

diseases such as inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders. Therefore, the

5 FPRL1 agonists can be promising therapeutic agent for these diseases.

[0007]

Known examples of the non-peptide low-molecular weight compound exhibiting FPRL1 agonist activity include quinazolinones (Non-Patent Literature 15), pyrazolones (Non-Patent Literature 16),
10 benzimidazoles (Non-Patent Literature 17), aminoazoles (Patent Literatures 1, 2, 3, 4, and 5), spiro[2,4]heptanes (Patent Literature 6), pyridazinones (Non-Patent Literature 18), cycloalkyl and cycloalkenyl-1,2-dicarboxylic acids (Patent Literature 7), dihydronaphthalenes (Patent Literature 8), pyrrolidine-2,5-diones
15 (Patent Literature 9), and phenyl urea derivatives (Patent Literatures 10, 11, 12, and 13) (Non-Patent Literatures 19 and 20).

[0008]

However, the basic chemical structures of these compounds are different from those of the compounds of the present invention.

20 It will be appreciated that the above compounds are not included in the claims of the present application.

Citation List

Non-Patent Literature

[0009]

25 Non-Patent Literature 1: Murphy P. M., et al., "The Journal

of Biological Chemistry," 1992, vol. 267, pp. 7637-7643

Non-Patent Literature 2: Gavins F. N. E, et al., "Trends in Pharmacological Sciences," 2010, vol. 31, pp. 266-276

Non-Patent Literature 3: Cattaneo F., et al., "International
5 Journal of Molecular Sciences," 2013, vol. 14, No. 4, pp. 7193-7230

Non-Patent Literature 4: Dufton N, et al., "The Journal of Immunology," 2010, vol. 184, pp. 2611-2619

Non-Patent Literature 4: Le Y, et al., "Trends in immunology," 2002, vol. 23, No. 11, pp. 541-548

10 Non-Patent Literature 5: Krishnamoorthy S, "Proceedings of the National Academy of Sciences," 2010, vol. 107, No. 4, pp. 1660-1665

Non-Patent Literature 7: Li B. Q, et al., "Blood," 2001, vol. 97, pp. 2941-2947

Non-Patent Literature 6: Sogawa Y, et al., "Immunology," 2011,
15 1416974161282, pp. 441-450

Non-Patent Literature 7: Summers C, et al., "Trends in Immunology," 2010, vol. 31, pp. 318-324

Non-Patent Literature 8: Kim S. D, et al., "Experimental & Molecular Medicine," 2013, vol. 13, No. 45: e40.

20 Non-Patent Literature 9: Tae Y. M, et al., "The Journal of Immunology," 2012, vol. 188, pp. 1799-808

Non-Patent Literature 10: Kim S. D, et al., "The Journal of Immunology," 2010, vol. 185, pp. 4302-4310

Non-Patent Literature 13: Kim S. D, et al., "PLOS ONE," vol.
25 7, No. 1: e30522.

Non-Patent Literature 11: Min H. E, et al., "Acta Pharmacologica Sinica" 2011, vol. 32, pp. 601-610

Non-Patent Literature 12: Nanamori M, et al., "Molecular Pharmacology," 2004, vol. 66, pp. 1213-1222

5 Non-Patent Literature 13: Burli R. W, et al., "Bioorganic & Medicinal Chemistry Letters," 2006, vol. 16, pp. 3713-3718

Non-Patent Literature 14: Frohn M, et al., "Bioorganic & Medicinal Chemistry Letters," 2007, vol. 17, pp. 6633-6637

10 Non-Patent Literature 15: Cilibrizzi A, et al., "Journal of Medicinal Chemistry," 2009, vol. 52, pp. 5044-5057

Non-Patent Literature 16: Kirpotina L. N, et al., "Molecular Pharmacology," 2010, vol. 77, pp. 159-170

Non-Patent Literature 17: Schepetkin I. A, et al., "Molecular Pharmacology," 2011, vol. 79, pp. 77-90

15 [Patent Literature]

[0010]

Patent Literature 1: W02009/077990

Patent Literature 2: W02009/077954

Patent Literature 3: W02010/143158

20 Patent Literature 4: W02012/077049

Patent Literature 5: W02012/077051

Patent Literature 6: W02012/066488

Patent Literature 7: W02011/163502

Patent Literature 8: W02012/125305

25 Patent Literature 9: US130018067

Patent Literature 10: W02005/047899

Patent Literature 11: W02012/074785

Patent Literature 12: W02012/109544

Patent Literature 13: W02013/062947

5 Summary of Invention

Technical Problem

[0011]

At present, no compound has been found which has a superior
FPRL1 agonist effect as a prophylactic or therapeutic agent for
10 various disease states described above and can be used as a
sufficiently satisfactory pharmaceutical.

[0012]

It is an object of the present invention to provide a compound
having an FPRL1 agonist effect.

15 Solution to Problem

[0013]

The present inventors have conducted extensive studies and
found that a urea compound represented by the general formula (I)
below (this compound may be referred to as a compound (I)) or a
20 pharmacologically acceptable salt thereof has a superior FPRL1
agonist effect and is sufficiently satisfactory as a pharmaceutical,
and thus the present invention has been completed.


[0014]

Accordingly, the present invention is as follows.

25 [0015]

[1] A compound represented by the general formula (I) or a pharmacologically acceptable salt thereof:

[Chemical Formula 1]



(I)

5 [wherein, in the formula (I), Ar¹ is a phenyl group optionally having
substituent(s), a 5-membered aromatic heterocyclic group optionally
having substituent(s), a 6-membered aromatic heterocyclic group
optionally having substituent(s), or a bicyclic aromatic
heterocyclic group having 8 or 9 atoms and optionally having
10 substituent(s);

Ar² is a phenyl group optionally having substituent(s) (except
for a phenyl group substituted only with halogen atom(s)), a
5-membered aromatic heterocyclic group optionally having
substituent(s), a 6-membered aromatic heterocyclic group optionally
15 having substituent(s), or a bicyclic aromatic heterocyclic group
having 8 or 9 atoms and optionally having substituent(s);

X is a group selected from the group consisting of the following
a), b), and c),

- 20 a) an oxygen atom or a sulfur atom,
b) NR⁴, and
c) NOR⁴, wherein,

when X is b) or c), R^4 is a hydrogen atom, a phenyl group optionally having substituent(s), a heterocyclic group optionally having substituent(s), or a C_1 to C_6 alkyl group optionally having substituent(s);

5 R^1 is a hydrogen atom, a hydroxy group, a C_1 to C_6 alkoxy group optionally having substituent(s), or a C_1 to C_6 alkyl group optionally having substituent(s);


R^2 and R^3 are each independently a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s) or together form
10 a C_2 to C_6 alkylene group; and

each carbon atom marked with an asterisk is an asymmetric carbon atom].

[2] The compound according to [1] or a pharmacologically acceptable salt thereof, wherein

15 in the formula (I), Ar^2 is a group selected from the group consisting of the following A1), A2), A3), A4), A5), A6), A7), A8), A9), and A10):

[Chemical Formula 2]



wherein, when Ar^2 is A2), W^1 is a nitrogen atom or CH optionally
20 substituted with a hydrogen atom, with a halogen atom, or with a

C₁ to C₆ alkyl group optionally having substituent(s);

when Ar² is A2), W² is CH or a nitrogen atom;

when Ar² is A3), A4), A5), or A6), W³ is an oxygen atom, a sulfur atom, or NH optionally substituted with a C₁ to C₆ alkyl group;

5 when Ar² is A3), A4), or A6), W⁴ is CH or a nitrogen atom;

when Ar² is A7), W⁵ is CH₂, an oxygen atom, or a sulfur atom;

when Ar² is A7), W⁶ is C=O, CH₂, CF₂, CHOH, NH optionally substituted with a C₁ to C₆ alkyl group, SO, SO₂, an oxygen atom, or a sulfur atom;

10 when Ar² is A8), W⁷ is NH optionally substituted with a C₁ to C₆ alkyl group or C=O;

when Ar² is A8), W⁸ is C=O with W⁷ being NH optionally substituted with a C₁ to C₆ alkyl group and W⁸ is NH optionally substituted with a C₁ to C₆ alkyl group with W⁷ being C=O;

15 when Ar² is A10), W⁹ is a nitrogen atom or N=O;

when Ar² is A1), A2), A3), A4), or A5), R⁶ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally
20 having substituent(s), a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl group optionally having substituent(s), a heterocyclic group optionally having substituent(s), -CONR¹⁰R¹¹, or -NR¹⁰R¹¹, wherein, when R⁶ is -CONR¹⁰R¹¹
25 or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group optionally

having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), or a C₁ to C₆ alkylsulfonyl group optionally having substituent(s) and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group optionally having substituent(s), or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group;

when Ar² is A1), A2), A3), A4), A5), A6), A7), A8), A9), or A10), R⁷ is a hydrogen atom, a halogen atom, a C₁ to C₆ alkyl group optionally having substituent(s), or a C₁ to C₆ alkoxy group optionally having substituent(s);

when Ar² is A1), A7), A8), or A10), R⁸ is a hydrogen atom, a halogen atom, or a C₁ to C₆ alkyl group optionally having substituent(s);

when Ar² is A9), R⁹ is a hydrogen atom or a C₁ to C₆ alkyl group;

when Ar² is A7), m is 0 or 1; and


when Ar² is A8), n is 0 or 1;

provided that, when Ar² is A1), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

[3] The compound according to [2] or a pharmacologically acceptable salt thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of the following B1), B2), B3), B4), B5), B6), B7), B8), B9), B10), B11), and B12):

[Chemical Formula 3]



wherein, when Ar¹ is B2), B3), B7), B8), B10), B11), or B12), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₃ to C₆ cycloalkyl group optionally having substituent(s), a C₃ to C₆ cycloalkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), a C₂ to C₆ alkenyl group optionally having substituent(s), a C₂ to C₆ alkynyl group optionally having substituent(s), a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl group optionally having substituent(s), -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), or a C₁ to C₆ alkylsulfonyl group optionally having substituent(s).

and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s), or R^{10} and R^{11} together form a C_3 to C_{10} heterocycloalkyl group;

when Ar^1 is B1), R^{12} is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 alkoxy group optionally having substituent(s), a C_3 to C_6 cycloalkyl group optionally having substituent(s), a C_3 to C_6 cycloalkoxy group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), a C_2 to C_6 alkenyl group optionally having substituent(s), a C_2 to C_6 alkynyl group optionally having substituent(s), a C_1 to C_6 alkoxycarbonyl group, a C_1 to C_6 alkylsulfanyl group optionally having substituent(s), a C_1 to C_6 alkylsulfinyl group optionally having substituent(s), a C_1 to C_6 alkylsulfonyl group optionally having substituent(s), $-CONR^{10}R^{11}$, $-NR^{10}R^{11}$, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R^{12} is $-CONR^{10}R^{11}$ or $-NR^{10}R^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), or a C_1 to C_6 alkylsulfonyl group optionally having substituent(s) and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s), or R^{10} and R^{11} together form a C_3 to C_{10} heterocycloalkyl group, and R^{13} is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C_1 to C_6 alkyl group, or R^{12} and R^{13} may together form a C_3 to C_5 alkylene group or a C_1 to C_2 alkylenedioxy group;

when Ar¹ is B6), R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group;

when Ar¹ is B9), R¹⁴ and R¹⁵ are each independently a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group, or a
5 C₁ to C₆ alkoxy group;

when Ar¹ is B2), one of W¹⁰ and W¹¹ is a nitrogen atom, and the other thereof is CH or a nitrogen atom;

when Ar¹ is B6), B7), B8), B9), B10), B11), or B12), W¹² is an oxygen atom, a sulfur atom, or N-R¹⁶, wherein, when W¹² is an N-R¹⁶,
10 R¹⁶ is a hydrogen atom or a C₁ to C₆ alkyl group; and

when Ar¹ is B6), B7), B8), or B9), W¹³ is CH or a nitrogen atom.
[4] The compound according to [3] or a pharmacologically acceptable salt thereof, wherein

when Ar² is A1), A2), A3), A4), or A5), R⁶ is a hydrogen atom,
15 a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ acyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, a heterocyclic group optionally having substituent(s), -CONR¹⁰R¹¹, or -NR¹⁰R¹¹, wherein, when R⁶ is
20 -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, a C₁ to C₆ acyl group, or a C₁ to C₆ alkylsulfonyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group;

when Ar² is A1), A2), A3), A4), A5), A6), A7), A8), A9), or
25 A10), R⁷ is a hydrogen atom, a halogen atom, a C₁ to C₆ alkyl group,

or a C₁ to C₆ alkoxy group;

when Ar² is A1), A7), A8), or A10), R⁸ is a hydrogen atom, a halogen atom, or a C₁ to C₆ alkyl group; and

when Ar² is A9), R⁹ is a hydrogen atom or a C₁ to C₆ alkyl group;

5 provided that, when Ar² is A1), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

[5] The compound according to [4] or a pharmacologically acceptable salt thereof, wherein

10 R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a carboxy C₁ to C₆ alkyl group, a carbamoyl C₁ to C₆ alkyl group, a mono C₁ to C₆ alkylcarbamoyl C₁ to C₆ alkyl group, a di-C₁ to C₆ alkylcarbamoyl C₁ to C₆ alkyl group, an aminosulfonyl
15 C₁ to C₆ alkyl group, an aromatic heterocyclic C₁ to C₃ alkyl group optionally having substituent(s), or a phenyl C₁ to C₃ alkyl group optionally having substituent(s);

R² and R³ are each independently a hydrogen atom or a C₁ to C₃ alkyl group or together form a C₂ to C₆ alkylene group;

20 when Ar¹ is B2), B3), B7), B8), B10), B11), or B12), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to
25 C₆ alkenyl group, a C₂ to C₆ alkynyl group, a C₁ to C₆ alkoxycarbonyl

group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, a C₁ to C₆ acyl group, or a C₁ to C₆ alkylsulfonyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group;

when Ar¹ is B1), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to C₆ alkenyl group, a C₂ to C₆ alkynyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, a C₁ to C₆ acyl group, or a C₁ to C₆ alkylsulfonyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group, and R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group, or R¹² and R¹³ may together form a C₃ to C₅ alkylene group or a C₁ to C₂ alkylenedioxy group; and

X is a group selected from the group consisting of the following

a), b), and c),

a) an oxygen atom,

b) NR^4 , and


c) NOR^4 , wherein,

when X is b) or c), R^4 is a hydrogen atom, a hydroxy C_1 to C_6 alkyl group, a phenyl group, a heterocyclic group, or a C_1 to C_6 alkyl group.

[6] The compound according to [5] or a pharmacologically acceptable salt thereof, wherein

in the formula (I), Ar^2 is a group selected from the group consisting of A1), A2a), A3), and A7a):

[Chemical Formula 4]



wherein, when Ar^2 is A2a), W^2 is the same as that defined in [2] when Ar^2 is A2);

when Ar^2 is A3), W^4 is the same as that defined in [2] when Ar^2 is A3);

when Ar^2 is A7a), W^6 is the same as that defined in [2] when Ar^2 is A7);

when Ar^2 is A1), A2a), A3), or A7a), R^7 is the same as that defined in [4] when Ar^2 is A1), A2), A3), or A7);

when Ar^2 is A1) or A7a), R^8 is the same as that defined in [4] when Ar^2 is A1) or A7);

when Ar^2 is A7a), m is the same as that defined in [2] when Ar^2 is A7);


when Ar² is A1), A2a), or A3), R⁶ is a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ acyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, -CONR¹⁰R¹¹, or
 5 -NR¹⁰R¹¹, wherein, when R⁶ is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group; and

10 when Ar² is A3), W³ is an oxygen atom or a sulfur atom;
 provided that, when Ar² is A1), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

[7] The compound according to [6] or a pharmacologically acceptable
 15 salt thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B2), B6a), B9a), and B9b):

[Chemical Formula 5]



wherein, when Ar¹ is B6a), R¹³ is the same as that defined in
 20 [3] when Ar¹ is B6);

when Ar¹ is B9a) or B9b), R¹⁴ and R¹⁵ are the same as those defined in [3] when Ar¹ is B9);

when Ar¹ is B2), W¹⁰ and W¹¹ are the same as those defined in

[3] when Ar¹ is B2);

when Ar¹ is B2), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to C₆ alkenyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group;

when Ar¹ is B1), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to C₆ alkenyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl


group, a piperidinyl group, a piperazinyl group, or a morpholinyl group, and R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group, or R¹² and R¹³ may together form a C₃ to C₅ alkylene group or a C₁ to C₂ alkylenedioxy group;
 5 and

when Ar¹ is B6a), B9a), or B9b), w¹² is an oxygen atom or a sulfur atom.

[8] The compound according to [7] or a pharmacologically acceptable salt thereof, wherein

10 in the formula (I), Ar² is a group selected from the group consisting of A1a), A2b), A3a), and A7b):

[Chemical Formula 6]



wherein, when Ar² is A1a), R⁶ is the same as that defined in [6] when Ar² is A1);

15 when Ar² is A1a) or A7b), R⁸ is the same as that defined in [4] when Ar² is A1) or A7);

when Ar² is A7b), m is the same as that defined in [2] when Ar² is A7);

R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₃ alkyl group, a C₁ to C₃ alkoxy group, a hydroxy C₁ to C₄ alkyl group, a carboxy C₁ to C₃ alkyl group, a carbamoyl C₁ to C₃ alkyl group, a mono-C₁ to C₂ alkylcarbamoyl C₁ to C₃ alkyl group, or a di-C₁ to C₂ alkylcarbamoyl C₁ to C₃ alkyl group;

20

R^2 and R^3 are each independently a hydrogen atom or a C_1 to C_3 alkyl group;

when X is b) or c), R^4 is a hydrogen atom, a hydroxy C_1 to C_4 alkyl group, or a C_1 to C_3 alkyl group;

5 when Ar^2 is A2b) or A3a), R^{6a} is a C_1 to C_3 alkoxy group;

when Ar^2 is A1a), A2b), A3a), or A7b), R^7 is a hydrogen atom, a fluorine atom, a chlorine atom, or a C_1 to C_3 alkyl group; and

when Ar^2 is A7b), W^6 is $C=O$, CH_2 , CF_2 , $CHOH$, or an oxygen atom;


provided that, when Ar^1 is A1a), combinations of the

10 substituents R^6 , R^7 and R^8 exclude a combination of a hydrogen atom and a halogen atom.

[9] The compound according to [8] or a pharmacologically acceptable salt thereof, wherein

in the formula (I), Ar^2 is a group selected from the group
15 consisting of A1a), A2b), and A7c):

[Chemical Formula 7]



wherein, when Ar^2 is A2b), R^{6a} is the same as that defined in
[8] when Ar^2 is A2b);

when Ar^2 is A1a) or A7c), R^8 is the same as that defined in
20 [4] when Ar^2 is A1) or A7);

when Ar^2 is A7c), m is the same as that defined in [8] when
 Ar^2 is A7b);

when Ar^2 is A1a), R^6 is a hydrogen atom, a fluorine atom, a

chlorine atom, a cyano group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, or -CONR¹⁰R¹¹, wherein, when R⁶ is -CONR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group; and


when Ar² is A1a), A2b), or A7c), R⁷ is a hydrogen atom, a fluorine atom, or a chlorine atom;

provided that, when Ar² is A1a), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

[10] The compound according to [9] or a pharmacologically acceptable salt thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B2), B6b), B9c), and B9d):

[Chemical Formula 8]



wherein R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₃ alkyl group, a C₁ to C₃ alkoxy group, or a hydroxy C₁ to C₄ alkyl group;

when Ar¹ is B1) or B2), R¹² is a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, a C₁ to C₃ alkyl group, or a C₁ to C₆ alkoxy group;

when Ar¹ is B1) or B6b), R¹³ is a hydrogen atom, a hydroxy group,

a fluorine atom, or a chlorine atom;

when Ar¹ is B9c) or B9d), R¹⁴ is a hydrogen atom, a fluorine atom, a chlorine atom, a C₁ to C₃ alkyl group, a methoxy group, or an ethoxy group;


5 when Ar¹ is B9c) or B9d), R¹⁵ is a hydrogen atom, a fluorine atom, or a chlorine atom; and

when Ar¹ is B2), one of W¹⁰ and W¹¹ is N, and the other thereof is CH.

11] The compound according to [10] or a pharmacologically acceptable
10 salt thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B6b), and B9c1):

[Chemical Formula 9]



wherein when Ar¹ is B1), R¹² is the same as that defined in
15 [10] when Ar¹ is B1);


when Ar¹ is B1) or B6b), R¹³ is the same as that defined in
[10] when Ar¹ is B1) or B6b); and

when Ar¹ is B9c1), R¹⁴ is the same as that defined in [10] when
Ar¹ is B9c).

20 [12] The compound according to [11] or a pharmacologically acceptable
salt thereof, wherein

in the formula (I), Ar² is a group selected from the group consisting of A1b), A2b), and A7c):

[Chemical Formula 10]



wherein, when Ar² is A2b), R^{6a} is the same as that defined in [8] when Ar² is A2b);

when Ar² is A7c), m is the same as that defined in [8] when Ar² is A7b);

when Ar² is A1b), R⁶ is a cyano group, an ethyl group, or a C₁ to C₃ alkoxy group;

R⁷ is a fluorine atom or a chlorine atom; and

when Ar² is A1b) or A7c), R⁸ is a hydrogen atom, a fluorine atom, a chlorine atom, or a C₁ to C₃ alkyl group.

[13] The compound according to [1] or a pharmacologically acceptable salt thereof, wherein

the compound represented by the formula (1) is

(-)-1-(4-fluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-chlorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(3,4-difluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-cyanophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-chlorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

5 (-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

1-[(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

10 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

15 (-)-1-[(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

20 (-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea,

25 (-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-met

hoxypyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methoxyphenyl)urea,

(-)-1-(5-chlorothiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(5-chloropyridin-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylthiophen-2-yl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea,

(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-2-oxo-4-phenylpyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethoxy)phenyl]urea,

5 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-phenoxyphenyl)urea,

10 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3,4-difluorophenyl)urea,

(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

15 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea,

(-)-1-[(3S*,4R*)-1-ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

20 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-1-(1-hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(+)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

25 (-)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

ethoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylacetamide,

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer A),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer B),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N,2-dimethylpropionamide,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(+)-1-(4-fluorophenyl)-3-[(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]benzamide,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

5 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

10 (-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(2-hydroxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxy-4-methylphenyl)urea,

15 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-hydroxyphenyl)urea,

(-)-1-(4-chloro-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

20 (-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(p-tolyl)urea, or

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-phenylurea.

25 [14] A pharmaceutical containing, as an active ingredient, the

compound according to any one of [1] to [13] or a pharmacologically acceptable salt thereof.

[15] An FPRL1 agonist containing, as an active ingredient, the compound according to any one of [1] to [13] or a pharmacologically acceptable salt thereof.

[16] A method of treatment or prevention of inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders, comprising administering the compound according to any one of [1] to [13] or a pharmacologically acceptable salt thereof.

[17] Use of the compound according to any one of [1] to [13] or a pharmacologically acceptable salt thereof to produce a pharmaceutical for treatment or prevention of inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders.

[18] A pharmaceutical composition containing the compound according to any one of [1] to [13] or a pharmacologically acceptable salt thereof and a pharmaceutically acceptable carrier, used for prevention or treatment of inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders.

Advantageous Effects of Invention

[0016]

The compound (I) or a pharmacologically acceptable salt thereof exhibited superior agonist activity in, for example, a test of calcium
5 influx into FPRL1-overexpressing cells. The compound (I) and salts thereof strongly suppressed lipopolysaccharide-induced neutrophilic infiltration into the lungs of mice. In addition, the compound (I) and salts thereof have low toxicity and are therefore safe. Therefore, the compound (I) according to the present invention
10 or a pharmacologically acceptable salt thereof is useful as a therapeutic or prophylactic agent for inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, immune
15 disorders and the like.

[0017]

In addition, the compound (I) according to the present invention or a pharmacologically acceptable salt thereof is highly useful for treatment, prevention, or suppression of various disease
20 states associated with the FPRL1 receptor (such as Behcet's disease, Sweet disease, systemic lupus erythematosus (SLE), Wegener's granulomatosis, virus infection, diabetes, amputations, cancers, bacterial infection, physical external injuries, physical disorders including exposure to radiation, vasoconstriction, anaphylactic
25 reactions, allergic reactions, rhinitis, shocks (endotoxic,

hemorrhagic, traumatic, splanchnic ischemia, and circulatory shocks), rheumatoid arthritis, gout, psoriasis, benign prostatic hyperplasia, myocardial ischemia, myocardial infarction, brain injuries, pulmonary diseases, COPD, COAD, COLD, acute lung injury, acute respiratory distress syndrome, chronic bronchitis, pulmonary emphysema, asthma (allergic asthma and non-allergic asthma), cystic pulmonary fibrosis, nephropathy, renal glomerular diseases, ulcerative colitis, IBD, Crohn's disease, periodontitis, pains, Alzheimer's disease, AIDS, uveitic glaucoma, conjunctivitis, Sjogren's syndrome, and rhinitis).

Description of Embodiments

[0018]

Terms in the present description will be described.

[0019]

The term "halogen atom" as used herein means a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom. Preferably, the halogen atom is a fluorine atom or a chlorine atom.

[0020]

The 5-membered aromatic heterocyclic group in the term "5-membered aromatic heterocyclic group optionally having substituent(s)" as used herein means a 5-membered aromatic heterocyclic group containing, in its ring, 1 to 4 atoms selected from sulfur, oxygen, and nitrogen atoms. Examples of the 5-membered aromatic heterocyclic group may include a furyl group, a thienyl group, a pyrrolyl group, an azepinyl group, a pyrazolyl group, an

imidazolyl group, an oxazolyl group, an isoxazolyl group, a thiazolyl group, an isothiazolyl group, a 1,2,3-oxadiazolyl group, a triazolyl group, a tetrazolyl group, a thiadiazolyl group and the like.

[0021]

5 The 6-membered aromatic heterocyclic group in the term "6-membered aromatic heterocyclic group optionally having substituent(s)" as used herein means a 6-membered aromatic heterocyclic group containing, in its ring, 1 to 4 nitrogen atoms. Examples of the 6-membered aromatic heterocyclic group may include
10 a pyridyl group, a pyridazinyl group, a pyrimidinyl group, a pyrazinyl group and the like.

[0022]

 The bicyclic aromatic heterocyclic group having 8 or 9 atoms in the term "bicyclic aromatic heterocyclic group having 8 or 9
15 atoms and optionally having substituent(s)" as used herein means a bicyclic aromatic heterocyclic group having 8 or 9 atoms containing 1 to 4 atoms selected from sulfur, oxygen, and nitrogen atoms. Examples of the bicyclic aromatic heterocyclic group having 8 or 9 atoms may include a benzofuranyl group, an isobenzofuranyl group,
20 a benzoxazolyl group, a benzisoxazolyl group, a benzothiazolyl group, a benzisothiazolyl group, a benzimidazolyl group, a benzothiophenyl group, an indolyl group, an isoindolyl group, an indazolyl group, a thiazolopyridyl group, an oxazolopyrazinyl group and the like.

[0023]

25 The C₁ to C₆ alkoxy group in the terms "C₁ to C₆ alkoxy group

optionally having substituent(s)" and "C₁ to C₆ alkoxy group" as used herein means a linear or branched alkoxy group having 1 to 6 carbon atoms. Examples of the C₁ to C₆ alkoxy group may include a methoxy group, an ethoxy group, a propoxy group, an isopropoxy group, an isobutoxy group, a butoxy group, a sec-butoxy group, a tert-butoxy group, a pentyloxy group, a hexyloxy group and the like. Preferred examples may include a methoxy group and an ethoxy group.

[0024]

The C₁ to C₆ alkyl group in the terms "C₁ to C₆ alkyl group optionally having substituent(s)" and "C₁ to C₆ alkyl group" as used herein means a linear or branched alkyl group having 1 to 6 carbon atoms and optionally having substituent(s). Examples of the C₁ to C₆ alkyl group may include a methyl group, an ethyl group, a propyl group, an isopropyl group, a butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an isopentyl group, a neopentyl group, a 1-methylbutyl group, a 2-methylbutyl group, a 1,2-dimethylpropyl group, a hexyl group, an isohexyl group and the like.

[0025]

The C₁ to C₆ acyl group in the terms "C₁ to C₆ acyl group optionally having substituent(s)" and "C₁ to C₆ acyl group" as used herein means an acyl group derived from a linear or branched aliphatic carboxylic acid having 1 to 6 carbon atoms. Examples of the C₁ to C₆ acyl group may include a formyl group, an acetyl group, a propanoyl group, a butanoyl group, a pentanoyl group, a hexanoyl group and the like.

[0026]

The C₁ to C₆ alkylsulfanyl group in the terms "C₁ to C₆ alkylsulfanyl group optionally having substituent(s)" and "C₁ to C₆ alkylsulfanyl group" as used herein means a linear or branched
5 alkylsulfanyl group having 1 to 6 carbon atoms or a cyclic alkylsulfanyl group having 3 to 6 carbon atoms. Examples of the C₁ to C₆ alkylsulfanyl group may include a methylsulfanyl group, an ethylsulfanyl group, a propylsulfanyl group, an isopropylsulfanyl group, a butylsulfanyl group, an isobutylsulfanyl group, a
10 sec-butylsulfanyl group, a tert-butylsulfanyl group, a cyclopropylsulfanyl group, a cyclobutylsulfanyl group, a cyclopentylsulfanyl group and the like.

[0027]

The C₁ to C₆ alkylsulfinyl group in the terms "C₁ to C₆ alkylsulfinyl group optionally having substituent(s)" and "C₁ to C₆ alkylsulfinyl group" as used herein means a linear or branched
15 alkylsulfinyl group having 1 to 6 carbon atoms or a cyclic alkylsulfinyl group having 3 to 6 carbon atoms. Examples of the C₁ to C₆ alkylsulfinyl group may include a methylsulfinyl group, an ethylsulfinyl group, a propylsulfinyl group, an isopropylsulfinyl group, a butylsulfinyl group, an isobutylsulfinyl group, a
20 sec-butylsulfinyl group, a tert-butylsulfinyl group, a cyclopropylsulfinyl group, a cyclobutylsulfinyl group, a cyclopentylsulfinyl group and the like.

[0028]

The C₁ to C₆ alkylsulfonyl group in the terms "C₁ to C₆ alkylsulfonyl group optionally having substituent(s)" and "C₁ to C₆ alkylsulfonyl group" as used herein means a linear or branched alkylsulfonyl group having 1 to 6 carbon atoms or a cyclic alkylsulfonyl group having 3 to 6 carbon atoms. Examples of the C₁ to C₆ alkylsulfonyl group may include a methylsulfonyl group, an ethylsulfonyl group, a propylsulfonyl group, an isopropylsulfonyl group, a butylsulfonyl group, an isobutylsulfonyl group, a sec-butylsulfonyl group, a tert-butylsulfonyl group, a cyclopropylsulfonyl group, a cyclobutylsulfonyl group, a cyclopentylsulfonyl group and the like.

[0029]

The heterocyclic group in the term "heterocyclic group optionally having substituent(s)" as used herein means a 5- to 7-membered heterocyclic group containing 1 to 4 atoms selected from sulfur, oxygen, and nitrogen atoms. Examples of the heterocyclic group may include: aromatic heterocyclic groups such as a furyl group, a thienyl group, a pyrrolyl group, an azepinyl group, a pyrazolyl group, an imidazolyl group, an oxazolyl group, an isoxazolyl group, a thiazolyl group, an isothiazolyl group, a 1,2,3-oxadiazolyl group, a triazolyl group, a tetrazolyl group, a thiadiazolyl group, a pyranyl group, a pyridyl group, a pyridazinyl group, a pyrimidinyl group, and a pyrazinyl group; unsaturated heterocyclic groups such as a pyrrolinyl group, an imidazolinyl group, a pyrazolinyl group, a dihydropyranyl group, a

dihydrothiopyranyl group, and a dihydropyridyl group; and saturated heterocyclic groups such as a morpholinyl group, a thiomorpholinyl group, a pyrrolidinyl group, an imidazolidinyl group, an imidazolinyl group, a pyrazolidinyl group, a pyrazolinyl group, a piperidinyl group, a piperazinyl group, a tetrahydrofuranyl group and the like.

The above "heterocyclic group" may be annelated with another cyclic group. Examples of the heterocyclic group annelated with another cyclic group may include an isobenzofuranyl group, a benzoxazolyl group, a benzisoxazolyl group, a benzothiazolyl group, a benzisothiazolyl group, a chromenyl group, a chromanonyl group, a xanthenyl group, a phenoxathiinyl group, an indolizinyll group, an isoindolizinyll group, an indolyl group, an indazolyl group, a purinyll group, a quinolizinyll group, an isoquinolyl group, a quinolyl group, a phthalazinyll group, a naphthyridinyl group, a quinoxalinyll group, a quinazolinyl group, a carbazolyl group, a carbolinyl group, an acridinyl group, an isoindolinyl group and the like.

[0030]

The term "C₃ to C₁₀ heterocycloalkyl group" as used herein means a monocyclic, bicyclic, or tricyclic non-aromatic heterocycloalkyl group which has a ring structure containing at least one nitrogen atom and optionally containing an oxygen atom or a sulfur atom and in which the number of carbon atoms forming the ring(s) of the cyclic group is 4 to 10. Examples of the C₃ to C₁₀ heterocycloalkyl group may include an azetidinyll group, a pyrrolidinyl group, a piperidyl

group, a piperazyl group, a morpholyl group and the like.

[0031]

The C₃ to C₆ cycloalkyl group in the terms "C₃ to C₆ cycloalkyl group optionally having substituent(s)" and "C₃ to C₆ cycloalkyl group" as used herein means a monocyclic saturated alicyclic hydrocarbon group having 3 to 6 carbon atoms. Examples of the C₃ to C₆ cycloalkyl group may include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group and the like.

[0032]

The C₃ to C₆ cycloalkoxy group in the terms "C₃ to C₆ cycloalkoxy group optionally having substituent(s)" and "C₃ to C₆ cycloalkoxy group" as used herein means a cyclic alkoxy group having 3 to 6 carbon atoms. Examples of the C₃ to C₆ cycloalkoxy group may include a cyclopropyloxy group, a cyclobutyloxy group, a cyclopentyloxy group, and a cyclohexyloxy group.

[0033]

The C₂ to C₆ alkenyl group in the terms "C₂ to C₆ alkenyl group optionally having substituent(s)" and "C₂ to C₆ alkenyl group" as used herein means a linear or branched unsaturated hydrocarbon group having 2 to 6 carbon atoms and having at least one double bond. Examples of the C₂ to C₆ alkenyl group may include a vinyl group, a 2-propenyl group, a 1-propenyl group, a 3-propenyl group, a 1-buten-1-yl group, a 1-buten-2-yl group, a 1-buten-3-yl group, a 1-buten-4-yl group, a 2-buten-1-yl group, a 2-buten-2-yl group, a 1-penten-1-yl group, a 1-penten-2-yl group, a 1-penten-3-yl group,

a 2-penten-1-yl group, a 2-penten-2-yl group, a 2-penten-3-yl group, a 1-hexen-1-yl group, a 1-hexen-2-yl group, a 1-hexen-3-yl group, a 2-methyl-1-propen-1-yl group and the like.

[0034]

5 The "C₂ to C₆ alkynyl group" in the term "C₂ to C₆ alkynyl group optionally having substituent(s)" as used herein means a linear or branched unsaturated hydrocarbon group having 2 to 6 carbon atoms and having at least one triple bond. Examples of the C₂ to C₆ alkynyl group may include an ethynyl group, a 1-propynyl group, a 2-propynyl
10 group, a 1-butynyl group, a 2-butynyl group, a 3-butynyl group, a 3-methyl-1-propynyl group, a 1-ethynyl-2-propynyl group, a 2-methyl-3-propynyl group, a 1-pentynyl group, a 1-hexynyl group, a 1,3-hexanediynyl group, a 1,5-hexanediynyl group and the like.

[0035]

15 The C₁ to C₆ alkoxy carbonyl group in the terms "C₁ to C₆ alkoxy carbonyl group optionally having substituent(s)" and "C₁ to C₆ alkoxy carbonyl group" as used herein means a linear or branched alkoxy carbonyl group having 1 to 6 carbon atoms. Examples of the C₁ to C₆ alkoxy carbonyl group may include a methoxy carbonyl group,
20 an ethoxy carbonyl group, a propoxy carbonyl group, an isopropoxy carbonyl group, an isobutoxy carbonyl group, a butoxy carbonyl group, a sec-butoxy carbonyl group, a tert-butoxy carbonyl group, a pentyloxy carbonyl group, a hexyloxy carbonyl group and the like. Preferred examples thereof
25 may include a methoxy carbonyl group and a tert-butoxy carbonyl group.

[0036]

The term "halo-C₁ to C₆ alkoxy group" as used herein means a C₁ to C₆ alkoxy group substituted with 1 to 5 halogen atoms of the same type or different types. Examples of the halo-C₁ to C₆ alkoxy group may include a fluoromethoxy group, a difluoromethoxy group, a trifluoromethoxy group, a 2-fluoroethoxy group, a 2-chloroethoxy group, a 2,2-difluoroethoxy group, a 1,1-difluoroethoxy group, a 1,2-difluoroethoxy group, a 2,2,2-trifluoroethoxy group, a 1,1,2,2,2-pentafluoroethoxy group, a 2,2,2-trichloroethoxy group, a 3-fluoropropoxy group, a 2-fluoropropoxy group, a 1-fluoropropoxy group, a 3,3-difluoropropoxy group, a 2,2-difluoropropoxy group, a 1,1-difluoropropoxy group, a 4-fluorobutoxy group, a 5-fluoropentoxy group, a 6-fluorohexyloxy group and the like.

[0037]

The term "hydroxy C₁ to C₆ alkyl group" as used herein means a C₁ to C₆ alkyl group substituted with a hydroxy group. Examples of the hydroxy C₁ to C₆ alkyl group may include a 2-hydroxyethyl group, a 1-hydroxyethyl group, a 3-hydroxypropyl group, a 2-hydroxypropyl group, a 1-hydroxypropyl group, a 4-hydroxybutyl group, a 3-hydroxybutyl group, a 2-hydroxybutyl group, a 1-hydroxybutyl group, a 5-hydroxypentyl group, a 6-hydroxyhexyl group and the like.

[0038]

Examples of the "C₁ to C₂ alkylenedioxy group" as used herein

may include a methylenedioxy group (-O-CH₂-O-) and an ethylenedioxy group (-O-CH₂CH₂-O-).

[0039]

The term "C₂ to C₆ alkylene group" as used herein means a divalent linear or branched saturated hydrocarbon chain having 2 to 6 carbon atoms. Examples of the C₂ to C₆ alkylene group may include -(CH₂)₂-, -(CH₂)₃-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, -(CH₂)₄-, -CH(CH₃)-(CH₂)₂-, -(CH₂)₂-CH(CH₃)-, -CH(CH₂CH₃)-CH₂-, -C(CH₃)₂CH₂-, -CH₂-C(CH₃)₂-, -CH(CH₃)-CH(CH₃)-, -(CH₂)₅-, -CH(CH₃)-(CH₂)₃-, -C(CH₃)₂CH₂CH₂-, -(CH₂)₆-, -C(CH₃)₂-(CH₂)₃- and the like. Preferred examples thereof may include -(CH₂)₂- and -(CH₂)₃-.

[0040]

The term "C₃ to C₅ alkylene group" as used herein means a divalent linear or branched saturated hydrocarbon chain having 3 to 5 carbon atoms. Examples of the C₃ to C₅ alkylene group may include -(CH₂)₃-, -CH(CH₃)CH₂-, -CH₂CH(CH₃)-, -(CH₂)₄-, -CH(CH₃)-(CH₂)₂-, -(CH₂)₂-CH(CH₃)-, -CH(CH₂CH₃)-CH₂-, -C(CH₃)₂CH₂-, -CH₂-C(CH₃)₂-, -CH(CH₃)-CH(CH₃)-, -(CH₂)₅-, -CH(CH₃)-(CH₂)₃-, -C(CH₃)₂CH₂CH₂- and the like.

[0041]

The term "aryloxy group" as used herein means an aromatic hydrocarbon alkoxy group having 6 to 14 carbon atoms. Examples of the aryloxy group may include a phenyloxy group, an indenyloxy group, a naphthyloxy group, a phenanthrenyloxy group, an anthracenyloxy group and the like.

[0042]

The term "carboxy C₁ to C₆ alkyl group" as used herein means a C₁ to C₆ alkyl group substituted with a carboxylic acid. Examples of the carboxy C₁ to C₆ alkyl group may include a carboxymethyl group, a 2-carboxyethyl group, a 2-carboxypropyl group, a 3-carboxypropyl group, a 4-carboxybutyl group, a 5-carboxypentyl group, a 6-carboxyhexyl group and the like.

[0043]

The term "carbamoyl C₁ to C₆ alkyl group" as used herein means a C₁ to C₆ alkyl group substituted with a carbamoyl group. Examples of the carbamoyl C₁ to C₆ alkyl group may include a carbamoylmethyl group, a 2-carbamoylethyl group, a 2-carbamoylpropyl group, a 3-carbamoylpropyl group, a 4-carbamoylbutyl group, a 5-carbamoylpentyl group, a 6-carbamoylhexyl group and the like.

[0044]

The term "mono C₁ to C₆ alkylcarbamoyl C₁ to C₆ alkyl group" as used herein means a C₁ to C₆ alkyl group substituted with a carbamoyl group in which one hydrogen atom in the amino group is substituted with a C₁ to C₆ alkyl group. Examples of the mono C₁ to C₆ alkylcarbamoyl C₁ to C₆ alkyl group may include an N-methylcarbamoylmethyl group, an N-ethylcarbamoylmethyl group, a 2-(N-methylcarbamoyl)ethyl group, a 2-(N-ethylcarbamoyl)ethyl group, a 2-(N-propylcarbamoyl)ethyl group, a 3-(N-methylcarbamoyl)propyl group, a 4-(N-ethylcarbamoyl)butyl group, a 5-(N-ethylcarbamoyl)pentyl group, a 6-(N-propylcarbamoyl)hexyl

group and the like.

[0045]

The term "di- C_1 to C_6 alkylcarbamoyl C_1 to C_6 alkyl group" as used herein means a C_1 to C_6 alkyl group substituted with a carbamoyl group in which two hydrogen atoms in the amino group are substituted with C_1 to C_6 alkyl groups. Examples of the di- C_1 to C_6 alkylcarbamoyl C_1 to C_6 alkyl group may include an N,N-dimethylcarbamoylmethyl group, an N-ethyl-N-methylcarbamoylmethyl group, a 2-(N,N-dimethylcarbamoyl)ethyl group, a 2-(N-ethyl-N-methylcarbamoyl)ethyl group, a 2-(N-methyl-N-propylcarbamoyl)ethyl group, a 3-(N,N-dimethylcarbamoyl)propyl group, a 4-(N,N-diethylcarbamoyl)butyl group, a 5-(N-ethyl-N-propylcarbamoyl)pentyl group, a 6-(N,N-dipropylcarbamoyl)hexyl group and the like.

[0046]

The term " C_1 to C_6 alkylamino group" as used herein means an amino group in which one or two hydrogen atoms in the amino group are substituted with linear or branched alkyl groups having 1 to 6 carbon atoms. Examples of the C_1 to C_6 alkylamino group may include a methylamino group, an ethylamino group, a propylamino group, an isopropylamino group, a butylamino group, an isobutylamino group, a sec-butylamino group, a tert-butylamino group, a pentylamino group, an isopentylamino group, a neopentylamino group, a 1-methylbutylamino group, a 2-methylbutylamino group, a

1,2-dimethylpropylamino group, a hexylamino group, an isohexylamino group, a dimethylamino group, a diethylamino group, an N-ethyl-N-methylamino group, an N-ethyl-N-propylamino group and the like.

5 [0047]

The term "C₁ to C₆ acylamino group" as used herein means an amino group substituted with C₁ to C₆ acyl. Examples of the C₁ to C₆ acylamino group may include a formylamino group, an acetylamino group, a propanoylamino group, a butanoylamino group, a
10 pentanoylamino group, a hexanoylamino group and the like.

[0048]

The term "C₁ to C₃ alkyl group" as used herein means a linear or branched alkyl group having 1 to 3 carbon atoms. Examples of the C₁ to C₃ alkyl group may include a methyl group, an ethyl group,
15 a propyl group, and an isopropyl group.

[0049]

The term "C₁ to C₃ alkoxy group" as used herein means a linear or branched alkoxy group having 1 to 3 carbon atoms. Examples of the C₁ to C₃ alkoxy group may include a methoxy group, an ethoxy
20 group, a propoxy group, and an isopropoxy group.

[0050]

The term "hydroxy C₁ to C₄ alkyl group" as used herein means a linear or branched alkyl group having 1 to 4 carbon atoms and substituted with a hydroxy group. Examples of the hydroxy C₁ to
25 C₄ alkyl group may include a 2-hydroxyethyl group, a 1-hydroxyethyl

group, a 3-hydroxypropyl group, a 2-hydroxypropyl group, a 1-hydroxypropyl group, a 4-hydroxybutyl group, a 3-hydroxybutyl group and the like.

[0051]

5 The term "carboxy C_1 to C_3 alkyl group" as used herein means a C_1 to C_3 alkyl group substituted with a carboxylic acid. Examples of the carboxy C_1 to C_3 alkyl group may include a carboxymethyl group, a 2-carboxyethyl group, a 2-carboxypropyl group, a 3-carboxypropyl group and the like.

10 [0052]

 The term "carbamoyl C_1 to C_3 alkyl group" as used herein means a C_1 to C_3 alkyl group substituted with a carbamoyl group. Examples of the carbamoyl C_1 to C_3 alkyl group may include a carbamoylmethyl group, a 2-carbamoylethyl group, a 2-carbamoylpropyl group, a 3-carbamoylpropyl group and the like.

[0053]

 The term "mono C_1 to C_2 alkylcarbamoyl C_1 to C_3 alkyl group" as used herein means a C_1 to C_3 alkyl group substituted with a carbamoyl group in which one hydrogen atom in the amino group is substituted with a C_1 to C_2 alkyl group. Examples of the mono C_1 to C_2 alkylcarbamoyl C_1 to C_3 alkyl group may include an N-methylcarbamoylmethyl group, an N-ethylcarbamoylmethyl group, a 2-(N-methylcarbamoyl)ethyl group, a 2-(N-ethylcarbamoyl)ethyl group, a 3-(N-methylcarbamoyl)propyl group and the like.

25 [0054]

The term "di- C_1 to C_2 alkylcarbamoyl C_1 to C_3 alkyl group" as used herein means a C_1 to C_3 alkyl group substituted with a carbamoyl group in which two hydrogen atoms in the amino group are substituted with C_1 to C_2 alkyl groups. Examples of the di- C_1 to C_2 alkylcarbamoyl C_1 to C_3 alkyl group may include an N,N-dimethylcarbamoylmethyl group, an N-ethyl-N-methylcarbamoylmethyl group, a 2-(N,N-dimethylcarbamoyl)ethyl group, a 2-(N-ethyl-N-methylcarbamoyl)ethyl group, a 3-(N,N-dimethylcarbamoyl)propyl group, a 3-(N,N-diethylcarbamoyl)propyl group and the like.

[0055]

Examples of an "aromatic hydrocarbon cyclic group" as used herein may include a phenyl group, an indenyl group, a 1-naphthyl group, a 2-naphthyl group, an azulenyl group, a heptalenyl group, a biphenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a benzocyclooctenyl group and the like.

[0056]

The term "aromatic heterocyclic group" as used herein means an aromatic cyclic structure containing a nitrogen atom, an oxygen atom, or a sulfur atom. Examples of the aromatic heterocyclic group may include a furyl group, a thienyl group, a pyrrolyl group, an azepinyl group, a pyrazolyl group, an imidazolyl group, an oxazolyl group, an isoxazolyl group, a thiazolyl group, an isothiazolyl group, a 1,2,3-oxadiazolyl group, a triazolyl group, a tetrazolyl group,

a thiadiazolyl group, a pyranyl group, a pyridyl group, a pyridazinyl group, a pyrimidinyl group, a pyrazinyl group and the like. The above "heterocyclic group" may be annelated with another cyclic group. Examples of such a heterocyclic group may include an
5 isobenzofuranyl group, a benzoxazolyl group, a benzisoxazolyl group, a benzothiazolyl group, a benzisothiazolyl group, a chromenyl group, a chromanonyl group, a xanthenyl group, a phenoxathiinyl group, an indolizinyll group, an isoindolizinyll group, an indolyl group, an indazolyl group, a purinyl group, a quinolizinyll group, an
10 isoquinolyl group, a quinolyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a carbolinyl group, an acridinyl group, an isoindolinyl group and the like.

[0057]

15 The term "halo-C₁ to C₆ alkyl group" as used herein means a C₁ to C₆ alkyl group substituted with 1 to 5 halogen atoms of the same type or different types. Examples of the halo-C₁ to C₆ alkyl group may include a fluoromethyl group, a difluoromethyl group, a trifluoromethyl group, a 2-fluoroethyl group, a 2-chloroethyl
20 group, a 2,2-difluoroethyl group, a 1,1-difluoroethyl group, a 1,2-difluoroethyl group, a 2,2,2-trifluoroethyl group, a 1,1,2,2,2-pentafluoroethyl group, a 2,2,2-trichloroethyl group, a 3-fluoropropyl group, a 2-fluoropropyl group, a 1-fluoropropyl group, a 3,3-difluoropropyl group, a 2,2-difluoropropyl group, a
25 1,1-difluoropropyl group, a 4-fluorobutyl group, a 5-fluoropentyl

group, a 6-fluorohexyl group and the like.

[0058]

No particular limitation is imposed on the groups acceptable as the "substituent(s)" in the "phenyl group optionally having substituent(s)," the "5-membered aromatic heterocyclic group optionally having substituent(s)," the "6-membered aromatic heterocyclic group optionally having substituent(s)," the "bicyclic aromatic heterocyclic group having 8 or 9 atoms and optionally having substituent(s)," and the "heterocyclic group optionally having substituent(s)," so long as the substituent(s) are generally known substituent(s). Examples of these substituent(s) may include halogen atoms, an amino group, a hydroxy group, a cyano group, a nitro group, a carboxy group, C₁ to C₆ alkoxy carbonyl groups, a formyl group, C₁ to C₆ acyl groups, C₁ to C₆ alkyl groups, C₁ to C₆ alkylamino groups, C₁ to C₆ alkoxy groups, C₁ to C₆ alkylthio groups, C₃ to C₆ cycloalkyl groups, 4- to 10-membered heterocycloalkyl groups, aromatic hydrocarbon cyclic groups optionally having a halogen atom, aromatic heterocyclic groups, C₁ to C₆ alkylcarbonylamino groups, C₃ to C₆ cycloalkylcarbonylamino groups, 4- to 10-membered heterocycloalkylcarbonylamino groups, aromatic hydrocarbon cyclic carbonylamino groups, aromatic heterocyclic carbonylamino groups and the like.

[0059]

No particular limitation is imposed on the groups acceptable as the "substituent(s)" in the "C₁ to C₆ alkoxy group optionally

having substituent(s)," the "C₁ to C₆ alkyl group optionally having substituent(s)," the "C₁ to C₆ acyl group optionally having substituent(s)," the "C₁ to C₆ alkylsulfanyl group optionally having substituent(s)," the "C₁ to C₆ alkylsulfinyl group optionally having substituent(s)," the "C₁ to C₆ alkylsulfonyl group optionally having substituent(s)," the "C₃ to C₆ cycloalkyl group optionally having substituent(s)," the "C₃ to C₆ cycloalkoxy group optionally having substituent(s)," the "C₂ to C₆ alkenyl group optionally having substituent(s)," the "C₁ to C₆ alkoxy carbonyl group optionally having substituent(s)," the "aromatic heterocyclic C₁ to C₃ alkyl group optionally having substituent(s)," and the "phenyl C₁ to C₃ alkyl group optionally having substituent(s)," so long as the substituent(s) are generally known substituent(s). Examples of these substituent(s) may include halogen atoms, an amino group, a hydroxy group, a cyano group, a nitro group, a carboxy group, C₁ to C₆ alkoxy carbonyl groups, a formyl group, C₁ to C₆ acyl groups, C₁ to C₆ alkyl groups, C₁ to C₆ alkylamino groups, di-C₁ to C₆ alkylamino groups, C₁ to C₆ alkoxy groups, C₁ to C₆ alkylthio groups, C₃ to C₆ cycloalkyl groups, 4- to 10-membered heterocycloalkyl groups, aromatic hydrocarbon cyclic groups optionally having a halogen atom, aromatic heterocyclic groups, C₁ to C₆ alkylcarbonylamino groups, C₃ to C₆ cycloalkylcarbonylamino groups, 4- to 10-membered heterocycloalkylcarbonylamino groups, aromatic hydrocarbon cyclic carbonylamino groups, aromatic heterocyclic carbonylamino groups and the like.

[0060]

Hereinafter, the present embodiment will be described in more detail.

[0061]

5 In the following, descriptions of the definitions of functional groups included in general formulas may be omitted, and the definitions already described may be quoted instead. The definitions quoted refer to definitions in the description of the following embodiment.


10 As for the definitions of functional groups included in the general formulas, the definition of a symbol is common to general formulas containing this symbol, unless otherwise mentioned.

[0062]

The present embodiment relates to a urea compound represented
15 by the following general formula (I) or a pharmacologically acceptable salt thereof.

[0063]

[Chemical Formula 11]



20 [0064]

In the formula (I), Ar¹ is a phenyl group optionally having

substituent(s), a 5-membered aromatic heterocyclic group optionally having substituent(s), a 6-membered aromatic heterocyclic group optionally having substituent(s), or a bicyclic aromatic heterocyclic group having 8 or 9 atoms and optionally having substituent(s);

Ar^2 is a phenyl group optionally having substituent(s) (except for a phenyl group substituted only with halogen atom(s)), a 5-membered aromatic heterocyclic group optionally having substituent(s), a 6-membered aromatic heterocyclic group optionally having substituent(s), or a bicyclic aromatic heterocyclic group having 8 or 9 atoms and optionally having substituent(s);

X is a group selected from the group consisting of the following a), b), and c),

- a) an oxygen atom or a sulfur atom,
- b) NR^4 , and
- c) NOR^4 , wherein,

when X is b) or c), R^4 is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s);

R^1 is a hydrogen atom, a hydroxy group, a C_1 to C_6 alkoxy group optionally having substituent(s), or a C_1 to C_6 alkyl group optionally having substituent(s);

R^2 and R^3 are each independently a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s) or R^2 and R^3 together form a C_2 to C_6 alkylene group; and

Each carbon atom marked with an asterisk is an asymmetric carbon

atom.

The term "independently" means that at least two substituents present may be the same or different.

[0065]


5 In the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof, preferred substituents are as follows.

[0066]

Preferably, Ar¹ is a group selected from the group consisting
10 of the following B1), B2), B3), B4), B5), B6), B7), B8), B9), B10), B11), and B12).

[0067]

[Chemical Formula 12]




[0068]

15 More preferably, Ar¹ is a group selected from the group consisting of the following B1), B2), B6a), B9a), and B9b).

[0069]

[Chemical Formula 13]



[0070]

R^{12} is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 alkoxy group optionally having substituent(s), a C_3 to C_6 cycloalkyl group optionally having substituent(s), a C_3 to C_6 cycloalkoxy group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), a C_2 to C_6 alkenyl group optionally having substituent(s), a C_2 to C_6 alkynyl group optionally having substituent(s), a C_1 to C_6 alkoxycarbonyl group, a C_1 to C_6 alkylsulfanyl group optionally having substituent(s), a C_1 to C_6 alkylsulfinyl group optionally having substituent(s), a C_1 to C_6 alkylsulfonyl group optionally having substituent(s), $-\text{CONR}^{10}\text{R}^{11}$, $-\text{NR}^{10}\text{R}^{11}$, an aryloxy group, or a heterocyclic group optionally having substituent(s). When R^{12} is $-\text{CONR}^{10}\text{R}^{11}$ or $-\text{NR}^{10}\text{R}^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), or a C_1 to C_6 alkylsulfonyl group optionally having substituent(s) and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s), or R^{10} and R^{11} may together form a C_3 to C_{10} heterocycloalkyl group.

R^{13} may be a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C_1 to C_6 alkyl group.

When Ar¹ is B1), each of R¹² and R¹³ is any of the above described functional groups, or R¹² and R¹³ may together form a C₃ to C₅ alkylene group or a C₁ to C₂ alkylenedioxy group.

R¹⁴ and R¹⁵ are each independently a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group, or a C₁ to C₆ alkoxy group.

One of W¹⁰ and W¹¹ is a nitrogen atom, and the other thereof is CH or a nitrogen atom.

W¹² is an oxygen atom, a sulfur atom, or N-R¹⁶. When W¹² is N-R¹⁶, R¹⁶ is a hydrogen atom or a C₁ to C₆ alkyl group.

W¹³ is CH or a nitrogen atom.

[0071]

Preferably, one of W¹⁰ and W¹¹ is N, and the other thereof is CH.

W¹² is preferably an oxygen atom or a sulfur atom. W¹² is more preferably a sulfur atom.

W¹³ is preferably CH.

[0072]

Preferably, R¹² is a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), or a C₁ to C₆ alkoxy group optionally having substituent(s). R¹² is more preferably a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, a C₁ to C₃ alkyl group, or a C₁ to C₆ alkoxy group and is particularly more preferably a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, or a C₁ to C₃ alkyl group.

[0073]

Preferably, R^{13} is a hydrogen atom, a hydroxy group, a halogen atom, or a C_1 to C_6 alkyl group. R^{13} is more preferably a hydrogen atom, a fluorine atom, or a chlorine atom.

5 [0074]

Preferably, R^{14} is a hydrogen atom, a fluorine atom, a chlorine atom, a C_1 to C_3 alkyl group, a methoxy group, or an ethoxy group.

[0075]


R^{15} is preferably a hydrogen atom, a fluorine atom, or a chlorine atom and particularly preferably a hydrogen atom.

[0076]

Preferably, Ar^2 is a group selected from the group consisting of the following A1), A2), A3), A4), A5), A6), A7), A8), A9), and A10).

15 [0077]

[Chemical Formula 14]




[0078]

Ar^2 is more preferably a group selected from the group consisting of the following A1), A2a), A3), and A7a).

20 [0079]

[Chemical Formula 15]



[0080]

W¹ is a nitrogen atom or CH optionally substituted with a hydrogen atom, with a halogen atom, or with a C₁ to C₆ alkyl group optionally having substituent(s). W² is CH or a nitrogen atom.

W³ is an oxygen atom, a sulfur atom, or NH optionally substituted with a C₁ to C₆ alkyl group.

W⁴ is CH or a nitrogen atom.

W⁵ is CH₂, an oxygen atom, or a sulfur atom.

W⁶ is C=O, CH₂, CF₂, CHOH, NH optionally substituted with a C₁ to C₆ alkyl group, SO, SO₂, an oxygen atom, or a sulfur atom.

W⁷ is NH optionally substituted with a C₁ to C₆ alkyl group or C=O.

W⁸ is C=O when W⁷ is NH optionally substituted with a C₁ to C₆ alkyl group and is NH optionally substituted with a C₁ to C₆ alkyl group when W⁷ is C=O.

W⁹ is a nitrogen atom or N=O.

R⁶ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl

group optionally having substituent(s), a heterocyclic group optionally having substituent(s), $-\text{CONR}^{10}\text{R}^{11}$, or $-\text{NR}^{10}\text{R}^{11}$. When R^6 is $-\text{CONR}^{10}\text{R}^{11}$ or $-\text{NR}^{10}\text{R}^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), or a C_1 to C_6 alkylsulfonyl group optionally having substituent(s) and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s), or R^{10} and R^{11} may together form a C_3 to C_{10} heterocycloalkyl group.

R^7 is a hydrogen atom, a halogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), or a C_1 to C_6 alkoxy group optionally having substituent(s).

R^8 is a hydrogen atom, a halogen atom, or a C_1 to C_6 alkyl group optionally having substituent(s).

R^9 is a hydrogen atom or a C_1 to C_6 alkyl group.

m is 0 or 1.

n is 0 or 1.

When Ar^2 is A1), combinations of the substituents R^6 , R^7 and R^8 exclude a combination of a hydrogen atom and a halogen atom.

[0081]

Preferably, W^1 is CH or N. W^1 is more preferably N.

Preferably, W^2 is CH or N. W^2 is more preferably CH.

Preferably, W^3 is an oxygen atom or a sulfur atom. W^3 is more preferably a sulfur atom.

Preferably, W^5 is CH_2 or O.

Preferably, W^6 is $\text{C}=\text{O}$, CH_2 , CF_2 , CHOH , or an oxygen atom. W^6

is more preferably C=O, CH₂, CHOH, or an oxygen atom and particularly preferably an oxygen atom.

Preferably, W⁹ is a nitrogen atom.

Preferably, m is 0.

5 [0082]

Preferably, R⁶ is a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ acyl group optionally having
10 substituent(s), a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), -CONR¹⁰R¹¹, or -NR¹⁰R¹¹.

Preferably, when R⁶ is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is
15 a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group.

R⁶ is more preferably a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, a C₁ to C₆ alkoxy group, a halo-C₁
20 to C₆ alkoxy group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkylsulfanyl group, or a C₁ to C₆ alkylsulfinyl group and is particularly preferably a cyano group, an ethyl group, or a C₁ to C₃ alkoxy group.

[0083]

Preferably, R⁷ is a hydrogen atom, a halogen atom, a C₁ to
25 C₆ alkyl group optionally having substituent(s), or a C₁ to C₆ alkoxy

group optionally having substituent(s). R^7 is more preferably a hydrogen atom, a halogen atom, or a C_1 to C_3 alkyl group and is particularly preferably a hydrogen atom, a fluorine atom, or a chlorine atom.

5 [0084]

Preferably, R^8 is a hydrogen atom, a halogen atom, or a C_1 to C_6 alkyl group optionally having substituent(s). R^8 is more preferably a hydrogen atom, a halogen atom, or a C_1 to C_6 alkyl group and is particularly preferably a hydrogen atom, a fluorine atom,
10 a chlorine atom, or a C_1 to C_3 alkyl group.

[0085]

Preferably, R^9 is a hydrogen atom, a methyl group, or an ethyl group.

[0086]

15 Preferably, X is an oxygen atom, NOH, N-(C_1 to C_3 alkyl group), N-O-(C_1 to C_3 alkyl group), or N-O-(hydroxy C_1 to C_4 alkyl group). X is more preferably an oxygen atom, NMe, NOME, NOH, or $\text{NOCH}_2\text{CH}_2\text{OH}$ and is particularly preferably an oxygen atom.

[0087]

20 Preferably, R^1 is a hydrogen atom, a hydroxy group, a C_1 to C_6 alkyl group, a C_1 to C_6 alkoxy group, a halo- C_1 to C_6 alkyl group, a hydroxy C_1 to C_6 alkyl group, a carboxy C_1 to C_6 alkyl group, a carbamoyl C_1 to C_6 alkyl group, a monoalkylcarbamoyl C_1 to C_6 alkyl group, or a dialkylcarbamoyl C_1 to C_6 alkyl group. R^1 is more
25 preferably a hydrogen atom, a hydroxy group, a C_1 to C_3 alkyl group,

a C₁ to C₃ alkoxy group, or a hydroxy C₁ to C₄ alkyl group and is particularly preferably a hydrogen atom, a methyl group, an ethyl group, or a hydroxyethyl group.

[0088]

5 Preferably, R² is a hydrogen atom or a C₁ to C₃ alkyl group. R² is more preferably a hydrogen atom or a methyl group and is particularly preferably a hydrogen atom.

[0089]

10 Preferably, R³ is a hydrogen atom or a C₁ to C₃ alkyl group. R³ is more preferably a hydrogen atom or a methyl group and is particularly preferably a hydrogen atom.

[0090]

Preferred examples of the compound of the present embodiment may include the following compounds:

15 (-)-1-(4-fluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-chlorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

20 (-)-1-(3,4-difluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(2,4-difluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-cyanophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

25 (-)-1-(4-chlorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-1

-methyl-2-oxopyrrolidin-3-yl]urea;

1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

5 (-)-1-[(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

1-[(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

10 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

15 (-)-1-[(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

20 (-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(7-fluorochroman-6-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

25 (-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxo

pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methylsulfonylphenyl)urea;

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methoxyphenyl)urea;

(-)-1-(benzo[d][1,3]dioxole-5-yl)-3-[(3R*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(5-chlorothiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrimidin-4-yl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyridin-2-yl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyridin-3-yl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrimidin-5-yl)urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(5-chloropyridin-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

5 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrazin-2-yl)urea;

(-)-1-(benzo[d]thiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

10 (-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(benzo[d]oxazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylisoxazol-3-yl)urea;

15 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(2-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylthiophen-2-yl)urea;

20 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

1-[(3S*,4R*)-4-(4-ethyl-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

25 (+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-2-oxo-4-phenylpyrroli

din-3-yl]urea;

(+)-1-(4-chlorophenyl)-3-[(3R^{*}, 4S^{*})-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

5 (+)-1-(4-chlorophenyl)-3-[(3R^{*}, 4S^{*})-4-(3-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-chlorophenyl)-3-[(3R^{*}, 4S^{*})-4-(2-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-[(3R^{*}, 4S^{*})-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

10 (+)-1-(4-chlorophenyl)-3-[(3R^{*}, 4S^{*})-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-fluorophenyl)-3-[(3R^{*}, 4S^{*}, 5S^{*})-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

15 (+)-1-(4-fluorophenyl)-3-[(3R^{*}, 4S^{*}, 5R^{*})-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-fluorophenyl)-3-[(3R^{*}, 4S^{*})-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-{(3R^{*}, 4S^{*})-4-[4-(difluoromethoxy)phenyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea;

20 (±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

25 (±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(+)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea;

(-)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea;

5 (+)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea;

(-)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea;

10 (+)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea;

(-)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylsulfinylphenyl)-2-oxopyrrolidin-3-yl]urea;

15 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylsulfonylphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethoxy)phenyl]urea;

20 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea;

(-)-1-[4-(tert-butyl)phenyl]-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}benzoic acid ethyl ester;

25 (-)-1-[(1,1'-biphenyl)-4-yl]-3-[(3S*,4R*)-4-(2,6-difluoro

-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-acetylphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-phenoxyphenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3,4-difluorophenyl)urea;

(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(hydroxymethyl)phenyl]urea;

(-)-4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}-N-methylbenzamide;

(-)-1-[(3S*,4R*)-1-ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid ethyl ester;

2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-2-methylpropionic acid ethyl ester;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-1-(1-hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester (isomer A);

5 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester (isomer B);

(±)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

10 (+)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid;

15 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylacetamide;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A);

20 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer A);

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B);

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer B);

25 2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A);

nyl)-2-oxopyrrolidin-1-yl}-2-methylpropionic acid;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N,2-dimethylpropionamide;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(+)-1-(4-fluorophenyl)-3-[(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-hydroxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]-N,N-dimethylbenzamide;

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]-N-methylbenzamide;

(-)-1-[(3S*,4R*)-4-(4-cyano-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]benzamide;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

5 (-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methylimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

10 (+)-1-(4-fluorophenyl)-3-[(3R*,4S*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(2-hydroxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxyphenyl)urea;

15 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-methylisothiazol-5-yl)urea;

(-)-1-(4-cyclopropylphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

20 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethyl)phenyl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxy-4-methylphenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-hydroxyphenyl)urea;

25 (-)-1-(4-chloro-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difl

uro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

1-(4-cyano-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

1-[(3S*,4R*)-4-[4-(difluoromethoxy)-2,6-difluorophenyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(6-fluorobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

1-[(3S*,4R*)-4-(4,6-difluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(benzo[b]thiophene-2-yl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-ethylphenyl)urea;

(-)-1-(3-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridine-2-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-1-(4-fluorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-2-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid;

5 (-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid;

(-)-2-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-N-methylacetamide;

10 (-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetamide;

(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-2-methylpropanoic acid;

15 (-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-(benzo[b]thiophene-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

20 (-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-phenylurea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea;

25 (-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea;
;

(-)-1-(benzo[b]thiophene-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-propylpyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-1-benzyl-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-(pyridin-3-ylmethyl)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(methylsulfonyl)methyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea;

(-)-{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl}methanesulfonamide;

(-)-2-{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl}acetic acid ethyl ester;

(-)-1-{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(5-methyloxazol-2-yl)methyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea;

(-)-1-{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl}-3-(4-

fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

5 (-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl}acetic acid ethyl ester;

(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-2-methylpropanoic acid ethyl ester;

10 (-)-1-[(3S*,4R*)-1-(cyclopropylmethyl)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-1-(cyanomethyl)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

15 (-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

20 1-[(3S*,4R*)-4-(2,6-difluoro-4-hydroxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(p-tolyl)urea;

25 (-)-1-(4-cyanophenyl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-me

thoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea;

(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(p-tolyl)urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]urea;

(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]urea;

(-)-1-[(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-[(1-methyl-1H-pyrazol-3-yl)amino]-3,4-dihydro-2H-pyrrole-4-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(morpholinoimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

1-[(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-(phenylamino)-3,4-dihydro-2H-pyrrole-4-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

3-((Z)-{(3S,4R)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]pyrrolidin-2-ylidene}amino)propanoic acid ethyl ester;

(-)-1-((3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(hydroxyimino)pyrrolidin-3-yl)-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-ethynylphenyl)urea;

(-)-1-[(3S*,4R*)-4-[2,6-difluoro-4-(methylamino)phenyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea; and

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea.

[0091]

More preferred examples of the compound of the present embodiment may include the following compounds:

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(3,4-difluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-

-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

1-[(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methoxyphenyl)urea;

5 (-)-1-(5-chlorothiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

10 (-)-1-(5-chloropyridin-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylthiophen-2-yl)urea;

15 (-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea;

20 (+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-2-oxo-4-phenylpyrrolidin-3-yl]urea;

(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

25 (±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea;

hyl-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethoxy)phenyl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-phenoxyphenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3,4-difluorophenyl)urea;

(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea;

(-)-1-[(3S*,4R*)-1-ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-1-(1-hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(±)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(+)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylacetamide;

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A);

5 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer A);

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B);

10 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer B);

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N,2-dimethylpropionamide;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

15 (-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

20 (+)-1-(4-fluorophenyl)-3-[(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]benzamide;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

25 (-)-1-[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxy

yimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-{(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl}urea;

(-)-1-{(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}-3-(4-fluorophenyl)urea;

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methylimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea;

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(2-hydroxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxy-4-methylphenyl)urea;

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-hydroxyphenyl)urea;

(-)-1-(4-chloro-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea;

(-)-1-{(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}-3-(p-tolyl)urea; and

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-phenylurea.

[0092]

If necessary, the compound (I) of the present embodiment can

be converted to a pharmacologically acceptable salt according to a usual method. The pharmacologically acceptable salt means a salt with a pharmacologically acceptable nontoxic base or acid (for example, an inorganic or organic base or an inorganic or organic acid).

[0093]

Examples of the salt derived from a pharmacologically acceptable nontoxic base may include: salts with inorganic bases such as sodium salts, potassium salts, calcium salts, magnesium salts and the like; and salts with organic bases such as piperidine, morpholine, pyrrolidine, arginine, lysine and the like.

[0094]

Examples of the salt derived from a pharmacologically acceptable nontoxic acid may include: acid addition salts with mineral acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid and the like; and acid addition salts with organic acids such as formic acid, acetic acid, maleic acid, fumaric acid, succinic acid, lactic acid, malic acid, tartaric acid, citric acid, methanesulfonic acid, p-toluenesulfonic acid, salicylic acid, stearic acid, palmitic acid and the like.

[0095]

The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof may be present as a hydrate or a solvate. Any hydrate and solvate formed from the urea derivative represented by the general formula (I) above, including any of the

preferred compounds specifically described above, or a salt thereof are included in the scope of the present invention. Examples of the solvent that can form the solvate may include methanol, ethanol, 2-propanol, acetone, ethyl acetate, dichloromethane, diisopropyl ether and the like.

[0096]

The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof may be a racemate and also includes their optically active substances, stereoisomers, and rotational isomers.

When the compound (I) of the present embodiment is one of its optical isomers having one or more asymmetric carbon atom, the configuration of each asymmetric carbon atoms in the compound (I) of the present embodiment is any of the R configuration and the S configuration. Any of the optical isomers is included in the present invention, and a mixture of these optical isomers is also included in the present invention. A mixture of optically active substances may be a racemate formed of equal amounts of the optical isomers, and this racemate is also included in the scope of the present invention. When the compound (I) of the present embodiment is a solid or crystalline racemate, the racemate, a racemic mixture, a racemic solid solution are included in the scope of the present invention.

[0097]

When the compound (I) of the present embodiment includes

geometrical isomers, all the geometrical isomers are included in the present invention.

When the compound (I) of the present embodiment includes tautomers, all the tautomers are included in the present invention.

5 Pharmacologically acceptable salts of the compound (I) include proton tautomers.

[0098]

The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof may be a compound labeled
10 with an isotope (for example, ^3H , ^{14}C , ^{35}S and the like). Such a compound is also included in the present invention.

The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof may be a deuterium-substituted compound in which ^1H is substituted with $^2\text{H}(\text{D})$.
15 Such a compound is also included in the present invention.

[0099]

The term "FPRL1 agonist effect" in the present embodiment means agonist activity obtained by the action on formyl peptide receptor like 1 (FPRL1). The compound (I) of the present embodiment or a
20 pharmacologically acceptable salt thereof exhibits superior agonist activity in, for example, a test of calcium influx into FPRL1-overexpressing cells. Therefore, it can be understood that the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof is useful as a therapeutic or prophylactic
25 agent for inflammatory diseases, chronic airway diseases, cancers,

septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, immune disorders and the like.

[0100]

5 Method of producing the compound (I) of the present embodiment

The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof can be produced, for example, in accordance with any of methods described in the following schemes 1 to 17, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0101]

Synthesis of compound (Ia)


The compound (I) of the present embodiment when X is O (this compound is hereinafter referred to as the compound (Ia)) can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0102]

Scheme 1

[0103]

[Chemical Formula 16]



[0104]

In the above formulas, Ar^1 , Ar^2 , R^1 , R^2 , and R^3 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

5 [0105]

Step 1-1

This step is a step of reacting the compound (1) with the compound (2) to produce the compound (Ia). The compound (Ia) can be produced by, for example, allowing diphenylphosphoryl azide (DPPA) and the
10 like to act on the compound (1) in a solvent in the presence or absence of a base and then reacting the compound (2) with the obtained product.

The above reaction is usually performed in a solvent that does not adversely affect the reaction, and examples of the solvent used
15 may include benzene, toluene, tetrahydrofuran, acetonitrile, dioxane, mixed solvents of them and the like. Examples of the base used may include trimethylamine, triethylamine, N-methylmorpholine and the like. The amount of the base used is usually about 0.5 to about 100 molar equivalents with respect to 1 mole of the compound
20 and preferably about 1 to about 5 molar equivalents. The reaction temperature can usually be performed at -10°C to the reflux temperature of the solvent and is performed preferably at 20°C to 120°C . The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature, or the like, but
25 is usually 10 minutes to 3 days.

The compound (1) used in this step can be produced in accordance with any of methods described below in detail, methods similar thereto, methods described in other literatures, and methods similar thereto.

Further, the compound (2) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0106]

Step 1-2

This step is a step of reacting the compound (3) with the compound (4) to produce the compound (Ia). The compound (Ia) can be produced by, for example, reacting the compound (3) with the compound (4) in a solvent in the presence or absence of a base. The amount of the compound (4) used is about 0.5 to about 10 molar equivalents with respect to 1 mole of the compound (3) and is preferably about 1 to about 2 molar equivalents.

The above reaction is usually performed in a solvent that does not adversely affect the reaction, and examples of the solvent used may include dichloromethane, 1,2-dichloroethane, benzene, toluene, tetrahydrofuran, ethyl acetate, methanol, water, mixed solvents of them and the like. Examples of the base used may include alkali metal hydrides such as lithium hydride, sodium hydride, potassium hydride and the like, alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like, hydrogen carbonates such as sodium hydrogen carbonate, potassium hydrogen carbonate and the like, carbonates such as sodium carbonate, potassium carbonate and the

like, organic acid salts such as sodium acetate and the like, tertiary amines such as trimethylamine, triethylamine, N-methylmorpholine and the like, and aromatic amines such as pyridine, picoline, N,N-dimethyl aniline and the like. The amount of the base used is usually about 1 to about 100 molar equivalents with respect to 1 mole of the compound and preferably about 1 to about 5 molar equivalents. The reaction temperature can usually be performed at -20°C to the reflux temperature of the solvent and is performed preferably at 0°C to 50°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 10 minutes to 48 hours.

The compound (3) used in this step can be produced in accordance with any of methods described below in detail, methods similar thereto, methods described in other literatures, and methods similar thereto.

Further, the compound (4) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0107]

Synthesis of compound (Ib)


The compound (I) of the present embodiment when X is O and R¹ is H (this compound may hereinafter be referred to as the compound (Ib)) can be produced in accordance with any of methods described in scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0108]

Scheme 2

[0109]

[Chemical Formula 17]



[0110]

- 5 In the above formulas, Ar¹, Ar², R², and R³ are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0111]

Step 2-1

- 10 This step is a step of producing the compound (Ib) from the compound (1a). The compound (Ib) by the step 2-1 can be produced in accordance with any of the method described in step 1-1 of scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

15 [0112]

Step 2-2

This step is a step of converting the carboxy group in the

compound (1a) to a primary amine to produce the compound (3a). The compound (3a) can be produced by, for example, allowing diphenylphosphoryl azide (DPPA) and the like to act on the compound (1a) in a solvent A in the presence or absence of a base and then
5 reacting the product obtained with an acid in a solvent B. Examples of the solvent A may include benzene, toluene, tetrahydrofuran, acetonitrile, dioxane, mixed solvents of them and the like. Examples of the solvent B may include dioxane, water and the like. Examples of the base used may include trimethylamine, triethylamine,
10 N-methylmorpholine and the like. Examples of the acid used may include conc. hydrochloric acid, conc. sulfuric acid and the like. The reaction temperature can usually be performed at -10°C to the reflux temperature of the solvent and is performed preferably at 0°C to 100°C. The reaction time varies depending on the starting
15 materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0113]

Step 2-3

This step is a step of producing the compound (5) from the
20 compound (1a). The compound (5) can be produced by, for example, allowing diphenylphosphoryl azide (DPPA) and the like to act on the compound (1a) in a solvent in the presence or absence of a base and then reacting the obtained product with benzyl alcohol in a solvent or without any solvent. Examples of the solvent used may
25 include benzene, toluene, tetrahydrofuran, acetonitrile, dioxane,

mixed solvents of them and the like. Examples of the base used may include trimethylamine, triethylamine, N-methylmorpholine and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably
5 at 10°C to 120°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0114]

Step 2-4

10 This step is a step of deprotecting the Cbz (benzyloxycarbonyl) group in the compound (5) to produce the compound (3a). The compound (3a) can be produced by, for example, hydrogenating the compound (5) in a solvent in the presence of a catalyst such as 10% palladium on carbon (10% Pd-C). Examples of the solvent used may include
15 methanol, ethanol, dichloromethane, tetrahydrofuran, ethyl acetate, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the
20 reaction temperature or the like, but is usually 30 minutes to 3 days.

[0115]

Step 2-5

This step is a step of producing the compound (Ib) from the
25 compound (3a). The compound (Ib) by the step 2-5 can be produced

in accordance with any of the method described in step 1-2 of scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0116]

5 Synthesis of compound (1b)


The compound (1b) among the compound (1) when R^3 is H can be produced from the compound (6) in accordance with any of methods described in scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

10 [0117]

Scheme 3

[0118]

[Chemical Formula 18]



[0119]

15 In the above formulas, Ar^2 and R^2 are as described above; Q^1 is a C_{1-6} alkyl group and the like, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0120]

Step 3-1

This step is a step of reacting the compound (6) with the compound (7) to produce the compound (8). The compound (8) can be produced by, for example, reacting the compound (6) with the compound (7) in acetic acid in the presence of ammonium acetate. The reaction temperature can usually be performed at 20°C to the reflux temperature of the solvent and is performed preferably at 80°C to 110°C. Alternatively, the compound (8) can be produced by reacting the compound (6) with the compound (7) in 2-hydroxyethylammonium formate (2-HEAF). The reaction temperature can usually be performed at 0°C to 80°C and is performed preferably at 10°C to 50°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (6) and the compound (7) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0121]

Step 3-2

This step is a step of reacting the compound (8) with a malonic ester (9) to produce the compound (10). The compound (10) can be produced by, for example, reacting the compound (8) with the compound (9) in a solvent in the presence of a catalyst. Examples of the solvent used may include benzene, toluene, tetrahydrofuran, methyl

t-butyl ether, acetonitrile, methanol, ethanol, ethyl acetate and the like. Examples of the catalyst used may include nickel(II) bis[(S,S)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide, nickel(II)

5 bis[(R,R)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide, 1-(3,5-bis(trifluoromethyl)phenyl)-3-((1S,2S)-2-(dimethylamino)cyclohexyl)thiourea, and 1-(3,5-bis(trifluoromethyl)phenyl)-3-((1R,2R)-2-(dimethylamino)cyclohexyl)thiourea that can be obtained according to methods
10 described in Non Patent Literature. The amount of the catalyst used is usually 0.001 to 0.2 moles with respect to 1 mole of the compound (8). The reaction temperature can usually be performed at -0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 50°C. The reaction time varies depending on the starting
15 materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 7 days.

Further, the compound (9) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto. Also,
20 the compound (8) may be a commercially available in addition to that obtained in step 3-1, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0122]

25 Step 3-3

This step is a step of producing the compound (11) from the compound (10). The compound (11) can be produced by, for example, allowing sodium borohydride (NaBH_4) to act on the compound (10) in a solvent in the presence of nickel(II) chloride hexahydrate
5 ($\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$). Examples of the solvent used may include methanol, ethanol, tetrahydrofuran, methyl t-butyl ether and the like. The reaction temperature can usually be performed at -30°C to the reflux temperature of the solvent and is performed preferably at 0°C to 80°C . The reaction time varies depending on the starting materials
10 used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0123]

Step 3-4

This step is a step of hydrolyzing the ester moiety of the
15 compound (11) to produce the compound (1b). The compound (1b) can be produced by, for example, treating the compound (11) with a base in a solvent. Examples of the solvent used may include water, methanol, ethanol, propanol, 2-propanol, butanol, tetrahydrofuran, mixed solvents of them and the like. Examples of the base used may
20 include lithium hydroxide, sodium hydroxide, potassium hydroxide and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 70°C . The reaction time varies depending on the starting materials used, the solvent used, the reaction
25 temperature or the like, but is usually 30 minutes to 3 days.

[0124]

Synthesis of compound (1c)


The compound (1) when R^2 and R^3 are not H (this compound may hereinafter be referred to as the compound (1c)) can be produced from the compound (6) in accordance with any of methods described in scheme 4, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0125]

Scheme 4

[0126]

[Chemical Formula 19]



[0127]

In the above formulas, Ar^2 and Q^1 are as described above, $R^{2'}$ and $R^{3'}$ are each independently a C_{1-6} alkyl group optionally having substituent(s) or together form a C_2 to C_6 alkylene group, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0128]

Step 4-1

This step is a step of reacting the compound (6) with the compound (9) to produce the compound (12). The compound (12) can be produced by, for example, reacting the compound (6) with the compound (9) in a solvent in the presence of a base. Examples of the solvent used may include benzene, toluene, dimethyl sulfoxide, N,N-dimethylformamide, mixed solvents of them and the like.

Examples of the base used may include aliphatic amines such as piperidine, pyrrolidine, triethylamine, N-methylmorpholine and the like, aromatic amines such as pyridine, picoline, N,N-dimethyl aniline and the like, and organic acid salts such as piperidinium acetate, ammonium acetate and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 70°C to 110°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like. and is usually 30 minutes to 3 days.

Further, the compound (6) and the compound (9) used in this step may be commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0129]

Step 4-2

This step is a step of reacting the compound (12) with the compound (13) to produce the compound (14). The compound (14) can be produced by, for example, reacting the compound (12) with the

compound (13) in a solvent in the presence of a base. Examples of the solvent used may include toluene, benzene, tetrahydrofuran, acetonitrile, dimethyl sulfoxide, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the base used may include alumina-treated potassium fluoride (KF-Al₂O₃) and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0130]

Step 4-3

This step is a step of producing the compound (15) from the compound (14). The compound (15) can be produced in accordance with any of the method described in step 3-3 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0131]

Step 4-4

This step is a step of hydrolyzing the ester moiety of the compound (15) to produce the compound (1c). The compound (1c) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0132]

Synthesis of compound (3b)


The compound (3) when R³ is H (this compound may hereinafter be referred to as the compound (3b)) can be produced from the compound (8) in accordance with any of methods described in scheme 5, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0133]

Scheme 5

[0134]

10 [Chemical Formula 20]



[0135]

In the above formulas, Ar², R² and Q¹ are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

15 [0136]

Step 5-1

This step is a step of reacting the compound (8) with the compound

(16) to produce the compound (17). The compound (17) can be produced by, for example, reacting the compound (16) with the compound (8) in a solvent in the presence of a base. Examples of the solvent used may include tetrahydrofuran, hexane, mixed solvents of them and the like. Examples of the base used may include organolithiums such as lithium diisopropylamide (LDA) lithium hexamethyl disilazide (LHMDS) and the like. The reaction temperature can usually be performed at -78°C to the reflux temperature of the solvent and is performed preferably at -78°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

Further, the compound (8) and the compound (16) used in this step may be commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0137]

Step 5-2

This step is a step of converting the imine moiety of the compound (17) to produce the compound (18). The compound (18) can be produced by, for example, reacting the compound (17) with water in acetic acid and then reacting the obtained product with di-tert-butyl dicarbonate ((Boc)₂O) in a solvent in the presence of a base. Examples of the solvent used may include water, methanol, ethanol, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide,

1,4-dioxane, acetonitrile, mixed solvents of them and the like. Examples of the base used may include triethylamine, sodium hydrogen carbonate, sodium carbonate and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 70°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0138]

10 Step 5-3

This step is a step of producing the compound (19) from the compound (18). The compound (19) can be produced in accordance with any of the method described in step 3-3 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0139]

Step 5-4

This step is a step of deprotecting the tert-butoxycarbonyl (Boc) group in the compound (19) to produce the compound (3b). The compound (3b) can be produced by, for example, reacting the compound (19) with an acid such as trifluoroacetic acid (TFA) or hydrogen chloride in a solvent. Examples of the solvent used may include dichloromethane, dioxane, ethyl acetate, methanol, water, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and

is performed preferably at 0°C to 60°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

5 [0140]

Synthesis of compound (1d)


The compound (1) when R¹ is a C₁ to C₆ alkyl group (this compound may hereinafter be referred to as the compound (1d)) can be produced from the compound (20) in accordance with any of methods described in scheme 6, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0141]

Scheme 6

[0142]

15 [Chemical Formula 21]



In the above formulas, Ar², R², R³, and Q¹ are as described above, R^a is a C₁₋₆ alkyl group, W is an leaving group such as a chlorine atom, a bromine atom, an iodine atom, a methane sulfonyloxy group, a trifluoromethane sulfonyloxy group and the like, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0143]

Step 6-1

This step is a step of chlorinating the compound (20) to produce the compound (21). The compound (21) can be produced by, for example, reacting the compound (20) with sulfuryl chloride (SO_2Cl_2) in a solvent.

5 Examples of the solvent used may include tetrahydrofuran and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 70°C . The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the
10 like, but is usually 30 minutes to 3 days.

The compound (20) used in this step can be produced in accordance with any of methods described in scheme 3 or 4, methods similar thereto, methods described in other literatures, and methods similar thereto.

15 [0144]

Step 6-2

This step is a step of reacting the compound (21) with the compound (22) to produce the compound (23). The compound (23) can be produced by, for example, reacting the compound (21) with the
20 compound (22) in a solvent in the presence of a base. Examples of the solvent used may include tetrahydrofuran, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the base used may include alkali metal hydrides such as lithium hydride, sodium hydride and the like, alkali metal hydroxides such as sodium hydroxide,
25 potassium hydroxide and the like, and carbonates such as potassium

carbonate, cesium carbonate and the like. The reaction temperature can usually be performed at -10°C to the reflux temperature of the solvent and is performed preferably at 0°C to 60°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (22) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0145]

Step 6-3

This step is a step of dechlorinating the compound (23) to produce the compound (24). The compound (24) can be produced by, for example, reacting the compound (23) with zinc powder (Zn powder) in a solvent in the presence of acetic acid. Examples of the solvent used may include methanol, ethanol, water, tetrahydrofuran, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0146]

Step 6-4

This step is a step of hydrolyzing the ester moiety of the

compound (24) to produce the compound (1d). The compound (1d) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

5 [0147]

Synthesis of compound (1e)


The compound (1) when R^1 is a $-O-C_1$ to C_6 alkyl group (this compound may hereinafter be referred to as the compound (1e)) can be produced from the compound (14) in accordance with any of methods
10 described in scheme 7, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0148]

Scheme 7

[0149]

15 [Chemical Formula 22]



[0150]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , Q^1 , and W are as described above, and each carbon atom marked with an asterisk is an asymmetric
20 carbon atom.

[0151]

Step 7-1

This step is a step of producing the compound (25) from the compound (14). The compound (25) can be produced by, for example, reacting the compound (14) with zinc powder in a solvent in the presence of ammonium chloride. Examples of the solvent used may
5 include methanol, ethanol, water, tetrahydrofuran, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the
10 reaction temperature or the like, but is usually 10 minutes to 3 days.

[0152]

Step 7-2

This step is a step of reacting the compound (25) with the
15 compound (22) to produce the compound (26). The compound (26) can be produced in accordance with any of the method described in step 6-2 of scheme 6, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0153]

20 Step 7-3

This step is a step of hydrolyzing the ester moiety of the compound (26) to produce the compound (1e). The compound (1e) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other
25 literatures, and methods similar thereto.

[0154]

Synthesis of compound (1f)


The compound (1) when R^1 is a hydrogen atom and Ar^2 is a group represented by A7) (this compound may hereinafter be referred to as the compound (1f)) can be produced from the compound (27) in accordance with any of methods described in scheme 8, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0155]

10 Scheme 8

[0156]

[Chemical Formula 23]



[0157]

15 In the above formulas, R^2 , R^3 , R^7 , R^8 , Q^1 , and m are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0158]

Step 8-1

20 This step is a step of deprotecting the tert-butyldimethylsilyl (TBS) group in the compound (27) to produce the compound (28). The compound (28) can be produced by, for example, reacting the compound

(27) with an aqueous hydrogen chloride solution in a solvent. Examples of the solvent used may include tetrahydrofuran, water, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (27) used in this step can be produced in accordance with any of methods described in scheme 3 or 4, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0159]

Step 8-2

This step is a step of reducing the hydroxy group in the compound (28) to produce the compound (29). The compound (29) can be produced by, for example, hydrogenating the compound (28) in a solvent in the presence of a catalyst such as 10% palladium on carbon (10% Pd-C). Examples of the solvent used may include methanol, ethanol, dichloromethane, tetrahydrofuran, ethyl acetate, acetic acid, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3

days.

[0160]

Step 8-3

This step is a step of hydrolyzing the ester moiety of the
5 compound (29) to produce the compound (1f). The compound (1f) can
be produced in accordance with any of the method described in step
3-4 of scheme 3, methods similar thereto, methods described in other
literatures, and methods similar thereto.

[0161]

10 Synthesis of compounds (3c) to (3e)


The compound (3) when R¹ is a C₁ to C₆ alkyl group optionally
having substituent(s) (these compounds may hereinafter be referred
to as the compound (3c), the compound (3d), and the compound (3e))
can be produced from the compound (3a) in accordance with any of
15 methods described in scheme 9, methods similar thereto, methods
described in other literatures, and methods similar thereto.

[0162]

Scheme 9

[0163]

20 [Chemical Formula 24]



[0164]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , Q^1 , and W are as described above, R^b and R^c are each independently a hydrogen atom or a C_{1-6} alkyl group, Y^1 is a C_{1-3} alkylene, Y^2 is a single bond or a C_{1-3} alkylene group, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0165]

Step 9-1

This step is a step of protecting the amino group in the compound (3a) with a tert-butoxycarbonyl (Boc) group to produce the compound (30). The compound (30) can be produced by, for example, reacting the compound (3a) with di-tert-butyl dicarbonate ($(Boc)_2O$) in a solvent in the presence or absence of a base. Examples of the solvent used may include water, methanol, ethanol, tetrahydrofuran, N,N-dimethylformamide, 1,4-dioxane, acetonitrile, mixed solvents of them and the like. Examples of the base used may include trimethylamine, triethylamine, N-methylmorpholine and the like.

The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 60°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0166]

Step 9-2

This step is a step of reacting the compound (30) with the compound (22) to produce the compound (31). The compound (31) can be produced by, for example, reacting the compound (30) with the compound (22) in a solvent in the presence of a base. Examples of the solvent used may include tetrahydrofuran, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the base used may include lithium hydride, sodium hydride, potassium tert-butoxide and the like. The reaction temperature can usually be performed at -10°C to the reflux temperature of the solvent and is performed preferably at 0°C to 60°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

Further, the compound (22) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0167]

Step 9-3

This step is a step of deprotecting the tert-butoxycarbonyl (Boc) group in the compound (31) to produce the compound (3c). The compound (3c) can be produced in accordance with any of the method described in step 5-4 of scheme 5, methods similar thereto, methods
5 described in other literatures, and methods similar thereto.

[0168]

Step 9-4

This step is a step of reacting the compound (30) with the compound (32) to produce the compound (33). The compound (33) can
10 be produced in accordance with any of the method described in step 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods similar thereto.

Further, the compound (32) used in this step may be a commercially available, or can be produced in accordance with any
15 of methods described in other literatures, and methods similar thereto.

[0169]

Step 9-5

This step is a step of deprotecting the TBS and Boc groups
20 in the compound (33) to produce the compound (3d). The compound (3d) can be produced by, for example, reacting the compound (33) with an acid such as trifluoroacetic acid (TFA) or hydrogen chloride in a water-containing solvent. Examples of the solvent used may include mixtures of water with methanol, ethanol, dioxane, and so
25 on. The reaction temperature can usually be performed at -10°C to

the reflux temperature of the solvent and is performed preferably at 0°C to 60°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

5 [0170]

Step 9-6

This step is a step of reacting the compound (30) with the compound (34) to produce the compound (35). The compound (35) can be produced in accordance with any of the method described in step
10 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods similar thereto.

Further, the compound (34) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar
15 thereto.

[0171]

Step 9-7

This step is a step of deprotecting the Boc group in the compound (35) to produce the compound (3e). The compound (3e) can be produced
20 in accordance with any of the method described in step 5-4 of scheme 5, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0172]

Alternative synthesis of compounds (3c) and (3d)

25 The compounds (3c) and (3d) can be produced from the compound


(5) in accordance with any of methods described in scheme 10, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0173]

5 Scheme 10

[0174]

[Chemical Formula 25]



[0175]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , W , and Y^1 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0176]

Step 10-1

This step is a step of reacting the compound (5) with the compound (22) to produce the compound (36). The compound (36) can be produced in accordance with any of the method described in step 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods similar thereto.

The compound (22) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0177]

5 Step 10-2

This step is a step of deprotecting the benzyloxycarbonyl (Cbz) group in the compound (36) to produce the compound (3c). The compound (3c) can be produced in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described
10 in other literatures, and methods similar thereto.

[0178]

Step 10-3

This step is a step of reacting the compound (5) with the compound (32) to produce the compound (37). The compound (37) can be produced
15 in accordance with any of the method described in step 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods similar thereto.

The compound (32) used in this step may be a commercially available, or can also be produced in accordance with any of methods
20 described in other literatures, and methods similar thereto.

[0179]

Step 10-4

This step is a step of deprotecting the TBS group in the compound (37) to produce the compound (38). The compound (38) can be produced
25 by, for example, reacting the compound (37) with tetrabutylammonium

fluoride (TBAF) in a solvent. Examples of the solvent used may include tetrahydrofuran, N,N-dimethylformamide, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0180]

Step 10-5

This step is a step of deprotecting the Cbz group in the compound (38) to produce the compound (3d). The compound (3d) can be produced in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0181]

Synthesis of compounds (3f) to (3g)


The compounds represented by the following formula (3f) (the compound may hereinafter be referred to as the compound (3f)) and represented by the following formula (3g) (the compound may hereinafter be referred to as the compound (3g)) among the compound (3) can be produced from the compound (39) in accordance with any of methods described in scheme 11, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0182]

Scheme 11

[0183]

[Chemical Formula 26]



[0184]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , Q^1 , and W are as described
5 above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0185]

Step 11-1

This step is a step of reacting the compound (39) with the
10 compound (22) to produce the compound (40). The compound (40) can be produced by, for example, reacting the compound (39) with the compound (22) in a solvent in the presence of a base. Examples of the solvent used may include tetrahydrofuran and the like. Examples of the base used may include organolithiums such as lithium
15 diisopropylamide (LDA), lithium hexamethyl disilazide (LHMDS) and the like. The reaction temperature can usually be performed at -78°C to the reflux temperature of the solvent and is performed preferably at -78°C to 30°C . The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the

like, but is usually 30 minutes to 3 days.

The compound (39) used in this step can be produced in accordance with any of the method described in step 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods
5 similar thereto.

Further, the compound (22) used in this step may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

10 [0186]

Step 11-2

This step is a step of reducing the ester moiety of the compound (40) to produce the compound (41). The compound (41) can be produced by, for example, reacting the compound (40) with a reducing agent
15 such as lithium borohydride (LiBH_4), lithium aluminum hydride (LiAlH_4), diisobutylaluminium hydride (DIBAL) and the like in a solvent. Examples of the solvent used may include tetrahydrofuran and the like. The reaction temperature can usually be performed at -20°C to the reflux temperature of the solvent and is performed
20 preferably at 0°C to 30°C . The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0187]

Step 11-3

25 This step is a step of deprotecting the Boc group in the compound

(41) to produce the compound (3f). The compound (3f) can be produced in accordance with any of the method described in step 5-4 of scheme 5, methods similar thereto, methods described in other literatures, and methods similar thereto.

5 [0188]

Step 11-4

This step is a step of deprotecting the Boc group in the compound (40) to produce the compound (3g). The compound (3g) can be produced in accordance with any of the method described in step 5-4 of scheme 5, methods similar thereto, methods described in other literatures, and methods similar thereto.

10 5, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0189]

The compound (3g) can be produced from the compound (42) in accordance with any of methods described in scheme 12, methods similar thereto, methods described in other literatures, and methods similar thereto.


15 thereto, methods described in other literatures, and methods similar thereto.

[0190]

Scheme 12

[0191]

20 [Chemical Formula 27]



[0192]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , and Q^1 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0193]

5 Step 12-1

This step is a step of reacting the compound (22) with the compound (42) to produce the compound (43). The compound (43) can be produced in accordance with any of the method described in step 11-1 of scheme 11, methods similar thereto, methods described in
10 other literatures, and methods similar thereto.

The compound (42) used in this step can be produced in accordance with any of the method described in step 10-1 of scheme 10, methods similar thereto, methods described in other literatures, and methods similar thereto.

15 [0194]

Step 12-2

This step is a step of deprotecting the Cbz group in the compound (43) to produce the compound (3g). The compound (3g) can be produced in accordance with any of the method described in step 2-4 of scheme
20 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0195]

Synthesis of compounds (3h) and (3i)

The compounds represented by the following formula (3h) (the
25 compound may hereinafter be referred to as the compound (3h)) and


represented by the following formula (3i) (the compound may hereinafter be referred to as the compound (3i)) among the compound (3) can be produced from the compound (5) in accordance with any of methods described in scheme 13, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0196]

Scheme 13

[0197]

[Chemical Formula 28]



[0198]

In the above formulas, Ar^2 , R^2 , R^3 , R^a , R^b , R^c , Q^1 , W , and Y^2 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0199]

Step 13-1

This step is a step of reacting the compound (5) with the compound (34) to produce the compound (44). The compound (44) can be produced in accordance with any of the method described in step 9-2 of scheme 9, methods similar thereto, methods described in other literatures, and methods similar thereto.

The compound (34) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0200]

Step 13-2

This step is a step of hydrolyzing the ester moiety of the compound (44) to produce the compound (45). The compound (45) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0201]

Step 13-3

This step is a step of reacting the compound (45) with the compound (46) to produce the compound (47). The compound (47) can be produced by, for example, reacting the compound (45) with the compound (46) in a solvent in the presence or absence of a base and in the presence of a condensing agent. Examples of the solvent used may include N,N-dimethylformamide, dichloromethane, 1,4-dioxane, tetrahydrofuran, mixed solvents of them and the like.

Examples of the condensing agent used may include

1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI),

dicyclohexylcarbodiimide (DCC),

2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium

5 hexafluorophosphate methanaminium (HATU) and the like. If

required, N,N-dimethylamino pyridine, pyridine,

1-hydroxybenzotriazole (HOBt) and the like can be used as a reaction

promotor. The reaction temperature can usually be performed at 0°C

to the reflux temperature of the solvent and is performed preferably

10 at 0°C to 30°C. Examples of the usable base may include potassium

carbonate, sodium carbonate, triethylamine,

N,N-diisopropylethylamine and the like. The reaction time varies

depending on the starting materials used, the solvent used, the

reaction temperature or the like, but is usually 30 minutes to 3

15 days.

The compound (46) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0202]

20 Step 13-4

This step is a step of producing the compound (48) from the compound (47). The compound (48) can be produced by, for example, reacting the compound (47) with triphenylphosphine, triethylamine,

and 1,1,1,2,2,2-hexachloroethane in a solvent. Examples of the

25 solvent used may include dichloromethane and the like. The reaction

temperature can usually be performed at -78°C to the reflux temperature of the solvent and is performed preferably at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0203]

Step 13-5

This step is a step of deprotecting the Cbz group in the compound (48) to produce the compound (3h). The compound (3h) can be produced in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0204]

Step 13-6

This step is a step of reacting the compound (45) with the compound (49) to produce the compound (50). The compound (50) can be produced in accordance with any of the method described in step 13-3 of scheme 13, methods similar thereto, methods described in other literatures, and methods similar thereto.

The compound (49) used in this step may be a commercially available, or can also be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0205]

Step 13-7

This step is a step of oxidizing the alcohol moiety of the

compound (50) to a ketone to produce the compound (51). The compound (51) can be produced by, for example, reacting the compound (50) with an oxidizer such as 2-iodoxybenzoic acid in a solvent. Examples of the solvent used may include dimethyl sulfoxide and the like.

5 The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

10 [0206]

Step 13-8

This step is a step of producing the compound (52) from the compound (51). The compound (52) can be produced by, for example, reacting the compound (51) with triphenylphosphine and triethylamine
15 in a solvent. Examples of the solvent used may include carbon tetrachloride and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 50°C to 90°C. The reaction time varies depending on the starting materials used, the solvent used, the
20 reaction temperature or the like, but is usually 30 minutes to 3 days.

[0207]

Step 13-9

This step is a step of deprotecting the Cbz group in the compound
25 (52) to produce the compound (3i). The compound (3i) can be produced

in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0208]

5 Synthesis of compound (3j)


The compound represented by the following formula (3j) (the compound may hereinafter be referred to as the compound (3j)) among the compound (3) can be produced from the compound (53) in accordance with any of methods described in scheme 14, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0209]

Scheme 14

[0210]

[Chemical Formula 29]



[0211]

In the above formulas, Ar^2 , R^2 , R^3 , and R^a are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0212]

Step 14-1

This step is a step of converting the sulfanyl group in the

compound (53) to a sulfonyl group to produce the compound (54). The compound (54) can be produced by, for example, treating the compound (53) with an oxidizer such as meta-chloroperoxybenzoic acid (m-CPBA) in a solvent. Examples of the solvent used may include
5 dichloromethane, acetonitrile, tetrahydrofuran, mixed solvents of them and the like. The amount of the oxidizer used is usually about 2 to 5 molar equivalents with respect to 1 mole of the compound (53). The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably
10 at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (53) used in this step can be produced in accordance with any of the method described in step 10-1 of scheme 10, methods
15 similar thereto, methods described in other literatures, and methods similar thereto.

[0213]

Step 14-2

This step is a step of deprotecting the Cbz group in the compound
20 (54) to produce the compound (3j). The compound (3j) can be produced in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0214]

25 Synthesis of compounds (Ic), (Id), and (Ie)

The compounds represented by the following formulas (Ic), (Id), and (Ie) (the compounds may hereinafter be referred to as the compounds (Ic), (Id), and (Ie)) among the compound (I) can be produced from the compound (3e) in accordance with any of methods described in


5 scheme 15, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0215]

Scheme 15

[0216]

10 [Chemical Formula 30]



[0217]

In the above formulas, Ar¹, Ar², R², R³, R^b, R^c, Y², and Q¹ are as described above, Q² and Q³ are each independently a hydrogen atom or a C₁₋₆ alkyl group optionally having substituent(s) or Q² and Q³

15 together form an alkylene group, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0218]

Step 15-1

This step is a step of reacting the compound (3e) with the

compound (4) to produce the compound (Ic). The compound (Ic) can be produced in accordance with any of the method described in step 1-2 of scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

5 [0219]

Step 15-2

This step is a step of hydrolyzing the ester moiety of the compound (Ic) to produce the compound (Id). The compound (Id) can be produced in accordance with any of the method described in step 10 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0220]

Step 15-3

This step is a step of reacting the compound (Id) with the 15 compound (55) to produce the compound (Ie). The compound (Ie) can be produced in accordance with any of the method described in step 13-3 of scheme 13, methods similar thereto, methods described in other literatures, and methods similar thereto.

Further, the compound (55) used in this step may be a 20 commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

[0221]

Synthesis of compounds (If), (Ig), (Ih), and (Ii)

25 The compounds represented by the following formulas (If), (Ig),


(Ih), and (Ii) (the compounds may hereinafter be referred to as the compounds (If), (Ig), (Ih), and (Ii)) among the compound (I) of the embodiment can be produced from the compound (56) in accordance with any of methods described in scheme 16, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0222]

Scheme 16

[0223]

[Chemical Formula 31]



[0224]

In the above formulas, Ar¹, R², R³, R⁷, R⁸, Q¹, Q², and Q³ are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0225]

Step 16-1

This step is a step of protecting the phenolic hydroxy group in the compound (56) with a benzyl (Bn) group to produce the compound (57). The compound (57) can be produced by, for example, reacting the compound (56) with benzyl bromide in a solvent in the presence of a base. Examples of the solvent used may include tetrahydrofuran, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the base used may include sodium carbonate, potassium carbonate, cesium carbonate and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (56) used in this step can be produced in accordance with any of methods described in scheme 3 or 4, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0226]

Step 16-2

This step is a step of hydrolyzing the ester moiety of the compound (57) to produce the compound (58). The compound (58) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0227]

Step 16-3

This step is a step of producing the compound (59) from the compound (58). The compound (59) can be produced in accordance with
5 any of the method described in step 1-1 of scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0228]

Step 16-4

10 This step is a step of deprotecting the Bn group in the compound (59) to produce the compound (If). The compound (If) can be produced by, for example, hydrogenating the compound (59) in a solvent in the presence of a catalyst such as 10% palladium on carbon (10% Pd-C) and the like. Examples of the solvent used may include methanol,
15 ethanol, dichloromethane, tetrahydrofuran, ethyl acetate, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the
20 reaction temperature or the like, but is usually 30 minutes to 3 days.

[0229]

Step 16-5

This step is a step of trifluoromethane-sulfonylating the
25 hydroxy group in the compound (If) to produce the compound (60).

The compound (60) can be produced by, for example, reacting the compound (If) with a trifluoromethane-sulfonylation agent in a solvent in the presence of a base. Examples of the solvent used may include dichloromethane, 1,2-dichloroethane, pyridine, tetrahydrofuran, N,N-dimethylformamide, 1,4-dioxane, acetonitrile, diethyl ether, mixed solvents of them and the like. Examples of the trifluoromethane-sulfonylation agent may include N-phenyl bis(trifluoromethanesulfonimide), trifluoromethanesulfonic anhydride and the like. Examples of the base used may include potassium carbonate, sodium carbonate, sodium hydride, potassium phosphate, N,N-diisopropylethylamine, triethylamine, 2,6-lutidine and the like. The reaction temperature can usually be performed at -78°C to the reflux temperature of the solvent and is performed preferably at -20°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0230]

Step 16-6

This step is a step of producing the compound (Ig) from the compound (60). The compound (Ig) can be produced by, for example, reacting the compound (60) with the compound (55) in a solvent in the presence of acetonitrile, cesium carbonate, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP), palladium acetate(II), and molybdenum hexacarbonyl ($\text{Mo}(\text{CO})_6$). Examples of the solvent used may include toluene and the like. The reaction

temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 50°C to 80°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0231]

Step 16-7

This step is a step of producing the compound (Ih) from the compound (60). The compound (Ih) can be produced by, for example, reacting the compound (60) with zinc cyanide ($\text{Zn}(\text{CN})_2$) in a solvent using a palladium reagent as a catalyst. Examples of the solvent used may include 1,4-dioxane, toluene, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the palladium reagent used may include tetrakis(triphenylphosphine) palladium (0) ($\text{Pd}(\text{PPh}_3)_4$) and the like. The reaction temperature can usually be performed at 20°C to the reflux temperature of the solvent and is performed preferably at 60°C to 100°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0232]

Step 16-8

This step is a step of hydrolyzing the cyano group in the compound (Ih) to an amido group to produce the compound (Ii). The compound (Ii) can be produced by, for example, reacting the compound (Ih)

in a solvent in the presence of a base. Examples of the solvent used may include water, methanol, ethanol, propanol, 2-propanol, butanol, tetrahydrofuran, mixed solvents of them and the like. Examples of the base used may include lithium hydroxide, sodium
5 hydroxide, potassium hydroxide and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 60°C to 100°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually
10 30 minutes to 3 days.

[0233]

Synthesis of compounds (Ij), (Ik), and compound (Il)


The compound represented by the following formula (Ij) (the compound may hereinafter be referred to as the compound (Ij)) among
15 the compound (I) of the embodiment can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

Furthermore, the compounds represented by the following formulas (Ik) and (Il) (the compounds may hereinafter be referred
20 to as the compounds (Ik) and (Il)) among the compound (I) of the embodiment can be produced from the compound (Ij) in accordance with any of methods described in scheme 17, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0234]

25 Scheme 17

[Chemical Formula 32]



[0237]

10 This step is a step of converting the sulfanyl group in the compound (Ij) to a sulfinyl group to produce the compound (Ik). The compound (Ik) can be produced by, for example, treating the compound (Ij) with an oxidizer such as meta-chloroperoxybenzoic acid (m-CPBA) in a solvent. Examples of the solvent used may include
15 dichloromethane, acetonitrile, tetrahydrofuran, mixed solvents of them and the like. The amount of the oxidizer used is usually about 1 to 2 molar equivalents with respect to 1 mole of the compound (Ij). The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably
20 at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0238]

Step 17-2

This step is a step of converting the sulfanyl group in the compound (Ij) to a sulfonyl group to produce the compound (Il).

5 The compound (Il) can be produced by, for example, treating the compound (Ij) with an oxidizer such as meta-chloroperoxybenzoic acid (m-CPBA) in a solvent. Examples of the solvent used may include dichloromethane, acetonitrile, tetrahydrofuran, mixed solvents of them and the like. The amount of the oxidizer used is usually about
10 2 to 5 molar equivalents with respect to 1 mole of the compound (Ij). The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the
15 like, but is usually 30 minutes to 3 days.

[0239]

Synthesis of compounds (Im), (In), compound (Io), and compound (Ip)

The compound represented by the following formula (Im) (the compound may hereinafter be referred to as the compound (Im)) among
20 the compound (I) of the embodiment can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

Furthermore, the compounds represented by the following formulas (In), (Io) and (Ip) (the compounds may hereinafter be
25 referred to as the compounds (In), (Io) and (Ip)) among the compound


(I) of the embodiment can be produced from the compound (Im) in accordance with any of methods described in scheme 18, methods similar thereto, methods described in other literatures, and methods similar thereto.

5 [0240]

Scheme 18

[0241]

[Chemical Formula 33]



10 [0242]

In the above formulas, Ar^2 , R^1 , R^2 , R^3 , R^{13} , Q^1 , Q^2 , and Q^3 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0243]

15 Step 18-1

This step is a step of reducing the ester moiety of the compound (Im) to produce the compound (In). The compound (In) can be produced by, for example, treating the compound (Im) with a reducing agent

such as diisobutylaluminum hydride in a solvent. Examples of the solvent used may include dichloromethane, tetrahydrofuran, toluene, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0244]

10 Step 18-2

This step is a step of hydrolyzing the ester moiety of the compound (Im) to produce the compound (Io). The compound (Io) can be produced in accordance with any of the method described in step 3-4 of scheme 3, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0245]

Step 18-3

This step is a step of reacting the compound (Io) with the compound (55) to produce the compound (Ip). The compound (Ip) can be produced in accordance with any of the method described in step 13-3 of scheme 13, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0246]

Synthesis of compound (Iq)

25 The compound represented by the following formula (Iq) (the


compound may hereinafter be referred to as the compound (Iq)) among the compound (I) of the embodiment can be produced from the compound (61) in accordance with any of method described in scheme 19, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0247]

Scheme 19

[0248]

[Chemical Formula 34]



[0249]

In the above formulas, Ar^1 , Ar^2 , R^2 , and R^3 are as described above, R^f is a C_{1-6} alkyl group, a C_{1-6} alkoxy group, or a TBSO group, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0250]

Step 19-1

This step is a step of producing the compound (Iq) from the compound (61). The compound (Iq) can be produced in accordance with any of the method described in step 1-2 of scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0251]


The compound (61) can be produced from the compound (5) in accordance with any of methods described in scheme 20, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0252]

Scheme 20

[0253]

[Chemical Formula 35]



[0254]

In the above formulas, Ar^2 , R^2 , R^3 , and R^f are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0255]

Step 20-1

This step is a step of producing the compound (62) from the compound (5). The compound (62) can be produced by, for example, reacting the compound (5) with triethyloxonium hexafluorophosphate (Et_3OPF_6) in a solvent. Examples of the solvent used may include dichloromethane and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and

is performed preferably at 10°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

5 [0256]

Step 20-2

This step is a step of reacting the compound (62) with the compound (63) to produce the compound (64). The compound (64) can be produced by, for example, reacting the compound (62) with the
10 compound (63) in a solvent in the presence or absence of an acid or a base. Examples of the solvent used may include methanol, ethanol, toluene, tetrahydrofuran, acetonitrile, dioxane, mixed solvents of them and the like. Examples of the acid that may be used may include hydrogen chloride, ammonium chloride and the like. Examples
15 of the base that may be used may include trimethylamine, triethylamine, N-methylmorpholine and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 0°C to 50°C. The reaction time varies depending on the starting materials used, the solvent used, the
20 reaction temperature or the like, but is usually 10 minutes to 3 days.

Further, the compound (63) may be a commercially available, or can be produced in accordance with any of methods described in other literatures, and methods similar thereto.

25 [0257]

Step 20-3

This step is a step of deprotecting the Cbz group in the compound (64) to produce the compound (61). The compound (61) can be produced in accordance with any of the method described in step 2-4 of scheme 2, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0258]

Synthesis of compound (Ir)


The compound (Ir) can be produced from the compound (65) in accordance with any of methods described in scheme 21, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0259]

Scheme 21

[0260]

[Chemical Formula 36]



[0261]

In the above formulas, Ar^1 , Ar^2 , R^2 , and R^3 are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0262]

Step 21-1

This step is a step of deprotecting the TBS group in the compound (65) to produce the compound (Ir). The compound (Ir) can be produced by, for example, reacting the compound (65) with an acid such as trifluoroacetic acid (TFA) or hydrogen chloride in a solvent mixture of water-dioxane.

The reaction temperature can usually be performed at -20°C to the reflux temperature of the solvent and is performed preferably at 0°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

The compound (65) used in this step can be produced in accordance with any of methods described in scheme 19, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0263]

Step 22

Synthesis of compounds (Is) and (It)

The compound represented by the following formula (Is) (the compound may hereinafter be referred to as the compound (Is)) among the compound (I) of the embodiment can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

Furthermore, the compound represented by the following formula (It) (the compound may hereinafter be referred to as the compound (It)) among the compound (I) of the embodiment can be produced,


for example, from the compound (Is) in accordance with any of methods described in scheme 22, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0264]

5 Scheme 22

[0265]

[Chemical Formula 37]



[0266]

10 In the above formulas, Ar^2 , R^1 , R^2 , R^3 , and R^{13} are as described above, X is a halogen atom, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0267]

Step 22-1

15 This step is a step of reacting the compound (Is) with trimethylsilylacetylene (66) to produce the compound (67). The compound (67) can be produced by, for example, reacting the compound (Is) with trimethylsilylacetylene (66) in a solvent in the presence of dichlorobis(triphenylphosphine)palladium(II), copper(I) iodide,

20 and triethylamine. Examples of the solvent used may include N,N-dimethylformamide and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 40°C to 60°C. The reaction time varies

depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0268]

5 Step 22-2

This step is a step of deprotecting the TMS group in the compound (67) to produce the compound (It). The compound (It) can be produced by, for example, reacting the compound (67) with potassium carbonate in a solvent. Examples of the solvent used may include methanol
10 and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

15 [0269]

Synthesis of compound (Iu)


The compound represented by the following formula (Iu) (the compound may hereinafter be referred to as the compound (Iu)) among the compound (I) of the embodiment can be produced, for example,
20 from the compound (68) in accordance with any of method described in scheme 23, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0270]

Scheme 23

25 [0271]

[Chemical Formula 38]



[0272]

In the above formulas, Ar¹, R¹, R², R³, R⁷, R⁸, and R^a are as
 5 described above, and each carbon atom marked with an asterisk is
 an asymmetric carbon atom.

[0273]

Step 23-1

This step is a step of deprotecting the Boc group in the compound
 10 (68) to produce the compound (Iu). The compound (Iu) can be produced
 in accordance with any of the method described in step 5-4 of scheme
 5, methods similar thereto, methods described in other literatures,
 and methods similar thereto.

The compound (68) used in this step can be produced in accordance
 15 with any of methods described in scheme 1, methods similar thereto,
 methods described in other literatures, and methods similar thereto.

[0274]

Synthesis of compounds (Iv) and (Iw)

The compound represented by the following formula (Iv) (the
 20 compound may hereinafter be referred to as the compound (Iv)) among
 the compound (I) of the embodiment can be produced in accordance
 with any of methods described in scheme 1, methods similar thereto,

methods described in other literatures, and methods similar thereto.


Furthermore, the compound represented by the following formula (Iw) (the compound may hereinafter be referred to as the compound (Iw)) among the compound (I) of the embodiment can be produced, for example, from the compound (Iv) in accordance with any of methods described in scheme 24, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0275]

Scheme 24

10 [0276]

[Chemical Formula 39]



[0277]

In the above formulas, Ar¹, R¹, R², R³, R⁷, R⁸, and m are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0278]

Step 24-1

This step is a step of converting the hydroxy group in the compound (Iv) to a ketone to produce the compound (Iw). The compound (Iw) can be produced by, for example, treating the compound (Iv) with an oxidizer such as Dess-Martin periodinane in a solvent. Examples of the solvent used may include dichloromethane,

acetonitrile, mixed solvents of them and the like. The amount of the oxidizer used is usually about 1 to 2 molar equivalents with respect to 1 mole of the compound. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 30°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

[0279]

10 Step 25

Synthesis of compound (Ix)


The compound represented by the following formula (Ix) (the compound may hereinafter be referred to as the compound (Ix)) among the compound (I) of the embodiment can be produced, for example, from the compound (69) in accordance with any of method described in scheme 25, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0280]

Scheme 25

20 [0281]

[Chemical Formula 40]



[0282]

In the above formulas, Ar¹, R¹, R², R³, R⁷, R⁸, and m are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

5 [0283]

Step 25-1

This step is a step of deprotecting the Bn group in the compound (69) to produce the compound (Ix). The compound (Ix) can be produced by, for example, hydrogenating the compound (69) in a solvent in
10 the presence of a catalyst such as 10% palladium on carbon (10% Pd-C). Examples of the solvent used may include methanol, ethanol, dichloromethane, tetrahydrofuran, ethyl acetate, N,N-dimethylformamide, mixed solvents of them and the like. Examples of the base used may include triethylamine acetate and
15 the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 20°C to 40°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

20 The compound (69) used in this step can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0284]

Synthesis of compound (Iz)

25 The compound represented by the following formula (Iy) (the

compound may hereinafter be referred to as the compound (Iy)) among the compound (I) of the embodiment can be produced in accordance with any of methods described in scheme 1, methods similar thereto, methods described in other literatures, and methods similar thereto.


The compound represented by the following formula (Iz) (the compound may hereinafter be referred to as the compound (Iz)) among the compound (I) of the embodiment can be produced, for example, from the compound (Iy) in accordance with any of methods described in scheme 26, methods similar thereto, methods described in other literatures, and methods similar thereto.

[0285]

Scheme 26

[0286]

[Chemical Formula 41]



[0287]

In the above formulas, Ar¹, Ar², R², R³, and Y¹ are as described above, and each carbon atom marked with an asterisk is an asymmetric carbon atom.

[0288]

Step 26-1

This step is a step of producing the compound (Iz) from the compound (Iy). The compound (Iz) can be produced by, for example,

reacting the compound (Iy) with sodium azide in a solvent in the presence of triethylamine and acetic acid. Examples of the solvent used may include toluene, N,N-dimethylformamide, isopropanol, water, mixed solvents of them and the like. The reaction temperature can usually be performed at 0°C to the reflux temperature of the solvent and is performed preferably at 100°C to 130°C. The reaction time varies depending on the starting materials used, the solvent used, the reaction temperature or the like, but is usually 30 minutes to 3 days.

10 [0289]

A pharmacologically acceptable salt of the compound (I) of the present embodiment can be produced using the compound (I) of the present embodiment according to a usual method.

[0290]

15 The schemes described above are examples of the method of producing the compound (I) of the present embodiment or a production intermediate thereof. These schemes can be modified to various schemes that can be readily understood by a person skilled in the art.

20 [0291]

Also, in the case that there is a need of a protective group according to the kind of the functional group, an appropriate combination of introduction and removal procedures may be performed according to a usual method. For the types of protective groups and introduction and removal of the protective groups, see, for

example, methods described in "Greene's Protective Groups in Organic Synthesis," Theodora W. Green & Peter G. M. Wuts, ed., fourth edition, Wiley-Interscience, 2006.

[0292]

5 The intermediates used for preparation of the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof can be isolated/purified, as necessary, by solvent extraction, crystallization, recrystallization, chromatography, or preparative high-performance liquid chromatography or the like,
10 that is an isolation/purification means well-known to a skilled person in the art.

[0293]

 The term "FPRL1 agonist effect" used in the present embodiment means that agonist activity exhibits through the action on formyl
15 peptide receptor like 1 (FPRL1).

 As described above, it is known that LXA4 and peptides reported as endogenous agonists of FPRL1 contribute to resolution of inflammation.

[0294]

20 The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof exhibits superior agonist activity in, for example, a test of calcium influx into FPRL1-overexpressing cells. Therefore, the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof
25 is useful as a therapeutic or prophylactic agent for inflammatory

diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, immune disorders and the like.

5 The compound (I) of the present embodiment or a pharmacologically acceptable salt thereof can also be used to produce pharmaceuticals for treatment or prevention of inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation,
10 nervous disorders, pains, prion diseases, amyloidosis, immune disorders and the like.

[0295]

 In addition, pharmaceuticals containing, as an active ingredient, the compound (I) of the present embodiment or a
15 pharmacologically acceptable salt thereof can be used as, for example, prophylactic or therapeutic agents for various disease states associated with the FPRL1 receptor (such as Behcet's disease, Sweet disease, systemic lupus erythematosus (SLE), Wegener's granulomatosis, virus infection, diabetes, amputations, cancers,
20 bacterial infection, physical external injuries, physical disorders including exposure to radiation, vasoconstriction, anaphylactic reactions, allergic reactions, rhinitis, shocks (endotoxic, hemorrhagic, traumatic, splanchnic ischemia, and circulatory shocks), rheumatoid arthritis, gout, psoriasis, benign prostatic
25 hyperplasia, myocardial ischemia, myocardial infarction, brain

injuries, pulmonary diseases, COPD, COAD, COLD, acute lung injury, acute respiratory distress syndrome, chronic bronchitis, pulmonary emphysema, asthma (allergic asthma and non-allergic asthma), cystic pulmonary fibrosis, nephropathy, renal glomerular diseases, ulcerative colitis, IBD, Crohn's disease, periodontitis, pains, Alzheimer's disease, AIDS, uveitic glaucoma, conjunctivitis, Sjogren's syndrome, rhinitis and the like).

[0296]

Pharmaceutical containing the compound (I) of the present embodiment or pharmacologically acceptable salt thereof

A pharmaceutical containing, as an active ingredient, the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof can have various forms according to the usages. Examples of the forms may include powders, granules, fine granules, dry syrups, tablets, capsules, injections, liquids, ointments, suppositories, patches, sublingual tablets and the like, which are administered orally or parenterally.

[0297]

Such a pharmaceutical can be formed as a pharmaceutical composition containing, as an active ingredient, the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof and a pharmacologically acceptable additive using a well-known method according to the form of the pharmaceutical. Examples of the additive contained in the pharmaceutical composition may include an excipient, a disintegrant, a binder, a lubricant,

a diluent, a buffering agent, an isotonicizing agent, an antiseptic, a humectant, an emulsifier, a dispersant, a stabilizer, a solubilizing agent and the like. The pharmaceutical composition can be prepared by appropriately mixing the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof with an additive or by diluting the compound (I) or a pharmacologically acceptable salt thereof with an additive and dissolving it in the additive. When the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof is used in combination with agents other than the FPRL1 receptor agonist, a pharmaceutical composition can be produced by forming active ingredients of these components into a formulation simultaneously or separately in the manner described above.

[0298]

The pharmaceutical according to the present embodiment can be systemically or locally administered orally or parenterally (transnasally, pulmonary, intravenously, intrarectally, hypodermically, intramuscularly, percutaneously and the like).

[0299]

When a pharmaceutical composition containing, as an active ingredient, the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof is used for practical treatment, the dose of the compound (I) of the present embodiment or the pharmacologically acceptable salt thereof used as the active ingredient is appropriately determined according to the age, sex,

and body weight of the patient, the disease of the patient, the degree of the treatment and the like. For example, in the case of oral administration, it may be appropriately administered to an adult (the body weight is assumed to be 60 kg) at a daily dose within the range of about 0.03 to about 1,000 mg/body in one portion or several divided portions. The dose per day as an oral administration is preferably 0.06 to 540 mg/body and more preferably 0.18 to 180 mg/body. In the case of parenteral administration, it may be appropriately administered to an adult at a daily dose within the range of about 0.01 to about 300 mg/body in one portion or several divided portions. The dose per day as a parenteral administration is preferably 0.01 to 100 mg/body and more preferably 0.06 to 60 mg/body. The dose of the compound (I) of the present embodiment or a pharmacologically acceptable salt thereof may be reduced according to the dose of agents other than the FPRL1 receptor agonist.

Examples

[0300]

Hereinafter, the present invention will be described in more detail on the basis of Test Examples, Examples, and Reference Examples. Starting materials used in production of the compound (I) include a novel compound, and therefore Production examples for the starting materials will be also described as Reference Examples. The present invention is not limited to compounds described in the following Examples, and may be modified without departing from the scope of the present invention.

[0301]

Among symbols used in each Reference Example, each Example, and each Table, Ref.No. represents Reference Example Number, Ex.No. represents Example Number, P.D. represents physical chemical data, 5 Str. represents a structural formula, and ^1H -NMR represents a proton nuclear magnetic resonance spectrum. CDCl_3 represents chloroform-d, and DMSO-d_6 represents dimethyl sulfoxide- d_6 . $\text{MS}(\text{ESI}^+)$ represents mass spectral data measured by electron-spray ionization. An optical rotation represents a specific optical 10 rotation, which measured in described solvent at described concentration and temperature using sodium D-line as light source.

[0302]

Wedge-shaped solid line and dashed line in a structural formula represent relative configuration in an optically active 15 substance, but do not represent absolute configuration. Thick solid line and dashed line represent relative configuration in a racemate and an optically active substance obtained by resolution of a racemate. A carbon atom marked with "*" represents an asymmetric carbon. A wavy line bond of carbon atom marked with 20 "*" represents the presence of a racemate.

[0303]

Both R^* and S^* in the name of a compound represent relative steric configuration about an asymmetric carbon atom.

When both a substituent and a hydrogen atom are bonded to 25 each of two positions of a pyrrolidine ring in a structural formula,

the relative configuration of the substituents is expressed as cis or trans, and cis or trans is sometimes followed by a hyphen and the name of a compound.

When the pyrrolidine ring is considered as a face, cis means that the two adjacent substituents are on the same side, and trans means that the two adjacent substituents are on the respective opposite sides.


[0304]

In order to represent isomers about a double bond and a double bond of imine in the name of a compound, a cis-isomer is expressed as "Z," and a trans-isomer is expressed as "E."

[0305]

<Reference Example 1-1>

[Chemical Formula 42]



[0306]

(E)-2-Fluoro-1-methoxy-4-(2-nitrovinyl)benzene

3-Fluoro-4-methoxybenzaldehyde (500 mg) was dissolved in 2-hydroxyethylammonium formate (1.7 g) under an argon atmosphere, and nitromethane (175 μ L) was added to produce a reaction solution. The reaction solution was stirred at room temperature for 9 hours. Water was added to the reaction solution, and the precipitated solid was collected by filtration, and washed with water. The resulting solid was dried to obtain the title compound as a yellow

solid (607 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.96 (3H, s), 7.01 (1H, t, $J = 8.5$ Hz), 7.27-7.32 (2H, m), 7.48 (1H, d, $J = 13.3$ Hz), 7.92 (1H, d, $J = 13.3$ Hz).

5 [0307]





The following Reference Examples 1-2 to 1-27 were obtained using each corresponding aromatic aldehyde in the same method as in Reference Example 1-1.

The structures and spectral data thereof are shown in Tables

10 1 to 5.







[0308]

[Table 1]

Ref.No	Str.	Chemical name	P.D.
1-2		(E)-4-methoxy-2-methyl-1-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.47 (3H, s), 3.85 (3H, s), 6.77-6.81 (2H, m), 7.46-7.51 (2H, m), 8.27 (1H, d, $J = 13.4$ Hz).
1-3		(E)-2-fluoro-4-methoxy-1-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.87 (3H, s), 6.71 (1H, dd, $J = 12.1, 2.4$ Hz), 6.78 (1H, dd, $J = 8.5, 2.4$ Hz), 7.43 (1H, t, $J = 8.5$ Hz), 7.66 (1H, d, $J = 13.9$ Hz), 8.02 (1H, d, $J = 13.9$ Hz).
1-4		(E)-2-chloro-4-methoxy-1-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.87 (3H, s), 6.87 (1H, dd, $J = 8.5, 2.4$ Hz), 7.02 (1H, d, $J = 2.4$ Hz), 7.53 (1H, d, $J = 8.5$ Hz), 7.56 (1H, d, $J = 13.9$ Hz), 8.38 (1H, d, $J = 13.9$ Hz).
1-5		(E)-1,3-difluoro-5-methoxy-2-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.87 (3H, s), 6.54-6.59 (2H, m), 7.77 (1H, d, $J = 13.4$ Hz), 8.11 (1H, d, $J = 13.4$ Hz).






[0309]

[Table 2]

Ref.No	Str.	Chemical name	P.D.
1-6		(E)-1,4-difluoro-2-methoxy-5-(2-nitrovinyl)benzene	¹ H-NMR (400 MHz, CDCl ₃) δ 3.95 (3H, s), 6.79 (1H, dd, J = 11.0, 6.7 Hz), 7.21 (1H, dd, J = 11.0, 6.7 Hz), 7.61 (1H, d, J = 13.4 Hz), 7.98 (1H, d, J = 13.4 Hz).
1-7		(E)-1,3-difluoro-2-methoxy-5-(2-nitrovinyl)benzene	¹ H-NMR (400 MHz, CDCl ₃) δ 4.10 (3H, t, J=1.8 Hz), 7.07-7.15 (2H, m), 7.46 (1H, d, J=13.4 Hz), 7.84 (1H, d, J = 13.4 Hz).
1-8		(E)-5-(2-nitrovinyl)-2,3-dihydrobenzofuran	¹ H-NMR (400 MHz, CDCl ₃) δ 3.26 (2H, t, J=8.8 Hz), 4.67 (2H, t, J = 8.8 Hz), 6.84 (1H, d, J = 8.3 Hz), 7.35 (1H, d, J = 8.3 Hz), 7.41 (1H, s), 7.51 (1H, d, J=13.4 Hz), 7.97 (1H, d, J = 13.4 Hz).
1-9		(E)-6-fluoro-5-(2-nitrovinyl)-2,3-dihydrobenzofuran	¹ H-NMR (400 MHz, CDCl ₃) δ 3.23 (2H, t, J=8.9 Hz), 4.72 (2H, t, J = 8.9 Hz), 6.60 (1H, d, J = 11.6 Hz), 7.30 (1H, d, J = 6.7 Hz), 7.62 (1H, d, J=13.4 Hz), 8.03 (1H, d, J=13.4 Hz).
1-10		(E)-6-fluoro-5-(2-nitrovinyl)chromane	¹ H-NMR (400 MHz, CDCl ₃) δ 2.00-2.05 (2H, m), 2.77 (2H, t, J = 6.4 Hz), 4.24 (2H, t, J = 5.5 Hz), 6.59 (1H, d, J = 11.6 Hz), 7.17 (1H, d, J = 8.6 Hz), 7.64 (1H, d, J=13.4 Hz), 7.97 (1H, d, J = 13.4 Hz).
1-11		(E)-3-fluoro-5-methoxy-2-(2-nitrovinyl)pyridine	¹ H-NMR (400 MHz, CDCl ₃) δ 3.93 (3H, s), 6.98 (1H, dd, J=11.0, 2.4 Hz), 7.93 (1H, d, J=13.4 Hz), 8.13 (1H, d, J=13.4 Hz), 8.22 (1H, d, J = 2.4 Hz).








[0310]

[Table 3]

Ref.No	Str.	Chemical name	P.D.
1-12		(E)-3,5-difluoro-4-(2-nitrovinyl)phenol	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 6.49–6.55 (2H, m), 7.76 (1H, d, $J = 14.1$ Hz), 8.10 (1H, d, $J = 14.1$ Hz).
1-13		(E)-methyl(4-(2-nitrovinyl)phenyl)sulfane	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.53 (3H, s), 7.27 (2H, d, $J = 8.5$ Hz), 7.45 (2H, d, $J = 8.5$ Hz), 7.57 (1H, d, $J = 13.3$ Hz), 7.97 (1H, d, $J = 13.3$ Hz).
1-14		(E)-2-methoxy-5-(2-nitrovinyl)thiophene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.98 (3H, s), 6.27 (1H, d, $J = 4.2$ Hz), 7.19 (1H, d, $J = 4.2$ Hz), 7.26 (1H, d, $J = 13.3$ Hz), 8.03 (1H, d, $J = 13.3$ Hz).
1-15		(E)-1-(difluoromethoxy)-4-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 6.59 (1H, t, $J = 73.0$ Hz), 7.20 (2H, d, $J = 8.5$ Hz), 7.53–7.59 (3H, m), 7.99 (1H, d, $J = 13.3$ Hz).
1-16		(E)-5-ethyl-1,3-difluoro-2-(2-nitrovinyl)benzene	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.26 (3H, t, $J = 7.6$ Hz), 2.69 (2H, q, $J = 7.6$ Hz), 6.86 (2H, d, $J = 9.8$ Hz), 7.82 (1H, d, $J = 13.4$ Hz), 8.13 (1H, d, $J = 13.4$ Hz).





[0311]

[Table 4]

Ref.No.	Str.	Chemical name	P.D
1-17		(E)-5-(difluoromethoxy)-1,3-difluoro-2-(2-nitrovinyl)benzene	¹ H-NMR (400 MHz, CDCl ₃) δ 6.60 (1H, t, J = 71.5 Hz), 6.82–6.87 (2H, m), 7.81 (1H, d, J = 13.9 Hz), 8.09 (1H, d, J = 13.9 Hz).
1-18		(E)-5-ethoxy-1,3-difluoro-2-(2-nitrovinyl)benzene	¹ H-NMR (400 MHz, CDCl ₃) δ 1.45 (3H, t, J = 7.1 Hz), 4.07 (2H, q, J = 7.1 Hz), 6.54 (2H, d, J = 10.4 Hz), 7.76 (1H, d, J = 13.8 Hz), 8.11 (1H, d, J = 13.8 Hz).
1-19		(E)-3,5-difluoro-N,N-dimethyl-4-(2-nitrovinyl)aniline	¹ H-NMR (400 MHz, CDCl ₃) δ 3.06 (6H, s), 6.22 (2H, d, J = 12.8 Hz), 7.71 (1H, d, J = 13.4 Hz), 8.13 (1H, d, J = 13.4 Hz).
1-20		(E)-6-fluoro-5-(2-nitrovinyl)benzofuran	¹ H-NMR (400 MHz, CDCl ₃) δ 6.81–6.82 (1H, m), 7.35 (1H, d, J = 11.0 Hz), 7.69–7.79 (3H, m), 8.17 (1H, d, J = 14.1 Hz).
1-21		(E)-2-methoxy-5-(2-nitrovinyl)pyridine	¹ H-NMR (400 MHz, CDCl ₃) δ 4.00 (3H, s), 6.83 (1H, d, J = 8.9 Hz), 7.53 (1H, d, J = 14.1 Hz), 7.76 (1H, dd, J = 8.9, 2.4 Hz), 7.98 (1H, d, J = 14.1 Hz), 8.35 (1H, d, J = 2.4 Hz).
1-22		(E)-N-(3,5-difluoro-4-(2-nitrovinyl)phenyl)acetamide	¹ H-NMR (400 MHz, CDCl ₃) δ 2.22 (3H, s), 7.30 (2H, d, J = 10.4 Hz), 7.42 (1H, brs), 7.79 (1H, d, J = 13.8 Hz), 8.10 (1H, d, J = 13.8 Hz),
1-23		(E)-[3,5-difluoro-4-(2-nitrovinyl)phenyl](methyl)carbamate	¹ H-NMR (400 MHz, CDCl ₃) δ 1.53 (9H, s), 3.31 (3H, s), 7.09 (2H, d, J = 11.0 Hz), 7.81 (1H, d, J = 14.1 Hz), 8.11 (1H, d, J = 14.1 Hz).

[0312]


[Table 5]

Ref.No.	Str.	Chemical name	P.D
1-24		(E)-4,6-difluoro-5-(2-nitrovinyl)-2,3-dihydrobenzofuran	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.29 (2H, t, $J = 8.9$ Hz), 4.76 (2H, t, $J = 8.9$ Hz), 6.48 (1H, d, $J = 10.4$ Hz), 7.75 (1H, d, $J = 14.1$ Hz), 8.12 (1H, d, $J = 14.1$ Hz).
1-25		(E)-tert-butyl {[6-fluoro-5-(2-nitrovinyl)-2,3-dihydro-1H-inden-1-yl]oxy}dimethylsilane	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.16 (3H, s), 0.19 (3H, s), 0.95 (9H, s), 1.92-2.05 (1H, m), 2.45-2.52 (1H, m), 2.72-2.80 (1H, m), 2.96 (1H, dd, $J = 15.6, 6.4$ Hz), 5.22 (1H, t, $J = 7.6$ Hz), 7.07 (1H, d, $J = 10.4$ Hz), 7.32 (1H, d, $J = 6.1$ Hz), 7.70 (1H, d, $J = 14.1$ Hz), 8.05 (1H, d, $J = 14.1$ Hz).
1-26		(E)-1-benzyl-6-fluoro-5-(2-nitrovinyl)indoline	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.04 (2H, t, $J = 8.5$ Hz), 3.61 (2H, t, $J = 8.5$ Hz), 4.38 (2H, s), 6.16 (1H, d, $J = 12.1$ Hz), 7.09 (1H, d, $J = 6.7$ Hz), 7.25-7.38 (5H, m), 7.55 (1H, d, $J = 13.9$ Hz), 8.05 (1H, d, $J = 13.9$ Hz).
1-27		(E)-3,5-difluoro-4-(2-nitrovinyl)pyridine	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 7.90 (1H, d, $J = 14.1$ Hz), 8.06 (1H, d, $J = 14.1$ Hz), 8.52 (2H, s).

[0313]

5 <Reference Example 2-1>

[Chemical Formula 43]



[0314]

(-)-(R^{*})-2-[1-(4-Methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester

Malonic acid dimethyl ester (0.36 mL) and nickel(II) bis[(S,S)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide (68 mg) were added to a solution of (E)-1-methoxy-4-(2-nitrovinyl)benzene (500 mg) in toluene (2.8 mL) under an argon atmosphere, and the mixture was stirred at room temperature for 6 hours. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1), to obtain the title compound as a colorless liquid (865 mg).

¹H-NMR (400 MHz, CDCl₃) δ 3.57 (3H, s), 3.76 (3H, s), 3.78 (3H, s), 3.83 (1H, d, J = 9.1 Hz), 4.16-4.22 (1H, m), 4.83 (1H, dd, J = 12.7, 9.1 Hz), 4.89 (1H, dd, J = 12.7, 5.1 Hz), 6.84 (2H, d, J = 9.1 Hz), 7.14 (2H, d, J = 9.1 Hz).

[α]_D²⁵ = -20 (c 0.26, EtOH)





[0315]

The following Reference Examples 2-2 to 2-28 were obtained using a corresponding nitrostyrene in the same method as in Reference Example 2-1.

The structures and spectral data thereof are shown in Tables 6 to 12.

[0316]

[Table 6]

Ref.No	Str.	Chemical name	P.D.
2-2		(R*)-2-[1-(4-methoxy-2-methylphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.40 (3H, s), 3.55 (3H, s), 3.76 (3H, s), 3.77 (3H, s), 3.80 (1H, J = 9.1 Hz), 4.47-4.53 (1H, m), 4.78-4.89 (2H, m), 6.69-6.72 (2H, m), 7.02-7.05 (1H, m).
2-3		(R*)-2-[1-(2-chloro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.64 (3H, s), 3.74 (3H, s), 3.78 (3H, s), 4.08 (1H, d, J = 9.7 Hz), 4.64-4.70 (1H, m), 4.92 (1H, dd, J = 13.4, 4.3 Hz), 5.07 (1H, dd, J = 13.4, 8.6 Hz), 6.77 (1H, dd, J = 8.6, 3.1 Hz), 6.94 (1H, d, J = 3.1 Hz), 7.14 (1H, d, J = 8.6 Hz).
2-4		(R*)-2-[1-(2-fluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.57 (3H, s), 3.77 (6H, s), 3.96 (1H, d, J = 9.7 Hz), 4.36 (1H, q, J = 9.7 Hz), 4.89 (2H, d, J = 6.7 Hz), 6.59-6.64 (2H, m), 7.11 (1H, t, J = 8.8 Hz).
2-5		(R*)-2-[1-(3-fluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.61 (3H, s), 3.77 (3H, s), 3.80 (1H, d, J = 9.2 Hz), 3.86 (3H, s), 4.14-4.20 (1H, m), 4.79-4.91 (2H, m), 6.87-6.99 (3H, m).






[0317]

[Table 7]

Ref.No	Str.	Chemical name	P.D.
2-6		(R*)-2-[1-(3,5-difluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.65 (3H, s), 3.76-3.79 (1H, m), 3.77 (3H, s), 3.98 (3H, s), 4.14-4.19 (1H, m), 4.79-4.91 (2H, m), 6.77-6.84 (2H, m).
2-7		(R*)-2-[1-(2,5-difluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.61 (3H, s), 3.78 (3H, s), 3.86 (3H, s), 3.92 (1H, d, J = 9.8 Hz), 4.30-4.37 (1H, m), 4.83-4.91 (2H, m), 6.69 (1H, dd, J = 11.0, 7.3 Hz), 6.96 (1H, dd, J = 11.0, 7.3 Hz).
2-8		(-)-(R*)-2-[1-(2,6-difluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.57 (3H, s), 3.77 (3H, s), 3.80 (3H, s), 3.93 (1H, d, J = 10.4 Hz), 4.63-4.70 (1H, m), 4.81 (1H, dd, J = 13.1, 10.1 Hz), 4.91 (1H, dd, J = 13.1, 4.7 Hz), 6.41-6.47 (2H, m). [α] _D ²⁴ = -25 (c 0.11, EtOH)
2-9		(R*)-2-[1-(2,3-dihydrobenzofuran-5-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.17 (2H, t, J = 8.8 Hz), 3.59 (3H, s), 3.76 (3H, s), 3.81 (1H, d, J = 9.2 Hz), 4.14-4.20 (1H, m), 4.55 (2H, t, J = 8.8 Hz), 4.78-4.90 (2H, m), 6.70 (1H, d, J = 7.9 Hz), 6.94 (1H, d, J = 7.9 Hz), 7.05 (1H, s).
2-10		(-)-(R*)-2-[1-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.13 (2H, t, J = 8.9 Hz), 3.58 (3H, s), 3.77 (3H, s), 3.96 (1H, d, J = 9.8 Hz), 4.29-4.35 (1H, m), 4.59 (2H, t, J = 8.9 Hz), 4.87 (2H, d, J = 7.3 Hz), 6.48 (1H, d, J = 11.6 Hz), 6.99 (1H, d, J = 7.3 Hz). [α] _D ²⁶ = -20 (c 0.31, EtOH)



[0318]

[Table 8]

Ref.No	Str.	Chemical name	P.D.
2-11		(R*)-2-[1-(7-fluoro-6,7-dihydrochroman-6-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.92-1.98 (2H, m), 2.68 (2H, t, J = 6.4 Hz), 3.58 (3H, s), 3.77 (3H, s), 3.94 (1H, d, J = 9.8 Hz), 4.14 (2H, t, J = 4.9 Hz), 4.30 (1H, q, J = 7.3 Hz), 4.87 (2H, d, J = 7.3 Hz), 6.48 (1H, d, J = 12.2 Hz), 6.84 (1H, d, J = 8.6 Hz).
2-12		(S*)-2-[1-(3-fluoro-5-methoxypyridin-2-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.66 (3H, s), 3.76 (3H, s), 3.84 (3H, s), 3.98 (1H, d, J = 7.9 Hz), 4.62-4.67 (1H, m), 4.89 (1H, dd, J = 14.1, 4.3 Hz), 5.09 (1H, dd, J = 14.1, 9.5 Hz), 6.94 (1H, dd, J = 11.0, 2.4 Hz), 8.05 (1H, d, J = 2.4 Hz).
2-13		(S*)-2-[1-(5-methoxythiophen-2-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.70 (3H, s), 3.77 (3H, s), 3.84 (3H, s), 3.86 (1H, d, J = 7.9 Hz), 4.31-4.36 (1H, m), 4.81-4.89 (2H, m), 5.98 (1H, d, J = 4.2 Hz), 6.55 (1H, d, J = 4.2 Hz).
2-14		(R*)-2-[1-(2,6-difluoro-4-hydroxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.58 (3H, s), 3.80 (3H, s), 3.92 (1H, d, J = 10.4 Hz), 4.62-4.69 (1H, m), 4.80 (1H, dd, J = 12.8, 10.4 Hz), 4.91 (1H, dd, J = 12.8, 4.9 Hz), 5.79 (1H, brs), 6.35-6.41 (2H, m).
2-15		(R*)-2-[1-(4-methylthiophenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.45 (3H, s), 3.59 (3H, s), 3.76 (3H, s), 3.83 (1H, d, J = 9.1 Hz), 4.17-4.23 (1H, m), 4.81-4.92 (2H, m), 7.14 (2H, d, J = 8.5 Hz), 7.18 (2H, d, J = 8.5 Hz).

[0319]

[Table 9]

Ref.No	Str.	Chemical name	P.D.
2-16		2-[(1R*)-1-(4-methoxyphenyl)-2-nitropropyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.35 (0.6H, d, J = 6.7 Hz), 1.42 (2.4H, d, J = 6.7 Hz), 3.40-3.52 (6H, m), 3.72-4.41 (5H, m), 5.01-5.15 (1H, m), 6.80-7.40 (4H, m).
2-17		(R*)-2-[1-(4-ethyl-2,6-difluorophenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.20 (3H, t, J = 7.6 Hz), 2.60 (2H, q, J = 7.6 Hz), 3.55 (3H, s), 3.80 (3H, s), 3.95 (1H, d, J = 10.4 Hz), 4.68-4.75 (1H, m), 4.84 (1H, dd, J = 13.4, 9.8 Hz), 4.93 (1H, dd, J = 13.4, 4.9 Hz), 6.73 (2H, dd, J = 14.1, 4.3 Hz).





[0320]

[Table 10]

Ref.No.	Str.	Chemical name	P.D
2-18		(R*)-2-[1-[4-(difluoromethoxy)-2,6-difluorophenyl]-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.58 (3H, s), 3.81 (3H, s), 3.93 (1H, d, J = 10.3 Hz), 4.71 (1H, dt, J = 10.3, 4.2 Hz), 4.82 (1H, dd, J = 13.3, 10.3 Hz), 4.93 (1H, dd, J = 13.3, 4.2 Hz), 6.50 (1H, t, J = 72.6 Hz), 6.69-6.74 (2H, m).
2-19		(R*)-2-[1-(4-ethoxy-2,6-difluorophenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.40 (3H, t, J = 7.1 Hz), 3.57 (3H, s), 3.80 (3H, s), 3.92 (1H, d, J = 10.2 Hz), 3.96 (2H, q, J = 7.1 Hz), 4.66 (1H, td, J = 10.2, 4.8 Hz), 4.81 (1H, dd, J = 13.0, 10.2 Hz), 4.91 (1H, dd, J = 13.0, 4.8 Hz), 6.39-6.45 (2H, m).
2-20		(R*)-2-[1-[4-(dimethylamino)-2,6-difluorophenyl]-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.92 (6H, s), 3.57 (3H, s), 3.79 (3H, s), 3.92 (1H, d, J = 10.4 Hz), 4.62 (1H, td, J = 10.4, 5.1 Hz), 4.80 (1H, t, J = 11.3 Hz), 4.89 (1H, dd, J = 12.8, 4.9 Hz), 6.13 (2H, d, J = 12.8 Hz).
2-21		(R*)-2-[1-(6-fluorobenzofuran-5-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.54 (3H, s), 3.78 (3H, s), 4.06 (1H, d, J = 9.8 Hz), 4.49-4.55 (1H, m), 4.95 (1H, dd, J = 13.2, 5.2 Hz), 4.99 (1H, dd, J = 13.2, 8.9 Hz), 6.71-6.72 (1H, m), 7.25 (1H, d, J = 12.8 Hz), 7.44 (1H, d, J = 7.3 Hz), 7.61 (1H, d, J = 2.4 Hz)




[0321]

[Table 11]

Ref.No.	Str.	Chemical name	P.D
2-22		(R*)-2-[1-(6-methoxypyridin-3-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.62 (3H, s), 3.78 (3H, s), 3.83 (1H, d, J = 9.2 Hz), 3.91 (3H, s), 4.17-4.23 (1H, m), 4.80-4.93 (2H, m), 6.71 (1H, d, J = 8.6 Hz), 7.47 (1H, dd, J = 8.6, 2.4 Hz), 8.05 (1H, d, J = 2.4 Hz).
2-23		(R*)-2-[1-(4-acetamido-2,6-difluorophenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.18 (3H, s), 3.57 (3H, s), 3.80 (3H, s), 3.94 (1H, d, J = 10.4 Hz), 4.66-4.73 (1H, m), 4.82 (1H, dd, J = 13.1, 10.1 Hz), 4.92 (1H, dd, J = 13.1, 4.9 Hz), 7.16 (2H, d, J = 10.2 Hz), 7.25 (1H, brs),
2-24		(R*)-2-(1-{4-[(tert-butoxycarbonyl)(methyl)amino]-2,6-difluorophenyl}-2-nitroethyl)malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.48 (9H, s), 3.23 (3H, s), 3.58 (3H, s), 3.80 (3H, s), 3.94 (1H, d, J = 10.4 Hz), 4.67-4.75 (1H, m), 4.83 (1H, dd, J = 13.4, 9.8 Hz), 4.93 (1H, dd, J = 13.4, 4.9 Hz), 6.92 (2H, d, J = 10.4 Hz).
2-25		(R*)-2-[1-(4,6-difluoro-2,3-dihydrobenzofuran-5-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.20 (2H, t, J = 8.6 Hz), 3.58 (3H, s), 3.80 (3H, s), 3.92 (1H, d, J = 11.0 Hz), 4.62-4.69 (3H, m), 4.80 (1H, dd, J = 12.8, 9.8 Hz), 4.90 (1H, dd, J = 12.8, 4.9 Hz), 6.34 (1H, d, J = 9.8 Hz).

[0322]

[Table 12]


Ref.No.	Str.	Chemical name	P.D
2-26		2-((1R*)-1-{1-[(tert-butyl dimethylsilyl)oxy]-6-fluoro-2,3-dihydro-1H-inden-5-yl}-2-nitroethyl)malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.14 (3H, s), 0.16 (3H, s), 0.94 (9H, d, J = 1.2 Hz), 1.85-1.95 (1H, m), 2.39-2.46 (1H, m), 2.64-2.72 (1H, m), 2.84-2.90 (1H, m), 3.50 (1.5H, s), 3.60 (1.5H, s), 3.76 (1.5H, s), 3.77 (1.5H, s), 3.98 (1H, dd, J = 9.8, 4.9 Hz), 4.36-4.45 (1H, m), 4.87-4.97 (2H, m), 5.15-5.19 (1H, m), 6.93-6.97 (1H, m), 7.01-7.04 (1H, m).
2-27		(R*)-2-[1-(1-benzyl-6-fluoroindolin-5-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.89 (2H, t, J = 8.5 Hz), 3.37 (2H, t, J = 8.5 Hz), 3.60 (3H, s), 3.76 (3H, s), 3.93 (1H, d, J = 9.7 Hz), 4.20 (2H, s), 4.28 (1H, dt, J = 9.7, 7.3 Hz), 4.85 (2H, d, J = 7.3 Hz), 6.11 (1H, d, J = 12.1 Hz), 6.82 (1H, d, J = 7.2 Hz), 7.28-7.36 (5H, m).
2-28		(R*)-2-[1-(3,5-difluoropyridin-4-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.60 (3H, s), 3.83 (3H, s), 3.96 (1H, d, J = 9.8 Hz), 4.76 (1H, td, J = 9.9, 4.5 Hz), 4.88 (1H, dd, J = 13.8, 10.1 Hz), 4.97 (1H, dd, J = 13.8, 4.6 Hz), 8.36 (2H, s).

[0323]

5 <Reference Example 3-1>

[0324]

[Chemical Formula 44]



[0325]

(+)-(S*)-2-[1-(4-Methoxyphenyl)-2-nitroethyl]malonic acid
dimethyl ester

Nickel(II)

5 bis[(R,R)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide was
used as a catalyst in place of nickel(II)
bis[(S,S)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide, and
the same method as in Reference Example 2-1 was performed to obtain
the title compound.

10 ¹H-NMR (400 MHz, CDCl₃) δ 3.57 (3H, s), 3.76 (3H, s), 3.78 (3H, s),
3.83 (1H, d, J = 9.1 Hz), 4.16-4.22 (1H, m), 4.83 (1H, dd, J = 12.7,
9.1 Hz), 4.89 (1H, dd, J = 12.7, 5.1 Hz), 6.84 (2H, d, J = 9.1 Hz),
7.14 (2H, d, J = 9.1 Hz).

[α]_D²⁵ = +14 (c 0.26, EtOH)






15 [0326]

The same method as in Reference Example 3-1 was performed
using a corresponding nitrostyrene to obtain the following
Reference Examples 3-2 to 3-8.

The structures and spectral data thereof are shown in Tables
20 13 to 14.



[0327]

[Table 13]

Ref.No	Str.	Chemical name	P.D.
3-2		(S*)-2-(2-nitro-1-phenylethyl)malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.57 (3H, s), 3.76 (3H, s), 3.87 (1H, d, J = 8.5 Hz), 4.22-4.28 (1H, m), 4.85-4.95 (2H, m), 7.22-7.35 (5H, m).
3-3		(S*)-2-[1-(3-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.60 (3H, s), 3.76 (3H, s), 3.78 (3H, s), 3.86 (1H, d, J = 9.1 Hz), 4.18-4.25 (1H, m), 4.84-4.94 (2H, m), 6.75-6.83 (3H, m), 7.23 (1H, t, J = 7.9 Hz).
3-4		(S*)-2-[1-(2-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.49 (3H, s), 3.73 (3H, s), 3.85 (3H, s), 4.16 (1H, d, J = 9.6 Hz), 4.34-4.41 (1H, m), 4.86 (1H, dd, J = 13.0, 4.8 Hz), 5.01 (1H, dd, J = 13.0, 9.6 Hz), 6.84-6.88 (2H, m), 7.12 (1H, dd, J = 7.6, 1.8 Hz), 7.21-7.26 (1H, m).
3-5		(+)-(S*)-2-[1-(2,6-difluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.57 (3H, s), 3.77 (3H, s), 3.80 (3H, s), 3.93 (1H, d, J = 10.4 Hz), 4.63-4.70 (1H, m), 4.81 (1H, dd, J = 13.1, 10.1 Hz), 4.91 (1H, dd, J = 13.1, 4.7 Hz), 6.41-6.47 (2H, m). [α] _D ²⁴ = +15 (c 0.07, EtOH)
3-6		(R*)-2-[1-(5-methoxythiophen-2-yl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.70 (3H, s), 3.77 (3H, s), 3.84 (3H, s), 3.86 (1H, d, J = 7.9 Hz), 4.31-4.36 (1H, m), 4.81-4.89 (2H, m), 5.98 (1H, d, J = 4.2 Hz), 6.55 (1H, d, J = 4.2 Hz).

[0328]

[Table 14]


Ref.No	Str.	Chemical name	P.D.
3-7		2-[(1S*)-1-(4-methoxyphenyl)-2-nitropropyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.35 (0.6H, d, J = 6.7 Hz), 1.42 (2.4H, d, J = 6.7 Hz), 3.40-3.52 (6H, m), 3.72-4.41 (5H, m), 5.01-5.15 (1H, m), 6.80-7.40 (4H, m).
3-8		(S*)-2-[1-(4-difluoromethoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.59 (3H, s), 3.77 (3H, s), 3.83 (1H, d, J = 9.1 Hz), 4.22-4.28 (1H, m), 4.82-4.94 (2H, m), 6.49 (1H, t, J = 73.6 Hz), 7.08 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz).

[0329]

5 <Reference Example 4-1>

[0330]

[Chemical Formula 45]



[0331]

(±)-2-[1-(4-Methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester

10

The same method as in Reference Example 2-1 was performed using (±)-Nickel(II) bis[N,N'-dibenzylcyclohexane-1,2-diamine]bromide as a catalyst in place of nickel(II)

15 bis[(S,S)-N,N'-dibenzylcyclohexane-1,2-diamine]bromide to

obtain the title compound.


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.57 (3H, s), 3.76 (3H, s), 3.78 (3H, s),
3.83 (1H, d, $J = 9.1$ Hz), 4.16-4.22 (1H, m), 4.83 (1H, dd, $J = 12.7$,
9.1 Hz), 4.89 (1H, dd, $J = 12.7$, 5.1 Hz), 6.84 (2H, d, $J = 9.1$ Hz),
5 7.14 (2H, d, $J = 9.1$ Hz).

[0332]

<Reference Example 5-1>

[0333]

[Chemical Formula 46]



10 2-(4-Methoxybenzylidene)malonic acid dimethyl ester

Piperidine (0.4 mL) and acetic acid (0.47 mL) were added to a solution of p-anisaldehyde (5.0 mL) and malonic acid dimethyl ester (4.7 mL) in benzene (0.9 mL) under an argon atmosphere, to produce a reaction solution. The reaction solution was heated
15 to reflux with a Dean-Stark apparatus for 3 hours. Ethyl acetate was added to the reaction solution, and the mixture was washed with a 1 mol/L hydrochloric acid, a saturated aqueous sodium hydrogen carbonate solution, and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under
20 reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1), to obtain the title compound as a white solid (9.5 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.83 (3H, s), 3.84 (3H, s), 3.87 (3H, s),


6.90 (2H, d, J = 8.6 Hz), 7.39 (2H, d, J = 8.6 Hz), 7.71 (s, 1H).

[0334]

<Reference Example 6-1>

[0335]

5 [Chemical Formula 47]



[0336]

(±)-2-[1-(4-Methoxyphenyl)-2-methyl-2-nitropropyl]malonic acid dimethyl ester

2-Nitropropane (0.14 mL) and potassium fluoride supported
10 on alumina (KF/Al₂O₃) (400 mg) were added to a solution of
2-(4-methoxybenzylidene)malonic acid dimethyl ester (250 mg) in
dimethyl sulfoxide (0.8 mL) under an argon atmosphere, to produce
a reaction solution. The reaction solution was stirred at room
temperature for 1 hour and a half. Diethyl ether was added to
15 the reaction solution, the insoluble was removed by filtration,
and the solution was washed with water and a brine successively,
and dried over anhydrous sodium sulfate. The solvent was removed
under reduced pressure, and the residue was then purified by silica
gel column chromatography (ethyl acetate : hexane = 4 : 1), to
20 obtain the title compound as a white solid (298 mg).


¹H-NMR (400 MHz, CDCl₃) δ 1.43 (3H, s), 1.64 (3H, s), 3.35 (3H, s),
3.75 (3H, s), 3.78 (3H, s), 4.22 (2H, s), 6.81 (2H, d, J = 9.1 Hz),
7.10 (2H, d, J = 9.1 Hz).

[0337]

<Reference Example 6-2>

[0338]

[Chemical Formula 48]



5 [0339]

(±)-2-[(4-Methoxyphenyl)-(1-nitrocyclopentyl)methyl]malonic acid dimethyl ester

The same method as in Reference Example 6-1 was performed using nitrocyclopentane in place of 2-nitropropane to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 1.43-1.86 (6H, m), 2.33-2.42 (1H, m), 2.45-2.53 (1H, m), 3.35 (3H, s), 3.77 (3H, s), 3.80 (3H, s), 4.14 (1H, d, J = 11.5 Hz), 4.31 (1H, d, J = 11.5 Hz), 6.80 (2H, d, J = 8.5 Hz), 7.04 (2H, d, J = 9.1 Hz).

15 [0340]

<Reference Example 6-3>

[0341]

[Chemical Formula 49]



[0342]

20 (±)-2-[(4-Methoxyphenyl)-(1-nitrocyclohexyl)methyl]malonic

acid dimethyl ester

The same method as in Reference Example 6-1 was performed using nitrocyclohexane in place of 2-nitropropane to obtain the title compound.


5 ¹H-NMR (400 MHz, CDCl₃) δ 1.10-1.17 (2H, m), 1.22-1.68 (6H, m), 2.35-2.44 (2H, m), 3.28 (3H, s), 3.78 (3H, s), 3.79 (3H, s), 3.94 (1H, d, J = 10.9 Hz), 4.32 (1H, d, J = 10.9 Hz), 6.79 (2H, d, J = 7.9 Hz), 7.02 (2H, brd, J = 7.9 Hz).

[0343]

10 <Reference Example 7-1>

[0344]

[Chemical Formula 50]



[0345]

(-)-(3S*,4R*)-4-(4-Methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester

Nickel(II) chloride hexahydrate (1.3 g) was added to a solution of (R*)-2-[1-(4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester (1.7 g) in methanol (110 mL) under an argon atmosphere, to produce a reaction solution. To the reaction solution, sodium borohydride (1.03 g) was added in several doses under ice-cooling, and the reaction solution was warmed to room temperature, and stirred for 2 hours. A saturated aqueous ammonium chloride solution and ethyl acetate were added to the reaction

solution, and the mixture was stirred at room temperature for 1 hour. The mixture was extracted with ethyl acetate, and the extract was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the crude product was washed with ethanol-diisopropyl ether, to obtain the title compound as a white solid (840 mg).

¹H-NMR (400 MHz, CDCl₃) δ 3.40 (1H, t, J = 9.1 Hz), 3.53 (1H, d, J = 9.7 Hz), 3.76-3.81 (1H, m), 3.78 (3H, s), 3.80 (3H, s), 4.08 (1H, q, J = 8.9 Hz), 5.85 (1H, brs), 6.88 (2H, d, J = 8.5 Hz), 7.18 (2H, d, J = 8.5 Hz).


[α]_D²⁵ = -96 (c 0.19, EtOH)

[0346]

The same method as in Reference Example 7-1 was performed using a corresponding nitro substance to obtain the following Reference Examples 7-2 to 7-42. The structures and spectral data thereof are shown in Tables 15 to 25.






[0347]

[Table 15]

Ref.No	Str.	Chemical name	P.D.
7-2		(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.38 (1H, t, J = 8.9 Hz), 3.50 (1H, d, J = 9.8 Hz), 3.76-3.81 (1H, m), 3.79 (3H, s), 3.88 (3H, s), 4.06 (1H, q, J = 8.8 Hz), 5.94 (1H, brs), 6.90-7.01 (3H, m).





[0348]

[Table 16]

Ref.No	Str.	Chemical name	P.D.
7-3		(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.44 (1H, dd, J = 9.7, 7.9 Hz), 3.68 (1H, d, J = 9.7 Hz), 3.75-3.79 (1H, m), 3.78 (3H, s), 3.79 (3H, s) 4.22 (1H, q, J = 8.9 Hz), 6.15 (1H, s), 6.62- 6.68 (2H, m), 7.15 (1H, t, J = 8.5Hz).
7-4		(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.35-3.41 (1H, m), 3.67 (1H, d, J = 8.6 Hz), 3.79 (3H, s), 3.80 (3H, s), 3.89 (1H, t, J = 8.6 Hz), 4.47 (1H, q, J = 7.9 Hz), 5.70 (1H, brs), 6.82 (1H, dd, J = 8.6, 2.4 Hz), 6.95 (1H, d, J = 2.4Hz), 7.20 (2H, t, J = 8.6 Hz).
7-5		(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.36 (3H, s), 3.29-3.36 (1H, m), 3.57 (1H, d, J = 9.1Hz), 3.76-3.81 (1H, m), 3.77 (3H, s), 3.78 (3H, s), 4.31 (1H, q, J = 8.4 Hz), 6.09 (1H, brs), 6.73 (1H, d, J = 3.0 Hz), 6.76 (1H, dd, J = 8.5, 3.0 Hz), 7.17 (1H, d, J = 8.5 Hz).
7-6		(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.37 (1H, t, J = 8.6 Hz), 3.47 (1H, d, J = 9.8 Hz), 3.77-3.81 (1H, m), 3.81 (3H, s), 3.98 (3H, s) 4.05 (1H, q, J = 8.8 Hz), 5.69 (1H, brs), 6.78-6.85 (2H, m).
7-7		(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.42 (1H, t, J = 8.9 Hz), 3.62 (1H, d, J = 9.8 Hz), 3.75-3.81 (1H, m), 3.79 (3H, s), 3.87 (3H, s) 4.20 (1H, q, J = 8.9 Hz), 6.14 (1H, brs) 6.71 (1H, dd, J = 11.0, 7.3 Hz), 6.98 (1H, dd, J = 11.0, 7.3 Hz).






[0349]

[Table 17]

Ref.No	Str.	Chemical name	P.D.
7-8		(-)-(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.51 (1H, t, J = 9.2 Hz), 3.64-3.70 (1H, m), 3.78 (6H, s), 3.78-3.81 (1H, m), 4.46 (1H, q, J = 9.4 Hz), 6.24 (1H, brs), 6.43-6.50 (2H, m). [α] _D ²³ = -120 (c 0.11, EtOH)
7-9		(3S*,4R*)-4-(2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.20 (2H, t, J = 8.8 Hz), 3.38 (1H, t, J = 8.9 Hz), 3.53 (1H, d, J = 9.8 Hz), 3.77 (1H, t, J = 9.0 Hz), 3.78 (3H, s), 4.06 (1H, q, J = 9.0 Hz), 4.57 (2H, t, J = 8.8 Hz), 5.94 (1H, brs), 6.74 (1H, d, J = 7.9 Hz), 6.99 (1H, d, J = 7.9 Hz), 7.09 (1H, s).
7-10		(-)-(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.15 (2H, t, J = 8.9 Hz), 3.41 (1H, t, J = 8.6 Hz), 3.66 (1H, d, J = 9.8 Hz), 3.74-3.79 (1H, m), 3.79 (3H, s), 4.19 (1H, q, J = 8.6 Hz), 4.61 (2H, t, J = 8.9 Hz), 5.68 (1H, brs), 6.52 (1H, d, J = 11.6 Hz), 7.03 (1H, d, J = 7.9 Hz). [α] _D ²⁶ = -121 (c 0.20, EtOH)
7-11		(3S*,4R*)-4-(7-fluorochroman-6-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.94-2.01 (2H, m), 2.71 (2H, t, J = 6.1 Hz), 3.42 (1H, t, J = 8.9 Hz), 3.67 (1H, d, J = 9.8 Hz), 3.73-3.78 (1H, m), 3.78 (3H, s), 4.11-4.17 (3H, m), 5.68 (1H, brs), 6.52 (1H, d, J = 11.6 Hz), 6.87 (1H, d, J = 7.9 Hz).





[0350]

[Table 18]

Ref.No	Str.	Chemical name	P.D.
7-12		(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.53 (1H, t, J = 8.6 Hz), 3.77-3.81 (1H, m), 3.79 (3H, s), 3.86 (3H, s), 4.04 (1H, d, J = 9.2 Hz), 4.52 (1H, q, J = 8.8 Hz), 5.72 (1H, brs), 6.94 (1H, dd, J = 11.0, 2.4 Hz), 8.11 (1H, d, J = 2.4 Hz).
7-13		(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.41 (1H, t, J = 9.1 Hz), 3.49 (1H, d, J = 10.3 Hz), 3.75-3.79 (1H, m), 3.81 (3H, s), 3.86 (3H, s), 4.22 (1H, q, J = 9.1 Hz), 5.78 (1H, brs), 6.01 (1H, d, J = 4.2 Hz), 6.54 (1H, d, J = 4.2 Hz).
7-14		(3S*,4R*)-4-(2,6-difluoro-4-hydroxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.25 (1H, t, J = 9.2 Hz), 3.52-3.59 (2H, m), 3.63 (3H, s), 4.12 (1H, q, J = 9.6 Hz), 6.43-6.49 (2H, m), 8.27 (1H, s), 10.41 (1H, brs).
7-15		(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.48 (3H, s), 3.41 (1H, t, J = 8.9 Hz), 3.54 (1H, d, J = 9.7 Hz), 3.78 (3H, s), 3.80 (1H, t, J = 8.9 Hz), 4.10 (1H, q, J = 8.9 Hz), 5.92 (1H, brs), 7.18 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz).
7-16		(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.67 (3H, d, J = 6.7 Hz), 3.60 (3H, s), 3.72 (3H, s), 3.80-3.90 (1H, m), 3.95-4.05 (2H, m), 6.88 (2H, d, J = 9.1 Hz), 7.21 (2H, d, J = 9.1 Hz), 8.21 (1H, s).






[0351]

[Table 19]

Ref.No	Str.	Chemical name	P.D.
7-17		(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.27 (3H, d, J = 6.1 Hz), 3.49 (1H, dd, J = 10.9, 8.5 Hz), 3.63 (1H, d, J = 10.9 Hz), 3.68-3.73 (1H, m), 3.74 (3H, s), 3.80 (3H, s), 5.65 (1H, s), 6.89 (2H, d, J = 9.1 Hz), 7.19 (2H, d, J = 9.1 Hz).
7-18		(3S*,4R*)-4-(4-ethyl-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.22 (3H, t, J = 7.6 Hz), 2.62 (2H, q, J = 7.6 Hz), 3.53 (1H, t, J = 9.2 Hz), 3.68 (1H, t, J = 9.2 Hz), 3.78 (3H, s), 3.83 (1H, d, J = 9.8 Hz), 4.52 (1H, q, J = 9.8 Hz), 6.03 (1H, brs), 6.75 (2H, dd, J = 12.2, 3.1 Hz).
7-19		(3R*,4S*)-2-oxo-4-phenylpyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.44 (1H, dd, J = 9.7, 8.5 Hz), 3.59 (1H, d, J = 9.7 Hz), 3.79 (3H, s), 3.80-3.85 (1H, m), 4.14 (1H, q, J = 8.9 Hz), 5.82 (1H, brs), 7.25-7.38 (5H, m).
7-20		(+)-(3R*,4S*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.40 (1H, t, J = 9.1 Hz), 3.53 (1H, d, J = 9.7 Hz), 3.76-3.81 (1H, m), 3.78 (3H, s), 3.80 (3H, s), 4.08 (1H, q, J = 8.9 Hz), 5.85 (1H, brs), 6.88 (2H, d, J = 8.5 Hz), 7.18 (2H, d, J = 8.5 Hz). [α] _D ²⁵ = +89 (c 0.16, EtOH)





[0352]

[Table 20]

Ref.No	Str.	Chemical name	P.D.
7-21		(3R*,4S*)-4-(3-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.43 (1H, t, J = 8.6 Hz), 3.59 (1H, d, J = 9.7 Hz), 3.79 (3H, s), 3.79-3.84 (1H, m), 3.81 (3H, s), 4.05-4.15 (1H, m), 6.05-6.25 (1H, br), 6.76-6.86 (3H, m), 7.27 (1H, t, J = 8.0 Hz).
7-22		(3R*,4S*)-4-(2-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.49 (1H, t, J = 8.5 Hz), 3.73 (1H, d, J = 9.2 Hz), 3.77 (3H, s), 3.83 (3H, s), 3.84 (1H, d, J = 9.2 Hz), 4.26 (1H, q, J = 8.5, Hz), 5.85 (1H, brs), 6.89-6.95 (2H, m), 7.20 (1H, dd, J = 7.3, 1.8 Hz), 7.27-7.30 (1H, m).
7-23		(+)-(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.51 (1H, t, J = 9.2 Hz), 3.64-3.70 (1H, m), 3.78 (6H, s), 3.78-3.81 (1H, m), 4.46 (1H, q, J = 9.4 Hz), 6.24 (1H, brs), 6.43-6.50 (2H, m). [α] _D ²⁴ = +105 (c 0.09, EtOH)
7-24		(3R*,4S*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.67 (3H, d, J = 6.7 Hz), 3.60 (3H, s), 3.72 (3H, s), 3.80-3.90 (1H, m), 3.95-4.05 (2H, m), 6.88 (2H, d, J = 9.1 Hz), 7.21 (2H, d, J = 9.1 Hz), 8.21 (1H, s).
7-25		(3R*,4S*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.27 (3H, d, J = 6.1 Hz), 3.49 (1H, dd, J = 10.9, 8.5 Hz), 3.63 (1H, d, J = 10.9 Hz), 3.68-3.73 (1H, m), 3.74 (3H, s), 3.80 (3H, s), 5.65 (1H, s), 6.89 (2H, d, J = 9.1 Hz), 7.19 (2H, d, J = 9.1 Hz).



[0353]

[Table 21]

Ref.No	Str.	Chemical name	P.D.
7-26		(3R*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.41 (1H, t, J = 9.1 Hz), 3.49 (1H, d, J = 10.3 Hz), 3.75-3.79 (1H, m), 3.81 (3H, s), 3.86 (3H, s), 4.22 (1H, q, J = 9.1 Hz), 5.78 (1H, s), 6.01 (1H, d, J = 4.2 Hz), 6.54 (1H, d, J = 4.2 Hz).
7-27		(3R*,4S*)-4-[4-(difluoromethoxy)phenyl]-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.39-3.44 (1H, m), 3.54 (1H, d, J = 9.7 Hz), 3.80 (3H, s), 3.80 (1H, t, J = 9.2 Hz), 4.12 (1H, q, J = 8.8 Hz), 5.98 (1H, brs), 6.50 (1H, t, J = 73.6 Hz), 7.12 (2H, d, J = 9.2 Hz), 7.22-7.26 (2H, m).
7-28		(±)-trans-4-[4-methoxyphenyl]-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.40 (1H, t, J = 9.1 Hz), 3.53 (1H, d, J = 9.7 Hz), 3.76-3.81 (1H, m), 3.78 (3H, s), 3.80 (3H, s), 4.08 (1H, q, J = 8.9 Hz), 5.85 (1H, brs), 6.88 (2H, d, J = 8.5 Hz), 7.18 (2H, d, J = 8.5 Hz).
7-29		(±)-trans-4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.92 (3H, s), 1.39 (3H, s), 3.72 (3H, s), 3.80 (1H, d, J = 12.1 Hz), 3.81 (3H, s), 3.93 (1H, d, J = 12.1 Hz), 5.90 (1H, s), 6.89 (2H, d, J = 8.6 Hz), 7.18 (2H, d, J = 8.6 Hz).





[0354]

[Table 22]

Ref.No	Str.	Chemical name	P.D.
7-30	 (±)-trans	(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4,4]nonane-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.21–1.67 (6H, m), 1.77–1.85 (1H, m), 1.88–1.95 (1H, m), 3.72 (3H, s), 3.80 (3H, s), 3.85 (1H, d, J = 11.0 Hz), 3.97 (1H, d, J = 11.0 Hz), 6.01 (1H, brs), 6.88 (2H, d, J = 8.6 Hz), 7.17 (2H, d, J = 8.6 Hz).
7-31	 (±)-trans	(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4,5]decane-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.78–0.89 (1H, m), 0.92–1.07 (1H, m), 1.16–1.34 (2H, m), 1.41–1.51 (2H, m), 1.52–1.80 (4H, m), 3.69 (1H, d, J = 12.1 Hz), 3.72 (3H, s), 3.81 (3H, s), 3.93 (1H, d, J = 12.1 Hz), 6.22 (1H, brs), 6.88 (2H, d, J = 9.1 Hz), 7.15 (2H, d, J = 9.1 Hz).





[0355]

[Table 23]

Ref.No.	Str.	Chemical name	P.D.
7-32		(3S*,4R*)-4-[4-(difluoromethoxy)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.52 (1H, t, J = 9.1 Hz), 3.68-3.73 (1H, m), 3.79-3.82 (4H, m), 4.49-4.56 (1H, m), 6.27 (1H, br), 6.51 (1H, t, J = 72.1 Hz), 6.72-6.78 (2H, m).
7-33		(3S*,4R*)-4-(4-ethoxy-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.40 (3H, t, J = 7.1 Hz), 3.51 (1H, t, J = 8.9 Hz), 3.67 (1H, t, J = 9.5 Hz), 3.78 (3H, s), 3.79 (1H, d, J = 11.0 Hz), 3.98 (2H, q, J = 7.1 Hz), 4.42-4.49 (1H, m), 6.33 (1H, s), 6.42-6.48 (2H, m).
7-34		(3S*,4R*)-4-[4-(dimethylamino)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.93 (6H, s), 3.50 (1H, t, J = 9.2 Hz), 3.64 (1H, t, J = 9.5 Hz), 3.77-3.80 (4H, m), 4.42 (1H, q, J = 9.6 Hz), 5.83 (1H, s), 6.18 (2H, d, J = 12.2 Hz).
7-35		(3S*,4R*)-4-(6-fluorobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.50 (1H, t, J = 8.9 Hz), 3.78 (1H, d, J = 10.4 Hz), 3.79 (3H, s), 3.86 (1H, t, J = 8.9 Hz), 4.34-4.40 (1H, m), 6.04 (1H, s), 6.74 (1H, d, J = 1.5 Hz), 7.25-7.28 (1H, m), 7.47 (1H, d, J = 7.3 Hz), 7.63 (1H, d, J = 1.5 Hz).




[0356]

[Table 24]

Ref.No.	Str.	Chemical name	P.D.
7-36		(3S*,4R*)-4-(6-methoxypyridin-3-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.39 (1H, t, J = 9.2 Hz), 3.51 (1H, d, J = 9.8 Hz), 3.80 (3H, s), 3.78-3.83 (1H, m), 3.93 (3H, s), 4.05-4.11 (1H, m), 6.23 (1H, brs), 6.76 (1H, d, J = 8.6 Hz), 7.49 (1H, dd, J = 8.6, 2.4 Hz), 8.08 (1H, d, J = 2.4 Hz).
7-37		(3S*,4R*)-4-(4-acetamide-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.19 (3H, s), 3.52 (1H, t, J = 9.2 Hz), 3.66-3.72 (1H, m), 3.78 (3H, s), 3.78-3.81 (1H, m), 4.50 (1H, q, J = 9.4 Hz), 5.88 (1H, s), 7.17 (2H, d, J = 10.4 Hz), 7.32 (1H, s).
7-38		(3S*,4R*)-4-{4-[(tert-butoxy carbonyl)(methyl)amino]-2,6-difluorophenyl}-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.49 (9H, s), 3.25 (3H, s), 3.53 (1H, t, J = 8.9 Hz), 3.66-3.72 (1H, m), 3.79 (3H, s), 3.82 (1H, d, J = 9.8 Hz), 4.52 (1H, q, J = 9.4 Hz), 5.99 (1H, brs), 6.92 (2H, d, J = 10.4 Hz).
7-39		(3S*,4R*)-4-(4,6-difluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.21 (2H, t, J = 8.6 Hz), 3.50 (1H, dd, J = 11.6, 6.7 Hz), 3.66 (1H, t, J = 8.9 Hz), 3.77-3.80 (4H, m), 4.46 (1H, q, J = 9.4 Hz), 4.65 (2H, t, J = 8.6 Hz), 6.16 (1H, s), 6.37 (1H, d, J = 10.4 Hz).

[0357]

[Table 25]


Ref.No.	Str.	Chemical name	P.D.
7-40		(3S*,4R*)-4-{1-[(tert-butyl dimethylsilyl)oxy]-6-fluoro-2,3-dihydro-1H-inden-5-yl}-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.14 (3H, s), 0.17 (3H, s), 0.94 (9H, s), 1.87-1.97 (1H, m), 2.41-2.49 (1H, m), 2.67-2.75 (1H, m), 2.91 (1H, dd, J = 15.9, 6.7 Hz), 3.41-3.46 (1H, m), 3.68-3.72 (1H, m), 3.75-3.81 (4H, m), 4.22-4.29 (1H, m), 5.20 (1H, t, J = 7.0 Hz), 5.82 (1H, s), 6.98 (1H, d, J = 10.4 Hz), 7.07 (1H, d, J = 6.7 Hz).
7-41		(3S*,4R*)-4-(1-benzyl-6-fluoroindolin-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.93 (2H, t, J = 8.6 Hz), 3.37-3.43 (3H, m), 3.65 (1H, d, J = 9.8 Hz), 3.70-3.75 (1H, m), 3.78 (3H, s), 4.11-4.19 (1H, m), 4.22 (2H, s), 5.81 (1H, br), 6.16 (1H, d, J = 11.6 Hz), 6.89 (1H, d, J = 7.9 Hz), 7.28-7.36 (5H, m).
7-42		(3S*,4R*)-4-(3,5-difluoropyridin-4-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.56 (1H, t, J = 9.2 Hz), 3.76-3.85 (5H, m), 4.60 (1H, q, J = 9.2 Hz), 5.91 (1H, s), 8.37 (2H, s).

[0358]

5 <Reference Example 7-43>

[0359]

[Chemical Formula 51]



[0360]

(3S*, 4R*)-4-(6-Fluoro-1-hydroxy-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester

5 A 1 mol/L hydrochloric acid (15 mL) was added to a solution of
of
(3S*, 4R*)-4-{1-[(tert-butyldimethylsilyl)oxy]-6-fluoro-2,3-dihydro-1H-inden-5-yl}-2-oxopyrrolidine-3-carboxylic acid methyl ester (2.0 g) in tetrahydrofuran (25 mL) under an argon atmosphere
10 to produce a reaction solution. The reaction solution was stirred at room temperature for 8 hours. The reaction solution was extracted with ethyl acetate, and the extract was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue
15 was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by 1 : 4), to obtain the title compound as a colorless amorphous (1.37 g).


¹H-NMR (400 MHz, CDCl₃) δ 1.80 (1H, dd, J = 7.3, 4.3 Hz), 1.92-2.01 (1H, m), 2.49-2.57 (1H, m), 2.73-2.81 (1H, m), 2.96-3.03 (1H, m),
20 3.44 (1H, t, J = 8.9 Hz), 3.70 (1H, dd, J = 9.5, 2.1 Hz), 3.78-3.83 (4H, m), 4.27 (1H, q, J = 8.8 Hz), 5.21 (1H, q, J = 6.3 Hz), 5.87 (1H, s), 7.11 (1H, d, J = 4.3 Hz), 7.13 (1H, s).

[0361]

<Reference Example 7-44>

[0362]

[Chemical Formula 52]



[0363]

(3S*, 4R*)-4-(6-Fluoro-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester

10 of

10% Palladium on carbon (100 mg) was added to a solution of (3S*, 4R*)-4-(6-fluoro-1-hydroxy-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidine-3-carboxylic acid methyl ester (200 mg) in methanol (5 mL) under an argon atmosphere to produce a reaction solution. The reaction solution was stirred at room temperature under a hydrogen atmosphere for 4 hours. The reaction solution was filtered over Celite and washed with ethyl acetate. The solvent was removed under reduced pressure to obtain the title compound as a colorless amorphous (170 mg).


20 ¹H-NMR (400 MHz, CDCl₃) δ 2.05-2.13 (2H, m), 2.83-2.89 (4H, m), 3.44 (1H, t, J = 8.9 Hz), 3.70 (1H, d, J = 9.8 Hz), 3.76-3.81 (4H, m), 4.25 (1H, q, J = 8.8 Hz), 5.67 (1H, s), 6.92 (1H, d, J = 10.4 Hz), 7.08 (1H, d, J = 6.7 Hz).

[0364]

<Reference Example 8-1>

[0365]

[Chemical Formula 53]



[0366]

5 (-)-(3S*,4R*)-4-(4-Methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid

A 2 mol/L sodium hydroxide aqueous solution (0.52 mL) was added to a solution of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester (130 mg) in methanol (2.6 mL) to produce a
10 reaction solution. The reaction solution was stirred at 60°C for 1 hour. A 1 mol/L hydrochloric acid was added to the reaction solution to make the reaction solution acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was
15 washed with water and a s brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the crude product was washed with ethyl acetate-diisopropyl ether, to obtain the title compound as a white solid (112 mg).

20 ¹H-NMR (400 MHz, DMSO-d₆) δ 3.16 (1H, t, J = 9.4 Hz), 3.42 (1H, d, J = 10.9 Hz), 3.55 (1H, t, J = 8.2 Hz), 3.72 (3H, s), 3.79 (1H, q, J = 9.5 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz), 8.03 (1H, s), 12.54 (1H, brs).






$[\alpha]_D^{27} = -68$ (c 0.15, EtOH)

[0367]

The same method as in Reference Example 8-1 was performed using a corresponding ester substance to obtain the following
5 Reference Examples 8-2 to 8-41. The structures and spectral data thereof are shown in Tables 26 to 35.





[0368]

[Table 26]

Ref.No	Str.	Chemical name	P.D.
8-2		(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.16 (1H, t, J = 9.4 Hz), 3.47 (1H, d, J = 10.9 Hz), 3.54 (1H, t, J = 8.2 Hz), 3.75-3.84 (1H, m), 3.80 (3H, s), 7.06-7.12 (2H, m), 7.24-7.28 (1H, m), 8.05 (1H, s), 12.65 (1H, brs).
8-3		(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.1 Hz), 3.48 (1H, d, J = 10.3 Hz), 3.55 (1H, t, J = 9.1 Hz), 3.74 (3H, s), 4.00 (1H, q, J = 10.3), 6.75-6.83 (2H, m), 7.37 (1H, t, J = 8.8 Hz), 8.09 (1H, s), 12.66 (1H, brs).
8-4		(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.11 (1H, t, J = 9.2 Hz), 3.54-3.61 (2H, m), 3.75 (3H, s), 4.20 (1H, q, J = 9.0 Hz), 6.93 (1H, dd, J = 8.9, 2.8 Hz), 7.02 (1H, d, J = 2.8 Hz), 7.49 (1H, d, J = 8.9 Hz), 8.10 (1H, s), 12.67 (1H, brs).
8-5		(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.28 (3H, s), 3.08 (1H, t, J = 9.1 Hz), 3.44 (1H, d, J = 10.3 Hz), 3.55 (1H, t, J = 9.1 Hz), 3.70 (3H, s), 3.97-4.04 (1H, m), 6.72-6.76 (2H, m), 7.33 (1H, d, J = 7.9 Hz), 8.04 (1H, s), 12.61 (1H, brs).
8-6		(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.15 (1H, t, J = 9.5 Hz), 3.52-3.57 (2H, m), 3.83 (1H, q, J = 9.6 Hz), 3.88 (3H, s), 7.15-7.22 (2H, m), 8.08 (1H, s), 12.68 (1H, brs).





[0369]

[Table 27]

Ref.No	Str.	Chemical name	P.D.
8-7		(3S*, 4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.5 Hz), 3.49-3.55 (2H, m), 3.82 (3H, s), 4.02 (1H, q, J = 9.5 Hz), 7.10 (1H, dd, J = 12.2, 7.3 Hz), 7.44 (1H, dd, J = 12.2, 7.3 Hz), 8.11 (1H, s), 12.67 (1H, brs).
8-8		(-)-(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25 (1H, t, J = 9.2 Hz), 3.43 (1H, d, J = 10.4 Hz), 3.56 (1H, t, J = 9.2 Hz), 3.76 (3H, s), 4.14 (1H, q, J = 9.4 Hz), 6.73-6.80 (2H, m), 8.20 (1H, s), 12.77 (1H, brs). [α] _D ²³ = -121 (c 0.10, EtOH)
8-9		(3S*, 4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.10-3.17 (3H, m), 3.41 (1H, d, J = 11.0 Hz), 3.53 (1H, t, J = 8.9 Hz), 3.77 (1H, q, J = 9.6 Hz), 4.48 (2H, t, J = 8.6 Hz), 6.68 (1H, d, J = 8.6 Hz), 7.02 (1H, d, J = 8.6 Hz), 7.20 (1H, s), 8.03 (1H, s), 12.62 (1H, brs).
8-10		(-)-(3S*, 4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.09-3.19 (3H, m), 3.46 (1H, d, J = 10.4 Hz), 3.52 (1H, t, J = 8.9 Hz), 3.94-4.03 (1H, m), 4.55 (2H, t, J = 8.6 Hz), 6.65 (1H, d, J = 11.0 Hz), 7.31 (1H, d, J = 7.9 Hz), 8.09 (1H, s), 12.65 (1H, brs). [α] _D ²⁷ = -114 (c 0.30, EtOH)






[0370]

[Table 28]

Ref.No	Str.	Chemical name	P.D.
8-11		(3S*,4R*)-4-(7-fluorochroman-6-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.84–1.90 (2H, m), 2.67 (2H, t, J = 6.1 Hz), 3.16 (1H, t, J = 9.2 Hz), 3.45 (1H, d, J = 10.4 Hz), 3.52 (1H, t, J = 8.9 Hz), 3.94 (1H, q, J = 9.4 Hz), 4.10 (2H, t, J = 5.2 Hz), 6.56 (1H, d, J = 12.2 Hz), 7.13 (1H, d, J = 8.6 Hz), 8.08 (1H, s), 12.67 (1H, brs).
8-12		(3S*,4S*)-4-(3-fluoro-5-methoxy-pyridin-2-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28 (1H, t, J = 8.9 Hz), 3.60–3.66 (2H, m), 3.83 (3H, s), 4.21 (1H, q, J = 9.0 Hz), 7.44 (1H, dd, J = 11.6, 2.4 Hz), 8.06 (1H, s), 8.17 (1H, d, J = 2.4 Hz), 12.70 (1H, brs).
8-13		(3S*,4S*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.4 Hz), 3.30 (1H, d, J = 10.3 Hz), 3.55–3.59 (1H, m), 3.79 (3H, s), 3.91 (1H, q, J = 10.3), 6.12 (1H, d, J = 4.2 Hz), 6.59 (1H, d, J = 4.2 Hz), 8.08 (1H, s), 12.78 (1H, brs).
8-14		(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.05 (3H, d, J = 6.1 Hz), 3.16 (1H, dd, J = 11.5, 9.1 Hz), 3.49–3.58 (2H, m), 3.72 (3H, s), 6.88 (2H, d, J = 8.5 Hz), 7.27 (2H, d, J = 8.5 Hz), 8.11 (1H, s), 12.54 (1H, s).






[0371]

[Table 29]

Ref.No	Str.	Chemical name	P.D.
8-15		(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.66 (3H, d, J = 6.7 Hz), 3.72 (3H, s), 3.77 (1H, d, J = 10.9 Hz), 3.81-3.89 (1H, m), 3.94 (1H, dd, J = 10.9, 7.3 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.20 (2H, d, J = 8.5 Hz), 8.12 (1H, s), 12.61 (1H, s).
8-16		(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.49 (3H, s), 3.22 (1H, t, J = 9.4 Hz), 3.50 (1H, d, J = 9.7 Hz), 3.62 (1H, t, J = 8.8 Hz), 3.86 (1H, q, J = 9.5 Hz), 7.26 (2H, d, J = 8.5 Hz), 7.32 (2H, d, J = 8.5 Hz), 8.10 (1H, brs).
8-17		(3S*,4R*)-4-(4-ethyl-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.15 (3H, t, J = 7.6 Hz), 2.59 (2H, q, J = 7.6 Hz), 3.27 (1H, t, J = 9.2 Hz), 3.46 (1H, d, J = 10.4 Hz), 3.58 (1H, t, J = 9.2 Hz), 4.20 (1H, q, J = 9.4 Hz), 6.99 (2H, dd, J = 12.8, 3.1 Hz), 8.22 (1H, s), 12.81 (1H, brs).
8-18		(3R*,4S*)-4-phenyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.47 (1H, t, J = 9.1 Hz), 3.57 (1H, d, J = 10.3 Hz), 3.82 (1H, t, J = 9.1 Hz), 4.05 (1H, q, J = 9.4 Hz), 6.71 (1H, s), 7.24-7.37 (5H, m).
8-19		(+)-(3R*,4S*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.16 (1H, t, J = 9.4 Hz), 3.42 (1H, d, J = 10.9 Hz), 3.55 (1H, t, J = 8.2 Hz), 3.72 (3H, s), 3.79 (1H, q, J = 9.5 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz), 8.03 (1H, s), 12.54 (1H, brs). [α] _D ²⁵ = +99 (c 0.19, EtOH)





[0372]

[Table 30]

Ref.No	Str.	Chemical name	P.D.
8-20		(3R*,4S*)-methyl-4-(3-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.46 (1H, t, J = 9.4 Hz), 3.56 (1H, d, J = 9.7 Hz), 3.75-3.83 (1H, m), 3.79 (3H, s), 4.01 (1H, q, J = 9.4 Hz), 6.58 (1H, s), 6.79-6.89 (3H, m), 7.26 (1H, t, J = 8.0 Hz).
8-21		(3R*,4S*)-methyl-4-(2-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.58 (1H, t, J = 9.1 Hz), 3.71 (1H, t, J = 9.1 Hz), 3.86 (3H, s), 3.98 (1H, d, J = 10.3 Hz), 4.06 (1H, t, J = 9.4 Hz), 6.31 (1H, brs), 6.91 (1H, d, J = 8.6 Hz), 6.93-6.98 (1H, m), 7.26-7.32 (2H, m).
8-22		(+)-(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25 (1H, t, J = 9.2 Hz), 3.43 (1H, d, J = 10.4 Hz), 3.56 (1H, t, J = 9.2 Hz), 3.76 (3H, s), 4.14 (1H, q, J = 9.4 Hz), 6.73-6.80 (2H, m), 8.20 (1H, s), 12.77 (1H, brs). [α] _D ²³ = +120 (c 0.10, EtOH)
8-23		(3R*,4S*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.05 (3H, d, J = 6.1 Hz), 3.16 (1H, dd, J = 11.5, 9.1 Hz), 3.49-3.58 (2H, m), 3.72 (3H, s), 6.88 (2H, d, J = 8.5 Hz), 7.27 (2H, d, J = 8.5 Hz), 8.11 (1H, s), 12.54 (1H, s).
8-24		(3R*,4S*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.66 (3H, d, J = 6.7 Hz), 3.72 (3H, s), 3.77 (1H, d, J = 10.9 Hz), 3.81-3.89 (1H, m), 3.94 (1H, dd, J = 10.9, 7.3 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.20 (2H, d, J = 8.5 Hz), 8.12 (1H, s), 12.61 (1H, s).



[0373]

[Table 31]

Ref.No	Str.	Chemical name	P.D.
8-25		(3R*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.4 Hz), 3.30 (1H, d, J = 10.3 Hz), 3.55-3.59 (1H, m), 3.79 (3H, s), 3.91 (1H, q, J = 10.3), 6.12 (1H, d, J = 4.2 Hz), 6.59 (1H, d, J = 4.2 Hz), 8.08 (1H, s), 12.78 (1H, brs).
8-26		(3R*,4S*)-4-(4-(difluoromethoxy)phenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.46 (1H, t, J = 9.7 Hz), 3.54 (1H, d, J = 9.7 Hz), 3.83 (1H, t, J = 9.4 Hz), 4.05 (1H, q, J = 9.4 Hz), 6.50 (1H, t, J = 73.9 Hz), 6.80 (1H, s), 7.13 (2H, d, J = 9.1 Hz), 7.31 (2H, d, J = 9.1 Hz).
8-27		(±)-trans-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.16 (1H, t, J = 9.4 Hz), 3.42 (1H, d, J = 10.9 Hz), 3.55 (1H, t, J = 8.2 Hz), 3.72 (3H, s), 3.79 (1H, q, J = 9.5 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz), 8.03 (1H, s), 12.54 (1H, brs).
8-28		(±)-trans-4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.78 (3H, s), 1.21 (3H, s), 3.50 (1H, d, J = 12.1 Hz), 3.73 (3H, s), 3.98 (1H, d, J = 12.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.27 (2H, d, J = 8.5 Hz), 8.09 (1H, s), 12.50 (1H, brs).






[0374]

[Table 32]

Ref.No	Str.	Chemical name	P.D.
8-29	 (±)-trans	(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonane-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ1.00–1.81 (8H, m), 3.69 (1H, d, J = 11.6 Hz), 3.72 (3H, s), 3.88 (1H, d, J = 11.6 Hz), 6.88 (2H, d, J = 8.6 Hz), 7.28 (2H, d, J = 8.6 Hz), 8.37 (1H, s), 12.48 (1H, brs).
8-30	 (±)-trans	(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decane-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ0.80–0.89 (1H, m), 0.94–1.09 (1H, m), 1.16–1.37 (2H, m), 1.41–1.49 (2H, m), 1.56–1.82 (4H, m), 3.55 (1H, d, J = 11.5 Hz), 3.81 (3H, s), 3.86 (1H, d, J = 11.5 Hz), 6.72 (1H, brs), 6.90 (2H, d, J = 9.1 Hz), 7.15 (2H, d, J = 9.1 Hz).

[0375]

[Table 33]

Ref.No.	Str.	Chemical name	P.D.
8-31		(3S*,4R*)-4-[4-(difluoromethoxy)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.58 (1H, t, J = 9.7 Hz), 3.69–3.74 (1H, m), 3.82 (1H, d, J = 10.9 Hz), 4.35 (1H, dt, J = 10.3, 9.7 Hz), 6.26 (1H, br), 6.51 (1H, t, J = 72.1 Hz), 6.74–6.80 (2H, m).
8-32		(3S*,4R*)-4-(4-ethoxy-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.29 (3H, t, J = 6.3 Hz), 3.24 (1H, t, J = 9.2 Hz), 3.42 (1H, d, J = 9.8 Hz), 3.56 (1H, t, J = 9.2 Hz), 4.03 (2H, q, J = 6.3 Hz), 4.14 (1H, ddd, J = 9.8, 9.2, 9.2 Hz), 6.74 (2H, d, J = 10.4 Hz), 8.19 (1H, s), 12.81 (1H, s).
8-33		(3S*,4R*)-4-[4-(dimethylamino)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.88 (6H, s), 3.21 (1H, t, J = 9.2 Hz), 3.37 (1H, d, J = 10.4 Hz), 3.50 (1H, t, J = 9.2 Hz), 4.08 (1H, q, J = 9.4 Hz), 6.36 (2H, d, J = 12.8 Hz), 8.11 (1H, s), 12.75 (1H, s).
8-34		(3S*,4R*)-4-(6-fluorobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.32 (1H, t, J = 9.1 Hz), 3.52 (1H, d, J = 9.2 Hz), 3.63 (1H, t, J = 9.1 Hz), 4.09–4.16 (1H, m), 6.95 (1H, d, J = 2.1 Hz), 7.56 (1H, d, J = 11.0 Hz), 7.71 (1H, d, J = 7.9 Hz), 8.00 (1H, d, J = 2.1 Hz), 8.06 (1H, s), 12.78 (1H, brs).
8-35		(3S*,4R*)-4-(6-methoxypyridin-3-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 3.43–3.51 (2H, m), 3.82 (1H, t, J = 9.2 Hz), 3.93 (3H, s), 4.01 (1H, q, J = 9.4 Hz), 6.37 (1H, brs), 6.77 (1H, d, J = 8.6 Hz), 7.56 (1H, dd, J = 8.6, 2.4 Hz), 8.15 (1H, d, J = 2.4 Hz).


[0376]

[Table 34]

Ref.No.	Str.	Chemical name	P.D.
8-36		(3S*,4R*)-4-(4-acetamido-2,6-difluorophenyl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.05 (3H, s), 3.26 (1H, t, J = 9.5 Hz), 3.44 (1H, d, J = 9.8 Hz), 3.57 (1H, t, J = 9.5 Hz), 4.16 (1H, q, J = 9.5 Hz), 7.30 (2H, d, J = 11.0 Hz), 8.21 (1H, s), 10.30 (1H, s), 12.80 (1H, brs).
8-37		(3S*,4R*)-4-{4-[(tert-butoxycarbonyl)(methyl)amino]-2,6-difluorophenyl}-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 1.49 (9H, s), 3.25 (3H, s), 3.58 (1H, t, J = 9.5 Hz), 3.70 (1H, t, J = 9.2 Hz), 3.84 (1H, d, J = 11.0 Hz), 4.34 (1H, q, J = 9.8 Hz), 6.16 (1H, brs), 6.94 (2H, d, J = 10.4 Hz).
8-38		(3S*,4R*)-4-(4,6-difluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17-3.26 (3H, m), 3.42 (1H, d, J = 10.4 Hz), 3.54 (1H, t, J = 9.2 Hz), 4.14 (1H, q, J = 9.6 Hz), 4.63 (2H, t, J = 8.6 Hz), 6.64 (1H, d, J = 11.0 Hz), 8.19 (1H, s), 12.53 (1H, brs).
8-39		(3S*,4R*)-4-(6-fluoro-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.97-2.04 (2H, m), 2.76-2.84 (4H, m), 3.18 (1H, t, J = 9.2 Hz), 3.47 (1H, d, J = 9.8 Hz), 3.55 (1H, t, J = 8.6 Hz), 3.98-4.05 (1H, m), 7.03 (1H, d, J = 11.0 Hz), 7.28 (1H, d, J = 7.3 Hz), 8.08 (1H, s), 12.71 (1H, brs).
8-40		(3S*,4R*)-4-(1-benzyl-6-fluoroindolin-5-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, CDCl ₃) δ 2.93-2.97 (2H, m), 3.43-3.51 (3H, m), 3.68-3.78 (2H, m), 3.97 (1H, q, J = 9.2 Hz), 4.26 (2H, s), 6.15 (1H, br), 6.28 (1H, d, J = 12.2 Hz), 7.01 (1H, d, J = 6.7 Hz), 7.30-7.38 (5H, m).

[0377]

[Table 35]


Ref.No.	Str.	Chemical name	P.D.
8-41		(3S*,4R*)-4-(3,5-difluoropyridin-4-yl)-2-oxopyrrolidine-3-carboxylic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28 (1H, t, J = 9.2 Hz), 3.43 (1H, d, J = 9.2 Hz), 3.64 (1H, t, J = 9.5 Hz), 4.30 (1H, q, J = 8.8 Hz), 8.18 (1H, s), 8.51 (2H, s), 13.00 (1H, brs).

[0378]

5 <Reference Example 8-42>

[0379]

[Chemical Formula 54]



[0380]

10 (3S*,4R*)-4-(6-Fluoro-1-hydroxy-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidine-3-carboxylic acid

A 2 mol/L sodium hydroxide aqueous solution (0.44 mL) was added to a solution of

(3S*,4R*)-4-{1-[(tert-butyldimethylsilyl)oxy]-6-fluoro-2,3-dihydro-1H-inden-5-yl}-2-oxopyrrolidine-3-carboxylic acid methyl ester (180 mg) in methanol (0.88 mL) to produce a reaction solution. The reaction solution was stirred at 60°C for 2 hours. A 1 mol/L hydrochloric acid was added to the reaction solution to make the

reaction solution acidic (pH: 1), and the reaction solution was stirred at room temperature for 1 hour, and extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed, and the crude product was washed with a mixed solution of ethyl acetate and methanol at a ratio of 6 : 1, to obtain the title compound as a white solid (110 mg).


¹H-NMR (400 MHz, DMSO-d₆) δ 1.74-1.81 (1H, m), 2.29-2.36 (1H, m), 2.61-2.69 (1H, m), 2.81-2.87 (1H, m), 3.19 (1H, t, J = 9.2 Hz), 3.49 (1H, d, J = 10.4 Hz), 3.56 (1H, t, J = 9.2 Hz), 4.04 (1H, q, J = 9.2 Hz), 4.99 (1H, q, J = 6.7 Hz), 5.32 (1H, d, J = 5.5 Hz), 7.06 (1H, d, J = 10.4 Hz), 7.30 (1H, d, J = 7.3 Hz), 8.11 (1H, s), 12.69 (1H, brs).

[0381]

<Reference Example 9-1>

[0382]

[Chemical Formula 55]



[0383]

(4R*)-3-Chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester

Sulfuryl chloride (200 μL) was added to a solution of (-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester (250 mg) in tetrahydrofuran (5 mL) under an

argon atmosphere to produce a reaction solution. The reaction solution was stirred at room temperature for 2 hours. A saturated aqueous sodium hydrogen carbonate solution was added to the reaction solution under ice-cooling, and the reaction solution
5 was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 9 : 1 followed by ethyl acetate), to obtain
10 (4R^{*})-3-chloro-4-(4-methoxyphenyl)2-oxopyrrolidine-3-carboxylic acid methyl ester as an intermediate compound.

To a solution of the obtained (4R^{*})-3-chloro-4-(4-methoxyphenyl)2-oxopyrrolidine-3-carboxylic acid methyl ester (222 mg) in N,N-dimethylformamide (3.4 mL),
15 sodium hydride (56 mg, 60% in oil) was added under ice-cooling to produce a reaction solution. The reaction solution was stirred for 5 minutes, then iodomethane (97 μ L) was added, and the reaction solution was stirred at room temperature for 1 hour. A 1 mol/L hydrochloric acid was added to the reaction solution, and the
20 mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 9 : 1 followed by ethyl
25 acetate), to obtain the title compound as a white solid (184 mg).


¹H-NMR (400 MHz, CDCl₃) δ 3.07 (3H, s), 3.50 (3H, s), 3.61-3.65 (1H, m), 3.80 (3H, s), 3.88-3.99 (2H, m), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 8.5 Hz).

[0384]

5 <Reference Example 9-2>

[0385]

[Chemical Formula 56]



[0386]

(4S*)-3-Chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-

10 3-carboxylic acid methyl ester

The same method as in Reference Example 9-1 was performed using

(+)-(3R*, 4S*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

15 (-)-(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 3.07 (3H, s), 3.50 (3H, s), 3.61-3.65 (1H, m), 3.80 (3H, s), 3.88-3.99 (2H, m), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 8.5 Hz).

20 [0387]

<Reference Example 9-3>

[0388]

[Chemical Formula 57]



(±)-3-Chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester

The same method as in Reference Example 9-1 was performed

5 using

(±)-trans-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.


10 ¹H-NMR (400 MHz, CDCl₃) δ 3.07 (3H, s), 3.50 (3H, s), 3.61-3.65 (1H, m), 3.80 (3H, s), 3.88-3.99 (2H, m), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 8.5 Hz).

[0390]

<Reference Example 10-1>

15 [0391]

[Chemical Formula 58]



(3S*,4R*)-4-(4-Methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid

20 Metal zinc (595 mg) and acetic acid (0.3 mL) were added to

a solution of

(4R^{*})-3-chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester (184 mg) in methanol (11 mL) to produce a reaction solution. The reaction solution was stirred
5 at room temperature for 1 hour. The reaction solution was filtered over Celite to remove the insoluble, and the solvent was removed under reduced pressure. A saturated aqueous sodium hydrogen carbonate solution was added to the residue, and the mixture was extracted with ethyl acetate. The organic layer was washed with
10 a 1 mol/L hydrochloric acid and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane followed by ethyl acetate : hexane = 1 :
1), to obtain

15 (3S^{*},4R^{*})-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester as an intermediate compound.

A 2 mol/L sodium hydroxide aqueous solution (0.57 mL) was added to a solution of the obtained

(3S^{*},4R^{*})-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester (130 mg) in methanol (2.8 mL) to produce
20 a reaction solution. The reaction solution was stirred at 70°C for 1 hour. A 1 mol/L hydrochloric acid was added to the reaction solution to make the reaction solution acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was
25 washed with water and a brine successively, and dried over anhydrous

sodium sulfate. The solvent was removed under reduced pressure, and the crude product was washed with ethyl acetate-diisopropyl ether, to obtain the title compound as a white solid (133 mg).


¹H-NMR (400 MHz, DMSO-d₆) δ 2.78 (3H, s), 3.27-3.34 (1H, m), 3.45 (1H, d, J = 10.3 Hz), 3.65 (1H, t, J = 8.8 Hz), 3.70-3.77 (1H, m), 3.72 (3H, s), 6.89 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz), 12.70 (1H, brs).

[0393]

<Reference Example 10-2>

10 [0394]

[Chemical Formula 59]



[0395]

(3R*, 4S*)-4-(4-Methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid

15 [0396]

The same method as in Reference Example 10-1 was performed using

(4S*)-3-chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

20 (4R*)-3-chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 2.78 (3H, s), 3.27-3.34 (1H, m), 3.45 (1H, d, J = 10.3 Hz), 3.65 (1H, t, J = 8.8 Hz), 3.70-3.77 (1H, m),


3.72 (3H, s), 6.89(2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz),
12.70 (1H, brs).

[0397]

<Reference Example 10-3>

5 [0398]

[Chemical Formula 60]



[0399]

(±)-trans-4-(4-Methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid

10 The same method as in Reference Example 10-1 was performed using

(±)-3-Chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

(4R^{*})-3-chloro-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidine-

15 3-carboxylic acid methyl ester to obtain the title compound.


¹H-NMR (400 MHz, DMSO-d₆) δ 2.78 (3H, s), 3.27-3.34 (1H, m), 3.45 (1H, d, J = 10.3 Hz), 3.65 (1H, t, J = 8.8 Hz), 3.70-3.77 (1H, m), 3.72 (3H, s), 6.89 (2H, d, J = 8.5 Hz), 7.24 (2H, d, J = 8.5 Hz), 12.70 (1H, brs).

20 [0400]

<Reference Example 11-1>

[0401]

[Chemical Formula 61]



[0402]

(-)-[(3S*,4R*)-4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbam
ic acid benzyl ester

5 Triethylamine (3.95 mL) and diphenylphosphoryl azide (6.2
mL) were added to a solution of
(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxyl
ic acid (6.04 g) in toluene (128 mL) to produce a reaction solution.
The reaction solution was stirred at room temperature for 4.5 hours.
10 The reaction temperature of the reaction solution was heated up
to 80°C, and the reaction solution was then stirred for 30 minutes.
To the reaction solution, benzyl alcohol (13.3 mL) was added, and
the reaction solution was stirred at 120°C for 5 hours. The reaction
solution was concentrated under reduced pressure, and the residue
15 was then purified by silica gel column chromatography (ethyl
acetate followed by ethyl acetate : methanol = 10 : 1), to obtain
the title compound as a white solid (6.3 g).

¹H-NMR (400 MHz, CDCl₃) δ 3.36 (1H, t, J = 9.1 Hz), 3.49-3.70 (2H,
m), 3.80 (3H, s), 4.42 (1H, dd, J = 11.5, 8.5 Hz), 5.07 (2H, s),
20 5.16 (1H, brs), 5.98 (1H, brs), 6.89 (2H, d, J = 7.9 Hz), 7.22 (2H,
d, J = 7.9 Hz), 7.20-7.40 (5H, m).


[α]_D²⁷ = -79 (c 0.17, EtOH)

[0403]

<Reference Example 11-2>

[0404]

[Chemical Formula 62]



5 [0405]

(+)-[(3R*,4S*)-4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbam
mic acid benzyl ester

The same method as in Reference Example 11-1 was performed
using

10 (+)-(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidi
ne-3-carboxylic acid in place of
(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxyl
ic acid to obtain the title compound.

¹H-NMR (400 MHz, CDCl₃) δ 3.36 (1H, t, J = 9.1 Hz), 3.49-3.70 (2H,
15 m), 3.80 (3H, s), 4.42 (1H, dd, J = 11.5, 8.5 Hz), 5.07 (2H, s),
5.16 (1H, brs), 5.98 (1H, brs), 6.89 (2H, d, J = 7.9 Hz), 7.22 (2H,
d, J = 7.9 Hz), 7.20-7.40 (5H, m).


[α]_D²⁷ = +81 (c 0.16, EtOH)

[0406]

20 <Reference Example 12-1>

[0407]

[Chemical Formula 63]



(-)-[(3S*,4R*)-4-(2-Fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester

Triethylamine (1.0 mL) and diphenylphosphoryl azide (1.5
 5 mL) were added to a solution of
 (3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid (1.6 g) in a mixture of toluene (63 mL) and acetonitrile (10 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. The reaction
 10 temperature of the reaction solution was heated up to 80°C, and the reaction solution was then stirred for 30 minutes. To the reaction solution, benzyl alcohol (3.3 mL) was added, and the reaction solution was stirred at 110°C for 3.5 hours. Ethyl acetate was added to the reaction solution, and the mixture was washed with
 15 a 1 mol/L hydrochloric acid, water, and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate: methanol = 10 : 1), to obtain the title compound
 20 as a white solid (767 mg).

¹H-NMR (400 MHz, CDCl₃) δ 3.40 (1H, t, J = 8.6 Hz), 3.63-3.82 (2H, m), 3.79 (3H, s), 4.62 (1H, dd, J = 11.0, 8.6 Hz), 5.07 (2H, brs), 5.16 (1H, brs), 6.01 (1H, brs), 6.58-6.73 (2H, m), 7.25-7.33 (6H,

m).




$[\alpha]_D^{24} = -134$ (c 0.16, EtOH)

[0409]

The same method as in Reference Example 12-1 was performed
5 using a corresponding carboxylic acid substance to obtain the
following Reference Examples 12-2 to 12-4. The structures and
spectral data thereof are shown in Table 36.

[0410]

[Table 36]


Ref.No	Str.	Chemical name	P.D.
12-2		(-)-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.48-3.62 (2H, m), 3.79 (3H, s), 3.80-3.96 (1H, m), 4.67-4.75 (1H, m), 5.05 (2H, s), 5.38 (1H, brd, J = 8.0 Hz), 6.42-6.53 (2H, m), 6.60 (1H, s), 7.26-7.36 (5H, m). [α] _D ²⁴ = -107 (c 0.10, EtOH)
12-3		(-)-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.05-3.25 (2H, m), 3.37 (1H, t, J = 9.2 Hz), 3.55-3.80 (2H, m), 4.50-4.65 (3H, m), 5.04-5.14 (3H, m), 5.83 (1H, s), 6.50 (1H, d, J = 10.4 Hz), 7.10-7.22 (1H, m), 7.26-40 (5H, m). [α] _D ²⁹ = -185 (c 0.16, EtOH)
12-4		(-)-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.36 (1H, m), 3.45 (1H, t, J = 8.9 Hz), 3.83 (3H, s), 3.91-3.98 (1H, m), 4.38 (1H, t, J = 9.8 Hz), 4.93 (1H, d, J = 12.8 Hz), 4.98 (1H, d, J = 12.8 Hz), 7.27-7.35 (5H, m), 7.40 (1H, dd, J = 11.6, 1.8 Hz), 7.61 (1H, d, J = 8.6 Hz), 7.94 (1H, s), 8.20 (1H, d, J = 1.8 Hz). [α] _D ²⁹ = -126 (c 0.12, EtOH)

[0411]

<Reference Example 13-1>

[0412]

[Chemical Formula 64]



5 [0413]

(-)-(3S*, 4R*)-3-Amino-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-2-one

10 10% Palladium on carbon (81 mg) was added to a solution of (-)-[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester (811 mg) in ethanol (30 mL) to produce a reaction solution. The reaction solution was stirred under a hydrogen atmosphere for 2 hours. The reaction solution was filtered over Celite, and the solvent was removed under reduced pressure. The residue was purified by silica gel column

15 chromatography (hexane : ethyl acetate = 1 : 1 followed by ethyl acetate: methanol = 9 : 1), to obtain the title compound as a white solid (520 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.22 (1H, t, J = 8.0 Hz), 3.34-3.43 (2H, m), 3.47 (1H, d, J = 9.8 Hz), 3.76 (3H, s), 6.74 (2H, d, J = 11.0 Hz), 7.88 (1H, s).

20


[α]_D²⁴ = -90 (c 0.11, EtOH)

[0414]

<Reference Example 13-2>

[0415]

[Chemical Formula 65]



[0416]

- 5 (3S*, 4R*)-3-Amino-4-(3-fluoro-5-methoxypyridin-2-yl)pyrrolidin-2-one

The same method as in Reference Example 13-1 was performed using

(-)-[(3S*, 4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester in place of
10 (-)-[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title compound.


¹H-NMR (400 MHz, DMSO-d₆) δ 1.84 (2H, s), 3.22 (1H, t, J = 9.2 Hz),
3.40 (1H, t, J = 8.6 Hz), 3.51-3.58 (1H, m), 3.61 (1H, d, J = 9.8
15 Hz), 3.83 (3H, s), 7.41 (1H, dd, J = 11.9, 1.4 Hz), 7.77 (1H, s),
8.19 (1H, d, J = 1.4 Hz).

[0417]

<Reference Example 14-1>

[0418]

- 20 [Chemical Formula 66]



[0419]

(-)-[(3S*,4R*)-4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester

Triethylamine (2.64 mL) and diphenylphosphoryl azide (4.07
5 mL) were added to a solution of
(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid (4.05 g) in toluene (80 mL) to produce a first reaction solution. The first reaction solution was stirred at room temperature for 5 hours. The reaction temperature of the first
10 reaction solution was heated up to 80°C, and the first reaction solution was then stirred for 20 minutes. After that, the first reaction solution was concentrated under reduced pressure. To the residue, dioxane (50 mL) and a 1 mol/L hydrochloric acid (50 mL) were added to produce a second reaction solution. The second
15 reaction solution was stirred at room temperature for 30 minutes. The second reaction solution was concentrated under reduced pressure, water (40 mL) was added, and the mixture was washed with diethyl ether. The aqueous layer was concentrated under reduced pressure, and methanol (77 mL), triethylamine (10 mL), and
20 di-tert-butyl dicarbonate (4.28 g) were added to the residue to produce a third reaction solution. The third reaction solution was stirred at room temperature for 3 hours. The third reaction solution was concentrated under reduced pressure, water was added to the residue, and the mixture was extracted with ethyl acetate.
25 The organic layer was washed with a brine, and dried over anhydrous

sodium sulfate. The solvent was removed under reduced pressure, and the crude product was washed with chloroform, to obtain the title compound as a white solid (2.97 g).

¹H-NMR (400 MHz, CDCl₃) δ 1.33 (9H, s), 3.10-3.17 (1H, m), 3.37-3.45 (2H, m), 3.71 (3H, s), 4.12 (1H, t, J = 7.0 Hz), 6.87 (2H, d, J = 8.6 Hz), 7.07 (1H, d, J = 9.2 Hz), 7.23 (2H, d, J = 8.6 Hz), 7.83 (1H, brs).


[α]_D²⁷ = -64 (c 0.14, EtOH)

[0420]

10 <Reference Example 14-2>

[0421]

[Chemical Formula 67]



[0422]

(±)-trans-[4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester

The same method as in Reference Example 14-1 was performed using

(±)-trans-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain the title compound.

¹H-NMR (400 MHz, CDCl₃) δ 1.33 (9H, s), 3.10-3.17 (1H, m), 3.37-3.45


(2H, m), 3.71 (3H, s), 4.12 (1H, t, J = 7.0 Hz), 6.87 (2H, d, J = 8.6 Hz), 7.07 (1H, d, J = 9.2 Hz), 7.23 (2H, d, J = 8.6 Hz), 7.83 (1H, brs).

[0423]

5 <Reference Example 15-1>

[0424]

[Chemical Formula 68]



0425]

[(3S*, 4R*)-1-Ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]c

10 arbamic acid tert-butyl ester

A solution of

(-)-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carba
mic acid tert-butyl ester (181 mg) in N,N-dimethylformamide (0.9
mL) and a solution of ethyl iodide (50 μ L) in tetrahydrofuran (1.2
15 mL) were added to a solution of sodium hydride (42 mg, 60% in oil)
in tetrahydrofuran (2.3 mL) under cooling, and the mixture was
stirred at room temperature for 4 hours. Water was added to the
reaction mixture, and the mixture was extracted with ethyl acetate.
The organic layer was washed with a brine, and dried over anhydrous
20 sodium sulfate. The solvent was removed under reduced pressure,
and the residue was then purified by silica gel column
chromatography (ethyl acetate : hexane = 4 : 1), to obtain the

title compound as a white solid (65 mg).


¹H-NMR (400 MHz, CDCl₃) δ 1.17 (3H, t, J = 7.4 Hz), 1.38 (9H, s), 3.31-3.60 (5H, m), 3.80 (3H, s), 4.30 (1H, t, J = 9.2 Hz), 4.84 (1H, brs), 6.89 (2H, d, J = 8.6 Hz), 7.25 (2H, d, J = 8.6 Hz).

5 [0426]

<Reference Example 15-2>

[0427]

[Chemical Formula 69]



[0428]

10 2-{{(3S*,4R*)-3-[(tert-Butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}acetic acid ethyl ester

The same method as in Reference Example 15-1 was performed using bromoacetic acid ethyl ester in place of ethyl iodide as an alkylating agent to obtain the title compound.


15 ¹H-NMR (400 MHz, CDCl₃) δ 1.30 (3H, t, J = 7.3 Hz), 1.39 (9H, s), 3.50-3.65 (3H, m), 3.80 (3H, s), 3.96 (1H, d, J = 17.8 Hz), 4.21 (2H, q, J = 7.3 Hz), 4.31 (1H, d, J = 17.8 Hz), 4.34 (1H, t, J = 8.6 Hz), 4.87 (1H, brs), 6.89 (2H, d, J = 8.6 Hz), 7.25 (2H, d, J = 8.6 Hz).

20 [0429]

<Reference Example 15-3>

[0430]

[Chemical Formula 70]



[0431]

2-[(3S*, 4R*)-3-[(tert-butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester

5 The same method as in Reference Example 15-1 was performed using 2-bromopropionic acid ethyl ester in place of ethyl iodide as an alkylating agent to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 1.27-1.48 (15H, m), 3.29-3.80 (3H, m), 3.80 (3H, s), 4.15-4.31 (3H, m), 4.57 (0.5H, brs), 4.80 (0.5H, brs),
10 4.88-4.99 (1H, m), 6.88 (2H, d, J = 8.6 Hz), 7.25 (2H, d, J = 9.2 Hz).

[0432]

<Reference Example 16-1>

[0433]

15 [Chemical Formula 71]



[0434]

(±)-trans-({1-[2-[(tert-butyldimethylsilyl)oxy]ethyl]-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl}carbamic acid tert-butyl ester

To a solution of
(±)-trans-(4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic
acid tert-butyl ester (1.03 g) and

(2-bromoethoxy)(tert-butyl)dimethylsilane (1.44 mL) in

5 N,N-dimethylformamide (10 mL), sodium hydride (141 mg, 60% in oil)
was added in several portions under cooling to produce a reaction
solution. The reaction solution was stirred at room temperature
for 2 hours. A saturated aqueous ammonium chloride solution was
added to the reaction solution, and the mixture was extracted with
10 ethyl acetate. The organic layer was washed with water and a brine
successively, and dried over anhydrous sodium sulfate. The
solvent was removed under reduced pressure, and the residue was
then purified by silica gel column chromatography (hexane : ethyl
acetate = 3 : 1), to obtain the title compound as a colorless oil
15 (1.1 g).


¹H-NMR (400 MHz, CDCl₃) δ 0.06 (6H, s), 0.89 (9H, s), 1.37 (9H, s),
3.30-3.47 (2H, m), 3.52 (1H, t, J = 9.7 Hz), 3.57-3.64 (1H, m),
3.71 (1H, t, J = 9.1 Hz), 3.76-3.80 (2H, m), 3.80 (3H, s), 4.33
(1H, t, J = 9.4 Hz), 4.81 (1H, brs), 6.89 (2H, d, J = 8.8 Hz), 7.22
20 (2H, d, J = 8.8 Hz).

[0435]

<Reference Example 17-1>

[0436]

[Chemical Formula 72]



[0437]

2-{{(3S*,4R*)-3-[(tert-Butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}-2-methylpropionic acid ethyl ester

Lithium hexamethyldisilazide (3.8 mL, 1.3 mol/L

5 tetrahydrofuran solution) was added to a solution of 2-{{(3S*,4R*)-3-[(tert-butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}acetic acid ethyl ester (384 mg) in tetrahydrofuran (10 mL) at -78°C, to produce a reaction solution. The reaction solution was stirred for 30 minutes. Methyl iodide 10 (0.31 mL) was added to the reaction solution, the temperature was heated up to room temperature, and the reaction solution was stirred for 24 hours. A 10% citric acid aqueous solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine 15 successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1 followed by ethyl acetate), to obtain the title compound as a white solid (222 mg).

20 ¹H-NMR (400 MHz, CDCl₃) δ 1.29 (3H, t, J = 7.4 Hz), 1.36 (9H, s), 1.50 (3H, s), 1.57 (3H, s), 3.34-3.42 (2H, m), 3.70-3.77 (1H, m), 3.80 (3H, s), 4.20 (2H, q, J = 7.4 Hz), 4.36 (1H, t, J = 10.4 Hz),


4.83 (1H, brs), 6.89 (2H, d, J = 8.6 Hz), 7.25 (2H, d, J = 8.6 Hz).

[0438]

<Reference Example 17-2>

[0439]

5 [Chemical Formula 73]



[0440]

2-((3S*, 4R*)-3-{{[(Benzyloxy)carbonyl]amino}-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-1-yl}-2-methylpropanoic acid ethyl ester

The same method as in Reference Example 17-1 was performed using

2-((3S*, 4R*)-3-{{[(benzyloxy)carbonyl]amino}-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-1-yl}acetic acid ethyl ester in place of

(-)-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 1.28 (3H, t, J = 7.0 Hz), 1.50 (3H, s), 1.56 (3H, s), 3.10-3.22 (2H, m), 3.37 (1H, t, J = 9.2 Hz), 3.52-3.66 (1H, m), 3.69-3.80 (1H, m), 4.20 (2H, q, J = 7.0 Hz), 4.57-4.65 (1H, m), 4.60 (2H, t, J = 8.6 Hz), 5.00-5.13 (3H, m), 6.51 (1H, d, J = 10.4 Hz), 7.17-7.23 (1H, m), 7.28-7.36 (5H, m).

[0441]

<Reference Example 18-1>

[0442]

[Chemical Formula 74]



5 [0443]

[(3S*, 4R*)-1-(1-Hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester

Lithium borohydride (1.5 mL, 3 mol/L tetrahydrofuran solution) was added to a solution of

10 2-[(3S*, 4R*)-3-[(tert-butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-2-methylpropionic acid ethylester (421 mg) in tetrahydrofuran (10 mL) under ice-cooling, to produce a reaction solution. The reaction solution was stirred at room temperature for 2 hours. A 10% citric acid aqueous solution was

15 added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate :

20 hexane = 4 : 1), to obtain the title compound as a white solid (88 mg).

¹H-NMR (400 MHz, CDCl₃) δ 1.30 (3H, s), 1.31 (3H, s), 1.40 (9H, s),


3.28-3.39 (2H, m), 3.65-3.75 (2H, m), 3.81 (3H, s), 3.92 (1H, brs),
4.32 (1H, t, J = 9.2 Hz), 4.57 (1H, brs), 4.86 (1H, brs), 6.89 (2H,
d, J = 8.6 Hz), 7.22 (2H, d, J = 8.6 Hz).

[0444]

5 <Reference Example 19-1>

[0445]

[Chemical Formula 75]



[0446]

[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-methyl-2-oxopyr-
10 rolidin-3-yl]carbamic acid benzyl ester

Potassium tert-butoxide (60 mg) and iodomethane (33 μ L) were
added to a solution of

(-)-[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidi-
n-3-yl]carbamic acid benzyl ester (200 mg) in

15 N,N-dimethylformamide (5.3 mL) under ice-cooling, to produce a
reaction solution. The reaction solution was stirred for 2 hours.
A saturated aqueous ammonium chloride solution was added to the
reaction solution, and the mixture was extracted with ethyl acetate.
The organic layer was washed with a brine, and dried over anhydrous
20 sodium sulfate. The solvent was removed under reduced pressure,
and the residue was then purified by silica gel column
chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl

acetate), to obtain the title compound as a white solid (100 mg).




$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.95 (3H, s), 3.40-3.62 (2H, m), 3.77 (3H, s), 3.80-3.84 (1H, m), 4.57-4.67 (1H, m), 5.04 (2H s), 5.20 (1H, brs), 6.40-6.51 (2H, m), 7.26-7.33 (5H, m).

5 [0447]

The same method as in Reference Example 19-1 was performed using a corresponding alkylating agent to obtain the following Reference Examples 19-2 to 19-9. The structures and spectral data thereof are shown in Tables 37 to 39.

[0448]

[Table 37]

Ref.No	Str.	Chemical name	P.D.
19-2		[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-propylpyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.94 (3H, t, J = 7.3 Hz), 1.52-1.65 (2H, m), 3.20-3.35 (1H, m), 3.36-3.63 (3H, m), 3.65-3.85 (1H, m), 3.77 (3H, s), 4.63-4.70 (1H, m), 5.03 (2H, s), 5.21 (1H, brs), 6.42-6.52 (2H, m), 7.25-7.35 (5H, m).
19-3		[(3S*,4R*)-1-benzyl-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.30-3.47 (2H, m), 3.73-3.82 (1H, m), 3.75 (3H, s), 4.50 (1H, d, J = 14.7 Hz), 4.61 (1H, d, J = 15.3 Hz), 4.70 (1H, t, J = 9.8 Hz), 5.04 (2H, d, J = 11.2 Hz), 5.27 (1H, brs), 6.42 (2H, d, J = 10.4 Hz), 7.26-7.37 (10H, m).
19-4		[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-(pyridin-3-ylmethyl)pyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.56 (2H, t, J = 8.0 Hz), 3.76-3.95 (4H, m), 4.61-4.79 (3H, m), 5.03-5.08 (2H, m), 5.26 (1H, brs), 6.42 (2H, d, J = 9.8 Hz), 7.20-7.31 (7H, m), 7.71 (1H, brs), 8.54 (1H, d, J = 4.9 Hz).



[0449]

[Table 38]

Ref.No	Str.	Chemical name	P.D.
19-5		2-((3S*,4R*)-3- -{[(benzyloxy) carbonyl]amino }-4-(2,6-diflu oro-4-methoxy phenyl)-2-oxopy rrolidin-1-yl) acetic acid ethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.30 (3H, t, J = 7.3 Hz), 3.47-3.60 (1H, m), 3.70-3.95 (2H, m), 3.78 (3H, s), 4.06 (1H, d, J = 17.8 Hz), 4.22 (2H, q, J = 7.3 Hz), 4.22-4.27 (1H, d, J = 17.8 Hz) 4.76 (1H, t, J = 9.2 Hz), 4.98-5.10 (2H, m), 5.19 (1H, brs), 6.40-6.55 (2H, m), 7.26-7.36 (5H, m).
19-6		2-((3S*,4R*)-3- -{[(benzyloxy) carbonyl]amino }-4-(6-fluoro- 2,3-dihydroben zofuran-5-yl)- 2-oxopyrrolidi n-1-yl)acetic acid ethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.29 (3H, t, J = 7.1 Hz), 3.09-3.23 (2H, m), 3.49-3.59 (1H, m), 3.59-3.76 (2H, m), 3.98 (1H, d, J = 17.7 Hz), 4.11-4.31 (1H, m), 4.21 (2H, q, J = 7.1 Hz), 4.58-4.66 (3H, m), 5.00-5.20 (1H, m), 5.08 (2H, s), 6.51 (1H, d, J = 11.0 Hz), 7.10-7.20 (1H, m), 7.26-7.36 (5H, m).
19-7		[(3S*,4R*)-1-(cyclopropylmet hyl)-4-(6-fluo ro-2,3-dihydro benzofuran-5-y l)-2-oxopyrrol idin-3-yl]carb amic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.22-0.26 (2H, m), 0.52-0.57 (2H, m), 0.85-0.99 (1H, m), 3.09 (1H, dd, J = 13.9, 7.3 Hz), 3.10-3.22 (2H, m), 3.37 (1H, dd, J = 13.9, 7.0 Hz), 3.43 (1H, t, J = 9.2 Hz), 3.55-3.77 (2H, m), 4.58-4.64 (3H, m), 5.02-5.17 (3H, m), 6.52 (1H, d, J = 11.0 Hz), 7.10-7.22 (1H, m), 7.28-7.36 (5H, m).

[0450]

[Table 39]


Ref.No	Str.	Chemical name	P.D.
19-8		[(3S*, 4R*)-1-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.14–3.19 (2H, m), 3.48–3.54 (1H, m), 3.70–3.80 (2H, m), 4.23–4.27 (1H, m), 4.44–4.52 (2H, m), 4.62 (2H, t, J = 9.1 Hz), 5.06–5.13 (3H, m), 6.52–6.54 (1H, m), 7.09–7.12 (1H, m), 7.32–7.34 (5H, m).
19-9		{(3S*, 4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl}carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.53 (3H, s), 3.12–3.16 (2H, m), 3.43–3.47 (1H, m), 3.67–3.73 (2H, m), 4.57–4.66 (4H, m), 4.80–4.84 (1H, m), 5.07 (2H, s), 5.42 (1H, br), 6.59 (1H, d, J = 10.9 Hz), 7.13–7.14 (1H, m), 7.27–7.32 (5H, m).

[0451]

5 <Reference Example 19-10>

[0452]

[Chemical Formula 76]



[0453]

10 {(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-[(methylthio)me

thyl]-2-oxopyrrolidin-3-yl}carbamic acid benzyl ester

Chloromethyl methyl sulfide (73 μ L), tetrabutylammonium iodide (12 mg), and potassium tert-butoxide (86 mg) were added to a solution of

5 [(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester (120 mg) in N,N-dimethylformamide (3.2 mL) under ice-cooling, to produce a reaction solution. The reaction solution was stirred at room temperature for 4 hours. A saturated aqueous ammonium chloride solution was added to the
10 reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1 followed
15 by 1 : 1), to obtain the title compound (56 mg).


¹H-NMR (400 MHz, CDCl₃) δ 2.14 (3H, brs), 3.60-3.69 (2H, m), 3.78 (3H, s), 3.85-3.93 (1H, m), 4.49 (2H, s), 4.58-4.66 (1H, m), 5.03 (1H, d, J = 12.2 Hz), 5.09 (1H, d, J = 12.2 Hz), 5.20 (1H, brs), 6.48 (2H, d, J = 9.8 Hz), 7.28-7.34 (5H, m).

20 [0454]

<Reference Example 19-11>

[0455]

[Chemical Formula 77]



[0456]

[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxo-1-(sulfamoylmethyl)pyrrolidin-3-yl]carbamic acid benzyl ester

5 The same method as in Reference Example 19-10 was performed using 1-chloromethanesulfonamide in place of chloromethyl methysulfide to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 3.60-3.70 (1H, m), 3.78 (3H, s), 4.21-4.24 (4H, m), 5.05 (1H, d, J = 12.2 Hz), 5.11-5.17 (2H, m), 5.29 (2H, 10 brs), 5.63 (1H, brs), 6.48 (2H, d, J = 10.4 Hz), 7.29-7.38 (5H, m).

[0457]

<Reference Example 19-12>

[0458]

15 [Chemical Formula 78]



[0459]

{(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-[(methylsulfonyl)methyl]-2-oxopyrrolidin-3-yl}carbamic acid benzyl ester

Meta-chloroperoxybenzoic acid (79 mg) was added to a solution of

{(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(methylthio)methyl]-2-oxopyrrolidin-3-yl}carbamic acid benzyl ester (56 mg) in dichloromethane (2.2 mL) under ice-cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 4 hours. A saturated aqueous sodium sulfite solution was added to the reaction solution, and the mixture was extracted with dichloromethane. The organic layer was washed with a saturated sodium hydrogen carbonate solution and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1 followed by ethyl acetate), to obtain the title compound (57 mg).


¹H-NMR (400 MHz, CDCl₃) δ 3.01 (3H, brs), 3.75-3.81 (4H, m), 4.06-4.22 (2H, m), 4.30-4.41 (2H, m), 4.86 (1H, d, J = 14.7 Hz), 5.04 (1H, d, J = 12.2 Hz), 5.12 (1H, d, J = 12.2 Hz), 5.29 (1H, d, J = 4.2 Hz), 6.48 (2H, d, J = 10.4 Hz), 7.28-7.37 (5H, m).

[0460]

<Reference Example 20-1>

[0461]

[Chemical Formula 79]



[0462]

((3S*,4R*)-1-{2-[(tert-Butyldimethylsilyl)oxy]ethyl}-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester

5 To a solution of

(-)-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester (346 mg) and

(2-bromoethoxy)(tert-butyl)dimethylsilane (2.0 mL) in

N,N-dimethylformamide (9.7 mL), potassium tert-butoxide (216 mg)

10 was added in several portions to produce a reaction solution. The reaction solution was stirred at room temperature for 15 hours. A saturated aqueous ammonium chloride solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively,
15 and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1 followed by ethyl acetate), to obtain the title compound as a colorless oil (204 mg).

20 ¹H-NMR (400 MHz, CDCl₃) δ 0.06 (6H, s), 0.88 (9H, s), 3.32-3.41 (1H, m), 3.48-3.63 (3H, m), 3.75- 3.81 (6H, m), 4.64 (1H, dd, J = 11.0, 8.0 Hz), 5.00-5.15 (3H, m), 6.58-6.73 (2H, m), 7.24-7.32 (6H, m).

[0463]




The same method as in Reference Example 20-1 was performed
25 using a corresponding 2-oxopyrrolidine substance to obtain the

following Reference Examples 20-2 to 20-4.

The structures and spectral data thereof are shown in Table 40.

[0464]

[Table 40]


Ref.No	Str.	Chemical name	P.D.
20-2		((3S*, 4R*)-1-{2-[(tert-butyl dimethylsilyl)oxy]ethyl}-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.00 (6H, s), 0.82 (9H, s), 3.24-3.33 (1H, m), 3.48-3.62 (3H, m), 3.66- 3.75 (6H, m), 4.64-4.70 (1H, m), 4.97 (2H, s), 5.16 (1H, brs), 6.35-6.45 (2H, m), 7.21-7.27 (5H, m).
20-3		(1-{2-[(tert-butyl dimethylsilyl)oxy]ethyl}-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid tert-butyl ester	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.08 (6H, s), 0.89 (9H, s), 1.36 (9H, s), 3.48 (1H, dt, J = 14.0, 4.9 Hz), 3.89 (1H, dt, J = 14.0, 4.9 Hz), 3.66 (1H, d, J = 10.4 Hz), 3.79 (3H, s), 3.84 (2H, t, J = 4.9 Hz), 3.92 (1H, dd, J = 10.4, 6.1 Hz), 4.57-4.61 (2H, m), 6.80 (2H, d, J = 8.6 Hz), 7.05 (2H, d, J = 8.6 Hz).
20-4		((3S*, 4R*)-1-{2-[(tert-butyl dimethylsilyl)oxy]ethyl}-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester	$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 0.06 (6H, s), 0.88 (9H, s), 3.10-3.20 (2H, m), 3.32-3.39 (1H, m), 3.47-3.61 (3H, m), 3.70-3.79 (3H, m), 4.58-4.64 (3H, m), 5.00-5.15 (3H, m), 6.51 (1H, d, J=11.0 Hz), 7.10-7.20 (1H, m), 7.26-7.40 (5H, m).

[0465]

<Reference Example 21-1>

[0466]

[Chemical Formula 80]



5 [0467]

[(3S*, 4R*)-4-(2-Fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester

To a solution of

10 ((3S*, 4R*)-1-{2-[(tert-butyl dimethylsilyl)oxy]ethyl}-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester (219 mg) in tetrahydrofuran (4.2 mL), tetrabutylammonium fluoride (465 μ L, 1.0 M tetrahydrofuran solution) was added to produce a reaction solution. The reaction solution was stirred at room temperature for 30 minutes. Water was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1, ethyl acetate, 15 and ethyl acetate : methanol = 10 : 1 in turn), to obtain the title compound as a white solid (150 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.73 (1H, brs), 3.40-3.64 (3H, m), 3.65-3.89




(7H, m), 4.53 (1H, t, J = 9.2 Hz), 5.00-5.12 (2H, brs), 5.25 (1H, brs), 6.57-6.73 (2H, m), 7.23-7.35 (6H, m).

[0468]

The same method as in Reference Example 21-1 was performed
5 using a corresponding [(tert-butyldimethylsilyl)oxy]ethyl
substance to obtain the following Reference Examples 21-2 to 21-4.
The structures and spectral data thereof are shown in Table 41.

[0469]

[Table 41]


Ref.No	Str.	Chemical name	P.D.
21-2		[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl] carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.70 (1H, brs), 3.40-3.48 (1H, m), 3.59-3.70 (3H, m), 3.78 (3H, s), 3.80-3.94 (3H, m), 4.50-4.61 (1H, m), 5.01-5.09 (2H, m), 5.29 (1H, brs), 6.41-6.52 (2H, m), 7.27-7.36 (5H, m).
21-3		[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl] carbamic acid tert-butyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.36 (9H, s), 2.71 (1H, brs), 3.50 (1H, dt, J = 14.0, 4.9 Hz), 3.62-3.68 (2H, m), 3.79-3.92 (7H, m), 4.57 (1H, t, J = 7.3 Hz), 4.64 (1H, d, J = 6.8 Hz), 6.83 (2H, d, J = 8.6 Hz), 7.07 (2H, d, J = 8.6 Hz).
21-4		[(3S*, 4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl] carbamic acid benzyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.10-3.18 (3H, m), 3.30-3.63 (6H, m), 4.42 (1H, t, J = 9.8 Hz), 4.56 (2H, t, J = 8.6 Hz), 4.74 (1H, t, J = 4.0 Hz), 4.99 (2H, s), 6.65 (1H, d, J = 11.0 Hz), 7.28-7.35 (6H, m), 7.62 (1H, d, J = 9.8 Hz).

[0470]

5 <Reference Example 22-1>

[0471]

[Chemical Formula 81]



[0472]

(3R*, 4S*)-1-Hydroxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester

5 Zinc powder (2.0 g) and an aqueous solution (5 mL) of ammonium chloride (0.9 g) was added to a solution of (+)-(S*)-2-[1-(4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester (1.04 g) in methanol (30 mL) to produce a reaction solution. The reaction solution was stirred for 1 hour. Ethyl acetate was added to the reaction solution, and the reaction solution was filtered over Celite. The filtrate was washed with a 1 mol/L hydrochloric acid, water, and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate), to obtain the title compound as a colorless oil (760 mg).


¹H-NMR (400 MHz, CDCl₃) δ 3.57 (1H, d, J = 8.5 Hz), 3.68 (1H, t, J = 8.1 Hz), 3.78 (3H, s), 3.79 (3H, s), 3.95 (1H, q, J = 8.1 Hz), 4.03 (1H, t, J = 8.5 Hz), 6.88 (2H, d, J = 8.8 Hz), 7.18 (2H, d, J = 8.8 Hz).

[0473]

<Reference Example 22-2>

[0474]

[Chemical Formula 82]



[0475]

(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-hydroxy-2-oxopyr-

5 rolidine-3-carboxylic acid methyl ester

The same method as in Reference Example 22-1 was performed using

(-)-(R)-2-[1-(2,6-difluoro-4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester in place of

10 (+)-(S*)-2-[1-(4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 3.72-3.82 (8H, m), 3.93 (1H, t, J = 8.8 Hz), 4.33 (1H, q, J = 8.8 Hz), 6.48 (2H, d, J = 8.6 Hz).

[0476]

15 <Reference Example 23-1>

[0477]

[Chemical Formula 83]



[0478]

(3R*, 4S*)-1-Methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-ca

20 rboxylic acid methyl ester

Potassium carbonate (276 mg) and iodomethane (0.12 mL) were

added to a solution of

(3R*,4S*)-1-hydroxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester (265 mg) in N,N-dimethylformamide (2 mL), to produce a reaction solution. The reaction solution was stirred at room temperature for 12 hours. Ethyl acetate was added to the reaction solution, and the mixture was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1), to obtain the title compound as a colorless oil (220 mg).


¹H-NMR (400 MHz, CDCl₃) δ 3.48 (1H, d, J = 9.1 Hz), 3.51 (1H, t, J = 8.5 Hz), 3.78 (3H, s), 3.80 (3H, s), 3.85 (3H, s), 3.90 (1H, dd, J = 8.5, 8.2 Hz), 3.98 (1H, t, J = 8.2 Hz), 6.89 (2H, d, J = 9.1 Hz), 7.16 (2H, d, J = 9.1 Hz).

[0479]

<Reference Example 23-2>

[0480]

[Chemical Formula 84]



[0481]

(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyrrolidine-3-carboxylic acid methyl ester

The same method as in Reference Example 23-1 was performed using

(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-hydroxy-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

5 (3R*,4S*)-1-hydroxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 3.62 (3H, s), 3.62-3.91 (10H, m), 6.43-6.50 (2H, m).

[0482]

10 <Reference Example 24-1>

[0483]

[Chemical Formula 85]



[0484]

(3R*,4S*)-1-Methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid

15 rboxylic acid

The same method as in Reference Example 8-1 was performed using

(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

20 (-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.

¹H-NMR (400 MHz, CDCl₃) δ 3.47 (1H, d, J = 9.1 Hz), 3.60 (1H, t, J = 8.5 Hz), 3.80 (3H, s), 3.88 (3H, s), 3.85-3.93 (1H, m), 3.99


(1H, t, J = 8.8 Hz), 6.90 (2H, d, J = 9.1 Hz), 7.23 (2H, d, J = 9.1 Hz).

[0485]

<Reference Example 24-2>

5 [0486]

[Chemical Formula 86]



[0487]

(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyr

10 rolidine-3-carboxylic acid

The same method as in Reference Example 8-1 was performed using

(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyrrolidine-3-carboxylic acid methyl ester in place of

15 (-)-(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 3.68-3.91 (9H, m), 4.18-4.26 (1H, m), 6.46-6.52 (2H, m).

[0488]

20 <Reference Example 25-1>

[0489]

[Chemical Formula 87]



[0490]

(3S*, 4R*)-4-[4-(Benzyloxy)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid methyl ester

Potassium carbonate (268 mg) and benzyl bromide (0.23 mL)

5 were added to a solution of

(3S*, 4R*)-4-(2,6-difluoro-4-hydroxyphenyl)-2-oxopyrrolidine-3-carboxylic acid methyl ester (500 mg) in N,N-dimethylformamide

(4 mL), to produce a reaction solution. The reaction solution was stirred at room temperature for 25 hours. A saturated aqueous

10 ammonium chloride solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic

layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under

reduced pressure, and the residue was then purified by silica gel

15 column chromatography (hexane : ethyl acetate = 4 : 1), to obtain the title compound as a colorless oil (643 mg).

¹H-NMR (400 MHz, CDCl₃) δ 3.50 (1H, t, J = 8.5 Hz), 3.66 (1H, t, J = 8.5 Hz), 3.76-3.80 (1H, m), 3.78 (3H, s), 4.46 (1H, q, J = 9.5 Hz), 5.02 (2H, s), 5.89 (1H, brs), 6.54 (2H, d, J = 10.3 Hz), 7.36-7.40

20 (5H, m).

[0491]

[0492]

Chemical structure of (S)-2-((2,6-difluorophenyl)methoxy)-2-oxo-1,3-dihydroisoindol-3-ylideneacetic acid.

5 (3S*,4R*)-4-[4-(Benzyloxy)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid


(3S*, 4R*)-4-[4-(benzyloxy)-2,6-difluorophenyl]-2-oxopyrrolidin

¹H-NMR (400 MHz, DMSO-d₆) δ 3.25 (1H, t, J = 9.4 Hz), 3.43 (1H, d, J = 10.3 Hz), 3.56 (1H, d, J = 9.1 Hz), 4.15 (1H, q, J = 9.4 Hz), 5.11 (2H, s), 6.85 (2H, d, J = 10.9 Hz), 7.32-7.44 (5H, m), 8.18 (1H, s), 12.76 (1H, brs).

15 [0494]

[0495]

246



[0496]

1-{(3S*, 4R*)-4-[4-(Benzyloxy)-2,6-difluorophenyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea

5 The same method as in Example 1-1 was performed using (3S*, 4R*)-4-[4-(benzyloxy)-2,6-difluorophenyl]-2-oxopyrrolidine-3-carboxylic acid in place of (-)-(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain the title compound.




10 ¹H-NMR (400 MHz, DMSO-d₆) δ 3.27-3.32 (1H, m), 3.44 (1H, t, J = 8.9 Hz), 3.79 (1H, q, J = 9.8 Hz), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 5.10 (2H, s), 6.46 (1H, d, J = 8.6 Hz), 6.82 (2H, d, J = 10.4Hz), 6.99-7.05 (2H, m), 7.30-7.44 (7H, m), 8.05 (1H, s), 8.66 (1H, s).

[0497]

15 The same method as in Reference Example 27-1 was performed using a corresponding carboxylic acid substance to obtain the following Reference Examples 27-2 to 27-4. The structures and spectral data thereof are shown in Table 42.

[0498]

[Table 42]


Ref.No	Str.	Chemical name	P.D.
27-2		1-[(3S*,4R*)-4-(4-benzyloxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	^1H NMR (400 MHz, CDCl_3) δ 3.56-3.61 (2H, m), 4.05-4.11 (1H, m), 4.67 (1H, t, J = 8.9 Hz), 4.99 (2H, s), 5.88 (1H, brs), 6.34 (1H, brs), 6.51 (2H, d, J = 10.4 Hz), 6.92 (1H, t, J = 7.6 Hz), 7.13 (1H, t, J = 7.6 Hz), 7.23 (2H, d, J = 7.6 Hz), 7.30-7.40 (5H, m), 7.46 (1H, brs).
27-3		(3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]phenyl)(methyl)carbamate tert-butyl ester	^1H -NMR (400 MHz, CDCl_3) δ 1.49 (9H, s), 3.25 (3H, s), 3.59-3.71 (2H, m), 4.18-4.26 (1H, m), 4.54-4.59 (1H, m), 5.75 (1H, brs), 6.03 (1H, brs), 6.80-6.85 (2H, m), 6.92 (2H, d, J = 10.4 Hz), 7.13-7.18 (2H, m), 7.29 (1H, brs).
27-4		(-)-1-[(3S*,4R*)-4-(1-benzyl-6-fluoroindolin-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	^1H -NMR (400 MHz, $\text{DMSO}-d_6$) δ 2.84 (2H, t, J = 8.5 Hz), 3.13 (1H, t, J = 9.7 Hz), 3.27-3.29 (2H, m), 3.38 (1H, t, J = 8.5 Hz), 3.56-3.64 (1H, m), 4.25 (2H, s), 4.48 (1H, dd, J = 11.5, 9.1 Hz), 6.36-6.40 (2H, m), 7.03 (2H, t, J = 9.1 Hz), 7.14 (1H, d, J = 7.9 Hz), 7.25-7.37 (7H, m), 7.88 (1H, s), 8.55 (1H, s). MS (ESI $^+$) m/z : 463 (MH^+). [α] $_D^{22}$ = -155 (c 0.10, EtOH)

[0499]

<Reference Example 28-1>

[0500]

[Chemical Formula 90]



5 [0501]

Trifluoromethanesulfonic acid 3,5-difluoro-4-((3^S,4^S)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl}phenyl ester

To a solution of

10 1-[(3^S,4^R)-4-(2,6-difluoro-4-hydroxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea (365 mg) in dichloromethane (5 mL), pyridine (0.4 mL) and trifluoromethanesulfonic anhydride (0.25 mL) were added under an argon atmosphere under ice-cooling to produce a reaction solution. The reaction solution was stirred
15 at room temperature for 3 hours. Water was added to the reaction solution, the mixture was extracted with ethyl acetate, and the extract was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column
20 chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title compound as a brown oil (388 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.36 (1H, t, J = 9.5 Hz), 3.52 (1H, t, J = 9.2 Hz), 3.93 (1H, q, J = 9.6 Hz), 4.55 (1H, dd, J = 10.4, 7.9 Hz), 6.51 (1H, d, J = 7.9 Hz), 6.98-7.04 (2H, m), 7.27-7.33 (2H, m), 7.58 (2H, d, J = 8.6 Hz), 8.13 (1H, s), 8.74 (1H, s).


5 MS (ESI⁺) m/z: 498 (MH⁺).

[0502]

<Reference Example 29-1>

[0503]

[Chemical Formula 91]



10 [0504]

[(3S*, 4R*, Z)-2-(Methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]carbamic acid benzyl ester

Triethyloxonium hexafluoro phosphate (446 mg) was added to a suspension of

15 (-)-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester (500 mg) in dichloromethane (3.7 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 20 hours. After that, a saturated aqueous sodium hydrogen carbonate solution was added, and the reaction solution
20 was stirred at room temperature for 30 minutes, and extracted with dichloromethane. The organic layer was dried over anhydrous sodium sulfate, and the solvent was removed under reduced pressure to obtain

[(3R*,4S*)-5-ethoxy-3-(4-methoxyphenyl)-3,4-dihydro-2H-pyrrol-4-yl]carbamic acid benzyl ester that was an intermediate as a yellow oil.

0-methylhydroxylamine hydrochloride (91 mg) was added to
5 a solution of

[(3R*,4S*)-5-ethoxy-3-(4-methoxyphenyl)-3,4-dihydro-2H-pyrrolo-4-yl]carbamic acid benzyl ester (325 mg) in methanol (2 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 3.5 hours. After that, a saturated
10 aqueous sodium hydrogen carbonate solution was added to the reaction solution under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica
15 gel column chromatography (ethyl acetate : hexane = 2 : 1), to obtain the title compound as a colorless oil (228 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.15 (1H, t, J = 9.1 Hz), 3.22-3.30 (1H, m), 3.45 (1H, t, J = 8.2 Hz), 3.59 (3H, s), 3.72 (3H, s), 4.60 (1H, t, J = 9.7 Hz), 4.96 (2H, s), 6.63 (1H, s), 6.86 (2H, d, J = 8.5
20 Hz), 7.20-7.34 (7H, m), 7.59 (1H, d, J = 9.1 Hz).

[0505]





The same method as in Reference Example 29-1 was performed using a corresponding Cbz substance and amine to obtain the following Reference Examples 29-2 to 29-7.

25 The structures and spectral data thereof are shown in Tables

43 to 44.



[0506]

[Table 43]

Ref.No	Str.	Chemical name	P.D.
29-2		[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.35 (1H, t, J = 9.2 Hz), 3.48-3.69 (2H, m), 3.70-3.81 (6H, m), 4.92-5.11 (5H, m), 6.51-6.72 (2H, m), 7.24-7.34 (6H, m).
29-3		[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.52-3.58 (2H, m), 3.60-3.81 (7H, m), 4.94-5.18 (5H, m), 6.40-6.48 (2H, m), 7.26-7.31 (5H, m).
29-4		{(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl}carbamic acid benzyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.16 (1H, t, J = 9.1 Hz), 3.24-3.30 (1H, m), 3.47 (1H, t, J = 8.2 Hz), 3.53-3.59 (2H, m), 3.72 (3H, s), 3.80 (2H, t, J = 5.1 Hz), 4.44 (1H, t, J = 5.8 Hz), 4.60 (1H, t, J = 9.7 Hz), 4.96 (2H, s), 6.63 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 7.20-7.34 (7H, m), 7.61 (1H, d, J = 9.1 Hz).
29-5		{(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.87 (1H, brs), 3.50-3.64 (2H, m), 3.65-3.80 (4H, m), 3.83-3.90 (2H, m), 4.08 (1H, dd, J = 4.3, 4.3 Hz), 4.80-5.15 (3H, m), 5.13 (1H, dd, J = 9.8, 8.0 Hz), 5.25 (1H, s), 6.40-6.50 (2H, m), 7.26-7.32 (5H, m).

5 [0507]

[Table 44]

Ref.No	Str.	Chemical name	P.D.
29-6		[(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-(methylamino)-3,4-dihydro-2H-pyrrolo-4-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.66 (3H, d, J = 4.9 Hz), 3.34-3.43 (1H, m), 3.45-3.56 (1H, m), 3.63-3.73 (1H, m), 3.75 (3H, s), 4.86 (1H, t, J = 9.5 Hz), 4.92 (1H, d, J = 12.8 Hz), 4.99 (1H, d, J = 12.8 Hz), 5.99-6.06 (1H, m), 6.70 (2H, d, J = 10.4 Hz), 7.22-7.36 (5H, m), 7.69 (1H, d, J = 9.2 Hz).
29-7		((3S*,4R*,Z)-2-[[[(tert-butyldimethylsilyl)oxy]imino}-4-(4-methoxyphenyl)pyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.05 (6H, s), 0.90 (9H, s), 3.18 (1H, t, J = 9.1 Hz), 3.22-3.40 (1H, m), 3.49 (1H, t, J = 8.2 Hz), 3.72 (3H, s), 4.55 (1H, t, J = 9.7 Hz), 4.91 (1H, d, J = 13.3 Hz), 5.03 (1H, d, J = 13.3 Hz), 6.49 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 7.20-7.33 (7H, m), 7.58 (1H, d, J = 9.1 Hz).





[0508]

The same method as in Reference Example 29-1 was performed using a corresponding Cbz substance and amine to obtain the following Reference Examples 29-8 to 29-13.

The structures and spectral data thereof are shown in Tables 45 to 46.



[0509]

[Table 45]

Ref.No	Str.	Chemical name	Spectrum data
29-8		((3S*,4R*,Z)-2-([(tert-butyl dimethylsilyl)oxy]imino)-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-3-yl)carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 0.15 (6H, s), 0.94 (9H, s), 3.50-3.62 (2H, m), 3.65-3.80 (4H, m), 5.01 (2H, brs), 5.06-5.13 (1H, m), 5.31 (1H, s), 6.45 (2H, d, J=10.4 Hz), 7.25-7.35 (5H, m).
29-9		[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(morpholinoimino)pyrrolidin-3-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 2.66-2.72 (4H, m), 3.52-3.82 (10H, m), 4.90-5.07 (3H, m), 5.27 (1H, brs), 5.70 (1H, brs), 6.40-6.50 (2H, m), 7.25-7.35 (5H, m).
29-10		{(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-[(1-methyl-1H-pyrazol-3-yl)amino]-3,4-dihydro-2H-pyrrol-4-yl}carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.50-3.80 (6H, m), 3.82 (3H, s), 4.95-5.15 (3H, m), 6.06 (1H, s), 6.42-6.52 (2H, m), 7.21 (1H, s), 7.25-7.40 (5H, m), 8.50 (1H, brs).
29-11		[(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-(phenylamino)-3,4-dihydro-2H-pyrrol-4-yl]carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.50-3.90 (5H, m), 4.18-4.32 (1H, m), 4.90-5.20 (3H, m), 6.40-6.50 (2H, m), 7.10-7.40 (10H, m), 7.55 (1H, brs).

[0510]

[Table 46]


Ref.No	Str.	Chemical name	Spectrum data
29-12		{(3S*,4R*,Z)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}carbamic acid benzyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 3.00 (1H, brs), 3.10-3.20 (2H, m), 3.32-3.36 (1H, m), 3.50-3.65 (2H, m), 3.86 (2H, s), 4.07-4.09 (1H, m), 4.93-5.11 (4H, m), 5.23 (1H, s), 6.50 (1H, d, J=10.4 Hz), 7.15 (1H, d, J = 6.12 Hz), 7.26-7.40 (5H, m)
29-13		3-((Z)-{(3S,4R)-3-[(benzyloxycarbonyl)amino]-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-2-ylidene}amino)propanoic acid ethyl ester	¹ H-NMR (CDCl ₃) δ: 1.28 (3H, t, J = 7.3 Hz), 2.64-2.78 (1H, m), 3.50-4.20 (9H, m), 4.02-4.09 (1H, m), 4.65-4.75 (1H, m), 4.95-5.20 (3H, m), 5.75 (1H, brs), 6.42-6.52 (2H, m), 7.27-7.37 (5H, m).

[0511]

5 <Reference Example 30-1>

[0512]

[Chemical Formula 92]



[0513]

[(3R*,4S*,Z)-2-(Methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3

10 -yl]carbamic acid benzyl ester

The same method as in Reference Example 29-1 was performed

using

(+)-[(3R*,4S*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester in place of

(-)-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title compound.


¹H-NMR (400 MHz, DMSO-d₆) δ 3.15 (1H, t, J = 9.1 Hz), 3.22-3.30 (1H, m), 3.45 (1H, t, J = 8.2 Hz), 3.59 (3H, s), 3.72 (3H, s), 4.60 (1H, t, J = 9.7 Hz), 4.96 (2H, s), 6.63 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 7.20-7.34 (7H, m), 7.59 (1H, d, J = 9.1 Hz).

[0514]

<Reference Example 31-1>

[0515]

[Chemical Formula 93]



[0516]

1-((3S*,4R*,Z)-2-[[tert-butyl(dimethyl)silyl]oxy]imino)-4-(4-methoxyphenyl)pyrrolidin-3-yl)-3-(4-fluorophenyl)urea

The same method as in Example 32-1 was followed using ((3S*,4R*,Z)-2-[[tert-butyl(dimethyl)silyl]oxy]imino)-4-(4-methoxyphenyl)pyrrolidin-3-yl)carbamic acid benzyl ester in place of

[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title compound.


¹H-NMR (400 MHz, DMSO-d₆) δ 0.04 (3H, s), 0.05 (3H, s), 0.89 (9H, s), 3.17 (1H, t, J = 8.8 Hz), 3.28-3.37 (1H, m), 3.53 (1H, t, J = 8.5 Hz), 3.71 (3H, s), 4.74 (1H, t, J = 9.1 Hz), 6.34 (1H, d, J = 9.1 Hz), 6.56 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 7.00-7.04 (2H, m), 7.27 (2H, d, J = 8.5 Hz), 7.30-7.34 (2H, m), 8.45 (1H, s).
 MS (FD⁺) m/z: 472 (M⁺).

[0517]

<Reference Example 31-2>

[0518]

10 [Chemical Formula 94]



[0519]

1-((3S^{*}, 4R^{*}, Z)-2-{[(tert-Butyldimethylsilyl)oxy]imino}-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-3-yl)-3-(4-fluorophenyl)
 15 urea

The same method as in Example 32-1 was followed using ((3S^{*}, 4R^{*}, Z)-2-{[(tert-butyl(dimethyl)silyl)oxy]imino}-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-3-yl)carbamic acid benzyl ester in place of

20 [(3S^{*}, 4R^{*}, Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title compound.

¹H-NMR (400 MHz, CDCl₃) δ 0.12 (6H, s), 0.93 (9H, s), 3.57-3.64 (2H,


m), 3.73-3.82 (1H, m), 3.76 (3H, s), 5.12-5.21 (1H, m), 5.38 (1H, brs), 5.47 (1H, s), 6.46 (2H, d, J = 10.8 Hz), 6.86-6.93 (2H, m), 7.17-7.22 (2H, m), 7.39 (1H, brs).

[0520]

5 <Reference Example 32-1>

[0521]

[Chemical Formula 95]



[0522]

10 3-(2,6-Difluoro-4-methoxyphenyl)-2-[(diphenylmethylene)amino]
-4-nitrobutanoic acid ethyl ester

Lithium diisopropylamide (1.1 mL, 1.05 M in tetrahydrofuran/hexane) was added to a solution of 2-[(diphenylmethylene)amino]acetic acid ethyl ester (300 mg) in
15 tetrahydrofuran (2.3 mL) at -78°C to produce a reaction solution. The reaction solution was stirred for 1 hour, a solution of (E)-1,3-difluoro-5-methoxy-2-(2-nitrovinyl)benzene (241 mg) in tetrahydrofuran was added, and the reaction solution was stirred at the same temperature for 15 minutes. The temperature of the
20 reaction solution was gradually heated up to room temperature, and the reaction solution was stirred for 30 minutes. Water was added to the reaction solution, and the mixture was extracted with

ethyl acetate. The organic layer was washed with a brine, and dried over sodium sulfate. The solvent was removed, and the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 2 : 1), to obtain the title compound as a yellow oil (543 mg).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.12 (3H, t, $J = 7.3$ Hz), 3.73 (3H, s), 4.00-4.06 (2H, m), 4.43 (1H, d, $J = 7.8$ Hz), 4.73-4.95 (3H, m), 6.36 (2H, d, $J = 10.4$ Hz), 7.00-7.05 (2H, m), 7.31-7.38 (6H, m), 7.81-7.85 (2H, m).

[0523]

<Reference Example 32-2>

[0524]

[Chemical Formula 96]



[0525]

2-[(Diphenylmethylene)amino]-3-(4-methoxyphenyl)-4-nitrobutanoic acid ethyl ester

The same method as in Reference Example 32-1 was performed using (E)-1-methoxy-4-(2-nitrovinyl)benzene in place of (E)-1,3-difluoro-5-methoxy-2-(2-nitrovinyl)benzene to obtain the title compound.

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.19 (3H, t, $J = 6.7$ Hz), 3.76 (3H, s),


3.90-4.36 (6H, m), 7.05 (2H, d, J = 8.6 Hz), 7.24-7.47 (10H, m),
7.63 (2H, d, J = 8.6 Hz).

[0526]

<Reference Example 33-1>

5 [0527]

[Chemical Formula 97]



[0528]

2-[(tert-Butoxycarbonyl)amino]-3-(2,6-difluoro-4-methoxyphenyl)-
10 4-nitrobutanoic acid ethyl ester

Acetic acid (4.2 mL) and water (0.8 mL) were added to
3-(2,6-difluoro-4-methoxyphenyl)-2-[(diphenylmethylene)amino]-
4-nitrobutyric acid ethyl ester (530 mg) at room temperature to
produce a first reaction solution. The first reaction solution
15 was stirred at 60°C for 7 hours. The reaction solution was
concentrated under reduced pressure, a saturated aqueous sodium
hydrogen carbonate solution was added to the residue until the
pH was 8, to produce a second reaction solution. After that,
tetrahydrofuran (5.0 mL) was added to the second reaction solution,
20 di-tert-butyl dicarbonate (290 mg) was then added, and the mixture
was stirred at room temperature for 16 hours. The second reaction
solution was extracted with ethyl acetate, and the organic layer
was washed with a brine, and dried over anhydrous sodium sulfate.

The solvent was removed, and the residue was purified by silica gel column chromatography (toluene : ethyl acetate = 2 : 1), to obtain the title compound as a white solid (315 mg).


¹H-NMR (400 MHz, CDCl₃) δ 1.08 (3H, brt), 1.46 (9H, s), 3.78 (3H, s), 4.00-4.10 (2H, m), 4.22-4.33 (1H, m), 4.71 (1H, brt), 4.88 (2H, d, J = 7.9 Hz), 5.22 (1H, brd), 6.45 (2H, d, J = 10.4 Hz).

[0529]

<Reference Example 33-2>

[0530]

10 [Chemical Formula 98]



[0531]

2-[(tert-Butoxycarbonyl)amino]-3-(4-methoxyphenyl)-4-nitrobutanoic acid ethyl ester

15 The same method as in Reference Example 33-1 was performed using

2-[(Diphenylmethylene)amino]-3-(4-methoxyphenyl)-4-nitrobutanoic acid ethyl ester in place of

3-(2,6-difluoro-4-methoxyphenyl)-2-[(diphenylmethylene)amino]-4-nitrobutanoic acid ethyl ester to obtain the title compound.

20 ¹H-NMR (400 MHz, CDCl₃) δ 1.08 (3H, brt), 1.46 (9H, s), 3.78 (3H, s), 3.82 (1H, q, J = 8.6 Hz), 4.02 (2H, d, J = 6.7 Hz), 4.55 (1H, t, J = 6.7 Hz), 4.77 (1H, dd, J = 13.4, 9.2 Hz), 4.85 (1H, dd, J


= 13.4, 5.5 Hz), 5.12 (1H, brs), 6.85 (2H, d, J = 8.6 Hz), 7.09 (2H, d, J = 8.6 Hz).

[0532]

<Reference Example 34-1>

5 [0533]

[Chemical Formula 99]



(-)-[4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester

10 (+)-[4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester

2-[(tert-Butoxycarbonyl)amino]-3-(2,6-difluoro-4-methoxyphenyl)-4-nitrobutanoic acid ethyl ester was used in place of (R*)-2-[1-(4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester, and

(±)-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester was obtained in the same method as in Reference Example 7-1. The obtained compound was subjected to optical resolution by high performance liquid chromatography (hexane : methyl tert-butyl ether : ethanol = 45 : 25 : 30, flow rate: 15.0 mL) using a column for separation of enantiomers (CHIRALPAK ID). An isomer A(-) with a retention time of 13.4

minutes and an isomer B(+) with a retention time of 22.5 minutes that were the title compounds were each obtained as a white solid by the optical resolution.

Isomer A(-):

5 $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.26 (9H, s), 3.40 (1H, d, $J = 10.4$ Hz), 3.78 (3H, s), 3.85 (1H, d, $J = 10.4, 8.6$ Hz), 4.46 (1H, t, $J = 6.6$ Hz), 4.60 (1H, dd, $J = 9.8, 6.1$ Hz), 4.94 (1H, d, $J = 5.5$ Hz), 5.75 (1H, s) 6.44 (2H, d, $J = 10.0$ Hz).

$[\alpha]_{\text{D}}^{25} = -131$ (c 0.10, EtOH)

10 Isomer B(+):

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.26 (9H, s), 3.40 (1H, d, $J = 10.4$ Hz), 3.78 (3H, s), 3.85 (1H, d, $J = 10.4, 8.6$ Hz), 4.46 (1H, t, $J = 6.6$ Hz), 4.60 (1H, dd, $J = 9.8, 6.1$ Hz), 4.94 (1H, d, $J = 5.5$ Hz), 5.90 (1H, s) 6.44 (2H, d, $J = 10.0$ Hz).


15 $[\alpha]_{\text{D}}^{25} = +144$ (c 0.10, EtOH)

[0534]


<Reference Example 34-2>

[0535]

[Chemical Formula 100]



(-)-cis



(+)-cis

[0536]

(-)-[4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid

tert-butyl ester

(+)-[4-(4-Methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid

tert-butyl ester

2-[(tert-Butoxycarbonyl)amino]-3-(4-methoxyphenyl)-4-ni

5 trobutanoic acid ethyl ester was used in place of

(R^{*})-2-[1-(4-methoxyphenyl)-2-nitroethyl]malonic acid dimethyl ester, and

(±)-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid

tert-butyl ester was obtained in the same method as in Reference

10 Example 7-1. The obtained compound was subjected to optical resolution by high performance liquid chromatography (hexane : methyl tert-butyl ether : ethanol = 45 : 25 : 30, flow rate: 15.0 mL) using a column for separation of enantiomers (CHIRALPAK ID). An isomer A(-) with a retention time of 7.5 minutes and an isomer
15 B(+) with a retention time of 10.2 minutes that were the title compounds were each obtained as a white solid by the optical resolution.

Isomer A(-):

¹H-NMR (400 MHz, CDCl₃) δ 1.37 (9H, s), 3.56 (1H, d, J = 9.8 Hz),
20 3.80 (3H, s), 3.84 (1H, dd, J = 10.4, 6.1 Hz), 3.92 (1H, brs), 4.56 (2H, s), 5.75 (1H, s), 6.83 (2H, d, J = 8.6 Hz), 7.10 (2H, d, J = 8.6 Hz).

[α]_D²² = -88 (c 0.10, EtOH)

Isomer B(+):

25 ¹H-NMR (400 MHz, CDCl₃) δ 1.37 (9H, s), 3.56 (1H, d, J = 9.8 Hz),

3.80 (3H, s), 3.84 (1H, dd, J = 10.4, 6.1 Hz), 3.92 (1H, brs), 4.56 (2H, s), 5.75 (1H, s), 6.83 (2H, d, J = 8.6 Hz), 7.10 (2H, d, J = 8.6 Hz).


$[\alpha]_D^{22} = +98$ (c 0.10, EtOH)

5 [0537]

<Reference Example 35-1>

[0538]

[Chemical Formula 101]



10 [0539]

1-[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-{4-[(trimethylsilyl)ethynyl]phenyl}urea

Trimethylsilylacetylene (115 μ L) and triethylamine (115 μ L)
15 were added to a solution of
1-[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-iodophenyl)urea (80 mg),
bis(triphenylphosphine)palladium(II) dichloride (11.5 mg), and
copper(I) iodide (15.6 mg) in N,N-dimethylformamide (2 mL) to
20 produce a reaction solution. The reaction solution was stirred
at 50°C for 9 hours. The reaction solution was concentrated under
reduced pressure, and the residue was then purified by silica gel

column chromatography (hexane : ethyl acetate = 1 : 1, ethyl acetate, and ethyl acetate : methanol = 4 : 1 in turn), to obtain the title compound as a colorless solid (67 mg).


¹H-NMR (400 MHz, CDCl₃) δ 0.22 (9H, s), 3.52-3.63 (2H, m), 3.75 (3H, s), 4.09 (1H, q, J = 9.2 Hz), 4.65 (1H, dd, J = 10.4, 7.4 Hz), 6.08 (1H, brs), 6.36-6.45 (3H, m), 7.16 (2H, d, J = 8.6 Hz), 7.21 (2H, d, J = 8.6 Hz), 7.81 (1H, brs).

[0540]

<Reference Example 36-1>

10 [0541]

[Chemical Formula 102]



[0542]

2-((3S*, 4R*)-3-{[(Benzyloxy)carbonyl]amino}-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-1-yl)acetic acid

A 2 mol/L sodium hydroxide aqueous solution (1.26 mL) was added to a solution of

2-((3S*, 4R*)-3-{[(benzyloxy)carbonyl]amino}-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-1-yl)acetic acid ethyl ester

20 (521 mg) in methanol (25 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 5 hours. A 1 mol/L Hydrochloric acid was added to the reaction solution

to make the reaction solution acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed to obtain the title compound as a white solid (465 mg).


¹H-NMR (400 MHz, DMSO-d₆) δ 3.50-3.58 (2H, m), 3.72-3.79 (4H, m), 3.87 (1H, d, J = 17.7 Hz), 4.11 (1H, d, J = 17.7 Hz), 4.48 (1H, t, J = 10.0 Hz), 4.94 (1H, d, J = 12.2 Hz), 4.99 (1H, d, J = 12.2 Hz), 6.78 (2H, d, J = 10.4 Hz), 7.25-7.35 (5H, m), 7.74 (1H, d, J = 9.2 Hz), 12.9 (1H, s).

[0543]

<Reference Example 37-1>

[0544]

[Chemical Formula 103]



[0545]

((3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-{2-[(2-hydroxypropyl)amino]-2-oxoethyl}-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester

1-Hydroxybenzotriazole (35 mg), 1-amino-2-propanol (20 μL), and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (57 mg) were added to a solution of

2-((3S*,4R*)-3-[[[(benzyloxy)carbonyl]amino]-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid (100 mg) in dichloromethane (1.0 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 4 hours.

5 Water was added to the reaction solution, and the mixture was extracted with dichloromethane. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate
10 followed by ethyl acetate : methanol = 4 : 1), to obtain the title compound (96 mg).


¹H-NMR (400 MHz, DMSO-d₆) δ 1.00 (3H, d, J = 6.1 Hz), 2.98-3.05 (2H, m), 3.54 (2H, t, J = 9.2 Hz), 3.60-3.66 (1H, m), 3.77-3.81 (4H, m), 4.04 (1H, d, J = 16.5 Hz), 4.47 (1H, t, J = 9.8 Hz), 4.68 (1H, q, J = 2.0 Hz), 4.97 (2H, dd, J = 17.4, 12.5 Hz), 6.76 (2H, d, J = 11.0 Hz), 7.26-7.33 (5H, m), 7.73 (1H, d, J = 9.2 Hz), 7.94 (1H, t, J = 5.5 Hz).

[0546]

<Reference Example 37-2>

20 [0547]

[Chemical Formula 104]



((3S*,4R*)-1-[2-(2-[(Acetylhydrazinyl)-2-oxoethyl]-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester


5 The same method as in Reference Example 37-1 was performed using acetylhydrazine in place of 1-amino-2-propanol as amine to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 1.84 (3H, s), 3.55 (2H, d, J = 10.4 Hz), 3.76 (4H, s), 3.85 (1H, d, J = 16.5 Hz), 4.12 (1H, d, J = 16.5 Hz), 4.50 (1H, t, J = 9.2 Hz), 4.97 (2H, dd, J = 16.5, 12.2 Hz), 6.75 (2H, d, J = 11.0 Hz), 7.25-7.33 (5H, m), 7.70 (1H, d, J = 9.2 Hz), 7.94 (2H, s).

<Reference Example 38-1>

15 [0550]

[Chemical Formula 105]



((3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxo-1-{2-oxo-2-[(2-oxopropyl)amino]ethyl}pyrrolidin-3-yl)carbamic acid benzyl ester

2-Iodoxybenzoic acid (131 mg) was added to a solution of

((3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-{2-[(2-hydroxypropyl)amino]-2-oxoethyl}-2-oxopyrrolidin-3-yl)carbamic acid benzyl ester (96 mg) in dimethyl sulfoxide (2.0 mL), to produce a reaction solution. The reaction solution was stirred at room temperature for 18 hours. A saturated aqueous sodium hydrogen carbonate solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed to obtain the title compound

(84 mg).


¹H-NMR (400 MHz, DMSO-d₆) δ 2.02 (3H, s), 3.46-3.54 (2H, m), 3.69 (1H, d, J = 9.8 Hz), 3.73 (3H, s), 3.82 (1H, d, J = 16.5 Hz), 3.91 (2H, d, J = 5.5 Hz), 4.07 (1H, d, J = 16.5 Hz), 4.45 (1H, t, J = 11.0 Hz), 4.92 (2H, dd, J = 16.5, 12.2 Hz), 6.72 (2H, d, J = 10.4 Hz), 7.22-7.31 (5H, m), 7.67 (1H, d, J = 9.2 Hz), 8.23 (1H, t, J = 5.5 Hz).

[0552]

<Reference Example 39-1>

[0553]

[Chemical Formula 106]



[0554]

{(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-[(5-methyloxazol-2-yl)methyl]-2-oxopyrrolidin-3-yl}carbamic acid benzyl ester

Triphenylphosphine (325 mg) and triethylamine (860 μ L) were added to a solution of

5 ((3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-{2-oxo-2-[(2-oxopropyl)amino]ethyl}pyrrolidin-3-yl)carbamic acid benzyl ester (76 mg) in carbon tetrachloride (7.8 mL) to produce a reaction solution. The reaction solution was stirred at 80°C for 30 minutes, dichloromethane (2 mL) was added, and the reaction solution was
10 stirred at 80°C for 4 hours. A saturated aqueous sodium hydrogen carbonate solution was added to the reaction solution, and the mixture was extracted with dichloromethane. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced
15 pressure, and the residue was then purified by silica gel column chromatography (chloroform : methanol = 9 : 1), to obtain the title compound (15 mg).


¹H-NMR (400 MHz, CDCl₃) δ 2.30 (3H, s), 3.58 (2H, brs), 3.77 (4H, s), 4.64 (2H, s), 4.75 (1H, dd, J = 10.4, 7.3 Hz), 5.02 (2H, d, J = 12.2 Hz), 5.22 (1H, brs), 6.45 (2H, d, J = 9.8 Hz), 6.67 (1H, s), 7.27-7.35 (5H, m).

[0555]

<Reference Example 39-2>

[0556]

25 [Chemical Formula 107]



[0557]

{(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl}carbamic acid
5 benzyl ester

Triphenylphosphine (151 mg), triethylamine (160 μ L), and 1,1,1,2,2,2-hexachloroethane (136 mg) were added to a solution of {(3S*, 4R*)-1-[2-(2-acetyl
hydrazinyl)-2-oxoethyl]-4-(2,6-difluoro-4-methoxyphenyl)-2-ox
10 opyrrolidin-3-yl}carbamic acid benzyl ester (113 mg) in dichloromethane (11.5 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. A saturated aqueous sodium hydrogen carbonate solution was added to the reaction solution, and the mixture was extracted with
15 dichloromethane. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title
20 compound (69 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.54 (3H, s), 3.61 (2H, d, J = 8.6 Hz), 3.77 (3H, s), 3.82-3.92 (1H, m), 4.70 (1H, t, J = 9.8 Hz), 4.76


(2H, s), 5.01-5.10 (2H, m), 5.29 (1H, brs), 6.45 (2H, d, J = 9.8 Hz), 7.26-7.35 (5H, m).

[0558]

<Reference Example 40-1>

5 [0559]

[Chemical Formula 108]



[0560]

4-(Difluoromethoxy)-2,6-difluorobenzaldehyde

10 A solution (reaction solution) of 2,6-difluoro-4-hydroxybenzaldehyde (1.41 g), potassium carbonate (1.48 g), and sodium chlorodifluoroacetate (2.73 g) in a mixture of N,N-dimethylformamide (15.8 mL) and water (1.84 mL) was stirred at 100°C for 1 hour and a half.

15 The reaction solution was ice-cooled, water (4.1 mL) and concentrated hydrochloric acid (2.8 mL) were added, and the reaction solution was stirred at room temperature overnight. A 2 mol/L sodium hydroxide aqueous solution was added to the reaction solution under ice-cooling to adjust the pH to 10, and the mixture
20 was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl

acetate = 95 : 5 followed by 3 : 1), to obtain the title compound as a light brown oil (1.61 g).


¹H-NMR (400 MHz, CDCl₃) δ 6.63 (1H, t, J = 71.5 Hz), 6.78 (2H, d, J = 9.1 Hz), 10.26 (1H, s).

5 [0561]

<Reference Example 41-1>

[0562]

[Chemical Formula 109]



10 [0563]

3,5-Difluoro-4-formylphenyl trifluoromethanesulfonate

Pyridine (0.8 mL) and trifluoromethanesulfonic anhydride (1.18 mL) were added to a solution of

2,6-difluoro-4-hydroxybenzaldehyde (1.03 g) in dichloromethane

15 (30 mL) under an argon atmosphere under ice-cooling to produce

a reaction solution. The reaction solution was stirred at room temperature for 3 hours. Water was added to the reaction solution,

the mixture was extracted with ethyl acetate. The extract was washed with water and a brine, and dried over anhydrous sodium sulfate.

20 The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), to obtain the title compound as a colorless oil (1.5 g).


¹H-NMR (400 MHz, CDCl₃) δ 6.99-7.04 (2H, m), 10.32 (1H, s).

[0564]

<Reference Example 42-1>

[0565]

5 [Chemical Formula 110]



[0566]

N-(3,5-Difluoro-4-formylphenyl)acetamide

Acetamide (225 mg),

10 tris(dibenzylideneacetone)dipalladium(0) (116 mg), xantphos (145 mg), and cesium carbonate (1.24 g) were added to a solution of 3,5-difluoro-4-formylphenyl trifluoromethanesulfonate (737 mg) in toluene (15 mL) to produce a reaction solution. The reaction solution was stirred at 100°C under an argon atmosphere for 5 hours.

15 The reaction solution was filtered over Celite and washed with ethyl acetate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title compound as a pale yellow solid (255

20 mg).


¹H-NMR (400 MHz, CDCl₃) δ 2.22 (3H, s), 7.24 (2H, d, J = 10.4 Hz), 7.44 (1H, brs), 10.23 (1H, s).

[0567]

<Reference Example 42-2>

[0568]

[Chemical Formula 111]



5 [0569]

(3,5-Difluoro-4-formylphenyl)carbamic acid tert-butyl ester

The same method as in Reference Example 42-1 was performed using tert-butyl carbamate in place of acetamide as a reagent to obtain the title compound.


¹H-NMR (400 MHz, CDCl₃) δ 1.53 (9H, s), 6.77 (1H, brs), 7.07 (2H, d, J = 11.0 Hz), 10.20 (1H, s).

[0570]

<Reference Example 43-1>

15 [0571]

[Chemical Formula 112]



[0572]

(3,5-Difluoro-4-formylphenyl)(methyl)carbamic acid tert-butyl ester

Iodomethane (110 μL) and sodium hydride (51 mg, 60% in oil) were added to a solution of (3,5-difluoro-4-formylphenyl)carbamic

acid tert-butyl ester (220 mg) in N,N-dimethylformamide (4 mL) under cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. A saturated aqueous ammonium chloride solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1), to obtain the title compound as a colorless oil (150 mg).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.53 (9H, s), 3.32 (3H, s), 7.05 (2H, d, $J = 11.0$ Hz), 10.25 (1H, s).

[0573]

<Reference Example 44-1>

[0574]

[Chemical Formula 113]



[0575]

5-Bromo-6-fluoro-2,3-dihydro-1H-inden-1-ol

Sodium borohydride (56 mg) was added to a solution of 5-bromo-6-fluoro-2,3-dihydro-1H-inden-1-one (310mg) in a mixture of tetrahydrofuran (7 mL) and ethanol (7 mL) under an argon atmosphere under ice-cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 1 hour. Water

was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), to obtain the title compound as a colorless solid (290 mg).


¹H-NMR (400 MHz, CDCl₃) δ 1.78 (1H, d, J = 7.3 Hz), 1.92-2.01 (1H, m), 2.50-2.58 (1H, m), 2.74-2.82 (1H, m), 2.96-3.03 (1H, m), 5.19 (1H, q, J = 6.7 Hz), 7.15 (1H, d, J = 7.9 Hz), 7.41 (1H, d, J = 6.7 Hz).

[0576]

<Reference Example 45-1>

[0577]

[Chemical Formula 114]



[0578]

[(5-Bromo-6-fluoro-2,3-dihydro-1H-inden-1-yl)oxy](tert-butyl)dimethylsilane

Triethylamine (0.34 mL), N,N-dimethyl-4-aminopyridine (7.4 mg), and tert-butyl dimethyl chlorosilane (274 mg) were added to a solution of 5-bromo-6-fluoro-2,3-dihydro-1H-inden-1-ol (280 mg) in N,N-dimethylformamide (2.4 mL) under an argon atmosphere, to produce a reaction solution. The reaction solution was stirred

at room temperature for 3 hours. Water was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1), to obtain the title compound as a colorless solid (410 mg).


¹H-NMR (400 MHz, CDCl₃) δ 0.14 (3H, s), 0.17 (3H, s), 0.94 (9H, s), 1.89-1.98 (1H, m), 2.40-2.48 (1H, m), 2.68-2.76 (1H, m), 2.88-2.94 (1H, m), 5.16 (1H, t, J = 7.0 Hz), 7.01 (1H, d, J = 7.9 Hz), 7.36 (1H, d, J = 6.1 Hz).

[0579]

<Reference Example 46-1>

[0580]

[Chemical Formula 115]



[0581]

1-[(tert-Butyldimethylsilyl)oxy]-6-fluoro-2,3-dihydro-1H-indene-5-carbaldehyde

n-Butyllithium (4.15 mL, 2.65 mol/L hexane solution) was slowly added to a solution of

[(5-bromo-6-fluoro-2,3-dihydro-1H-inden-1-yl)oxy](tert-butyl)dimethyl silane (3.45 g) in tetrahydrofuran (50 mL) at -78°C under an argon atmosphere to produce a reaction solution. The reaction

solution was stirred for 40 minutes. After that,
N,N-dimethylformamide (1.26 mL) was added to the reaction solution,
and the reaction solution was stirred at -78°C for 1 hour. A saturated
aqueous sodium chloride solution was added to the reaction solution,
5 and the mixture was stirred at 0°C for 1 hour and extracted with
ethyl acetate. The organic layer was washed with water and a brine,
and dried over anhydrous sodium sulfate. The solvent was removed
under reduced pressure, and the residue was then purified by silica
gel column chromatography (hexane : ethyl acetate = 4 : 1), to
10 obtain the title compound as a yellow oil (2.63 g).


¹H-NMR (400 MHz, CDCl₃) δ 0.16 (3H, s), 0.19 (3H, s), 0.95 (9H, s),
1.91-2.01 (1H, m), 2.44-2.52 (1H, m), 2.71-2.79 (1H, m), 2.96 (1H,
dd, J = 14.7, 7.9 Hz), 5.22 (1H, t, J = 7.6 Hz), 7.05 (1H, d, J
= 10.4 Hz), 7.67 (1H, d, J = 6.7 Hz), 10.32 (1H, s).

15 [0582]

<Reference Example 47-1>

[0583]

[Chemical Formula 116]



20 [0584]

2-(2-Bromo-3,5-difluorophenoxy)ethanol

Ethylene glycol (2.65 mL) and N-methyl-2-pyrrolidone (2 mL)
were added to 2-bromo-1,3,5-trifluorobenzene (2.0 g) under an argon

atmosphere, and potassium tert-butoxide (1.28 g) was added under ice-cooling to produce a reaction solution. The reaction solution was stirred under heating at 90°C for 2 hours. The reaction solution was cooled to room temperature, water was added, and the mixture
5 was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1 followed by 1 : 1), to obtain the title compound
10 as a colorless oil (1.43 g).


¹H-NMR (400 MHz, CDCl₃) δ 2.10 (1H, t, J = 6.4 Hz), 4.00-4.04 (2H, m), 4.13 (2H, t, J = 4.6 Hz), 6.48-6.52 (1H, m), 6.55-6.61 (1H, m).

[0585]

15 <Reference Example 48-1>

[0586]

[Chemical Formula 117]



[0587]

20 2-Bromo-1-(2-bromoethoxy)-3,5-difluorobenzene

Carbon tetrabromide (606 mg) and triphenyl phosphine (575 mg) were added to a solution of 2-(2-bromo-3,5-difluorophenoxy)ethanol (370 mg) in

dichloromethane (3.7 mL) under an argon atmosphere under ice-cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. The reaction solution was concentrated under reduced pressure, ethyl acetate was added, and the precipitated solid was removed by filtration. The filtrate was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane followed by hexane : ethyl acetate = 9 : 1), to obtain the title compound as a colorless solid (453 mg).


¹H-NMR (400 MHz, CDCl₃) δ 3.69 (2H, t, J = 6.4 Hz), 4.32 (2H, t, J = 6.4 Hz), 6.47 (1H, dt, J = 10.0, 2.3 Hz), 6.59 (1H, td, J = 8.4, 2.3 Hz).

[0588]

<Reference Example 49-1>

[0589]

[Chemical Formula 118]



[0590]

4,6-Difluoro-2,3-dihydrobenzofuran

n-Butyllithium (0.55 mL, 2.65 mol/L hexane solution) was added to a solution of 2-bromo-1-(2-bromoethoxy)-3,5-difluorobenzene (440 mg) in tetrahydrofuran (5 mL) at -78°C under an argon atmosphere to produce a reaction solution. The reaction solution was stirred

for 2 hours, and stirred at room temperature for 1 hour. A saturated aqueous ammonium chloride solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over
5 anhydrous sodium sulfate. The solvent was removed under reduced pressure to obtain the title compound (205 mg).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.20 (2H, t, $J = 8.9$ Hz), 4.65 (2H, t, $J = 8.9$ Hz), 6.28-6.34 (2H, m).

[0591]

10 <Reference Example 50-1>

[0592]

[Chemical Formula 119]



[0593]

15 4,6-Difluoro-2,3-dihydrobenzofuran-5-carbaldehyde

Phosphorus oxychloride (778 μL) was added to a solution of 4,6-difluoro-2,3-dihydrobenzofuran (650 mg) in N,N-dimethylformamide (710 μL) under an argon atmosphere, to produce a reaction solution. The reaction solution was stirred
20 at 100°C for 20 hours. The reaction solution was allowed to cool to room temperature, and added to iced water, and the mixture was extracted with ethyl acetate. The organic layer was washed with a saturated sodium hydrogen carbonate aqueous solution and a brine,

and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1), to obtain the title compound as a pale yellow oil (121 mg).


5 $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.27 (2H, t, $J = 8.9$ Hz), 4.77 (2H, t, $J = 8.9$ Hz), 6.41 (1H, d, $J = 11.0$ Hz), 10.16 (1H, s).

[0594]

<Reference Example 51-1>

[0595]

10 [Chemical Formula 120]



[0596]

1-Benzyl-6-fluoroindoline

Potassium carbonate (4.45 g) and benzyl bromide (3.83 mL)
15 were added to a solution of 6-fluoroindoline (4.2 g) in methanol (38 mL), to produce a reaction solution. The reaction solution was heated to reflux for 3 hours. The reaction solution was allowed to cool to room temperature, and the insoluble was removed by filtration. The filtrate was concentrated under reduced pressure,
20 and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1), to obtain the title compound as a pale brown oil (6.14 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 2.92 (2H, t, $J = 8.5$ Hz), 3.38 (2H, t,


J = 8.5 Hz), 4.23 (2H, s), 6.18 (1H, dd, J = 10.9, 2.4 Hz), 6.27-6.32 (1H, m), 6.93-6.97 (1H, m), 7.25-7.34 (5H, m).

[0597]

<Reference Example 52-1>

5 [0598]

[Chemical Formula 121]



[0599]

1-Benzyl-6-fluoroindoline-5-carbaldehyde

10 Phosphorus oxychloride (518 μ L) was added to a solution of 1-benzyl-6-fluoroindoline (1.0 g) in N,N-dimethylformamide (4.9 mL) under ice-cooling, to produce a reaction solution. The reaction solution was stirred at 100°C for 2 hours. The reaction solution was allowed to cool to room temperature, and added to
15 iced water. A 2 mol/L sodium hydroxide aqueous solution was added to the iced water to adjust the pH to 10, and the mixture was extracted with chloroform. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel
20 column chromatography (hexane : ethyl acetate = 8 : 1 followed by 2 : 1), to obtain the title compound as a pale yellow oil (1.07 g).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 3.02 (2H, t, J = 8.5 Hz), 3.61 (2H, t,


J = 8.5 Hz), 4.39 (2H, s), 6.05 (1H, d, J = 12.1 Hz), 7.25-7.38 (5H, m), 7.48-7.49 (1H, m), 10.02 (1H, s).

[0600]

<Example 1-1>

5 [0601]

[Chemical Formula 122]



[0602]

(-)-1-(4-Fluorophenyl)-3-[(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea

10 Triethylamine (131 μ L) and diphenylphosphoryl azide (171 μ L) were added to a solution of (-)-(3S*, 4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid (170 mg) in toluene (3.6 mL) under an argon atmosphere to produce a reaction solution. The reaction solution was stirred
15 at room temperature for 1 hour. The temperature was heated up to 90°C and the reaction solution was stirred for 1 hour. After that, 4-fluoroaniline (140 μ L) was added to the reaction solution, and the reaction solution was stirred at 120°C for 3 hours. The reaction solution was cooled to room temperature, a 1 mol/L hydrochloric
20 acid was added, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine successively,

and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1, ethyl acetate, and ethyl acetate : methanol = 10 : 1 in turn), to obtain
5 the title compound as a white solid (44 mg).

^1H -NMR (400 MHz, DMSO- d_6) δ 3.11-3.17 (1H, m), 3.38-3.48 (2H, m), 3.69 (3H, s), 4.45 (1H, dd, $J = 10.9, 9.1$ Hz), 6.38 (1H, d, $J = 9.1$ Hz), 6.86 (2H, d, $J = 8.5$ Hz), 7.01 (2H, t, $J = 9.1$ Hz), 7.28 (2H, d, $J = 8.5$ Hz), 7.32-7.35 (2H, m), 7.88 (1H, s), 8.53 (1H,
10 s).

MS (ESI $^+$) m/z : 344 (MH^+).

$[\alpha]_{\text{D}}^{28} = -156$ (c 0.10, EtOH)




[0603]

The same method as in Example 1-1 was performed using a
15 corresponding carboxylic acid and aromatic amine to obtain the following Examples 1-2 to 1-87.

The structures and spectral data thereof are shown in Tables
47 to 75.




[0604]

[Table 47]

Ex.No	Str.	Chemical name	P.D.
1-2		(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.11-3.21 (1H, m), 3.41-3.50 (2H, m), 3.71 (3H, s), 4.46 (1H, dd, J = 10.9, 9.1 Hz), 6.47 (1H, d, J = 9.1 Hz), 6.87 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 7.90 (1H, s), 8.67 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺). [α] _D ²⁸ = -130 (c 0.10, EtOH)
1-3		(-)-1-(3,4-difluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13-3.20 (1H, m), 3.42-3.51 (2H, m), 3.71 (3H, s), 4.45 (1H, t, J = 9.7 Hz), 6.52 (1H, d, J = 8.5 Hz), 6.87 (2H, d, J = 8.5 Hz), 6.98-7.03 (1H, m), 7.21-7.28 (1H, m), 7.29 (2H, d, J = 8.5 Hz), 7.53-7.60 (1H, m), 7.91 (1H, s), 8.76 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁸ = -149 (c 0.10, EtOH)
1-4		(-)-1-(2,4-difluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17 (1H, t, J = 9.1 Hz), 3.36-3.49 (2H, m), 3.71 (3H, s), 4.47 (1H, dd, J = 11.2, 8.8 Hz), 6.80 (1H, d, J = 8.5 Hz), 6.88 (2H, d, J = 8.5 Hz), 6.93-6.98 (1H, m), 7.19-7.25 (1H, m), 7.30 (2H, d, J = 8.5 Hz), 7.93 (1H, s), 7.94-8.01 (1H, m), 8.28 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁸ = -126 (c 0.10, EtOH)




[0605]

[Table 48]

Ex.No	Str.	Chemical name	P.D.
1-5		(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.14-3.21 (1H, m), 3.43-3.52 (2H, m), 3.71 (3H, s), 4.48 (1H, t, J = 9.7 Hz), 6.67 (1H, d, J = 9.1 Hz), 6.87 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.53 (2H, dd, J = 7.0, 2.1 Hz), 7.64 (2H, dd, J = 7.0, 2.1 Hz), 7.93 (1H, s), 9.09 (1H, s). MS (ESI ⁺) m/z: 351 (MH ⁺). [α] _D ²⁸ = -166 (c 0.10, EtOH)
1-6		(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.79 (3H, s), 3.29 (1H, t, J = 9.7 Hz), 3.43 (1H, q, J = 9.5 Hz), 3.54 (1H, t, J = 8.5 Hz), 3.71 (3H, s), 4.46 (1H, t, J = 10.0 Hz), 6.53 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s). MS (ESI ⁺) m/z: 374 (MH ⁺). [α] _D ²⁶ = -170 (c 0.10, EtOH)
1-7		1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.79 (3H, s), 3.28 (1H, t, J = 10.0 Hz), 3.43 (1H, q, J = 9.5 Hz), 3.54 (1H, t, J = 8.8 Hz), 3.71 (3H, s), 4.47 (1H, t, J = 10.0 Hz), 6.46 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.02 (2H, t, J = 8.8 Hz), 7.29 (2H, d, J = 8.5 Hz), 7.35 (2H, dd, J = 8.8, 4.8 Hz), 8.58 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺).




[0606]

[Table 49]

Ex.No	Str.	Chemical name	P.D.
1-8		(-)-1-[(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13-3.20 (1H, m), 3.43-3.50 (2H, m), 3.79 (3H, s), 4.47 (1H, t, J = 9.7 Hz), 6.42 (1H, d, J = 8.5 Hz), 7.00-7.14 (4H, m), 7.29-7.37 (3H, m), 7.91 (1H, s), 8.58 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁸ = -130 (c 0.10, EtOH)
1-9		(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17 (1H, t, J = 9.8 Hz), 3.45 (1H, t, J = 8.9 Hz), 3.68 (1H, q, J = 9.8 Hz), 3.73 (3H, s), 4.56 (1H, dd, J = 11.3, 8.9 Hz), 6.41 (1H, d, J = 8.6 Hz), 6.74-6.80 (2H, m), 7.00-7.05 (2H, m), 7.32-7.37 (2H, m), 7.45 (1H, t, J = 8.9 Hz), 7.94 (1H, s), 8.59 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁸ = -99 (c 0.10, EtOH)
1-10		1-[(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.07 (1H, t, J = 9.8 Hz), 3.50 (1H, t, J = 8.5 Hz), 3.74 (3H, s), 3.91 (1H, q, J = 9.8 Hz), 4.66 (1H, dd, J = 11.3, 8.9 Hz), 6.42 (1H, d, J = 8.6 Hz), 6.93 (1H, dd, J = 8.6, 2.4 Hz), 7.00-7.06 (3H, m), 7.33-7.37 (2H, m), 7.59 (1H, d, J = 8.6 Hz), 7.94 (1H, s), 8.57 (1H, s). MS (ESI ⁺) m/z: 378 (MH ⁺).




[0607]

[Table 50]

Ex.No	Str.	Chemical name	P.D.
1-11		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.27 (3H, s), 3.05 (1H, t, J = 9.7 Hz), 3.46 (1H, t, J = 8.4 Hz), 3.69 (3H, s), 3.74 (1H, dd, J = 9.8, 8.4 Hz), 4.51 (1H, dd, J = 11.5, 9.1 Hz), 6.37 (1H, d, J = 9.1 Hz), 6.71 (1H, d, J = 3.0 Hz), 6.75 (1H, dd, J = 8.5, 3.0 Hz), 7.00-7.04 (2H, m), 7.32-7.36 (2H, m), 7.40 (1H, d, J = 8.5 Hz), 7.89 (1H, s), 8.51 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺). [α] _D ²³ = -187 (c 0.35, EtOH)
1-12		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.33 (1H, m), 3.44 (1H, t, J = 9.5 Hz), 3.75 (3H, s), 3.79 (1H, q, J = 9.8 Hz), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 10.4 Hz), 6.98-7.04 (2H, m), 7.30-7.34 (2H, m), 8.05 (1H, s), 8.66 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁸ = -156 (c 0.10, EtOH)
1-13		(-)-1-[(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.8 Hz), 3.43 (1H, t, J = 8.9 Hz), 3.72 (1H, q, J = 9.8 Hz), 3.80 (3H, s), 4.53 (1H, dd, J = 11.6, 8.6 Hz), 6.42 (1H, d, J = 8.6 Hz), 7.00-7.10 (3H, m), 7.32-7.36 (2H, m), 7.52 (1H, dd, J = 11.6, 8.6 Hz), 7.96 (1H, s), 8.62 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁸ = -123 (c 0.10, EtOH)

[0608]

[Table 51]

Ex.No	Str.	Chemical name	P.D.
1-14		(-)-1-[(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13-3.19 (1H, m), 3.45-3.56 (2H, m), 3.87 (3H, s), 4.49 (1H, dd, J = 11.3, 8.9 Hz), 6.44 (1H, d, J = 9.2 Hz), 7.01-7.07 (2H, m), 7.19-7.26 (2H, m), 7.33-7.38 (2H, m), 7.95 (1H, s), 8.61 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁸ = -111 (c 0.10, EtOH)
1-15		(-)-1-[(3S*,4R*)-4-(2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.10-3.18 (3H, m), 3.39-3.47 (2H, m), 4.40-4.49 (3H, m), 6.40 (1H, d, J = 8.6 Hz), 6.68 (1H, d, J = 8.6 Hz), 7.00-7.08 (3H, m), 7.24 (1H, s), 7.33-7.37 (2H, m), 7.88 (1H, s), 8.55 (1H, s). MS (ESI ⁺) m/z: 356 (MH ⁺). [α] _D ²⁸ = -130 (c 0.10, EtOH)
1-16		(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.09-3.18 (3H, m), 3.39-3.46 (1H, m), 3.67 (1H, q, J = 10.0 Hz), 4.50-4.57 (3H, m), 6.41 (1H, d, J = 9.2 Hz), 6.62 (1H, d, J = 11.0 Hz), 7.02 (2H, t, J = 8.9 Hz), 7.33-7.38 (3H, m), 7.93 (1H, s), 8.60 (1H, s). MS (ESI ⁺) m/z: 374 (MH ⁺). [α] _D ²⁸ = -152 (c 0.10, EtOH)




[0609]

[Table 52]

Ex.No	Str.	Chemical name	P.D.
1-17		(-)-1-[(3S*,4R*)-4-(7-fluorochroman-6-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.84-1.90 (2H, m), 2.60-2.73 (2H, m), 3.16 (1H, t, J = 9.8 Hz), 3.42 (1H, t, J = 8.9 Hz), 3.64 (1H, q, J = 9.8 Hz), 4.09 (2H, t, J = 4.9 Hz), 4.51 (1H, dd, J = 11.3, 8.9 Hz), 6.40 (1H, d, J = 8.6 Hz), 6.54 (1H, d, J = 11.6 Hz), 7.00-7.06 (2H, m), 7.20 (1H, d, J = 8.6 Hz), 7.33-7.37 (2H, m), 7.92 (1H, s), 8.58 (1H, s). MS (ESI ⁺) m/z: 388 (MH ⁺). $[\alpha]_D^{28} = -220$ (c 0.10, EtOH)
1-18		(-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.38 (1H, t, J = 9.2 Hz), 3.47 (1H, t, J = 8.9 Hz), 3.83 (3H, s), 3.96 (1H, q, J = 9.5 Hz), 4.51 (1H, dd, J = 10.4, 7.9 Hz), 6.48 (1H, d, J = 7.9 Hz), 7.01 (2H, t, J = 8.9 Hz), 7.29-7.35 (2H, m), 7.38 (1H, dd, J = 11.6, 2.4 Hz), 7.95 (1H, s), 8.20 (1H, d, J = 2.4 Hz), 8.62 (1H, s). MS (ESI ⁺) m/z: 363 (MH ⁺). $[\alpha]_D^{28} = -142$ (c 0.10, EtOH)
1-19		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.7 Hz), 3.49 (1H, t, J = 9.5 Hz), 3.60 (1H, q, J = 9.5 Hz), 3.78 (3H, s), 4.28 (1H, dd, J = 10.9, 9.1 Hz), 6.11 (1H, d, J = 3.6 Hz), 6.47 (1H, d, J = 9.1 Hz), 6.62 (1H, d, J = 3.6 Hz), 7.01-7.08 (2H, m), 7.36-7.41 (2H, m), 7.93 (1H, s), 8.64 (1H, s). MS (ESI ⁺) m/z: 350 (MH ⁺). $[\alpha]_D^{28} = -129$ (c 0.10, EtOH)




[0610]

[Table 53]

Ex.No	Str.	Chemical name	P.D.
1-20		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.49 (3H, s), 3.20-3.27 (1H, m), 3.50-3.58 (2H, m), 4.54 (1H, t, J = 9.7 Hz), 6.47 (1H, d, J = 9.1 Hz), 7.05-7.11 (2H, m), 7.26 (2H, d, J = 8.5 Hz), 7.35-7.42 (4H, m), 7.97 (1H, s), 8.61 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺). [α] _D ²⁷ = -544 (c 0.26, DMSO)
1-21		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methylsulfonylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.10 (3H, s), 3.29-3.31 (1H, m), 3.46 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.79-3.87 (1H, m), 4.58 (1H, dd, J = 11.0, 8.0 Hz), 6.70-6.78 (3H, m), 7.57 (2H, d, J = 9.2 Hz), 7.72 (2H, d, J = 9.2 Hz), 8.09 (1H, s), 9.25 (1H, s). MS (ESI ⁺) m/z: 440 (MH ⁺). [α] _D ²⁷ = -138 (c 0.10, EtOH)
1-22		(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.31 (1H, m), 3.44 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.75-3.83 (1H, m), 4.56 (1H, t, J = 9.6 Hz), 6.51 (1H, d, J = 8.0 Hz), 6.73 (2H, d, J = 12.0 Hz), 7.22 (2H, d, J = 9.2 Hz), 7.35 (2H, d, J = 9.2 Hz), 8.06 (1H, s), 8.79 (1H, s). MS (ESI ⁺) m/z: 396 (MH ⁺). [α] _D ²⁷ = -195 (c 0.10, EtOH)




[0611]

[Table 54]

Ex.No	Str.	Chemical name	P.D.
1-23		(-)-1-(4-cyano phenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.33 (1H, m), 3.45 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.78-3.86 (1H, m), 4.58 (1H, dd, J = 11.0, 8.0 Hz), 6.70-6.77 (3H, m), 7.51 (2H, d, J = 9.2 Hz), 7.63 (2H, d, J = 9.2 Hz), 8.09 (1H, s), 9.24 (1H, s). MS (ESI ⁺) m/z: 387 (MH ⁺). [α] _D ²⁶ = -163 (c 0.10, EtOH)
1-24		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methoxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29 (1H, t, J = 8.6 Hz), 3.43 (1H, t, J = 8.9 Hz), 3.66 (s, 3H), 3.70-3.82 (1H, m), 3.75 (3H, s), 4.55 (1H, dd, J = 10.4, 8.6 Hz), 6.34 (1H, d, J = 8.6 Hz), 6.71-6.79 (4H, m), 7.22 (2H, d, J = 8.6 Hz), 8.03 (1H, s), 8.40 (1H, s). MS (ESI ⁺) m/z: 392 (MH ⁺). [α] _D ²⁸ = -168 (c 0.29, EtOH)
1-25		(-)-1-(benzo[d][1,3]dioxol-5-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29 (1H, t, J = 8.6 Hz), 3.43 (1H, t, J = 8.9 Hz), 3.72-3.82 (1H, m), 3.75 (3H, s), 4.55 (1H, dd, J = 11.0, 8.6 Hz), 5.90 (2H, s), 6.37 (1H, d, J = 7.9 Hz), 6.64 (1H, dd, J = 8.3, 2.1 Hz), 6.70-6.76 (3H, m), 7.06 (1H, d, J = 2.1 Hz), 8.04 (1H, s), 8.50 (1H, s). MS (ESI ⁺) m/z: 406 (MH ⁺). [α] _D ²⁸ = -150 (c 0.30, EtOH)




[0612]

[Table 55]

Ex.No	Str.	Chemical name	P.D.
1-26		(-)-1-(5-chlorothiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.31 (1H, m), 3.46 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.82-3.91 (1H, m), 4.55 (1H, dd, J = 9.8, 6.8 Hz), 6.75 (2H, d, J = 11.0 Hz), 6.88 (1H, d, J = 6.8 Hz), 7.31 (1H, s), 8.11 (1H, s), 10.9 (1H, s). MS (ESI ⁺) m/z: 403 (MH ⁺). [α] _D ²⁷ = -88 (c 0.10, EtOH)
1-27		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrimidin-4-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.31 (1H, m), 3.46 (1H, t, J = 8.6 Hz), 3.75 (3H, s), 3.80-3.88 (1H, m), 4.60 (1H, dd, J = 10.4, 8.0 Hz), 6.75 (2H, d, J = 11.0 Hz), 7.46 (1H, d, J = 6.1 Hz), 7.86 (1H, brs), 8.14 (1H, s), 8.44 (1H, d, J = 5.5 Hz), 8.70 (1H, s), 9.68 (1H, brs). MS (ESI ⁺) m/z: 364 (MH ⁺). [α] _D ²⁹ = -180 (c 0.10, DMSO)
1-28		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyridin-2-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.30 (1H, m), 3.40-3.47 (1H, m), 3.75 (3H, s), 3.79-3.88 (1H, m), 4.58-4.63 (1H, m), 6.73 (2H, d, J = 10.4 Hz), 6.90-6.93 (1H, m), 7.28 (1H, d, J = 8.0 Hz), 7.62-7.67 (1H, m), 8.10 (1H, m), 8.14-8.16 (1H, m), 8.37-8.43 (1H, m), 9.22 (1H, s). MS (ESI ⁺) m/z: 363 (MH ⁺). [α] _D ²⁷ = -188 (c 0.10, EtOH)




[0613]

[Table 56]

Ex.No	Str.	Chemical name	P.D.
1-29		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyridin-3-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.31 (1H, m), 3.45 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.82 (1H, q, J = 9.5 Hz), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.66 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 7.21 (1H, dd, J = 8.6, 4.9 Hz), 7.78-7.82 (1H, m), 8.07 (1H, s), 8.08-8.10 (1H, m), 8.47 (1H, d, J = 2.4 Hz), 8.87 (1H, s). MS (ESI ⁺) m/z: 363 (MH ⁺). [α] _D ²⁷ = -168 (c 0.10, EtOH)
1-30		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrimidin-5-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.50 (2H, m), 3.74 (3H, s), 3.75-3.90 (1H, m), 4.58 (1H, dd, J = 9.8, 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 6.92 (1H, d, J = 8.6 Hz), 8.08 (1H, s), 8.71 (1H, s), 8.78 (2H, s), 9.11 (1H, s). MS (ESI ⁺) m/z: 364 (MH ⁺). [α] _D ²⁸ = -139 (c 0.10, DMSO)
1-31		(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.31 (1H, m), 3.45 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.83 (1H, q, J = 9.2 Hz), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.70-6.79 (3H, m), 7.32 (1H, d, J = 8.6 Hz), 7.86 (1H, dd, J = 8.6, 3.0 Hz), 8.07 (1H, s), 8.34 (1H, d, J = 3.0 Hz), 9.07 (1H, s). MS (ESI ⁺) m/z: 397 (MH ⁺). [α] _D ²⁷ = -152 (c 0.10, EtOH)




[0614]

[Table 57]

Ex.No	Str.	Chemical name	P.D.
1-32		(-)-1-(5-chloropyridin-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.35 (1H, m), 3.47 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.81 (1H, q, J = 9.2 Hz), 4.59 (1H, dd, J = 11.0, 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 7.49 (1H, d, J = 8.6 Hz), 7.73-7.79 (2H, m), 8.11 (1H, s), 8.20 (1H, d, J = 3.0 Hz), 9.35 (1H, s). MS (ESI ⁺) m/z: 397 (MH ⁺). [α] _D ²⁷ = -180 (c 0.10, EtOH)
1-33		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(pyrazin-2-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.36-3.30 (1H, m), 3.48 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.85 (1H, q, J = 9.5 Hz), 4.60 (1H, dd, J = 10.4, 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 7.70 (1H, d, J = 8.6 Hz), 8.12 (1H, s), 8.16-8.22 (2H, m), 8.79 (1H, s), 9.51 (1H, s). MS (ESI ⁺) m/z: 364 (MH ⁺). [α] _D ²⁸ = -178 (c 0.10, DMSO)
1-34		(-)-1-(benzo[d]thiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.36 (1H, m), 3.48 (1H, t, J = 8.6 Hz), 3.75 (3H, s), 3.89 (1H, q, J = 9.2, Hz), 4.60 (1H, dd, J = 10.4, 9.2 Hz), 6.75 (2H, d, J = 11.0 Hz), 7.06-7.23 (2H, m), 7.30-7.37 (1H, m), 7.54-7.62 (1H, m), 7.83 (1H, d, J = 6.7 Hz), 8.14 (1H, s), 11.0 (1H, brs). MS (ESI ⁺) m/z: 419 (MH ⁺). [α] _D ²⁸ = -164 (c 0.10, DMSO)

[0615]

[Table 58]

Ex.No	Str.	Chemical name	P.D.
1-35		(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.31 (1H, m), 3.46 (1H, t, J = 8.6 Hz), 3.75 (3H, s), 3.86 (1H, q, J = 9.2 Hz), 4.59 (1H, dd, J = 11.0, 8.6 Hz), 6.66 (1H, s), 6.75 (2H, d, J = 11.0 Hz), 6.82 (1H, d, J = 8.6 Hz), 7.09 (1H, t, J = 8.0 Hz), 7.21 (1H, t, J = 8.0 Hz), 7.52 (1H, d, J = 8.0 Hz), 7.69 (1H, d, J = 8.0 Hz), 8.09 (1H, s), 10.1 (1H, s). MS (ESI ⁺) m/z: 418 (MH ⁺). [α] _D ²⁷ = -74 (c 0.10, EtOH)
1-36		(-)-1-(benzo[d]oxazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.32-3.38 (1H, m), 3.51 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.97 (1H, q, J = 9.5 Hz), 4.67 (1H, dd, J = 9.8, 8.0 Hz), 6.74 (2H, d, J = 11.0 Hz), 7.17-7.30 (2H, m), 7.46-7.57 (2H, m), 8.16 (1H, s), 8.57 (1H, d, J = 7.3 Hz), 11.14 (1H, brs). MS (ESI ⁺) m/z: 403 (MH ⁺). [α] _D ²⁸ = -158 (c 0.10, EtOH)
1-37		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylisoxazol-3-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.29 (3H, s), 3.27-3.32 (1H, m), 3.45 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.76-3.84 (1H, m), 4.55 (1H, t, J = 9.8 Hz), 6.31 (1H, s), 6.70-6.81 (3H, m), 8.09 (1H, s), 9.44 (1H, s). MS (ESI ⁺) m/z: 367 (MH ⁺). [α] _D ²⁷ = -154 (c 0.10, EtOH)




[0616]

[Table 59]

Ex.No	Str.	Chemical name	P.D.
1-38		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(2-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.35 (1H, m), 3.46 (1H, t, J = 9.2 Hz), 3.67-3.74 (1H, m), 3.75 (3H, s), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.75 (2H, d, J = 10.4 Hz), 6.88-6.96 (2H, m), 7.03 (1H, t, J = 7.2 Hz), 7.12-7.18 (1H, m), 7.93-7.98 (1H, m), 8.11 (1H, s), 8.41 (1H, d, J = 2.4 Hz). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁷ = -152 (c 0.10, EtOH)
1-39		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylthiophen-2-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.25 (3H, s), 3.26-3.32 (1H, m), 3.43 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.76-3.84 (1H, m), 4.53 (1H, dd, J = 9.2, 8.0 Hz), 6.17 (1H, d, J = 3.2 Hz), 6.38 (1H, br s), 6.52 (1H, d, J = 8.0 Hz), 6.73 (2H, d, J = 10.3 Hz), 8.03 (1H, s), 9.42 (1H, s). MS (ESI ⁺) m/z: 382 (MH ⁺). [α] _D ²⁷ = -162 (c 0.10, EtOH)
1-40		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.68 (3H, d, J = 6.7 Hz), 3.64-3.71 (1H, m), 3.71 (3H, s), 3.74-3.80 (1H, m), 4.79 (1H, dd, J = 12.1, 9.1 Hz), 6.37 (1H, d, J = 9.1 Hz), 6.89 (2H, d, J = 9.1 Hz), 7.01-7.06 (2H, m), 7.25 (2H, d, J = 8.5 Hz), 7.35-7.39 (2H, m), 8.05 (1H, s), 8.51 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺). [α] _D ²⁵ = -149 (c 0.22, EtOH).




[0617]

[Table 60]

Ex.No	Str.	Chemical name	P.D.
1-41		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.04 (3H, d, J = 6.1 Hz), 2.86 (1H, dd, J = 11.5, 9.1 Hz), 3.45-3.54 (1H, m), 3.71 (3H, s), 4.53 (1H, dd, J = 11.5, 8.5 Hz), 6.34 (1H, d, J = 8.5 Hz), 6.87 (2H, d, J = 8.5 Hz), 6.99-7.03 (2H, m), 7.27 (2H, d, J = 8.5 Hz), 7.31-7.34 (2H, m), 7.98 (1H, s), 8.50 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺). [α] _D ²⁵ = -108 (c 0.39, EtOH).
1-42		1-[(3S*,4R*)-4-(4-ethyl-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.15 (3H, t, J = 7.6 Hz), 2.58 (2H, q, J = 7.6 Hz), 3.29-3.36 (1H, m), 3.41-3.48 (1H, m), 3.86 (1H, q, J = 9.8 Hz), 4.59 (1H, dd, J = 10.4, 8.6 Hz), 6.48 (1H, d, J = 8.6 Hz), 6.96 (2H, d, J = 9.8 Hz), 7.01 (2H, t, J = 8.9 Hz), 7.30-7.34 (2H, m), 8.06 (1H, s), 8.68 (1H, s). MS (ESI ⁺) m/z: 378 (MH ⁺).
1-43		(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-2-oxo-4-phenylpyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17-3.23 (1H, m), 3.49-3.58 (2H, m), 4.51 (1H, dd, J = 11.2, 8.5 Hz), 6.51 (1H, d, J = 8.5 Hz), 7.21-7.25 (3H, m), 7.29-7.33 (2H, m), 7.36-7.39 (4H, m), 7.94 (1H, s), 8.70 (1H, s). MS (ESI ⁺) m/z: 330 (MH ⁺). [α] _D ²⁵ = +122 (c 0.35, DMSO)




[0618]

[Table 61]

Ex.No	Str.	Chemical name	P.D.
1-44		(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.11-3.21 (1H, m), 3.41-3.50 (2H, m), 3.71 (3H, s), 4.46 (1H, dd, J = 10.9, 9.1 Hz), 6.47 (1H, d, J = 9.1 Hz), 6.87 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 7.90 (1H, s), 8.67 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺). [α] _D ²⁵ = +146 (c 0.30, EtOH)
1-45		(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-4-(3-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.16-3.23 (1H, m), 3.47-3.56 (2H, m), 3.72 (3H, s), 4.52 (1H, t, J = 9.7 Hz), 6.49 (1H, d, J = 9.1 Hz), 6.80 (1H, dd, J = 8.5, 2.1 Hz), 6.93 (1H, d, J = 7.4 Hz), 6.97 (1H, s), 7.20-7.24 (3H, m), 7.38 (2H, d, J = 8.5 Hz), 7.92 (1H, s), 8.68 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺). [α] _D ²⁵ = +115 (c 0.35, DMSO)
1-46		(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-4-(2-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.09 (1H, t, J = 9.4 Hz), 3.40-3.50 (1H, m), 3.69-3.75 (1H, m), 3.78 (3H, s), 4.72 (1H, dd, J = 11.2, 8.8 Hz), 6.45 (1H, d, J = 9.1 Hz), 6.88-6.93 (1H, m), 6.98 (1H, d, J = 7.3 Hz), 7.20-7.24 (3H, m), 7.34-7.39 (3H, m), 7.88 (1H, s), 8.66 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺). [α] _D ²⁵ = +110 (c 0.35, EtOH)




[0619]

[Table 62]

Ex.No	Str.	Chemical name	P.D.
1-47		(+)-1-[(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.33 (1H, m), 3.44 (1H, t, J = 9.5 Hz), 3.75 (3H, s), 3.79 (1H, q, J = 9.8 Hz), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 10.4 Hz), 6.98-7.04 (2H, m), 7.30-7.34 (2H, m), 8.05 (1H, s), 8.66 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁸ +175 (c 0.10, EtOH)
1-48		(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.79 (3H, s), 3.29 (1H, t, J = 9.7 Hz), 3.43 (1H, q, J = 9.5 Hz), 3.54 (1H, t, J = 8.5 Hz), 3.71 (3H, s), 4.46 (1H, t, J = 10.0 Hz), 6.53 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s). MS (ESI ⁺) m/z: 374 (MH ⁺). [α] _D ²⁵ = +162 (c 0.31, EtOH)
1-49		(+)-1-(4-fluorophenyl)-3-[(3R*,4S*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.04 (3H, d, J = 6.1 Hz), 2.86 (1H, dd, J = 11.5, 9.1 Hz), 3.45-3.54 (1H, m), 3.71 (3H, s), 4.53 (1H, dd, J = 11.5, 8.5 Hz), 6.34 (1H, d, J = 8.5 Hz), 6.87 (2H, d, J = 8.5 Hz), 6.99-7.03 (2H, m), 7.27 (2H, d, J = 8.5 Hz), 7.31-7.34 (2H, m), 7.98 (1H, s), 8.50 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺). [α] _D ²⁵ +116 (c 0.38, EtOH).




[0620]

[Table 63]

Ex.No	Str.	Chemical name	P.D.
1-50		(+)-1-(4-fluorophenyl)-3-[(3R*,4S*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.68 (3H, d, J = 6.7 Hz), 3.64-3.71 (1H, m), 3.71 (3H, s), 3.74-3.80 (1H, m), 4.79 (1H, dd, J = 12.1, 9.1 Hz), 6.37 (1H, d, J = 9.1 Hz), 6.89 (2H, d, J = 9.1 Hz), 7.01-7.06 (2H, m), 7.25 (2H, d, J = 8.5 Hz), 7.35-7.39 (2H, m), 8.05 (1H, s), 8.51 (1H, s). MS (ESI ⁺) m/z: 358 (MH ⁺). [α] _D ²⁵ +167 (c 0.45, EtOH).
1-51		(+)-1-(4-fluorophenyl)-3-[(3R*,4S*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (1H, t, J = 9.7 Hz), 3.49 (1H, t, J = 9.5 Hz), 3.60 (1H, q, J = 9.5 Hz), 3.78 (3H, s), 4.28 (1H, dd, J = 10.9, 9.1 Hz), 6.11 (1H, d, J = 3.6 Hz), 6.47 (1H, d, J = 9.1 Hz), 6.62 (1H, d, J = 3.6 Hz), 7.04 (2H, m), 7.37-7.40 (2H, m), 7.93 (1H, s), 8.64 (1H, s). MS (ESI ⁺) m/z: 350 (MH ⁺). [α] _D ²⁵ = +122 (c 0.10, EtOH)
1-52		(+)-1-[(3R*,4S*)-4-[4-(difluoromethoxy)phenyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17-3.22 (1H, m), 3.47-3.57 (2H, m), 4.50 (1H, dd, J = 10.9, 9.1 Hz), 6.44 (1H, d, J = 9.1 Hz), 7.00-7.05 (2H, m), 7.00-7.37 (1H, m), 7.12 (2H, d, J = 8.5 Hz), 7.33-7.37 (2H, m), 7.44 (2H, d, J = 8.5 Hz), 7.94 (1H, s), 8.57 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²³ = +107 (c 0.35, EtOH)




[0621]

[Table 64]

Ex.No	Str.	Chemical name	P.D.
1-53		(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.11-3.21 (1H, m), 3.41-3.50 (2H, m), 3.71 (3H, s), 4.46 (1H, dd, J = 10.9, 9.1 Hz), 6.47 (1H, d, J = 9.1 Hz), 6.87 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 7.90 (1H, s), 8.67 (1H, s). MS (ESI ⁺) m/z: 360 (MH ⁺).
1-54		(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.11-3.17 (1H, m), 3.38-3.48 (2H, m), 3.69 (3H, s), 4.45 (1H, dd, J = 10.9, 9.1 Hz), 6.38 (1H, d, J = 9.1 Hz), 6.86 (2H, d, J = 8.5 Hz), 7.01 (2H, t, J = 9.1 Hz), 7.28 (2H, d, J = 8.5 Hz), 7.32-7.35 (2H, m), 7.88 (1H, s), 8.53 (1H, s). MS (ESI ⁺) m/z: 344 (MH ⁺).
1-55		(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.79 (3H, s), 3.29 (1H, t, J = 9.7 Hz), 3.43 (1H, q, J = 9.5 Hz), 3.54 (1H, t, J = 8.5 Hz), 3.71 (3H, s), 4.46 (1H, t, J = 10.0 Hz), 6.53 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz), 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s). MS (ESI ⁺) m/z: 374 (MH ⁺).




[0622]

[Table 65]

Ex.No.	Str.	Chemical name	P.D.
1-56		1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-hydroxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26–3.31 (1H, m), 3.43 (1H, t, J = 9.7 Hz), 3.72–3.79 (4H, m), 4.64 (1H, dd, J = 10.3, 7.8 Hz), 6.29 (1H, d, J = 8.5 Hz), 6.57–6.60 (2H, m), 6.73 (2H, d, J = 10.9 Hz), 7.06–7.09 (2H, m), 8.02 (1H, s), 8.25 (1H, s), 8.92 (1H, s). MS (ESI ⁺) m/z: 378 (MH ⁺).
1-57		1-[(3S*,4R*)-4-(2,6-difluoro-3-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27–3.32 (1H, m), 3.44 (1H, t, J = 9.7 Hz), 3.72–3.80 (4H, m), 4.56 (1H, dd, J = 10.9, 8.5 Hz), 6.27 (1H, dd, J = 7.3, 1.8 Hz), 6.37 (1H, d, J = 8.5 Hz), 6.65 (1H, dd, J = 7.2, 1.8 Hz), 6.73 (2H, d, J = 10.9 Hz), 6.91–6.96 (2H, m), 8.05 (1H, s), 8.49 (1H, s), 9.16 (1H, s). MS (ESI ⁺) m/z: 378 (MH ⁺).
1-58		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-methylisothiazol-5-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.21 (3H, s), 3.27–3.32 (1H, m), 3.45 (1H, t, J = 9.6 Hz), 3.75 (3H, s), 3.87 (1H, q, J = 9.6 Hz), 4.54 (1H, dd, J = 9.6, 8.6 Hz), 6.46 (1H, s), 6.75 (2H, d, J = 10.4 Hz), 7.16 (1H, d, J = 8.6 Hz), 8.09 (1H, s), 10.5 (1H, s). MS (ESI ⁺) m/z: 383 (MH ⁺). [α] _D ²⁶ = -264 (c 0.03, EtOH)




[0623]

[Table 66]

Ex.No.	Str.	Chemical name	P.D.
1-59		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(1H-indol-5-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.30 (1H, t, J = 9.2 Hz), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.77 (1H, q, J = 9.2 Hz), 4.58 (1H, dd, J = 10.4, 8.6 Hz), 6.27 (1H, brs), 6.32 (1H, d, J = 8.6 Hz), 6.71-6.77 (2H, d, J = 10.4 Hz), 6.94 (1H, dd, J = 8.6, 1.8 Hz), 7.20 (1H, d, J = 8.6 Hz), 7.23 (1H, t, J = 3.0 Hz), 7.52 (1H, d, J = 1.8 Hz), 8.04 (1H, s), 8.31 (1H, s), 10.9 (1H, s). MS (ESI ⁺) m/z: 401 (MH ⁺). [α] _D ²⁷ = -156 (c 0.11, EtOH)
1-60		1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-iodophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.34 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.79 (1H, q, J = 9.2 Hz), 4.55 (1H, dd, J = 11, 8.6 Hz), 6.53-6.65 (1H, m), 6.79-6.77 (2H, m), 7.18 (2H, d, J = 8.6 Hz), 7.49 (2H, d, J = 8.6 Hz), 8.05 (1H, s), 8.79-8.88 (1H, m). MS (ESI ⁺) m/z: 488 (MH ⁺).
1-61		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(oxazol-2-yl)phenyl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.33 (1H, m), 3.46 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.82 (1H, q, J = 9.2 Hz), 4.58 (1H, dd, J = 10.4, 8.0 Hz), 6.64 (1H, brs), 6.71-6.78 (2H, m), 7.29 (1H, s), 7.49 (2H, d, J = 8.6 Hz), 7.80 (2H, d, J = 8.6 Hz), 8.08 (1H, s), 8.11 (1H, s), 9.02 (1H, brs). MS (ESI ⁺) m/z: 429 (MH ⁺). [α] _D ²⁹ = -166 (c 0.15, EtOH)

[0624]

[Table 67]

Ex.No.	Str.	Chemical name	P.D.
1-62		(-)-1-(4-cyclopropylphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.52-0.56 (2H, m), 0.81-0.87 (2H, m), 1.76-1.82 (1H, m), 3.29 (1H, t, J = 9.2 Hz), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.77 (1H, q, J = 9.2 Hz), 4.55 (1H, dd, J = 10.4, 8.6 Hz), 6.39 (1H, d, J = 8.6 Hz), 6.70-6.76 (2H, m), 6.89 (2H, d, J = 8.6 Hz), 7.18 (2H, d, J = 8.6 Hz), 8.04 (1H, s), 8.50 (1H, s). MS (ESI ⁺) m/z: 402 (MH ⁺). [α] _D ²⁵ = -157 (c 0.14, EtOH)
1-63		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethyl)phenyl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.35 (1H, m), 3.47 (1H, t, J = 9.2 Hz), 3.76 (3H, s), 3.80-3.88 (1H, m), 4.60 (1H, dd, J = 11.0, 8.6 Hz), 6.65 (1H, d, J = 7.9 Hz), 6.75 (2H, d, J = 11.0 Hz), 7.50-7.55 (4H, m), 8.09 (1H, s), 9.11 (1H, s). MS (ESI ⁺) m/z: 430 (MH ⁺). [α] _D ²⁶ = -95 (c 0.13, EtOH)
1-64		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluoro-4-methylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.10 (3H, s), 3.25-3.35 (1H, m), 3.41-3.47 (1H, m), 3.75 (3H, s), 3.77-3.83 (1H, m), 4.55 (1H, dd, J = 11.0, 8.6 Hz), 6.52 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.91 (1H, dd, J = 8.3, 2.1 Hz), 7.06 (1H, t, J = 8.9 Hz), 7.28 (1H, dd, J = 12.5, 2.1 Hz), 8.05 (1H, s), 8.77 (1H, s). MS (ESI ⁺) m/z: 394 (MH ⁺). [α] _D ²⁵ = -125 (c 0.13, EtOH)

[0625]

[Table 68]

Ex.No.	Str.	Chemical name	P.D.
1-65		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-methylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.14 (3H, s), 3.26-3.33 (1H, m), 3.44 (1H, t, J = 8.9 Hz), 3.75 (3H, s), 3.77-3.83 (1H, m), 4.55 (1H, dd, J = 11.0, 8.6 Hz), 6.44 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.94 (1H, t, J = 9.2 Hz), 7.07-7.12 (1H, m), 7.23-7.26 (1H, m), 8.04 (1H, s), 8.57 (1H, s). MS (ESI ⁺) m/z: 394 (MH ⁺). [α] _D ²⁶ = -151 (c 0.14, EtOH)
1-66		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(6-methylpyridin-3-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.34 (3H, s), 3.25-3.34 (1H, m), 3.44 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.77-3.85 (1H, m), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.57 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 7.06 (1H, d, J = 8.6 Hz), 7.68 (1H, dd, J = 8.6, 2.4 Hz), 8.05 (1H, s), 8.34 (1H, d, J = 2.4 Hz), 8.72 (1H, s). MS (ESI ⁺) m/z: 377 (MH ⁺). [α] _D ²⁴ = -168 (c 0.22, EtOH)
1-67		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[6-(trifluoromethyl)pyridin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.35 (1H, m), 3.40-3.50 (1H, m), 3.75 (3H, s), 3.81-3.90 (1H, m), 4.59 (1H, dd, J = 10.7, 8.3 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.84 (1H, d, J = 8.6 Hz), 7.71 (1H, d, J = 9.2 Hz), 8.06-8.11 (2H, m), 8.62 (1H, d, J = 2.4 Hz), 9.36 (1H, s). MS (ESI ⁺) m/z: 431 (MH ⁺). [α] _D ²⁴ = -236 (c 0.20, EtOH)




[0626]

[Table 69]

Ex.No.	Str.	Chemical name	P.D.
1-68		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxy-4-methylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.98 (3H, s), 3.26-3.33 (1H, m), 3.40-3.47 (1H, m), 3.70-3.79 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.31 (1H, d, J = 7.9 Hz), 6.54 (1H, dd, J = 8.6, 2.4 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.81 (1H, d, J = 8.6 Hz), 6.97 (1H, d, J = 2.4 Hz), 8.05 (1H, s), 8.39 (1H, s), 9.08 (1H, s). MS (ESI ⁺) m/z: 392 (MH ⁺). [α] _D ²⁵ = -198 (c 0.15, EtOH)
1-69		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(2-fluoro-4-methylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.21 (3H, s), 3.26-3.33 (1H, m), 3.46 (1H, t, J = 9.5 Hz), 3.66-3.74 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.82-6.86 (2H, m), 6.97 (1H, d, J = 12.8 Hz), 7.78 (1H, t, J = 8.3 Hz), 8.10 (1H, s), 8.27 (1H, s). MS (ESI ⁺) m/z: 394 (MH ⁺). [α] _D ²⁶ = -188 (c 0.12, EtOH)
1-70		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(2,3-dihydro-1H-inden-5-yl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.90-1.99 (2H, m), 2.74 (4H, q, J = 7.7 Hz), 3.26-3.33 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.72-3.80 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.37 (1H, td, J = 8.6 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.97-7.03 (2H, m), 7.25 (1H, s), 8.04 (1H, s), 8.45 (1H, s). MS (ESI ⁺) m/z: 402 (MH ⁺). [α] _D ²⁶ = -186 (c 0.26, EtOH)




[0627]

[Table 70]

Ex.No.	Str.	Chemical name	P.D.
1-71		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-hydroxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.33 (1H, m), 3.43 (1H, t, J = 9.5 Hz), 3.73-3.81 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.36 (1H, d, J = 8.6 Hz), 6.58-6.62 (1H, m), 6.73 (2H, d, J = 11.0 Hz), 6.90 (1H, dd, J = 11.0, 9.2 Hz), 7.12 (1H, dd, J = 7.9, 2.4 Hz), 8.05 (1H, s), 8.54 (1H, s), 9.65 (1H, s). MS (ESI ⁺) m/z: 396 (MH ⁺). [α] _D ²⁶ = -188 (c 0.12, EtOH)
1-72		(-)-1-(4-chloro-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.33 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.76-3.82 (1H, m), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.42 (1H, d, J = 8.6 Hz), 6.66 (1H, dd, J = 8.6, 2.4 Hz), 6.73 (2H, d, J = 11.0 Hz), 7.07 (1H, d, J = 8.6 Hz), 7.21 (1H, d, J = 2.4 Hz), 8.07 (1H, s), 8.67 (1H, s), 9.95 (1H, s). MS (ESI ⁺) m/z: 412 (MH ⁺). [α] _D ²⁶ = -189 (c 0.18, EtOH)
1-73		1-(4-cyano-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.33 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.76-3.84 (1H, m), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.61 (1H, d, J = 7.9 Hz), 6.71-6.76 (3H, m), 7.29 (1H, s), 7.35 (1H, d, J = 8.6 Hz), 8.09 (1H, s), 9.08 (1H, s), 10.81 (1H, brs). MS (ESI ⁺) m/z: 403 (MH ⁺).




[0628]

[Table 71]

Ex.No.	Str.	Chemical name	P.D.
1-74		1-{{(3S*,4R*)-4-[4-(difluoroethoxy)-2,6-difluorophenyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.34 (1H, m), 3.46-3.50 (1H, m), 3.86 (1H, q, J = 9.7 Hz), 4.57 (1H, dd, J = 10.9, 7.9 Hz), 6.48 (1H, d, J = 8.5 Hz), 6.99-7.08 (4H, m), 7.30-7.33 (2H, m), 7.32 (1H, t, J = 73.3 Hz), 8.09 (1H, br), 8.70 (1H, s). MS (ESI ⁺) m/z: 416 (MH ⁺).
1-75		(-)-1-[(3S*,4R*)-4-(4-ethoxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, CDCl ₃) δ 1.29 (3H, t, J = 7.0 Hz), 3.25-3.34 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.75-3.82 (1H, m), 4.02 (2H, q, J = 7.0 Hz), 4.54 (1H, dd, J = 10.4, 8.3 Hz), 6.68-6.75 (1H, m), 6.71 (2H, d, J = 11.0 Hz), 7.01 (2H, t, J = 9.0 Hz), 7.34 (2H, dd, J = 9.0, 4.9 Hz), 8.03 (1H, s), 8.89 (1H, s) MS (ESI ⁺) m/z: 394 (MH ⁺). [α] _D ²⁶ = -149 (c 0.13, EtOH)
1-76		1-{{(3S*,4R*)-4-[4-(dimethylamino)-2,6-difluorophenyl]-2-oxopyrrolidin-3-yl}-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.87 (6H, s), 3.27 (1H, t, J = 9.8 Hz), 3.39 (1H, t, J = 8.9 Hz), 3.71 (1H, q, J = 10.0 Hz), 4.54 (1H, dd, J = 11.0, 8.6 Hz), 6.34 (2H, d, J = 12.2 Hz), 6.52 (1H, d, J = 9.2 Hz), 6.98-7.04 (2H, m), 7.31-7.37 (2H, m), 7.99 (1H, s), 8.71 (1H, s). MS (ESI ⁺) m/z: 393 (MH ⁺).




[0629]

[Table 72]

Ex.No.	Str.	Chemical name	P.D.
1-77		(-)-1-[(3S*,4R*)-4-(6-fluorobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.21-3.26 (1H, m), 3.51-3.55 (1H, m), 3.80-3.87 (1H, m), 4.65 (1H, dd, J = 11.3, 8.8 Hz), 6.77 (1H, d, J = 8.8 Hz), 6.94 (1H, d, J = 2.4 Hz), 7.00 (2H, t, J = 9.0 Hz), 7.34 (2H, dd, J = 9.0, 4.9 Hz), 7.54 (1H, d, J = 11.0 Hz), 7.80 (1H, d, J = 7.3 Hz), 7.97 (1H, s), 7.99 (1H, d, J = 2.4 Hz), 8.87 (1H, s). MS (ESI ⁺) m/z: 372 (MH ⁺). [α] _D ²⁵ = -201 (c 0.10, EtOH)
1-78		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(6-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.17-3.24 (1H, m), 3.42-3.52 (2H, m), 3.80 (3H, s), 4.47-4.52 (1H, m), 6.60 (1H, brs), 6.79 (1H, d, J = 8.6 Hz), 7.02 (2H, dd, J = 9.2, 8.6 Hz), 7.35 (2H, dd, J = 9.2, 4.9 Hz), 7.80 (1H, dd, J = 8.6, 2.4 Hz), 7.95 (1H, s), 8.10 (1H, d, J = 2.4 Hz), 8.75 (1H, brs). MS (ESI ⁺) m/z: 345 (MH ⁺). [α] _D ²⁶ = -103 (c 0.12, EtOH)
1-79		(-)-N-(3,5-difluoro-4-{(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl}phenyl)acetamide	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.03 (3H, s), 3.28-3.35 (1H, m), 3.42-3.48 (1H, m), 3.81 (1H, q, J = 9.8 Hz), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.97-7.04 (2H, m), 7.25-7.34 (4H, m), 8.06 (1H, s), 8.67 (1H, s), 10.26 (1H, s). MS (ESI ⁺) m/z: 407 (MH ⁺). [α] _D ²⁴ = -31 (c 0.12, EtOH)




[0630]

[Table 73]

Ex.No.	Str.	Chemical name	P.D.
1-80		1-[(3S*,4R*)-4-(4,6-difluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18 (2H, t, J = 8.6 Hz), 3.28 (1H, t, J = 9.5 Hz), 3.42 (1H, t, J = 9.2 Hz), 3.79 (1H, q, J = 9.8 Hz), 4.55 (1H, dd, J = 10.4, 8.6 Hz), 4.62 (2H, t, J = 8.6 Hz), 6.48 (1H, d, J = 8.6 Hz), 6.62 (1H, d, J = 11.0 Hz), 6.99-7.05 (2H, m), 7.31-7.37 (2H, m), 8.03 (1H, s), 8.68 (1H, s). MS (ESI ⁺) m/z: 392 (MH ⁺).
1-81		1-[(3S*,4R*)-4-(6-fluoro-1-hydroxy-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.74-1.81 (1H, m), 2.28-2.36 (1H, m), 2.62-2.69 (1H, m), 2.79-2.86 (1H, m), 3.18 (1H, t, J = 9.5 Hz), 3.46 (1H, t, J = 8.9 Hz), 3.74 (1H, q, J = 10.0 Hz), 4.57 (1H, dd, J = 11.3, 8.9 Hz), 4.98 (1H, q, J = 6.3 Hz), 5.28 (1H, t, J = 5.2 Hz), 6.66 (1H, d, J = 8.6 Hz), 6.99-7.05 (3H, m), 7.33-7.38 (3H, m), 7.95 (1H, s), 8.79 (1H, s). MS (ESI ⁺) m/z: 388 (MH ⁺).
1-82		1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.96-2.04 (2H, m), 2.80 (4H, q, J = 8.2 Hz), 3.18 (1H, t, J = 9.8 Hz), 3.45 (1H, d, J = 8.9 Hz), 3.72 (1H, q, J = 9.8 Hz), 4.56 (1H, dd, J = 11.3, 8.9 Hz), 6.54 (1H, d, J = 7.9 Hz), 6.99-7.05 (3H, m), 7.32-7.37 (3H, m), 7.94 (1H, s), 8.69 (1H, s). MS (ESI ⁺) m/z: 372 (MH ⁺).



[0631]

[Table 74]

Ex.No.	Str.	Chemical name	P.D.
1-83		(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.19 (3H, s), 3.11 (2H, t, J = 8.6 Hz), 3.15 (1H, t, J = 9.5 Hz), 3.42 (1H, t, J = 8.9 Hz), 3.61-3.68 (1H, m), 4.50-4.56 (3H, m), 6.46 (1H, brs), 6.62 (1H, d, J = 11.0 Hz), 6.99 (2H, d, J = 8.6 Hz), 7.22 (2H, d, J = 8.6 Hz), 7.37 (1H, d, J = 7.3 Hz), 7.92 (1H, s), 8.52 (1H, brs). MS (ESI ⁺) m/z: 370 (MH ⁺). [α] _D ²⁸ = -158 (c 0.20, EtOH)
1-84		(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.04-3.21 (3H, m), 3.43 (1H, t, J = 8.9 Hz), 3.66-3.73 (1H, m), 4.51-4.56 (3H, m), 6.63 (1H, d, J = 11.6 Hz), 7.29 (1H, brs), 7.36 (1H, d, J = 7.9 Hz), 7.55 (2H, d, J = 8.6 Hz), 7.62 (2H, d, J = 8.6 Hz), 7.94 (1H, s), 9.70 (1H, brs). MS (ESI ⁺) m/z: 381 (MH ⁺). [α] _D ²⁷ = -138 (c 0.12, EtOH)
1-85		(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.12 (2H, t, J = 8.6 Hz), 3.18 (1H, t, J = 9.8 Hz), 3.44 (1H, t, J = 9.8 Hz), 3.69-3.76 (1H, m), 4.52-4.56 (3H, m), 6.62-6.64 (2H, m), 7.01 (1H, brs), 7.07 (1H, dd, J = 7.9, 7.4 Hz), 7.20 (1H, dd, J = 7.9, 7.4 Hz), 7.39 (1H, d, J = 7.9 Hz), 7.50 (1H, d, J = 7.9 Hz), 7.68 (1H, d, J = 7.9 Hz), 7.95 (1H, s), 10.21 (1H, brs). MS (ESI ⁺) m/z: 412 (MH ⁺). [α] _D ²⁹ = -128 (c 0.14, EtOH)

[0632]

[Table 75]


Ex.No.	Str.	Chemical name	P.D.
1-86		(+)-1-[(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.18 (3H, s), 3.29-3.49 (2H, m), 3.71-3.78 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.41 (1H, brd, J = 8.6 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.98 (2H, d, J = 8.6 Hz), 7.20 (2H, d, J = 8.6 Hz), 8.04 (1H, s), 8.51 (1H, s). MS (ESI ⁺) m/z: 376 (MH ⁺). [α] _D ²⁵ +108 (c 0.13, EtOH)
1-87		(+)-1-[(3R*,4S*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.30 (1H, t, J = 9.8 Hz), 3.45 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.76-3.82 (m, 1H), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.86 (1H, t, J = 7.3 Hz), 7.18 (2H, dd, J = 8.6, 7.3 Hz), 7.32 (2H, d, J = 8.6 Hz), 8.06 (1H, s), 8.62 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁴ = +153 (c 0.10, EtOH)

[0633]

5 <Example 2-1>

[0634]

[Chemical Formula 123]



[0635]

(+)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea

(-)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea

The same method as in Example 1-1 was performed using (±)-trans-4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidine-3-carboxylic acid in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea.

The obtained

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-5,5-dimethyl-2-oxopyrrolidin-3-yl]urea was subjected to optical resolution by high performance liquid chromatography (methyl tert-butyl ether : ethanol = 4 : 1, flow rate: 15.0 mL) using a column for separation of enantiomers (Daicel Corporation, CHIRALPAK ID). An isomer A(+) with a retention time of 6.94 minutes and an isomer B(-) with a retention time of 16.14 minutes that were the title compounds were each obtained as a white solid.

Isomer A(+):

¹H-NMR (400 MHz, DMSO-d₆) δ 0.78 (3H, s), 1.22 (3H, s), 3.26 (1H, d, J = 12.1 Hz), 3.72 (3H, s), 4.88 (1H, dd, J = 12.1, 9.1 Hz), 6.31 (1H, d, J = 9.1 Hz), 6.89 (2H, d, J = 8.5 Hz), 7.02 (2H, t, J = 9.1 Hz), 7.25 (2H, d, J = 8.5 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.01 (1H, s), 8.48 (1H, s).

MS (ESI⁺) m/z: 372 (MH⁺).

[α]_D²⁵ = +133 (c 0.21, CHCl₃)

Isomer B(-):

¹H-NMR (400 MHz, DMSO-d₆) δ 0.78 (3H, s), 1.22 (3H, s), 3.26 (1H, d, J = 12.1 Hz), 3.72 (3H, s), 4.88 (1H, dd, J = 12.1, 9.1 Hz), 6.31 (1H, d, J = 9.1 Hz), 6.89 (2H, d, J = 8.5 Hz), 7.02 (2H, t, J = 9.1 Hz), 7.25 (2H, d, J = 8.5 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.01 (1H, s), 8.48 (1H, s).

MS (ESI⁺) m/z: 372 (MH⁺).



[α]_D²⁵ = -147 (c 0.21, CHCl₃)

[0636]

<Example 3-1>

[0637]

[Chemical Formula 124]



[0638]

(+)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea

(-)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea

5 The same method as in Example 1-1 was performed using

(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonane-3-carboxylic acid in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain

10 (±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea. The obtained

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.4]nonan-3-yl]urea was subjected to optical resolution by high performance liquid chromatography (methyl tert-butyl

15 ether : ethanol = 4 : 1, flow rate: 20.0 mL) using a column for separation of enantiomers (CHIRALPAK ID). An isomer A(+) with a retention time of 9.02 minutes and an isomer B(-) with a retention time of 14.27 minutes that were the title compounds were each obtained as a white solid.

20 Isomer A(+):

¹H-NMR (400 MHz, DMSO-d₆) δ 1.05-1.81 (8H, m), 3.45 (1H, d, J = 12.1 Hz), 3.71 (3H, s), 4.82 (1H, dd, J = 12.1, 9.1 Hz), 6.31 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 9.1 Hz), 7.02 (2H, t, J = 9.1 Hz), 7.25 (2H, d, J = 9.1 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.28 (1H, s), 8.47 (1H, s).

25

MS (ESI⁺) m/z: 398 (MH⁺).

HRMS (ESI⁺) for C₂₂H₂₅FN₃O₃ (MH⁺): calcd, 398.18799; found, 398.18738.

[α]_D²⁴ = +124 (c 0.31, CHCl₃).

Isomer B(-):

5 ¹H-NMR (400 MHz, DMSO-d₆) δ 1.05-1.81 (8H, m), 3.45 (1H, d, J = 12.1 Hz), 3.71 (3H, s), 4.82 (1H, dd, J = 12.1, 9.1 Hz), 6.31 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 9.1 Hz), 7.02 (2H, t, J = 9.1 Hz), 7.25 (2H, d, J = 9.1 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.28 (1H, s), 8.47 (1H, s).

10 MS (ESI⁺) m/z: 398 (MH⁺).

HRMS (ESI⁺) for C₂₂H₂₅FN₃O₃ (MH⁺): calcd, 398.18799; found, 398.18880.


[α]_D²⁴ = -125 (c 0.31, CHCl₃).

[0639]

<Example 4-1>

15 [0640]

[Chemical Formula 125]



[0641]

(+)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea

20 (-)-trans-1-(4-Fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea

The same method as in Example 1-1 was performed using
(±)-trans-4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decane-3-carboxylic acid in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea. The obtained

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxo-1-azaspiro[4.5]decan-3-yl]urea was subjected to optical resolution

by high performance liquid chromatography (methyl tert-butyl ether : ethanol = 85 : 15, flow rate: 15.0 mL) using a column for separation of enantiomers (CHIRALPAK ID). An isomer A(+) with a retention time of 12.16 minutes and an isomer B(-) with a retention time of 17.07 minutes that were the title compounds were each obtained as a white solid.

Isomer A(+):

¹H-NMR (400 MHz, DMSO-d₆) δ 0.55-0.67 (1H, m), 0.76-0.92 (1H, m), 1.29-1.71 (8H, m), 3.14 (1H, d, J = 12.2 Hz), 3.72 (3H, s), 4.89 (1H, dd, J = 12.2, 8.6 Hz), 6.31 (1H, d, J = 8.6 Hz), 6.88 (2H, d, J = 8.6 Hz), 7.02 (2H, t, J = 8.8 Hz), 7.25 (2H, d, J = 8.5 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.01 (1H, s), 8.48 (1H, s).

MS (ESI⁺) m/z: 412 (MH⁺).

HRMS (ESI⁺) for C₂₃H₂₇N₃O₃ (MH⁺): calcd, 412.20364; found, 412.20364.

[α]_D²⁵ = +133 (c 0.32, CHCl₃).

Isomer B(-):

¹H-NMR (400 MHz, DMSO-d₆) δ 0.55-0.67 (1H, m), 0.76-0.92 (1H, m), 1.29-1.71 (8H, m), 3.14 (1H, d, J = 12.2 Hz), 3.72 (3H, s), 4.89 (1H, dd, J = 12.2, 8.6 Hz), 6.31 (1H, d, J = 8.6 Hz), 6.88 (2H, d, J = 8.6 Hz), 7.02 (2H, t, J = 8.8 Hz), 7.25 (2H, d, J = 8.5 Hz), 7.34 (2H, dd, J = 9.1, 4.8 Hz), 8.01 (1H, s), 8.48 (1H, s).

MS (ESI⁺) m/z: 412 (MH⁺).

HRMS (ESI⁺) for C₂₃H₂₇FN₃O₃ (MH⁺): calcd, 412.20364; found, 412.20416.


[α]_D²⁵ = -137 (c 0.31, CHCl₃).

[0642]

10 <Example 5-1>

[0643]

[Chemical Formula 126]



[0644]

(-)-1-(4-Fluorophenyl)-3-[(3S*,4R*)-4-(4-methylsulfinylphenyl)-2-oxopyrrolidin-3-yl]urea

Meta-chloroperoxybenzoic acid (mCPBA) (70 mg) was added to a solution of 1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea (100 mg) in dichloromethane (5 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 1 hour. A saturated sodium hydrogen carbonate

solution was added to the reaction solution, and the mixture was extracted with a dichloromethane-methanol mixed solution. The extract was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : methanol = 5 : 1), to obtain the title compound as a white solid (85 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 2.71 (3H, s), 3.23 (1H, t, $J = 9.4$ Hz), 3.50-3.68 (2H, m), 4.55 (1H, dd, $J = 11.5, 8.5$ Hz), 6.47 (1H, d, $J = 8.5$ Hz), 7.00-7.05 (2H, m), 7.32-7.37 (2H, m), 7.58-7.64 (4H, m), 7.97 (1H, s), 8.59 (1H, s).

MS (ESI^+) m/z : 376 (MH^+).


$[\alpha]_{\text{D}}^{26} = -168$ (c 0.14, EtOH).

[0645]

<Example 6-1>

[0646]

[Chemical Formula 127]



[0647]

(-)-1-(4-Fluorophenyl)-3-[(3S*,4R*)-4-(4-methylsulfonylphenyl)-2-oxopyrrolidin-3-yl]urea

Meta-chloroperoxybenzoic acid (mCPBA) (140 mg) was added

to a solution of

1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea (84 mg) in dichloromethane (5 mL) to produce a reaction solution. The reaction solution was stirred at room
5 temperature for 1 hour. A saturated sodium hydrogen carbonate solution was added to the reaction solution, and the mixture was extracted with a dichloromethane-methanol mixed solution. The extract was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under
10 reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : methanol = 5 : 1), to obtain the title compound as a white solid (67 mg).

¹H-NMR (400 MHz, DMSO- d₆) δ 3.19 (3H, s), 3.25 (1H, t, J = 9.7 Hz), 3.55 (1H, t, J = 8.5 Hz), 3.64-3.72 (1H, m), 4.57 (1H, dd, J = 10.9, 8.5 Hz), 6.49 (1H, d, J = 8.5 Hz), 7.00-7.05 (2H, m), 7.32-7.36 (2H, m), 7.68 (2H, d, J = 8.5 Hz), 7.87 (2H, d, J = 8.5 Hz), 8.00 (1H, s), 8.61 (1H, s).

MS (ESI⁺) m/z: 392 (MH⁺).


[α]_D²⁶ = -189 (c 0.14, EtOH).

[0648]

<Example 7-1>

[0649]

[Chemical Formula 128]



[0650]

(-)-1-[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethoxy)phenyl]urea

4-(Trifluoromethoxy)phenyl isocyanate (36 μ L) was added to
5 a solution of

(-)-(3S*, 4R*)-3-amino-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-2-one (58 mg) in tetrahydrofuran (2.4 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 30 minutes. The reaction solution was concentrated, and the
10 residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 9 : 1 followed by ethyl acetate), to obtain the title compound as a white solid (94 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.27-3.31 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.74 (3H, s), 3.76-3.84 (1H, m), 4.57 (1H, dd, J = 10.4, 8.6
15 Hz), 6.56 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 7.18 (2H, d, J = 8.6 Hz), 7.14 (2H, d, J = 8.6 Hz), 8.06 (1H, s), 8.89 (1H, s).

MS (ESI $^+$) m/z : 446 (MH^+).

$[\alpha]_D^{27}$ = -150 (c 0.10, EtOH).

20 [0651]




The same method as in Example 7-1 was performed using a corresponding isocyanate to obtain the following Examples 7-2 to

7-14.

The structures and spectral data thereof are shown in Tables 76 to 80.




[0652]

[Table 76]

Ex.No	Str.	Chemical name	P.D.
7-2		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.30 (1H, t, J = 9.8 Hz), 3.45 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.76-3.82 (m, 1H), 4.57 (1H, dd, J = 11.0, 8.6 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.86 (1H, t, J = 7.3 Hz), 7.18 (2H, dd, J = 8.6, 7.3 Hz), 7.32 (2H, d, J = 8.6 Hz), 8.06 (1H, s), 8.62 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). $[\alpha]_D^{27} = -162$ (c 0.10, EtOH)
7-3		(-)-1-[4-(tert-butyl)phenyl]-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.21 (9H, s), 3.27-3.48 (2H, m), 3.71-3.80 (1H, m), 3.74 (3H, s), 4.56 (1H, dd, J = 11.0, 8.0 Hz), 6.39 (1H, d, J = 8.0 Hz), 6.73 (2H, d, J = 11.0 Hz), 7.17-7.23 (4H, m), 8.04 (1H, s), 8.51 (1H, s). MS (ESI ⁺) m/z: 418 (MH ⁺). $[\alpha]_D^{27} = -117$ (c 0.11, EtOH)
7-4		(-)-4-{[(3S*,4R*)-3-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido]benzoic acid ethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.28 (3H, t, J = 6.8 Hz), 3.28-3.39 (1H, m), 3.45 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.78-3.86 (1H, m), 4.23 (2H, q, J = 6.8 Hz), 4.57 (1H, dd, J = 10.4, 8.6 Hz), 6.66 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 7.46 (2H, d, J = 8.6 Hz), 7.79 (2H, d, J = 8.6 Hz), 8.08 (1H, s), 9.13 (1H, s). MS (ESI ⁺) m/z: 434 (MH ⁺). $[\alpha]_D^{27} = -161$ (c 0.10, EtOH)



[0653]

[Table 77]

Ex.No	Str.	Chemical name	P.D.
7-5		(-)-1-[(1,1'-biphenyl)-4-yl]-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.34 (1H, m), 3.46 (1H, t, J = 8.6 Hz), 3.75 (3H, s), 3.81 (1H, q, J = 8.6 Hz), 4.59 (1H, dd, J = 8.6, 8.6 Hz), 6.51 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 7.28 (1H, t, J = 7.3 Hz), 7.38-7.43 (4H, m), 7.50 (2H, d, J = 9.2 Hz), 7.58 (2H, d, J = 8.0 Hz), 8.07 (1H, s), 8.75 (1H, s). MS (ESI ⁺) m/z: 438 (MH ⁺). [α] _D ²⁷ = -168 (c 0.10, EtOH)
7-6		(-)-1-(4-acetylphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.45 (3H, s), 3.26-3.31 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.74 (3H, s), 3.81 (1H, q, J = 9.2 Hz), 4.57 (1H, dd, J = 9.2, 8.6 Hz), 6.65 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 10.4 Hz), 7.45 (2H, d, J = 8.6 Hz), 7.80 (2H, d, J = 8.6 Hz), 8.07 (1H, s), 9.11 (1H, s). MS (ESI ⁺) m/z: 404 (MH ⁺). [α] _D ²⁷ = -153 (c 0.10, EtOH)
7-7		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-phenoxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.30 (1H, t, J = 9.2 Hz), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.79 (1H, q, J = 9.2 Hz), 4.57 (1H, t, J = 8.6 Hz), 6.47 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.88-6.91 (4H, m), 7.05 (1H, t, J = 7.3 Hz), 7.30-7.35 (4H, m), 8.05 (1H, s), 8.67 (1H, s). MS (ESI ⁺) m/z: 454 (MH ⁺). [α] _D ²⁷ = -163 (c 0.10, EtOH)

[0654]

[Table 78]

Ex.No	Str.	Chemical name	P.D.
7-8		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.28-3.33 (1H, m), 3.40-3.49 (1H, m), 3.75 (3H, s), 3.81 (1H, q, J = 9.5 Hz), 4.56 (1H, dd, J = 11.0, 8.0 Hz), 6.57 (1H, d, J = 8.0 Hz), 6.64-6.70 (1H, m), 6.74 (2H, d, J = 10.4 Hz), 7.00 (1H, d, J = 8.6 Hz), 7.17-7.23 (1H, m), 7.32-7.37 (1H, m), 8.07 (1H, s), 8.90 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). $[\alpha]_D^{27} = -152$ (c 0.10, EtOH)
7-9		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3,4-difluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.36 (1H, m), 3.40-3.48 (1H, m), 3.75 (3H, s), 3.81 (1H, q, J = 9.2 Hz), 4.56 (1H, dd, J = 9.2, 8.6 Hz), 6.60 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 11.0 Hz), 6.97-7.03 (1H, m), 7.20-7.27 (1H, m), 7.49-7.56 (1H, m), 8.06 (1H, s), 8.91 (1H, s). MS (ESI ⁺) m/z: 398 (MH ⁺). $[\alpha]_D^{27} = -138$ (c 0.10, EtOH)



[0655]

[Table 79]

Ex.No	Str.	Chemical name	P.D.
7-10		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-ethylphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.11 (3H, t, J = 7.4 Hz), 2.49 (2H, q, J = 7.4 Hz), 3.30 (1H, t, J = 9.2 Hz), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.77 (1H, q, J = 9.2 Hz), 4.56 (1H, dd, J = 10.4, 8.6 Hz), 6.42 (1H, d, J = 8.6 Hz), 6.73 (2H, d, J = 10.4 Hz), 7.01 (2H, d, J = 8.6 Hz), 7.22 (2H, d, J = 8.6 Hz), 8.05 (1H, s), 8.53 (1H, s). MS (ESI ⁺) m/z: 390 (MH ⁺). $[\alpha]_D^{27} = -134$ (c 0.10, EtOH)
7-11		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(m-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.20 (3H, s), 3.27-3.33 (1H, m), 3.44 (1H, t, J = 9.5 Hz), 3.75 (3H, s), 3.76-3.82 (1H, m), 4.56 (1H, dd, J = 10.7, 8.3 Hz), 6.43 (1H, d, J = 7.9 Hz), 6.69 (1H, d, J = 6.7 Hz), 6.73 (2H, d, J = 11.0 Hz), 7.02-7.10 (2H, m), 7.18 (1H, s), 8.05 (1H, s), 8.53 (1H, s). MS (ESI ⁺) m/z: 376 (MH ⁺). $[\alpha]_D^{25} = -135$ (c 0.22, EtOH)
7-12		(-)-1-(3-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.35 (1H, m), 3.40-3.50 (1H, m), 3.75 (3H, s), 3.78-3.87 (1H, m), 4.57 (1H, dd, J = 10.7, 8.3 Hz), 6.74 (2H, d, J = 11.0 Hz), 7.30-7.33 (1H, m), 7.40 (1H, t, J = 7.9 Hz), 7.52-7.57 (1H, m), 7.85 (1H, s), 8.07 (1H, s), 9.08 (1H, s). MS (ESI ⁺) m/z: 387 (MH ⁺). $[\alpha]_D^{25} = -86$ (c 0.13, EtOH)

[0656]

[Table 80]


Ex.No	Str.	Chemical name	P.D.
7-13		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-methoxyphenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.33 (1H, m), 3.44 (1H, t, J = 9.5 Hz), 3.67 (3H, s), 3.75 (3H, s), 3.80 (1H, t, J = 9.8 Hz), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.43-6.47 (2H, m), 6.74 (2H, d, J = 11.0 Hz), 6.80-6.83 (1H, m), 7.04-7.10 (2H, m), 8.05 (1H, s), 8.63 (1H, s). MS (ESI ⁺) m/z: 392 (MH ⁺). [α] _D ²⁶ = -173 (c 0.14, EtOH)
7-14		(-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.18 (3H, s), 3.38 (1H, t, J = 9.5 Hz), 3.47 (1H, t, J = 9.2 Hz), 3.83 (3H, s), 3.90-4.01 (1H, m), 4.50 (1H, dd, J = 10.1, 8.3 Hz), 6.51 (1H, d, J = 7.3 Hz), 6.98 (2H, J = 8.3 Hz), 7.20 (2H, d, J = 8.3 Hz), 7.38 (1H, dd, J = 12.2, 2.4 Hz), 7.93 (1H, s), 8.20 (1H, d, J = 2.4 Hz), 8.53 (1H, s). MS (ESI ⁺) m/z: 359 (MH ⁺). [α] _D ²⁶ = -177 (c 0.10, EtOH)

[0657]

5 <Example 8-1>

[0658]

[Chemical Formula 129]



[0659]

(-)-1-(5-Chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea

Triethylamine (51 μ L) and diphenylphosphoryl azide (83 μ L) were added to a solution of 5-chlorothiophene-2-carboxylic acid (60 mg) in toluene (3.7 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. The reaction temperature was heated up to 100°C, and the reaction solution was then stirred for 1 hour.

(-)-(3S*,4R*)-3-amino-4-(2,6-difluoro-4-methoxyphenyl)pyrrolidin-2-one (71 mg) was added and the reaction solution was stirred at 100°C for 1 hour. The reaction solution was concentrated, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title compound as a white solid (45 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.27-3.31 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.80-3.88 (1H, m), 4.52 (1H, dd, J = 11.0, 9.2 Hz), 6.20 (1H, d, J = 4.3 Hz), 6.71-6.81 (4H, m), 8.06 (1H, s), 9.94 (1H, s).

MS (ESI $^+$) m/z : 402 (MH^+).



$[\alpha]_D^{27}$ = -135 (c 0.11, EtOH)

[0660]

The same method as in Example 8-1 was performed using a corresponding carboxylic acid substance and amine to obtain the following Examples 8-2 to 8-3. The structures and spectral data
5 thereof are shown in Table 81.

[0661]

[Table 81]


Ex.No	Str.	Chemical name	P.D.
8-2		(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.30-3.37 (1H, m), 3.46 (1H, t, J = 8.9 Hz), 3.83 (3H, s), 3.96-4.03 (1H, m), 4.48 (1H, d, J = 10.4 Hz), 6.05-6.15 (1H, m), 6.66 (1H, d, J = 4.3 Hz), 7.00-7.30 (1H, m), 7.38 (1H, dd, J = 11.6, 2.4 Hz), 7.89 (1H, s), 8.20 (1H, d, J = 2.4 Hz), 10.3-10.6 (1H, m). MS (ESI ⁺) m/z: 385 (MH ⁺). [α] _D ²⁶ = -166 (c 0.10, EtOH)
8-3		(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.39 (1H, t, J = 9.2 Hz), 3.49 (1H, t, J = 9.2 Hz), 3.83 (3H, s), 4.00-4.07 (1H, m), 4.57 (1H, t, J = 9.2 Hz), 6.65 (1H, s), 6.98-7.06 (1H, m), 7.08 (1H, t, J = 7.6 Hz), 7.21 (1H, t, J = 7.6 Hz), 7.38-7.42 (1H, m), 7.52 (1H, d, J = 7.9 Hz), 7.69 (1H, d, J = 7.9 Hz), 7.98 (1H, s), 8.20-8.23 (1H, m), 10.19 (1H, s). MS (ESI ⁺) m/z: 401 (MH ⁺). [α] _D ²⁷ = -157 (c 0.11, EtOH)

[0662]

<Example 9-1>

[0663]

[Chemical Formula 130]



[0664]

- 5 (-)-1-[(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea

The same method as in Example 8-1 was performed using 4-methyl benzoic acid in place of 5-chlorothiophene-2-carboxylic acid to obtain the title compound.

- 10 $^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 2.18 (3H, s), 3.29-3.49 (2H, m), 3.71-3.78 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, $J = 11.0, 8.6$ Hz), 6.41 (1H, brd, $J = 8.6$ Hz), 6.73 (2H, d, $J = 11.0$ Hz), 6.98 (2H, d, $J = 8.6$ Hz), 7.20 (2H, d, $J = 8.6$ Hz), 8.04 (1H, s), 8.51 (1H, s).

MS (ESI^+) m/z : 376 (MH^+).


- 15 $[\alpha]_{\text{D}}^{27} = -144$ (c 0.10, EtOH)

[0665]

<Example 10-1>

[0666]

[Chemical Formula 131]



[0667]

(-)-1-[(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(hydroxymethyl)phenyl]urea

Diisobutylaluminum hydride (1.0 mL, 1.04 mol/L hexane solution) was added to a solution of (-)-4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}benzoic acid ethyl ester (87 mg) in dichloromethane (2.5 mL) under ice-cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 2.5 hours. A 10% potassium sodium tartrate aqueous solution and ethyl acetate were added to the reaction solution, and the mixture was stirred for 30 minutes and extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : methanol = 50 : 1 followed by ethyl acetate : methanol = 4 : 1), to obtain the title compound as a white solid (66 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.30 (1H, dd, J = 9.8, 9.2 Hz), 3.44 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.75-3.81 (1H, m), 4.36 (2H, d, J = 5.5 Hz), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 4.99 (1H, t, J = 5.5 Hz), 6.43 (1H, d, J = 7.9 Hz), 6.73 (2H, d, J = 11.0 H), 7.12

(2H, d, J = 8.6 Hz), 7.27 (2H, d, J = 8.6 Hz), 8.05 (1H, s), 8.58 (1H, s).

MS (ESI⁺) m/z: 392 (MH⁺).


$[\alpha]_D^{27} = -159$ (c 0.10, EtOH)

5 [0668]

<Example 11-1>

[0669]

[Chemical Formula 132]



[0670]

10 (-)-4-{3-[(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}-N-methylbenzamide

A 2 mol/L sodium hydroxide aqueous solution (0.28 mL) was added to a solution of

(-)-4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}benzoic acid ethyl ester (100 mg) in methanol (2.4 mL) to produce a reaction solution. The reaction solution was stirred at 60°C for 9 hours. The reaction solution was concentrated under reduced pressure, a 1 mol/L hydrochloric acid was added to the residue under ice-cooling to make the mixture acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure

15

20

to obtain

4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}benzoic acid as an intermediate compound.

2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluroniu
5 m hexafluorophosphate methanaminium (HATU) (92 mg) was added to
a solution of the obtained

4-{3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]ureido}benzoic acid (45 mg), methylamine (45 μ L, 12 M/L
aqueous solution), and diisopropyl ethyl amine (DIEA) (40 μ L) in

10 N,N-dimethylformamide (1.1 mL) to produce a reaction solution.
The reaction solution was stirred at room temperature for 1 hour.

Water was added to the reaction solution, and the mixture was
extracted with ethyl acetate. The organic layer was washed with
a brine, and dried over anhydrous sodium sulfate. The solvent
15 was removed under reduced pressure to obtain the title compound
as a white solid (17 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 2.72 (3H, s), 3.55-3.25 (2H, m), 3.75
(3H, s), 3.76-3.86 (1H, m), 4.52-4.61 (1H, m), 6.61 (1H, brs),
6.70-6.79 (2H, m), 7.34-7.41 (2H, m), 7.63-7.71 (2H, m), 8.07 (1H,
20 s), 8.16-8.23 (1H, brs), 8.93 (1H, brs).

MS (ESI^-) m/z : 417 (M-H^-).


$[\alpha]_{\text{D}}^{28} = -159$ (c 0.10, DMSO)

[0671]

<Example 12-1>

25 [0672]

[Chemical Formula 133]



[0673]

(-)-1-[(3S*,4R*)-1-Ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

5 Trifluoroacetic acid (0.4 mL) was added to a solution of [(3S*,4R*)-1-ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester (63 mg) in dichloromethane (2 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, ethyl acetate and a saturated aqueous sodium hydrogen carbonate solution were added to the residue to adjust the pH of the aqueous layer to 9. 4-Fluorophenyl isocyanate (20 μ L) was added to the solution of two layers, and the mixture was stirred at room temperature for 15 minutes and extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 4 : 1 followed by ethyl acetate), to obtain the title compound as a white solid (48 mg).

10

15

20

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 1.21 (3H, t, $J = 7.3$ Hz), 3.31-3.45 (2H, m), 3.52-3.61 (1H, m), 3.71-3.77 (2H, m), 3.80 (3H, s), 4.24 (1H,

dd, J = 9.8, 6.7 Hz), 6.13 (1H, brs), 6.76 (2H, t, J = 8.6 Hz), 6.89 (2H, d, J = 8.6 Hz), 7.06-7.10 (2H, m), 7.24 (2H, d, J = 8.6 Hz), 7.84 (1H, brs).

MS (ESI⁺) m/z: 372 (MH⁺).

5 $[\alpha]_D^{24} = -135$ (c 0.30, EtOH).


[0674]

The same method as in Example 12-1 was performed using a corresponding Boc substance to obtain the following Examples 12-2 to 12-15.

10 The structures and spectral data thereof are shown in Tables 82 to 87.



[0675]

[Table 82]

Ex.No	Str.	Chemical name	P.D.
12-2		2-((3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid ethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.29 (3H, t, J = 7.4 Hz), 3.61 (1H, t, J = 9.2 Hz), 3.72 (1H, t, J = 9.2 Hz), 3.79 (3H, s), 3.81-3.89 (1H, m), 3.95 (1H, d, J = 17.8 Hz), 4.17-4.25 (2H, m), 4.37 (1H, d, J = 17.8 Hz), 4.39-4.43 (1H, m), 6.12 (1H, brs), 6.73 (2H, t, J = 8.6 Hz), 6.88 (2H, d, J = 8.6 Hz), 7.05-7.09 (2H, m), 7.27 (2H, d, J = 8.6 Hz), 7.80 (1H, brs). MS (ESI ⁺) m/z: 430 (MH ⁺).




[0676]

[Table 83]

Ex.No	Str.	Chemical name	P.D.
12-3		2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-2-methylpropionic acid ethyl ester	¹ H-NMR (400 MHz, CDCl ₃) δ 1.18 (3H, t, J = 7.4 Hz), 1.52 (3H, s), 1.57 (3H, s) 3.38 (1H, t, J = 9.2 Hz), 3.68 (1H, q, J = 9.2 Hz), 3.79 (3H, s), 3.83 (1H, t, J = 8.6 Hz), 4.10 (2H, q, J = 7.4 Hz), 4.80 (1H, dd, J = 11.6, 9.2 Hz), 5.87 (1H, brd, J = 8.6 Hz), 6.73 (2H, t, J = 9.2 Hz), 6.89 (2H, d, J = 8.6 Hz), 7.10 (2H, dd, J = 9.2, 4.9 Hz), 7.30 (2H, d, J = 8.6 Hz), 7.88 (1H, brs). MS (ESI ⁺) m/z: 458 (MH ⁺).
12-4		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-1-(1-hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, CDCl ₃) δ 1.39 (3H, s), 1.41 (3H, s), 3.42-3.53 (2H, m), 3.71 (1H, q, J = 9.6 Hz), 3.80 (3H, s), 3.98-4.06 (2H, m), 4.17-4.24 (2H, m), 6.49 (1H, brs), 6.70-6.75 (2H, m), 6.89 (2H, d, J = 8.6 Hz), 7.03-7.08 (2H, m), 7.21 (2H, d, J = 8.6 Hz), 7.56 (1H, s). MS (ESI ⁺) m/z: 416 (MH ⁺). $[\alpha]_D^{27} = -113$ (c 0.31, EtOH).



[0677]

[Table 84]

Ex.No	Str.	Chemical name	P.D.
12-5		(-)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13 (1H, d, J = 10.4 Hz), 3.31 (3H, s), 3.72-3.76 (1H, m), 4.18-4.23 (1H, m), 4.61 (1H, dd, J = 9.8, 6.7 Hz), 6.04 (1H, d, J = 6.7 Hz), 6.66 (2H, d, J = 11.0 Hz), 6.97-7.05 (2H, m), 7.22-7.26 (2H, m), 8.06 (1H, s), 8.65 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁵ -239 (c 0.10, EtOH)
12-6		(+)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13 (1H, d, J = 10.4 Hz), 3.31 (3H, s), 3.72-3.76 (1H, m), 4.18-4.23 (1H, m), 4.61 (1H, dd, J = 9.8, 6.7 Hz), 6.04 (1H, d, J = 6.7 Hz), 6.66 (2H, d, J = 11.0 Hz), 6.97-7.05 (2H, m), 7.22-7.26 (2H, m), 8.06 (1H, s), 8.65 (1H, s). MS (ESI ⁺) m/z: 380 (MH ⁺). [α] _D ²⁵ +188 (c 0.10, EtOH)
12-7		(-)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.17 (3H, s), 3.12 (1H, d, J = 10.4 Hz), 3.71 (3H, s), 3.72-3.76 (1H, m), 4.17-4.23 (1H, m), 4.61 (1H, dd, J = 9.8, 6.7 Hz), 6.01 (1H, d, J = 6.7 Hz), 6.66 (2H, d, J = 11.0 Hz), 6.97 (2H, d, J = 8.6 Hz), 7.11 (2H, d, J = 8.6 Hz), 8.04 (1H, s), 8.49 (1H, s). MS (ESI ⁺) m/z: 376 (MH ⁺). [α] _D ²⁶ -142 (c 0.12, EtOH)




[0678]

[Table 85]

Ex.No	Str.	Chemical name	P.D.
12-8		(+)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.17 (3H, s), 3.12 (1H, d, J = 10.4 Hz), 3.71 (3H, s), 3.72-3.76 (1H, m), 4.17-4.23 (1H, m), 4.61 (1H, dd, J = 9.8, 6.7 Hz), 6.01 (1H, d, J = 6.7 Hz), 6.66 (2H, d, J = 11.0 Hz), 6.97 (2H, d, J = 8.6 Hz), 7.11 (2H, d, J = 8.6 Hz), 8.04 (1H, s), 8.49 (1H, s). MS (ESI ⁺) m/z: 376 (MH ⁺). [α] _D ²⁶ +125 (c 0.11, EtOH)
12-9		(-)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13 (1H, d, J = 10.4 Hz), 3.71 (3H, s), 3.74 (1H, dd, J = 10.4, 8.6 Hz), 4.21 (1H, t, J = 8.0 Hz), 4.62 (1H, dd, J = 9.8, 6.7 Hz), 6.08 (1H, d, J = 6.7 Hz), 6.66 (2H, d, J = 13.8 Hz), 6.85 (1H, t, J = 7.4 Hz), 7.17 (2H, t, J = 7.3 Hz), 7.23 (2H, t, J = 7.3 Hz), 8.06 (1H, s), 8.62 (1H, s). MS (ESI ⁺) m/z: 362 (MH ⁺). [α] _D ²⁶ -201 (c 0.10, EtOH)




[0679]

[Table 86]

Ex.No	Str.	Chemical name	P.D.
12-10		(+)-1-[4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.13 (1H, d, $J = 10.4$ Hz), 3.71 (3H, s), 3.74 (1H, dd, $J = 10.4, 8.6$ Hz), 4.21 (1H, t, $J = 8.0$ Hz), 4.62 (1H, dd, $J = 9.8, 7.3$ Hz), 6.08 (1H, d, $J = 7.3$ Hz), 6.66 (2H, d, $J = 13.8$ Hz), 6.85 (1H, t, $J = 7.4$ Hz), 7.17 (2H, t, $J = 7.3$ Hz), 7.23 (2H, t, $J = 7.3$ Hz), 8.05 (1H, s), 8.62 (1H, s). MS (ESI $^+$) m/z : 362 (MH $^+$). $[\alpha]_D^{26} +242$ (c 0.10, EtOH)
12-11		(-)-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.29 (1H, t, $J = 7.4$ Hz), 3.67-3.72 (5H, m), 4.59 (1H, t, $J = 6.7$ Hz), 5.80 (1H, d, $J = 6.7$ Hz), 6.85 (2H, d, $J = 8.6$ Hz), 7.05 (2H, d, $J = 8.6$ Hz), 7.23 (2H, d, $J = 8.6$ Hz), 7.32 (2H, d, $J = 8.6$ Hz), 8.10 (1H, s), 8.74 (1H, s). MS (ESI $^+$) m/z : 360 (MH $^+$). $[\alpha]_D^{25} -260$ (c 0.10, EtOH)
12-12		(+)-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	$^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.29 (1H, t, $J = 7.4$ Hz), 3.67-3.72 (5H, m), 4.59 (1H, t, $J = 6.7$ Hz), 5.80 (1H, d, $J = 6.7$ Hz), 6.85 (2H, d, $J = 8.6$ Hz), 7.05 (2H, d, $J = 8.6$ Hz), 7.23 (2H, d, $J = 8.6$ Hz), 7.32 (2H, d, $J = 8.6$ Hz), 8.10 (1H, s), 8.74 (1H, s). MS (ESI $^+$) m/z : 360 (MH $^+$). $[\alpha]_D^{25} +235$ (c 0.10, EtOH)

[0680]

[Table 87]


Ex.No	Str.	Chemical name	P.D.
12-13	 (+)-cis	(+)-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.29 (1H, m), 3.68-3.72 (5H, m), 4.59 (1H, t, J = 6.7 Hz), 5.69 (1H, d, J = 6.7 Hz), 6.85 (2H, d, J = 8.6 Hz), 7.00-7.06 (4H, m), 7.28-7.31 (2H, m), 8.09 (1H, s), 8.59 (1H, s). MS (ESI ⁺) m/z: 344 (MH ⁺). [α] _D ²⁵ +288 (c 0.10, EtOH)
12-14	 (+)-cis	(+)-1-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.19 (3H, s), 3.26-3.29 (1H, m), 3.69-3.71 (5H, m), 4.59 (1H, t, J = 6.7 Hz), 5.67 (1H, d, J = 6.7 Hz), 6.85 (2H, d, J = 8.6 Hz), 6.99 (2H, d, J = 8.6 Hz), 7.04 (2H, d, J = 8.6 Hz), 7.17 (2H, m), 8.08 (1H, s), 8.44 (1H, s). MS (ESI ⁺) m/z: 340 (MH ⁺). [α] _D ²⁵ +177 (c 0.12, EtOH)
12-15	 (+)-cis	(+)-1-(4-fluorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.29-3.39 (2H, m), 3.46 (1H, d, J = 9.8 Hz), 3.57 (2H, q, J = 5.5 Hz), 3.66 (1H, t, J = 8.0 Hz), 3.69 (3H, s), 3.86 (1H, dd, J = 10.4, 6.7 Hz), 4.69 (1H, t, J = 8.0 Hz), 4.86 (1H, brs), 5.76 (1H, brs), 6.84 (2H, d, J = 8.6 Hz), 7.00-7.06 (4H, m), 7.29 (2H, dd, J = 9.2, 5.5 Hz), 8.65 (1H, s). MS (ESI ⁺) m/z: 388 (MH ⁺). [α] _D ²⁵ +155 (c 0.11, EtOH)

[0681]

5 <Example 13-1>

[0682]

[Chemical Formula 134]



[0683]

2- $\{(3S^*, 4R^*)$ -3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}propionic acid ethyl ester

5 Trifluoroacetic acid (3 mL) was added to a solution of 2- $\{(3S^*, 4R^*)$ -3-[(tert-butoxycarbonyl)amino]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester (651 mg) in dichloromethane (12 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 1 hour. The

10 reaction solution was concentrated under reduced pressure, and ethyl acetate and a saturated aqueous sodium hydrogen carbonate solution were added to the residue to adjust the pH of the aqueous layer to 9. 4-Fluorophenyl isocyanate (180 μ L) was added to the solution of two layers, and the mixture was stirred at room

15 temperature for 15 minutes and extracted with ethyl acetate. The organic layer was washed with water and a brine successively, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : hexane = 1 : 1 followed

20 by 1 : 3), to obtain an isomer A (298 mg) as a low polar fraction and an isomer B (276 mg) as a high polar fraction that were the

title compounds as white solids.

Isomer A:

¹H-NMR (400 MHz, CDCl₃) δ 1.26 (3H, t, J = 7.3 Hz), 1.48 (3H, d, J = 7.4 Hz), 3.35 (1H, t, J = 11.0 Hz), 3.56-3.64 (1H, m), 3.75
5 (1H, d, J = 9.2 Hz), 3.79 (3H, s), 4.13-4.21 (2H, m), 4.84 (1H, q, J = 7.4 Hz), 5.02 (1H, dd, J = 11.6, 9.2 Hz), 5.65 (1H, d, J = 9.2 Hz), 6.78 (2H, d, J = 9.2 Hz), 6.88 (2H, d, J = 9.2 Hz), 7.14 (2H, dd, J = 4.9, 9.2 Hz), 7.28 (2H, d, J = 9.2 Hz), 7.87 (1H, brs).
MS (ESI⁺) m/z: 444 (MH⁺).

10 $[\alpha]_D^{25} = -102$ (c 0.21, EtOH).

Isomer B:

¹H-NMR (400 MHz, CDCl₃) δ 1.28 (3H, t, J = 6.7 Hz), 1.52 (3H, d, J = 7.9 Hz), 3.52 (1H, t, J = 8.6 Hz), 3.80 (3H, s), 3.77-3.88 (2H, m), 4.16-4.22 (3H, m), 4.90 (1H, q, J = 7.9 Hz), 6.13 (1H, brd, J = 6.7 Hz), 6.72-6.77 (2H, m), 6.89 (2H, d, J = 8.6 Hz), 7.04-7.08
15 (2H, m), 7.26 (2H, d, J = 8.6 Hz), 7.72 (1H, brs).

MS (ESI⁺) m/z: 444 (MH⁺).


$[\alpha]_D^{25} = -94$ (c 0.21, EtOH).

[0684]

20 <Example 14-1>

[0685]

[Chemical Formula 135]



[0686]

(±)-trans-1-(4-Chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea

Trifluoroacetic acid (1.5 mL) and water (0.5 mL) were added

5 to a solution of

(±)-trans-1-{2-[(tert-butyldimethylsilyl)oxy]ethyl}-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]carbamic acid tert-butyl ester (1.2 g) in dichloromethane (2 mL) to produce a reaction solution.

The reaction solution was stirred at room temperature for 3 hours.

10 The reaction solution was concentrated under reduced pressure, a saturated aqueous sodium hydrogen carbonate solution was added to the residue to adjust the pH of the aqueous layer to 9, and an ethyl acetate-methanol (5 : 1) mixed liquid (10 mL) was added. 4-Chlorophenyl isocyanate (396 mg) was added to the solution of
15 two layers, and the mixture was stirred at room temperature for 1 hour and extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography
20 (ethyl acetate followed by ethyl acetate : methanol = 4 : 1), to obtain the title compound as a white solid (500 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.13-3.20 (1H, m), 3.35-3.45 (3H, m), 3.52 (2H, q, J = 5.7 Hz), 3.61 (1H, t, J = 6.7 Hz), 3.71 (3H, s), 4.57 (1H, t, J = 10.0 Hz), 4.75 (1H, t, J = 5.8 Hz), 6.52 (1H, d, J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz),

25

7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s).


MS (ESI⁺) m/z: 404 (MH⁺).

[0687]

<Example 15-1>

5 [0688]

[Chemical Formula 136]



[0689]

(+)-trans-1-(4-Chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea

10 (-)-trans-1-(4-Chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea

(±)-trans-1-(4-Chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea was subjected to optical resolution by high performance liquid chromatography (hexane : ethanol = 1 : 1, flow rate: 15.0 mL) using a column for separation of enantiomers (CHIRALPAK ID). An isomer A(-) with a retention time of 7.91 minutes and an isomer B(+) with a retention time of 17.78 minutes that were the title compounds were each obtained as a white solid.

20 Isomer A(-):

¹H-NMR (400 MHz, DMSO-d₆) δ 3.13-3.20 (1H, m), 3.35-3.45 (3H, m),

3.52 (2H, q, J = 5.7 Hz), 3.61 (1H, t, J = 6.7 Hz), 3.71 (3H, s),
 4.57 (1H, t, J = 10.0 Hz), 4.75 (1H, t, J = 5.8 Hz), 6.52 (1H, d,
 J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz),
 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s).

5 MS (ESI⁺) m/z: 404 (MH⁺).

$[\alpha]_D^{25} = -140$ (c 0.1, EtOH)

Isomer B(+):

¹H-NMR (400 MHz, DMSO-d₆) δ 3.13-3.20 (1H, m), 3.35-3.45 (3H, m),
 3.52 (2H, q, J = 5.7 Hz), 3.61 (1H, t, J = 6.7 Hz), 3.71 (3H, s),
 10 4.57 (1H, t, J = 10.0 Hz), 4.75 (1H, t, J = 5.8 Hz), 6.52 (1H, d,
 J = 9.1 Hz), 6.88 (2H, d, J = 8.5 Hz), 7.23 (2H, d, J = 9.1 Hz),
 7.29 (2H, d, J = 9.1 Hz), 7.38 (2H, d, J = 8.5 Hz), 8.71 (1H, s).

MS (ESI⁺) m/z: 404 (MH⁺).


$[\alpha]_D^{26} = +140$ (c 0.1, EtOH)

15 [0690]

<Example 16-1>

[0691]

[Chemical Formula 137]



[0692]

20 (-)-2-[(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid

A 2 mol/L sodium hydroxide aqueous solution (0.6 mL) was

added to a solution of

2- $\{(3S^*, 4R^*)$ -3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}acetic acid ethylester (258 mg) in methanol (3.0 mL) to produce a reaction solution. The reaction solution
5 was stirred at room temperature for 30 minutes. A 1 mol/L hydrochloric acid was added to the reaction solution to make the reaction solution acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed
10 under reduced pressure, and the crude product was washed with diisopropyl ether, to obtain the title compound as a white solid (228 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 3.34 (1H, t, $J = 9.8$ Hz), 3.44-3.51 (1H, m), 3.63 (1H, t, $J = 8.6$ Hz), 3.71 (3H, s), 3.93 (1H, d, $J = 17.7$
15 Hz), 4.05 (1H, d, $J = 17.7$ Hz), 4.51-4.56 (1H, m), 6.56 (1H, d, $J = 9.2$ Hz), 6.89 (2H, d, $J = 8.6$ Hz), 7.03 (2H, t, $J = 8.6$ Hz), 7.28 (2H, d, $J = 8.6$ Hz), 7.33-7.37 (2H, m), 8.58 (1H, s), 12.9 (1H, brs).

MS (ESI $^+$) m/z : 402 (MH^+).



20 $[\alpha]_{\text{D}}^{24} = -90$ (c 0.31, EtOH).

[0693]

The same method as in Example 16-1 was performed using a corresponding ester substance to obtain the following Examples 16-2 to 16-3. The structures and spectral data thereof are shown
25 in Table 88.

[0694]

[Table 88]


Ref.No	Str.	Chemical name	P.D.
16-2		(-)-2-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.51-3.60 (2H, m), 3.72-3.82 (1H, m), 3.75 (3H, s), 3.87 (1H, d, J = 17.0 Hz), 4.12 (1H, d, J = 17.0 Hz), 4.70 (1H, dd, J = 10.4, 8.6 Hz), 6.61 (1H, brs), 6.71-6.6.78 (2H, m), 7.01 (2H, t, J = 9.2 Hz), 7.33 (2H, dd, J = 9.2, 4.9 Hz), 8.68 (1H, s), 12.9 (1H, brs). MS (ESI ⁺) m/z: 438 (MH ⁺). [α] _D ²⁸ = -115 (c 0.08, EtOH)
16-3		(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.12 (2H, t, J = 8.6 Hz), 3.40-3.52 (1H, m), 3.59 (1H, t, J = 8.6 Hz), 3.66 (1H, t, J = 9.8 Hz), 3.90 (1H, d, J = 17.7 Hz), 4.03 (1H, d, J = 17.7 Hz), 4.55 (2H, t, J = 8.6 Hz), 4.61-4.65 (1H, m), 6.60-6.70 (1H, m), 6.65 (1H, d, J = 11.0 Hz), 7.03 (2H, t, J = 8.6 Hz), 7.32-7.37 (3H, m), 8.67 (1H, s), 12.99 (1H, brs). MS (ESI ⁺) m/z: 432 (MH ⁺). [α] _D ²⁶ = -120 (c 0.13, DMSO)

[0695]

5 <Example 17-1>

[0696]

[Chemical Formula 138]



[0697]

(-)-2-{{(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}}-N-methylacetamide

1-Hydroxybenzotriazole (46 mg), methylamine (0.5 mL, 2 mol/L, tetrahydrofuran solution), and

1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (53 mg) were added to a solution of

(-)-2-{{(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}acetic acid (103 mg) in tetrahydrofuran (1.3 mL) under cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. Water was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, the residue was purified by preparative TLC (chloroform : methanol = 9 : 1), and the crude product was washed with diisopropyl ether, to obtain the title compound as a white solid (81 mg).

¹H-NMR (400 MHz, CDCl₃) δ 2.83 (3H, d, J = 4.9 Hz), 3.50 (1H, dd, J = 9.8, 6.7 Hz), 3.57 (1H, d, J = 17.1 Hz), 3.77 (1H, t, J = 7.4

Hz), 3.80 (3H, s), 3.88-3.94 (1H, m), 4.00 (1H, t, J = 9.8 Hz), 4.64 (1H, d, J = 17.1 Hz), 6.67 (2H, t, J = 8.6 Hz), 6.89 (2H, d, J = 8.6 Hz), 6.92-6.96 (2H, m), 7.04 (1H, d, J = 6.7 Hz), 7.15 (2H, d, J = 8.6 Hz), 7.80 (1H, brs), 7.87 (1H, d, J = 4.9 Hz).

5 MS (ESI⁺) m/z: 415 (MH⁺).



$[\alpha]_D^{25} = -152$ (c 0.31, EtOH).

[0698]

The same method as in Example 17-1 was performed using a corresponding carboxylic acid substance and amine to obtain the
10 following Examples 17-2 to 17-3. The structures and spectral data thereof are shown in Table 89.

[0699]

[Table 89]


Ref.No	Str.	Chemical name	P.D.
17-2		(-)-2-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-N-methylacetamide	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.61 (3H, d, J = 4.9 Hz), 3.49-3.58 (2H, m), 3.76 (3H, s), 3.82 (1H, q, J = 10.4 Hz), 3.88 (2H, s), 4.60 (1H, dd, J = 10.4, 8.6 Hz), 6.68 (1H, brs), 6.71-6.79 (2H, m), 7.01 (2H, t, J = 9.2 Hz), 7.33 (2H, dd, J = 9.2, 4.9 Hz), 7.92 (1H, q, J = 4.9 Hz), 8.85 (1H, s). MS (ESI ⁺) m/z: 451 (MH ⁺). [α] _D ²⁷ = -156 (c 0.11, EtOH)
17-3		(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetamide	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13 (2H, t, J = 8.6 Hz), 3.27-3.39 (1H, m), 3.56 (1H, t, J = 8.6 Hz), 3.68-3.75 (1H, m), 3.78 (1H, d, J = 16.5 Hz), 3.86 (1H, d, J = 16.5 Hz), 4.55 (2H, t, J = 8.6 Hz), 4.65 (1H, dd, J = 10.7, 8.9 Hz), 6.59-6.64 (1H, m), 6.64 (1H, d, J = 11.0 Hz), 7.01-7.05 (2H, m), 7.19 (1H, s), 7.34-7.43 (4H, m), 8.77 (1H, s). MS (ESI ⁺) m/z: 431 (MH ⁺). [α] _D ²⁷ = -145 (c 0.11, EtOH)

[0700]

5 <Example 18-1>

[0701]

[Chemical Formula 139]



[0702]

(-)-2-[(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A)

5 The same method as in Example 16-1 was performed using (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester (isomer A) in place of

2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid ethyl ester to obtain the title compound as a white solid.

¹H-NMR (400 MHz, DMSO-d₆) δ 1.35 (3H, d, J = 7.4 Hz), 3.27-3.41 (2H, m), 3.61 (1H, t, J = 9.1 Hz), 3.71 (3H, s), 4.58 (1H, q, J = 7.4 Hz), 4.66 (1H, dd, J = 11.6, 9.2 Hz), 6.55 (1H, d, J = 9.2 Hz),
15 6.88 (2H, d, J = 8.6 Hz), 7.03 (2H, t, J = 8.6 Hz), 7.31-7.36 (4H, m), 8.46 (1H, s), 12.8 (1H, brs).

MS (ESI⁺) m/z: 416 (MH⁺).


[α]_D²⁵ = -103 (c 0.33, EtOH).

[0703]

20 <Example 19-1>

[0704]

[Chemical Formula 140]



[0705]

(-)-2-[(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer A)

The same method as in Example 17-1 was performed using

5 (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A) in place of (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid to obtain the title compound as a white solid.

10 ¹H-NMR (400 MHz, DMSO-d₆) δ 1.29 (3H, d, J = 7.3 Hz), 2.62 (3H, d, J = 4.9 Hz), 3.28 (1H, t, J = 9.8 Hz), 3.43-3.51 (1H, m), 3.68 (1H, t, J = 8.6 Hz), 3.71 (3H, s), 4.48-4.55 (2H, m), 6.54 (1H, d, J = 9.2 Hz), 6.87 (2H, d, J = 8.6 Hz), 7.04 (2H, t, J = 8.6 Hz), 7.31-7.38 (4H, m), 7.83 (1H, d, J = 4.9 Hz), 8.67 (1H, brs).

15 MS (ESI⁺) m/z: 429(MH⁺).


[α]_D²⁵ = -199 (c 0.34, EtOH).

[0706]

<Example 20-1>

[0707]

20 [Chemical Formula 141]



[0708]

(-)-2-[(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B)

The same method as in Example 16-1 was performed using

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid ethyl ester (isomer B) in place of

2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid ethyl ester to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 1.36 (3H, d, J = 7.4 Hz), 3.20 (1H, t, J = 9.2 Hz), 3.43-3.50 (1H, m), 3.68-3.74 (1H, m), 3.72 (3H, s), 4.49 (1H, dd, J = 11.0, 8.6 Hz), 4.62 (1H, q, J = 7.4 Hz), 6.53 (1H, d, J = 8.6 Hz), 6.90 (2H, d, J = 8.6 Hz), 7.03 (2H, t, J = 9.2 Hz), 7.27 (2H, d, J = 8.6 Hz), 7.32-7.38 (2H, m), 8.61 (1H, s), 12.9 (1H, brs).

MS (ESI⁺) m/z: 416 (MH⁺).


[α]_D²⁵ = -111 (c 0.32, EtOH).

[0709]

<Example 21-1>

[0710]

[Chemical Formula 142]



[0711]

(-)-2-[(3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer B)

The same method as in Example 17-1 was performed using

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B) in place of (-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 1.29 (3H, d, J = 7.3 Hz), 2.57 (3H, d, J = 4.9 Hz), 3.24 (1H, t, J = 9.8 Hz), 3.34-3.44 (1H, m), 3.66 (1H, t, J = 8.6 Hz), 3.72 (3H, s), 4.56 (1H, q, J = 7.3 Hz), 4.69 (1H, dd, J = 11.6, 9.2 Hz), 6.50 (1H, d, J = 9.2 Hz), 6.89 (2H, d, J = 8.6 Hz), 7.03 (2H, t, J = 9.2 Hz), 7.30 (2H, d, J = 8.6 Hz), 7.34-7.38 (2H, m), 7.87 (1H, d, J = 4.9 Hz), 8.60 (1H, s).

MS (ESI⁺) m/z: 429 (MH⁺).


[α]_D²⁵ = -94 (c 0.30, EtOH).

[0712]

<Example 22-1>

[0713]

[Chemical Formula 143]



[0714]

2-((3S*,4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}-2-methylpropionic acid

The same method as in Example 16-1 was performed using
 5 2-((3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}-2-methylpropionic acid ethyl ester in place of

2-((3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl}propionic acid ethyl ester to obtain the
 10 title compound.

¹H-NMR (400 MHz, CDCl₃) δ 1.52 (3H, s), 1.53 (3H, s), 3.36 (1H, t, J = 9.2 Hz), 3.60-3.80 (1H, m), 3.76 (3H, s), 3.78-3.86 (1H, m), 4.60-4.68 (1H, m), 6.06 (1H, brs), 6.72-6.78 (2H, m), 6.84 (2H, d, J = 8.6 Hz), 7.07-7.13 (2H, m), 7.23 (2H, d, J = 8.6 Hz), 7.61
 15 (1H, brs).


MS (ESI⁺) m/z: 430 (MH⁺).

[0715]

Example 22-2

[0716]

20 [Chemical Formula 144]



[0717]

(-)-2-[(3S*,4R*)-4-(6-Fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-2-methylpropanoic acid

A 2 mol/L sodium hydroxide aqueous solution (0.2 mL) was added to a solution of

2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-2-methylpropanoic

acid ethyl ester (18 mg) in methanol (0.2 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 18 hours. A 1 mol/L hydrochloric acid was added to the reaction solution to adjust the pH of the reaction solution to 1, and the reaction solution was concentrated under reduced pressure. Water

was added to the resulting residue, and the precipitate was collected by filtration and purified by preparative TLC

(chloroform : methanol = 10 : 1) to obtain the title compound as a colorless solid (4.4 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 1.40 (3H, s), 1.42 (3H, s), 3.12 (2H, t, J = 8.6 Hz), 3.28-3.37 (1H, m), 3.47-3.59 (1H, m), 3.70 (1H, t, J = 8.3 Hz), 4.53-4.62 (3H, m), 6.64 (1H, d, J = 11.6 Hz), 6.70 (1H, brs), 7.02 (2H, t, J = 8.9 Hz), 7.31-7.40 (3H, m), 8.73 (1H, brs), 12.47 (1H, brs).

MS (ESI⁺) m/z: 460 (MH⁺).


[α]_D²⁷ = -163 (c 0.09, DMSO)

[0718]

<Example 23-1>

5 [0719]

[Chemical Formula 145]



[0720]

(-)-2-[(3S*, 4R*)-3-[3-(4-Fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N,2-dimethylpropionamide

10 The same method as in Example 17-1 was performed using 2-[(3S*, 4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-2-methylpropionic acid in place of (-)-2-[(3S*, 4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]acetic acid to obtain the title
15 compound.

¹H-NMR (400 MHz, CDCl₃) δ 1.53 (3H, s), 1.72 (3H, s), 2.69 (3H, d, J = 4.9 Hz), 3.45-3.49 (1H, m), 3.80 (3H, s), 3.85 (2H, dd, J = 7.3, 4.3 Hz), 3.93-3.98 (1H, m), 6.73 (1H, brs), 6.78 (2H, t, J = 9.2 Hz), 6.89 (2H, t, J = 8.6 Hz), 7.08 (2H, dd, J = 9.2, 4.9 Hz), 7.17 (2H, d, J = 8.6 Hz), 7.36 (1H, d, J = 4.3 Hz), 7.88 (1H, brs).

MS (ESI⁺) m/z: 443 (MH⁺).


$[\alpha]_D^{29} = -109$ (c 0.31, EtOH).

[0721]

<Example 24-1>

[0722]

5 [Chemical Formula 146]



[0723]

(-)-1-[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

10 10% Palladium on carbon (10 mg) was added to a solution of [(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester (100 mg) in ethanol (5 mL) to produce a reaction solution. The reaction solution was stirred under a hydrogen atmosphere for 1 hour. The reaction solution was filtered over Celite, and the solvent was removed
15 under reduced pressure to obtain

(3S*, 4R*)-3-amino-4-(2,6-difluoro-4-methoxyphenyl)-1-methylpyrrolidin-2-one as a colorless oily intermediate compound.

4-Fluorophenyl isocyanate (28 μ L) was added to a solution of the resulting

20 (3S*, 4R*)-3-amino-4-(2,6-difluoro-4-methoxyphenyl)-1-methylpyrrolidin-2-one (66 mg) in tetrahydrofuran (2.6 mL) to produce a reaction solution. The reaction solution was stirred at room

temperature for 20 minutes. The reaction solution was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title
5 compound as a white solid (91 mg).

$^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ 2.81 (3H, s), 3.40 (1H, dd, $J = 9.8, 9.2$ Hz), 3.53 (1H, t, $J = 9.2$ Hz), 3.73-3.81 (1H, m), 3.75 (3H, s), 4.57 (1H, dd, $J = 9.8, 8.6$ Hz), 6.52 (1H, d, $J = 8.6$ Hz), 6.74 (2H, d, $J = 11.0$ Hz), 6.97-7.05 (2H, m), 7.30-7.36 (2H, m), 8.72
10 (1H, s).

MS (ESI^+) m/z : 394 (MH^+).

$[\alpha]_{\text{D}}^{27} = -143$ (c 0.10, EtOH)




[0724]

The same method as in Example 24-1 was performed using a
15 corresponding Cbz substance and isocyanate to obtain the following Examples 24-2 to 24-24.

The structures and spectral data thereof are shown in Tables 90 to 100.



[0725]

[Table 90]

Ex.No	Structure	Chemical name	P.D.
24-2		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.18 (3H, s), 2.81 (3H, s), 3.40 (1H, t, J = 9.2 Hz), 3.52 (1H, t, J = 7.8 Hz), 3.71-3.80 (1H, m), 3.75 (3H, s), 4.56 (1H, dd, J = 10.4, 8.6 Hz), 6.51 (1H, d, J = 8.6 Hz), 6.70-6.78 (2H, m), 6.98 (2H, d, J = 8.0 Hz), 7.21 (2H, d, J = 8.0 Hz), 8.59 (1H, s). MS (ESI ⁺) m/z: 390 (MH ⁺). [α] _D ²⁷ = -120 (c 0.14, EtOH)
24-3		(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.81 (3H, s), 3.41 (1H, t, J = 9.8 Hz), 3.54 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.80 (1H, q, J = 9.8 Hz), 4.58 (1H, t, J = 9.8 Hz), 6.71-6.78 (2H, m), 6.85 (1H, brd, J = 7.9 Hz), 7.51 (2H, d, J = 8.0 Hz), 7.62 (2H, d, J = 8.0 Hz), 9.34 (1H, s). MS (ESI ⁺) m/z: 401 (MH ⁺). [α] _D ²⁷ = -176 (c 0.11, EtOH)
24-4		(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.81 (3H, s), 3.41 (1H, t, J = 9.8 Hz), 3.53 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.80 (1H, q, J = 9.8 Hz), 4.57 (1H, t, J = 9.8 Hz), 6.70-6.79 (2H, m), 6.94 (1H, brs), 7.32 (1H, d, J = 8.6 Hz), 7.87 (1H, dd, J = 8.6, 3.1 Hz), 8.35 (1H, d, J = 3.1 Hz), 9.24 (1H, brs). MS (ESI ⁺) m/z: 411 (MH ⁺). [α] _D ²⁷ = -197 (c 0.11, EtOH)

[0726]

[Table 91]

Ex.No	Structure	Chemical name	P.D.
24-5		(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.83 (3H, s), 3.43 (1H, t, J = 9.8 Hz), 3.55 (1H, t, J = 9.8 Hz), 3.75 (3H, s), 3.83 (1H, q, J = 9.8 Hz), 4.60 (1H, t, J = 9.8 Hz), 6.67 (1H, s), 6.72-6.79 (2H, m), 6.84 (1H, brd, J = 8.0 Hz), 7.09 (1H, td, J = 7.9, 1.2 Hz), 7.21 (1H, td, J = 7.9, 1.2 Hz), 7.52 (1H, d, J = 7.9 Hz), 7.69 (1H, d, J = 7.9 Hz), 10.1 (1H, s). MS (ESI ⁺) m/z: 432 (MH ⁺). [α] _D ²⁹ = -189 (c 0.10, EtOH)
24-6		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-phenylurea	¹ H-NMR (400 MHz, CDCl ₃) δ 3.21-3.28 (1H, m), 3.35-3.44 (1H, m), 3.47-3.63 (4H, m), 3.68-3.75 (1H, m), 3.75 (3H, s), 4.64-4.68 (1H, m), 4.77 (1H, t, J = 5.5 Hz), 6.51 (1H, d, J = 8.6 Hz), 6.74 (2H, d, J = 10.4 Hz), 6.87 (1H, t, J = 7.3 Hz), 7.18 (2H, t, J = 7.9 Hz), 7.32 (2H, d, J = 7.3 Hz), 8.64 (1H, s). MS (ESI ⁺) m/z: 406 (MH ⁺). [α] _D ²⁴ = -88 (c 0.13, EtOH)



[0727]

[Table 92]

Ex.No	Structure	Chemical name	P.D.
24-7		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.18 (3H, s), 3.20-3.30 (1H, m), 3.33-3.43 (1H, m), 3.45-3.62 (4H, m), 3.67-3.75 (1H, m), 3.75 (3H, s), 4.65 (1H, dd, J = 9.2, 8.6 Hz), 4.77 (1H, t, J = 5.5 Hz), 6.46 (1H, d, J = 8.6 Hz), 6.70-6.77 (2H, m), 6.98 (2H, d, J = 8.6 Hz), 7.20 (2H, d, J = 8.6 Hz), 8.53 (1H, s). MS (ESI ⁺) m/z: 420 (MH ⁺). [α] _D ²⁷ = -199 (c 0.12, EtOH)
24-8		(-)-1-(4-cyano phenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.20-3.28 (1H, m), 3.37-3.46 (1H, m), 3.48-3.63 (4H, m), 3.72-3.82 (1H, m), 3.75 (3H, s), 4.66 (1H, dd, J = 10.4, 8.6 Hz), 4.78 (1H, t, J = 4.9 Hz), 6.70-6.78 (2H, m), 6.83 (1H, d, J = 8.6 Hz), 7.49 (2H, d, J = 8.6 Hz), 7.62 (2H, d, J = 8.6 Hz), 9.29 (1H, s). MS (ESI ⁺) m/z: 431 (MH ⁺). [α] _D ²⁸ = -148 (c 0.11, EtOH)

[0728]

[Table 93]

Ex.No	Structure	Chemical name	P.D.
24-9		(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.18-3.28 (1H, m), 3.38-3.46 (1H, m), 3.47-3.63 (4H, m), 3.73-3.81 (1H, m), 3.75 (3H, s), 4.66 (1H, dd, J = 11.0, 8.6 Hz), 4.77 (1H, t, J = 5.5 Hz), 6.71-6.78 (2H, m), 6.83 (1H, d, J = 8.6 Hz), 7.32 (1H, d, J = 8.6 Hz), 7.87 (1H, dd, J = 8.6, 3.0 Hz), 8.34 (1H, d, J = 3.0 Hz), 9.09 (1H, s). MS (ESI ⁺) m/z: 441 (MH ⁺). [α] _D ²⁶ = -153 (c 0.10, EtOH)
24-10		(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.20-3.27 (1H, m), 3.40-3.48 (1H, m), 3.50-3.64 (4H, m), 3.76 (3H, s), 3.80 (1H, q, J = 9.8 Hz), 4.68 (1H, dd, J = 9.8, 8.6 Hz), 4.78 (1H, t, J = 5.5 Hz), 6.67 (1H, s), 6.72-6.79 (2H, m), 6.87 (1H, brd, J = 8.0 Hz), 7.09 (1H, td, J = 7.9, 1.2 Hz), 7.21 (1H, td, J = 7.9, 1.2 Hz), 7.52 (1H, d, J = 7.9 Hz), 7.69 (1H, d, J = 7.9 Hz), 10.1 (1H, s). MS (ESI ⁺) m/z: 462 (MH ⁺). [α] _D ²⁸ = -170 (c 0.11, EtOH)



[0729]

[Table 94]

Ex.No	Structure	Chemical name	P.D.
24-11		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-propylpyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, CDCl ₃) δ 0.96 (3H, t, J = 7.3 Hz), 1.57-1.67 (2H, m), 3.25-3.33 (1H, m), 3.42-3.50 (1H, m), 3.57 (1H, t, J = 9.2 Hz), 3.62-3.70 (1H, m), 3.78 (3H, s), 4.14-4.23 (1H, m), 4.39-4.46 (1H, m), 6.20 (1H, brs), 6.47 (2H, d, J = 10.4 Hz), 6.70-6.77 (2H, m), 7.03-7.10 (2H, m), 7.68 (1H, brs). MS (ESI ⁺) m/z: 422 (MH ⁺). [α] _D ²⁴ = -126 (c 0.19, EtOH)
24-12		(-)-1-[(3S*,4R*)-1-benzyl-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, CD ₃ CN) δ 3.35-3.45 (2H, m), 3.72 (3H, s), 3.88 (1H, q, J = 9.2 Hz), 4.47 (2H, s), 4.65 (1H, dd, J = 10.4, 8.0 Hz), 5.66 (1H, d, J = 8.0 Hz), 6.53 (2H, d, J = 11.0 Hz), 6.91 (2H, t, J = 9.2 Hz), 7.23-7.36 (7H, m), 7.48 (1H, s). MS (ESI ⁺) m/z: 470 (MH ⁺). [α] _D ²⁹ = -109 (c 0.17, EtOH)
24-13		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxo-1-(pyridin-3-ylmethyl)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.46 (1H, t, J = 9.8 Hz), 3.55 (1H, t, J = 9.8 Hz), 3.74 (3H, s), 3.88 (1H, q, J = 9.6 Hz), 4.51 (1H, d, J = 15.9 Hz), 4.61-4.69 (2H, m), 6.64 (1H, d, J = 9.5 Hz), 6.74 (2H, d, J = 9.5 Hz), 7.03 (2H, td, J = 7.6, 2.2 Hz), 7.29-7.37 (4H, m), 7.81 (1H, td, J = 7.6, 1.8 Hz), 8.54 (1H, q, J = 2.2 Hz), 8.74 (1H, s). MS (ESI ⁺) m/z: 470 (MH ⁺). [α] _D ³⁰ = -142 (c 0.10, EtOH)

[0730]

[Table 95]

Ex.No	Structure	Chemical name	P.D.
24-14		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(methylsulfonyl)methyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.31 (3H, s), 3.67 (1H, t, J = 9.5 Hz), 3.76 (3H, s), 3.85 (1H, t, J = 8.9 Hz), 4.02 (1H, q, J = 9.8 Hz), 4.55 (1H, dd, J = 11.0, 7.9 Hz), 4.71 (1H, d, J = 14.7 Hz), 4.82 (1H, d, J = 14.7 Hz), 6.72 (1H, d, J = 10.0 Hz), 6.77 (2H, d, J = 10.0 Hz), 7.03 (2H, dt, J = 14.5, 3.7 Hz), 7.33-7.36 (2H, m), 8.89 (1H, s). MS (ESI ⁺) m/z: 472 (MH ⁺). [α] _D ²⁸ = -203 (c 0.10, EtOH)
24-15		(-)-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]methanesulfonamide	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.68-3.90 (6H, m), 4.53 (1H, d, J = 13.5 Hz), 4.69 (1H, d, J = 13.5 Hz), 4.77 (1H, t, J = 10.0 Hz), 6.51 (1H, d, J = 8.6 Hz), 6.76 (2H, d, J = 10.4 Hz), 7.03 (2H, t, J = 8.6 Hz), 7.09 (2H, s), 7.31-7.35 (2H, m), 8.81 (1H, s). MS (ESI ⁺) m/z: 473 (MH ⁺). [α] _D ²⁵ = -120 (c 0.10, EtOH)



[0731]

[Table 96]

Ex.No	Structure	Chemical name	P.D.
24-16		(-)-2-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid ethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.21 (3H, t, J = 6.7 Hz), 3.51-3.61 (2H, m), 3.76 (3H, s), 3.82 (1H, q, J = 10.4 Hz), 3.97 (1H, d, J = 17.0 Hz), 4.13 (2H, q, J = 6.7 Hz), 4.27 (1H, d, J = 17.0 Hz), 4.67 (1H, dd, J = 10.4, 8.6 Hz), 6.68-6.80 (3H, m), 7.01 (2H, t, J = 9.2 Hz), 7.33 (2H, dd, J = 9.2, 4.9 Hz), 8.79 (1H, s). MS (ESI ⁺) m/z: 466 (MH ⁺). [α] _D ²⁶ = -128 (c 0.11, EtOH)
24-17		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-3-[(5-methyloxazol-2-yl)methyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.27 (3H, s), 3.49 (1H, t, J = 9.2 Hz), 3.59 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.82 (1H, q, J = 9.8 Hz), 4.44 (1H, d, J = 16.5 Hz), 4.64-4.71 (2H, m), 6.61 (1H, d, J = 8.6 Hz), 6.75 (2H, d, J = 10.4 Hz), 6.80 (1H, s), 7.02 (2H, t, J = 9.2 Hz), 7.33 (2H, dd, J = 8.6, 4.9 Hz), 8.71 (1H, s). MS (ESI ⁺) m/z: 475 (MH ⁺). [α] _D ²⁵ = -108 (c 0.12, EtOH)



[0732]

[Table 97]

Ex.No	Structure	Chemical name	P.D.
24-18		(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.49 (3H, s), 3.51 (1H, t, J = 9.2 Hz), 3.62 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.84 (1H, q, J = 9.2 Hz), 4.64-4.68 (2H, m), 4.81 (1H, d, J = 16.5 Hz), 6.61 (1H, d, J = 8.6 Hz), 6.76 (2H, d, J = 11.0 Hz), 7.02 (2H, t, J = 9.2 Hz), 7.33 (2H, dd, J = 8.6, 4.9 Hz), 8.71 (1H, s). MS (ESI ⁺) m/z: 476 (MH ⁺). [α] _D ²⁹ = -106 (c 0.11, EtOH)
24-19		(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.12 (2H, t, J = 8.6 Hz), 3.16-3.21 (1H, m), 3.29-3.34 (2H, m), 3.49-3.53 (2H, m), 3.56-3.66 (2H, m), 4.55 (2H, t, J = 8.6 Hz), 4.62 (1H, t, J = 9.5 Hz), 4.76 (1H, t, 5.5 Hz), 6.53-6.66 (1H, m), 6.64 (1H, d, J = 11.0 Hz), 7.03 (2H, t, J = 8.6 Hz), 7.34-7.39 (3H, m), 8.67-8.76 (1H, m). MS (ESI ⁺) m/z: 418 (MH ⁺). [α] _D ²⁶ = -164 (c 0.12, EtOH)

[0733]

[Table 98]

Ex.No	Structure	Chemical name	P.D.
24-20		(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]acetic acid ethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.21 (3H, t, J = 6.7 Hz), 3.10-3.16 (2H, m), 3.27-3.38 (1H, m), 3.60 (1H, t, J = 8.6 Hz), 3.65-3.70 (1H, m), 4.02 (1H, d, J = 17.7 Hz), 4.11-4.19 (3H, m), 4.55 (2H, t, J = 8.3 Hz), 4.62 (1H, t, J = 9.8 Hz), 6.66 (1H, d, J = 11.0 Hz), 6.81 (1H, brs), 7.03 (2H, t, J = 8.6 Hz), 7.31-7.38 (3H, m), 8.81-8.84 (1H, m). MS (ESI ⁺) m/z: 460 (MH ⁺). [α] _D ²⁷ = -109 (c 0.11, EtOH)
24-21		(-)-2-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-3-[3-(4-fluorophenyl)ureido]-2-oxopyrrolidin-1-yl]-2-methylpropanoic acid ethyl ester	¹ H-NMR (400 MHz, DMSO-d ₆) δ 1.18 (3H, t, J = 7.0 Hz), 1.39 (3H, s), 1.44 (3H, s), 3.12 (2H, t, J = 8.9 Hz), 3.28-3.38 (1H, m), 3.60 (1H, t, J = 10.1 Hz), 3.74 (1H, t, J = 8.9 Hz), 4.03-4.09 (2H, m), 4.53-4.57 (3H, m), 6.62-6.67 (1H, m), 6.66 (1H, d, J = 11.0 Hz), 7.02 (2H, t, J = 8.9 Hz), 7.33-7.36 (3H, m), 8.70 (1H, s). MS (ESI ⁺) m/z: 488 (MH ⁺). [α] _D ²⁷ = -140 (c 0.12, EtOH)

[0734]

[Table 99]

Ex.No	Structure	Chemical name	P.D.
24-22		(-)-1-[(3S*,4R*)-1-(cyclopropylmethyl)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 0.21-0.29 (2H, m), 0.49-0.55 (2H, m), 0.92-1.02 (1H, m), 3.03 (1H, dd, J = 13.9, 7.3), 3.17 (2H, t, J = 8.8 Hz), 3.28 (1H, dd, J = 13.9, 7.0 Hz), 3.33-3.43 (1H, m), 3.64-3.73 (2H, m), 4.58-4.67 (1H, m), 4.60 (2H, t, J = 8.8 Hz), 6.64-6.71 (1H, m), 6.70 (1H, J = 11.0 Hz), 7.07 (2H, t, J = 8.9 Hz), 7.38-7.46 (3H, m), 8.77 (1H, s). MS (ESI ⁺) m/z: 428 (MH ⁺). [α] _D ²⁷ = -159 (c 0.11, EtOH)
24-23		(-)-1-[(3S*,4R*)-1-(cyanomethyl)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.12 (2H, t, J = 9.1 Hz), 3.29-3.34 (1H, m), 3.65 (1H, t, J = 9.1 Hz), 3.73-3.81 (1H, m), 4.44 (2H, s), 4.53-4.60 (3H, m), 6.65 (2H, d, J = 10.9 Hz), 7.25-7.32 (1H, m), 7.37-7.40 (3H, m), 9.32 (1H, brs). MS (ESI ⁺) m/z: 413 (MH ⁺). [α] _D ²⁶ = -136 (c 0.11, EtOH)

[0735]

[Table 100]


Ex.No	Structure	Chemical name	P.D.
24-24		(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-1-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.49 (3H, s), 3.12 (2H, t, J = 9.1 Hz), 3.29–3.37 (1H, m), 3.63 (1H, t, J = 9.1 Hz), 3.69–3.77 (1H, m), 4.55 (2H, t, J = 8.5 Hz), 4.62–4.76 (3H, m), 6.57 (1H, d, J = 9.1 Hz), 6.65 (1H, d, J = 10.9 Hz), 7.01–7.06 (2H, m), 7.34–7.38 (3H, m), 8.64 (1H, s). MS (ESI ⁺) m/z: 470 (MH ⁺). [α] _D ²⁷ = -126 (c 0.10, EtOH)

[0736]

5 <Example 25-1>

[0737]

[Chemical Formula 147]



[0738]

(-)-1-[(3S*,4R*)-4-(2-Fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

The same method as in Example 24-1 was performed using [(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester in place of

[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 3.14-3.20 (1H, m), 3.28-3.42 (2H, m),
5 3.48-3.52 (2H, m), 3.57-3.66 (2H, m), 3.72 (3H, s), 4.66 (1H, dd,
J = 11.0, 8.6 Hz), 4.73 (1H, t, J = 5.5 Hz), 6.45 (1H, d, J = 8.6
Hz), 6.75-6.80 (2H, m), 6.98-7.04 (2H, m), 7.30-7.36 (2H, m), 7.44
(1H, dd, J = 9.2, 9.2 Hz), 8.60 (1H, s).

MS (ESI⁺) m/z: 406 (MH⁺).


10 [α]_D²⁷ = -118 (c 0.10, EtOH)

[0739]

<Example 26-1>

[0740]

[Chemical Formula 148]



15 [0741]

(-)-1-[(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

The same method as in Example 24-1 was performed using
[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl
20)-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester in place of
[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]carbamic acid benzyl ester to obtain the title

compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 3.21-3.30 (1H, m), 3.36-3.45 (1H, m), 3.48-3.62 (4H, m), 3.71-3.78 (1H, m), 3.76 (3H, s), 4.66 (1H, dd, J = 10.4, 8.6 Hz), 4.77 (1H, t, J = 5.5 Hz), 6.51 (1H, d, J = 8.6 Hz), 6.75 (2H, d, J = 11.0 Hz), 6.99-7.06 (2H, m), 7.31-7.37 (2H, m), 8.68 (1H, s).

MS (ESI⁺) m/z: 424 (MH⁺).


[α]_D²⁷ = -139 (c 0.11, EtOH)

[0742]

10 <Example 27-1>

[0743]

[Chemical Formula 149]



[0744]

(+)-1-(4-Fluorophenyl)-3-[(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea
15 nyl)-2-oxopyrrolidin-3-yl]urea

The same method as in Example 1-1 was performed using (3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid in place of

(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain the title compound.
20 ic acid to obtain the title compound.

¹H-NMR (400 MHz, CDCl₃) δ 3.55 (1H, t, J = 8.5 Hz), 3.74-3.82 (1H, m), 3.80 (3H, s), 3.89 (3H, s), 4.01 (1H, t, J = 8.2 Hz), 4.10 (1H,

dd, J = 8.5, 8.2 Hz), 6.08 (1H, brs), 6.79 (2H, t, J = 8.8 Hz),
6.89 (2H, d, J = 9.1 Hz), 7.12 (2H, q, J = 4.6 Hz), 7.20 (2H, d,
J = 9.1 Hz), 7.50 (1H, brs).

MS (ESI⁺) m/z: 374 (MH⁺).


5 $[\alpha]_D^{27} = +144$ (c 0.12, EtOH)

[0745]

<Example 27-2>

[0746]

[Chemical Formula 150]



[0747]

(-)-1-[(3S*,4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

The same method as in Example 1-1 was performed using
15 (3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methoxy-2-oxopyrrolidine-3-carboxylic acid in place of
(-)-(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidine-3-carboxylic acid to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 3.54-3.61 (1H, m), 3.73 (3H, s), 3.76
20 (3H, s), 3.80-3.89 (2H, m), 4.46 (1H, t, J = 8.9 Hz), 6.67 (1H,
d, J = 8.6 Hz), 6.77 (2H, d, J = 8.6 Hz), 7.00-7.05 (2H, m), 7.32-7.36
(2H, m), 8.83 (1H, s).

MS (ESI⁺) m/z: 410 (MH⁺).


$[\alpha]_D^{25} = -208$ (c 0.20, EtOH)

[0748]

<Example 28-1>

5 [0749]

[Chemical Formula 151]



[0750]

(-)-1-[(3S*, 4R*)-4-(2,6-Difluoro-4-hydroxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

10 10% Palladium on carbon (49 mg) was added to a solution of 1-[(3S*, 4R*)-4-[4-(benzyloxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea (490 mg) in ethanol (10 mL) under an argon atmosphere to produce a reaction solution. The reaction solution was stirred at room temperature under a hydrogen
15 atmosphere for 4 hours. The reaction solution was filtered over Celite, and the solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title compound as a white solid (383 mg).

20 ¹H-NMR (400 MHz, DMSO-d₆) δ 3.23-3.28 (1H, m), 3.41 (1H, t, J = 9.2 Hz), 3.74 (1H, q, J = 9.6 Hz), 4.52 (1H, dd, J = 10.7, 8.3 Hz), 6.41 (2H, d, J = 10.4 Hz), 6.45 (1H, d, J = 7.9 Hz), 6.99-7.04 (2H,

m), 7.31-7.35 (2H, m), 8.01 (1H, s), 8.66 (1H, s).

MS (ESI⁺) m/z: 366 (MH⁺).


[α]_D²⁸ = -151 (c 0.1, EtOH)

[0751]

5 <Example 28-2>

[0752]

[Chemical Formula 152]



[0753]

10 1-[(3S*, 4R*)-4-(2,6-Difluoro-4-hydroxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea

The same method as in Example 28-1 was performed using 1-[(3S*, 4R*)-4-(4-(benzyloxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl)-3-phenylurea in place of

15 1-[(3S*, 4R*)-4-(4-benzyloxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea to obtain the title compound.

¹H NMR (400 MHz, DMSO-d₆) δ 3.25-3.30 (1H, m), 3.42 (1H, t, J = 8.6 Hz), 3.74 (1H, q, J = 9.6 Hz), 4.53 (1H, dd, J = 11.0, 8.6 Hz), 6.41-6.47 (3H, m), 6.86 (1H, t, J = 7.3 Hz), 7.18 (2H, t, J = 7.9 Hz), 7.32 (2H, d, J = 7.9 Hz), 8.03 (1H, s), 8.60 (1H, s), 10.35 (1H, brs).


MS (ESI⁺) m/z: 348 (MH⁺).

[0754]

<Example 28-3>

[0755]

[Chemical Formula 153]



5

[0756]

(-)-1-[(3S*,4R*)-4-(6-Fluoroindolin-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

The same method as in Example 28-1 was performed using

10 1-[(3S*,4R*)-4-(1-benzyl-6-fluoroindolin-5-yl)-2-

oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea in place of

1-[(3S*,4R*)-4-(4-benzyloxy-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 2.83 (2H, t, J = 8.5 Hz), 3.11 (1H, t, J = 9.7 Hz), 3.35-3.44 (3H, m), 3.53-3.61 (1H, m), 4.47 (1H, dd, J = 11.5, 9.1 Hz), 5.68 (1H, br), 6.20 (1H, d, J = 11.5 Hz), 6.36 (1H, d, J = 8.5 Hz), 7.00-7.05 (2H, m), 7.11 (1H, d, J = 7.9 Hz), 7.33-7.37 (2H, m), 7.87 (1H, s), 8.55 (1H, s).

MS (ESI⁺) m/z: 373 (MH⁺).


20 [α]_D²⁷ = -202 (c 0.12, EtOH)

[0757]

<Example 28-4>

[0758]

[Chemical Formula 154]



[0759]

- 5 (-)-1-[(3S*, 4R*)-4-(2,6-Difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-ethynylphenyl)urea

Potassium carbonate (1 mg) was added to a solution of 1-[(3S*, 4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-{4-[(trimethylsilyl)ethynyl]phenyl}urea (66 mg) in
10 methanol (1.4 mL) under ice-cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 8 hours. The reaction solution was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1, ethyl acetate,
15 and ethyl acetate : methanol = 4 : 1 in turn), to obtain the title compound as a colorless solid (44 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.27-3.32 (1H, m), 3.45 (1H, t, J = 9.2 Hz), 3.75 (3H, s), 3.80 (1H, ddd, J = 11.0, 9.2, 9.2 Hz), 3.98 (1H, s), 4.56 (1H, dd, J = 11.0, 8.6 Hz), 6.57 (1H, d, J = 8.6 Hz), 6.70-6.77
20 (2H, m), 7.28 (2H, d, J = 8.6 Hz), 7.35 (2H, d, J = 8.6 Hz), 8.07 (1H, s), 8.89 (1H, s).

MS (ESI⁺) m/z: 386 (MH⁺).


$[\alpha]_D^{28} = -147$ (c 0.11, EtOH)

[0760]

<Example 28-5>

[0761]

5 [Chemical Formula 155]



[0762]

(-)-1-[(3S*,4R*)-4-[2,6-Difluoro-4-(methanimino)phenyl]-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

10 Trifluoroacetic acid (1 mL) was added to
(3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]phenyl)(methyl)carbamic acid tert-butyl ester
(80 mg) to produce a reaction solution. The reaction solution was
stirred at 60°C for 1 hour. The reaction solution was concentrated
15 under reduced pressure, a saturated aqueous sodium hydrogen
carbonate solution was added to the residue, and the mixture was
extracted with ethyl acetate. The organic layer was washed with
water and a brine successively, and dried over anhydrous sodium
sulfate. The solvent was removed under reduced pressure, and the
20 residue was then purified by silica gel column chromatography
(ethyl acetate), to obtain the title compound as a colorless solid
(30 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 2.63 (3H, d, J = 4.9 Hz), 3.25 (1H, t, J = 10.1 Hz), 3.37 (1H, t, J = 8.9 Hz), 3.67 (1H, q, J = 9.8 Hz), 4.52 (1H, dd, J = 11.0, 8.6 Hz), 6.16 (2H, d, J = 12.0 Hz), 6.21 (1H, d, J = 4.9 Hz), 6.40 (1H, d, J = 8.6 Hz), 6.99-7.04 (2H, m),
5 7.32-7.36 (2H, m), 7.98 (1H, s), 8.61 (1H, s).

MS (ESI⁺) m/z: 379 (MH⁺).


[α]_D²⁴ = -54 (c 0.14, EtOH)

[0763]

<Example 28-6>

10 [0764]

[Chemical Formula 156]



[0765]

1-[(3S*, 4R*)-4-(6-fluoro-1-oxo-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea
15

A Dess-Martin reagent (14 mg) was added to a solution of 1-[(3S*, 4R*)-4-(6-fluoro-1-hydroxy-2,3-dihydro-1H-inden-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea (13 mg) in a mixture of dichloromethane (0.66 mL) and acetonitrile (0.66 mL)
20 under an argon atmosphere under cooling to produce a reaction solution. The reaction solution was stirred at room temperature for 3 hours. A saturated aqueous sodium hydrogen carbonate

solution was added to the reaction solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 1 : 1, ethyl acetate, and ethyl acetate : methanol = 9 : 1 in turn), to obtain the title compound as a colorless solid (11 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 2.64-2.67 (2H, m), 3.15 (2H, t, J = 5.2 Hz), 3.27 (1H, t, J = 8.6 Hz), 3.53 (1H, t, J = 8.9 Hz), 3.89 (1H, q, J = 9.8 Hz), 4.65 (1H, dd, J = 11.6, 8.6 Hz), 6.53 (1H, d, J = 8.6 Hz), 7.00-7.04 (2H, m), 7.32-7.37 (3H, m), 7.82 (1H, d, J = 6.1 Hz), 8.04 (1H, s), 8.69 (1H, s).


MS (ESI⁺) m/z: 386 (MH⁺).

[0766]

<Example 29-1>

[0767]

[Chemical Formula 157]



[0768]

(-)-3,5-Difluoro-4-((3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl)-N,N-dimethylbenzamide

Acetonitrile (3 mL), palladium(II) acetate (14 mg),
(±)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (rac-BINAP)
(38 mg), hexacarbonyl molybdenum (Mo(CO)₆) (240 mg), cesium
carbonate (236 mg), and dimethylamine (3 mL, 2.0 M tetrahydrofran
5 solution) were added to a solution of

3,5-difluoro-4-{{(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxop
yrrolidin-3-yl}phenyl trifluoromethanesulfonate (300 mg) in
toluene (6 mL) to produce a reaction solution. The reaction
solution was stirred at 70°C under an argon atmosphere for 3 hours.

10 The reaction solution was filtered over Celite and washed with
ethyl acetate. The solvent was removed under reduced pressure,
and the residue was then purified by silica gel column
chromatography (hexane : ethyl acetate = 4 : 1, ethyl acetate,
and ethyl acetate : methanol = 9 : 1 in turn), to obtain the title
15 compound as a white solid (192 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 2.88 (3H, s), 2.95 (3H, s), 3.35-3.39
(1H, m), 3.51 (1H, t, J = 8.9 Hz), 3.94 (1H, q, J = 9.6 Hz), 4.58
(1H, dd, J = 10.7, 8.3 Hz), 6.50 (1H, d, J = 7.9 Hz), 6.98-7.03
(2H, m), 7.17 (2H, d, J = 9.2 Hz), 7.29-7.33 (2H, m), 8.11 (1H,
20 s), 8.71 (1H, s).

MS (ESI⁺) m/z: 421 (MH⁺).


[α]_D²⁸ = -111(c 0.1, EtOH)

[0769]

<Example 29-2>

25 [0770]

[Chemical Formula 158]



[0771]

(-)-3,5-Difluoro-4-{(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl}-N-methylbenzamide

5 The same method as in Example 29-1 was performed using methyl amine in place of dimethyl amine to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 2.77 (3H, d, J = 4.9 Hz), 3.34-3.40 (1H, m), 3.51 (1H, t, J = 9.2 Hz), 3.93 (1H, q, J = 9.6 Hz), 4.60 (1H, dd, J = 10.4, 7.9 Hz), 6.53 (1H, d, J = 7.9 Hz), 6.98-7.03 (2H, m), 7.29-7.32 (2H, m), 7.52 (2H, d, J = 9.8 Hz), 8.13 (1H, s), 8.56-8.60 (1H, m), 8.74 (1H, s).

MS (ESI⁺) m/z: 407 (MH⁺).


[α]_D²⁸ = -167 (c 0.1, EtOH)

[0772]

15 <Example 30-1>

[0773]

[Chemical Formula 159]



[0774]

(-)-1-[(3S*,4R*)-4-(4-Cyano-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea

Zinc cyanide (53 mg) and

5 tetrakis(triphenylphosphine)palladium (18 mg) were added to a solution of

3,5-difluoro-4-[(3R*,4S*)-4-(3-(4-fluorophenyl)ureido)-5-oxopyrrolidin-3-yl]phenyl trifluoromethanesulfonate (150 mg) in N,N-dimethylformamide (3 mL) under an argon atmosphere to produce
10 a reaction solution. The reaction solution was stirred at 80°C for 5 hours. The reaction solution was filtered over Celite, the solvent was removed under reduced pressure, and the residue was purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1 followed by ethyl acetate), to obtain the title
15 compound as a white solid (100 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.33-3.38 (1H, m), 3.52 (1H, t, J = 10.1 Hz), 3.95 (1H, q, J = 9.8 Hz), 4.58 (1H, dd, J = 10.4, 7.3 Hz), 6.51 (1H, d, J = 7.9 Hz), 6.98-7.04 (2H, m), 7.27-7.32 (2H, m), 7.81 (2H, d, J = 8.6 Hz), 8.18 (1H, s), 8.75 (1H, s).

20 MS (ESI⁺) m/z: 375 (MH⁺).


[α]_D²⁸ = -145 (c 0.1, EtOH)

[0775]

<Example 31-1>

[0776]

25 [Chemical Formula 160]



[0777]

(-)-3,5-Difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]benzamide

A 2 mol/L sodium hydroxide aqueous solution (0.21 mL) was added to a solution of (-)-1-[(3S*,4R*)-4-(4-cyano-2,6-difluorophenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea (160 mg) in methanol (4 mL) under an argon atmosphere to produce a reaction solution. The reaction solution was stirred under heating at room temperature for 1 hour. A 1 mol/L hydrochloric acid (1 mL) was added to the reaction solution to make the reaction solution acidic (pH: 1), and the mixture was extracted with ethyl acetate. The organic layer was washed with water and a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was then purified by silica gel column chromatography (hexane : ethyl acetate = 4 : 1, ethyl acetate, and ethyl acetate : methanol = 95 : 5 in turn), to obtain the title compound as a white solid (37 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.34-3.41 (1H, m), 3.52 (1H, t, J = 9.2 Hz), 3.94 (1H, q, J = 9.6 Hz), 4.62 (1H, dd, J = 10.7, 8.3 Hz), 6.52 (1H, d, J = 7.9 Hz), 6.99-7.05 (2H, m), 7.30-7.34 (2H, m),

7.58 (2H, d, J = 9.8 Hz), 7.64 (1H, s), 8.10 (1H, s), 8.14 (1H, s), 8.74 (1H, s).

MS (ESI⁻) m/z: 391 (M-H)⁻.

HRESIMS (-) : 391.10226 (calculated as C₁₈H₁₄F₃N₄O₃: 391.10180).


5 [α]_D²⁸ = -130 (c 0.1, EtOH)

[0778]

<Example 32-1>

[0779]

[Chemical Formula 161]



10 [0780]

(-)-1-(4-Fluorophenyl)-3-[(3S*, 4R*, Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea

10% Palladium on carbon (22 mg) was added to a solution of [(3S*, 4R*, Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]carbamic acid benzylester (223 mg) in ethanol (12 mL) to produce a reaction solution. The reaction solution was stirred under a hydrogen atmosphere for 1 hour. The reaction solution was filtered over Celite, and the filtrate was removed under reduced pressure to obtain

20 (3S*, 4R*, Z)-3-amino-4-(4-methoxyphenyl)pyrrolidin-2-one

O-methyloxime as a white solid. 4-Fluorophenyl isocyanate (68.3 μL) was added to a solution of the obtained

(3S*,4R*,Z)-3-amino-4-(4-methoxyphenyl)pyrrolidin-2-one

O-methyloxime (140 mg) in tetrahydrofuran (2 mL) to produce a reaction solution. The reaction solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, and the residue was then purified by silica gel column chromatography (ethyl acetate : methanol = 10 : 1), to obtain the title compound as a white solid (156 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.13 (1H, t, J = 9.1 Hz), 3.22-3.30 (1H, m), 3.48 (1H, t, J = 8.2 Hz), 3.59 (3H, s), 3.70 (3H, s), 4.82 (1H, t, J = 9.4 Hz), 6.40 (1H, d, J = 9.7 Hz), 6.69 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 6.99- 7.05 (2H, m), 7.26 (2H, d, J = 8.5 Hz), 7.32-7.36 (2H, m), 8.45 (1H, s).

MS (ESI⁺) m/z: 373 (MH⁺).

[α]_D²⁸ = -72 (c 0.31, DMSO)

[0781]

The same method as in Example 32-1 was performed using a corresponding imino substance and isocyanate to obtain the following Examples 32-2 to 32-20.

The structures and spectral data thereof are shown in Tables 101 to 107.




[0782]

[Table 101]

Ex.No	Str.	Chemical name	P.D.
32-2		(-)-1-[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.08-3.15 (1H, m), 3.40-3.49 (2H, m), 3.58 (3H, s), 3.71 (3H, s), 4.97 (1H, t, J = 9.2 Hz), 6.39 (1H, d, J = 9.2 Hz), 6.71-6.77 (3H, m), 6.97-7.03 (2H, m), 7.29-7.34 (2H, m), 7.38 (1H, dd, J = 8.6, 8.6 Hz), 8.48 (1H, s). MS (ESI ⁺) m/z: 391 (MH ⁺). [α] _D ²⁹ = -50 (c 0.10, EtOH)
32-3		(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.26-3.46 (2H, m), 3.53-3.59 (1H, m), 3.60 (3H, s), 3.74 (3H, s), 5.08 (1H, t, J = 9.2 Hz), 6.43 (1H, d, J = 9.2 Hz), 6.71 (2H, d, J = 11.0 Hz), 6.80 (1H, s), 6.98-7.03 (2H, m), 7.29-7.33 (2H, m), 8.56 (1H, s). MS (ESI ⁺) m/z: 409 (MH ⁺). [α] _D ²⁷ = -67 (c 0.10, EtOH)
32-4		(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.15 (1H, t, J = 9.1 Hz), 3.22-3.30 (1H, m), 3.48-3.58 (3H, m), 3.70 (3H, s), 3.80 (2H, t, J = 5.4 Hz), 4.45 (1H, t, J = 5.8 Hz), 4.82 (1H, t, J = 9.4 Hz), 6.39 (1H, d, J = 8.5 Hz), 6.69 (1H, s), 6.86 (2H, d, J = 9.1 Hz), 6.99-7.05 (2H, m), 7.27 (2H, d, J = 9.1 Hz), 7.31-7.36 (2H, m), 8.45 (1H, s). MS (ESI ⁺) m/z: 403 (MH ⁺). [α] _D ²⁸ = -86 (c 0.36, EtOH)




[0783]

[Table 102]

Ex.No	Str.	Chemical name	P.D.
32-5		(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethyl)imino]pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.31 (1H, m), 3.45 (1H, t, J = 9.8 Hz), 3.50-3.62 (3H, m), 3.74 (3H, s), 3.80-3.82 (2H, m), 4.47 (1H, t, J = 5.5 Hz), 5.07 (1H, dd, J = 10.4, 9.2 Hz), 6.42 (1H, d, J = 9.2 Hz), 6.71 (2H, d, J = 11.0 Hz), 6.79 (1H, s), 6.97-7.04 (2H, m), 7.26-7.34 (2H, m), 8.56 (1H, s). MS (ESI ⁺) m/z: 439 (MH ⁺). [α] _D ²⁷ = -105 (c 0.10, EtOH)
32-6		1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methylimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.68 (3H, d, J = 4.3 Hz), 3.35-3.51 (2H, m), 3.66-3.80 (4H, m), 5.04 (1H, t, J = 8.6 Hz), 5.96-6.03 (1H, m), 6.59 (1H, d, J = 9.8 Hz), 6.71 (2H, d, J = 10.4 Hz), 7.02 (2H, t, J = 9.2 Hz), 7.32-7.39 (2H, m), 8.53 (1H, s). MS (ESI ⁺) m/z: 393 (MH ⁺). [α] _D ²⁸ = -205 (c 0.10, EtOH).
32-7		(+)-1-(4-fluorophenyl)-3-[(3R*,4S*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.13 (1H, t, J = 9.1 Hz), 3.22-3.30 (1H, m), 3.48 (1H, t, J = 8.2 Hz), 3.59 (3H, s), 3.70 (3H, s), 4.82 (1H, t, J = 9.4 Hz), 6.40 (1H, d, J = 9.7 Hz), 6.69 (1H, s), 6.86 (2H, d, J = 8.5 Hz), 6.99-7.05 (2H, m), 7.26 (2H, d, J = 8.5 Hz), 7.32-7.36 (2H, m), 8.45 (1H, s). MS (ESI ⁺) m/z: 373 (MH ⁺). [α] _D ²⁷ = +100 (c 0.15, EtOH)




[0784]

[Table 103]

Ex.No	Structure	Chemical name	P.D.
32-8		(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(p-tolyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 2.18 (3H, s), 3.25-3.31 (1H, m), 3.43 (1H, t, J = 9.2 Hz), 3.53 (1H, q, J = 9.2 Hz), 3.60 (3H, s), 3.74 (3H, s), 5.07 (1H, dd, J = 9.2, 8.6 Hz), 6.38 (1H, d, J = 8.6 Hz), 6.67-6.74 (2H, m), 6.79 (1H, s), 6.97 (2H, d, J = 8.6 Hz), 7.18 (2H, d, J = 8.6 Hz), 8.40 (1H, s). MS (ESI ⁺) m/z: 405 (MH ⁺). [α] _D ²⁷ = -102 (c 0.11, EtOH)
32-9		(-)-1-(4-cyanophenyl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.34 (1H, m), 3.44 (1H, t, J = 9.2 Hz), 3.56-3.65 (1H, m), 3.60 (3H, s), 3.74 (3H, s), 5.08 (1H, dd, J = 10.4, 8.6 Hz), 6.68-6.76 (2H, m), 6.77-6.84 (2H, m), 7.49 (2H, d, J = 8.6 Hz), 7.62 (2H, d, J = 8.6 Hz), 9.18 (1H, s). MS (ESI ⁺) m/z: 416 (MH ⁺). [α] _D ²⁸ = -69 (c 0.10, EtOH)
32-10		(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.25-3.35 (1H, m), 3.43 (1H, t, J = 8.6 Hz), 3.58-3.66 (1H, m), 3.60 (3H, s), 3.74 (3H, s), 5.08 (1H, dd, J = 10.4, 8.6 Hz), 6.68-6.80 (3H, m), 6.81 (1H, s), 7.31 (1H, d, J = 8.6 Hz), 7.85 (1H, dd, J = 8.6, 3.0 Hz), 8.32 (1H, d, J = 3.0 Hz), 8.94 (1H, s). MS (ESI ⁺) m/z: 426 (MH ⁺). [α] _D ²⁶ = -66 (c 0.13, EtOH)




[0785]

[Table 104]

Ex.No	Structure	Chemical name	P.D.
32-11		(-)-1-(benzo[<i>b</i>]thiophen-2-yl)-3-[(3 <i>S</i> [*] ,4 <i>R</i> [*] , <i>Z</i>)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO- <i>d</i> ₆) δ 3.27-3.33 (1H, m), 3.60 (3H, s), 3.44 (1H, t, <i>J</i> = 9.2 Hz), 3.62-3.71 (1H, m), 3.74 (3H, s), 5.09 (1H, dd, <i>J</i> = 9.8, 9.2 Hz), 6.66 (1H, s), 6.70-6.76 (2H, m), 6.81-6.87 (2H, m), 7.08 (1H, td, <i>J</i> = 7.9, 1.2 Hz), 7.20 (1H, td, <i>J</i> = 7.9, 1.2 Hz), 7.51 (1H, d, <i>J</i> = 7.9 Hz), 7.69 (1H, d, <i>J</i> = 7.9 Hz), 9.93 (1H, s). MS (ESI ⁺) <i>m/z</i> : 447 (MH ⁺). [α] _D ²⁸ = -78 (c 0.10, EtOH)
32-12		(-)-1-[(3 <i>S</i> [*] ,4 <i>R</i> [*] , <i>Z</i>)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(<i>p</i> -tolyl)urea	¹ H-NMR (400 MHz, DMSO- <i>d</i> ₆) δ 2.18 (3H, s), 3.27-3.34 (1H, m), 3.45 (1H, t, <i>J</i> = 8.6 Hz), 3.50-3.60 (3H, m), 3.74 (3H, s), 3.79-3.82 (2H, m), 4.47 (1H, t, <i>J</i> = 6.1 Hz), 5.07 (1H, dd, <i>J</i> = 10.6, 8.6 Hz), 6.35 (1H, d, <i>J</i> = 8.6 Hz), 6.67-6.75 (2H, m), 6.78 (1H, s), 6.97 (2H, d, <i>J</i> = 8.6 Hz), 7.17 (2H, d, <i>J</i> = 8.6 Hz), 8.40 (1H, s). MS (ESI ⁺) <i>m/z</i> : 435 (MH ⁺). [α] _D ²⁸ = -52 (c 0.17, EtOH)
32-13		(-)-1-(4-cyanophenyl)-3-[(3 <i>S</i> [*] ,4 <i>R</i> [*] , <i>Z</i>)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]urea	¹ H-NMR (400 MHz, DMSO- <i>d</i> ₆) δ 3.28-3.34 (1H, m), 3.46 (1H, t, <i>J</i> = 8.6 Hz), 3.51-3.67 (3H, m), 3.74 (3H, s), 3.79-3.82 (2H, m), 4.47 (1H, t, <i>J</i> = 5.5 Hz), 5.07 (1H, dd, <i>J</i> = 10.6, 8.6 Hz), 6.68-6.77 (3H, m), 6.82 (1H, s), 7.48 (2H, d, <i>J</i> = 9.2 Hz), 7.62 (2H, d, <i>J</i> = 9.2 Hz), 9.14 (1H, s). MS (ESI ⁺) <i>m/z</i> : 446 (MH ⁺). [α] _D ²⁵ = -88 (c 0.10, EtOH)




[0786]

[Table 105]

Ex.No	Structure	Chemical name	P.D.
32-14		(-)-1-(6-chloropyridin-3-yl)-3-{{(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}}urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.32 (1H, m), 3.46 (1H, t, J = 8.6 Hz), 3.52-3.67 (3H, m), 3.74 (3H, s), 3.81 (2H, t, J = 5.5 Hz) 4.45 (1H, t, J = 5.5 Hz), 5.07 (1H, dd, J = 10.4, 8.6 Hz), 6.68-6.76 (2H, m), 6.78-6.83 (2H, m), 7.31 (1H, d, J = 8.6 Hz), 7.84 (1H, dd, J = 8.6, 3.0 Hz), 8.32 (1H, d, J = 3.0 Hz), 9.00 (1H, s). MS (ESI ⁺) m/z: 456 (MH ⁺). [α] _D ²⁷ = -71 (c 0.10, EtOH)
32-15		(-)-1-(benzo[b]thiophen-2-yl)-3-{{(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl}}urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.27-3.35 (1H, m), 3.55 (1H, t, J = 9.8 Hz), 3.52-3.60 (2H, m), 3.67 (1H, q, J = 9.8 Hz), 3.74 (3H, s), 3.81 (2H, t, J = 4.9 Hz), 4.46 (1H, t, J = 5.5 Hz), 5.09 (1H, dd, J = 10.4, 9.2 Hz), 6.66 (1H, s), 6.70-6.77 (2H, m), 6.80 (1H, brd, J = 9.2 Hz), 6.821 (1H, s), 7.08 (1H, td, J = 7.9, 1.2 Hz), 7.20 (1H, td, J = 7.9, 1.2 Hz), 7.51 (1H, d, J = 7.9 Hz), 7.68 (1H, d, J = 7.9 Hz), 9.89 (1H, s). MS (ESI ⁺) m/z: 477 (MH ⁺). [α] _D ²⁹ = -91 (c 0.12, EtOH)
32-16		(-)-1-{{(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-[(1-methyl-1H-pyrazol-3-yl)amino]-3,4-dihydro-2H-pyrrol-4-yl}}-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.50-3.73 (3H, m), 3.76 (3H, s), 3.76 (3H, s), 5.04 (1H, t, J = 8.9 Hz), 5.85 (1H, d, J = 2.4 Hz), 6.43 (1H, d, J = 7.9 Hz), 6.73 (2H, d, J = 11.0 Hz), 6.98-7.04 (2H, m), 7.30-7.36 (2H, m), 7.50 (1H, d, J = 2.4 Hz), 8.41 (1H, s), 8.72 (1H, s). MS (ESI ⁺) m/z: 459 (MH ⁺). [α] _D ²⁴ = -30 (c 0.16, EtOH)


[0787]

[Table 106]

Ex.No	Structure	Chemical name	P.D.
32-17		(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(morpholinoimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, CDCl ₃) δ 2.74-2.84 (4H, m), 3.57-3.83 (10H, m), 5.46 (1H, brs), 5.95 (1H, brs), 6.46 (2H, d, J=11.0 Hz), 6.75-6.82 (2H, m), 7.06-7.12 (2H, m), 9.11 (1H, brs). MS (ESI ⁺) m/z: 464 (MH ⁺). [α] _D ²⁴ = -31 (c 0.14, EtOH)
32-18		1-[(3R*,4S*)-3-(2,6-difluoro-4-methoxyphenyl)-5-(phenylamino)-3,4-dihydro-2H-pyrrrol-4-yl]-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.52-3.63 (2H, m), 3.75 (3H, s), 3.99-4.08 (1H, m), 5.16 (1H, t, J = 8.9 Hz), 6.72 (2H, d, J = 11.0 Hz), 6.75 (1H, d, J = 6.4 Hz), 6.88 (1H, t, J = 7.9 Hz), 7.00-7.05 (2H, m), 7.22 (2H, t, J = 7.9 Hz), 7.36-7.40 (2H, m), 7.83 (2H, d, J = 7.9 Hz), 8.56 (1H, s), 8.60 (1H, s). MS (ESI ⁺) m/z: 455 (MH ⁺).
32-19		(-)-1-((3S*,4R*,Z)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl)-3-(4-fluorophenyl)urea	¹ H-NMR (400 MHz, DMSO-d ₆) δ 3.08-3.17 (3H, m), 3.42-3.49 (2H, m), 3.54-3.56 (2H, m), 3.79-3.81 (2H, m), 4.47 (1H, t, J = 5.2 Hz), 4.54 (2H, t, J = 8.6 Hz), 4.95 (1H, t, J = 8.9 Hz), 6.60-6.70 (3H, m), 7.02 (2H, t, J = 8.9 Hz), 7.30-7.36 (3H, m), 8.73 (1H, s). MS (ESI ⁺) m/z: 433 (MH ⁺). [α] _D ²⁵ = -137 (c 0.14, EtOH)

[0788]

[Table 107]


Ex.No	Structure	Chemical name	P.D.
32-20		3-((Z)-{[(3S,4R)-4-(2,6-difluoro-4-methoxyphenyl)-3-[(4-fluorophenyl)ureido]pyrrolidin-2-ylidene}amino)propanoic acid ethyl ester	¹ H-NMR (DMSO-D ₆) δ: 1.17 (3H, t, J = 7.3 Hz), 2.65 (2H, t, J = 7.3 Hz), 3.42-3.53 (2H, m), 3.57-3.66 (1H, m), 3.75 (3H, s), 3.86-3.93 (1H, m), 3.97-4.10 (3H, m), 5.18 (1H, t, J = 8.6 Hz), 6.70 (1H, d, J = 8.6 Hz), 6.77 (2H, d, J = 11.0 Hz), 7.04 (2H, t, J = 8.6 Hz), 7.33-7.37 (2H, m), 8.99 (1H, s). MS (ESI ⁺) m/z: 479 (MH ⁺).

[0789]

5 <Example 33-1>

[0790]

[Chemical Formula 162]



[0791]

10 (-)-1-(4-Fluorophenyl)-3-[(3S*,4R*,Z)-2-(2-hydroxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea

Water (1.35 mL) and trifluoroacetic acid (135 μL) were added to a solution of

1-((3S*,4R*,Z)-2-[[tert-butyldimethylsilyl]oxy]imino)-4-(4-methoxyphenyl)pyrrolidin-3-yl)-3-(4-fluorophenyl)urea (290 mg)

15 in 1,4-dioxane (1.5 mL) to produce a reaction solution. The

reaction solution was stirred at room temperature for 30 minutes. After that, a saturated aqueous sodium hydrogen carbonate solution was added to the reaction solution under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was washed with a brine, and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure, and the residue was washed with diisopropyl ether, to obtain the title compound as a white solid (212 mg).

¹H-NMR (400 MHz, DMSO-d₆) δ 3.19 (1H, t, J = 9.1 Hz), 3.30 (1H, q, J = 8.7 Hz), 3.55 (1H, t, J = 8.2 Hz), 3.76 (3H, s), 4.80 (1H, t, J = 9.4 Hz), 6.37 (1H, d, J = 9.1 Hz), 6.43 (1H, s), 6.92 (2H, d, J = 8.5 Hz), 7.04-7.10 (2H, m), 7.32 (2H, d, J = 8.5 Hz), 7.36-7.40 (2H, m), 8.49 (1H, s), 8.89 (1H, s).

MS (ESI⁺) m/z: 359 (MH⁺).


[α]_D²⁹ = -95 (c 0.30, EtOH)

[0792]

<Example 33-2>

[0793]

[Chemical Formula 163]



[0794]

(-)-1-((3S*,4R*,Z)-4-(2,6-Difluoro-4-methoxyphenyl)-2-(hydroxylamino)-5-oxo-1-phenyl-1H-pyrazol-5-yl)-1H-pyrazole-5-carboxamide

yimino)pyrrolidin-3-yl)-3-(4-fluorophenyl)urea

The same method as in Example 33-1 was performed using
1-((3S*,4R*,Z)-2-[[tert-butyl(dimethylsilyl)oxy]imino]-4-(2,6-
difluoro-4-methoxyphenyl)pyrrolidin-3-yl)-3-(4-fluorophenyl)
5 urea in place of

1-((3S*,4R*,Z)-2-[[tert-butyl(dimethylsilyl)oxy]imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl)-3-(4-fluorophenyl)urea to obtain the title compound.

¹H-NMR (400 MHz, DMSO-d₆) δ 3.26-3.33 (1H, m), 3.42 (1H, t, J = 8.6
10 Hz), 3.49-3.57 (1H, m), 3.75 (3H, s), 5.02 (1H, t, J = 9.8 Hz),
6.33 (1H, d, J = 8.6 Hz), 6.48 (1H, s), 6.72 (2H, d, J = 10.4 Hz),
6.97-7.04 (2H, m), 7.26-7.32 (2H, m), 8.52 (1H, s), 8.90 (1H, s).
MS (ESI⁺) m/z: 395 (MH⁺).


[α]_D²⁶ = -21 (c 0.10, EtOH)

15 [0795]

< Example 34-1 >

[0796]

[Chemical Formula 164]



20 [0797]

(-)-1-{{(3S*,4R*)-1-[(2H-Tetrazol-5-yl)methyl]-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl}}-3-(4-fluorophenyl)urea

Sodium azide (22 mg), triethylamine (47 μ L), and acetic acid (19 μ L) were added to a solution of

1-[(3S*,4R*)-1-(cyanomethyl)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea (35 mg) in

5 a mixture of toluene (850 μ L) and N,N-dimethylformamide (500 μ L)

to produce a reaction solution. The reaction solution was stirred

at 125°C for 2 hours. The reaction solution was allowed to cool

to room temperature, a 10% citric acid aqueous solution was added,

and the mixture was extracted with ethyl acetate. The organic

10 layer was washed with water and a brine successively, and dried

over anhydrous sodium sulfate. The solvent was removed under

reduced pressure, and the resulting crude product was washed with

diethyl ether, to obtain the title compound as a pale brown solid

(24 mg).

15 $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ 3.12 (2H, t, J = 8.6 Hz), 3.31-3.35 (1H, m), 3.64 (1H, t, J = 9.2 Hz), 3.71-3.78 (1H, m), 4.55 (2H, t, J = 9.2 Hz), 4.70-4.76 (2H, m), 4.85-4.89 (1H, m), 6.51 (1H, d, J = 9.2 Hz), 6.64 (1H, d, J = 11.0 Hz), 7.01-7.06 (2H, m), 7.34-7.40 (3H, m), 8.67 (1H, s), 16.47 (1H, brs).

20 MS (ESI $^+$) m/z : 456(MH $^+$).

$[\alpha]_D^{26}$ = -120 (c 0.09, EtOH)

[0798]

Next, results in support of availability of the compound of the present invention will be shown with reference to Test

25 Examples.

[0799]

<Test Example 1>

Measurement test of agonist activity on human FPRL1

(1-1) Construction of human FPRL1 expression vector

5 Human FPRL (SEQ ID NO: 3) was amplified in a PCR reaction from cDNA derived from a monocytic leukemia cell line THP-1 (TIB-202, ATCC) as a template using a forward primer shown in SEQ ID NO: 1, a reverse primer shown in SEQ ID NO: 2, and KOD-plus- ver. 2 (KOD-211, TOYOBO CO., LTD.). The amplified PCR product and
10 pCMV-script vector (212220, STRATAGENE) were digested with Hind III (1060A, Takara Bio Inc.) and XhoI (1094A, Takara Bio Inc.), and the resultant digest was ligated with Ligation high ver. 2 (LGK-201, TOYOBO CO., LTD.). The ligation product was transformed into DH5 α (DNA-901, TOYOBO CO., LTD.), cultured on a 100 μ g/mL
15 kanamycin-containing LB medium, and purified with HiSpeed Plasmid Maxi Kit (12662, QIAGEN).

[0800]

(1-2) Construction of human G α 15 expression vector

Human G α 15 (SEQ ID NO: 6) was amplified in a PCR reaction
20 from cDNA derived from a myeloid leukemia cell line HL-60 (CCL-240, ATCC) as a template using a forward primer shown in SEQ ID NO: 4, a reverse primer shown in SEQ ID NO: 5, and KOD-plus- ver. 2. The amplified PCR product and pCMV-script vector were digested with Hind III and XhoI, and the resultant digest was ligated with
25 Ligation high ver. 2. The ligation product was transformed into

DH5 α , cultured on a 100 μ g/mL kanamycin-containing LB medium, and purified with HiSpeed Plasmid Maxi Kit.

[0801]

(2-1) Method for culturing and subculturing HEK293

5 HEK293 (JCRB9068, NIBIO) was cultured in an incubator at 5% CO₂ and 37°C using DMEM (11885-092, GIBCO) containing 10% FBS and 1 $\times$ Penicillin-Streptomycin (15140-122, GIBCO). Subculture was carried out as followings: The cells that reached 80 to 90% confluency were washed with PBS(-), separated using 0.25%
10 Trypsin-EDTA (25200-072, GIBCO), centrifuged, re-suspended in a fresh medium, and then seeded in Collagen Type 1 Coated dish (4020-010, IWAKI) at a split ratio of 1 : 8 (cultured for 3 days).

[0802]

(2-2) Introduction of human FPRL1 and G α 15 expression vectors

15 HEK293 that reached 80 to 90% confluency was washed with PBS(-), separated using 0.25% Trypsin-EDTA, centrifuged, and re-suspended in a fresh medium excluding
1 $\times$ Penicillin-Streptomycin. The cells were inoculated in a Collagen Type 1 coated 6-well plate (4810-010, IWAKI) to 5 $\times$ 10⁵
20 cells/2.5 mL/well and cultured overnight. On the next day, human FPRL1 and G α 15 expression vectors were introduced using Lipofectamine 2000 transfection reagent (11668-019, Life
technologies). First, the human FPRL1 and G α 15 expression vectors were diluted with Opti-MEM I Reduced Serum Medium (31985-070,
25 GIBCO) to 2 μ g/250 μ L/well and Lipofectamine 2000 transfection

reagent was diluted with Opti-MEM I Reduced Serum Medium to be 4 μ L/250 μ L/well. The vectors and reagent were softly diffused, and incubated at room temperature for 5 minutes. The vector solution was mixed with Lipofectamine 2000 transfection reagent in equal amounts. In order to form a composite of the vectors and Lipofectamine 2000 transfection reagent, the mixture was incubated at room temperature for 20 minutes, and added at 500 μ L/well to the medium of inoculated cells. The treated cells were cultured for 24 hours, inoculated in Poly-D-Lysine coated 96-well plate (356640, BD Biosciences) at a cell density of 7×10^4 cells/100 μ L/well, and cultured for another 24 hours. The resultant cells were used in a measurement test of calcium mobilization in the cells.

[0803]

(3) Evaluation of agonist activity on human FPRL1 (Test of calcium mobilization in cell)

An appropriate amount of each test compound was first weighed, and dissolved to 10^{-2} M by addition of dimethyl sulfoxide (DMSO). For calculation of an EC_{50} value for agonist activity, each compound solution was serially diluted with DMSO by 10-fold increments to make eight solutions having a concentration of 10^{-2} M to 10^{-9} M. The formed compound solution having each concentration was diluted 100 times with an assay buffer that was contained in Fluo-4 NW Calcium Assay Kit (F36206, Life technologies), and dispensed in an amount of 100 μ L into a 96-well plate with a V-bottom shape.

The plate dispensed with compound solutions was set in Flexstation (Molecular Devices, LLC.) until measurement.

Subsequently, 10 mL of assay buffer and 100 μ L of probenecid solution (dissolved by addition of 1 mL of assay buffer to a 250 mM stock) were sufficiently mixed and dissolved in Fluo-4 NW dye mix. The medium of cells inoculated on the previous day was removed, the dissolved Fluo-4 NW dye mix was added in an amount of 90 μ L/well, and a reaction was caused in the dark at 37°C for 45 minutes. The cells after the reaction and chips for addition of the compound were set in Flexstation, and variation in fluorescence intensity over time after addition of the compound was measured [amount of added compound = 10 μ L (final concentration : 10^{-5} M to 10^{-12} M), excitation wavelength: 485 nm, measured wavelength: 525 nm, 1.5 sec $\times$ 54 read]. A value was calculated by subtracting a base value during addition of DMSO from the maximum value of relative fluorescence unit, and analyzed. All the measurement data were analyzed with Prism 4 that was a data analysis tool. As an EC₅₀ value, a molar concentration that resulted in 50% maximum activation was calculated. The EC₅₀ values of the resultant test compounds are shown in Tables I to V.

[0804]

[Table I]

COMPOUND TO BE TESTED	EC ₅₀ (nM)
EXAMPLE 1-1	0.09
EXAMPLE 1-2	0.09
EXAMPLE 1-3	2.57
EXAMPLE 1-5	2.88
EXAMPLE 1-6	0.20
EXAMPLE 1-7	0.22
EXAMPLE 1-8	0.90
EXAMPLE 1-9	0.45
EXAMPLE 1-10	3.98

[0805]

[Table II]

COMPOUND TO BE TESTED	EC ₅₀ (nM)
EXAMPLE 1-11	1.59
EXAMPLE 1-12	0.06
EXAMPLE 1-13	0.15
EXAMPLE 1-14	4.12
EXAMPLE 1-16	0.24
EXAMPLE 1-18	0.36
EXAMPLE 1-19	0.72
EXAMPLE 1-20	1.75
EXAMPLE 1-22	0.11
EXAMPLE 1-23	0.78
EXAMPLE 1-24	3.20
EXAMPLE 1-26	1.91
EXAMPLE 1-31	2.67
EXAMPLE 1-32	4.54
EXAMPLE 1-35	0.76
EXAMPLE 1-39	3.51
EXAMPLE 1-40	1.44
EXAMPLE 1-41	0.73
EXAMPLE 1-43	5.13
EXAMPLE 1-53	0.18
EXAMPLE 1-54	0.11
EXAMPLE 1-55	1.02
EXAMPLE 7-1	1.09
EXAMPLE 7-2	0.45
EXAMPLE 7-7	1.56
EXAMPLE 7-8	0.95
EXAMPLE 7-9	0.16
EXAMPLE 8-1	0.29
EXAMPLE 9-1	0.86
EXAMPLE 12-1	0.13
EXAMPLE 12-4	0.05
EXAMPLE 14-1	0.32
EXAMPLE 15-1A	0.01
EXAMPLE 15-1B	1.99
EXAMPLE 17-1	0.11
EXAMPLE 18-1	2.36
EXAMPLE 19-1	0.21

[0806]

[Table III]

COMPOUND TO BE TESTED	EC ₅₀ (nM)
EXAMPLE 20-1	0.47
EXAMPLE 21-1	0.04
EXAMPLE 23-1	0.09
EXAMPLE 24-1	0.09
EXAMPLE 25-1	0.04
EXAMPLE 26-1	0.07
EXAMPLE 27-1	3.14
EXAMPLE 31-1	1.94
EXAMPLE 32-1	0.13
EXAMPLE 32-2	0.03
EXAMPLE 32-3	0.05
EXAMPLE 32-4	0.17
EXAMPLE 32-5	0.03
EXAMPLE 32-6	0.37
EXAMPLE 33-1	1.14

[0807]

[Table IV]

COMPOUND TO BE TESTED	EC ₅₀ (nM)
EXAMPLE 1-57	0.16
EXAMPLE 1-58	0.32
EXAMPLE 1—62	1.05
EXAMPLE 1-63	0.92
EXAMPLE 1-68	0.08
EXAMPLE 1-71	0.09
EXAMPLE 1-72	0.05
EXAMPLE 1-73	1.2
EXAMPLE 1-74	1.8
EXAMPLE 1-77	5.2
EXAMPLE 1-80	0.69
EXAMPLE 1-84	1.92
EXAMPLE 1-85	3.5
EXAMPLE 7-10	2.31
EXAMPLE 7-12	3.9
EXAMPLE 8-2	1.6
EXAMPLE 12-11	4.57
EXAMPLE 12-12	0.9
EXAMPLE 12-13	1.12
EXAMPLE 12-15	0.43
EXAMPLE 16-2	0.49
EXAMPLE 16-3	0.75
EXAMPLE 17-2	0.07
EXAMPLE 17-3	0.29
EXAMPLE 22-2	0.38
EXAMPLE 24-3	1.1
EXAMPLE 24-4	0.81
EXAMPLE 24-5	1.6
EXAMPLE 24-6	0.4
EXAMPLE 24-7	0.69
EXAMPLE 24-8	0.39
EXAMPLE 24-9	0.88
EXAMPLE 24-10	1.0
EXAMPLE 24-11	0.06
EXAMPLE 24-12	2.2
EXAMPLE 24-13	0.11
EXAMPLE 24-14	0.12
EXAMPLE 24-15	0.03
EXAMPLE 24-16	0.15

[0808]
[Table V]

COMPOUND TO BE TESTED	EC ₅₀ (nM)
EXAMPLE 24-17	0.02
EXAMPLE 24-18	0.03
EXAMPLE 24-19	0.33
EXAMPLE 24-20	0.15
EXAMPLE 24-21	0.45
EXAMPLE 24-22	1.32
EXAMPLE 24-23	0.8
EXAMPLE 24-24	0.46
EXAMPLE 27-2	0.1
EXAMPLE 28-4	1.64
EXAMPLE 28-5	1.42
EXAMPLE 32-8	0.44
EXAMPLE 32-9	0.3
EXAMPLE 32-10	0.53
EXAMPLE 32-11	0.88
EXAMPLE 32-12	0.27
EXAMPLE 32-13	0.25
EXAMPLE 32-14	0.82
EXAMPLE 32-15	0.27
EXAMPLE 32-16	0.54
EXAMPLE 32-17	0.57
EXAMPLE 32-18	0.54
EXAMPLE 32-19	0.13
EXAMPLE 32-20	2.86
EXAMPLE 33-2	0.03

[0809]

- 5 As seen from Tables I to V, the compounds (I) of the present invention or pharmacologically acceptable salts thereof show a superior FPRL1 agonist effect.

[0810]

<Test Example 2>

- 10 Effect of lipopolysaccharide induction on neutrophilic infiltration in mouse lung

A compound to be tested was orally administrated to a mouse

(BALB/c, male), and after 30 minutes, the mouse was placed in a plastic container. Lipopolysaccharide (0.3 mg/mL) dissolved in physiological saline was aerosolized with an ultrasonic wave nebulizer (NE-U17, OMRON Corporation), and exposed to the mouse
5 for 10 minutes. After 5 hours, the anesthetized mouse was sacrificed by exsanguination. A cannula was inserted in the respiratory tract and bronchoalveolar lavage (BAL) with 1 mL of 0.85% NaCl liquid containing 0.4% sodium citrate was carried out. This operation was repeated 3 times, to obtain a BAL fluid. The
10 BAL fluid was centrifuged at 4°C and $\times 200$ g for 5 minutes, and the pellet was suspended in a physiological saline containing 0.1% BSA. The number of white blood cells was counted using Turks solution with a microscope, and the total white blood cell count was calculated. The white blood cells were fixed on a glass slide
15 using Cytospin 3 (Thermo BioAnalysis Japan K. K.). The cells were stained with Diff-Quik (SYSMEX INTERNATIONAL REAGENTS CO., LTD.), and the number thereof was counted with a microscope, and the neutrophil ratio was calculated. The neutrophil ratio was multiplied by the total white blood cell count to calculate the
20 total neutrophil count. An effect of the compound to be tested represents a percentage (%) of suppression ratio relative to the neutrophil count in a control. The suppression ratios of the resultant test compounds are shown in Tables VI to VIII.

[0811]

[Table VI]

COMPOUND TO BE TESTED	SUPPRESSION RATIO	DOZE (mg/kg)
EXAMPLE 1-1	95.1	1
EXAMPLE 1-2	99.2	1
EXAMPLE 1-7	96.9	1
EXAMPLE 1-9	98.5	3

[0812]

[Table VII]

COMPOUND TO BE TESTED	SUPPRESSION RATIO	DOZE (mg/kg)
EXAMPLE 1-12	96.8	1
EXAMPLE 1-16	95.8	1
EXAMPLE 1-18	98.3	1
EXAMPLE 1-22	93.1	1
EXAMPLE 1-23	98.9	3
EXAMPLE 1-31	69.7	1
EXAMPLE 1-53	78.9	1
EXAMPLE 1-54	98.6	3
EXAMPLE 1-55	83.7	3
EXAMPLE 12-1	97.4	3
EXAMPLE 25-1	99.1	3
EXAMPLE 26-1	99.6	1
EXAMPLE 32-1	99.0	3
EXAMPLE 32-3	99.1	3
EXAMPLE 32-5	99.8	1

5 [0813]

[Table VIII]

COMPOUND TO BE TESTED	SUPPRESSION RATIO (%)	DOZE(mg/kg)
EXAMPLE 1-68	85.6	3
EXAMPLE 1-71	79.7	3
EXAMPLE 1-72	93.3	3
EXAMPLE 1-84	99.2	1
EXAMPLE 7-2	87.7	1
EXAMPLE 9-1	57.0	3
EXAMPLE 32-12	96.8	1

[0814]

As seen from Tables VI to VIII, the compounds (I) of the present invention or pharmacologically acceptable salts thereof had a superior action of suppressing neutrophil infiltration.

5 Industrial Applicability

[0815]

The compound of the present invention has a superior action of suppressing neutrophil infiltration due to a superior FPRL1 agonist effect, and therefore is useful as a therapeutic or
10 prophylactic agent for inflammatory disease, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, immune disorders and the like.

15 Sequence Listing Free Text

[0816]

<Sequence Listing 1>

SEQ ID NO: 1 is a sequence of a forward primer used for amplification of DNA of human FPRL1 (SEQ ID NO: 3), and is
20 supplemented with a Hind III recognition site.

<Sequence Listing 2>

SEQ ID NO: 2 is a sequence of a reverse primer used for amplification of DNA of human FPRL1 (SEQ ID NO: 3), and is supplemented with an XhoI recognition site.

25 <Sequence Listing 3>

SEQ ID NO: 3 is an open reading frame (ORF) of human FPRL1,
and is a DNA sequence of a site translated into an amino acid.

<Sequence Listing 4>

5 SEQ ID NO: 4 is a sequence of a forward primer used for
amplification of DNA of human G α 15 (SEQ ID NO: 6), and is supplemented
with a Hind III recognition site.

<Sequence Listing 5>

10 SEQ ID NO: 5 is a sequence of a reverse primer used for
amplification of DNA of human G α 15 (SEQ ID NO: 6), and is supplemented
with an XhoI recognition site.

<Sequence Listing 6>

SEQ ID NO: 6 is an open reading frame (ORF) of human G α 15,
and is a DNA sequence of a site translated into an amino acid.

12737 Sequence listing.txt
SEQUENCE LISTING

<110> KYORIN PHARMACEUTICAL CO., LTD.
<120> UREA DERIVATIVES, OR THE PHARMACEUTICALLY ACCEPTABLE SALTS
<130> 201315
<160> 6
<170> PatentIn version 3.1
<210> 1
<211> 43
<212> DNA
<213> Artificial
<220>
<223> 5' primer
<400> 1
cgaagcttca@ccatggaaac@caacttctcc@actcctctga@atg@ @ 43

<210> 2
<211> 38
<212> DNA
<213> Artificial
<220>
<223> 3' primer
<400> 2
cgctcgagtc@atattgcctt@tatttcaatg@tcttcagg @ 38

<210> 3
<211> 1056
<212> DNA
<213> Human
<400> 3
atggaaacca@acttctccac@tcctctgaat@gaatatgaag@aagtgtccta@tgagtctgct@ 60@
ggctacactg@ttctgcggat@cctcccattg@gtgggtgcttg@ggggtcacctt@tgctcctcggg 120@
gtcctgggca@atgggcttgt@gatctgggtg@gctggattcc@ggatgacacg@cacagtcacc@ 180
accatctgtt@acctgaacct@gggcctggct@gacttttctt@tcacggccac@attaccattc@ 240
ctcattgtct@ccatggccat@gggagaaaaa@tggccttttg@gctggttcct@gtgtaagtta@ 300
attcacatcg@tggtggacat@caacctcttt@ggaagtgtct@tcttgattgg@tttcattgca@ 360
ctggaccgct@gcatttgtgt@cctgcatcca@gtctggggccc@agaaccaccg@cactgtgagt@ 420
ctggccatga@aggtgatcgt@cggaccttgg@attcttgctc@tagtccttac@cttgccagtt@ 480
ttcctctttt@tgactacagt@aaactattcca@aatggggaca@catactgtac@tttcaacttt@ 540
gcatcctggg@gtggcaccac@tgaggagagg@ctgaagggtgg@ccattaccat@gctgacagcc@ 600
agagggatta@tccggtttgt@cattggcttt@agcttgccga@tgtccattgt@tgccatctgc@ 660
tatgggctca@ttgcagccaa@gatccacaaa@aagggcatga@ttaaatccag@ccgtccctta@ 720
cgggtcctca@ctgctgtggt@ggcttctttc@ttcatctgtt@ggtttccctt@tcaactggtt@ 780

12737 Sequence listing.txt

gcccttctgg@gcacgtctg@gctcaaagag@atgttggtct@atggcaagta@caaaatcatt@ 840
gacatcctgg@ttaaccaac@gagctccctg@gccttcttca@acagctgcct@caaccccatg@ 900
ctttacgtct@ttgtggggcca@agacttccga@gagagactga@tccactccct@gcccaccagt@ 960
ctggagaggg@ccctgtctga@ggactcagcc@ccaactaatg@acacggctgc@caattctgct@1020
tcacctcctg@cagagactga@gttacaggca@atgtga 1056

<210> 4
<211> 29
<212> DNA
<213> Artificial

<220>
<223> 5' primer

<400> 4
cgaagcttca@ccatggcccg@ctcgctgac

29

<210> 5
<211>
<212> DNA
<213> Artificial

<400> 5
cgctcgagtc@acagcagggt@gatctcgctcc@@@@@@@@@@@@@@@@@@@@@30

<210> 6
<211> 1125
<212> DNA
<213> Human

<400> 6
atggcccgct@cgctgacctg@gcgctgctgc@ccctggtgcc@tgacggagga@tgagaaggcc@@60
@@
gccgcccggg@tgaccagga@gatcaacagg@atcctcttgg@agcagaagaa@gcaggaccgc 120
@
ggggagctga@agctgctgct@tttggggcca@ggcgagagcg@ggaagagcac@cttcatcaag@ 180
cagatgcgga@tcatccacgg@cgccggctac@tcggaggagg@agcgcaaggg@cttccggccc@ 240
ctggtctacc@agaacatctt@cggtgcatg@cgggccatga@tcgaggccat@ggagcggctg@ 300
cagattccat@tcagcaggcc@cgagagcaag@caccacgcta@gcctggtcat@gagccaggac@ 360
ccctataaag@tgaccacgtt@tgagaagcgc@tacgtgcgg@ccatgcagtg@gctgtggagg@ 420
gatgccggca@tccgggccta@ctatgagcgt@cggcgggaat@tccacctgct@cgattcagcc@ 480
gtgtactacc@tgtccacct@ggagcgcata@accgaggagg@gctacgtccc@cacagctcag@ 540
gacgtgctcc@gcagccgcat@gcccaccact@ggcatcaacg@agtactgctt@ctccgtgcag@ 600
aaaaccaacc@tgcggtcgt@ggacgtcggg@ggccagaagt@cagagcgtaa@gaaatggatc@ 660
cattgtttcg@agaacgtgat@cgccctcatc@tacctggcct@actgagtga@atacgaccag@ 720
tgcctggagg@agaacaacca@ggagaaccgc@atgaaggaga@gcctcgatt@gtttgggact@ 780
atcctggaac@taccctgggt@caaaagcaca@tccgtcatcc@tctttctcaa@caaaaccgac@ 840
atcctggagg@agaaaatccc@cacctcccac@ctggctacct@atttcccag@tttccagggc@ 900


12737 Sequence listing.txt

cctaagcagg@atgctgaggc@agccaagagg@ttcatcctgg@acatgtacac@gaggatgtac@ 960
accgggtgcg@tggacggccc@cgagggcagc@aagaagggcg@cacgatcccg@acgcctcttc@1020
agccactaca@catgtgccac@agacacacag@aacatccgca@aggtcttcaa@ggacgtgcgg@1080
gactcgggtgc@tcgcccgcta@cctggacgag@atcaacctgc@tgtga 1125

CLAIMS

1. A compound represented by the general formula (I) or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof:

[Chemical Formula 1]



wherein, in the formula (I), Ar¹ is a phenyl group optionally having substituent(s), a 5-membered aromatic heterocyclic group optionally having substituent(s), a 6-membered aromatic heterocyclic group optionally having substituent(s), or a bicyclic aromatic heterocyclic group having 8 or 9 atoms and optionally having substituent(s);

Ar² is a phenyl group optionally having substituent(s) (except for a phenyl group substituted only with halogen atom(s)), a 5-membered aromatic heterocyclic group optionally having substituent(s), a 6-membered aromatic heterocyclic group optionally having substituent(s), or a bicyclic aromatic heterocyclic group having 8 or 9 atoms and optionally having substituent(s);

X is a group selected from the group consisting of the following a), b), and c),

a) an oxygen atom or a sulfur atom,

b) NR^4 , and

c) NOR⁴, wherein,

when X is b) or c), R⁴ is a hydrogen atom, a phenyl group optionally having substituent(s), a heterocyclic group optionally having substituent(s), or a C₁ to C₆ alkyl group optionally having substituent(s);

R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₆ alkoxy group optionally having substituent(s), or a C₁ to C₆ alkyl group optionally having substituent(s);

R² and R³ are each independently a hydrogen atom or a C₁ to C₆ alkyl group optionally having substituent(s) or together form a C₂ to C₆ alkylene group; and

each carbon atom marked with an asterisk is an asymmetric carbon atom;

wherein the term “optionally having substituent(s)” means optionally substituted with one or more substituents selected from the group consisting of:


halogen atoms, an amino group, a hydroxy group, a cyano group, a nitro group, a carboxy group, C₁ to C₆ alkoxy carbonyl groups, a formyl group, C₁ to C₆ acyl groups, C₁ to C₆ alkyl groups, C₁ to C₆ alkylamino groups, di-C₁ to C₆ alkylamino groups, C₁ to C₆ alkoxy groups, C₁ to C₆ alkylthio groups, C₃ to C₆ cycloalkyl groups, 4- to 10-membered heterocycloalkyl groups, aromatic hydrocarbon cyclic groups optionally having a halogen atom, aromatic heterocyclic groups, C₁ to C₆ alkylcarbonylamino groups, C₃ to C₆ cycloalkylcarbonylamino groups, 4- to 10-membered heterocycloalkylcarbonylamino groups, aromatic hydrocarbon cyclic carbonylamino groups, and aromatic heterocyclic carbonylamino groups,

with the proviso that when the group optionally having substituent(s) is selected from the group consisting of phenyl, a 5-membered aromatic heterocyclic group, a 6-membered aromatic heterocyclic group, a bicyclic aromatic heterocyclic group having 8 or 9 atoms and a heterocyclic group, the optional substituent is not a di-C₁ to C₆ alkylamino group.

2. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the term “optionally having substituent(s)” has the meaning defined in claim 1, and wherein

in the formula (I), Ar² is a group selected from the group consisting of the following A1), A2), A3), A4), A5), A6), A7), A8), A9), and A10):

[Chemical Formula 2]



wherein, when Ar² is A2), W¹ is a nitrogen atom or CH optionally substituted with a hydrogen atom, with a halogen atom, or with a C₁ to C₆ alkyl group optionally having substituent(s);

when Ar² is A2), W² is CH or a nitrogen atom;

when Ar² is A3), A4), A5), or A6), W³ is an oxygen atom, a sulfur atom, or NH optionally substituted with a C₁ to C₆ alkyl group;

when Ar² is A3), A4), or A6), W⁴ is CH or a nitrogen atom;

when Ar² is A7), W⁵ is CH₂, an oxygen atom, or a sulfur atom;

when Ar² is A7), W⁶ is C=O, CH₂, CF₂, CHOH, NH optionally substituted with a C₁ to C₆ alkyl group, SO, SO₂, an oxygen atom, or a sulfur atom;

when Ar² is A8), W⁷ is NH optionally substituted with a C₁ to C₆ alkyl group or C=O;

when Ar² is A8), W⁸ is C=O with W⁷ being NH optionally substituted with a C₁ to C₆ alkyl group and W⁸ is NH optionally substituted with a C₁ to C₆ alkyl group with W⁷ being C=O;

when Ar² is A10), W⁹ is a nitrogen atom or N=O;

when Ar² is A1), A2), A3), A4), or A5), R⁶ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl group optionally having substituent(s), a heterocyclic group

optionally having substituent(s), $-\text{CONR}^{10}\text{R}^{11}$, or $-\text{NR}^{10}\text{R}^{11}$, wherein, when R^6 is $-\text{CONR}^{10}\text{R}^{11}$ or $-\text{NR}^{10}\text{R}^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), a C_1 to C_6 acyl group optionally having substituent(s), or a C_1 to C_6 alkylsulfonyl group optionally having substituent(s) and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group optionally having substituent(s), or R^{10} and R^{11} together form a C_3 to C_{10} heterocycloalkyl group;

when Ar^2 is A1), A2), A3), A4), A5), A6), A7), A8), A9), or A10), R^7 is a hydrogen atom, a halogen atom, a C_1 to C_6 alkyl group optionally having substituent(s), or a C_1 to C_6 alkoxy group optionally having substituent(s);

when Ar^2 is A1), A7), A8), or A10), R^8 is a hydrogen atom, a halogen atom, or a C_1 to C_6 alkyl group optionally having substituent(s);

when Ar^2 is A9), R^9 is a hydrogen atom or a C_1 to C_6 alkyl group;


when Ar^2 is A7), m is 0 or 1; and

when Ar^2 is A8), n is 0 or 1;

provided that, when Ar^2 is A1), combinations of the substituents R^6 , R^7 and R^8 exclude a combination of a hydrogen atom and a halogen atom.

3. The compound according to claim 2 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the term “optionally having substituent(s)” has the meaning defined in claim 1, and wherein in the formula (I), Ar^1 is a group selected from the group consisting of the following B1), B2), B3), B4), B5), B6), B7), B8), B9), B10), B11), and B12):

[Chemical Formula 3]



wherein, when Ar¹ is B2), B3), B7), B8), B10), B11), or B12), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₃ to C₆ cycloalkyl group optionally having substituent(s), a C₃ to C₆ cycloalkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), a C₂ to C₆ alkenyl group optionally having substituent(s), a C₂ to C₆ alkynyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl group optionally having substituent(s), -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), or a C₁ to C₆ alkylsulfonyl group optionally having substituent(s) and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group optionally having substituent(s), or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group;

when Ar¹ is B1), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ alkoxy group optionally having substituent(s), a C₃ to C₆ cycloalkyl group optionally having substituent(s), a C₃ to C₆ cycloalkoxy group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), a C₂ to C₆ alkenyl group optionally having substituent(s), a C₂ to C₆ alkynyl group, a C₁ to C₆ alkoxycarbonyl

group, a C₁ to C₆ alkylsulfanyl group optionally having substituent(s), a C₁ to C₆ alkylsulfinyl group optionally having substituent(s), a C₁ to C₆ alkylsulfonyl group optionally having substituent(s), -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group optionally having substituent(s), a C₁ to C₆ acyl group optionally having substituent(s), or a C₁ to C₆ alkylsulfonyl group optionally having substituent(s) and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group optionally having substituent(s), or R¹⁰ and R¹¹ together form a C₃ to C₁₀ heterocycloalkyl group, and R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group, or R¹² and R¹³ may together form a C₃ to C₅ alkylene group or a C₁ to C₂ alkylendioxy group;

when Ar¹ is B6), R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group;

when Ar¹ is B9), R¹⁴ and R¹⁵ are each independently a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group, or a C₁ to C₆ alkoxy group;

when Ar¹ is B2), one of W¹⁰ and W¹¹ is a nitrogen atom, and the other thereof is CH or a nitrogen atom;

when Ar¹ is B6), B7), B8), B9), B10), B11), or B12), W¹² is an oxygen atom, a sulfur atom, or N-R¹⁶, wherein, when W¹² is an N-R¹⁶, R¹⁶ is a hydrogen atom or a C₁ to C₆ alkyl group; and

when Ar¹ is B6), B7), B8), or B9), W¹³ is CH or a nitrogen atom.

4. The compound according to claim 3 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the term “optionally having substituent(s)” has the meaning defined in claim 1, and wherein

when Ar² is A1), A2), A3), A4), or A5), R⁶ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ acyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, a heterocyclic group optionally having substituent(s), -CONR¹⁰R¹¹, or -NR¹⁰R¹¹, wherein, when R⁶

is $-\text{CONR}^{10}\text{R}^{11}$ or $-\text{NR}^{10}\text{R}^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group, a C_1 to C_6 acyl group, or a C_1 to C_6 alkylsulfonyl group and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group, or R^{10} and R^{11} together form a C_3 to C_{10} heterocycloalkyl group;

when Ar^2 is A1), A2), A3), A4), A5), A6), A7), A8), A9), or A10), R^7 is a hydrogen atom, a halogen atom, a C_1 to C_6 alkyl group, or a C_1 to C_6 alkoxy group;

when Ar^2 is A1), A7), A8), or A10), R^8 is a hydrogen atom, a halogen atom, or a C_1 to C_6 alkyl group; and

when Ar^2 is A9), R^9 is a hydrogen atom or a C_1 to C_6 alkyl group;

provided that, when Ar^2 is A1), combinations of the substituents R^6 , R^7 and R^8 exclude a combination of a hydrogen atom and a halogen atom.

5. The compound according to claim 4 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the term “optionally having substituent(s)” has the meaning defined in claim 1, and wherein R^1 is a hydrogen atom, a hydroxy group, a C_1 to C_6 alkyl group, a C_1 to C_6 alkoxy group, a halo- C_1 to C_6 alkyl group, a hydroxy C_1 to C_6 alkyl group, a carboxy C_1 to C_6 alkyl group, a carbamoyl C_1 to C_6 alkyl group, a mono C_1 to C_6 alkylcarbamoyl C_1 to C_6 alkyl group, a di- C_1 to C_6 alkylcarbamoyl C_1 to C_6 alkyl group, an aminosulfonyl C_1 to C_6 alkyl group, an aromatic heterocyclic C_1 to C_3 alkyl group optionally having substituent(s), or a phenyl C_1 to C_3 alkyl group optionally having substituent(s);

R^2 and R^3 are each independently a hydrogen atom or a C_1 to C_3 alkyl group or together form a C_2 to C_6 alkylene group;

when Ar^1 is B2), B3), B7), B8), B10), B11), or B12), R^{12} is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C_1 to C_6 alkyl group, a halo- C_1 to C_6 alkyl group, a hydroxy C_1 to C_6 alkyl group, a C_1 to C_6 alkoxy group, a C_3 to C_6 cycloalkyl group, a C_3 to C_6 cycloalkoxy group, a C_1 to C_6 acyl group, a C_2 to C_6 alkenyl group, a C_2 to C_6 alkynyl group, a C_1 to C_6 alkoxycarbonyl group, a C_1 to C_6 alkylsulfanyl group, a C_1 to C_6 alkylsulfinyl group, a C_1 to C_6 alkylsulfonyl group, $-\text{CONR}^{10}\text{R}^{11}$, $-\text{NR}^{10}\text{R}^{11}$, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R^{12} is $-\text{CONR}^{10}\text{R}^{11}$ or $-\text{NR}^{10}\text{R}^{11}$, R^{10} is a hydrogen

atom, a C₁ to C₆ alkyl group, a C₁to C₆ acyl group, or a C₁to C₆ alkylsulfonyl group and R¹¹ is a hydrogen atom or a C₁to C₆ alkyl group, or R¹⁰ and R¹¹ together form a C₃to C₁₀ heterocycloalkyl group;

when Ar¹ is B1), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁to C₆ alkyl group, a halo-C₁to C₆ alkyl group, a hydroxy C₁to C₆ alkyl group, a C₁to C₆ alkoxy group, a C₃to C₆ cycloalkyl group, a C₃to C₆ cycloalkoxy group, a C₁to C₆ acyl group, a C₂to C₆ alkenyl group, a C₂to C₆ alkynyl group, a C₁to C₆ alkoxycarbonyl group, a C₁to C₆ alkylsulfanyl group, a C₁to C₆ alkylsulfinyl group, a C₁to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁to C₆ alkyl group, a C₁to C₆ acyl group, or a C₁to C₆ alkylsulfonyl group and R¹¹ is a hydrogen atom or a C₁to C₆ alkyl group, or R¹⁰ and R¹¹ together form a C₃to C₁₀ heterocycloalkyl group, and R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁to C₆ alkyl group, or R¹² and R¹³ may together form a C₃to C₅ alkylene group or a C₁to C₂ alkylenedioxy group; and

X is a group selected from the group consisting of the following a), b), and c),

a) an oxygen atom,

b) NR⁴, and


c) NOR⁴, wherein,

when X is b) or c), R⁴ is a hydrogen atom, a hydroxy C₁to C₆ alkyl group, a phenyl group, a heterocyclic group, or a C₁to C₆ alkyl group.

6. The compound according to claim 5 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar² is a group selected from the group consisting of A1), A2a), A3), and A7a):

[Chemical Formula 4]



wherein, when Ar² is A2a), W² is the same as that defined in claim 2 when Ar² is A2);

when Ar² is A3), W⁴ is the same as that defined in claim 2 when Ar² is A3);

when Ar² is A7a), W⁶ is the same as that defined in claim 2 when Ar² is A7);

when Ar² is A1), A2a), A3), or A7a), R⁷ is the same as that defined in claim 4 when Ar² is A1), A2), A3), or A7);

when Ar² is A1) or A7a), R⁸ is the same as that defined in claim 4 when Ar² is A1) or A7);

when Ar² is A7a), m is the same as that defined in claim 2 when Ar² is A7);

when Ar² is A1), A2a), or A3), R⁶ is a hydrogen atom, a halogen atom, a cyano group, a C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a halo-C₁ to C₆ alkoxy group, a C₁ to C₆ acyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, -CONR¹⁰R¹¹, or -NR¹⁰R¹¹, wherein, when R⁶ is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group; and


when Ar² is A3), W³ is an oxygen atom or a sulfur atom;

provided that, when Ar² is A1), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

7. The compound according to claim 6 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the term “optionally having substituent(s)” has the meaning defined in claim 1, and wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B2), B6a), B9a), and B9b):

[Chemical Formula 5]



wherein, when Ar¹ is B6a), R¹³ is the same as that defined in claim 3 when Ar¹ is B6);

when Ar¹ is B9a) or B9b), R¹⁴ and R¹⁵ are the same as those defined in claim 3 when Ar¹ is B9);

when Ar¹ is B2), W¹⁰ and W¹¹ are the same as those defined in claim 3 when Ar¹ is B2);

when Ar¹ is B2), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to C₆ alkenyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group;

when Ar¹ is B1), R¹² is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, a C₁ to C₆ alkyl group, a halo-C₁ to C₆ alkyl group, a hydroxy C₁ to C₆ alkyl group, a C₁ to C₆ alkoxy group, a C₃ to C₆ cycloalkyl group, a C₃ to C₆ cycloalkoxy group, a C₁ to C₆ acyl group, a C₂ to C₆ alkenyl group, a C₁ to C₆ alkoxycarbonyl group, a C₁ to C₆ alkylsulfanyl group, a C₁ to C₆ alkylsulfinyl group, a C₁ to C₆ alkylsulfonyl group, -CONR¹⁰R¹¹, -NR¹⁰R¹¹, an aryloxy group, or a heterocyclic group optionally having substituent(s), wherein, when R¹² is -


CONR¹⁰R¹¹ or -NR¹⁰R¹¹, R¹⁰ is a hydrogen atom, a C₁ to C₆ alkyl group, or a C₁ to C₆ acyl group and R¹¹ is a hydrogen atom or a C₁ to C₆ alkyl group, or R¹⁰ and R¹¹ may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group, and R¹³ is a hydrogen atom, a halogen atom, a hydroxy group, a cyano group, or a C₁ to C₆ alkyl group, or R¹² and R¹³ may together form a C₃ to C₅ alkylene group or a C₁ to C₂ alkylenedioxy group; and

when Ar¹ is B6a), B9a), or B9b), W¹² is an oxygen atom or a sulfur atom.

8. The compound according to claim 7 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar² is a group selected from the group consisting of A1a), A2b), A3a), and A7b):

[Chemical Formula 6]



wherein, when Ar² is A1a), R⁶ is the same as that defined in claim 6 when Ar² is A1);

when Ar² is A1a) or A7b), R⁸ is the same as that defined in claim 4 when Ar² is A1) or A7);

when Ar² is A7b), m is the same as that defined in claim 2 when Ar² is A7);

R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₃ alkyl group, a C₁ to C₃ alkoxy group, a hydroxy C₁ to C₄ alkyl group, a carboxy C₁ to C₃ alkyl group, a carbamoyl C₁ to C₃ alkyl group, a mono-C₁ to C₂ alkylcarbamoyl C₁ to C₃ alkyl group, or a di-C₁ to C₂ alkylcarbamoyl C₁ to C₃ alkyl group;

R² and R³ are each independently a hydrogen atom or a C₁ to C₃ alkyl group;

when X is b) or c), R⁴ is a hydrogen atom, a hydroxy C₁ to C₄ alkyl group, or a C₁ to C₃ alkyl group;

when Ar^2 is A2b) or A3a), R^{6a} is a C_1 to C_3 alkoxy group;

when Ar^2 is A1a), A2b), A3a), or A7b), R^7 is a hydrogen atom, a fluorine atom, a chlorine atom, or a C_1 to C_3 alkyl group; and


when Ar^2 is A7b), W^6 is $\text{C}=\text{O}$, CH_2 , CF_2 , CHOH , or an oxygen atom;

provided that, when Ar^1 is A1a), combinations of the substituents R^6 , R^7 and R^8 exclude a combination of a hydrogen atom and a halogen atom.

9. The compound according to claim 8 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar^2 is a group selected from the group consisting of A1a), A2b), and A7c):

[Chemical Formula 7]



wherein, when Ar^2 is A2b), R^{6a} is the same as that defined in claim 8 when Ar^2 is A2b);

when Ar^2 is A1a) or A7c), R^8 is the same as that defined in claim 4 when Ar^2 is A1) or A7);

when Ar^2 is A7c), m is the same as that defined in claim 8 when Ar^2 is A7b);

when Ar^2 is A1a), R^6 is a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, a C_1 to C_6 alkyl group, a C_1 to C_6 alkoxy group, a halo- C_1 to C_6 alkoxy group, a C_1 to C_6 alkylsulfanyl group, a C_1 to C_6 alkylsulfinyl group, or $-\text{CONR}^{10}\text{R}^{11}$, wherein, when R^6 is $-\text{CONR}^{10}\text{R}^{11}$, R^{10} is a hydrogen atom, a C_1 to C_6 alkyl group, or a C_1 to C_6 acyl group and R^{11} is a hydrogen atom or a C_1 to C_6 alkyl group, or R^{10} and R^{11} may together form a pyrrolidinyl group, a piperidinyl group, a piperazinyl group, or a morpholinyl group; and


when Ar² is A1a), A2b), or A7c), R⁷ is a hydrogen atom, a fluorine atom, or a chlorine atom;

provided that, when Ar² is A1a), combinations of the substituents R⁶, R⁷ and R⁸ exclude a combination of a hydrogen atom and a halogen atom.

10. The compound according to claim 9 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B2), B6b), B9c), and B9d):

[Chemical Formula 8]



wherein R¹ is a hydrogen atom, a hydroxy group, a C₁ to C₃ alkyl group, a C₁ to C₃ alkoxy group, or a hydroxy C₁ to C₄ alkyl group;

when Ar¹ is B1) or B2), R¹² is a hydrogen atom, a fluorine atom, a chlorine atom, a cyano group, a C₁ to C₃ alkyl group, or a C₁ to C₆ alkoxy group;

when Ar¹ is B1) or B6b), R¹³ is a hydrogen atom, a hydroxy group, a fluorine atom, or a chlorine atom;

when Ar¹ is B9c) or B9d), R¹⁴ is a hydrogen atom, a fluorine atom, a chlorine atom, a C₁ to C₃ alkyl group, a methoxy group, or an ethoxy group;


when Ar¹ is B9c) or B9d), R¹⁵ is a hydrogen atom, a fluorine atom, or a chlorine atom; and

when Ar¹ is B2), one of W¹⁰ and W¹¹ is N, and the other thereof is CH.

11. The compound according to claim 10 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar¹ is a group selected from the group consisting of B1), B6b), and B9c1):

[Chemical Formula 9]



wherein when Ar¹ is B1), R¹² is the same as that defined in claim 10 when Ar¹ is B1);


when Ar¹ is B1) or B6b), R¹³ is the same as that defined in claim 10 when Ar¹ is B1) or B6b); and

when Ar¹ is B9c1), R¹⁴ is the same as that defined in claim 10 when Ar¹ is B9c).

12. The compound according to claim 11 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

in the formula (I), Ar² is a group selected from the group consisting of A1b), A2b), and A7c):

[Chemical Formula 10]



wherein, when Ar² is A2b), R^{6a} is the same as that defined in claim 8 when Ar² is A2b);

when Ar² is A7c), m is the same as that defined in claim 8 when Ar² is A7b);

when Ar² is A1b), R⁶ is a cyano group, an ethyl group, or a C₁ to C₃ alkoxy group;

R⁷ is a fluorine atom or a chlorine atom; and

when Ar² is A1b) or A7c), R⁸ is a hydrogen atom, a fluorine atom, a chlorine atom, or a C₁ to C₃ alkyl group.

13. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein

the compound represented by the formula (1) is

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(3,4-difluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(3-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

1-[(3S*,4R*)-4-(2-chloro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methoxy-2-methylphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(3,5-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(3-fluoro-5-methoxypyridin-2-yl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(5-methoxythiophen-2-yl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-4-(4-methylthiophenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-chlorophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-methoxyphenyl)urea,

(-)-1-(5-chlorothiazol-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(6-chloropyridin-3-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(5-chloropyridin-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(benzo[b]thiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(5-methylthiophen-2-yl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5S*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,5R*)-4-(4-methoxyphenyl)-5-methyl-2-oxopyrrolidin-3-yl]urea,

(+)-1-(4-chlorophenyl)-3-[(3R*,4S*)-2-oxo-4-phenylpyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-fluorophenyl)-3-[4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[4-(4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-[4-(trifluoromethoxy)phenyl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-phenylurea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-phenoxyphenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3,4-difluorophenyl)urea,

(-)-1-(5-chlorothiophen-2-yl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(p-tolyl)urea,

(-)-1-[(3S*,4R*)-1-ethyl-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*)-1-(1-hydroxy-2-methylpropan-2-yl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(±)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(+)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-trans-1-(4-chlorophenyl)-3-[1-(2-hydroxyethyl)-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylacetamide,

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer A),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer A),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]propionic acid (isomer B),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N-methylpropionamide (isomer B),

(-)-2-[(3S*,4R*)-3-[3-(4-fluorophenyl)ureido]-4-(4-methoxyphenyl)-2-oxopyrrolidin-1-yl]-N,2-dimethylpropionamide,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-methyl-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2-fluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(+)-1-(4-fluorophenyl)-3-[(3R*,4S*)-1-methoxy-4-(4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-3,5-difluoro-4-[(3R*,4S*)-4-[3-(4-fluorophenyl)ureido]-5-oxopyrrolidin-3-yl]benzamide,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(methoxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*,Z)-4-(2-fluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methoxyimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-[(2-hydroxyethoxy)imino]-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-(methylimino)pyrrolidin-3-yl]-3-(4-fluorophenyl)urea,

(-)-1-(4-fluorophenyl)-3-[(3S*,4R*,Z)-2-(2-hydroxyimino)-4-(4-methoxyphenyl)pyrrolidin-3-yl]urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(3-hydroxy-4-methylphenyl)urea,

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]-3-(4-fluoro-3-hydroxyphenyl)urea,


(-)-1-(4-chloro-3-hydroxyphenyl)-3-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-2-oxopyrrolidin-3-yl]urea,

(-)-1-(4-cyanophenyl)-3-[(3S*,4R*)-4-(6-fluoro-2,3-dihydrobenzofuran-5-yl)-2-oxopyrrolidin-3-yl]urea,


(-)-1-[(3S*,4R*,Z)-4-(2,6-difluoro-4-methoxyphenyl)-2-[(2-hydroxyethoxy)imino]pyrrolidin-3-yl]-3-(p-tolyl)urea, or

(-)-1-[(3S*,4R*)-4-(2,6-difluoro-4-methoxyphenyl)-1-(2-hydroxyethyl)-2-oxopyrrolidin-3-yl]-3-phenylurea.


14. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




15. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




16. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




17. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




18. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




19. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




20. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




21. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




22. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:




23. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:



24. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:



25. The compound according to claim 1 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, wherein the compound is:



26. A pharmaceutical comprising, as an active ingredient, the compound according to any one of claims 1 to 25 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof.

27. A pharmaceutical composition containing the compound according to any one of claims 1 to 25 or a pharmacologically acceptable salt thereof, or a solvate or hydrate thereof, and a pharmaceutically acceptable carrier, for use in the prevention or treatment of inflammatory diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders.