

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5

T Cell Mediated Lysis of Prostate Cancer Cell Line
22Rv1 with EGFR Targeting T Cell Engagers

T Cell Mediated Lysis of Prostate Cancer Cell Line
22Rv1 with PSMA Targeting T Cell Engagers



	Trispecific (EGFR)	BITE (EGFR)	Trispecific (PSMA)	BITE (PSMA)
EC ₅₀ (pM)	3.9	6.4	113	189

Figure 6




Figure 7





Figure 8


Activity of EGFR-scFv Trispecific and EGFR BITE
in an NCI-1563 TDCC Assay



	EGFR-scFv.C:A	EGFR BITE
EC50 (M)	1.6E-12	1.3E-12
EC50 with HSA (M)	1.3E-11	1.7E-12
Fold change	7.8	1.3

Figure 9

Activiy of Fynomer Based, Her2 Targeting
Trispecific in SK-OV-3 TDCC Assay



	Trispecific (Her2)
EC ₅₀ (nM)	4.1

Figure 10





Figure 11



	EC50 TDCC (M)	EC50 TNF-α (M)	EC50 IFN-γ (M)
αPSMA:αCD3:αALB (P.C.A)	1.7E-10	2.6E-10	2.3E-10
αPSMA:αALB:αCD3 (P.A.C)	6.9E-11	1.5E-10	1.3E-10

Figure 12




Figure 13

