



159159

QUINOLONE CARBOXYLIC ACIDS, DERIVATIVES THEREOF, PROCESSES FOR ITS PREPARATION AND ITS USE AS ANTIBACTERIAL AGENTS

BACKGROUND OF THE INVENTION

The present invention relates to quinolone carboxylic acids, derivatives thereof, and methods of making and using the same. In particular, the present invention relates to fluoroquinolone carboxylic acids, derivatives thereof, methods of making and using the same.

Bacterial pathogens continue to pose a serious threat to public health as indicated by a worldwide resurgence of bacterial diseases. One aspect of this resurgence appears to be the result of prior widespread, and largely effective, therapeutic and prophylactic use of antibiotics, which, unfortunately, over time has also selected for resistant strains of various bacterial pathogens. Of particular concern to the public health has been the emergence and proliferation of bacterial strains that are resistant to multiple antibiotics in the current arsenal of antimicrobial agents. Such multiantibiotic-resistant ("MAR") bacterial strains include species of Gram-positive bacteria, such as, antibiotic-resistant strains of *Staphylococcus aureus*, *Enterococcus faecalis*, and *Enterococcus fecium*, which, along with antibiotic-resistant Gram-negative strains of *Escherichia coli*, constitute the most frequent etiological agents of nosocomial (hospital-acquired) diseases, such as septicemia, endocarditis, and infections of wounds and the urinary tract. *S. aureus* is currently the most frequent cause of nosocomial bacteremia and skin or wound infection. *Streptococcus pneumoniae* causes several serious and life-threatening diseases, including a contagious meningitis, bacteremia, and otitis media. Annual



G00002

mortality from *S. pneumoniae* infection alone is estimated at between 3-5 million persons globally. More recently, clinical accounts of highly aggressive skin and tissue infections by "flesh-eating" strains of Group-A *streptococcus* bacteria, such as *Streptococcus pyogenes*, has heightened the concern and need for new or improved antibacterial agents.

Quinolones constitute a group of antibiotics that have been available since the early 1960s and have proved to be valuable antibacterial agents. Quinolone carboxylic acid derivatives having various chemical structures have been synthesized, developed, and marketed. Nalidixic acid (1,4-dihydro-1-ethyl-7-methyl-1,8-naphthyridin-4-one-3-carboxylic acid), the progenitor of the series, was used primarily as a urinary-tract antiseptic. Later development provided agents with broader activity, increased potency against selected pathogens and improved pharmacokinetic and pharmacodynamic properties.

From a medical utility viewpoint, the quinolones are classified as first-, second-, and third-generation compounds. First-generation compounds like piromidic acid (8-ethyl-5,8-dihydro-5-oxo-2-(1-pyrrolidinyl)pyrido(2,3-d)pyrimidine-6-carboxylic acid) and pipemidic acid (8-ethyl-5,8-dihydro-5-oxo-2-(1-piperazinyl)pyrido(2,3-d)pyrimidine-6-carboxylic acid) provided coverage for Gram-negative *Enterobacteriaceae*. The second-generation compounds are divided into those with enhanced but predominant Gram-negative activity, against pathogens like *Escherischia coli* and *Pseudomonas aeruginosa*, and those with balanced broad-spectrum activity (norfloxacin, pefloxacin, enoxacin, fleroxacin, lomefloxacin, ciprofloxacin, ofloxacin, rufloxacin, nadifloxacin). Norfloxacin, ofloxacin, and ciprofloxacin have, therefore, been used mainly for treatment of diseases including urinary tract infections,

gastrointestinal infections, sexually transmitted diseases and the like. Third-generation antibiotics (levofloxacin, pazufloxacin, sparfloxacin, clinafloxacin, sitafloxacin, trovafloxacin, tosufloxacin, temafloxacin, grepafloxacin, balofloxacin, moxifloxacin, gatifloxacin) are those with enhanced activity against Gram-positive cocci (notably clinafloxacin, sitafloxacin, trovafloxacin for *Streptococcus pneumoniae*) and, for essentially all the third-generation quinolones, activity also against Gram-negative *Haemophilus influenzae* and *Legionella pneumophila*, and against anaerobes and atypical pathogens. Levofloxacin, moxifloxacin, and gatifloxacin have, therefore, found use for community-acquired infections such as those of the upper and lower respiratory tract infections (“RTI”) like pneumonia, sinusitis and pharyngitis, and for skin and soft tissue infections (“STI”) caused by Gram-positive strains of *staphylococci*, *pneumococci*, *streptococci*, and *enterococci*.


The improvements seen in most of the third-generation antibiotics in current use are generally attributed to their uniqueness in inhibiting DNA gyrase and topoisomerase IV of the bacterial targets. Three categories of quinolone inhibition have been suggested. Type I quinolones (norfloxacin, enoxacin, fleroxacin, ciprofloxacin, lomefloxacin, trovafloxacin, grepafloxacin, ofloxacin and levofloxacin) indicate a preference for topoisomerase IV inhibition. Type II quinolones (nadifloxacin and sparfloxacin) indicate a preference for DNA gyrase inhibition. Type III quinolones to which some of the third-generation quinolones belong (e.g., gatifloxacin, pazufloxacin, moxifloxacin and clinafloxacin) display, however, a dual-targeting property, and equally influence DNA gyrase inhibition and topoisomerase IV inhibition. (M. Takei, et al., *Antimicrobial Agents and Chemotherapy*, Vol. 45, 3544-49 (2000)). DNA gyrase is the primary target in bacteria, and thus is explained the weaker activity in Gram-positive

bacteria of the topoisomerase IV-targeting second-generation quinolones like norfloxacin, ciprofloxacin, ofloxacin, and levofloxacin. The unusual activity of nadifloxacin described in the literature, especially against Gram-positive *S. aureus*, now can be explained by its ability to target DNA gyrase (N. Oizumi, et al., *J. Infect. Chemother.*, Vol. 7, 191-194 (2001)). That some third-generation quinolones are primarily capable of targeting topoisomerase IV in Gram-positive *staphylococci*, and DNA gyrase in Gram-positive *S. pneumoniae*, explains the advantages provided by the dual-targeting third-generation quinolones like moxifloxacin and gatifloxacin. However, because of continuing threat of new strains of antibiotic-resistant bacteria that may surface in the future, continued effort has been devoted to develop new broad-spectrum antibiotics.

A family of fluoroquinolones was recently developed, and some compounds of this family show good antimicrobial activity against a wide range of Gram-positive and Gram-negative bacteria. See U.S. Patents 5,385,900; 5,447,926; 6,685,958; and 6,699,492; all of which are incorporated herein by reference in their entirety. Because of the promise of their therapeutic value, it is very desirable, in one aspect, to develop improved processes for preparing this family of fluoroquinolones in order to allow for a more widespread availability of these compounds.

SUMMARY OF THE INVENTION


In general, the present invention provides an improved process for preparing fluoroquinolones that have Formula I or salts thereof.



wherein R^1 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, and groups that can be hydrolyzed in living bodies; R^2 is selected from the group consisting of hydrogen, unsubstituted amino group, and amino groups substituted with one or two lower alkyl groups; R^3 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted lower alkoxy groups, substituted lower alkoxy groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, unsubstituted C_5 - C_{24} aryloxy groups, substituted C_5 - C_{24} aryloxy groups, unsubstituted C_5 - C_{24} heteroaryloxy groups, substituted C_5 - C_{24} heteroaryloxy groups, and groups that can be hydrolyzed in

living bodies; X is selected from the group consisting of halogen atoms; Y is selected from the group consisting of CH₂, O, S, SO, SO₂, and NR⁴, wherein R⁴ is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, and cycloalkyl groups; and Z is selected from the group consisting of oxygen and two hydrogen atoms.





In one aspect, a process of preparing fluoroquinolones having Formula I comprises contacting a first compound having Formula II with a second compound having Formula III to produce a fluoroquinolone having Formula I, wherein the first compound and the second compound are represented by



wherein R¹, R², R³, X, Y, and Z have the meanings as disclosed above.

In another aspect, a process of preparing fluoroquinolones having Formula IV comprises: (a) contacting a first compound having Formula II with a third compound having Formula V to produce a fourth compound having Formula VI, wherein the

fluoroquinolones having Formula IV, the first compound, the third compound, and the fourth compound are represented by



wherein R^1 , R^3 , X, Y, and Z have the meanings as disclosed above and R^5 comprises a protected amino group having a formula of $-NR^6$, wherein R^6 comprises a protecting group that is capable of leaving the protected amino group $-NR^6$; and (b) contacting the fourth compound with a catalyst to effect a cleavage of the protecting group from the $-NR^6$ group, to produce a fluoroquinolone having Formula IV.

In still another aspect, the present invention provides quinolone carboxylic acids prepared by any process disclosed herein and their derivatives (such as their salts or esters), and methods of using such quinolone carboxylic acids and derivatives.

Other features and advantages of the present invention will become apparent from the following detailed description and claims.

DETAILED DESCRIPTION

As used herein, the term "lower alkyl" or "lower alkyl group" means a C_1 - C_{15} linear- or branched-chain saturated aliphatic hydrocarbon monovalent group, which may be unsubstituted or substituted. The group may be partially or completely substituted with halogen atoms (F, Cl, Br, or I). Non-limiting examples of lower alkyl groups include methyl, ethyl, n-propyl, 1-methylethyl(isopropyl), n-butyl, n-pentyl, 1,1-dimethylethyl (t-butyl), and the like. It may be abbreviated as "Alk".

As used herein, the term "lower alkoxy" or "lower alkoxy group" means a C_1 - C_{15} linear- or branched-chain saturated aliphatic alkoxy monovalent group, which may be unsubstituted or substituted. The group may be partially or completely substituted with halogen atoms (F, Cl, Br, or I). Non-limiting examples of lower alkoxy

groups include methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, n-pentoxy, t-butoxy, and the like.


The term "cycloalkyl" or "cycloalkyl group" means a stable aliphatic saturated 3- to 15-membered monocyclic or polycyclic monovalent radical consisting solely of carbon and hydrogen atoms which may comprise one or more fused or bridged ring(s), preferably a 3- to 7-membered monocyclic rings. Other exemplary embodiments of cycloalkyl groups include 7- to 10-membered bicyclic rings. Unless otherwise specified, the cycloalkyl ring may be attached at any carbon atom which results in a stable structure and, if substituted, may be substituted at any suitable carbon atom which results in a stable structure. Exemplary cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, norbornyl, adamantyl, tetrahydronaphthyl (tetralin), 1-decalinyl, bicyclo[2.2.2]octanyl, 1-methylcyclopropyl, 2-methylcyclopentyl, 2-methylcyclooctyl, and the like.

As used herein, the term "aryl" or "aryl group" means an aromatic carbocyclic monovalent or divalent radical. In some embodiments, the aryl group has a number of carbon atoms from 5 to 24 and has a single ring (e.g., phenyl or phenylene), multiple condensed rings (e.g., naphthyl or anthranyl), or multiple bridged rings (e.g., biphenyl). Unless otherwise specified, the aryl ring may be attached at any suitable carbon atom which results in a stable structure and, if substituted, may be substituted at any suitable carbon atom which results in a stable structure. Non-limiting examples of aryl groups include phenyl, naphthyl, anthryl, phenanthryl, indanyl, indenyl, biphenyl, and the like. It may be abbreviated as "Ar".

The term "heteroaryl" or "heteroaryl group" means a stable aromatic monocyclic or polycyclic monovalent or divalent radical, which may comprise one or more fused or bridged ring(s). In some embodiments, the heteroaryl group has 5-24 members, preferably a 5- to 7-membered monocyclic or 7- to 10-membered bicyclic radical. The heteroaryl group can have from one to four heteroatoms in the ring(s) independently selected from nitrogen, oxygen, and sulfur, wherein any sulfur heteroatoms may optionally be oxidized and any nitrogen heteroatom may optionally be oxidized or be quaternized. Unless otherwise specified, the heteroaryl ring may be attached at any suitable heteroatom or carbon atom which results in a stable structure and, if substituted, may be substituted at any suitable heteroatom or carbon atom which results in a stable structure. Non-limiting examples of heteroaryls include furanyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, tetrazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolizinyl, azaindolizinyl, indolyl, azaindolyl, diazaindolyl, dihydroindolyl, dihydroazaindolyl, isoindolyl, azaisoindolyl, benzofuranyl, furanopyridinyl, furanopyrimidinyl, furanopyrazinyl, furanopyridazinyl, dihydrobenzofuranyl, dihydrofuranopyridinyl, dihydrofuranopyrimidinyl, benzothienyl, thienopyridinyl, thienopyrimidinyl, thienopyrazinyl, thienopyridazinyl, dihydrobenzothienyl, dihydrothienopyridinyl, dihydrothienopyrimidinyl, indazolyl, azaindazolyl, diazaindazolyl, benzimidazolyl, imidazopyridinyl, benzthiazolyl, thiazolopyridinyl, thiazolopyrimidinyl, benzoxazolyl, benzoxazinyl, benzoxazinonyl, oxazolopyridinyl, oxazolopyrimidinyl, benzisoxazolyl, purinyl, chromanyl, azachromanyl, quinolizinyl, quinolinyl, dihydroquinolinyl, tetrahydroquinolinyl, isoquinolinyl, dihydroisoquinolinyl, tetrahydroisoquinolinyl, cinnolinyl, azacinnolinyl,

phthalazinyl, azaphthalazinyl, quinazolinyl, azaquinazolinyl, quinoxalinyl, azaquinoxalinyl, naphthyridinyl, dihydronaphthyridinyl, tetrahydronaphthyridinyl, pteridiny, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, and phenoxazinyl, and the like.

In general, the present invention provides an improved process for preparing fluoroquinolones that have Formula I or salts thereof.



wherein R^1 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, and groups that can be hydrolyzed in living bodies; R^2 is selected from the group consisting of hydrogen, unsubstituted amino group, and amino groups substituted with one or two lower alkyl groups; R^3 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted lower alkoxy groups, substituted lower alkoxy groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, unsubstituted C_5 - C_{24}

aryloxy groups, substituted C₅-C₂₄ aryloxy groups, unsubstituted C₅-C₂₄ heteroaryloxy groups, substituted C₅-C₂₄ heteroaryloxy groups, and groups that can be hydrolyzed in living bodies; X is selected from the group consisting of halogen atoms; Y is selected from the group consisting of CH₂, O, S, SO, SO₂, and NR⁴, wherein R⁴ is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, and cycloalkyl groups; and Z is selected from the group consisting of oxygen and two hydrogen atoms.

In one aspect, R¹ is selected from the group consisting of hydrogen, C₁-C₅ (or alternatively, C₁-C₃) substituted and unsubstituted alkyl groups, C₃-C₁₀ (or alternatively, C₃-C₅) cycloalkyl groups, C₅-C₁₄ (or alternatively, C₆-C₁₄, or C₅-C₁₀, or C₆-C₁₀) substituted and unsubstituted aryl groups, C₅-C₁₄ (or alternatively, C₆-C₁₄, or C₅-C₁₀, or C₆-C₁₀) substituted and unsubstituted heteroaryl groups, and groups that can be hydrolyzed in living bodies. In one embodiment, R¹ is selected from the group consisting of C₁-C₅ (or alternatively, C₁-C₃) substituted and unsubstituted alkyl groups.

In another aspect, R² is selected from the group consisting of unsubstituted amino group and amino groups substituted with one or two C₁-C₅ (or alternatively, C₁-C₃) alkyl groups.


In still another aspect, R³ is selected from the group consisting of hydrogen, C₁-C₅ (or alternatively, C₁-C₃) substituted and unsubstituted alkyl groups, C₃-C₁₀ (or alternatively, C₃-C₅) cycloalkyl groups, C₁-C₅ (or alternatively, C₁-C₃) substituted and unsubstituted alkoxy groups, C₅-C₁₄ (or alternatively, C₆-C₁₄, or C₅-C₁₀, or C₆-C₁₀) substituted and unsubstituted aryl groups, C₅-C₁₄ (or alternatively, C₆-C₁₄, or C₅-C₁₀, or C₆-C₁₀) substituted and unsubstituted heteroaryl groups, and C₅-C₁₄ (or alternatively, C₆-

C₁₄, or C₅-C₁₀, or C₆-C₁₀) substituted and unsubstituted aryloxy groups. In one embodiment, R³ is selected from the group consisting of C₃-C₁₀ (or alternatively, C₃-C₅) cycloalkyl groups.


In yet another aspect, X is selected from the group consisting of Cl, F, and Br. In one embodiment, X is Cl. In another embodiment, X is F.

In a further aspect, Y is hydrogen. In still another aspect, Z comprises two hydrogen atoms.

In one embodiment, the fluoroquinolone carboxylic acid has a Formula Ia.





In one aspect, the present invention provides an improved process of preparing fluoroquinolones having Formula I. The process comprises contacting a first compound having Formula II with a second compound having Formula III to produce a fluoroquinolone having Formula I, wherein the first compound and the second compound are represented by




wherein R^1 , R^2 , R^3 , X, Y, and Z have the meanings as disclosed above.

In another aspect, the first compound having Formula II, which is used in a process of the present invention as disclosed above, can be prepared according a procedure disclosed in published European Patent Application EP 0230946 A2, which is incorporated in its entirety by reference. For example, the first compound having Formula II is prepared by a process comprising: (a) reacting a compound having Formula X with an equimolar or excess amount of orthoformic acid ester in acetic anhydride (1 to 20-fold volume per total volume of the other reagents) at a temperature in the range from about room temperature to about 200 °C (preferably, from about 100 °C to about 150 °C) for a time from about 30 minutes to 24 hours to produce a compound having Formula XI; (b) treating the compound having Formula XI with an equalmolar or excess amount of an amine having a formula of NH_2R^3 in a solvent comprising an alcohol (preferably, ethanol or propanol), to convert the compound having Formula XI to a compound having Formula XII; (c) treating the compound having Formula XII with a fluoride salt (such as

one selected from the group consisting of sodium fluoride, potassium fluoride, and lithium fluoride) in a solvent selected from the group consisting of dioxane, dimethylformamide, dimethylsulfoxide, and sulfolane a temperature in the range from about 0 °C to about 200 °C (preferably, from about 50 °C to about 150 °C) for a time in the range from about 30 minutes to about 24 hours, to produce the compound having Formula II. The compounds having Formulae X, XI, and XII are shown below.



wherein R⁷ is unsubstituted lower alkyl groups, substituted lower alkyl groups, unsubstituted C₅-C₂₄ aryl groups (or alternatively, C₅-C₁₄, or C₅-C₁₀, or C₆-C₁₀), cycloalkyl groups, substituted C₅-C₂₄ aryl groups (or alternatively, C₅-C₁₄, or C₅-C₁₀, or C₆-C₁₀), unsubstituted C₅-C₂₄ heteroaryl groups (or alternatively, C₅-C₁₄, or C₅-C₁₀, or C₆-C₁₀), and substituted C₅-C₂₄ heteroaryl groups (or alternatively, C₅-C₁₄, or C₅-C₁₀, or C₆-C₁₀); and




In another aspect, the second compound having Formula III can be prepared by cyclization of various amino acids. For examples, such compounds having Formula III can be prepared according to the methods disclosed in D.W. Adamson, *J. Chem. Soc.*, p. 39 (1943); R. Pellegata et al., *Synthesis*, p.614 (1978); and M. Saburi et al., *Bull. Chem. Soc. Japan*, Vol. 60, pp 141-48 (1987). These references are incorporated herein by reference. Alternatively, various azepines having general Formula III can be prepared according to the methods disclosed in H. Chong et al., *J. Chem. Soc., Perkin Trans.*, Vol. 1, 2080-86 (2002); J. Barluenga, *Pure Appl. Chem.*, Vol. 74, No. 8, 1317-25 (2002); and T. Naito et al. (available at <http://www.ch.ic.ac.uk/ectoc/echet96/papers/054/index.htm>), using appropriate starting materials. The references by H. Chong et al., and by J. Barluenga are incorporated herein by reference.

In one embodiment of the present invention, a fluoroquinolone having Formula I is prepared as follows. One mole of the compound having Formula II is reacted with about 1-5 moles of the compound having Formula III in a solvent such as acetonitrile, dimethylsulfoxide, or the like, at a temperature in the range from about room temperature to about 100 °C for a time in the range from about 10 minutes to about 7 days. After the reaction, the precipitate is collected by filtration and washed, for example at room temperature, with a sufficient quantity of a suitable solvent, such as methanol, chloroform, ether, or the like, to obtain a crude product. The crude product is


purified, for example, by silica gel column chromatography or by recrystallization to obtain the fluoroquinolone having Formula I.

In another embodiment of the present invention, a fluoroquinolone having Formula IV is prepared as follows. One mole of the compound having Formula II is reacted with about 1-5 moles of the compound having Formula V in a solvent such as acetonitrile, dimethylsulfoxide, or the like, at a temperature in the range from about room temperature to about 100 °C for a time in the range from about 10 minutes to about 7 days to produce a compound having Formula VI. An amount of an acid or base (depending on whether the cleavage of the protecting group is acid- or base-catalyzable), such as from about 0.1 to about 5 moles per mole of of the compound having Formula V, is added to the reaction mixture to allow for the splitting of the protecting group R^6 from the protected amino $-NR^5$ group. In one embodiment, after this reaction, a base is added to the reaction mixture to convert free HF and HX acids to their salts (resulting pH is about 7), which are washed, for example at room temperature, from the mixture to produce a crude product. The crude product is purified, for example, by silica gel column chromatography or by recrystallization to obtain the fluoroquinolone having Formula IV.

In one embodiment, the compound having Formula V has Formula VII.




This compound can be prepared by the following reaction.




The nitrophenylalkylidene protecting group is disclosed in the above scheme only for illustrative purposes. Other protecting groups can be used in place of the nitrophenylalkylidene group, as can be recognized by people having skill in the art of organic synthesis. For example, another commonly used protecting group for the amine moiety is the t-butoxycarbonyl ("t-Boc"), which may be finally cleaved by an anhydrous acid catalyst, such as HCl to yield the amino group. Still another example of a protecting group for the amine moiety is the fluorenylmethoxycarbonyl ("Fmoc"), which can be cleaved by an anhydrous base catalyst, such as ammonia, piperidine, or morpholine.


In still another aspect, a process for preparing a fluoroquinolone carboxylic acid having Formula Ia comprises: (a) contacting a compound having Formula IIa with a compound having Formula VIIa at a temperature in the range from about room temperature to about 100 °C for a time from about 10 minutes to about 7 days, to produce a compound having Formula VIa



(IIa)



(VIIa)



(b) contacting the compound having Formula VIa with an amount of HCl equal to about 0.1 to about 5 moles per mole of the compound having Formula VIIa at a temperature in the range from about room temperature to about 100 °C, in a presence of methanol, to produce the fluoroquinolone carboxylic acid having Formula Ia; and (c) recovering the fluoroquinolone carboxylic acid having Formula Ia.

In yet another aspect, the crude product can comprise a mixture of enantiomers of the compound having Formula I or enantiomers of the compound having Formula IV, as the case may be. One of the enantiomers is often more soluble in water than the other. Therefore, another aspect of the present invention comprises the separation of one of the enantiomers of a crude product by washing or dissolving the crude product with water, and recovering such an enantiomer from the aqueous phase.

Therefore, in another aspect of the present invention, a process of preparing an enantiomer of a fluoroquinolone having Formula I comprises: (a) contacting a first compound having Formula II with a second compound having Formula III to produce a

crude enantiomeric mixture comprising enantiomers of the fluoroquinolone having Formula I; (b) recovering the crude enantiomeric mixture; (c) contacting the crude enantiomeric mixture thus recovered with water to produce an aqueous solution; and (d) recover the enantiomer of the fluoroquinolone having Formula I from the aqueous solution. In one embodiment, the step of contacting the crude enantiomeric mixture with water is carried out at a temperature in a range from about room temperature to about 80 °C, or from about room temperature to about 50 °C. In another embodiment, the step of contacting the crude enantiomeric mixture with water is carried out at about room temperature.

In still another aspect, a process of preparing an enantiomer of a fluoroquinolone having Formula IV comprises: (a) contacting a first compound having Formula II with a third compound having Formula V to produce a fourth compound having Formula VI; (b) contacting the fourth compound with a catalyst capable of assisting a cleavage of a protecting group from the R⁵ group, to produce a crude enantiomeric mixture of fluoroquinolones having Formula IV; (c) recovering the crude enantiomeric mixture; (c) contacting the crude enantiomeric mixture thus recovered with water to produce an aqueous solution; and (d) recover the enantiomer of the fluoroquinolone having Formula IV from the aqueous solution; wherein X has the meaning disclosed above. In one embodiment, the step of contacting the crude enantiomeric mixture with water is carried out at a temperature in a range from about room temperature to about 80 °C, or from about room temperature to about 50 °C. In another embodiment, the step of contacting the crude enantiomeric mixture with water is carried out at about room temperature.

In yet another aspect, the present invention provides a fluoroquinolone having Formula I, Ia, or IV prepared by any appropriate process disclosed herein.

The processes of the present invention has advantages over the process disclosed in U.S. Patents 5,385,900 and 5,447,926 in that the present processes are simpler and do not require the last step of U.S. Patents 5,385,900 and 5,447,926 for the attachment of a halogen atom to the position 8 on the compounds having Formulae I, Ia, and IV. This step requires the use of an excess amount of a halogenating agent such as sulfonyl chloride, chlorine, bromine, iodine, fluorine, N-chlorosuccinic acid imide, N-bromosuccinic acid imide, or the like. The use of such halogenating agents, especially in the gas phase, requires installation of precautionary measures in the manufacturing process, which would increase the complexity and cost of the manufacture.

Compounds of this family of fluoroquinolones can be used effectively against the survival of microbial pathogens. For example, the compounds having Formula I, Ia, or IV are potent antimicrobial agents and are found to be effective against the survival of Gram-positive bacteria, such as *Bacillus subtilis*, *Staphylococcus aureus*, *Staphylococcus epidermis*, *Sarcina lutea*, *Streptococcus faecalis*, and *Micrococcus lysodeikticus*; Gram-negative bacteria, such as *Escherichia coli*, *Salmonella typhi*, *Shigella flexneri*, *Pseudomonas aeruginosa*, *Klebsiella pneumonias*, *Proteus vulgaris*, *Proteus rettgeri*, and *Serratia marcescens*; and a methicillin-resistant strain of *Streptococcus aureus*. See; e.g., U.S. Patents 5,385,900 and 5,447,926; which are incorporated herein by reference in their entirety.



A fluoroquinolone compound prepared by any method disclosed herein can be formulated into an antimicrobial composition for topical, oral, systemic, ocular, or

intraocular administration. Such a composition comprises a fluoroquinolone compound and an excipient appropriate for the administration, as can be determined by a person having skill in the art of pharmaceutical formulation for the applications disclosed above. For example, various excipients known in the art can be used to formulate a solution, suspension, dispersion, ointment, gel, capsule, or tablet. A fluoroquinolone compound prepared by any method disclosed herein is particularly suitable for a treatment, reduction, amelioration, or prevention of infections of the eye, ear, nose, throat, or respiratory system caused by bacteria, including, but not being limited to, those bacteria disclosed above. In one embodiment, such a fluoroquinolone is formulated into an ophthalmic solution, ointment, suspension, dispersion, or gel.

While specific embodiments of the present invention have been described in the foregoing, it will be appreciated by those skilled in the art that many equivalents, modifications, substitutions, and variations may be made thereto without departing from the spirit and scope of the invention as defined in the appended claims.

WHAT IS CLAIMED IS:

1. A process of preparing a fluoroquinolone having Formula I or salts thereof, the process comprising contacting a first compound having Formula II with a second compound having Formula III to produce a fluoroquinolone having Formula I, wherein the fluoroquinolone, the first compound, and the second compound are represented by



wherein R¹ is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted C₅-C₂₄ aryl

groups, substituted C₅-C₂₄ aryl groups, unsubstituted C₅-C₂₄ heteroaryl groups, substituted C₅-C₂₄ heteroaryl groups, and groups that can be hydrolyzed in living bodies; R² is selected from the group consisting of hydrogen, unsubstituted amino group, and amino groups substituted with one or two lower alkyl groups; R³ is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted lower alkoxy groups, substituted lower alkoxy groups, unsubstituted C₅-C₂₄ aryl groups, substituted C₅-C₂₄ aryl groups, unsubstituted C₅-C₂₄ heteroaryl groups, substituted C₅-C₂₄ heteroaryl groups, unsubstituted C₅-C₂₄ aryloxy groups, substituted C₅-C₂₄ aryloxy groups, unsubstituted C₅-C₂₄ heteroaryloxy groups, substituted C₅-C₂₄ heteroaryloxy groups, and groups that can be hydrolyzed in living bodies; X is selected from the group consisting of halogen atoms; Y is selected from the group consisting of CH₂, O, S, SO, SO₂, and NR⁴, wherein R⁴ is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, and cycloalkyl groups; and Z is selected from the group consisting of oxygen and two hydrogen atoms.

2. The process of claim 1, wherein R¹ is selected from the group consisting of hydrogen, C₁-C₅ substituted and unsubstituted alkyl groups, C₃-C₁₀ cycloalkyl groups, C₆-C₁₄ substituted and unsubstituted aryl groups, C₅-C₁₄ substituted and unsubstituted heteroaryl groups, and groups that can be hydrolyzed in living bodies; R² is selected from the group consisting of unsubstituted amino group and amino groups substituted with one or two C₁-C₅ alkyl groups; R³ is selected from the group consisting of hydrogen, C₁-C₅ substituted and unsubstituted alkyl groups, C₃-C₁₀ cycloalkyl groups, C₁-C₅ substituted and unsubstituted alkoxy groups, C₅-C₁₄ substituted and unsubstituted aryl groups, C₅-C₁₄ substituted and unsubstituted heteroaryl groups, and C₅-C₁₄

substituted and unsubstituted aryloxy groups; and X is selected from the group consisting of Cl, F, and Br.

3. The process of claim 1, wherein R^1 is selected from the group consisting of hydrogen, C_1 - C_5 substituted and unsubstituted alkyl groups and groups that can be hydrolyzed in living bodies; R^2 is selected from the group consisting of unsubstituted amino group and amino groups substituted with one or two C_1 - C_5 alkyl groups; R^3 is selected from the group consisting of C_3 - C_{10} cycloalkyl groups; X is selected from the group consisting of Cl and F; Y comprises hydrogen; and Z comprises two hydrogen atoms.

4. The process of claim 1, wherein the step of contacting is carried out at a temperature in a range from about room temperature to about 100 °C.



5. The process of claim 3, wherein the step of contacting is carried out at a temperature in a range from about room temperature to about 100 °C.


6. The process of claim 1, wherein the step of contacting the first compound having Formula II with the second compound having Formula III results in a crude product comprising an enantiomeric mixture of a fluoroquinolone having Formula I, and the process further comprises washing or dissolving the crude product with water to produce an aqueous mixture, and substantially recovering an enantiomer from the aqueous mixture.

7. The process of claim 6, wherein the step of recovering the enantiomer from the aqueous mixture is carried out by recrystallization.

8. A process of preparing a fluoroquinolone having Formula IV or salts thereof, the method comprising:

- (a) contacting a first compound having Formula II with a third compound having Formula V to produce a fourth compound having Formula VI, wherein the fluoroquinolone having Formula IV, the first compound, the third compound, and the fourth compound are represented by





wherein R^1 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, and groups that can be hydrolyzed in living bodies; R^3 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, cycloalkyl groups, unsubstituted lower alkoxy groups, substituted lower alkoxy groups, unsubstituted C_5 - C_{24} aryl groups, substituted C_5 - C_{24} aryl groups, unsubstituted C_5 - C_{24} heteroaryl groups, substituted C_5 - C_{24} heteroaryl groups, unsubstituted C_5 - C_{24} aryloxy groups, substituted C_5 - C_{24} aryloxy groups, unsubstituted C_5 - C_{24} heteroaryloxy groups, substituted C_5 - C_{24} heteroaryloxy groups, and groups that can be hydrolyzed in living bodies; X is selected from the group consisting of halogen atoms; Y is selected from the group consisting of CH_2 , O, S, SO, SO_2 , and NR^4 , wherein R^4 is selected from the group consisting of hydrogen, unsubstituted lower alkyl groups, substituted lower alkyl groups, and cycloalkyl groups; Z is selected from the group consisting of oxygen and two hydrogen atoms; and R^5 comprises a protected amino group having a formula of

–NR⁶, wherein R⁶ comprises a protecting group that is capable of leaving the protected amino group –NR⁶; and




(b) contacting the fourth compound with a sufficient amount of a catalyst and at a condition sufficient to effect a cleavage of the protecting group R⁶ from the –NR⁶ group, to produce a fluoroquinolone having Formula IV.


9. The process of claim 8, wherein R¹ is selected from the group consisting of hydrogen, C₁-C₅ substituted and unsubstituted alkyl groups, C₃-C₁₀ cycloalkyl groups, C₆-C₁₄ substituted and unsubstituted aryl groups, C₆-C₁₄ substituted and unsubstituted heteroaryl groups, and groups that can be hydrolyzed in living bodies; R³ is selected from the group consisting of hydrogen, C₁-C₅ substituted and unsubstituted alkyl groups, C₃-C₁₀ cycloalkyl groups, C₁-C₅ substituted and unsubstituted alkoxy groups, C₅-C₁₄ substituted and unsubstituted aryl groups, C₅-C₁₄ substituted and unsubstituted heteroaryl groups, and C₅-C₁₄ substituted and unsubstituted aryloxy groups; R⁶ is selected from the group consisting of nitrophenylalkylidene, t-Boc, and Fmoc; and X is selected from the group consisting of Cl, F, and Br.

10. The process of claim 8, wherein R¹ is selected from the group consisting of hydrogen, C₁-C₅ substituted and unsubstituted alkyl groups and groups that can be hydrolyzed in living bodies; R³ is selected from the group consisting of C₃-C₁₀ cycloalkyl groups; R⁶ comprises a nitrophenylalkylidene group; X is selected from the group consisting of Cl and F; Y comprises hydrogen; and Z comprises two hydrogen atoms.

11. The process of claim 8, wherein the catalyst is selected from the group consisting of acids and bases.
12. The process of claim 8, wherein the step of contacting is carried out at a temperature in a range from about room temperature to about 100 °C.
13. The process of claim 10, wherein the step of contacting is carried out at a temperature in a range from about room temperature to about 100 °C.
14. The process of claim 10, wherein the catalyst is hydrochloric acid.
15. The process of claim 8, wherein the step of contacting the fourth compound with a catalyst produces a crude product comprising an enantiomeric mixture of the fluoroquinolone having Formula IV, and the process further comprises washing or dissolving the crude product with water to produce an aqueous mixture, and substantially recovering an enantiomer from the aqueous mixture.
16. The process of claim 15, wherein the step of recovering the enantiomer from the aqueous mixture is carried out by recrystallization.
17. A process for preparing a fluoroquinolone carboxylic acid having Formula Ia or salts thereof, the process comprising:
 - (a) contacting a compound having Formula IIa with a compound having Formula VIIa at a temperature in the range from about room temperature to about 100 °C for a time from about 10 minutes to about 7 days, to produce a compound

having Formula VIa, wherein the fluoroquinolone having Formula Ia and the compounds having Formulae IIa, VIa, and VIIa are represent by





(b) contacting the compound having Formula VIa with an amount of HCl equal to about 0.1 to about 5 moles per mole of the compound having Formula VIIa at a temperature in the range from about room temperature to about 100 °C, in a presence of methanol, to produce the fluoroquinolone carboxylic acid having Formula Ia; and

(c) recovering the fluoroquinolone carboxylic acid having Formula Ia.

18. The process of claim 17, wherein the step of contacting the compound having Formula VIa with HCl produces a crude product comprising an enantiomeric mixture of the fluoroquinolone having Formula Ia, and the process further comprises washing or dissolving the crude product with water to produce an aqueous mixture, and substantially recovering an enantiomer from the aqueous mixture.

19. The process of claim 18, wherein the step of substantially recovering the enantiomer from the aqueous mixture is carried out by recrystallization.

20. An enantiomer of a fluoroquinolone having Formula I prepared by the process of claim 6.
21. An enantiomer of a fluoroquinolone having Formula I prepared by the process of claim 7.
22. An enantiomer of a fluoroquinolone having Formula IV prepared by the process of claim 15.
23. An enantiomer of a fluoroquinolone having Formula IV prepared by the process of claim 16.
24. An enantiomer of a fluoroquinolone having Formula Ia prepared by the process of claim 18.
25. An enantiomer of a fluoroquinolone having Formula Ia prepared by the process of claim 19.
26. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 20.
27. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 21.

28. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 22.
29. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 23.
30. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 24.
31. A method of effecting an antimicrobial activity in a subject, comprising administering into said subject a composition that comprises an enantiomer of a fluoroquinolone of claim 25.
32. The method of claim 26, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.
33. The method of claim 27, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.
34. The method of claim 28, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.

35. The method of claim 29, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.
36. The method of claim 30, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.
37. The method of claim 31, wherein said administering is carried out to effect a treatment, reduction, amelioration, or prevention of an infection of an eye, ear, nose, throat, or respiratory system.
38. A composition comprising an enantiomer of a fluoroquinolone having Formula I prepared by the process of claim 6.
39. A composition comprising an enantiomer of a fluoroquinolone having Formula I prepared by the process of claim 7.
40. A composition comprising an enantiomer of a fluoroquinolone having Formula IV prepared by the process of claim 15.
41. A composition comprising an enantiomer of a fluoroquinolone having Formula IV prepared by the process of claim 16.
42. A composition comprising an enantiomer of a fluoroquinolone having Formula Ia prepared by the process of claim 18.

43. A composition comprising an enantiomer of a fluoroquinolone having Formula Ia prepared by the process of claim 19.