

Description

FRUIT JUICE BEVERAGE HAVING IMPROVED QUALITY OF TASTE
EXHIBITED BY SUGAR AND SWEETENER

Technical Field

[0001]

The present invention relates to a fruit juice beverage having a sugar and a sweetener with an improved taste quality, and a method for producing the fruit juice beverage.

Background Art

[0002]

Humans have five sensory systems, and the sense of taste is one of the sensory systems of humans. The taste receptor organ to receive tastes is called taste buds, which exist on the fungiform papillae existing over a wide area, mainly on the tip of the tongue, on the vallate papillae existing on a limited area of the back of the tongue, and on the foliate papillae. The taste buds are a cell assembly composed of elongate cells, called taste cells, and basal cells. The taste cells protrude microvilli toward the tongue surface, and form synapses at bottom of the cells with taste nerve fibers entering the taste buds. Tastes we usually sense are transmitted as taste information via the taste nerves to the brain, where the tastes are perceived. Known taste receptors of

sweetness include T1R2 and T1R3. T1R2 and T1R3 are reported to form hetero-dimers (Non-patent Literatures 1 to 3).

[0003]

Although various studies have been made on the sense of taste, little has been revealed yet in this field. We usually experience various tastes of foods. Foods that seem to be tasty have appropriately mixed and well-harmonized tastes. The taste of foods may be tasted as a single taste in some cases, but is often tasted as a mixed taste of various tastes, which are associated with one another.

[0004]

Meanwhile, foods have been required to have lower calories in addition to a good taste in recent years. This relates to a fact that lifestyle-related diseases such as obesity and diabetes are regarded as a problem.

However, to produce lower-calorie foods, their natural sugar concentration has to be maintained low. This is an obstruction in the case of providing foods that exhibit low calories and a good taste.

[0005]

As an example of a contrast effect, which is an interaction of tastes, there has been long known a phenomenon in which addition of salt to sweet red-bean soup enhances sweetness. There is an example that reports the interaction between saltiness and sweetness by

focusing on this phenomenon, and it is concluded that the interaction between sweetness and saltiness requires sweetness that is strong to a certain degree (a 15% solution) and a salt concentration that is high to a certain degree (0.1 to 0.2%) (Non Patent Literature 4).

[0006]

In addition, foods and drinks that exhibit sweetness by using a high-intensity sweetener of low calorie, such as Rebaudioside A, as a substitute for a natural sugar have also been developed, but the sweetness exhibited by a high-intensity sweetener has distinctive aftertaste, which hinders the provision of good taste.

Citation List

Non Patent Literature

[0007]

[Non Patent Literature 1] Zhao G. Q., Zhang Y., Hoon M. A., Chandrashekar J., Erlenbach I., Ryba N. J. P., and Zukerl C. S., Cell, 2003, Vol. 115, 255-266

[Non Patent Literature 2] Li X, Staszewski L, Xu H, Durick K, Zoller M, Adler E., Proc Natl Acad Sci U S A. 2002,99(7), 4692-4696.

[Non Patent Literature 3] Fernstrom J. D., Munger S. D., Sclafani A., de Araujo I. E., Roberts A., and Molinary S., J. Nutr. 2012. Vol. 142: 1134S-1141S

[Non Patent Literature 4] Ayumi Uchida, Nao Takagi, Rieko Horikiri, Miho Matsue, Yumiko Uchiyama and Masashi Omori,

Research Bulletin of Otsuma Women's University for Home
Economics -No.49 (2013. 3)

Summary of Invention

Problem to be Solved by the Invention

[0008]

The development of a method for well improving the taste quality exhibited by a sugar and a sweetener while having an adequate sweetness intensity and maintaining the amount of the natural sugar used in a fruit juice beverage low has been much awaited.

Means for Solving the Problems

[0009]

The present inventors succeeded for the first time in enhancing a sweetness based on a combined use of a natural sugar and a specific high-intensity sweetener and improving a taste quality by containing not only the natural sugar but also the specific high-intensity sweetener and adding a low concentration of sodium.

[0010]

That is, the present invention is as follows.

[1]

A fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 ; and

(c) less than 70 mg/100ml of sodium,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, a luohan guo extract, mogroside V, and thaumatin, and

$0.1 < (X_1 + X_2) \leq 20$ is satisfied.

[2]

The fruit juice beverage according to [1], wherein a sodium content is 5 to 60 mg/100ml.

[3]

The fruit juice beverage according to [1] or [2], wherein energy is 50 Kcal/100 ml or less.

[4]

The fruit juice beverage according to any one of [1] to [3], wherein X_1 is 0.1 to 5.9.

[5]

The fruit juice beverage according to any one of [1] to [4], wherein a fruit juice percentage of the fruit juice beverage is 1 to 100%.

[6]

The fruit juice beverage according to any one of [1] to [5], wherein a fruit juice contained in the fruit juice beverage is one or more selected from an orange, a mikan, a lemon, a grapefruit, a lime, a pineapple, a strawberry,

a blueberry, a black currant, a cranberry, a blueberry, a guava, a banana, an acerola, a papaya, a passion fruit, a mango, an apple, a grape, a peach, an ume, a pear, an apricot, a sumomo, a melon, a kiwi fruit, and a Japanese quince.

[7]

The fruit juice beverage according to any one of [1] to [6], wherein the natural sugar is at least one selected from the group consisting of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, lactose, psicose, allose, tagatose, and a combination thereof.

[8]

The fruit juice beverage according to any one of [1] to [7], wherein the high-intensity sweetener b1 is at least one selected from the group consisting of rebaudioside M, rebaudioside D, a luo han guo extract, mogroside V, and a combination thereof.

[9]

The fruit juice beverage according to any one of [1] to [8], wherein the sodium is at least one selected from the group consisting of sodium chloride, sodium hydroxide, sodium malate, sodium sulfate, sodium citrate, sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptonate, sodium gluconate, sodium glutamate, sodium

tartrate, sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and mixtures thereof.

[10]

The fruit juice beverage according to any one of [1] to [9], wherein the fruit juice beverage comprises 5 to 60 mg/100ml of sodium, energy is 50 Kcal/100 ml or less, and $X1 + X2$ is 6 or more.

[11]

The fruit juice beverage according to any one of [1] to [10], which is packed in a container.

[12]

A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity $X1$ and (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity $X2$; and

(ii) adding (c) sodium such that a sodium content in the beverage is less than 70 mg/100ml,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, a luohanguo extract, mogroside V, and thaumatin, and

$0.1 < (X1 + X2) \leq 20$ is satisfied.

[13]

The method according to [12], wherein in the fruit juice beverage, the sodium content is 5 to 60 mg/100ml.

[14]

The method according to [12] or [13], wherein energy of the fruit juice beverage is 50 Kcal/100 ml or less.

[15]

The method according to any one of [12] to [14], wherein X1 is 0.1 to 5.9.

[16]

The method according to any one of [12] to [15], wherein a fruit juice percentage of the fruit juice beverage is 1 to 100%.

[17]

The method according to any one of [12] to [16], wherein a fruit juice contained in the fruit juice beverage is one or more selected from an orange, a mikan, a lemon, a grapefruit, a lime, a pineapple, a strawberry, a blueberry, a black currant, a cranberry, a blueberry, a guava, a banana, an acerola, a papaya, a passion fruit, a mango, an apple, a grape, a peach, an ume, a pear, an apricot, a sumomo, a melon, a kiwi fruit, and a Japanese quince.

[18]

The method according to any one of [12] to [17], wherein the natural sugar is at least one selected from the group consisting of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup,

lactose, psicose, allose, tagatose, and a combination thereof.

[19]

The method according to any one of [12] to [18], wherein the high-intensity sweetener b1 is at least one selected from the group consisting of rebaudioside M, rebaudioside D, a luo han guo extract, mogroside V, and a combination thereof.

[20]

The method according to any one of [12] to [19], wherein the sodium is at least one selected from the group consisting of sodium chloride, sodium hydroxide, sodium malate, sodium sulfate, sodium citrate, sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptonate, sodium gluconate, sodium glutamate, sodium tartrate, sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and mixtures thereof.

[21]

The method according to any one of [12] to [20], wherein in the fruit juice beverage, the sodium content is 15 to 50 mg/100ml, energy is 50 Kcal/100 ml or less, and $X1 + X2$ is 6 or more.

[22]

A concentrate for providing the fruit juice beverage according to [1], comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X4;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X5; and

(c) less than 700 mg/100ml of sodium

wherein $(X4 + X5) \leq 200$ is satisfied.

[23]

A method for enhancing a sweetness of a fruit juice beverage, comprising containing:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2; and

(c) less than 70 mg/100ml of sodium
in the fruit juice beverage.

Advantageous Effects of Invention

[0011]

The present invention provides a method for increasing the sweetness of a fruit juice beverage, which is not a simple sweetness that is obtained by increasing the amounts of a natural sugar and a high-intensity sweetener used, and providing good taste quality. In addition, the method of the present invention provides a fruit juice beverage having good taste quality with increased sweetness by a means other than control of the used amounts of sugar and sweetener.

Description of Embodiments

[0012]

Hereinafter, the present invention will be described in detail. The following embodiments are examples to describe the present invention and do not intend to limit the present invention only to these embodiments. The present invention can be carried out in various embodiments without departing from the spirit of the present invention.

Note that all of the literatures, laid-open publications, patent publications, and other patent literatures cited herein are deemed to be incorporated by reference into the present specification.

In the present specification, for example, the phrase "the content of the A component is X mg/100ml" means that "X mg of the A component is contained in 100 ml of the beverage". In addition, for example, the phrase "the content of the B component is Y ppm" means that "the B component is contained in Y ppm with respect to the total amount (100 mass%) of the beverage".

[0013]

1. Fruit juice beverage having a natural sugar and a high-intensity sweetener with an improved taste quality

The present invention provides the following fruit juice beverage (hereinafter, referred to as "the fruit juice beverage of the present invention") as a first embodiment.

A fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 ;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 ; and

(c) less than 70 mg/100ml of sodium,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, a luohan guo extract, mogroside V, and thaumatin, and

$0.1 < (X_1 + X_2) \leq 20$ is satisfied.

The fruit juice beverage of the present invention has a sweetness of a sweetness intensity X_3 exhibited by the components (a) to (c).

[0014]

That is, in the fruit juice beverage of the present invention, the components having a sweetness are (a) a specific amount of a natural sugar and (b) a specific amount of a predetermined high-intensity sweetener. The sweetness exhibited by the fruit juice beverage of the present invention should computationally be the total value of the sweetness intensity obtained when the component (a) is added to the fruit juice beverage and the sweetness intensity obtained when the component (b) is added to the fruit juice beverage. However, (c) sodium is present in the fruit juice beverage although the

concentration thereof is low, and thus the sweetness exhibited by (a) the natural sugar and (b) the high-intensity sweetener is enhanced, and the fruit juice beverage of the present invention has a sweetness more than the simple sum of the sweetness intensities. The present invention means to possibly further include additional components such as an acidulant, a flavor, a vitamin, a color, an antioxidant, an emulsifier, a preservative, a seasoning, an extract, a pH adjuster, and a quality stabilizer in addition to these components (a) to (c). However, the fruit juice beverage of the present invention does not include a substance having a sweetness other than the components (a) and (b) as a sweetener.

[0015]

Further, in the fruit juice beverage according to a preferable embodiment of the present invention, improving effect on taste qualities is exhibited in addition to sweetness enhancement. For example, in the fruit juice beverage according to an embodiment of the present invention, it is preferable for at least one of flavor (aroma) intensity and suppression of an odd taste (bitterness, astringency, or the like) to be improved. The effect of eliciting a flavor (aroma) can be confirmed not only by sensory evaluation but also by analysis of the amount of a volatilized aroma component by gas chromatography.

[0016]

(Fruit juice)

In the present invention, the "fruit juice beverage" means a beverage containing a fruit juice, which is a squeezed juice of a fruit.

Specific examples of the "fruit juice beverage" include "fruit beverages" defined in the "Fair Competition Code for Labeling of Fruit Beverages and the like" enforced in 2018 and "fruit juices", "fruit mixed juices", "fruit grains-containing fruit juices", "fruit/vegetable mixed juices", and "fruit juice-containing beverages" defined in the "Ordinance for Enforcement of the Fair Competition Code for Labeling of Fruit Beverages and the like" enforced in 2016.

A beverage in which the proportion of a fruit juice used is less than 10%, which is classified into "other beverages" in the above Code, is also included in the "fruit juice beverage" in the present invention as long as the beverage contains a fruit juice.

[0017]

The fruit juice contained in the fruit juice beverage of the present invention is not particularly limited, and examples thereof include one or more selected from an orange, a mikan, a lemon, a grapefruit, a lime, a pineapple, a strawberry, a blueberry, a black currant, a cranberry, a blueberry, a guava, a banana, an acerola, a papaya, a passion fruit, a mango, an apple, a grape, a peach, an ume,

a pear, an apricot, a sumomo, a melon, a kiwi fruit, and a Japanese quince.

[0018]

The fruit juice contained in the fruit juice beverage of the present invention can be used regardless of the production method, in the form of straight fruit juice, concentrated fruit juice, or the like.

The concentrated fruit juice can be one prepared by either of a heat concentration method and a freeze concentration method.

[0019]

The fruit juice percentage of the present invention may be 1 to 100%, 5 to 100%, 10 to 100%, 15 to 100%, 20 to 100%, 25 to 100%, 30 to 100%, 35 to 100%, 40 to 100%, 45 to 100%, 50 to 100%, 55 to 100%, 60 to 100%, 65 to 100%, 70 to 100%, 1 to 95%, 5 to 95%, 10 to 95%, 15 to 95%, 20 to 95%, 25 to 95%, 30 to 95%, 35 to 95%, 40 to 95%, 45 to 95%, 50 to 95%, 55 to 95%, 60 to 95%, 65 to 95%, 70 to 95%, 1 to 90%, 5 to 90%, 10 to 90%, 15 to 90%, 20 to 90%, 25 to 90%, 30 to 90%, 35 to 90%, 40 to 90%, 45 to 90%, 50 to 90%, 55 to 90%, 60 to 90%, 65 to 90%, 70 to 90%, 1 to 85%, 5 to 85%, 10 to 85%, 15 to 85%, 20 to 85%, 25 to 85%, 30 to 85%, 35 to 85%, 40 to 85%, 45 to 85%, 50 to 85%, 55 to 85%, 60 to 85%, 65 to 85%, 70 to 85%, 1 to 80%, 5 to 80%, 10 to 80%, 15 to 80%, 20 to 80%, 25 to 80%, 30 to 80%, 35 to 80%, 40 to 80%, 45 to 80%, 50 to 80%, 55 to 80%, 60 to 80%, 65 to 80%, 70 to 80%, 1 to 75%, 5 to 75%, 10 to 75%, 15 to

75%, 20 to 75%, 25 to 75%, 30 to 75%, 35 to 75%, 40 to 75%, 45 to 75%, 50 to 75%, 55 to 75%, 60 to 75%, 65 to 75%, 1 to 70%, 5 to 70%, 10 to 70%, 15 to 70%, 20 to 70%, 25 to 70%, 30 to 70%, 35 to 70%, 40 to 70%, 45 to 70%, 50 to 70%, 55 to 70%, 60 to 70%, 1 to 65%, 5 to 65%, 10 to 65%, 15 to 65%, 20 to 65%, 25 to 65%, 30 to 65%, 35 to 65%, 40 to 65%, 45 to 65%, 50 to 65%, 55 to 65%, 1 to 60%, 5 to 60%, 10 to 60%, 15 to 60%, 20 to 60%, 25 to 60%, 30 to 60%, 35 to 60%, 40 to 60%, 45 to 60%, 50 to 60%, 1 to 55%, 5 to 55%, 10 to 55%, 15 to 55%, 20 to 55%, 25 to 55%, 30 to 55%, 35 to 55%, 40 to 55%, 45 to 55%, 1 to 50%, 5 to 50%, 10 to 50%, 15 to 50%, 20 to 50%, 25 to 50%, 30 to 50%, 35 to 50%, 40 to 50%, 1 to 45%, 5 to 45%, 10 to 45%, 15 to 45%, 20 to 45%, 25 to 45%, 30 to 45%, 35 to 45%, 1 to 40%, 5 to 40%, 10 to 40%, 15 to 40%, 20 to 40%, 25 to 40%, 30 to 40%, 1 to 35%, 5 to 35%, 10 to 35%, 15 to 35%, 20 to 35%, 25 to 35%, 1 to 30%, 5 to 30%, 10 to 30%, 15 to 30%, 20 to 30%, 1 to 25%, 5 to 25%, 10 to 25%, 15 to 25%, 1 to 20%, 5 to 20%, 10 to 20%, 1 to 15%, or 5 to 15%, depending on embodiments.

[0020]

In the present invention, the "fruit juice percentage" refers to a relative concentration with that of a straight fruit juice obtained by squeezing a fruit taken as 100% and can be converted based on a reference level of sugar refractometer index (°Bx) or a reference level of acidity (%) as defined in the JAS standard (Japanese agricultural standard for fruit beverages).

The values of the reference levels of sugar refractometer index of typical fruits (unit: °Bx) are orange: 11, mikan: 9, grapefruit: 9, pineapple: 11, cranberry: 7, blueberry and guava: 8, banana: 23, papaya: 9, passion fruit: 14, mango: 13, apple: 10, grape: 11, peach: 8, Japanese pear: 8, Western pear: 11, apricot: 7, sumomo: 6, and melon and kiwi fruit: 10.

The reference levels of acidity (unit: %) of typical fruits are lemon: 4.5, lime: 6, ume: 3.5, and kabosu: 3.5.
[0021]

Specifically, the fruit juice percentage can be calculated from the amount of a fruit juice blended (g) in 100 mL of a fruit juice beverage and the concentration rate of the fruit juice calculated from the reference level of sugar refractometer index (°Bx) or the reference level of acidity (%) described above by the following conversion expression.

$$[\text{Fruit juice percentage (\%)}] = [\text{amount of fruit juice blended (g)}] \times [\text{concentration rate}] / 100 \text{ mL} \times 100$$

For example, the JAS standard for an orange fruit juice is Bx 11°, and thus when 6.0% by weight of a concentrated orange fruit juice having Bx 55° is blended in a beverage, the fruit juice percentage is 30%. When converting the fruit juice percentage of a fruit juice based on sugar refractometer index of the JAS standard, the sugar refractometer index of a sugar, honey, or the like added to the fruit juice shall be excluded.

[0022]

The form of the fruit juice beverage of the present invention is not limited, and can be, for example, the form of a beverage in which a concentrated fruit juice extract is dissolved, or the form of a packed-in-container fruit juice beverage obtained by enclosing the fruit juice in a container such as a can or a PET bottle so that the same is packed in the container.

[0023]

(Natural Sugar)

In the present invention, a "natural sugar", which is represented by the general formula of $C_m(H_2O)_n$, wherein m and n independently represent a natural number, contains a D-form carbohydrate, digestible by humans. Examples of the D-form carbohydrate include glucose, sucrose, fructose, maltose, oligosaccharides (for example, fructooligosaccharide, maltooligosaccharide, isomaltooligosaccharide, and galactooligosaccharide), high-fructose corn syrup (isomerized sugars), lactose, psicose, allose, tagatose, and combinations thereof.

Examples of the combination of glucose, sucrose, fructose, maltose, oligosaccharides, high-fructose corn syrup, and lactose include the following. Specifically, the combination includes sucrose and high-fructose corn syrup, sucrose and glucose, sucrose and fructose, sucrose and maltose, sucrose and an oligosaccharide, sucrose and lactose, high-fructose corn syrup and an oligosaccharide,

sucrose and psicose; sucrose, high-fructose corn syrup, and glucose; sucrose, high-fructose corn syrup, and fructose; sucrose, high-fructose corn syrup, and maltose; sucrose, high-fructose corn syrup, and an oligosaccharide; sucrose, high-fructose corn syrup, glucose, and an oligosaccharide; sucrose, glucose, fructose, and an oligosaccharides; and glucose, sucrose, fructose, maltose, an oligosaccharide, high-fructose corn syrup, lactose, and tagatose.

[0024]

Since a natural sugar has high energy, the energy (calorie) of the fruit juice beverage can be greatly reduced by decreasing the content of the natural sugar. The present invention keeps the concentration of the natural sugar low to decrease the energy (i.e., low calorie), but strong sweetness is felt when ingested by a combination of a natural sugar, a high-intensity sweetener, and sodium. Accordingly, the amount of the natural sugar of a sweetness intensity X1 is preferably an amount of providing an energy of 50 Kcal/100 ml or less. The energy of the fruit juice beverage of the present invention may be 0 to 50 Kcal/100 ml, 0 to 45 Kcal/100 ml, 0 to 40 Kcal/100 ml, 0 to 35 Kcal/100 ml, 0 to 30 Kcal/100 ml, 0 to 25 Kcal/100 ml, 0 to 22 Kcal/100 ml, 0 to 20 Kcal/100 ml, 0 to 15 Kcal/100 ml, 0 to 10 Kcal/100 ml, 0 to 5 Kcal/100 ml, 5 to 50 Kcal/100 ml, 5 to 45 Kcal/100 ml, 5 to 40 Kcal/100 ml, 5 to 35 Kcal/100 ml, 5 to 30 Kcal/100

ml, 5 to 25 Kcal/100 ml, 5 to 20 Kcal/100 ml, 5 to 15 Kcal/100 ml, 5 to 10 Kcal/100 ml, 10 to 50 Kcal/100 ml, 10 to 45 Kcal/100 ml, 10 to 40 Kcal/100 ml, 10 to 35 Kcal/100 ml, 10 to 30 Kcal/100 ml, 10 to 25 Kcal/100 ml, 10 to 20 Kcal/100 ml, 10 to 15 Kcal/100 ml, 15 to 50 Kcal/100 ml, 15 to 45 Kcal/100 ml, 15 to 40 Kcal/100 ml, 15 to 35 Kcal/100 ml, 15 to 30 Kcal/100 ml, 15 to 25 Kcal/100 ml, 15 to 20 Kcal/100 ml, 20 to 50 Kcal/100 ml, 20 to 45 Kcal/100 ml, 20 to 40 Kcal/100 ml, 20 to 35 Kcal/100 ml, 20 to 30 Kcal/100 ml, 20 to 25 Kcal/100 ml, 25 to 50 Kcal/100 ml, 25 to 45 Kcal/100 ml, 25 to 40 Kcal/100 ml, 25 to 35 Kcal/100 ml, or 25 to 30 Kcal/100 ml, depending on embodiments. The energy of the fruit juice beverage of the present invention may also be 0 to 32 Kcal/100ml, 0 to 24 Kcal/100ml, 0 to 8 Kcal/100ml, 0 to 4 Kcal/100ml, 4 to 32 Kcal/100ml, 4 to 24 Kcal/100ml, 4 to 8 Kcal/100ml, 8 to 32 Kcal/100ml, 8 to 24 Kcal/100ml, or 24 to 32 Kcal/100ml, depending on embodiments. When a high calorie component such as a milk component, a total calorie of a natural sugar and a component such as a milk component is preferably 50 Kcal/100ml or less.

[0025]

Additionally, X1 of "the sweetness intensity X1" may be 0 to 0.5, 0 to 1.0, 0 to 1.5, 0 to 2.0, 0 to 2.5, 0 to 3.0, 0 to 3.5, 0 to 4.0, 0 to 4.5, 0 to 5.0, 0 to 5.5, 0 to 6.0, 0 to 6.5, 0 to 7.0, 0 to 7.5, 0 to 8.0, 0 to 8.25, 0 to 8.5, 0 to 8.75, 0 to 9.0, 0 to 9.25, 0 to 9.5, 0 to

9.75, 0 to 10.0, 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5,
0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05
to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.05 to 6.0,
0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05
to 8.25, 0.05 to 8.5, 0.05 to 8.75, 0.05 to 9.0, 0.05 to
9.25, 0.05 to 9.5, 0.05 to 9.75, 0.05 to 10.0, 0.1 to 0.5,
0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0,
0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5,
0.1 to 5.9, 0.1 to 6.0, 0.1 to 6.5, 0.1 to 7.0, 0.1 to 7.5,
0.1 to 8.0, 0.1 to 8.25, 0.1 to 8.5, 0.1 to 8.75, 0.1 to
9.0, 0.1 to 9.25, 0.1 to 9.5, 0.1 to 9.75, 0.1 to 10.0,
0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0,
0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5,
0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0,
0.5 to 8.25, 0.5 to 8.5, 0.5 to 8.75, 0.5 to 9.0, 0.5 to
9.25, 0.5 to 9.5, 0.5 to 9.75, 0.5 to 10.0, 1.0 to 0.5,
1.0 to 1.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0,
1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5,
1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0,
1.0 to 8.25, 1.0 to 8.5, 1.0 to 8.75, 1.0 to 9.0, 1.0 to
9.25, 1.0 to 9.5, 1.0 to 9.75, 1.0 to 10.0, 1.5 to 0.5,
1.5 to 1.0, 1.5 to 1.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0,
1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5,
1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0,
1.5 to 8.25, 1.5 to 8.5, 1.5 to 8.75, 1.5 to 9.0, 1.5 to
9.25, 1.5 to 9.5, 1.5 to 9.75, 1.5 to 10.0, 2.0 to 0.5,
2.0 to 1.0, 2.0 to 1.5, 2.0 to 2.0, 2.0 to 2.5, 2.0 to 3.0,

2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5,
2.0 to 6.0, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 8.0,
2.0 to 8.25, 2.0 to 8.5, 2.0 to 8.75, 2.0 to 9.0, 2.0 to
9.25, 2.0 to 9.5, 2.0 to 9.75, 2.0 to 10.0, 2.5 to 0.5,
2.5 to 1.0, 2.5 to 1.5, 2.5 to 2.0, 2.5 to 2.5, 2.5 to 3.0,
2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5,
2.5 to 6.0, 2.5 to 6.5, 2.5 to 7.0, 2.5 to 7.5, 2.5 to 8.0,
2.5 to 8.25, 2.5 to 8.5, 2.5 to 8.75, 2.5 to 9.0, 2.5 to
9.25, 2.5 to 9.5, 2.5 to 9.75, or 2.5 to 10.0.

[0026]

X1 may also be 0 to 10.5, 0 to 11.0, 0 to 11.5, 0 to
12.0, 0 to 12.5, 0 to 13.0, 0 to 13.5, 0 to 14.0, 0 to
14.5, 0 to 15.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5,
0.05 to 12.0, 0.05 to 12.5, 0.05 to 13.0, 0.05 to 13.5,
0.05 to 14.0, 0.05 to 14.5, 0.05 to 15.0, 0.1 to 10.5, 0.1
to 11.0, 0.1 to 11.5, 0.1 to 12.0, 0.1 to 12.5, 0.1 to
13.0, 0.1 to 13.5, 0.1 to 14.0, 0.1 to 14.5, 0.1 to 15.0,
0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to
12.5, 0.5 to 13.0, 0.5 to 13.5, 0.5 to 14.0, 0.5 to 14.5,
0.5 to 15.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to
12.0, 1.0 to 12.5, 1.0 to 13.0, 1.0 to 13.5, 1.0 to 14.0,
1.0 to 14.5, 1.0 to 15.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to
11.5, 1.5 to 12.0, 1.5 to 12.5, 1.5 to 13.0, 1.5 to 13.5,
1.5 to 14.0, 1.5 to 14.5, 1.5 to 15.0, 2.0 to 10.5, 2.0 to
11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 12.5, 2.0 to 13.0,
2.0 to 13.5, 2.0 to 14.0, 2.0 to 14.5, 2.0 to 15.0, 2.5 to
10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 12.5,

2.5 to 13.0, 2.5 to 13.5, 2.5 to 14.0, 2.5 to 14.5, or 2.5 to 15.0.

The amount corresponding to the sweetness intensity X1 of the natural sugar refers to an amount (a concentration) that exhibits the sweetness of the sweetness intensity X1 under conditions where the natural sugar is dissolved in water at 20°C having a volume equivalent to that of the fruit juice beverage of the present invention.

[0027]

Here, the sweetness intensity means sweetness exhibited by a substance. For example, in the case where the sweetness intensity exhibited by sucrose per unit concentration Brix 1 is defined as a degree of sweetness of 1, glucose has a degree of sweetness of 0.6 to 0.7 (center value: 0.65). A numerical value obtained by multiplying this degree of sweetness by the concentration Brix value of glucose is the sweetness intensity of glucose. Thus, when the concentration of glucose is Brix 1.5, the sweetness intensity of glucose is $0.65 \times 1.5 = 0.975$. When calculating the sweetness intensity in the present invention, the sweetness intensity of a sugar showing a range in the degree of sweetness, the center value is used unless otherwise noted.

Table 1

Sugar (D-form)	Degree of sweetness
Sucrose	1
Glucose	0.6 to 0.7

Fructose	1.3 to 1.7
Maltose	0.4
Fructooligosaccharide	0.6
Maltooligosaccharide	0.3
Isomaltooligosaccharide	0.4 to 0.5
Galactooligosaccharide	0.7
Isomerized sugar	0.8 to 0.9
Lactose	0.2 to 0.3
Psicose	0.7
Allose	0.8
Tagatose	0.9

[0028]

(High-Intensity Sweetener)

In the present invention, the "high-intensity sweetener" refers to a compound having sweetness more intense than that of sucrose, and may be a naturally occurring compound, a synthetic compound, and a combination of a naturally occurring compound and a synthetic compound. The high-intensity sweetener, in an amount equivalent to sucrose, exhibits sweetness 5 times or more, 10 times or more, 50 times or more, 100 times or more, 500 times or more, 1000 times or more, 5000 times or more, 10000 times or more, 50000 times or more, or 100000 times or more as that of sucrose. The high-intensity sweetener exhibits extremely intense sweetness in this way. When the existence of the natural sugar and high-intensity sweetener in the fruit juice beverage of the present invention is expressed in a weight ratio, "natural sugar:high-intensity sweetener" is 5:1 to 10:1, 50:1 to

100:1, 50:1 to 200:1, 500:1 to 1000:1, 5000:1 to 10000:1, and 50000:1 to 100000:1.

[0029]

In the present invention, as the high-intensity sweetener, at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, a Luo han guo extract, mogroside V, and thaumatin is used.

The content of the high-intensity sweetener b1 based on the total amount (100 mass%) of the high-intensity sweetener as the component (b) is preferably 50 mass% or more, more preferably 60 mass% or more, further preferably 70 mass% or more, more further preferably 80 mass% or more, and particularly preferably 90 mass% or more. In an aspect of the present invention, the high-intensity sweetener as the component (b) may substantially consist of the high-intensity sweetener b1. In the present specification, the phrase "substantially consist of the high-intensity sweetener b1" means that impurities, such as other steviol glycoside and mogroside, which are inevitably included in the process (e.g., purification of a stevia extract or a Luo han guo extract or biosynthesis) of preparing the high-intensity sweetener b1 may be included.

[0030]

Rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, and rebaudioside E may be directly

extracted from stevia or may be obtained by attaching glucose to a compound having another structure contained in a stevia extract.

[0031]

The Luo han guo extract as a sweetener is an extract of Luo han guo containing a sweet substance derived from Luo han guo, and has been approved as a food additive in various countries including Japan and is commercially available. Examples of the sweet substance derived from Luo han guo include mogroside V, mogroside IV, 11-oxo-mogroside V, and Siamenoside I.

[0032]

Mogroside V is one of the major mogrol glycosides contained in Luo han guo, and it has been reported that it has a good-quality sweetness property close to sucrose when compared with rebaudioside A. In addition, the degree of sweetness of mogroside V is about 300 times that of sucrose (Murata Y. et al., Nippon Shokuhin Kagaku Kogaku Kaishi, Vol. 53, No. 10, 527 to 533, (2006)). Mogroside V can be obtained from a Luo han guo extract (e.g., an alcohol extract of Luo han guo) through purification by chromatography or the like. Alternatively, mogroside V may be obtained by attaching glucose to a compound having another structure contained in a Luo han guo extract.

[0033]

The Luo han guo extract preferably contains mogroside V and the ratio thereof is not limited and can be 10 wt%

or more, 15 wt% or more, 20 wt% or more, 25 wt% or more, 30 wt% or more, 35 wt% or more, 40 wt% or more, 45 wt% or more, 50 wt% or more, 55 wt% or more, 60 wt% or more, 65 wt% or more, 70 wt% or more, and 75 wt% or more, of the total dry weight of a Luo han guo extract. The content of mogroside V can be determined by a known technique such as liquid chromatography. The Luo han guo extract can be obtained by extracting a fruit of Luo han guo (*Siraitia grosvenorii*) with a suitable solvent (for example, an aqueous solvent such as water, an alcohol solvent such as ethanol or methanol, a mixed solvent of an aqueous solvent and an alcohol solvent such as water-containing ethanol or water-containing methanol), and then optionally carrying out a treatment such as degreasing, purification, concentration, and drying.

[0034]

Mogroside V may be one having a high purity, for example, a purity of 80% or more, 85% or more, 90% or more, 91% or more, 92% or more, 93% or more, 94% or more, 95% or more, 96% or more, 97% or more, or 98% or more. In mogroside V obtained by purifying a Luo han guo extract, understandably, the amount of Luo han guo extract components other than mogroside V is decreased with an increase in purity.

[0035]

Thaumatococcus is a protein-based sweetener extracted from a plant.

[0036]

In the present invention, an optional high-intensity sweetener b2 may be used in combination with the above-described high-intensity sweetener b1, as needed. Specific examples of the optional high-intensity sweetener b2 include peptide-based sweeteners, such as aspartame, neotame, and alitame; sucrose derivatives, such as sucralose; synthetic sweeteners, such as acesulfame K, saccharine, advantame, cyclamate, and dulcin; protein-based sweeteners extracted from plants, such as monellin, curculin, and brazzein, other than thaumatin; high-intensity sweeteners other than protein-based sweetener extracted from plants; tourmaline; and neohesperidin dihydrochalcone.

[0037]

The sucrose derivative is that obtained by substituting the OH group or the H group of sucrose with another substituent, and examples thereof include halogen derivatives of sucrose (sucralose), oxathiazinonedioxide derivatives, sugar alcohols, aldonic acid, and uronic acid.

[0038]

Examples of the high-intensity sweetener other than the protein-based sweeteners extracted from plants include sweeteners present in a stevia extract, other than rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, and rebaudioside E; sweeteners present in a Luo han guo extract, other than mogroside V; sweeteners

present in a Glycyrrhiza extract; and glycosides thereof, and also include monatin and glycyrrhizin.

[0039]

Examples of the sweetener present in a stevia extract, other than rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, and rebaudioside E and the glycoside thereof include steviol, stevioside, rebaudioside A, rebaudioside B, and rebaudioside C.

[0040]

Examples of the sweetener present in a Luo han guo extract, other than mogroside V and the glycoside thereof include mogroside IV.

[0041]

The Glycyrrhiza extract refers to that obtained from roots or rhizomes of Glycyrrhiza uralensis Fisher, Glycyrrhiza inflata Batalin, or Glycyrrhiza glabra Linne and having glycyrrhizic acid as a main component. Examples of the Glycyrrhiza extract include a Glycyrrhiza extract, glycyrrhizin, and a licorice extract.

[0042]

In the present invention, the high-intensity sweetener b1 is suitable at least one selected from the group consisting of rebaudioside M, rebaudioside D, a Luo han guo extract, mogroside V, and a combination thereof. In particular, rebaudioside D, rebaudioside M, or a mixture of rebaudioside D and rebaudioside M is suitable as the high-intensity sweetener b1.

[0043]

Rebaudioside D and rebaudioside M have less negative flavor, such as astringent taste and metallic taste, which is observed in rebaudioside A and have characteristics, such as good-quality sweetness, and they are expected to be used in the fields of food and drink (NIPPON KAGAKU KAISHI, (5), (1981), 726-735, "Sweet Diterpene-Glycoside of Leaves of *Stevia rebaudiana* Bertoni Synthesis and Structure-Sweetness Relationship of Rebaudiosides-A, -D, -E and Their Related glycosides", Kasai, Kaneda, Tanaka, Yamasaki, Sakamoto, Morimoto, Okada, Kitahata, and Furukawa). Thus, rebaudioside D and rebaudioside M when used alone are excellent in that they have less odd taste compared to rebaudioside A and sweetness close to sucrose. Rebaudioside D and rebaudioside M each have sweetness about 200 times that of sucrose.

[0044]

Examples of the combination of the high-intensity sweeteners b1 and the combination of the high-intensity sweetener b1 and the optional high-intensity sweetener b2 include the followings. Specifically, the examples include combinations of rebaudioside D and rebaudioside M; rebaudioside D and rebaudioside A; rebaudioside M and rebaudioside A; rebaudioside M and mogroside V; rebaudioside D and mogroside V; rebaudioside D and advantame; rebaudioside D and acesulfame K; rebaudioside D and sucralose; rebaudioside M, rebaudioside D, and

rebaudioside A; rebaudioside M, rebaudioside D, and mogroside V; rebaudioside M, rebaudioside D, and advantame; rebaudioside M, rebaudioside D, and acesulfame K; rebaudioside M, rebaudioside D, and sucralose; rebaudioside A, rebaudioside M, rebaudioside D, and mogroside V; rebaudioside A, rebaudioside M, mogroside V, and sucralose; rebaudioside D, rebaudioside M, mogroside V, and acesulfame K; rebaudioside M, rebaudioside A, mogroside V, and neohesperidin dihydrochalcone; and rebaudioside M, rebaudioside D, sucralose, neohesperidin dihydrochalcone, and brazzein.

[0045]

In addition, the examples include combinations of rebaudioside D and mogroside V; rebaudioside D and a Luo han guo extract; rebaudioside M and a Luo han guo extract; rebaudioside M, rebaudioside D, and a Luo han guo extract; rebaudioside A, rebaudioside M, rebaudioside D, and a Luo han guo extract; rebaudioside A, rebaudioside M, a Luo han guo extract, and sucralose; rebaudioside D, rebaudioside M, a Luo han guo extract, and acesulfame K; and rebaudioside M, rebaudioside A, a Luo han guo extract, and neohesperidin dihydrochalcone.

[0046]

X2 of "the sweetness intensity X2" may be 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to

2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, 3.0 to 5.5, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 6.0, 2.5 to 7.0, 2.5 to 7.5, 2.5 to 6.0, 2.5 to 6.5, 3.0 to 6.0, 3.0 to 6.5, 3.0 to 7.0, 3.0 to 7.5, 3.0 to 8.0, 3.0 to 8.5, 3.0 to 9.0, 3.0 to 9.5, 3.5 to 7.0, 3.5 to 7.5, 3.5 to 8.0, 4.5 to 8.5, 3.5 to 9.0, 3.5 to 9.5, 4.0 to 7.5, 4.0 to 8.0, 4.0 to 8.5, 4.0 to 9.0, 4.0 to 9.5, 3.5 to 8.5, 3.5 to 10.0, 3.5 to 10.5, 3.5 to 11.0, 3.5 to 11.5, or 4.0 to 11.5.

[0047]

X2 may also be 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.5, 0.05 to 9.0, 0.05 to 9.5, 0.05 to 10.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5, 0.05 to 12.0, 0.05 to 13.0, 0.05 to 14.0, 0.05 to 15.0, 0.05 to 16.0, 0.05 to 17.0, 0.05 to 18.0, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.5, 0.5 to 9.0, 0.5 to 9.5, 0.5 to 10.0, 0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to 13.0, 0.5 to 14.0, 0.5 to 15.0, 0.5 to 16.0, 0.5 to 17.0, 0.5 to 18.0, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0

to 8.5, 1.0 to 9.0, 1.0 to 9.5, 1.0 to 10.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to 12.0, 1.0 to 13.0, 1.0 to 14.0, 1.0 to 15.0, 1.0 to 16.0, 1.0 to 17.0, 1.0 to 18.0, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.5, 1.5 to 9.0, 1.5 to 9.5, 1.5 to 10.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to 11.5, 1.5 to 12.0, 1.5 to 13.0, 1.5 to 14.0, 1.5 to 15.0, 1.5 to 16.0, 1.5 to 17.0, 1.5 to 18.0, 2.0 to 8.0, 2.0 to 8.5, 2.0 to 9.0, 2.0 to 9.5, 2.0 to 10.0, 2.0 to 10.5, 2.0 to 11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 13.0, 2.0 to 14.0, 2.0 to 15.0, 2.0 to 16.0, 2.0 to 17.0, 2.0 to 18.0, 2.5 to 8.0, 2.5 to 8.5, 2.5 to 9.0, 2.5 to 9.5, 2.5 to 10.0, 2.5 to 10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 13.0, 2.5 to 14.0, 2.5 to 15.0, 2.5 to 16.0, 2.5 to 17.0, 2.5 to 18.0, 3.0 to 10.0, 3.0 to 10.5, 3.0 to 11.0, 3.0 to 11.5, 3.0 to 12.0, 3.0 to 13.0, 3.0 to 14.0, 3.0 to 15.0, 3.0 to 16.0, 3.0 to 17.0, 3.0 to 18.0, 3.5 to 4.0, 3.5 to 4.5, 3.5 to 5.0, 3.5 to 5.5, 3.5 to 6.0, 3.5 to 6.5, 3.5 to 12.0, 3.5 to 13.0, 3.5 to 14.0, 3.5 to 15.0, 3.5 to 16.0, 3.5 to 17.0, 3.5 to 18.0, 4.0 to 4.5, 4.0 to 5.0, 4.0 to 5.5, 4.0 to 6.0, 4.0 to 6.5, 4.0 to 7.0, 4.0 to 10.0, 4.0 to 10.5, 4.0 to 11.0, 4.0 to 12.0, 4.0 to 13.0, 4.0 to 14.0, 4.0 to 15.0, 4.0 to 16.0, 4.0 to 17.0, or 4.0 to 18.0.

The amount corresponding to a sweetness intensity X2 of a high-intensity sweetener refers to an amount which provides sweetness of a sweetness intensity X2 under the conditions when the high-intensity sweetener is dissolved

in water having the same volume as the fruit juice beverage of the present invention at 20°C.

[0048]

As in a natural sugar, also here, the sweetness intensity means the sweetness exhibited by a substance. For example, when the sweetness intensity exhibited by sucrose per unit concentration Brix 1 is defined as a degree of sweetness of 1, the degree of sweetness of rebaudioside D is 200 to 250 (center value 225), the degree of sweetness of rebaudioside M is 200 to 260 (center value 230), the degree of sweetness of rebaudioside N is 200 to 250 (center value 225), the degree of sweetness of rebaudioside O is 200 to 250 (center value 225), the degree of sweetness of rebaudioside E is 70 to 80 (center value 75), the degree of sweetness of a Luo han guo extract is 110 to 150 (center value 130), the degree of sweetness of mogroside V is 240 to 300 (center value 270), and the degree of sweetness of thaumatin is 2,000. The numerical value obtained by multiplying these degrees of sweetness by the concentration (w/v% (considered to be the same as w/w% in the case of a beverage)) of the high-intensity sweetener in the fruit juice beverage is the sweetness intensity of the high-intensity sweetener. When calculating the sweetness intensity in the present invention, the sweetness intensity of a high-intensity sweetener showing a range in the degree of sweetness, the center value is used unless otherwise noted.

[0049]

Further, the amount of a high-intensity sweetener can be P2 ppm and P2 ppm herein refers to an amount corresponding to a sweetness intensity X2. The P2 herein can be a value of 20 to 550, 25 to 550, 30 to 550, 35 to 550, 40 to 550, 45 to 550, 50 to 550, 55 to 550, 20 to 540, 25 to 540, 30 to 540, 35 to 540, 40 to 540, 45 to 540, 50 to 540, 55 to 540, 20 to 530, 25 to 530, 30 to 530, 35 to 530, 40 to 530, 45 to 530, 50 to 530, 55 to 530, 20 to 520, 25 to 520, 30 to 520, 35 to 520, 40 to 520, 45 to 520, 50 to 520, 55 to 520, 20 to 510, 25 to 510, 30 to 510, 35 to 510, 40 to 510, 45 to 510, 50 to 510, 55 to 510, 20 to 505, 25 to 505, 30 to 505, 35 to 505, 40 to 505, 45 to 505, 50 to 505, 55 to 505, 20 to 500, 25 to 500, 30 to 500, 35 to 500, 40 to 500, 45 to 500, 50 to 500, 55 to 500, 20 to 495, 25 to 495, 30 to 495, 35 to 495, 40 to 495, 45 to 495, 50 to 495, 55 to 495, 20 to 490, 25 to 490, 30 to 490, 35 to 490, 40 to 490, 45 to 490, 50 to 490, or 55 to 490.

[0050]

The P2 can also be a value of 1 to 1500, 1 to 1200, 5 to 1200, 1 to 1000, 5 to 1000, 10 to 1000, 1 to 900, 5 to 900, 10 to 900, 15 to 900, 20 to 900, 25 to 900, 30 to 900, 35 to 900, 40 to 900, 45 to 900, 50 to 900, 55 to 900, 1 to 800, 5 to 800, 10 to 800, 15 to 800, 20 to 800, 25 to 800, 30 to 800, 35 to 800, 40 to 800, 45 to 800, 50 to 800, 55 to 800, 1 to 700, 5 to 700, 10 to 700, 15 to 700, 20 to 700, 25 to 700, 30 to 700, 35 to 700, 40 to 700, 45 to 700,

50 to 700, 55 to 700, 1 to 600, 5 to 600, 10 to 600, 15 to 600, 20 to 600, 25 to 600, 30 to 600, 35 to 600, 40 to 600, 45 to 600, 50 to 600, 55 to 600, 1 to 550, 1 to 540, 1 to 530, 1 to 520, 1 to 510, 1 to 505, 1 to 500, 1 to 495, 1 to 490, 5 to 550, 5 to 540, 5 to 530, 5 to 520, 5 to 510, 5 to 505, 5 to 500, 5 to 495, 5 to 490, 10 to 550, 10 to 540, 10 to 530, 10 to 520, 10 to 510, 10 to 505, 10 to 500, 10 to 495, 10 to 490, 15 to 550, 15 to 540, 15 to 530, 15 to 520, 15 to 510, 15 to 505, 15 to 500, 15 to 495, or 15 to 490.

[0051]

(Sodium)

The present invention contains (c) less than 70 mg/100ml of sodium, which means that the amount of sodium atoms is less than 70 mg/100ml of sodium. The amount of sodium (or sodium content) may be 0.1 to less than 70mg/100ml, 0.1 to 65mg/100ml, 0.1 to 60mg/100ml, 0.1 to 55mg/100ml, 0.1 to 50mg/100ml, 0.1 to 45mg/100ml, 0.1 to 40mg/100ml, 0.1 to 35mg/100ml, 0.1 to 30mg/100ml, 0.1 to 25mg/100ml, 0.1 to 20mg/100ml, 0.1 to 19mg/100ml, 0.1 to 18mg/100ml, 0.1 to 17mg/100ml, 0.1 to 16mg/100ml, 0.1 to 15mg/100ml, 0.1 to 14mg/100ml, 0.1 to 13mg/100ml, 0.1 to 12mg/100ml, 0.1 to 11mg/100ml, 0.1 to 10mg/100ml, 1g/100ml to less than 70mg/100ml, 1 to 65mg/100ml, 1 to 60mg/100ml, 1 to 55mg/100ml, 1 to 50mg/100ml, 1 to 45mg/100ml, 1 to 40mg/100ml, 1 to 35mg/100ml, 1 to 30mg/100ml, 1 to 25mg/100ml, 1 to 20mg/100ml, 1 to 19mg/100ml, 1 to

18mg/100ml, 1 to 17mg/100ml, 1 to 16mg/100ml, 1 to 15mg/100ml, 1 to 14mg/100ml, 1 to 13mg/100ml, 1 to 12mg/100ml, 1 to 11mg/100ml, 1 to 10mg/100ml, 5g/100ml to less than 70mg/100ml, 5 to 65mg/100ml, 5 to 60mg/100ml, 5 to 55mg/100ml, 5 to 50mg/100ml, 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 35mg/100ml, 5 to 30mg/100ml, 5 to 25mg/100ml, 5 to 20mg/100ml, 5 to 19mg/100ml, 5 to 18mg/100ml, 5 to 17mg/100ml, 5 to 16mg/100ml, 5 to 15mg/100ml, 5 to 14mg/100ml, 5 to 13mg/100ml, 5 to 12mg/100ml, 5 to 11mg/100ml, 5 to 10mg/100ml, 7 to less than 70mg/100ml, 7 to 65mg/100ml, 7 to 60mg/100ml, 7 to 55mg/100ml, 7 to 50mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 35mg/100ml, 7 to 30mg/100ml, 7 to 25mg/100ml, 7 to 20mg/100ml, 7 to 19mg/100ml, 7 to 18mg/100ml, 7 to 17mg/100ml, 7 to 16mg/100ml, 7 to 15mg/100ml, 10 to 70mg/100ml, 10 to 65mg/100ml, 10 to 60mg/100ml, 10 to 55mg/100ml, 10 to 50mg/100ml, 10 to 45mg/100ml, 10 to 40mg/100ml, 10 to 35mg/100ml, 10 to 30mg/100ml, 10 to 25mg/100ml, 10 to 20mg/100ml, 10 to 19mg/100ml, 10 to 18mg/100ml, 10 to 17mg/100ml, 10 to 16mg/100ml, 10 to 15mg/100ml, 15g/100ml to less than 70mg/100ml, 15 to 65mg/100ml, 15 to 60mg/100ml, 15 to 55mg/100ml, 15 to 50mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 35mg/100ml, 15 to 30mg/100ml, 15 to 25mg/100ml, 15 to 20mg/100ml, 20 to less than 70mg/100ml, 20 to 65mg/100ml, 20 to 60mg/100ml, 20 to 55mg/100ml, 20 to 50mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to

35mg/100ml, 20 to 30mg/100ml, 20 to 25mg/100ml, 25 to less than 70mg/100ml, 25 to 65mg/100ml, 25 to 60mg/100ml, 25 to 55mg/100ml, 25 to 50mg/100ml, 25 to 45mg/100ml, 25 to 40mg/100ml, 25 to 35mg/100ml, or 25 to 30mg/100ml.

[0052]

The amount of sodium may be 0.1 to 22mg/100ml, 0.1 to 21mg/100ml, 1 to 22mg/100ml, 1 to 21mg/100ml, 4 to 40mg/100ml, 4 to 35mg/100ml, 4 to 34mg/100ml, 4 to 33mg/100ml, 4 to 32mg/100ml, 4 to 31mg/100ml, 4 to 30mg/100ml, 4 to 29mg/100ml, 4 to 26mg/100ml, 4 to 25mg/100ml, 4 to 22mg/100ml, 4 to 21mg/100ml, 4 to 20mg/100ml, 4 to 19mg/100ml, 4 to 18mg/100ml, 4 to 17mg/100ml, 4 to 16mg/100ml, 4 to 15mg/100ml, 4 to 14mg/100ml, 4 to 13mg/100ml, 4 to 12mg/100ml, 4 to 11mg/100ml, 4 to 10mg/100ml, 5 to 34mg/100ml, 5 to 33mg/100ml, 5 to 32mg/100ml, 5 to 31mg/100ml, 5 to 29mg/100ml, 5 to 22mg/100ml, 5 to 21mg/100ml, 10 to 34mg/100ml, 10 to 33mg/100ml, 10 to 32mg/100ml, 10 to 31mg/100ml, 10 to 29mg/100ml, 10 to 22mg/100ml, 10 to 21mg/100ml, 11.5 to 34mg/100ml, 11.5 to 33mg/100ml, 11.5 to 32mg/100ml, 11.5 to 31mg/100ml, 11.5 to 30mg/100ml, 11.5 to 29mg/100ml, 11.5 to 22mg/100ml, 11.5 to 21mg/100ml, 11.5 to 20mg/100ml, 11.5 to 19mg/100ml, 11.5 to 18mg/100ml, 11.5 to 17mg/100ml, 11.5 to 16mg/100ml, 11.5 to 15mg/100ml, 11.5 to 14mg/100ml, 11.5 to 13mg/100ml, 11.5 to 12mg/100ml, 5.75 to 34.5mg/100ml, 5.75 to 28.75mg/100ml, 5.75 to 23mg/100ml, 5.75 to 17.25mg/100ml, 5.75 to 11.5mg/100ml,

11.5 to 34.5mg/100ml, 11.5 to 28.75mg/100ml, 11.5 to 23mg/100ml, 11.5 to 17.25mg/100ml, 17.25 to 34.5mg/100ml, 17.25 to 28.75mg/100ml, 17.25 to 23mg/100ml, 23 to 34.5mg/100ml, 23 to 28.75mg/100ml, or 28.75 to 34.5mg/100ml.

[0053]

In addition, the amount of sodium that is added to a fruit juice beverage may be 0.1 to 50mg/100ml, 0.1 to 45mg/100ml, 0.1 to 40mg/100ml, 0.1 to 35mg/100ml, 0.1 to 30mg/100ml, 0.1 to 25mg/100ml, 0.1 to 20mg/100ml, 0.1 to 19mg/100ml, 0.1 to 18mg/100ml, 0.1 to 17mg/100ml, 0.1 to 16mg/100ml, 0.1 to 15mg/100ml, 0.1 to 14mg/100ml, 0.1 to 13mg/100ml, 0.1 to 12mg/100ml, 0.1 to 11mg/100ml, 0.1 to 10mg/100ml, 1 to 50mg/100ml, 1 to 45mg/100ml, 1 to 40mg/100ml, 1 to 35mg/100ml, 1 to 30mg/100ml, 1 to 25mg/100ml, 1 to 20mg/100ml, 1 to 19mg/100ml, 1 to 18mg/100ml, 1 to 17mg/100ml, 1 to 16mg/100ml, 1 to 15mg/100ml, 1 to 14mg/100ml, 1 to 13mg/100ml, 1 to 12mg/100ml, 1 to 11mg/100ml, 1 to 10mg/100ml, 5 to 50mg/100ml, 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 35mg/100ml, 5 to 30mg/100ml, 5 to 25mg/100ml, 5 to 20mg/100ml, 5 to 19mg/100ml, 5 to 18mg/100ml, 5 to 17mg/100ml, 5 to 16mg/100ml, 5 to 15mg/100ml, 5 to 14mg/100ml, 5 to 13mg/100ml, 5 to 12mg/100ml, 5 to 11mg/100ml, 5 to 10mg/100ml, 7 to 50mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 35mg/100ml, 7 to 30mg/100ml, 7 to 25mg/100ml, 7 to 20mg/100ml, 7 to

19mg/100ml, 7 to 18mg/100ml, 7 to 17mg/100ml, 7 to 16mg/100ml, 7 to 15mg/100ml, 10 to 50mg/100ml, 10 to 45mg/100ml, 10 to 40mg/100ml, 10 to 35mg/100ml, 10 to 30mg/100ml, 10 to 25mg/100ml, 10 to 20mg/100ml, 10 to 19mg/100ml, 10 to 18mg/100ml, 10 to 17mg/100ml, 10 to 16mg/100ml, 10 to 15mg/100ml, 15 to 50mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 35mg/100ml, 15 to 30mg/100ml, 15 to 25mg/100ml, 15 to 20mg/100ml, 20 to 50mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to 35mg/100ml, 20 to 30mg/100ml, 20 to 25mg/100ml, 25 to 50mg/100ml, 25 to 45mg/100ml, 25 to 40mg/100ml, 25 to 35mg/100ml, or 25 to 30mg/100ml.

[0054]

Forms of sodium is not particularly limited provided that sodium in an ingestible state is contained in the fruit juice beverage of the present invention, and may be in at least one form selected from the group consisting of sodium chloride, sodium hydroxide, sodium malate, sodium sulfate, sodium citrate, sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptanoate, sodium gluconate, sodium glutamate, sodium tartrate, sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and a mixture thereof, for example.

In addition, when a fruit juice contains a sodium, the sodium contained in the fruit juice is also encompassed

in sodium in the fruit juice beverage of the present invention.

In an aspect of the present invention, sodium derived from a sodium component that is used as a preservative (e.g., sodium benzoate, sodium sulfite, sodium hyposulfite, sodium dehydroacetate, sodium pyrosulfite, or sodium propionate) is not substantially included in the component (c).

[0055]

In the present specification, the sodium content in a beverage can be measured by atomic absorption spectrometry. Incidentally, when the amount of a sodium-containing compound contained in the beverage is known, the value calculated from the amount may be used.

[0056]

The fruit juice beverage of the present invention has an enhanced sweetness as having been already mentioned. Whether or not the sweetness of the fruit juice beverage of the present invention is enhanced can be evaluated by panelists who received sensory trainings. Further, for the sweetness intensity X3 of the fruit juice beverage of the present invention, standard fruit juice beverages to be the sweetness standards are prepared with standards are prepared with sucrose concentrations assigned as sweetness intensities of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, and 15 and panelists compare the sweetness of the fruit juice beverage of the present invention with the sweetness

of these standard fruit juice beverages thereby to measure the sweetness intensity of the fruit juice beverage of the present invention. Note that the standard fruit juice beverages having sweetness intensities of 1, 2, ... 15 are prepared by adding sucrose in such a way that a sucrose content is 1 g/100 g, 2 g/100 g, ... and 15 g/100 g to a fruit juice beverage to which sucrose is not added.

Furthermore, of the standard fruit juice beverages having a lower sweetness than that of the fruit juice beverage of the present invention in the above measurement, the standard fruit juice beverage having the closest sweetness to that of the fruit juice beverage of the present invention is selected and adjusted in such a way as to have the same sweetness as that of the fruit juice beverage of the present invention by adding sucrose to the selected standard fruit juice beverage, during which a sweetness intensity X3 of the fruit juice beverage of the present invention can also be measured from a sucrose content in the adjusted standard fruit juice beverage.

[0057]

Other examples of the methods for measuring the sweetness of the fruit juice beverage of the present invention include a sweetness intensity rating using Visual Analogue Scale (VAS method). For the VAS method, literatures in J. Jpn. Soc. Stomatognath. Funct. (2014) 20 pp. 115-129 ("Construction of a Screening Test for Gustatory Function in Four Basic Tastes", Toyota et al.)

or the like can be referred. Specifically, in the measurement of sweetness intensity by the VAS method, for example, the evaluator defines the intensity of sweetness as "not sweet at all" at the lower end and "nothing could be sweeter than this" at the upper end, and makes evaluation by using a paper sheet on which a vertical line indicating the intensities of sweetness on a straight line is drawn and expressing the sweetness intensity felt at that time by a position on the straight line.

[0058]

The sweetness intensity X3 of the fruit juice beverage of the present invention may be 4.0 to 20, 4.0 to 15, 4.0 to 12.5, 4.0 to 10, 4.5 to 20, 4.5 to 15, 4.5 to 12.5, 4.5 to 10, 5.0 to 20, 5.0 to 15, 5.0 to 12.5, 5.0 to 10, 5.5 to 20, 5.5 to 15, 5.5 to 12.5, 5.5 to 10, 6.0 to 20, 6.0 to 15, 6.0 to 12.5, 6.0 to 10, 6.5 to 20, 6.5 to 15, 6.5 to 12.5, 6.5 to 10, 7.0 to 20, 7.0 to 15, 7.0 to 12.5, 7.0 to 10, 7.5 to 20, 7.5 to 15, 7.5 to 12.5, 7.5 to 10, 7.5 to 9, 7.5 to 8, 8.0 to 20, 8.0 to 15, 8.0 to 12.5, 8.0 to 10, 8.5 to 20, 8.5 to 15, 8.5 to 12.5, 8.5 to 10, 9.0 to 20, 9.0 to 15, 9.0 to 12.5, 9.0 to 10, 9.5 to 20, 9.5 to 15, 9.5 to 12.5, 9.5 to 10, 10.0 to 20, 10.0 to 15, 10.0 to 12.5, 10.5 to 20, 10.5 to 15, or 10.5 to 12.5. Alternatively, X3 may be 4.0 to 18, 4.0 to 16, 4.0 to 15.5, 4.0 to 14, 4.5 to 18, 4.5 to 16, 4.5 to 15.5, 4.5 to 14, 5.0 to 18, 5.0 to 16, 5.0 to 15.5, 5.0 to 14, 5.5 to 18, 5.5 to 16, 5.5 to 15.5, 5.5 to 14, 6.0 to 18, 6.0

to 16, 6.0 to 15.5, 6.0 to 14, 6.5 to 18, 6.5 to 16, 6.5 to 15.5, 6.5 to 14, 7.0 to 18, 7.0 to 16, 7.0 to 15.5, 7.0 to 14, 7.5 to 18, 7.5 to 16, 7.5 to 15.5, 7.5 to 14, 7.5 to 9, 7.5 to 8, 8.0 to 18, 8.0 to 18, 8.0 to 16, 8.0 to 15.5, 8.0 to 14, 8.5 to 18, 8.5 to 16, 8.5 to 15.5, 8.5 to 14, 9.0 to 18, 9.0 to 16, 9.0 to 15.5, 9.0 to 14, 9.5 to 18, 9.5 to 16, 9.5 to 15.5, 9.5 to 14, 10.0 to 18, 10.0 to 16, 10.0 to 15.5, 10.5 to 18, 10.5 to 16, or 10.5 to 15.5.
[0059]

In a certain embodiment, the fruit juice beverage of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1 of 0.1 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2 of 0.1 to 5; and

(c) less than 70 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 of 5.5 to 12.5 exhibited by the components (a) to (c).

[0060]

In another embodiment, the fruit juice beverage of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1 of 3 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2 of 1 to 5; and

(c) 5 to 60 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 of 5.5 to 12.5 exhibited by the components (a) to (c). Preferably, in the fruit juice beverage of the present invention, further, the energy is 50 Kcal/100 ml or less and $X1 + X2$ is 6 or more.

[0061]

In the fruit juice beverage of the present invention, the sweetness intensity X1 of the natural sugar, the sweetness intensity X2 of the high-intensity sweetener, the sodium content, the sweetness intensity X3 exhibited by the fruit juice beverage, and the energy of the fruit juice beverage can each be any value as long as $0.1 < (X1 + X2) \leq 20$ is satisfied, and for example, any of the numerical values of the sweetness intensity X1, the sweetness intensity X2, the sodium content, the sweetness intensity X3, and the energy shown below can also be combined.

[0062]

"The sweetness intensity X1": 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.5 to 3.0, 2.5 to 3.5,

2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, or 3.0 to 5.0.

[0063]

"The sweetness intensity X2": 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, or 3.0 to 5.0.

[0064]

"The amount of sodium": 5 to 60mg/100ml, 5 to 55mg/100ml, 5 to 50mg/100ml, 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 39mg/100ml, 5 to 38mg/100ml, 5 to 37mg/100ml, 5 to 36mg/100ml, 5 to 35mg/100ml, 5 to 34mg/100ml, 5 to 33mg/100ml, 5 to 32mg/100ml, 5 to 31mg/100ml, 5 to 30mg/100ml, 7 to 60mg/100ml, 7 to 55mg/100ml, 7 to 50mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 39mg/100ml, 7 to 38mg/100ml, 7 to 37mg/100ml, 7 to 36mg/100ml, 7 to 35mg/100ml, 7 to 34mg/100ml, 7 to 33mg/100ml, 7 to 32mg/100ml, 7 to 31mg/100ml, 7 to 30mg/100ml, 10 to 60mg/100ml, 10 to 55mg/100ml, 10 to 50mg/100ml, 10 to 45mg/100ml, 10 to

40mg/100ml, 10 to 39mg/100ml, 10 to 38mg/100ml, 10 to 37mg/100ml, 10 to 36mg/100ml, 10 to 35mg/100ml, 10 to 34mg/100ml, 10 to 33mg/100ml, 10 to 32mg/100ml, 10 to 31mg/100ml, 10 to 30mg/100ml, 15 to 60mg/100ml, 15 to 55mg/100ml, 15 to 50mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 39mg/100ml, 15 to 38mg/100ml, 15 to 37mg/100ml, 15 to 36mg/100ml, 15 to 35mg/100ml, 15 to 34mg/100ml, 15 to 33mg/100ml, 15 to 32mg/100ml, 15 to 31mg/100ml, 15 to 30mg/100ml, 20 to 60mg/100ml, 20 to 55mg/100ml, 20 to 50mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to 39mg/100ml, 20 to 38mg/100ml, 20 to 37mg/100ml, 20 to 36mg/100ml, 20 to 35mg/100ml, 25 to 60mg/100ml, 25 to 55mg/100ml, 25 to 50mg/100ml, 25 to 45mg/100ml, or 25 to 40mg/100ml.

[0065]

"The sweetness intensity X3": 4.0 to 20, 4.0 to 15, 4.0 to 12.5, 4.0 to 10, 4.5 to 20, 4.5 to 15, 4.5 to 12.5, 4.5 to 10, 5.0 to 20, 5.0 to 15, 5.0 to 12.5, 5.0 to 10, 5.5 to 20, 5.5 to 15, 5.5 to 12.5, 5.5 to 10, 6.0 to 20, 6.0 to 15, 6.0 to 12.5, 6.0 to 10, 6.5 to 20, 6.5 to 15, 6.5 to 12.5, 6.5 to 10, 7.0 to 20, 7.0 to 15, 7.0 to 12.5, 7.0 to 10, 7.5 to 20, 7.5 to 15, 7.5 to 12.5, 7.5 to 10, 7.5 to 9, 7.5 to 8, 8.0 to 20, 8.0 to 20, 8.0 to 15, 8.0 to 12.5, 8.0 to 10, 8.5 to 20, 8.5 to 15, 8.5 to 12.5, 8.5 to 10, 9.0 to 20, 9.0 to 15, 9.0 to 12.5, 9.0 to 10, 9.5 to 20, 9.5 to 15, 9.5 to 12.5, 9.5 to 10, 10.0 to 20, 10.0

to 15, 10.0 to 12.5, 10.5 to 20, 10.5 to 15, or 10.5 to 12.5.

[0066]

The energy: 0 to 50Kcal/100ml, 0 to 45Kcal/100ml, 0 to 40Kcal/100ml, 0 to 35Kcal/100ml, 0 to 30Kcal/100ml, 0 to 25Kcal/100ml, 0 to 20Kcal/100ml, 0 to 15Kcal/100ml, 0 to 10Kcal/100ml, 0 to 5Kcal/100ml, 5 to 50Kcal/100ml, 5 to 45Kcal/100ml, 5 to 40Kcal/100ml, 5 to 35Kcal/100ml, 5 to 30Kcal/100ml, 5 to 25Kcal/100ml, 5 to 20Kcal/100ml, 5 to 15Kcal/100ml, 5 to 10Kcal/100ml, 10 to 50Kcal/100ml, 10 to 45Kcal/100ml, 10 to 40Kcal/100ml, 10 to 35Kcal/100ml, 10 to 30Kcal/100ml, 10 to 25Kcal/100ml, 10 to 20Kcal/100ml, 10 to 15Kcal/100ml, 15 to 50Kcal/100ml, 15 to 45Kcal/100ml, 15 to 40Kcal/100ml, 15 to 35Kcal/100ml, 15 to 30Kcal/100ml, 15 to 25Kcal/100ml, 15 to 20Kcal/100ml, 20 to 50Kcal/100ml, 20 to 45Kcal/100ml, 20 to 40Kcal/100ml, 20 to 35Kcal/100ml, 20 to 30Kcal/100ml, 20 to 25Kcal/100ml, 25 to 50Kcal/100ml, 25 to 45Kcal/100ml, 25 to 40Kcal/100ml, 25 to 35Kcal/100ml, or 25 to 30Kcal/100ml.

[0067]

In addition, in the present invention, the sweetness intensity X1 of the natural sugar, the sweetness intensity X2 of the high-intensity sweetener, the sodium content, the sweetness intensity X3 exhibited by the fruit juice beverage, and the energy of the fruit juice beverage are not limited to combinations of the numerical values exemplified above, and any of the numerical values of the

sweetness intensity X1, the sweetness intensity X2, the sodium content, the sweetness intensity X3, and the energy described herein can be combined in a range that satisfies $0.1 < (X1 + X2) \leq 20$.

[0068]

Into the fruit juice beverage of the present invention, an acidulant (phosphoric acid, or the like), a flavor (an ester, or the like), a vitamin, a color, an antioxidant (sodium erythorbate, or the like), and an emulsifier (sucrose fatty acid ester, sorbitan fatty acid ester, polyglycerin fatty acid ester, or the like), a preservative, a seasoning, an extract, a pH adjuster, a quality stabilizer, or the like can further be appropriately blended as long as the effects of the present invention are not impaired.

[0069]

In a specific embodiment, the present invention provides the following fruit juice beverage (hereinafter, referred to as "fruit juice beverage A of the present invention").

A fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1;

(b) a high-intensity sweetener selected from the group consisting of rebaudioside M, rebaudioside D, and a combination thereof in an amount corresponding to a sweetness intensity X2; and

(c) 5 to 50 mg/100ml of sodium, wherein the fruit juice beverage has a sweetness of a sweetness intensity of X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0070]

In fruit juice beverage A of the present invention, the preferable ranges of the sweetness intensity X1, the sweetness intensity X2, the sodium content, the sweetness intensity X3, the high-intensity sweetener concentration P2, and the energy are, for example, as follows. These numerical values can be arbitrarily combined in a range that satisfies $0.1 < (X1 + X2) \leq 20$.

[0071]

"The sweetness intensity X1": 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.25, 0.05 to 8.5, 0.05 to 8.75, 0.05 to 9.0, 0.05 to 9.25, 0.05 to 9.5, 0.05 to 9.75, 0.05 to 10.0, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.1 to 6.0, 0.1 to 6.5, 0.1 to 7.0, 0.1 to 7.5, 0.1 to 8.0, 0.1 to 8.25, 0.1 to 8.5, 0.1 to 8.75, 0.1 to 9.0, 0.1 to 9.25, 0.1 to 9.5, 0.1 to 9.75, 0.1 to 10.0, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0,

0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.25, 0.5 to 8.5, 0.5 to 8.75, 0.5 to 9.0, 0.5 to 9.25, 0.5 to 9.5, 0.5 to 9.75, 0.5 to 10.0, 1.0 to 0.5, 1.0 to 1.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0 to 8.25, 1.0 to 8.5, 1.0 to 8.75, 1.0 to 9.0, 1.0 to 9.25, 1.0 to 9.5, 1.0 to 9.75, 1.0 to 10.0, 1.5 to 0.5, 1.5 to 1.0, 1.5 to 1.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.25, 1.5 to 8.5, 1.5 to 8.75, 1.5 to 9.0, 1.5 to 9.25, 1.5 to 9.5, 1.5 to 9.75, 1.5 to 10.0, 2.0 to 0.5, 2.0 to 1.0, 2.0 to 1.5, 2.0 to 2.0, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.0 to 6.0, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 8.0, 2.0 to 8.25, 2.0 to 8.5, 2.0 to 8.75, 2.0 to 9.0, 2.0 to 9.25, 2.0 to 9.5, 2.0 to 9.75, 2.0 to 10.0, 2.5 to 0.5, 2.5 to 1.0, 2.5 to 1.5, 2.5 to 2.0, 2.5 to 2.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 2.5 to 6.0, 2.5 to 6.5, 2.5 to 7.0, 2.5 to 7.5, 2.5 to 8.0, 2.5 to 8.25, 2.5 to 8.5, 2.5 to 8.75, 2.5 to 9.0, 2.5 to 9.25, 2.5 to 9.5, 2.5 to 9.75, or 2.5 to 10.0. More preferably, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0,

0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0,
1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5,
1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0,
1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0,
2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5,
2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0,
2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0,
or 3.0 to 5.5.

[0072]

"The sweetness intensity X2": 0.05 to 0.5, 0.05 to
1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0,
0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05
to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5,
0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0,
0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0,
1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5,
1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0,
1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0,
2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5,
2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0,
2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0,
3.0 to 5.5, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 6.0,
2.5 to 7.0, 2.5 to 7.5, 2.5 to 6.0, 2.5 to 6.5, 3.0 to 6.0,
3.0 to 6.5, 3.0 to 7.0, 3.0 to 7.5, 3.0 to 8.0, 3.0 to 8.5,
3.0 to 9.0, 3.0 to 9.5, 3.5 to 7.0, 3.5 to 7.5, 3.5 to 8.0,
4.5 to 8.5, 3.5 to 9.0, 3.5 to 9.5, 4.0 to 7.5, 4.0 to 8.0,
4.0 to 8.5, 4.0 to 9.0, 4.0 to 9.5, 3.5 to 8.5, 3.5 to

10.0, 3.5 to 10.5, 3.5 to 11.0, 3.5 to 11.5, 4.0 to 11.5, 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.5, 0.05 to 9.0, 0.05 to 9.5, 0.05 to 10.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.5, 0.5 to 9.0, 0.5 to 9.5, 0.5 to 10.0, 0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0 to 8.5, 1.0 to 9.0, 1.0 to 9.5, 1.0 to 10.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.5, 1.5 to 9.0, 1.5 to 9.5, 1.5 to 10.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to 11.5, 2.0 to 8.0, 2.0 to 8.5, 2.0 to 9.0, 2.0 to 9.5, 2.0 to 10.0, 2.0 to 10.5, 2.0 to 11.0, 2.0 to 11.5, 2.5 to 8.0, 2.5 to 8.5, 2.5 to 9.0, 2.5 to 9.5, 2.5 to 10.0, 2.5 to 10.5, 2.5 to 11.0, 2.5 to 11.5, 3.0 to 10.0, 3.0 to 10.5, 3.0 to 11.0, 3.0 to 11.5, 3.5 to 4.0, 3.5 to 4.5, 3.5 to 5.0, 3.5 to 5.5, 3.5 to 6.0, 3.5 to 6.5, 4.0 to 4.5, 4.0 to 5.0, 4.0 to 5.5, 4.0 to 6.0, 4.0 to 6.5, 4.0 to 7.0, 4.0 to 10.0, 4.0 to 10.5, or 4.0 to 11.0. More preferably, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, or 3.0 to 5.5.

[0073]

"The amount of sodium": 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 39mg/100ml, 5 to 38mg/100ml, 5 to 37mg/100ml, 5 to 36mg/100ml, 5 to 35mg/100ml, 5 to 34mg/100ml, 5 to 33mg/100ml, 5 to 32mg/100ml, 5 to 31mg/100ml, 5 to 30mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 39mg/100ml, 7 to 38mg/100ml, 7 to 37mg/100ml, 7 to 36mg/100ml, 7 to 35mg/100ml, 7 to 34mg/100ml, 7 to 33mg/100ml, 7 to 32mg/100ml, 7 to 31mg/100ml, 7 to 30mg/100ml, 10 to 45mg/100ml, 10 to 40mg/100ml, 10 to 39mg/100ml, 10 to 38mg/100ml, 10 to 37mg/100ml, 10 to 36mg/100ml, 10 to 35mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 39mg/100ml, 15 to 38mg/100ml, 15 to 37mg/100ml, 15 to 36mg/100ml, 15 to 35mg/100ml, 15 to 34mg/100ml, 15 to 33mg/100ml, 15 to 32mg/100ml, 15 to 31mg/100ml, 15 to 30mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to 39mg/100ml, 20 to 38mg/100ml, 20 to 37mg/100ml, 20 to 36mg/100ml, 20 to 35mg/100ml, 20 to 34mg/100ml, 20 to 33mg/100ml, 20 to 32mg/100ml, 20 to 31mg/100ml, or 20 to 30mg/100ml.

[0074]

"The sweetness intensity X3": 4.0 to 20, 4.0 to 15, 4.0 to 12.5, 4.0 to 10, 4.5 to 20, 4.5 to 15, 4.5 to 12.5, 4.5 to 10, 5.0 to 20, 5.0 to 15, 5.0 to 12.5, 5.0 to 10, 5.5 to 20, 5.5 to 15, 5.5 to 12.5, 5.5 to 10, 6.0 to 20, 6.0 to 15, 6.0 to 12.5, 6.0 to 10, 6.5 to 20, 6.5 to 15, 6.5 to 12.5, 6.5 to 10, 7.0 to 20, 7.0 to 15, 7.0 to 12.5,

7.0 to 10, 7.5 to 20, 7.5 to 15, 7.5 to 12.5, 7.5 to 10, 7.5 to 9, 7.5 to 8, 8.0 to 20, 8.0 to 20, 8.0 to 15, 8.0 to 12.5, 8.0 to 10, 8.5 to 20, 8.5 to 15, 8.5 to 12.5, 8.5 to 10, 9.0 to 20, 9.0 to 15, 9.0 to 12.5, 9.0 to 10, 9.5 to 20, 9.5 to 15, 9.5 to 12.5, 9.5 to 10, 10.0 to 20, 10.0 to 15, 10.0 to 12.5, 10.5 to 20, 10.5 to 15, or 10.5 to 12.5.

[0075]

"The amount of a high intensity sweetener P2 (ppm)":
20 to 550, 25 to 550, 30 to 550, 35 to 550, 40 to 550, 45 to 550, 50 to 550, 55 to 550, 20 to 540, 25 to 540, 30 to 540, 35 to 540, 40 to 540, 45 to 540, 50 to 540, 55 to 540, 20 to 530, 25 to 530, 30 to 530, 35 to 530, 40 to 530, 45 to 530, 50 to 530, 55 to 530, 20 to 520, 25 to 520, 30 to 520, 35 to 520, 40 to 520, 45 to 520, 50 to 520, 55 to 520, 20 to 510, 25 to 510, 30 to 510, 35 to 510, 40 to 510, 45 to 510, 50 to 510, 55 to 510, 20 to 505, 25 to 505, 30 to 505, 35 to 505, 40 to 505, 45 to 505, 50 to 505, 55 to 505, 20 to 500, 25 to 500, 30 to 500, 35 to 500, 40 to 500, 45 to 500, 50 to 500, 55 to 500, 20 to 495, 25 to 495, 30 to 495, 35 to 495, 40 to 495, 45 to 495, 50 to 495, 55 to 495, 20 to 490, 25 to 490, 30 to 490, 35 to 490, 40 to 490, 45 to 490, 50 to 490, or 55 to 490.

[0076]

The energy: 0 to 25Kcal/100ml, 0 to 20Kcal/100ml, 0 to 15Kcal/100ml, 0 to 10Kcal/100ml, 0 to 5Kcal/100ml, 5 to 25Kcal/100ml, 5 to 20Kcal/100ml, 5 to 15Kcal/100ml, 5 to

10Kcal/100ml, 10 to 25Kcal/100ml, 10 to 20Kcal/100ml, 10 to 15Kcal/100ml, 15 to 25Kcal/100ml, 15 to 20Kcal/100ml, 20 to 25Kcal/100ml, 0 to 24Kcal/100ml, 0 to 8Kcal/100ml, 0 to 4Kcal/100ml, 4 to 24Kcal/100ml, 4 to 8Kcal/100ml, or 8 to 24Kcal/100ml.

[0077]

The natural sugar (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose) and the sodium form in fruit juice beverage A of the present invention are defined as described in the section of the fruit juice of the present invention.

[0078]

In another specific embodiment, the present invention provides the following fruit juice beverage (hereinafter, referred to as "fruit juice beverage B of the present invention").

A fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X1;

(b) a luo han guo extract in an amount corresponding to a sweetness intensity X2; and

(c) 5 to 50 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0079]

In fruit juice beverage B of the present invention, the preferable ranges of the sweetness intensity X1, the sweetness intensity X2, the sodium content, the sweetness intensity X3, the luo han guo extract content P2, and the energy are, for example, as follows. These numerical values can be arbitrarily combined in a range that satisfies $0.1 < (X1 + X2) \leq 20$.

[0080]

"The sweetness intensity X1": 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.25, 0.05 to 8.5, 0.05 to 8.75, 0.05 to 9.0, 0.05 to 9.25, 0.05 to 9.5, 0.05 to 9.75, 0.05 to 10.0, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.1 to 6.0, 0.1 to 6.5, 0.1 to 7.0, 0.1 to 7.5, 0.1 to 8.0, 0.1 to 8.25, 0.1 to 8.5, 0.1 to 8.75, 0.1 to 9.0, 0.1 to 9.25, 0.1 to 9.5, 0.1 to 9.75, 0.1 to 10.0, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.25, 0.5 to 8.5, 0.5 to 8.75, 0.5 to 9.0, 0.5 to 9.25, 0.5 to 9.5, 0.5 to 9.75, 0.5 to 10.0, 1.0 to 0.5, 1.0 to 1.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.0 to 6.0, 1.0 to 6.5,

1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0 to 8.25, 1.0 to 8.5, 1.0 to 8.75, 1.0 to 9.0, 1.0 to 9.25, 1.0 to 9.5, 1.0 to 9.75, 1.0 to 10.0, 1.5 to 0.5, 1.5 to 1.0, 1.5 to 1.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.25, 1.5 to 8.5, 1.5 to 8.75, 1.5 to 9.0, 1.5 to 9.25, 1.5 to 9.5, 1.5 to 9.75, 1.5 to 10.0, 2.0 to 0.5, 2.0 to 1.0, 2.0 to 1.5, 2.0 to 2.0, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.0 to 6.0, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 8.0, 2.0 to 8.25, 2.0 to 8.5, 2.0 to 8.75, 2.0 to 9.0, 2.0 to 9.25, 2.0 to 9.5, 2.0 to 9.75, 2.0 to 10.0, 2.5 to 0.5, 2.5 to 1.0, 2.5 to 1.5, 2.5 to 2.0, 2.5 to 2.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 2.5 to 6.0, 2.5 to 6.5, 2.5 to 7.0, 2.5 to 7.5, 2.5 to 8.0, 2.5 to 8.25, 2.5 to 8.5, 2.5 to 8.75, 2.5 to 9.0, 2.5 to 9.25, 2.5 to 9.5, 2.5 to 9.75, 2.5 to 10.0, 0 to 10.5, 0 to 11.0, 0 to 11.5, 0 to 12.0, 0 to 12.5, 0 to 13.0, 0 to 13.5, 0 to 14.0, 0 to 14.5, 0 to 15.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5, 0.05 to 12.0, 0.05 to 12.5, 0.05 to 13.0, 0.05 to 13.5, 0.05 to 14.0, 0.05 to 14.5, 0.05 to 15.0, 0.1 to 10.5, 0.1 to 11.0, 0.1 to 11.5, 0.1 to 12.0, 0.1 to 12.5, 0.1 to 13.0, 0.1 to 13.5, 0.1 to 14.0, 0.1 to 14.5, 0.1 to 15.0, 0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to 12.5, 0.5 to 13.0, 0.5 to 13.5, 0.5 to 14.0, 0.5 to 14.5, 0.5 to 15.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to

12.0, 1.0 to 12.5, 1.0 to 13.0, 1.0 to 13.5, 1.0 to 14.0, 1.0 to 14.5, 1.0 to 15.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to 11.5, 1.5 to 12.0, 1.5 to 12.5, 1.5 to 13.0, 1.5 to 13.5, 1.5 to 14.0, 1.5 to 14.5, 1.5 to 15.0, 2.0 to 10.5, 2.0 to 11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 12.5, 2.0 to 13.0, 2.0 to 13.5, 2.0 to 14.0, 2.0 to 14.5, 2.0 to 15.0, 2.5 to 10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 12.5, 2.5 to 13.0, 2.5 to 13.5, 2.5 to 14.0, 2.5 to 14.5, or 2.5 to 15.0. More preferably, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, or 3.0 to 5.5.

[0081]

"The sweetness intensity X2": 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0,

0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0,
1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5,
1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0,
1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0,
2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5,
2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0,
2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0,
3.0 to 5.5, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 6.0,
2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.5 to 7.0, 2.5 to 7.5,
2.5 to 6.0, 2.5 to 6.5, 2.5 to 7.0, 2.5 to 7.5, 3.0 to 7.5,
3.0 to 6.0, 3.0 to 6.5, 3.0 to 7.0, 3.0 to 7.5, 3.0 to 8.0,
3.0 to 8.5, 3.0 to 9.0, 3.0 to 9.5, 3.5 to 7.0, 3.5 to 7.5,
3.5 to 8.0, 4.5 to 8.5, 3.5 to 9.0, 3.5 to 9.5, 4.0 to 7.5,
4.0 to 8.0, 4.0 to 8.5, 4.0 to 9.0, 4.0 to 9.5, 3.5 to 8.5,
3.5 to 9.0, 3.5 to 9.5, 3.5 to 10.0, 3.5 to 10.5, 3.5 to
11.0, 3.5 to 11.5, 4.0 to 11.5, 0.05 to 6.0, 0.05 to 6.5,
0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.5, 0.05
to 9.0, 0.05 to 9.5, 0.05 to 10.0, 0.05 to 10.5, 0.05 to
11.0, 0.05 to 11.5, 0.05 to 12.0, 0.05 to 13.0, 0.05 to
14.0, 0.05 to 15.0, 0.05 to 16.0, 0.05 to 17.0, 0.05 to
18.0, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5
to 8.0, 0.5 to 8.5, 0.5 to 9.0, 0.5 to 9.5, 0.5 to 10.0,
0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to
13.0, 0.5 to 14.0, 0.5 to 15.0, 0.5 to 16.0, 0.5 to 17.0,
0.5 to 18.0, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to
7.5, 1.0 to 8.0, 1.0 to 8.5, 1.0 to 9.0, 1.0 to 9.5, 1.0
to 10.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to

12.0, 1.0 to 13.0, 1.0 to 14.0, 1.0 to 15.0, 1.0 to 16.0, 1.0 to 17.0, 1.0 to 18.0, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.5, 1.5 to 9.0, 1.5 to 9.5, 1.5 to 10.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to 11.5, 1.5 to 12.0, 1.5 to 13.0, 1.5 to 14.0, 1.5 to 15.0, 1.5 to 16.0, 1.5 to 17.0, 1.5 to 18.0, 2.0 to 8.0, 2.0 to 8.5, 2.0 to 9.0, 2.0 to 9.5, 2.0 to 10.0, 2.0 to 10.5, 2.0 to 11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 13.0, 2.0 to 14.0, 2.0 to 15.0, 2.0 to 16.0, 2.0 to 17.0, 2.0 to 18.0, 2.5 to 8.0, 2.5 to 8.5, 2.5 to 9.0, 2.5 to 9.5, 2.5 to 10.0, 2.5 to 10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 13.0, 2.5 to 14.0, 2.5 to 15.0, 2.5 to 16.0, 2.5 to 17.0, 2.5 to 18.0, 3.0 to 10.0, 3.0 to 10.5, 3.0 to 11.0, 3.0 to 11.5, 3.0 to 12.0, 3.0 to 13.0, 3.0 to 14.0, 3.0 to 15.0, 3.0 to 16.0, 3.0 to 17.0, 3.0 to 18.0, 3.5 to 4.0, 3.5 to 4.5, 3.5 to 5.0, 3.5 to 5.5, 3.5 to 6.0, 3.5 to 6.5, 3.5 to 12.0, 3.5 to 13.0, 3.5 to 14.0, 3.5 to 15.0, 3.5 to 16.0, 3.5 to 17.0, 3.5 to 18.0, 4.0 to 4.5, 4.0 to 5.0, 4.0 to 5.5, 4.0 to 6.0, 4.0 to 6.5, 4.0 to 7.0, 4.0 to 10.0, 4.0 to 10.5, 4.0 to 11.0, 4.0 to 12.0, 4.0 to 13.0, 4.0 to 14.0, 4.0 to 15.0, 4.0 to 16.0, 4.0 to 17.0, or 4.0 to 18.0. More preferably, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0,

1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5,
1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0,
1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0,
2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5,
2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5,
3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, or 3.0 to 5.5.

[0082]

"The amount of sodium": 5 to 45mg/100ml, 5 to
40mg/100ml, 5 to 39mg/100ml, 5 to 38mg/100ml, 5 to
37mg/100ml, 5 to 36mg/100ml, 5 to 35mg/100ml, 5 to
34mg/100ml, 5 to 33mg/100ml, 5 to 32mg/100ml, 5 to
31mg/100ml, 5 to 30mg/100ml, 7 to 45mg/100ml, 7 to
40mg/100ml, 7 to 39mg/100ml, 7 to 38mg/100ml, 7 to
37mg/100ml, 7 to 36mg/100ml, 7 to 35mg/100ml, 7 to
34mg/100ml, 7 to 33mg/100ml, 7 to 32mg/100ml, 7 to
31mg/100ml, 7 to 30mg/100ml, 10 to 45mg/100ml, 10 to
40mg/100ml, 10 to 39mg/100ml, 10 to 38mg/100ml, 10 to
37mg/100ml, 10 to 36mg/100ml, 10 to 35mg/100ml, 15 to
45mg/100ml, 15 to 40mg/100ml, 15 to 39mg/100ml, 15 to
38mg/100ml, 15 to 37mg/100ml, 15 to 36mg/100ml, 15 to
35mg/100ml, 15 to 34mg/100ml, 15 to 33mg/100ml, 15 to
32mg/100ml, 15 to 31mg/100ml, 15 to 30mg/100ml, 20 to
45mg/100ml, 20 to 40mg/100ml, 20 to 39mg/100ml, 20 to
38mg/100ml, 20 to 37mg/100ml, 20 to 36mg/100ml, 20 to
35mg/100ml, 20 to 34mg/100ml, 20 to 33mg/100ml, 20 to
32mg/100ml, 20 to 31mg/100ml, or 20 to 30mg/100ml.

[0083]

"The sweetness intensity X3": 4.0 to 20, 4.0 to 15, 4.0 to 12.5, 4.0 to 10, 4.5 to 20, 4.5 to 15, 4.5 to 12.5, 4.5 to 10, 5.0 to 20, 5.0 to 15, 5.0 to 12.5, 5.0 to 10, 5.5 to 20, 5.5 to 15, 5.5 to 12.5, 5.5 to 10, 6.0 to 20, 6.0 to 15, 6.0 to 12.5, 6.0 to 10, 6.5 to 20, 6.5 to 15, 6.5 to 12.5, 6.5 to 10, 7.0 to 20, 7.0 to 15, 7.0 to 12.5, 7.0 to 10, 7.5 to 20, 7.5 to 15, 7.5 to 12.5, 7.5 to 10, 7.5 to 9, 7.5 to 8, 8.0 to 20, 8.0 to 20, 8.0 to 15, 8.0 to 12.5, 8.0 to 10, 8.5 to 20, 8.5 to 15, 8.5 to 12.5, 8.5 to 10, 9.0 to 20, 9.0 to 15, 9.0 to 12.5, 9.0 to 10, 9.5 to 20, 9.5 to 15, 9.5 to 12.5, 9.5 to 10, 10.0 to 20, 10.0 to 15, 10.0 to 12.5, 10.5 to 20, 10.5 to 15, 10.5 to 12.5, 4.0 to 18, 4.0 to 16, 4.0 to 15.5, 4.0 to 14, 4.5 to 18, 4.5 to 16, 4.5 to 15.5, 4.5 to 14, 5.0 to 18, 5.0 to 16, 5.0 to 15.5, 5.0 to 14, 5.5 to 18, 5.5 to 16, 5.5 to 15.5, 5.5 to 14, 6.0 to 18, 6.0 to 16, 6.0 to 15.5, 6.0 to 14, 6.5 to 18, 6.5 to 16, 6.5 to 15.5, 6.5 to 14, 7.0 to 18, 7.0 to 16, 7.0 to 15.5, 7.0 to 14, 7.5 to 18, 7.5 to 16, 7.5 to 15.5, 7.5 to 14, 7.5 to 9, 7.5 to 8, 8.0 to 18, 8.0 to 18, 8.0 to 16, 8.0 to 15.5, 8.0 to 14, 8.5 to 18, 8.5 to 16, 8.5 to 15.5, 8.5 to 14, 9.0 to 18, 9.0 to 16, 9.0 to 15.5, 9.0 to 14, 9.5 to 18, 9.5 to 16, 9.5 to 15.5, 9.5 to 14, 10.0 to 18, 10.0 to 16, 10.0 to 15.5, 10.5 to 18, 10.5 to 16, or 10.5 to 15.5.

[0084]

"The amount of Luo han guo extract P2 (ppm)": 20 to 550, 25 to 550, 30 to 550, 35 to 550, 40 to 550, 45 to 550,

50 to 550, 55 to 550, 20 to 540, 25 to 540, 30 to 540, 35 to 540, 40 to 540, 45 to 540, 50 to 540, 55 to 540, 20 to 530, 25 to 530, 30 to 530, 35 to 530, 40 to 530, 45 to 530, 50 to 530, 55 to 530, 20 to 520, 25 to 520, 30 to 520, 35 to 520, 40 to 520, 45 to 520, 50 to 520, 55 to 520, 20 to 510, 25 to 510, 30 to 510, 35 to 510, 40 to 510, 45 to 510, 50 to 510, 55 to 510, 20 to 505, 25 to 505, 30 to 505, 35 to 505, 40 to 505, 45 to 505, 50 to 505, 55 to 505, 20 to 500, 25 to 500, 30 to 500, 35 to 500, 40 to 500, 45 to 500, 50 to 500, 55 to 500, 20 to 495, 25 to 495, 30 to 495, 35 to 495, 40 to 495, 45 to 495, 50 to 495, 55 to 495, 20 to 490, 25 to 490, 30 to 490, 35 to 490, 40 to 490, 45 to 490, 50 to 490, 55 to 490, 1 to 1500, 1 to 1200, 5 to 1200, 1 to 1000, 5 to 1000, 10 to 1000, 1 to 900, 5 to 900, 10 to 900, 15 to 900, 20 to 900, 25 to 900, 30 to 900, 35 to 900, 40 to 900, 45 to 900, 50 to 900, 55 to 900, 1 to 800, 5 to 800, 10 to 800, 15 to 800, 20 to 800, 25 to 800, 30 to 800, 35 to 800, 40 to 800, 45 to 800, 50 to 800, 55 to 800, 1 to 700, 5 to 700, 10 to 700, 15 to 700, 20 to 700, 25 to 700, 30 to 700, 35 to 700, 40 to 700, 45 to 700, 50 to 700, 55 to 700, 1 to 600, 5 to 600, 10 to 600, 15 to 600, 20 to 600, 25 to 600, 30 to 600, 35 to 600, 40 to 600, 45 to 600, 50 to 600, 55 to 600, 1 to 550, 1 to 540, 1 to 530, 1 to 520, 1 to 510, 1 to 505, 1 to 500, 1 to 495, 1 to 490, 5 to 550, 5 to 540, 5 to 530, 5 to 520, 5 to 510, 5 to 505, 5 to 500, 5 to 495, 5 to 490, 10 to 550, 10 to 540, 10 to 530, 10 to 520, 10 to 510, 10 to 505, 10 to 500, 10 to 495,

10 to 490, 15 to 550, 15 to 550, 15 to 530, 15 to 520, 15 to 510, 15 to 505, 15 to 500, 15 to 495, or 15 to 490.

[0085]

The energy: 0 to 50Kcal/100ml, 0 to 45Kcal/100ml, 0 to 40Kcal/100ml, 0 to 35Kcal/100ml, 0 to 30Kcal/100ml, 0 to 25Kcal/100ml, 0 to 20Kcal/100ml, 0 to 15Kcal/100ml, 0 to 10Kcal/100ml, 0 to 5Kcal/100ml, 5 to 50Kcal/100ml, 5 to 45Kcal/100ml, 5 to 40Kcal/100ml, 5 to 35Kcal/100ml, 5 to 30Kcal/100ml, 5 to 25Kcal/100ml, 5 to 20Kcal/100ml, 5 to 15Kcal/100ml, 5 to 10Kcal/100ml, 10 to 50Kcal/100ml, 10 to 45Kcal/100ml, 10 to 40Kcal/100ml, 10 to 35Kcal/100ml, 10 to 30Kcal/100ml, 10 to 25Kcal/100ml, 10 to 20Kcal/100ml, 10 to 15Kcal/100ml, 15 to 50Kcal/100ml, 15 to 45Kcal/100ml, 15 to 40Kcal/100ml, 15 to 35Kcal/100ml, 15 to 30Kcal/100ml, 15 to 25Kcal/100ml, 15 to 20Kcal/100ml, 20 to 50Kcal/100ml, 20 to 45Kcal/100ml, 20 to 40Kcal/100ml, 20 to 35Kcal/100ml, 20 to 30Kcal/100ml, 20 to 25Kcal/100ml, 25 to 50Kcal/100ml, 25 to 45Kcal/100ml, 25 to 40Kcal/100ml, 25 to 35Kcal/100ml, 25 to 30Kcal/100ml, 0 to 32Kcal/100ml, 0 to 24Kcal/100ml, 0 to 8Kcal/100ml, 0 to 4Kcal/100ml, 4 to 32Kcal/100ml, 4 to 24Kcal/100ml, 4 to 8Kcal/100ml, 8 to 32Kcal/100ml, 8 to 24Kcal/100ml, or 24 to 32Kcal/100ml.

[0086]

In addition, the natural sugar (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose) and the sodium form in fruit juice beverage B of

the present invention are defined as described in the section of the fruit juice beverage of the present invention.

[0087]

In another specific embodiment, the present invention provides the following fruit juice beverage (hereinafter, referred to as "fruit juice beverage C of the present invention").

A fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 ;

(b) mogroside V in an amount corresponding to a sweetness intensity X_2 ; and

(c) 5 to 50 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X_3 exhibited by the components (a) to (c), and $0.1 < (X_1 + X_2) \leq 20$ is satisfied.

[0088]

In fruit juice beverage C of the present invention, the preferable ranges of the sweetness intensity X_1 , the sweetness intensity X_2 , the sodium content, the sweetness intensity X_3 , the mogroside V content P2 (ppm), and the energy are, for example, as follows. These numerical values can be arbitrarily combined in a range that satisfies $0.1 < (X_1 + X_2) \leq 20$.

[0089]

"The sweetness intensity X1": 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.25, 0.05 to 8.5, 0.05 to 8.75, 0.05 to 9.0, 0.05 to 9.25, 0.05 to 9.5, 0.05 to 9.75, 0.05 to 10.0, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.1 to 6.0, 0.1 to 6.5, 0.1 to 7.0, 0.1 to 7.5, 0.1 to 8.0, 0.1 to 8.25, 0.1 to 8.5, 0.1 to 8.75, 0.1 to 9.0, 0.1 to 9.25, 0.1 to 9.5, 0.1 to 9.75, 0.1 to 10.0, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.25, 0.5 to 8.5, 0.5 to 8.75, 0.5 to 9.0, 0.5 to 9.25, 0.5 to 9.5, 0.5 to 9.75, 0.5 to 10.0, 1.0 to 0.5, 1.0 to 1.0, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0 to 8.25, 1.0 to 8.5, 1.0 to 8.75, 1.0 to 9.0, 1.0 to 9.25, 1.0 to 9.5, 1.0 to 9.75, 1.0 to 10.0, 1.5 to 0.5, 1.5 to 1.0, 1.5 to 1.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.25, 1.5 to 8.5, 1.5 to 8.75, 1.5 to 9.0, 1.5 to 9.25, 1.5 to 9.5, 1.5 to 9.75, 1.5 to 10.0, 2.0 to 0.5, 2.0 to 1.0, 2.0 to 1.5,

2.0 to 2.0, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0,
2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.0 to 6.0, 2.0 to 6.5,
2.0 to 7.0, 2.0 to 7.5, 2.0 to 8.0, 2.0 to 8.25, 2.0 to
8.5, 2.0 to 8.75, 2.0 to 9.0, 2.0 to 9.25, 2.0 to 9.5, 2.0
to 9.75, 2.0 to 10.0, 2.5 to 0.5, 2.5 to 1.0, 2.5 to 1.5,
2.5 to 2.0, 2.5 to 2.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0,
2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 2.5 to 6.0, 2.5 to 6.5,
2.5 to 7.0, 2.5 to 7.5, 2.5 to 8.0, 2.5 to 8.25, 2.5 to
8.5, 2.5 to 8.75, 2.5 to 9.0, 2.5 to 9.25, 2.5 to 9.5, 2.5
to 9.75, 2.5 to 10.0, 0 to 10.5, 0 to 11.0, 0 to 11.5, 0
to 12.0, 0 to 12.5, 0 to 13.0, 0 to 13.5, 0 to 14.0, 0 to
14.5, 0 to 15.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5,
0.05 to 12.0, 0.05 to 12.5, 0.05 to 13.0, 0.05 to 13.5,
0.05 to 14.0, 0.05 to 14.5, 0.05 to 15.0, 0.1 to 10.5, 0.1
to 11.0, 0.1 to 11.5, 0.1 to 12.0, 0.1 to 12.5, 0.1 to
13.0, 0.1 to 13.5, 0.1 to 14.0, 0.1 to 14.5, 0.1 to 15.0,
0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to
12.5, 0.5 to 13.0, 0.5 to 13.5, 0.5 to 14.0, 0.5 to 14.5,
0.5 to 15.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to
12.0, 1.0 to 12.5, 1.0 to 13.0, 1.0 to 13.5, 1.0 to 14.0,
1.0 to 14.5, 1.0 to 15.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to
11.5, 1.5 to 12.0, 1.5 to 12.5, 1.5 to 13.0, 1.5 to 13.5,
1.5 to 14.0, 1.5 to 14.5, 1.5 to 15.0, 2.0 to 10.5, 2.0 to
11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 12.5, 2.0 to 13.0,
2.0 to 13.5, 2.0 to 14.0, 2.0 to 14.5, 2.0 to 15.0, 2.5 to
10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 12.5,
2.5 to 13.0, 2.5 to 13.5, 2.5 to 14.0, 2.5 to 14.5, or 2.5

to 15.0. More preferably, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.1 to 5.9, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, or 3.0 to 5.5.

[0090]

"The sweetness intensity X2": 0.05 to 0.5, 0.05 to 1.0, 0.05 to 1.5, 0.05 to 2.0, 0.05 to 2.5, 0.05 to 3.0, 0.05 to 3.5, 0.05 to 4.0, 0.05 to 4.5, 0.05 to 5.0, 0.05 to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, 3.0 to 5.5, 2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.0 to 6.0,

2.0 to 6.5, 2.0 to 7.0, 2.0 to 7.5, 2.5 to 7.0, 2.5 to 7.5, 2.5 to 6.0, 2.5 to 6.5, 2.5 to 7.0, 2.5 to 7.5, 3.0 to 7.5, 3.0 to 6.0, 3.0 to 6.5, 3.0 to 7.0, 3.0 to 7.5, 3.0 to 8.0, 3.0 to 8.5, 3.0 to 9.0, 3.0 to 9.5, 3.5 to 7.0, 3.5 to 7.5, 3.5 to 8.0, 4.5 to 8.5, 3.5 to 9.0, 3.5 to 9.5, 4.0 to 7.5, 4.0 to 8.0, 4.0 to 8.5, 4.0 to 9.0, 4.0 to 9.5, 3.5 to 8.5, 3.5 to 9.0, 3.5 to 9.5, 3.5 to 10.0, 3.5 to 10.5, 3.5 to 11.0, 3.5 to 11.5, 4.0 to 11.5, 0.05 to 6.0, 0.05 to 6.5, 0.05 to 7.0, 0.05 to 7.5, 0.05 to 8.0, 0.05 to 8.5, 0.05 to 9.0, 0.05 to 9.5, 0.05 to 10.0, 0.05 to 10.5, 0.05 to 11.0, 0.05 to 11.5, 0.05 to 12.0, 0.05 to 13.0, 0.05 to 14.0, 0.05 to 15.0, 0.05 to 16.0, 0.05 to 17.0, 0.05 to 18.0, 0.5 to 6.0, 0.5 to 6.5, 0.5 to 7.0, 0.5 to 7.5, 0.5 to 8.0, 0.5 to 8.5, 0.5 to 9.0, 0.5 to 9.5, 0.5 to 10.0, 0.5 to 10.5, 0.5 to 11.0, 0.5 to 11.5, 0.5 to 12.0, 0.5 to 13.0, 0.5 to 14.0, 0.5 to 15.0, 0.5 to 16.0, 0.5 to 17.0, 0.5 to 18.0, 1.0 to 6.0, 1.0 to 6.5, 1.0 to 7.0, 1.0 to 7.5, 1.0 to 8.0, 1.0 to 8.5, 1.0 to 9.0, 1.0 to 9.5, 1.0 to 10.0, 1.0 to 10.5, 1.0 to 11.0, 1.0 to 11.5, 1.0 to 12.0, 1.0 to 13.0, 1.0 to 14.0, 1.0 to 15.0, 1.0 to 16.0, 1.0 to 17.0, 1.0 to 18.0, 1.5 to 6.0, 1.5 to 6.5, 1.5 to 7.0, 1.5 to 7.5, 1.5 to 8.0, 1.5 to 8.5, 1.5 to 9.0, 1.5 to 9.5, 1.5 to 10.0, 1.5 to 10.5, 1.5 to 11.0, 1.5 to 11.5, 1.5 to 12.0, 1.5 to 13.0, 1.5 to 14.0, 1.5 to 15.0, 1.5 to 16.0, 1.5 to 17.0, 1.5 to 18.0, 2.0 to 8.0, 2.0 to 8.5, 2.0 to 9.0, 2.0 to 9.5, 2.0 to 10.0, 2.0 to 10.5, 2.0 to 11.0, 2.0 to 11.5, 2.0 to 12.0, 2.0 to 13.0, 2.0 to

14.0, 2.0 to 15.0, 2.0 to 16.0, 2.0 to 17.0, 2.0 to 18.0, 2.5 to 8.0, 2.5 to 8.5, 2.5 to 9.0, 2.5 to 9.5, 2.5 to 10.0, 2.5 to 10.5, 2.5 to 11.0, 2.5 to 11.5, 2.5 to 12.0, 2.5 to 13.0, 2.5 to 14.0, 2.5 to 15.0, 2.5 to 16.0, 2.5 to 17.0, 2.5 to 18.0, 3.0 to 10.0, 3.0 to 10.5, 3.0 to 11.0, 3.0 to 11.5, 3.0 to 12.0, 3.0 to 13.0, 3.0 to 14.0, 3.0 to 15.0, 3.0 to 16.0, 3.0 to 17.0, 3.0 to 18.0, 3.5 to 4.0, 3.5 to 4.5, 3.5 to 5.0, 3.5 to 5.5, 3.5 to 6.0, 3.5 to 6.5, 3.5 to 12.0, 3.5 to 13.0, 3.5 to 14.0, 3.5 to 15.0, 3.5 to 16.0, 3.5 to 17.0, 3.5 to 18.0, 4.0 to 4.5, 4.0 to 5.0, 4.0 to 5.5, 4.0 to 6.0, 4.0 to 6.5, 4.0 to 7.0, 4.0 to 10.0, 4.0 to 10.5, 4.0 to 11.0, 4.0 to 12.0, 4.0 to 13.0, 4.0 to 14.0, 4.0 to 15.0, 4.0 to 16.0, 4.0 to 17.0, or 4.0 to 18.0. More preferably, 0.1 to 0.5, 0.1 to 1.0, 0.1 to 1.5, 0.1 to 2.0, 0.1 to 2.5, 0.1 to 3.0, 0.1 to 3.5, 0.1 to 4.0, 0.1 to 4.5, 0.1 to 5.0, 0.1 to 5.5, 0.5 to 1.0, 0.5 to 1.5, 0.5 to 2.0, 0.5 to 2.5, 0.5 to 3.0, 0.5 to 3.5, 0.5 to 4.0, 0.5 to 4.5, 0.5 to 5.0, 0.5 to 5.5, 1.0 to 1.5, 1.0 to 2.0, 1.0 to 2.5, 1.0 to 3.0, 1.0 to 3.5, 1.0 to 4.0, 1.0 to 4.5, 1.0 to 5.0, 1.0 to 5.5, 1.5 to 2.0, 1.5 to 2.5, 1.5 to 3.0, 1.5 to 3.5, 1.5 to 4.0, 1.5 to 4.5, 1.5 to 5.0, 1.5 to 5.5, 2.0 to 2.5, 2.0 to 3.0, 2.0 to 3.5, 2.0 to 4.0, 2.0 to 4.5, 2.0 to 5.0, 2.0 to 5.5, 2.5 to 3.0, 2.5 to 3.5, 2.5 to 4.0, 2.5 to 4.5, 2.5 to 5.0, 2.5 to 5.5, 3.0 to 3.5, 3.0 to 4.0, 3.0 to 4.5, 3.0 to 5.0, or 3.0 to 5.5.

[0091]

"The amount of sodium": 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 39mg/100ml, 5 to 38mg/100ml, 5 to 37mg/100ml, 5 to 36mg/100ml, 5 to 35mg/100ml, 5 to 34mg/100ml, 5 to 33mg/100ml, 5 to 32mg/100ml, 5 to 31mg/100ml, 5 to 30mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 39mg/100ml, 7 to 38mg/100ml, 7 to 37mg/100ml, 7 to 36mg/100ml, 7 to 35mg/100ml, 7 to 34mg/100ml, 7 to 33mg/100ml, 7 to 32mg/100ml, 7 to 31mg/100ml, 7 to 30mg/100ml, 10 to 45mg/100ml, 10 to 40mg/100ml, 10 to 39mg/100ml, 10 to 38mg/100ml, 10 to 37mg/100ml, 10 to 36mg/100ml, 10 to 35mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 39mg/100ml, 15 to 38mg/100ml, 15 to 37mg/100ml, 15 to 36mg/100ml, 15 to 35mg/100ml, 15 to 34mg/100ml, 15 to 33mg/100ml, 15 to 32mg/100ml, 15 to 31mg/100ml, 15 to 30mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to 39mg/100ml, 20 to 38mg/100ml, 20 to 37mg/100ml, 20 to 36mg/100ml, 20 to 35mg/100ml, 20 to 34mg/100ml, 20 to 33mg/100ml, 20 to 32mg/100ml, 20 to 31mg/100ml, or 20 to 30mg/100ml.

[0092]

"The sweetness intensity X3": 4.0 to 20, 4.0 to 15, 4.0 to 12.5, 4.0 to 10, 4.5 to 20, 4.5 to 15, 4.5 to 12.5, 4.5 to 10, 5.0 to 20, 5.0 to 15, 5.0 to 12.5, 5.0 to 10, 5.5 to 20, 5.5 to 15, 5.5 to 12.5, 5.5 to 10, 6.0 to 20, 6.0 to 15, 6.0 to 12.5, 6.0 to 10, 6.5 to 20, 6.5 to 15, 6.5 to 12.5, 6.5 to 10, 7.0 to 20, 7.0 to 15, 7.0 to 12.5, 7.0 to 10, 7.5 to 20, 7.5 to 15, 7.5 to 12.5, 7.5 to 10,

7.5 to 9, 7.5 to 8, 8.0 to 20, 8.0 to 20, 8.0 to 15, 8.0 to 12.5, 8.0 to 10, 8.5 to 20, 8.5 to 15, 8.5 to 12.5, 8.5 to 10, 9.0 to 20, 9.0 to 15, 9.0 to 12.5, 9.0 to 10, 9.5 to 20, 9.5 to 15, 9.5 to 12.5, 9.5 to 10, 10.0 to 20, 10.0 to 15, 10.0 to 12.5, 10.5 to 20, 10.5 to 15, 10.5 to 12.5, 4.0 to 18, 4.0 to 16, 4.0 to 15.5, 4.0 to 14, 4.5 to 18, 4.5 to 16, 4.5 to 15.5, 4.5 to 14, 5.0 to 18, 5.0 to 16, 5.0 to 15.5, 5.0 to 14, 5.5 to 18, 5.5 to 16, 5.5 to 15.5, 5.5 to 14, 6.0 to 18, 6.0 to 16, 6.0 to 15.5, 6.0 to 14, 6.5 to 18, 6.5 to 16, 6.5 to 15.5, 6.5 to 14, 7.0 to 18, 7.0 to 16, 7.0 to 15.5, 7.0 to 14, 7.5 to 18, 7.5 to 16, 7.5 to 15.5, 7.5 to 14, 7.5 to 9, 7.5 to 8, 8.0 to 18, 8.0 to 18, 8.0 to 16, 8.0 to 15.5, 8.0 to 14, 8.5 to 18, 8.5 to 16, 8.5 to 15.5, 8.5 to 14, 9.0 to 18, 9.0 to 16, 9.0 to 15.5, 9.0 to 14, 9.5 to 18, 9.5 to 16, 9.5 to 15.5, 9.5 to 14, 10.0 to 18, 10.0 to 16, 10.0 to 15.5, 10.5 to 18, 10.5 to 16, or 10.5 to 15.5.

[0093]

"The amount of mogroside V P2 (ppm)": 20 to 550, 25 to 550, 30 to 550, 35 to 550, 40 to 550, 45 to 550, 50 to 550, 55 to 550, 20 to 540, 25 to 540, 30 to 540, 35 to 540, 40 to 540, 45 to 540, 50 to 540, 55 to 540, 20 to 530, 25 to 530, 30 to 530, 35 to 530, 40 to 530, 45 to 530, 50 to 530, 55 to 530, 20 to 520, 25 to 520, 30 to 520, 35 to 520, 40 to 520, 45 to 520, 50 to 520, 55 to 520, 20 to 510, 25 to 510, 30 to 510, 35 to 510, 40 to 510, 45 to 510, 50 to 510, 55 to 510, 20 to 505, 25 to 505, 30 to 505, 35 to 505,

40 to 505, 45 to 505, 50 to 505, 55 to 505, 20 to 500, 25 to 500, 30 to 500, 35 to 500, 40 to 500, 45 to 500, 50 to 500, 55 to 500, 20 to 495, 25 to 495, 30 to 495, 35 to 495, 40 to 495, 45 to 495, 50 to 495, 55 to 495, 20 to 490, 25 to 490, 30 to 490, 35 to 490, 40 to 490, 45 to 490, 50 to 490, 55 to 490, 1 to 700, 5 to 700, 10 to 700, 15 to 700, 20 to 700, 25 to 700, 30 to 700, 35 to 700, 40 to 700, 45 to 700, 50 to 700, 55 to 700, 1 to 600, 5 to 600, 10 to 600, 15 to 600, 20 to 600, 25 to 600, 30 to 600, 35 to 600, 40 to 600, 45 to 600, 50 to 600, 55 to 600, 1 to 550, 1 to 540, 1 to 530, 1 to 520, 1 to 510, 1 to 505, 1 to 500, 1 to 495, 1 to 490, 5 to 550, 5 to 540, 5 to 530, 5 to 520, 5 to 510, 5 to 505, 5 to 500, 5 to 495, 5 to 490, 10 to 550, 10 to 540, 10 to 530, 10 to 520, 10 to 510, 10 to 505, 10 to 500, 10 to 495, 10 to 490, 15 to 550, 15 to 550, 15 to 530, 15 to 520, 15 to 510, 15 to 505, 15 to 500, 15 to 495, or 15 to 490.

[0094]

The energy: 0 to 50Kcal/100ml, 0 to 45Kcal/100ml, 0 to 40Kcal/100ml, 0 to 35Kcal/100ml, 0 to 30Kcal/100ml, 0 to 25Kcal/100ml, 0 to 20Kcal/100ml, 0 to 15Kcal/100ml, 0 to 10Kcal/100ml, 0 to 5Kcal/100ml, 5 to 50Kcal/100ml, 5 to 45Kcal/100ml, 5 to 40Kcal/100ml, 5 to 35Kcal/100ml, 5 to 30Kcal/100ml, 5 to 25Kcal/100ml, 5 to 20Kcal/100ml, 5 to 15Kcal/100ml, 5 to 10Kcal/100ml, 10 to 50Kcal/100ml, 10 to 45Kcal/100ml, 10 to 40Kcal/100ml, 10 to 35Kcal/100ml, 10 to 30Kcal/100ml, 10 to 25Kcal/100ml, 10 to 20Kcal/100ml,

10 to 15Kcal/100ml, 15 to 50Kcal/100ml, 15 to 45Kcal/100ml, 15 to 40Kcal/100ml, 15 to 35Kcal/100ml, 15 to 30Kcal/100ml, 15 to 25Kcal/100ml, 15 to 20Kcal/100ml, 20 to 50Kcal/100ml, 20 to 45Kcal/100ml, 20 to 40Kcal/100ml, 20 to 35Kcal/100ml, 20 to 30Kcal/100ml, 20 to 25Kcal/100ml, 25 to 50Kcal/100ml, 25 to 45Kcal/100ml, 25 to 40Kcal/100ml, 25 to 35Kcal/100ml, 25 to 30Kcal/100ml, 0 to 32Kcal/100ml, 0 to 24Kcal/100ml, 0 to 8Kcal/100ml, 0 to 4Kcal/100ml, 4 to 32Kcal/100ml, 4 to 24Kcal/100ml, 4 to 8Kcal/100ml, 8 to 32Kcal/100ml, 8 to 24Kcal/100ml, or 24 to 32Kcal/100ml.

[0095]

In addition, the forms of the natural sugar (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose) and sodium in fruit juice beverage C of the present invention are defined as described in the section of the fruit juice beverage of the present invention.

[0096]

Further, another aspect of the present invention provides a fruit juice beverage comprising:

(a) a natural sugar in an amount of 0.1 to 0.5 w/v%, 0.1 to 1.0 w/v%, 0.1 to 1.5 w/v%, 0.1 to 2.0 w/v%, 0.1 to 2.5 w/v%, 0.1 to 3.0 w/v%, 0.1 to 3.5 w/v%, 0.1 to 4.0 w/v%, 0.1 to 4.5 w/v%, 0.1 to 5.0 w/v%, 0.1 to 5.5 w/v%, 0.1 to 5.9 w/v%, 0.5 to 1.0 w/v%, 0.5 to 1.5 w/v%, 0.5 to 2.0 w/v%, 0.5 to 2.5 w/v%, 0.5 to 3.0 w/v%, 0.5 to 3.5 w/v%, 0.5 to 4.0 w/v%, 0.5 to 4.5 w/v%, 0.5 to 5.0 w/v%,

0.5 to 5.5 w/v%, 0.5 to 5.9 w/v%, 1.0 to 1.5 w/v%, 1.0 to 2.0 w/v%, 1.0 to 2.5 w/v%, 1.0 to 3.0 w/v%, 1.0 to 3.5 w/v%, 1.0 to 4.0 w/v%, 1.0 to 4.5 w/v%, 1.0 to 5.0 w/v%, 1.0 to 5.5 w/v%, 1.0 to 5.9 w/v%, 1.5 to 2.0 w/v%, 1.5 to 2.5 w/v%, 1.5 to 3.0 w/v%, 1.5 to 3.5 w/v%, 1.5 to 4.0 w/v%, 1.5 to 4.5 w/v%, 1.5 to 5.0 w/v%, 1.5 to 5.5 w/v%, 1.5 to 5.9 w/v%, 2.0 to 2.5 w/v%, 2.0 to 3.0 w/v%, 2.0 to 3.5 w/v%, 2.0 to 4.0 w/v%, 2.0 to 4.5 w/v%, 2.0 to 5.0 w/v%, 2.0 to 5.5 w/v%, or 2.0 to 5.9 w/v%;

(b) at least one high-intensity sweetener selected from the group consisting of rebaudioside M, rebaudioside D, a Luo han guo extract, mogroside V, and a combination thereof in an amount of 20 to 550ppm, 25 to 550 ppm, 30 to 550 ppm, 35 to 550 ppm, 40 to 550 ppm, 45 to 550 ppm, 50 to 550 ppm, 55 to 550 ppm, 20 to 540 ppm, 25 to 540 ppm, 30 to 540 ppm, 35 to 540 ppm, 40 to 540 ppm, 45 to 540 ppm, 50 to 540 ppm, 55 to 540 ppm, 20 to 530 ppm, 25 to 530 ppm, 30 to 530 ppm, 35 to 530 ppm, 40 to 530 ppm, 45 to 530 ppm, 50 to 530 ppm, 55 to 530 ppm, 20 to 520 ppm, 25 to 520 ppm, 30 to 520 ppm, 35 to 520 ppm, 40 to 520 ppm, 45 to 520 ppm, 50 to 520 ppm, 55 to 520 ppm, 20 to 510 ppm, 25 to 510 ppm, 30 to 510 ppm, 35 to 510 ppm, 40 to 510 ppm, 45 to 510 ppm, 50 to 510 ppm, 55 to 510ppm, 20 to 505ppm, 25 to 505ppm, 30 to 505ppm, 35 to 505ppm, 40 to 505ppm, 45 to 505ppm, 50 to 505ppm, 55 to 505ppm, 20 to 500ppm, 25 to 500ppm, 30 to 500ppm, 35 to 500ppm, 40 to 500ppm, 45 to 500ppm, 50 to 500ppm, 55 to 500ppm, 20 to 495ppm, 25 to 495ppm, 30 to

495ppm, 35 to 495ppm, 40 to 495ppm, 45 to 495ppm, 50 to 495ppm, 55 to 495ppm, 20 to 490ppm, 25 to 490ppm, 30 to 490ppm, 35 to 490ppm, 40 to 490ppm, 45 to 490ppm, 50 to 490ppm, or 55 to 490ppm; and

(c) less than 70mg/100ml, 0.1 to less than 70mg/100ml, 0.1 to 65mg/100ml, 0.1 to 60mg/100ml, 0.1 to 55mg/100ml, 0.1 to 50mg/100ml, 0.1 to 45mg/100ml, 0.1 to 40mg/100ml, 0.1 to 35mg/100ml, 0.1 to 30mg/100ml, 0.1 to 25mg/100ml, 0.1 to 20mg/100ml, 0.1 to 19mg/100ml, 0.1 to 18mg/100ml, 0.1 to 17mg/100ml, 0.1 to 16mg/100ml, 0.1 to 15mg/100ml, 0.1 to 14mg/100ml, 0.1 to 13mg/100ml, 0.1 to 12mg/100ml, 0.1 to 11mg/100ml, 0.1 to 10mg/100ml, 1 to less than 70mg/100ml, 1 to 65mg/100ml, 1 to 60mg/100ml, 1 to 55mg/100ml, 1 to 50mg/100ml, 1 to 45mg/100ml, 1 to 40mg/100ml, 1 to 35mg/100ml, 1 to 30mg/100ml, 1 to 25mg/100ml, 1 to 20mg/100ml, 1 to 19mg/100ml, 1 to 18mg/100ml, 1 to 17mg/100ml, 1 to 16mg/100ml, 1 to 15mg/100ml, 1 to 14mg/100ml, 1 to 13mg/100ml, 1 to 12mg/100ml, 1 to 11mg/100ml, 1 to 10mg/100ml, 5 to less than 70mg/100ml, 5 to 65mg/100ml, 5 to 60mg/100ml, 5 to 55mg/100ml, 5 to 50mg/100ml, 5 to 45mg/100ml, 5 to 40mg/100ml, 5 to 35mg/100ml, 5 to 30mg/100ml, 5 to 25mg/100ml, 5 to 20mg/100ml, 5 to 19mg/100ml, 5 to 18mg/100ml, 5 to 17mg/100ml, 5 to 16mg/100ml, 5 to 15mg/100ml, 5 to 14mg/100ml, 5 to 13mg/100ml, 5 to 12mg/100ml, 5 to 11mg/100ml, 5 to 10mg/100ml, 7 to less than 70mg/100ml, 7 to 65mg/100ml, 7 to 60mg/100ml, 7 to

55mg/100ml, 7 to 50mg/100ml, 7 to 45mg/100ml, 7 to 40mg/100ml, 7 to 35mg/100ml, 7 to 30mg/100ml, 7 to 25mg/100ml, 7 to 20mg/100ml, 7 to 19mg/100ml, 7 to 18mg/100ml, 7 to 17mg/100ml, 7 to 16mg/100ml, 7 to 15mg/100ml, 10 to less than 70mg/100ml, 10 to 65mg/100ml, 10 to 60mg/100ml, 10 to 55mg/100ml, 10 to 50mg/100ml, 10 to 45mg/100ml, 10 to 40mg/100ml, 10 to 35mg/100ml, 10 to 30mg/100ml, 10 to 25mg/100ml, 10 to 20mg/100ml, 10 to 19mg/100ml, 10 to 18mg/100ml, 10 to 17mg/100ml, 10 to 16mg/100ml, 10 to 15mg/100ml, 15 to less than 70mg/100ml, 15 to 65mg/100ml, 15 to 60mg/100ml, 15 to 55mg/100ml, 15 to 50mg/100ml, 15 to 45mg/100ml, 15 to 40mg/100ml, 15 to 35mg/100ml, 15 to 30mg/100ml, 15 to 25mg/100ml, 15 to 20mg/100ml, 20 to less than 70mg/100ml, 20 to 65mg/100ml, 20 to 60mg/100ml, 20 to 55mg/100ml, 20 to 50mg/100ml, 20 to 45mg/100ml, 20 to 40mg/100ml, 20 to 35mg/100ml, 20 to 30mg/100ml, 20 to 25mg/100ml, 25 to less than 70mg/100ml, 25 to 65mg/100ml, 25 to 60mg/100ml, 25 to 55mg/100ml, 25 to 50mg/100ml, 25 to 45mg/100ml, 25 to 40mg/100ml, 25 to 35mg/100ml, or 25 to 30mg/100ml of sodium.

[0097]

2. Method for producing a fruit juice beverage

Even for a fruit juice beverage having an energy (Kcal/100 ml) controlled to be low by setting the amounts of the natural sugar and the high-intensity sweetener blended each to a low level, the sweetness based on the natural sugar and the high-intensity sweetener can be

enhanced by adding sodium in a concentration so low as to not be detectable by the human.

Therefore, the present invention provides, as yet another embodiment, the following method for producing a fruit juice beverage with an enhanced sweetness (hereinafter, referred to as "the method of the present invention").

A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2; and

(ii) adding (c) sodium such that a sodium concentration in the beverage is less than 70 mg/100ml,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, a luohan guo extract, mogroside V, and thaumatin, and

the fruit juice beverage has a sweetness of a sweetness intensity X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0098]

The fruit juice beverage produced by the method of the present invention is the fruit juice beverage of the present invention described in the above section "1. Fruit

juice beverage having a natural sugar and a high-intensity sweetener with an improved taste quality." In addition, the "raw material" in the method of the present invention can be each material needed to produce a fruit juice beverage or a mixture thereof, and can further contain an additional component such as a preservative, a flavor, a carrier, or a milk component. In addition, the "raw material" can be composed of a plurality of materials. However, regardless of the type of the raw material, the fruit juice beverage of the present invention finally produced does not include a substance having a sweetness other than (a) and (b) as a sweetener.

[0099]

In the method of the present invention, any of the following Steps (i) and (ii) can be carried out first.

(i) Adding (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2

(ii) Adding (c) sodium such that a sodium concentration in the beverage is less than 70 mg/100ml

[0100]

In Step (i), (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2 are added to a raw material, but (a) a natural sugar in an amount corresponding to a

sweetness intensity X1 and (b) a high-intensity sweetener in an amount corresponding to sweetness intensity X2 can be added separately.

Further, even when (a) a natural sugar in an amount corresponding to a sweetness intensity X1 is added, the natural sugar in an amount corresponding to a sweetness intensity X1 does not need to be added at once, and the natural sugar can be added in several divided batches. Similarly, even when a high-intensity sweetener in an amount corresponding to a sweetness intensity X2 is added, the high-intensity sweetener in an amount corresponding to a sweetness intensity X2 does not need to be added at once, and the natural sugar can be added in several divided batches.

In addition, as another embodiment, a mixture of the natural sugar and the high-intensity sweetener can also be added in several divided batches to make an adjustment such that the amount of the natural sugar and the amount of the high-intensity sweetener contained in the fruit juice beverage finally produced are the amount corresponding to the sweetness intensity X1 and the amount corresponding to the sweetness intensity X2, respectively.

[0101]

Even when (c) adding sodium such that the sodium concentration in the beverage is less than 70 mg/100ml in Step (ii), sodium does not need to be added at once, and sodium can be added in several divided batches.

The sodium added to the raw material in Step (ii) can be, for example, in the form of at least one selected from the group consisting of sodium chloride, sodium chloride, sodium malate, sodium sulfate, sodium citrate, sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptonate, sodium gluconate, sodium glutamate, sodium tartrate, sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and mixtures thereof.

In addition, when sodium is contained in the fruit juice used, the total content of sodium derived from the fruit juice and additional sodium is adjusted to less than 70 mg/100ml.

[0102]

The "addition" herein means not only the actual operation of adding any of the components (a), (b), and (c) to the raw material, but also the operation of adjusting the amounts of the components (a), (b), and (c) in the fruit juice beverage finally produced through the production process of the fruit juice beverage of the present invention to the amount corresponding to the sweetness intensity X1, the amount corresponding to the sweetness intensity X2, and less than 70 mg/100ml of sodium, respectively.

For example, if a first raw material contains milk, a grain, a bean, or an extract thereof, and accordingly the raw material contains any one or more of the components

(a), (b), and (c) in advance, a second raw material for being mixed with the first raw material also contains the components (a), (b), and (c), and the fruit juice beverage of the present invention can be produced by mixing the first and second raw materials, then there is no operation of individually adding the components (a), (b), and (c) to the raw materials, but in the method of the present invention, Steps (i) and (ii) are considered to have been carried out as long as the fruit juice beverage of the present invention finally produced contains (a) a natural sugar in an amount corresponding to a sweetness intensity X1, (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2, and (c) less than 70 mg/100ml of sodium.

[0103]

When the fruit juice beverage of the present invention is made into a packed-in-container beverage, the method for producing a beverage according to the present invention includes filling the container with the fruit juice beverage. In addition, when the fruit juice beverage of the present invention is made into a packed-in-container beverage, sterilization of the fruit juice beverage before or after filling the container with the fruit juice beverage is preferable because the sterilization allows long-term storage of the packed-in-container beverage. For example, when the fruit juice beverage of the present invention is made into a canned fruit juice beverage, a

can can be filled with a predetermined amount of the fruit juice beverage and subjected to retort sterilization, for example, at 120 to 125°C for about 5 to 20 minutes to perform heat sterilization. In addition, when the fruit juice beverage of the present invention is made into a PET bottled, paper packed, or bottled beverage, the fruit juice beverage can be subjected to UHT sterilization involving holding, for example, at 130 to 145°C for about 2 to 120 seconds or the like and a predetermined amount thereof can be subjected to hot packing or aseptic packaging at a low temperature to obtain a packed-in-container beverage.

[0104]

In the method of the present invention, the "fruit juice beverage", "natural sugar", "sweetness intensity X1", "high-intensity sweetener", "sweetness intensity X2", sodium content, sodium form in the fruit juice beverage, "sweetness intensity X3", and energy are defined as described in the above section for the fruit juice beverage, and the numerical values described in the above section for the fruit juice beverage are applicable to the numerical values thereof as they are. In addition, examples of the "combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose" and the "combination of high-intensity sweeteners" are also the same as the examples described in the above section for the fruit juice beverage.

[0105]

In a certain embodiment, the fruit juice beverage produced by the method of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 of 0.1 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 of 0.1 to 5; and

(c) less than 70 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X_3 of 5.5 to 12.5 exhibited by the components (a) to (c), and $0.2 < (X_1 + X_2) \leq 12.5$ is satisfied.

[0106]

In another embodiment, the fruit juice beverage produced by the method of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 of 3 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 of 1 to 5; and

(c) 5 to 60 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X_3 of 5.5 to 12.5 exhibited by the components (a) to (c), and $0.1 < (X_1 + X_2) \leq 12.5$ is satisfied.

[0107]

In yet another embodiment, the fruit juice beverage produced by the method of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 of 0.1 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 of 0.1 to 5; and

(c) 5 to 50 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X_3 of 2.0 to 12.0 exhibited by the components (a) to (c), and $0.1 < (X_1 + X_2) \leq 12.5$ is satisfied.

[0108]

In yet another embodiment, the fruit juice beverage produced by the method of the present invention is a fruit juice beverage comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 of 3 to 5;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 of 1 to 5; and

(c) 5 to 50 mg/100ml of sodium,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X_3 of 2.0 to 12.0 exhibited by the components (a) to (c), and $0.1 < (X_1 + X_2) \leq 12.5$ is satisfied.

[0109]

In a specific embodiment, the present invention provides the following method (hereinafter, referred to as "method A of the present invention").

A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) a high-intensity sweetener selected from the group consisting of rebaudioside M, rebaudioside D, and a combination thereof in an amount corresponding to a sweetness intensity X2; and

(ii) adding (c) sodium such that a sodium content in the beverage is 5 to 50 mg/100ml,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0110]

In method A of the present invention, the "fruit juice beverage", "natural sugar" (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose), "sweetness intensity X1", "high-intensity sweetener", "sweetness intensity X2", sodium content, sodium form in the fruit juice beverage, "sweetness intensity X3", and energy are defined as described in the above section for fruit juice beverage A, and the numerical values and the like described in the above section for fruit juice

beverage A are applicable to the numerical values and the like thereof as they are. In addition, the fruit juice beverage in method A of the present invention corresponds to fruit juice beverage A of the present invention, and the items described above for fruit juice beverage A of the present invention are applicable to the fruit juice beverage in method A of the present invention as they are. Further, the "raw material", "addition", the order of Steps (i) and (ii), and the mode of addition of each component in method A of the present invention are defined as described above for the method of the present invention.

[0111]

In another specific embodiment, the present invention provides the following method (hereinafter, referred to as "method B of the present invention").

A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) a luohan guo extract in an amount corresponding to a sweetness intensity X2; and

(ii) adding (c) sodium such that a sodium content in the beverage is 5 to 50 mg/100ml,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0112]

In method B of the present invention, the "fruit juice beverage", "natural sugar" (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose), "sweetness intensity X1", "luo han guo extract", "sweetness intensity X2", sodium content, sodium form in the fruit juice beverage, "sweetness intensity X3", and energy are defined as described above in the above section for fruit juice beverage B, and the numerical values and the like described in the above section for fruit juice beverage B are applicable to the numerical values and the like thereof as they are. In addition, the fruit juice beverage in method B of the present invention corresponds to fruit juice beverage B of the present invention, and the items described above for fruit juice beverage B of the present invention are applicable to the fruit juice beverage in method B of the present invention as they are. Further, the "raw material", "addition", the order of Steps (i) and (ii), and the mode of addition of each component in method B of the present invention are defined as described above for the method of the present invention.

[0113]

In another specific embodiment, the present invention provides the following method (hereinafter, referred to as "method C of the present invention").

A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity X1 and (b) mogroside V in an amount corresponding to a sweetness intensity X2; and

(ii) adding (c) sodium such that a sodium content in the beverage is 5 to 50 mg/100ml,

wherein the fruit juice beverage has a sweetness of a sweetness intensity X3 exhibited by the components (a) to (c), and $0.1 < (X1 + X2) \leq 20$ is satisfied.

[0114]

In method C of the present invention, the "fruit juice beverage", "natural sugar" (including an example of a combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose), "sweetness intensity X1", "mogroside V", "sweetness intensity X2", sodium content, sodium form in the fruit juice beverage, "sweetness intensity X3", and energy are defined as described in the above section for fruit juice beverage C, and the numerical values and the like described in the above section for fruit juice beverage C are applicable to the numerical values and the like thereof as they are. In addition, the fruit juice beverage in method C of the present invention corresponds to fruit juice beverage C of the present invention, and the items described above for fruit juice beverage C of the present invention are applicable to the fruit juice beverage in method C of the present invention as they are. Further,

the "raw material", "addition", the order of Steps (i) and (ii), and the mode of addition of each component in method C of the present invention are defined as described above for the method of the present invention.

[0115]

In the fruit juice beverage produced by the method of the present invention,

the sweetness intensity X1 of the natural sugar, the sweetness intensity X2 of the high-intensity sweetener, the sodium content, and the sweetness intensity X3 exhibited by the fruit juice beverage can each be any value as long as $0.1 < (X1 + X2) \leq 20$ is satisfied, and for example, any of the numerical values of the sweetness intensity X1, the sweetness intensity X2, the sodium content, the sweetness intensity X3, and energy shown in the section of the fruit juice beverage of the present invention can be combined.

[0116]

3. Concentrate for providing the fruit juice beverage

The present invention provides, as another embodiment, a concentrate for providing the fruit juice beverage of the present invention (hereinafter, referred to as "the concentrate of the present invention"). According to an embodiment of the present invention, the concentrate of the present invention comprises:

(a) a natural sugar in an amount corresponding to a sweetness intensity X4;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X5; and

(c) less than 700 mg/100ml of sodium,

wherein $(X4 + X5) \leq 200$, preferably $1 < (X4 + X5) \leq 200$, and more preferably $50 < (X4 + X5) \leq 200$ is satisfied.
[0117]

The concentrate of the present invention is used to provide a fruit juice beverage by diluting the concentrate in any ratio. The "fruit juice beverage" is the same as described in "1. Fruit juice beverage having a natural sugar and a high-intensity sweetener with an improved taste quality." For example, the concentrate of the present invention can be used as a syrup or an undiluted solution in a beverage. In this instance, the concentrate can be diluted 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold and used. In addition, the concentrate of the present invention is preferable in the aspects of preservability and transportability because it is concentrated. The concentrate of the present invention can be solid or liquid.

[0118]

The concentrate of the present invention is 2 to 10-fold concentrate, preferably 3 to 9-fold concentrate, more preferably 4 to 8-fold concentrate, and further preferably 5 to 7-fold concentrate, of the fruit juice beverage of the present invention.

[0119]

The concentrate according to an embodiment of the present invention is 5-fold concentrate of the fruit juice beverage of the present invention and comprises:

(a) a natural sugar in an amount corresponding to a sweetness intensity X6;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X7; and

(c) less than 350 mg/100ml of sodium,

wherein $(X6 + X7) \leq 100$, preferably $0.5 < (X6 + X7) \leq 100$, and more preferably $20 < (X6 + X7) \leq 100$ is satisfied.
[0120]

The concentrate according to an embodiment of the present invention is 6-fold concentrate of the fruit juice beverage of the present invention and comprises:

(a) a natural sugar in an amount corresponding to a sweetness intensity X8;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X9; and

(c) less than 420 mg/100ml of sodium,

wherein $(X8 + X9) \leq 120$, preferably $0.6 < (X8 + X9) \leq 120$, and more preferably $30 < (X8 + X9) \leq 120$ is satisfied.
[0121]

The concentrate according to another embodiment of the present invention is 8-fold concentrate of the fruit juice beverage of the present invention and comprises:

(a) a natural sugar in an amount corresponding to a sweetness intensity X10;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_{11} ; and

(c) less than 560 mg/100ml of sodium,

wherein $(X_{10} + X_{11}) \leq 160$, preferably $0.8 < (X_{10} + X_{11}) \leq 160$, and more preferably $40 < (X_{10} + X_{11}) \leq 160$ is satisfied.

[0122]

4. Method for enhancing the sweetness of the fruit juice beverage

The present invention provides, as another embodiment, a method for enhancing the sweetness of the fruit juice beverage (hereinafter, referred to as "the sweetness enhancement method of the present invention"). According to an embodiment of the present invention, the sweetness enhancement method of the present invention is characterized by containing:

(a) a natural sugar in an amount corresponding to a sweetness intensity X_1 ;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 ; and

(c) less than 70 mg/100ml of sodium
in a fruit juice beverage.

[0123]

According to the sweetness enhancement method of the present invention, the sweetness of the fruit juice beverage is enhanced and a fruit juice beverage having a sweetness more than the sweetness intensity obtained when the component (a) is added to the fruit juice beverage and

the sweetness intensity obtained when the component (b) is added to the fruit juice beverage can be provided. In the sweetness enhancement method of the present invention, the "fruit juice beverage", "natural sugar", "sweetness intensity X1", "high-intensity sweetener", "sweetness intensity X2", sodium content, sodium form in the fruit juice beverage, "sweetness intensity X3", and energy are defined as described in the above section for the fruit juice beverage, and the numerical values described in the above section for the fruit juice beverage are applicable to the numerical values thereof as they are. In addition, examples of the "combination of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, and lactose" and the "combination of high-intensity sweeteners" are also the same as the examples described in the above section for the fruit juice beverage.

[0124]

As used herein, the term "at least" means that the number of a specific item can be greater than or equal to the number expressed. In addition, in the present application, the term "about" means that a subject matter is in ranges of $\pm 25\%$, $\pm 10\%$, $\pm 5\%$, $\pm 3\%$, $\pm 2\%$, or $\pm 1\%$, of the numerical value following the "about." For example, "about 10" means a range from 7.5 to 12.5.

Examples

[0125]

Hereinafter, the present invention will be specifically described with reference to Examples but the present invention is not limited to the following Examples.

In the Examples shown below, the sodium content in the base beverage before the addition of various additives was measured by atomic absorption spectroscopy using SpectrAA240FS (manufactured by Agilent Technologies, Inc.). In addition, the sodium content of a beverage sample after blending various additives was obtained by adding the sodium content in the various additives calculated from the amounts blended to the sodium content in the base beverage measured by the above method.

[0126]

[Example 1] Evaluation of the improving effect of sodium on taste qualities

Experimental method

Natural sugars {sucrose (manufactured by Pacific Sugar Mfg. Co., Ltd.) and glucose (manufactured by Showa Sangyo Co., Ltd.)}, rebaudioside D (RebD) (purity 95% or more), and sodium gluconate were dissolved in an orange diluted fruit juice (fruit juice percentage: 10%, sodium content: 0 (mg/100ml), energy: 4 (kcal/100 ml), and Brix = 1.0) in the ratios shown in Table 2 below to prepare beverage samples. In addition, a solution to which sodium gluconate was not added was used as sample 1.

Brix in Table 2 was calculated from the concentrations of the natural sugars and the Brix derived from the orange

diluted fruit juice, the sodium content in the beverage was calculated from the sodium gluconate concentration, and the energy (kcal/100 ml) was calculated with the energy of RebD and the sodium-derived components being 0 (kcal/100 ml) .

Table 2

Content	Sample 1	1-1	1-2	1-3	1-4
Sucrose (w/v%)	1.0	1.0	1.0	1.0	1.0
Glucose (w/v%)	3.5	3.5	3.5	3.5	3.5
Brix (derived from sucrose + glucose + derived from orange diluted fruit juice)	5.5	5.5	5.5	5.5	5.5
RebD (ppm)	208	208	208	208	208
Sodium gluconate (g/L)	0	0.95	1.90	2.85	3.80
Sodium content in beverage sample (mg/100ml)	0	10	20	30	40
Energy (Kcal/100ml)	22	22	22	22	22

[0127]

The taste quality improvement effect by sodium was verified by sensory comparison of taste qualities of these beverage samples. The taste qualities as sensory evaluation items were "intensity of sweetness", "intensity of flavor (aroma)", "intensity of odd taste (such as bitter taste and astringent taste)", and "intensity of saltiness", and the verification was carried out by those (four persons) who received sensory training as panelists.

First, each panelist calculated the "sensory evaluation score" for each taste quality of each beverage sample based on the common taste quality evaluation criteria possessed by the panelists through daily training.

The "sensory evaluation score" a difference in the taste quality from sample 1 quantified in a range of -3.0 to +3.0 based on the degree of taste quality of sample 1 defined as "0" (reference). The criteria of the "sensory evaluation score" of each item are as follows.

(Sensory evaluation score)

" +3.0": it is felt that the taste quality as an object is very strong compared with sample 1;

" +2.0": it is felt that the taste quality as an object is strong compared with sample 1;

" +1.0": it is felt that the taste quality as an object is slightly strong compared with sample 1;

"0": it is felt that the taste quality as an object is equal to sample 1;

" -1.0": it is felt that the taste quality as an object is slightly weak compared with sample 1;

" -2.0": it is felt that the taste quality as an object is weak compared with sample 1; and

" -3.0": it is felt that the taste quality as an object is very weak compared with sample 1.

Incidentally, in light of the above criteria, for example, when it was judged that the taste quality was between "+1.0" and "+2.0, it was scored as "+1.5" in 0.5 increments.

[0125]

The "sensory evaluation score" calculated by each panelist was converted to a "converted score" based on the

following criteria, and the total value of the converted scores of four panelists was calculated for each taste quality.

(Converted score)

Converted score "3": a sensory evaluation score of +1.5 or more;

Converted score "2": a sensory evaluation score of +1.0 or more and less than +1.5;

Converted score "1": a sensory evaluation score of +0.5 or more and less than +1.0;

Converted score "0": a sensory evaluation score of higher than -0.5 and less than +0.5;

Converted score "-1": a sensory evaluation score of higher than -1.0 and -0.5 or less;

Converted score "-2": a sensory evaluation score of higher than -1.5 and -1.0 or less; and

Converted score "-3": a sensory evaluation score of -1.5 or less.

Incidentally, regarding the "intensity of sweetness" and the "intensity of flavor", a larger total value of converted scores is preferable, and regarding the "intensity of odd taste" and the "intensity of saltiness", a smaller total value of converted scores is preferable.

[0129]

Results

Table 3 shows the calculated results of the total conversion scores of each taste quality.

Table 3

Taste quality	Sample 1	1-1	1-2	1-3	1-4
Intensity of sweetness	0	3	5	7	7
Intensity of flavor	0	1	3	5	5
Intensity of odd taste	0	0	0	0	0
Intensity of saltiness	0	0	0	0	2

[0130]

[Example 2] Evaluation of the improving effect of various high-intensity sweeteners and sodium on taste qualities

[Example 2-1] Evaluation of the improving effect of mogroside V and sodium on taste qualities

Experimental method

In the same way as in Example 1, natural sugars {sucrose (manufactured by Pacific Sugar Mfg. Co., Ltd.) and glucose (manufactured by Showa Sangyo Co., Ltd.)}, a high-intensity sweetener {mogroside V (MogV) (purity 95% or more)}, and sodium gluconate were dissolved in an orange diluted fruit juice (fruit juice percentage: 10%, sodium content: 0 (mg/100ml), energy: 4 (kcal/100 ml), and Brix = 1.0) in the ratios shown in Table 4-1 below to prepare beverage samples. In addition, a solution to which sodium gluconate was not added was used as sample 2 (containing MogV). Brix in Table 4-1 was calculated from the concentrations of the natural sugars and the Brix derived from the orange diluted fruit juice, the sodium content in the beverage was calculated from the sodium gluconate concentration, and the energy (kcal/100 ml) was calculated with the energy of MogV and the sodium-derived components being 0 (kcal/100 ml).

Table 4-1

Content	Sample 2	2-1
Sucrose (w/v%)	1.0	1.0
Glucose (w/v%)	3.5	3.5
Brix (derived from sucrose + glucose + derived from orange diluted fruit juice)	5.5	5.5
MogV(ppm)	208	208
Sodium gluconate (g/L)	0	1.90
Sodium content in beverage sample (mg/100ml)	0	20
Energy (Kcal/100ml)	22	22

[0131]

The improving effect of sodium on taste qualities was verified by the sensory comparison of the taste qualities of these beverage samples. The taste qualities that were the sensory evaluation items were "intensity of sweetness", "intensity of flavor (aroma)", "intensity of odd taste (bitterness, astringency, or the like)", and "intensity of salty taste", and those who received sensory trainings (4 persons) carried out verification as panelists. Specifically, the same evaluation as in Example 1 was carried out, and the "sensory evaluation score" calculated by each panelist was converted to the "conversion score", and the total conversion score of the four panelists for each taste quality was calculated. For samples 2 and 2-1, the score of sample 2 was used as a reference (0 points).

[0132]

Results

Table 4-2 shows the calculated results of the total conversion scores of each taste quality.

Table 4-2

Taste quality	Sample 2	2-1
Intensity of sweetness	0	6
Intensity of flavor	0	6
Intensity of odd taste	0	-3
Intensity of saltiness	0	0

[0133]

[Example 2-2] Evaluation of the improving effect of rebaudioside M and sodium on taste qualities

Experimental method

In the same way as in Example 2-1, natural sugars {sucrose (manufactured by Pacific Sugar Mfg. Co., Ltd.) and glucose (manufactured by Showa Sangyo Co., Ltd.)}, a high-intensity sweetener {rebaudioside M (RebM) (purity 99% or more)}, and sodium gluconate were dissolved in an orange diluted fruit juice (fruit juice percentage: 10%, sodium content: 0 (mg/100ml), energy: 4 (kcal/100 ml), and Brix = 1.0) in the ratios shown in Table 5-1 below to prepare beverage samples. In addition, a solution to which sodium gluconate was not added was used as sample 2A (containing RebM). Brix in Table 5-1 was calculated from the concentrations of the natural sugars and the Brix derived from the orange diluted fruit juice, the sodium content in the beverage was calculated from the sodium gluconate concentration, and the energy (kcal/100 ml) was calculated with the energy of RebM and the sodium-derived components being 0 (kcal/100 ml).

Table 5-1

Content	Sample 2A	2-2
Sucrose (w/v%)	1.0	1.0
Glucose (w/v%)	3.5	3.5
Brix (derived from sucrose + glucose + derived from orange diluted fruit juice)	5.5	5.5
RebM (ppm)	208	208
Sodium gluconate (g/L)	0	1.90
Sodium content in beverage sample (mg/100ml)	0	20
Energy (Kcal/100ml)	22	22

[0134]

The improving effect of sodium on taste qualities was verified by the sensory comparison of the taste qualities of these beverage samples. The taste qualities that were the sensory evaluation items were "intensity of sweetness", "intensity of flavor (aroma)", "intensity of odd taste (bitterness, astringency, or the like)", and "intensity of salty taste", and those who received sensory trainings (4 persons) carried out verification as panelists. Specifically, the same evaluation as in Example 1 was carried out, and the "sensory evaluation score" calculated by each panelist was converted to the "conversion score", and the total conversion score of the four panelists for each taste quality was calculated. For samples 2A and 2-2, the score of sample 2A was used as a reference (0 points).

[0135]

Results

Table 5-2 shows the calculated results of the total conversion scores of each taste quality.

Table 5-2

Taste quality	Sample 2A	2-2
Intensity of sweetness	0	5
Intensity of flavor	0	3
Intensity of odd taste	0	-4
Intensity of saltiness	0	0

[0136]

[Example 3] Various high-intensity sweeteners and evaluation of the improving effect of sodium on taste qualities

Experimental method

In the same way as in Example 1, natural sugars {sucrose (manufactured by Pacific Sugar Mfg. Co., Ltd.) and glucose (manufactured by Showa Sangyo Co., Ltd.)}, rebaudioside D (RebD) (purity 95% or more), and sodium gluconate were dissolved in a black currant diluted fruit juice (fruit juice percentage: 10%, sodium content: 0 (mg/100ml), energy: 3 (kcal/100 ml), and Brix = 0.8) in the ratios shown in Table 6 below to prepare beverage samples. In addition, a solution to which sodium gluconate was not added was used as sample 3. Brix in Table 6 was calculated from the concentrations of the natural sugars and the Brix derived from the black currant diluted fruit juice, the sodium content in the beverage was calculated from the sodium gluconate concentration, and the energy (kcal/100 ml) was calculated with the energy of RebD and the sodium-derived components being 0 (kcal/100 ml).

Table 6

Content	Sample 3	3-1
Sucrose (w/v%)	1.0	1.0
Glucose (w/v%)	3.5	3.5
Brix (derived from sucrose + glucose + derived from black currant diluted fruit juice)	5.3	5.3
RebD (ppm)	208	208
Sodium gluconate (g/L)	0	1.90
Sodium content in beverage Sample (mg/100ml)	0	20
Energy (Kcal/100ml)	21	21

[0137]

The improving effect of sodium on taste qualities was verified by the sensory comparison of the taste qualities of these beverage samples. The taste qualities that were the sensory evaluation items were "intensity of sweetness", "intensity of flavor (aroma)", "intensity of odd taste (bitterness, astringency, or the like)", and "intensity of salty taste", and those who received sensory trainings (4 persons) carried out verification as panelists. Specifically, the same evaluation as in Example 1 was carried out, and the "sensory evaluation score" calculated by each panelist was converted to the "conversion score", and the total conversion score of the four panelists for each taste quality was calculated.

[0138]

Results

Table 7 shows the calculated results of the total conversion scores of each taste quality.

Table 7

Taste quality	Sample 3	3-1
Intensity of sweetness	0	8
Intensity of flavor	0	5
Intensity of odd taste	0	-1
Intensity of saltiness	0	0

[0139]

[Example 4] Aroma component analysis using gas chromatography

Experimental method

In the same way as in Example 1, natural sugars {sucrose (manufactured by Pacific Sugar Mfg. Co., Ltd.) and glucose (manufactured by Showa Sangyo Co., Ltd.)}, rebaudioside D (RebD) (purity 95% or more), and sodium gluconate were dissolved in an orange diluted fruit juice (fruit juice percentage: 10%, sodium content: 0 (mg/100ml), energy: 4 (kcal/100 ml), and Brix = 1.0) in the ratios shown in Table 8 below to prepare beverage samples. In addition, a solution to which sodium gluconate was not added was used as sample 4.

Brix in Table 8 was calculated from the concentrations of the natural sugars and the Brix derived from the orange diluted fruit juice, the sodium content in the beverage was calculated from the sodium gluconate concentration, and the energy (kcal/100 ml) was calculated with the energy of RebD and the sodium-derived components being 0 (kcal/100 ml).

Table 8

Content	Sample 4	4-1
---------	----------	-----

Sucrose (w/v%)	1.0	1.0
Glucose (w/v%)	3.5	3.5
Brix (derived from sucrose + glucose + derived from orange diluted fruit juice)	5.5	5.5
RebD(ppm)	208	208
Sodium gluconate (g/L)	0	1.90
Sodium content in beverage sample (mg/100ml)	0	20
Energy (Kcal/100ml)	22	22

[0140]

Method for collecting samples for aroma component analysis

About 50 ml of the beverage sample was placed in an impinger. While causing clean nitrogen to flow from one opening of the impinger at a flow rate of 100 ml/min, the gas flowing into the Tedlar bag attached to the other opening was collected for 10 minutes.

[0141]

Apparatus and analysis conditions

The apparatus and measurement of gas chromatography (GC) analysis were carried out under the following conditions.

- Apparatus: Agilent Technologies 6890N
- Column: DB-1701 (0.25 mm I.D. × 30 m df = 0.25 μm)
- Column temperature: 50°C → 280°C (20°C/min, hold)
- Inlet temperature: 260°C
- Carrier gas: Helium 1.5 mL/min
- Detector: Hydrogen flame ionization detector (FID)

280°C

- Injection method: Split method (1:5)
- Injection volume: 0.15 mL

The peaks obtained by the analysis were compared between sample 4-1 and sample 4, and the area value of the peak having a peak appearance time of 4.2 minutes, which was considered to be the peak of an aroma component, was calculated. Results thereof are shown in Table 9. As a result, it was found that the addition of sodium has the effect of improving the aroma.

Table 9

	Sample 4	4-1
Peak area value of peak appearance time 4.2 min	282.25	365.65

Industrial Applicability

[0142]

The method of the present invention provides a method for enhancing the sweetness of a fruit juice beverage or a sweetness composition to provide a good taste, rather than a simple sweetness obtained by increasing the amounts of a natural sugar and a high-intensity sweetener used.

Claims

1. A fruit juice beverage comprising:
 - (a) a natural sugar in an amount corresponding to a sweetness intensity X_1 ;
 - (b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X_2 ; and
 - (c) less than 70 mg/100ml of sodium,wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, and a luohan guo extract, and
$$0.1 < (X_1 + X_2) \leq 20$$
is satisfied.
2. The fruit juice beverage according to claim 1, wherein a sodium content is 5 to 60 mg/100ml.
3. The fruit juice beverage according to claim 1 or 2, wherein energy is 50 Kcal/100 ml or less.
4. The fruit juice beverage according to any one of claims 1 to 3, wherein X_1 is 0.1 to 5.9.
5. The fruit juice beverage according to any one of claims 1 to 4, wherein a fruit juice percentage of the fruit juice beverage is 1 to 100%.

6. The fruit juice beverage according to any one of claims 1 to 5, wherein a fruit juice contained in the fruit juice beverage is one or more selected from an orange, a mikan, a lemon, a grapefruit, a lime, a pineapple, a strawberry, a blueberry, a black currant, a cranberry, a guava, a banana, an acerola, a papaya, a passion fruit, a mango, an apple, a grape, a peach, an ume, a pear, an apricot, a sumomo, a melon, a kiwi fruit, and a Japanese quince.

7. The fruit juice beverage according to any one of claims 1 to 6, wherein the natural sugar is at least one selected from the group consisting of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, lactose, psicose, allose, tagatose, and a combination thereof.

8. The fruit juice beverage according to any one of claims 1 to 7, wherein the high-intensity sweetener b1 is at least one selected from the group consisting of rebaudioside M, rebaudioside D, a Luo Han Guo extract, and a combination thereof.

9. The fruit juice beverage according to any one of claims 1 to 8, wherein the sodium is at least one selected from the group consisting of sodium chloride, sodium hydroxide, sodium malate, sodium sulfate, sodium citrate,

sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptonate, sodium gluconate, sodium glutamate, sodium tartrate, sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and mixtures thereof.

10. The fruit juice beverage according to any one of claims 1 to 9, wherein the fruit juice beverage comprises 5 to 60 mg/100ml of sodium, energy is 50 Kcal/100 ml or less, and $X1 + X2$ is 6 or more.

11. The fruit juice beverage according to any one of claims 1 to 10, which is packed in a container.

12. The fruit juice beverage according to claim 1, wherein the content of the high-intensity sweetener b1 based on the amount of the high-intensity sweetener (b) is 50 mass% or more.

13. The fruit juice beverage according to claim 1, wherein a sodium content is 10 mg/100ml or more and less than 70 mg/100ml.

14. A method for producing a fruit juice beverage, comprising: to a raw material,

(i) adding (a) a natural sugar in an amount corresponding to a sweetness intensity $X1$ and (b) a high-

intensity sweetener in an amount corresponding to a sweetness intensity X2; and

(ii) adding (c) sodium such that a sodium content in the beverage is less than 70 mg/100ml,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, and a luo han guo extract, and

$0.1 < (X1 + X2) \leq 20$ is satisfied.

15. The method according to claim 14, wherein in the fruit juice beverage, the sodium content is 5 to 60 mg/100ml.

16. The method according to claim 14 or 15, wherein energy of the fruit juice beverage is 50 Kcal/100 ml or less.

17. The method according to any one of claims 14 to 16, wherein X1 is 0.1 to 5.9.

18. The method according to any one of claims 14 to 17, wherein a fruit juice percentage of the fruit juice beverage is 1 to 100%.

19. The method according to any one of claims 14 to 18, wherein a fruit juice contained in the fruit juice beverage is one or more selected from an orange, a mikan, a lemon,

a grapefruit, a lime, a pineapple, a strawberry, a blueberry, a black currant, a cranberry, a guava, a banana, an acerola, a papaya, a passion fruit, a mango, an apple, a grape, a peach, an ume, a pear, an apricot, a sumomo, a melon, a kiwi fruit, and a Japanese quince.

20. The method according to any one of claims 14 to 19, wherein the natural sugar is at least one selected from the group consisting of glucose, sucrose, fructose, maltose, an oligosaccharide, a high-fructose corn syrup, lactose, psicose, allose, tagatose, and a combination thereof.

21. The method according to any one of claims 14 to 20, wherein the high-intensity sweetener b1 is at least one selected from the group consisting of rebaudioside M, rebaudioside D, a Luo Han Guo extract, and a combination thereof.

22. The method according to any one of claims 14 to 21, wherein the sodium is at least one selected from the group consisting of sodium chloride, sodium hydroxide, sodium malate, sodium sulfate, sodium citrate, sodium phosphate, sodium carbonate, sodium disulfide, sodium bicarbonate, sodium alginate, sodium arginate, sodium glucoheptonate, sodium gluconate, sodium glutamate, sodium tartrate,

sodium aspartate, sodium lactate, sodium caseinate, sodium ascorbate, and mixtures thereof.

23. The method according to any one of claims 14 to 22, wherein in the fruit juice beverage, the sodium content is 15 to 50 mg/100ml, energy is 50 Kcal/100 ml or less, and $X1 + X2$ is 6 or more.

24. A concentrate for providing the fruit juice beverage according to claim 1, comprising:

(a) a natural sugar in an amount corresponding to a sweetness intensity $X4$;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity $X5$; and

(c) less than 700 mg/100ml of sodium

wherein $(X4 + X5) \leq 200$ is satisfied, and

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, and a luo han guo extract.

25. A method for enhancing a sweetness of a fruit juice beverage, comprising containing:

(a) a natural sugar in an amount corresponding to a sweetness intensity $X1$;

(b) a high-intensity sweetener in an amount corresponding to a sweetness intensity X2; and

(c) less than 70 mg/100ml of sodium in the fruit juice beverage,

wherein the high-intensity sweetener comprises at least one high-intensity sweetener b1 selected from the group consisting of rebaudioside M, rebaudioside D, rebaudioside N, rebaudioside O, rebaudioside E, and a luohan guo extract.