

Claims:

1. A compound selected from 2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one and pharmaceutically acceptable salts thereof.

2. A compound selected from (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one and pharmaceutically acceptable salts thereof.

3. The compound of claim 2, wherein the compound is substantially free of (R)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one and pharmaceutical acceptable salts thereof.

4. The compound of claim 2, wherein the compound is (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one 4-methylbenzenesulfonate.

5. The compound of claim 2, wherein the compound is selected from

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one sulphate;

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one hydrochloride;

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one benzenesulfonate;

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one maleate; and

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one camphor sulfonate.

6. A compound selected from

(S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-3-(3-fluorophenyl)-4H-chromen-4-one,

(R)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-3-(3-fluorophenyl)-4H-chromen-4-one,
and pharmaceutically acceptable salts thereof.

7. The compound of claim 6, wherein the compound is (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-3-(3-fluorophenyl)-4H-chromen-4-one or a pharmaceutically acceptable salt thereof.

8. The compound of claim 7, wherein the compound is substantially free of (R)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-3-(3-fluorophenyl)-4H-chromen-4-one and pharmaceutically acceptable salts thereof.

9. A pharmaceutical composition comprising a compound of any one of claims 1-8 and at least one pharmaceutically acceptable carrier.

10. A method of inhibiting a catalytic activity of a PI3 δ kinase present in a cell, comprising contacting the cell with an effective amount of a compound of any one of claims 1-8.

11. The method of claim 10, wherein the inhibition takes place in a subject suffering from a disease or disorder which is cancer, bone disorder, inflammatory disease, immune disease, nervous system disease, metabolic disease, respiratory disease, thrombosis, or cardiac disease.

12. Use of a compound of any one of claims 1-8, in the manufacture of a medicament for the treatment of a disease, disorder, or condition that would benefit from inhibiting catalytic activity of a PI3 δ kinase.

13. A method for the treatment of a PI3K associated disease or disorder, comprising the step of administering to a subject in need thereof an effective amount of the compound of any one of claims 1-8.

14. The method of claim 13, further comprising the step of administering simultaneously or sequentially to a subject in need thereof at least one other anti-cancer agent, anti-inflammatory agent, immunosuppressive agent, steroid, non-steroidal anti-inflammatory agent, antihistamine, analgesic, or a mixture thereof.

15. The method of claim 13, wherein the PI3K associated disease, disorder or condition is an immune system-related disease, a disease or disorder involving inflammation, cancer or other proliferative disease, a hepatic disease or disorder, or a renal disease or disorder.

16. The method of claim 13, wherein the PI3K associated disease, disorder or condition is selected from inflammation, glomerulonephritis, uveitis, hepatic diseases or disorders, renal diseases or disorders, chronic obstructive pulmonary disease, rheumatoid arthritis, inflammatory bowel disease, vasculitis, dermatitis, osteoarthritis, inflammatory muscle disease, allergic rhinitis, vaginitis, interstitial cystitis, scleroderma, osteoporosis, eczema, allogeneic or xenogeneic transplantation, graft rejection, graft-versus-host disease, lupus erythematosus, pulmonary fibrosis, dermatomyositis, thyroiditis, myasthenia gravis, autoimmune hemolytic anemia, cystic fibrosis, chronic relapsing hepatitis, primary biliary cirrhosis, allergic conjunctivitis, hepatitis, atopic dermatitis, asthma, Sjogren's syndrome, organ transplant rejection, multiple sclerosis, Guillain-Barre, autoimmune uveitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's granulomatosis, Behcet's disease, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, vitiligo, Crohn's disease, colitis, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, Type 1 or immune-mediated diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, autoimmune disorder of the adrenal gland, systemic lupus erythematosus, polymyositis, dermatomyositis, ankylosing spondylitis, transplant rejection, skin graft rejection, arthritis, bone diseases associated with increased bone resorption, ileitis, Barrett's syndrome, adult respiratory distress syndrome, chronic obstructive airway disease, corneal dystrophy, trachoma, onchocerciasis, sympathetic ophthalmitis, endophthalmitis, gingivitis, periodontitis, tuberculosis, leprosy, uremic complications, nephrosis, sclerodermitis, psoriasis, chronic demyelinating diseases of the nervous system, AIDS-related neurodegeneration, Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematoses; systemic lupus erythematosus (SLE); cardiomyopathy, ischemic heart disease hypercholesterolemia, atherosclerosis, preeclampsia; chronic liver failure, brain and spinal cord trauma, and cancer.


17. The method of claim 13, wherein the PI3K associated disease, disorder or condition is selected from hematopoietic tumors of lymphoid lineage, leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burkett's lymphoma; hematopoietic tumors of myeloid lineage, acute myelogenous leukemias, chronic myelogenous leukemias, myelodysplastic syndrome, promyelocytic leukemia; carcinoma of the bladder, carcinoma of the breast, carcinoma of the colon, carcinoma of the kidney, carcinoma of the liver, carcinoma of the lung, small cell lung cancer, esophageal cancer, gall bladdercancer, ovarian cancer, pancreatic cancer, stomachcancer, cervical cancer, thyroidcancer, prostatecancer, skincancer, squamous cell carcinoma; tumors of mesenchymal origin, fibrosarcoma, rhabdomyosarcoma; tumors of the central and peripheral nervous system, astrocytoma, neuroblastoma, glioma, schwannoma; melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoctanthoma, thyroid follicular cancer and Kaposi's sarcoma.

18. The method of claim 13, wherein the PI3K associated disease, disorder or condition is selected from chronic obstructive pulmonary disease, asthma, rheumatoid arthritis, chronic bronchitis, atopic dermatitis, multiple sclerosis, inflammatory bowel disease, allergic rhinitis, lupus erythematosus and ulcerative colitis.

19. The method of claim 13, wherein the PI3K associated disease, disorder or condition is selected from hematopoietic tumors of lymphoid lineage, leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkin's lymphoma, non-Hodgkins lymphoma, chronic lymphocytic leukemia, hairy cell lymphoma and Burkett's lymphoma, hematopoietic tumors of myeloid lineage, acute myelogenous leukemias, chronic myelogenous leukemias, myelodysplastic syndrome, promyelocytic leukemia or multiple myelomas which includes smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma, and extramedullary plasmacytoma.

20. The method of claim 13, wherein the PI3K associated disease, disorder or condition is selected from Chronic Lymphocytic Leukemia (CLL), Lymphoma Non-Hodgkin (NHL), Acute Myeloid Leukemia (AML), Multiple Myeloma (MM), Small Lymphocytic Lymphoma (SLL), Indolent Non-Hodgkin's Lymphoma (I-NHL), acute lymphocytic leukemia (ALL), mantle cell lymphoma (MCL), follicular lymphoma, Waldestrom's

macroglobulinemia (WM), T-cell lymphoma, B-cell lymphoma, and diffuse large B-cell lymphoma (DLBCL).



The invention claimed is:

1. A compound selected from (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one and pharmaceutically acceptable salts thereof.

2. A pharmaceutical composition comprising (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one or a pharmaceutically acceptable salt thereof, wherein the pharmaceutical composition is substantially free of (R)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one or a pharmaceutical acceptable salt thereof.

3. The compound of claim 1, wherein the compound is (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one 4-methylbenzenesulfonate.

4. The compound of claim 1, wherein the compound is selected from

- (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one sulphate;
- (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one hydrochloride;
- (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one benzenesulfonate;
- (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one maleate; and
- (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one camphor sulfonate.

5. A pharmaceutical composition comprising a compound of claim 3 and at least one pharmaceutically acceptable carrier.
6. A pharmaceutical composition comprising a compound of claim 1 and at least one pharmaceutically acceptable carrier.
7. A pharmaceutical composition comprising a compound of claim 4 and at least one pharmaceutically acceptable carrier.
8. The compound of claim 1, wherein the compound has an enantiomeric excess of at least 98%.
9. The compound of claim 3, wherein the compound has an enantiomeric excess of at least 98%.
10. The composition of claim 2, wherein the (S)-2-(1-(4-amino-3-(3-fluoro-4-isopropoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-6-fluoro-3-(3-fluorophenyl)-4H-chromen-4-one or a pharmaceutically acceptable salt thereof has an enantiomeric excess of at least 98%.