

DESCRIPTION

Title of Invention: CRYSTAL OF AMINO ACID SALT OF 3-HYDROXYISOVALERIC
ACID AND PRODUCTION METHOD THEREOF

5 Technical Field

[0001]

The present invention relates to a crystal of an amino acid salt of a 3-
hydroxyisovaleric acid (β -hydroxy- β -methylbutyrate) (hereinafter, referred to as HMB)
10 which is useful, for example, as a product, a raw material, an intermediate or the like of
health food, medicines, cosmetics, or the like, and a method for producing the crystal.

Background Art

[0002]

15 HMB is useful, for example, as a product, a raw material, an intermediate or the
like of health food, medicines, cosmetics, or the like. HMB is an organic acid obtained
by leucine metabolism in the body and is considered to have effect of anabolizing muscle
or preventing degradation of muscle (Non-Patent Documents 1 and 2).

[0003]

20 From a commercial perspective, HMB is distributed in the market only in the form
of either a free carboxylic acid or a calcium salt. Particularly, in supplement/health food
applications, a calcium salt is used in most cases because the calcium salt is a powder
which is excellent in handling (Non-Patent Document 3). Patent Document 4 discloses
that a crystal of an arginine salt is obtained but does not disclose properties of the crystal
25 obtained.

[0004]

Calcium is an essential mineral playing a role in the bone formation, nerve
activity, muscle movement, and the like. However, it has been recently reported that a
calcium overdose leads to an increased risk of death due to cardiovascular disease or
30 ischemic heart disease (Non-Patent Document 4).

Related Art

Patent Document

[0005]

35 [Patent Document 1] WO 2014/166273

[Patent Document 2] U.S. Patent No. 6,248,922

[Patent Document 3] WO 2013/025775

[Patent Document 4] U.S. Patent Application No. 2004/0176449

Non-Patent Document

[0006]

[Non-Patent Document 1] Journal of Applied Physiology, Vol. 81, p 2095, 1996

[Non-Patent Document 2] Nutrition & Metabolism, Vol. 5, p 1, 2008

5 [Non-Patent Document 3] Journal of the International Society of Sports Nutrition
Vol. 10, p 6, 2013

[Non-Patent Document 4] The BMJ., Vol. 346, p 228, 2013

Disclosure of Invention

10 Problems to Be Solved by the Invention

[0007]

In the pharmaceutical preparation field, there is a problem that calcium derived from a calcium salt binds to another component such as phosphate to form an insoluble salt, and a high-concentration solution cannot be prepared. Patent Documents 1 to 3,
15 Patent Document 1, and Patent Document 4 disclose a method for producing calcium salt of HMB (Patent Document 1 to 3), magnesium salt of HMB (Patent Document 1), and arginine salt of HMB (Patent Document 4), respectively, but a crystal cannot be obtained by any methods. That is, there is no known crystal for any salt forms, and an industrially useful crystal of an HMB salt and a production method thereof are demanded.

20 [0008]

An object of the present invention is to provide a crystal of an amino acid salt of HMB, which is excellent in solubility and easy to handle, and to provide a production method thereof.

25 Means for Solving the Problems

[0009]

The present invention relates to the following (1) to (25)

(1) A crystal of an amino acid salt of a 3-hydroxyisovaleric acid (hereinafter referred to as HMB).

30 (2) The crystal described in (1) above, wherein the amino acid salt of HMB is a basic amino acid salt of HMB.

(3) The crystal described in (2) above, wherein the basic amino acid salt of HMB is arginine salt of HMB.

(4) The crystal described in (2) above, wherein the basic amino acid salt of HMB is lysine
35 salt of HMB.

(5) The crystal described in (2) above, wherein the basic amino acid salt of HMB is ornithine salt of HMB.

- (6) The crystal described in (3) above, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $7.5\pm0.2^\circ$, $14.5\pm0.2^\circ$, $15.1\pm0.2^\circ$, $19.2\pm0.2^\circ$, and $20.2\pm0.2^\circ$.
- 5 (7) The crystal described in (6) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $11.6\pm0.2^\circ$, $12.7\pm0.2^\circ$, $17.9\pm0.2^\circ$, $21.5\pm0.2^\circ$, and $23.3\pm0.2^\circ$.
- (8) The crystal described in (7) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $19.6\pm0.2^\circ$, $21.9\pm0.2^\circ$, $25.2\pm0.2^\circ$, $25.5\pm0.2^\circ$, and $33.6\pm0.2^\circ$.
- 10 (9) The crystal described in (4) above, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $8.5\pm0.2^\circ$, $17.0\pm0.2^\circ$, $18.1\pm0.2^\circ$, $18.5\pm0.2^\circ$, and $19.5\pm0.2^\circ$.
- (10) The crystal described in (9) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $22.2\pm0.2^\circ$, $25.5\pm0.2^\circ$, $25.8\pm0.2^\circ$, $26.6\pm0.2^\circ$,
15 and $34.4\pm0.2^\circ$.
- (11) The crystal described in (10) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $4.8\pm0.2^\circ$, $20.4\pm0.2^\circ$, $31.0\pm0.2^\circ$, $33.8\pm0.2^\circ$, and $36.5\pm0.2^\circ$.
- (12) The crystal described in (5) above, wherein in powder X-ray diffraction, the crystal
20 has peaks at diffraction angles (2θ) of $5.1\pm0.2^\circ$, $14.0\pm0.2^\circ$, $15.3\pm0.2^\circ$, $20.4\pm0.2^\circ$, and $21.9\pm0.2^\circ$.
- (13) The crystal described in (12) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $16.4\pm0.2^\circ$, $16.8\pm0.2^\circ$, $19.4\pm0.2^\circ$, $21.4\pm0.2^\circ$, and $25.5\pm0.2^\circ$.
- 25 (14) The crystal described in (13) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $10.9\pm0.2^\circ$.
- (15) A method for producing a crystal of an amino acid salt of HMB, the method comprising a step of dissolving an amorphous amino acid salt of HMB in a solvent containing alcohol, a step of precipitating a crystal of an amino acid salt of HMB by
30 stirring or allowing the solvent to left stand, and a step of collecting a crystal of an amino acid salt of HMB from the solvent.
- (16) A method for producing a crystal of an amino acid salt of HMB, the method comprising a step of concentrating an aqueous HMB solution containing an amino acid-containing compound and having a pH of 2.5 to 11.0 to precipitate a crystal of an amino
35 acid salt of HMB, and a step of collecting a crystal of an amino acid salt of HMB from the aqueous solution.

(17) The production method described in (16) above, wherein the step of precipitating a crystal of an amino acid salt of HMB includes a step of adding or adding dropwise at least one solvent selected from the group consisting of alcohol, nitrile, and ketone.

(18) The production method described in (15) or (17) above, wherein the alcohol is at least one alcohol selected from the group consisting of C1 to C6 alcohols.

(19) The production method described in (17) or (18) above, wherein the nitrile is acetonitrile.

(20) The production method described in any one of (17) to (19) above, wherein the ketone is at least one ketone selected from the group consisting of acetone, methyl ethyl ketone, methyl isobutyl ketone, and diethyl ketone.

(21) The production method described in any one of (15) to (20) above, wherein the amino acid salt of HMB is a basic amino acid salt of HMB.

(22) The production method described in (21) above, wherein the basic amino acid salt of HMB is arginine salt of HMB, lysine salt of HMB, or ornithine salt of HMB.

(23) The crystal described in (5) above, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $4.9\pm 0.2^\circ$, $5.2\pm 0.2^\circ$, $5.5\pm 0.2^\circ$, $10.9\pm 0.2^\circ$ and $15.5\pm 0.2^\circ$.

(24) The crystal described in (23) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $15.9\pm 0.2^\circ$, $16.4\pm 0.2^\circ$, $17.4\pm 0.2^\circ$, $19.2\pm 0.2^\circ$, and $20.4\pm 0.2^\circ$.

(25) The crystal described in (24) above, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $20.8\pm 0.2^\circ$, $21.3\pm 0.2^\circ$, $21.8\pm 0.2^\circ$, $22.2\pm 0.2^\circ$, and $22.8\pm 0.2^\circ$.

Effects of the Invention

[0010]

According to the present invention, a crystal of an amino acid salt of HMB, which is easy to handle, and a production method thereof are provided. The crystal of an amino acid salt of HMB of the present invention is a salt crystal having superiority such as exhibiting high solubility, not forming an insoluble salt, and not inducing electrolyte abnormality, as compared to a calcium salt of HMB. In addition, the crystal of an amino acid salt of HMB of the present invention has high solubility and the excellent effect of improving flavor as compared to a calcium salt of HMB.

Brief Description of the Drawings

[0011]

[Fig. 1] Fig. 1 illustrates the results of powder X-ray diffraction of a seed crystal of arginine salt nonhydrate of HMB obtained in Example 2.

[Fig. 2] Fig. 2 illustrates the results of powder X-ray diffraction of a crystal of arginine salt nonhydrate of HMB obtained in Example 3.

[Fig. 3] Fig. 3 illustrates the results of infrared spectroscopic (IR) analysis of the crystal of arginine salt nonhydrate of HMB obtained in Example 3.

5 [Fig. 4] Fig. 4 illustrates the results of powder X-ray diffraction of a crystal of lysine salt nonhydrate of HMB obtained in Example 4.

[Fig. 5] Fig. 5 illustrates the results of infrared spectroscopic (IR) analysis of the crystal of lysine salt nonhydrate of HMB obtained in Example 4.

10 [Fig. 6] Fig. 6 illustrates the results of powder X-ray diffraction of a crystal of ornithine salt nonhydrate of HMB obtained in Example 6.

[Fig. 7] Fig. 7 illustrates the results of infrared spectroscopic (IR) analysis of the crystal of ornithine salt nonhydrate of HMB obtained in Example 6.

[Fig. 8] Fig. 8 illustrates the results of powder X-ray diffraction of the crystal of ornithine salt nonhydrate of HMB obtained in Example 7.

15

Embodiments for Carrying Out the Invention

[0012]

1. Crystal of Present Invention

20 A crystal of the present invention is a crystal of an amino acid salt of HMB (hereinafter referred to as “the crystal of the present invention” in some cases). Examples of the crystal of an amino acid salt of HMB preferably include a crystal of a basic amino acid salt of HMB, more preferably a crystal of arginine salt of HMB, a crystal of lysine salt of HMB, a crystal of histidine salt of HMB, and a crystal of ornithine salt of HMB, and most preferably a crystal of arginine salt of HMB, a crystal of lysine salt of HMB, and a
25 crystal of ornithine salt of HMB.

[0013]

30 The crystal of the present invention can be confirmed to be a crystal of HMB by the method using HPLC described in analysis examples later. As an amino acid in the crystal of the present invention, any one of an L form and a D form may be used, but the L form is preferable.

[0014]

The crystal of the present invention can be confirmed to be a crystal of an amino acid salt by measuring the amino acid content in the crystal by using HPLC described in the analysis examples later.

35

[0015]

For example, the crystal of the present invention can be confirmed to be a crystal of a mono-arginine salt by the fact that the arginine content in the crystal is generally 59.6 ± 5.0 wt%, preferably 59.6 ± 4.0 wt%, and most preferably 59.6 ± 3.0 wt%.

[0016]

For example, the crystal of the present invention can be confirmed to be a crystal of a monolysine salt by the fact that the lysine content in the crystal is generally 55.3 ± 5.0 wt%, preferably 55.3 ± 4.0 wt%, and most preferably 55.3 ± 3.0 wt%.

5 [0017]

For example, the crystal of the present invention can be confirmed to be a crystal of a mono-ornithine salt by the fact that the ornithine content in the crystal is generally 52.8 ± 5.0 wt%, preferably 52.8 ± 4.0 wt%, and most preferably 52.8 ± 3.0 wt%.

[0018]

10 The crystal of the present invention can be confirmed to be a crystal of a nonhydrate by the fact that the water content measured by using a Karl-Fischer method described in the analysis examples later is generally 2.5 wt% or less, preferably 2.3 wt% or less, and most preferably 2.0 wt% or less.

[0019]

15 The crystal of arginine salt nonhydrate of HMB includes a crystal of arginine salt nonhydrate of HMB of which powder X-ray diffraction pattern using $\text{CuK}\alpha$ as the X-ray source is specified by the values shown in Figs. 1 and 2 and Tables 1 and 2. Here, Fig. 1 and Fig. 2 correspond to the diffraction results of the crystal of arginine salt nonhydrate of HMB of Table 1 and Table 2, respectively.

20 [0020]

The crystal of arginine salt nonhydrate of HMB of which the powder X-ray diffraction pattern is specified by the values shown in Fig. 2 and Table 2 includes a crystal of arginine salt nonhydrate of HMB which shows the infrared absorption spectrum illustrated in Fig. 3 when subjected to the infrared (IR) analysis described in the analysis examples later.

25 [0021]

Specifically, as the crystal of arginine salt nonhydrate of HMB, a crystal of arginine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (i) in the powder X-ray diffraction using $\text{CuK}\alpha$ as the X-ray source is preferable, 30 a crystal of arginine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (ii) as well as the diffraction angles (2θ) described in the following (i) is more preferable, and a crystal of arginine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (iii) as well as the diffraction angles (2θ) described in the following (i) and (ii) is furthermore preferable.

35 (i) $7.5 \pm 0.2^\circ$, preferably $7.5 \pm 0.1^\circ$, $14.5 \pm 0.2^\circ$, preferably $14.5 \pm 0.1^\circ$, $15.1 \pm 0.2^\circ$, preferably $15.1 \pm 0.1^\circ$, $19.2 \pm 0.2^\circ$, preferably $19.2 \pm 0.1^\circ$, and $20.2 \pm 0.2^\circ$, preferably $20.2 \pm 0.1^\circ$.

(ii) $11.6\pm0.2^\circ$, preferably $11.6\pm0.1^\circ$, $12.7\pm0.2^\circ$, preferably $12.7\pm0.1^\circ$, $17.9\pm0.2^\circ$, preferably $17.9\pm0.1^\circ$, $21.5\pm0.2^\circ$, preferably $21.5\pm0.1^\circ$, and $23.3\pm0.2^\circ$, preferably $23.3\pm0.1^\circ$.

(iii) $19.6\pm0.2^\circ$, preferably $19.6\pm0.1^\circ$, $21.9\pm0.2^\circ$, preferably $21.9\pm0.1^\circ$, $25.2\pm0.2^\circ$, preferably $25.2\pm0.1^\circ$, $25.5\pm0.2^\circ$, preferably $25.5\pm0.1^\circ$, and $33.6\pm0.2^\circ$, preferably $33.6\pm0.1^\circ$.

[0022]

The crystal of lysine salt nonhydrate of HMB includes a crystal of lysine salt nonhydrate of HMB of which powder X-ray diffraction pattern using $\text{CuK}\alpha$ as the X-ray source is specified by the values shown in Fig. 4 and Table 4.

[0023]

Examples of the crystal of lysine salt nonhydrate of HMB of which the powder X-ray diffraction pattern is specified by the values shown in Fig. 4 and Table 4 include a crystal of lysine salt nonhydrate of HMB which shows the infrared absorption spectrum illustrated in Fig. 5 when subjected to the infrared spectroscopic (IR) analysis described in the analysis examples later.

[0024]

Specifically, as the crystal of lysine salt nonhydrate of HMB, a crystal of lysine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following

(i) in the powder X-ray diffraction using $\text{CuK}\alpha$ as the X-ray source is preferable, a crystal of lysine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (ii) as well as the diffraction angles (2θ) described in the following (i) is more preferable, and a crystal of lysine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (iii) as well as the diffraction angles (2θ) described in the following (i) and (ii) is furthermore preferable.

(i) $8.5\pm0.2^\circ$, preferably $8.5\pm0.1^\circ$, $17.0\pm0.2^\circ$, preferably $17.0\pm0.1^\circ$, $18.1\pm0.2^\circ$, preferably $18.1\pm0.1^\circ$, $18.5\pm0.2^\circ$, preferably $18.5\pm0.1^\circ$, and $19.5\pm0.2^\circ$, preferably $19.5\pm0.1^\circ$.

(ii) $22.2\pm0.2^\circ$, preferably $22.2\pm0.1^\circ$, $25.5\pm0.2^\circ$, preferably $25.5\pm0.1^\circ$, $25.8\pm0.2^\circ$, preferably $25.8\pm0.1^\circ$, $26.6\pm0.2^\circ$, preferably $26.6\pm0.1^\circ$, and $34.4\pm0.2^\circ$, preferably $34.4\pm0.1^\circ$.

(iii) $4.8\pm0.2^\circ$, preferably $4.8\pm0.1^\circ$, $20.4\pm0.2^\circ$, preferably $20.4\pm0.1^\circ$, $31.0\pm0.2^\circ$, preferably $31.0\pm0.1^\circ$, $33.8\pm0.2^\circ$, preferably $33.8\pm0.1^\circ$, and $36.5\pm0.2^\circ$, preferably $36.5\pm0.1^\circ$.

[0025]

The crystal of ornithine salt nonhydrate of HMB includes a crystal of ornithine salt nonhydrate of HMB of which powder X-ray diffraction pattern using $\text{CuK}\alpha$ as the X-ray source is specified by the values shown in Fig. 6 and Table 6, and a crystal of ornithine salt

nonhydrate of HMB of which the pattern is specified by the values shown in Fig. 8 and Table 9. Here, Fig. 6 and Fig. 8 correspond to the diffraction results of the crystal of ornithine salt nonhydrate of HMB of Table 6 and Table 9, respectively.

[0026]

Examples of the crystal of ornithine salt nonhydrate of HMB of which the powder X-ray diffraction pattern is specified by the values shown in Fig. 6 and Table 6 include a crystal of ornithine salt nonhydrate of HMB which shows the infrared absorption spectrum illustrated in Fig. 7 when subjected to the infrared spectroscopic (IR) analysis described in the analysis examples later.

[0027]

Specifically, as the crystal of ornithine salt nonhydrate of HMB, a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (i) in the powder X-ray diffraction using $\text{CuK}\alpha$ as the X-ray source is preferable, a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (ii) as well as the diffraction angles (2θ) described in the following (i) is more preferable, and a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (iii) as well as the diffraction angles (2θ) described in the following (i) and (ii) is furthermore preferable.

(i) $5.1\pm 0.2^\circ$, preferably $5.1\pm 0.1^\circ$, $14.0\pm 0.2^\circ$, preferably $14.0\pm 0.1^\circ$, $15.3\pm 0.2^\circ$, preferably $15.3\pm 0.1^\circ$, $20.4\pm 0.2^\circ$, preferably $20.4\pm 0.1^\circ$, and $21.9\pm 0.2^\circ$, preferably $21.9\pm 0.1^\circ$.

(ii) $16.4\pm 0.2^\circ$, preferably $16.4\pm 0.1^\circ$, $16.8\pm 0.2^\circ$, preferably $16.8\pm 0.1^\circ$, $19.4\pm 0.2^\circ$, preferably $19.4\pm 0.1^\circ$, $21.4\pm 0.2^\circ$, preferably $21.4\pm 0.1^\circ$, and $25.5\pm 0.2^\circ$, preferably $25.5\pm 0.1^\circ$.

(iii) $0.9\pm 0.2^\circ$, preferably $0.9\pm 0.1^\circ$.

[0028]

Furthermore, as the crystal of ornithine salt nonhydrate of HMB, a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (i) in the powder X-ray diffraction using $\text{CuK}\alpha$ as the X-ray source is preferable, a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (ii) as well as the diffraction angles (2θ) described in the following (i) is more preferable, and a crystal of ornithine salt nonhydrate of HMB having peaks at diffraction angles (2θ) described in the following (iii) as well as the diffraction angles (2θ) described in the following (i) and (ii) is furthermore preferable.

(i) $4.9\pm 0.2^\circ$, preferably $4.9\pm 0.1^\circ$, $5.2\pm 0.2^\circ$, preferably $5.2\pm 0.1^\circ$, $5.5\pm 0.2^\circ$, preferably $5.5\pm 0.1^\circ$, $10.9\pm 0.2^\circ$, preferably $10.9\pm 0.1^\circ$, and $15.5\pm 0.2^\circ$, preferably $15.5\pm 0.1^\circ$.

(ii) $15.9\pm 0.2^\circ$, preferably $15.9\pm 0.1^\circ$, $16.4\pm 0.2^\circ$, preferably $16.4\pm 0.1^\circ$, $17.4\pm 0.2^\circ$, preferably $17.4\pm 0.1^\circ$, $19.2\pm 0.2^\circ$, preferably $19.2\pm 0.1^\circ$, and $20.4\pm 0.2^\circ$, preferably $20.4\pm 0.1^\circ$.

(iii) $20.8 \pm 0.2^\circ$, preferably $20.8 \pm 0.1^\circ$, $21.3 \pm 0.2^\circ$, preferably $21.3 \pm 0.1^\circ$, $21.8 \pm 0.2^\circ$, preferably $21.8 \pm 0.1^\circ$, $22.2 \pm 0.2^\circ$, preferably $22.2 \pm 0.1^\circ$, and $22.8 \pm 0.2^\circ$, preferably $22.8 \pm 0.1^\circ$.

[0029]

2. Method for Producing Crystal of Present Invention

A method for producing the crystal of the present invention is a production method described in the following (1) or (2) (hereinafter will be referred to as “the method for producing the crystal” in some cases).

[0030]

(1) Method 1 for Producing Crystal of Present Invention

Examples of the method for producing the crystal of the present invention include a method for producing a crystal of an amino acid salt of HMB, including a step of dissolving an amorphous of amino acid salt of HMB in a solvent containing alcohol, a step of precipitating a crystal of an amino acid salt of HMB by stirring or allowing the solvent to left stand, and a step of collecting a crystal of an amino acid salt of HMB from the solvent.

[0031]

Examples of the amino acid preferably include a basic amino acid, more preferably arginine, ornithine, lysine, and histidine, furthermore preferably arginine, ornithine, and lysine, and most preferably arginine. As an amino acid, any one of the L form and the D form may be used, but the L form is preferable.

[0032]

Examples of the alcohol preferably include at least one alcohol selected from the group consisting of C1 to C6 alcohols, more preferably at least one alcohol selected from the group consisting of C1 to C3 alcohols, furthermore preferably at least one alcohol selected from the group consisting of methanol, ethanol, n-propanol, and isopropyl alcohol, and still furthermore preferably at least one alcohol selected from the group consisting of methanol and ethanol, and most preferably ethanol.

[0033]

The alcohol can be used by mixing one or more kinds thereof. In addition, water may be contained in the solvent containing alcohol. The water content in the solvent containing alcohol can generally be 40 wt% or less, preferably 20 wt% or less, furthermore preferably 10 wt% or less, and most preferably 5 wt% or less.

[0034]

Examples of a method of dissolving an amorphous of amino acid salt of HMB in the solvent containing alcohol include a method in which an amorphous of amino acid salt of HMB is suspended in the solvent and then heated to obtain a solution, or a method in which the solvent is filtrated to obtain a filtrate.

[0035]

The examples of a heating temperature in the method in which an amorphous of amino acid salt of HMB is suspended in the solvent, and then heated to obtain a solution include generally 0°C to 80°C, preferably 20°C to 70°C, and most preferably 40°C to 60°C. The examples of heating time can generally include 10 minutes to 6 hours, preferably 20 minutes to 4 hours, and most preferably 30 minutes to 2 hours.

[0036]

The amorphous of amino acid salt of HMB can be obtained by a method in Example 1 described below. The solvent obtained by dissolving the amorphous of amino acid salt of HMB therein is stirred or left to stand, and thereby a crystal of an amino acid salt of HMB can be precipitated.

[0037]

Examples of a concentration of the amorphous of amino acid salt of HMB to be dissolved in the solvent containing alcohol preferably include 50 g/L or more, more preferably 100 g/L or more, and furthermore preferably 150 g/L or more.

[0038]

A crystal of an amino acid salt of HMB may be precipitated by adding a crystal of an amino acid salt of HMB as a seed crystal to the solvent containing alcohol in which the amorphous of amino acid salt of HMB is dissolved, and then stirring or allowing the solvent to left stand. The addition of the seed crystal to the solvent can accelerate a precipitation rate. The seed crystal in the solvent is added to become a concentration of generally 0.05 to 15 wt%, preferably 0.5 to 10 wt%, and most preferably 2 to 7 wt%.

The crystal of an amino acid salt of HMB can be obtained by a method in Examples 2, 4, or 5 described below.

[0039]

Examples of a temperature for stirring or allowing the solvent to left stand include generally 0°C to 80°C, preferably 5°C to 50°C, and most preferably 10°C to 30°C. Examples of time for stirring or allowing the solvent to left stand include generally 1 to 100 hours, preferably 3 to 48 hours, and most preferably 5 to 24 hours.

[0040]

Examples of a method for collecting a crystal of an amino acid salt of HMB from the solvent are not particularly limited, but include filtration, pressurized filtration, suction filtration, centrifugal separation, and the like. Furthermore, in order to reduce attachment of the mother liquid and enhance the crystal quality, the crystal can be appropriately washed.

[0041]

The solution used for washing the crystal is not particularly limited, but a solution in which one or a plurality of kinds selected from water, methanol, ethanol, acetone, n-propanol, and isopropyl alcohol are mixed in an arbitrary ratio, can be used.

[0042]

The crystal of an amino acid salt of HMB can be obtained by drying the wet crystal obtained as above. As for the drying condition, any method may be used as long as the form of the crystal of an amino acid salt of HMB can be maintained, and reduced-
5 pressure drying, vacuum drying, fluidized-bed drying, forced air drying, or the like can be applied. The drying temperature may be any temperature as long as the attached water or solvent can be removed, but the temperature is preferably 80°C or less, more preferably 60°C or less.

[0043]

10 Under the above-described crystallization conditions, a high-purity crystal of an amino acid salt of HMB can be obtained. The purity of the crystal is generally 95% or more, preferably 96% or more, more preferably 97% or more, and most preferably 97.5% or more.

[0044]

15 The crystal of an amino acid salt of HMB, which can be produced by the above-described production method, includes a crystal of arginine salt nonhydrate of HMB of which powder X-ray diffraction pattern using CuK α as the X-ray source is specified by the values shown in Figs. 1 and 2 and Tables 1 and 2.

[0045]

20 (2) Method 2 for Producing Crystal of Present Invention

Examples a method for producing the crystal of the present invention include a method for producing a crystal of an amino acid salt of HMB, which includes a step of concentrating an aqueous HMB solution containing an amino acid-containing compound and having a pH of 2.5 to 11.0 and thereby precipitating a crystal of an amino acid salt of
25 HMB in the aqueous solution, and a step of collecting the crystal of an amino acid salt of HMB from the aqueous solution.

[0046]

HMB contained in the aqueous HMB solution may be a compound produced by any production method such as fermentation method, enzyme method, extraction method
30 from natural products, or chemical synthesis method.

[0047]

In the case where a solid matter hindering the crystallization is contained in the aqueous HMB solution, the solid matter can be removed using centrifugal separation, filtration, ceramic filter, or the like. In the case where a water-soluble impurity or salt
35 hindering the crystallization is contained in the aqueous HMB solution, the water-soluble impurity or salt can be removed, for example, by passing the solution through a column filled with an ion exchange resin, or the like.

[0048]

In the case where a hydrophobic impurity hindering the crystallization is contained in the aqueous HMB solution, the hydrophobic impurity can be removed, for example, by passing the solution through a column filled with a synthetic adsorption resin, activated carbon, or the like. It is possible to prepare the aqueous solution to have an HMB
5 concentration of usually 500 g/L or more, preferably 600 g/L or more, more preferably 700 g/L or more, and most preferably 800 g/L or more.

[0049]

Examples of the amino acid-containing compound preferably include a basic amino acid-containing compound, more preferably an arginine-containing compound, a
10 lysine-containing compound, an ornithine-containing compound, and a histidine-containing compound, and most preferably an arginine-containing compound, a lysine-containing compound, and an ornithine-containing compound. As an amino acid in the amino acid-containing compound, any one of the L form and the D form can be used, but the L form is preferable.

[0050]

Examples of the arginine-containing compound include free arginine and arginine hydrochloride. In a case of using free arginine as the arginine-containing compound, it is possible to obtain an aqueous HMB solution containing the arginine-containing compound having a pH of generally 2.5 to 11.0, preferably 2.8 to 10.5, and most preferably 3.0 to 10.0
20 by adjusting a pH of the aqueous HMB solution by using arginine.

[0051]

Examples of the lysine-containing compound include free lysine and lysine hydrochloride. In a case of using free lysine as the lysine-containing compound, it is possible to obtain an aqueous HMB solution containing the lysine-containing compound
25 having a pH of generally 2.5 to 11.0, preferably 2.8 to 10.5, and most preferably 3.0 to 10.0 by adjusting a pH of the aqueous HMB solution by using lysine.

[0052]

Examples of the ornithine-containing compound include free ornithine and ornithine hydrochloride. In a case of using free ornithine as the ornithine-containing
30 compound, it is possible to obtain an aqueous HMB solution containing the ornithine-containing compound having a pH of generally 2.5 to 11.0, preferably 2.8 to 10.5, and most preferably 3.0 to 10.0 by adjusting a pH of the aqueous HMB solution by using ornithine.

[0053]

Examples of a method for concentrating the aqueous HMB solution and thereby precipitating a crystal of an amino acid salt of HMB in the aqueous solution include a method for concentrating the aqueous solution under reduced pressure.

[0054]

In the method for concentrating the aqueous HMB solution under reduced pressure, the temperature of the aqueous solution is generally from 0°C to 100°C, preferably from 10°C to 90°C, most preferably from 20°C to 60°C. In the method for concentrating the aqueous solution under reduced pressure, the pressure reduction time is generally from 1 to 120 hours, preferably from 2 to 60 hours, most preferably from 3 to 50 hours.

[0055]

In the step of precipitating the crystal of an amino acid salt of HMB in the aqueous solution, a crystal of an amino acid salt of HMB may be added to the aqueous HMB solution as a seed crystal. The seed crystal is added to the aqueous solution such that the concentration thereof becomes generally 0.05 to 15 wt%, preferably 0.5 to 10 wt%, and most preferably 2 to 7 wt%. Specifically, the crystal of an amino acid salt of HMB can be obtained by, for example, a method in Examples 2, 4, or 5 described below.

[0056]

In the step of precipitating a crystal of an amino acid salt of HMB in the aqueous solution, the crystal of an amino acid salt of HMB may be precipitated by adding or adding dropwise, in the aqueous HMB solution, at least one solvent selected from the group consisting of alcohol, nitrile, and ketone. Alcohol, nitrile, and ketone may be used alone, or a plurality of kinds thereof may be used in combination.

[0057]

When adding or adding dropwise the solvent, the seed crystal may be added before precipitation of the crystal of an amino acid salt of HMB. The timing of adding the seed crystal can be generally within 0 to 5 hours, preferably within 0 to 4 hours, and most preferably within 0 to 3 hours, after starting adding dropwise or adding the solvent.

Furthermore, the seed crystal may be added before adding or adding dropwise the solvent in the aqueous HMB solution.

[0058]

Examples of the alcohol include the examples same as those described in 2. (1). Examples of the nitrile are preferably acetonitrile. Examples of the ketone are preferably at least one ketone selected from the group consisting of acetone, methyl ethyl ketone, methyl isobutyl ketone and diethyl ketone, more preferably at least one ketone selected from the group consisting of acetone and methyl ethyl ketone, and most preferably acetone.

[0059]

When adding or adding dropwise the solvent in the aqueous HMB solution, the temperature of the aqueous solution may be any temperature as long as it is a temperature not causing degradation of HMB, but in order to enhance the crystallization ratio of a crystal of amino acid salt nonhydrate of HMB by lowering the solubility, the temperature is

generally 80°C or less, preferably 70°C or less, more preferably 60°C or less, and most preferably 50°C or less. A lower limit value of the temperature is generally 0°C or more, and preferably 10°C or more.

[0060]

5 The amount in which the solvent is added or added dropwise to the aqueous HMB solution is generally from 0.5 to 30 times, preferably from 0.5 to 25 times, and most preferably from 0.5 to 10 times the amount of the aqueous solution.

[0061]

10 The time for which the solvent is added or added dropwise to the aqueous HMB solution is generally from 30 minutes to 48 hours, preferably from 2 to 30 hours, and most preferably from 3 to 20 hours.

[0062]

15 After a crystal of an amino acid salt of HMB is precipitated as described above, the precipitated crystal may be further ripened generally for 1 to 48 hours, preferably for 1 to 24 hours, and most preferably for 1 to 12 hours. The term “be ripened” means to grow the crystal by once stopping the step of precipitating a crystal of amino acid salt nonhydrate of HMB. After ripening the crystal, the step of precipitating a crystal of an amino acid salt of HMB may be restarted.

[0063]

20 The method for collecting a crystal of an amino acid salt of HMB is not particularly limited and examples thereof include collection by filtration, pressurized filtration, suction filtration, centrifugal separation, and the like. Furthermore, in order to reduce adhesion of the mother liquid and enhance the crystal quality, the crystal can be appropriately washed.

25 [0064]

A solution used for washing the crystal is not particularly limited and it is possible to use a solution in which one or a plurality of kinds thereof selected from water, methanol, ethanol, acetone, n-propanol, isopropyl alcohol, acetonitrile, acetone, methyl ethyl ketone, methyl isobutyl ketone, and diethyl ketone, are mixed in an arbitrary ratio.

30 [0065]

The thus-obtained wet crystal is dried, whereby a crystal of an amino acid salt of HMB of the present invention can be obtained. As for the drying condition, reduced-pressure drying, vacuum drying, fluidized-bed drying, or forced air drying may be applied. The drying temperature may be any temperature as long as the attached water or solvent
35 can be removed, but the temperature is preferably 80°C or less, and more preferably 60°C or less.

[0066]

Under the above-described crystallization conditions, a high-purity crystal of an amino acid salt of HMB can be obtained. The purity of the crystal of an amino acid salt of HMB is generally 93% or more, preferably 94% or more, more preferably 95% or more, and most preferably 96% or more.

5 [0067]

Examples of the crystal of an amino acid salt of HMB produced by the above production method include a crystal of lysine salt nonhydrate of HMB of which powder X-ray diffraction pattern using CuK α as the X-ray source is specified by the values shown in Fig. 4 and Table 4, a crystal of ornithine salt nonhydrate of HMB of which the pattern is
10 specified by the values shown in Fig. 6 and Table 6, and a crystal of ornithine salt nonhydrate of HMB of which the pattern is specified by the values shown in Fig. 8 and Table 9.

[0068]

[Analysis Examples]

15 (1) Powder X-Ray Diffraction

A powder X-ray diffraction apparatus (XRD), Ultima IV (manufactured by Rigaku Corporation), was used, and the measurement was performed according to the instruction book.

(2) Measurement of Concentration and Purity

20 The concentration of HMB and purity of various salts were measured using the following HPLC analysis conditions.

Guard column: Shodex SUGAR SH-G $\phi 6.0 \times 50$ mm

Column: SUGAR SH1011 $\phi 8.0 \times 300$ mm $\times 2$ columns in series

Column temperature: 60°C

25 Buffer: 0.005 mol/L of an aqueous sulfuric acid solution

Flow velocity: 0.6 mL/min

Detector: UV detector (wavelength: 210 nm)

(3) Measurement of Water Content of Crystal by Karl-Fischer Method

30 An automatic water measurement apparatus, AQV-2200 (manufactured by Hiranuma Sangyo Co., Ltd.), was used, and the water content of the crystal was measured according to the instruction book.

(4) Measurement of Amino Acid Content in Crystal

The amino acid content was measured by a phthalaldehyde (OPA) method by using HPLC having a fluorescence detector.

35 (5) Measurement of Melting Point

Melting Point M-565 (manufactured by BÜCHI Labortechnik AG) was used, and the melting point was measured using the following conditions according to the instruction book.

100°C to 250°C, 2°C/min

(6) Infrared Spectroscopic (IR) Analysis

Model FTIR-8400 (manufactured by Shimadzu Corporation) was used, and the analysis was performed according to the instruction book.

5 [0069]

[Reference Example 1]

Manufacturing of Free HMB Solution

Reagent of HMB calcium salt of 76.5 g in terms of the free form was dissolved in 850 mL of water. The aqueous solution passed through 640 mL of strong cation
10 exchange resin, XUS-40232.01(H⁺) (manufactured by The DOW Chemical Company) to remove Ca, and therefore 1.25 L of a solution containing 76.4 g of the free HMB was obtained.

[0070]

[Reference Example 2]

15 Examination on Crystallization of Arginine salt of HMB

Crystallization of arginine salt of HMB was examined with reference to U.S. Patent Application No. 2004/0176449. 48 mL of the aqueous solution containing 2.31 g of the free HMB was obtained in the same manner as in Reference Example 1. The obtained aqueous solution was concentrated to become 12 mL, and then 24 mL of
20 isobutanol was added thereto. Furthermore, when free arginine of 2.41 g was added and stirred for 1 hour at room temperature, a white transparent solution having two phases was obtained. When only an aqueous phase was collected and the aqueous phase was concentrated for 20 hours under conditions of 50°C and 15 mbar, a white precipitate of 6.5 g was obtained. When the powder was observed by using a polarizing microscope, it was
25 confirmed that the powder is a formless candy-like solid showing no polarization, and therefore is found to be noncrystalline and amorphous (melting point: 98°C to 126°C, 10°C/min). As above, the crystal of arginine salt of HMB was obtained by the method described in U.S. Patent Application No. 2004/0176449.

Examples

30 [0071]

Examples are described below, but the present invention is not limited to the following Examples.

[0072]

[Example 1]

35 Acquisition of Noncrystalline and Amorphous Arginine salt of HMB

To 200 mL of the aqueous free HMB solution obtained in Reference Example 1, 160 mL of an aqueous 128 g/L L-arginine solution was added to adjust the pH to 7.05. The obtained aqueous solution was provided for the next step.

[0073]

360 mL of the aqueous solution was concentrated at 50°C under reduced pressure at 15 mbar to remove the solvent, and thereby 31.1 g of white powder was obtained.

From the fact that no polarization was shown when the powder was observed by using a polarizing microscope, it was found that the powder is noncrystalline and amorphous.

[0074]

[Example 2]

Acquisition of Seed Crystal of Arginine salt Nonhydrate of HMB

1.5 mL of 100%-EtOH was added to 150 mg of the noncrystalline and amorphous arginine salt of HMB obtained in Reference Example 1 and heated at 50°C to be completely dissolved.

[0075]

A crystal was caused to naturally develop by stirring the aqueous solution at 25°C for 12 hours. The crystal slurry was further ripened for 12 hours, and the crystal was then collected by filtration, and forced air-dried at 25°C for 12 hours to obtain 60 mg of a crystal.

[0076]

The results of powder X-ray diffraction of the crystal are shown in Fig. 1 and Table 1. In the Table, “ 2θ ” indicates a diffraction angle ($2\theta^\circ$), and “Relative Intensity” indicates a relative intensity ratio (I/I_0). The relative intensity ratio is shown when the ratio is 1 or more.

[0077]

[Table 1]

2 θ	Relative Intensity
7.5	100
11.6	4
12.7	8
14.5	14
15.1	69
17.9	4
19.2	11
19.6	4
20.2	25
21.5	4
21.8	2
22.7	4
23.3	5
25.2	3
25.5	2
29.1	2
30.4	3
33.6	2
35.3	2
39.9	1

[0078]

[Example 3]

5 Acquisition of Crystal of Arginine salt Nonhydrate of HMB

160 mL of 100%-EtOH was added to 27.8 g of the noncrystalline and amorphous arginine salt of HMB obtained in Reference Example 1 and heated at 50°C to be completely dissolved.

[0079]

10 A crystal was caused to naturally develop by adding 0.1 g of the seed crystal of arginine salt of HMB obtained according to Example 2 to the aqueous solution. The crystal slurry was stirred at 25°C for 12 hours, ripened, and the crystal was then collected by filtration, and forced air-dried at 25°C to obtain 23.6 g of a crystal.

[0080]

15 11.6 g of the obtained crystal was further dried at 40°C for 24 hours under reduced pressure (at 15 mbar), and therefore 9.0 g of a crystal was obtained. It was possible to acquire 97.8% (area%) or more of the crystal of arginine salt nonhydrate of HMB in the measurement of the purity using HPLC.

[0081]

20 The results of powder X-ray diffraction of the crystal are shown in Fig. 2 and Table 2. The results of infrared spectroscopic (IR) analysis are shown in Fig. 3. In the

Table, “2 θ ” indicates a diffraction angle (2 θ°), and “Relative Intensity” indicates a relative intensity ratio (I/I₀). The relative intensity ratio is shown when the ratio is 5 or more.

[0082]

[Table 2]

2 θ	Relative Intensity
7.5	81
11.6	14
12.7	33
13.5	5
14.5	57
15.1	49
17.9	15
19.2	55
19.6	14
20.2	100
21.5	15
21.9	10
23.3	17
25.2	13
25.5	9
26.3	7
29.1	6
30.5	6
33.6	10
35.3	7
39.9	6

5

[0083]

As a result of measuring the L-arginine content in the crystal by HPLC, the content was 61.7 wt%, which substantially coincided with the theoretical value (59.6 wt%) of a mono-arginine salt. In addition, as a result of measuring the water amount contained in the crystal by the Karl-Fischer method, the content was 0.5 wt%.

10

[0084]

Accordingly, it was found that the crystal is a crystal of arginine salt nonhydrate of HMB. Various physical properties of the crystal acquired in Example 3 are shown in Table 3.

15

[0085]

[Table 3]

Water	Arginine content	Melting point
%	%	°C
0.5	61.7	177.9

[0086]

[Example 4]

Acquisition of Crystal of Lysine salt Nonhydrate of HMB

To 200 mL of the aqueous free HMB solution obtained in Reference Example 1,
5 94 mL of an aqueous 200 g/L L-lysine solution was added to adjust the pH to 6.03. The
obtained aqueous solution was provided for the next step. When the aqueous solution
was concentrated under conditions of 50°C and 15 mbar such that the weight of the
concentrated solution becomes 33.4 g, a crystal was caused to develop naturally, and
therefore white slurry was obtained. 20 mL of acetonitrile was added to the slurry, stirred
10 at 50°C for 1 hour, ripened, and the crystal was then collected by filtration.

[0087]

The obtained crystal was washed with 20 mL of acetonitrile, and then dried at
40°C for 24 hours under reduced pressure (at 15 mbar), and therefore 5.0 g of a crystal was
obtained. It was possible to acquire 97.4% (area%) or more of the crystal of lysine salt
15 nonhydrate of HMB in the measurement of the purity using HPLC.

[0088]

The results of powder X-ray diffraction of the crystal are shown in Fig. 4 and
Table 4. The results of infrared spectroscopic (IR) analysis are shown in Fig. 5. In the
Table, “2 θ ” indicates a diffraction angle (2 θ°), and “Relative Intensity” indicates a relative
20 intensity ratio (I/I_0). The relative intensity ratio is shown when the ratio is 2 or more.

[0089]

[Table 4]

2θ	Relative Intensity	2θ	Relative Intensity	2θ	Relative Intensity
4.8	9	26.6	23	37.3	4
8.5	72	27.5	4	37.6	3
14.7	5	27.8	6	37.9	3
14.8	3	27.8	5	38.7	3
17.0	65	28.2	2	39.6	5
17.6	5	28.9	5	39.7	5
18.1	45	31.0	12	40.5	3
18.5	48	31.2	8	41.1	3
19.5	100	32.5	5	41.7	3
20.4	12	33.0	5	42.8	3
21.8	9	33.3	4	43.9	3
22.2	23	33.5	5	44.8	3
22.6	4	33.8	10	51.8	3
23.4	6	34.4	14		
23.9	5	34.9	5		
24.5	7	35.0	4		
25.3	6	35.1	5		
25.5	21	35.8	4		
25.8	13	36.5	10		
26.0	6	36.6	7		

[0090]

As a result of measuring the L-lysine content in the crystal by HPLC, the content was 54.8 wt%, which substantially coincided with the theoretical value (55.3 wt%) of a monolysine salt. In addition, as a result of measuring the water amount contained in the crystal by the Karl-Fischer method, the content was 0.1 wt%.

[0091]

Accordingly, it was found that the crystal is a crystal of lysine salt nonhydrate of HMB. Various physical properties of the crystal acquired in Example 4 are shown in Table 5.

[0092]

[Table 5]

Water	Lysine content	Melting point
%	%	°C
0.1	54.8	186.2

[0093]

[Example 5]

Acquisition of Seed Crystal of Ornithine salt Nonhydrate of HMB

To 20 mL of the aqueous free HMB solution obtained in Reference Example 1, an aqueous 510 g/L L-ornithine solution was added to adjust the pH to 7.80. The obtained aqueous solution was provided for the next step. When the aqueous solution was concentrated under conditions of 50°C and 15 mbar, 2.8 g of a white precipitate was obtained. From the fact that no polarization was shown when the powder was observed by using a polarizing microscope, it was found that the powder is a crystal.

[0094]

[Example 6]

Acquisition 1 of Crystal of Ornithine salt Nonhydrate of HMB

To 20 mL of the aqueous free HMB solution obtained in Reference Example 1, 2.1 mL of an aqueous 510 g/L L-ornithine solution was added to adjust the pH to 7.08. The obtained aqueous solution was provided for the next step. After the aqueous solution was concentrated under conditions of 50°C and 15 mbar such that the weight of the concentrated solution becomes 3.9 g, and then 0.1 g of the seed crystal of ornithine salt nonhydrate of HMB obtained according to Example 5 was added thereto, 9 mL of 100%-ethanol was added dropwise over 30 minutes, and therefore the crystal was precipitated.

[0095]

The crystal slurry was ripened for 12 hours and the crystal was then collected by filtration, washed with an aqueous 100%-ethanol solution, forced air-dried at 25°C, and further dried for 24 hours at room temperature under reduced pressure (at 15 mbar), and therefore 250 mg of a crystal was obtained.

[0096]

It was possible to acquire 96.3% (area%) or more of the crystal of ornithine salt nonhydrate of HMB in the measurement of the purity using HPLC. The results of powder X-ray diffraction of the crystal are shown in Fig. 6 and Table 6. The results of infrared spectroscopic (IR) analysis are shown in Fig. 7. In the Table, "2 θ " indicates a diffraction angle (2 θ°), and "Relative Intensity" indicates a relative intensity ratio (I/I_0). The relative intensity ratio is shown when the ratio is 1 or more.

[0097]

[Table 6]

2 θ	Relative Intensity
5.1	100
10.5	1
10.9	2
14.0	2
15.3	8
16.4	2
16.8	2
19.4	2
20.4	3
21.4	2
21.9	3
23.9	1
24.9	1
25.5	2
26.1	1
28.1	1
30.7	1
34.7	1
35.4	1

[0098]

As a result of measuring the L-ornithine content in the crystal by HPLC, the content was 53.6 wt%, which substantially coincided with the theoretical value (52.8 wt%) of a mono-ornithine salt. In addition, as a result of measuring the water amount contained in the crystal by the Karl-Fischer method, the content was 2.0 wt%.

[0099]

Accordingly, it was found that the crystal is a crystal of ornithine salt nonhydrate of HMB. Various physical properties of the crystal acquired in Example 6 are shown in Table 7.

[0100]

[Table 7]

Water	Ornithine content	Melting point
%	%	°C
2.0	53.6	171.6

15 [0101]

[Example 7]

Acquisition 2 of Crystal of Ornithine salt Nonhydrate of HMB

4.6 L of the aqueous solution containing 280.3 g of the free HMB was obtained in the same manner as in Reference Example 1. 615 mL of an aqueous 510 g/L ornithine

solution was added to the obtained solution to adjust the pH to 6.2. The obtained aqueous solution was concentrated under conditions of 50°C and 15 mbar such that the weight of the concentrated solution becomes 696 g. 1.8 g of the seed crystal of ornithine salt nonhydrate of HMB obtained according to Example 5 was added thereto while maintaining the solution at room temperature, and thereby the crystal was precipitated. The crystal slurry was ripened for 24 hours at room temperature, and then subjected to centrifugal separation, and the crystal was then collected by filtration. The crystal was further vacuum-dried for 24 hours under conditions of 40°C and 30 hPa, and therefore 180 g of a crystal was obtained.

[0102]

As a result of measuring the L- ornithine content in the crystal by HPLC, the content was 51.6 wt%, which substantially coincided with the theoretical value (52.8 wt%) of a mono-ornithine salt. In addition, as a result of measuring the water amount contained in the crystal by the Karl-Fischer method, the content was 0.5 wt%.

[0103]

Accordingly, it was found that the crystal is a crystal of ornithine salt nonhydrate of HMB. Various physical properties of the crystal acquired in Example 7 are shown in Table 8.

[0104]

[Table 8]

Water	Ornithine content	Melting point
%	%	°C
0.5	51.6	169-172

[0105]

The results of powder X-ray diffraction of the crystal are shown in Fig. 8 and Table 9. In the Table, “2 θ ” indicates a diffraction angle (2 θ°), and “Relative Intensity” indicates a relative intensity ratio (I/I_0). The relative intensity ratio is shown when the ratio is 2 or more.

[0106]

[Table 9]

2 θ	Relative Intensity	2 θ	Relative Intensity
4.9	68	25.8	2
5.2	66	27.2	3
5.5	100	33.3	3
10.9	6	33.6	2
15.5	7	34.3	2
15.9	3	35.9	3
16.4	20	36.1	2
17.4	2	36.7	2
19.2	4	38.5	2
20.4	3	41.0	2
20.8	4		
21.3	6		
21.8	6		
22.2	2		
22.8	2		
23.5	3		
23.9	5		
24.5	2		
24.9	2		
25.4	2		

[0107]

When Fig. 8 which is a chart diagram of the crystal acquired in Example 7 was compared with Fig. 6 which is a chart diagram of the crystal acquired in Example 6, these do not coincide with each other. Accordingly, it was confirmed that the crystal has a different crystal form compared to the crystal obtained in Example 6.

[0108]

[Example 8]

10 Measurement of Solubility

The crystals of arginine salt nonhydrate of HMB obtained in Example 3 and calcium salt hydrate of HMB (manufactured by Tokyo Chemical Industry Co., Ltd.) were added respectively at room temperature until they dissolved in water, and after keeping the solution for a sufficient time under stirring, the supernatant containing no crystal was sampled and measured for the HMB concentration by using HPLC. The measurement results are shown in Table 10.

[0109]

[Table 10]

	Solubility (g/L)	Solubility (in terms of free form) (g/L)
Arginine salt nonhydrate of HMB	829	335
Calcium salt hydrate of HMB (*1)	150	129

*1: Purchased from Tokyo Chemical Industry Co., Ltd.

[0110]

As shown in Table 10, it was found that the acquired crystal of arginine salt nonhydrate of HMB is greatly enhanced in the solubility in water, as compared to an existing calcium salt of HMB.

[0111]

[Example 9]

Mixing of Crystals of Arginine salt of HMB and Lysine salt of HMB with Phosphate Buffer

A 100 g/L solution, in terms of free form, was prepared using the crystal of arginine salt nonhydrate of HMB obtained in Example 3, the crystal of lysine salt nonhydrate of HMB obtained in Example 4, and calcium salt hydrate of HMB (manufactured by Tokyo Chemical Industry Co., Ltd.), and mixed with 0.2 M phosphate buffer (pH: 6.80) in an arbitrary mixing ratio. The solution after mixing was measured for the light transmittance (660 nm) to evaluate the presence or absence of insoluble salt formation. The results are shown in Table 11.

[0112]

[Table 11]

Transmittance T% (660 nm)

	Mixing Ratio (Vol/Vol) (*1)			
	0.01	0.17	0.33	0.57
Arginine salt	-	100	100	100
Lysine salt	-	100	100	100
Calcium salt	60	0.11	0.02	0

(*1) Vol/Vol = HMB 100 g/L aqueous solution (Vol)/0.2 M phosphate buffer (Vol)

[0113]

As shown in Table 11, it was found that in the mixing with the phosphate buffer, the existing calcium salt of HMB forms an insoluble salt but the acquired crystals of arginine salt nonhydrate of HMB and lysine salt nonhydrate of HMB do not form an insoluble salt.

[0114]

[Example 10]

Mixing of Crystal of arginine salt of HMB and Glucose-Amino Acids-Electrolytes Infusion Solution

The arginine salt nonhydrate of HMB obtained in Example 3 and calcium salt hydrate of HMB (manufactured by Tokyo Chemical Industry Co., Ltd.) were mixed with a glucose-amino acids-electrolytes infusion solution for peripheral vein nutrition [pH: about 6.7; Product Name: Aminofluid Infusion Solution (Otsuka Pharmaceutical Factory, Inc.)] to obtain 0, 0.11, 0.21 and 0.42 weight/volume % solutions at final concentration, respectively, in terms of free form. The light transmittance T% (660 nm) was measured by ultraviolet and visible spectrophotometer immediately after the mixing or 24 hours after being left at room temperature to evaluate the presence or absence of insoluble salt formation. The results are shown in Table 12.

[0115]

[Table 12]

Transmittance T% (660 nm)

Time after addition (h)	T (%)			
	0	24	0	24
HMB final concentration (weight/volume%)	Calcium salt		Arginine salt	
0	100	100	100	100
0.11	100	70	100	100
0.21	90	28	100	100
0.42	65	20	100	100

15 [0116]

As shown in Table 12, it was found that in the mixing with the Aminofluid infusion solution, the existing calcium salt of HMB forms an insoluble salt but the acquired arginine salt nonhydrate of HMB does not form an insoluble salt.

[0117]

20 [Example 11]

Effect on Body Electrolyte When Glucose-Electrolytes Infusion Solution Containing Crystal of Arginine salt of HMB Is Administered

The arginine salt nonhydrate of HMB obtained in Example 3 and calcium salt hydrate of HMB (manufactured by Tokyo Chemical Industry Co., Ltd.) were mixed with a glucose- electrolytes infusion solution [Product Name: SOLITA-T No.3 Infusion Solution (AY Pharmaceuticals Co., Ltd.)] that does not contain phosphate ion to obtain 0 and 0.42 weight/volume % solutions at final concentration, respectively, in terms of free form. The solution was continuously administered to surgically stressed rats underwent intestinal tract scratch operation at a normal dose (240 mL/kg/day) for 3 days. On the final day of administration, urine was collected by 24-hour urine collection and measured for the urinary electrolyte concentration. The results are shown in Tables 13 and 14.

[0118]

[Table 13]

Urinary calcium excretion amount (mg/day)

	Urinary calcium excretion amount (mg/day)	
HMB final concentration (weight/volume%)	Calcium salt	Arginine salt
0	0.30	0.30
0.42	3.64	0.24

[0119]

[Table 14]

Urinary phosphate excretion amount (mg/day)

	Urinary phosphate excretion amount (mg/day)	
HMB final concentration (weight/volume%)	Calcium salt	Arginine salt
0	18.4	18.4
0.42	4.4	19.1

[0120]

As shown in Tables 13 and 14, it was found that in the combined administration with the SOLITA-T No.3 infusion solution, the existing calcium salt of HMB induces the increase in urinary calcium and the decrease in urinary phosphate excretion but the acquired crystal of arginine salt nonhydrate of HMB does not induce the above electrolyte abnormality.

[0121]

[Example 12]

15 Sensory Evaluation Test 1

[Preparation of Beverage (1)]

6 g of ornithine salt nonhydrate of HMB obtained in Example 7, 4 g of maltitol (manufactured by Mitsubishi Shoji Foodtech Co., Ltd.), 120 μ L of apple essence (manufactured by TAKATA KORYO CO., LTD.), 120 μ L of sugar flavor (manufactured by Ogawa & Co., Ltd.), and 200 mg of aspartame (manufactured by AJINOMOTO CO., INC.) were added to 200 mL of water and dissolved. An appropriated amount of a citric acid (manufactured by MC Food Specialties Inc) was added to this solution, and after adjusting a pH to 3.70, each 100 mL was dispensed into a glass bottle, and an aluminum cap was put. The glass bottle was heated at 90°C for 5 minutes, and then left to stand at room temperature to be cooled, and thereby Beverage (1) was produced. As a result of visually confirming the presence or absence of precipitation and turbidity immediately after being left to stand at room temperature, precipitation and turbidity were not found.

[0122]

[Preparation of Comparative Beverage (1)]

Comparative Beverage (1) was prepared in the same manner as in Beverage (1) except that calcium salt of HMB (HMB Kyowa, manufactured by KYOWA HAKKO BIO CO., LTD.) used instead of ornithine salt nonhydrate of HMB. The presence or absence of precipitation and turbidity immediately after being left to stand at room temperature was visually confirmed. As a result, white precipitation was found in a large amount.

[0123]

Eight panelists evaluated which one of Beverage (1) and Comparative Beverage (1) is preferably in flavor through a two-point discrimination test. As a result, eight out of eight evaluated that Beverage (1) is clearly preferable than Comparative Beverage (1) in flavor. From the result, it was found that the obtained ornithine salt of HMB has the excellent effect of enhancing the solubility and flavor compared to calcium salt of HMB.

[0124]

[Example 13]

Sensory Evaluation Test 2

[Preparation of Beverage (2)]

6 g of arginine salt nonhydrate of HMB obtained in Example 3, 4 g of maltitol (manufactured by Mitsubishi Shoji Foodtech Co., Ltd.), 120 μ L of black currant essence (manufactured by Ogawa & Co., Ltd.), 60 mg of molasses flavor (manufactured by Mitsui Sugar Co., Ltd.), and 200 mg of stevia (manufactured by B Food Science Co., Ltd.) were added to 200 mL of water and dissolved. An appropriated amount of 75% phosphate (manufactured by Taiyo Chemical Industry Co., Ltd.) was added to this solution, and after adjusting a pH to 3.70, each 100 mL was dispensed into a glass bottle, and an aluminum cap was put. The glass bottle was heated at 90°C for 5 minutes, and then left to stand at room temperature to be cooled, and immediately after that, the presence or absence of precipitation and turbidity was visually confirmed. As a result, no precipitation and turbidity were found.

[0125]

[Preparation of Comparative Beverage (2)]

Comparative Beverage (2) was prepared in the same manner as in Beverage (2) except that calcium salt of HMB (HMB Kyowa, manufactured by KYOWA HAKKO BIO CO., LTD.) used instead of arginine salt nonhydrate of HMB. The presence or absence of precipitation and turbidity immediately after being left to stand at room temperature was visually confirmed. As a result, white precipitation was found as same as before heating.

[0126]

Eight panelists evaluated which one of Beverage (2) and Comparative Beverage (2) is preferably in flavor through a two-point discrimination test. As a result, eight out of eight evaluated that Beverage (2) is clearly preferable than Comparative Beverage (2) in

flavor. From the result, it was found that the obtained arginine salt of HMB has the excellent effect of enhancing the solubility and flavor compared to calcium salt of HMB.

[0127]

While the present invention has been described in detail and with reference to
5 specific embodiments thereof, it will be apparent to one skilled in the art that various
changes and modifications can be made therein without departing from the spirit and scope
of the present invention. This application is based on Japanese Patent Application (Patent
Application No. 2016-125280) filed on June 24, 2016, the entirety of which is incorporated
herein by way of reference. All references cited herein are incorporated herein in its
10 entirety.

Industrial Applicability

[0128]

According to the present invention, a crystal of an amino acid salt of HMB
nonhydrate, which is useful, for example, as a product, a raw material, an intermediate, or
15 the like of health food, medicines, cosmetics, or the like, and a production method thereof
are provided.

Claims

[Claim 1] A crystal of an amino acid salt of a 3-hydroxyisovaleric acid (hereinafter referred to as HMB).

5

[Claim 2] The crystal according to claim 1, wherein the amino acid salt of HMB is a basic amino acid salt of HMB.

[Claim 3] The crystal according to claim 2, wherein the basic amino acid salt of HMB is arginine salt of HMB.

10

[Claim 4] The crystal according to claim 2, wherein the basic amino acid salt of HMB is lysine salt of HMB.

[Claim 5] The crystal according to claim 2, wherein the basic amino acid salt of HMB is ornithine salt of HMB.

15

[Claim 6] The crystal according to claim 3, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $7.5\pm 0.2^\circ$, $14.5\pm 0.2^\circ$, $15.1\pm 0.2^\circ$, $19.2\pm 0.2^\circ$, and $20.2\pm 0.2^\circ$.

20

[Claim 7] The crystal according to claim 6, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $11.6\pm 0.2^\circ$, $12.7\pm 0.2^\circ$, $17.9\pm 0.2^\circ$, $21.5\pm 0.2^\circ$, and $23.3\pm 0.2^\circ$.

25

[Claim 8] The crystal according to claim 7, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $19.6\pm 0.2^\circ$, $21.9\pm 0.2^\circ$, $25.2\pm 0.2^\circ$, $25.5\pm 0.2^\circ$, and $33.6\pm 0.2^\circ$.

[Claim 9] The crystal according to claim 4, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $8.5\pm 0.2^\circ$, $17.0\pm 0.2^\circ$, $18.1\pm 0.2^\circ$, $18.5\pm 0.2^\circ$, and $19.5\pm 0.2^\circ$.

30

[Claim 10] The crystal according to claim 9, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $22.2\pm 0.2^\circ$, $25.5\pm 0.2^\circ$, $25.8\pm 0.2^\circ$, $26.6\pm 0.2^\circ$, and $34.4\pm 0.2^\circ$.

35

[Claim 11] The crystal according to claim 10, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $4.8\pm0.2^\circ$, $20.4\pm0.2^\circ$, $31.0\pm0.2^\circ$, $33.8\pm0.2^\circ$, and $36.5\pm0.2^\circ$.

5 [Claim 12] The crystal according to claim 5, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $5.1\pm0.2^\circ$, $14.0\pm0.2^\circ$, $15.3\pm0.2^\circ$, $20.4\pm0.2^\circ$, and $21.9\pm0.2^\circ$.

[Claim 13] The crystal according to claim 12, wherein in powder X-ray diffraction,
10 the crystal further has peaks at diffraction angles (2θ) of $16.4\pm0.2^\circ$, $16.8\pm0.2^\circ$, $19.4\pm0.2^\circ$, $21.4\pm0.2^\circ$, and $25.5\pm0.2^\circ$.

[Claim 14] The crystal according to claim 13, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $10.9\pm0.2^\circ$.

15

[Claim 15] A method for producing a crystal of an amino acid salt of HMB, the method comprising:

a step of dissolving an amorphous amino acid salt of HMB in a solvent containing alcohol, a step of precipitating a crystal of an amino acid salt of HMB by stirring or
20 allowing the solvent to left stand, and a step of collecting a crystal of an amino acid salt of HMB from the solvent.

[Claim 16] A method for producing a crystal of an amino acid salt of HMB, the method comprising:

25 a step of concentrating an aqueous HMB solution containing an amino acid-containing compound and having a pH of 2.5 to 11.0 to precipitate a crystal of an amino acid salt of HMB, and a step of collecting a crystal of an amino acid salt of HMB from the aqueous solution.

30 [Claim 17] The production method according to claim 16, wherein the step of precipitating a crystal of an amino acid salt of HMB includes a step of adding or adding dropwise at least one solvent selected from the group consisting of alcohol, nitrile, and ketone.

35 [Claim 18] The production method according to claim 15 or 17, wherein the alcohol is at least one alcohol selected from the group consisting of C1 to C6 alcohols.

[Claim 19] The production method according to claim 17 or 18, wherein the nitrile is acetonitrile.

5 [Claim 20] The production method according to any one of claims 17 to 19, wherein the ketone is at least one ketone selected from the group consisting of acetone, methyl ethyl ketone, methyl isobutyl ketone, and diethyl ketone.

[Claim 21] The production method according to any one of claims 15 to 20, wherein the amino acid salt of HMB is a basic amino acid salt of HMB.

10

[Claim 22] The production method according to claim 21, wherein the basic amino acid salt of HMB is arginine salt of HMB, lysine salt of HMB, or ornithine salt of HMB.

15 [Claim 23] The crystal according to claim 5, wherein in powder X-ray diffraction, the crystal has peaks at diffraction angles (2θ) of $4.9\pm0.2^\circ$, $5.2\pm0.2^\circ$, $5.5\pm0.2^\circ$, $10.9\pm0.2^\circ$ and $15.5\pm0.2^\circ$.

20 [Claim 24] The crystal according to claim 23, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $15.9\pm0.2^\circ$, $16.4\pm0.2^\circ$, $17.4\pm0.2^\circ$, $19.2\pm0.2^\circ$, and $20.4\pm0.2^\circ$.

[Claim 25] The crystal according to claim 24, wherein in powder X-ray diffraction, the crystal further has peaks at diffraction angles (2θ) of $20.8\pm0.2^\circ$, $21.3\pm0.2^\circ$, $21.8\pm0.2^\circ$, $22.2\pm0.2^\circ$, and $22.8\pm0.2^\circ$.

25