


Claims:


1. Pyridone disulfide derivatives and its stereoisomers of formula (I)




(I)

wherein, R_1 , R_2 and R_3 are independently alkyl, alkoxy, halogen, halogenated alkoxy, halogenated alkyl, hydrogen and could be same or different and X is CH or N.


2. Pyridone disulfide derivatives of formula (I) according to claim 1, wherein R_1 is selected from the group comprising of methyl, methoxy, fluorine, trifluoromethyl, difluoromethoxy and hydrogen.
3. Pyridone disulfide derivatives of formula (I) according to claim 1, wherein R_2 is selected from the group comprising of methyl, methoxy and hydrogen.
4. Pyridone disulfide derivatives of formula (I) according to claim 1, wherein R_3 is selected from the group comprising of methyl and hydrogen.
5. ~~The A~~ process for preparation of pyridone disulfide derivatives and its stereoisomers of formula (I) as claimed in claim 1, the process comprising:
 (i) ~~treatment of~~ treating a compound of the following formula (IV)



with a dealkylating agent to give a compound of the following formula (V),



(ii) oxidizing the compound of formula (V) with an oxidizing agent ~~which was then oxidized~~ to give a compound of the following formula (VI),



and

(iii) treating the compound of formula (VI) further treatment with an acid in presence of a solvent, in the pH range of 4.5 to 8.5, provided to obtain the compound of formula (I) conforming to regulatory specifications.

6. The process according to claim 5, wherein the dealkylating agent is selected from the group comprising of sodium sulfide, hydrobromic acid and aluminium chloride.

7. The process according to claim 5, wherein the oxidizing agents is selected from 10-camphorsulfonyl oxaziridine or its stereoisomers and sodium hypochlorite.

8. The process according to claim 5, wherein the compound of formula (I) is obtained by treating the compound of fomula (VI) with an acid selected from the group ~~comprising~~ consisting of an organic acid, ~~like such as~~ acetic acid, citric acid, propionic acid and lactic acid or a mineral acid such as hydrochloric acid, sulphuric acid and nitric acid, and the solvent is selected from the group ~~comprising~~ consisting of esters, alcohols, ketones, hydrocarbons, halogenated hydrocarbons, water and mixtures thereof.

9. A pharmaceutical composition comprising an active pharmaceutical ingredient as claimed in claim 1 along with acceptable pharmaceutical excipients for use in the ~~and its use for treatment~~ of gastrointestinal disorders.

10. Use of a pharmaceutical composition comprising an active pharmaceutical ingredient as claimed in claim 1 along with acceptable pharmaceutical excipients in the manufacture of a medicament for the treatment of gastrointestinal disorders.