

Abstract

A mixture comprising one or more compounds of the general formula (II)



where

10 R⁶, R⁷ and R⁸ is independently of each other H, C₁ to C₂₀ alkyl, where C₁ to C₂₀ alkyl comprises straight-chain or branched C₁ to C₂₀ alkyl groups or cyclic C₃ to C₂₀ alkyl groups, alkanoyl having C₁ to C₈ alkyl, where C₁ to C₈ alkyl comprises straight-chain or branched C₁ to C₈ alkyl groups or cyclic C₃ to C₈ alkyl groups, cyanophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,4-dichlorophenyl, 3,4-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 15 dimethylphenyl, such as 3,4-dimethylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-ethylphenyl, 3-ethylphenyl, 4-ethylphenyl, 2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 4-isopropylphenyl, benzyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, (carboxymethyl)phenyl, in which the carboxyl group may also be esterified with C₁ to C₆ alkyl, where C₁ to C₆ alkyl comprises straight-chain or branched C₁ to C₆ alkyl groups or 20 cyclic C₃ to C₆ alkyl groups, (N,N-dimethylsulfonamido)phenyl, sulfonamidophenyl or 4-isopropoxyphenyl,

where R⁷ and R⁸ is independently of each other additionally also phenyl is carboxylic esters having C₁ to C₈ alkyl, where C₁ to C₈ alkyl comprises straight-chain or branched C₁ to C₈ 25 alkyl groups or cyclic C₃ to C₈ alkyl groups,

and (meth)acrylic acid and/or C₁ to C₈ alkyl (meth)acrylates.