


THIS SHEET HAS BEEN
REPLACED

Amadori reaction compounds and products, process for their manufacture, and their use

5

The present invention relates to novel Amadori reaction compounds and products, to the production thereof and to a new use of these compounds and products, having at least partly entered an Amadori rearrangement as per the following reaction scheme and/or a Maillard reaction:

10



25

Amadori reaction products are known; they are reaction products, for example, of an aminoacid or a peptide with a sugar, oligo- or polysaccharide having entered an Amadori rearrangement (J. Biol. Soc. 215 (1955), Henri Borsook et al.). Thus, in DE-C-3914354, a water-soluble glycoprotein of an aminoacid and a sugar is described which is isolated from an extract of Avena sativa sees. Further, EP-A-406087 describes water-soluble polysaccharide-glycopeptide complexes which are derived from the cell wall of Gram positive bacterium, and J. Biol. Chem. 1985, 260/9 states that NMR-spectroscopy has been used to characterize Amadori reaction products formed by reaction of glucose with free amino groups of protein.

35

The invention now describes specific novel Amadori rearrangement compounds as described in claim 1. preferred embodiments and improvements are described in claims 2 to 4. These compounds

40

This Sheet Has Been Replaced

SUBSTITUTE SHEET





reduce potassium ferricyanide, a test reaction for biological active substances formed in a reaction of sugar and aminoacid.

5 The invention further relates to a novel use of such compounds and simultaneously a novel use of Amadori reaction products of sugars and aminoacids in general, as described in claims 11 to 17. In the past, nobody has tested the various steps of purification of the extract of an Amadori reaction product (in order to get rid of ballast substances and impurities) for biological
10 activity so far.

Immunostimulating drugs are indeed known from natural sources such as mistletoe extracts, peat extracts etc. with the drawback of expensive treatment of large quantities of raw material to
15 obtain a few grams of active substance; uncontrollable impurities that might lead to toxicity and side effects and therefore, further, to problems with administration during practical use due to the complex nature and the hardly reproducible composition.

20 Comparable products or product mixes from artificial sources, such as interferon or other genetic engineering methods are even more expensive to make. Furthermore, the molecules of human interferon are very often too big to penetrate the human cell
25 wall so that only a fraction of the administered dose is effectively becoming active. Also, genetic engineering products usually have side effects and some are even toxic. Furthermore, some of them act effectively on one day and not on the next day, for reasons unknown so far.

30 Suprisingly, it has now been found that nearly any simple amino-acid/sugar complex after having at least partly entered an Amadori rearrangement does not show any of the above-mentioned disadvantages but on the contrary has a surprisingly high immunological activity. They can therefore be used in pharmaceutical
35 formulations and in cosmetics. These small molecules easily penetrate the cell wall and virtually act as a nutrient. They

induce the formation of natural interferon and other cytokines, including tumor necrosis factor. Even three days after the administration, they still show this stimulating effect on biological activity. This effect increases with increasing completion of the Amadori rearrangement and decreases again with increasing decomposition of the complex.

Instead of simple sugars also - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides may be used, for instance dextrane, which react similarly. Polysaccharides show biological activity and may retain some of this activity after they become oligosaccharides.

Very little has been known heretofore about the biological activity of these compounds. It was now found that combinations of these substances in contact with human leukocytes produce interferon and other cytokines. This is called polyclonal activation of the cells.

It is possible to test the substances produced under the influence of these compounds and to determine the biological activities in international units relevant to specific cytokines. These compounds are of especially high biological activity within a range of pure substance concentrations from 1-100 µg/ml. Within the molecule, the specific nature of the aminoacid is more important than the nature of the sugar part of the molecule.

Reaction products of L-aspartic acid with glucose or galactose - after having gone through the Amadori rearrangement - when contacted with human leukocytes and incubated in tissue culture media at 37°C for 20 hours in an atmosphere of 5% CO₂ will produce from 30 - 1000 antiviral units of interferon. The interferon is measured in a bioassay using human cancer cells.

Under the influence of these compounds, tumor necrosis compounds may also be produced.

This Sheet Has Been Replaced

~~SUBSTITUTE SHEET~~



The products which allow such an unexpected use have the general formula



wherein

- 5 R_1' represents a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, and
- 10 R_2' represents an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical.

Thus the group of biologically active compounds may cover either the specific Amadori rearrangement compounds described above or

15 the N-substituted derivatives of a number of different aminoacid compounds and one simple sugar, oligo- or - preferably low molecular weight, especially of less than 1000 daltons - polysaccharide, or N-substituted derivatives of one aminoacid compound and a number of simple sugars, oligo- and/or such polysaccharides, or else any combination of such derivatives, every single

20 one of them having sufficient biological activity.

Preferably, R_1' in the above formula may be a radical selected from the D-form of simple sugars, especially (but not exclusively)

25 ly) from the D-form of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine; R_2' may be a radical selected from the L-form of aminoacid compounds such as serine, glycine, histidine, arginine, glutamine, asparagine, alanine, aspartic acid, glutamic acid,

30 phenylalanine, treonine, cysteine, cystine, methionine, hydroxyproline, tryptophane, proline, tyrosine, valine, isoleucine, leucine and lysine or else peptides of these aminoacids in any combination.

35 The invention further relates to a process for obtaining the above mentioned compounds and products as described in claim 5 and whereby an intermediate is formed of the formula

~~SUBSTITUTE SHEET~~



wherein R' is a C_1 -1-deoxy radical in a straight carbon chain or any O-bridged form of a simple sugar or an oligo- or polysaccharide, and

5 R'' represents an aminoacid or a peptide radical, said intermediate being at least partly subjected to an Amadori rearrangement and/or to the Maillard reaction by continued heating of the reaction mixture - preferably under pressure - and simultaneously or subsequently removing the solvents. Preferred
10 embodiments of the process are described in claims 6 to 10.

In some cases, especially when an aminoacid having two carboxyl groups is used, it is advantageous to add to the process a buffer salt, such as sodium bicarbonate, preferably in a molar
15 ratio of 1:1.

It was further found that the Amadori rearrangement products are relatively susceptible to decomposition, and that decomposition products have the nature of dark brown and tar-like, unidentified
20 compounds having lost their biological activity. Therefore it is preferred to stop the Amadori rearrangement reaction at a stage at which the reaction mixture becomes light orange-brown in colour.

25 It is interesting to note that the intermediate reaction products, formed when the originally opaque solution of the aminoacid becomes clear (before the Amadori rearrangement), are easily hydrolysed, i.e. the reaction is reversible. With increasing rearrangement, the reversibility diminishes, i.e. the products
30 become more stable, and the colour gradually changes from light yellow to light orange, and then finally to orange-brown when the Amadori rearrangement seems to be complete. Samples taken during such rearrangement reaction and tested (according to various procedures described later) proved that the biological
35 activity increases with the Amadori reaction progressing, and decreases when further heating results in decomposition, for which a colour change to dark brown is a sign. Immediate re-



duction of ferricyanide and the resulting colour change will occur if the reaction mixture contains other keto groups and/or sulfur containing aminoacids such as cysteine; otherwise it will occur within 3 to 5 min, which is a good check for the degree of Amadori rearrangement developed. Unreacted sugars would show the colour change after half an hour or several hours only.

Isolation of the pure Amadori rearrangement products is carried out according to known methods based on binding the mixture on a strong cation-exchanger (such as Amberlite^(R) or Dowex^(R)), successively eluting with ammonia water, evaporating a chosen defined fraction of the eluate under reduced pressure and crystallising the pure compound from anhydrous methanol (J.E. Hodge and B.E. Fisher, Methods in Carbohydrate Chemistry, Vol.II, Reactions of Carbohydrates, 1963 Academic Press, N.Y., London, 1963, page 105-106; or Borsook et al. as quoted earlier; or J. Dubourg and P. Devilliers as quoted earlier).

In the process according to the present invention, all the above-mentioned preferences regarding the kind of radicals derived from simple sugar and aminoacid compounds remain unchanged. Additionally, a preferred mixture of sugar substrates comprises the D-forms of glucose, xylose, galactose, rhamnose and fructose in a weight ratio of about 20:10:4:1:1, while the preferred mixture of aminoacid substrates comprises the L-forms of serine, glycine, histidine, arginine, alanine, proline, tyrosine, valine, leucine, isoleucine and lysine in a weight ratio of 20.4:35.8:35.8:132: 180:360:216:160:72:68:780.

As already mentioned, the Amadori rearrangement products are able to reduce potassium ferricyanide, such a chemical test reaction providing a basis for quick determination of the biological activity of the composition formed in a reaction of sugar and aminoacid.

It has been found that Amadori reaction products are especially active if a mixture of simple sugars of the same composition and



in the same weight ratio as occurring in natural peat extracts are reacted with a mixture of aminoacid compounds of the same composition and in the same weight ratio as occurring in natural peat extracts in the presence of an aqueous solvent, preferably adding a lower alcohol - and optionally inorganic trace elements occurring in such natural peat extracts - at elevated temperature (and optionally under pressure), and subsequently prolonging the heating in order to cause an Amadori rearrangement of the obtained products, simultaneously or subsequently expelling the solvents, stopping the rearrangement reaction at the point when the reaction mixture becomes light orange-brown in colour, drying the products thus obtained and purifying the same by means of column chromatography and collecting the fractions that cause maximum reduction of potassium ferricyanide.

In one embodiment, the instant pharmaceutical formulations contain as an active ingredient at least one reaction product of the formula $R_1'-NH-R_2'$ or a specific compound of the formula R_1-NH-R_2 together with a pharmaceutically acceptable carrier and/or an adjuvant and/or optionally a lubricant in a weight ratio of active ingredient to the remaining components of between 1:1 and 1:100, preferably 1:8 to 1:20 and most preferably about 1:9.

Another advantageous pharmaceutical formulation contains - in addition to the active ingredient - lactose and a lubricant, the weight ratio of lactose to the lubricant being between 20:1 and 100:1, preferably 50:1.

These pharmaceutical preparations are used to treat and/or prevent hematological and/or immunological diseases and/or stimulate the immunosystem of humans and/or mammals by the induction of cytokine formation.

Another use for the active ingredients is in cosmetic preparations. The active ingredient is present in such preparations in amounts of 0,01-10% by weight, preferably 0,01-1% by weight and especially in amounts of 0,05-0,1%. These cosmetic preparat-



ions contain - besides the active ingredient - usual carriers, adjuvants, enriching components and/or fragrances.

5 The present invention will be further explained and demonstrated in the following examples, which do not limit in any respect the scope of the present invention.

Example 1:

10 The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with: 1.47 g (0.01 M) L-glutamic acid
0.84 g (0.01 M) NaHCO₃
0.91 g D-glucose
0.91 g galactose and
3.00 ml of redistilled water

15

The mixture was heated up to 80°C whereby - preferably under reduced pressure - 50% of the water were evaporated and then - under atmospheric pressure - heated to a temperature of 80-85°C, at which it was kept for 120 min until it became orange in
20 colour. The syrup-like concentrated aqueous solution was then evaporated to dryness under reduced pressure. The orange-red solid reaction product thus obtained was marked with the symbol D-10 and saved for biological tests.

25 Example 2:

The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with: 1.33 g (0.01 M) L-aspartic acid
0.84 g (0.01 M) NaHCO₃
0.91 g of D-glucose
30 0.91 g of galactose and
3.00 ml of redistilled water

35 The mixture was heated up to 80°C with stirring (by means of rotation), concentrated under pressure, whereby a total volume of 1.5 ml water was vaporized, and then treated under atmospheric pressure at a temperature of 80-85°C until it became light orange in colour; this took place within approx. 60 min.



The reaction mixture was a syrup-like concentrated aqueous solution and was dried under reduced pressure. The resulting reaction product was a dry, yellow-orange powder. It was marked with the symbol D-11 and saved for biological tests.

5

Example 3:

The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with:

1.05 g (0.01 M) L-serine
0.91 g D-glucose
0.91 g galactose and
2.50 ml redistilled water

10

The mixture was heated to 85-92°C with stirring (by means of rotation). After 100 min, the solution became orange in colour.

15 The pressure was reduced and the mixture was evaporated to dryness. The reaction product on the walls of the flask formed an orange transparent layer. The reaction product was scraped off and powdered. It was marked with the symbol D-12 and saved for biological tests.

20

Example 4:

The 25 ml flask of a rotary evaporator placed in a heated waterbath was charged with:

0.66 g (0.005 M) of D-aspartic acid
0.42 g (0.005 M) of NaHCO₃
0.45 g of D-glucose
0.45 g of galactose and
3.00 ml of redistilled water

25

The mixture was heated up to 80°C, evaporated - under pressure - whereby a volume of 1.5 ml water was vaporized, and then treated - under atmospheric pressure - at a temperature of 85°C. After 60 min. heating the mixture became orange in colour; the pressure was reduced and the resulting syrup-like concentrated aqueous solution evaporated to dryness. Before the residue became definitely dry, two times 10 ml water-free ethanol were introduced into the flask and evaporated in order to eliminate residual moisture. The dry reaction product thus obtained was

30

35

powdered, marked with the symbol D-13 and saved for biological tests.

Example 5:

- 5 In order to obtain - in a synthetic way - an equivalent of the biologically active fraction of a certain natural peat extract, the flask of the rotary evaporator placed in the heated water bath was charged with:
- | | |
|----|--------------------------|
| | 20.5 mg L-serine |
| | 35.8 mg L-glycine |
| 10 | 35.8 mg L-histidine |
| | 132.0 mg L-arginine |
| | 180.0 mg L-alanine |
| | 360.0 mg L-proline |
| | 216.0 mg L-tyrosine |
| 15 | 160.0 mg L-valine |
| | 68.0 mg L-isoleucine |
| | 72.0 mg L-leucine |
| | 780.0 mg L-lysine |
| | 2000.0 mg D-glucose |
| 20 | 1000.0 mg D-xylose |
| | 400.0 mg D-galactose |
| | 100.0 mg D-rhamnose |
| | 100.0 mg D-fructose |
| | 6.0 ml redistilled water |

- The mixture was stirred by means of rotation and heated under pressure for 45 min at a temperature rising from 75°C to 86°C. During that period, approx. 3 ml of water were evaporated and the substrates were totally dissolved. The mixture was then
- 30 treated for 30 min under atmospheric pressure at a temperature of 85-86°C for an Amadori rearrangement. During that period the solution quickly became red-brown in colour. The pressure was reduced, and heating at 84°C was continued, thus simultaneously evaporating the solvents. At the end of evaporation, two times
- 35 15 ml water-free ethanol were introduced and the reaction mixture was brought to dryness. The flask with the dried reaction product was placed in a desiccator over calcium chloride for 18

~~SUBSTITUTE SHEET~~



hours; then the reaction product was powdered. Approx. 4.5g of a powdered product were obtained and marked with the symbol EK₂-S.

5 A portion of 4g of this reaction product was dissolved in 20 ml of distilled water and placed on a chromatographic column of 25 mm x 330 mm size, filled with a sorbent Amberlite(R) XAD-2, analytical grade. The column was eluted with 0.4 ml/min distilled water. Fractions of 10 ml volume were collected to a total volume of 450 ml. The content of the fractions was monitored chromatographically. Fractions of consecutive numbers 11-13 were combined and evaporated under reduced pressure. These fractions were characterised by a high content of Amadori rearrangement products (confirmed with the potassium ferricyanide reduction test). The product was saved for biological tests under the
15 symbol of EK₂-S-11.

Biological tests for determining the biological activity were carried out with immunised Balb/C mice of both sexes, at the age of 8-10 weeks. Immunisation of mice is achieved by peritoneal
20 administration of 0.2 ml of a 10% suspension of sheep erythrocytes (SRBC), i.e. of 6×10^8 cells. The erythrocytes are fixed in a sterile Alsever's solution of the following composition:

glucose	2.05 g
sodium citrate	0.8 g
25 sodium chloride	0.42 g
citric acid	0.055 g
redistilled water to	100 ml

30 Into such Alsever's solution, a sheep blood cell aseptic sample is introduced in a ratio of 1:1 and the mixture is kept for at least 3 days at +4°C. The thus stabilised erythrocytes are then sampled aseptically and introduced into a phosphate buffered salt solution (PBS) in order to wash them out. Erythrocytes are rinsed with PBS twice, and are centrifugated for 10 min at 2000
35 rpm. The washed out cells are used in the form of a 10% suspension in PBS. Such a suspension is used for the immunisation of Balb/C mice.

SUBSTITUTE SHEET



The reaction product to be tested was administered intraperitoneally (i.p.) or orally (p.o.) four times at chosen doses, the first administration taking place 2 hours before immunisation of the mouse with SRBC, while the remaining three dosages were administered after immunisation at 24 h intervals.

Each tested group of animals was treated with different doses of the tested reaction product: 10 mg/kg, 1 mg/kg, 0.1 mg/kg and 0.01 mg/kg. A control group of animals was also immunised with SRBC, but instead of the substance to be tested, 0.2 ml of PBS were administered at the same time intervals.

Each group of animals, control and tested groups, in all experiments consisted of 8-12 mice.

On the fourth or (in case of determination of antibodies type 7S) tenth day after immunisation, mice were slightly anesthetized with ether and exsanguinated by eyeball extirpation. The blood was collected into test tubes. Next, the spinal cord was broken and spleen removed. The blood was used for obtaining the serum needed for determination of hemagglutinating antibodies of the 19S+7S and 7S types, while spleens were used to prepare the cells useful for determination of the percentage of cells able to form E-rosettes and of hemolytic activity. For such uses, mouse spleens were comminuted. The splenocyte cells obtained were suspended in approx. 2 ml of Hanks' medium at +4°C, layered on the Ficoll-Uropolin gradient of a density of 1.077, and then centrifugated for 15 min at 3000 rpm at +4°C. After separation from the interphase, the lymphocyte buffy coat was placed in the Hanks' medium at +4°C and rinsed twice with centrifugation for 7-10 min each time at 1800 rpm. The splenocytes were then suspended in 1 ml of Hanks' medium at such a ratio that it contained 1×10^6 cells.

For each test, the percentage of dead cells is determined by mixing a drop of a tested suspension of splenocytes with a drop



of ex tempore prepared dyestuff solution containing 4 parts of a 0.2% trypan blue solution and 1 part of a 4.25% NaCl solution. Under the microscope, the percentage of dead splenocytes is determined for each 100 cells. Dead cells are navy blue, while the bright cells are live cells. The presence of more than 10% of dead cells is critical; such a sample has to be eliminated from further use.

All steps carried out with the cells to be tested should be performed in a sterile, siliconized laboratory glass apparatus placed in an ice bath.

Example 6:

In the first test, an effect of the tested reaction products on the number of cells producing hemolytic antibodies (PFC-IgM) was determined. The test was carried out as follows: To 0.5 ml of an 0.5% agarose solution placed in a test tube and kept in a heated water bath at 45°C, 0.1 ml of a 10% suspension of SRBC (prepared as described above) were admixed. Then 0.1 ml of a splenocyte suspension having a density of 1×10^6 cells/ml was added, the mixture stirred rapidly and immediately poured out on slides previously covered with agarose. The slides are incubated at 37°C for two hours. Next, the tested samples are covered with guinea pig complement diluted at a ratio of 1:20 for a further 2h. After the incubation of the tested samples with the complement, the number of plaque forming cells (PFC) was counted and recalculated for 1×10^6 splenocytes. Each test was performed twice.

The strongest amplification of the response to SRBC expressed in terms of increase of the number of splenocytes producing hemolysins IgM (PFC) was observed after administration of the D-11 substance at a dose of 0.1 mg/kg. The amplification was 119%. When the daily dose was increased ten times up to 1 mg/kg, amplification of the response was lowered to 53%.



Reaction product D-12 showed the strongest activity - an increase of 58% - at a dose of 1 mg/kg.

5 Reaction product EK₂-S-11 in this test showed the strongest activity at a dose of 0.1 mg/kg (increase of 65%). At a dose ten times higher, i.e. 1 mg/kg, the increase was slightly lower, i.e. 52%.

10 Reaction product D-13 tested at a dose of 1 mg/kg caused an increase of 40%. When the dose was increased to 10 mg/kg, i.e. ten times, the response was only a 14% increase.

Example 7:

15 An active hemagglutination test was also carried out, determining antibodies of the anti-SRBC type 19S+7S and 7S antibodies. In order to determine the 19S - IgM type antibodies level, mouse serum was prepared on the fourth day after immunisation with SRBC, while for the determination of 7S-IgG type antibodies level such preparation took place on the tenth day after mice
20 immunisation with SRBC, which is related to the day of maximum count of antibodies of a given type in mice immunised with SRBC.

A. Determination of 19S+7S antibodies count.

25 A sample of blood was centrifuged for 30 min at 3500 rpm. From each sample thus prepared, serum was collected and placed in a heated water bath at a temperature of 56°C for 30 min in order to deactivate the complement. Next, a number of solutions at several different dilutions of each tested serum was prepared
30 (from 1:1 to 1:4096) employing a microtitrator and U-shaped microplates having a volume of 250 µl each. The diluted sera were incubated for 1 hour at room temperature. A drop of a 1% suspension of SRBC in PBS (prepared as described above) was added to each serum; the mixtures were incubated for a further 2
35 h at a temperature of 37°C and then stored at a temperature of +4°C. The results were taken next day. The maximum dilution at which hemagglutination is still caused was considered the anti-



body count. A ring at the bottom of the plate is a sign that hemagglutination occurs. A button-like formation at the bottom of the plate is considered as a negative result - no hemagglutination.

5

For statistical analysis of the results, the increase of dilution of serum in a tested substance was compared to the one in a control group.

- 10 Reaction product D-11 at a dose of 1 mg/kg increased the IgM count 2.57 times. At a ten times higher dose, the stimulation effect in comparison to the control group increased by 3.5 times.
- 15 Reaction product D-12 in this test showed a weaker activity. At a dose of 0.1 mg/kg, it increased the IgM count 2 times, at a dose of 1 mg/kg 1.4 times.

- 20 Reaction product EK₂-S-11 showed the strongest activity in this test. At a dose of 0.1 mg/kg, it increased the IgM count 4.3 times, at a dose of 1 mg/kg 3.6 times.

Reaction product D-13 at a dose of 1 mg/kg increased the IgM count 3 times, at a ten times higher dose 1.5 times.

25

B. Determination of 7S antibody count

- 30 The tested inactivated sera were combined - at a ratio of 1:1 - with a 0.1 M solution of 2-mercaptoethanol and the mixtures incubated for 30 min at a temperature of 37°C. 2-Mercaptoethanol destroys immunoglobulines of the 19S-(IgM) type, while immunoglobulines of the 7S-(IgG) type are not susceptible to the action of 2-mercaptoethanol.

- 35 After 30 min of incubation the reduction reaction was stopped by means of cooling down to a temperature of +4°C for 15 min. Next, a number of dilutions was prepared in a similar manner as de-



scribed above with respect to determining the 19S-antibody count and combined with a 1% suspension of SRBC; after 2 hours of incubation at 37°C, the samples were stored at a temperature of +4°C. The results were evaluated on the following day, according to the criteria of determining the hemagglutination count described above. Simultaneously, a control test was carried out with a combination of a 1% suspension of SRBC with PBS in a ratio of 1:1.

10 When the substance D-11 was tested as described above, at a dose of 1 mg/kg it increased the production of antibodies IgG 3.16 times. At a dose of 10 mg/kg, the increase was 2.2 times.

15 Reaction product D-12 tested at a dose of 0.1 mg/kg and 1 mg/kg respectively stimulated the production of antibodies IgG 1.3 times, at a dose of 10 mg/kg 1.5 times.

20 Reaction product EK₂-S-11 at a dose of 0.1 mg/kg stimulates the production of IgG 1.9 times, at a dose of 1 mg/kg 2.89 times (in comparison with the control).

25 The results of tests A and B obtained for each production lot or for each fraction of the biologically active reaction products synthesized according to the invention and giving the above-mentioned immunological response were subjected to statistical analysis by the T-student method, $\alpha=0.05$. Results obtained for each dose tested were compared with a parallel control test and showed an increase of biological activity.

30 Example 8:

The group of biologically active reaction products obtained according to Examples 1 to 5 have also been submitted to the test in which the percentage of splenocytes forming E-rosettes was determined.

35

250 μ l of a 1% suspension of SRBC and 250 μ l of cells to be tested at a concentration of 1×10^6 cells/ml were added to 550



5 μ l of Hanks' medium. Each such sample was incubated in a heated water bath equipped with a shaker for 15 min at a temperature of 37°C. Then, it was stored at a temperature of +4°C for a further 20 h. The percentage of splenocytes forming E-rosettes with SRBC was determined after the suspension was coloured with 1 to 3 drops of crystal violet.

10 Each sample was subjected to determination of the percentage of splenocytes three times, counting at each instance 400 splenocytes. A splenocyte surrounded with at least 3 erythrocytes was considered an E-rosette.

15 For statistical evaluation, the percentage increase of the number of splenocytes with E-rosettes was compared between the substances to be tested and a control group.

20 In this test, the strongest stimulating effect was shown by the reaction products D-11 (63%) and EK₂-S-11 (70%) at a dose of 1 mg/kg. At a dose ten times smaller, i.e. 0.1 mg/kg, the values decreased to 45% and 57% respectively.

25 Reaction product D-12 at a dose of 1 mg/kg caused an increase of the ability to form E-rosettes of 22% in comparison to the control group. The respective value for a dose ten times smaller, i.e. of 0.1 mg/kg, was 29%.

30 Reaction product D-13 shows the maximum effect at a dose of 1 mg/kg (58%), while at higher doses the effect is slightly smaller.

Biological activity of synthesised compounds was evaluated according to the following tests:

- 35 1. Test for determination of the percentage of splenocytes forming E-rosettes, carried out according to Bach and Dardenne (Cell. Immunol. 3, 1-16, 1972)
2. Test for determination of the number of cells producing hemolytic antibodies of an IgM type, carried out according



to the Jerne method, modified by Mishell and Dutton (J. Exp. Med. 126, 423-442, 1967) and

3. Test for determination of a hemagglutination 19S + 7S and 7S antibody count, carried out according to Adler's active hemagglutination methods (J. Immunol. 95, 26-38, 39-47, 1965) with the use of microplates (J. Immunopharmacol. 4, 43-52, 1982).

Example 9:

- 10 A rotary evaporator flask placed in a heated water bath was charged with:
- | | |
|-----------------|--|
| 1.33 g (0.01 M) | L-aspartic acid |
| 0.84 g (0.01 M) | NaHCO ₃ |
| 10.00 g | hydrolysed dextrane of an average molecular weight of 3000 daltons |
| 10.00 ml | redistilled water. |
- 15

- The mixture was heated under pressure at a temperature of 70°C until the solid substances were completely dissolved, expelling by means of distillation during that period approx. 3 ml of water (heating time was approx. 30min). The flask with the solution was loosely covered, placed in a steam steriliser and heated under pressure at a temperature of 121°C for 40 min. After cooling down, the resulting yellow-orange solution was diluted with 15 ml of water, clarified by means of centrifugation and spray dried by air having an inlet temperature of +160°C and an outlet temperature of +85°C. 10.5 g of a light beige reaction product resulted that was easily soluble in water.
- 20
- 25

- 30 The presence of Amadori rearrangement products in this reaction product was confirmed by a test by the potassium ferricyanide method described by Borsook, Abrams and Lowy, J. Biol. Chem 215, (1955), 111-124 and by chromatographic methods.

- 35 Biological tests as described in the preceding Examples confirmed an immunotropic activity of the above product similar to the one exhibited by preparations obtained with simple sugars.

Example 10:

A conical flask was charged with:

- 5.0 g hydrolysed dextrane of an average molecular weight of approx. 5000 daltons
1.1 g L-proline
4.0 ml redistilled water.

The content was dissolved by means of stirring. The uniform mixture thus obtained was placed in a steam steriliser and heated under pressure at a temperature of 110°C for 40 min. The resulting transparent orange solution was diluted with 20 ml of redistilled water and clarified by means of centrifugation. The clear solution was spray dried.

5.3 g of a reaction product easily soluble in water was obtained. Immunotropic activity was similar to the one observed in other experiments according to the preceding examples.

Example 11:

Conventional methods test the biological activity of the compounds in mice, but not in humans. For this reason, new bioassays have been introduced, which measure the amounts and activity of cytokines released from the human peripheral blood leukocytes (PBL) treated with the reaction products according to the Examples 1 to 5, 9 and 10. The cytokines are the hormone-like proteins that play an important role in practically all of the immunological reactions, as well as in the regulatory processes responsible for the maintenance of homeostasis.

Cytotoxicity assays. Cytotoxicity of the reaction products was determined in human lung adenocarcinoma cell line A549 (included in the American Type Culture Collection - ATCC CCL 185). The cell monolayers were trypsinized, suspensions of 2×10^5 cells/ml in Dulbecco's-modified Eagle's minimum essential medium (DMEM) plus 10% calf serum (CS) were mixed with various doses of the drugs, seeded in the plastic microplates, and incubated for 48 h



at 37°C. CD_{50} was the minimal concentration of the compound which caused approximately 50% destruction of the cell culture, as measured by staining with 0.015% solution of neutral red in DMEM.

5

Cytokine induction. Buffy coats from healthy blood donors were obtained from the regional transfusion center. The erythrocytes were lysed by NH_4Cl treatment according to Cantell et al.

(Cantell, K., Hirvonen, S., Kauppinen, H.L.: Production and
10 Partial Purification of Human Immune Interferon. Meth. Enzymol.,
119, 54, 1986). The leukocytes from a single donor containing
approximately 8×10^6 leukocytes/ml in RPMI 1640 medium supple-
mented with 10% fetal calf serum (FCS), L-glutamine, and anti-
15 biotics were used. All lots of FCS were pretested. Only non-
mitogenic FCS for PBL cultures was used. The cytokine inducers
were added to 1 ml volumes of the cultures. The reference cyto-
kine inducers were phytohemagglutinin (PHA) (Pharmacia Fine Che-
micals, Sweden) and LPS from E. coli 0111:B4 (Difco Laborator-
ies). The induced cultures of PBL were incubated in an atmo-
20 sphere of 5% CO_2 in air at 37°C for 20 h and centrifuged. Super-
natants were stored at 4°C and assayed for TNF and IFN activity
within one week.

Interferon (IFN) assay. The confluent monolayer of A549 cells
25 was prepared in the microplates in DMEM with 10% CS, L-glut-
amine, and antibiotics (100 units/ml penicillin and 100 μg /ml
streptomycin). IFN samples diluted in plates were added to the
cell monolayer and incubated at 37°C for 20 h in 5% CO_2 in air.
The cells were then washed and challenged with encephalomyo-
30 carditis virus (EMCV). The titer of IFN was defined as the di-
lution of IFN sample that reduced the virus cytopathogenic
effect by 50% after 48 h of incubation. The MTT (3-[4,5-
dimethylthiazol-2-yl]-2,5- diphenyltetrazolium bromide) method
(Hansen, M.B., Nielsen, S.E., and Berg, K.: Re-examination and
35 Further Development of a Precise and Rapid Dye Method for
Measuring Cell Growth/ Cell Kill. J. Immuno. Meth., 1989, 119,
203-210) to measure the cell kill in the ELISA scanner was also



used. Laboratory standards of IFNs were included in all assays: Recombinant human IFN- γ (Genentech Inc., USA, specific activity 2×10^8 units/mg), the natural human leukocyte IFN- α (3×10^6 IU/ml) and IFN- γ (2×10^6 IU/ml) obtained from Dr. K. Cantell, Helsinki, Finland.

Tumor Necrosis Factor (TNF) assay. The cytotoxic activity of TNF was measured in L929 cells according to Flick and Gifford (Flick, D.A., Gifford, G.E.: Comparison of in Vitro Cell Cytotoxic Assays for Tumor Necrosis Factor. J. Immunol. Meth., 68, 167, 1984).

The samples and actinomycin D solution were added to monolayer cultures of the cells. After incubation at 37°C for 20 h, the cultures were stained with crystal violet and toxic effects were determined. The amount causing approximately 50% destruction of the cell cultures was defined as one unit of TNF activity. Comparison with a preparation of TNF- α (Genentech Inc., USA) showed that 1 unit in our assays was equal to 100 - 200 pg/ml TNF.

Cytokine neutralization assays. The antisera used were: rabbit anti-human TNF- α , lot 2958-40 and rabbit anti-human IFN- γ , lot 2891-56 (Genentech Inc., USA), sheep anti-human IFN- α, β , and sheep anti-human IFN- γ (obtained from Dr. K. Cantell, Finland).

The cytokine preparations were treated with the sera diluted 1:200 in culture medium and incubated for one hour at room temperature. Then, the residual IFN or TNF activities were determined as described.

Five different batches (L_1 to L_5) of PBL prepared from the blood of healthy blood donors were used. The optimal PBL concentration for the assays was found to be 8×10^6 cells per one ml of medium (RPMI 1640 supplemented with 10% fetal calf serum and antibiotics).

Incubation of human PBL with the new reaction products I - XI (Table 1) resulted in IFN and TNF synthesis. The observed re-



sponses were dose related in the range of 3 - 100 µg/ml of the compounds (Table 2). The compounds used in the indicated concentrations were non-cytotoxic. In all of the bioassays, the negative and positive controls were included. The negative controls measured the amounts of the cytokines (IFN and TNF) produced spontaneously without the addition of any exogenous stimulants. The positive controls indicated the amounts of the cytokine produced in response to a known reference inducer; in our case this was phytohemagglutinin (PHA, Pharmacia, Sweden, 10 µg per ml).

It should be pointed out that the cytokine induction in human PBL cultures obtained from different individuals usually gives considerable variation of the results. The phenomenon may be explained in terms of genetic differentiation of human population. Furthermore, PBL cultures often produce IFN and TNF spontaneously.

In other words, high responders and low responders or even non-responders are commonly observed among the healthy donors of PBL.

Despite the presented limitations, the results of the bioassays showed that PBL (L₁ to L₅) treated with the reaction products (I - XI) produced IFN and/or TNF that could be measured quantitatively.

In the case of L₁ which contained leukocytes of the high responder, the reaction product II (containing L-form of aspartic acid) was found to be considerably more active as a cytokine inducer than the reaction product III (containing the D-form of aspartic acid which also is more expensive by two orders of magnitude). The observation is significant because mainly L-forms of the amino-acids are recognized by cells as natural substrates in biochemical reactions.

Furthermore, for the expression of biological activity of the reaction products, the amino-acid part of the molecule is much more important than the form of sugar. Instead of single sugars, - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides (such as dextrans, reaction products X - XI) can be used and they react similarly.

And vice versa, polysaccharides containing the amino-acid residues may have biological activity, and this activity is retained when they are decomposed to oligosaccharides with the bound amino-acid (data not shown).

Similar results may also be observed if a short peptide is taken instead of a single amino-acid and is used to stimulate the leukocytes to produce cytokines (data not shown).

Seven reference batches of the unfractionated TTP assayed in over 100 PBL cultures from different donors produced from 10 to 1.000 units of IFN per ml, and from 9 to 750 units of TNF per ml. The fraction EK₂-S prepared from a mixture corresponding to the contents of natural peat extract (Example 5) has been assayed in eight PBL cultures from eight different blood donors. It was found to be the most active preparation in inducing both IFN and TNF (data not shown).

25

Possible applications of the reaction products are as immunomodulators and such activity was clinically useful. Tissue regeneration is another proven activity. Anti-cancer activity probably connected with the presence of the induced interferon and tumor necrosis factor was also observed. Anti-viral activity was also noted.

30

The main use of the above reaction products involves oral administration, but parenteral treatment is also possible, as is topical application. The products appear relatively stable.

35

This Sheet Has Been Replaced

~~SUBSTITUTE SHEET~~

Table 1. List of Reaction Products

No.	Substrates
5	I (D-10) L-glutamic acid, glucose, galactose
	II (D-11) L-aspartic acid, glucose, galactose
	III (D-13) D-aspartic acid, glucose, galactose
	IV (D-12) L-serine, glucose, galactose
	V EK ₂ -S (fractions 11 - 13)
10	VI EK ₂ -S (fractions 6 - 7)
	VII EK ₂ -S (fractions 8 - 10)
	VIII EK ₂ -S (fractions 28 - 34)
	IX L-proline, glucose
	X L-aspartic acid + dextrane (variety 1)
15	XI L-aspartic acid + dextrane (variety 2)

Example 12:

Pharmaceutical formulations containing as an active ingredient the reaction products according to Examples 1 to 5, 9 and 10, were prepared using the following reaction products:

A. Tablets/Granules:

- 5.0 g of the reaction product obtained according to Example 1 or 10 (active substance),
- 444.0 g of pharmaceutically acceptable lactose
- 1.0 g of lubricant, e.g. MYVATEX^(R), trademark of Eastman Kodak)

The ingredients were mixed and granulated with 30% aqueous ethanol in a conventional way, then dried at 40°C. The granules were used to prepare capsules, each containing approx. 450 mg of granules, i.e. 5 mg of the active substance. Alternatively, the granules were used to form tablets, each weighing approx. 450 mg and containing 5 mg of the active substance.

B. In the same conventional manner, the active substances obtained according to the preceding examples 1 to 5, 9 and 10 were formulated into other pharmaceutical formulations using suitable carriers.

SUBSTITUTE SHEET

This Sheet Has Been Replaced




Table 2. Biological Activity of the Reaction Products I-XI in Human PBL

		Dose µg/ml	IFN units/ml					TFN units/ml				
			L ₁	L ₂	L ₃	L ₄	L ₅	L ₁	L ₂	L ₃	L ₄	L ₅
5	Control	-	10	<10	<10	20	<10	80	9	27	27	<9
	PHA	10	100	30	30	60	100	250	80	250	250	250
10	I	100	100	<10	<10	20	-	250	9	9	160	-
		30	-	-	-	30	-	-	-	-	500	-
		10	30	<10	<10	30	-	80	9	18	160	-
		3	-	-	-	10	-	-	-	-	160	-
15	II	100	100	10	<10	10	<10	250	18	18	250	<9
		30	-	-	-	10	<10	-	-	-	250	<9
		10	1000	10	30	10	10	250	50	27	250	<9
		3	-	-	-	10	<10	-	-	-	160	<9
	III	100	<10	<10	10	10	<10	250	9	18	250	<9
		30	-	-	-	30	<10	-	-	-	250	<9
20		10	<10	<10	<10	10	<10	80	27	18	250	<9
		3	-	-	-	10	<10	-	-	-	80	<9
	IV	100	<10	10	10	20	<10	80	60	9	160	<9
		30	-	-	-	20	<10	-	-	-	27	<9
		10	<10	30	<10	20	<10	80	50	18	80	<9
25		3	-	-	-	20	<10	-	-	-	80	<9
	V	100	30	<10	<10	60	-	80	50	18	250	-
		30	-	-	-	60	-	-	-	-	80	-
		10	<10	<10	<10	10	-	80	27	9	80	-
		3	-	-	-	10	-	-	-	-	80	-
30	VI	100	<10	<10	<10	20	-	80	27	18	80	-
		30	-	-	-	10	-	-	-	-	80	-
		10	10	<10	<10	20	-	50	27	18	160	-
		3	-	-	-	10	-	-	-	-	250	-
35	VII	100	<10	<10	<10	-	-	80	27	27	-	-
		30	-	-	-	-	-	-	-	-	-	-
		10	10	<10	<10	-	-	80	27	27	-	-
		3	-	-	-	-	-	-	-	-	-	-
	VIII	100	<10	<10	<10	-	-	80	160	27	-	-
		30	-	-	-	-	-	-	-	-	-	-
40		10	<10	<10	<10	-	-	750	80	27	-	-
		3	-	-	-	-	-	-	-	-	-	-
	IX	100	10	<10	<10	10	-	27	27	18	80	-
		30	-	-	-	20	-	-	-	-	80	-
		10	100	30	<10	20	-	80	27	18	80	-
45		3	-	-	-	10	-	-	-	-	50	-
	X	100	100	<10	20	20	-	27	27	18	250	-
		30	-	-	-	30	-	-	-	-	80	-
		10	<10	10	<10	30	-	18	50	27	50	-
		3	-	-	-	30	-	-	-	-	250	-
50	XI	100	<10	10	<10	30	-	18	27	27	50	-
		30	-	-	-	10	-	-	-	-	250	-
		10	<10	30	<10	10	-	18	80	80	80	-
		3	-	-	-	20	-	-	-	-	750	-



Example 13:

The active substances obtained according to preceding Examples 1 to 5, 9 and 10 were used as a beneficial additive to cosmetic preparations intended for everyday hair and body care, the content of the substances being within a range of 0.01 - 10% by weight, depending on the type of the preparation, the method of application and the frequency of use intended for the particular preparation.

This Sheet Has Been Replaced~~SUBSTITUTE SHEET~~

C L A I M S

1. An Amadori rearrangement compound of the formula



wherein



- 5 R_1 comprises a 1-amino-1-deoxy-2-ketose derived from a sugar radical, preferably in its D-form, selected from the group consisting of glucose, xylose, galactose, rhamnose, fructose, mannose, 2-deoxy-glucose, 6-deoxy-glucose, glucosamine and galactosamine, and
- 10 R_2 comprises an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, preferably its L-form, selected from the group consisting of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid,
- 15 phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine.
- 20 2. A compound according to claim 1, wherein R_1 comprises a sugar radical selected from the group of D-glucose and D-galactose.
3. A compound according to claim 1 or 2, wherein R_2 comprises an aminoacid radical selected from the group of L-glutamic acid, L-aspartic acid and L-serine.
4. A mixture of compounds according to claim 1.
- 25 5. Process for the manufacture of at least one reaction product of the formula $R_1'-NH-R_2'$, wherein R_1' comprises a 1-amino-1-deoxy-2-ketose derived from a simple sugar, an oligo- or a - preferably low molecular weight, especially of less than 1000 daltons - polysaccharide and R_2' comprises an aminoacid
- 30 or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, wherein at least one substance selected from the group of aminoacids and -





preferably low molecular weight, especially of less than 1000 daltons - peptides, and at least one second substance selected from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, are reacted with each other at elevated temperature - optionally under pressure - and/or in the presence of a lower alcohol, whereafter the intermediate(s) is (are) subjected to an Amadori rearrangement and/or to the Maillard reaction.

6. Process according to claim 5, wherein the reaction is carried out in the presence of an aminoacid having two carboxyl groups and a buffer salt, preferably sodium bicarbonate in a molar ratio of 1:1.
7. Process according to claim 5 or 6, wherein at least one simple sugar is reacted with at least one monobasic aminoacid, and the molar ratio of the simple sugar(s) to the aminoacid(s) is between 2:1 and 1:1, preferably 1.5:1.
8. Process according to any one of claims 5 to 7, wherein the reaction of the sugar(s) and/or saccharide(s) with the aminoacid(s) and/or peptide(s) is carried out in a concentrated aqueous solution at a temperature of 75-90°C, and subsequent Amadori rearrangement is carried out at the same temperature - optionally under pressure, with simultaneous or subsequent removal of the solvents - until the reaction mixture becomes light orange-brown in colour.
9. Process according to any one of claims 5 to 8, wherein a mixture of simple sugars of the same composition and in the same weight ratio as occurring in natural peat extract is combined with a mixture of aminoacid compounds of the same composition and in the same weight ratio as occurring in said extract - and optionally with inorganic trace elements occurring with said extract - the resulting substrate mixture being reacted in the presence of water as a solvent at elevated temperature - and optionally under pressure

~~SUBSTITUTE SHEET~~

This Sheet Has Been Replaced





and/or in the presence of a lower alcohol - whereafter the resulting products are dissolved and the reaction is continued to cause an Amadori rearrangement and/or a Maillard reaction which is stopped when the reaction mixture becomes orange-brown in colour, whereafter the reaction mixture is dried.

10. Process according to any one of claims 5 to 9, wherein the dried product is purified by column chromatography, whereby specific fractions causing reduction of potassium ferricyanide are collected.
11. Use of at least one compound according to any one of claims 1 to 4, or of at least one N-substituted reaction product of the general formula $R_1'-NH-R_2'$ wherein R_1' comprises a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, and R_2' comprises an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, for the manufacture of pharmaceutical preparations to induce cytokine formation in human or mammal cells.
12. Use according to claim 11, wherein said reaction product is synthesized according to any one of claims 5 to 9 and has at least partly entered an Amadori-rearrangement and/or a Maillard-reaction, to treat and/or prevent hematological and/or immunological diseases and/or stimulate the immunosystem of humans and/or mammals.
13. Use according to claim 11 or 12, wherein R_1' is a radical of the D-form of a simple sugar, preferably selected from the group comprising glucose, xylose, galactose, rhamnose, fructose, mannose, 2-deoxy-glucose, 6-deoxy-glucose, glucosamine and galactosamine.
14. Use according to any one of claims 11 to 13, wherein R_2' is a radical of the L-form of an aminoacid or peptide,





preferably selected from the group comprising serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine.

15. Use according to any one of claims 11 to 14, wherein said reaction product is present together with a pharmaceutically acceptable carrier and/or adjuvant and/or optionally a lubricant in a weight ratio of reaction product to the remaining components of between 1:1 to 1:100, preferably 1:8 to 1:20 and most preferably about 1:9.

16. Use according to claim 15, wherein said reaction products are present in admixture with lactose and a lubricant, the weight ratio of lactose to the lubricant being between 20:1 and 100:1, preferably approx. 50:1.

17. Use according to any one of claims 11 to 16, wherein said reaction products are present in an amount of 0.01 - 10% by weight, preferably 0.01 - 1% by weight, especially 0.05 - 0.10% by weight, in a preparation adapted for cosmetic application.

This Sheet Has Been Replaced



~~SUBSTITUTE SHEET~~

THIS SHEET HAS BEEN
REPLACED

920227

- 31 -

LP2618EU

ABSTRACT

Novel Amadori reaction compounds have the formula R_1-NH-R_2 , wherein R_1 comprises the D-form of a 1-amino-1-deoxy-2-ketose radical derived from a sugar radical selected from the group of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine, and R_2 comprises the L-form of an aminoacid or peptide radical selected from the group of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine. Compounds and combinations of compounds having the general formula $R_1'-NH-R_2'$, wherein R_1' comprises a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and polysaccharides, and R_2' comprises an aminoacid or a peptide radical, are used to produce pharmaceutical preparations which in contact with human leukocytes produce interferon and other cytokines.

This Sheet Has Been Replaced



REPLACEMENT SHEET


02 OCT 1996

THIS SHEET HAS BEEN
REPLACED

- 1 -

Amadori reaction compounds and products, process for their manufacture, and their use

The present invention relates to novel Amadori reaction compounds and products, to the production thereof and to a new use of these compounds and products, having at least partly entered an Amadori rearrangement as per the following reaction scheme and/or a Maillard reaction:




Amadori reaction products are known; they are reaction products, for example, of an aminoacid or a peptide with a sugar, oligo- or polysaccharide having entered an Amadori rearrangement (J. Biol. Soc. 215 (1955), Henri Borsook et al.). Thus, in DE-C-3914354, a water-soluble glycoprotein of an aminoacid and a sugar is described which is isolated from an extract of *Avena sativa* seeds. Further, EP-A-406087 describes water-soluble polysaccharide-glycopeptide complexes which are derived from the cell wall of Gram positive bacterium, and J. Biol. Chem. 1985, 260/9 states that NMR-spectroscopy has been used to characterize Amadori reaction products formed by reaction of glucose with free amino groups of protein.

The invention now describes specific novel Amadori rearrangement compounds as described in claim 1. preferred embodiments and improvements are described in claims 2 to 4. These compounds

This Sheet Has Been Replaced





- 2 -

This Sheet Has Been Replaced



reduce potassium ferricyanide, a test reaction for biological active substances formed in a reaction of sugar and aminoacid.

The invention further relates to a novel use of such compounds and simultaneously a novel use of Amadori reaction products of sugars and aminoacids in general, as described in claims 11 to 17. In the past, nobody has tested the various steps of purification of the extract of an Amadori reaction product (in order to get rid of ballast substances and impurities) for biological activity so far.

Immunostimulating drugs are indeed known from natural sources such as mistletoe extracts, peat extracts etc. with the drawback of expensive treatment of large quantities of raw material to obtain a few grams of active substance; uncontrollable impurities that might lead to toxicity and side effects and therefore, further, to problems with administration during practical use due to the complex nature and the hardly reproducible composition.

Comparable products or product mixes from artificial sources, such as interferon or other genetic engineering methods are even more expensive to make. Furthermore, the molecules of human interferon are very often too big to penetrate the human cell wall so that only a fraction of the administered dose is effectively becoming active. Also, genetic engineering products usually have side effects and some are even toxic. Furthermore, some of them act effectively on one day and not on the next day, for reasons unknown so far.

Suprisingly, it has now been found that nearly any simple amino-acid/sugar complex after having at least partly entered an Amadori rearrangement does not show any of the above-mentioned disadvantages but on the contrary has a surprisingly high immunological activity. They can therefore be used in pharmaceutical formulations and in cosmetics. These small molecules easily penetrate the cell wall and virtually act as a nutrient. They

REPLACEMENT SHEET

02 OCT 1996

- 3 -


induce the formation of natural interferon and other cytokines, including tumor necrosis factor. Even three days after the administration, they still show this stimulating effect on biological activity. This effect increases with increasing completion of the Amadori rearrangement and decreases again with increasing decomposition of the complex.

Instead of simple sugars also - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides may be used, for instance dextrane, which react similarly. Polysaccharides show biological activity and may retain some of this activity after they become oligosaccharides.


Very little has been known heretofore about the biological activity of these compounds. It was now found that combinations of these substances in contact with human leukocytes produce interferon and other cytokines. This is called polyclonal activation of the cells.

It is possible to test the substances produced under the influence of these compounds and to determine the biological activities in international units relevant to specific cytokines. These compounds are of especially high biological activity within a range of pure substance concentrations from 1-100 µg/ml. Within the molecule, the specific nature of the aminoacid is more important than the nature of the sugar part of the molecule.

Reaction products of L-aspartic acid with glucose or galactose - after having gone through the Amadori rearrangement - when contacted with human leukocytes and incubated in tissue culture media at 37°C for 20 hours in an atmosphere of 5% CO₂ will produce from 30 - 1000 antiviral units of interferon. The interferon is measured in a bioassay using human cancer cells. Under the influence of these compounds, tumor necrosis compounds may also be produced.



THIS SHEET HAS BEEN
REPLACED



- 4 -

The products which allow such an unexpected use have the general formula



wherein


R_1' represents a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, and

R_2' represents an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical.


Thus the group of biologically active compounds may cover either the specific Amadori rearrangement compounds described above or the N-substituted derivatives of a number of different aminoacid compounds and one simple sugar, oligo- or - preferably low molecular weight, especially of less than 1000 daltons - polysaccharide, or N-substituted derivatives of one aminoacid compound and a number of simple sugars, oligo- and/or such polysaccharides, or else any combination of such derivatives, every single one of them having sufficient biological activity.

Preferably, R_1' in the above formula may be a radical selected from the D-form of simple sugars, especially (but not exclusively) from the D-form of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine; R_2' may be a radical selected from the L-form of aminoacid compounds such as serine, glycine, histidine, arginine, glutamine, asparagine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, methionine, hydroxyproline, tryptophane, proline, tyrosine, valine, isoleucine, leucine and lysine or else peptides of these aminoacids in any combination.

The invention further relates to a process for obtaining the above mentioned compounds and products as described in claim 5 and whereby an intermediate is formed of the formula



- 5 -





wherein R' is a C_1 -1-deoxy radical in a straight carbon chain or any O-bridged form of a simple sugar or an oligo- or polysaccharide, and

R'' represents an aminoacid or a peptide radical, said intermediate being at least partly subjected to an Amadori rearrangement and/or to the Maillard reaction by continued heating of the reaction mixture - preferably under pressure - and simultaneously or subsequently removing the solvents. Preferred embodiments of the process are described in claims 6 to 10.

In some cases, especially when an aminoacid having two carboxyl groups is used, it is advantageous to add to the process a buffer salt, such as sodium bicarbonate, preferably in a molar ratio of 1:1.

It was further found that the Amadori rearrangement products are relatively susceptible to decomposition, and that decomposition products have the nature of dark brown and tar-like, unidentified compounds having lost their biological activity. Therefore it is preferred to stop the Amadori rearrangement reaction at a stage at which the reaction mixture becomes light orange-brown in colour.

It is interesting to note that the intermediate reaction products, formed when the originally opaque solution of the aminoacid becomes clear (before the Amadori rearrangement), are easily hydrolysed, i.e. the reaction is reversible. With increasing rearrangement, the reversibility diminishes, i.e. the products become more stable, and the colour gradually changes from light yellow to light orange, and then finally to orange-brown when the Amadori rearrangement seems to be complete. Samples taken during such rearrangement reaction and tested (according to various procedures described later) proved that the biological activity increases with the Amadori reaction progressing, and decreases when further heating results in decomposition, for which a colour change to dark brown is a sign. Immediate re-



- 6 -


duction of ferricyanide and the resulting colour change will occur if the reaction mixture contains other keto groups and/or sulfur containing aminoacids such as cysteine; otherwise it will occur within 3 to 5 min, which is a good check for the degree of Amadori rearrangement developed. Unreacted sugars would show the colour change after half an hour or several hours only.

Isolation of the pure Amadori rearrangement products is carried out according to known methods based on binding the mixture on a strong cation-exchanger (such as Amberlite^(R) or Dowex^(R)), successively eluting with ammonia water, evaporating a chosen defined fraction of the eluate under reduced pressure and crystallising the pure compound from anhydrous methanol (J.E. Hodge and B.E. Fisher, Methods in Carbohydrate Chemistry, Vol.II, Reactions of Carbohydrates, 1963 Academic Press, N.Y., London, 1963, page 105-106; or Borsook et al. as quoted earlier; or J. Dubourg and P. Devilliers as quoted earlier).


In the process according to the present invention, all the above-mentioned preferences regarding the kind of radicals derived from simple sugar and aminoacid compounds remain unchanged. Additionally, a preferred mixture of sugar substrates comprises the D-forms of glucose, xylose, galactose, rhamnose and fructose in a weight ratio of about 20:10:4:1:1, while the preferred mixture of aminoacid substrates comprises the L-forms of serine, glycine, histidine, arginine, alanine, proline, tyrosine, valine, leucine, isoleucine and lysine in a weight ratio of 20.4:35.8:35.8:132:180:360:216:160:72:68:780.

As already mentioned, the Amadori rearrangement products are able to reduce potassium ferricyanide, such a chemical test reaction providing a basis for quick determination of the biological activity of the composition formed in a reaction of sugar and aminoacid.

It has been found that Amadori reaction products are especially active if a mixture of simple sugars of the same composition and



- 7 -




in the same weight ratio as occurring in natural peat extracts are reacted with a mixture of aminoacid compounds of the same composition and in the same weight ratio as occurring in natural peat extracts in the presence of an aqueous solvent, preferably adding a lower alcohol - and optionally inorganic trace elements occurring in such natural peat extracts - at elevated temperature (and optionally under pressure), and subsequently prolonging the heating in order to cause an Amadori rearrangement of the obtained products, simultaneously or subsequently expelling the solvents, stopping the rearrangement reaction at the point when the reaction mixture becomes light orange-brown in colour, drying the products thus obtained and purifying the same by means of column chromatography and collecting the fractions that cause maximum reduction of potassium ferricyanide.

In one embodiment, the instant pharmaceutical formulations contain as an active ingredient at least one reaction product of the formula $R_1'-NH-R_2'$ or a specific compound of the formula R_1-NH-R_2 together with a pharmaceutically acceptable carrier and/or an adjuvant and/or optionally a lubricant in a weight ratio of active ingredient to the remaining components of between 1:1 and 1:100, preferably 1:8 to 1:20 and most preferably about 1:9.

Another advantageous pharmaceutical formulation contains - in addition to the active ingredient - lactose and a lubricant, the weight ratio of lactose to the lubricant being between 20:1 and 100:1, preferably 50:1.

These pharmaceutical preparations are used to treat and/or prevent hematological and/or immunological diseases and/or stimulate the immunosystem of humans and/or mammals by the induction of cytokine formation.

Another use for the active ingredients is in cosmetic preparations. The active ingredient is present in such preparations in amounts of 0,01-10% by weight, preferably 0,01-1% by weight and especially in amounts of 0,05-0,1%. These cosmetic preparat-



- 8 -

ions contain - besides the active ingredient - usual carriers, adjuvants, enriching components and/or fragrances.

The present invention will be further explained and demonstrated in the following examples, which do not limit in any respect the scope of the present invention.

Example 1:

The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with:

- 1.47 g (0.01 M) L-glutamic acid
- 0.84 g (0.01 M) NaHCO_3
- 0.91 g D-glucose
- 0.91 g galactose and
- 3.00 ml of redistilled water


The mixture was heated up to 80°C whereby - preferably under reduced pressure - 50% of the water were evaporated and then - under atmospheric pressure - heated to a temperature of 80-85°C, at which it was kept for 120 min until it became orange in colour. The syrup-like concentrated aqueous solution was then evaporated to dryness under reduced pressure. The orange-red solid reaction product thus obtained was marked with the symbol D-10 and saved for biological tests.

Example 2:

The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with:

- 1.33 g (0.01 M) L-aspartic acid
- 0.84 g (0.01 M) NaHCO_3
- 0.91 g of D-glucose
- 0.91 g of galactose and
- 3.00 ml of redistilled water

The mixture was heated up to 80°C with stirring (by means of rotation), concentrated under pressure, whereby a total volume of 1.5 ml water was vaporized, and then treated under atmospheric pressure at a temperature of 80-85°C until it became light orange in colour; this took place within approx. 60 min.



- 9 -

The reaction mixture was a syrup-like concentrated aqueous solution and was dried under reduced pressure. The resulting reaction product was a dry, yellow-orange powder. It was marked with the symbol D-11 and saved for biological tests.

Example 3:

The 25 ml flask of a rotary evaporator placed in a heated water bath was charged with:

- 1.05 g (0.01 M) L-serine
- 0.91 g D-glucose
- 0.91 g galactose and
- 2.50 ml redistilled water


The mixture was heated to 85-92°C with stirring (by means of rotation). After 100 min, the solution became orange in colour. The pressure was reduced and the mixture was evaporated to dryness. The reaction product on the walls of the flask formed an orange transparent layer. The reaction product was scraped off and powdered. It was marked with the symbol D-12 and saved for biological tests.

Example 4:

The 25 ml flask of a rotary evaporator placed in a heated waterbath was charged with:

- 0.66 g (0.005 M) of D-aspartic acid
- 0.42 g (0.005 M) of NaHCO_3
- 0.45 g of D-glucose
- 0.45 g of galactose and
- 3.00 ml of redistilled water

The mixture was heated up to 80°C, evaporated - under pressure - whereby a volume of 1.5 ml water was vaporized, and then treated - under atmospheric pressure - at a temperature of 85°C. After 60 min. heating the mixture became orange in colour; the pressure was reduced and the resulting syrup-like concentrated aqueous solution evaporated to dryness. Before the residue became definitely dry, two times 10 ml water-free ethanol were introduced into the flask and evaporated in order to eliminate residual moisture. The dry reaction product thus obtained was



- 10 -


powdered, marked with the symbol D-13 and saved for biological tests.

Example 5:

In order to obtain - in a synthetic way - an equivalent of the biologically active fraction of a certain natural peat extract, the flask of the rotary evaporator placed in the heated water bath was charged with:

20.5 mg	L-serine
35.8 mg	L-glycine
35.8 mg	L-histidine
132.0 mg	L-arginine
180.0 mg	L-alanine
360.0 mg	L-proline
216.0 mg	L-tyrosine
160.0 mg	L-valine
68.0 mg	L-isoleucine
72.0 mg	L-leucine
780.0 mg	L-lysine
2000.0 mg	D-glucose
1000.0 mg	D-xylose
400.0 mg	D-galactose
100.0 mg	D-rhamnose
100.0 mg	D-fructose
6.0 ml	redistilled water

The mixture was stirred by means of rotation and heated under pressure for 45 min at a temperature rising from 75°C to 86°C. During that period, approx. 3 ml of water were evaporated and the substrates were totally dissolved. The mixture was then treated for 30 min under atmospheric pressure at a temperature of 85-86°C for an Amadori rearrangement. During that period the solution quickly became red-brown in colour. The pressure was reduced, and heating at 84°C was continued, thus simultaneously evaporating the solvents. At the end of evaporation, two times 15 ml water-free ethanol were introduced and the reaction mixture was brought to dryness. The flask with the dried reaction product was placed in a desiccator over calcium chloride for 18



- 11 -

hours; then the reaction product was powdered. Approx. 4.5g of a powdered product were obtained and marked with the symbol EK₂-S.

A portion of 4g of this reaction product was dissolved in 20 ml of distilled water and placed on a chromatographic column of 25 mm x 330 mm size, filled with a sorbent Amberlite^(R) XAD-2, analytical grade. The column was eluted with 0.4 ml/min distilled water. Fractions of 10 ml volume were collected to a total volume of 450 ml. The content of the fractions was monitored chromatographically. Fractions of consecutive numbers 11-13 were combined and evaporated under reduced pressure. These fractions were characterised by a high content of Amadori rearrangement products (confirmed with the potassium ferricyanide reduction test). The product was saved for biological tests under the symbol of EK₂-S-11.

Biological tests for determining the biological activity were carried out with immunised Balb/C mice of both sexes, at the age of 8-10 weeks. Immunisation of mice is achieved by peritoneal administration of 0.2 ml of a 10% suspension of sheep erythrocytes (SRBC), i.e. of 6×10^8 cells. The erythrocytes are fixed in a sterile Alsever's solution of the following composition:

glucose	2.05 g
sodium citrate	0.8 g
sodium chloride	0.42 g
citric acid	0.055 g
redistilled water to	100 ml

Into such Alsever's solution, a sheep blood cell aseptic sample is introduced in a ratio of 1:1 and the mixture is kept for at least 3 days at +4°C. The thus stabilised erythrocytes are then sampled aseptically and introduced into a phosphate buffered salt solution (PBS) in order to wash them out. Erythrocytes are rinsed with PBS twice, and are centrifugated for 10 min at 2000 rpm. The washed out cells are used in the form of a 10% suspension in PBS. Such a suspension is used for the immunisation of Balb/C mice.

REPLACEMENT SHEET

02 OCT 1996

- 12 -


The reaction product to be tested was administered intraperitoneally (i.p.) or orally (p.o.) four times at chosen doses, the first administration taking place 2 hours before immunisation of the mouse with SRBC, while the remaining three dosages were administered after immunisation at 24 h intervals.

Each tested group of animals was treated with different doses of the tested reaction product: 10 mg/kg, 1 mg/kg, 0.1 mg/kg and 0.01 mg/kg. A control group of animals was also immunised with SRBC, but instead of the substance to be tested, 0.2 ml of PBS were administered at the same time intervals.

Each group of animals, control and tested groups, in all experiments consisted of 8-12 mice.

On the fourth or (in case of determination of antibodies type 7S) tenth day after immunisation, mice were slightly anesthetized with ether and exsanguinated by eyeball extirpation. The blood was collected into test tubes. Next, the spinal cord was broken and spleen removed. The blood was used for obtaining the serum needed for determination of hemagglutinating antibodies of the 19S+7S and 7S types, while spleens were used to prepare the cells useful for determination of the percentage of cells able to form E-rosettes and of hemolytic activity. For such uses, mouse spleens were comminuted. The splenocyte cells obtained were suspended in approx. 2 ml of Hanks' medium at +4°C, layered on the Ficoll-Uroplon gradient of a density of 1.077, and then centrifugated for 15 min at 3000 rpm at +4°C. After separation from the interphase, the lymphocyte buffy coat was placed in the Hanks' medium at +4°C and rinsed twice with centrifugation for 7-10 min each time at 1800 rpm. The splenocytes were then suspended in 1 ml of Hanks' medium at such a ratio that it contained 1×10^6 cells.

For each test, the percentage of dead cells is determined by mixing a drop of a tested suspension of splenocytes with a drop



- 13 -

of ex tempore prepared dyestuff solution containing 4 parts of a 0.2% trypan blue solution and 1 part of a 4.25% NaCl solution. Under the microscope, the percentage of dead splenocytes is determined for each 100 cells. Dead cells are navy blue, while the bright cells are live cells. The presence of more than 10% of dead cells is critical; such a sample has to be eliminated from further use.

All steps carried out with the cells to be tested should be performed in a sterile, siliconized laboratory glass apparatus placed in an ice bath.

Example 6:

In the first test, an effect of the tested reaction products on the number of cells producing hemolytic antibodies (PFC-IgM) was determined. The test was carried out as follows: To 0.5 ml of an 0.5% agarose solution placed in a test tube and kept in a heated water bath at 45°C, 0.1 ml of a 10% suspension of SRBC (prepared as described above) were admixed. Then 0.1 ml of a splenocyte suspension having a density of 1×10^6 cells/ml was added, the mixture stirred rapidly and immediately poured out on slides previously covered with agarose. The slides are incubated at 37°C for two hours. Next, the tested samples are covered with guinea pig complement diluted at a ratio of 1:20 for a further 2h. After the incubation of the tested samples with the complement, the number of plaque forming cells (PFC) was counted and recalculated for 1×10^6 splenocytes. Each test was performed twice.

The strongest amplification of the response to SRBC expressed in terms of increase of the number of splenocytes producing hemolysins IgM (PFC) was observed after administration of the D-11 substance at a dose of 0.1 mg/kg. The amplification was 119%. When the daily dose was increased ten times up to 1 mg/kg, amplification of the response was lowered to 53%.

REPLACEMENT SHEET

02 OCT 1996

- 14 -

Reaction product D-12 showed the strongest activity - an increase of 58% - at a dose of 1 mg/kg.

Reaction product EK₂-S-11 in this test showed the strongest activity at a dose of 0.1 mg/kg (increase of 65%). At a dose ten times higher, i.e. 1 mg/kg, the increase was slightly lower, i.e. 52%.


Reaction product D-13 tested at a dose of 1 mg/kg caused an increase of 40%. When the dose was increased to 10 mg/kg, i.e. ten times, the response was only a 14% increase.

Example 7:

An active hemagglutination test was also carried out, determining antibodies of the anti-SRBC type 19S+7S and 7S antibodies. In order to determine the 19S - IgM type antibodies level, mouse serum was prepared on the fourth day after immunisation with SRBC, while for the determination of 7S-IgG type antibodies level such preparation took place on the tenth day after mice immunisation with SRBC, which is related to the day of maximum count of antibodies of a given type in mice immunised with SRBC.

A. Determination of 19S+7S antibodies count.

A sample of blood was centrifuged for 30 min at 3500 rpm. From each sample thus prepared, serum was collected and placed in a heated water bath at a temperature of 56°C for 30 min in order to deactivate the complement. Next, a number of solutions at several different dilutions of each tested serum was prepared (from 1:1 to 1:4096) employing a microtitrator and U-shaped microplates having a volume of 250 µl each. The diluted sera were incubated for 1 hour at room temperature. A drop of a 1% suspension of SRBC in PBS (prepared as described above) was added to each serum; the mixtures were incubated for a further 2 h at a temperature of 37°C and then stored at a temperature of +4°C. The results were taken next day. The maximum dilution at which hemagglutination is still caused was considered the anti-



- 15 -

body count. A ring at the bottom of the plate is a sign that hemagglutination occurs. A button-like formation at the bottom of the plate is considered as a negative result - no hemagglutination.

For statistical analysis of the results, the increase of dilution of serum in a tested substance was compared to the one in a control group.

Reaction product D-11 at a dose of 1 mg/kg increased the IgM count 2.57 times. At a ten times higher dose, the stimulation effect in comparison to the control group increased by 3.5 times.

Reaction product D-12 in this test showed a weaker activity. At a dose of 0.1 mg/kg, it increased the IgM count 2 times, at a dose of 1 mg/kg 1.4 times.


Reaction product EK₂-S-11 showed the strongest activity in this test. At a dose of 0.1 mg/kg, it increased the IgM count 4.3 times, at a dose of 1 mg/kg 3.6 times.

Reaction product D-13 at a dose of 1 mg/kg increased the IgM count 3 times, at a ten times higher dose 1.5 times.

B. Determination of 7S antibody count

The tested inactivated sera were combined - at a ratio of 1:1 - with a 0.1 M solution of 2-mercaptoethanol and the mixtures incubated for 30 min at a temperature of 37°C. 2-Mercaptoethanol destroys immunoglobulines of the 19S-(IgM) type, while immunoglobulines of the 7S-(IgG) type are not susceptible to the action of 2-mercaptoethanol.

After 30 min of incubation the reduction reaction was stopped by means of cooling down to a temperature of +4°C for 15 min. Next, a number of dilutions was prepared in a similar manner as de-



- 16 -

scribed above with respect to determining the 19S-antibody count and combined with a 1% suspension of SRBC; after 2 hours of incubation at 37°C, the samples were stored at a temperature of +4°C. The results were evaluated on the following day, according to the criteria of determining the hemagglutination count described above. Simultaneously, a control test was carried out with a combination of a 1% suspension of SRBC with PBS in a ratio of 1:1.

When the substance D-11 was tested as described above, at a dose of 1 mg/kg it increased the production of antibodies IgG 3.16 times. At a dose of 10 mg/kg, the increase was 2.2 times.

Reaction product D-12 tested at a dose of 0.1 mg/kg and 1 mg/kg respectively stimulated the production of antibodies IgG 1.3 times, at a dose of 10 mg/kg 1.5 times.


Reaction product EK₂-S-11 at a dose of 0.1 mg/kg stimulates the production of IgG 1.9 times, at a dose of 1 mg/kg 2.89 times (in comparison with the control).

The results of tests A and B obtained for each production lot or for each fraction of the biologically active reaction products synthesized according to the invention and giving the above-mentioned immunological response were subjected to statistical analysis by the T-student method, $\alpha=0.05$. Results obtained for each dose tested were compared with a parallel control test and showed an increase of biological activity.

Example 8:

The group of biologically active reaction products obtained according to Examples 1 to 5 have also been submitted to the test in which the percentage of splenocytes forming E-rosettes was determined.

250 μ l of a 1% suspension of SRBC and 250 μ l of cells to be tested at a concentration of 1×10^6 cells/ml were added to 550



- 17 -

μl of Hanks' medium. Each such sample was incubated ~~in~~ a heated water bath equipped with a shaker for 15 min at a temperature of 37°C. Then, it was stored at a temperature of +4°C for a further 20 h. The percentage of splenocytes forming E-rosettes with SRBC was determined after the suspension was coloured with 1 to 3 drops of crystal violet.

Each sample was subjected to determination of the percentage of splenocytes three times, counting at each instance 400 splenocytes. A splenocyte surrounded with at least 3 erythrocytes was considered an E-rosette.

For statistical evaluation, the percentage increase of the number of splenocytes with E-rosettes was compared between the substances to be tested and a control group.

In this test, the strongest stimulating effect was shown by the reaction products D-11 (63%) and EK₂-S-11 (70%) at a dose of 1 mg/kg. At a dose ten times smaller, i.e. 0.1 mg/kg, the values decreased to 45% and 57% respectively.

Reaction product D-12 at a dose of 1 mg/kg caused an increase of the ability to form E-rosettes of 22% in comparison to the control group. The respective value for a dose ten times smaller, i.e. of 0.1 mg/kg, was 29%.

Reaction product D-13 shows the maximum effect at a dose of 1 mg/kg (58%), while at higher doses the effect is slightly smaller.

Biological activity of synthetised compounds was evaluated according to the following tests:

1. Test for determination of the percentage of splenocytes forming E-rosettes, carried out according to Bach and Dardenne (Cell. Immunol. 3, 1-16, 1972)
2. Test for determination of the number of cells producing hemolytic antibodies of an IgM type, carried out according

REPLACEMENT SHEET

02 OCT 1996

- 18 -

to the Jerne method, modified by Mishell and Dutton (J. Exp. Med. 126, 423-442, 1967) and

3. Test for determination of a hemagglutination 19S + 7S and 7S antibody count, carried out according to Adler's active hemagglutination methods (J. Immunol. 95, 26-38, 39-47, 1965) with the use of microplates (J. Immunopharmacol. 4, 43-52, 1982).

Example 9:


A rotary evaporator flask placed in a heated water bath was

charged with:	1.33 g (0.01 M)	L-aspartic acid
	0.84 g (0.01 M)	NaHCO ₃
	10.00 g	hydrolysed dextrane of an
		average molecular weight of
		3000 daltons
	10.00 ml	redistilled water.

The mixture was heated under pressure at a temperature of 70°C until the solid substances were completely dissolved, expelling by means of distillation during that period approx. 3 ml of water (heating time was approx. 30min). The flask with the solution was loosely covered, placed in a steam steriliser and heated under pressure at a temperature of 121°C for 40 min. After cooling down, the resulting yellow-orange solution was diluted with 15 ml of water, clarified by means of centrifugation and spray dried by air having an inlet temperature of +160°C and an outlet temperature of +85°C. 10.5 g of a light beige reaction product resulted that was easily soluble in water.

The presence of Amadori rearrangement products in this reaction product was confirmed by a test by the potassium ferricyanide method described by Borsook, Abrams and Lowy, J. Biol. Chem 215, (1955), 111-124 and by chromatographic methods.

Biological tests as described in the preceding Examples confirmed an immunotropic activity of the above product similar to the one exhibited by preparations obtained with simple sugars.



- 19 -

Example 10:

A conical flask was charged with:

- 5.0 g hydrolysed dextrane of an average molecular weight of approx. 5000 daltons
- 1.1 g L-proline
- 4.0 ml redistilled water.


The content was dissolved by means of stirring. The uniform mixture thus obtained was placed in a steam steriliser and heated under pressure at a temperature of 110°C for 40 min. The resulting transparent orange solution was diluted with 20 ml of redistilled water and clarified by means of centrifugation. The clear solution was spray dried.

5.3 g of a reaction product easily soluble in water was obtained. Immunotropic activity was similar to the one observed in other experiments according to the preceding examples.

Example 11:

Conventional methods test the biological activity of the compounds in mice, but not in humans. For this reason, new bioassays have been introduced, which measure the amounts and activity of cytokines released from the human peripheral blood leukocytes (PBL) treated with the reaction products according to the Examples 1 to 5, 9 and 10. The cytokines are the hormone-like proteins that play an important role in practically all of the immunological reactions, as well as in the regulatory processes responsible for the maintenance of homeostasis.

Cytotoxicity assays. Cytotoxicity of the reaction products was determined in human lung adenocarcinoma cell line A549 (included in the American Type Culture Collection - ATCC CCL 185). The cell monolayers were trypsinized, suspensions of 2×10^5 cells/ml in Dulbecco's-modified Eagle's minimum essential medium (DMEM) plus 10% calf serum (CS) were mixed with various doses of the drugs, seeded in the plastic microplates, and incubated for 48 h




- 20 -

at 37°C. CD₅₀ was the minimal concentration of the compound which caused approximately 50% destruction of the cell culture, as measured by staining with 0.015% solution of neutral red in DMEM.

Cytokine induction. Buffy coats from healthy blood donors were obtained from the regional transfusion center. The erythrocytes were lysed by NH₄Cl treatment according to Cantell et al. (Cantell, K., Hirvonen, S., Kauppinen, H.L.: Production and Partial Purification of Human Immune Interferon. Meth. Enzymol., 119, 54, 1986). The leukocytes from a single donor containing approximately 8x10⁶ leukocytes/ml in RPMI 1640 medium supplemented with 10% fetal calf serum (FCS), L-glutamine, and antibiotics were used. All lots of FCS were pretested. Only non-mitogenic FCS for PBL cultures was used. The cytokine inducers were added to 1 ml volumes of the cultures. The reference cytokine inducers were phytohemagglutinin (PHA) (Pharmacia Fine Chemicals, Sweden) and LPS from E. coli 0111:B4 (Difco Laboratories). The induced cultures of PBL were incubated in an atmosphere of 5% CO₂ in air at 37°C for 20 h and centrifuged. Supernatants were stored at 4°C and assayed for TNF and IFN activity within one week.

Interferon (IFN) assay. The confluent monolayer of A549 cells was prepared in the microplates in DMEM with 10% CS, L-glutamine, and antibiotics (100 units/ml penicillin and 100 µg/ml streptomycin). IFN samples diluted in plates were added to the cell monolayer and incubated at 37°C for 20 h in 5% CO₂ in air. The cells were then washed and challenged with encephalomyocarditis virus (EMCV). The titer of IFN was defined as the dilution of IFN sample that reduced the virus cytopathogenic effect by 50% after 48 h of incubation. The MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) method (Hansen, M.B., Nielsen, S.E., and Berg, K.: Re-examination and Further Development of a Precise and Rapid Dye Method for Measuring Cell Growth/ Cell Kill. J. Immuno. Meth., 1989, 119, 203-210) to measure the cell kill in the ELISA scanner was also



- 21 -

used. Laboratory standards of IFNs were included in all assays. Recombinant human IFN- γ (Genentech Inc., USA, specific activity 2×10^8 units/mg), the natural human leukocyte IFN- α (3×10^6 IU/ml) and IFN- γ (2×10^6 IU/ml) obtained from Dr. K. Cantell, Helsinki, Finland.

Tumor Necrosis Factor (TNF) assay. The cytotoxic activity of TNF was measured in L929 cells according to Flick and Gifford (Flick, D.A., Gifford, G.E.: Comparison of in Vitro Cell Cytotoxic Assays for Tumor Necrosis Factor. J. Immunol. Meth., 68, 167, 1984).

The samples and actinomycin D solution were added to monolayer cultures of the cells. After incubation at 37°C for 20 h, the cultures were stained with crystal violet and toxic effects were determined. The amount causing approximately 50% destruction of the cell cultures was defined as one unit of TNF activity. Comparison with a preparation of TNF- α (Genentech Inc., USA) showed that 1 unit in our assays was equal to 100 - 200 pg/ml TNF.

Cytokine neutralization assays. The antisera used were: rabbit anti-human TNF- α , lot 2958-40 and rabbit anti-human IFN- γ , lot 2891-56 (Genentech Inc., USA), sheep anti-human IFN- α, β , and sheep anti-human IFN- γ (obtained from Dr. K. Cantell, Finland). The cytokine preparations were treated with the sera diluted 1:200 in culture medium and incubated for one hour at room temperature. Then, the residual IFN or TNF activities were determined as described.

Five different batches (L_1 to L_5) of PBL prepared from the blood of healthy blood donors were used. The optimal PBL concentration for the assays was found to be 8×10^6 cells per one ml of medium (RPMI 1640 supplemented with 10% fetal calf serum and antibiotics).

Incubation of human PBL with the new reaction products I - XI (Table 1) resulted in IFN and TNF synthesis. The observed re-

02 OCT 1996

sponses were dose related in the range of 3 - 100 µg/ml of the compounds (Table 2). The compounds used in the indicated concentrations were non-cytotoxic. In all of the bioassays, the negative and positive controls were included. The negative controls measured the amounts of the cytokines (IFN and TNF) produced spontaneously without the addition of any exogenous stimulants. The positive controls indicated the amounts of the cytokine produced in response to a known reference inducer; in our case this was phytohemagglutinin (PHA, Pharmacia, Sweden, 10µg per ml).

It should be pointed out that the cytokine induction in human PBL cultures obtained from different individuals usually gives considerable variation of the results. The phenomenon may be explained in terms of genetic differentiation of human population. Furthermore, PBL cultures often produce IFN and TNF spontaneously.

In other words, high responders and low responders or even non-responders are commonly observed among the healthy donors of PBL.

Despite the presented limitations, the results of the bioassays showed that PBL (L_1 to L_5) treated with the reaction products (I - XI) produced IFN and/or TNF that could be measured quantitatively.

In the case of L_1 which contained leukocytes of the high responder, the reaction product II (containing L-form of aspartic acid) was found to be considerably more active as a cytokine inducer than the reaction product III (containing the D-form of aspartic acid which also is more expensive by two orders of magnitude). The observation is significant because mainly L-forms of the amino-acids are recognized by cells as natural substrates in biochemical reactions.

02 OCT 1996

- 23 -

Furthermore, for the expression of biological activity of the reaction products, the amino-acid part of the molecule is much more important than the form of sugar. Instead of single sugars, - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides (such as dextrans, reaction products X - XI) can be used and they react similarly.

And vice versa, polysaccharides containing the amino-acid residues may have biological activity, and this activity is retained when they are decomposed to oligosaccharides with the bound amino-acid (data not shown).

Similar results may also be observed if a short peptide is taken instead of a single amino-acid and is used to stimulate the leukocytes to produce cytokines (data not shown).

Seven reference batches of the unfractionated TTP assayed in over 100 PBL cultures from different donors produced from 10 to 1.000 units of IFN per ml, and from 9 to 750 units of TNF per ml. The fraction EK₂-S prepared from a mixture corresponding to the contents of natural peat extract (Example 5) has been assayed in eight PBL cultures from eight different blood donors. It was found to be the most active preparation in inducing both IFN and TNF (data not shown).

Possible applications of the reaction products are as immuno-modulators and such activity was clinically useful. Tissue regeneration is another proven activity. Anti-cancer activity probably connected with the presence of the induced interferon and tumor necrosis factor was also observed. Anti-viral activity was also noted.

The main use of the above reaction products involves oral administration, but parenteral treatment is also possible, as is topical application. The products appear relatively stable.

02 OCT 1996

Table 1. List of Reaction Products

No.	Substrates
I (D-10)	L-glutamic acid, glucose, galactose
II (D-11)	L-aspartic acid, glucose, galactose
III (D-13)	D-aspartic acid, glucose, galactose
IV (D-12)	L-serine, glucose, galactose
V	EK ₂ -S (fractions 11 - 13)
VI	EK ₂ -S (fractions 6 - 7)
VII	EK ₂ -S (fractions 8 - 10)
VIII	EK ₂ -S (fractions 28 - 34)
IX	L-proline, glucose
X	L-aspartic acid + dextrane (variety 1)
XI	L-aspartic acid + dextrane (variety 2)

Example 12:

Pharmaceutical formulations containing as an active ingredient the reaction products according to Examples 1 to 5, 9 and 10, were prepared using the following reaction products:

A. Tablets/Granules:

5.0 g of the reaction product obtained according to Example 1 or 10 (active substance),
444.0 g of pharmaceutically acceptable lactose
1.0 g of lubricant, e.g. MYVATEX^(R), trademark of Eastman Kodak)

The ingredients were mixed and granulated with 30% aqueous ethanol in a conventional way, then dried at 40°C. The granules were used to prepare capsules, each containing approx. 450 mg of granules, i.e. 5 mg of the active substance. Alternatively, the granules were used to form tablets, each weighing approx. 450 mg and containing 5 mg of the active substance.

B. In the same conventional manner, the active substances obtained according to the preceding examples 1 to 5, 9 and 10 were formulated into other pharmaceutical formulations using suitable carriers.


REPLACEMENT SHEET

02 OCT 1996

- 25 -

Table 2. Biological Activity of the Reaction Products I-XI in Human PBL


	Dose µg/ml	IFN units/ml					TFN units/ml				
		L ₁	L ₂	L ₃	L ₄	L ₅	L ₁	L ₂	L ₃	L ₄	L ₅
Control	-	10	<10	<10	20	<10	80	9	27	27	<9
PHA	10	100	30	30	60	100	250	80	250	250	250
I	100	100	<10	<10	20	-	250	9	9	160	-
	30	-	-	-	30	-	-	-	-	500	-
	10	30	<10	<10	30	-	80	9	18	160	-
	3	-	-	-	10	-	-	-	-	160	-
II	100	100	10	<10	10	<10	250	18	18	250	<9
	30	-	-	-	10	<10	-	-	-	250	<9
	10	1000	10	30	10	10	250	50	27	250	<9
	3	-	-	-	10	<10	-	-	-	160	<9
III	100	<10	<10	10	10	<10	250	9	18	250	<9
	30	-	-	-	30	<10	-	-	-	250	<9
	10	<10	<10	<10	10	<10	80	27	18	250	<9
	3	-	-	-	10	<10	-	-	-	80	<9
IV	100	<10	10	10	20	<10	80	60	9	160	<9
	30	-	-	-	20	<10	-	-	-	27	<9
	10	<10	30	<10	20	<10	80	50	18	80	<9
	3	-	-	-	20	<10	-	-	-	80	<9
V	100	30	<10	<10	60	-	80	50	18	250	-
	30	-	-	-	60	-	-	-	-	80	-
	10	<10	<10	<10	10	-	80	27	9	80	-
	3	-	-	-	10	-	-	-	-	80	-
VI	100	<10	<10	<10	20	-	80	27	18	80	-
	30	-	-	-	10	-	-	-	-	80	-
	10	10	<10	<10	20	-	50	27	18	160	-
	3	-	-	-	10	-	-	-	-	250	-
VII	100	<10	<10	<10	-	-	80	27	27	-	-
	30	-	-	-	-	-	-	-	-	-	-
	10	10	<10	<10	-	-	80	27	27	-	-
	3	-	-	-	-	-	-	-	-	-	-
VIII	100	<10	<10	<10	-	-	80	160	27	-	-
	30	-	-	-	-	-	-	-	-	-	-
	10	<10	<10	<10	-	-	750	80	27	-	-
	3	-	-	-	-	-	-	-	-	-	-
IX	100	10	<10	<10	10	-	27	27	18	80	-
	30	-	-	-	20	-	-	-	-	80	-
	10	100	30	<10	20	-	80	27	18	80	-
	3	-	-	-	10	-	-	-	-	50	-
X	100	100	<10	20	20	-	27	27	18	250	-
	30	-	-	-	30	-	-	-	-	80	-
	10	<10	10	<10	30	-	18	50	27	50	-
	3	-	-	-	30	-	-	-	-	250	-
XI	100	<10	10	<10	30	-	18	27	27	50	-
	30	-	-	-	10	-	-	-	-	250	-
	10	<10	30	<10	10	-	18	80	80	80	-
	3	-	-	-	20	-	-	-	-	750	-



- 26 -

Example 13:

The active substances obtained according to preceding Examples 1 to 5, 9 and 10 were used as a beneficial additive to cosmetic preparations intended for everyday hair and body care, the content of the substances being within a range of 0.01 - 10% by weight, depending on the type of the preparation, the method of application and the frequency of use intended for the particular preparation.



THIS SHEET HAS BEEN
REPLACED

- 27 -



C L A I M S

1. An Amadori rearrangement compound of the formula



wherein

- R_1 comprises a 1-amino-1-deoxy-2-ketose derived from a sugar radical, preferably in its D-form, selected from the group consisting of glucose, xylose, galactose, rhamnose, fructose, mannose, 2-deoxy-glucose, 6-deoxy-glucose, glucosamine and galactosamine, and
- R_2 comprises an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, preferably its L-form, selected from the group consisting of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine.
2. A compound according to claim 1, wherein R_1 comprises a sugar radical selected from the group of D-glucose and D-galactose.
3. A compound according to claim 1 or 2, wherein R_2 comprises an aminoacid radical selected from the group of L-glutamic acid, L-aspartic acid and L-serine.
4. A mixture of compounds according to claim 1.
5. Process for the manufacture of at least one reaction product of the formula $R_1'-NH-R_2'$, wherein R_1' comprises a 1-amino-1-deoxy-2-ketose derived from a simple sugar, an oligo- or a - preferably low molecular weight, especially of less than 1000 daltons - polysaccharide and R_2' comprises an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, wherein at least one substance selected from the group of aminoacids and -

This Sheet Has Been Replaced




REPLACEMENT SHEET

02 OCT 1996




- 28 -

This Sheet Has Been Replaced



preferably low molecular weight, especially of less than 1000 daltons - peptides, and at least one second substance selected from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, are reacted with each other at elevated temperature - optionally under pressure - and/or in the presence of a lower alcohol, whereafter the intermediate(s) is (are) subjected to an Amadori rearrangement and/or to the Maillard reaction.

6. Process according to claim 5, wherein the reaction is carried out in the presence of an aminoacid having two carboxyl groups and a buffer salt, preferably sodium bicarbonate in a molar ratio of 1:1.
7. Process according to claim 5 or 6, wherein at least one simple sugar is reacted with at least one monobasic aminoacid, and the molar ratio of the simple sugar(s) to the aminoacid(s) is between 2:1 and 1:1, preferably 1.5:1.
8. Process according to any one of claims 5 to 7, wherein the reaction of the sugar(s) and/or saccharide(s) with the aminoacid(s) and/or peptide(s) is carried out in a concentrated aqueous solution at a temperature of 75-90°C, and subsequent Amadori rearrangement is carried out at the same temperature - optionally under pressure, with simultaneous or subsequent removal of the solvents - until the reaction mixture becomes light orange-brown in colour.
9. Process according to any one of claims 5 to 8, wherein a mixture of simple sugars of the same composition and in the same weight ratio as occurring in natural peat extract is combined with a mixture of aminoacid compounds of the same composition and in the same weight ratio as occurring in said extract - and optionally with inorganic trace elements occurring with said extract - the resulting substrate mixture being reacted in the presence of water as a solvent at elevated temperature - and optionally under pressure




- 29 -

This Sheet Has Been Replaced



and/or in the presence of a lower alcohol - whereafter the resulting products are dissolved and the reaction is continued to cause an Amadori rearrangement and/or a Maillard reaction which is stopped when the reaction mixture becomes orange-brown in colour, whereafter the reaction mixture is dried.

10. Process according to any one of claims 5 to 9, wherein the dried product is purified by column chromatography, whereby specific fractions causing reduction of potassium ferricyanide are collected.
11. Use of at least one compound according to any one of claims 1 to 4, or of at least one N-substituted reaction product of the general formula $R_1'-NH-R_2'$ wherein R_1' comprises a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and - preferably low molecular weight, especially of less than 1000 daltons - polysaccharides, and R_2' comprises an aminoacid or a - preferably low molecular weight, especially of less than 1000 daltons - peptide radical, for the manufacture of pharmaceutical preparations to induce cytokine formation in human or mammal cells.
12. Use according to claim 11, wherein said reaction product is synthesized according to any one of claims 5 to 9 and has at least partly entered an Amadori-rearrangement and/or a Maillard-reaction, to treat and/or prevent hematological and/or immunological diseases and/or stimulate the immunosystem of humans and/or mammals.
13. Use according to claim 11 or 12, wherein R_1' is a radical of the D-form of a simple sugar, preferably selected from the group comprising glucose, xylose, galactose, rhamnose, fructose, mannose, 2-deoxy-glucose, 6-deoxy-glucose, glucosamine and galactosamine.
14. Use according to any one of claims 11 to 13, wherein R_2' is a radical of the L-form of an aminoacid or peptide,




This Sheet Has Been Replaced



preferably selected from the group comprising serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine.

15. Use according to any one of claims 11 to 14, wherein said reaction product is present together with a pharmaceutically acceptable carrier and/or adjuvant and/or optionally a lubricant in a weight ratio of reaction product to the remaining components of between 1:1 to 1:100, preferably 1:8 to 1:20 and most preferably about 1:9.
16. Use according to claim 15, wherein said reaction products are present in admixture with lactose and a lubricant, the weight ratio of lactose to the lubricant being between 20:1 and 100:1, preferably approx. 50:1.
17. Use according to any one of claims 11 to 16, wherein said reaction products are present in an amount of 0.01 - 10% by weight, preferably 0.01 - 1% by weight, especially 0.05 - 0.10% by weight, in a preparation adapted for cosmetic application.



This Sheet Has Been Replaced



- 31 -

AMADORI REACTION COMPOUNDS AND PRODUCTS, PROCESS FOR THEIR MANUFACTURE, AND THEIR USE

A B S T R A C T

Novel Amadori reaction compounds have the formula $R_1\text{-NH-R}_2$, wherein R_1 comprises the D-form of a 1-amino-1-deoxy-2-ketose radical derived from a sugar radical selected from the group of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine, and R_2 comprises the L-form of an aminoacid or peptide radical selected from the group of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine. Compounds and combinations of compounds having the general formula $R_1'\text{-NH-R}_2'$, wherein R_1' comprises a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and polysaccharides, and R_2' comprises an aminoacid or a peptide radical, are used to produce pharmaceutical preparations which in contact with human leukocytes produce interferon and other cytokines.

9603877-3


8-7703000

17 SEP 1999

- 1 -

Amadori reaction compounds and products, process for their manufacture, and their use

The present invention relates to novel Amadori reaction compounds and products, to the production thereof and to a new use of these compounds and products, having at least partly entered an Amadori rearrangement as per the following reaction scheme and/or a Maillard reaction:



Amadori reaction products are known; they are reaction products, for example, of an amino acid or a peptide with a sugar, oligo- or polysaccharide having entered an Amadori rearrangement (J. Biol. Soc. 215 (1955), Henri Borsook et al.). Thus, in DE-C-3914354, a water-soluble glycoprotein of an amino acid and a sugar is described which is isolated from an extract of *Avena sativa* seeds. Further, EP-A-406087 describes water-soluble polysaccharide-glycopeptide complexes which are derived from the cell wall of Gram positive bacterium, and J. Biol. Chem. 1985, 260/9 states that NMR-spectroscopy has been used to characterize Amadori reaction products formed by reaction of glucose with free amino groups of protein.

The invention now describes specific novel Amadori rearrangement compounds. These compounds

9603877-3
17/9/99

17 SEP 1999

- 2 -

reduce potassium ferricyanide, a test reaction for biological active substances formed in a reaction of sugar and aminoacid.

The invention further relates to a novel use of such compounds and simultaneously a novel use of Amadori reaction products of sugars and aminoacids in general.

In the past, nobody has tested the various steps of purification of the extract of an Amadori reaction product (in order to get rid of ballast substances and impurities) for biological activity.

Immunostimulating drugs are indeed known from natural sources such as mistletoe extracts, peat extracts etc. with the drawback of expensive treatment of large quantities of raw material to obtain a few grams of active substance; uncontrollable impurities that might lead to toxicity and side effects and therefore, further, to problems with administration during practical use due to the complex nature and the hardly reproducible composition.


Comparable products or product mixes from artificial sources, such as interferon or other genetic engineering methods are even more expensive to make. Furthermore, the molecules of human interferon are very often too big to penetrate the human cell wall so that only a fraction of the administered dose is effectively becoming active. Also, genetic engineering products usually have side effects and some are even toxic. Furthermore, some of them act effectively on one day and not on the next day, for reasons unknown so far.

Suprisingly, it has now been found that nearly any simple amino-acid/sugar complex after having at least partly entered an Amadori rearrangement does not show any of the above-mentioned disadvantages but on the contrary has a surprisingly high immunological activity. They can therefore be used in pharmaceutical formulations and in cosmetics. These small molecules easily penetrate the cell wall and virtually act as a nutrient. They

17 SEP 1999

- 4 -

The products which allow such an unexpected use have the general formula



wherein

R_1 represents a 1-deoxy-2-ketose radical derived from a simple sugar, oligo- or polysaccharide, and

R_2 represents an amino acid radical or peptide radical (or better "residue").


Thus the group of biologically active compounds may cover either the specific Amadori rearrangement compounds described above or the N-substituted derivatives of a number of different aminoacid compounds and one simple sugar, oligo- or - preferably low molecular weight, especially of less than 1000 daltons - polysaccharide, or N-substituted derivatives of one aminoacid compound and a number of simple sugars, oligo- and/or such polysaccharides, or else any combination of such derivatives, every single one of them having sufficient biological activity.

Preferably, R_1 in the above formula may be a radical selected from the D-form of simple sugars, especially (but not exclusively) from the D-form of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine; R_2 may be a radical selected from the L-form of aminoacid compounds such as serine, glycine, histidine, arginine, glutamine, asparagine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, methionine, hydroxyproline, tryptophane, proline, tyrosine, valine, isoleucine, leucine and lysine or else peptides of these aminoacids in any combination.

The invention further relates to a process for obtaining the above mentioned compounds and products whereby an intermediate is formed of the formula

17 SEP 1999

- 5 -



wherein R'_1 is a ~~1-deoxy-2-keto~~ radical in a straight carbon chain or any O-bridged form of a simple sugar or an oligo- or polysaccharide, and

R'_2 represents an amino acid radical or a peptide radical (or better "residue"), said intermediate being at least partly subjected to an Amadori rearrangement and/or to the Maillard reaction by continued heating of the reaction mixture - preferably under pressure - and simultaneously or subsequently removing the solvents.

In some cases, especially when an amino acid having two carboxyl groups is used, it is advantageous to add to the process a buffer salt, such as sodium bicarbonate, preferably in a molar ratio of 1:1.

It was further found that the Amadori rearrangement products are relatively susceptible to decomposition, and that decomposition products have the nature of dark brown and tar-like, unidentified compounds having lost their biological activity. Therefore it is preferred to stop the Amadori rearrangement reaction at a stage at which the reaction mixture becomes light orange-brown in colour.

It is interesting to note that the intermediate reaction products, formed when the originally opaque solution of the amino acid becomes clear (before the Amadori rearrangement), are easily hydrolysed, i.e. the reaction is reversible. With increasing rearrangement, the reversibility diminishes, i.e. the products become more stable, and the colour gradually changes from light yellow to light orange, and then finally to orange-brown when the Amadori rearrangement seems to be complete. Samples taken during such rearrangement reaction and tested (according to various procedures described later) proved that the biological activity increases with the Amadori reaction progressing, and decreases when further heating results in decomposition, for which a colour change to dark brown is a sign. Immediate re-



27

REPLACEMENT SHEET

17 SEP 1999

C L A I M S

1. Use of one or more Amadori rearrangement compounds of the general formula



wherein R_1 comprises a 1-deoxy-2-ketose radical derived from a simple sugar, oligo- or polysaccharide and R_2 comprises an amino acid radical or peptide radical, for the manufacture of a pharmaceutical composition to induce cytokine formation and/or to treat or prevent hematological diseases.

2. Use of one or more Amadori rearrangement compounds of the general formula



wherein R_1 comprises a 1-deoxy-2-ketose radical derived from a simple sugar, oligo- or polysaccharide and R_2 comprises an amino acid radical or peptide radical, for the manufacture of a cosmetic composition for tissue regeneration and/or action as a cell nutrient.

3. Use according to claim 1 or 2, characterized in that the composition comprises a mixture of at least two Amadori rearrangement compounds having different 1-deoxy-2-ketose radicals and/or different amino acid or peptide radicals.

4. Use according to anyone of claims 1 to 3, wherein the sugar radical is in its D-form, preferably selected from the group consisting of glucose, xylose, galactose, rhamnose, fructose, mannose, and 6-deoxy-glucose.

5. Use according to anyone of claims 1 to 4, wherein the amino acid radical is in its L-form, preferably selected from the group consisting of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, threonine, cysteine, cystine, glutamine, valine, asparagine, methionine, tyrosine, hydroxyproline, lysine, tryptophane, isoleucine, and leucine.



17 SEP 1999

6. Use according to any one of claims 1 to 5, wherein the oligo- or polysaccharide radical has a molecular weight of 5000 dalton or less, especially of less than 1000 dalton and/or the peptide radical has a molecular weight of less than 1000 dalton.
7. Use according to any one of claims 1 to 6, wherein the composition further comprises at least one pharmaceutically acceptable additive selected from the group consisting of a carrier, an adjuvant and a lubricant, in a weight ratio of said Amadori rearrangement compounds to said additive of between 1:1 to 1:100, preferably 1:8 to 1:20, and most preferably about 1:9.
8. Use according to any one of claims 1 to 6, wherein the composition further comprises at least one acceptable additive selected from the group consisting of a carrier, an adjuvant and a lubricant, and wherein the Amadori rearrangement compounds are present in an amount of 0.01 - 10% by weight.
9. Use according to claim 8, wherein the Amadori rearrangement compounds are present in an amount of 0.01 - 1%, preferably 0.05-0.1 % by weight.
10. Process for the manufacture of at least one Amadori rearrangement compound, preferably a mixture of Amadori rearrangement compounds, of the general formula
- $$R_1-NH-R_2,$$
- wherein R_1 comprises a 1-deoxy-2-ketose radical derived from a simple sugar, oligo- or polysaccharide and R_2 comprises an amino acid radical or peptide radical, characterized in that said process comprises
- a) reacting one or more amino acids or peptides with one or more simple sugars, oligo- or polysaccharides, wherein the molar ratio of sugars and saccharides to the amino



17 SEP 1999

acids and peptides is between 2:1 and 1:1, and preferably is 1.5:1, in a concentrated aqueous solution at a temperature of about 70-121°C,

- b) continuing heating to allow the resulting products to undergo Amadori rearrangement while simultaneously or subsequently removing the aqueous solvent, until - preferably after about 30 to 120 minutes - the reaction mixture turns light orange-brown in colour and biological activity and ferricyanide reducing capacity can be detected in samples taken from the mixture,
 - c) stopping the Amadori rearrangement substantially before decomposition occurs yielding compounds that have lost their biological activity, and
 - d) drying the reaction mixture and/or further subjecting it to a purification by column chromatography wherein biologically active fractions causing reduction of potassium ferricyanide are collected.
11. Process according to claim 10, wherein the reaction is carried out in a concentrated aqueous solution of about 0.67 to about 2.75 pbw of solids per 1 pbw of aqueous solvent
12. Process according to claim 10 or 11, wherein the reaction is carried out under pressure and/or in the presence of a lower alcohol.
13. Process according to any one of claims 10 to 12, wherein the oligo- or polysaccharides have a molecular weight of 5000 daltons or less and/or the peptides have a molecular weight of less than 1000 daltons.
14. Process according to anyone of claims 10 to 13, wherein at least one amino acid has two carboxyl groups and the reaction is carried out in the presence of a buffer salt,



17 SEP 1999

preferably sodium bicarbonate, in a molar ratio with the amino acid of 1:1.

15. A mixture of Amadori rearrangement compounds of the general formula



wherein R_1 comprises a 1-deoxy-2-ketose radical derived from a simple sugar, oligo- or polysaccharide and R_2 comprises an amino acid radical or peptide radical, characterized in that it is biologically active and preferably has cytokine inducing activity, and is obtainable in a process as defined in any one of claims 10 to 14.

16. A mixture according to claim 15, characterized in that it is part of the reaction mixture obtainable in step d) of the process of claim 10.

17. A mixture according to claim 15 or 16, for medical use.

18. A mixture according to claim 15 or 16, for cosmetic use, particularly for tissue regeneration and/or action as a cell nutrient.

19. A pharmaceutical or cosmetic composition comprising a mixture of Amadori rearrangement compounds as defined in anyone of claims 15 to 18, in combination with at least one pharmaceutically or cosmetically acceptable additive selected from the group consisting of a carrier, an adjuvant and a lubricant.

20. A composition according to claim 19, characterized in that the Amadori rearrangement compounds are present in an amount of 0.01 - 10% by weight.

21. A composition according to claim 20, characterized in that the Amadori rearrangement compounds are present in an amount of 0.01 - 1%, preferably 0.05-0.1%, by weight.



REPLACEMENT SHEET

17 SEP 1999

22. Use according to any one of claims 1 to 9, wherein said Amadori rearrangement compounds are part of a biologically active reaction mixture obtainable in step d) of the process of claim 10.

17 SEP 1999

AMADORI REACTION COMPOUNDS AND PRODUCTS, PROCESS FOR THEIR
MANUFACTURE, AND THEIR USE

A B S T R A C T

Novel Amadori reaction compounds have the formula $R_1\text{-NH-R}_2$, wherein R_1 comprises the D-form of a 1-amino-1-deoxy-2-ketose radical derived from a sugar radical selected from the group of glucose, xylose, galactose, rhamnose, fructose, mannose, 6-deoxyglucose, glucosamine and galactosamine, and R_2 comprises the L-form of an aminoacid or peptide radical selected from the group of serine, glycine, proline, histidine, arginine, alanine, aspartic acid, glutamic acid, phenylalanine, treonine, cysteine, cystine, glutamine, asparagine, methionine, tyrosine, hydroxyproline, tryptophane, valine, isoleucine, lysine and leucine. Compounds and combinations of compounds having the general formula $R_1'\text{-NH-R}_2'$, wherein R_1' comprises a 1-amino-1-deoxy-2-ketose radical derived from the group of simple sugars, oligo- and polysaccharides, and R_2' comprises an aminoacid or a peptide radical, are used to produce pharmaceutical preparations which in contact with human leukocytes produce interferon and other cytokines.