kifunensine a virus packaging cell comprising: (1) a lentiviral vector genome comprising an exogenous polynucleotide ~~encoding an exogenous antigen~~of interest, (2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN, and (3) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and (b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN. In some embodiments, the E2 glycoprotein is 90% identical to SEQ ID NO: 30 [SIN-Var1]. In some embodiments, (i) residue 160 of the E2 glycoprotein is absent or is an amino acid other than glutamic acid, (ii) one or more of residues 70, 76, or 159 of the E2 glycoprotein variant is a non-basic residue, and (iii) the E2 glycoprotein variant is not part of a fusion protein with Sindbis virus E3 glycoprotein. In some embodiments, the E2 glycoprotein is SEQ ID NO: 30 [SIN-Var1].

[0010] In some or any of the embodiments described herein, the Vpx protein comprises an amino acid sequence that is at least 80% identical to SIVmac Vpx (SEQ ID NO: 44).

[0011] In some or any of the embodiments described herein, the Vpx protein comprises an amino acid sequence at least 90% identical to SIVmac Vpx (SEQ ID NO: 44), SIVsm Vpx (SEQ ID NO: 45), SIVrcm Vpx (SEQ ID NO: 46), or HIV-2 Vpx (SEQ ID NO: 47).

[0012] In some or any of the embodiments described herein, the Vpr protein comprises an amino acid sequence at least 90% identical to SIVdeb Vpr (SEQ ID NO: 48) or SIVmus Vpr (SEQ ID NO: 49).

[0013] In some or any of the embodiments described herein, the antigen is a tumor-specific antigen or a virus-specific antigen. In some or any of the embodiments described herein, the tumor-specific antigen is selected from the group consisting of NY-ESO-1, MAGE, e.g., MAGE-A3 and MAGE-A1, MART-1/Melan-A, BAGE, RAGE, gp100, gp75, mda-7, tyrosinase, tyrosinase-related protein, e.g., TRP2, renal cell carcinoma antigen, 5T4, SM22-alpha, carbonic anhydrase I, carbonic anhydrase IX (also known as G250), HIF-1alpha, HIF-2alpha, VEGF, prostate specific membrane antigen (PSMA), prostate-specific antigen (PSA), prostatic acid phosphates, six-transmembrane epoithelial antigen of the prostate (STEAP), NKX3.1, telomerase enzyme, survivin, mesothelin, mutated ras, bcr/abl rearrangement, Her2/neu, mutated p53, wild-type p53, cytochrome P450 1B1, N-acetylglucosaminyltransferase-V, human papilloma virus protein E6, human papilloma virus protein E7, carcinoembryonic antigen, merkel cell virus T-antigen oncoproteins and alpha-fetoprotein. In some or any of the embodiments described herein, the virus-specific antigen is an HIV antigen, an SIV antigen, an adenovirus antigen, an enterovirus antigen, a

wherein expression of MAGE-A3 and NY-ESO-1 is controlled by a UbiC promoter lacking an intron, (2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN, (3) a polynucleotide comprising human codon optimized gag and pol genes, wherein the polynucleotide lacks a functional rev responsive element (RRE) and wherein the pol gene encodes an inactive integrase enzyme, (4) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and (b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN.

[0023] Related aspects of the disclosure provide a lentiviral vector particle produced by any of the methods recited above.

Compositions comprising pseudotyped lentiviral vector particles

[0023A] Another aspect of the disclosure provides a composition comprising pseudotyped lentiviral vector particles comprising (a) a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity, (b) a lentiviral vector genome comprising an exogenous polynucleotide ~~encoding an antigen~~of interest, and (c) a plurality of Sindbis envelope glycoproteins that preferentially bind cells expressing DC-SIGN, wherein said composition is more highly mannosylated compared to a control composition of the same pseudotyped lentiviral vector particles prepared in the absence of a mannosidase inhibitor, wherein high mannosylation is characterized by being more ~~EndoH-sensitve~~ susceptible to cleavage by EndoH than a control composition in the absence of a mannosidase inhibitor.

[0024] Also disclosed is a composition comprising pseudotyped lentiviral vector particles comprising (a) a SAMHD1 inhibitor, (b) an exogenous

WHAT IS CLAIMED:

1. A composition comprising pseudotyped lentiviral vector particles comprising (a) a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity, (b) an exogenous polynucleotide encoding an antigen, and (c) a plurality of Sindbis envelope glycoproteins that preferentially bind cells expressing DC-SIGN, wherein said composition is more highly mannosylated compared to a control composition of the same pseudotyped lentiviral vector particles prepared in the absence of a mannosidase inhibitor,

wherein high mannosylation is characterized by being more EndoH-sensitive than a control composition prepared in the absence of a mannosidase inhibitor.

2. The composition of claim 1, wherein EndoH sensitivity is determined by assessing the molecular weight of the envelope glycoproteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) after EndoH treatment.

3. The composition of claim 1, wherein the molecular weight of the envelope glycoproteins after treatment with EndoH has shifted about 45% or more, 70% or more, or 90% or more of the distance between (a) envelope glycoproteins not treated with endoglycosidase, and (b) envelope glycoproteins treated with PNGase F.

4. The composition of claim 1, wherein at least 30% of the envelope glycoproteins in said composition have an increased amount of EndoH-sensitive glycan as compared to control glycoproteins having the same amino acid sequence(s) in a control composition of pseudotyped lentiviral vector particles prepared in the absence of a mannosidase inhibitor.

5. The composition of claim 1, wherein the pseudotyped lentiviral vector particles are integration deficient.

6. The composition of claim 1, wherein the composition comprises glycoprotein selected from the group consisting of a Sindbis E2 glycoprotein, a Sindbis E2 glycoprotein is that 90% identical to SEQ ID NO: 30 [SIN-Var1] a Sindbis E2 glycoprotein is that 90% identical to SEQ ID NO: 30 [SIN-Var1] wherein (i) residue 160 of the E2 glycoprotein is absent or is an amino acid other than glutamic acid, (ii) one or more of residues 70, 76, or 159 of the E2 glycoprotein variant is a non-basic residue, and (iii) the E2 glycoprotein variant is not part of a

fusion protein with Sindbis virus E3 glycoprotein or a Sindbis E2 glycoprotein as set out in SEQ ID NO: 30 [SIN-Var1].

7. The composition of claim 1, wherein the Vpx protein comprises an amino acid sequence that is at least 80% identical to SIVmac Vpx [SEQ ID NO: 44], or wherein the Vpx protein comprises an amino acid sequence at least 90% identical to SIVmac Vpx (SEQ ID NO: 44), SIVsm Vpx (SEQ ID NO: 45), SIVrcm Vpx (SEQ ID NO: 46), or HIV-2 Vpx (SEQ ID NO: 47), or

wherein the Vpx protein comprises an amino acid sequence at least 90% identical to SIVdeb Vpr (SEQ ID NO: 48) or SIVmus Vpr (SEQ ID NO: 49).

8. The composition of claim 1, wherein the antigen (a) is a tumor-specific antigen or a virus-specific antigen; (b) is selected from the group consisting of NY-ESO-1, MAGE, MART-1/Melan-A, BAGE, RAGE, gp100, gp75, mda-7, tyrosinase, tyrosinase-related protein, renal cell carcinoma antigen, 5T4, SM22-alpha, carbonic anhydrase I, carbonic anhydrase IX (also known as G250), HIF-1alpha, HIF-2alpha, VEGF, prostate specific membrane antigen (PSMA), prostate-specific antigen (PSA), prostatic acid phosphates, six-transmembrane epithelial antigen of the prostate (STEAP), NKX3.1, telomerase enzyme, survivin, mesothelin, mutated ras, bcr/abl rearrangement, Her2/neu, mutated p53, wild-type p53, cytochrome P450 1B1, N-acetylglucosaminyltransferase-V, human papilloma virus protein E6, human papilloma virus protein E7, carcinoembryonic antigen, and alpha-fetoprotein; (c) is an HIV antigen, an SIV antigen, an adenovirus antigen, an enterovirus antigen, a coronavirus antigen, a calicivirus antigen, a distemper virus antigen, an Ebola virus antigen, a flavivirus antigen, a hepatitis virus antigen, a herpesvirus antigen, an infectious peritonitis virus antigen, an influenza virus antigen, a leukemia virus antigen, a Marburg virus antigen, an orthomyxovirus antigen, a papilloma virus antigen, a parainfluenza virus antigen, a paramyxovirus antigen, a parvovirus antigen, a pestivirus antigen, a picorna virus antigen, a poliovirus antigen, a pox virus antigen, a rabies virus antigen, a reovirus antigen, a retrovirus antigen, or a rotavirus antigen.

9. The composition of claim 1, wherein the lentiviral vector further comprises a nucleotide sequence encoding a second antigen.

10. The composition of claim 9 wherein the first and second antigen are expressed as a fusion protein that comprises a self-cleaving A2 peptide between the two antigens.

11. The composition of claim 10, wherein self-cleaving A2 peptide comprises the amino acid sequence of SEQ ID NO: 56 or SEQ ID NO: 57.

12. The composition of claim 10, wherein the first antigen is NY-ESO-1 and the second antigen is MAGE-A3.

13. The composition of claim 9, wherein the first and second antigen are expressed from a bi-directional promoter.

14. The composition of claim 1 wherein the envelope glycoproteins bind cells expressing mouse SIGNR1.

15. The composition of claim 1, wherein the pseudotyped lentiviral vector particles more efficiently transduce cells expressing mouse SIGNR1 compared to cells not expressing mouse SIGNR1.

16. A method of generating a pseudotyped lentiviral vector particle comprising:

(a) culturing in a culture medium comprising kifunensine a virus packaging cell comprising:

(1) a lentiviral vector genome comprising a polynucleotide encoding an exogenous antigen,

(2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN, and

(3) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and

(b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN.

17. The method of claim 16, wherein (a) the E2 glycoprotein is 90% identical to SEQ ID NO: 30 [SIN-Var1] (b) (i) residue 160 of the E2 glycoprotein is absent or is an amino acid other than glutamic acid, (ii) one or more of residues 70, 76, or 159 of the E2 glycoprotein variant is a non-basic residue, and (iii) the E2 glycoprotein variant is not part of a fusion protein with Sindbis virus E3 glycoprotein; or (c) the E2 glycoprotein is SEQ ID NO: 30 [SIN-Var1].

18. The method of claim 16, wherein (a) the Vpx protein comprises an amino acid sequence that is at least 80% identical to SIVmac Vpx [SEQ ID NO: 44]; (b) the Vpx protein comprises an amino acid sequence at least 90% identical to SIVmac Vpx (SEQ ID NO: 44), SIVsm Vpx (SEQ ID NO: 45), SIVrcm Vpx (SEQ ID NO: 46), or HIV-2 Vpx (SEQ ID NO: 47);

or (c) the Vpx protein comprises an amino acid sequence at least 90% identical to SIVdeb Vpr (SEQ ID NO: 48) or SIVmus Vpr (SEQ ID NO: 49).

19. The method of claim 16, wherein the antigen is (a) a tumor-specific antigen or a virus-specific antigen; (b) selected from the group consisting of NY-ESO-1, MAGE, MART-1/Melan-A, BAGE, RAGE, gp100, gp75, mda-7, tyrosinase, tyrosinase-related protein, renal cell carcinoma antigen, 5T4, SM22-alpha, carbonic anhydrase I, carbonic anhydrase IX (also known as G250), HIF-1alpha, HIF-2alpha, VEGF, prostate specific membrane antigen (PSMA), prostate-specific antigen (PSA), prostatic acid phosphates, six-transmembrane epoithelial antigen of the prostate (STEAP), NKX3.1, telomerase enzyme, survivin, mesothelin, mutated ras, bcr/abl rearrangement, Her2/neu, mutated p53, wild-type p53, cytochrome P450 1B1, N-acetylglucosaminyltransferase-V, human papilloma virus protein E6, human papilloma virus protein E7, carcinoembryonic antigen, and alpha-fetoprotein; or (c) an HIV antigen, an SIV antigen, an adenovirus antigen, an enterovirus antigen, a coronavirus antigen, a calicivirus antigen, a distemper virus antigen, an Ebola virus antigen, a flavivirus antigen, a hepatitis virus antigen, a herpesvirus antigen, an infectious peritonitis virus antigen, an influenza virus antigen, a leukemia virus antigen, a Marburg virus antigen, an orthomyxovirus antigen, a papilloma virus antigen, a parainfluenza virus antigen, a paramyxovirus antigen, a parvovirus antigen, a pestivirus antigen, a picorna virus antigen, a poliovirus antigen, a pox virus antigen, a rabies virus antigen, a reovirus antigen, a retrovirus antigen, or a rotavirus antigen.

20. The method of claim 16, wherein the lentiviral vector genome further comprises a nucleotide sequence encoding a second antigen.

21. The method of claim 20 wherein the first and second antigen are expressed as a fusion protein that comprises a self-cleaving A2 peptide between the two antigens.

22. The method of claim 21, wherein self-cleaving A2 peptide comprises the amino acid sequence of SEQ ID NO: 56 or SEQ ID NO: 57.

23. The method of claim 20, wherein the first antigen is NY-ESO-1 and the second antigen is MAGE-A3.

24. The method of claim 20, wherein the first and second antigen are expressed from a bi-directional promoter.

25. The method of claim 16, wherein the kifunensine is present in the culture medium at a concentration of about 0.1 μg/ml to about 10 μg/ml, or

wherein the kifunensine is present in the culture medium at a concentration of about 0.25 μg/ml to about 2 μg/ml.

26. The method of claim 16, wherein the virus packaging cell further comprises:

(i) a polynucleotide comprising gag and pol genes; and

(ii) a polynucleotide encoding a rev protein.

27. The method of claim 26, wherein the gag and pol genes are human codon optimized and comprise a non-optimized window around position 1228 to 1509 of SEQ ID NO: 54.

28. The method of claim 26, wherein the polynucleotide comprising gag and pol genes lacks a functional rev responsive element (RRE).

29. The method of claim 26, wherein the pol gene encodes an inactive integrase enzyme.

30. The method of claim 29, wherein the integrase enzyme has a D64V mutation.

31. The method of claim 16, wherein the lentiviral vector genome is derived from HIV-1.

32. The method of claim 16, wherein the lentiviral vector genome has an inactivated 3' long terminal repeat (LTR) or a self-inactivating 3' long terminal repeat (LTR).

33. The method of claim 32, wherein the lentiviral vector genome comprises a U3 element lacking at least one of an enhancer sequence, a TATA box, an Sp1 site, an NK-kappa B site, or a polypurine tract (PPT).

34. The method of claim 16, wherein the lentiviral vector genome comprises the nucleotide sequence of any one of SEQ ID NOs: 21, 22, or 23.

35. The method of claim 16, wherein the lentiviral vector genome further comprises a nucleotide sequence encoding a dendritic cell maturation/stimulatory factor.

36. The method of claim 35, wherein the dendritic cell maturation/stimulatory factor is selected from the group consisting of GM-CSF, IL-2, IL-4, IL-6, IL-7, IL-15, IL-21, IL-23, TNFα, B7.1, B7.2, 4-1BB, CD40 ligand, and drug-inducible CD40.

37. The method of claim 16, wherein the nucleotide sequence encoding an antigen is operably linked to a promoter selected from the group consisting of the human Ubiquitin-C promoter (UbiC), the cytomegalovirus immediate early promoter (CMV), the Rous sarcoma virus promoter (RSV), and the tetracycline-responsive promoter.

38. The method of claim 37, wherein the promoter is an intron-deficient promoter.

39. The method of claim 38, wherein the intron-deficient promoter is a UbiC promoter.

40. The lentiviral vector particle produced by the method of claim 16.

41. The lentiviral vector particle produced by the method of claim 26.

42. A method of generating a pseudotyped lentiviral vector particle comprising:

(a) culturing in a culture medium comprising kifunensine a virus packaging cell comprising:

(1) a lentiviral vector genome comprising a polynucleotide encoding MAGE-A3 and NY-ESO-1, wherein a polynucleotide encoding a self-cleaving TA2 peptide is positioned between the polynucleotide encoding MAGE-A3 and NY-ESO-1, wherein the lentiviral genome lacks a polypurine tract (PPT), and wherein expression of MAGE-A3 and NY-ESO-1 is controlled by a UbiC promoter lacking an intron,

(2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN,

(3) a polynucleotide comprising human codon optimized gag and pol genes, wherein the polynucleotide lacks a functional rev responsive element (RRE) and wherein the pol gene encodes an inactive integrase enzyme,

(4) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and

(b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN.

WHAT IS CLAIMED:

1. A composition comprising pseudotyped lentiviral vector particles comprising (a) a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity, (b) a lentiviral vector genome comprising an exogenous polynucleotide of interest, and (c) a plurality of Sindbis envelope glycoproteins that preferentially bind cells expressing DC-SIGN, wherein said composition is more highly mannosylated compared to a control composition of the same pseudotyped lentiviral vector particles prepared in the absence of a mannosidase inhibitor,

wherein high mannosylation is characterized by being more susceptible to cleavage by EndoH than a control composition prepared in the absence of a mannosidase inhibitor.

2. The composition of claim 1, wherein susceptibility to cleavage by EndoH is determined by assessing the molecular weight of the envelope glycoproteins by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) after EndoH treatment.

3. The composition of claim 1, wherein the molecular weight of the envelope glycoproteins after treatment with EndoH has shifted about 45% or more, 70% or more, or 90% or more of the distance between (a) envelope glycoproteins not treated with endoglycosidase, and (b) envelope glycoproteins treated with PNGase F.

4. The composition of claim 1, wherein at least 30% of the envelope glycoproteins in said composition have an increased amount of highly mannosylated glycans as compared to control glycoproteins having the same amino acid sequence(s) in a control composition of pseudotyped lentiviral vector particles prepared in the absence of a mannosidase inhibitor.

5. The composition of claim 1, wherein the pseudotyped lentiviral vector particles are integration deficient.

6. The composition of claim 1, wherein the composition comprises glycoprotein selected from the group consisting of a Sindbis E2 glycoprotein, a Sindbis E2 glycoprotein is that 90% identical to SEQ ID NO: 30 [SIN-Var1] a Sindbis E2 glycoprotein is that 90% identical to SEQ ID NO: 30 [SIN-Var1] wherein (i) residue 160 of the E2 glycoprotein is absent or is an amino acid other than glutamic acid, (ii) one or more of residues 70, 76, or 159 of the E2 glycoprotein variant is a non-basic residue, and (iii) the E2 glycoprotein variant is not part of a

fusion protein with Sindbis virus E3 glycoprotein or a Sindbis E2 glycoprotein as set out in SEQ ID NO: 30 [SIN-Var1].

7. The composition of claim 1, wherein the Vpx protein comprises an amino acid sequence that is at least 80% identical to SIVmac Vpx [SEQ ID NO: 44], or wherein the Vpx protein comprises an amino acid sequence at least 90% identical to SIVmac Vpx (SEQ ID NO: 44), SIVsm Vpx (SEQ ID NO: 45), SIVrcm Vpx (SEQ ID NO: 46), or HIV-2 Vpx (SEQ ID NO: 47), or

wherein the Vpx protein comprises an amino acid sequence at least 90% identical to SIVdeb Vpr (SEQ ID NO: 48) or SIVmus Vpr (SEQ ID NO: 49).

8. The composition of claim 1, wherein the exogenous polynucleotide of interest encodes an antigen that (a) is a tumor-specific antigen or a virus-specific antigen; (b) is selected from the group consisting of NY-ESO-1, MAGE, MART-1/Melan-A, BAGE, RAGE, gp100, gp75, mda-7, tyrosinase, tyrosinase-related protein, renal cell carcinoma antigen, 5T4, SM22-alpha, carbonic anhydrase I, carbonic anhydrase IX (also known as G250), HIF-1alpha, HIF-2alpha, VEGF, prostate specific membrane antigen (PSMA), prostate-specific antigen (PSA), prostatic acid phosphates, six-transmembrane epoithelial antigen of the prostate (STEAP), NKX3.1, telomerase enzyme, survivin, mesothelin, mutated ras, bcr/abl rearrangement, Her2/neu, mutated p53, wild-type p53, cytochrome P450 1B1, N-acetylglucosaminyltransferase-V, human papilloma virus protein E6, human papilloma virus protein E7, carcinoembryonic antigen, and alpha-fetoprotein; (c) is an HIV antigen, an SIV antigen, an adenovirus antigen, an enterovirus antigen, a coronavirus antigen, a calicivirus antigen, a distemper virus antigen, an Ebola virus antigen, a flavivirus antigen, a hepatitis virus antigen, a herpesvirus antigen, an infectious peritonitis virus antigen, an influenza virus antigen, a leukemia virus antigen, a Marburg virus antigen, an orthomyxovirus antigen, a papilloma virus antigen, a parainfluenza virus antigen, a paramyxovirus antigen, a parvovirus antigen, a pestivirus antigen, a picorna virus antigen, a poliovirus antigen, a pox virus antigen, a rabies virus antigen, a reovirus antigen, a retrovirus antigen, or a rotavirus antigen.

9. The composition of claim 1, wherein the lentiviral vector genome further comprises a second exogenous polynucleotide of interest.

10. The composition of claim 9 wherein the first and second exogenous polynucleotides of interest are expressed as a fusion protein that comprises a self-cleaving A2 peptide between the two proteins encoded by the first and second exogenous polynucleotides of interest.

11. The composition of claim 10, wherein self-cleaving A2 peptide comprises the amino acid sequence of SEQ ID NO: 56 or SEQ ID NO: 57.

12. The composition of claim 10, wherein the first exogenous polynucleotide of interest encodes NY-ESO-1 and the second exogenous polynucleotide of interest encodes MAGE-A3.

13. The composition of claim 9, wherein the first and second exogenous polynucleotides of interest are expressed using a bi-directional promoter.

14. The composition of claim 1 wherein the envelope glycoproteins bind cells expressing mouse SIGNR1.

15. The composition of claim 1, wherein the pseudotyped lentiviral vector particles more efficiently transduce cells expressing mouse SIGNR1 compared to cells not expressing mouse SIGNR1.

16. A method of generating a pseudotyped lentiviral vector particle comprising:

(a) culturing in a culture medium comprising kifunensine a virus packaging cell comprising:

(1) a lentiviral vector genome comprising an exogenous polynucleotide of interest,

(2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN, and

(3) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and

(b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN.

17. The method of claim 16, wherein (a) the E2 glycoprotein is 90% identical to SEQ ID NO: 30 [SIN-Var1] (b) (i) residue 160 of the E2 glycoprotein is absent or is an amino acid other than glutamic acid, (ii) one or more of residues 70, 76, or 159 of the E2 glycoprotein variant is a non-basic residue, and (iii) the E2 glycoprotein variant is not part of a fusion protein with Sindbis virus E3 glycoprotein; or-(c)-the E2 glycoprotein is SEQ ID NO: 30 [SIN-Var1].

18. The method of claim 16, wherein (a) the Vpx protein comprises an amino acid sequence that is at least 80% identical to SIVmac Vpx [SEQ ID NO: 44]; (b) the Vpx protein comprises an amino acid sequence at least 90% identical to SIVmac Vpx (SEQ ID NO: 44), SIVsm Vpx (SEQ ID NO: 45), SIVrcm Vpx (SEQ ID NO: 46), or HIV-2 Vpx (SEQ ID NO: 47);

or (c) the Vpx protein comprises an amino acid sequence at least 90% identical to SIVdeb Vpr (SEQ ID NO: 48) or SIVmus Vpr (SEQ ID NO: 49).

19. The method of claim 16, wherein the exogenous polynucleotide of interest encodes an antigen that is (a) a tumor-specific antigen or a virus-specific antigen; (b) selected from the group consisting of NY-ESO-1, MAGE, MART-1/Melan-A, BAGE, RAGE, gp100, gp75, mda-7, tyrosinase, tyrosinase-related protein, renal cell carcinoma antigen, 5T4, SM22-alpha, carbonic anhydrase I, carbonic anhydrase IX (also known as G250), HIF-1alpha, HIF-2alpha, VEGF, prostate specific membrane antigen (PSMA), prostate-specific antigen (PSA), prostatic acid phosphates, six-transmembrane epoithelial antigen of the prostate (STEAP), NKX3.1, telomerase enzyme, survivin, mesothelin, mutated ras, bcr/abl rearrangement, Her2/neu, mutated p53, wild-type p53, cytochrome P450 1B1, N-acetylglucosaminyltransferase-V, human papilloma virus protein E6, human papilloma virus protein E7, carcinoembryonic antigen, and alpha-fetoprotein; or (c) an HIV antigen, an SIV antigen, an adenovirus antigen, an enterovirus antigen, a coronavirus antigen, a calicivirus antigen, a distemper virus antigen, an Ebola virus antigen, a flavivirus antigen, a hepatitis virus antigen, a herpesvirus antigen, an infectious peritonitis virus antigen, an influenza virus antigen, a leukemia virus antigen, a Marburg virus antigen, an orthomyxovirus antigen, a papilloma virus antigen, a parainfluenza virus antigen, a paramyxovirus antigen, a parvovirus antigen, a pestivirus antigen, a picorna virus antigen, a poliovirus antigen, a pox virus antigen, a rabies virus antigen, a reovirus antigen, a retrovirus antigen, or a rotavirus antigen.

20. The method of claim 16, wherein the lentiviral vector genome further comprises a second exogenous polynucleotide of interest.

21. The method of claim 20 wherein the first and second exogenous polynucleotides of interest are expressed as a fusion protein that comprises a self-cleaving A2 peptide between the two proteins encoded by the first and second exogenous polynucleotides of interest.

22. The method of claim 21, wherein self-cleaving A2 peptide comprises the amino acid sequence of SEQ ID NO: 56 or SEQ ID NO: 57.

23. The method of claim 20, wherein the first exogenous polynucleotide of interest encodes NY-ESO-1 and the second exogenous polynucleotide of interest encodes MAGE-A3.

24. The method of claim 20, wherein the first and second exogenous polynucleotides of interest are expressed using a bi-directional promoter.

25. The method of claim 16, wherein the kifunensine is present in the culture medium at a concentration of about 0.1 μg/ml to about 10 μg/ml, or

wherein the kifunensine is present in the culture medium at a concentration of about 0.25 μg/ml to about 2 μg/ml.

26. The method of claim 16, wherein the virus packaging cell further comprises:

(i) a polynucleotide comprising gag and pol genes; and

(ii) a polynucleotide encoding a rev protein.

27. The method of claim 26, wherein the gag and pol genes are human codon optimized and comprise a non-optimized window around position 1228 to 1509 of SEQ ID NO: 54.

28. The method of claim 26, wherein the polynucleotide comprising gag and pol genes lacks a functional rev responsive element (RRE).

29. The method of claim 26, wherein the pol gene encodes an inactive integrase enzyme.

30. The method of claim 29, wherein the integrase enzyme has a D64V mutation.

31. The method of claim 16, wherein the lentiviral vector genome is derived from HIV-1.

32. The method of claim 16, wherein the lentiviral vector genome has an inactivated 3' long terminal repeat (LTR) or a self-inactivating 3' long terminal repeat (LTR).

33. The method of claim 32, wherein the lentiviral vector genome comprises a U3 element lacking at least one of an enhancer sequence, a TATA box, an Sp1 site, an NK-kappa B site, or a polypurine tract (PPT).

34. The method of claim 16, wherein the lentiviral vector genome comprises the nucleotide sequence of any one of SEQ ID NOs: 21, 22, or 23.

35. The method of claim 16, wherein the lentiviral vector genome further comprises a nucleotide sequence encoding a dendritic cell maturation/stimulatory factor.

36. The method of claim 35, wherein the dendritic cell maturation/stimulatory factor is selected from the group consisting of GM-CSF, IL-2, IL-4, IL-6, IL-7, IL-15, IL-21, IL-23, TNFα, B7.1, B7.2, 4-1BB, CD40 ligand, and drug-inducible CD40.

37. The method of claim 16, wherein the exogenous polynucleotide of interest is operably linked to a promoter selected from the group consisting of the human Ubiquitin-C promoter (UbiC), the cytomegalovirus immediate early promoter (CMV), the Rous sarcoma virus promoter (RSV), and the tetracycline-responsive promoter.

38. The method of claim 37, wherein the promoter is an intron-deficient promoter.

39. The method of claim 38, wherein the intron-deficient promoter is a UbiC promoter.

40. The lentiviral vector particle produced by the method of claim 16.

41. The lentiviral vector particle produced by the method of claim 26.

42. A method of generating a pseudotyped lentiviral vector particle comprising:

(a) culturing in a culture medium comprising kifunensine a virus packaging cell comprising:

(1) a lentiviral vector genome comprising a polynucleotide encoding MAGE-A3 and NY-ESO-1, wherein a polynucleotide encoding a self-cleaving TA2 peptide is positioned between the polynucleotide encoding MAGE-A3 and NY-ESO-1, wherein the lentiviral genome lacks a polypurine tract (PPT), and wherein expression of MAGE-A3 and NY-ESO-1 is controlled by a UbiC promoter lacking an intron,

(2) a polynucleotide encoding a Sindbis E2 glycoprotein that preferentially binds dendritic cells expressing DC-SIGN,

(3) a polynucleotide comprising human codon optimized gag and pol genes, wherein the polynucleotide lacks a functional rev responsive element (RRE) and wherein the pol gene encodes an inactive integrase enzyme,

(4) a polynucleotide encoding a Vpx protein or a Vpr protein that retains SAMHD1-inhibiting activity; and

(b) isolating a pseudotyped lentiviral vector particle that preferentially binds dendritic cells expressing DC-SIGN.