

CYCLOPAMINE ANALOGS

RELATED APPLICATIONS

[0001] This application claims priority to USSN 60/878,018, filed December 28, 2006, and USSN 60/941,596, filed June 1, 2007, both of which are hereby incorporated by reference in their entirety.

BACKGROUND OF THE INVENTION

[0002] The present invention generally relates to cyclopamine analogs and pharmaceutical compositions thereof, and methods for preparing such analogs and compositions. These compounds and compositions are useful for the treatment of disorders mediated by the hedgehog pathway, such as cancer.


[0003] Inhibition of the hedgehog pathway in certain cancers has been shown to result in inhibition of tumor growth. For example, anti-hedgehog antibodies have been shown to antagonize the function of the hedgehog pathway and inhibit the growth of tumors. Small molecule inhibition of hedgehog pathway activity has also been shown to result in cell death in a number of cancer types.

[0004] Research in this area has focused primarily on the elucidation of hedgehog pathway biology and the discovery of new hedgehog pathway inhibitors. Although inhibitors of the hedgehog pathway have been identified, there still exists the need to identify more potent inhibitors of the hedgehog pathway.

[0005] PCT publication WO 2006/026430 published 9 March 2006 and assigned to the same assignee as the present application, discloses a wide variety of cyclopamine analogs, focusing on those with unsaturation in the A or B ring. In the present application, the surprisingly potent analogs contain completely saturated A and B rings.

SUMMARY OF THE INVENTION

[0006] The present invention relates to analogs of steroidal alkaloids, such as cyclopamine, pharmaceutical compositions containing these compounds, and methods for preparing these compounds. In one embodiment, the present invention relates to a compound represented by the following structure:



or a pharmaceutically acceptable salt thereof;

wherein R¹ is H, alkyl, -OR, amino, sulfonamido, sulfamido, -OC(O)R⁵, -N(R⁵)C(O)R⁵, or a sugar;

R² is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, nitrile, or heterocycloalkyl;

or R^1 and R^2 taken together form =O, =S, =N(OR), =N(R), =N(NR₂), =C(R)₂;

R³ is H, alkyl, alkenyl, or alkynyl;

R^4 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl, haloalkyl, $-OR^5$, $-C(O)R^5$, $-CO_2R^5$, $-SO_2R^5$, $-C(O)N(R^5)(R^5)$, $-[C(R)_2]_qR^5$, $-[(W)-N(R)C(O)]_qR^5$, $-[(W)-C(O)]_qR^5$, $-[(W)-C(O)O]_qR^5$, $-[(W)-OC(O)]_qR^5$, $-[(W)-SO_2]_qR^5$, $-[(W)-N(R^5)SO_2]_qR^5$, $-[(W)-C(O)N(R^5)]_qR^5$, $-[(W)-O]_qR^5$, $-[(W)-N(R)]_qR^5$, $-W-NR_3^+X^-$ or $-[(W)-S]_qR^5$;

wherein each W is independently a diradical;

each q is independently 1, 2, 3, 4, 5, or 6;

X^- is a halide:

each R⁵ is independently H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl or -[C(R)₂]_p-R⁶; wherein p is 0-6; or any two occurrences of R⁵ can be taken together to form a 4-8 membered optionally substituted ring which contains 0-3 heteroatoms selected from N, O, S, and P;

each R⁶ is independently hydroxyl, -N(R)COR, -N(R)C(O)OR, -N(R)SO₂(R), -C(O)N(R)₂, -OC(O)N(R)(R), -SO₂N(R)(R), -N(R)(R), -COOR, -C(O)N(OH)(R), -OS(O)₂OR, -S(O)₂OR, -OP(O)(OR)(OR), -NP(O)(OR)(OR), or -P(O)(OR)(OR); and


each R is independently H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl or aralkyl;


provided that when R², R³, and R⁴ are H; R¹ is not hydroxyl or a sugar; further


provided that when R⁴ is hydroxyl, then R¹ is not a sugar or hydroxyl, and R¹ and R² are not C=O.


[0007] In certain embodiments, R^1 is H, hydroxyl, alkoxy, aryloxy, or amino. In other embodiments, R^1 and R^2 taken together along with the carbon to which they are bonded, form $=O$, $=N(OR)$, or $=S$. In other embodiments, R^3 is H and/or R^4 is H, alkyl, hydroxyl, aralkyl, $-[C(R)_2]_qR^5$, $-[(W)-N(R)C(O)]_qR^5$, $-[(W)-N(R)SO_2]_qR^5$, $-[(W)-C(O)N(R)]_qR^5$, $-[(W)-O]_qR^5$, $-[(W)-C(O)]_qR^5$, or $-[(W)-C(O)O]_qR^5$. In other embodiments, R^1 is H or $-OR$, R^2 is H or alkyl, and R^4 is H. In others, R^2 is H or alkyl, R^3 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, or aralkyl; and/or R^4 is H, alkyl, aralkyl, $-[(W)-N(R)C(O)]_qR^5$, $-[(W)-N(R)SO_2]_qR^5$, $-[(W)-C(O)N(R)]_qR^5$, $-[(W)-O]_qR^5$, $-[(W)-C(O)]_qR^5$, or $-[(W)-C(O)O]_qR^5$. In other embodiments, R^1 is sulfonamido.

[0008] In another embodiment, the present invention relates to a compound selected from the group consisting of:










and

or a pharmaceutically acceptable salt thereof.

[0009] In certain embodiments, the compounds mentioned above are isolated.


[0010] In another embodiment, the present invention relates to an isolated compound selected from the group consisting of:



and

or a pharmaceutically acceptable salt thereof.

[0011] In another embodiment, the present invention relates to a compound represented by the following structure:



or a pharmaceutically acceptable salt thereof;

wherein R^1 is H, alkyl, -OR, amino, sulfonamido, sulfamido, $-OC(O)R^5$, $-N(R^5)C(O)R^5$, or a sugar;

R^2 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, nitrile, or heterocycloalkyl;

or R^1 and R^2 taken together form $=O$, $=S$, $=N(OR)$, $=N(R)$, $=N(NR_2)$, $=C(R)_2$;

R^3 is H, alkyl, alkenyl, or alkynyl;

R^4 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl, haloalkyl, $-OR^5$, $-C(O)R^5$, $-CO_2R^5$, $-SO_2R^5$, $-C(O)N(R^5)(R^5)$, $-[C(R)_2]_qR^5$, $-[(W)-N(R)C(O)]_qR^5$, $-[(W)-C(O)]_qR^5$, $-[(W)-C(O)O]_qR^5$, $-[(W)-OC(O)]_qR^5$, $-[(W)-SO_2]_qR^5$, $-[(W)-N(R^5)SO_2]_qR^5$, $-[(W)-C(O)N(R^5)]_qR^5$, $-[(W)-O]_qR^5$, $-[(W)-N(R)]_qR^5$, $-W-NR_3^+X^-$, or $-[(W)-S]_qR^5$;

wherein each W is, independently, a diradical;

each q is, independently, 1, 2, 3, 4, 5, or 6;

X^- is a halide;

each R^5 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl or $-[C(R)_2]_pR^6$; wherein p is 0-6; or any two occurrences of R^5 can be taken together to form a 4-8 membered optionally substituted ring which contains 0-3 heteroatoms selected from N, O, S, and P;

each R^6 is, independently, hydroxyl, $-N(R)COR$, $-N(R)C(O)OR$, $-N(R)SO_2(R)$, $-C(O)N(R)_2$, $-OC(O)N(R)(R)$, $-SO_2N(R)(R)$, $-N(R)(R)$, $-COOR$, $-C(O)N(OH)(R)$, $-OS(O)_2OR$, $-S(O)_2OR$, $-OP(O)(OR)(OR)$, $-NP(O)(OR)(OR)$, or $-P(O)(OR)(OR)$;

each R is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl or aralkyl;

each of R^7 and $R^{7'}$ is H; or R^7 and $R^{7'}$ taken together form $=O$;

R^8 and R^9 are H; or

R^8 and R^9 taken together form a bond; and

provided that when R^3, R^4, R^8, R^9 are H and R^7 and $R^{7'}$ taken together form $=O$;
 R^1 can not be hydroxyl and R^2 can not be H;


provided that when R^3, R^4, R^8, R^9 are H and, R^7 and $R^{7'}$ taken together form $=O$;
 R^1 can not be acetate and R^2 can not be H;

provided that when R^3, R^4, R^8, R^9 are H and, R^7 and $R^{7'}$ are H; R^1 and R^2 taken together can not be $=O$; and

provided that when R^3, R^4, R^8, R^9 are H and, R^7 and $R^{7'}$ are H; R^1 and R^2 can not be H.

[0012] In some embodiments, the compound is epimerically pure and/or isolated. In other embodiments, R^4 is alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl, haloalkyl, $-OR^5$, $-[C(R)_2]_qR^5$, $-(W)-N(R)C(O)]_qR^5$, $-(W)-C(O)]_qR^5$, $-(W)-C(O)O]_qR^5$, $-(W)-OC(O)]_qR^5$, $-(W)-SO_2]_qR^5$, $-(W)-N(R^5)SO_2]_qR^5$, $-(W)-C(O)N(R^5)]_qR^5$, $-(W)-O]_qR^5$, $-(W)-N(R)]_qR^5$, or $-(W)-S]_qR^5$. Each of R^7 and $R^{7'}$ can be H. In addition, R^1 and R^2 taken together form $=O$.


[0013] In another embodiment, the present invention relates to a compound selected from the group consisting of:



or a pharmaceutically acceptable salt thereof.

[0014] In certain embodiments, the above compounds are epimerically pure and/or isolated.

[0015] In another embodiment, the present invention relates to an epimerically pure compound selected from the group consisting of:




and

or a pharmaceutically acceptable salt thereof.

[0016] Another aspect of the present invention relates to a pharmaceutical composition including any of the aforementioned compounds, and a pharmaceutically acceptable excipient.

[0017] In one embodiment, the present invention relates to a process for preparing cyclopropyl derivatives of cyclopamine and related analogs having the formula 136:



136

wherein

Y is CR^7R^8 ;

R^1 is H, alkyl, amino, hydroxyl, carboxyl, carbamoyl, alkoxy, hydroxyl, sugar or a protected hydroxyl group;

R^2 is H, alkyl, alkenyl, alkynyl, nitrile, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; or R^1 and R^2 taken together form $=\text{O}$, $=\text{S}$, $=\text{N}(\text{OR}^9)$, $=\text{N}(\text{R}^9)$, $=\text{C}(\text{R}^9)_2$, or $=\text{N}(\text{N}(\text{R}^9)_2)$;


each of R^3 , R^4 , and R^5 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; or R^3 and R^4 or R^4 and R^5 taken together form a bond;

R^6 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl, haloalkyl, $-OR^9$, $-C(O)R^9$, $-CO_2R^9$, $-SO_2R^9$, $-C(O)N(R^9)(R^9)$, $-[C(R^9)_2]_qR^9$, $-[(W)-N(R^9)C(O)]_qR^9$, $-[(W)-C(O)]_qR^9$, $-[(W)-C(O)O]_qR^9$, $-[(W)-OC(O)]_qR^9$, $-[(W)-SO_2]_qR^9$, $-[(W)-N(R^9)SO_2]_qR^9$, $-[(W)-C(O)N(R^9)]_qR^9$, $-[(W)-O]_qR^9$, $-[(W)-N(R^9)]_qR^9$, $-[(W)-S]_qR^9$, or a nitrogen protecting group; wherein each W is independently a diradical; each q is independently 1, 2, 3, 4, 5, or 6;

each of R^7 and R^8 is, independently, H, alkyl, alkenyl, aryl, nitrile, amido, halide, or ester; and

each R^9 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl.

[0018] The process includes the steps of contacting a compound of formula **136a** with a haloalkylzinc phosphate cyclopropanating agent to yield a compound of formula **136**:




136a

wherein

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 are as defined in compound **136**.

[0019] In another embodiment, the present invention provides methods for preparing a compound of formula **137**:



137

wherein

Y is CR^7R^8 ;

R^1 is H, alkyl, amino, hydroxyl, carboxyl, carbamoyl, alkoxy, hydroxyl, sugar or a protected hydroxyl group;

R^2 is H, alkyl, alkenyl, alkynyl, nitrile, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; or R^1 and R^2 taken together form $=\text{O}$, $=\text{S}$, $=\text{N}(\text{OR}^9)$, $=\text{N}(\text{R}^9)$, $=\text{C}(\text{R}^9)_2$, or $=\text{N}(\text{N}(\text{R}^9)_2)$;

each of R^3 , R^4 , and R^5 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; or R^3 and R^4 or R^4 and R^5 taken together form a bond;


R^6 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, heteroaralkyl, haloalkyl, $-\text{OR}^9$, $-\text{C}(\text{O})\text{R}^9$, $-\text{CO}_2\text{R}^9$, $-\text{SO}_2\text{R}^9$, $-\text{C}(\text{O})\text{N}(\text{R}^9)(\text{R}^9)$, $-\text{[C}(\text{R}^9)_2\text{]}_q\text{R}^9$, $-\text{[(W)-N}(\text{R}^9)\text{C}(\text{O})\text{]}_q\text{R}^9$, $-\text{[(W)-C}(\text{O})\text{]}_q\text{R}^9$, $-\text{[(W)-C}(\text{O})\text{O}\text{]}_q\text{R}^9$, $-\text{[(W)-OC}(\text{O})\text{]}_q\text{R}^9$, $-\text{[(W)-SO}_2\text{]}_q\text{R}^9$, $-\text{[(W)-N}(\text{R}^9)\text{SO}_2\text{]}_q\text{R}^9$, $-\text{[(W)-C}(\text{O})\text{N}(\text{R}^9)\text{]}_q\text{R}^9$, $-\text{[(W)-O}\text{]}_q\text{R}^9$, $-\text{[(W)-N}(\text{R}^9)\text{]}_q\text{R}^9$, $-\text{[(W)-S}\text{]}_q\text{R}^9$, or a nitrogen protecting group;

wherein each W is, independently, a diradical;

each q is independently 1, 2, 3, 4, 5, or 6;

each of R^7 and R^8 is, independently, H, alkyl, alkenyl, aryl, nitrile, amido, halide, or ester; and


each R^9 is independently H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl. The process includes the steps of: first contacting a compound of formula **137a** with a haloalkylzinc phosphate cyclopropanating agent;



137a

wherein

$R^1, R^2, R^3, R^4, R^5, R^6$ are as defined in compound 137; to form a compound with formula **137b**



**137b**

wherein

$R^1, R^2, R^3, R^4, R^5, R^6$ and Y are as defined in compound 137; and then contacting the compound of formula **137b** with an acid to give a compound of formula **137**.

[0020] In certain embodiments, R^7 and R^8 can both be H; in other embodiments R^1 can be a protected hydroxyl; and/or R^6 is a nitrogen protecting group.

[0021] In certain embodiments, the haloalkylzinc phosphate cyclopropanating agent is formed by combining a phosphoric acid of formula **141a**, a dialkylzinc, and a dihaloalkylane of formula **141b**:


**141a****141b**

wherein

each of X and X' is, independently, chloride, bromide, or iodide;

each of R^7 and R^8 is, independently, H, alkyl, halide, amido, nitro, or ester;


each of R^{10} and R^{11} is, independently, alkyl, alkenyl, aralkyl, aryl, heteroaryl, heteroaralkyl; or R^{10} and R^{11} taken together have the formula **141c**, **141d**, or **141e**;




wherein


m is, independently for each occurrence, 0, 1, 2, 3, or 4; n is, independently for each occurrence, 0, 1, or 2; and each of R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} and R^{18} is, independently, alkyl, aryl, aralkyl, or halide.

[0022] In another embodiment, the present invention relates to a process for preparing a compound of formula 142:



[0023] The process includes the steps of contacting a compound of formula 142a with a cyclopropanating agent to form a compound formula 142b; and





142b

combining the compound of formula **142b** with an acid to give the compound of formula **142**;

wherein

Y is CR^7R^8 ; R^1 is a protected hydroxyl group;

R^2 is H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; each of R^3 , R^4 , and R^5 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl; or


R^3 and R^4 or R^4 and R^5 taken together form a bond; R^6 is a nitrogen protecting group;

each of R^7 and R^8 is, independently, H, alkyl, alkenyl, aryl, nitrile, amido, halide, or ester; and

each R^9 is, independently, H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, or heteroaralkyl.

In certain embodiments, R^7 and R^8 can both be H; in other embodiments the protected hydroxyl group can be an ester or a carbonate; the nitrogen protecting can be a carbamate or an amide; R^7 and R^8 can both be H and the nitrogen protecting can be a carbamate or an amide; R^2 and R^3 can be H and R^4 and R^5 taken together can form a bond; and/or the cyclopropanating agent is generated from a dihaloalkane and a metal species (e.g., dialkyl zinc or a zinc copper couple).


[0024] In certain embodiments the cyclopropanating agent is generated from a dihaloalkane species and a dialkyl zinc species, and a phosphoric acid species or a salt thereof. The phosphoric acid species can have a structure of formula **151**:

**151**

or a salt thereof;

wherein

each of R^{10} and R^{11} is independently alkyl, alkenyl, aralkyl, aryl, heteroaryl, heteroaralkyl; or R^{10} and R^{11} taken together have the formula **151a**, **151b**, or **151c**;




wherein

m independently for each occurrence is 0, 1, 2, 3, or 4; n independently for each occurrence is 0, 1, or 2; each of R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} and R^{18} is, independently, alkyl, aryl, aralkyl, or halide.


[0025] In certain embodiments the acid is a Bronsted acid (e.g., acetic acid, trifluoromethanesulfonic acid, phosphoric acid, methanesulfonic acid or HCl). In other embodiments the acid is a Lewis acid (e.g., BF_3 , zinc chloride, zinc methanesulfonate, or a zinc salt).

[0026] The present invention also relates to a process for preparing a compound of formula **156**:


**156**

[0027] The process includes the steps of:

contacting a compound of formula **156a** with a cyclopropanating agent to form a compound formula **156b**; and



156a




156b

combining the compound of formula **156b** with an acid to give the compound of formula **156**;


where R^1 is an oxygen protecting group selected from the group consisting of formate, acetate, chloroacetate, dichloroacetate, trichloroacetate, pivaloate, benzoate, alkyl carbonate, alkenyl carbonate, aryl carbonates, aralkyl carbonate, 2,2,2-trichloroethyl carbonate, alkoxymethyl ether, aralkoxymethyl ether, alkylthiomethyl ether, aralkylthio ether, arylthio ether, trialkylsilyl ether, alkylarylsilyl ether, benzyl ether, arylmethyl ether, allyl ether; and R^2 is a nitrogen protecting group selected from the group consisting of formyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, phenyl acetyl, benzoyls, alkyl carbamates, aralkyl carbamates, aryl carbamates, allyl, aralkyl, triarylmethyl, alkoxymethyl, aralkoxymethyl, *N*-2-cyanoethyl, diarylphosphinamides, dialkylphosphinamidates, diarylphosphinamidates, and trialkylsilyl.

[0028] In certain embodiments the cyclopropanating agent is formed by combining a phosphoric acid of formula **58a**, a dialkylzinc, and a dihaloalkylane of formula **158b**:



wherein

each of X and X' is, independently, chloride, bromide, or iodide; each of R⁷ and R⁸ is, independently, H, alkyl, halide, amido, or ester; each of R¹⁰ and R¹¹ is, independently, alkyl, alkenyl, aralkyl, aryl, heteroaryl, heteroaralkyl; or R¹⁰ and R¹¹ taken together have the formula **158c**, **158d**, or **158e**;



wherein

m independently for each occurrence is 0, 1, 2, 3, or 4; n independently for each occurrence is 0, 1, or 2; each of R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷ and R¹⁸ is, independently, alkyl, aryl, aralkyl, or halide.

[0029] The oxygen protecting group can be, in some embodiments, selected from alkyl carbonates, aralkyl carbonates (e.g., benzylcarbonate), benzoates, pivaloate, or formate. The nitrogen protecting group can be selected from benzoyl, trichloroacetyl, trifluoroacetyl, formyl, alkyl carbamates, aralkyl carbamates (e.g., benzylcarbamate), aryl carbamates, diarylphosphinamides, dialkylphosphinamidates, or diarylphosphinamidates.

DETAILED DESCRIPTION OF THE INVENTION


Definitions

[0030] The definitions of terms used herein are meant to incorporate the present state-of-the-art definitions recognized for each term in the chemical and pharmaceutical

fields. Where appropriate, exemplification is provided. The definitions apply to the terms as they are used throughout this specification, unless otherwise limited in specific instances, either individually or as part of a larger group.

[0031] As used herein, the definition of each expression, e.g., alkyl, m, n, etc., when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

[0032] The term "acylamino" refers to a moiety that may be represented by the general formula:



wherein R50 and R54 independently represent a hydrogen, an alkyl, an alkenyl or $\text{-(CH}_2\text{)}_m\text{-R61}$,

where R61 represents an aryl, a cycloalkyl, a cycloalkenyl, a heterocycle or a polycycle; and m is zero or an integer in the range of 1 to 8.

[0033] The terms "alkenyl" and "alkynyl" refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively.


[0034] The terms "alkoxyl" or "alkoxy" refers to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl groups include methoxy, ethoxy, propyloxy, tert-butoxy and the like.

[0035] The term "alkyl" refers to the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (e.g., $\text{C}_1\text{-C}_{30}$ for straight chain, $\text{C}_3\text{-C}_{30}$ for branched chain), 20 or fewer. Likewise, certain cycloalkyls have from 3-10 carbon atoms in their ring structure, and others have 5, 6 or 7 carbons in the ring structure.

[0036] The term "alkylthio" refers to an alkyl group, as defined above, having a sulfur radical attached thereto. In certain embodiments, the "alkylthio" moiety is


represented by one of -S-alkyl, -S-alkenyl, -S-alkynyl, and -S-(CH₂)_m-R61, wherein m and R61 are defined above. Representative alkylthio groups include methylthio, ethylthio, and the like.

[0037] The term "amido" is art recognized as an amino-substituted carbonyl and includes a moiety that may be represented by the general formula:



wherein R50 and R51 each independently represent a hydrogen, an alkyl, an alkenyl, -(CH₂)_m-R61, R61 represents an aryl, a cycloalkyl, a cycloalkenyl, a heterocycle or a polycycle; and m is zero or an integer in the range of 1 to 8, or R50 and R51, taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure. Certain embodiments of the amide in the present invention will not include imides which may be unstable.

[0038] The terms "amine" and "amino" are art-recognized and refer to both unsubstituted and substituted amines, e.g., a moiety that may be represented by the general formulas:



wherein R50 and R51 (and optionally R52) each independently represent a hydrogen, an alkyl, an alkenyl, or -(CH₂)_m-R61, where R61 is defined as above. Thus, the term "alkylamine" includes an amine group, as defined above, having a substituted or unsubstituted alkyl attached thereto, i.e., at least one of R50 and R51 is an alkyl group.


[0039] The term "aralkyl", as used herein, refers to an alkyl group substituted with an aryl group (e.g., an aromatic or heteroaromatic group).

[0040] The term "aryl" as used herein includes 5-, 6- and 7-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, anthracene, naphthalene, pyrene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole,

triazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles" or "heteroaromatics." The aromatic ring may be substituted at one or more ring positions with such substituents as described above, for example, halogen, azide, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, alkoxy, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, sulfonamido, ketone, aldehyde, ester, heterocyclyl, aromatic or heteroaromatic moieties, $-\text{CF}_3$, $-\text{CN}$, or the like. The term "aryl" also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are "fused rings") wherein at least one of the rings is aromatic, e.g., the other cyclic rings may be cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls.

[0041] The term "Bronsted acid" refers to any substance that can act as a hydrogen ion (proton) donor.

[0042] The term "carboxyl" is defined to include esters, thiocarboxyl, aldehydes, ketones and the like and thus includes such moieties as may be represented by the general formulas:




wherein X50 is a bond or represents an oxygen or a sulfur, and each of R55 and R56 represents independently a hydrogen, an alkyl, an alkenyl, $-(\text{CH}_2)_m\text{-R61}$, where m and R61 are defined above.

[0043] The term "diradical" refers to any of a series of divalent groups from alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycloalkyl, aralkyl, heteroaryl, and heteroaralkyl

groups. For example, is an alkyl diradical; is also an alkyl diradical; is an aralkyl diradical; and is an (alkyl)heteroaralkyl diradical. Typical examples include alkylenes of general structure $(\text{CH}_2)_x$ where x is 1-6, and corresponding alkenylene and alkynylene linkers having 2-6

carbon atoms and one or more double or triple bonds; cycloalkylene groups having 3-8 ring members; and aralkyl groups wherein one open valence is on the aryl ring and one is

on the alkyl portion such as  and its isomers.

[0044] The term "haloalkyl", as used herein, refers to an alkyl group where anywhere from 1 to all hydrogens have been replaced with a halide. A "perhaloalkyl" is where all of the hydrogens have been replaced with a halide.

[0045] The term "heteroatom" as used herein means an atom of any element other than carbon or hydrogen. Examples of heteroatoms include boron, nitrogen, oxygen, phosphorus, sulfur and selenium.

[0046] The terms "heterocyclyl" or "heterocyclic group" refer to 3- to 10-membered ring structures, in some instances from 3- to 7-membered rings, whose ring structures include one to four heteroatoms. Heterocycles can also be polycycles. Heterocyclyl groups include, for example, thiophene, thianthrene, furan, pyran, isobenzofuran, chromene, xanthene, phenoxathiin, pyrrole, imidazole, pyrazole, isothiazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, pyrimidine, phenanthroline, phenazine, phenarsazine, phenothiazine, furazan, phenoxazine, pyrrolidine, oxolane, thiolane, oxazole, piperidine, piperazine, morpholine, lactones, lactams such as azetidinones and pyrrolidinones, sultams, sultones, and the like. The heterocyclic ring may be substituted at one or more positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like.

[0047] The term "isolated" in connection with a compound of the present invention means the compound is not in a cell or organism and the compound is separated from some or all of the components that typically accompany it in nature.

[0048] The term "Lewis acid" refers to any substance that can act as an electron pair acceptor.


[0049] Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to ten carbons, in some embodiments from one to six carbon atoms in its backbone structure. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths. Certain alkyl groups are lower alkyls. In some embodiments, a substituent designated herein as alkyl is a lower alkyl.

[0050] As used herein, the term "nitro" means -NO_2 ; the term "halogen" designates -F , -Cl , -Br or -I ; the term "sulfhydryl" means -SH ; the term "hydroxyl" means -OH ; and the term "sulfonyl" means $\text{-SO}_2\text{-}$.


[0051] The term "oxo" refers to a carbonyl oxygen ($=\text{O}$).

[0052] The terms "polycyclyl" or "polycyclic group" refer to two or more rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings, e.g., the rings are "fused rings". Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle may be substituted with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, nitro, sulfhydryl, imino, amido, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, sulfonyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF_3 , -CN , or the like.

[0053] The term "epimerically pure" in connection with a compound of the present invention means that the compound is substantially free of stereoisomers of the compound wherein the configuration of the stereogenic center that R^3 is bonded to is inverted. For example an epimerically pure compound represented by the following formula:




wherein R^1 , R^2 , R^3 , R^4 , R^7 , R^8 , and R^9 are as defined below, is substantially free of compounds represented by the following formula:




wherein R^1 , R^2 , R^3 , R^4 , R^7 , R^8 , and R^9 are as defined below. Epimerically pure compounds contain less than about 20 % by mass, less than about 15% by mass, less than about 10% by mass, less than about 5% by mass, or less than about 3% by mass of stereoisomeric compounds wherein the configuration of the stereogenic center that R^3 is bonded to is inverted relative to the compound.

[0054] The phrase "protecting group" as used herein means temporary substituents which protect a potentially reactive functional group from undesired chemical transformations. Examples of such protecting groups include esters of carboxylic acids, silyl ethers of alcohols, and acetals and ketals of aldehydes and ketones, respectively. The field of protecting group chemistry has been reviewed (Greene, T.W.; Wuts, P.G.M. *Protective Groups in Organic Synthesis*, 2nd ed.; Wiley: New York, 1991). In some cases, the functional group being protected and the protecting group are together referred to as one moiety. For example, the fragment shown below is sometimes referred to as a benzyl carbonate; i.e., the protected (underlined) O makes up part of the carbonate.




[0055] Similarly, the fragment shown below, in which the protected N makes up part of the carbamate, is referred to as a benzyl carbamate.



[0056] The term "sugar" as used herein refers to a natural or an unnatural monosaccharide, disaccharide or oligosaccharide comprising one or more pyranose or furanose rings. The sugar may be covalently bonded to the steroidal alkaloid of the present invention through an ether linkage or through an alkyl linkage. In certain embodiments the saccharide moiety may be covalently bonded to a steroidal alkaloid of the present invention at an anomeric center of a saccharide ring. Sugars may include, but are not limited to ribose, arabinose, xylose, lyxose, allose, altrose, glucose, mannose, gulose, idose, galactose, talose, glucose, and trehalose.

[0057] The term "sulfonamido" or "sulfonamide" as used herein includes a moiety having either of the following formulae:



wherein R50 and R56 are as defined above.

[0058] The terms "triflyl", "tosyl", "mesyl", and "nonaflyl" refer to trifluoromethanesulfonyl, *p*-toluenesulfonyl, methanesulfonyl, and nonafluorobutanesulfonyl groups, respectively. The terms "triflate", "tosylate", "mesylate", and "nonaflate" to trifluoromethanesulfonate ester, *p*-toluenesulfonate ester, methanesulfonate ester, and nonafluorobutanesulfonate ester functional groups and molecules that contain the groups, respectively.

[0059] The term "thioxo" refers to a carbonyl sulfur (=S).

[0060] It will be understood that "substitution" or "substituted with" includes the implicit proviso that such substitution is in accordance with permitted valence of the substituted atom and the substituent, and that the substitution results in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

[0061] Certain compounds of the present invention may exist in particular geometric or stereoisomeric forms. The present invention contemplates all such compounds, including *cis*- and *trans*-isomers, *R*- and *S*-enantiomers, diastereomers, (D)-isomers, (L)-isomers, the racemic mixtures thereof, and other mixtures thereof, as falling

within the scope of the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers, as well as mixtures thereof, are intended to be included in this invention.

[0062] As set out above, certain embodiments of the present compounds may contain a basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable acids. The term "pharmaceutically-acceptable salts" in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* in the administration vehicle or the dosage form manufacturing process, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed during subsequent purification. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, for example, Berge *et al.* (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19)

[0063] The pharmaceutically acceptable salts of the compounds of the present invention include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from non-toxic organic or inorganic acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like.

[0064] In other cases, the compounds of the present invention may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can

likewise be prepared *in situ* in the administration vehicle or the dosage form manufacturing process, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia, or with a pharmaceutically-acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. (See, for example, Berge *et al.*, *supra*)

Synthesis of Steroidal Alkaloid Compounds

[0065] The ring expanded steroidal alkaloid derivatives described above can be prepared directly from naturally occurring steroidal alkaloids or synthetic analogs thereof. In certain instances, the steroidal alkaloid starting materials can be cyclopamine or jervine. These steroidal alkaloids can be purchased commercially or extracted from *Veratrum Californicum*. Briefly, the process of the present invention comprises the steps of cyclopropanating suitable starting steroidal alkaloid derivatives followed by ring expansion rearrangement of the cyclopropyl derivatives. In some instances, it may be desirable to suitably protect or otherwise transform reactive functionalities present on the molecule prior to cyclopropanation. For example, an alcohol present at R¹ and a secondary nitrogen present on the fused furano-piperidine ring can both be protected prior to cyclopropanation. In certain embodiments, protecting groups that are efficiently added and removed from the alkaloid, yield intermediates in the synthetic process with improved handling properties and which allow for the efficient purification of the synthetic intermediates formed may be preferred.

[0066] Examples of oxygen protecting groups include, but are not limited to formate, acetate, chloroacetate, dichloroacetate, trichloroacetate, pivaloate, benzoates, alkyl carbonate, alkenyl carbonate, aryl carbonates, aralkyl carbonate (e.g., benzyl carbonate), 2,2,2-trichloroethyl carbonate, alkoxymethyl ether, aralkoxymethyl ether, alkylthiomethyl ether, aralkylthio ether, arylthio ether, trialkylsilyl ether, alkylarylsilyl ether, benzyl ether, arylmethyl ether, and allyl ether.

[0067] Examples of nitrogen protecting groups include, but are not limited to formyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, phenyl acetyl, benzoyls, benzamides, alkyl carbamates, aralkyl carbamates (e.g., benzyl carbamates), aryl carbamates, allyl, aralkyl, alkoxymethyl, aralkoxymethyl, *N*-2-cyanoethyl, diarylphosphinamides, dialkylphosphinamidates, diarylphosphinamidates, and trialkylsilyl.

[0068] Additional protecting groups that may be used in the methods of the present invention are described in Green T.W.; Wuts P.G. *Protective Groups in Organic Synthesis 3rd Edition*, John Wiley & Sons, Inc. 1999.

[0069] A variety of cyclopropanating agents can be used to cyclopropanate the steroidal alkaloid. 1,1-haloalkylmetal complexes and reactive species referred to as carbenoids, are commonly used to cyclopropanate olefins. These reagents are typically made using a diiodoalkane or diazoalkane and a metal or organometallic species such as Et₂Zn, *i*Bu₃Al, samarium, copper, rhodium, or palladium. In certain embodiments, Et₂Zn and diiodomethane are used to generate the 1,1-haloalkylmetal species.

[0070] The reactivity and the ease of handling of the 1,1-haloalkylzinc complexes can be modified by the addition of certain reagents, such as acids. It is believed that the addition of an acid to the 1,1-haloalkylzinc species generates an alkyl zinc mixed salt. In the examples described below a biarylphosphoric acid is combined with diiodomethane and diethylzinc to generate a putative haloalkyl zinc phosphate cyclopropanating agent. A variety of phosphoric acids can be used to generate the putative haloalkylzinc phosphate.

[0071] Other known cyclopropanation methods such as those utilizing sulfur ylides to react with an olefin conjugated to a carbonyl to add a CH₂ or CH-alkyl or CH-aryl group, and metal-catalyzed decomposition of diazoalkyl and α -diazo-carbonyl compounds, such as diazomethane and ethyl diazoacetate, can also be used: these methods readily provide cyclopropanes having alkyl, aryl, alkoxycarbonyl (-COOR), or acyl substituents. Additional cyclopropanating agents are described in Masalov et al., *Organic Letters* (2004) Vol. 6, pp. 2365-8 and Hansen et al., *Chem. Comm.* (2006) 4838-40.

[0072] The cyclopropyl ring may be substituted or unsubstituted. In cases where the cyclopropyl ring is substituted, the groups attached to the methylene of the cyclopropane will be installed onto the D ring after rearrangement and ring expansion.

[0073] The cyclopropanation reactions may be conducted in an aprotic solvent. Suitable solvents include ethers, such as diethyl ether, 1,2-dimethoxyethane, diglyme, t-butyl methyl ether, tetrahydrofuran and the like; halogenated solvents, such as chloroform, dichloromethane, dichloroethane, and the like; aliphatic or aromatic hydrocarbon solvents, such as benzene, xylene, toluene, hexane, pentane and the like; esters and ketones, such as ethyl acetate, acetone, and 2-butanone; polar aprotic solvents, such as acetonitrile, dimethylsulfoxide, dimethylformamide, and the like; or combinations of two or more solvents. In a certain embodiments, dichloromethane is the solvent used for the cyclopropanation when a dialkyl zinc and diiodomethane is used.

[0074] In the examples described below, a solution containing the cyclopropanating agent is prepared by first adding a solution of a phosphoric acid to a solution of diethylzinc, followed by addition of diiodomethane to the reaction solution. The cyclopropanation substrate is then added to this solution. Alternatively, the cyclopropanation agent can be prepared in the presence of the cyclopropanation substrate by changing the order of addition of the reagents. In certain embodiments, the cyclopropanation reaction is conducted by first adding the phosphoric acid to a solution of dialkylzinc, followed by the addition of the cyclopropanation substrate, and finally the dihaloalkane is added. Using this method the cyclopropanating agent is generated under controlled conditions and immediately reacts with the cyclopropanation substrate. The cyclopropanation methods described herein can also be used to cyclopropanate other polycyclic compounds, for example, those with steroidal backbones.

[0075] Following synthesis of the cyclopropanated steroidal alkaloid core, the compound may be derivatized using a variety of functionalization reactions known in the art. Representative examples include palladium coupling reactions to alkenylhalides or aryl halides, oxidations, reductions, reactions with nucleophiles, reactions with electrophiles, pericyclic reactions, radical reactions, installation of protecting groups, removal of protecting groups, and the like.

[0076] In the presence of Lewis or Bronsted acids the cyclopropyl analogs undergo a rearrangement and ring expansion to afford steroidal alkaloid analogs in which the D ring has been expanded by one carbon.

[0077] The cyclopropanation and ring expansion can take place in a two-step one reaction vessel process or in a two-step two reaction vessel process. When the cyclopropanation and ring expansion are conducted in the same reaction vessel the acid used to initiate the ring expansion rearrangement is added after completion of the cyclopropanation reaction. Under certain conditions, the zinc salts that are generated in the course of cyclopropanating the steroidal alkaloid can themselves act as Lewis acids to catalyze the ring expansion rearrangement. The reactivity of the zinc salts generated after the cyclopropanation can be modified by the addition of acids to generate more active Lewis acids.

[0078] As described below in the examples section, the methanesulfonic acid is added to the cyclopropanation reaction vessel after completion of the cyclopropanation. Additional examples of suitable acids include, but are not limited to zinc salts, boron compounds, magnesium salts, titanium salts, indium salts, aluminum salts, tin salts, lanthanum salts, trifluoromethanesulfonic acid, diaryloxyphosphoric acids, acetic acid, and HCl. In a certain embodiments of the invention the Lewis acid used is a zinc salt or BF_3 .

[0079] These ring expanded analogs may be further functionalized using a variety of functionalization reactions known in the art. Representative examples include palladium coupling reactions to alkenylhalides or aryl halides, oxidations, reductions, reactions with nucleophiles, reactions with electrophiles, pericyclic reactions, radical reactions, installation of protecting groups, removal of protecting groups, and the like.

Utility of Compounds

[0080] Hedgehog signaling is essential in many stages of development, especially in formation of left-right symmetry. Loss or reduction of hedgehog signaling leads to multiple developmental deficits and malformations, one of the most striking of which is cyclopia.

[0081] Many tumors and proliferative conditions have been shown to depend on the hedgehog pathway. The growth of such cells and survival can be affected by treatment with the compounds of the present invention. Recently, it has been reported that activating hedgehog pathway mutations occur in sporadic basal cell carcinoma (Xie *et al.* (1998) Nature 391: 90-2) and primitive neuroectodermal tumors of the central nervous

system (Reifenberger *et al.* (1998) Cancer Res 58: 1798-803). Uncontrolled activation of the hedgehog pathway has also been shown in numerous cancer types such as GI tract cancers including pancreatic, esophageal, gastric cancer (Berman *et al.* (2003) Nature 425: 846-51, Thayer *et al.* (2003) Nature 425: 851-56) lung cancer (Watkins *et al.* (2003) Nature 422: 313-317, prostate cancer (Karhadkar *et al.* (2004) Nature 431: 707-12, Sheng *et al.* (2004) Molecular Cancer 3: 29-42, Fan *et al.* (2004) Endocrinology 145: 3961-70), breast cancer (Kubo *et al.* (2004) Cancer Research 64: 6071-74, Lewis *et al.* (2004) Journal of Mammary Gland Biology and Neoplasia 2: 165-181) and hepatocellular cancer (Sicklick *et al.* (2005) ASCO conference, Mohini *et al.* (2005) AACR conference).

[0082] For example, small molecule inhibition of the hedgehog pathway has been shown to inhibit the growth of basal cell carcinoma (Williams, *et al.*, 2003 PNAS 100: 4616-21), medulloblastoma (Berman *et al.*, 2002 Science 297: 1559-61), pancreatic cancer (Berman *et al.*, 2003 Nature 425: 846-51), gastrointestinal cancers (Berman *et al.*, 2003 Nature 425: 846-51, published PCT application WO 05/013800), esophageal cancer (Berman *et al.*, 2003 Nature 425: 846-51), lung cancer (Watkins *et al.*, 2003. Nature 422: 313-7), and prostate cancer (Karhadkar *et al.*, 2004. Nature 431: 707-12).

[0083] In addition, it has been shown that many cancer types have uncontrolled activation of the hedgehog pathway, for example, breast cancer (Kubo *et al.*, 2004. Cancer Research 64: 6071-4), hepatocellular cancer (Patil *et al.*, 2005. 96th Annual AACR conference, abstract #2942 Sicklick *et al.*, 2005. ASCO annual meeting, abstract #9610), hematological malignancies (Watkins and Matsui, unpublished results), basal carcinoma (Bale & Yu, 2001. Human Molec. Genet. 10:757-762 Xie *et al.*, 1998 Nature 391: 90-92), medulloblastoma (Pietsch *et al.*, 1997. Cancer Res. 57: 2085-88), and gastric cancer (Ma *et al.*, 2005 Carcinogenesis May 19, 2005 (Epub)). As shown in the Examples, the compounds disclosed herein have been shown to modulate the hedgehog pathway, and selected compounds have been shown to inhibit tumor growth. It is therefore believed that these compounds can be useful to treat a variety of conditions, such as various cancers.

Pharmaceutical Compositions

[0084] In another embodiment, the present invention provides pharmaceutically acceptable compositions which comprise a therapeutically-effective amount of one or more of the compounds described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. The pharmaceutical compositions of the present invention may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, *e.g.*, those targeted for buccal, sublingual, and systemic absorption, capsules, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (3) topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually; (6) ocularly; (7) transdermally; (8) pulmonarily, or (9) nasally.

[0085] Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0086] These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, dispersing agents, lubricants, and/or antioxidants. Prevention of the action of microorganisms upon the compounds of the present invention may be ensured by the inclusion of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form

may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0087] Methods of preparing these formulations or compositions include the step of bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

[0088] When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, about 0.1 to 99%, or about 10 to 50%, or about 10 to 40%, or about 10 to 30, or about 10 to 20%, or about 10 to 15% of active ingredient in combination with a pharmaceutically acceptable carrier.

[0089] Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

[0090] The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion or metabolism of the particular compound being employed, the rate and extent of absorption, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0091] In general, a suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Generally, oral, intravenous and subcutaneous doses of the compounds of the present invention for a patient, when used for the indicated effects, will range from about 0.0001


to about 100 mg, or about 0.001 to about 100 mg, or about 0.01 to about 100 mg, or about 0.1 to about 100 mg per, or about 1 to about 50 mg per kilogram of body weight per day.

[0092] The subject receiving this treatment is any animal in need, including primates, in particular humans, and other mammals such as equines, cattle, swine and sheep; and poultry and pets in general.

Exemplification


[0093] The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1



1

Step A




[0094] Recrystallized cyclopamine **2** (14.1 g, 34.0 mmol, 1 eq) is dissolved in anhydrous DCM (70 mL) and anhydrous MeOH (29 mL). The clear solution is cooled, and triethylamine (10.4 g, 102.7 mmol, 3 eq) followed by benzyl chloroformate (6.20 g, 36.3 mmol, 1.1 eq) is added. After the addition is complete, the solution is stirred in the ice bath for 30 min. Three portions of benzyl chloroformate (3 X 0.35 g, 3.46 mmol, 0.03 eq) are added over the 3 h. The reaction is slowly quenched with water (71 mL), while maintaining the temperature below 20 °C. The mixture is stirred for 15 min before the layers are settled and separated. The organic layer is dried over sodium sulfate and

filtered. The combined filtrate is buffered with anhydrous pyridine (30 mL), concentrated, and solvent exchanged with additional anhydrous pyridine (43 mL) and concentrated.

[0095] The solution of the compound in pyridine (43 mL) is further diluted with additional anhydrous pyridine (85 mL). Trimethylacetyl chloride (8.3 g, 68.7 mmol, 2 eq) is added slowly to the reaction mixture, and the reaction is heated to 45 °C. The reaction is stirred at 45 °C for 30 min. The reaction is cooled and quenched by the addition of anhydrous MeOH (4.5 mL). The quenched reaction mixture is stirred at rt for 40 min and then diluted with toluene (97 mL) and is treated sequentially with water (35 mL) and a 10 wt % aqueous sodium carbonate solution (100 mL). After vigorous stirring, the layers are separated and the organic layer is washed twice with water (2 x 100 mL), dried over sodium sulfate, and filtered. The filter cake is rinsed with toluene (49 mL) and discarded. The combined filtrates are concentrated, and solvent exchanged with concentration to toluene (145 mL) and further concentrating to dryness. The product is recrystallized from toluene and heptane. The crystalline product is isolated by suction filtration, washed with cold heptane and dried to a constant weight to afford 15.1 g of the desired product.


Step B



[0096] Bis(2,6-dimethylphenyl)phosphate (10.65 g, 34.8 mmol, 3.1 eq) is dried by concentration from anhydrous DCM (42 mL) and held under a nitrogen atmosphere. The phosphate is then redissolved in anhydrous DCM (110 mL). In a separate flask, a solution of neat diethylzinc (4.17 g, 33.8 mmol, 3.0 eq) in anhydrous DCM (35 mL) is prepared and cooled to -25 °C. The phosphate solution is slowly transferred to the vessel containing the diethylzinc solution over 1 h, maintaining the temperature at or below -10 °C. The clear ethylzinc phosphate solution is warmed to 0 °C and stirred for 15 min. Diiodomethane (9.25 g, 34.5 mmol, 3.0 eq) is slowly added to the ethylzinc phosphate solution, maintaining the reaction temperature between 0 and 5 °C. After the addition is complete, the zinc carbenoid solution is stirred for an additional 20 min.

[0097] In a separate flask, compound **3** (7.20 g, 11.4 mmol, 1 eq) is dissolved in anhydrous DCM (36 mL) and transferred to the reaction flask. After the addition is complete, the ice bath is removed and the reaction mixture is allowed to warm to rt. After 6 h the contents of the flask are cooled to -53°C . A solution of methanesulfonic acid (3.38 g, 35.2 mmol, 3.1 eq) in anhydrous DCM (3 mL) is added, maintaining the reaction temperature below -45°C . After 10 min morpholine (20 g, 230 mmol, 20 eq) is added to the reaction mixture, maintaining the reaction temperature below -40°C . The reaction is allowed to warm to rt overnight. The morpholine salts are removed by filtration and the filter cake rinsed with DCM (22 mL). The combined filtrates are washed with 2N aqueous hydrochloric acid (2 x 140 mL), 5 % aqueous sodium bicarbonate (140 mL), 5 % aqueous sodium bicarbonate (70 mL) and 5 % aqueous sodium bisulfite (70 mL), and brine (144 mL). The organic layer is dried over magnesium sulfate and filtered. Without going to dryness, the DCM solution is concentrated and solvent exchanged with methanol (280 mL). The suspension are chilled with an ice bath and stirred for 40 minutes. The solids are isolated by filtration, washed twice with cold methanol (2 x 25 mL), and dried to a constant weight to afford 5.94 g of the desired product.

Step C



[0098] In a round bottom flask, compound **4** (11.67 g, 18.1 mmol, 1 eq) and 20 % palladium hydroxide on wet carbon (2.40 g, 1.71 mmol, 0.09 eq) are placed under a nitrogen atmosphere and diluted with EtOAc (115 mL) and toluene (60 mL). The solution is degassed with nitrogen (3X) with evacuation/purge cycles, and the process is repeated for hydrogen. The suspension is vigorously stirred at rt for 1.5 h. The hydrogen atmosphere is replaced with nitrogen. Ethylenediamine (0.57 g, 9.5 mmol, 0.52 eq) is added to the reaction, and the resulting mixture stirred for 20 min. The solution is filtered under nitrogen, and the filtrate is washed with a 2 % (wt/wt) aqueous solution of ethylenediamine (125 mL) then water (130 mL), and then dried over sodium sulfate. The drying agent is removed by filtration and the filtrate is concentrated to dryness under


vacuum. The solids that remained are chased with toluene (2 x 55 mL) on the rotary evaporator and the resulting material used without further purification in the next step

[0099] The material from the previous step is dissolved in anhydrous DCM (26 mL). The resulting clear solution is added to a 1 M solution of DIBAL in DCM (65 mL, 65 mmol, 3.6 eq) while maintaining the reaction temperature between -10 and -25 °C. After 30 min the reaction is quenched with acetone (13 mL), maintaining the reaction temperature at or below 0 °C. After stirring the quenched reaction mixture for 17 min, it is added in portions to a flask containing a cold, stirred solution of 20 % (wt/wt) aqueous Rochelle salt (200 mL). The resulting gelatinous suspension is stirred at rt for 15 h. After stirring, the clean layers are separated and the aqueous layer back extracted with DCM (30 mL). The combined organic layers are washed with water (60 mL) and dried over sodium sulfate. The drying agent is removed by filtration and discarded. The filtrate is concentrated under vacuum and solvent exchanged to toluene (225 mL added in portions). The resulting solution is further concentrated to a suspension (50 mL) and diluted with heptane (115 mL). The resulting mixture is heated until turning homogeneous (92 °C). The solution is cooled slowly over 12 h to 15 °C, and then held for 16 additional h. The crystalline product is isolated by suction filtration, washed with heptane (2 x 75 mL) and dried to a constant weight to afford 7.70 g of the desired product.

[0100] A round bottom flask is sequentially charged with the homo-allylic alcohol (7.50 g, 17.6 mmol, 1 eq), aluminum tri-*tert*-butoxide (6.10 g, 24.8 mmol, 1.4 eq), anhydrous toluene (115 mL), and 2-butanone (90 g, 1.24 mol, 7 eq). The suspension is heated under a nitrogen atmosphere to 75 °C for 16 h. The reaction temperature is then allowed to cool to 49 °C. Aqueous 20 % (w/w) potassium sodium tartrate solution (226 g) is added to the stirred suspension. The suspension is stirred at rt for 3.5 h. The layers are separated. The organic layer washed with aqueous 20 % Rochelle salt (2 x 250 mL) and water (225 mL), then dried over sodium sulfate and filtered. The residue is rinsed with toluene (30 mL) and discarded. The combined organics are concentrated to dryness. Residual reaction solvents are removed from the material by concentrating from 2-propanol (250 mL added portion-wise) to a final solution mass of 44 g. Solvent exchange from 2-propanol to heptane (275 mL added portion-wise) to a final solution mass is 41 g


fully precipitated the desired product. The suspension is diluted with of additional heptane (40 mL), stirred at rt for 1 h, and filtered. The product is washed with n-heptane (17 mL) and dried to afford 5.4 g of the desired product.

Step D




[0101] A round-bottom flask is charged with starting material (110 mg, 0.26 mmol, 1 eq) and 10% palladium on carbon (106 mg). The solids are suspended in pyridine (4 mL). The suspension is placed under hydrogen atmosphere (1 atm) and the mixture is stirred overnight at rt. The reaction mixture is filtered through Celite® and the filtrate concentrated *in vacuo*. The crude material is purified using silica gel flash chromatography (MeOH/DCM 5:95) to afford 93 mg of the desired compound. ($[M+H] = 426.6$ m/z).

Example 2




Step A



[0102] Cyclopamine 2 (5.02 g, 12.2 mmol, 1.0 eq) is dissolved in anhydrous pyridine (25 mL). DMAP (300 mg, 2.44 mmol, 0.2 eq.) and triethyl amine (5.5 mL, 39.1 mmol, 3.2 eq) are added, followed by BtO-CBz (10.5 g, 39.1 mmol, 3.2 eq) and heated at 40 °C for 2h. The mixture is cooled to rt, treated with 30 mL water, heated to get a


homogeneous solution and allowed to cool to room temp. The white precipitate that formed is collected by filtration, the filter cake is washed with portions of water (3 X 50 mL), and dried in air to afford 9.53 g of crude material which is crystallized from toluene/heptanes (1:9, 70 mL) to give 6.75 g of the desired product.

Step B




[0103] To a solution of diethyl zinc (572 mg, 482 μL , 4.63 mmol, 3.00 eq) in 5.0 mL DCM at -20°C is added a solution of bis-(2,6-Dimethylphenyl)phosphoric acid (1.42 g, 4.63 mmol, 3.00 eq) in DCM (15 mL) maintaining the reaction temperature below -8°C . The solution is aged for 15 min. at 0°C , neat diiodomethane (1.24 g, 374 μL , 3.00 eq) is added, aged for 15 min. at 0°C before adding a solution of (BisCBzcyclopamine, 1.05 g, 1.54 mmol, 1.0 eq), in DCM (10 mL). The cooling bath is replaced by a water bath at rt and maintained at rt for 4.5 h. The mixture is cooled to -76°C with a dry ice-acetone bath and treated drop wise with methanesulfonic acid DCM solution (0.6 mL 50% v/v solution 4.63 mmol, 3.0 eq) maintaining the reaction temperature below -74°C . The mixture is aged for 15-20 min. and quenched drop wise with morpholine (2.69 g, 2.70 mL, 20 eq) maintaining the reaction temperature below -65°C . The cooling bath is removed, the reaction mixture is stirred for 16-18 h., the white precipitate is filtered off, and the filtrate is successively washed with 2.0 M HCl (2 x 20 mL), satd. sodium bicarbonate (2 x 20 mL), water (2 x 20 mL) and brine (20 mL). Dried over magnesium sulfate, concentrated *in vacuo* to dryness and the crude is purified by silica gel flash chromatography (hexanes/EtOAc 17:3 $\rightarrow$ 4:1) to afford 924 mg (1.33 mmol, 86%) of the desired product.

Step C



[0104] To a solution of compound **7** (4.05 g, 5.83 mmol, 1 eq) in a solution of EtOAc:toluene (2:1, 60 mL) is added of 20% palladium hydroxide on carbon (823 mg, 0.583 mmol, 0.1 eq.). The flask is evacuated and filled with hydrogen three times. The mixture is stirred under an atmosphere of hydrogen for 1h. Neat ethylene diamine (0.38 mL) is added, stirred for 1h., and the catalyst is filtered off. The filter cake is washed twice with EtOAc:toluene (2:1, 12 mL). The combined filtrates are washed with a 2% aqueous solution of ethylene diamine (3 X 20 mL), dried over sodium sulfate and concentrated *in vacuo* to give 2.46 g as a white crystalline solid.


Step D



[0105] A round bottom flask is sequentially charged with the homo-allylic alcohol **8** (7.50 g, 17.6 mmol, 1 eq), aluminum tri-*tert*-butoxide (6.10 g, 24.8 mmol, 1.4 eq), anhydrous toluene (115 mL), and 2-butanone (90 g, 1.24 mol, 7 eq). The suspension is heated under a nitrogen atmosphere to 75 °C for 16 h. The reaction temperature is then allowed to cool to 49 °C. Aqueous 20 % (w/w) potassium sodium tartrate solution (226 g) is added to the stirred suspension. The suspension is stirred at rt for 3.5 h. The layers are separated. The organic layer washed with aqueous 20 % Rochelle's salt (2 x 250 mL) and water (225 mL), then dried over sodium sulfate and filtered. The residue is rinsed with toluene (30 mL) and discarded. The combined organics are concentrated to dryness. Residual reaction solvents are removed from the material by concentrating from 2-propanol (250 mL added portion-wise) to a final solution mass of 44 g. Solvent exchange from 2-propanol to *n*-heptane (275 mL added portion-wise) to a final solution mass of 41 g fully precipitated the desired product. The


suspension is diluted with additional *n*-heptane (40 mL), stirred at rt for 1 h, and filtered. The product is washed with *n*-heptane (17 mL) and dried to afford 5.4 g of the desired product.

Step E



[0106] A round-bottom flask is charged with starting material (110 mg, 0.26 mmol, 1 eq) and 10% palladium on carbon (106 mg). The solids are suspended in pyridine (4 mL). The suspension is placed under hydrogen atmosphere (1 atm) and the mixture is stirred overnight at rt. The reaction mixture is filtered through Celite® and the filtrate concentrated *in vacuo*. The crude material is purified using silica gel flash chromatography (MeOH/DCM 5:95) to afford 93 mg of the desired compound. ($[M+H]^+ = 426.6$ m/z).


Example 3



[0107] In a seal tube, ketone **6** (85 mg, 0.199 mmol, 1 equiv.) was charged and triethyleneglycol (2 mL) was added followed by hydrazine monohydrate (500 mg, 10 mmol, 50 equiv.) and potassium carbonate (138 mg, 1 mmol, 5 equiv.). The tube was sealed and the reaction was heated at 150 °C for 16 h. The reaction was cooled to rt and water was added. The residue was extracted with chloroform (3X). The combined organic layers are washed with water, dried over Na₂SO₄, and concentrated to dryness. The colorless oil was purified

using silica gel flash chromatography (DCM/MeOH 96:4). The purified fractions are pooled and concentrated to dryness. The resulting oil was dissolved in MTBE and washed with water (2X), 2N NaOH, and then brine. The combined organic layers are dried over Na₂SO₄, filtered and evaporated to afford 64 mg of the desired material as a white foam. ([M+H] = 412.7 m/z).

Example 4




[0108] A sealed tube was charged with compound **5** (223 mg, 0.52 mmol, 1 eq) and DMF (1mL). 2-bromopropane (1.3 g, 10.5 mmol, 20 eq) and Na₂CO₃ (73 mg, 0.68 mmol, 1.3 eq) were added and the flask was sealed and heated to 50 °C. The mixture was stirred for 16 h at which point ~70% conversion was observed. Additional (0.26 g, 2.12 mmol, 4 eq) was added. The reaction was stirred for 2 h and additional 2-bromopropane (0.13 g, 1.1 mmol, 2 eq) was added. The reaction was stirred for another 1h. The reaction was cooled to rt and water was added. The residue was extracted with MTBE (3X). The organic layers were combined washed with brine, dried over Na₂SO₄, filtered, and concentrated to dryness. The white foam was purified using silica gel flash chromatography (DCM/MeOH 99:1) to give 206 mg of the *N*-isopropyl derivative as a white foam.

[0109] The *N*-isopropyl derivative (205 mg, 0.44 mmol, 1 eq) was dissolved in of 4-methoxypyridine (1.5 mL). The flask was placed under inert atmosphere and Pd/C 10% (wet, Aldrich Degussa type E101, 40 mg) was added. The flask was sealed and purged three times with hydrogen and left 16 h under 1 atm of hydrogen. Celite® was added to the reaction mixture. The mixture was filtered through a small pad of Celite® and washed with EtOAc. The organic layer was washed with 1N HCl aq. (2x) then with water. The organic layer was

dried over Na_2SO_4 , filtered through cotton and evaporated to give 34 mg of crude. The aqueous layer was neutralized with 2N KOH and extracted with DCM (3X). The combined organic layers were washed with water, dried over Na_2SO_4 , filtered through cotton and combined with the initial 34 mg of crude. The crude material was purified using silica gel flash chromatography hexane/EtOAc (6:4) to afford 80 mg of desired product. ($[M+H]^+ = 468.7 \text{ m/z}$).


Example 5



[0110] In a round-bottom flask, compound **6** (88 mg, 0.21 mmol, 1 eq) was dissolved in anhydrous THF (1 mL). The mixture was cooled to 0 °C. Pyridine (84 μL , 1 mmol, 5 eq) and benzoylperoxide (150 mg, 0.62 mmol, 3 eq) were added successively. The homogeneous mixture was gradually warmed to rt over 2 h and stirred overnight at rt. The reaction was quenched by adding saturated NaHCO_3 . The residue was extracted with MTBE. The combined organic layers were washed with water, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude was purified using silica gel flash chromatography (hexane/EtOAc (9:1 to 4:1)) to give the N-O derivative product (60 mg, 0.11 mmol) as a white foam. This foam was dissolved in 2 mL of MeOH followed by 2N aqueous KOH (0.4 mL). The reaction mixture was stirred for 1h. Most of the MeOH was evaporated under a stream of nitrogen and 1N HCl (500 μL) was added. The material was extracted with DCM (3X). The combined organic layers were washed with water, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude was purified using silica gel


flash chromatography (hexanes/EtOAc (from 88:12→1:1)) to yield 11 mg of the desired product. ($[M+H] = 442.5$ m/z).

Example 6




12

Step A




[0111] In a round bottom flask, compound 6 (89 mg, 0.209 mmol, 1 eq) and *N*-(benzyloxycarbonyl)-aminoacetaldehyde (148 mg, 0.85 mmol, 4 eq) were dissolved in DCM (2 mL). Sodium triacetoxyborohydride (177 mg, 0.85 mmol, 4 eq) was added and the reaction was stirred for 3 h at rt. The mixture was poured in saturated aqueous NaHCO_3 solution and the residue was extracted with DCM (3x). The combined organic layers were washed with water, dried over Na_2SO_4 , filtered through cotton and evaporated to give a foamy solid (247 mg). The crude was dissolved in EtOAc (2 mL) and treated with 4M HCl (156 μL). After 30 min a white precipitate slowly formed. The resulting slurry was stirred for 15 min. Filtration gave 120 mg of white solid. The material was dissolved in EtOAc and treated with a saturated aqueous NaHCO_3 solution. The organic layer was collected and the aqueous layer was extracted with EtOAc (2X). The combined organic layers were washed with brine, dried over Na_2SO_4 . Filtration and evaporation gave the desired intermediate. This material was used in the next step without purification.

Step B




[0112] All of the material from Step A was dissolved in EtOAc (3 mL) and treated with of Pd/C 10% (30 mg, wet, Aldrich Degussa type E101). The flask was sealed and purged three times with hydrogen and left overnight under 1 atm of hydrogen. After 16 h, the mixture was filtered through a small pad of Celite® and washed with EtOAc to afford 52 mg of the amine as a white foam.

Step C




[0113] A round-bottom flask containing the amine **14** (52 mg, 0.11 mmol, 1 eq) was charged with the 1H-tetrazole-5-acetic acid (21 mg, 0.166 mmol, 1.5 eq), DCM (2 mL), EDCI (42 mg, 0.22 mmol, 2 eq) and N,N-diisopropylethylamine (57 mg, 0.44 mmol, 4 eq). The resulting yellow solution was stirred at rt for 4 h. The reaction was quenched by the addition of saturated aqueous NaHCO₃ solution and the residue was extracted with DCM (3X). The combined organic layers were dried over Na₂SO₄, filtered through cotton and evaporated to give 62 mg of off-white solid. This material was purified using silica gel flash chromatography (MeOH/DCM 5:95→10:90) to afford 31 mg of the desired product. ([M+H]⁺ = 579.7 m/z).

Example 7




15




[0114] A round-bottom flask was charged with starting material (47 mg, 0.110 mmol, 1 eq) and potassium carbonate (150 mg, 1.09 mmol, 10 eq). The solids were suspended in 2 mL of DCM. Iodomethane (14 μ L, 0.22 mmol, 2 eq) was added and the mixture was stirred for 2 at rt. TLC (DCM/MeOH 95:5) indicate >90% completion. Iodomethane (14 μ L, 0.22 mmol, 2 eq) was added to the reaction mixture, which was stirred for 2 h. The reaction mixture was added water. The phases were separated and the organics were dried and concentrated to dryness. The residue was purified using silica gel flash chromatography (DCM/MeOH 100:0 $\rightarrow$ 98:2) afford 34 mg of the desired product ([M+H] = 440.5 m/z).

Example 8




16




14

16


[0115] A round-bottom flask was charged with starting material (59 mg, 0.126 mmol, 1 eq) and potassium carbonate (350 mg, 2.5 mmol, 20 eq). The solids were suspended in 3 mL of DCM. The reaction was charged with iodomethane (80 μ L, 1.29 mmol, 10 eq) and the mixture was stirred overnight at rt. The reaction mixture was charged with water. The organic phase was separated and the aqueous layer was back extracted with DCM. The combined organic layers were dried and concentrated to dryness. The residue was purified using silica gel flash chromatography. DCM/MeOH (95:5 $\rightarrow$ 90:10) to afford 52 mg of the desired product. ([M+H] = 639.5 m/z).

Example 9


17

Step A


[0116] In a round bottom flask, compound **5** (50 mg, 0.12 mmol, 1 eq) and *N*-(*t*-butoxycarbonyl)-aminoacetaldehyde (6 mg, 0.38 mmol, 3.1 eq) were dissolved in DCM (2 mL). Sodium triacetoxyborohydride (8 mg, 0.38 mmol, 3.1 eq) was added and the reaction was stirred for 2 h at rt. The mixture was poured in saturated aqueous NaHCO₃ solution and the residue was extracted with DCM (3x). The combined organic layers were washed with water, dried over Na₂SO₄, filtered through cotton and evaporated to give a foamy solid (95 mg). The crude material was purified using silica gel flash chromatography (EtOAc/Hexanes 1:1) to yield 55 mg of compound **18**.

Step B


[0117] A round-bottom flask was charged with starting material **18** (800 mg, 1.4 mmol, 1 eq). The solid was dissolved in a solution of DCM and TFA (10 mL, 1:1). The solution was stirred for 45 min at rt. The reaction was partitioned between a solution of 10% sodium carbonate and DCM. The organic was separated and washed with 10% sodium carbonate. The organic phase was concentrated to dryness. The residue was used without further purification for the next step.


Step C

[0118] A round-bottom flask was charged with starting material (300 mg, 0.64 mmol, 1 eq) was dissolved in THF/ACN (1:1, 4 mL). The reaction was charged 37% formaldehyde in water (240 μ L, 3.22 mmol, 5 eq) and sodium cyanoborohydride (64 mg, 1 mmol, 1.6 eq). The mixture was stirred for 30 min at rt. The reaction was then partitioned between a solution a saturated aqueous solution of sodium bicarbonate and DCM. The organic was separated, dried and concentrated to dryness. The crude material was purified using silica gel flash chromatography (MeOH/DCM 5:95 $\rightarrow$ 10:90) to give the desired material.

Step D


[0119] A round-bottom flask was charged with starting material **20** (30 mg, 0.06 mmol, 1 eq) and 10% palladium on carbon (30 mg). The solids were suspended in pyridine (2 mL). The suspension was placed under hydrogen atmosphere and the mixture was stirred overnight at rt. The reaction mixture was filtered on Celite® and the filtrate concentrated to dryness. The crude material was purified using silica gel flash chromatography (MeOH/DCM 5:95 $\rightarrow$ 10:90) to gave the desired material. ([M+H] = 497.7 m/z).

Example 10**21**




[0120] A round-bottom flask was charged with starting material (85 mg, 0.20 mmol, 1 eq) was dissolved in DCM (4 mL). The reaction was charged with *N*-(2-oxoethyl)acetamide (80 mg, 0.70 mmol, 3.5 eq) and sodium triacetoxyborohydride (170 mg, 0.80, 4 eq). The mixture was stirred for 1 hour at rt. The reaction was partitioned between a solution a saturated aqueous solution of sodium bicarbonate and DCM. The organic was separated, dried and concentrated to dryness. The crude material was purified using silica gel flash chromatography (MeOH/DCM 5:95) to give the desired material. ($[M+H] = 511.7$ m/z).

Example 11




[0121] Compound 22 was synthesized according to the procedure described in example 9, using *N*-methyl-*N*-(2-oxoethyl)acetamide in place of *N*-(2-oxoethyl)acetamide. ($[M+H] = 525.7$ m/z).



Example 12



[0122] Compound 23 was synthesized according to the procedure described in example 10, using *N*-(2-oxoethyl)-3-phenylpropanamide in place of *N*-(2-oxoethyl)acetamide. ($[M+H] = 601.8$ m/z).


Example 13**24**

[0123] Compound **23** was synthesized according to the procedure described in example 10, using *N*-methyl-*N*-(2-oxoethyl)-3-phenylpropanamide in place of *N*-(2-oxoethyl)acetamide. ([M+H] 615.9 m/z)

Example 14**25**Step A


[0124] A round-bottom flask was charged with compound **6** (4.23 g, 9.94 mmol, 1 eq) and THF (60 mL). Triethylamine (6.92 mL, 49.7 mmol, 5.0 eq) and benzyl chloroformate (1.54 mL, 10.93 mmol, 1.1 eq) were added and the mixture was stirred for 1 hour at rt. The reaction mixture was partitioned between saturated aqueous bicarbonate (100 mL) and EtOAc (100 mL). The phases were separated and the organics were dried (Na_2SO_4) and concentrated to dryness. The crude material was purified using silica gel flash chromatography (EtOAc/Hexanes 2:98 $\rightarrow$ 14:86) to give 3.75g of material.

Step B




[0125] A MeOH solution (10 ml) of cerium trichloride heptahydrate (260 mg, 0.69 mmol, 1.3 eq.) at 0 °C was treated with sodium borohydride (24 mg, 0.65 mmol, 1.2 eq), stirred for 15 min, and then cooled to -78 °C. A THF solution (10 ml) of ketone **26** (300 mg, 0.54 mmol, 1 eq) was added, and the mixture was stirred for 1 h and then warmed to rt. Water (50 ml) and EtOAc (50 ml) were added, mixed, and the layers split. The organic layer was collected, washed with brine (30 ml), dried over sodium sulfate, and concentrated to a white residue. The crude product was purified by silica gel flash chromatography (ether/hexanes 2:3→1:1) to give 235 mg of 3-beta alcohol **27**.


Step C




[0126] Compound **27** (235 mg, 0.42 mmol, 1 eq) was dissolved in EtOAc (7 ml) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 50 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen, filtered through a 0.45 µm polyethylene membrane and concentrated to a clear oil. The oil was purified by silica gel flash chromatography (NH₄OH(aq)/MeOH/DCM 0.5:2:97.5→0.5:6:93.5) to give 130 mg of compound **25** as a white powder. ([M+H] = 427.4 m/z)

Example 15

28

Step A


[0127] A THF solution (10 ml) of ketone **26** (300 mg, 0.54 mmol, 1 eq) at -78 °C was treated with K-Selectride® (Potassium tri-sec-butylborohydride) (0.58 ml, 0.58 mmol, 1.1 eq) and stirred for 60 min. Methanol (1 ml) was added and the solution warmed to rt. Water (50 ml) and EtOAc (50 ml) were added, mixed, and the layers split. The organic layer was washed with brine (30 ml), dried over sodium sulfate, and concentrated to a white residue. The crude product was purified by silica gel flash chromatography (Ether/Hexanes 2:3 $\rightarrow$ 1:14) to give 170 mg of pure 3- α alcohol **29**.

Step B

[0128] Compound **29** (170 mg, 0.30 mmol, 1 eq) was dissolved in EtOAc (5 ml) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 35 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen, filtered through a 0.45 μ m polyethylene membrane and concentrated to a clear oil. The oil was purified by silica gel flash chromatography (NH₄OH(aq)/MeOH/DCM


0.5:2:97.5→0.5:6:93.5) to afford 76 mg of compound **28** as a white powder ($[M+H] = 427.4$ m/z).

Example 16




30

Step A



[0129] Compound **27** (100 mg, 0.18 mmol, 1 eq) with benzyltriethylammonium chloride (8 mg, 0.36 mmol, 0.2 eq) was dissolved in DCM (5 ml) and stirred vigorously with dimethyl sulfate (130 μ L, 1.43 mmol, 8 eq) and 50% aqueous potassium hydroxide (0.5 ml) at rt for 18h. The mixture was partitioned between water (30 ml) and EtOAc (30 ml), and the organic layer was then washed with brine, dried over sodium sulfate, and concentrated to a clear oil. The crude ether was purified by silica gel flash chromatography (Ether/Hexanes 3:7→9:113) to give 75 mg of the methyl ether as a clear oil.


Step B



[0130] Compound **31** (66 mg, 0.115 mmol, 1 eq) was dissolved in EtOAc (5 ml) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 20 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen, filtered through a 0.45 μ m polyethylene membrane and concentrated to a clear oil. The oil was


purified by silica gel flash chromatography ($\text{NH}_4\text{OH(aq)/MeOH/DCM}$ 0.5:2:97.5 $\rightarrow$ 0.5:6:93.5) to give 22 mg of compound **30** as a white powder ($[\text{M}+\text{H}] = 441.4$ m/z).

Example 17




32

Step A



[0131] Compound **27** (100 mg, 0.18 mmol, 1 eq) was dissolved in DCM (5 ml), and 4-dimethylaminopyridine (4 mg, 0.35 mmol, 0.2 eq), *N,N*-diisopropylethylamine (0.15 ml, 0.9 mmol, 5 eq), and acetic anhydride (0.070 ml, 0.72 mmol, 4 eq) were added. After stirring for 12h at rt, the solution was split between EtOAc (30 ml) and 5% aqueous sodium bicarbonate (15 ml). The organic layer was washed with brine, dried over sodium sulfate, and concentrated to a clear oil. The crude ester was purified by silica gel chromatography (Ether/Hexanes 3:7 $\rightarrow$ 9:113) to give 100 mg of the ester as a clear oil.


Step B



[0132] Compound **33** (100 mg, 0.18 mmol, 1 eq) was dissolved in EtOAc (5 ml) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 20 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen,

filtered through a 0.45 μm polyethylene membrane and concentrated to a clear oil. The oil was purified by silica gel flash chromatography ($\text{NH}_4\text{OH(aq)}/\text{MeOH}/\text{DCM}$ 0.5:2:97.5 $\rightarrow$ 0.5:6:93.5) to give 45 mg of compound **32** as a white powder ($[\text{M}+\text{H}] = 469.4$ m/z).


Example 18



34

[0133] Compound **34** was synthesized according to the procedure described in example 16, using compound **29** in place of compound **27**. ($[\text{M}+\text{H}] = 441.4$ m/z).


Example 19




35

[0134] Compound **34** was synthesized according to the procedure described in example 17, using compound **29** in place of compound **27**. MS ($[\text{M}+\text{H}] = 469.4$ m/z)


Example 20




36


Step A

[0135] An ethanol solution (5 ml) of compound **26** (185 mg, 0.3 mmol, 1 eq) was treated with hydroxylamine hydrochloride (140 mg, 2 mmol, 6 eq), sodium acetate (160 mg, 2 mmol, 6 eq), and water (0.5 mL), and the mixture was stirred at rt for 1 hr. The mixture was split between EtOAc and water (50 mL each). The organic layer was washed with brine (30 mL), dried over sodium sulfate, and concentrated to a white residue. The crude product was purified by silica gel chromatography (ether/hexanes 2:3→1:1) to give 193 mg of oxime **37**.


Step B

[0136] Compound **37** (65 mg, 0.113 mmol) was dissolved in EtOAc (7 ml) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 20 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen, filtered through a 0.45 μ m polyethylene membrane and concentrated to a clear oil. The oil was purified by silica gel flash chromatography ($\text{NH}_4\text{OH(aq)}/\text{MeOH}/\text{DCM}$ 0.5:2:97.5→0.5:6:93.5) to give 15 mg of compound **36** as a white powder, a mixture of cis and trans oxime isomers ($[\text{M}+\text{H}] = 440.3 \text{ m/z}$).


Example 21

Step A

[0137] Compound **27** (42 mg, 0.075 mmol, 1 eq) was dissolved in DCM (5 ml), and 4-dimethylaminopyridine (2 mg, 0.02 mmol, 0.2 eq), N-Cbz glycine (23 mg, 0.110 mmol, 1.5 eq), and diisopropylcarbodiimide (0.023 ml, 0.150 mmol, 2 eq) were added. After stirring for 12h at rt, the solution was split between EtOAc (30 ml) and 5% aqueous sodium bicarbonate (15 ml). The organic layer was washed with brine, dried over sodium sulfate, and concentrated to a clear oil. The crude ester was purified by silica gel flash chromatography (ether/hexanes 2:3→1:1) to give 35 mg of the ester as a clear oil

Step B


[0138] Compound **39** (235 mg, 0.42 mmol, 1 eq) was dissolved in EtOAc (7 mL) in a flask with stir bar and rubber septum. The solution was sparged with nitrogen, and Pd/C 10% (wet, Aldrich Degussa type E101, 50 mg) was added. This mixture was sparged with nitrogen and then hydrogen gas and stirred at rt for 3h. The mixture was then sparged with nitrogen, filtered through a 0.45 μ m polyethylene membrane and concentrated to a clear oil. The oil was purified by silica gel flash chromatography (NH₄OH(aq)/MeOH/DCM 0.5:2:97.5→0.5:6:93.5) to give 17 mg of the desire product as a white powder ([M+H] = 452.4 m/z).

Example 22

40

[0139] Compound 40 was synthesized according to the procedure described in example 21, using compound 29 in place of compound 27. ([M+H] = 452.4 m/z)


Example 23



41


[0140] Compound 41 was synthesized according to the procedure described in example 10, using *N*-(2-oxoethyl)-2-phenylacetamide in place of *N*-(2-oxoethyl)acetamide. ([M+H] = 587.7 m/z).

Example 24



42

Step A




[0141] A round-bottom flask was charged with alcohol 29 (7.60 g, 13.53 mmol, 1 eq) and was dissolved in DCM (115 mL). The reaction was charged with triethylamine (8.21 g, 81 mmol, 6.0 eq). The mixture was cooled to 0°C and charged with methanesulfonylchloride (1.86 g, 16.2 mmol, 1.2 eq). After 30 min, the reaction mixture was partitioned between a saturated aqueous solution of sodium bicarbonate and EtOAc. The organic layer was separated, dried over sodium sulfate and concentrated to dryness. The residue was purified

using silica gel flash chromatography (EtOAc/hexanes 10 $\rightarrow$ 25%) gave the desired material mesylate.


[0142] A round-bottom flask was charged with the mesylate (9.1 g, 14.22 mmol, 1 eq) and was dissolved in 50 mL of DMPU. The reaction was charged with sodium azide (4.62 g, 71.1 mmol, 5.0 eq) and heated to 60°C. The mixture was stirred for 17 h. The reaction mixture was then cooled to rt and charged with water. The mixture was stirred for 30 min. The mixture was filtered under vacuum, rinsed with water and air dried and used directly without purification in the next step.

Step B




[0143] A round-bottom flask was charged with azide **43** (8.35 g, 14.23 mmol, 1 eq) and THF (120 mL) was added. The reaction was then charged with triphenylphosphine (11.2 g, 42.7 mmol, 3.0 eq). The mixture was heated to 50°C and stirred for 20 h. The reaction mixture was then cooled to rt and the solvent removed under vacuum. The residue purified using silica gel flash chromatography (MeOH/DCM 10% $\rightarrow$ 20%) to afford the amine.


[0144] A round-bottom flask was charged with the amine (5.10 g, 9.09 mmol, 1 eq) and was dissolved in DCM (60 mL). The reaction was charged with N,N-diisopropylethylamine (5.88 g, 45.5 mmol, 5.0 eq). The mixture was cooled to 0°C and charged with methanesulfonylchloride (2.08 g, 18.2 mmol, 2.0 eq). After 30 minutes, the reaction mixture was partitioned between a saturated aqueous solution of sodium bicarbonate and EtOAc. The organic layer was collected, dried over sodium sulfate and concentrated to dryness. The residue was purified using silica gel flash chromatography (EtOAc/hexanes 10 $\rightarrow$ 30%) to afford the Cbz protected methanesulfonamide.

Step C


[0145] A round-bottom flask was charged with the Cbz protected methanesulfonamide (5.37 g, 8.41 mmol, 1 eq) and 10% palladium on carbon (1.0 g). The solids were suspended in 2-propanol (50 mL). The suspension was placed under hydrogen atmosphere and the mixture was stirred for 4 h at 25°C. The reaction mixture was then filtered on Celite® and the filtrate concentrated to dryness. The residue was then purified using silica gel flash chromatography (DCM/MeOH 0 → 5%) to afford the desired product. [M+H] = 505.6 m/z.

Alternate Synthesis of Compound 42

[0146] Recrystallized cyclopamine (2.07 g) is charged to an appropriately sized reaction vessel and placed under an inert atmosphere. EtOAc (7.6 g), triethylamine (1.53 g), and DMAP (307 mg) are added sequentially. The suspension is warmed to 40°C. Cbz-OBt is added in three portions over 90 minutes, keeping the internal temperature below 45°C. The reaction mixture is stirred at 40°C for 90 minutes. The temperature is maintained while methanol (26.4 g) is slowly added to the reaction mixture. The resulting suspension is cooled to room temperature and stirred for at least 15 hours. The crude product is collected by filtration and rinsed with methanol (5 g). The white solid is dried under vacuum to a constant weight and recrystallized from heptane (30.3 g) and toluene (3.2 g) to afford Compound 24a (3.0 g).




[0147] Solid bis(2,6-dimethylphenyl) hydrogenphosphate and **24a** are pre-dried and placed under a nitrogen atmosphere. Neat diethyl zinc (722 mg) is charged to an appropriately sized reaction vessel containing DCM (9.0 g). DCM solutions of the phosphate (1.83 g in 17.9 g) and IPI-332690 (1.34 g in 3.6 g) are added sequentially at or below 25 °C. Diiodomethane (1.58 g) is charged and the reaction is stirred at 28°C for 4-6 hours. The reaction is cooled to -45°C and a solution of methanesulfonic acid in DCM (566 mg in 1.5 g) is charged. After 15 minutes, morpholine (1.711 g) is added and the mixture is allowed to warm to room temperature overnight. The organic layer is washed twice with 2N HCl (2 x 13.6 g) then sequentially with 4.8 wt % sodium carbonate (aq), 4.8 wt% sodium sulfite (aq), and 4.8 wt% brine (13.6 g each). The organic layer is dried, filtered, concentrated to 4 g and diluted with isopropanol (4 g). The product is crystallized from solution by the slow addition of methanol (9.3 g). Filtration with a methanol rinse (2.6 g) and drying afford 1.09 g of **24b** (79% isolated yield).




[0148] Johnson Matthey Pd/C catalyst A-305038-5 (890 mg) is charged to an appropriately sized reaction vessel, followed by **24b** (2.24 g). The reaction vessel is purged with N₂ and toluene (21.8 g) and 2-propanol (6.7 g) are added sequentially. The system is degassed and placed under a nitrogen atmosphere, and the process is repeated with hydrogen. The system is stirred vigorously and the hydrogen blanket is maintained at one atmosphere for 4-5 hours. The reaction is monitor by either TLC or HPLC. If incomplete, the reaction is


inerted, additional catalyst (145 mg) is charged, and the hydrogen atmosphere is returned for another hour. Ethylenediamine (12.9 mg) is charged and the mixture was stirred for 15 minutes. The catalyst is removed by filtration with a toluene:IPA (3:1) rinse. The filtrate and rinses are concentrated and solvent exchanged to toluene. The product is crystallized from toluene (19.0 g) and heptane (18.0 g) to afford **24c** as a white crystalline solid (1.34 g, 98% yield).



[0149] **24c** (644 mg) is charged to an appropriately sized reaction vessel followed by aluminum t-butoxide (525 mg), toluene (8.34 g, 15 vol), and 2-butanone (7.83 g, 15 vol). The contents of the flask are degassed with evacuation/nitrogen purge cycles to remove oxygen and the reaction mixture is heated at 75°C with vigorous stirring for 16-18 hours. The reaction is quenched by the addition of aqueous Rochelle's salt (2.6 g in 10.3 g water) and the mixture vigorously stirred for one hour at 45 °C. The aqueous and organic layers are separated. The aqueous layer is back extracted with a mixture of toluene (2.9 g) and EtOAc (2.9 g). The organic layers are combined and washed with fresh Rochelle's salt solution (2.6 g in 10.3 g water) and then with water (12.9 g). The resulting organic layer is dried over sodium sulfate (1.97 g), filtered, and concentrated in vacuo. The product is crystallized via a charge and concentration solvent exchange first to IPA (6.5 g) and then Heptane (7.7 g). The thick heptane slurry (~2.7 g) is stirred overnight and solids are collected by filtration. Vacuum drying affords **24d** (550 mg) in an 85% yield.




[0150] The enone **24d** (459 mg) and Johnson-Matthey 5% palladium on carbon (A503023-5, 101 mg) are charged to an appropriately sized multi neck reaction vessel. The vessel is purged with nitrogen and 3-picoline (2.2 g) is charged as the solvent. Stirring is started and the vessel is first degassed using nitrogen and then stirred under hydrogen at atmospheric pressure for 8 hours. At the end of the reaction, the catalyst is removed by filtration through 0.2 micron media, rinsing with ACN (1.4 ml). The filtrate and rinse are combined in a clean reaction vessel equipped with mechanical stirring, an internal temperature probe, and a nitrogen atmosphere. A solution of citric acid (3.7 g) in water (9.2 ml) is charged to the reaction vessel at or below 30°C, and IPI-335589 is allowed to slowly crystallize from solution as the citrate salt at 20 and then 0°C. The crystalline product is recovered by suction filtration and washed with water (3.7 ml). After drying, the citrate salt, **24e**, is isolated as a hydrate (3-5 wt% water) in 89.5% yield (622 mg) with a $\beta:\alpha$ ratio approaching 90:1.




[0151] **24e** (1.50 g) is charged to the appropriately sized reactor along with 2-methyltetrahydrofuran (7.7 g) and 1M sodium carbonate (9.0 ml). A solution of benzyl chloroformate (454 mg) in 2-methyltetrahydrofuran (300 mg) is added via addition funnel and the reaction is ambient temperature for 1-2 hours. When the reaction is complete, the stirring is stopped, the layers are separated and the organic layer is washed twice with water (2 x 6 g). The organic layer is dried over of sodium sulfate (3 g), filtered and concentrated. Residual

water is reduced further by concentration from fresh 2-methyltetrahydrofuran (6.5 g) and the material is transferred as solution in anhydrous 2-methyltetrahydrofuran to the next reaction.




[0152] Commercial 1 M K-Selectride in THF (1.20 g) is charged to a dry reaction vessel under a nitrogen atmosphere, diluted with anhydrous 2-methyltetrahydrofuran (2.10 g) and cooled to -65°C . The solution of **24f** (0.41 g) in 2-methyltetrahydrofuran (1.5 g), is then slowly added to the reaction vessel to control the internal temperature at $-65\pm 5^{\circ}\text{C}$. The reaction is stirred for 2 hours and warmed to -20°C over approximately 1 hour and stirred for an additional hour. The reaction is monitored by HPLC and reactions that are incomplete are driven to completion with additional K-selectride. The reaction is quenched at low temperature with MeOH (0.33 g), then 3M NaOH (2.4 g) at -20°C and 15% hydrogen peroxide in water (1.04 g) at or below 5°C , then stirring overnight at ambient temperatures. The layers are split and the organic layer is washed sequentially with 1M aqueous NaOH (2 ml), 0.5 M aqueous Na_2SO_3 (2 ml), and water (2 ml) adjusted to a pH of 3 with HCl. The organic layer is dried over sodium sulfate (0.82 g), filtered and concentrated. The product **24g** (0.457 g) is re-concentrated from DCM (0.9 g) and used in the next reaction.




[0153] **24g** (1.36 g) is charged with anhydrous DCM (18.1 g) to an appropriately size reaction vessel, place under an inert atmosphere and cooled to -20°C . Triethylamine (0.61 mg) is charged followed by the slow addition of methanesulfonyl chloride (373 mg) in anhydrous DCM (300 mg). The reaction is stirred for 1 hour at -20°C . The reaction is monitored by

HPLC. Incomplete reactions are driven to completion with additional methanesulfonyl chloride. When complete, the reaction is quenched with water (13.6 g) and allowed to warm. The layers are separated and the organic layer is washed with 2.5 wt% sodium bicarbonate (13.8 g) and then water (10.9 g). The organic layer is dried over of sodium sulfate (4 g), filtered, and concentrated. The product solution is solvent exchanged via charge and concentration to t-butyl methyl ether (10.9 ml) and then 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (DMPU, 4.7 ml). The DMPU solution is used directly in the next reaction.




[0154] Sodium azide (0.74 g) is charged to an appropriately sized reaction vessel. The solution of **24h** (1.46 g) in DMPU (5.9 g) is charged to the reaction vessel, rinsing with additional DMPU (1.9 g). The suspension is heated to 60°C for 15 hours, maintaining a nitrogen sweep for the entire reaction. The reaction is cooled to ambient temperature and diluted with MTBE (11.7 g). The organic solution is washed 3 times with 2% saline (3 x 8 g), dried over sodium sulfate (4.4 g), filtered, and concentrated. The product is concentrated from THF (6.4 g) and used directly in the next reaction.




[0155] The crude **24i** (1.34 g) is dissolved and transferred to a suitably sized reaction vessel with THF (12.6 g). Triphenylphosphine (0.70 g) and water (0.44 g) are charged and the reaction is heated to 55°C for 15-24 hours. When complete, the reaction is cooled to ambient

temperature, dried with magnesium sulfate (1.4 g), filtered and concentrated. The solids are dissolved and concentrated from three portions of DCM (3 x 9 g) and purified by silica gel chromatography using DCM/MeOH/Et₃N gradients to remove reagent based impurities. The pooled fractions are concentrated to dryness, dissolved in DCM (6.8 g) and concentrated to dryness again to afford an amorphous solid (1.12 g) which is used in the next reaction.




[0156] **24j** (1.09 g) is dissolved and transferred to an appropriately sized reaction vessel with anhydrous DCM (15.8 g) and placed under a nitrogen atmosphere. The solution is cooled to 0°C. N,N-diisopropylethylamine (357 mg) and neat methanesulfonyl chloride (0.165 ml) are charged sequentially while maintaining temperature between below 5 °C. The reaction is monitored by HPLC. Incomplete reactions are driven to completion with additional methanesulfonyl chloride. The reaction is quenched with 0.4 M aqueous sodium bicarbonate (11.4 g) and warmed to ambient temperature. The layers are separated and the aqueous phase is back extracted with DCM (5.8 g). The combined organic layers are dried over magnesium sulfate (0.55 g), filtered and concentrated. The product **24k** is dissolved and stripped from 2-propanol (4.0 g) to remove residual DCM and used directly in the next reaction.



[0157] Aldrich Degussa type E101 NE/W 10% Pd/C (249 mg) is charged to an appropriately sized reaction vessel and placed under a nitrogen atmosphere. A 2-propanol (9.8 g) solution of **24k** (1.24 g) is charged to the reaction vessel. The system is degassed and placed under a nitrogen atmosphere, and the process is repeated with hydrogen. The reaction


is stirred under a 1 atm of hydrogen at ambient temperature for 8 hours. An inert atmosphere is returned to the vessel and a second charge of catalyst (125 mg) slurried in 2-propanol (0.5 g) is added to the reaction. The reaction mixture is degassed and placed under a nitrogen atmosphere, and the process is repeated with hydrogen. The reaction is stirred under 1 atm of hydrogen for another 15 hours at ambient temperature. The reaction is monitored by HPLC. Incomplete reactions are treated with additional catalyst and hydrogen. When complete, the reaction is filtered, treated with steam activated carbon (200 mg), and filtered again. The solution is dried by partial concentration transferred to a reaction vessel and diluted with anhydrous 2-propanol to 0.09 M based on the theoretical yield. A 1.25 M HCl solution in 2-propanol (1.64 g) is charged over 20 minutes. The hydrochloride salt crystallizes slowly with gentle stirring and is isolated by filtration. The crystals are washed with 2-propanol (2.5 g) and vacuum dried to afford Compound 42 (916 mg, 80% yield) as a 1:1 IPA solvate.

Example 25



47

Step A




42

48

[0158] A round-bottom flask was charged with the amine 42 (1.1 g, 2.1 mmol, 1 equiv.), dry tetrahydrofuran (10 ml) and pyridine (880 μL , 10.9 mmol, 5 equiv.). The cooled (0°C) mixture was treated with benzoylperoxide (1.6 g, 6.5 mmol, 3 equiv.). The mixture was


stirred for 2 hours at 0°C then overnight at 25°C. Reaction mixture diluted with MTBE and washed with a mixture of saturated aqueous NaHCO₃ with 1 N NaOH until the layer split. The organic layer was collected and the aqueous was re-extracted once with MTBE. Combined organic layers were washed with brine, dry over Na₂SO₄, filtered and concentrated to dryness. The crude oil was dissolved in 5 mL of CH₂Cl₂, loaded onto SiO₂ (40 g) column and eluted from hexanes/EtOAc (10% to 50%) to give the benzoyl derivative **48** (380 mg) ([M+H]⁺ = 625.4 m/z).

Step B




[0159] A round-bottom flask was charged with **48** (374 mg, 0.6 mmol, 1 equiv.) and MeOH (5 mL). The solution was treated at 25°C in presence of 2 N KOH (0.3 mL, 0.6 mmol, 1 equiv.). The mixture was stirred for 3h. The solvent was removed under vacuum. MTBE was added to the residue, which was neutralized with 1N HCl. The layers were cut and the aqueous layer was extracted with two portions of CH₂Cl₂. Combined organic layers were dried over Na₂SO₄, filtered, and concentrated to dryness. The crude material (380 mg) was dissolved with CH₂Cl₂, loaded onto a SiO₂ column (12 g) and eluted with hexanes/EtOAc (0% to 100%) to give the hydroxylamine **47**. The material was lyophilized from t-BuOH/7% H₂O to give 213 mg of **47** as a white powder ([M+H]⁺ = 521.4 m/z).


Example 26




50

Step A



[0160] A heat-gun dried flask was charged with dry CH_2Cl_2 (5 mL) and benzyl alcohol (785 μL , 7.58 mmol, 1.3 equiv.). The cooled (0°C) solution was treated with chlorosulfonyl isocyanate (506 μL , 5.83 mmol, 1 equiv.). Then, DMAP (1.4 g, 11.6 mmol, 2 equiv.) was added and the mixture was stirred for 1 h at 25°C . After complete dissolution of DMAP, the reaction was clear for a short period. Then, a white fluffy precipitate formed. The mixture was diluted with CH_2Cl_2 (30 mL) and washed with three portions (30 mL each) of water. The organic layer was dried over Na_2SO_4 , filtered, and evaporated to dryness. The desired white solid **51** was taken to the next step without purification.

Step B

[0161] A round-bottom flask was charged with **52** (30 mg, 0.053 mmol, 1 equiv.) and **51** (18 mg, 0.053 mmol, 1 equiv.). Both reagents were dissolved in CH_2Cl_2 (2 mL) and the solution was stirred at 25°C . The crude material was loaded onto a SiO_2 column (4 g) and eluted with hexanes/EtOAc (0% to 50%) to give 16 mg of the sulfamoylated derivative **53** ($[\text{M}+\text{Na}] = 796.4 \text{ m/z}$).


Step C

[0162] A round-bottom flask was charged with **53** (16 mg, 0.021 mmol, 1 equiv.) and 11 mg of 10% Pd/C (wet, Aldrich Degussa type E101). The material was suspended in 2-propanol (3 mL). The flask was sealed and purged three times with hydrogen and left overnight under 1 atm of hydrogen. The slurry was filtered through 0.2 micron Acrodisc, washed with 2-propanol, and the solvent was removed under vacuum. The residue was purification by SiO₂ column (1 g) eluting with CH₂Cl₂/MeOH (5% to 10%). The major product was lyophilized from t-BuOH/7% H₂O to give 9 mg of sulfamide **50** ([M+H] = 506.4 m/z).


Example 27Step A

[0163] A round-bottom flask was charged with cycloamine 4-en-3-one (3.5 g, 8.5 mmol, 1 equiv.) and pyridine (70 mL). The reactor was charged with Pd/C (10% Pd, 500 mg). The reaction was placed under 1 atmosphere of hydrogen. After 3.5 hrs, LCMS showed complete consumption of starting material. The catalyst was filtered off on an Acrodisk 0.2 micron filter and washed with toluene. The solvent was removed by azeotropic removal with toluene (2 x 10 mL). The desired material **56**, 3.5 g ($[M+H] = 412.5$ m/z) was used as it for the next step.


Step B



[0164] A round-bottom flask was charged with **56** (1.2 g, 2.8 mmol, 1 equiv.), CH_2Cl_2 (10 mL) and triethylamine (1.9 mL, 14.2 mmol, 5 equiv.). The cooled (0°C) solution was treated with CBz-Cl (440 μL , 2.8 mmol, 1 equiv.). After 1 hr, LCMS showed complete consumption of starting material. The mixture was diluted with water. The layers were cut and the organic layer was washed twice with water. The organic layer was dried over sodium sulfate, filtered, and concentrated to dryness. The product was purified by column chromatography (SiO_2 , 40 g) eluting with hexane/EtOAc (0 to 20%) to give **57** (891 mg) ($[M+\text{Na}] = 468.4$ m/z).

Step C


[0165] In a round-bottom flask, the ketone **57** was azeotroped a couple times with CH_2Cl_2 and dried under vacuum for 1 h. Under nitrogen, the ketone **2** (693 mg, 1.27 mmol, 1 equiv.) was dissolved in anhydrous THF (20 mL) and the solution was cooled to -78°C . A 1 M solution of K-selectride in THF (1.9 mL, 1.9 mmol, 1.5 equiv.) was added dropwise. After 1 h, the reaction was complete by TLC. The reaction was quenched by addition of 2.6 mL of 5 N NaOH followed by slow addition of 2.6 mL of 30% wt H_2O_2 . The resulting mixture was allowed to stir overnight. The mixture was partitioned between water and EtOAc. The aqueous layer was back extracted with EtOAc. The combined organic were washed first with water (buffered with a small portion of ammonium chloride) then with brine. The organic were dried, filtered, and concentrated to a crude foam (840 mg). The crude material was dissolved in CH_2Cl_2 , loaded on a SiO_2 column (40g) and eluted with hexanes/EtOAc (0 to 50%) to give **58** (565 mg).

Step D


58**59**

[0166] In a round-bottom flask under nitrogen, the alcohol **58** (530 mg, 0.98 mmol, 1 equiv.) was dissolved in 5 mL of anhydrous CH_2Cl_2 and triethylamine (800 μL , 5.81 mmol, 6 equiv.). The reaction mixture was cooled to 0°C and Ms-Cl (112 μL , 1.45 mmol, 1.5 equiv) was added dropwise. The mixture was stirred at 0°C for 30 min. TLC (hexane: EtOAc, 7:3) showed ~ 70% conversion. 70 μL of triethylamine (70 μL , 0.5 equiv.) and Ms-Cl (10 μL , 0.1 equiv) were charged to the reaction vessel. After 90 min, a solution of saturated bicarbonate was charged and the residue was extracted with CH_2Cl_2 . The organic layer was washed with water, dried and concentrated to a off-white foam. The material was dissolved in CH_2Cl_2 and purified with SiO_2 (40 g) eluting with hexanes/EtOAc (0% to 50%) to give **59** (430 mg).


Step E



[0167] In a round-bottom flask, the mesylate **59** (420 mg, 0.67 mmol, 1 equiv.) was dissolved in 2 mL of DMPU. The solution was treated with sodium azide (218 mg, 3.4 mmol, 5 equiv.) at 60°C for 5 h. The mixture was cooled to 25°C , then poured into ice-water to generate a white solid. The compound was extracted with MTBE (3 times). The combined organic layers were washed with water (2X), then brine. The organic layers were dried over Na_2SO_4 , filtered, and concentrated to a white foam (342 mg). The desired material **60** was used as is for the next step.

Step F


[0168] In a round-bottom flask equipped with a condenser, the azide **60** (336 mg, 0.58 mmol, 1 equiv.) was dissolved in 7 mL of THF and 140 μ L of water and treated with triphenylphosphine (462 mg, 1.76 mmol, 3 equiv.). The mixture was heated to 70°C overnight. TLC (hexane/EtOAc, 7:3) confirmed the reaction was complete. The reaction was concentrated to dryness. The crude material was dissolved in CH_2Cl_2 , loaded onto 12 g of SiO_2 and eluted with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (0 to 20%) to give the amine **61** (254 mg).

Step G

[0169] In a round-bottom flask under nitrogen, the amine **61** (248 mg, 0.45 mmol, 1 equiv.) was dissolved in 7 mL of anhydrous CH_2Cl_2 and *N,N*-diisopropylethylamine (237 μ L, 0.91 mmol, 2 equiv.). The reaction mixture was cooled to 0°C and Ms-Cl (70 μ L, 1.45 mmol, 1.5 equiv) was added dropwise. The mixture was stirred at 0°C for 2 h. TLC (hexane/EtOAc, 7:3) showed a little amount of amine. The mixture was charged with 10 μ L of

Ms-Cl (0.2 equiv.), and warmed to 25°C for 1h. The reaction mixture was diluted with CH₂Cl₂ then a saturated solution of NaHCO₃. The layers were cut. The aqueous layer was extracted with one portion of CH₂Cl₂. The combined organic layers were washed with water, dried over Na₂SO₄, filtered and concentrated to dryness. The crude (326 mg) was added to a SiO₂ column (12 g) and was eluted with hexanes/EtOAc (0 to 50%) to give the sulfonamide **62** (256 mg).

Step H



[0170] A round-bottom flask was charged with the sulfonamide **62** (250 mg, 0.4 mmol, 1 equiv.) and 50 mg of 10% Pd/C (wet, Aldrich Degussa type E101 lot 08331KC). The material was suspended in EtOAc (5 mL). The flask was sealed and purged three times with hydrogen and stirred under 1 atm of hydrogen. After 3 h some conversion was observed, but a lot of starting material remained. The slurry was filtered through 0.2 micron Acrodisc, washed with 2-propanol. The filtrate solution was re-subjected to the reaction condition by adding 54 mg of catalyst. The reaction was completed after 3 h. The slurry was filtered through 0.2 micron Acrodisc, washed with 2-propanol, and the solvent was concentrated to dryness. The crude material (200 mg) was loaded onto a SiO₂ column (12 g) and the compound was eluted using a gradient CH₂Cl₂/MeOH (0 to 10%) to give the free amine. The material was lyophilized from t-BuOH/7% H₂O to give 175 mg of **55** as a white powder ([M+H] = 491.3 m/z).

Example 28: Inhibition of the Hedgehog pathway in cell culture

[0171] Hedgehog pathway specific cancer cell killing effects may be ascertained using the following assay. C3H10T1/2 cells differentiate into osteoblasts when contacted with the

sonic hedgehog peptide (Shh-N). Upon differentiation; these osteoblasts produce high levels of alkaline phosphatase (AP) which can be measured in an enzymatic assay (Nakamura *et al.*, 1997 *BBRC* 237: 465). Compounds that block the differentiation of C3H10T1/2 into osteoblasts (a Shh dependent event) can therefore be identified by a reduction in AP production (van der Horst *et al.*, 2003 *Bone* 33: 899). The assay details are described below. The results approximate (EC_{50} for inhibition) of the differentiation assay is shown below in Table 1.

Assay Protocol

Cell Culture

[0172] Mouse embryonic mesoderm fibroblasts C3H10T1/2 cells (obtained from ATCC) were cultured in Basal MEM Media (Gibco/Invitrogen) supplemented with 10% heat inactivated FBS (Hyclone), 50 units/ml penicillin and 50ug/ml streptomycin (Gibco/Invitrogen) at 37C with 5% CO₂ in air atmosphere.

Alkaline Phosphatase Assay

[0173] C3H10T1/2 cells were plated in 96 wells with a density of 8×10^3 cells/well. Cells were grown to confluence (72hrs). After sonic Hedgehog (250ng/ml), and/or compound treatment, the cells were lysed in 110 μ L of lysis buffer (50 mM Tris pH 7.4, 0.1% TritonX100), plates were sonicated and lysates spun through 0.2 μ m PVDF plates (Corning). 40 μ L of lysates was assayed for AP activity in alkaline buffer solution (Sigma) containing 1mg/ml p-Nitrophenyl Phosphate. After incubating for 30 min at 37 °C, the plates were read on an Envision plate reader at 405 nm. Total protein was quantified with a BCA protein assay kit from Pierce according to manufacturer's instructions. AP activity was normalized against total protein. Note that "A" indicates that the IC_{50} is less than 20 nM, "B" indicates that the IC_{50} is 20-100 nM, "C" indicates that the IC_{50} is > 100 nM.

**Table 1 -- Approximate EC₅₀ for Inhibition
Differentiation Assay EC₅₀**

Compound	
1	A
7	C
8	C
9	C
10	C
13	A
20	A
21	B
22	A
23	A
24	A
27	B
29	B
31	B
33	C
35	A
37	A
39	B
40	A
42	A
55	A

Examples 29: Pancreatic Cancer Model

[0174] The activity of Compound 42 was tested in a human pancreatic model: BXPC-3 cells were implanted subcutaneously into the flanks of the right legs of mice. On day 42 post-tumor implant, the mice were randomized into two groups to receive either Vehicle (30%HPBCD) or Compound 42. Compound 42 was dosed at 40mg/kg/day. After receiving 25 daily doses, Compound 42 statistically reduced tumor volume growth by 40% when compared to the vehicle control (p=0.0309). At the end of the study, the tumors were harvested 4 hours post the last dose to evaluate an on target response by q-RT-PCR analysis of the HH pathway genes. Analysis of human Gli-1 resulted in no modulation. Analysis of murine Gli-1 mRNA levels resulted in a robust down-regulation in the Compound treated group, when compared to the Vehicle treated group.

Example 30: Medulloblastoma Model

[0175] The activity of Compound 42 was also evaluated in a transgenic mouse model of medulloblastoma. Mice that are heterozygous for loss of function mutations in the tumor suppressors Patched1 (Ptch1) and Hypermethylated in Cancer (Hic1) develop

spontaneous medulloblastoma. Similar to human medulloblastoma, these tumors demonstrate complete promoter hypermethylation of the remaining Hic1 allele, as well as loss of expression of the wild type Ptch1 allele. When passaged as subcutaneous allografts, these tumors grow aggressively and are Hedgehog pathway-dependent. This model was employed to evaluate the efficacy of orally administered Compound, and to correlate activity with drug exposure in plasma and tumors. Oral administration (PO) of a single dose of Compound 42 led to dose-dependent down-regulation of the HH pathway in subcutaneously implanted tumors, as measured by decreased Gli-1 mRNA expression 8 hours post dose administration.

[0176] Daily (QD) administration of the Compound PO led to a dose dependent inhibition of tumor growth, with frank tumor regression seen at higher doses. The approximate effective daily oral dose for 50% inhibition of tumor growth (ED50) is 4mg/kg. When animals were treated QD for 21 days, long term survival was observed following cessation of treatment (>60 days), with little to no tumor re-growth.

Example 31: Lung Cancer Model

[0177] To test the activity of Compound 42 in a human SCLC tumor model, LX22 cells were implanted subcutaneously into the flank of the right leg. LX22 is primary xenograft model of SCLC derived from chemo-naive patients, which has been maintained by mouse to mouse passaging. This tumor responds to etoposide/carboplatin chemotherapy in way that closely resembles a clinical setting. LX22 regresses during chemotherapy treatment, goes through a period of remission, and then begins to recur. In the LX22 model, Compound single agent activity and its ability to modulate the chemoresistant recurrence was tested. On day 32 post tumor implant, mice were randomized into three dosing groups to receive Vehicle (30% HBPCD), Compound, or the chemotherapy combination of etoposide and carboplatin (E/P). Compound 42 was administered at a dose of 40mg/kg/day, and after 16 consecutive doses there was no measurable difference between the treated and vehicle groups. Etoposide was administered i.v at 12mg/kg on days 34, 35, 36, and 48, while Carboplatin was administered i.v. at 60mg/kg on days 34, 41, and 48, post tumor implant. On day 50, the E/P treated mice were further randomized to receive either Vehicle (30%HPBCD) or Compound follow up treatment. The Compound was administered at the oral multi-dose MTD of 40mg/kg/day, and after 35 consecutive doses there was a substantial delay in tumor recurrence in the treated group, compared to the vehicle group (p=0.0101).

Incorporation by Reference


[0178] All of the U.S. patents and U.S. published patent applications cited herein are hereby incorporated by reference.

Equivalents

[0179] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.


CLAIMS

1. A compound having the structure:



or a pharmaceutically acceptable salt thereof.

2. A pharmaceutical composition comprising a compound having the structure:



or a pharmaceutically acceptable salt thereof,
and at least one pharmaceutically acceptable excipient.