


ABSTRACT

A compound of Formula II, or a salt, ester, solvate, hydrate or prodrug thereof:



where:

X^1 is a branched or unbranched C_{1-10} alkyl, $(CH_2)_s-NH-(CH_2)_t$, $(CH_2)_s-O-(CH_2)_t$, or $(CH_2)_s-C(NH_2)-(CH_2)_v-NH-(CH_2)_t$, where s, t and v are each, independently an integer from 1 to 5;

10 A and B are each, independently, C_{6-10} aryl, C_{6-10} aryl- C_{1-6} alkyl, C_{3-9} heteroaryl, or C_{3-9} heteroaryl- C_{1-6} alkyl, each optionally substituted with 1, 2 or 3 independently selected R^g groups;

C and D are each, independently, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-6} alkyl, C_{3-7} heterocycloalkyl, C_{3-7} heterocycloalkyl- C_{1-6} alkyl, C_{6-10} aryl, C_{6-10} aryl- C_{1-6} alkyl, C_{3-9} heteroaryl, or C_{3-9} heteroaryl- C_{1-6} alkyl, each optionally substituted with 1, 2 or 3 independently selected R^g groups;

15 each R^g is, independently, halogen, C_{1-6} alkyl, C_{1-6} haloalkyl, hydroxyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, amino, C_{1-6} alkylamino, or di- C_{1-6} alkylamino; and n^1 and p^1 are each, independently, integers from 1 to 10.

[No Figure]