

the contents of GLP-1 and glucagon may be more effective.

However, for the combined therapy, it is required to improve the physical characteristics of glucagon, which aggregates at a neutral pH and precipitates with time, thus showing poor solubility.

Under these circumstances, the present inventors have developed glucagon derivatives with partial modifications of the amino acid sequence of glucagon for the improvement of the therapeutic effects of glucagon on hypoglycemia and obesity by improving the physical properties of glucagon, and have discovered that these glucagon derivatives have improved solubility and higher stability at a neutral pH, thereby completing the present invention.

[Disclosure]

[Technical Problem]

An object of the present invention is to provide a novel glucagon derivative with improved physical properties.

Another object of the present invention is to provide a composition for preventing or treating hypoglycemia containing the glucagon derivative as an active ingredient.

A further object of the present invention is to provide a composition for preventing or treating obesity containing the glucagon derivative as an active ingredient.

[Technical Solution]

In order to achieve the above objects, in an aspect, the present invention provides a novel glucagon derivative as defined in the appending claim set. Also disclosed herein is a glucagon derivative which includes the amino acid sequence of the following General Formula 1 and has an isoelectric point (pI), which is not the same as, i.e., different from, that of native glucagon:

X1-X2-QGTF-X7-SDYS-X12-X13-X14-X15-X16-X17-X18-X19-X20-X21-F-X23-X24-W-L-X27-X28-T (General Formula 1)

wherein, in General Formula 1,

X1 is histidine, desamino-histidine, N-dimethyl-histidine, β -hydroxyimidazopropionic acid, 4-imidazoacetic acid, β -carboxyimidazopropionic acid, tryptophan, tyrosine, or deleted;

X2 is α -methyl-glutamic acid, aminoisobutyric acid (Aib), D-alanine, glycine, Sar(N-methylglycine), serine, or D-serine;

X7 is threonine or valine;

X12 is lysine or cysteine;

X13 is tyrosine or cysteine;

X14 is leucine or cysteine;

X15 is aspartic acid, glutamic acid, or cysteine;

X16 is glutamic acid, aspartic acid, serine, α -methyl-glutamic acid, or cysteine;

X17 is aspartic acid, glutamine, glutamic acid, lysine, arginine, serine, valine, or cysteine;

X18 is aspartic acid, glutamine, glutamic acid, arginine, or cysteine;

X19 is alanine or cysteine;

X20 is lysine, glutamic acid, glutamine, aspartic acid, lysine, or cysteine;

X21 is aspartic acid, glutamic acid, valine, or cysteine;

X23 is valine or arginine;

X24 is valine, leucine, glutamine, or arginine;

X27 is isoleucine or methionine; and

X28 is arginine or asparagine

(with the proviso that when the amino acid sequence of General Formula 1 is identical to SEQ ID NO: 1, it is excluded).

The glucagon derivative as described herein includes a peptide, a peptide derivative, or a peptide mimetic, which has improved physical properties by having a different pI from that of native glucagon by modifying a part of the amino acid(s) of native glucagon.

As used herein, the term "native glucagon" refers to native human glucagon having the sequence of His-Ser-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Lys-Tyr-Leu-Asp-Ser-Arg-Arg-Ala-Gln-Asp-Phe-Val-Gln-Trp-Leu-Met-Asn-Thr (SEQ ID NO: 1).

Referring to the sequence of General Formula 1 as described herein, the amino acids proceed from the N-terminus on the left to the C-terminus on the right according to the conventional method of description. Accordingly, when the "position" of any particular residue is referred to in the sequence of General Formula 1, it should be interpreted in the same manner as when any position in native glucagon or other molecules is referred to.

Over the entire specification of the present invention, not only the conventional one-letter or three-letter codes for naturally occurring amino acids, but also those three-letter codes generally allowed for other amino acids, such as α -aminoisobutyric acid (Aib), Sar(N-methylglycine), and α -methyl-glutamic acid, are used.

Additionally, the amino acids mentioned herein are abbreviated according to the nomenclature rules of IUPAC-IUB as follows:

Alanine	A;	Arginine	R;
Asparagine	N;	Aspartic acid	D;
Cysteine	C;	Glutamic acid	E;
Glutamine	Q;	Glycine	G;
Histidine	H;	Isoleucine	I;
Leucine	L;	Lysine	K;
Methionine	M;	Phenylalanine	F;
Proline	P;	Serine	S;
Threonine	T;	Tryptophan	W;
Tyrosine	Y; and	Valine	V.

As used herein, the term “peptide” refers to a compound of two or more native and non-native amino acids or amino acid derivatives such as α -amino acids linked by a peptide bond. As used herein, the term “glucagon derivative” refers to a peptide including the sequence of General Formula 1 or a derivative thereof, an analog, or a modified product thereof. The peptide as described herein includes peptidomimetics, which have a change in pI compared to that of native glucagon by modifying a part of the amino acid(s) of glucagon in the form of a substitution, etc. In an exemplary example, the glucagon derivative has an isoelectric point different from that of native glucagon while maintaining the activity of a glucagon receptor. In a more specific exemplary example, the glucagon derivative refers to a peptide with improved solubility of native glucagon at a physiological pH while maintaining the glucagon receptor activity.

As used herein, the term “pI” or “isoelectric point” refers to the pH value at which a macromolecule such as a polypeptide has no net charge (0). In the case of a polypeptide with

various charged functional groups, the net charge of the total polypeptide is “0” at a point where the pH value is the same as that of pI. The net charge of the polypeptide at a pH higher than the pI will be negative while the net charge of the polypeptide at a pH lower than the pI will be positive.

The pI values may be measured or estimated by a conventional method used in the art. For example, the pI values may be measured on an immobilized pH gradient gel consisting of polyacrylamide, starch, or agarose by isoelectric electrophoresis, or, for example, may be estimated from an amino acid sequence using a pI/MW tool (http://expasy.org/tools/pi_tool.html; Gasteiger et al., 2003) in an ExPASy server.

In a specific example, the glucagon derivative containing an amino acid sequence of General Formula 1 encompasses any peptide that is prepared by the substitution, addition, deletion, or post-translational modification (e.g., methylation, acylation, ubiquitination, or intramolecular covalent bonding) of amino acid(s) in the amino acid sequence of native glucagon represented by SEQ ID NO: 1, which exhibits improved solubility according to the pH of a solution due to the difference of its pI from that of native glucagon while maintaining the glucagon receptor activities as they are, thereby having improved *in vivo* chemical stability.

During the substitution or addition of amino acids, not only the 20 amino acids commonly found in human proteins, but also atypical or non-naturally occurring amino acids and derivatives thereof can be used. Commercial sources of atypical amino acids may include Sigma-Aldrich, ChemPep Inc., Genzyme Pharmaceuticals, etc. The peptides including these amino acids and atypical peptide sequences may be synthesized and purchased from commercial suppliers, e.g., American Peptide Company, Bachem (USA), or Anygen (Korea). Amino acid derivatives, e.g., desamino-histidine, β -hydroxyimidazopropionic acid, 4-imidazoacetic acid, or β -carboxyimidazopropionic acid, may be obtained in the same manner.

Since glucagon has a pI of about 7, it is insoluble in a solution having a pH of 6 to 8 and tends to precipitate at a neutral pH. In an aqueous solution with a pH of 3 or below, glucagon is dissolved at the initial stage but precipitates within one hour by forming a gel.

Since the gelated glucagon mainly consists of β -sheet fibrils, the administration of the thus-precipitated glucagon via an injection needle or intravenous injection will block blood vessels, and thus is not suitable for use as an injection agent. In order to delay the precipitation process, acidic (pH of 2 to 4) formulations are commonly used, and by doing so, glucagon can be maintained in a relatively non-aggregated state for a short period of time. However, glucagon can form fibrils very rapidly at a low pH, and thus these acidic formulations must be injected upon preparation.

It is widely known in the art that the solubility, activity, and stability of a protein in a solution can vary according to pI (Shaw, K. L. et al., Protein Science 10, pp 1206-1215, 2001).

As such, the present inventors have developed glucagon derivatives with extended stability and functional effects by modifying the sequence of native glucagon, thereby altering its pI. The glucagon derivatives of the present invention, by having an altered pI compared to that of native glucagon, are characterized in that they have improved solubility and stability according to the pH of a given solution, compared to that of native glucagon.

In a specific example, the glucagon derivative may be a peptide in which, in the amino acid sequence of General Formula 1,

X1 is histidine or tryptophan, tyrosine, or deleted;

X2 is serine or aminoisobutyric acid (Aib);

X7 is threonine or valine;

X12 is lysine or cysteine;

X13 is tyrosine or cysteine;

X14 is leucine or cysteine;

X15 is aspartic acid or cysteine;

X16 is glutamic acid, aspartic acid, serine, or cysteine;

X17 is aspartic acid, glutamic acid, lysine, arginine, valine, or cysteine;

X18 is aspartic acid, glutamic acid, arginine, or cysteine;

X19 is alanine or cysteine;

X20 is lysine, glutamic acid, glutamine, aspartic acid, lysine, or cysteine;

X21 is aspartic acid, glutamic acid, valine, or cysteine;

X23 is valine or arginine;

X24 is valine, leucine, or glutamine;

X27 is isoleucine or methionine; and

X28 is arginine or asparagine

(with the proviso that when the amino acid sequence of General Formula 1 is identical to SEQ ID NO: 1, it is excluded).

More preferably, the glucagon derivative as described herein may be a peptide including any one amino acid sequence among the amino acid sequences represented by SEQ ID NOS: 2 to 34.

The peptide including the glucagon derivative as described herein may be prepared by a standard synthesis method, a recombinant expression system, or any other method in the art. Accordingly, the glucagon analog as described herein may be synthesized by many methods including the methods described below:

(a) a method of synthesizing a peptide by a solid-phase or liquid-phase method stepwise or by fragment assembly, followed by isolation and purification of the final peptide product; or

(b) a method of expressing a nucleic acid construct encoding a peptide in a host cell and recovering the expression product from the host cell culture; or

(c) a method of performing an *in vitro* cell-free expression of a nucleic acid construct encoding a peptide and recovering the expression product therefrom; or

(d) a method of obtaining peptide fragments by any combination of the methods (a), (b), and (c), obtaining the peptide by linking the peptide fragments, and then recovering the peptide.

In an exemplary example, it was confirmed that the glucagon derivative of the present invention has a different pI compared to that of native glucagon (see Table 1). As a result, the glucagon derivative as described herein has improved solubility and higher stability according to the pH of a given solution, compared to that of native glucagon. Accordingly, the glucagon derivative as described herein can increase patient compliance when used as a hypoglycemic agent and is also suitable for the combined administration to be administered in combination with other anti-obesity agents, and thus can be effectively used for the prevention and treatment of hypoglycemia and obesity.

In this regard, the glucagon derivative as described herein can provide an attractive therapeutic selection regarding hypoglycemia, obesity, or associated diseases thereof.

For example, the glucagon derivative as described herein is a major insulin response-controlling hormone, and can be effectively used for the treatment of severe hypoglycemia in diabetic patients.

Additionally, the glucagon derivative as described herein may be used as a pharmaceutical drug not only for preventing body weight increase, promotion of body weight decrease, reduction of overweight, and obesities including morbid obesity (e.g., by controlling appetite, ingestion, food intake, calorie intake, and/or energy consumption), but also for treating obesity-related inflammation, obesity-related gallbladder disease, and obesity-induced sleep apnea, but is not limited thereto, and may be used for treating the associated diseases or health conditions thereof. The glucagon derivative as described herein may also be used for treating the health conditions that may be associated with obesity, such as metabolic syndrome, hypertension, atherosclerosis-induced dyslipidemia, arteriosclerosis, arterial chlerosis, coronary heart disease, strokes, etc. However, the effects of the glucagon derivative according to the present invention may be mediated entirely or partially by the body weight-related effects described above or may be independent of the same.

Meanwhile, for the improvement of the therapeutic effects of the glucagon derivative as described herein, the glucagon derivative may be modified using the typical techniques in the art, including a modification with polymers such as polyethylene glycol (PEG) and sugar chains, or a fusion with albumin, transferrin, fatty acid, and immunoglobulin, etc. For example, at least one amino acid side chain within the compounds as described herein may be attached to a polymer in order to increase *in vivo* solubility and/or half-lives, and/or increase bioavailabilities thereof. These modifications are known to reduce the clearance of therapeutic proteins and peptides.

For these polymers, soluble (amphipathic or hydrophilic), non-toxic, and pharmaceutically inert polymers are appropriate, and preferably, they may include PEG, homopolymers or copolymers of PEG, monomethyl-substituted polymers (mPEG), and poly-amino acids such as poly-lysine, poly-aspartic acid, and poly-glutamic acid.

In another example, described herein are polynucleotides encoding the glucagon derivatives described above.

The term “homology”, as used herein for the polynucleotide, indicates the degree of similarity to a wild-type amino acid sequence and a wild-type nucleic acid sequence, and includes a gene sequence that is 75% or higher, preferably 85% or higher, more preferably 90% or higher, and even more preferably 95% or higher identical to the polynucleotide sequence encoding the glucagon derivatives. The homology evaluation may be done with the naked eye or using a commercially available program. Using a commercially available computer program, the homology between two or more sequences may be expressed as a percentage (%), and the homology (%) between adjacent sequences may be evaluated. The polynucleotide encoding the glucagon derivatives may be inserted into a vector and expressed to obtain a large amount of the glucagon derivatives.

For these recombinant expressions, the polynucleotides as described herein are generally inserted into appropriate vectors to construct cloning vectors or recombinant vectors containing these polynucleotides, and these vectors also belong to the scope of the present invention.

As used herein, the term “recombinant vector” refers to a DNA construct including the sequence of a polynucleotide encoding a target peptide, which is operably linked to an appropriate regulatory sequence to enable the expression of the target peptide, e.g., a glucagon derivative, in a host cell. The regulatory sequence includes a promoter capable of initiating transcription, any operator sequence for the regulation of the transcription, a sequence encoding an appropriate mRNA ribosome-binding domain, and a sequence regulating the termination of transcription and translation. The recombinant vector, after being transformed into a suitable host cell, may be replicated or function irrespective of the host genome, or may be integrated into the host genome itself.

The recombinant vector used herein may not be particularly limited as long as the vector is replicable in the host cell, and it may be constructed using any vector known in the art. Examples of the vector may include natural or recombinant plasmids, cosmids, viruses, and bacteriophages. For example, as a phage vector or cosmid vector, pWE15, M13, MBL3, MBL4, IXII, ASHII, APII, t10, t11, Charon4A, Charon21A, etc., may be used; and as a

plasmid vector, those based on pBR, pUC, pBluescriptII, pGEM, pTZ, pCL, pET, etc., may be used. The vectors that can be used herein are not particularly limited but any known expression vector may be used.

The recombinant vector is used for the transformation of a host cell for producing glucagon derivatives as described herein. Additionally, these transformed cells, may be used for the amplification of nucleic acid fragments and vectors, or may be cultured cells or cell lines used in the recombinant production of glucagon derivatives of the present invention.

As used herein, the term “transformation” refers to a process of introducing a recombinant vector including a polynucleotide encoding a target protein into a host cell, thereby enabling the expression of the polynucleotide encoded by the protein in the host cell. For the transformed polynucleotide, it does not matter whether it is inserted into the chromosome of a host cell and located therein or located outside the chromosome, as long as it can be expressed in the host cell, and both cases are included.

Additionally, the polynucleotide includes DNA and RNA which encode the target protein. The polynucleotide may be inserted in any form insofar as it can be introduced into a host cell and expressed therein. For example, the polynucleotide may be introduced into a host cell in the form of an expression cassette. The expression cassette may conventionally include a promoter operably linked to the polynucleotide, a transcription termination signal, a ribosome-binding domain, and a translation termination signal. The expression cassette may be in the form of an expression vector capable of self-replication. Additionally, the polynucleotide may be introduced into a host cell as it is and operably linked to a sequence essential for its expression in the host cell, but is not limited thereto. Additionally, as used herein, the term “operably linked” refers to a functional connection between a promoter sequence, which initiates and mediates the transcription of the polynucleotide encoding the target protein as described herein, and the above gene sequence.

An appropriate host to be used herein may not be particularly limited as long as it can express the polynucleotide of the present invention. Examples of the appropriate host may include bacteria belonging to the genus *Escherichia* such as *E. coli*, bacteria belonging to the genus *Bacillus* such as *Bacillus subtilis*, bacteria belonging to the genus *Pseudomonas* such as *Pseudomonas putida*, yeasts such as *Pichia pastoris*, *Saccharomyces cerevisiae*, and

Schizosaccharomyces pombe, insect cells such as *Spodoptera frugiperda* (Sf9), and animal cells such as CHO, COS, and BSC.

In another example, described herein is a pharmaceutical composition for preventing or treating hypoglycemia or obesity containing the glucagon derivatives.

As used herein, the term “prevention” refers to any action resulting in suppression or delay of the onset of obesity by the administration of the glucagon derivatives or the pharmaceutical composition, and the term “treatment” refers to any action resulting in improvement in symptoms of obesity or the beneficial alteration by the administration of the glucagon derivatives or the pharmaceutical composition.

As used herein, the term “administration” refers to introduction of a particular material to a patient by an appropriate manner, and the composition may be administered via any of the common routes as long as the composition can arrive at a target tissue. For example, administration may be performed intraperitoneally, intravenously, intramuscularly, subcutaneously, intradermally, orally, topically, intranasally, intrapulmonarily, intrarectally, etc.

As used herein, the term “hypoglycemia” refers to an acute symptom of diabetes, in which blood glucose levels are lower than those of normal people, and in general, refers to a state when the blood glucose levels are 50 mg/dL or less. Hypoglycemia is frequently caused when a person who takes an oral hypoglycemic agent has eaten less than usual or has performed activities or exercised more than usual. In addition, hypoglycemia may occur due to the use of glucose level-lowering drugs, severe physical diseases, hormone deficiency such as adrenocortical hormones and glucagon, tumor in insulin-producing pancreas, insulin autoimmune syndrome, gastrectomy patients, inborn error of carbohydrate metabolism disorder, etc.

Symptoms of hypoglycemia include weakness, trembling, pale skin, cold sweat, dizziness, excitement, anxiety, pounding heart, empty stomach, headache, fatigue, etc. In the case of persistent hypoglycemia, it may lead to convulsion or seizure, and may cause shock and thus fainting.

As used herein, the term “obesity” refers to a medical condition with excess body fat in the body, and when a person having a body mass index (BMI; body mass (kg) divided by the square of the body height (m)) of 25 or higher is diagnosed as having obesity. Obesity generally

occurs due to a long-term energy imbalance in which energy intake exceeds energy expenditure. Obesity is a metabolic disease that affects the entire body, which increases the risk of diabetes, hyperlipidemia, sexual dysfunction, arthritis, and cardiovascular disease, and in some cases, it is also associated with the occurrence of cancers.

The pharmaceutical composition as described herein may contain a pharmaceutically acceptable carrier, excipient, or diluent. As used herein, the term “pharmaceutically acceptable” refers to the properties of having a sufficient amount to exhibit a therapeutic effect and not causing adverse effects, and may be easily determined by a skilled person in the art based on the factors well known in the medical field, such as the kind of disease, age, body weight, health status, sex, drug sensitivity of a patient, administration route, administration method, administration frequency, duration of treatment, a drug to be mixed or administered simultaneously in combination, etc.

The pharmaceutical composition of the present invention containing the glucagon derivative of the present invention may further contain a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier may include, for oral administration, a binder, a glidant, a disintegrant, an excipient, a solubilizing agent, a dispersant, a stabilizing agent, a suspending agent, a coloring agent, a flavoring agent, etc.; for injections, a buffering agent, a preserving agent, an analgesic, a solubilizing agent, an isotonic agent, a stabilizing agent, etc., which may be combined to be used; and for topical administrations, a base, an excipient, a lubricant, a preserving agent, etc., although it is not limited thereto.

The formulation type of the composition as described herein may be prepared variously by combining with a pharmaceutically acceptable carrier as described above. For example, for oral administration, the composition may be formulated into tablets, troches, capsules, elixirs, suspensions, syrups, wafers, etc. For injections, the composition may be formulated into single-dose ampoules or multidose containers. The composition may be also formulated into solutions, suspensions, tablets, capsules, and sustained-release formulations.

Meanwhile, examples of suitable carriers, excipients, and diluents may include lactose, dextrose, sucrose, sorbitol, mannitol, xylitol, erythritol, maltitol, starch, acacia rubber, alginate, gelatin, calcium phosphate, calcium silicate, cellulose, methyl cellulose, microcrystalline cellulose, polyvinylpyrrolidone, water, methyl hydroxybenzoate, propyl hydroxybenzoate, talc, magnesium stearate, mineral oil, etc. Additionally, the composition may further contain a filler,

an anti-coagulant, a lubricant, a humectant, a flavoring agent, a preservative, etc.

Additionally, the pharmaceutical composition as described herein may be prepared in any formulation type selected from the group consisting of tablets, pills, powders, granules, capsules, suspensions, liquid medicine for internal use, emulsions, syrups, sterile injection solutions, non-aqueous solvents, lyophilized formulations, and suppositories.

Additionally, the composition may be formulated into a single dosage form suitable for the patient's body, and preferably is formulated into a preparation useful for peptide drugs according to the typical method in the pharmaceutical field so as to be administered by an oral or parenteral route, such as through skin, intravenously, intramuscularly, intra-arterially, intramedullarily, intrathecally, intraventricularly, pulmonarily, transdermally, subcutaneously, intraperitoneally, intranasally, intragastrically, topically, sublingually, vaginally, or rectally, but is not limited thereto.

Additionally, the glucagon derivative may be used by blending with a variety of pharmaceutically acceptable carriers such as physiological saline or organic solvents. In order to increase the stability or absorptivity, carbohydrates such as glucose, sucrose, or dextrans, antioxidants such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins, or other stabilizers may be used.

The administration dose and frequency of the pharmaceutical composition as described herein are determined by the type of active ingredient(s), together with various factors such as the disease to be treated, administration route, patient's age, gender, and body weight, and severity of the disease.

The total effective dose of the composition as described herein may be administered to a patient in a single dose, or may be administered for a long period of time in multiple doses according to a fractionated treatment protocol. In the pharmaceutical composition as described herein, the content of active ingredient(s) may vary depending on the disease severity. Preferably, the total daily dose of the peptide as described herein may be approximately 0.0001 µg to 500 mg per 1 kg of body weight of a patient. However, the effective dose of the glucagon derivative is determined considering various factors including patient's age, body weight, health conditions, gender, disease severity, diet, and excretion rate, in addition to administration route and treatment frequency of the pharmaceutical composition. In this regard,

those skilled in the art may easily determine the effective dose suitable for the particular use of the pharmaceutical composition as described herein. The pharmaceutical composition as described herein is not particularly limited to the formulation and administration route and mode, as long as it shows the effects as described herein.

The pharmaceutical composition as described herein shows excellent *in vivo* duration of efficacy and titer, thereby remarkably reducing the number and frequency of administration thereof.

In particular, since the pharmaceutical composition as described herein contains, as an active ingredient, a glucagon derivative having an altered pI different from that of native glucagon, it shows improved solubility and high stability according to the pH of a given solution, and thus the pharmaceutical composition as described herein can be effectively used in the preparation of a stable glucagon formulation for treating hypoglycemia or obesity.

Furthermore, the pharmaceutical composition as described herein may be administered alone or in combination with other pharmaceutical formulation(s) showing prophylactic or therapeutic effects on obesity. Additionally, the pharmaceutical composition as described herein may further contain a pharmaceutical formulation showing prophylactic or therapeutic effects on obesity.

The pharmaceutical formulations showing prophylactic or therapeutic effects on obesity are not particularly limited, and may include a GLP-1 receptor agonist, a glucose-dependent insulintropic peptide (GIP) receptor antagonist, a leptin receptor agonist, a DPP-IV inhibitor, a Y5 receptor antagonist, a melanin-concentrating hormone (MCH) receptor antagonist, a Y2/3/4 receptor agonist, an MC3/4 receptor agonist, a gastric/pancreatic lipase inhibitor, a 5HT2c agonist, a β 3A receptor agonist, an amylin receptor agonist, a ghrelin antagonist and a ghrelin receptor antagonist, FGF1, an FGF21 receptor agonist, a cholecystokinin (CCK) receptor agonist, a pancreatic polypeptide (PP) receptor agonist, a dopamine reabsorption inhibitor, etc.

In still another example, the present disclosure provides a method for preventing or treating hypoglycemia or obesity, including administration of the glucagon derivative or the pharmaceutical composition containing the same to a subject.

As used herein, the term "subject" refers to those suspected of having hypoglycemia or

obesity, which are mammals including humans, mice, and livestock animals having hypoglycemia or obesity, or with the possibility of having hypoglycemia or obesity. However, any subject to be treated with the glucagon derivatives or the pharmaceutical composition as described herein is included without limitation. The pharmaceutical composition containing the glucagon derivative as described herein may be administered to a subject suspected of having hypoglycemia or obesity, thereby treating the subject effectively. The hypoglycemia and obesity are the same as described above.

The therapeutic method as described herein may include administration of the pharmaceutical composition containing the glucagon derivative at a pharmaceutically effective amount. Preferably, the total daily dose should be determined based on appropriate medical judgment by a physician and administered once or several times. In view of the objects of the present invention, the specific therapeutically effective dose for any particular patient may vary depending on various factors, such as the kind and degree of the response to be achieved, specific compositions including whether other agent(s) is(are) used therewith or not, the patient's age, body weight, health condition, gender, and diet, the time and route of administration, the secretion rate of the composition, duration of therapy, other drug(s) used in combination or simultaneously with the specific compositions, and similar factors well known in the medical art.

In still another example, the present disclosure provides a use of the glucagon derivatives in the preparation of pharmaceutical drugs for the prevention or treatment of hypoglycemia or obesity.

[Advantageous Effects]

The novel glucagon derivatives of the present invention have excellent stability and solubility according to the pH of a given solution by having a pI different from that of native glucagon. Therefore, when the novel glucagon derivatives of the present invention are used as a therapeutic agent for treating hypoglycemia, they can increase patient compliance. Additionally, the novel glucagon derivatives of the present invention are suitable for administration in combination with other anti-obesity agents, and thus they can be effectively used for preventing or treating hypoglycemia and obesity.

[CLAIMS]

[Claim 1]

A glucagon derivative, which comprises the amino acid sequence of the following General Formula 1 and has a different isoelectric point (pI) from that of native glucagon:

X1-X2-QGTF-X7-SDYS-X12-X13-X14-X15-X16-X17-X18-X19-X20-X21-F-X23-X24-W-L-X27-X28-T (General Formula 1)

wherein, in General Formula 1,

X1 is histidine, desamino-histidine, N-dimethyl-histidine, β -hydroxyimidazopropionic acid, 4-imidazoacetic acid, β -carboxyimidazopropionic acid, tryptophan, tyrosine, or deleted;

X2 is α -methyl-glutamic acid, aminoisobutyric acid (Aib), D-alanine, glycine, Sar(N-methylglycine), serine, or D-serine;

X7 is threonine or valine;

X12 is lysine or cysteine;

X13 is tyrosine or cysteine;

X14 is leucine or cysteine;

X15 is aspartic acid, glutamic acid, or cysteine;

X16 is glutamic acid, aspartic acid, serine, α -methyl-glutamic acid, or cysteine;

X17 is aspartic acid, glutamine, glutamic acid, lysine, arginine, serine, valine, or cysteine;

X18 is aspartic acid, glutamine, glutamic acid, arginine, or cysteine;

X19 is alanine or cysteine;

X20 is lysine, glutamic acid, glutamine, aspartic acid, lysine, or cysteine;

X21 is aspartic acid, glutamic acid, valine, or cysteine;

X23 is valine or arginine;

X24 is valine, leucine, glutamine, or arginine;

X27 is isoleucine or methionine; and

X28 is arginine or asparagine

(with the proviso that when the amino acid sequence of General Formula 1 is identical to SEQ ID NO: 1 or SEQ ID NO: 12, it is excluded).

[Claim 2]

The glucagon derivative of claim 1, wherein, in the amino acid sequence of General Formula 1,

X1 is histidine or tryptophan, tyrosine, or deleted;

X2 is serine or aminoisobutyric acid (Aib);

X7 is threonine or valine;

X12 is lysine or cysteine;

X13 is tyrosine or cysteine;

X14 is leucine or cysteine;

X15 is aspartic acid or cysteine;

X16 is glutamic acid, aspartic acid, serine, or cysteine;

X17 is aspartic acid, glutamic acid, lysine, arginine, valine, or cysteine;

X18 is aspartic acid, glutamic acid, arginine, or cysteine;

X19 is alanine or cysteine;

X20 is lysine, glutamic acid, glutamine, aspartic acid, lysine, or cysteine;

X21 is aspartic acid, glutamic acid, valine, or cysteine;

X23 is valine or arginine;

X24 is valine, leucine, or glutamine;

X27 is isoleucine or methionine; and

X28 is arginine or asparagine

(with the proviso that when the amino acid sequence of General Formula 1 is identical to SEQ ID NO: 1 or SEQ ID NO: 12, it is excluded).

[Claim 3]

The glucagon derivative of claim 1, wherein the glucagon derivative has glucagon receptor-stimulating activity.

[Claim 4]

The glucagon derivative of claim 1, wherein the glucagon derivative comprises an amino acid sequence selected from the group consisting of amino acid sequences represented by SEQ

ID NOS: 2 to 11 and 13 to 34.

[Claim 5]

A polynucleotide encoding the glucagon derivative according to any one of claims 1 to 4.

[Claim 6]

Use of a glucagon derivative according to any one of claims 1 to 4 as an active ingredient in the manufacture of a pharmaceutical composition for preventing or treating hypoglycemia.

[Claim 7]

Use of a glucagon derivative according to any one of claims 1 to 4 as an active ingredient in the manufacture of a pharmaceutical composition for preventing or treating obesity.

[Claim 8]

The use according to claim 6, further comprising a pharmaceutically acceptable carrier.

[Claim 9]

The use according to claim 7, further comprising a pharmaceutically acceptable carrier.

[Claim 10]

The use according to claim 7, wherein the pharmaceutical composition is to be administered alone or in combination with a pharmaceutical agent having the effect of preventing or treating obesity.

[Claim 11]

The use according to claim 7, further comprising a pharmaceutical agent having the effect of preventing or treating obesity.

[Claim 12]