

FIGURE 1




FIGURE 2A




FIGURE 2B




FIGURE 3




FIGURE 4




FIGURE 5A




FIGURE 5B




FIGURE 5C





FIGURE 5D

Pre-Dose C5 Levels in Mice Expressing Human C5



Overall Average: 48.98µg/mL (± 15.09 S.D.)

Female Average: 28.80µg/mL (± 6.17 S.D.)

Male Average (excluding #25): 58.46µg/mL (± 9.98 S.D.)

FIGURE 6




FIGURE 7




FIGURE 8A (CONT)




FIGURE 8B




FIGURE 8B (CONT)

Humanized C5 mouse serum treated with anti-human C5 mAb
Dilution 1:5 at different time points




FIGURE 9A




FIGURE 9B




FIGURE 9B (CONT)




FIGURE 10A




FIGURE 10B





FIGURE 11

CH₅₀ assay for classical pathway hemolytic activity



- C3 KO
- C3KO + C3 100ug/ml
- △ C3KO + C3 500ug/ml
- ◇ C3KO + C3 1200ug/ml
- WT
- ▨ C3depl Hum
- ▩ C3depl Hum + 100ug/ml
- ▧ C3depl Hum + 500ug/ml
- ▦ C3depl Hum + 1200ug/ml

	Human (ug/ml)	Mouse (ug/ml)
C3	1200	150
C5	50	90

FIGURE 12

