


**CITRIC ACID SALT OF A THERAPEUTIC COMPOUND AND PHARMACEUTICAL
COMPOSITIONS THEREOF**

The present invention is directed to a polymorph of the citric acid salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one:



and pharmaceutical compositions thereof.

The compound, 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one, is an antagonist of the serotonin-1D (5-HT_{1D}) receptor and is useful in the treatment of a number of disorders, diseases and conditions of the central nervous system. This compound is particularly useful in the treatment of hypertension, all forms of depression, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, post partum depression, dysthymia; mild, moderate, or severe depressions with or without atypical features, melancholic features, psychotic features, catatonic features; seasonal affective disorder, geriatric depression, chronic depression; adjustment disorder with depressed mood or with anxiety and depressed mood; mixed anxiety and depression; substance induced mood disorder; and mood disorder secondary to a general medical condition, bipolar disorder, bipolar disorder-depressed phase, generalized anxiety disorder, phobias, agoraphobia, social anxiety, social phobia, simple phobias, separation anxiety disorder, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, eating disorders, binge eating disorder, anorexia nervosa, bulimia nervosa, obesity; chemical dependencies and addictions to alcohol, cocaine, heroin, phenobarbital, nicotine, marijuana and benzodiazepines; cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, panic disorder with agoraphobia, memory disorders, dementia, amnesic disorders, and age-related cognitive decline (ARCD), Parkinson's diseases, dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias, endocrine disorders, hyperprolactinaemia, vasospasm, vasospasm in the cerebral vasculature, cerebellar ataxia; gastrointestinal tract disorders involving changes in motility and secretion; negative symptoms of schizophrenia, premenstrual

-2-

syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, cancer, small cell lung carcinoma, chronic paroxysmal hemicrania, headache associated with vascular disorders, autism, pervasive developmental disorder NOS, Asperger's disorder, selective mutism, chronic motor or vocal tic disorder, somatization disorder, insomnia, intermittent explosive disorder, pyromania, pathological gambling, impulse-control disorder, premenstrual dysphoric disorder and attention-deficit/hyperactivity disorder (ADHD) in a mammal, particularly a human. The citrate salt of this invention may also be used in a pharmaceutical composition in combination with a serotonin reuptake inhibiting antidepressant (SRI), in order to treat a number of these conditions.

Compounds that are antagonists of the serotonin-1D receptor, including 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one, specifically including its hydrochloride salt, are referred to in WO 98/14433, published April 9, 1998 (corresponding to U.S. Ser. No. 09/254,999, filed September 8, 1997, 09/733346, filed December 8, 2000 and PCT/IB01/02139, filed November 12, 2001). The foregoing applications, owned in common with the present application, generically recite pharmaceutically acceptable acid addition salts for the compounds referred to therein.

The citrate salt of the present invention exhibits properties, including those of solid-state stability and compatibility with certain drug product formulation excipients, that render it superior to previously known salts of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is the differential scanning calorimetric (DSC) trace of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.

Figure 2 is the observed powder X-ray diffraction pattern of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one (y axis is linear counts per second; X in degrees 2 theta).

Figure 3 is the calculated powder X-ray diffraction pattern of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one (y axis is linear counts per second; X in degrees 2 theta).

Figure 4 is the X-ray crystal structure of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.

Figure 5 is the ¹³C NMR spectrum of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one in the solid phase as measured by cross-polarization magic angle spinning (CPMAS) at 295 K on a Bruker 7mm wide-bore magic angle spinning (WB MAS) probe positioned in a Bruker Avance DRX 500 MHz NMR Spectrometer.

SUMMARY OF THE INVENTION

A first aspect of the present invention provides the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.

The citrate salt of one embodiment can be an anhydrous or nearly anhydrous polymorph.

This citrate salt can also be characterized by the principal x-ray diffraction pattern peaks expressed in terms of 2θ and d-spacings as measured with copper radiation (within the margins of error indicated):

Angle 2θ (± 0.2)	d-value ($\text{\AA}$) (± 0.2)
13.0	6.8
17.4	5.1
18.0	4.9
18.9	4.7
20.0	4.4
21.2	4.2
22.2	4.0
24.0	3.7
27.1	3.3
32.4	2.8

The citrate salt can be characterized in that it generally forms flakes. Further, the citrate salt may also be characterized in that it can form monoclinic crystals belonging to the Pc space group. The citrate salt may further be characterized in having an onset of melting transition/decomposition point at about 198-199 °C as measured by differential scanning calorimetry (DSC). Further, the citrate salt can also be characterized in having an aqueous solubility of 1.3 mg/ml and a native pH of 3.37 in aqueous solution. In addition, the citrate salt has a hygroscopicity of approximately 1.27 % at 90% relative humidity.

The citrate salt of one embodiment is also characterized in that when examined by solid state ^{13}C NMR cross-polarization magic angle spinning techniques it exhibits the following principal resonance peaks downfield from 100 parts per million (± 0.1 ppm; relative to an adamantane standard at 29.5 ppm): δ 179.3, 177.0, 171.6, 164.0, 151.0 and 144.1.

A second aspect of the present invention provides a pharmaceutical composition comprising a compound according to the first aspect and a pharmaceutically acceptable carrier.

-3a-

An embodiment can provide a pharmaceutical composition comprising the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one, and a pharmaceutically acceptable carrier or excipient, particularly, one for use in the treatment of hypertension, all forms of depression, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, post partum depression, dysthymia; mild, moderate, or severe depressions with or without atypical

- features, melancholic features, psychotic features, catatonic features; seasonal affective disorder, geriatric depression, chronic depression; adjustment disorder with depressed mood or with anxiety and depressed mood; mixed anxiety and depression; substance induced mood disorder; and mood disorder secondary to a general medical condition, bipolar disorder,
- 5 bipolar disorder-depressed phase, generalized anxiety disorder, phobias, agoraphobia, social anxiety, social phobia, simple phobias, separation anxiety disorder, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, eating disorders, binge eating disorder, anorexia nervosa, bulimia nervosa, obesity; chemical dependencies and addictions to alcohol, cocaine, heroin, phenobarbital, nicotine, marijuana and benzodiazepines; cluster
- 10 headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, panic disorder with agoraphobia, memory disorders, dementia, amnesic disorders, and age-related cognitive decline (ARCD), Parkinson's diseases, dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias, endocrine disorders, hyperprolactinaemia, vasospasm, vasospasm in the cerebral vasculature, cerebellar ataxia;
- 15 gastrointestinal tract disorders involving changes in motility and secretion; negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, cancer, small cell lung carcinoma, chronic paroxysmal hemicrania, headache associated with vascular disorders, autism, pervasive developmental disorder NOS, Asperger's disorder, selective mutism,
- 20 chronic motor or vocal tic disorder, somatization disorder, insomnia, intermittent explosive disorder, pyromania, pathological gambling, impulse-control disorder, premenstrual dysphoric disorder and attention-deficit/hyperactivity disorder (ADHD) in a mammal, preferably a human.

There can be provided a method of treating hypertension, all forms of depression, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction

25 depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, post partum depression, dysthymia; mild, moderate, or severe depressions with or without atypical features, melancholic features, psychotic features, catatonic features; seasonal affective disorder, geriatric depression, chronic depression;

30 adjustment disorder with depressed mood or with anxiety and depressed mood; mixed anxiety and depression; substance induced mood disorder; and mood disorder secondary to a general medical condition, bipolar disorder, bipolar disorder-depressed phase, generalized anxiety disorder, phobias, agoraphobia, social anxiety, social phobia, simple phobias, separation anxiety disorder, posttraumatic stress syndrome, avoidant personality disorder,

35 premature ejaculation, eating disorders, binge eating disorder, anorexia nervosa, bulimia nervosa, obesity; chemical dependencies and addictions to alcohol, cocaine, heroin, phenobarbital, nicotine, marijuana and benzodiazepines; cluster headache, migraine, pain,

Alzheimer's disease, obsessive-compulsive disorder, panic disorder, panic disorder with agoraphobia, memory disorders, dementia, amnesic disorders, and age-related cognitive decline (ARCD), Parkinson's diseases, dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias, endocrine disorders, hyperprolactinaemia, vasospasm, vasospasm in the cerebral vasculature, cerebellar ataxia; gastrointestinal tract disorders involving changes in motility and secretion; negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, kleptomania, male impotence, cancer, small cell lung carcinoma, chronic paroxysmal hemicrania, headache associated with vascular disorders, autism, pervasive developmental disorder NOS, Asperger's disorder, selective mutism, chronic motor or vocal tic disorder, somatization disorder, insomnia, intermittent explosive disorder, pyromania, pathological gambling, impulse-control disorder, premenstrual dysphoric disorder and attention-deficit/hyperactivity disorder (ADHD) in a mammal, preferably a human, comprising the administration of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one to the subject in need thereof.

There may also be provided a process for the preparation of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one comprising the steps of

- (i) contacting 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one dissolved in a suitable solvent with citric acid; and
- (ii) collecting the crystals formed.

A preferred embodiment is wherein the suitable solvent is selected from the group consisting of a (C₁-C₆)alkyl alcohol, a (C₁-C₆)alkyl ketone or a (C₁-C₆)alkyl ether. More preferably, the suitable solvent is 2-propanol. Preferably, the process of the invention is wherein the contacting of step (i) is carried out by contacting 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one in solution phase with a solution of citric acid. More preferably, contacting of step (i) is carried out by adding solid citric acid to the solution of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.

Preferably, the contacting step is carried out over a period of between 1 and 24 hours, more preferably between 10 and 20 hours, and comprising stirring or mixing the resulting mixture. A preferred embodiment of the process is wherein step (i) is run between ambient temperature and the refluxing temperature of the solvent; more preferably, between ambient temperature and the refluxing temperature of 2-propanol, i.e., about 80 °C; most preferably, the process is run between 30 and 60 °C. Preferably, the reaction mixture is allowed to cool to ambient temperature once the addition of citric acid is complete and permitted to stir for the remainder of the reaction period.

There may also be provided the citric acid salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one, prepared in accordance with the process as defined above.

A third aspect of the present invention provides the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one for use as a medicament.

A fourth aspect of the present invention provides the use the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one in the manufacture of a medicament for the treatment of hypertension, all forms of depression, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, post partum depression, dysthymia; mild, moderate, or severe depressions with or without atypical features, melancholic features, psychotic features, catatonic features; seasonal affective disorder, geriatric depression, chronic depression; adjustment disorder with depressed mood or with anxiety and depressed mood; mixed anxiety and depression; substance induced mood disorder; and mood disorder secondary to a general medical condition, bipolar disorder, bipolar disorder-depressed phase, generalized anxiety disorder, phobias, agoraphobia, social anxiety, social phobia, simple phobias, separation anxiety disorder, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, eating disorders, binge eating disorder, anorexia nervosa, bulimia nervosa, obesity; chemical dependencies and addictions to alcohol, cocaine, heroin, phenobarbital, nicotine, marijuana and benzodiazepines; cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, panic disorder with agoraphobia, memory disorders, dementia, amnestic disorders, and age-related cognitive decline (ARCD). Parkinson's diseases, dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias, endocrine disorders, hyperprolactinaemia, vasospasm, vasospasm in the cerebral vasculature, cerebellar ataxia; gastrointestinal tract disorders involving changes in motility and secretion; negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, lipothymia, male impotence,

cancer, small cell lung carcinoma, chronic paroxysmal hemicrania, headache associated with vascular disorders, autism, pervasive developmental disorder NOS, Asperger's disorder, selective mutism, chronic motor or vocal tic disorder, somatization disorder, insomnia, intermittent explosive disorder, pyromania, pathological gambling, impulse-control disorder, premenstrual dysphoric disorder and attention-deficit/hyperactivity disorder (ADHD) in a mammal.

DETAILED DESCRIPTION OF THE INVENTION

5 The compound, 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one is an antagonist of the serotonin-1D receptor (5-HT_{1D} receptor) and useful in the treatment of a number of CNS diseases, disorders and conditions. The free base of the compound and its hydrochloride salt may be prepared in accordance with the methods set forth in International Patent Publication No. WO 98/14433, published April 9,
10 1998 herein incorporated by reference in its entirety.

 The citrate salt may be prepared under a variety of different conditions. However, in accordance with the present invention, the free base of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one is preferably dissolved in a suitable solvent until completely dissolved, whereupon citric acid is added to the solution thereby
15 prepared to create the citrate addition salt of the invention. Preferably, the suitable solvent is selected from the group consisting of a (C₁-C₆)alkyl alcohol, a (C₁-C₆)alkyl ketone or a (C₁-C₆)alkyl ether; more preferably, a (C₁-C₆)alkyl alcohol; most preferably 2-propanol. Preferably, the process of the invention is wherein the contacting of step (i) is carried out by contacting
20 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one in solution phase with either a solution of citric acid or the solid form of citric acid.

 Preferably, the contacting step is carried out over a period of between 1 and 24 hours, more preferably between 10 and 20 hours, and comprising stirring or mixing the resulting mixture. A preferred embodiment of the process is wherein step (i) is run between ambient temperature and the refluxing temperature of the solvent; more preferably, between
25 ambient temperature and the refluxing temperature of 2-propanol, i.e., about 80 °C; most preferably, the process is run between 30 and 60 °C. Preferably, the reaction mixture is allowed to cool to ambient temperature once the addition of citric acid is complete and permitted to stir for the remainder of the reaction period.

 The citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one is only slightly hygroscopic and has high aqueous solubility. These characteristics combined with its relative inertness towards common excipients used in pharmaceutical formulation make it highly suitable for pharmaceutical formulation use.

 Although in general the known acid addition salts of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one are all crystalline, the majority of those
35 salts are so hygroscopic as to render them poor candidates for pharmaceutical formulation use. The citrate salt of the present invention exhibits a hygroscopicity of approximately 1.27 % wt/wt on exposure to 90% relative humidity in a moisture chamber. The aqueous solubility of

-7-

the citrate salt is 1.3 mg/ml having a pH of 3.37. Further, the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one exhibits excellent solid state stability both in light and elevated temperatures as well as high humidity challenges.

5 Differential Scanning Calorimetry

The solid state thermal behavior of the citrate salt of the invention was investigated by differential scanning calorimetry (DSC). The trace for the salt is shown in Figure 1. The DSC thermograms were obtained on a Mettler Toledo DSC 821^e (STAR^e System). Generally, samples between 1 and 10 mg were prepared in crimped aluminum pans with a small
10 pinhole. The measurements were run at a heating rate of 5 °C per minute in the range of 30 to 300 °C.

As seen in Figure 1, the citrate salt exhibits onset of melt transition at about 198-199 °C. One of skill in the art will however note that in DSC measurement there is a certain degree of variability in actual measured onset and peak temperatures which occur depending
15 on rate of heating, crystal shape and purity, and other measurement parameters.

Powder X-ray Diffraction Patterns

The power x-ray diffraction patterns for the citrate salt of the invention was collected using a Bruker D5000 diffractometer (Bruker AXS, Madison, Wisconsin) equipped with copper radiation CuK_α, fixed slits (1.0, 1.0, 0.6 mm), and a Kevex solid state detector. Data was
20 collected from 3.0 to 40.0 degrees in two theta (2θ) using a step size of 0.04 degrees and a step time of 1.0 seconds.

The x-ray powder diffraction pattern of the citrate salt was conducted with a copper anode with wavelength 1 at 1.54056 and wavelength 2 at 1.54439 (relative intensity: 0.500). The range for 2θ was between 3.0 to 40.0 degrees with a step size of 0.04 degrees, a step
25 time of 1.00 second, a smoothing width of 0.300 and a threshold of 1.0.

The diffraction peaks at diffraction angles (2θ) in a measured powder X-ray diffraction analysis for the salt are shown in Table I. The relative intensities, however, may change depending on the crystal size and morphology. The actual measured powder diffractogram is displayed in Figure 2.

Table I. Powder X-ray Diffraction Pattern for Citrate Salt with Intensities and Peak Locations of Diffraction Lines.

Angle 2 θ	d-value (Å)	I (rel.)	Angle 2 θ	d-value (Å)	I (rel.)	Angle 2 θ	d-value (Å)	I (rel.)
8.2	10.7	3.7	23.1	3.9	13.6	32.4	2.8	18.6
11.5	7.7	4.9	24.0	3.7	33.5	32.8	2.7	7.4
13.0	6.8	14.5	24.7	3.6	5.3	33.6	2.7	3.3
13.7	6.5	5.6	25.1	3.5	9.9	34.5	2.6	6.3
14.9	5.9	9.8	26.2	3.4	6.9	34.9	2.6	4.7
16.3	5.4	7.0	26.6	3.4	11.2	35.8	2.5	4.1
16.6	5.3	13.1	27.1	3.3	20.4	36.3	2.5	7.5
17.4	5.1	100.0	27.7	3.2	6.2	36.8	2.4	4.3
18.0	4.9	40.2	28.3	3.2	6.3	37.3	2.4	5.7
18.9	4.7	17.3	29.6	3.0	10.7	38.3	2.4	6.3
20.0	4.4	71.6	30.4	2.9	8.3	38.9	2.3	4.3
21.2	4.2	23.4	31.0	2.9	3.4			
22.2	4.0	20.0	31.6	2.8	2.8			

Table II sets forth the 2 θ , d-spacings and relative intensities and peak locations for the powder x-ray diffraction pattern representative for the citrate salt. The numbers as listed are computer-generated.

Table II. Powder X-ray Diffraction Intensities and Peak Locations Representative of the Citrate Salt.

Angle 2 θ	d-value (Å)	I (rel.)
13.0	6.8	14.5
17.4	5.1	100.0
18.0	4.9	40.2
18.9	4.7	17.3
20.0	4.4	71.6
21.2	4.2	23.4
22.2	4.0	20.0
24.0	3.7	33.5
27.1	3.3	20.4
32.4	2.8	18.6

Single Crystal X-ray Analysis

A single crystal for the citrate salt of the invention was obtained and investigated by x-ray diffraction. A representative crystal was surveyed and a 1Å data set (maximum $\sin \Theta/\lambda = 0.5$) was collected on a Siemens R4RA/v diffractometer. Atomic scattering factors were taken from the International Tables for X-Ray Crystallography, Vol. IV, pp. 55, 99, 149 (Birmingham: Kynoch Press, 1974). From the data gathered on the single crystal, a powder X-ray diffraction pattern was calculated to offer comparison against the actual measured diffraction pattern.

Structures were solved using direct methods. The SHELXTL™ computer library provided by Bruker AXS, Inc. facilitated all necessary crystallographic computations and molecular displays (SHELXTL™ Reference Manual, Version 5.1, Bruker AXS, Madison, WI 1997). Pertinent crystal, data collection, and refinement are summarized in Table III.

A trial structure was obtained by direct methods and was then refined routinely. Hydrogen positions were calculated wherever possible. The methyl hydrogens and the hydrogens on nitrogen and oxygen were located by difference Fourier techniques. The hydrogen parameters were added to the structure factor calculations but were not refined. The shifts calculated in the final cycles of least squares refinement were all less than 0.1 of the corresponding standard deviations. The final R-index was 4.72%. A final difference Fourier revealed no missing or misplaced electron density. The refined structure was plotted using the SHELXTL plotting package and is shown in Figure 4.

Table IV sets forth the atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for the salt. Table V lists the observed bond lengths [Å] and angles [°] for the citrate salt. In Table VI, the anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for the citrate salt are set forth to allow calculation of the anisotropic displacement factor exponent which has the form: $-2\pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$. Finally, in Table VII, below, hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for the salt are listed.

Table III. Crystal Structure Data And Measurement Parameters For The Citrate

Salt

Parameter	For Citrate Salt	
Empirical formula	$C_{22}H_{24}N_3OSCl_2^+C_6H_7O_7^-$	
Formula weight	640.52	
Temperature	293(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	Pc	
Unit cell dimensions	$a = 6.5940(10)$ Å	$\alpha = 90^\circ$.
	$b = 15.257(2)$ Å	$\beta = 101.440(10)^\circ$.
	$c = 15.099(2)$ Å	$\gamma = 90^\circ$.
Volume	1488.9(4) Å ³	
Z	2	
Density (calculated)	1.429 Mg/m ³	
Absorption coefficient	3.081 mm ⁻¹	
F(000)	668	
Crystal size	0.24 x 0.04 x 0.04 mm ³	
Reflections collected	1707	
Independent reflections	1707 [R(int) = 0.0000]	
Completeness to theta = 49.98°	100.0 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	1707 / 0 / 391	
Goodness-of-fit on F ²	1.065	
Final R indices [I > 2sigma(I)]	R1 = 0.0472, wR2 = 0.1028	
Absolute structure parameter	-0.01(3)	
Extinction coefficient	0.0038(6)	
Largest diff. peak and hole	0.223 and -0.268 e.Å ⁻³	

Table IV. Atomic coordinates ($\times 10^4$) And Equivalent Isotropic Displacement Parameters ($\text{\AA}^2 \times 10^3$) For The Citrate Salt. ($U(\text{eq})$ is defined as one third of the trace of the orthogonalized U_{ij} tensor.)

	x	y	x	U(eq)
S(1X)	1600	3502(2)	8401	53(1)
Cl(1X)	7950(8)	4063(2)	13244(3)	83(1)
Cl(2X)	11404(8)	2613(2)	13403(3)	73(1)
N(1)	8720(15)	1936(5)	6037(6)	42(2)
C(2)	9164(19)	2836(7)	5723(7)	50(3)
C(3)	7216(19)	3358(7)	5475(7)	48(3)
N(4)	6189(14)	3417(5)	6246(5)	37(2)
C(5)	5665(19)	2527(6)	6513(7)	44(3)
C(6)	7568(18)	1994(7)	6786(6)	41(3)
C(7)	4480(17)	4015(6)	6160(6)	34(3)
C(8)	3464(19)	4335(7)	5318(7)	45(3)
C(9)	1780(20)	4884(8)	5270(8)	57(4)
C(10)	1145(19)	5140(7)	6041(9)	55(3)
C(11)	2141(19)	4849(7)	6862(7)	47(3)
C(12)	3874(17)	4282(6)	6954(6)	36(3)
C(13)	5020(16)	4044(6)	7850(6)	35(3)
C(14)	4255(15)	3731(6)	8528(6)	32(3)
C(15)	5686(18)	3625(7)	9400(7)	41(3)
N(16)	5221(14)	3032(5)	10007(5)	38(2)
C(17)	3520(20)	2409(8)	9721(10)	77(4)
C(18)	1591(19)	2834(8)	9378(8)	58(3)
C(19)	6643(17)	2904(7)	10837(6)	33(3)
C(20)	8123(17)	2257(7)	10928(7)	40(3)
C(21)	9521(19)	2151(7)	11710(7)	51(3)
C(22)	9542(18)	2713(7)	12428(7)	45(3)
C(23)	8029(19)	3353(7)	12367(6)	43(3)
C(24)	6611(19)	3453(7)	11563(7)	45(3)
O(25)	7262(14)	4069(6)	9602(5)	70(3)
C(26)	10622(18)	1388(7)	6296(8)	57(3)
C(1X)	10057(19)	-243(7)	3822(8)	53(3)
O(2X)	11060(17)	-820(6)	3541(6)	100(4)
O(3X)	10378(16)	-38(6)	4658(6)	84(3)
C(4X)	8610(18)	320(6)	3209(6)	40(3)

WO 03/048141

PCT/IB02/04972

-12-

	x	y	x	U(eq)
C(5X)	6358(19)	214(6)	3317(7)	35(3)
O(6X)	5754(14)	-675(5)	3152(5)	52(2)
C(7X)	6105(16)	491(8)	4257(6)	39(3)
O(8X)	5421(15)	-95(6)	4709(5)	75(3)
O(9X)	6600(14)	1254(5)	4507(5)	55(2)
C(10X)	4897(18)	794(7)	2640(6)	46(3)
C(11X)	4463(19)	533(6)	1651(6)	32(2)
O(12X)	2826(13)	708(5)	1170(5)	50(2)
O(13X)	5938(13)	140(6)	1372(5)	64(3)

Table V. Observed Bond Lengths [Å] And Angles [°] For Citrate Salt.

S(1X)-C(14)	1.758(10)	C(15)-N(16)	1.365(12)
S(1X)-C(18)	1.794(11)	N(16)-C(19)	1.422(12)
Cl(1X)-C(23)	1.719(10)	N(16)-C(17)	1.470(13)
Cl(2X)-C(22)	1.727(10)	C(17)-C(18)	1.429(17)
N(1)-C(6)	1.486(12)	C(19)-C(20)	1.375(13)
N(1)-C(2)	1.500(12)	C(19)-C(24)	1.385(14)
N(1)-C(26)	1.493(13)	C(20)-C(21)	1.356(14)
C(2)-C(3)	1.495(16)	C(21)-C(22)	1.380(14)
C(3)-N(4)	1.461(12)	C(22)-C(23)	1.386(14)
N(4)-C(7)	1.435(13)	C(23)-C(24)	1.386(14)
N(4)-C(5)	1.478(12)	C(1X)-O(2X)	1.227(13)
C(5)-C(6)	1.483(14)	C(1X)-O(3X)	1.278(13)
C(7)-C(8)	1.402(13)	C(1X)-C(4X)	1.468(14)
C(7)-C(12)	1.398(13)	C(4X)-C(5X)	1.534(15)
C(8)-C(9)	1.384(16)	C(5X)-O(6X)	1.423(11)
C(9)-C(10)	1.369(15)	C(5X)-C(7X)	1.522(14)
C(10)-C(11)	1.357(14)	C(5X)-C(10X)	1.537(14)
C(11)-C(12)	1.418(15)	C(7X)-O(9X)	1.247(12)
C(12)-C(13)	1.458(13)	C(7X)-O(8X)	1.261(12)
C(13)-C(14)	1.317(12)	C(10X)-C(11X)	1.517(13)
C(14)-C(15)	1.470(14)	C(11X)-O(12X)	1.205(11)
C(15)-O(25)	1.227(12)	C(11X)-O(13X)	1.283(11)
C(14)-S(1X)-C(18)	100.9(5)	C(18)-C(17)-N(16)	112.7(10)
C(6)-N(1)-C(2)	110.3(8)	C(17)-C(18)-S(1X)	114.1(9)
C(6)-N(1)-C(26)	111.3(8)	C(20)-C(19)-C(24)	118.5(9)
C(2)-N(1)-C(26)	112.9(9)	C(20)-C(19)-N(16)	121.2(9)
N(1)-C(2)-C(3)	110.8(9)	C(24)-C(19)-N(16)	120.3(9)
N(4)-C(3)-C(2)	110.2(8)	C(21)-C(20)-C(19)	121.3(10)
C(7)-N(4)-C(3)	116.6(8)	C(20)-C(21)-C(22)	120.6(10)
C(7)-N(4)-C(5)	112.8(8)	C(21)-C(22)-C(23)	119.4(10)
C(3)-N(4)-C(5)	109.3(8)	C(21)-C(22)-Cl(2X)	120.6(9)
N(4)-C(5)-C(6)	110.4(9)	C(23)-C(22)-Cl(2X)	119.9(8)
C(5)-C(6)-N(1)	110.7(8)	C(24)-C(23)-C(22)	119.2(9)
C(8)-C(7)-C(12)	120.7(10)	C(24)-C(23)-Cl(1X)	119.1(9)

WO 03/048141

PCT/IB02/04972

-14-

C(8)-C(7)-N(4)	121.9(9)	C(22)-C(23)-Cl(1X)	121.6(8)
C(12)-C(7)-N(4)	117.4(8)	C(23)-C(24)-C(19)	120.9(10)
C(9)-C(8)-C(7)	119.6(10)	O(2X)-C(1X)-O(3X)	121.5(11)
C(10)-C(9)-C(8)	120.3(10)	O(2X)-C(1X)-C(4X)	122.0(11)
C(11)-C(10)-C(9)	120.6(11)	O(3X)-C(1X)-C(4X)	116.1(10)
C(10)-C(11)-C(12)	121.6(11)	C(1X)-C(4X)-C(5X)	113.2(9)
C(7)-C(12)-C(11)	117.1(9)	O(6X)-C(5X)-C(7X)	110.4(9)
C(7)-C(12)-C(13)	122.6(10)	O(6X)-C(5X)-C(4X)	109.0(8)
C(11)-C(12)-C(13)	120.2(9)	C(7X)-C(5X)-C(4X)	110.9(8)
C(14)-C(13)-C(12)	127.1(9)	O(6X)-C(5X)-C(10X)	108.4(8)
C(13)-C(14)-C(15)	117.4(9)	C(7X)-C(5X)-C(10X)	106.9(8)
C(13)-C(14)-S(1X)	120.9(7)	C(4X)-C(5X)-C(10X)	111.1(8)
C(15)-C(14)-S(1X)	121.6(8)	O(9X)-C(7X)-O(8X)	126.8(9)
O(25)-C(15)-N(16)	119.5(9)	O(9X)-C(7X)-C(5X)	118.0(10)
O(25)-C(15)-C(14)	121.5(10)	O(8X)-C(7X)-C(5X)	115.2(10)
N(16)-C(15)-C(14)	119.0(10)	C(11X)-C(10X)-C(5X)	118.4(8)
C(15)-N(16)-C(19)	119.1(9)	O(12X)-C(11X)-O(13X)	123.6(9)
C(15)-N(16)-C(17)	119.5(9)	O(12X)-C(11X)-C(10X)	120.4(9)
C(19)-N(16)-C(17)	120.1(9)	O(13X)-C(11X)-C(10X)	116.0(9)

Table VI. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for Citrate Salt. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$

	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
S(1X)	40(2)	76(2)	39(2)	6(2)	2(1)	-14(2)
Cl(1X)	128(3)	69(2)	46(2)	-17(2)	1(2)	24(2)
Cl(2X)	81(2)	82(2)	47(2)	7(2)	-13(2)	13(2)
N(1)	53(6)	44(6)	30(5)	-11(4)	10(5)	-16(5)
C(2)	53(8)	57(8)	46(7)	-9(6)	28(6)	-22(7)
C(3)	56(8)	56(7)	35(6)	1(5)	16(6)	-14(6)
N(4)	45(6)	45(6)	23(5)	-1(4)	12(4)	-2(5)
C(5)	61(8)	39(6)	36(6)	15(5)	24(6)	-7(6)
C(6)	63(8)	41(6)	22(6)	2(5)	17(6)	-2(6)
C(7)	43(7)	37(6)	21(6)	3(5)	4(5)	-23(6)
C(8)	51(8)	49(7)	33(7)	6(5)	6(6)	-15(7)
C(9)	64(10)	52(7)	45(8)	22(6)	-15(7)	-17(8)
C(10)	49(9)	43(7)	66(9)	11(7)	-5(7)	-1(6)
C(11)	55(8)	46(7)	36(7)	10(5)	-1(6)	4(6)
C(12)	47(7)	27(6)	33(7)	6(5)	5(5)	-9(6)
C(13)	38(7)	43(6)	24(6)	0(5)	6(5)	-4(5)
C(14)	35(6)	39(6)	19(5)	-6(5)	2(5)	-12(5)
C(15)	35(7)	51(7)	38(7)	-5(6)	5(6)	-8(6)
N(16)	53(6)	28(5)	30(5)	6(4)	-1(5)	-8(5)
C(17)	77(10)	58(8)	78(9)	22(7)	-26(8)	-39(8)
C(18)	46(8)	56(8)	71(8)	18(7)	7(7)	-22(7)
C(19)	39(7)	35(6)	22(6)	5(5)	-2(5)	-1(6)
C(20)	51(8)	41(7)	29(6)	-1(5)	9(6)	7(6)
C(21)	61(8)	50(7)	45(7)	10(6)	22(7)	19(6)
C(22)	49(8)	45(7)	39(7)	18(6)	3(6)	19(6)
C(23)	70(8)	47(7)	14(6)	-2(5)	12(6)	1(7)
C(24)	60(9)	35(7)	41(7)	-2(6)	13(7)	1(6)
O(25)	60(6)	104(7)	39(4)	29(4)	-6(4)	-35(6)
C(26)	57(9)	55(7)	60(8)	-12(6)	14(7)	2(7)
C(1X)	54(8)	41(7)	61(9)	-9(6)	2(7)	4(7)
O(2X)	106(9)	93(7)	87(7)	-21(6)	-13(6)	43(7)
O(3X)	99(8)	107(7)	33(5)	-7(5)	-21(5)	52(6)
C(4X)	53(8)	30(6)	34(6)	-1(5)	4(6)	14(6)

WO 03/048141

PCT/IB02/04972

-16-

	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C(5X)	49(8)	31(6)	24(5)	1(5)	7(5)	3(6)
O(6X)	71(6)	43(5)	40(5)	-4(3)	8(4)	-5(4)
C(7X)	38(7)	62(8)	15(6)	-7(6)	-2(5)	9(6)
O(8X)	106(8)	108(7)	18(4)	-7(4)	29(5)	-51(6)
O(9X)	75(6)	51(5)	34(4)	-18(4)	0(4)	10(5)
C(10X)	54(8)	60(7)	23(6)	-4(5)	9(5)	18(6)
C(11X)	40(7)	33(6)	27(6)	10(5)	13(6)	6(6)
O(12X)	43(5)	75(5)	28(4)	-12(4)	-7(4)	15(4)
O(13X)	68(6)	104(7)	19(4)	9(4)	5(4)	44(5)

Table VII. Hydrogen coordinates ($\times 10^4$) and Isotropic Displacement Parameters ($\text{\AA}^2 \times 10^3$) for the Citrate Salt.

	x	y	z	U(eq)
H(1X)	7880(190)	1710(70)	5570(80)	80
H(2A)	9814	2789	5202	80
H(2B)	10118	3134	6199	80
H(3A)	7538	3942	5290	80
H(3B)	6298	3082	4971	80
H(5A)	4753	2246	6011	80
H(5B)	4944	2564	7013	80
H(6A)	7199	1410	6952	80
H(6B)	8445	2258	7310	80
H(8)	3923	4179	4796	80
H(9)	1065	5080	4711	80
H(10)	23	5518	6002	80
H(11)	1675	5026	7377	80
H(13)	6447	4120	7953	80
H(17A)	3859	2028	9257	80
H(17B)	3373	2045	10232	80
H(18A)	1235	3198	9851	80
H(18B)	524	2391	9226	80
H(20)	8167	1886	10444	80
H(21)	10473	1695	11764	80
H(24)	5623	3895	11512	80
H(26A)	11422	1595	6859	80
H(26B)	11432	1429	5835	80
H(26C)	10234	789	6361	80
H(3XX)	11700(300)	-220(100)	5150(100)	140(60)
H(4XA)	8699	184	2591	80
H(4XB)	9020	927	3321	80
H(6XX)	6600(200)	-1000(80)	3490(80)	80
H(10A)	3582	822	2834	80
H(10B)	5465	1382	2683	80
H(13X)	5480(180)	190(70)	610(80)	80

The powder X-ray diffraction pattern was calculated from the single crystal data gathered for the citrate salt via the use of the XFOG and XPOW computer programs provided as part of the SHELXTL™ computer library. The calculated powder pattern is shown in Figure 3.

5 Solid State NMR

The citrate salt was characterized by solid state NMR techniques. Approximately 300 mg of a sample was tightly packed into 7mm ZrO spinner. The ^{13}C NMR spectra were collected using cross-polarization magic angle spinning (CPMAS) at 295 K on Bruker 7mm WB MAS probe positioned into a wide-bore Bruker Avance DRX 500 MHz NMR spectrometer. The sample was spun at 15.0 kHz. The cross-polarization contact time was set to 1 ms. The total of 512 scans were acquired for most of the samples resulting in approximately 30 minute acquisition times. The spectra were referenced using external sample of adamantane (δ 29.5 ppm) with the most upfield methyl signal set to 29.5 ppm.

The resulting ^{13}C NMR CPMAS spectrum for the citrate salt is shown in Figure 5.

15 The samples of the citrate salt behaved reasonably well from the point of view of solid state spectra quality. The resolution was good and the sensitivity was acceptable.

The major resonance peaks for the solid state carbon spectrum of the citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one downfield from 100 ppm are listed in Table VIII.

20 **Table VIII. The Major Solid State ^{13}C -NMR Resonance Peaks
For Citrate Salt (adamantane standard 29.5 ppm).**

^{13}C (ppm)
179.3
177.0
171.6
164.0
151.0
144.1

The citrate salt of the invention (hereafter "the active salt") can be administered via either the oral, transdermal (e.g., through the use of a patch), intranasal, sublingual, rectal, parenteral or topical routes. Transdermal and oral administration are preferred. The active salt is, most desirably, administered in dosages ranging from about 0.01 mg up to about 1500 mg per day, preferably from about 0.1 to about 300 mg per day in single or divided doses, although variations will necessarily occur depending upon the weight and condition of the subject being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 0.001 mg to about 10 mg per kg of body weight per day is most desirably employed.

30 Variations may nevertheless occur depending upon the weight and condition of the persons

being treated and their individual responses to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval during which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effects, provided that such larger doses are first divided into several small doses for administration throughout the day.

The active salt can be administered alone or in combination with pharmaceutically acceptable carriers or diluents by any of the several routes previously indicated. More particularly, the active salt can be administered in a wide variety of different dosage forms, e.g., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, transdermal patches, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents. In addition, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the active salt is present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc can be used for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar, as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration the active ingredient may be combined with various sweetening or flavoring agents, coloring matter and, if so desired, emulsifying and/or suspending agents, together with such diluents as water, ethanol, propylene glycol, glycerin and various combinations thereof.

For parenteral administration, a solution of an active salt in either sesame or peanut oil or in aqueous propylene glycol can be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8), if necessary, and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

-20-

It is also possible to administer the active salt topically and this can be done by way of creams, a patch, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

Example

- 5 The following example illustrate the method and compound of the present invention. It will be understood, however, that the invention is not limited to this specific Example.


Citrate Salt of 4-(3,4-Dichlorophenyl)-2-

[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one

- 10 In a 1-liter round-bottomed flask equipped with an overhead mechanical stirrer, the free base of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one (30.14 g; 0.067 mol; prepared in accordance with the procedures set forth in International Patent Publication No. WO 98/14433) was dissolved in 525 ml 2-propanol. The solution was stirred and heated to 50°C. Citric acid (16.2 g; 0.084 mol) was added in portions to the resulting clear solution and the reaction mixture was allowed to cool and was granulated at room
- 15 temperature for 18 hours. A sample of the precipitated solid was examined by DSC to ascertain whether any unreacted free base were present prior to collection of the white crystalline solid product. After filtration, the solid product was washed with 2-propanol (100 ml) and dried at 45 °C under vacuum with a nitrogen purge. The title citrate salt was yielded in 94% yield (40.6 g; 0.063 mol).

CLAIMS

1. The citrate salt of 4-(3,4-dichlorophenyl)-2-[2-(4-methylpiperazin-1-yl)-benzylidene]-thiomorpholin-3-one.
2. The citrate salt of a compound of formula :



3. A compound according to claim 1 having an x-ray diffraction pattern characterized substantially by an x-ray diffraction pattern peak as measured with copper radiation of a 2θ of about 17.4.

4. A compound according to claim 1 having an x-ray diffraction pattern characterized substantially by an x-ray diffraction pattern peak as measured with copper radiation of a 2θ of about 20.0.

5. A compound according to claim 1 having an x-ray diffraction pattern characterized substantially by x-ray diffraction pattern peaks as measured with copper radiation of a 2θ of about 17.4 and 20.0.

6. A compound according to claim 1 having an x-ray diffraction pattern characterized substantially by the following principal x-ray diffraction pattern peaks expressed in terms of 2θ and d-spacings as measured with copper radiation:

Angle 2θ (± 0.2)	d-value ($\text{\AA}$) (± 0.2)
13.0	6.8
17.4	5.1
18.0	4.9
18.9	4.7
20.0	4.4
21.2	4.2
22.2	4.0
24.0	3.7
27.1	3.3
32.4	2.8

7. A compound according to claim 1 characterized by an onset of melting/decomposition transition at 198-199 °C.

8. A compound according to claim 1 characterized in that when examined by solid state ^{13}C NMR cross-polarization magic angle spinning techniques it exhibits the following principal resonance peak: δ 179.3.

5 9. A compound according to claim 1 characterized in that when examined by solid state ^{13}C NMR cross-polarization magic angle spinning techniques it exhibits the following principal resonance peak: δ 177.0.

10. A compound according to claim 1 characterized in that when examined by solid state ^{13}C NMR cross-polarization magic angle spinning techniques it exhibits the following principal resonance peak: δ 171.6.

10 11. A compound according to claim 1 characterized in that when examined by solid state ^{13}C NMR cross-polarization magic angle spinning techniques it exhibits the following principal resonance peaks: δ 179.3, 177.0, 171.6, 164.0, 151.0 and 144.1.

12. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier.

13. A compound according to Claim 1 for use as a medicament.

14. The use of a compound according to Claim 1 in the manufacture of a medicament for the treatment of hypertension, all forms of depression, depression in cancer patients, depression in Parkinson's patients, postmyocardial infarction depression, subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depressive disorder, single episode depression, recurrent depression, child abuse induced depression, post partum depression, dysthymia; mild, moderate, or severe depressions with or without atypical features, melancholic features, psychotic features, catatonic features; seasonal affective disorder, geriatric depression, chronic depression; adjustment disorder with depressed mood or with anxiety and depressed mood; mixed anxiety and depression; substance induced mood disorder; and mood disorder secondary to a general medical condition, bipolar disorder, bipolar disorder-depressed phase, generalized anxiety disorder, phobias, agoraphobia, social anxiety, social phobia, simple phobias, separation anxiety disorder, posttraumatic stress syndrome, avoidant personality disorder, premature ejaculation, eating disorders, binge eating disorder, anorexia nervosa, bulimia nervosa, obesity; chemical dependencies and addictions to alcohol, cocaine, heroin, phenobarbital, nicotine, marijuana and benzodiazepines; cluster headache, migraine, pain, Alzheimer's disease, obsessive-compulsive disorder, panic disorder, panic disorder with agoraphobia, memory disorders, dementia, amnesic disorders, and age-related cognitive decline (ARCD). Parkinson's diseases, dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias, endocrine disorders, hyperprolactinaemia, vasospasm, vasospasm in the cerebral vasculature, cerebellar ataxia; gastrointestinal tract disorders involving changes in motility and secretion; negative symptoms of schizophrenia, premenstrual syndrome, fibromyalgia syndrome, stress incontinence, Tourette's syndrome, trichotillomania, lideptomenia,

male impotence, cancer, small cell lung carcinoma, chronic paroxysmal hemicrania, headache associated with vascular disorders, autism, pervasive developmental disorder NOS, Asperger's disorder, selective mutism, chronic motor or vocal tic disorder, somatization disorder, insomnia, intermittent explosive disorder, pyromania, pathological gambling, impulse-control disorder, premenstrual dysphoric disorder and attention-deficit/hyperactivity disorder (ADHD) in a mammal.