

This Sheet Has Been Replaced



1

REPLACED SHEETS		SG
☒	Description	: paged ...1... to paged ...17...
☒	Claim(s)	: paged ...18... to paged ...23...
☐	Drawing(s)	: paged to paged
☒	Abstract	: paged ...24... to paged ...25...
DATE OF RECEIPT		

Substituted β -diketones and their use.


The present invention relates to substituted β -diketones and their physiologically acceptable salts and esters, which are useful in the treatment of inflammatory bowel diseases.

EP-A-0357403 discloses β -diketones which possess cytoprotective efficacy in the stomach and in the duodenum and are useful in the treatment or prophylaxis of gastric and duodenal ulcers.

It has now been found that some substituted β -diketones are especially useful in the treatment of inflammatory bowel diseases (IBD).


The term IBD encompasses chronic inflammatory conditions of the gastrointestinal tract such as Crohn's disease and ulcerative colitis. The exact cause of the disease is still unknown. Infectious agents, immunological abnormalities, permeability disorders and leukotrienes are some of the factors suggested to play a role in the pathogenesis of IBD. The present medical therapy (sulfasalazine, 5-ASA, corticosteroids) is focused only on the diminishing of the inflammatory response, and there is still a great requirement for a more specific and effective drug for the treatment of IBD (Sutherland, CMAJ 137:799-802, 1987).

The present invention provides use of a compound of general formula I






wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;




wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;




wherein $m=0$, 1 or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; Y is oxygen or sulfur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.

The present invention also provides a compound according to the general formula II




wherein $n=0$ or 1 , R_1 is methyl, ethyl or cyclopropyl and R_2 is ethyl or cyclopropyl and R is






wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;




wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl or R_1 and R_2 are methyl and R is




where X_1 , X_2 and X_3 are as defined above provided that one of X_2 and X_3 is carbamoyl which is optionally substituted by alkyl, aryl or aralkyl, or a pharmaceutically acceptable salt or ester thereof.

The invention also provides a new compound according to the general formula III



wherein n is 0 or 1 , R_1 is methyl and X_1 is as defined above and X_2 is



wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl and X_3 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both



nitro or cyano; or X_2 is nitro and X_3 is formyl or halo or, when X_1 is hydroxy, X_2 may be carboxy, or X_2 and X_3 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl;

or a pharmaceutically acceptable salt or ester thereof but excluding

3-[(2-chloro-5-nitrophenyl)methylene]-2,4-pentanedione.

The term "alkyl" as employed herein refers to an alkyl group having preferably 1 to 2 carbon atoms.

The term "alkoxy" as employed herein refers to an alkyl residue as defined above linked to an oxygen atom.

The term "aralkyl" as employed herein refers to an alkyl group having aryl, preferably phenyl as the substituent.

The term "aryl" as employed herein refers to a carbocyclic aromatic group containing from 6 to 10 carbon atoms in the ring portion. The group may be a single ring or fused rings. Specific examples are phenyl and naphthyl.

The term "heteroaryl" as employed herein refers to a monocyclic or bicyclic aromatic group of carbon atoms containing one or more heteroatoms such as nitrogen, oxygen or sulfur.

The preparation of compounds according to the formulae I and II may be carried out in a similar way as described in EP-A-0357403. For example the compounds of formula I may be prepared by reacting a compound of the formula IV




in which R_1 and R_2 are as defined above with a compound of formula V






in which n and R are as defined above and Z is CHO or $-\text{CH}_2-\text{Q}$, wherein Q is halogen or some other activated group, in the presence of acidic or basic catalyst to produce a compound of formula I or a compound of formula VI



in which n , R_1 , R_2 and R are as defined above, which compound is halogenated to give a compound of the formula VII



in which n , R_1 , R_2 and R are as defined above and X is halogen and dehydrohalogenating compound VII to produce a compound of the formula I.

The basic catalyst may be for example an organic or inorganic base. The acidic catalyst may be for example a mineral or sulfonic acid. The activated group Q may be a halogen or an alkyl or aryl sulfonate.

When halogenating a compound of formula VI elementary halogen, preferably chlorine or bromine may be used, or another known halogenating agent such as sulfonyl chloride may be used.

Salts of these compounds may be prepared by known methods. Most commonly used physiologically acceptable salts are sodium, potassium, ammonium, calcium and magnesium salts and hydrochlorides or hydrobromides. Preferred esters of the



compounds of formula I are acyl or aroyl derivatives which will hydrolyse under physiological conditions.

The above compounds may be formulated to dosage forms using the principles which are known to those having average skill in the art. Suitable solvents, gel forming ingredients, dispersion agents, antioxidants and colourants may be added in a normal way.

It is preferred to administer the compounds enterally. When they are given orally, they may be in the form of tablets, granules, capsules, emulsions, suspensions or solutions. It is often advisable to use coated tablets or granules, i.e. so called enterocoated preparations, to ensure that the medicine reaches the desired part of the gastrointestinal tract. Alternatively, the compositions may be given rectally in the form of enemas or suppositories.

The effective dose varies considerably depending on location, degree and severity of the disease being treated as well as the age and the general condition of the patient. The effective dose is generally from about 20 to 1000 mg per day, preferably from 50 to 250 mg per day for an adult, once a day or divided into two to five doses.

In vivo tests

The efficacy of the compounds can be demonstrated by using a model of chronic colonic inflammation in the rat, TNB-induced chronic colitis. TNB-induced chronic colonic inflammation in rats is an animal model which has been shown to resemble highly human IBD.



20 mg of TNB (2,4,6-trinitrobenzenesulfonic acid) in 50 % ethanol was administered into the lumen of the colon via a rubber catheter inserted rectally to Wistar rats. The test compounds were given rectally 1 h prior and 24, 48, 72 and 95 h after TNB. The rats were sacrificed 96 h after TNB and their colons were scored according to occurrence of ulcers and inflammation. The number of animals in each group was at least 8. 5-ASA (5-aminosalicylic acid), a clinically used drug for the treatment of IBD was used as a reference compound. The results are presented in the Table 1.

Table 1.

The effect of test compounds on the damage score of chronic inflammation induced by TNB.

Test compound	Dose mg/kg	% inhibition against control
5-ASA	100	24
1	30	37
2	30	28
3	10	18
	30	57
4	30	54
5	30	43
6	30	41
7	10	60
	30	48
8	10	51
	30	62
9	30	40
10	10	27
11	30	32
12	30	52
13	30	58
14	30	30



Test compounds

- 1 3-(4-hydroxyphenyl)methylene-2,4-pentanedione
- 2 3-(4-methoxyphenyl)methylene-2,4-pentanedione
- 3 3-(3-nitrophenyl)methylene-2,4-pentanedione
- 4 3-(4-trifluoromethylphenyl)methylene-2,4-pentanedione
- 5 3-(4-nitrophenyl)methylene-2,4-pentanedione
- 6 3-(2-nitrophenyl)methylene-2,4-pentanedione
- 7 3-(4-cyanophenyl)methylene-2,4-pentanedione
- 8 3-(3-cyanophenyl)methylene-2,4-pentanedione
- 9 3-[(4-(N-phenethyl)carboxamido-phenyl)methylene]-
2,4-pentanedione
- 10 3-[(5-nitrofuryl)methylene]-2,4-pentanedione
- 11 4-[(4-cyanophenyl)methylene]-3,5-heptanedione
- 12 2-[(4-cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione
- 13 2-[(3-trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione
- 14 2-[(4-carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione

The test compounds significantly decreased the ulceration and inflammation of the colons. The test compounds were much more potent than the reference compound 5-ASA and should therefore be more effective than 5-ASA in the treatment of human IBD.



The following examples illustrate the production of compounds of use according to the invention.

Example 1

3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione

To a solution containing 2.1 g of 4-phenylsulfonylbenzaldehyde and 1.5 g of 2,4-pentanedione in 10 ml of 2-propanol was added 1 ml of thionyl chloride with stirring at 20°C. The solution was stirred for 1h at 20°C and evaporated to dryness in vacuo. The residue was crystallized from toluene, mp 102-105°C.

Example 2

3-[(4-Methylsulfonylphenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 1.1 g of 4-methylsulfonylbenzaldehyde and 1.5 g of 2,4-pentanedione. The product was crystallized from ether, mp 139-140°C.

Example 3

3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione

A mixture containing 1.8 g of 4-carboxamidobenzaldehyde, 1.1 g of 2,4-pentanedione, 0.2 g of piperidine and 0.2 g of acetic acid in 40 ml of toluene was refluxed for 3 h by using a Dean-Stark separator. The warm clean toluene solution was decanted from the tarry residue and cooled. The crystalline product was filtered and dried. Mp 154-155°C.



Example 4

4-[(4-Nitrophenyl)methylene]-3,5-heptanedione

The procedure described in Example 1 was repeated by using 1.5 g of 4-nitrobenzaldehyde and 2.0 g of 3,5-heptanedione. Yield 1.7 g, yellow oil.

Example 5

3-[(3,4-Dichlorophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 4.26 g of 3,4-dichlorobenzaldehyde and 3.0 g of 2,4-pentanedione. Yield 1.3 g, mp 73°C.

Example 6

3-[(4-Chloro-3-nitrophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 1.85 g of 4-chloro-3-nitrobenzaldehyde and 1.5 g of 2,4-pentanedione. Yield 1.32 g, mp 110-111°C.

Example 7

3-[(4-(N-Phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.

The procedure described in Example 1 was repeated by using 2.5 g of 4-(N-phenethyl)carboxamidobenzaldehyde and 1.0 g of 2,4-pentanedione. Yield 0.5 g, mp 146-149°C.

Example 8

3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione

To a solution containing 1.84 g of 2-nitroterephthalaldehyde and 1.00 g of 2,4-pentanedione in 10 ml trifluoroacetic acid were added 0.9 ml of thionyl chloride and a catalytic amount of water. The solution was stirred for 2 h at 20°C. 30 ml of water was added and the mixture was extracted with ether. The ether extract was washed several times with Na₂CO₃-solution and dried over K₂CO₃. After filtration the solvent was evaporated in vacuo. The product was purified by column chromatography. Yield 0.2 g, mp 102-104°C.

Example 9

3-[(3,5-Dinitro-4-hydroxyphenyl)methylene]-2,4-pentanedione.

The procedure described in Example 1 was repeated by using 2.12 g of 3,5-dinitro-4-hydroxybenzaldehyde and 1.5 g of 2,4-pentanedione. Yield 1.9 g, mp 158°C.

Example 10

3-[(3,5-Dinitrophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 2.53 g 3,5-dinitrobenzaldehyde and 2.43 g of 2,4-pentanedione. Yield 0.8 g, mp 101-105°C.

Example 11

3-[3-(4-Methoxyphenyl)propenylene]-2,4-pentanedione

To a solution containing 2.45 g of 4-methoxycinnamaldehyde in 90 ml of toluene was added 2.2 g of 2,4-pentanedione, 0.26 g of piperidine and 0.45 g of acetic acid. The solution was refluxed in N₂ atmosphere for 1 h and the water formed in the reaction was azeotropically distilled off by using a Dean-Stark water separator. The solution was evaporated to dryness in vacuo. The residue was trifurated with ether. The crystals were filtered. Yield 0.65 g, mp 108-110°C.

Example 12

3-[(5-Nitrofuryl)methylene]-2,4-pentanedione

A solution containing 2.43 g of 5-nitro-2-furanecarboxaldehyde diacetate, 1.5 g of 2,4-pentanedione and 0.1 g of water in 5 ml of tetrahydrofuran was saturated with hydrogen chloride gas at 20°C. The mixture was stirred for 1 h and evaporated to dryness in vacuo. The product was recrystallized from acetic acid. Yield 0.1 g, mp 117-120°C.

Example 13

3-[(5-Nitrothienyl)methylene]-2,4-pentanedione

The procedure described in Example 12 was repeated by using 2.59 g of 5-nitro-2-thiophenecarboxaldehyde diacetate, 1.5 g of 2,4-pentanedione. Yield 0.5 g, yellow viscous oil.

Example 14

3-(3-Pyridylmethylene)-2,4-pentanedione

The procedure described in Example 3 was repeated by using 3.21 g of pyridine-3-carboxaldehyde and 3.3 g of 2,4-pentanedione. Yield 2.95 g, mp 52-57°C.

Example 15

3-(4-Pyridylmethylene)-2,4-pentanedione hydrochloride

A solution containing 1.1 g of pyridine-4-carboxaldehyde and 1.0 g of 2,4-pentanedione in 10 ml of N,N-dimethylformamide was saturated with hydrogen chloride gas at 20°C. The solution was stirred for 1 h at 20°C and then for 2 h at 80°C. The solvent was evaporated in vacuo and the residue was crystallized from ethanol. Yield 0.17 g, mp 175-186°C.

Example 16

3-(3-Quinolylmethylene)-2,4-pentanedione hydrochloride

The procedure described in Example 3 was repeated by refluxing 2.4 g quinoline-3-aldehyde, 6.6 g 2,4-pentanedione, 0.4 g piperidine and 0.8 g acetic acid in 100 ml toluene. Yield 1.4 g, mp 99-101°C.

Example 17

3-[(3-Carboxy-4-hydroxyphenyl)methylene]-2,4-pentanedione

A mixture containing 2 g 3-carboxy-4-hydroxybenzaldehyde, 4 ml 2,4-pentanedione and 20 ml tetrahydrofuran was saturated with hydrogen chloride gas. The solution was stirred for 20 h



at 50°C and evaporated to dryness in vacuo. The residue was crystallized from ethyl ether, mp. 112-114°C.

Example 18

3-[[(4-(2-Phenylethoxy)phenyl)methylene]-2,4-pentanedione

The procedure described in example 8 was repeated by using 5.9 g of 4-(2-phenylethoxy)benzaldehyde, 3.9 ml of 2,4-pentanedione, 26 ml of trifluoroacetic acid and 2.1 ml of thionyl chloride. Yield 4.7 g, an viscous oil.

Example 19

2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

A mixture containing 1.6 g 4-acetamidobenzaldehyde, 1.5 g 1,3-dicyclopropyl-1,3-propanedione, 0.2 ml of piperidine and 0.4 ml of acetic acid in 50 ml of toluene was refluxed for 3 h by using a Dean-Stark separator. After standing over night at room temperature the crystals were filtered and washed with toluene. The product was recrystallized from ethanol. Yield 2.0 g, mp 166-167°C.

Example 20

2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 0.68 g 4-methoxybenzaldehyde and 0.77 g 1,3-dicyclopropyl-1,3-propanedione. Yield 0.46 g, mp 60.5-61.5°C.

Example 21

4-[(3-Cyanophenyl)methylene]-3,5-heptanedione

The procedure described in Example 19 was repeated by using 1.82 g 3-cyanobenzaldehyde and 3.5 g 3,5-heptanedione. Yield 0.4 g, mp 51-52°C.

Example 22

4-[(4-Cyanophenyl)methylene]-3,5-heptanedione

The procedure described in Example 19 was repeated by using 2.6 g 4-cyanobenzaldehyde and 5.0 g 3,5-heptanedione. Yield 2.8 g, mp 59-60°C.

Example 23

3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione

The procedure described in Example 19 was repeated by using 2.6 g 4-cyanobenzaldehyde and 2.5 g 4-cyclopropyl-2,4-butanedione. The product was purified by column chromatography. Yield 0.37 g, mp 83-85°C.

Example 24

2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 3.78 g 4-cyanobenzaldehyde and 2.23 g 1,3-dicyclopropyl-1,3-propanedione. Yield 1.5 g, mp 100-102°C.

Example 25

3-[(5-Bromothiophenyl)methylene]-2,4-pentanedione

To a solution containing 5.73 g 5-bromothiophene-2-carboxaldehyde and 5.0 g 2,4-pentanedione in 25 ml of 2-propanol, 2.9 ml thionylchloride was gradually added. The mixture was stirred for 2 h at 20°C. 120 ml of conc. Na₂SO₄-solution was then added and the mixture was extracted with ether. The solvent was evaporated in vacuo and the crystalline residue was washed with cold ether. Yield 3.9 g, mp 70-74°C.

Example 26

2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 2.15 ml 3-trifluoromethylbenzaldehyde and 2.42 g 1,3-dicyclopropyl-1,3-propanedione. Yield 2.71 g, mp 56-58°C.

Example 27


2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 3.0 g 4-carboxybenzaldehyde and 3.06 g 1,3-dicyclopropyl-1,3-propanedione. Yield 1.7 g, mp 175-178°C.




CLAIMS


1. Use of a compound of formula I



wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;



wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; Y is oxygen or sulfur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.

2. The use according to claim 1, in which, in the compound of formula I, X_1 is hydroxy or alkoxy.



3. The use according to claim 1 or 2, in which, in the compound of formula I, one of X_2 and X_3 is nitro, cyano, trifluoromethyl, carboxy or carbamoyl which is substituted by aryl.

4. The use according to claims 1 to 3, in which the compound is

3-(4-hydroxyphenyl)methylene-2,4-pentanedione,

3-(4-methoxyphenyl)methylene-2,4-pentanedione,

3-(3-nitrophenyl)methylene-2,4-pentanedione,

3-(4-nitrophenyl)methylene-2,4-pentanedione,

3-(2-nitrophenyl)methylene-2,4-pentanedione,

3-[(4-(N-phenethyl)carboxamido-phenyl)methylene]-
2,4-pentanedione,

3-[(5-nitrofuryl)methylene]-2,4-pentanedione, or

2-[(4-carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-

propanedione or a pharmaceutically acceptable salt or ester thereof.

5. The use according to claims 1 or 3, in which, the compound is 3-(4-trifluoromethylphenyl)methylene-2,4-pentanedione.

6. The use according to claims 1 or 3, in which, the compound is 3-(4-cyanophenyl)methylene-2,4-pentanedione.

7. The use according to claims 1 or 3, in which, the compound is 3-(3-cyanophenyl)methylene-2,4-pentanedione.


8. The use according to claims 1 or 3, in which, the compound is 4-[(4-cyanophenyl)methylene]-3,5-heptanedione.

9. The use according to claims 1 or 3, in which, the compound is 2-[(4-cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.




10. The use according to claims 1 or 3, in which, the compound is 2-[(3-trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.


11. A compound according to the general formula II




wherein $n=0$ or 1, R_1 is methyl, ethyl or cyclopropyl and R_2 is ethyl or cyclopropyl and R is



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;



wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl or R_1 and R_2 are methyl and R is



where X_1 , X_2 and X_3 are as defined above provided that one of X_2 and X_3 is carbamoyl which is optionally substituted by



alkyl, aryl or aralkyl, or a pharmaceutically acceptable salt or ester thereof.

12. A compound according to claim 11 wherein in the compound of formula II one of X_2 and X_3 is cyano or trifluoromethyl.

13. 2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

14. 2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

15. 3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione.

16. 2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

17. 2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

18. 2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione or a pharmaceutically acceptable salt or ester thereof.

19. 4-[(4-Nitrophenyl)methylene]-3,5-heptanedione.

20. 4-[(3-Cyanophenyl)methylene]-3,5-heptanedione.


21. 4-[(4-Cyanophenyl)methylene]-3,5-heptanedione.

22. 3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione.


23. 3-[(4-(N-phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.



24. A compound according to the general formula III



wherein R_1 is methyl, n is 0 or 1, X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl, and X_2 is;



wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl, and X_3 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both nitro or both cyano; or X_2 is nitro and X_3 is formyl or halo; or when X_1 is hydroxy, X_2 may be carboxy or X_2 and X_3 may both be hydrogen when $n=0$ and X_1 is hydroxy or alkoxy optionally substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl; or a pharmaceutically acceptable salt or ester thereof but excluding

3-[(2-chloro-5-nitrophenyl)methylene]-2,4-pentanedione.

25. 3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione.

26. 3-[(4-Methylsulphonylphenyl)methylene]-2,4-pentanedione.

27. 3-[(4-Chloro-3-nitrophenyl)methylene]-2,4-pentanedione.

28. 3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione.

29. 3-[(3,5-Dinitro-4-hydroxyphenyl)methylene]-2,4-pentanedione or a pharmaceutically acceptable salt or ester thereof.




30. 3-[(3,5-Dinitrophenyl)methylene]-2,4-pentanedione.
31. 3-[(3-Carboxy-4-hydroxy)methylene]-2,4-pentanedione a pharmaceutically acceptable salt or ester thereof.
32. 3-[[[4-(2-Phenethyloxy)phenyl)methylene]-2,4-pentanedione.
33. 3-[3-(4-Methoxyphenyl)propenylene]-2,4-pentanedione.
34. A compound according to any one of claims 11 to 33 for use as a medicament.
35. A compound according to any one of claims 11 to 33 for the use in the treatment of inflammatory bowel disease.
36. A pharmaceutical composition comprising a compound according to any one of claims 11 to 33 and a pharmaceutically acceptable carrier or diluent.
37. Use of a compound as defined in any one of claims 11 to 33 for the manufacture of a medicament for use in the treatment of inflammatory bowel disease.
38. The use according to any one of claims 1 to 10 or 37, in which the inflammatory bowel disease is ulcerative colitis.
39. The use according to any one of claims 1 to 10 or 37, in which the inflammatory bowel disease is Crohn's disease.

ABSTRACT

SUBSTITUTED β -DIKETONES AND THEIR USE

Compounds of formula



wherein n is 0 or 1, R₁ and R₂ are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group or salt or ester thereof are useful in the treatment of inflammatory bowel disease.

This Sheet Has Been Replaced



REPLACED SHEET	SG
☒ Description : paged 4 to paged 16, 21-23	
☒ Claim(s) : paged 16, 21-23 to paged 16, 21-23	
☐ Drawing(s) : paged 16, 21-23 to paged 16, 21-23	
☒ Abstract : paged 16, 21-23 to paged 16, 21-23	4

REPLACEMENT SHEET	SG
☒ Description : paged 4 to paged 16, 21-23	
☒ Claim(s) : paged 16, 21-23 to paged 16, 21-23	
☐ Drawing(s) : paged 16, 21-23 to paged 16, 21-23	
☒ Abstract : paged 16, 21-23 to paged 16, 21-23	
DATE OF RECEIPT	

nitro or cyano, or X_1 is nitro and X_2 is formyl or halo or, when X_1 is hydroxy, X_2 may be carboxy, or X_1 and X_2 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl;

or a pharmaceutically acceptable salt or ester thereof but excluding

3-[(2-chloro-5-nitrophenyl)methylene]-2,4-pentanedione.

The term "alkyl" as employed herein refers to an alkyl group having preferably 1 to 2 carbon atoms.

The term "alkoxy" as employed herein refers to an alkyl residue as defined above linked to an oxygen atom.

The term "aralkyl" as employed herein refers to an alkyl group as defined above having aryl, preferably phenyl as the substituent.

The term "aryl" as employed herein refers to a carbocyclic aromatic group containing from 6 to 10 carbon atoms in the ring portion. The group may be a single ring or fused rings. Specific examples are phenyl and naphthyl.

The term "heteroaryl" as employed herein refers to a monocyclic or bicyclic aromatic group of carbon atoms containing one or more heteroatoms such as nitrogen, oxygen or sulfur.

The preparation of compounds according to the formulae I and II may be carried out in a similar way as described in EP-A-0357403. For example the compounds of formula I may be prepared by reacting a compound of the formula IV



IV

in which R_1 and R_2 are as defined above with a compound of formula V



REPLACEMENT SHEET
12 MAR 1998

V

This Sheet Has Been Replaced










18

CLAIMS


1. Use of a compound of formula I



wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;



wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; in which the terms "alkyl", "alkoxy" and "aralkyl" are as defined herein; Y is oxygen or sulfur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.

2. The use according to claim 1, in which, in the compound of formula I, X_1 is hydroxy or alkoxy.

This Sheet Has Been Replaced



REPLACEMENT SHEET

12 MAR 1998



alkyl, aryl or aralkyl, in which "alkyl", "alkoxy" and "aralkyl" are as defined herein; or a pharmaceutically acceptable salt or ester thereof, but excluding 4-(phenylmethylene)-3,5-heptanedione, 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione and 3-(phenylmethylene)-2,4-hexanedione.

12. A compound according to claim 11 wherein in the compound of formula II one of X_2 and X_3 is cyano or trifluoromethyl.

13. 2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

14. 2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

15. 3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione.

16. 2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

17. 2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

18. 2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione or a pharmaceutically acceptable salt or ester thereof.

19. 4-[(4-Nitrophenyl)methylene]-3,5-heptanedione.


20. 4-[(3-Cyanophenyl)methylene]-3,5-heptanedione.

21. 4-[(4-Cyanophenyl)methylene]-3,5-heptanedione.

22. 3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione.


23. 3-[(4-(N-phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.

This Sheet Has Been Replaced






24. A compound according to the general formula III



wherein R_1 is methyl, n is 0 or 1, X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl, and X_2 is;



wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl, and X_3 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both cyano; or X_2 is nitro and X_3 is formyl; or when X_1 is hydroxy, X_2 may be carboxy or X_2 and X_3 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl; in which "alkyl", "alkoxy" and "aralkyl" are as defined herein; or a pharmaceutically acceptable salt or ester thereof.

25. 3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione.

26. 3-[(4-Methylsulphonylphenyl)methylene]-2,4-pentanedione.

27. 3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione.

REPLACEMENT SHEET


12 MAR 1998



- 23 -


28. 3-[(3-Carboxy-4-hydroxy)methylene]-2,4-pentanedione a pharmaceutically acceptable salt or ester thereof.
29. 3-[[4-(2-Phenethyloxy)phenyl)methylene]-2,4-pentanedione.
- 5 30. 3-[3-(4-Methoxyphenyl)propenylene]-2,4-pentanedione.
31. A compound according to any one of claims 11 to 30 for use as a medicament.
- 10 32. A compound according to any one of claims 11 to 30 for use in the treatment of inflammatory bowel disease.
33. 3-[(4-methylsulphonylphenyl)methylene]-2,4-pentanedione for use as a medicament.
- 15 34. A pharmaceutical composition comprising a compound according to any one of claims 11 to 30 and a pharmaceutically acceptable carrier or diluent.
- 20 35. Use of a compound as defined in any one of claims 11 to 30 for the manufacture of a medicament for use in the treatment of inflammatory bowel disease.
- 25 36. Use of a 3-[(4-methylsulphonylphenyl)methylene]-2,4-pentanedione in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.
- 30 37. The use according to any one of claims 1 to 10 or 35 or 36, in which the inflammatory bowel disease is ulcerative colitis.
38. The use according to any one of claims 1 to 10 or 35 or 36, in which the inflammatory bowel disease is Crohn's disease.

This Sheet Has Been Replaced




ABSTRACTSUBSTITUTED β -DIKETONES AND THEIR USE

Compounds of formula



wherein n is 0 or 1, R₁ and R₂ are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group or salt or ester thereof are useful in the treatment of inflammatory bowel disease.

This Sheet Has Been Replaced



REPLACED SHEETS		SG
☒ Description	: paged 1, 1a, 4 to paged -	
☒ Claim(s)	: paged 18, 20-22 to paged -	
☐ Drawing(s)	: paged to paged	
☐ Abstract	: paged to paged	
DATE OF RECEIPT		- 1 -

9105242-8 2905

REPLACEMENT SHEETS		SG
☒ Description	: paged 1, 1a, 4 to paged -	
☒ Claim(s)	: paged 18, 20-22 to paged -	
☐ Drawing(s)	: paged to paged	
☐ Abstract	: paged to paged	

~~SUBSTITUTED β -DIKETONES AND THEIR USE~~ DATE OF RECEIPT 29/6/99

The present invention relates to substituted β -diketones and their physiologically acceptable salts and esters, which are useful in the treatment of inflammatory bowel diseases.

DE-A-2,210,663 discloses 3'-nitro-4'-chlorobenzylidene-acetyl-acetone as an α, β -unsaturated dioxo compound which exhibits a cardiac relaxing and hypertensive effect.

DE-A-3,303,066 discloses substituted 1,4-dihydropyridines for the prevention of cardiovascular disorders and for treating troubles of the cerebral and peripheral blood circulation.

Bull. Chem. Soc. Japan, 48(3), 1975, pp. 882-884, T. Kawato et al., discloses the property of chelation of the compounds 3-(p-hydroxybenzylidene)-2,4-pentadione and 3-(m-hydroxybenzylidene)-2,4-pentadione.

Bull. Chem. Soc. Japan, 49(59), 1976, pp. 1368-1374, M. Iwata et al., refers to the condensation products of aromatic aldehydes with ketone promoted by Cu-II, including 3-(2-methoxybenzylidene)-2,4-pentanedione.


EP-A-0,008,458 discloses insecticidal triazole compounds useful in the protection of plants and crops.

US-A-3,280,069 is concerned with the stability of plastic against UV-light and mentions the compound 3-(o-hydroxybenzylidene)-2,4-pentanedione.

Bull. Chim. Soc. France, No. 7-8, 1974, pp. 1683-1690, J.-P. Vecchionacci et al., discloses the compound benzylidene-3-hexane-2,4-dione (cf. page 1688, table IV).

Eur. J. Med. Chem. 22(2), 1988, pp. 111-117, J.R. Dimmock et al., discloses in its experiments protocol the compounds 4-(phenylmethylene)-3,5-heptanedione and 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione.

This Sheet Has Been Replaced




EP-A-0357403 discloses β -diketones which possess cytoprotective efficacy in the stomach and in the duodenum and are useful in the treatment or prophylaxis of gastric and duodenal ulcers.

It has now been found that some substituted β -diketones are especially useful in the treatment of inflammatory bowel diseases (IBD).


The term IBD encompasses chronic inflammatory conditions of the gastrointestinal tract such as Crohn's disease and ulcerative colitis. The exact cause of the disease is still unknown. Infectious agents, immunological abnormalities, permeability disorders and leukotrienes are some of the factors suggested to play a role in the pathogenesis of IBD. The present medical therapy (sulfasalazine, 5-ASA, corticosteroids) is focused only on the diminishing of the inflammatory response, and there is still a great requirement for a more specific and effective drug for the treatment of IBD (Sutherland, CMAJ 137:799-802, 1987).

The present invention provides use of a compound of general formula I



I

This Sheet Has Been Replaced



nitro or cyano; or X_2 is nitro and X_1 is formyl or halo or, when X_1 is hydroxy, X_2 may be carboxy, or X_2 and X_1 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl;

or a pharmaceutically acceptable salt or ester thereof but excluding 4-(phenylmethylene)-3,5-heptanedione, 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione, 3-[(2-chloro-5-nitrophenyl)methylene]-2,4-pentanedione, and 3-(phenylmethylene)-2,4-hexanedione.

The term "alkyl" as employed herein refers to an alkyl group having preferably 1 to 2 carbon atoms.

The term "alkoxy" as employed herein refers to an alkyl residue as defined above linked to an oxygen atom.

The term "aralkyl" as employed herein refers to an alkyl group as defined above having aryl, preferably phenyl as the substituent.

The term "aryl" as employed herein refers to a carbocyclic aromatic group containing from 6 to 10 carbon atoms in the ring portion. The group may be a single ring or fused rings. Specific examples are phenyl and naphthyl.

The term "heteroaryl" as employed herein refers to a monocyclic or bicyclic aromatic group of carbon atoms containing one or more heteroatoms such as nitrogen, oxygen or sulfur.

The preparation of compounds according to the formulae I and II may be carried out in a similar way as described in EP-A-0357403. For example the compounds of formula I may be prepared by reacting a compound of the formula IV




IV

in which R_1 and R_2 are as defined above with a compound of formula V



V

This Sheet Has Been Replaced






CLAIMS





This Sheet Has Been Replaced




1. Use of a compound of formula I



wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;




wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;



wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; in which "alkyl" is an alkyl group having preferably 1 to 2 carbon atoms, "alkoxy" is an alkyl residue as defined above linked to an oxygen atom, and "aralkyl" is an alkyl group as defined above having aryl, preferably phenyl as the substituent; Y is oxygen or sulphur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.

2. The use according to claim 1, in which, in the compound of formula I, X_1 is hydroxy or alkoxy.







- 20 -

10. This use according to claims 1 or 3, in which, the compound is 2-[(3-trifluoromethylphenyl)methylene]-1,3-dicyclopentyl-1,3-propanedione.


11. A compound according to the general formula II




wherein $n=0$ or 1, R_1 is methyl, ethyl or cyclopropyl and R_2 is ethyl or cyclopropyl and R is



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido, provided that when $n=0$, X_1 , X_2 and X_3 are not all independently selected from the group consisting of hydrogen, hydroxy and nitro; or R is




wherein $m=0$, 1 or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl or R_1 and R_2 are methyl and R is



where X_1 , X_2 and X_3 are as defined above provided that one of X_2 and X_3 is carbamoyl which is optionally substituted by

This Sheet Has Been Replaced





alkyl, aryl or aralkyl, in which "alkyl", "alkoxy" and "aralkyl" are as defined herein; or a pharmaceutically acceptable salt or ester thereof, but excluding 4-(phenylmethylene)-3,5-heptanedione, 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione and 3-(phenylmethylene)-2,4-hexanedione.

12. A compound according to claim 11 wherein in the compound of formula II one of X_2 and X_3 is cyano or trifluoromethyl.

13. 2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

14. 2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

15. 3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione.

16. 2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

17. 2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

18. 2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione or a pharmaceutically acceptable salt or ester thereof.

19. 4-[(4-Nitrophenyl)methylene]-3,5-heptanedione.

20. 4-[(3-Cyanophenyl)methylene]-3,5-heptanedione.

21. 4-[(4-Cyanophenyl)methylene]-3,5-heptanedione.

22. 3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione.

23. 3-[(4-(N-phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.

This Sheet Has Been Replaced




REPLACEMENT SHEET


29 JUN 1999



24. A compound according to the general formula III



wherein R_1 is methyl, n is 0 or 1, X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl, and X_2 is;




wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl, and X_3 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both cyano; or X_2 is nitro and X_3 is formyl; or when X_1 is hydroxy, X_2 may be carboxy or X_2 and X_3 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl; in which "alkyl" is an alkyl group having preferably 1 to 2 carbon atoms, "alkoxy" is an alkyl residue as defined above linked to an oxygen atom, and "aralkyl" is an alkyl group as defined above having aryl, preferably phenyl as the substituent; or a pharmaceutically acceptable salt or ester thereof.

25. 3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione.

26. 3-[(4-Methylsulphonylphenyl)methylene]-2,4-pentanedione.

27. 3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione.

This Sheet Has Been Replaced



SUBSTITUTED β -DIKETONES AND THEIR USE RECEIPT 17/8/99

The present invention relates to substituted β -diketones and their physiologically acceptable salts and esters, which are useful in the treatment of inflammatory bowel diseases.

DE-A-2,210,663 discloses 3'-nitro-4'-chlorobenzylidene-acetyl-acetone as an α,β -unsaturated dioxo compound which exhibits a cardiac relaxing and hypertensive effect.

DE-A-3,303,066 discloses substituted 1,4-dihydropyridines for the prevention of cardiovascular disorders and for treating troubles of the cerebral and peripheral blood circulation.

Bull. Chem. Soc. Japan, 48(3), 1975, pp. 882-884, T. Kawato et al., discloses the property of chelation of the compounds 3-(p-hydroxybenzylidene)-2,4-pentadione and 3-(m-hydroxybenzylidene)-2,4-pentadione.


Bull. Chem. Soc. Japan, 49(59), 1976, pp. 1368-1374, M. Iwata et al., refers to the condensation products of aromatic aldehydes with ketone promoted by Cu-II, including 3-(2-methoxybenzylidene)-2,4-pentanedione.

EP-A-0,008,458 discloses insecticidal triazole compounds useful in the protection of plants and crops.

US-A-3,280,069 is concerned with the stability of plastic against UV-light and mentions the compound 3-(o-hydroxybenzylidene)-2,4-pentanedione.

Bull. Chim. Soc. France, No. 7-8, 1974, pp. 1683-1690, J.-P. Vecchionacci et al., discloses the compound benzylidene-3-hexane-2,4-dione (cf. page 1688, table IV).

Eur. J. Med. Chem. 22(2), 1988, pp. 111-117, J.R. Dimmock et al., discloses in its experiments protocol the compounds 4-(phenylmethylene)-3,5-heptanedione and 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione.





EP-A-0357403 discloses β -diketones which possess cytoprotective efficacy in the stomach and in the duodenum and are useful in the treatment or prophylaxis of gastric and duodenal ulcers.


It has now been found that some substituted β -diketones are especially useful in the treatment of inflammatory bowel diseases (IBD).

The term IBD encompasses chronic inflammatory conditions of the gastrointestinal tract such as Crohn's disease and ulcerative colitis. The exact cause of the disease is still unknown. Infectious agents, immunological abnormalities, permeability disorders and leukotrienes are some of the factors suggested to play a role in the pathogenesis of IBD. The present medical therapy (sulfasalazine, 5-ASA, corticosteroids) is focused only on the diminishing of the inflammatory response, and there is still a great requirement for a more specific and effective drug for the treatment of IBD (Sutherland, CMAJ 137:799-802, 1987).


The present invention provides use of a compound of general formula I



wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;




wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;





wherein $m=0$, 1 or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; Y is oxygen or sulfur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.


The present invention also provides a compound according to the general formula II




wherein $n=0$ or 1 , R_1 is methyl, ethyl or cyclopropyl and R_2 is ethyl or cyclopropyl and R is



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;




wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl or R_1 and R_2 are methyl and R is




where X_1 , X_2 and X_3 are as defined above provided that one of X_2 and X_3 is carbamoyl which is optionally substituted by alkyl, aryl or aralkyl, or a pharmaceutically acceptable salt or ester thereof.

The invention also provides a new compound according to the general formula III



wherein n is 0 or 1 , R_1 is methyl and X_1 is as defined above and X_2 is



wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl and X_1 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both

17 AUG 1999

- 5 -

nitro or cyano; or X_1 is nitro and X_2 is formyl or halo or, when X_1 is hydroxy, X_2 may be carboxy, or X_1 and X_2 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl;

or a pharmaceutically acceptable salt or ester thereof but excluding 4-(phenylmethylene)-3,5-heptanedione, 4-(3,4-dichloroprenylmethylene)-3,5-heptanedione, 3-[(2-chloro-5-nitrophenyl)methylene]-2,4-pentanedione, and 3-(phenylmethylene)-2,4-hexanedione.

The term "alkyl" as employed herein refers to an alkyl group having preferably 1 to 2 carbon atoms.

The term "alkoxy" as employed herein refers to an alkyl residue as defined above linked to an oxygen atom.

The term "aralkyl" as employed herein refers to an alkyl group as defined above having aryl, preferably phenyl as the substituent.

The term "aryl" as employed herein refers to a carbocyclic aromatic group containing from 6 to 10 carbon atoms in the ring portion. The group may be a single ring or fused rings. Specific examples are phenyl and naphthyl.

The term "heteroaryl" as employed herein refers to a monocyclic or bicyclic aromatic group of carbon atoms containing one or more heteroatoms such as nitrogen, oxygen or sulfur.

The preparation of compounds according to the formulae I and II may be carried out in a similar way as described in EP-A-0357403. For example the compounds of formula I may be prepared by reacting a compound of the formula IV




in which R_1 and R_2 are as defined above with a compound of formula V




17 AUG 1969

- 6 -

in which n and R are as defined above and Z is CHO or $-\text{CH}_2-\text{Q}$, wherein Q is halogen or some other activated group, in the presence of acidic or basic catalyst to produce a compound of formula I or a compound of formula VI



in which n , R_1 , R_2 and R are as defined above, which compound is halogenated to give a compound of the formula VII




in which n , R_1 , R_2 and R are as defined above and X is halogen and dehydrohalogenating compound VII to produce a compound of the formula I.

The basic catalyst may be for example an organic or inorganic base. The acidic catalyst may be for example a mineral or sulfonic acid. The activated group Q may be a halogen or an alkyl or aryl sulfonate.

When halogenating a compound of formula VI elementary halogen, preferably chlorine or bromine may be used, or another known halogenating agent such as sulfuryl chloride may be used.

Salts of these compounds may be prepared by known methods. Most commonly used physiologically acceptable salts are sodium, potassium, ammonium, calcium and magnesium salts and hydrochlorides or hydrobromides. Preferred esters of the



- 7 -

compounds of formula I are acyl or aroyl derivatives which will hydrolyse under physiological conditions.

The above compounds may be formulated to dosage forms using the principles which are known to those having average skill in the art. Suitable solvents, gel forming ingredients, dispersion agents, antioxidants and colourants may be added in a normal way.

It is preferred to administer the compounds enterally. When they are given orally, they may be in the form of tablets, granules, capsules, emulsions, suspensions or solutions. It is often advisable to use coated tablets or granules, i.e. so called enterocoated preparations, to ensure that the medicine reaches the desired part of the gastrointestinal tract. Alternatively, the compositions may be given rectally in the form of enemas or suppositories.

The effective dose varies considerably depending on location, degree and severity of the disease being treated as well as the age and the general condition of the patient. The effective dose is generally from about 20 to 1000 mg per day, preferably from 50 to 250 mg per day for an adult, once a day or divided into two to five doses.

In vivo tests

The efficacy of the compounds can be demonstrated by using a model of chronic colonic inflammation in the rat, TNB-induced chronic colitis. TNB-induced chronic colonic inflammation in rats is an animal model which has been shown to resemble highly human IBD.

17 AUG 1999

- 8 -

20 mg of TNB (2,4,6-trinitrobenzenesulfonic acid) in 50 % ethanol was administered into the lumen of the colon via a rubber catheter inserted rectally to Wistar rats. The test compounds were given rectally 1 h prior and 24, 48, 72 and 95 h after TNB. The rats were sacrificed 96 h after TNB and their colons were scored according to occurrence of ulcers and inflammation. The number of animals in each group was at least 8. 5-ASA (5-aminosalicylic acid), a clinically used drug for the treatment of IBD was used as a reference compound. The results are presented in the Table 1.

Table 1.

The effect of test compounds on the damage score of chronic inflammation induced by TNB.

Test compound	Dose mg/kg	% inhibition against control
5-ASA	100	24
1	30	37
2	30	28
3	10	18
	30	57
4	30	54
5	30	43
6	30	41
7	10	60
	30	48
8	10	51
	30	62
9	30	40
10	10	27
11	30	32
12	30	52
13	30	58
14	30	30

17 AUG 1999

- 10 -

Test compounds

- 1 3-(4-hydroxyphenyl)methylene-2,4-pentanedione
- 2 3-(4-methoxyphenyl)methylene-2,4-pentanedione
- 3 3-(3-nitrophenyl)methylene-2,4-pentanedione
- 4 3-(4-trifluoromethylphenyl)methylene-2,4-pentanedione
- 5 3-(4-nitrophenyl)methylene-2,4-pentanedione
- 6 3-(2-nitrophenyl)methylene-2,4-pentanedione
- 7 3-(4-cyanophenyl)methylene-2,4-pentanedione
- 8 3-(3-cyanophenyl)methylene-2,4-pentanedione
- 9 3-[(4-(N-phenethyl)carboxamido-phenyl)methylene]-
2,4-pentanedione
- 10 3-[(5-nitrofuryl)methylene]-2,4-pentanedione
- 11 4-[(4-cyanophenyl)methylene]-3,5-heptanedione
- 12 2-[(4-cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione
- 13 2-[(3-trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione
- 14 2-[(4-carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione

The test compounds significantly decreased the ulceration and inflammation of the colons. The test compounds were much more potent than the reference compound 5-ASA and should therefore be more effective than 5-ASA in the treatment of human IBD.

17 AUG 1999

- 11 -

The following examples illustrate the production of compounds of use according to the invention.

Example 1

3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione

To a solution containing 2.1 g of 4-phenylsulfonylbenzaldehyde and 1.5 g of 2,4-pentanedione in 10 ml of 2-propanol was added 1 ml of thionyl chloride with stirring at 20°C. The solution was stirred for 1h at 20°C and evaporated to dryness in vacuo. The residue was crystallized from toluene, mp 102-105°C.

Example 2

3-[(4-Methylsulfonylphenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 1.1 g of 4-methylsulfonylbenzaldehyde and 1.5 g of 2,4-pentanedione. The product was crystallized from ether, mp 139-140°C.

Example 3

3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione

A mixture containing 1.8 g of 4-carboxamidobenzaldehyde, 1.1 g of 2,4-pentanedione, 0.2 g of piperidine and 0.2 g of acetic acid in 40 ml of toluene was refluxed for 3 h by using a Dean-Stark separator. The warm clean toluene solution was decanted from the tarry residue and cooled. The crystalline product was filtered and dried. Mp 154-155°C.

17 AUG 1997

- 12 -

Example 4

4-[(4-Nitrophenyl)methylene]-3,5-heptanedione

The procedure described in Example 1 was repeated by using 1.5 g of 4-nitrobenzaldehyde and 2.0 g of 3,5-heptanedione. Yield 1.7 g, yellow oil.

Example 5

3-[(3,4-Dichlorophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 4.26 g of 3,4-dichlorobenzaldehyde and 3.0 g of 2,4-pentanedione. Yield 1.3 g, mp 73°C.

Example 6

3-[(4-Chloro-3-nitrophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 1.85 g of 4-chloro-3-nitrobenzaldehyde and 1.5 g of 2,4-pentanedione. Yield 1.32 g, mp 110-111°C.

Example 7

3-[(4-(N-Phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.

The procedure described in Example 1 was repeated by using 2.5 g of 4-(N-phenethyl)carboxamidobenzaldehyde and 1.0 g of 2,4-pentanedione. Yield 0,5 g, mp 146-149°C.

Example 8

3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione

To a solution containing 1.84 g of 2-nitroterephthalaldehyde and 1.00 g of 2,4-pentanedione in 10 ml trifluoroacetic acid were added 0.9 ml of thionyl chloride and a catalytic amount of water. The solution was stirred for 2 h at 20°C. 30 ml of water was added and the mixture was extracted with ether. The ether extract was washed several times with Na₂CO₃-solution and dried over K₂CO₃. After filtration the solvent was evaporated in vacuo. The product was purified by column chromatography. Yield 0.2 g, mp 102-104°C.

Example 9


3-[(3,5-Dinitro-4-hydroxyphenyl)methylene]-2,4-pentanedione.

The procedure described in Example 1 was repeated by using 2.12 g of 3,5-dinitro-4-hydroxybenzaldehyde and 1.5 g of 2,4-pentanedione. Yield 1.9 g, mp 158°C.

Example 10

3-[(3,5-Dinitrophenyl)methylene]-2,4-pentanedione

The procedure described in Example 1 was repeated by using 2.53 g 3,5-dinitrobenzaldehyde and 2.43 g of 2,4-pentanedione. Yield 0.8 g, mp 101-105°C.



- 14 -

Example 11

3-[3-(4-Methoxyphenyl)propenylene]-2,4-pentanedione

To a solution containing 2.45 g of 4-methoxycinnamaldehyde in 90 ml of toluene was added 2.2 g of 2,4-pentanedione, 0.26 g of piperidine and 0.45 g of acetic acid. The solution was refluxed in N₂ atmosphere for 1 h and the water formed in the reaction was azeotropically distilled off by using a Dean-Stark water separator. The solution was evaporated to dryness in vacuo. The residue was trifurated with ether. The crystals were filtered. Yield 0.65 g, mp 108-110°C.

Example 12


3-[(5-Nitrofuryl)methylene]-2,4-pentanedione

A solution containing 2.43 g of 5-nitro-2-furanecarboxaldehyde diacetate, 1.5 g of 2,4-pentanedione and 0.1 g of water in 5 ml of tetrahydrofuran was saturated with hydrogen chloride gas at 20°C. The mixture was stirred for 1 h and evaporated to dryness in vacuo. The product was recrystallized from acetic acid. Yield 0.1 g, mp 117-120°C.

Example 13

3-[(5-Nitrothienyl)methylene]-2,4-pentanedione

The procedure described in Example 12 was repeated by using 2.59 g of 5-nitro-2-thiophenecarboxaldehyde diacetate, 1.5 g of 2,4-pentanedione. Yield 0.5 g, yellow viscous oil.



- 15 -

Example 14

3-(3-Pyridylmethylene)-2,4-pentanedione

The procedure described in Example 3 was repeated by using 3.21 g of pyridine-3-carboxaldehyde and 3.3 g of 2,4-pentanedione. Yield 2.95 g, mp 52-57°C.

Example 15

3-(4-Pyridylmethylene)-2,4-pentanedione hydrochloride

A solution containing 1.1 g of pyridine-4-carboxaldehyde and 1.0 g of 2,4-pentanedione in 10 ml of N,N-dimethylformamide was saturated with hydrogen chloride gas at 20°C. The solution was stirred for 1 h at 20°C and then for 2 h at 80°C. The solvent was evaporated in vacuo and the residue was crystallized from ethanol. Yield 0.17 g, mp 175-186°C.

Example 16


3-(3-Quinolylmethylene)-2,4-pentanedione hydrochloride

The procedure described in Example 3 was repeated by refluxing 2.4 g quinoline-3-aldehyde, 6.6 g 2,4-pentanedione, 0.4 g piperidine and 0.8 g acetic acid in 100 ml toluene. Yield 1.4 g, mp 99-101°C.

Example 17

3-[(3-Carboxy-4-hydroxyphenyl)methylene]-2,4-pentanedione

A mixture containing 2 g 3-carboxy-4-hydroxybenzaldehyde, 4 ml 2,4-pentanedione and 20 ml tetrahydrofuran was saturated with hydrogen chloride gas. The solution was stirred for 20 h



- 16 -

at 50°C and evaporated to dryness in vacuo. The residue was crystallized from ethyl ether, mp. 112-114°C.

Example 18

3-[[4-(2-Phenylethoxy)phenyl]methylene]-2,4-pentanedione

The procedure described in example 8 was repeated by using 5.9 g of 4-(2-phenylethoxy)benzaldehyde, 3.9 ml of 2,4-pentanedione, 26 ml of trifluoroacetic acid and 2.1 ml of thionyl chloride. Yield 4.7 g, an viscous oil.

Example 19

2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

A mixture containing 1.6 g 4-acetamidobenzaldehyde, 1.5 g 1,3-dicyclopropyl-1,3-propanedione, 0.2 ml of piperidine and 0.4 ml of acetic acid in 50 ml of toluene was refluxed for 3 h by using a Dean-Stark separator. After standing over night at room temperature the crystals were filtered and washed with toluene. The product was recrystallized from ethanol. Yield 2.0 g, mp 166-167°C.

Example 20

2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 0.68 g 4-methoxybenzaldehyde and 0.77 g 1,3-dicyclopropyl-1,3-propanedione. Yield 0.46 g, mp 60.5-61.5°C.

17 AUG 1969

- 17 -

Example 21

4-[(3-Cyanophenyl)methylene]-3,5-heptanedione

The procedure described in Example 19 was repeated by using 1.82 g 3-cyanobenzaldehyde and 3.5 g 3,5-heptanedione. Yield 0.4 g, mp 51-52°C.

Example 22

4-[(4-Cyanophenyl)methylene]-3,5-heptanedione

The procedure described in Example 19 was repeated by using 2.6 g 4-cyanobenzaldehyde and 5.0 g 3,5-heptanedione. Yield 2.8 g, mp 59-60°C.

Example 23

3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione

The procedure described in Example 19 was repeated by using 2.6 g 4-cyanobenzaldehyde and 2.5 g 4-cyclopropyl-2,4-butanedione. The product was purified by column chromatography. Yield 0.37 g, mp 83-85°C.

Example 24

2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 3.78 g 4-cyanobenzaldehyde and 2.23 g 1,3-dicyclopropyl-1,3-propanedione. Yield 1.5 g, mp 100-102°C.

17 AUG 1999

- 18 -

Example 25

3-[(5-Bromothieryl)methylene]-2,4-pentanedione

To a solution containing 5.73 g 5-bromothiophene-2-carboxaldehyde and 5.0 g 2,4-pentanedione in 25 ml of 2-propanol, 2.9 ml thionylchloride was gradually added. The mixture was stirred for 2 h at 20°C. 120 ml of conc. Na₂SO₄-solution was then added and the mixture was extracted with ether. The solvent was evaporated in vacuo and the crystalline residue was washed with cold ether. Yield 3.9 g, mp 70-74°C.

Example 26

2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 2.15 ml 3-trifluoromethylbenzaldehyde and 2.42 g 1,3-dicyclopropyl-1,3-propanedione. Yield 2.71 g, mp 56-58°C.

Example 27

2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione

The procedure described in Example 19 was repeated by using 3.0 g 4-carboxybenzaldehyde and 3.06 g 1,3-dicyclopropyl-1,3-propanedione. Yield 1.7 g, mp 175-178°C.




REPLACEMENT SHEET

17 AUG 1999





- 19 -

CLAIMS

1. Use of a compound of formula I



wherein $n=0$ or 1 , R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group;




wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido or;



wherein $m=0, 1$ or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl; in which "alkyl" is an alkyl group having preferably 1 to 2 carbon atoms, "alkoxy" is an alkyl residue as defined above linked to an oxygen atom, and "aralkyl" is an alkyl group as defined above having aryl, preferably phenyl as the substituent; Y is oxygen or sulphur and X_4 is hydrogen, nitro or halo; or a pharmaceutically acceptable salt or ester thereof in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.

2. The use according to claim 1, in which, in the compound of formula I, X_1 is hydroxy or alkoxy.



- 20 -

3. The use according to claim 1 or 2, in which, in the compound formula I, one of X_1 and X_2 is nitro, cyano, trifluoromethyl, carboxy or carbamoyl which is substituted by aryl.

4. The use according to claims 1 to 3, in which the compound is

3-(4-hydroxyphenyl)methylene-2,4-pentanedione,

3-(4-methoxyphenyl)methylene-2,4-pentanedione,

3-(3-nitrophenyl)methylene-2,4-pentanedione,

3-(4-nitrophenyl)methylene-2,4-pentanedione,

3-(2-nitrophenyl)methylene-2,4-pentanedione,

3-[(4-(N-phenethyl)carboxamido-phenyl)methylene]-
2,4-pentanedione,

3-[(5-nitrofuryl)methylene]-2,4-pentanedione, or

2-[(4-carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione or a pharmaceutically acceptable salt or ester
thereof.

5. The use according to claims 1 or 3, in which, the compound is 3-(4-trifluoromethylphenyl)methylene-2,4-pentanedione.

6. The use according to claims 1 or 3, in which, the compound is 3-(4-cyanophenyl)methylene-2,4-pentanedione.

7. The use according to claims 1 or 3, in which, the compound is 3-(3-cyanophenyl)methylene-2,4-pentanedione.

8. The use according to claims 1 or 3, in which, the compound is 4-[(4-cyanophenyl)methylene]-3,5-heptanedione.

9. The use according to claims 1 or 3, in which, the compound is 2-[(4-cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-
propanedione.




17 AUG 1999


- 21 -

10. This use according to claims 1 or 3, in which, the compound is 2-[(3-trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.


11. A compound according to the general formula II




wherein $n=0$ or 1, R_1 is methyl, ethyl or cyclopropyl and R_2 is ethyl or cyclopropyl and R is



wherein X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl; X_2 and X_3 are independently hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy, acetamido, provided that when $n=0$, X_1 , X_2 and X_3 are not all independently selected from the group consisting of hydrogen, hydroxy and nitro; or R is



wherein $m=0$, 1 or 2 and R_3 is alkyl, aryl or aralkyl; or carbamoyl which is optionally substituted by alkyl, aryl or aralkyl or R_1 and R_2 are methyl and R is



where X_1 , X_2 and X_3 are as defined above provided that one of X_2 and X_3 is carbamoyl which is optionally substituted by



17 AUG 1999

alkyl, aryl or aralkyl, in which "alkyl", "alkoxy" and "aralkyl" are as defined herein; or a pharmaceutically acceptable salt or ester thereof, but excluding 4-(phenylmethylene)-3,5-heptanedione, 4-(3,4-dichlorophenylmethylene)-3,5-heptanedione and 3-(phenylmethylene)-2,4-hexanedione.

12. A compound according to claim 11 wherein in the compound of formula II one of X_2 and X_3 is cyano or trifluoromethyl.

13. 2-[(4-Acetamidophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

14. 2-[(4-Methoxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

15. 3-[(4-Cyanophenyl)methylene]-4-cyclopropyl-2,4-butanedione.

16. 2-[(4-Cyanophenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

17. 2-[(3-Trifluoromethylphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione.

18. 2-[(4-Carboxyphenyl)methylene]-1,3-dicyclopropyl-1,3-propanedione or a pharmaceutically acceptable salt or ester thereof.

19. 4-[(4-Nitrophenyl)methylene]-3,5-heptanedione.

20. 4-[(3-Cyanophenyl)methylene]-3,5-heptanedione.


21. 4-[(4-Cyanophenyl)methylene]-3,5-heptanedione.

22. 3-[(4-Carboxamidophenyl)methylene]-2,4-pentanedione.


23. 3-[(4-(N-phenethyl)carboxamidophenyl)methylene]-2,4-pentanedione.



24. A compound according to the general formula III



wherein R_1 is methyl, n is 0 or 1, X_1 is hydrogen, hydroxy or alkoxy which is optionally substituted by aryl, and X_2 is;




wherein m is 1 or 2 and R_3 is alkyl, aryl or aralkyl, and X_3 is hydrogen, nitro, cyano, halo, trifluoromethyl, formyl, carboxy or a group as defined for X_2 ; or X_2 and X_3 are both cyano; or X_2 is nitro and X_3 is formyl; or when X_1 is hydroxy, X_2 may be carboxy or X_2 and X_3 may both be hydrogen when $n=0$ and X_1 is alkoxy substituted by aryl, or when $n=1$ and X_1 is hydroxy or alkoxy optionally substituted by aryl; in which "alkyl" is an alkyl group having preferably 1 to 2 carbon atoms, "alkoxy" is an alkyl residue as defined above linked to an oxygen atom, and "aralkyl" is an alkyl group as defined above having aryl, preferably phenyl as the substituent; or a pharmaceutically acceptable salt or ester thereof.


25. 3-[(4-Phenylsulfonylphenyl)methylene]-2,4-pentanedione.

26. 3-[(4-Methylsulphonylphenyl)methylene]-2,4-pentanedione.

27. 3-[(4-Formyl-3-nitrophenyl)methylene]-2,4-pentanedione.



28. 3-[(3-Carboxy-4-hydroxy)methylene]-2,4-pentanedione a pharmaceutically acceptable salt or ester thereof.
29. 3-[[[4-(2-Phenethyloxy)phenyl)methylene]-2,4-pentanedione.
30. 3-[3-(4-Methoxyphenyl)propenylene]-2,4-pentanedione.
31. A compound according to any one of claims 11 to 30 for use as a medicament.
32. A compound according to any one of claims 11 to 30 for use in the treatment of inflammatory bowel disease.
33. 3-[(4-methylsulphonylphenyl)methylene]-2,4-pentanedione for use as a medicament.
34. A pharmaceutical composition comprising a compound according to any one of claims 11 to 30 and a pharmaceutically acceptable carrier or diluent.
35. Use of a compound as defined in any one of claims 11 to 30 for the manufacture of a medicament for use in the treatment of inflammatory bowel disease.
36. Use of a 3-[(4-methylsulphonylphenyl)methylene]-2,4-pentanedione in the manufacture of a medicament for use in the treatment of inflammatory bowel disease.
37. The use according to any one of claims 1 to 10 or 35 or 36, in which the inflammatory bowel disease is ulcerative colitis.
38. The use according to any one of claims 1 to 10 or 35 or 36, in which the inflammatory bowel disease is Crohn's disease.




- 25 -

ABSTRACT

SUBSTITUTED β -DIKETONES AND THEIR USE

Compounds of formula



wherein n is 0 or 1, R_1 and R_2 are independently methyl, ethyl or cyclopropyl and R is an optionally substituted phenyl or heteroaryl group or salt or ester thereof are useful in the treatment of inflammatory bowel disease.