

METHODS FOR TREATING OSTEOGENESIS IMPERFECTA

RELATED APPLICATIONS

This application is related to U.S. Provisional Patent Application No.
5 61/803,647, filed March 20, 2013, U.S. Provisional Patent Application No.
61/875,399, filed September 9, 2013, and U.S. Provisional Patent Application No.
61/883,151, filed October 26, 2013, the contents of each of which is incorporated
herein by reference in its entirety.

GOVERNMENT FUNDING

This invention was made with government support under P01 HD070394, P01
HD22657& R01 DE01771 awarded by the National Institutes of Health. The United
States government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to methods for treating osteogenesis imperfecta
(OI). More specifically, the invention relates to methods for treating OI using a
binding protein, e.g., an antibody or antigen binding fragment thereof, which
specifically binds to human transforming growth factor beta (TGF β) or isoforms
20 thereof.

BACKGROUND

Osteogenesis imperfecta (OI), also known as “brittle bone disease” or
Lobstein syndrome, is a debilitating and rare congenital bone disease that affects
25 about one in every 15,000 people. Though phenotypes vary among OI types,
common symptoms include incomplete ossification of bones and teeth, reduced bone
mass, brittle bones, and pathologic fractures. These common symptoms of OI are
thought to be caused by gene mutations which result in deficiencies in Type-I
collagen or other proteins involved in bone matrix deposition or homeostasis. As a
30 result of these symptoms and the propensity for fatal bone fractures and
complications, life expectancy of OI patients is reduced as compared to the general
population. Accordingly, there clearly exists an urgent need in the art to develop
effective treatments for OI.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A (Western blot) and **1B-1F** (graphs) demonstrate that TGF β signaling is elevated in bones from *Crtap*^{-/-} mice, as compared to wild type controls.

Figures 2A (luminescent picture) and **2B-C** (graphs) show that *Crtap*^{-/-} mice
5 crossed with TGF β reporter mice exhibit higher TGF β activity compared to wild type controls.

Figure 3 is a μ CT image of vertebrae from *Crtap*^{-/-} mice treated with the mouse pan-specific anti-TGF β antibody 1D11.

Figure 4 is a graph that contains quantitative measurements of vertebrae from
10 *Crtap*^{-/-} mice treated with the mouse pan-specific anti-TGF β antibody 1D11.

Figure 5 is a histomorphometric analysis of vertebrae from *Crtap*^{-/-} mice treated with the mouse pan-specific anti-TGF β antibody 1D11.

Figure 6A and **6B** are graphs that demonstrates the biomechanical properties (Figure 6A – maximum load, and Figure 6B – stiffness) of femurs from *Crtap*^{-/-} mice
15 treated with the mouse pan-specific anti-TGF β antibody 1D11, as determined by a three-point bending test.

Figure 7A and **7B** (micrograph images) and **7C** (graph) demonstrate the lung phenotype of *Crtap*^{-/-} mice treated with the mouse pan-specific anti-TGF β antibody 1D11.

Figure 8 are micrograph images of *Crtap*^{-/-} and WT lungs stained with an anti-decorin antibody.
20

Figure 9 is a graph depicting a decorin binding assay.

Figure 10 is a group of graphs, photos, and images that show increased TGF β signaling in *Crtap*^{-/-} mice. **Figure 10A** is a series of three graphs that show the results
25 of quantitative RT-PCR of TGF β target genes *p21*, *PAI-1*, and *Col1a1*. The graphs indicate increased TGF β signaling in calvarial bone of P3 WT and *Crtap*^{-/-} mice. Results are shown as fold change of the mean of WT group $\pm$ SD; n=5 per group, *p<0.05. **Figure 10B** is a photograph of a Western blot analysis of P3 calvarial protein extracts, which shows increased amounts of activated Smad2 (pSmad2)
30 relative to total Smad2 protein in *Crtap*^{-/-} versus WT mice, suggesting increased TGF β -signaling; n=3 per group. **Figure 10C** is a graph showing the quantification of the Western blot shown in Figure 10B. Results are shown as fold change of the mean of WT group $\pm$ SD, *p<0.05. **Figure 10D** is an image showing increased

bioluminescence in regions that overlap with skeletal structures in *Crtap*^{-/-} compared with WT mice that were intercrossed to TGFβ-reporter mice (SBE-Luc mice). Representative image of 3 litters at P10 is shown. In 3 litters *Crtap*^{-/-} mice show a mean 2.86 fold (SD±0.34) bioluminescence signal at the head/calvaria compared with WT mice (scale bar=1cm). **Figure 10E** is a graph that shows, using a TGFβ reporter cell line, TGFβ activity was assessed in conditioned medium from WT and *Crtap*^{-/-} bone marrow stromal cells cultured under osteogenic conditions for 3 days, demonstrating greater TGFβ activity compared with the medium from WT cells. Results are shown as fold change of the mean of WT group±SD, n=5 per group, *p<0.05. **Figure 10F** is two pictures showing immunostaining of lungs (P10) for pSmad2, which shows increased intracellular staining in WT and *Crtap*^{-/-} mice (40X magnification). Representative images of n=3 mice per group are shown (scale bar=20μm).

Figure 11 is a group of photos and graphs showing phenotypic correction of *Crtap*^{-/-} mice after treatment with the TGFβ neutralizing antibody 1D11. **Figure 11A** is a series of three MicroCT images of L4 vertebral bodies of 16-week-old wildtype (WT), control antibody-treated *Crtap*^{-/-}, and 1D11-treated *Crtap*^{-/-} mice after treatment for 8 weeks. **Figure 11B** is a group of three graphs showing the results of MicroCT of L4 vertebral bodies, which demonstrates increased bone volume/total volume (BV/TV), trabecular number (Tb.N), and thickness (Tb.Th) in WT, control *Crtap*^{-/-} and 1D11 treated *Crtap*^{-/-} mice. Results are shown as means±SDs, n=8 per group, *p<0.05 for *Crtap*^{-/-} 1D11 vs. *Crtap*^{-/-} control, +p<0.05 for *Crtap*^{-/-} vs. WT. **Figure 11C** is a group of three graphs showing the results of histomorphometric analysis of L4 vertebrae, which shows increased osteoclast (N.Oc/BS) and osteoblast (N.Ob/BS) numbers per bone surface in *Crtap*^{-/-} mice compared with WT. Reduced osteoblast and osteoclast numbers after treatment with 1D11 indicates effective suppression of accelerated bone remodeling in *Crtap*^{-/-} mice. Increased numbers of osteocytes per bone area (N.Ot/B.Ar) in *Crtap*^{-/-} mice are reduced to WT level after 1D11 treatment. Results are shown as means±SDs, n=6 per group, *p<0.05 for *Crtap*^{-/-} 1D11 vs. *Crtap*^{-/-} control, +p<0.05 for *Crtap*^{-/-} vs. WT. **Figure 11D** is a series of three pictures showing hematoxylin/eosin staining of inflated lungs of 16-week-old wildtype (WT), control *Crtap*^{-/-}, and 1D11-treated *Crtap*^{-/-} mice after treatment for 8 weeks. *Crtap*^{-/-} control mice show an increase in distal airway space compared with WT mice. After

treatment with 1D11, there is a reduction of the distal airway diameter compared with control *Crtap*^{-/-} mice. Representative images of n=8 mice per group are shown (scale bar=100 μm). **Figure 11E** is a graph showing quantification of the distance between alveolar structures by the mean-linear-intercept (MLI) method, which demonstrates a significant reduction of the distal airway space in 1D11-treated *Crtap*^{-/-} mice compared with control antibody-treated *Crtap*^{-/-} and WT mice. Results are shown as means±SDs, n=8 mice per group, 10 images analyzed per mouse, *p<0.05 for *Crtap*^{-/-} 1D11 vs. *Crtap*^{-/-} control, +p<0.05 for *Crtap*^{-/-} vs. WT.

Figure 12 is a series of graphs that show that decorin binding to type I collagen overlaps the P986 3Hyp site and is reduced in type I collagen of *Crtap*^{-/-} mice. **Figure 12A** is a group of three graphs that show the results of quantitative RT-PCR of calvarial bone of P3 mice, which shows no difference in RNA expression of the small leucine-rich proteoglycans decorin (Dcn), biglycan (Bgn), and asporin (Aspn) in calvarial bone of *Crtap*^{-/-} mice compared with WT. Results are given as fold change of the mean of WT group±SD, n=5 per group. **Figure 12B** is a graph that shows the results of surface plasmon resonance analysis, which indicates that binding of recombinant decorin core protein to type I collagen of *Crtap*^{-/-} mice is approximately 45% less compared with WT type I collagen. Three independent experiments using 3, 5, and 12 μM of decorin were performed. Response units (RU) of total amount decorin bound normalized to type I collagen immobilized on the chip are shown. Mean reduction of decorin binding to *Crtap*^{-/-} type I collagen is 44.6 ± 7.9 %.

Figure 13 is a series of graphs and photos showing inhibition of increased TGFβ signaling improves the bone phenotype in a mouse model of dominant OI resulting from a G610C mutation in the *Coll1a2* gene (*Coll1a2*^{tm1.1Mcb}). **Figure 13A** is two graphs showing the results of quantitative RT-PCR of TGFβ target genes *p21* and *PAI-1*, which indicates increased TGFβ signaling in calvarial bone of P3 WT and *Coll1a2*^{tm1.1Mcb} mice. Results are shown as fold change of the mean of WT group±SD; n=3 per group, *p<0.05. **Figure 13B** is a photo of the results of Western blot analysis, which shows increased levels of activated Smad2 (pSmad2) relative to total levels of Smad2 protein in P3 calvaria of WT and *Coll1a2*^{tm1.1Mcb} mice compared with WT, suggesting increased TGFβ-signaling; n=3 per group. **Figure 13C** is a graph showing the quantification of the Western blot seen in Figure 13B. Results are

shown as fold change of the mean of WT group $\pm$ SD; * $p < 0.05$. **Figure 13D** are a series of photos of MicroCT images of L4 vertebral bodies of 16-week-old wildtype (WT), control antibody-treated *Colla2^{tm1.1Mcb}* and 1D11-treated *Colla2^{tm1.1Mcb}* mice after treatment for 8 weeks. **Figure 13E** is a series of graphs of data from MicroCT of L4 vertebral bodies, which shows increased bone volume/total volume (BV/TV), trabecular number (Tb.N) and thickness (Tb.Th) in *Colla2^{tm1.1Mcb}* mice after treatment with 1D11. Results are shown as means $\pm$ SDs, $n = 6$ per group, * $p < 0.05$ for *Colla2^{tm1.1Mcb}* 1D11 vs. *Colla2^{tm1.1Mcb}* control, + $p < 0.05$ for *Colla2^{tm1.1Mcb}* vs. WT.

Figure 14 is a weight curve graph, which shows a reduced weight of *Crtap^{-/-}* mice compared with WT during the study period ($p < 0.05$ for all time points, means $\pm$ SEs are shown). No statistically significant difference in weight was observed in 1D11-treated *Crtap^{-/-}* mice compared to control *Crtap^{-/-}* mice.

Figure 15 is a series of graphs and tables showing no effect of TGF β inhibition on the abnormal type I collagen post-translational modification in *Crtap^{-/-}* mice. **Figure 15A** is a series of three graphs showing tandem mass spectra of extracted type I collagen from tibia of WT, control *Crtap^{-/-}*, and 1D11-treated *Crtap^{-/-}* mice (16 week old mice, after treatment for 8 weeks). The sequence in the top graph is SEQ ID NO: 19, the sequence in the middle graph is SEQ ID NO: 20, and the sequence in the bottom graph is SEQ ID NO: 21. **Figure 15B** is a table showing a summary of tandem mass spectra analyses. 1D11 treatment did not significantly affect 3-hydroxylation status of collagen residue Pro986 alpha 1(I) in bone samples. Mean of percentage of 3-hydroxylated residues ($\pm$ SD) is shown, $n = 5$ per group.

Figure 15C is a group of three graphs showing that bone type I collagen of control *Crtap^{-/-}* and 1D11-treated *Crtap^{-/-}* mice exhibit changes in hydroxylysyl pyridinoline (HP) and lysyl pyridinoline crosslinks (LP) levels and an increased HP/LP ratio compared with WT mice. 1D11 treatment of *Crtap^{-/-}* mice did not significantly affect these parameters compared to control *Crtap^{-/-}* mice. Results are given as means $\pm$ SDs, $n = 4$ mice per group, + $p < 0.05$ for *Crtap^{-/-}* vs. WT.

Figure 16 is a series of graphs showing serum bone-turnover markers osteocalcin (OCN) and C-terminal cross-linked telopeptide of bone collagen (CTX) at start (Figure 16A = 8 weeks of age) and end of the treatment study (Figure 16B = 16 weeks of age). **Figure 16A** is two graphs showing increased OCN and CTX serum levels in 8 week old *Crtap^{-/-}* compared with WT mice at the start of the study indicate

increased bone turnover in *Crtap*^{-/-} mice. Results are given as means±SDs, n=8 for WT, n=14 for *Crtap*^{-/-} mice, +p<0.05 for *Crtap*^{-/-} vs. WT. **Figure 16B** is two graphs that show that at 16 weeks of age 1D11-treated *Crtap*^{-/-} mice show a trend to reduced serum OCN and significantly reduced CTX serum levels compared with control *Crtap*^{-/-} mice, indicating a suppression of increased bone turnover by inhibition of TGFβ. Results are given as means±SDs, n=8 for WT, n=7 per *Crtap*^{-/-} group; *p<0.05 for *Crtap*^{-/-} 1D11 vs. *Crtap*^{-/-} control, +p<0.05 for *Crtap*^{-/-} vs. WT.

Figure 17 is a table showing the results of MicroCT analyses of vertebral body L4 of WT, control *Crtap*^{-/-}, and 1D11 treated *Crtap*^{-/-} mice (16 week old mice, after treatment for 8 weeks). Means±SDs are shown for bone volume/tissue volume (BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), and bone mineral density of bone volume (BMD BV); n=8 per group, + indicates Kruskal-Wallis one-way ANOVA on ranks where the equal variance test failed. n.s.=not statistically significant.

Figure 18 is a table showing the results of MicroCT analyses of trabecular bone in proximal femurs for WT, control *Crtap*^{-/-}, and 1D11- treated *Crtap*^{-/-} mice (16 week old mice, after treatment for 8 weeks). Means±SDs are shown for bone volume/tissue volume (BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), and bone mineral density of bone volume (BMD BV); n=8 per group. + indicates Kruskal-Wallis one-way ANOVA on ranks where the equal variance test failed. n.s.=not statistically significant.

Figure 19 is a table showing the results of MicroCT analysis of cortical bone at the femur midshaft for WT, control *Crtap*^{-/-}, and 1D11-treated *Crtap*^{-/-} mice (16 week old mice, after treatment for 8 weeks). Means±SDs are shown for cortical thickness, bone mineral density of bone volume (BMD BV), anterior-posterior (a.p.) diameter, cross-sectional area (CSA), and cross-sectional moments of inertia (CSMI) for medio-lateral (m.l.) and anterior-posterior (a.p.) axis; n=8 per group. n.s.=not statistically significant.

Figure 20 is a table showing the results of biomechanical testing of femurs by 3-point bending (16 week old mice, after treatment for 8 weeks). Compared with WT mice, control *Crtap*^{-/-} mice exhibit significantly reduced biomechanical parameters except elastic modulus and elastic displacement. Anti TGFβ-treatment with 1D11 resulted in significant improvements of maximum load and ultimate strength in *Crtap*^{-/-}

^{-/-} femurs, indicating increased whole bone and tissue strength. However, no significant changes in post-yield displacement were observed, indicating that 1D11 did not affect the increased brittleness of the OI bone. N=6 for WT, n=4 for control *Crtap*^{-/-} and n=3 for 1D11 treated *Crtap*^{-/-} mice. n.s.=not statistically significant.

5 **Figure 21** is a table showing the results of histomorphometry analyses of L4 vertebral bodies of WT, control *Crtap*^{-/-}, and 1D11- treated *Crtap*^{-/-} mice (16 week old mice, after treatment for 8 weeks). Means±SDs are shown for bone volume/tissue volume (BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), number of osteoclasts/bone surface (N.Oc/BS), osteoclast
10 surface/bone surface (Oc.S/BS), number of osteoblasts/bone surface (N.Ob/BS), osteoblast surface/bone surface (Oc.S/BS), and number of osteocytes/bone area (N.Ot/B.Ar); n=6 per group. + indicates Kruskal-Wallis one way ANOVA on ranks where equal variance test failed. n.s.=not statistically significant.

Figure 22 is a table showing the results of MicroCT analyses of vertebral
15 body L4 of WT, control *Collα2*^{tm1.1Mcb} and 1D11 treated *Collα2*^{tm1.1Mcb} mice (16 week old mice, after treatment for 8 weeks). Means±SDs are shown for bone volume/tissue volume (BV/TV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular separation (Tb.Sp), and bone mineral density of bone volume (BMD BV); n=6 per group, n.s.=not statistically significant.

20 **Figure 23** is a graph showing surface plasmon resonance analysis measuring the binding of recombinant decorin core protein to type I collagen of WT and *Crtap*^{-/-} mice. Three technical replicates at each of the indicated concentrations of decorin were performed from two independent biological replicates (◆ replicate 1, ▲ replicate 2). Results are shown as the percentage of the mean of WT (bars indicate
25 mean per group).

Figure 24 (micrograph images) demonstrate immunostaining for decorin in the distal femur metaphysis of WT and *Crtap*^{-/-} mice at 20X, n=3 per genotype, scale bars=100µm (**Figures 24A-24C**) and 40X magnification, n=3 per genotype, scale bars=50µm (**Figures 24D-24F**). Control femurs were incubated in secondary
30 antibody only.

Figure 25 (micrograph images) demonstrate immunostaining for TGFβ1 in the distal femur metaphysis of WT and *Crtap*^{-/-} mice at 20X, n=3 per genotype, scale bars=100µm (**Figures 25A-25C**) and 40X magnification, n=3 per genotype, scale

bars=50µm (**Figures 25D-25F**). Control femurs were incubated in secondary antibody only.

Figure 26 (micrograph images) demonstrate immunostaining for TGFβ1 in the distal femur metaphysis of WT and *Collα2^{tm1.1Mcb}* mice at 20X, n=3 per genotype, scale bars=100µm (**Figures 26A-26C**) and 40X magnification, n=3 per genotype, scale bars=50µm (**Figures 26D-26F**). Control femurs were incubated in secondary antibody only.

SUMMARY OF THE INVENTION

The present invention relates to methods for effectively treating osteogenesis imperfecta (OI). More specifically, the invention relates to methods for treating OI using a binding protein such as antibody or an antigen binding fragment thereof that specifically binds to transforming growth factor beta (TGFβ) or an isoform thereof. Preferably, the binding protein is “pan-specific” and binds to all three human isoforms of TGFβ, i.e., TGFβ1, TGFβ2, and TGFβ3. More preferably, the binding protein specifically binds to and neutralizes human TGFβ1, TGFβ2, and TGFβ3. In one aspect, the invention provides a method for treating OI in a subject in need thereof comprising administering to the subject a therapeutically effective amount of an antibody or an antigen binding fragment thereof that specifically binds to TGFβ.

In one embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising three complementarity determining regions (CDRs) having amino acid sequences selected from the group consisting of SEQ ID NOs: 4, 5, and 6; and a light chain variable region comprising three CDRs having amino acid sequences selected from the group consisting of SEQ ID NOs: 7, 8, and 9.

In another embodiment, the antibody or antigen binding fragment thereof comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 10, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 11.

In one embodiment, the antibody or antigen binding fragment thereof further comprises a human IgG4 constant region. In one embodiment, the human IgG4 constant region comprises the amino acid sequence of SEQ ID NO: 12. In another embodiment, the antibody or antigen binding fragment thereof further comprises a

human κ light chain constant region. In another embodiment, the human κ light chain constant region comprises the amino acid sequence of SEQ ID NO: 13. In another embodiment, the antibody or antigen binding fragment thereof further comprises a human IgG4 constant region, and a human κ light chain constant region.

5 In another embodiment, the human IgG4 constant region comprises the amino acid sequence of SEQ ID NO: 12, and the human κ light chain constant region comprises the amino acid sequence of SEQ ID NO: 13. In another embodiment, the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14. In another embodiment, the antibody comprises a light chain comprising the amino acid sequence of SEQ ID NO: 15. In another embodiment, the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15.

 In another embodiment, the antibody or antigen binding fragment thereof binds to human TGF β 1, TGF β 2, and TGF β 3. In another embodiment, the antibody or antigen binding fragment thereof neutralizes human TGF β 1, TGF β 2, and TGF β 3.

 In another embodiment, the antibody or antigen binding fragment thereof improves a bone parameter selected from the group consisting of bone volume density (BV/TV), total bone surface (BS), bone surface density (BS/BV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular spacing (Tb.Sp), and total volume (Dens TV).

 In another embodiment, the antibody or antigen binding fragment thereof inhibits bone resorption.

 In another embodiment, the antibody or antigen binding fragment thereof reduces a serum biomarker of bone resorption selected from the group consisting of urinary hydroxyproline, urinary total pyridinoline (PYD), urinary free deoxypyridinoline (DPD), urinary collagen type-I cross-linked N-telopeptide (NTX), urinary or serum collagen type-I cross-linked C-telopeptide (CTX), bone sialoprotein (BSP), osteopontin (OPN), and tartrate-resistant acid phosphatase 5b (TRAP).

 In another embodiment, the antibody or antigen binding fragment thereof increases a serum biomarker of bone deposition selected from the group consisting of as total alkaline phosphatase, bone-specific alkaline phosphatase, osteocalcin, and type-I procollagen (C-terminal/N-terminal).

In another embodiment, the antibody or antigen binding fragment thereof inhibits bone resorption. In another embodiment, the antibody or antigen binding fragment thereof promotes bone deposition. In another embodiment, the antibody or antigen binding fragment thereof improves the function of a non-skeletal organ affected by OI selected from the group consisting of hearing function, lung function, and kidney function.

In another aspect, the invention provides a method for treating OI in a subject in need thereof comprising administering to the subject a therapeutically effective amount of an antibody or an antigen binding fragment thereof that binds to TGF β , wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15.

In another aspect, the invention provides a method for treating OI in a subject in need thereof comprising administering to the subject a therapeutically effective amount of an antibody or an antigen binding fragment thereof that binds to TGF β in combination with at least one therapeutic agent. In another embodiment, the agent is a bisphosphonate.

DETAILED DESCRIPTION OF THE INVENTION

A. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art.

It is noted here that as used in this specification and the appended claims, the singular forms "a", "an", and "the" also include plural reference, unless the context clearly dictates otherwise.

The term "about" or "approximately" means within 10%, and more preferably within 5% (or 1% or less), of a given value or range.

The terms "administer" or "administration" refer to the act of injecting or otherwise physically delivering a substance as it exists outside the body (*e.g.*, an antibody) into a patient, such as by mucosal, intradermal, intravenous, subcutaneous, intramuscular delivery, and/or any other method of physical delivery described herein or known in the art. When a disease, or a symptom thereof, is being treated, administration of the substance typically occurs after the onset of the disease or

symptoms thereof. When a disease or symptoms thereof, are being prevented, administration of the substance typically occurs before the onset of the disease or symptoms thereof.

5 An "antagonist" or "inhibitor" of TGF β refers to a molecule that is capable of inhibiting or otherwise decreasing one or more of the biological activities of TGF β , such as in a cell expressing TGF β or in a cell expressing a TGF β ligand, or expressing a TGF β receptor. In certain exemplary embodiments, antibodies of the invention are antagonist antibodies that inhibit or otherwise decrease the activity of TGF β in a cell having a cell surface-expressed TGF β receptor (*e.g.*, TGF β receptor 1, 2, or 3) when
10 said antibody is contacted with said cell. In some embodiments, an antagonist of TGF β (*e.g.*, an antibody of the invention) may, for example, act by inhibiting or otherwise decreasing the activation and/or cell signaling pathways of the cell expressing a TGF β receptor, thereby inhibiting a TGF β -mediated biological activity of the cell relative to the TGF β -mediated biological activity in the absence of
15 antagonist. In certain embodiments of the invention, the anti-TGF β antibodies are antagonistic anti-TGF β antibodies, preferably fully human, monoclonal, antagonistic anti-TGF β antibodies.

The terms "antibody", "immunoglobulin", or "Ig" may be used interchangeably herein. The term antibody includes, but is not limited to, synthetic
20 antibodies, monoclonal antibodies, recombinantly produced antibodies, multispecific antibodies (including bi-specific antibodies), human antibodies, humanized antibodies, chimeric antibodies, intrabodies, single-chain Fvs (scFv) (*e.g.*, including monospecific, bispecific, etc.), camelized antibodies, Fab fragments, F(ab') fragments, disulfide-linked Fvs (sdFv), anti-idiotypic (anti-Id) antibodies, and epitope-binding
25 fragments of any of the above. In particular, antibodies include immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, antigen binding domains or molecules that contain an antigen-binding site that specifically binds to a TGF β antigen (*e.g.*, one or more complementarity determining regions (CDRs) of an anti-TGF β antibody). The anti-TGF β antibodies can be of any
30 type (*e.g.*, IgG, IgE, IgM, IgD, IgA, and IgY), any class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2), or any subclass (*e.g.*, IgG2a and IgG2b) of an immunoglobulin molecule. In certain embodiments, the anti-TGF β antibodies are humanized, such as humanized monoclonal anti-TGF β antibodies. In other

embodiments, the anti-TGF β antibodies are fully human, such as fully human monoclonal anti-TGF β antibodies. In preferred embodiments, the anti-TGF β antibodies are IgG antibodies, such as IgG4 antibodies.

5 The terms "composition" and "formulation" are intended to encompass a product containing specified ingredients (*e.g.*, an anti-TGF β antibody) in, optionally, specified amounts, as well as any product which results, directly or indirectly, from the combination of specified ingredients in, optionally, specified amounts.

10 The terms "constant region" or "constant domain" refer to a carboxy terminal portion of the light and heavy chain that is not directly involved in binding of the antibody to antigen, but exhibits various effector functions, such as interaction with the Fc receptor. The terms refer to the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable domain, which contains the antigen binding site. The constant domain contains the CH1, CH2, and CH3 domains of the heavy chain, and
15 the CHL domain of the light chain.

The term "epitope" refers to a localized region on the surface of an antigen, such as a TGF β polypeptide or TGF β polypeptide fragment, that is capable of being bound to one or more antigen binding regions of an antibody, and that has antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a
20 human, that is capable of eliciting an immune response. An epitope having immunogenic activity is a portion of a polypeptide that elicits an antibody response in an animal. An epitope having antigenic activity is a portion of a polypeptide to which an antibody specifically binds, as determined by any method well known in the art, for example, such as an immunoassay. Antigenic epitopes need not necessarily be
25 immunogenic. Epitopes usually consist of chemically active surface groupings of molecules, such as amino acids or sugar side chains, and have specific three-dimensional structural characteristics, as well as specific charge characteristics. A region of a polypeptide contributing to an epitope may be contiguous amino acids of the polypeptide or the epitope may come together from two or more non-contiguous
30 regions of the polypeptide. The epitope may or may not be a three-dimensional surface feature of the antigen. In certain embodiments, a TGF β epitope is a three-dimensional surface feature of a TGF β polypeptide (*e.g.*, in a trimeric form of a TGF β polypeptide). In other embodiments, a TGF β epitope is a linear feature of a TGF β

polypeptide (*e.g.*, in a dimeric form or monomeric form of the TGF β polypeptide). Anti-TGF β antibodies may specifically bind to an epitope of the monomeric form of TGF β , an epitope of the dimeric form of TGF β , or both the monomeric form and the dimeric form of TGF β .

5 The term "excipients" refers to inert substances that are commonly used as a diluent, vehicle, preservative, binder, stabilizing agent, etc. for drugs and includes, but is not limited to, proteins (*e.g.*, serum albumin, etc.), amino acids (*e.g.*, aspartic acid, glutamic acid, lysine, arginine, glycine, histidine, etc.), fatty acids and phospholipids (*e.g.*, alkyl sulfonates, caprylate, etc.), surfactants (*e.g.*, SDS, polysorbate, nonionic
10 surfactant, etc.), saccharides (*e.g.*, sucrose, maltose, trehalose, etc.) and polyols (*e.g.*, mannitol, sorbitol, etc.). See, also, Remington's Pharmaceutical Sciences (1990) Mack Publishing Co., Easton, Pa., which is hereby incorporated by reference in its entirety.

 In the context of a peptide or polypeptide, the term "fragment" refers to a
15 peptide or polypeptide that comprises less than the full length amino acid sequence. Such a fragment may arise, for example, from a truncation at the amino terminus, a truncation at the carboxy terminus, and/or an internal deletion of a residue(s) from the amino acid sequence. Fragments may, for example, result from alternative RNA
splicing or from *in vivo* protease activity. In certain embodiments, TGF β fragments
20 include polypeptides comprising an amino acid sequence of at least 50, at 100 amino acid residues, at least 125 contiguous amino acid residues, at least 150 contiguous amino acid residues, at least 175 contiguous amino acid residues, at least 200 contiguous amino acid residues, or at least 250 contiguous amino acid residues of the amino acid sequence of a TGF β polypeptide. In a specific embodiment, a fragment of
25 a TGF β polypeptide or an antibody that specifically binds to a TGF β antigen retains at least 1, at least 2, or at least 3 functions of the full-length polypeptide or antibody.

 The terms "fully human antibody" or "human antibody" are used interchangeably herein and refer to an antibody that comprises a human variable region and, most preferably a human constant region. In specific embodiments, the
30 terms refer to an antibody that comprises a variable region and constant region of human origin. "Fully human" anti-TGF β antibodies, in certain embodiments, can also encompass antibodies that bind TGF β polypeptides and are encoded by nucleic acid sequences that are naturally occurring somatic variants of a human germline

immunoglobulin nucleic acid sequence. In a specific embodiment, the anti-TGF β antibodies are fully human antibodies. The term "fully human antibody" includes antibodies having variable and constant regions corresponding to human germline immunoglobulin sequences as described by Kabat et al. (See Kabat *et al.* (1991)

- 5 Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Methods of producing fully human antibodies are known in the art.

The phrase "recombinant human antibody" includes human antibodies that are prepared, expressed, created, or isolated by recombinant means, such as antibodies
10 expressed using a recombinant expression vector transfected into a host cell, antibodies isolated from a recombinant, combinatorial human antibody library, antibodies isolated from an animal (*e.g.*, a mouse or cow) that is transgenic and/or transchromosomal for human immunoglobulin genes (see, *e.g.*, Taylor, L. D. et al. (1992) Nucl. Acids Res. 20:6287-6295) or antibodies prepared, expressed, created, or
15 isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies can have variable and constant regions derived from human germline immunoglobulin sequences (See Kabat, E. A. *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH
20 Publication No. 91-3242). In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may
25 not naturally exist within the human antibody germline repertoire *in vivo*.

The term "heavy chain" when used in reference to an antibody refers to five distinct types, called alpha (α), delta (Δ), epsilon (ϵ), gamma (γ) and mu (μ), based on the amino acid sequence of the heavy chain constant domain. These distinct types of heavy chains are well known in the art and give rise to five classes of antibodies, IgA,
30 IgD, IgE, IgG, and IgM, respectively, including four subclasses of IgG, namely IgG1, IgG1, IgG3, and IgG4. Preferably the heavy chain is a human heavy chain.

An "isolated" or "purified" antibody is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the antibody is

derived, or substantially free of chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of an antibody in which the antibody is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, an antibody that is substantially free of cellular material includes preparations of antibody having less than about 30%, 20%, 10%, or 5% (by dry weight) of heterologous protein (also referred to herein as a "contaminating protein"). When the antibody is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, 10%, or 5% of the volume of the protein preparation. When the antibody is produced by chemical synthesis, it is preferably substantially free of chemical precursors or other chemicals, *i.e.*, it is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. Accordingly, such preparations of the antibody have less than about 30%, 20%, 10%, 5% (by dry weight) of chemical precursors or compounds other than the antibody of interest. In a preferred embodiment, anti-TGF β antibodies are isolated or purified.

The terms "Kabat numbering," and like terms are recognized in the art and refer to a system of numbering amino acid residues that are more variable (*i.e.* hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat *et al.* (1971) Ann. NY Acad. Sci. 190:382-391 and, Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region typically ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region typically ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

The term "light chain" when used in reference to an antibody refers to two distinct types, called kappa (κ) or lambda (λ), based on the amino acid sequence of the constant domains. Light chain amino acid sequences are well known in the art. In preferred embodiments, the light chain is a human light chain.

The terms "manage", "managing", and "management" refer to the beneficial effects that a subject derives from a therapy (*e.g.*, a prophylactic or therapeutic agent), which does not result in a cure of the disease or disorder. In certain embodiments, a subject is administered one or more therapies (*e.g.*, prophylactic or therapeutic agents) to "manage" a TGF β -mediated disease (*e.g.*, OI), or one or more symptoms thereof, so as to prevent the progression or worsening of the disease.

The term "monoclonal antibody" refers to an antibody obtained from a population of homogenous or substantially homogeneous antibodies, and each monoclonal antibody will typically recognize a single epitope on the antigen. In preferred embodiments, a "monoclonal antibody" is an antibody produced by a single hybridoma or other cell. The term "monoclonal" is not limited to any particular method for making the antibody. For example, monoclonal antibodies may be made by the hybridoma method as described in Kohler *et al.*; Nature, 256:495 (1975) or may be isolated from phage libraries. Other methods for the preparation of clonal cell lines and of monoclonal antibodies expressed thereby are well known in the art (see, for example, Chapter 11 in: Short Protocols in Molecular Biology, (2002) 5th Ed.; Ausubel *et al.*, eds., John Wiley and Sons, New York).

The term "pharmaceutically acceptable" means being approved by a regulatory agency of the Federal or a State government or listed in the U.S. Pharmacopeia, European Pharmacopeia, or other generally recognized Pharmacopeia for use in animals, and more particularly in humans.

The term "pharmaceutically acceptable excipient" means any inert substance that is combined with an active molecule, such as a monoclonal antibody, for preparing an agreeable or convenient dosage form. The "pharmaceutically acceptable excipient" is an excipient that is non-toxic to recipients at the dosages and concentrations employed, and is compatible with other ingredients of the formulation comprising the monoclonal antibody.

The terms "prevent", "preventing", and "prevention" refer to the total or partial inhibition of the development, recurrence, onset, or spread of a TGF β -mediated disease and/or symptom related thereto, resulting from the administration of a therapy or combination of therapies provided herein (*e.g.*, a combination of prophylactic or therapeutic agents).

The term "TGF β antigen" refers to that portion of a TGF β polypeptide to which an antibody specifically binds. A TGF β antigen also refers to an analog or derivative of a TGF β polypeptide or fragment thereof to which an antibody specifically binds. In some embodiments, a TGF β antigen is a monomeric TGF β antigen or a dimeric TGF β antigen. A region of a TGF β polypeptide contributing to an epitope may be contiguous amino acids of the polypeptide, or the epitope may come together from two or more non-contiguous regions of the polypeptide. The epitope may or may not be a three-dimensional surface feature of the antigen. A localized region on the surface of a TGF β antigen that is capable of eliciting an immune response is a TGF β epitope. The epitope may or may not be a three-dimensional surface feature of the antigen. As used herein, an "analog" of the TGF β antigen refers to a polypeptide that possesses a similar or identical function as a TGF β polypeptide, a fragment of a TGF β polypeptide, or a TGF β epitope described herein. For example, the analog may comprise a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to the amino acid sequence of a TGF β polypeptide (*e.g.*, SEQ ID NO: 1, 2, or 3), a fragment of a TGF β polypeptide, a TGF β epitope, or an anti-TGF β antibody described herein. Additionally or alternatively, the polypeptide is encoded by a nucleotide sequence that hybridizes under stringent conditions to a nucleotide sequence encoding a TGF β polypeptide, a fragment of a TGF β polypeptide, or a TGF β epitope described herein

The term "human TGF β ," "hTGF β ," or "hTGF β polypeptide" and similar terms refer to the polypeptides ("polypeptides," "peptides," and "proteins" are used interchangeably herein) comprising the amino acid sequence of SEQ ID NO: 1, 2, or 3, and related polypeptides, including SNP variants thereof. Related polypeptides include allelic variants (*e.g.*, SNP variants); splice variants; fragments; derivatives; substitution, deletion, and insertion variants; fusion polypeptides; and interspecies homologs, preferably, which retain TGF β activity and/or are sufficient to generate an anti-TGF β immune response. Also encompassed are soluble forms of TGF β that are sufficient to generate an anti-TGF β immunological response. As those skilled in the art will appreciate, an anti-TGF β antibody can bind to a TGF β polypeptide, polypeptide fragment, antigen, and/or epitope, as an epitope is part of the larger antigen, which is part of the larger polypeptide fragment, which, in turn, is part of the larger polypeptide. hTGF β can exist in a dimeric or monomeric form.

The terms "TGF β -mediated disease" and "TGF β -mediated disorder" are used interchangeably and refer to any disease or disorder that is completely or partially caused by or is the result of TGF β , *e.g.*, hTGF β . In certain embodiments, TGF β is aberrantly expressed. In some embodiments, TGF β may be aberrantly upregulated in a particular cell type. In other embodiments, normal, aberrant, or excessive cell signaling is caused by binding of TGF β to a TGF β receptor. In certain embodiments, the TGF β receptor (*e.g.*, TGF β receptor 1, 2, or 3), is expressed on the surface of a cell, such as an osteoblast, osteoclast, or a bone marrow stromal cell. In certain embodiments, the TGF β -mediated disease is a degenerative bone disease, such as osteogenesis imperfecta.

The terms "specifically binds" or "specifically binding" mean specifically binding to an antigen or a fragment thereof (*e.g.*, TGF β) and not specifically binding to other antigens. An antibody that specifically binds to an antigen may bind to other peptides or polypeptides with lower affinity, as determined by, *e.g.*, radioimmunoassays (RIA), enzyme-linked immunosorbent assays (ELISA), BIACORE, or other assays known in the art. In certain embodiments, an anti-TGF β antibody of the invention may specifically bind to TGF β (*e.g.*, hTGF β) with more than two-fold greater affinity than a different, non-TGF β antigen. Antibodies or variants or fragments thereof that specifically bind to an antigen may be cross-reactive with related antigens. For example, in certain embodiments an anti-TGF β antibody may cross-react with hTGF β and another TGF β antigen (*e.g.*, a rodent or non-human primate TGF β antibody). Preferably, antibodies or variants or fragments thereof that specifically bind to an antigen do not cross-react with other non-TGF β antigens. An antibody or a variant or a fragment thereof that specifically binds to a TGF β antigen can be identified, for example, by immunoassays, BIAcore, or other techniques known to those of skill in the art. Typically a specific or selective reaction will be at least twice background signal or noise, and more typically more than 10 times background. In some embodiments, the binding protein or antibody will bind to its antigen, *e.g.* TGF β , with a dissociation constant of between 1×10^{-6} M and 1×10^{-7} . In other embodiments, the dissociation constant is between 1×10^{-6} M and 1×10^{-8} . See, *e.g.*, Paul, ed., 1989, Fundamental Immunology Second Edition, Raven Press, New York at pages 332-336 for a discussion regarding antibody specificity.

The terms "subject" and "patient" are used interchangeably. As used herein, a subject is preferably a mammal, such as a non-primate (*e.g.*, cows, pigs, horses, cats, dogs, rats, etc.) or a primate (*e.g.*, monkey and human), most preferably a human. In one embodiment, the subject is a mammal, preferably a human, having a TGF β -mediated disease. In another embodiment, the subject is a mammal, preferably a human, at risk of developing a TGF β -mediated disease.

The term "therapeutic agent" refers to any agent that can be used in the treatment, management, or amelioration of a TGF β -mediated disease and/or a symptom related thereto. In certain embodiments, the term "therapeutic agent" refers to a TGF β antibody. In certain other embodiments, the term "therapeutic agent" refers to an agent other than a TGF β antibody. Preferably, a therapeutic agent is an agent that is known to be useful for, or has been, or is currently being used for the treatment, management, or amelioration of a TGF β -mediated disease, or one or more symptoms related thereto.

The term "therapy" refers to any protocol, method, and/or agent that can be used in the prevention, management, treatment, and/or amelioration of a TGF β -mediated disease (*e.g.*, OI). In certain embodiments, the terms "therapies" and "therapy" refer to a biological therapy, supportive therapy, and/or other therapies useful in the prevention, management, treatment, and/or amelioration of a TGF β -mediated disease known to one of skill in the art, such as medical personnel.

The terms "treat", "treatment", and "treating" refer to the reduction or amelioration of the progression, severity, and/or duration of a TGF β -mediated disease (*e.g.*, OI) resulting from the administration of one or more therapies (including, but not limited to, the administration of one or more prophylactic or therapeutic agents). In specific embodiments, such terms refer to the reduction or inhibition of the binding of TGF β to a TGF β receptor, the reduction or inhibition of the production or secretion of TGF β from a cell expressing a TGF β receptor of a subject, the reduction or inhibition of the production or secretion of TGF β from a cell not expressing a TGF β receptor of a subject, and/or the inhibition or reduction of one or more symptoms associated with a TGF β -mediated disease, such as OI.

The terms "variable region" or "variable domain" refer to a portion of the light and heavy chains, typically about the amino-terminal 120 to 130 amino acids in the heavy chain and about 100 to 110 amino acids in the light chain, which differ

extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. The variability in sequence is concentrated in those regions called complementarity determining regions (CDRs), while the more highly conserved regions in the variable domain are called framework regions (FR). The CDRs of the light and heavy chains are primarily responsible for the interaction of the antibody with antigen. Numbering of amino acid positions is according to the EU Index, as in Kabat *et al.* (1991) Sequences of proteins of immunological interest. (U.S. Department of Health and Human Services, Washington D.C.) 5th ed. ("Kabat *et al.*"). In preferred embodiments, the variable region is a human variable region.

B. Osteogenesis Imperfecta (OI)

OI encompasses a group of congenital bone disorders characterized by deficiencies in one or more proteins involved in bone matrix deposition or homeostasis. There are eight types of OI that are defined by their specific gene mutation, and the resulting protein deficiency and phenotype of the affected individual. Though phenotypes vary among OI types, common symptoms include incomplete ossification of bones and teeth, reduced bone mass, brittle bones, and pathologic fractures.

Type-I collagen is one of the most abundant connective tissue proteins in both calcified and non-calcified tissues. Accurate synthesis, post-translational modification, and secretion of type-I collagen are necessary for proper tissue development, maintenance, and repair. Most mutations identified in individuals with OI result in reduced synthesis of type-I collagen, or incorrect synthesis and/or processing of type-I collagen.

In addition to mutations to the type-I collagen gene, other mutations in genes that participate in the intracellular trafficking and processing of collagens have been identified in OI affected individuals. These genes include molecular chaperones, such as FK506 binding protein 10 (FKBP10) and heat shock protein 47 (HSP47) (Alanay *et al.*, 2010; Christiansen *et al.*, 2010; Kelley *et al.*, 2011). Additional mutations have been identified in intermolecular collagen cross-linking genes, such as procollagen-lysine, 2-oxoglutarate 5-dioxygenase 2 (PLOD2), and in members of the collagen prolyl hydroxylase family of genes, including leucine proline-enriched proteoglycan

(leprecan) (LEPRE1), peptidylprolyl isomerase B (cyclophilin B) (CYPB), and cartilage associated protein (CRTAP) (Morello et al., 2006; Cabral et al., 2007; Baldrige et al., 2008; van Dijk et al., 2009; Choi et al., 2009; Barnes et al., 2010; Pyott et al., 2011). Mutations aside, proteins such as bone morphogenetic protein (BMP) and transforming growth factor β (TGF β) and their respective receptors are thought to participate in the various OI phenotypes, though the exact mechanisms of their actions are unknown (Gebken et al., 2000).

In an embodiment, TGF β expression is regulated by molecules that bind type-I and type-II collagen. In certain s embodiment, a small leucine rich proteoglycan (SLRP) regulates TGF β expression. In a specific embodiment, decorin regulates TGF β synthesis. In a certain embodiment, decorin does not bind type-I or type-II collagen in which the 3-hydroxyproline site is absent at position 986 of the type-I and/or type-II collagen molecules.

15 **C. Bone Biology**

The vertebrate skeleton is comprised of bone, which is a living, calcified tissue that provides structure, support, protection, and a source of minerals for regulating ion transport. Bone is a specialized connective tissue that is comprised of both cellular and acellular components. The acellular extracellular matrix (ECM) contains both collagenous and non-collagenous proteins, both of which participate in the calcification process. A correctly secreted and aligned ECM is critical for proper bone formation. Pathology results when any of the ECM proteins are absent, malformed or misaligned, as is evidenced in osteogenesis imperfecta.

The term “cortical bone” or “compact bone” refers to the outer layer of bone, which is dense, rigid, and tough. The term “trabecular bone” or “cancellous bone” is the spongy inner layer of bone, which is lighter and less dense than cortical bone. The term “trabecula” refers to the microscopic structural unit of spongy bone, which is of a rod-like shape and collagenous composition.

Bone is a dynamic tissue that undergoes constant remodeling. The term “osteoblast” refers to a terminally-differentiated bone forming cell that deposits osteoid. The term “osteoid” refers to immature, unmineralized bone that is comprised primarily of type-I collagen. The term “pre-osteoblast” refers to a proliferating immature osteoblast that is not fully differentiated. The term “osteoprogenitor” refers

to a pluripotent cell that gives rise to several stromal cell types, including osteoblasts. Osteoprogenitor cells, which are commonly referred to as “mesenchymal stem cells,” arise in the bone marrow and can be isolated in small numbers from circulating blood. The term “osteoclast” refers to a terminally-differentiated bone resorbing cell that is descended from a bone marrow monocyte. Osteoclasts can be identified by their expression of tartrate resistant acid phosphatase (TRAP).

Under normal homeostatic conditions, osteoblasts and osteoclasts work in unison to maintain bone integrity. Pathology results when bone deposition and bone resorption become uncoupled. For example, osteopetrosis is a bone disease characterized by overly dense, hard bone that is a result of unresorptive osteoclasts, while osteoporosis is a bone disorder characterized by brittle, porous bones which can result from increased osteoclast activity. Evidence suggests that osteoclast activity may be increased in osteogenesis imperfecta, implicating these cell types as a potential target for therapeutic intervention. The present disclosure includes methods of inhibiting osteoclasts with a TGF β antibody.

Several methods can be used to measure and characterize the structure, density, and quality of bone, including histology and histomorphometry, atomic force microscopy, confocal Raman microscopy, nanoindentation, three-point bending test, X-ray imaging, and micro computed tomography (μ -CT). In an exemplified embodiment, bones are measured and characterized by at least one of these methods.

The term “bone volume density” refers to the fraction of a given volume of bone (total volume or TV) that is comprised of calcified matter (bone volume or BV). Therefore, bone volume density is calculated as BV/TV and reported as a percentage. The term “specific bone surface” refers to the total bone surface (BS) per given volume of bone. Therefore, specific bone surface is calculated as BS/TV. Other common bone measurements include: bone area (B.Ar), trabecular number (Tb.N); trabecular spacing (Tb.Sp); N.Oc (osteoclast number); Oc.S (osteoclast surface area); Oc.S/BS; osteoblast number (N.Ob), osteoblast surface area (Ob.S), osteoblast perimeter (Ob.Pm), and derivatives of any of said measurements. A larger Oc.S/BS is an indicator of increased bone resorption by osteoclasts.

D. Transforming growth factor beta (TGFβ)

TGFβs are multifunctional cytokines that are involved in cell proliferation and differentiation, embryonic development, extracellular matrix formation, bone development, wound healing, hematopoiesis, and immune and inflammatory responses (Roberts et al., 1981; Border et al., 1995a). Secreted TGFβ protein is cleaved into a latency-associated peptide (LAP) and a mature TGFβ peptide, and is found in latent and active forms. The mature TGFβ peptide forms both homodimers and heterodimers with other TGFβ family members. TGFβ may be purified from any natural source, or may be produced synthetically (*e.g.*, by use of recombinant DNA technology). Preferably, the TGFβ molecule is from a human, known herein as "hTGFβ".

There are three human TGFβ isoforms: TGFβ1, TGFβ2, and TGFβ3 (Swiss Prot accession numbers P01137, P08112, and P10600, respectively) which, in their biologically active state, are 25 kDa homodimers comprising two 112 amino acid monomers joined by an inter-chain disulfide bridge. TGFβ1 differs from TGFβ2 by 27, and from TGFβ3 by 22, mainly conservative amino acid changes. These differences have been mapped on the 3D structure of TGFβ determined by X-ray crystallography (Schlunegger et al., 1992; Peer et al., 1996) and the receptor binding regions have been defined (Griffith et al., 1996; Qian et al., 1996).

20 hTGFβ1 (SEQ ID NO: 1)

MPPSGLRLLL LLLPLLWLLV LTPGRPAAGL STCKTIDMEL VKRKRIEAIR
GQILSKLRLA 60
SPPSQGEVPP GPLPEAVLAL YNSTRDRVAG ESAEPEPEPE ADYYAKEVTR
VLMVETHNEI 120
25 YDKFKQSTHS IYMFNTSEL REAVPEPVLL SRAELRLLRL KKKVEQHVEL
YQKYSNNSWR 180
YLSNRLLAPS DSPEWLSFDV TGVVRQWLSR GGEIEGFRLS AHCSCDSRDN
TLQVDINGFT 240
TGRRGDLATI HGMNRPFLLL MATPLERAQH LQSSRHRRAL DTNYCFSSTE
30 KNCCVRQLYI 300
DFRKDLGWKW IHEPKGYHAN FCLGPCPYIW SLDTQYSKVL ALYNQHNPQA
SAAPCCVPQA 360
LEPLPIVYYV GRKPKVEQLS NMIVRSCKCS 390 (SEQ ID NO: 1)

hTGFβ2 (SEQ ID NO: 2)

MHYCVLSAFL ILHLVTVALS LSTCSTLDMD QFMRKRIEAI RGQILSKLKL
TSPPEDYPEP 60
EEVPPEVISI YNSTRDLLQE KASRRAAACE RERSDEEYYA KEVYKIDMPP
5 FFPSENAIPP 120
TFYRPFYFRIV RFDVSAMEKN ASNLVKAEFR VFRLQNPKEAR VPEQRIELYQ
ILKSKDLTSP 180
TQRYIDSKVV KTRAEGEWLS FDVTDAVHEW LHHKDRNLGF KISLHPCCT
FVPSNNYIIP 240
10 NKSEELERF AGIDGTSTYT SGDQKTIKST RKKNSGKTPH LLLMLLPSYR
LESQQTNRK 300
KRALDAAYCF RNVQDNCCLR PLYIDFKRDL GWKWIHEPKG YNANFCAGAC
PYLWSSDTQH 360
SRVLSLYNTI NPEASASPCC VSQDLEPLTI LYYIGKTPKI EQLSNMIVKS CKCS
15 414 (**SEQ ID NO: 2**)

hTGFβ3 (SEQ ID NO: 3)

MKMHLQRALV VLALLNFATV SLSLSTCTTL DFGHIKKKRV EAIRGQILSK
LRLTSPPEPT 60
20 VMTHVPYQVL ALYNSTRELL EEMHGEREEG CTQENTESEY YAKEIHKFDM
IQGLAEHNEL 120
AVCPKGITSK VFRFNVSSVE KNRTNLFRAE FRVLRVPNPS SKRNEQRIEL
FQILRPDEHI 180
AKQRYIGGKN LPTRGTAEWL SFDVTDTVRE WLLRRESNLG LEISIHCPCH
25 TFQPNGDILE 240
NIHEVMEIKF KGVDNEDDHG RGDGLRLKKQ KDHHNPHLIL MMIPPHRLDN
PGQGGQRKKR 300
ALDTNYCFRN LEENCCVRPL YIDFRQDLGW KVVHEPKGYY ANFCSGPCPY
LRSADTTHST 360
30 VLGLYNTLNP EASASPCCVP QDLEPLTILY YVGRTPKVEQ LSNMVKSCS CS 412
(**SEQ ID NO: 3**)

There are three TGFβ receptors in humans, TGFβ receptor 1, 2, and 3, which
can be distinguished by their structural and functional properties, including affinity
for TGFβ protein family members. Binding of a TGFβ protein to a homodimeric or
35 heterodimeric TGFβ transmembrane receptor complex activates the canonical TGFβ
signaling pathway mediated by intracellular SMAD proteins.

The deregulation of TGFβs leads to pathological processes that, in humans, have been implicated in numerous conditions, such as, birth defects, cancer, chronic inflammatory, autoimmune diseases, and fibrotic diseases (Border et al., 1994; Border et al., 1995b).

5 Human TGFβs are very similar to mouse TGFβs: human TGFβ1 has only one amino acid difference from mouse TGFβ1; human TGFβ2 has only three amino acid differences from mouse TGFβ2; and human TGFβ3 is identical to mouse TGFβ3.

E. Molecules that bind to transforming growth factor beta (TGFβ)

10 The present invention includes methods that comprise administering to a subject a molecule that binds to TGFβ. The TGFβ binder may be any binding molecule, such as an antibody, a fusion protein (e.g., an immunoadhesin), an siRNA, a nucleic acid, an aptamer, a protein, or a small molecule organic compound.

In certain embodiments, the invention includes an antibody that binds to TGFβ
15 (an anti-TGFβ antibody), or a variant thereof, or an antigen binding fragment thereof. Anti-TGFβ antibodies specifically bind to a TGFβ protein, polypeptide fragment, or epitope. The molecule that binds to TGFβ may be from any species.

In certain exemplary embodiments, the antibody that binds to TGFβ is a humanized antibody, a fully human antibody, or a variant thereof, or an antigen-
20 binding fragment thereof. Preferred anti-TGFβ antibodies prevent binding of TGFβ with its receptors and inhibit TGFβ biological activity (e.g., the TGFβ receptor-mediated intracellular SMAD signaling and resulting cellular activity).

In certain embodiments, the antibody, or antigen-binding fragment thereof, is Lerdelimumab (CAT-152), Metelimumab (CAT-192), Fresolimumab (GC-1008),
25 LY2382770, STX-100, or IMC-TR1.

In certain specific embodiments, the antibody that binds to TGFβ comprises a heavy chain variable region (VH) comprising the amino acid sequence of any one or more of the following complementarity determining regions (CDRs):

HCDR1 – SNVIS (**SEQ ID NO: 4**);
30 HCDR2 – GVIPIVDIANYAQRFKG (**SEQ ID NO: 5**); or
HCDR3 – TLGLVLDAMDY (**SEQ ID NO: 6**).

In other specific embodiments, the antibody that binds to TGFβ comprises a light chain variable region (VL) comprising the amino acid sequence of any one or more of the following complementarity determining regions (CDRs):

LCDR1 – RASQSLGSSYLA (SEQ ID NO: 7);

5 LCDR2 – GASSRAP (SEQ ID NO: 8); or

LCDR3 – QQYADSPIT (SEQ ID NO: 9).

In a specific embodiment, the antibody that binds to TGFβ comprises a heavy chain variable region (VH) comprising the amino acid sequences of SEQ ID NOs: 4, 5, and 6.

10 In another specific embodiment, the antibody that binds to TGFβ comprises a light chain variable region (VL) comprising the amino acid sequences of SEQ ID NOs: 7, 8, and 9.

In more specific embodiments, the antibody that binds to TGFβ comprises a heavy chain variable region comprising the amino acid sequences of SEQ ID NOs: 4, 5, and 6; and a light chain variable region comprising the amino acid sequences of SEQ ID NOs: 7, 8, and 9.

In a specific embodiment, the antibody that binds to TGFβ comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 10:

20 QVQLVQSGAE VKKPGSSVKV SCKASGYTFS SNVISWVRQA PGQGLEWMGG
VIPIVDIANY AQRFKGRVTI TADESTSTTY MELSSLRSED TAVYYCASTL
GLVLDAMDYW GQGTTLVTVSS (SEQ ID NO: 10) .

In another specific embodiment, the antibody that binds to TGFβ comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 11:

25 ETVLTQSPGT LSLSPGERAT LSCRASQSLG SSYLAWYQQK PGQAPRLLIY
GASSRAPGIP DRFSGSGSGT DFTLTISRLE PEDFAVYYCQ QYADSPITFG
QGTRLEIK (SEQ ID NO: 11) .

In more specific embodiments, the antibody that binds to TGFβ comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 10; and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 11.

In some embodiments, the antibody that binds to TGFβ further comprises a constant region, *e.g.*, a human IgG constant region. In some embodiments, the constant region is a human IgG4 constant region. In additional embodiments, the

constant region is a modified human IgG4 constant region. Preferably, the IgG4 constant region comprises the amino acid sequence of SEQ ID NO: 12:

ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV
HTFPAVLQSS GLYSLSSVVT VPSSSLGTKT YTCNVDHKPS NTKVDKRVES
5 KYGPPCPSCP APEFLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSQED
PEVQFNWYVD GVEVHNAKTK PREEQFNSTY RVVSVLTVLH QDWLNGKEYK
CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK NQVSLTCLVK
GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG
NVFSCSV MHE ALHNHYTQKS LSLSLGK (**SEQ ID NO: 12**) .

10 In other embodiments, the constant region is a human Ck constant region. Preferably, the Ck constant region comprises the amino acid sequence of SEQ ID NO: 13:

RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNALQSG
NSQESVTEQD SKDSTYSLSS TLTLKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC
15 (**SEQ ID NO: 13**) .

In specific embodiments, the antibody that binds to TGF β comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14:

QVQLVQSGAE VKKPGSSVKV SCKASGYTFS SNVISWVRQA PGQGLEWMGG
VIPIVDIANY AQRFKGRVTI TADESTSTTY MELSSLRSED TAVYYCASTL
20 GLVLDAMDYW GQGTLVTVSS ASTKGPSVFP LAPCSRSTSE STAALGCLVK
DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTKT
YTCNVDHKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV FLFPPKPKDT
LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY
RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT
25 LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS
DGSFFLYSRL TVDKSRWQEG NVFSCSV MHE ALHNHYTQKS LSLSLGK (**SEQ ID NO: 14**) .

Positions 1-120: variable region of the heavy chain (VH). The CDRs (complementarity determining regions, according to Kabat
30 definition) are underlined.

Positions 121-447: constant region of human IgG4 (SwissProt IGHG4_HUMAN).

In other specific embodiments, the antibody that binds to TGF β comprises a light chain comprising the amino acid sequence of SEQ ID NO: 15:

ETVLTQSPGT LSLSPGERAT LSCRASQSLG SSYLAWYQQK PGQAPRLLIY
35 GASSRAPGIP DRFSGSGSGT DFTLTISRLE PEDFAVYYCQ QYADSPITFG

QGTRLEI**K**RRT VAAPSVFIFP PSDEQLKSGT ASVVCLLNNF YPBREAKVQWK
VDNALQSGNS QESVTEQDSK DSTYLSSTL TLSKADYEKH KVVACEVTHQ
GLSSPVTKSF NRGEC (**SEQ ID NO: 15**) .

5 Positions 1-108: variable region of the light chain (VL). The CDRs
(complementarity determining regions, according to Kabat
definition) are underlined.

Positions 109-215: constant region of human Cκ.

In further embodiments, the antibody that binds to TGFβ comprises a heavy
chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain
10 comprising the amino acid sequence of SEQ ID NO: 15.

In some embodiments, the antibody that binds to TGFβ is expressed by a host
cells as comprising leader sequences. The leader sequence preferably comprises an
amino acid sequence from 1-30 amino acids in length, more preferably 25-25 amino
acids, and most preferably 19 amino acids. The heavy chain, light chain, or both the
15 heavy and light chain may comprise a leader sequence.

For example, the light or heavy chain leader sequence may comprise the
amino acid sequence of SEQ ID NO: 16: MGWSCIILFL VATATGVHS (**SEQ ID NO:**
16) . Accordingly, a host cell may expressing a unprocessed heavy chain may
comprise the amino acid sequence of SEQ ID NO: 17:

20 MGWSCIILFL VATATGVHS**Q** VQLVQSGAEV KKP GSSVKVS CKASGYTFSS 50
NVISWVRQAP GQGLEWMGGV IPIVDIANYA QRFKGRVTIT ADESTSTTYM 100
ELSSLRSEDV AVYYCASTLG LVLDAMDYWG QGTLVTVSS**A** STKGPSVFPL 150
APCSRSTSES TAALGCLVKD YFPEPVTVSW NSGALTSGVH TFP AVLQSSG 200
LYSLSSVTV PSSSLGTKTY TCNV D HKPSN TKVDKR VESK YGPPCPSCPA 250
25 PEFLGGPSVF LFPPKPKDTL MISRTPEVTC VVVDVSQEDP EVQFNWYVDG 300
VEVHNAKTKP REEQFNSTYR VVSVLTVLHQ DWLNGKEYKC KVS NKGLPSS 350
IEKTISKAKG QPREPQVYTL PPSQEEMTKN QVSLTCLVKG FYPSDIAVEW 400
ESNGQPENNY KTTTPVLDSD GSFFLYSRLT VDKSRWQEGN VFSCSV MHEA 450
LHNHYTQKSL SLSLGK 466 (**SEQ ID NO: 17**) .

30 wherein

positions 1-19: leader sequence

Positions 20-139: variable region of the heavy chain (VH). The CDRs
(complementarity determining regions, according to Kabat
definition) are underlined.

Positions 140-466: constant region of human IgG4 (SwissProt IGHG4_HUMAN).

In other exemplary embodiments, a host cell expressing a unprocessed light chain may comprise the amino acid of SEQ ID NO: 18:

5 MGWSCIILFL VATATGVHSE TVLTQSPGTL SLSPGERATL SCRASQSLGS 50
SYLAWYQQKP GQAPRLLIYG ASSRAPGIPD RFSGSGSGTD FTLTISRLEP 100
EDFAVYYCQQ YADSPITFGQ GTRLEIKRTV AAPSVFIFPP SDEQLKSGTA 150
SVVCLLNNFY PREAKVQWKV DNALQSGNSQ ESVTEQDSKD STYSLSTLT 200
LSKADYEKHK VYACEVTHQG LSSPVTKSFN RGEK 234 (SEQ ID NO: 18) .

10 wherein

Positions 1-19: leader sequence

Positions 20-127: variable region of the light chain (VL). The CDRs (complementarity determining regions, according to Kabat definition) are underlined.

15 Positions 128-234: constant region of human Cκ.

In an exemplary embodiment of the invention, the antibody that binds to TGFβ is a humanized or fully human antibody. Examples of humanized and fully human antibody isotypes include IgA, IgD, IgE, IgG, and IgM. Preferably, the anti-TGFβ antibody is an IgG antibody. There are four forms of IgG. Preferably, the anti-TGFβ antibody is an IgG4 antibody. In one embodiment of the invention, the anti-TGFβ antibody is a humanized IgG4 antibody. In another embodiment of the invention, the anti-TGFβ antibody is a fully human IgG4 antibody.

In a most preferred embodiment of the invention, the anti-TGFβ antibody is an IgG4 anti-TGFβ antibody comprising a heavy chain comprising the amino acid sequence of SEQ ID NO: 14 and a light chain comprising the amino acid sequence of SEQ ID NO: 15. In an alternative most preferred embodiment of the invention, the anti-TGFβ antibody is an IgG4 anti-TGFβ antibody comprising a heavy chain variable region and a light chain variable region, the heavy chain variable region comprising 3 complementarity determining regions (CDRs) comprising the amino acid sequences of SEQ ID NOs: 4, 5, and 6, and the light chain variable region comprising 3 CDRs comprising the amino acid sequences of SEQ ID NOs: 7, 8, and 9. Identification, isolation, preparation, and characterization of anti-TGFβ antibodies, including the anti-TGFβ antibody comprising a heavy chain amino acid sequence comprising SEQ

ID NO: 14 and a light chain amino acid sequence comprising SEQ ID NO: 15, and the CDR sequences corresponding with SEQ ID NOs: 4-9 have been described in detail in U.S. Patent No. 7,723,486, and U.S. Patent No. 8,383,780, each of which is incorporated herein by reference in its entirety.

5 Preferably, the antibody or antigen binding fragment thereof is “pan-specific” and binds to human TGFβ1, TGFβ2, and TGFβ3. More preferably, the antibody or antigen binding fragment thereof binds to human TGFβ1, TGFβ2, and TGFβ3, and acts as an antagonist. Most preferably, the antibody or antigen binding fragment thereof binds to human TGFβ1, TGFβ2, and TGFβ3, and neutralizes human TGFβ1,
10 TGFβ2, and TGFβ3. Exemplary pan-specific anti-TGFβ monoclonal antibodies (mAbs) suitable for use in the methods of the invention are described in US Patent Nos. 7,723,476 and 8,383,780, each of which is incorporated by reference herein in its entirety.

 1D11.16 is an exemplary murine pan-specific anti-TGFβ antibody that
15 neutralizes human and mouse TGFβ1, TGFβ2, and TGFβ3 in a wide range of in vitro assays (Dasch et al., 1989; Dasch et al., 1996; R&D System product sheet for MAB1835, each of which is incorporated herein by reference in their entirety) and is efficacious in proof-of principle studies in animal models of fibrosis (Ling et al., 2003; Miyajima et al., 2000; Schneider et al., 1999; Khanna et al., 1999; Shenkar et
20 al., 1994). However, since 1D11.16 is a murine monoclonal antibody (Dasch et al., 1989; Dasch et al., 1996), it is not a preferred for therapeutic use in humans. Accordingly, in certain embodiments, variants or derivatives of the 1D11.16 antibody are employed in the methods of the invention.

 As indicated above, certain embodiments of the invention also include variants
25 or derivatives of anti-TGFβ antibodies. Specifically, the invention may include variants of the anti-TGFβ antibody that is an IgG4 anti-TGFβ antibody comprising a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15. In other embodiment, the invention includes variants or derivatives of the 1D11.16 antibody. Variants of anti-
30 TGFβ antibodies may have similar physicochemical properties based on their high similarity, and therefore are also included within the scope of the invention. Variants are defined as antibodies with an amino acid sequence that is at least 80%, at least 90%, at least 95%, or at least 97%, e.g., least 98% or 99% homologous to an anti-

TGF β antibody described herein, and capable of competing for binding to a TGF β polypeptide, a TGF β polypeptide fragment, or a TGF β epitope. Preferably, the variants will ameliorate, neutralize, or otherwise inhibit binding of TGF β with its receptors and TGF β biological activity (*e.g.*, TGF β receptor-mediated intracellular SMAD signaling and resulting cellular activity). Determining competition for binding to the target can be done by routine methods known to the skilled person in the art. Preferably the variants are human antibodies, and preferably are IgG4 molecules. In preferred embodiments, a variant is at least 90%, 95%, 96%, 97%, 98%, or 99% identical in amino acid sequence with the IgG4 anti-TGF β antibody comprising a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15. The term "variant" refers to an antibody that comprises an amino acid sequence that is altered by one or more amino acids compared to the amino acid sequences of the anti-TGF β antibody. The variant may have conservative sequence modifications, including amino acid substitutions, modifications, additions, and deletions.

Examples of modifications include, but are not limited to, glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, and linkage to a cellular ligand or other protein. Amino acid modifications can be introduced by standard techniques known in the art, such as site-directed mutagenesis, molecular cloning, oligonucleotide-directed mutagenesis, and random PCR-mediated mutagenesis in the nucleic acid encoding the antibodies. Conservative amino acid substitutions include the ones in which the amino acid residue is replaced with an amino acid residue having similar structural or chemical properties. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (*e.g.*, glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (*e.g.*, threonine, valine, isoleucine), and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan). It will be clear to the skilled artisan that classifications of amino acid residue families other than the one used above can also be employed. Furthermore, a variant may have non-conservative amino acid

substitutions, *e.g.*, replacement of an amino acid with an amino acid residue having different structural or chemical properties. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, modified, inserted, or deleted without abolishing immunological activity may be found using computer programs well known in the art. Computer algorithms, such as, *inter alia*, Gap or Bestfit, which are known to a person skilled in the art, can be used to optimally align amino acid sequences to be compared and to define similar or identical amino acid residues. Variants may have the same or different, either higher or lower, binding affinities compared to an anti-TGF β antibody, but are still capable of specifically binding to TGF β , and may have the same, higher or lower, biological activity as the anti-TGF β antibody.

Embodiments of the invention also include antigen binding fragments of the anti-TGF β antibodies. The term "antigen binding domain," "antigen binding region," "antigen binding fragment," and similar terms refer to that portion of an antibody that comprises the amino acid residues that interact with an antigen and confer on the binding agent its specificity and affinity for the antigen (*e.g.*, the complementarity determining regions (CDR)). The antigen binding region can be derived from any animal species, such as rodents (*e.g.*, rabbit, rat or hamster) and humans. Preferably, the antigen binding region will be of human origin. Non-limiting examples of antigen binding fragments include: Fab fragments, F(ab')₂ fragments, Fd fragments, Fv fragments, single chain Fv (scFv) molecules, dAb fragments, and minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of the antibody.

F. Therapeutic Administration

The methods described herein comprise administering a therapeutically effective amount of an antibody that binds to TGF β to a subject. As used herein, the phrase "therapeutically effective amount" means a dose of antibody that binds to TGF β that results in a detectable improvement in one or more symptoms associated with OI or which causes a biological effect (*e.g.*, a decrease in the level of a particular biomarker) that is correlated with the underlying pathologic mechanism(s) giving rise to the condition or symptom(s) of osteogenesis imperfecta. For example, a dose of antibody that binds to TGF β that increases bone mineral density, increases bone mass

and/or bone strength, reduces bone and/or tooth fractures, and/or improves any diagnostic measurement of OI is deemed a therapeutically effective amount.

In an embodiment, bone mineral density, bone mass, and/or bone strength are increased by about 5% to about 200% following treatment with an antibody that binds to TGF β . In certain embodiments, bone mineral density, bone mass, and/or bone strength are increased by about 5% to about 10%, 10% to about 15%, 15% to about 20%, 20% to about 25%, 25% to about 30%, 30% to about 35%, 35% to about 40%, 40% to about 45%, 45% to about 50%, 50% to about 55%, 55% to about 60%, 60% to about 65%, 65% to about 70%, 70% to about 75%, 75% to about 80%, 80% to about 85%, 85% to about 90%, 90% to about 95%, 95% to about 100%, 100% to about 105%, 105% to about 110%, 110% to about 115%, 115% to about 120%, 120% to about 125%, 125% to about 130%, 130% to about 135%, 135% to about 140%, 140% to about 145%, 145% to about 150%, 150% to about 155%, 155% to about 160%, 160% to about 165%, 165% to about 170%, 170% to about 175%, 175% to about 180%, 180% to about 185%, 185% to about 190%, 190% to about 195%, or 195% to about 200% , following treatment with an antibody that binds to TGF β .

In certain embodiments, a dose of an antibody which reduces serum biomarkers of bone resorption, such as urinary hydroxyproline, urinary total pyridinoline (PYD), urinary free deoxypyridinoline (DPD), urinary collagen type-I cross-linked N-telopeptide (NTX), urinary or serum collagen type-I cross-linked C-telopeptide (CTX), bone sialoprotein (BSP), osteopontin (OPN), and tartrate-resistant acid phosphatase 5b (TRAP), is deemed a therapeutically effective amount. In an embodiment, serum biomarkers of bone resorption are reduced by about 5% to about 200% following treatment with an antibody that binds to TGF β

In an embodiment, serum biomarkers of bone resorption, such as urinary hydroxyproline, urinary total pyridinoline (PYD), urinary free deoxypyridinoline (DPD), urinary collagen type-I cross-linked N-telopeptide (NTX), urinary or serum collagen type-I cross-linked C-telopeptide (CTX), bone sialoprotein (BSP), osteopontin (OPN), and tartrate-resistant acid phosphatase 5b (TRAP), are decreased by about 5% to about 10%, 10% to about 15%, 15% to about 20%, 20% to about 25%, 25% to about 30%, 30% to about 35%, 35% to about 40%, 40% to about 45%, 45% to about 50%, 50% to about 55%, 55% to about 60%, 60% to about 65%, 65% to about 70%, 70% to about 75%, 75% to about 80%, 80% to about 85%, 85% to about

90%, 90% to about 95%, 95% to about 100%, 100% to about 105%, 105% to about 110%, 110% to about 115%, 115% to about 120%, 120% to about 125%, 125% to about 130%, 130% to about 135%, 135% to about 140%, 140% to about 145%, 145% to about 150%, 150% to about 155%, 155% to about 160%, 160% to about 165%,
5 165% to about 170%, 170% to about 175%, 175% to about 180%, 180% to about 185%, 185% to about 190%, 190% to about 195%, or 195% to about 200% , following treatment with an antibody that binds to TGF β .

In certain embodiments, a dose of an antibody which increase serum biomarkers of bone deposition, such as total alkaline phosphatase, bone-specific
10 alkaline phosphatase, osteocalcin, and type-I procollagen (C-terminal/N-terminal), is deemed a therapeutically effective amount. In an embodiment, serum biomarkers of bone deposition are increased by about 5% to about 200% following treatment with an antibody that binds to TGF β

In an embodiment, serum biomarkers of bone deposition, such as total alkaline
15 phosphatase, bone-specific alkaline phosphatase, osteocalcin, and type-I procollagen (C-terminal/N-terminal), are increased by about 5% to about 10%, 10% to about 15%, 15% to about 20%, 20% to about 25%, 25% to about 30%, 30% to about 35%, 35% to about 40%, 40% to about 45%, 45% to about 50%, 50% to about 55%, 55% to about 60%, 60% to about 65%, 65% to about 70%, 70% to about 75%, 75% to about 80%,
20 80% to about 85%, 85% to about 90%, 90% to about 95%, 95% to about 100%, 100% to about 105%, 105% to about 110%, 110% to about 115%, 115% to about 120%, 120% to about 125%, 125% to about 130%, 130% to about 135%, 135% to about 140%, 140% to about 145%, 145% to about 150%, 150% to about 155%, 155% to about 160%, 160% to about 165%, 165% to about 170%, 170% to about 175%, 175%
25 to about 180%, 180% to about 185%, 185% to about 190%, 190% to about 195%, or 195% to about 200% , following treatment with an antibody that binds to TGF β .

Other embodiments include administering a therapeutically effective dose of an antibody which improves the function of non-skeletal organs affected by OI. For example, a dose of antibody that binds to TGF β that improves hearing, lung, and/or
30 kidney function is deemed a therapeutically effective amount.

In accordance with the methods of the present invention, a therapeutically effective amount of an antibody that binds to TGF β that is administered to a subject will vary depending upon the age and the size (*e.g.*, body weight or body surface area)

of the subject, as well as the route of administration, and other factors well known to those of ordinary skill in the art.

In certain exemplary embodiments, the anti-TGF β antibody is administered to the subject as a subcutaneous dose. Other exemplary modes of administration
5 include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, intranasal, epidural, and oral routes. The composition may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, *etc.*) and may be administered together with other biologically
10 active agents. Administration can be systemic or local. The TGF β antibody can be administered parenterally or subcutaneously.

Various delivery systems are known and can be used to administer the pharmaceutical composition, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, receptor mediated endocytosis (see, *e.g.*, Wu *et al.* (1987) J. Biol.
15 Chem. 262:4429-4432). The therapeutic compositions will be administered with suitable carriers, excipients, and other agents that are incorporated into formulations to provide improved transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton,
20 PA. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax.
25 See also Powell *et al.* "Compendium of excipients for parenteral formulations" PDA (1998) J Pharm Sci Technol 52:238-311.

Pharmaceutical compositions may be prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, *etc.*

30 Pharmaceutical compositions can also be administered to the subject using any acceptable device or mechanism. For example, the administration can be accomplished using a syringe and needle or with a reusable pen and/or autoinjector delivery device. The methods of the present invention include the use of numerous

reusable pen and/or autoinjector delivery devices to administer a TGF β binder (or pharmaceutical formulation comprising the binder). Examples of such devices include, but are not limited to AUTOPENTM (Owen Mumford, Inc., Woodstock, UK), DISETRONICTM pen (Disetronic Medical Systems, Bergdorf, Switzerland),
5 HUMALOG MIX 75/25TM pen, HUMALOGTM pen, HUMALIN 70/30TM pen (Eli Lilly and Co., Indianapolis, IN), NOVOPENTM I, II and III (Novo Nordisk, Copenhagen, Denmark), NOVOPEN JUNIORTM (Novo Nordisk, Copenhagen, Denmark), BDTM pen (Becton Dickinson, Franklin Lakes, NJ), OPTIPENTM, OPTIPEN PROTM, OPTIPEN STARLETTM, and OPTICLIKTM (sanofi-aventis,
10 Frankfurt, Germany), to name only a few. Examples of disposable pen and/or autoinjector delivery devices having applications in subcutaneous delivery of a pharmaceutical composition include, but are not limited to the SOLOSTARTM pen (sanofi-aventis), the FLEXPENTM (Novo Nordisk), and the KWIKPENTM (Eli Lilly), the SURECLICKTM Autoinjector (Amgen, Thousand Oaks, CA), the PENLETTM (Haselmeier, Stuttgart, Germany), the EPIPEN (Dey, L.P.), and the HUMIRATM Pen
15 (Abbott Labs, Abbott Park, IL), to name only a few.

The use of a microinfusor to deliver a TGF β binder (or pharmaceutical formulation comprising the binder) to a subject is also contemplated herein. As used herein, the term "microinfusor" means a subcutaneous delivery device designed to
20 slowly administer large volumes (*e.g.*, up to about 2.5 mL or more) of a therapeutic formulation over a prolonged period of time (*e.g.*, about 10, 15, 20, 25, 30 or more minutes). See, *e.g.*, U.S. 6,629,949; US 6,659,982; and Meehan *et al.*, J. Controlled Release 46:107-116 (1996). Microinfusors are particularly useful for the delivery of large doses of therapeutic proteins contained within high concentration (*e.g.*, about
25 100, 125, 150, 175, 200 or more mg/mL) and/or viscous solutions.

G. Combination Therapies

In certain aspects, the invention includes methods for treating OI that comprise administering to a subject in need of such treatment an antibody that binds to TGF β in
30 combination with at least one additional therapeutic agent. Examples of additional therapeutic agents that can be administered in combination with an anti-TGF β antibody in the practice of the methods of the present invention include, but are not limited to, bisphosphonates, calcitonin, teriparatide, and any other compound known

to treat, prevent, or ameliorate osteogenesis imperfecta in a subject. In the present methods, the additional therapeutic agent(s) can be administered concurrently or sequentially with the antibody that binds to TGF β . For example, for concurrent administration, a pharmaceutical formulation can be made that contains both an antibody that binds to TGF β and at least one additional therapeutic agent. In an embodiment, the antibody that binds to TGF β is administered in combination with pharmaceutical bisphosphonates (e.g., Etidronate, Clodronate, Tiludronate, Pamidronate, Neridronate, Olpadronate, Alendronate, Ibandronate, Zoledronate, and Risedronate,). In another embodiment, the antibody that binds to TGF β is administered in combination with a drug that stimulates bone formation, such as parathyroid hormone analogs and calcitonin. In yet another embodiment, the antibody that binds to TGF β is administered in combination with a selective estrogen receptor modulator (SERM). The amount of the additional therapeutic agent that is administered in combination with the antibody that binds to TGF β in the practice of the methods of the present invention can be easily determined using routine methods known and readily available in the art.

EXAMPLES

OI is a generalized connective tissue disease in which affected individuals display an abnormality in forming type-I collagen fibrils due to mutations in the primary sequence of the alpha 1 or alpha 2 chain of type-I collagen, as well abnormalities in the post-translational modification of type-I collagen and proteins that bind type-I collagen fibrils. CRTAP encodes a protein called the cartilage associated protein, a member of the prolyl-3-hydroxylation complex whose function is to assist with the proper folding, post-translational modification and secretion of type-I collagen. Mutations to CRTAP are responsible for type VII osteogenesis imperfecta. Mice lacking the *Crtap* gene (*Crtap*^{-/-}) display a phenotype that mimics osteogenesis imperfecta, and are used as a model of this disease (Morello et al., 2006). *Crtap*^{-/-} mice and age-matched wild type (WT) littermate controls were used in the following examples.

Material and methods

Animals, anti-TGF β treatment and tissue collection

Crtap^{-/-} mice were generated and maintained on a mixed C57Black/6J /129Sv genetic background. Mice harboring a *G610C* mutation in the *Col1a2* gene (*Col1a2*^{tm1.1Mcb}) were obtained and bred to wildtype C57Bl/6J mice. Mice that were heterozygous for the *Col1a2*^{tm1.1Mcb} allele were used for experiments. TGF β -reporter mice that express luciferase in response to the Smad2/3 dependent TGF β signaling pathway (SBE-Luc mice) were obtained and bred to *Crtap*^{+/-} mice for 2 generations to generate *Crtap*^{-/-} mice and wildtype littermates expressing the reporter transgene. All mice were housed in a vivarium and animal experiments were performed following the approved protocol of the Animal Care and Use Committee (IACUC).

For protein and RNA analyses, calvaria of P3 mice were isolated, cleaned of extraskeletal tissue, and snap frozen in liquid nitrogen. For immunostaining of *Crtap*^{-/-} P10 mice lungs, the lungs of each mouse were equally inflated immediately after sacrifice by gravity with 4% paraformaldehyde at a constant pressure of 25cm H₂O and then suture closed at the trachea. Lungs were then gently dissected from the thorax and fixed in 4% paraformaldehyde overnight.

Eight week old female *Crtap*^{-/-} and *Col1a2*^{tm1.1Mcb} mice were treated with the pan-TGF β neutralizing antibody 1D11 for 8 weeks (10 mg/kg body weight, I.P. injections 3 times each week). Control *Crtap*^{-/-}, *Col1a2*^{tm1.1Mcb}, and WT mice received a control antibody (13C4) of the same IgG1 isotype. After treatment, mice were sacrificed, and lumbar spines and femurs were collected and fixed in 10% formalin for microCT and bone histomorphometry. Contralateral femurs of *Crtap*^{-/-} mice were stored in -20°C wrapped in saline soaked gauze until biomechanical testing was performed. Lungs of these *Crtap*^{-/-} mice were equally inflated, collected and fixed as described for P10 mice. No blinding was possible during treatment, because 1D11 or control antibody were injected according to group allocation. In all subsequent analyses the investigators were blinded to genotype and treatment group.

Immunoblotting

Protein was extracted from snap frozen P3 calvaria samples, transferred to 300 μ l lysis buffer (0.0625 M Tris-HCl pH 7.5, 2% SDS, 5 mM NaF, 2 mM Na_3VO_4 and RocheComplete proteinase inhibitor) and homogenized for 1 minute, followed by incubation at 95°C for 40 minutes. The supernatant was transferred to Centrifugal Filter Units/Amicon Ultra 3K (Millipore) and centrifuged to concentrate the protein. The total protein concentration of the lysate was measured using the Micro BCA reagent (Pierce) following the manufacturer's directions. 40 μ g of calvaria protein extracts were suspended in laemmeli buffer containing 5% β -mercaptoethanol and separated on Mini Protean TGX SDS-PAGE gels (gradient 4-20%; Bio-Rad) and transferred onto PVDF membranes for western blot analyses. PVDF membranes were incubated with pSmad2 monoclonal antibody (Cell Signaling #3108, 1:750 in TBST containing 5% BSA overnight), followed by secondary HRP-linked anti-rabbit antibody (GE, 1:5000 in TBST containing 5% BSA for 2 hours), treated with ECL Plus Western Blotting Detection System (GE) and exposed to X-ray film. Subsequently, antibodies were stripped from membranes using ReBlot Plus reagent (Millipore), and incubated with Smad2 monoclonal antibody (Cell Signaling #5339, 1:2000 in TBST containing 5% BSA overnight), followed by similar secondary antibody incubation and ECL mediated visualization. X-ray films were scanned and the density of each band was quantified using ImageJ software (National Institutes of Health).

Quantitative realtime PCR

Total RNA was extracted from snap frozen P3 mouse calvaria using Trizol reagent (Invitrogen). The Superscript III RT system (Invitrogen) was used to synthesize cDNA from total RNA according to the manufacturer's protocol. Quantitative RT-PCR was performed on a LightCycler.v 1.5 (Roche) using gene-specific primers and SYBR Green I reagent (Roche). β 2-Microglobulin was used as the reference gene for normalizing cDNA concentrations.

In vivo bioluminescence imaging

P10 *Trap*^{-/-} mice and wildtype littermates that expressed the TGF β -reporter transgene (SBE-Luc mice) were injected with D-luciferin (Goldbio, 150 mg/kg, IP),

anaesthetized with isoflurane, and imaged 10 minutes after injection using a bioluminescence imaging system (Xenogen).

Primary osteoblast culture, TGF β -reporter cells

5 Bone marrow cells were isolated from tibias and femurs of approximately 2 month old *Crtap*^{-/-} and wildtype mice and cultured in α -MEM supplied with 10% FBS, 100 U/mL penicillin and 100 ug/mL streptomycin. Media was changed every second day and unattached cells were discarded. After 7 days, the attached cells, defined as bone marrow stromal cells (BMSCs), were reseeded to 24-well plates at
10 2.5 x10⁴ cells per cm² and cultured in osteogenic medium (α -MEM, 10% FBS, 500 μ M ascorbic acid, and 10 mM β -glycerophosphate) for 3 days. Conditioned medium was collected and incubated with PAI-luciferase reporter mink lung epithelial cells. After 24 hours, the cell lysates were collected for luciferase activity assays, which were measured using the Dual-Luciferase Reporter System (Promega). The results
15 were normalized to the total protein amount quantified using the Micro BCA reagent (Pierce).

MicroCT, bone histomorphometry

Lumbar vertebrae and femurs were scanned using a Scanco μ CT-40 microCT
20 for quantification of trabecular and cortical bone parameters. Vertebral and femoral trabecular bone parameters were analyzed using the Scanco analysis software by manually contouring the trabecular bone of vertebral body L4 as well as the distal metaphyseal section of the femur. The cortical bone parameters at the center of the femoral midshaft were quantified using the automated thresholding algorithm
25 included in the software.

Scanned undecalcified *Crtap*^{-/-} mouse spine samples were then embedded in plastic for sectioning. Toluidine blue staining and TRAP staining was performed using standard protocols for visualization and quantification of Ob's and Oc's, respectively, using the Bioquant Osteo Image Analysis System.

30

Immunostaining and histology

For immunohistochemistry, hind limbs of P5 mice were collected, fixed overnight in 4% paraformaldehyde and embedded in paraffin. After deparaffinization

and rehydration, heat-induced antigen retrieval was performed (Dako, S1700) followed by treatment with hyaluronidase for 30 min (2mg/ml; Sigma). Endogenous peroxidase was blocked using 3% hydrogen peroxide for 10 min. After incubation with blocking solution (3% normal goat serum, 0.1% BSA, 0.1% Triton X-100 in PBS), sections were incubated in antibodies for TGF β 1 (G1221, Promega) and decorin (LF-113, kindly provided from Larry Fisher, National Institute of Dental and Craniofacial Research, Bethesda, MD, USA) for 60 min (1:25 dilution each in PBS, control samples were incubated in PBS only) at 37C, and subsequently incubated with secondary antibody (SuperPicTure Polymer Detection kit, Invitrogen). Substrate DAB was added according to the manufacturer's recommendations and samples were dehydrated and mounted using Cytoseal XYL xylene based mounting medium (Thermo Scientific). Sections of WT and mutant littermates were processed at the same time. Images of the trabecular bone were taken with a light microscope (Axioplan 2, Zeiss) using identical exposure times for WT and mutant littermates.

Lungs of P10 and 16 week old *Crtap*^{-/-} mice were equally inflated during tissue collection, fixed in 4% paraformaldehyde, and were paraffin embedded. Lungs of P10 *Crtap*^{-/-} and wildtype mice were used for immunostaining for pSmad2. Briefly, paraffin sections were treated with xylene, rehydrated, and heated for 20 minutes for antigen retrieval (pH 6; Dako). Sections were then incubated in blocking solution (3% normal Donkey serum, 0.1% BSA, 0.1% Triton X-100 in PBS), and subsequently incubated with rabbit anti-pSmad2 antibody (1:500) (Cell signaling, #3108), donkey anti-rabbit secondary antibody conjugated to Alexa flour 594 (1:600) (Invitrogen), and mounted with Prolong Gold anti-fade reagent with DAPI (Invitrogen). Fluorescent images from these sections were taken using a Zeiss microscope (Axiovision Software) using identical exposure times.

For lung histology and morphometry of 16 week old mice, parasagittal sections were stained using a standard protocol for Hematoxylin and Eosin staining. The mean linear intercept (MLI) method was used to quantify the distance between alveolar structures. Briefly, 10 histological fields were captured per mouse at 20X magnification from all lobes of both lungs using a light microscope (Axioplan 2, Zeiss). The MLI was measured using modified ImageJ software (National Institutes of Health, modified by Paul Thompson). After manual removal of blood vessels, large airways and other nonalveolar structures, the software automatically thresholds

the alveolar tissue in each image and overlays a line grid comprised of 1,353 lines with each line measuring 21 pixels over the image. The number of lines that intercepted alveolar structures was used to calculate the MLI.

5 Biomechanical testing by 3-point bending

Crtap^{-/-} and WT femurs were tested by three point bending using a span of 6mm with an Instron 5848 device (Instron Inc., Norwood MA). All the femurs were tested wet at room temperature. They were preloaded to 1N at a rate of 0.05 N/s for 5 seconds. Following the pre-loading, the femurs were loaded to failure at a rate of 10 0.1mm/sec. Load and displacement data was captured at rate of 40Hz by using BLUEHILL Software (Instron 5848).

To determine the Yield Point, a region was identified after the preload and before the maximum load on the load-displacement curve. This region was separated into 5 segments from which the fitted line of the segment with greatest slope was 15 taken. Next, a 0.012mm offset was implemented on the line. The point of intersection between the offset line and the load-displacement curve was the 0.012 Offset Yield Point. This yield point corresponded more closely to a 0.2% offset strain, which is commonly chosen in the literature. The elastic region was identified as the region from the completion of the preload to the Yield Point. The Post-Yield 20 region was identified as the region from the Yield Point until the point at which the change in load exceeded -1N, indicating failure. The Elastic Displacement was the displacement during which specimen remained in the elastic region. The Post-Yield Displacement was the displacement during which the specimen remained in the Post-Yield region. The Total Displacement was calculated as the sum of the Elastic 25 Displacement and the Post-Yield Displacement. Using the trapezoidal numerical integration method, Energy to failure was calculated as the area under the Load-Displacement curve. The Maximum Load was determined by finding the highest load value recorded by BLUEHILL, before the specimen failed. To calculate Stiffness, the Least Square fit method was applied to the steepest segment of the elastic region of 30 the load-displacement curve. Stiffness was the slope of least square fit line. Geometric data (diameter and moment of inertia) obtained from microCT analysis of the femoral midshaft were utilized to calculate the intrinsic material properties: ultimate strength, toughness to failure and elastic modulus.

Serum bone turnover markers

Serum osteocalcin (OCN) was quantified using the Mouse Osteocalcin EIA Kit from Biomedical Technologies Inc. C-terminal cross-linked telopeptide of bone collagen (CTX) was quantified using the RatLapsTM EIA Kit from Immunodiagnostic Systems Ltd. Both analyses were performed according to the manufacturer's protocols.

Collagen SDS-PAGE, mass spectrometry and crosslinks analyses

For mass spectrometry, type I collagen was prepared from *Crtap*^{-/-} and wildtype tibiae. Bone was defatted with chloroform/methanol (3:1 v/v) and demineralized in 0.5 M EDTA, 0.05 M Tris-HCl, pH 7.5, all steps at 4°C. Bone were finely minced and collagen solubilized by heat denaturation (90°C) in SDS-PAGE sample buffer. Collagen α -chains were cut from SDS-PAGE gels and subjected to in-gel trypsin digestion. Electrospray MS was performed on the tryptic peptides using an LCQ Deca XP ion-trap mass spectrometer equipped with in-line liquid chromatography (LC) (ThermoFinnigan) using a C8 capillary column (300 μ m x 150 mm; Grace Vydac 208 MS5.315) eluted at 4.5 μ l min. Sequest search software (ThermoFinnigan) was used for peptide identification using the NCBI protein database.

Pyridinoline cross-links (HP and LP) were quantified by HPLC after hydrolyzing demineralized bone in 6N HCl.

Surface plasmon resonance analysis

Surface plasmon resonance experiments were carried out using a BIAcore X instrument (GE Healthcare Bio-Science Corp.). Purified native mouse tendon type I collagen from wild type and *Crtap*^{-/-} mice were immobilized on a CM5 sensor chip by amide coupling at a concentration of about 0.05 ng/mm² (500 RU) and 0.08 ng/mm² (800 RU), respectively. The experiments were conducted at a flow rate of 10 μ l/min and 20 °C in HBS-P buffer (10 mM Hepes buffer, pH 7.4, containing 150 mM NaCl and 0.005 % Surfactant P20). Recombinant human decorin core protein (R&D systems) was injected onto both type I CM5 chips. The concentration of the stock solution of human decorin was determined by amino acid analysis. The binding

response of decorin to wild type and *Crtap*^{-/-} mouse type I collagen was normalized by the amounts of immobilized type I collagen on the CM5 sensor chips. Three concentrations of decorin were used (3, 5 and 12 μM), for each concentration the analysis was repeated three times. This experiment was performed twice with
5 collagen isolated from different mice each time

Statistical methods

Comparisons between two groups were performed using unpaired, two-tailed Student's t-tests. For comparisons between 3 groups, One Way Analysis of Variance
10 (ANOVA) was performed if equal variance of groups was confirmed, followed by all pairwise multiple comparison using the Holm-Sidak method. If the equal variance test failed, Kruskal-Wallis One Way ANOVA on Ranks was performed, followed by all pairwise multiple comparison using the Tukey Test. A P value less than 0.05 was considered statistically significant for Student's t-test, ANOVA and Kruskal-Wallis
15 One Way ANOVA on Ranks. For posthoc pairwise multiple comparisons, each P value was compared to a critical level depending on the rank of the P value and the total number of comparisons made to determine if differences between groups are significant. Sigma Plot V11.0 (Systat Software Inc.) was used for statistical analyses.

The effects of 1D11 on bone and lungs of OI mice were unknown at study
20 start. To determine the initial sample size per group of mice we calculated that to detect a minimal difference of 20% in bone mass (BV/TV) by MicroCT between 1D11 and control treated OI mice with a 90% power, a group size of 8 mice is required.

25 **Example 1: Altered TGFβ signaling in *Crtap*^{-/-} calvaria**

Crtap^{-/-} mice and age-matched wild type (WT) littermate controls were analyzed for expression of activated pSmad 2, a member of the TGFβ signaling pathway, as well as other downstream targets of TGFβ. Calvaria bones were excised and RNA and protein were extracted and analyzed by Realtime-PCR and Western
30 blot, respectively. As can be seen in **Figure 1A and 1B**, *Crtap*^{-/-} mice had a 100% higher ratio of activated pSmad2 to total Smad2 compared to WT mice, as measured by Western blot and quantified by densitometry, indicating that TGFβ signaling is elevated in *Crtap*^{-/-} mice. Transcriptional targets of TGFβ, such as Col1a1 and p21,

were elevated compared to WT controls, as measured by RT-PCR and demonstrated in **Figure 1C** and **Figure 1D**, respectively. The pro-fibrotic ECM protein connective tissue growth factor (CTGF) was measured and found to be approximately 50% higher in *Crtap*^{-/-} mice compared to WT controls, as determined by RT-PCR and demonstrated in **Figure 1E**. As shown in **Figure 1F**, RT-PCR analysis revealed that expression of the cyclin-dependent kinase inhibitor p27 was not altered in either the *Crtap*^{-/-} or the WT mice.

Example 2: Increased TGFb activity in *Crtap*^{-/-} mice *in vivo* and in *crtap*^{-/-} osteoblastic cells

Crtap^{-/-} mice were crossed to TGFβ reporter mice that express luciferase in response to activation of TGFβ signaling (Jackson Laboratory; B6.Cg-Tg(SBE/TK-luc)7Twc/J). P9 mice were injected with the substrate D-Luciferin (150 mg/kg) 10 minutes before imaging (Xenogen; IVIS camera system). As demonstrated in **Figure 2A**, *Crtap*^{-/-} mice had considerably higher luminescence in their tails, long bones, and calvaria compared to WT controls, indicating increased TGFb activity in *crtap*^{-/-} mice. **Figure 2B**, quantification of luciferase activity at calvaria.

Bone marrow stromal cells (BMSCs) were isolated from *Crtap*^{-/-} mice and WT mice, cultured under osteogenic conditions *ex vivo*, and the conditioned culture medium was analyzed for TGFβ activity using a cell line that expresses luciferase in response to activation of TGFβ signaling. As shown in **Figure 2C**, conditioned medium from *Crtap*^{-/-} BMSCs resulted in nearly a two-fold greater luciferase activity of the reporter cell line compared to BMSCs from WT mice. Together, these data indicate that TGFβ secretion and activity is elevated in the bones and osteoblastic cells of *Crtap*^{-/-} mice.

Example 3: μCT analysis of *Crtap*^{-/-} vertebrae

Adult 8 week old *Crtap*^{-/-} mice (N = 6 per group) were administered 1D11 (10mg/kg, I.P., 3 times/week, 8 weeks total), a murine surrogate of the pan-specific antibody that binds to TGFβ comprising a heavy chain comprising the amino acid sequence of SEQ ID NO: 14 and a light chain comprising the amino acid sequence of SEQ ID NO: 15. An unrelated 13C4 antibody was administered to a separate group of *Crtap*^{-/-} mice and WT mice as a control (N = 6). The L4 vertebral bodies of 16

week old mice (treated from week 8-16) were imaged by μ -CT. MicroCT data of vertebral body L4 from 8 week old *Crtap*^{-/-} mice (n=6 per group) that were treated with the TGF β neutralizing antibody 1D11 (Genzyme; 10mg/kg, I.P., 3 times/week) for 8 weeks and wild-type (WT) and control *Crtap*^{-/-} mice that were treated with a control antibody (13C4-placebo) is shown in **Figure 3**. As shown in **Figure 3**, *Crtap*^{-/-} vertebrae were cavernous compared to WT control vertebrae. However, 1D11 treatment result in a skeletal phenotype that was comparable to the WT condition.

The data from **Figure 3** was quantified in **Figure 4**, treatment with the pan-specific anti-TGF β antibody rescued the skeletal phenotype of *Crtap*^{-/-} mice.

10 Vertebrae from treated *Crtap*^{-/-} mice were statistically similar to WT control mice in measured parameters, including bone volume density (BV/TV), total bone surface (BS), bone surface density (BS/BV), trabecular number (Tb.N), trabecular thickness (Tb.Th), trabecular spacing (Tb.Sp), and total volume (Dens TV).

15 **Example 4: Histomorphometry of anti-TGF β treated *Crtap*^{-/-} vertebrae**

In addition to μ -CT, the vertebral bones of antibody-treated, placebo-treated, and WT mice were analyzed by histomorphometry. As shown in **Figure 5A** and **5B**, μ -CT results were confirmed by histomorphometric analysis. In addition, tissue sections were stained for expression of the osteoclast marker TRAP. Analysis
20 revealed that there were more osteoclasts covering more of the bone surface in *Crtap*^{-/-} mice compared to WT controls (N.Oc/BS and Oc.S/BS), indicating increased osteoclastic activity. Treatment with the 1D11 anti-TGF β reduced all osteoclast-specific parameters to below WT numbers. Thus, osteoclasts were identified as a potential target for TGF β antibodies, and more specifically, pan-specific anti-TGF β
25 antibodies.

Example 5: Three-point bending test of anti-TGF β treated *Crtap*^{-/-} femurs

Biomechanical testing was performed on the excised femurs of 16 week old mice (after treatment from week 8-16) using a standard three-point bending test with
30 an Instron 5848 device (Instron Inc., Norwood MA) with a 6mm span, preloaded to 1N at a rate of 1N/s for 5 seconds. Following the pre-loading, femurs were compressed to failure at a rate of 0.1mm/sec. Load and displacement data were captured at rate of 40Hz by using BLUEHILL Software (Instron 5848).

As demonstrated in **Figure 6**, *Crtap*^{-/-} mice femurs were less stiff and were able to withstand a significantly smaller maximal load compared to WT control mice. Femurs of 1D11 treated *Crtap*^{-/-} mice showed a significant improvement of the maximum load, and a trend to an increased stiffness compared to control *crtap*^{-/-} mice.

Thus, treatment with the 1D11 pan-specific anti-TGFβ antibody quantitatively, qualitatively, and biomechanically restored the skeletal phenotype of *Crtap*^{-/-} mice.

Example 6: Inhibition of TGFβ signaling with 1D11 ameliorates the lung phenotype

Crtap^{-/-} mice have a generalized connective tissue disease manifested by low bone mass, glomerulosclerosis, and pulmonary dysplasia (Baldridge *et al.*; PLoSone, 5(5):e10560 (2010)). Increased TGFβ expression was seen the lungs of *Crtap*^{-/-} mice, as evidenced by positive immunostaining for pSmad2 and demonstrated in **Figure 7A**. Histologically, *Crtap*^{-/-} mice exhibited increased distal airway space compared to WT mice, as shown in **Figure 7B**. 1D11 treatment (10mg/kg, IP, 3x/week for 8 weeks) reduced pSmad2 expression in *Crtap*^{-/-} mice and reduced the distal airway space and ameliorated the lung phenotype, as demonstrated in **Figure 7A** and **Figure 7B** and quantified in **Figure 7C** (*P<0.05 vs. control *Crtap*^{-/-}; 10 images analyzed per mouse, n=8 mice per group).

Example 7: Decorin expression in *Crtap*^{-/-} lungs

Transcriptional regulators of TGFβ expression were investigated in order to understand the basis of dysregulated TGFβ signaling in the bones and lungs of *Crtap*^{-/-} mice. A major class of extracellular proteins that can regulate TGFβ in ECM include the small leucine rich proteoglycans (SLRP), such as decorin. Immunostaining revealed increased expression of decorin in *Crtap*^{-/-} lungs compared to WT control lungs, as shown in **Figure 8**.

As decorin is a regulator of mature TGFβ, this finding suggests that altered post-translational modification of collagen, as occurs in OI, alters the interactions of ECM proteins, including SLRPs. Decorin binds to hydroxyproline sites, such as the one located at amino acid residue 396 of type-I and type-II collagens, which are absent in *Crtap*^{-/-} mice. A decorin binding assay was performed to determine whether decorin binding may be altered in OI, and whether this may be at least partly

responsible for the phenotypes observed in *Crtap*^{-/-} bones and mice. As shown in **Figure 9**, decorin binding to 3-hydroxylated collagen peptides (as in type I and II collagen) was greater than collagen peptides without 3-hydroxylation (as in type III collagen).

5

Example 8: Increased TGFβ signaling is a common mechanism in Osteogenesis Imperfecta

OI is characterized by brittle bones, low bone mass, bone deformities and fractures. In addition, extraskeletal manifestations including lung abnormalities contribute substantially to morbidity and mortality. Most cases of OI are caused by autosomal dominant mutations in the genes encoding type I collagen (*COL1A1* and *COL1A2*). In recent years, mutations in additional genes encoding the proteins involved in the post-translational modification of collagen have been identified as causing recessive forms of OI. The first described was in cartilage associated protein (*CRTAP*), a member of the prolyl-3-hydroxylase complex that is responsible for 3-hydroxylation of proline residue 986 α1(I) in type I collagen. Hypomorphic *CRTAP* mutations lead to partial loss of 3-hydroxyproline (3Hyp) in fibrillar collagen as well as overmodification of other residues and result in recessive OI type VII, which clinically overlaps with dominant forms of severe OI. The physiological function of 3Hyp is not completely understood, but biochemical studies suggest that it may be involved in collagen-protein interactions, rather than negatively affecting collagen stability.

The ECM is an important reservoir for signaling molecules and their regulators. In bone, TGFβ acts as a central coordinator of bone remodeling by coupling the localized activity of bone resorbing osteoclasts and bone forming osteoblasts. TGFβ is abundantly produced by osteoblasts, is secreted predominantly in inactive latent forms, and is deposited into the bone matrix. Here, it can be released and activated during bone resorption by osteoclasts. As an additional level of regulation, active TGFβ can be bound by proteoglycans, which modulate its bioactivity in association with collagen fibrils. Because type I collagen is the most abundant component of the ECM in bone, this raises the intriguing hypothesis that the alteration of collagen structure observed in OI not only increases bone fragility, but also affects the signaling reservoir function of the bone matrix. Interestingly, *Crtap*^{-/-}

mice show phenotypic overlap with animal models of increased TGF β signaling. For example, TGF β overexpression results in low bone mass. In addition, *Crtap*^{-/-} mice exhibit an enlargement in alveolar airway space in lungs, which is similar to that observed in a mouse model of Marfan syndrome, where increased TGF β signaling has been shown to be a major contributor to the lung pathology. Therefore, the status of TGF β signaling was studied in the *Crtap*^{-/-} mouse model of recessive OI.

To assess the status of TGF β signaling in bone, the expression levels of TGF β target genes in calvarial bone of *Crtap*^{-/-} mice were evaluated. Compared with wild type (WT) samples, *Crtap*^{-/-} bone showed an increased expression of the TGF β downstream targets *p21* (cyclin-dependent kinase inhibitor 1), *PAI-1* (plasminogen activator inhibitor-1), and *Col1a1*, consistent with elevated TGF β activity (**Figure 10A**). To confirm increased activation of the intracellular TGF β signaling pathway, the status of Smad2, an intracellular second messenger protein, which becomes phosphorylated after activation of TGF β receptors, was evaluated. Consistent with target gene expression, immunoblot analyses demonstrated a higher ratio of phosphorylated Smad2 (pSmad2) to total Smad2 in bone samples of *Crtap*^{-/-} mice, indicating increased TGF β signaling (**Figures 10B and 10C**).

To determine whether these static measures reflect increased TGF β activity *in vivo*, *Crtap*^{-/-} mice were intercrossed with TGF β -reporter mice expressing luciferase under control of TGF β -responsive Smad binding elements (SBE-Luc mice). Compared with WT/SBE-Luc littermates, *Crtap*^{-/-}/SBE-Luc mice showed an increase in bioluminescence of areas over skeletal structures, indicating increased TGF β activity *in vivo* (**Figure 10D**). In 3 litters, *Crtap*^{-/-} mice show a mean 2.86 fold (SD $\pm$ 0.34) bioluminescence signal at the head/calvaria compared with WT mice. Moreover, in order to test whether the increased TGF β /Smad signaling associated with loss of *Crtap* is intrinsic to bone, *i.e.*, tissue autonomous, bone marrow stromal cells (BMSCs) were differentiated to osteoblastic cell *in vitro*. By using a TGF β reporter cell line, it was found that conditioned medium from *Crtap*^{-/-} BMSCs exhibited higher TGF β activity compared with medium from WT BMSCs (**Figure 10E**). Together, these findings indicate that loss of *Crtap* enhances TGF β signaling in bone in a tissue autonomous fashion.

Patients with severe OI can also exhibit intrinsic lung abnormalities, and respiratory failure is one of the leading causes of death in these individuals.

Interestingly, *Crtap*^{-/-} mice show a diffuse increase in alveolar airway space, a feature associated with increased TGFβ signaling in other developmental models.

Accordingly, lungs of *Crtap*^{-/-} mice showed increased intracellular staining for pSmad2 in alveolar cells, indicating that the increased TGFβ activity is also present in extraskkeletal tissues (**Figure 10F**).

To understand whether increased TGFβ signaling represents a causal mechanism contributing to the bone and lung phenotypes in *Crtap*^{-/-} mice, a rescue experiment was performed with a pan-TGFβ neutralizing antibody (1D11). Eight week old *Crtap*^{-/-} mice were treated with 1D11 for 8 weeks; control *Crtap*^{-/-} and WT mice received a non-specific control antibody (13C4). 1D11 did not significantly change body weight of the treated *Crtap*^{-/-} mice, indicating that TGFβ inhibition did not affect the general nutritioinal status (**Figure 14**). In addition, mass spectrometric and cross-links analyses showed that 1D11 did not significantly change the status of type I collagen P986 3-hydroxylation or collagen crosslinks in *Crtap*^{-/-} mice, suggesting that dysregulated TGFβ signaling is a consequence of the altered molecular collagen structure, and not directly involved in intracellular collagen processing or extracellular fibril assembly (**Figure 15**). *Crtap*^{-/-} mice exhibit a reduced bone mass and abnormal trabecular bone parameters (**Figures 11A and 11B**). MicroCT imaging analysis of vertebrae demonstrated that compared with control *Crtap*^{-/-} mice, TGFβ inhibition significantly improved trabecular bone parameters, including bone volume/total volume, trabecular number and trabecular thickness to near WT levels (**Figures 11A and 11B, and Figure 17**). Similar beneficial effects were observed in femoral trabecular bone in *Crtap*^{-/-} mice, where TGFβ inhibition significantly improved trabecular bone parameters (**Figure 18**). The effects of TGFβ inhibition on the skeleton with 1D11 have been reported previously in WT mice and in *Esl-1*^{-/-} mice¹¹, a model with increased TGFβ activity due to a defect in normal TGFβ maturation. While 1D11 moderately increased trabecular BV/TV by 33% in the spine in WT mice, *Esl-1*^{-/-} mice exhibited an 106% increase in BV/TV. This suggested that targeting TGFβ in a pathophysiological situation where it is increased in the skeleton, could lead to a relatively more pronounced positive effect. In the present study, 1D11 increased the trabecular BV/TV at the spine by 235% in *Crtap*^{-/-} mice, supporting that the dysregulated TGFβ signaling is an important contributor of the low bone mass in *Crtap*^{-/-} mice. At the femur midshaft, the parameters of cortical

architecture including cortical thickness, diameter, cross-sectional area and cross-sectional moments of inertia in *Crtap*^{-/-} mice were significantly reduced compared to WT mice. Following 1D11 treatment, these parameters were no longer significantly different from WT mice (**Figure 19**). To test if these changes in cortical and trabecular bone translated into improved bone strength, biomechanical testing of the femurs by 3-point bending was performed. It was found that TGFβ inhibition was able to increase maximum load and ultimate strength in the treated *Crtap*^{-/-} mice, indicating improved whole bone and tissue strength and improved resistance to fracture. However, 1D11 treatment had no effects on the increased brittleness of the OI bone, as indicated by the reduced post-yield displacement in both control and 1D11 treated *Crtap*^{-/-} mice (**Figure 20**). This likely reflects the inherent abnormal mineralization associated with altered collagen structure. Taken together, these findings indicate that increased TGFβ signaling is major contributor to the bone phenotype in recessive OI resulting from *Crtap* deficiency and that inhibition of dysregulated TGFβ signaling restores bone mass, microstructural parameters and improves whole bone strength.

To understand the effects of TGFβ inhibition in *Crtap*^{-/-} mice at the cellular level, histomorphometric analyses on treated mice was performed. In sections of vertebral bodies in this study it was found increased osteoclast (Oc) and osteoblast (Ob) numbers per bone surface in control *Crtap*^{-/-} compared to WT mice, indicating increased bone remodeling in the spine (**Figure 11C** and **Figure 21**). Consistently, the serum bone turnover markers osteocalcin (OCN) and C-terminal cross-linked telopeptide of bone collagen (CTX) were elevated in 8 week old (OCN and CTX) and 16 week old (CTX only) control *Crtap*^{-/-} mice (**Figure 19**). Similar changes in the cellular composition of bone have been described in patients with dominant and recessive OI, showing increased Oc and Ob numbers consistent with an increased bone turnover. Interestingly, mouse models of increased TGFβ signaling also show low bone mass with increased osteoclastic bone resorption and abnormal bone remodeling. Most reports of the effects of TGFβ on bone cells are consistent with a model where TGFβ can stimulate the recruitment and initial differentiation of Oc and Ob precursors at the site of bone repair, followed by insulin-like growth factor 1 (IGF-1) mediated Ob differentiation. However, at persistently high doses, TGFβ can inhibit Ob differentiation by repressing the differentiation factor RUNX2. Given the

crucial effects on Oc/Ob interaction, fine tuning of TGF β availability is a key factor for the local coupling of bone resorption with formation during bone remodeling and its imbalance can lead to significant bone pathology.

In contrast to the findings in control *Crtap*^{-/-} mice, bone sections of *Crtap*^{-/-} mice treated with 1D11 revealed reduced Oc and Ob numbers, which were even lower than the values measured in the WT mice, indicating a supraphysiologic suppression of dysregulated bone remodeling as a result of TGF β inhibition at the dose of 1D11 used in this experiment (**Figure 11C**). Consistent with an earlier report, the observation of a reduction of Oc's and Ob's below WT levels also underscores the physiological requirement of local amounts of TGF β to normally coordinate Oc's and Ob's during the bone remodeling process. Our findings are different from previous studies in WT mice, where 1D11 treatment reduced Oc numbers but increased Ob numbers. This may reflect distinct cellular effects of TGF β inhibition in a pathophysiological situation with increased TGF β signaling and increased bone remodeling compared with normal bone in WT mice. TGF β has been shown to inhibit differentiation of osteoblast precursor cells, and increased TGF β signaling could thereby lead to a higher proportion of immature osteoblast lineage cells. On the other hand, an increased number or higher proportion of immature Ob's on the bone surface could result in an increased amount of secreted TGF β by these cells. The finding that TGF β inhibition with 1D11 significantly reduces the increased Ob numbers in *Crtap*^{-/-} mice suggests that the increased TGF β signaling causally contributes to the increase in osteoblast lineage cells.

In addition to the findings regarding Oc and Ob numbers, greater osteocyte (Ot) numbers per bone area in control *Crtap*^{-/-} mice were observed, which were reduced to levels comparable with those of WT mice in 1D11 treated *Crtap*^{-/-} mice (**Figure 11C** and **Figure 21**). In OI patients, an increased Ot density has been observed in individuals with more severe forms of the disease, likely reflecting the presence of immature primary bone due to a defect in physiological maturation in OI bone. Consistent with our hypothesis that increased TGF β signaling contributes to the bone pathology in OI, overexpression of TGF β in WT mice similarly results in an increased Ot density. As a possible explanation, TGF β can inhibit Ob apoptosis during the transition of Ob's to Ot's, and thereby lead to an increased Ot density. Collectively, these findings indicate that increased TGF β signaling contributes to a

high bone turnover status and impaired bone maturation in *Crtap*^{-/-} mice and that inhibition of dysregulated TGFβ signaling reverses these cellular alterations.

Given these crucial effects on Oc/Ob interaction, fine tuning of TGFβ availability is a key factor for the local coupling of bone resorption with bone formation during bone remodeling and its imbalance can lead to significant bone pathology. Our findings indicate that inhibition of dysregulated TGFβ signaling in *Crtap*^{-/-} mice restores bone mass as well as microstructural parameters, improves whole bone strength, and reverses the cellular alterations observed in *Crtap*^{-/-} mice. Therefore, dysregulation of TGFβ signaling is an important contributor to the bone phenotype in this mouse model of recessive OI.

We were also interested in whether TGFβ inhibition affected the lung phenotype of *Crtap*^{-/-} mice. Lungs of control *Crtap*^{-/-} mice show an increase in the distal airway space compared with WT mice (**Figure 11D**). Interestingly, lungs of *Crtap*^{-/-} mice treated with the TGFβ neutralizing antibody showed a 60% improvement in the distance between alveolar structures (**Figures 11D and 11E**). This finding indicates that excessive TGFβ signaling is also an important pathogenic contributor to the lung abnormalities present in *Crtap*^{-/-} mice. Increased TGFβ signaling has been linked to developmental pulmonary abnormalities as well as disease in mature lungs. For example, TGFβ overexpression in lungs results in impaired lung development with areas of enlarged airway space and increased TGFβ signaling is a contributing pathomechanism in lung abnormalities in Marfan syndrome as well as in the development of emphysema and bronchial asthma. Our results indicate that excessive TGFβ signaling is an important pathogenic contributor to the lung abnormalities present in *Crtap*^{-/-} mice. Given the partial rescue of the lung phenotype with 1D11 in *Crtap*^{-/-} mice, it is possible that dysregulated TGFβ signaling affects pulmonary tissue development when the anatomic structures are established, in addition to maintaining lung tissue at later stages when TGFβ inhibition is able to ameliorate the phenotype.

The next question asked was how alterations in collagen due to loss of *Crtap* (leading to the loss of 3Hyp at P986 and post-translational over modification of collagen) result in dysregulated TGFβ signaling. Biochemical analyses indicate that collagen prolyl-3-hydroxylation does not fundamentally affect the stability of collagen molecules, but instead it may affect collagen-protein interactions. An

attractive hypothesis is that loss of 3Hyp could affect collagen interaction with small leucine-rich proteoglycans (SLRPs). SLRPs are known to bind to both type I collagen as well as TGF β , and thereby modulate TGF β activity. For example, the SLRP decorin is able to inhibit distinct effects of TGF β in osteosarcoma cells whereas it
5 enhances TGF β activity in preosteoblastic cells. The binding region of decorin on type I collagen is suggested to center at residues 961/962 of the triple helical domain, which is located in close proximity to the P986 residue, that is unhydroxylated in OI due to *Crtap* deficiency. Therefore, it is possible that the P986 3Hyp position marks an interacting site for the binding of decorin to type I collagen, thereby mediating the
10 sequestration of mature TGF β to collagen.

Hence, it was hypothesized that decorin binding to collagen is critical for TGF β regulation and that this binding is disrupted with altered collagen structure, for example by loss of post-translational 3-prolyl-hydroxylation modification of P986 in the α 1 chain of type I collagen in the case of recessive OI. It was identified that
15 although loss of *Crtap* did not alter the RNA expression of decorin and other SLRPs in calvarial bone (**Figure 12A**), nor the qualitative abundance of decorin in trabecular bone (**Figure 24**), it did reduce binding of recombinant decorin core protein to type I collagen isolated from *Crtap*^{-/-} mice versus WT mice (**Figure 12B**). Surface plasmon resonance analysis measurements of the binding of recombinant decorin core protein
20 to type I collagen of WT and *Crtap*^{-/-} mice demonstrated reduced binding in *Crtap*^{-/-} mice at the three concentrations tested (**Figure 23**). Three technical replicates at each of the indicated concentrations of decorin were performed from two independent biological replicates (◆ replicate 1, ▲ replicate 2). Results are shown as the percentage of the mean of WT (bars indicate mean per group). The mean reductions
25 of decorin binding to *Crtap*^{-/-} type I collagen at 3, 5 and 12 μ M of decorin were 28.5%, 33.5% and 38.1%, respectively.

This finding suggests that alterations of collagen-proteoglycan interactions may contribute to the dysregulated TGF β signaling in bone and other collagen rich tissues in OI. Based on the reported requirement of decorin-collagen binding for
30 decorin to effectively reduce TGF β bioactivity, it is possible that the defects in OI collagen lead to altered binding of decorin, and hence, its ability to sequester TGF β in the matrix and modulate TGF β functions. Hence, altered proteoglycan-collagen interactions may contribute to dysregulated TGF β signaling in bone and other

collagen rich tissues in OI, even if no major changes in absolute TGF β levels are present (**Figure 24 and Figure 25**). This notion is supported by the finding that *COL1A1* and *COL1A2* mutations in more severe forms of dominant OI cluster in specific regions that are known to bind proteoglycans, further supporting a physiological relevance of proteoglycan-collagen interactions for normal bone homeostasis. This would also imply that other proteoglycans that are competing with decorin for the collagen binding site may also contribute to dysregulated TGF β activity, and that additional signaling pathways could be altered.

Because of the clinical overlap of some recessive and dominant forms of OI where defective structure of collagen fibers leads to brittle bones and extraskeletal manifestations, it is possible that dysregulation of TGF β signaling is a common pathophysiological disease mechanism. To address this hypothesis, the status of TGF β signaling in a mouse model of dominant OI was investigated. Knock-in mice carrying a G610C mutation in the *Col1a2* gene (*Col1a2^{tm1.1Mcb}*) phenocopy a dominantly inherited, moderate form of OI that was originally identified in an Amish population. In bone samples of *Col1a2^{tm1.1Mcb}* mice, increased expression of the TGF β target genes *p21* and *PAI-1* was found, indicating upregulation of TGF β signaling (**Figure 13A**). Consistently, immunoblot analyses of bone extracts from *Col1a2^{tm1.1Mcb}* mice also showed an increased ratio of activated pSmad2/total Smad2, similar to our observation in the *Crtap^{-/-}* mice (**Figures 13B and 13C**).

To test if the increased TGF β signaling in this model of dominant OI also represents a causal mechanism, 8 week old *Col1a2^{tm1.1Mcb}* mice were treated with the TGF β -neutralizing antibody 1D11 for 8 weeks; control *Col1a2^{tm1.1Mcb}* and WT mice were treated with the control antibody 13C4. Similar to the findings in *Crtap^{-/-}* mice, 1D11-treatment restored the trabecular bone parameters at the spine to WT levels (**Figures 13D, 13E, and 22**). Taken together, these findings indicate that dysregulated TGF β signaling is also an important contributor to the pathogenesis of dominant forms of OI, and that anti-TGF β therapy corrects the bone phenotype in dominant OI.

From a clinical-translational perspective, potential negative effects of systemic TGF β inhibition in OI patients have to be considered. While *TGF- β 1^{-/-}* mice develop a severe multifocal inflammatory disease and dysregulation of the immune system within the first weeks of life, in both *Crtap^{-/-}* and *Col1a2^{tm1.1Mcb}* mice treated with 1D11 we did not observe obvious negative effects on general health, behavior or

growth, suggesting that the effects of a partial pharmacological inhibition of TGF β ligands in adult mice are different from a complete loss of TGF β 1 during development. In humans, Fresolimumab (GC1008, Genzyme), which is similar to 1D11 in its affinity and specificity to the 3 isoforms of TGF β , was used in phase I
5 studies in patients with treatment-resistant primary focal segmental glomerulosclerosis, idiopathic pulmonary fibrosis and malignant melanoma or renal cell carcinoma. In these studies, Fresolimumab was in general well-tolerated, with possible, dose-related adverse events including skin rashes or lesions, epistaxis, gingival bleeding and fatigue.

10 The molecular mechanisms of OI are incompletely understood. As a result, current treatment options for patients with OI are mainly limited to anti-resorptive therapies as used for the treatment of osteoporosis. Interestingly, a recent randomized, placebo controlled trial of the anabolic agent teriparatide in adults with OI showed that severe OI type III/IV responded differently than did those with mild
15 OI type I (Orwoll et al., 2014). This suggests genotypic differences in response to therapies targeted at modifying cell signaling and that TGF β -targeted treatment may be a promising option to further study in severe OI due to collagen and collagen post-translational modification gene mutations. Overall, our data support the concept of dysregulated matrix-cell signaling as a mechanism in the pathogenesis of different
20 genetically inherited forms of brittle bone disease and point to a disease-specific mechanism-based strategy for the treatment of OI by neutralizing the overactive TGF β activity in skeletal and extraskelatal tissues.

Claims

1. Use of an antibody or an antigen binding fragment thereof that specifically binds to and neutralizes TGF β 1, TGF β 2, and TGF β 3 in the preparation of a medicament for treating osteogenesis imperfecta (OI) in a human subject in need thereof, wherein

the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 10, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 11, and wherein the medicament is to be administered in combination with a bisphosphonate.

2. The use according to claim 1, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15.

3. The use according to claim 1 or 2, wherein the bisphosphonate is alendronate.

4. The use according to claim 1 or 2, wherein the bisphosphonate is pamidronate.

5. The use according to claim 1 or 2, wherein the bisphosphonate is zoledronate.

6. The use according to claim 1 or 2, wherein the bisphosphonate is risedronate.

7. The use according to claim 1 or 2, wherein the bisphosphonate is etidronate.

8. The use according to claim 1 or 2, wherein the bisphosphonate is clodronate.
9. The use according to claim 1 or 2, wherein the bisphosphonate is tiludronate.
10. The use according to claim 1 or 2, wherein the bisphosphonate is neridronate.
11. The use according to claim 1 or 2, wherein the bisphosphonate is olpadronate.
12. Use of an antibody or an antigen binding fragment thereof that specifically binds to and neutralizes TGF β 1, TGF β 2, and TGF β 3 in the preparation of a medicament for treating osteogenesis imperfecta (OI) in a human subject in need thereof, wherein
the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 10, and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 11, and wherein the medicament is to be administered in combination with teriparatide.
13. The use according to claim 12, wherein the antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 14, and a light chain comprising the amino acid sequence of SEQ ID NO: 15.