

THE DESCRIPTION

POLYMORPHIC FORMS OF ICOTINIB AND USES THEREOF

FIELD OF THE INVENTION

The present invention relates to the polymorphic forms of Icotinib, processes for preparing
5 these polymorphic forms, pharmaceutical compositions thereof, and use of new polymorphic
forms and pharmaceutical compositions for the treatment of cancer and cancer
occurrence-related diseases.

BACKGROUND OF THE INVENTION

Tyrosine kinase receptors are trans-membrane proteins that, in response to an
10 extracellular stimulus, propagate a signaling cascade to control cell proliferation, angiogenesis,
apoptosis and other important features of cell growth. One class of such receptors, epidermal
growth factor receptor (EGFR) tyrosine kinases, are overly expressed in many human cancers,
including brain, lung, liver, bladder, breast, head and neck, esophagus, gastrointestinal, ovary,
cervix or thyroid cancer.

15 EGFR is expressed in many types of tumor cells. Binding of cognate ligands (including
EGF, TGF α (i.e., Transforming Growth Factor- α) and neuregulins) to the extracellular domain
causes homo- or heterodimerization between family members. The juxtaposition of
cytoplasmic tyrosine kinase domains results in transphosphorylation of specific tyrosine, serine
and threonine residues within each cytoplasmic domain. The formed phosphotyrosines act as
20 docking sites for various adaptor molecules and subsequent activation of signal transduction
cascades (Ras/mitogen-activated, PI3K/Akt and Jak/STAT) that trigger proliferative cellular
responses.

Various molecular and cellular biology and clinical studies have demonstrated that EGFR
tyrosine kinase inhibitors can block cancer cell proliferation, metastasis and other EGFR-related
25 signal transduction responses to achieve clinical anti-tumor therapeutic effects. Two oral EGFR

kinase inhibitors with similar chemical structures are Gefitinib (Iressa, AstraZeneca), approved by the U.S. FDA for advanced non-small cell lung cancer in 2003 (and later withdrawn), and Erlotinib Hydrochloride (Tarceva, Roche and OSI), approved by the U.S. FDA for advanced non-small cell lung cancer and pancreatic cancer treatment in 2004.


Many pharmaceutically active organic compounds can crystallize in more than one type of three-dimensional crystal structure. That is, the compounds may crystallize in different crystalline forms. This phenomenon (identical chemical structure but different crystalline structure) is referred to as polymorphism, and the species having different molecular structures are referred to as polymorphs.

Polymorphs of a particular organic pharmaceutical compound may have different physical properties, such as solubility and hygroscopicity, due to their distinct three-dimensional crystal structures. However, it is generally not possible to predict whether a particular organic compound will form different crystalline forms, let alone predict the structure and properties of the crystalline forms themselves. The discovery of a new crystalline or polymorph form of a pharmaceutically useful compound may provide a new opportunity for improving the overall characteristics of a pharmaceutical product. It enlarges the repertoire of materials that a formulation scientist has available for designing. It may be advantageous when this repertoire is enlarged by the discovery of new polymorphs of a useful compound.

Chinese Patent Publication No. CN1305860C discloses the structure of Icotinib(free base) on page 29, Example 15, Compound 23, and WO 2010/003313 disclosed Icotinib hydrochloride and its new crystalline polymorphs.

DESCRIPTION OF THE INVENTION

The present invention relates to Icotinib (i.e., the compound of Formula I), approximately pure polymorph forms, and pharmaceutical acceptable salt thereof.



Formula I .

The compound of Formula I of the present invention exists in one or more polymorphic forms. These polymorphic forms are referred as Form I, Form II, Form III and Form IV. The compound of Formula I and its polymorphic forms have better solubility and chemical stability, making them preferable for clinical applications.

5 In one aspect, the polymorph form is of Polymorph Form I and its X-ray powder diffraction pattern having characteristic peaks at diffraction angles 2θ of approximately 7.6° , 14.7° , 20.9° , 23.1° and $27.5^\circ \pm 0.2^\circ$.

In some other embodiments of Polymorph Form I, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $11.6\text{\AA}$, $6.0\text{\AA}$, $4.3\text{\AA}$, $3.9\text{\AA}$
10 and $3.3\text{\AA}$.

In some embodiments of Polymorph Form I, its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 7.6° , 14.7° , 20.9° , 23.1° , 24.2° , 27.5° , 29.8° and $30.9^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form I, its X-ray powder diffraction
15 pattern has characteristic peaks, expressed in terms of the interplanar distance, of $11.6\text{\AA}$, $6.0\text{\AA}$, $4.3\text{\AA}$, $3.9\text{\AA}$, $3.7\text{\AA}$, $3.3\text{\AA}$, $3.0\text{\AA}$ and $2.9\text{\AA}$.

Preferably, the X-ray powder diffraction pattern is shown as in Figure 1.

Preferably, the melting point of the Polymorph Form I is $175\text{--}177^\circ\text{C}$.

The present also provides a method of preparing the polymorph, comprising the steps of
20 dissolving Icotinib hydrochloride in the mixture of $\text{C}_1\text{--}\text{C}_4$ alcohol and water, the mixture was warmed to 60°C and a solution of base was dropped. The reaction solution was stirred for 1h and cooled down to room temperature. The precipitate was filtered and washed with purified water and dried 8h in vacuum below 60°C to obtain the resulted Polymorph Form I.

In another aspect, the polymorph form is of Polymorph Form II and its X-ray powder
25 diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 5.5° , 19.7° and $22.6^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form II, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $16.0\text{\AA}$, $4.5\text{\AA}$ and $3.9\text{\AA}$.

30 In some embodiments of Polymorph Form II, its X-ray powder diffraction pattern has

characteristic peaks at diffraction angles 2θ of approximately 5.5° , 11.0° , 19.7° , 20.9° and $22.6^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form II, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $16.0\text{\AA}$, $8.0\text{\AA}$, $4.5\text{\AA}$, $4.3\text{\AA}$, and $3.9\text{\AA}$.

In some other embodiments of Polymorph Form II, its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 5.5° , 8.8° , 11.0° , 16.5° , 19.7° , 20.9° , 22.6° and $23.7^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form II, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance of $16.0\text{\AA}$, $10.1\text{\AA}$, $8.0\text{\AA}$, $5.4\text{\AA}$, $4.5\text{\AA}$, $4.3\text{\AA}$, $3.9\text{\AA}$ and $3.8\text{\AA}$.

Preferably, the X-ray powder diffraction pattern is shown as in Figure 2.

Preferably, the melting point of the Polymorph Form II is $176\text{--}178^\circ\text{C}$.

The present invention also provides a method of preparing the polymorph comprising the steps of heating the polymorph of Form I to about 165°C under N_2 , and cooling down to room temperature naturally to recover the resulted Polymorph Form II.

In another aspect, the polymorph form is of Polymorph Form III and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 5.1° , 16.7° , 20.9° , 21.7° and $23.9^\circ \pm 0.2^\circ$.

In some other embodiments of Polymorph Form III, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $17.3\text{\AA}$, $5.3\text{\AA}$, $4.3\text{\AA}$, $4.1\text{\AA}$ and $3.7\text{\AA}$.

In some embodiments of Polymorph Form III, its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 5.1° , 7.6° , 16.7° , 19.8° , 20.9° , 21.7° , 23.9° and $24.7^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form III, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $17.3\text{\AA}$, $11.7\text{\AA}$, $5.3\text{\AA}$, $4.5\text{\AA}$, $4.3\text{\AA}$, $4.1\text{\AA}$, $3.7\text{\AA}$ and $3.6\text{\AA}$.

Preferably, the X-ray powder diffraction pattern is shown as in Figure 3.

Preferably, the melting point of the Polymorph Form III is $175\text{--}178^\circ\text{C}$.

The present also provides a method of preparing the polymorph, comprising the steps of putting the polymorph of Form I in the saturated vapor of acetonitrile at room temperature for a week, and recovering the resulted Polymorph Form III.

In another aspect, the polymorph form is of Polymorph Form IV and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 6.1° , 11.2° , 15.8° , 19.5° and $22.5^\circ \pm 0.2^\circ$.

In some other embodiments of Polymorph Form IV, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $14.5\text{\AA}$, $7.9\text{\AA}$, $5.6\text{\AA}$, $4.6\text{\AA}$ and $4.0\text{\AA}$.

In some embodiments of Polymorph Form IV, its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of approximately 6.1° , 11.2° , 12.9° , 14.2° , 15.8° , 18.3° , 19.5° and $22.5^\circ \pm 0.2^\circ$.

In still some other embodiments of Polymorph Form IV, its X-ray powder diffraction pattern has characteristic peaks, expressed in terms of the interplanar distance, of $14.5\text{\AA}$, $7.9\text{\AA}$, $6.9\text{\AA}$, $6.2\text{\AA}$, $5.6\text{\AA}$, $4.9\text{\AA}$, $4.6\text{\AA}$ and $4.0\text{\AA}$.

Preferably, the X-ray powder diffraction pattern is shown as in Figure 4.

Preferably, the melting point of the Polymorph Form IV is $175-177^\circ\text{C}$.

The present invention also provides a method of preparing the polymorph comprising the steps of putting the polymorph of Form I in the saturated vapor of methanol at room temperature for a week, and recovering the resulted Polymorph Form IV.

Polymorph Form I, II, III or IV of this invention can have a purity of $\geq 85\%$, $\geq 95\%$, or even $\geq 99\%$.

In still another aspect, the invention provides processes for preparing a polymorph form of Icotinib, comprising the step of reacting Icotinib hydrochloride with a base in a media to produce Icotinib in the polymorph form. This reaction can be carried out, for example, at the temperature of $40-80^\circ\text{C}$.

Examples of a suitable base include sodium hydroxide, potassium hydroxide, sodium carbonate, and potassium carbonate.

The second reaction media can include, e.g., water and ethanol, water and isopropanol, water and methanol, or water and tetrahydrofuran.

The polymorph forms of this invention, particularly Polymorph Form I, Polymorph Form II, Polymorph Form III and Polymorph Form IV, unexpectedly exhibited significantly better bioavailability and chemical stability than Icotinib hydrochloride, the active ingredient of a drug currently in the marketplace. Accordingly, the invention further provides pharmaceutical compositions comprising a therapeutically effective amount of a polymorph form of this invention and a pharmaceutically acceptable excipient, adjuvant or carrier.

The pharmaceutical composition can further include a second therapeutically active ingredient, and can be in a form suitable for oral administration (e.g., a tablet or capsule).

Also within the scope of this invention is the use of a polymorph form or a pharmaceutical composition of this invention in the manufacturing of a medicament for the prevention or treatment in mammals of an excessive non-malignant disease, pancreatitis, kidney disease, cancer, angiogenesis or vascular disease, or for mammalian embryo cell transplantation.

The excessive non-malignant disease can be, for example, benign skin hyperplasia or benign prostatic hyperplasia. Or, the excessive non-malignant disease, pancreatitis, kidney disease, cancer, angiogenesis or vascular occurrence-related illness is selected from: tumor angiogenesis, chronic inflammatory disease such as rheumatoid arthritis, atherosclerosis, atherosclerosis, skin diseases such as psoriasis, and scleroderma, diabetes-induced skin diseases, diabetic retinopathy, premature retinopathy, age-related degeneration stains, hemangioma, glioma, Kaposi's sarcoma, ovarian cancer, breast cancer, lung cancer, pancreatic cancer, lymphoma, prostate, colon and skin tumors and their complications.

All the polymorphs of the present invention are approximately pure.

The term “approximately pure” as herein used refers to at least 85wt%, preferably at least 95wt%, more preferably at least 99wt% of the compound of Formula I exists in a polymorphic form of the present invention, particularly in the polymorphic forms of Form I, Form II, Form III or Form IV.

The main peaks described in the polymorphs above are reproducible and are within the error limit (the specified value ± 0.2).

In the present invention, “the X-ray powder diffraction pattern is shown as in Figure 1” refers to the X-ray powder diffraction pattern that show major peaks as in Figure 1, wherein major peaks refer to those with the relative intensity greater than 10%, preferably greater than

30%, relative to the highest peak (with its relative intensity designated to be 100%) in Figure 1. Likewise, in the present invention, the X-ray powder diffraction pattern shown as in Figure 2, 3 or 4 refers to the X-ray powder diffraction pattern that show major peaks as in Figure 2, 3 or 4 wherein major peaks refer to those with the relative intensity greater than 10%, preferably greater than 30%, relative to the highest peak (with its relative intensity designated to be 100%) in Figure 2, 3 or 4 respectively.

Crystallization in the present invention is related to dynamics and equilibria among different polymorphic forms under certain conditions. Therefore, those skilled in the art will realize that the resulting polymorphic form depends on the kinetics and thermodynamics of the crystallization process. Under certain conditions (solvent system, temperature, pressure, and the concentration of the compound of the present invention), a polymorphic form may be more stable than another one (or, actually be more stable than any other polymorphic forms). However, the polymorphs that are less stable thermodynamically may be favorable in kinetics. The polymorph form may also be affected by factors other than kinetics, such as time, impurity distribution, agitation, presence or absence of polymorphic seed. For purposes of this invention, various hydrate and solvate forms are included in the scope of "polymorph".

The present invention further provides a pharmaceutical composition, comprising a therapeutically effective amount of one or more polymorphs of Form I, Form II, Form III, or Form IV of the compound of Formula I, and a pharmaceutically acceptable excipient, adjuvant or carrier. Wherein, the pharmaceutical composition contains 0.01wt%-99wt%, preferably 1wt%-70wt%, more preferably 10wt%-50wt% of any one polymorph of Form I, Form II, Form III, or Form IV of the compound of Formula I.

The term "therapeutically effective amount" as herein used, refers to the amount of a compound that, when administered to a subject for treating a disease, or at least one of the clinical symptoms of a disease or disorder, is sufficient to affect such treatment for the disease, disorder, or symptom. The "therapeutically effective amount" can vary with the compound, the disease, disorder, and/or symptoms of the disease or disorder, severity of the disease, disorder, and/or symptoms of the disease or disorder, the age of the subject to be treated, and/or the weight of the subject to be treated. An appropriate amount in any given instance can be apparent to those skilled in the art or can be determined by routine experiments. In the case of combination

therapy, the "therapeutically effective amount" refers to the total amount of the combination objects for the effective treatment of a disease, a disorder or a condition.

The pharmaceutical composition comprising the compound of the present invention can be administrated via oral, inhalation, rectal, parenteral or topical administration to a subject who
5 needs treatment. For oral administration, the pharmaceutical composition may be a regular solid formulation such as tablets, powder, granule, capsules and the like, a liquid formulation such as water or oil suspension or other liquid formulation such as syrup, solution, suspension or the like; for parenteral administration, the pharmaceutical composition may be solution, water solution, oil suspension concentrate, lyophilized powder or the like. Preferably, the formulation
10 of the pharmaceutical composition is selected from tablet, coated tablet, capsule, suppository, nasal spray or injection, more preferably tablet or capsule. The pharmaceutical composition can be a single unit administration with an accurate dosage. In addition, the pharmaceutical composition may further comprise additional active ingredients.

All formulations of the pharmaceutical composition of the present invention can be
15 produced by the conventional methods in the pharmaceutical field. For example, the active ingredient can be mixed with one or more excipients, then to make the desired formulation. The "pharmaceutically acceptable carrier" refers to conventional pharmaceutical carriers suitable for the desired pharmaceutical formulation, for example: a diluent, a vehicle such as water, various organic solvents, etc, a filler such as starch, sucrose, etc; a binder such as cellulose
20 derivatives, alginates, gelatin and polyvinylpyrrolidone (PVP) ; a wetting agent such as glycerol; a disintegrating agent such as agar, calcium carbonate and sodium bicarbonate; an absorption enhancer such as quaternary ammonium compound; a surfactant such as hexadecanol; an absorption carrier such as Kaolin and soap clay; a lubricant such as talc, calcium stearate, magnesium stearate, polyethylene glycol, etc. In addition, the pharmaceutical composition
25 further comprises other pharmaceutically acceptable excipients such as a decentralized agent, a stabilizer, a thickener, a complexing agent, a buffering agent, a permeation enhancer, a polymer, aromatics, a sweetener, and a dye. Preferably, the excipient is suitable for desired formulation and administration type.

The term "disease" or "disorder" or "condition" refers to any disease, discomfort, illness,
30 symptoms or indications.

In another aspect, the present invention provides use of the compounds (Icotinib and its polymorphs) and/or the pharmaceutical compositions in manufacturing a medicament for the treatment or prevention in mammals of excessive non-malignant hyperplasia disease, pancreatitis, kidney disease, cancer, angiogenesis or vascular occurrence-related illness, or for the mammalian embryo cell transplantation. The excessive non-malignant hyperplasia disease can be a benign skin hyperplasia or a benign prostatic hyperplasia.

Preferably, the polymorphs or pharmaceutical composition of the present invention can be used in manufacturing a medicament for the treatment or prevention in mammals of excessive non-malignant hyperplasia disease, pancreatitis, kidney disease, cancer, angiogenesis or vascular occurrence-related illness selected from tumor angiogenesis, chronic inflammatory diseases such as rheumatoid arthritis, atherosclerosis, skin diseases such as psoriasis, and scleroderma, diabetes included skin diseases, diabetic retinopathy, premature retinopathy, age-related degeneration stains, hemangioma, glioma, Kaposi internal tumor, ovarian cancer, breast cancer, lung cancer, pancreatic cancer, lymphoma, prostate, colon and skin tumors and their complications.

Among the mammals mentioned herein, human beings are preferred.

This invention provides a method for treating malignant tissue hyperplasia in mammals. This treatment method includes application of an effective amount of Icotinib and/or its polymorph forms and/or the pharmaceutical compositions to mammalian patients with hyperplasia disease. In some embodiments, the treatment method also includes use of MMP (matrix metalloproteinase) inhibitor, VEGFR (vascular endothelial growth factor receptor) kinase inhibitors, HER2 inhibitor, VEGFR antibody drugs, and/or endostatin drugs. In some embodiments, the treatment method also includes using one or more anti-tumor agents such as mitotic inhibitors, alkylating agents, anti-metabolites, tumor antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, enzyme inhibitors, biological response modifiers, anti-hormone drugs and so on. The anti-tumor agents can be selected from carboplatin, paclitaxel, gemcitabine, methotrexate, 5-FU, camptothecin, cyclophosphamide, BCNU and other medications.

Another purpose of this invention is to provide the use of the compounds (Icotinib and its polymorphs) and/or the pharmaceutical compositions in manufacturing a medicament for treating a disease related to tyrosine kinase dysfunction.

Preferably, this invention provides a method for the treatment of disease caused by tyrosine kinase dysfunction. This treatment method includes administering to a patient with the disease caused by tyrosine kinase dysfunction an effective amount of the compounds (Icotinib and its polymorphs) and/or the pharmaceutical compositions. The tyrosine kinase dysfunction-related disease include, but are not limited to disease of brain, lung, liver, bladder, breast, head and neck, esophagus, gastrointestinal tract, ovary, cervix or thyroid tumors and their complications.

Target disease for the just described treatment method are preferably selected from brain cancer, lung cancer (such as non-small cell lung cancer (NSCLC)), kidney cancer, bone cancer, liver cancer, bladder cancer, chest cancer, lymphoma, or thyroid tumors and their complications.

The above-described methods can be applied in combination with any chemical therapy, biological therapy or radiation therapy.

The above-described treatment methods can further include application of anti-EGFR antibodies, anti-EGF antibodies, or both, in the same treatment.

The dosage of the active ingredient or compound when administered will be determined by the individual needs of the patient to treated, administration route, severity of disease or illness, dosing schedule, as well as evaluation and judgment of the designated doctor. However, based on the active compound, the preferred range of the effective dosage can be approximately 0.01-120mg daily per kilogram of body weight; or more preferably 1-50mg per day per kilogram of body weight in single or separate doses. In some cases, it is more suitable to apply the lower end of the above-described dosage range, while in other cases the higher dosage may be used without causing harmful side effects.

Another aspect of the present invention is to provide Icotinib for clinical applications. In particular, the present invention related to clinical treatment with Icotinib with the following treatment options for cancer patients: the dosage of Icotinib and/or polymorph Form I , II , III, or IV can be 25-2100mg/day with the administration frequency being 1-3 times a day; a preferred dosage is 75-1200mg/day with the administration frequency being 2-3 times a day; an even more preferred dosage is 100-1200mg/day with the administration frequency being 2-3 times a day.

DESCRIPTIONS OF THE FIGURES

Figure 1: The X-ray powder diffraction pattern of Polymorph Form I of compound of Formula I.

Figure 2: The X-ray powder diffraction pattern of Polymorph Form II of compound of Formula I.

Figure 3: The X-ray powder diffraction pattern of Polymorph Form III of compound of Formula I.

Figure 4: The X-ray powder diffraction pattern of Polymorph Form IV of compound of Formula I.

Figure 5: The plasma concentration-time curves of Crystalline Form I Icotinib hydrochloride and Polymorph Form II of the compound of Formula I.


EXAMPLES

The present invention is further exemplified, but not limited, by the following examples that illustrate the invention. The techniques or methods used in these examples, unless expressly stated otherwise, were conventional techniques or methods well known in the art.

The X-ray powder diffraction (XRPD) patterns for the crystalline forms of Icotinib were generated on a PANalytical X-ray Diffraction System with Empyrean console. The diffraction peak positions were calibrated using silicon powder which had a 2θ value of 28.443 degree. An Empyrean Cu LEF X-ray tube K-Alpha radiation was used as the source.

Example 1

Preparation of Form I




100g Icotinib hydrochloride was dissolved in the mixture of 300ml ethanol and 200ml water. A solution of 11.2g sodium hydroxide in 100ml water was added dropwisely at 60°C to the Icotinib hydrochloride solution until the pH value of the reaction solution reached 13. The reaction solution was then stirred for an hour and then cooled down to the room temperature.

The precipitate was filtered and washed with purified water and dried for 8 hours under vacuum below 60°C to obtain 90 g the desired Form I . And the melting point of the Polymorph Form I is 175-177°C.

Example 2


5 Preparation of Form I




10g Icotinib hydrochloride was first dissolved in the mixture of 30ml isopropanol and 20ml water, a solution of 1.6g potassium hydroxide in 10 ml water was added to the Icotinib hydrochloride solution until the pH value of the reaction mixture reached 13. The reaction solution was then stirred for 1-2hrs and cooling down to the room temperature. The precipitate was filtered and washed with purified water and dried under vacuum for 8-10 hours at a temperature below 50°C to give 7.9g the desired Form I .

Example 3

Preparation of Form I



5g Icotinib hydrochloride was dissolved in the mixture of 20ml methanol and 15ml water. To this Icotinib hydrochloride solution was then added dropwisely a solution of 1.5g sodium carbonate in 10ml water at 40°C until the pH value of the resultant mixture reached 13. The reaction solution was then stirred for 1-2hrs before cooling down to the room temperature. The precipitate was filtered and washed with purified water and then dried under vacuum for 8-10hrs below 60°C to obtain 4g the desired Form I .

Example 4**Preparation of Form I**

5g Icotinib hydrochloride was dissolved in a mixture of 20ml tetrahydrofuran and 15ml water, and then to this Icotinib solution was added a solution of 1.9g potassium carbonate in 10ml water dropped at 50°C until the pH value of resultant reaction mixture reached 13. The reaction solution was then stirred for 1-2hrs before cooling down to the room temperature to give a precipitate. The precipitate was filtered and washed with purified water and then dried for 8-10hrs under vacuum at a temperature below 60°C to give 4g the desired Form I.

Example 5**Preparation of Form II**

Form I prepared from the method of Example 1 was heated to 150°C under N₂, then cooled down to room temperature naturally to obtain the desired Form II. And the melting point of the Polymorph Form is II 176-178°C.

Example 6**Preparation of Form II**

Form I prepared from the method of Example 1 was heated to 160°C under N₂, then cooled down to room temperature naturally to obtain the desired Form II.

Example 7**Preparation of Form II**

Form I prepared from the method of Example 1 was heated to 170°C under N₂, then cooled down to room temperature naturally to obtain the desired Form II.

Example 8**Preparation of Form II**

Form I prepared from the method of Example 1 was heated to 180°C under N₂, then

cooled down to room temperature naturally to obtain the desired Form II.

Example 9

Preparation of Form II

Form I prepared from the method of Example 1 was heated to 165°C under N₂, then
5 cooled down to room temperature naturally to obtain the desired Form II.

Example 10

Preparation of Form III

15mg Form I prepared from the method of Example 1 was put into a 3-ml vial. This vial
was placed inside a 20ml vial filled with saturated vapor of acetonitrile. The large vial was
10 sealed and kept at room temperature for one week, recovered the desired polymorph. And the
melting point of the Polymorph Form III is 175-178°C.

Example 11

Preparation of Form IV

15mg Form I prepared from the method of Example 1 was put into a 3-ml vial. This vial
15 was placed inside a 20ml vial filled with saturated vapor of methanol. The larger vial was
sealed and kept at room temperature for one week, and recovered the desired polymorph. And
the melting point of the Polymorph Form is IV 175-177°C.

Example 12

Pharmacokinetic study of Icotinib Hydrochloride and the Polymorph Form II of 20 Icotinib

Drugs and reagents: the Icotinib hydrochloride used in this study was of Crystalline Form
I disclosed by the WO2010/003313. Polymorph Form II of Icotinib and Icotinib
hydrochloride were ground to fine particles. The material content (purity) was not less than
99.0%. Sodium carboxymethyl cellulose was medical supply graded.

25 Experimental animals: SD rats were divided to the Icotinib hydrochloride group and a
Polymorph Form II group.

Pharmaceutical preparation: the amount of each compound was weighed and then sodium
carboxymethyl cellulose was added to result in the test compound's concentration of 0.5%. The

solid mixture was then added to prepare a suspension thereof at a final concentration of 10mg/ml in water.

Administration and sample collection: each suspension was administered orally to fasted SD rats at a dose equivalent to 50 mg/kg Icotinib in a dose volume of 5ml/kg. 0.4ml of blood was collected in EDTA-K pre anticoagulant tubes at time intervals of 0.5, 1, 1.5, 2, 4, 6, 8 and 24 hours after the administration of the test compound, centrifuged at 3000 rpm for 10 minutes, and 120µl plasma was collected and kept in cold storage.

Samples were analyzed by high performance liquid chromatography. The chromatographic conditions utilized C18-silane bonded silica as stationary phase, 0.02mol/L of sodium dihydrogen phosphate in acetonitrile (40:60, using sodium hydroxide solution to adjust pH to 5.0) as the mobile phase and a detection wavelength of 334nm. PK profile comparison of Polymorph Form II of Icotinib and Crystalline Form I of Icotinib hydrochloride was summarized in Table 1 and Figure 5. Polymorph Form II of Icotinib showed higher bioavailability than Crystalline Form I of Icotinib hydrochloride.

Table 1

Group	Gender	AUC ₍₀₋₂₄₎ (mg/L*h)	AUC _(0-∞) (mg/L*h)	T _{1/2} (h)	T _{max} (h)	C _{max} (mg/L)
Crystalline Form I of Icotinib hydrochloride	Male	39.2±27.6	39.8±28.5	3.1±1.4	1.3±1.1	4.9±2.6
Polymorph Form II of Icotinib	Male	62.4±30.1	63.0±30.0	3.5±0.6	4.5±2.6	5.8±2.6

Example 13


Formulation of a Hard Gel Capsule

As a specific embodiment of an oral composition, about 100mg of the Polymorph Form of Example 1-11 is formulated with sufficient finely divided lactose to provide a total amount of about 580mg to about 590mg to fill a size 0 hard gel capsule.

Although the present invention has been fully described in connection with embodiments thereof with reference to the accompanying drawings, it is to be noted that various change and modifications will become apparent to those skilled in the art. Such changes and modifications are to be understood as being included within the scope of the present invention as defined by the appended Claims.

WHAT IS CLAIMED IS

1. A polymorph form of the compound of Formula I,



Formula I


wherein the polymorph form is of Form I and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 7.6° , 14.7° , 20.9° , 23.1° and $27.5^\circ \pm 0.2^\circ$.

2. The polymorph form of Claim 1, wherein the X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 7.6° , 14.7° , 20.9° , 23.1° , 24.2° , 27.5° , 29.8° and $30.9^\circ \pm 0.2^\circ$.

3. The polymorph form of either Claim 1 or Claim 2, wherein the X-ray powder diffraction pattern is as shown in Figure 1.

4. The polymorph form of any one of Claims 1 to 3, wherein the melting point is $175-177^\circ\text{C}$.

5. A polymorph form of the compound of Formula I,




Formula I

wherein the polymorph form is of Form II and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 5.5° , 11.0° , 19.7° , 20.9° and $22.6^\circ \pm 0.2^\circ$.

6. The polymorph form of Claim 5, wherein its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 5.5° , 8.8° , 11.0° , 16.5° , 19.7° , 20.9° , 22.6° and $23.7^\circ \pm 0.2^\circ$.

7. The polymorph form of either Claim 5 or Claim 6, wherein the X-ray powder diffraction pattern is as shown in Figure 2.

8. The polymorph form of any one of Claims 5 to 7, wherein the melting point is 176-178°C.
9. A polymorph form of the compound of Formula I,



Formula I


wherein the polymorph form is of Form III and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 5.1°, 16.7°, 20.9°, 21.7° and 23.9°±0.2°.

10. The polymorph form of Claim 9, wherein its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 5.1°, 7.6°, 16.7°, 19.8°, 20.9°, 21.7°, 23.9° and 24.7°±0.2°.

11. The polymorph form of either Claim 9 or Claim 10, wherein the X-ray powder diffraction pattern is as shown in Figure 3.

12. The polymorph form of any one of Claims 9 to 11, wherein the melting point is 175-178°C.

13. A polymorph form of the compound of Formula I,



Formula I

wherein the polymorph form is of Form IV and its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 6.1°, 11.2°, 15.8°, 19.5° and 22.5°±0.2°.

14. The polymorph form of Claim 13, wherein its X-ray powder diffraction pattern has characteristic peaks at diffraction angles 2θ of 6.1°, 11.2°, 12.9°, 14.2°, 15.8°, 18.3°, 19.5° and 22.5°±0.2°.

15. The polymorph form of either Claim 13 or Claim 14, wherein the X-ray powder diffraction pattern is as shown in Figure 4.

16. The polymorph form of any one of Claims 13 to 15, wherein the melting point is

175-177°C.

17. A method of preparing a polymorph form of Claims 1 to 16, comprising:

a) dissolving Icotinib hydrochloride in a mixture of ethanol and water, warming the mixture to 40-60°C and adding a solution of base; then stirring the solution for 1hr, cooling down to room temperature and recovering the resultant polymorph form of any one of Claims 1 to 4; or

b) heating the polymorph form of any one of Claims 1 to 4 to a temperature of 150-180°C under N₂, then cooling down to room temperature naturally to obtain the resultant polymorph form of any one of Claims 5 to 8; or

c) placing the polymorph form of any one of Claims 1 to 4 in a saturated vapor of acetonitrile at room temperature, and recovering the resultant polymorph form of any one of Claims 9 to 12; or

d) placing the polymorph form of any one of Claims 1 to 4 in a saturated vapor of methanol at room temperature, and recovering the resultant polymorph form of any one of Claims 13 to 16.

18. The method of Claim 17, wherein the temperature of b) is 160-170°C.

19. The method of Claim 18, wherein the temperature is 165°C.

20. A pharmaceutical composition comprising a therapeutically effective amount of the polymorph of any one of Claims 1 to 16, and a pharmaceutically acceptable excipient, adjuvant or carrier.

21. The pharmaceutical composition of Claim 20, wherein the polymorph form of any one of Claims 1 to 16 has a purity of $\geq 85\text{wt}\%$.

22. The pharmaceutical composition of Claim 20, wherein the polymorph form of any one of Claims 1 to 16 has a purity of $\geq 99\text{wt}\%$.

23. The pharmaceutical composition of any one of Claims 20 to 22, further comprising a second therapeutically active ingredient.

24. The pharmaceutical composition of any one of Claims 20 to 23, wherein the pharmaceutical composition is suitable for oral administration.

25. The pharmaceutical composition according to any one of Claims 20 to 24, wherein the pharmaceutical composition is in the form of tablet or capsule.

26. The pharmaceutical composition of any one of Claims 20 to 25, wherein the composition comprises 0.01wt% to 99wt% of the polymorph form of any one of Claims 1 to 16.

27. The pharmaceutical composition of Claim 26, wherein the composition comprises 10wt%

to 50wt% of the polymorph form of any one of Claims 1 to 16.

28. Use of a polymorph form of any one of Claims 1 to 16 or the pharmaceutical composition of any one of Claims 20 to 27 in the manufacture of a medicament for treating a disease related to EGFR-mediated tyrosine kinase dysfunction.

29. The use of Claim 28, wherein the disease related to EGFR-mediated tyrosine kinase dysfunction is brain, lung, liver, bladder, chest, head and neck, esophagus, gastrointestinal tract, breast, ovary, cervix or thyroid tumors or their complications.

30. Use of a polymorph form of any one of Claims 1 to 16 or the pharmaceutical composition of any one of Claims 20 to 27 in the manufacture of a medicament for the prevention or treatment of cancer associated with EGFR-mediated tyrosine kinase dysfunction in mammals.

31. The use of Claim 30, wherein the cancer is selected from: hemangionma, glioma, Kaposiinternal tumor, ovarian cancer, breast cancer, lung cancer, pancreatic cancer, lymphoma, prostate, colon and skin tumors and their complications.