

ANALGESIC COMPOUNDS

INCORPORATION BY REFERENCE TO ANY PRIORITY APPLICATIONS

[0001] Any and all applications for which a foreign or domestic priority claim is identified, for example, in the Application Data Sheet or Request as filed with the present application, are hereby incorporated by reference under 37 CFR 1.57, and Rules 4.18 and 20.6, such as U.S. Provisional Application No. 62/506,512, filed May 15, 2017 is hereby incorporated by reference in its entirety.

BACKGROUND

Field

[0002] The present application relates to the fields of chemistry, biochemistry and medicine. More particularly, disclosed herein are substituted bicyclo[1.1.1]pentyl compounds. Also disclosed herein are methods of using substituted bicyclo[1.1.1]pentyl compounds as an analgesic.

Description

[0003] Nonsteroidal anti-inflammatory compounds, or NSAIDs, are an extremely useful group of small molecule drugs, typified by acetylsalicylic acid, ibuprofen and naproxen. Acetaminophen, also known as paracetamol or APAP, is also an effective pain reliever often sold over the counter (without prescription). Opioids is another class of drugs used to treat pain, such as short-term pain. Each of these classes of drugs is associated with one or more undesirable side effects.

SUMMARY

[0004] Some embodiments described herein generally relate to a compound selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing. Other embodiments described herein generally relate to a pharmaceutical composition that include

one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing.

[0005] Some embodiments described herein generally relate to a method of ameliorating and/or treating pain that can include administering to a subject suffering from pain an effective amount of one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, or a pharmaceutical composition that includes one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing. Other embodiments described herein generally relate to using one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, or a pharmaceutical composition that includes one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, in the manufacture of a medicament for ameliorating and/or treating pain. Still other embodiments described herein generally relate to a compound selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, or a pharmaceutical composition that includes one or more compounds selected from Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, that can be used for ameliorating and/or treating pain.

BRIEF DESCRIPTION OF THE DRAWINGS


[0006] FIG. 1 graphically depicts the results of a conditioned place preference (CPP) test performed on mice comparing the administration of morphine to some compounds described herein.

DETAILED DESCRIPTION

[0007] NSAIDs are often sold without prescription, and are variously used to treat pain, inflammation, and fever. However, NSAIDs can have undesirable side effects, including gastric upset and/or gastric bleeding. Some of these undesirable side effects of NSAIDs are known to be associated with inhibition of the COX-1 and/or COX-2 receptors. (See Suleyman et al. “Anti-inflammatory and side effects of cyclo-oxygenase inhibitors.” *Pharmacological reports* 59.3 (2007): 247). Furthermore, both acetaminophen and NSAIDs are known to only be mildly effective in treating neuropathic pain. (See McPherson, “Neuropathic pain: an update on effective management strategies.” *American Pharmacists Association 2006 Annual Meeting*, 2006).

[0008] Acetaminophen shares analgesic and antipyretic properties with NSAIDs. However, APAP has only weak anti-inflammatory properties, and is thus not an NSAID. Unlike many NSAIDs, acetaminophen does not cause gastric upset or bleeding in prescribed doses. Thus, APAP is an extremely useful drug for those wishing analgesia without adverse gastric side effects.

[0009] Acetaminophen has the structure:



[0010] Acetaminophen is often combined with other drugs for relief of symptoms of influenza and the common cold, among other indications. It is particularly useful in combination with opioid analgesics, where it exhibits synergistic analgesic properties and allows patients to achieve adequate pain relief with lower doses of opioids. The most widely prescribed drug in the United States is a combination of acetaminophen and hydrocodone, with over 130 million prescriptions in the year 2010. Other acetaminophen-opioid combinations, including combinations with oxycodone, are also widely prescribed.

[0011] Acetaminophen poisoning is the most common cause of acute liver failure in the Western world, and acetaminophen accounts for the most drug overdoses in the English-speaking world. Acetaminophen is metabolized to form N-acetyl-p-benzoquinoneimine (NAPQI), which depletes glutathione in the liver, and if the glutathione is sufficiently depleted, as is the case with an acetaminophen overdose, the NAPQI

metabolite injures hepatocytes leading to acute liver failure and often death. The acetaminophen-opioid combination drugs are commonly implicated in such toxicity, for various reasons. First, patients might not recognize that the prescribed pain relievers contain acetaminophen, and may supplement with additional acetaminophen if pain relief is inadequate. Second, continued administration of opioids can lead to tolerance and the need for increased dosages to obtain a comparable opioid analgesic effect, and users or abusers of the combination drugs may exceed safe dosages of acetaminophen as a consequence.

[0012] This has led the U.S. FDA to seek reduced amounts of acetaminophen in the opioid combination drugs and has also led an FDA advisory panel to recommend banning such drugs all together. Although the acetaminophen-opioid drugs remain on the market, there is a strong need for a less toxic replacement without the same hepatotoxicity risks.

[0013] Opioids are generally known to be associated with a number of common side effects and long term negative consequences. Common side effects of opioid administration may include, for example, sedation, dizziness, nausea, vomiting, constipation, physical dependence, tolerance and respiratory depression; and long term negative consequences of opioids may include, for example, psychological addiction and abuse. (*See* Benyamin, et al. "Opioid Complications and Side Effects" *Pain Physician* 11 (2008): S105-S120). Some of these side effects of opioids are known to be associated with the activation of an opiate receptor, such as opiate receptors δ_1 , δ_2 , κ , and/or μ .

[0014] With the side effects and dangers associated with NSAIDs, acetaminophen and opioids, there is a desire in the medical community for compounds with comparable or improved analgesic properties that minimize and/or avoid one or more of the undesirable effects associated with NSAIDs, acetaminophen and/or opioids.


Definitions

[0015] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art. All patents, applications, published applications and other publications referenced herein are incorporated by reference in their entirety unless stated otherwise. In the event that there are a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

[0016] Whenever a group is described as being “optionally substituted” that group may be unsubstituted or substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) may be selected from one or more of the indicated substituents. If no substituents are indicated, it is meant that the indicated “optionally substituted” or “substituted” group may be substituted with one or more group(s) individually and independently selected from deuterium (D), halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₈ alkyl, C₃₋₂₀ cycloalkyl, aryl, heteroaryl, heterocyclyl, C₁₋₆ haloalkyl, cyano, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₃₋₂₀ cycloalkenyl, aryl(alkyl), heteroaryl(alkyl), heterocyclyl(alkyl), acyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, C-thioamido, N-thioamido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, sulfenyl, sulfinyl, sulfonyl, haloalkoxy, an amino, a mono-substituted amine group and a di-substituted amine group.

[0017] As used herein, “C_a to C_b” in which “a” and “b” are integers refer to the number of carbon atoms in a group. The indicated group can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C₁ to C₄ alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, CH₃-, CH₃CH₂-, CH₃CH₂CH₂-, (CH₃)₂CH-, CH₃CH₂CH₂CH₂-, CH₃CH₂CH(CH₃)- and (CH₃)₃C-. If no “a” and “b” are designated, the broadest range described in these definitions is to be assumed.

[0018] If two “R” groups are described as being “taken together” the R groups and the atoms they are attached to can form a cycloalkyl, cycloalkenyl, aryl, heteroaryl or heterocycle. For example, without limitation, if R^a and R^b of an NR^aR^b group are indicated to be “taken together,” it means that they are covalently bonded, either indirectly through intermediate atoms, or directly to one another, to form a ring, for example:



[0019] As used herein, the term “alkyl” refers to a fully saturated aliphatic hydrocarbon group. The alkyl moiety may be branched or straight chain. Examples of branched alkyl groups include, but are not limited to, iso-propyl, sec-butyl, t-butyl and the like. Examples of straight chain alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl and the like. The alkyl group may have 1 to 30

carbon atoms (whenever it appears herein, a numerical range such as “1 to 30” refers to each integer in the given range; *e.g.*, “1 to 30 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 30 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium alkyl having 1 to 12 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 6 carbon atoms. An alkyl group may be substituted or unsubstituted.



[0020] The term “alkenyl” used herein refers to a monovalent straight or branched chain radical of from two to thirty carbon atoms containing a carbon double bond(s) including, but not limited to, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like. An alkenyl group may be unsubstituted or substituted.

[0021] The term “alkynyl” used herein refers to a monovalent straight or branched chain radical of from two to thirty carbon atoms containing a carbon triple bond(s) including, but not limited to, 1-propynyl, 1-butyne, 2-butyne and the like. An alkynyl group may be unsubstituted or substituted.

[0022] As used herein, “cycloalkyl” refers to a completely saturated (no double or triple bonds) mono- or multi- cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. As used herein, the term “fused” refers to two rings which have two atoms and one bond in common.

For example, in the following structure, rings A and B are fused . As used herein, the term “bridged cycloalkyl” refers to compounds wherein the cycloalkyl contains a linkage

of one or more atoms connecting non-adjacent atoms. The following structures 

 and  are examples of “bridged” rings. As used herein, the term “spiro” refers to two rings which have one atom in common and the two rings are not linked by a bridge. Cycloalkyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to

10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). A cycloalkyl group may be unsubstituted or substituted. Examples of mono-cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. Examples of fused cycloalkyl groups are decahydronaphthalenyl, dodecahydro-1H-phenalenyl and tetradecahydroanthracenyl; examples of bridged cycloalkyl groups are bicyclo[1.1.1]pentyl, adamantanyl, and norbornanyl; and examples of spiro cycloalkyl groups include spiro[3.3]heptane and spiro[4.5]decane.

[0023] As used herein, “cycloalkenyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more double bonds in at least one ring; although, if there is more than one, the double bonds cannot form a fully delocalized pi-electron system throughout all the rings (otherwise the group would be “aryl,” as defined herein). Cycloalkenyl groups can contain 3 to 30 atoms in the ring(s), 3 to 20 atoms in the ring(s), 3 to 10 atoms in the ring(s), 3 to 8 atoms in the ring(s) or 3 to 6 atoms in the ring(s). When composed of two or more rings, the rings may be connected together in a fused, bridged or spiro fashion. A cycloalkenyl group may be unsubstituted or substituted.

[0024] As used herein, “cycloalkynyl” refers to a mono- or multi- cyclic hydrocarbon ring system that contains one or more triple bonds in at least one ring. If there is more than one triple bond, the triple bonds cannot form a fully delocalized pi-electron system throughout all the rings. Cycloalkynyl groups can contain 8 to 30 atoms in the ring(s), 8 to 20 atoms in the ring(s) or 8 to 10 atoms in the ring(s). When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro fashion. A cycloalkynyl group may be unsubstituted or substituted.

[0025] As used herein, “aryl” refers to a carbocyclic (all carbon) monocyclic or multicyclic aromatic ring system (including fused ring systems where two carbocyclic rings share a chemical bond) that has a fully delocalized pi-electron system throughout all the rings. The number of carbon atoms in an aryl group can vary. For example, the aryl group can be a C₆-C₁₄ aryl group, a C₆-C₁₀ aryl group, or a C₆ aryl group. Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted.

[0026] As used herein, “heteroaryl” refers to a monocyclic or multicyclic aromatic ring system (a ring system with fully delocalized pi-electron system) that contain(s) one or more heteroatoms (for example, 1, 2 or 3 heteroatoms), that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur. The number of atoms in the ring(s) of a heteroaryl group can vary. For example, the heteroaryl group can contain 4 to 14 atoms in the ring(s), 5 to 10 atoms in the ring(s) or 5 to 6 atoms in the ring(s). Furthermore, the term “heteroaryl” includes fused ring systems where two rings, such as at least one aryl ring and at least one heteroaryl ring, or at least two heteroaryl rings, share at least one chemical bond. Examples of heteroaryl rings include, but are not limited to, furan, furazan, thiophene, benzothiophene, phthalazine, pyrrole, oxazole, benzoxazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, thiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, benzothiazole, imidazole, benzimidazole, indole, indazole, pyrazole, benzopyrazole, isoxazole, benzoisoxazole, isothiazole, triazole, benzotriazole, thiadiazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, purine, pteridine, quinoline, isoquinoline, quinazoline, quinoxaline, cinnoline and triazine. A heteroaryl group may be substituted or unsubstituted.

[0027] As used herein, “heterocyclyl” or “heteroalicyclyl” refers to three-, four-, five-, six-, seven-, eight-, nine-, ten-, up to 18-membered monocyclic, bicyclic and tricyclic ring system wherein carbon atoms together with from 1 to 5 heteroatoms constitute said ring system. A heterocycle may optionally contain one or more unsaturated bonds situated in such a way, however, that a fully delocalized pi-electron system does not occur throughout all the rings. The heteroatom(s) is an element other than carbon including, but not limited to, oxygen, sulfur and nitrogen. A heterocycle may further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo-systems and thio-systems such as lactams, lactones, cyclic imides, cyclic thioimides and cyclic carbamates. When composed of two or more rings, the rings may be joined together in a fused or spiro fashion. Additionally, any nitrogens in a heteroalicyclyl may be quaternized. Heterocyclyl or heteroalicyclyl groups may be unsubstituted or substituted. Examples of such “heterocyclyl” or “heteroalicyclyl” groups include but are not limited to, 1,3-dioxin, 1,3-dioxane, 1,4-dioxane, 1,2-dioxolane, 1,3-dioxolane, 1,4-dioxolane, 1,3-oxathiane, 1,4-oxathiin, 1,3-oxathiolane, 1,3-dithiole, 1,3-dithiolane, 1,4-oxathiane, tetrahydro-1,4-thiazine, 2H-1,2-

oxazine, maleimide, succinimide, barbituric acid, thiobarbituric acid, dioxopiperazine, hydantoin, dihydrouracil, trioxane, hexahydro-1,3,5-triazine, imidazoline, imidazolidine, isoxazoline, isoxazolidine, oxazoline, oxazolidine, oxazolidinone, thiazoline, thiazolidine, morpholine, oxirane, piperidine N-Oxide, piperidine, piperazine, pyrrolidine, pyrrolidone, pyrrolidione, 4-piperidone, pyrazoline, pyrazolidine, 2-oxopyrrolidine, tetrahydropyran, 4H-pyran, tetrahydrothiopyran, thiamorpholine, thiamorpholine sulfoxide, thiamorpholine sulfone and their benzo-fused analogs (e.g., benzimidazolidinone, tetrahydroquinoline and/or 3,4-methylenedioxyphenyl).

[0028] As used herein, “aralkyl” and “aryl(alkyl)” refer to an aryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, 2-phenylalkyl, 3-phenylalkyl and naphthylalkyl.

[0029] As used herein, “heteroaralkyl” and “heteroaryl(alkyl)” refer to a heteroaryl group connected, as a substituent, via a lower alkylene group. The lower alkylene and heteroaryl group of heteroaralkyl may be substituted or unsubstituted. Examples include but are not limited to 2-thienylalkyl, 3-thienylalkyl, furylalkyl, thienylalkyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl and imidazolylalkyl and their benzo-fused analogs.

[0030] A “heteroalicycyl(alkyl)” and “heterocycyl(alkyl)” refer to a heterocyclic or a heteroalicyclic group connected, as a substituent, via a lower alkylene group. The lower alkylene and heterocycyl of a (heteroalicycyl)alkyl may be substituted or unsubstituted. Examples include but are not limited tetrahydro-2H-pyran-4-yl(methyl), piperidin-4-yl(ethyl), piperidin-4-yl(propyl), tetrahydro-2H-thiopyran-4-yl(methyl) and 1,3-thiazinan-4-yl(methyl).

[0031] “Lower alkylene groups” are straight-chained $\text{-CH}_2\text{-}$ tethering groups, forming bonds to connect molecular fragments via their terminal carbon atoms. Examples include but are not limited to methylene ($\text{-CH}_2\text{-}$), ethylene ($\text{-CH}_2\text{CH}_2\text{-}$), propylene ($\text{-CH}_2\text{CH}_2\text{CH}_2\text{-}$) and butylene ($\text{-CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-}$). A lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene group and/or by substituting both

hydrogens on the same carbon with a cycloalkyl group (e.g., ).

[0032] As used herein, the term “hydroxy” refers to a -OH group.

[0033] As used herein, “alkoxy” refers to the formula –OR wherein R is an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl) is defined herein. A non-limiting list of alkoxy is methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy, phenoxy and benzoxy. An alkoxy may be substituted or unsubstituted.

[0034] As used herein, “acyl” refers to a hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aryl(alkyl), heteroaryl(alkyl) and heterocyclyl(alkyl) connected, as substituents, via a carbonyl group. Examples include formyl, acetyl, propanoyl, benzoyl and acryl. An acyl may be substituted or unsubstituted.

[0035] A “cyano” group refers to a “-CN” group.

[0036] The term “halogen atom” or “halogen” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, such as, fluorine, chlorine, bromine and iodine.

[0037] A “thiocarbonyl” group refers to a “-C(=S)R” group in which R can be the same as defined with respect to O-carboxy. A thiocarbonyl may be substituted or unsubstituted.

[0038] An “O-carbamyl” group refers to a “-OC(=O)N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-carbamyl may be substituted or unsubstituted.

[0039] An “N-carbamyl” group refers to an “ROC(=O)N(R_A)-” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-carbamyl may be substituted or unsubstituted.

[0040] An “O-thiocarbamyl” group refers to a “-OC(=S)-N(R_AR_B)” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An O-thiocarbamyl may be substituted or unsubstituted.

[0041] An “N-thiocarbamyl” group refers to an “ $\text{ROC(=S)N(R}_A\text{)-}$ ” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-thiocarbamyl may be substituted or unsubstituted.

[0042] A “C-amido” group refers to a “ $\text{-C(=O)N(R}_A\text{R}_B\text{)-}$ ” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A C-amido may be substituted or unsubstituted.

[0043] An “N-amido” group refers to a “ $\text{RC(=O)N(R}_A\text{)-}$ ” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-amido may be substituted or unsubstituted.

[0044] A “C-thioamido” group refers to a “ $\text{-C(=S)N(R}_A\text{R}_B\text{)-}$ ” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A C-thioamido may be substituted or unsubstituted.

[0045] An “N-thioamido” group refers to a “ $\text{RC(=S)N(R}_A\text{)-}$ ” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-thioamido may be substituted or unsubstituted.

[0046] An “S-sulfonamido” group refers to a “ $\text{-SO}_2\text{N(R}_A\text{R}_B\text{)-}$ ” group in which R_A and R_B can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An S-sulfonamido may be substituted or unsubstituted.

[0047] An “N-sulfonamido” group refers to a “ $\text{RSO}_2\text{N(R}_A\text{)-}$ ” group in which R and R_A can be independently hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). An N-sulfonamido may be substituted or unsubstituted.

[0048] An “O-carboxy” group refers to a “RC(=O)O-” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, an alkoxy, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. An O-carboxy may be substituted or unsubstituted.

[0049] The terms “ester” and “C-carboxy” refer to a “-C(=O)OR” group in which R can be the same as defined with respect to O-carboxy. An ester and C-carboxy may be substituted or unsubstituted.

[0050] A “sulfenyl” group refers to an “-SR” group in which R can be hydrogen, an alkyl, an alkenyl, an alkynyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl). A sulfenyl may be substituted or unsubstituted.

[0051] A “sulfinyl” group refers to an “-S(=O)-R” group in which R can be the same as defined with respect to sulfenyl. A sulfinyl may be substituted or unsubstituted.

[0052] A “sulfonyl” group refers to an “SO₂R” group in which R can be the same as defined with respect to sulfenyl. A sulfonyl may be substituted or unsubstituted.

[0053] As used herein, “haloalkyl” refers to an alkyl group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkyl, di-haloalkyl and tri-haloalkyl). Such groups include but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 1-chloro-2-fluoromethyl and 2-fluoroisobutyl. A haloalkyl may be substituted or unsubstituted.

[0054] As used herein, “haloalkoxy” refers to an alkoxy group in which one or more of the hydrogen atoms are replaced by a halogen (e.g., mono-haloalkoxy, di-haloalkoxy and tri-haloalkoxy). Such groups include but are not limited to, chloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 1-chloro-2-fluoromethoxy and 2-fluoroisobutoxy. A haloalkoxy may be substituted or unsubstituted.

[0055] The term “amino” as used herein refers to a -NH₂ group.

[0056] A “mono-substituted amine” group refers to a “-NHR” group in which R can be an alkyl, an alkenyl, an alkynyl, a haloalkyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. A mono-substituted amino may be substituted or unsubstituted. Examples

of mono-substituted amino groups include, but are not limited to, -NH(methyl) , -NH(phenyl) and the like.

[0057] A “di-substituted amine” group refers to a “ $\text{-NR}_\text{A}\text{R}_\text{B}$ ” group in which R_A and R_B can be independently an alkyl, an alkenyl, an alkynyl, a haloalkyl, a cycloalkyl, a cycloalkenyl, aryl, heteroaryl, heterocyclyl, cycloalkyl(alkyl), aryl(alkyl), heteroaryl(alkyl) or heterocyclyl(alkyl), as defined herein. A di-substituted amino may be substituted or unsubstituted. Examples of di-substituted amino groups include, but are not limited to, -N(methyl)_2 , $\text{-N(phenyl)(methyl)}$, -N(ethyl)(methyl) and the like.

[0058] As used herein, the term “amino acid” refers to α -amino acids. Examples of suitable α -amino acids include, but are not limited to, alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, proline, serine, selenocysteine, tyrosine, arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine. Additional examples of suitable amino acids include, but are not limited to, ornithine, hypusine, 2-aminoisobutyric acid, dehydroalanine, citrulline, alpha-ethyl-glycine, alpha-propyl-glycine and norleucine.

[0059] Where the numbers of substituents is not specified (e.g. haloalkyl), there may be one or more substituents present. For example, “haloalkyl” may include one or more of the same or different halogens. As another example, “ $\text{C}_1\text{-C}_3$ alkoxyphenyl” may include one or more of the same or different alkoxy groups containing one, two or three atoms.



[0060] As used herein, a radical indicates species with a single, unpaired electron such that the species containing the radical can be covalently bonded to another species. Hence, in this context, a radical is not necessarily a free radical. Rather, a radical indicates a specific portion of a larger molecule. The term “radical” can be used interchangeably with the term “group.”

[0061] The term “pharmaceutically acceptable salt” refers to a salt of a compound that does not cause significant irritation to an organism to which it is administered and does not abrogate the biological activity and properties of the compound. In some embodiments, the salt is an acid addition salt of the compound. Pharmaceutical salts can be obtained by reacting a compound with inorganic acids such as hydrohalic acid (e.g., hydrochloric acid or hydrobromic acid), a sulfuric acid, a nitric acid and a phosphoric acid (such as 2,3-

dihydroxypropyl dihydrogen phosphate). Pharmaceutical salts can also be obtained by reacting a compound with an organic acid such as aliphatic or aromatic carboxylic or sulfonic acids, for example formic, acetic, succinic, lactic, malic, tartaric, citric, ascorbic, nicotinic, methanesulfonic, ethanesulfonic, p-toluenesulfonic, trifluoroacetic, benzoic, salicylic, 2-oxopentanedioic, or naphthalenesulfonic acid. Pharmaceutical salts can also be obtained by reacting a compound with a base to form a salt such as an ammonium salt, an alkali metal salt, such as a sodium, a potassium or a lithium salt, an alkaline earth metal salt, such as a calcium or a magnesium salt, a salt of a carbonate, a salt of a bicarbonate, a salt of organic bases such as dicyclohexylamine, N-methyl-D-glucamine, tris(hydroxymethyl)methylamine, C₁-C₇ alkylamine, cyclohexylamine, triethanolamine, ethylenediamine, and salts with amino acids such as arginine and lysine. For compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), those skilled in the art understand that when a salt is formed by protonation of a nitrogen-based group (for example, NH₂), the nitrogen-based group can be associated with a positive charge (for example, NH₂ can become NH₃⁺) and the positive charge can be balanced by a negatively charged counterion (such as Cl⁻).

[0062] It is understood that, in any compound described herein having one or more chiral centers, if an absolute stereochemistry is not expressly indicated, then each center may independently be of R-configuration or S-configuration or a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, enantiomerically enriched, racemic mixture, diastereomerically pure, diastereomerically enriched, or a stereoisomeric mixture. In addition, it is understood that in any compound described herein having one or more double bond(s) generating geometrical isomers that can be defined as E or Z, each double bond may independently be E or Z, or a mixture thereof.

[0063] In some embodiments, in any compound described, all tautomeric forms are also intended to be included. For example, without limitation, a reference to the

compound  may be interpreted to include tautomer .

[0064] It is to be understood that where compounds disclosed herein have unfilled valencies, then the valencies are to be filled with hydrogens or isotopes thereof, e.g., hydrogen-1 (protium) and hydrogen-2 (deuterium).

[0065] It is understood that the compounds described herein can be labeled isotopically. Substitution with isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements. Each chemical element as represented in a compound structure may include any isotope of said element. For example, in a compound structure a hydrogen atom may be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom may be present, the hydrogen atom can be any isotope of hydrogen, including but not limited to hydrogen-1 (protium) and hydrogen-2 (deuterium). Thus, reference herein to a compound encompasses all potential isotopic forms unless the context clearly dictates otherwise.

[0066] It is understood that the methods and combinations described herein include crystalline forms (also known as polymorphs, which include the different crystal packing arrangements of the same elemental composition of a compound), amorphous phases, salts, solvates, and hydrates. In some embodiments, the compounds described herein exist in solvated forms with pharmaceutically acceptable solvents such as water, ethanol, or the like. In other embodiments, the compounds described herein exist in unsolvated form. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, or the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.


[0067] Where a range of values is provided, it is understood that the upper and lower limit, and each intervening value between the upper and lower limit of the range is encompassed within the embodiments.

[0068] Terms and phrases used in this application, and variations thereof, especially in the appended claims, unless otherwise expressly stated, should be construed as open ended as opposed to limiting. As examples of the foregoing, the term ‘including’ should be read to mean ‘including, without limitation,’ ‘including but not limited to,’ or the like; the term ‘comprising’ as used herein is synonymous with ‘including,’ ‘containing,’ or ‘characterized by,’ and is inclusive or open-ended and does not exclude additional, unrecited elements or method steps; the term ‘having’ should be interpreted as ‘having at least;’ the term ‘includes’ should be interpreted as ‘includes but is not limited to;’ the term ‘example’ is used to provide exemplary instances of the item in discussion, not an exhaustive or limiting list thereof; and use of terms like ‘preferably,’ ‘preferred,’ ‘desired,’ or ‘desirable,’ and words of similar meaning should not be understood as implying that certain features are critical, essential, or even important to the structure or function, but instead as merely intended to highlight alternative or additional features that may or cannot be utilized in a particular embodiment. In addition, the term “comprising” is to be interpreted synonymously with the phrases "having at least" or "including at least". When used in the context of a process, the term "comprising" means that the process includes at least the recited steps, but may include additional steps. When used in the context of a compound, composition or device, the term "comprising" means that the compound, composition or device includes at least the recited features or components, but may also include additional features or components.

[0069] With respect to the use of substantially any plural and/or singular terms herein, those having skill in the art can translate from the plural to the singular and/or from the singular to the plural as is appropriate to the context and/or application. The various singular/plural permutations may be expressly set forth herein for sake of clarity. The indefinite article “a” or “an” does not exclude a plurality. The mere fact that certain measures are recited in mutually different dependent claims does not indicate that a combination of these measures cannot be used to advantage. Any reference signs in the claims should not be construed as limiting the scope.

Compounds

[0070] Some embodiments described herein generally relate to a compound, or a pharmaceutically acceptable salt thereof, selected from:




wherein: each X^1 , each X^2 , each X^3 , each X^4 , each X^5 , each X^6 , each X^7 , each X^{11} , each X^{12} , each X^{13} and each X^{14} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^1 is fluoro or chloro, provided that at least two of X^2 is fluoro or chloro, provided that at least two of X^3 is fluoro or chloro, provided that at least two of X^4 is fluoro or chloro, provided that at least two of X^5 is fluoro or chloro, provided that at least two of X^6 is fluoro or chloro, provided that at least two of X^7 is fluoro or chloro, provided that at least two of X^{11} is fluoro or chloro, provided that at least two of X^{12} is fluoro or chloro, provided that at least two of X^{13} is fluoro or chloro, and provided that at least two of X^{14} is fluoro or chloro; each X^8 , each X^9 and X^{10} can be independently

deuterium, fluoro or chloro; R^1 can be an unsubstituted α -amino acid; R^2 and R^3 can be independently an unsubstituted C_{1-4} alkyl; R^{9a} and R^{9b} can be independently an unsubstituted C_{1-4} alkyl; R^{5A} , R^{5B} , R^{5D} , R^{5E} , R^{5F} , R^{5G} , R^{5H} , R^{5J} , R^{5K} and R^{5L} can be independently hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; R^{5C} can be selected from hydrogen, deuterium, an unsubstituted C_{1-4} alkyl and $C(=O)R^{12}$; R^{4a} and R^{4b} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, a hydroxy substituted C_{1-4} alkyl or $-C(X^{16})_3$, provided that at least one of R^{4a} and R^{4b} is $-C(X^{16})_3$; R^6 , R^7 and R^8 can be independently $-(CH_2)_m$; R^{10} can be selected from hydrogen, deuterium, halogen, hydroxy, an unsubstituted C_{1-8} alkyl, an unsubstituted C_{3-20} cycloalkyl and an unsubstituted C_{1-8} haloalkyl; R^{11} can be selected from hydrogen, deuterium, halogen, an unsubstituted C_{1-30} alkyl, an unsubstituted C_{2-30} alkenyl and an unsubstituted C_{1-8} haloalkyl; R^{12} can be selected from of hydrogen, deuterium, an unsubstituted C_{1-30} alkyl and an unsubstituted C_{2-30} alkenyl; each X^{16} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{16} is fluoro or chloro; Z^1 , Z^2 and Z^3 can be independently nitrogen, oxygen or sulfur; m can be 1, 2, 3 or 4; each X^A , each X^B , each X^C , each X^D , each X^E , each X^F , each X^G , each X^H , each X^I , each X^K , each X^L , each X^M , each X^N , each X^P , each X^Q , each X^R and each X^S can be independently deuterium, chloro or fluoro; $p1$, $p2$, $p3$, $p4$, $p5$, $p6$, $p7$, $p8$, $p9$, $p10$, $p11$, $p12$, $p13$, $p14$, $p15$ and $p16$ can be independently 0, 1, 2, 3, 4, 5 or 6; and $p17$ can be 1, 2, 3, 4, 5 or 6.

Formula (Ia)


[0071] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ia), or pharmaceutically acceptable salt thereof, having the structure:



[0072] In some embodiments of Formula (Ia), each X^1 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^1 is fluoro or chloro; R^1 can be an unsubstituted α -amino acid; each X^A can be independently deuterium, chloro or fluoro; and $p1$ can be independently 0, 1, 2, 3, 4, 5 or 6.


[0073] In some embodiments of Formula (Ia), each X^1 can be fluoro. In some embodiments, each X^1 can be chloro. In some embodiments, two of X^1 can be fluoro, and one of X^1 can be chloro. In some embodiments, two of X^1 can be chloro, and one of X^1 can be fluoro. In some embodiments, two of X^1 can be fluoro, and one of X^1 can be hydrogen. In some embodiments, two of X^1 can be fluoro, and one of X^1 can be deuterium. In some embodiments, two of X^1 can be fluoro, and one of X^1 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, two of X^1 can be chloro, and one of X^1 can be hydrogen. In some embodiments, two of X^1 can be chloro, and one of X^1 can be deuterium. In some embodiments, two of X^1 can be chloro, and one of X^1 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0074] In some embodiments of Formula (Ia), R^1 can be an unsubstituted α -amino acid. Examples of unsubstituted α -amino acids are provided herein. In some embodiments, R^1 can be selected from alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, proline, serine, selenocysteine, tyrosine, arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan and valine. When R^1 is an unsubstituted α -amino acid, the carboxylic acid portion of the unsubstituted α -amino acid can be transformed to a $-NH$ group such that the unsubstituted α -amino acid is attached to the bicyclo[1.1.1]pentyl ring via the $-NH$ group. For example, when R^1 is valine, R^1 has the




structure . Other examples of unsubstituted α -amino acids wherein the carboxylic acid portion of the unsubstituted α -amino acid is transformed to a $-NH$ group

include, but are not limited to, the following:






and . In some embodiments, R¹ can be H₂N-CH(R'')-C(=O)-NH-, wherein R'' is a side chain of an α-amino acid. Alpha-amino acids are known to those skilled in the art and include but are not limited to, alanine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, proline, serine, selenocysteine, tyrosine, arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, ornithine, hypusine, 2-aminoisobutyric acid, dehydroalanine, citrulline, alpha-ethyl-glycine, alpha-propyl-glycine and norleucine.

[0075] In some embodiments of Formula (Ia), p1 can be 0. In other embodiments, each X^A can be deuterium, and p1 can be 6. In still other embodiments, each X^A can be deuterium, and p1 can be 5. In yet still other embodiments, each X^A can be deuterium, and p1 can be 4. In some embodiments, each X^A can be deuterium, and p1 can be 3. In other

embodiments, each X^A can be deuterium, and p1 can be 2. In still other embodiments, each X^A can be deuterium, and p1 can be 1. In some embodiments, each X^A can be chloro, and p1 can be 6. In other embodiments, each X^A can be chloro, and p1 can be 5. In still other embodiments, each X^A can be chloro, and p1 can be 4. In yet still other embodiments, each X^A can be chloro, and p1 can be 3. In some embodiments, each X^A can be chloro, and p1 can be 2. In other embodiments, each X^A can be chloro, and p1 can be 1. In some embodiments, each X^A can be fluoro, and p1 can be 6. In other embodiments, each X^A can be fluoro, and p1 can be 5. In still other embodiments, each X^A can be fluoro, and p1 can be 4. In yet still other embodiments, each X^A can be fluoro, and p1 can be 3. In some embodiments, each X^A can be fluoro, and p1 can be 2. In other embodiments, each X^A can be fluoro, and p1 can be 1.

Formula (Ib)

[0076] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ib), or pharmaceutically acceptable salt thereof, having the structure:



[0077] In some embodiments of Formula (Ib), each X^2 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^2 is fluoro or chloro; R^2 can be an unsubstituted C_{1-4} alkyl; each X^B can be independently deuterium, chloro or fluoro; and p2 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0078] In some embodiments of Formula (Ib), each X^2 can be fluoro. In some embodiments, each X^2 can be chloro. In some embodiments, two of X^2 can be fluoro, and one of X^2 can be chloro. In some embodiments, two of X^2 can be chloro, and one of X^2 can be fluoro. In some embodiments, two of X^2 can be fluoro, and one of X^2 can be hydrogen. In some embodiments, two of X^2 can be fluoro, and one of X^2 can be deuterium. In some embodiments, two of X^2 can be fluoro, and one of X^2 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^2 can be chloro, and one of X^2 can be hydrogen. In some


embodiments, two of X^2 can be chloro, and one of X^2 can be deuterium. In some embodiments, two of X^2 can be chloro, and one of X^2 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0079] In some embodiments of Formula (Ib), R^2 can be methyl. In other embodiments, R^2 can be ethyl. In still other embodiments, R^2 can be propyl, such as n-propyl or iso-propyl. In yet still other embodiments, R^2 can be butyl, for example, n-butyl, iso-butyl or tert-butyl.

[0080] In some embodiments of Formula (Ib), p2 can be 0. In other embodiments, each X^B can be deuterium, and p2 can be 6. In still other embodiments, each X^B can be deuterium, and p2 can be 5. In yet still other embodiments, each X^B can be deuterium, and p2 can be 4. In some embodiments, each X^B can be deuterium, and p2 can be 3. In other embodiments, each X^B can be deuterium, and p2 can be 2. In still other embodiments, each X^B can be deuterium, and p2 can be 1. In some embodiments, each X^B can be chloro, and p2 can be 6. In other embodiments, each X^B can be chloro, and p2 can be 5. In still other embodiments, each X^B can be chloro, and p2 can be 4. In yet still other embodiments, each X^B can be chloro, and p2 can be 3. In some embodiments, each X^B can be chloro, and p2 can be 2. In other embodiments, each X^B can be chloro, and p2 can be 1. In some embodiments, each X^B can be fluoro, and p2 can be 6. In other embodiments, each X^B can be fluoro, and p2 can be 5. In still other embodiments, each X^B can be fluoro, and p2 can be 4. In yet still other embodiments, each X^B can be fluoro, and p2 can be 3. In some embodiments, each X^B can be fluoro, and p2 can be 2. In other embodiments, each X^B can be fluoro, and p2 can be 1.


[0081] In some embodiments of Formula (Ib), when at least two of X^2 are halogen, such as fluoro and chloro, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when at least two of X^2 are halogen, such as fluoro and chloro, and p2 is 0, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when at least two of X^2 are fluoro and p2 is 0, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In some embodiments, when one of X^2 is an unsubstituted C_{1-4} alkyl, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when one of X^2 is an unsubstituted C_{1-4} alkyl

and p_2 is 0, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when one of X^2 is methyl and p_2 is 0, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In some embodiments, when p_2 is 0, then each of X^2 cannot be halogen, such as fluoro and chloro. In still other embodiments, when p_2 is 0, then two of X^2 cannot be fluoro. In yet still other embodiments, when p_2 is 0, then $-C(X^2)_3$ cannot be $-CF_3$. In some embodiments, when p_2 is 0, then R^2 cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when p_2 is 0, then R^2 cannot be an unsubstituted C_4 alkyl. In some embodiments, when two of X^2 are fluoro, one of X^2 is methyl and p_2 is 0, then R^2 cannot be an unsubstituted C_4 alkyl. In other embodiments, when three of X^2 is fluoro and p_2 is 0, then R^2 cannot be an unsubstituted C_4 alkyl. In some embodiments, p_2 cannot be 0. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. Examples of C_4 alkyl groups include n-butyl, iso-butyl and tert-butyl. In still other embodiments, a compound of Formula (Ib), or a pharmaceutically acceptable salt thereof, cannot one or more of the following compounds,



Formula (Ic)

[0082] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ic), or pharmaceutically acceptable salt thereof, having the structure:



[0083] In some embodiments of Formula (Ic), R^3 can be an unsubstituted C_{1-4} alkyl; R^{5A} can be independently hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^C can be independently deuterium, chloro or fluoro; and p_3 can be independently 0, 1, 2, 3, 4, 5 or 6.


[0084] In some embodiments of Formula (Ic), R^3 can be methyl. In other embodiments, R^3 can be ethyl. In still other embodiments, R^3 can be propyl, such as n-propyl or iso-propyl. In yet still other embodiments, R^3 can be butyl, for example, n-butyl, iso-butyl or tert-butyl.

[0085] In some embodiments of Formula (Ic), R^{5A} can be hydrogen. In other embodiments, R^{5A} can be deuterium. In still other embodiments, R^{5A} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0086] In some embodiments of Formula (Ic), p3 can be 0. In other embodiments, each X^C can be deuterium, and p3 can be 6. In still other embodiments, each X^C can be deuterium, and p3 can be 5. In yet still other embodiments, each X^C can be deuterium, and p3 can be 4. In some embodiments, each X^C can be deuterium, and p3 can be 3. In other embodiments, each X^C can be deuterium, and p3 can be 2. In still other embodiments, each X^C can be deuterium, and p3 can be 1. In some embodiments, each X^C can be chloro, and p3 can be 6. In other embodiments, each X^C can be chloro, and p3 can be 5. In still other embodiments, each X^C can be chloro, and p3 can be 4. In yet still other embodiments, each X^C can be chloro, and p3 can be 3. In some embodiments, each X^C can be chloro, and p3 can be 2. In other embodiments, each X^C can be chloro, and p3 can be 1. In some embodiments, each X^C can be fluoro, and p3 can be 6. In other embodiments, each X^C can be fluoro, and p3 can be 5. In still other embodiments, each X^C can be fluoro, and p3 can be 4. In yet still other embodiments, each X^C can be fluoro, and p3 can be 3. In some embodiments, each X^C can be fluoro, and p3 can be 2. In other embodiments, each X^C can be fluoro, and p3 can be 1.

Formula (Id)

[0087] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Id), or pharmaceutically acceptable salt thereof, having the structure:



[0088] In some embodiments of Formula (Id), each X^3 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^3 is fluoro or chloro; R^{5B} can be independently hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^D can be independently deuterium, chloro or fluoro; and p_4 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0089] In some embodiments of Formula (Id), each X^3 can be fluoro. In some embodiments, each X^3 can be chloro. In some embodiments, two of X^3 can be fluoro, and one of X^3 can be chloro. In some embodiments, two of X^3 can be chloro, and one of X^3 can be fluoro. In some embodiments, two of X^3 can be fluoro, and one of X^3 can be hydrogen. In some embodiments, two of X^3 can be fluoro, and one of X^3 can be deuterium. In some embodiments, two of X^3 can be fluoro, and one of X^3 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^3 can be chloro, and one of X^3 can be hydrogen. In some embodiments, two of X^3 can be chloro, and one of X^3 can be deuterium. In some embodiments, two of X^3 can be chloro, and one of X^3 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0090] In some embodiments of Formula (Id), R^{5B} can be hydrogen. In other embodiments, R^{5B} can be deuterium. In still other embodiments, R^{5B} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0091] In some embodiments of Formula (Id), p_4 can be 0. In other embodiments, each X^D can be deuterium, and p_4 can be 6. In some embodiments, each X^D can be deuterium, and p_4 can be 5. In still other embodiments, each X^D can be deuterium,

and p4 can be 4. In yet still other embodiments, each X^D can be deuterium, and p4 can be 3. In some embodiments, each X^D can be deuterium, and p4 can be 2. In other embodiments, each X^D can be deuterium, and p4 can be 1. In some embodiments, each X^D can be chloro, and p4 can be 6. In other embodiments, each X^D can be chloro, and p4 can be 5. In still other embodiments, each X^D can be chloro, and p4 can be 4. In yet still other embodiments, each X^D can be chloro, and p4 can be 3. In some embodiments, each X^D can be chloro, and p4 can be 2. In other embodiments, each X^D can be chloro, and p4 can be 1. In some embodiments, each X^D can be fluoro, and p4 can be 6. In other embodiments, each X^D can be fluoro, and p4 can be 5. In still other embodiments, each X^D can be fluoro, and p4 can be 4. In yet still other embodiments, each X^D can be fluoro, and p4 can be 3. In some embodiments, each X^D can be fluoro, and p4 can be 2. In other embodiments, each X^D can be fluoro, and p4 can be 1.

Formula (Ie)

[0092] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ie), or pharmaceutically acceptable salt thereof, having the structure:



[0093] In some embodiments of Formula (Ie), R^{4a} and R^{4b} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, a hydroxy substituted C_{1-4} alkyl or $-C(X^{16})_3$, provided that at least one of R^{4a} and R^{4b} is $-C(X^{16})_3$; each X^{16} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{16} is fluoro or chloro; R^{5C} can be selected from hydrogen, deuterium, an unsubstituted C_{1-4} alkyl and $C(=O)R^{12}$; R^{12} can be selected from hydrogen, deuterium, an unsubstituted C_{1-30} alkyl and an unsubstituted C_{2-30} alkenyl; each X^E can be independently deuterium, chloro or fluoro; p5 can be independently 0, 1, 2, 3, 4, 5 or 6; and Z^1 can be nitrogen, oxygen or sulfur.

[0094] In some embodiments of Formula (Ie), R^{4a} can be hydrogen, and R^{4b} can be $-C(X^{16})_3$. In some embodiments, R^{4a} can be deuterium, and R^{4b} can be $-C(X^{16})_3$. In some embodiments, R^{4a} can be methyl, and R^{4b} can be $-C(X^{16})_3$. In some embodiments, R^{4a} can be

ethyl, and R^{4b} can be $-C(X^{16})_3$. In some embodiments, R^{4a} can be propyl, such as n-propyl or iso-propyl, and R^{4b} can be $-C(X^{16})_3$. In some embodiments, R^{4a} can be butyl, for example, n-butyl, iso-butyl or tert-butyl, and R^{4b} can be $-C(X^{16})_3$. In other embodiments, R^{4a} can be a hydroxy substituted C_{1-4} alkyl (for example, $HO-CH_2-$), and R^{4b} can be $-C(X^{16})_3$. In any embodiments of this paragraph, each X^{16} can be fluoro. In any embodiments of this paragraph, each X^{16} can be chloro. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be chloro. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be fluoro. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be hydrogen. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be deuterium. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be an unsubstituted C_{1-4} alkyl. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be hydrogen. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be deuterium. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be an unsubstituted C_{1-4} alkyl.

[0095] In some embodiments of Formula (Ie), R^{4b} can be hydrogen, and R^{4a} can be $-C(X^{16})_3$. In some embodiments, R^{4b} can be deuterium, and R^{4a} can be $-C(X^{16})_3$. In some embodiments, R^{4b} can be methyl, and R^{4a} can be $-C(X^{16})_3$. In some embodiments, R^{4b} can be ethyl, and R^{4a} can be $-C(X^{16})_3$. In some embodiments, R^{4b} can be propyl, such as n-propyl or iso-propyl, and R^{4a} can be $-C(X^{16})_3$. In some embodiments, R^{4b} can be butyl, for example, n-butyl, iso-butyl or tert-butyl, and R^{4a} can be $-C(X^{16})_3$. In other embodiments, R^{4b} can be a hydroxy substituted C_{1-4} alkyl (such as, $HO-CH_2-$), and R^{4a} can be $-C(X^{16})_3$. In any embodiments of this paragraph, each X^{16} can be fluoro. In any embodiments of this paragraph, each X^{16} can be chloro. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be chloro. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be fluoro. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be hydrogen. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be deuterium. In any embodiments of this paragraph, two of X^{16} can be fluoro, and one of X^{16} can be an unsubstituted C_{1-4} alkyl. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be hydrogen. In

any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be deuterium. In any embodiments of this paragraph, two of X^{16} can be chloro, and one of X^{16} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups, for example for X^{16} , include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0096] In some embodiments of Formula (Ie), R^{5C} can be hydrogen. In other embodiments, R^{5C} can be deuterium. In still other embodiments, R^{5C} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In still other embodiments, R^{5C} can be $C(=O)R^{12}$.


[0097] When R^{5C} is $C(=O)R^{12}$, R^{12} can be a variety of groups. In some embodiments, R^{12} can be hydrogen. In other embodiments, R^{12} can be deuterium. When R^{12} is an alkyl group, the number of carbon atoms of the alkyl group can vary. The alkyl group can be a long alkyl having 1 to 30 carbons, a medium alkyl having 1 to 12 carbon atoms or a lower alkyl having 1 to 6 carbon atoms. In some embodiments, R^{12} can be an unsubstituted C_{1-6} alkyl. In other embodiments, R^{12} can be an unsubstituted C_{1-12} alkyl. In still other embodiments, R^{12} can be an unsubstituted C_{1-30} alkyl. In yet still other embodiments, R^{12} can be an unsubstituted C_{7-26} alkyl. In even still other embodiments, R^{12} can be an unsubstituted C_{12-26} alkyl. Examples of lower alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, a tert-butyl, pentyl (straight and branched) and hexyl (straight and branched). Examples of unsubstituted C_{1-30} alkyls include, but are not limited to, $-(CH_2)_6CH_3$, $-(CH_2)_8CH_3$, $-(CH_2)_{10}CH_3$, $-(CH_2)_{12}CH_3$, $-(CH_2)_{14}CH_3$, $-(CH_2)_{16}CH_3$, $-(CH_2)_{18}CH_3$, $-(CH_2)_{20}CH_3$, $-(CH_2)_{22}CH_3$ and $-(CH_2)_{24}CH_3$.

[0098] Similar to alkyls, alkenyls can be a long alkenyl having 2 to 30 carbons, a medium alkenyl having 2 to 12 carbon atoms or a lower alkenyl having 2 to 6 carbon atoms. In some embodiments, R^{12} can be an unsubstituted C_{2-30} alkenyl. In other embodiments, R^{12} can be an unsubstituted C_{14-22} alkenyl. Examples of unsubstituted C_{2-30} alkenyls include, but are not limited to, $-(CH_2)_7CH=CH(CH_2)_3CH_3$, $-(CH_2)_7CH=CHCH_2CH=CH(CH_2)_4CH_3$, $-(CH_2)_7CH=CH(CH_2)_7CH_3$, $-(CH_2)_7CH=CHCH_2CH=CH(CH_2)_4CH_3$, $-(CH_2)_7CH=CH(CH_2)_7CH_3$, $-(CH_2)_7CH=CHCH_2CH=CHCH_2CH=CHCH_2CH_3$, $-(CH_2)_9CH=CH(CH_2)_5CH_3$, $-(CH_2)_3CH=CHCH_2CH=CHCH_2CH=CHCH_2CH=CH(CH_2)_4CH_3$, $-(CH_2)_{11}CH=CH(CH_2)_7CH_3$, $-(CH_2)_3CH=CHCH_2CH=CHCH_2CH=CHCH_2CH=CHCH_2CH=CHCH_2CH_3$,

$-(\text{CH}_2)_4\text{CH}=\text{CHCH}(\text{CH}_3)_2$ and

$-(\text{CH}_2)_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$.

[0099] In some embodiments, R^{12} can be the aliphatic tail of a saturated or an unsaturated fatty acid. As an example, R^{12} can be the aliphatic tail of caprylic acid ($\text{HOOC}\underline{\text{C}}(\text{CH}_2)_6\text{CH}_3$). In this example of caprylic acid, the aliphatic tail is bolded and italicized. When the saturated or an unsaturated fatty acid becomes part of a compound of Formula (Ie), the carbon of the carboxylic acid of the saturated or an unsaturated fatty acid becomes the carbon of $\underline{\text{C}}(=\text{O})\text{R}^{12}$ that is bold and underlined. For example, when R^{12} is the aliphatic tail of caprylic acid, the compound of Formula (Ie) can have the following structure:



. A non-limiting list of suitable saturated or an unsaturated fatty acids are myristoleic acid, palmitoleic, sapienic acid, linoleic acid, oleic acid, linoleiaidic acid, elaidic acid, alpha-linolenic acid, vaccenic acid, arachidonic acid, erucic acid, eicosapentaenoic acid, (E)-8-methylnon-6-enoic acid, docosahexaenoic acid, caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, arachidic acid, behenic acid, lignoceric acid and cerotic acid.


[0100] In some embodiments of Formula (Ie), p_5 can be 0. In other embodiments, each X^{E} can be deuterium, and p_5 can be 6. In still other embodiments, each X^{E} can be deuterium, and p_5 can be 5. In yet still other embodiments, each X^{E} can be deuterium, and p_5 can be 4. In some embodiments, each X^{E} can be deuterium, and p_5 can be 3. In other embodiments, each X^{E} can be deuterium, and p_5 can be 2. In still other embodiments, each X^{E} can be deuterium, and p_5 can be 1. In some embodiments, each X^{E} can be chloro, and p_5 can be 6. In other embodiments, each X^{E} can be chloro, and p_5 can be 5. In still other embodiments, each X^{E} can be chloro, and p_5 can be 4. In yet still other embodiments, each X^{E} can be chloro, and p_5 can be 3. In some embodiments, each X^{E} can be chloro, and p_5 can be 2. In other embodiments, each X^{E} can be chloro, and p_5 can be 1. In some embodiments, each X^{E} can be fluoro, and p_5 can be 6. In other embodiments, each X^{E} can be fluoro, and p_5 can be 5. In still other embodiments, each X^{E} can be fluoro, and p_5 can be 4. In yet still other embodiments, each X^{E} can be fluoro, and p_5 can be 3. In some embodiments, each X^{E}

can be fluoro, and p5 can be 2. In other embodiments, each X^E can be fluoro, and p5 can be 1.

[0101] In some embodiments of Formula (Ie), Z^1 can be nitrogen. In other embodiments, Z^1 can be oxygen. In still other embodiments, Z^1 can be sulfur.

Formula (If)

[0102] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (If), or pharmaceutically acceptable salt thereof, having the structure:



[0103] In some embodiments of Formula (If), each X^4 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^4 is fluoro or chloro; R^{5D} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^F can be independently deuterium, chloro or fluoro; and p6 can be independently 0, 1, 2, 3, 4, 5 or 6.


[0104] In some embodiments of Formula (If), each X^4 can be fluoro. In some embodiments, each X^4 can be chloro. In some embodiments, two of X^4 can be fluoro, and one of X^4 can be chloro. In some embodiments, two of X^4 can be chloro, and one of X^4 can be fluoro. In some embodiments, two of X^4 can be fluoro, and one of X^4 can be hydrogen. In some embodiments, two of X^4 can be fluoro, and one of X^4 can be deuterium. In some embodiments, two of X^4 can be fluoro, and one of X^4 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^4 can be chloro, and one of X^4 can be hydrogen. In some embodiments, two of X^4 can be chloro, and one of X^4 can be deuterium. In some embodiments, two of X^4 can be chloro, and one of X^4 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0105] In some embodiments of Formula (If), R^{5D} can be hydrogen. In some embodiments, R^{5D} can be deuterium. In some embodiments, R^{5D} can be an unsubstituted C_{1-4} alkyl, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0106] In some embodiments of Formula (If), p6 can be 0. In other embodiments, each X^F can be deuterium, and p6 can be 6. In still other embodiments, each X^F can be deuterium, and p6 can be 5. In yet still other embodiments, each X^F can be deuterium, and p6 can be 4. In some embodiments, each X^F can be deuterium, and p6 can be 3. In other embodiments, each X^F can be deuterium, and p6 can be 2. In still other embodiments, each X^F can be deuterium, and p6 can be 1. In some embodiments, each X^F can be chloro, and p6 can be 6. In other embodiments, each X^F can be chloro, and p6 can be 5. In still other embodiments, each X^F can be chloro, and p6 can be 4. In yet still other embodiments, each X^F can be chloro, and p6 can be 3. In some embodiments, each X^F can be chloro, and p6 can be 2. In other embodiments, each X^F can be chloro, and p6 can be 1. In some embodiments, each X^F can be fluoro, and p6 can be 6. In other embodiments, each X^F can be fluoro, and p6 can be 5. In still other embodiments, each X^F can be fluoro, and p6 can be 4. In yet still other embodiments, each X^F can be fluoro, and p6 can be 3. In some embodiments, each X^F can be fluoro, and p6 can be 2. In other embodiments, each X^F can be fluoro, and p6 can be 1.


[0107] In some embodiments of Formula (If), when at least two of X^4 are halogen, such as fluoro and chloro, then R^{5D} cannot be hydrogen. In other embodiments, when at least two of X^4 are halogen, such as fluoro and chloro, and p6 is 0, then R^{5D} cannot be hydrogen. In still other embodiments, when at least two of X^4 are fluoro and p6 is 0, then R^{5D} cannot be hydrogen. In some embodiments, when at least two of X^4 are halogen, such as fluoro and chloro, then R^{5D} cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when at least two of X^4 are halogen, such as fluoro and chloro, and p6 is 0, then R^{5D} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when at least two of X^4 are fluoro, and p6 is 0, then R^{5D} cannot be an unsubstituted C_{1-4} alkyl. In some embodiments, when one of X^4 is an unsubstituted C_{1-4} alkyl then R^{5D} cannot be hydrogen. In other embodiments, when one of X^4 is an unsubstituted C_{1-4} alkyl and p6 is 0, then R^{5D} cannot be hydrogen. In still other embodiments, when one of X^4 is methyl and p6 is 0, then R^{5D} cannot be hydrogen. In some embodiments, when one of X^4 is hydrogen, then R^{5D} cannot be hydrogen. In other embodiments, when one of X^4 is hydrogen and p6 is 0, then R^{5D} cannot be hydrogen. In some embodiment, when p6 is 0, then R^{5D} cannot be hydrogen. In other embodiments, when

p6 is 0, then R^{5D} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when p6 is 0, then $-C(X^4)_3$ cannot be $-CF_3$. In yet still other embodiments, when p6 is 0, then $-C(X^4)_3$ cannot be $-CCl_3$. In some embodiments, when $-C(X^4)_3$ is $-CF_3$ and p6 is 0, then R^{5D} cannot be an unsubstituted C_4 alkyl. In some embodiments, p6 cannot be 0. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. Examples of C_4 alkyl groups include n-butyl, iso-butyl and tert-butyl. In other embodiments, a compound of Formula (If), or a pharmaceutically acceptable salt thereof, cannot be one or more of the following compounds,



Formula (Ig)

[0108] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ig), or pharmaceutically acceptable salt thereof, having the structure:



[0109] In some embodiments of Formula (Ig), each X^5 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^5 is fluoro or chloro; R^6 can be $-(CH_2)_m-$; m can be 1, 2, 3 or 4; each X^G can be independently deuterium, chloro or fluoro; and p7 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0110] In some embodiments of Formula (Ig), each X^5 can be fluoro. In some embodiments, each X^5 can be chloro. In some embodiments, two of X^5 can be fluoro, and one of X^5 can be chloro. In some embodiments, two of X^5 can be chloro, and one of X^5 can be fluoro. In some embodiments, two of X^5 can be fluoro, and one of X^5 can be hydrogen. In some embodiments, two of X^5 can be fluoro, and one of X^5 can be deuterium. In some


embodiments, two of X^5 can be fluoro, and one of X^5 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^5 can be chloro, and one of X^5 can be hydrogen. In some embodiments, two of X^5 can be chloro, and one of X^5 can be deuterium. In some embodiments, two of X^5 can be chloro, and one of X^5 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0111] In some embodiments of Formula (Ig), R^6 can be $-(CH_2)_m-$, and m can be 1. In other embodiments, R^6 can be $-(CH_2)_m-$, and m can be 2. In still other embodiments, R^6 can be $-(CH_2)_m-$, and m can be 3. In yet still other embodiments, R^6 can be $-(CH_2)_m-$, and m can be 4.

[0112] In some embodiments of Formula (Ig), p7 can be 0. In other embodiments, each X^G can be deuterium, and p7 can be 6. In still other some embodiments, each X^G can be deuterium, and p7 can be 5. In yet still other embodiments, each X^G can be deuterium, and p7 can be 4. In some embodiments, each X^G can be deuterium, and p7 can be 3. In other embodiments, each X^G can be deuterium, and p7 can be 2. In still other embodiments, each X^G can be deuterium, and p7 can be 1. In some embodiments, each X^G can be chloro, and p7 can be 6. In other embodiments, each X^G can be chloro, and p7 can be 5. In still other embodiments, each X^G can be chloro, and p7 can be 4. In yet still other embodiments, each X^G can be chloro, and p7 can be 3. In some embodiments, each X^G can be chloro, and p7 can be 2. In other embodiments, each X^G can be chloro, and p7 can be 1. In some embodiments, each X^G can be fluoro, and p7 can be 6. In other embodiments, each X^G can be fluoro, and p7 can be 5. In still other embodiments, each X^G can be fluoro, and p7 can be 4. In yet still other embodiments, each X^G can be fluoro, and p7 can be 3. In some embodiments, each X^G can be fluoro, and p7 can be 2. In other embodiments, each X^G can be fluoro, and p7 can be 1.

Formula (Ih)

[0113] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ih), or pharmaceutically acceptable salt thereof, having the structure:



[0114] In some embodiments of Formula (Ih), each X^6 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^6 is fluoro or chloro; R^7 can be $-(CH_2)_m-$; m can be 1, 2, 3 or 4; each X^H can be independently deuterium, chloro or fluoro; and p_8 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0115] In some embodiments of Formula (Ih), each X^6 can be fluoro. In some embodiments, each X^6 can be chloro. In some embodiments, two of X^6 can be fluoro, and one of X^6 can be chloro. In some embodiments, two of X^6 can be chloro, and one of X^6 can be fluoro. In some embodiments, two of X^6 can be fluoro, and one of X^6 can be hydrogen. In some embodiments, two of X^6 can be fluoro, and one of X^6 can be deuterium. In some embodiments, two of X^6 can be fluoro, and one of X^6 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^6 can be chloro, and one of X^6 can be hydrogen. In some embodiments, two of X^6 can be chloro, and one of X^6 can be deuterium. In some embodiments, two of X^6 can be chloro, and one of X^6 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0116] In some embodiments of Formula (Ih), R^7 can be $-(CH_2)_m-$, and m can be 1. In other embodiments, R^7 can be $-(CH_2)_m-$, and m can be 2. In still other embodiments, R^7 can be $-(CH_2)_m-$, and m can be 3. In yet still embodiments, R^7 can be $-(CH_2)_m-$, and m can be 4.

[0117] In some embodiments of Formula (Ih), p_8 can be 0. In other embodiments, each X^H can be deuterium, and p_8 can be 6. In still other embodiments, each X^H can be deuterium, and p_8 can be 5. In yet still other embodiments, each X^H can be deuterium, and p_8 can be 4. In some embodiments, each X^H can be deuterium, and p_8 can be

3. In other embodiments, each X^H can be deuterium, and p8 can be 2. In still other embodiments, each X^H can be deuterium, and p8 can be 1. In some embodiments, each X^H can be chloro, and p8 can be 6. In other embodiments, each X^H can be chloro, and p8 can be 5. In still other embodiments, each X^H can be chloro, and p8 can be 4. In yet still other embodiments, each X^H can be chloro, and p8 can be 3. In some embodiments, each X^H can be chloro, and p8 can be 2. In other embodiments, each X^H can be chloro, and p8 can be 1. In some embodiments, each X^H can be fluoro, and p8 can be 6. In other embodiments, each X^H can be fluoro, and p8 can be 5. In still other embodiments, each X^H can be fluoro, and p8 can be 4. In yet still other embodiments, each X^H can be fluoro, and p8 can be 3. In some embodiments, each X^H can be fluoro, and p8 can be 2. In other embodiments, each X^H can be fluoro, and p8 can be 1.

Formula (Ik)

[0118] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ik), or pharmaceutically acceptable salt thereof, having the structure:



[0119] In some embodiments of Formula (Ik), each X^7 can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^7 is fluoro or chloro; R^8 can be $-(CH_2)_m-$; m can be 1, 2, 3 or 4; each X^J can be independently deuterium, chloro or fluoro; and p9 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0120] In some embodiments of Formula (Ik), each X^7 can be fluoro. In some embodiments, each X^7 can be chloro. In some embodiments, two of X^7 can be fluoro, and one of X^7 can be chloro. In some embodiments, two of X^7 can be chloro, and one of X^7 can be fluoro. In some embodiments, two of X^7 can be fluoro, and one of X^7 can be hydrogen. In some embodiments, two of X^7 can be fluoro, and one of X^7 can be deuterium. In some embodiments, two of X^7 can be fluoro, and one of X^7 can be an unsubstituted C_{1-4} alkyl. In some embodiments, two of X^7 can be chloro, and one of X^7 can be hydrogen. In some


embodiments, two of X^7 can be chloro, and one of X^7 can be deuterium. In some embodiments, two of X^7 can be chloro, and one of X^7 can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0121] In some embodiments of Formula (Ik), R^8 can be $-(CH_2)_m-$, and m can be 1. In other embodiments, R^8 can be $-(CH_2)_m-$, and m can be 2. In still other embodiments, R^8 can be $-(CH_2)_m-$, and m can be 3. In yet still other embodiments, R^8 can be $-(CH_2)_m-$, and m can be 4.

[0122] In some embodiments of Formula (Ik), p9 can be 0. In other embodiments, each X^J can be deuterium, and p9 can be 6. In still other embodiments, each X^J can be deuterium, and p9 can be 5. In yet still other embodiments, each X^J can be deuterium, and p9 can be 4. In some embodiments, each X^J can be deuterium, and p9 can be 3. In other embodiments, each X^J can be deuterium, and p9 can be 2. In still other embodiments, each X^J can be deuterium, and p9 can be 1. In some embodiments, each X^J can be chloro, and p9 can be 6. In other embodiments, each X^J can be chloro, and p9 can be 5. In still other embodiments, each X^J can be chloro, and p9 can be 4. In yet still other embodiments, each X^J can be chloro, and p9 can be 3. In some embodiments, each X^J can be chloro, and p9 can be 2. In other embodiments, each X^J can be chloro, and p9 can be 1. In some embodiments, each X^J can be fluoro, and p9 can be 6. In other embodiments, each X^J can be fluoro, and p9 can be 5. In still other embodiments, each X^J can be fluoro, and p9 can be 4. In yet still other embodiments, each X^J can be fluoro, and p9 can be 3. In some embodiments, each X^J can be fluoro, and p9 can be 2. In other embodiments, each X^J can be fluoro, and p9 can be 1.

Formula (Im)

[0123] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Im), or pharmaceutically acceptable salt thereof, having the structure:



[0124] In some embodiments of Formula (Im), each X^8 can be independently deuterium, fluoro or chloro; R^{5E} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^K can be independently deuterium, chloro or fluoro; and $p10$ can be independently 0, 1, 2, 3, 4, 5 or 6.

[0125] In any embodiments of Formula (Im) of this paragraph, five of X^8 can be deuterium. In any embodiments of this paragraph, four of X^8 can be deuterium. In any embodiments of this paragraph, three of X^8 can be deuterium. In any embodiments of this paragraph, two of X^8 can be deuterium. In any embodiments of this paragraph, one of X^8 can be deuterium. In any embodiments of this paragraph, five of X^8 can be fluoro. In any embodiments of this paragraph, four of X^8 can be fluoro. In any embodiments of this paragraph, three of X^8 can be fluoro. In any embodiments of this paragraph, two of X^8 can be fluoro. In any embodiments of this paragraph, one of X^8 can be fluoro. In any embodiments of this paragraph, five of X^8 can be chloro. In any embodiments of this paragraph, four of X^8 can be chloro. In any embodiments of this paragraph, three of X^8 can be chloro. In any embodiments of this paragraph, two of X^8 can be chloro. In any embodiments of this paragraph, one of X^8 can be chloro.

[0126] In some embodiments of Formula (Im), R^{5E} can be hydrogen. In other embodiments, R^{5E} can be deuterium. In still other embodiments, R^{5E} can be an unsubstituted C_{1-4} alkyl, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0127] In some embodiments of Formula (Im), $p10$ can be 0. In other embodiments, each X^K can be deuterium, and $p10$ can be 6. In still other embodiments, each X^K can be deuterium, and $p10$ can be 5. In yet still other embodiments, each X^K can be deuterium, and $p10$ can be 4. In some embodiments, each X^K can be deuterium, and $p10$ can

be 3. In other embodiments, each X^K can be deuterium, and p10 can be 2. In still other embodiments, each X^K can be deuterium, and p10 can be 1. In some embodiments, each X^K can be chloro, and p10 can be 6. In other embodiments, each X^K can be chloro, and p10 can be 5. In still other embodiments, each X^K can be chloro, and p10 can be 4. In yet still other embodiments, each X^K can be chloro, and p10 can be 3. In some embodiments, each X^K can be chloro, and p10 can be 2. In other embodiments, each X^K can be chloro, and p10 can be 1. In some embodiments, each X^K can be fluoro, and p10 can be 6. In other embodiments, each X^K can be fluoro, and p10 can be 5. In still other embodiments, each X^K can be fluoro, and p10 can be 4. In yet still other embodiments, each X^K can be fluoro, and p10 can be 3. In some embodiments, each X^K can be fluoro, and p10 can be 2. In other embodiments, each X^K can be fluoro, and p10 can be 1.


[0128] In some embodiments of Formula (Im), when at least one of X^8 is halogen, such as fluoro and chloro, then R^{5E} cannot be hydrogen. In other embodiments, when at least one of X^8 is halogen, such as fluoro and chloro, and p10 is 0, then R^{5E} cannot be hydrogen. In still other embodiments, when at least one of X^8 is fluoro and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least two of X^8 are halogen, such as fluoro and/or chloro, then R^{5E} cannot be hydrogen. In other embodiments, when at least two of X^8 are halogen, such as fluoro and/or chloro, and p10 is 0, then R^{5E} cannot be hydrogen. In still other embodiments, when at least two of X^8 are fluoro and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least three of X^8 are halogen, such as fluoro and/or chloro, then R^{5E} cannot be hydrogen. In other embodiments, when at least three of X^8 are halogen, such as fluoro and/or chloro, and p10 is 0, then R^{5E} cannot be hydrogen. In still other embodiments, when at least three of X^8 are fluoro and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least one of X^8 is deuterium, then R^{5E} cannot be hydrogen. In other embodiments, when at least one of X^8 is deuterium and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least two of X^8 is deuterium, then R^{5E} cannot be hydrogen. In other embodiments, when at two one of X^8 is deuterium and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least three of X^8 is deuterium, then R^{5E} cannot be hydrogen. In other embodiments, when at least three of X^8 is deuterium and p10 is 0, then R^{5E} cannot be hydrogen. In some embodiments, when at least one of X^8 is

halogen, such as fluoro and chloro, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when at least one of X^8 is halogen, such as fluoro and chloro, and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when at least one of X^8 is fluoro and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In some embodiments, when at least two of X^8 is halogen, such as fluoro and chloro, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when at least two of X^8 is halogen, such as fluoro and chloro, and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when at least two of X^8 is fluoro and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In some embodiments, when at least three of X^8 is halogen, such as fluoro and chloro, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In other embodiments, when at least three of X^8 is halogen, such as fluoro and chloro, and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when at least three of X^8 is fluoro and p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In some embodiment, when p_{10} is 0, then R^{5E} cannot be hydrogen. In other embodiments, when p_{10} is 0, then R^{5E} cannot be an unsubstituted C_{1-4} alkyl. In still other embodiments, when p_{10} is 0, then R^{5E} cannot be deuterium. In still other embodiments, p_{10} cannot be 0. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, a compound of Formula (In), or a pharmaceutically acceptable salt thereof, cannot be,



Formula (In)

[0129] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (In), or pharmaceutically acceptable salt thereof, having the structure:



[0130] In some embodiments of Formula (In), each X^9 can be independently deuterium, fluoro or chloro; R^{5F} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl;

each X^L can be independently deuterium, chloro or fluoro; p_{11} can be independently 0, 1, 2, 3, 4, 5 or 6; and Z^2 can be nitrogen, oxygen or sulfur.

[0131] In any embodiments of Formula (In) of this paragraph, four of X^9 can be deuterium. In any embodiments of this paragraph, three of X^9 can be deuterium. In any embodiments of this paragraph, two of X^9 can be deuterium. In any embodiments of this paragraph, one of X^9 can be deuterium. In any embodiments of this paragraph, four of X^9 can be fluoro. In any embodiments of this paragraph, three of X^9 can be fluoro. In any embodiments of this paragraph, two of X^9 can be fluoro. In any embodiments of this paragraph, one of X^9 can be fluoro. In any embodiments of this paragraph, four of X^9 can be chloro. In any embodiments of this paragraph, three of X^9 can be chloro. In any embodiments of this paragraph, two of X^9 can be chloro. In any embodiments of this paragraph, one of X^9 can be chloro.


[0132] In some embodiments of Formula (In), R^{5F} can be hydrogen. In other embodiments, R^{5F} can be deuterium. In still other embodiments, R^{5F} can be an unsubstituted C_{1-4} alkyl, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0133] In some embodiments of Formula (In), p_{11} can be 0. In other embodiments, each X^L can be deuterium, and p_{11} can be 6. In still other embodiments, each X^L can be deuterium, and p_{11} can be 5. In yet still other embodiments, each X^L can be deuterium, and p_{11} can be 4. In some embodiments, each X^L can be deuterium, and p_{11} can be 3. In other embodiments, each X^L can be deuterium, and p_{11} can be 2. In still other embodiments, each X^L can be deuterium, and p_{11} can be 1. In some embodiments, each X^L can be chloro, and p_{11} can be 6. In other embodiments, each X^L can be chloro, and p_{11} can be 5. In still other embodiments, each X^L can be chloro, and p_{11} can be 4. In yet still other embodiments, each X^L can be chloro, and p_{11} can be 3. In some embodiments, each X^L can be chloro, and p_{11} can be 2. In other embodiments, each X^L can be chloro, and p_{11} can be 1. In some embodiments, each X^L can be fluoro, and p_{11} can be 6. In other embodiments, each X^L can be fluoro, and p_{11} can be 5. In still other embodiments, each X^L can be fluoro, and p_{11} can be 4. In yet still other embodiments, each X^L can be fluoro, and p_{11} can be 3. In some embodiments, each X^L can be fluoro, and p_{11} can be 2. In other embodiments, each X^L can be fluoro, and p_{11} can be 1.

[0134] In some embodiments of Formula (In), Z^2 can be nitrogen. In other embodiments, Z^2 can be oxygen. In still other embodiments, Z^2 can be sulfur.

Formula (Io)

[0135] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Io), or pharmaceutically acceptable salt thereof, having the structure:



[0136] In some embodiments of Formula (Io), X^{10} can be deuterium, fluoro or chloro; R^{9a} and R^{9b} can be independently an unsubstituted C_{1-4} alkyl; R^{5G} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^M can be independently deuterium, chloro or fluoro; and p_{12} can be independently 0, 1, 2, 3, 4, 5 or 6.

[0137] In some embodiments of Formula (Io), R^{9a} can be methyl. In other embodiments, R^{9a} can be ethyl. In still other embodiments, R^{9a} can be propyl, such as n-propyl or iso-propyl. In yet still other embodiments, R^{9a} can be butyl, for example, n-butyl, iso-butyl or tert-butyl. In some embodiments, including those of this paragraph, R^{9b} can be methyl. In other embodiments, including those of this paragraph, R^{9b} can be ethyl. In still other embodiments, including those of this paragraph, R^{9b} can be propyl, such as n-propyl or iso-propyl. In yet still other embodiments, including those of this paragraph, R^{9b} can be butyl, for example, n-butyl, iso-butyl or tert-butyl.

[0138] In some embodiments of Formula (Io), X^{10} can be deuterium. In other embodiments, X^{10} can be fluoro. In still other embodiments, X^{10} can be chloro.


[0139] In some embodiments of Formula (Io), R^{5G} can be hydrogen. In other embodiments, R^{5G} can be deuterium. In still other embodiments, R^{5G} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0140] In some embodiments of Formula (Io), p_{12} can be 0. In other embodiments, each X^M can be deuterium, and p_{12} can be 6. In still other embodiments, each

X^M can be deuterium, and p12 can be 5. In yet still other embodiments, each X^M can be deuterium, and p12 can be 4. In some embodiments, each X^M can be deuterium, and p12 can be 3. In other embodiments, each X^M can be deuterium, and p12 can be 2. In still other embodiments, each X^M can be deuterium, and p12 can be 1. In some embodiments, each X^M can be chloro, and p12 can be 6. In other embodiments, each X^M can be chloro, and p12 can be 5. In still other embodiments, each X^M can be chloro, and p12 can be 4. In yet still other embodiments, each X^M can be chloro, and p12 can be 3. In some embodiments, each X^M can be chloro, and p12 can be 2. In other embodiments, each X^M can be chloro, and p12 can be 1. In some embodiments, each X^M can be fluoro, and p12 can be 6. In other embodiments, each X^M can be fluoro, and p12 can be 5. In still other embodiments, each X^M can be fluoro, and p12 can be 4. In yet still other embodiments, each X^M can be fluoro, and p12 can be 3. In some embodiments, each X^M can be fluoro, and p12 can be 2. In other embodiments, each X^M can be fluoro, and p12 can be 1.

Formula (Ip)

[0141] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ip), or pharmaceutically acceptable salt thereof, having the structure:



[0142] In some embodiments of Formula (Ip), each X^{11} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{11} is fluoro or chloro; R^{5H} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^N can be independently deuterium, chloro or fluoro; and p13 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0143] In some embodiments of Formula (Ip), each X^{11} can be fluoro. In some embodiments, each X^{11} can be chloro. In some embodiments, two of X^{11} can be fluoro, and one of X^{11} can be chloro. In some embodiments, two of X^{11} can be chloro, and one of X^{11} can be fluoro. In some embodiments, two of X^{11} can be fluoro, and one of X^{11} can be


hydrogen. In some embodiments, two of X^{11} can be fluoro, and one of X^{11} can be deuterium. In some embodiments, two of X^{11} can be fluoro, and one of X^{11} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, two of X^{11} can be chloro, and one of X^{11} can be hydrogen. In some embodiments, two of X^{11} can be chloro, and one of X^{11} can be deuterium. In some embodiments, two of X^{11} can be chloro, and one of X^{11} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0144] In some embodiments of Formula (Ip), R^{5H} can be hydrogen. In other embodiments, R^{5H} can be deuterium. In still other embodiments, R^{5H} can be an unsubstituted C_{1-4} alkyl, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0145] In some embodiments of Formula (Ip), p13 can be 0. In other embodiments, each X^N can be deuterium, and p13 can be 6. In still other embodiments, each X^N can be deuterium, and p13 can be 5. In yet still other embodiments, each X^N can be deuterium, and p13 can be 4. In some embodiments, each X^N can be deuterium, and p13 can be 3. In other embodiments, each X^N can be deuterium, and p13 can be 2. In still other embodiments, each X^N can be deuterium, and p13 can be 1. In some embodiments, each X^N can be chloro, and p13 can be 6. In other embodiments, each X^N can be chloro, and p13 can be 5. In still other embodiments, each X^N can be chloro, and p13 can be 4. In yet still other embodiments, each X^N can be chloro, and p13 can be 3. In some embodiments, each X^N can be chloro, and p13 can be 2. In other embodiments, each X^N can be chloro, and p13 can be 1. In some embodiments, each X^N can be fluoro, and p13 can be 6. In other embodiments, each X^N can be fluoro, and p13 can be 5. In still other embodiments, each X^N can be fluoro, and p13 can be 4. In yet still other embodiments, each X^N can be fluoro, and p13 can be 3. In some embodiments, each X^N can be fluoro, and p13 can be 2. In other embodiments, each X^N can be fluoro, and p13 can be 1.

Formula (Iq)

[0146] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Iq), or pharmaceutically acceptable salt thereof, having the structure:



[0147] In some embodiments of Formula (Iq), each X^{12} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{12} is fluoro or chloro; R^{5J} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^P can be independently deuterium, chloro or fluoro; and p_{14} can be independently 0, 1, 2, 3, 4, 5 or 6.

[0148] In some embodiments of Formula (Iq), each X^{12} can be fluoro. In some embodiments, each X^{12} can be chloro. In some embodiments, two of X^{12} can be fluoro, and one of X^{12} can be chloro. In some embodiments, two of X^{12} can be chloro, and one of X^{12} can be fluoro. In some embodiments, two of X^{12} can be fluoro, and one of X^{12} can be hydrogen. In some embodiments, two of X^{12} can be fluoro, and one of X^{12} can be deuterium. In some embodiments, two of X^{12} can be fluoro, and one of X^{12} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, two of X^{12} can be chloro, and one of X^{12} can be hydrogen. In some embodiments, two of X^{12} can be chloro, and one of X^{12} can be deuterium. In some embodiments, two of X^{12} can be chloro, and one of X^{12} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0149] In some embodiments of Formula (Iq), R^{5J} can be hydrogen. In other embodiments, R^{5J} can be deuterium. In still other embodiments, R^{5J} can be an unsubstituted C_{1-4} alkyl, such as, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0150] In some embodiments of Formula (Iq), p_{14} can be 0. In other embodiments, each X^P can be deuterium, and p_{14} can be 6. In still other embodiments, each X^P can be deuterium, and p_{14} can be 5. In yet still other embodiments, each X^P can be

deuterium, and p14 can be 4. In some embodiments, each X^P can be deuterium, and p14 can be 3. In other embodiments, each X^P can be deuterium, and p14 can be 2. In still other embodiments, each X^P can be deuterium, and p14 can be 1. In some embodiments, each X^P can be chloro, and p14 can be 6. In other embodiments, each X^P can be chloro, and p14 can be 5. In still other embodiments, each X^P can be chloro, and p14 can be 4. In yet still other embodiments, each X^P can be chloro, and p14 can be 3. In some embodiments, each X^P can be chloro, and p14 can be 2. In other embodiments, each X^P can be chloro, and p14 can be 1. In some embodiments, each X^P can be fluoro, and p14 can be 6. In other embodiments, each X^P can be fluoro, and p14 can be 5. In still other embodiments, each X^P can be fluoro, and p14 can be 4. In yet still other embodiments, each X^P can be fluoro, and p14 can be 3. In some embodiments, each X^P can be fluoro, and p14 can be 2. In other embodiments, each X^P can be fluoro, and p14 can be 1.

Formula (Ir)

[0151] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Ir), or pharmaceutically acceptable salt thereof, having the structure:



[0152] In some embodiments of Formula (Ir), each X^{13} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{13} is fluoro or chloro; R^{5K} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^Q can be independently deuterium, chloro or fluoro; and p15 can be independently 0, 1, 2, 3, 4, 5 or 6.

[0153] In some embodiments of Formula (Ir), each X^{13} can be fluoro. In some embodiments, each X^{13} can be chloro. In some embodiments, two of X^{13} can be fluoro, and one of X^{13} can be chloro. In some embodiments, two of X^{13} can be chloro, and one of X^{13} can be fluoro. In some embodiments, two of X^{13} can be fluoro, and one of X^{13} can be hydrogen. In some embodiments, two of X^{13} can be fluoro, and one of X^{13} can be deuterium.


In some embodiments, two of X^{13} can be fluoro, and one of X^{13} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, two of X^{13} can be chloro, and one of X^{13} can be hydrogen. In some embodiments, two of X^{13} can be chloro, and one of X^{13} can be deuterium. In some embodiments, two of X^{13} can be chloro, and one of X^{13} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0154] In some embodiments of Formula (Ir), R^{5K} can be hydrogen. In other embodiments, R^{5K} can be deuterium. In still other embodiments, R^{5K} can be an unsubstituted C_{1-4} alkyl, for example, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0155] In some embodiments of Formula (Ir), p15 can be 0. In other embodiments, each X^Q can be deuterium, and p15 can be 6. In still other embodiments, each X^Q can be deuterium, and p15 can be 5. In yet still other embodiments, each X^Q can be deuterium, and p15 can be 4. In some embodiments, each X^Q can be deuterium, and p15 can be 3. In other embodiments, each X^Q can be deuterium, and p15 can be 2. In still other embodiments, each X^Q can be deuterium, and p15 can be 1. In some embodiments, each X^Q can be chloro, and p15 can be 6. In other embodiments, each X^Q can be chloro, and p15 can be 5. In still other embodiments, each X^Q can be chloro, and p15 can be 4. In yet still other embodiments, each X^Q can be chloro, and p15 can be 3. In some embodiments, each X^Q can be chloro, and p15 can be 2. In other embodiments, each X^Q can be chloro, and p15 can be 1. In some embodiments, each X^Q can be fluoro, and p15 can be 6. In other embodiments, each X^Q can be fluoro, and p15 can be 5. In still other embodiments, each X^Q can be fluoro, and p15 can be 4. In yet still other embodiments, each X^Q can be fluoro, and p15 can be 3. In some embodiments, each X^Q can be fluoro, and p15 can be 2. In other embodiments, each X^Q can be fluoro, and p15 can be 1.

Formula (Is)

[0156] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (Is), or pharmaceutically acceptable salt thereof, having the structure:



[0157] In some embodiments of Formula (Is), each X^{14} can be independently hydrogen, deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of X^{14} is fluoro or chloro; R^{5L} can be hydrogen, deuterium or an unsubstituted C_{1-4} alkyl; each X^R can be independently deuterium, chloro or fluoro; p_{16} can be independently 0, 1, 2, 3, 4, 5 or 6; and Z^3 can be nitrogen, oxygen or sulfur.

[0158] In some embodiments of Formula (Is), each X^{14} can be fluoro. In some embodiments, each X^{14} can be chloro. In some embodiments, two of X^{14} can be fluoro, and one of X^{14} can be chloro. In some embodiments, two of X^{14} can be chloro, and one of X^{14} can be fluoro. In some embodiments, two of X^{14} can be fluoro, and one of X^{14} can be hydrogen. In some embodiments, two of X^{14} can be fluoro, and one of X^{14} can be deuterium. In some embodiments, two of X^{14} can be fluoro, and one of X^{14} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl. In some embodiments, two of X^{14} can be chloro, and one of X^{14} can be hydrogen. In some embodiments, two of X^{14} can be chloro, and one of X^{14} can be deuterium. In some embodiments, two of X^{14} can be chloro, and one of X^{14} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

[0159] In some embodiments of Formula (Is), R^{5L} can be hydrogen. In other embodiments, R^{5L} can be deuterium. In still other embodiments, R^{5L} can be an unsubstituted C_{1-4} alkyl. Examples of C_{1-4} alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0160] In some embodiments of Formula (Is), p_{16} can be 0. In other embodiments, each X^R can be deuterium, and p_{16} can be 6. In still other embodiments, each

X^R can be deuterium, and p16 can be 5. In yet still other embodiments, each X^R can be deuterium, and p16 can be 4. In some embodiments, each X^R can be deuterium, and p16 can be 3. In other embodiments, each X^R can be deuterium, and p16 can be 2. In still other embodiments, each X^R can be deuterium, and p16 can be 1. In some embodiments, each X^R can be chloro, and p16 can be 6. In other embodiments, each X^R can be chloro, and p16 can be 5. In still other embodiments, each X^R can be chloro, and p16 can be 4. In yet still other embodiments, each X^R can be chloro, and p16 can be 3. In some embodiments, each X^R can be chloro, and p16 can be 2. In other embodiments, each X^R can be chloro, and p16 can be 1. In some embodiments, each X^R can be fluoro, and p16 can be 6. In other embodiments, each X^R can be fluoro, and p16 can be 5. In still other embodiments, each X^R can be fluoro, and p16 can be 4. In yet still other embodiments, each X^R can be fluoro, and p16 can be 3. In some embodiments, each X^R can be fluoro, and p16 can be 2. In other embodiments, each X^R can be fluoro, and p16 can be 1.

[0161] In some embodiments of Formula (Is), Z^3 can be nitrogen. In other embodiments, Z^3 can be oxygen. In still other embodiments, Z^3 can be sulfur.

Formula (It)

[0162] In some embodiments, the compound, or pharmaceutically acceptable salt thereof, can be a compound of Formula (It), or pharmaceutically acceptable salt thereof, having the structure:



[0163] In some embodiments of Formula (It), R^{10} can be selected from hydrogen, deuterium, halogen, hydroxy, an unsubstituted C_{1-8} alkyl, an unsubstituted C_{3-20} cycloalkyl and an unsubstituted C_{1-8} haloalkyl; R^{11} can be selected from hydrogen, deuterium, halogen, an unsubstituted C_{1-30} alkyl, an unsubstituted C_{2-30} alkenyl and an unsubstituted C_{1-8} haloalkyl; each X^S can be independently deuterium, chloro or fluoro; and p17 can be 0, 1, 2, 3, 4, 5 or 6.

[0164] In some embodiments of Formula (It), R^{10} can be selected from hydrogen, deuterium, halogen, hydroxy, an unsubstituted C_{1-8} alkyl, an unsubstituted C_{3-20} cycloalkyl

and an unsubstituted C₁₋₈ haloalkyl; R¹¹ can be selected from hydrogen, deuterium, halogen, an unsubstituted C₁₋₃₀ alkyl, an unsubstituted C₂₋₃₀ alkenyl and an unsubstituted C₁₋₈ haloalkyl; each X^S can be independently deuterium, chloro or fluoro; and p17 can be 1, 2, 3, 4, 5 or 6.

[0165] In some embodiments of Formula (It), R¹⁰ can be hydrogen. In other embodiments, R¹⁰ can be deuterium. In still other embodiments, R¹⁰ can be halogen, such as fluoro, chloro, bromo or iodo. In yet still other embodiments, R¹⁰ can be hydroxy. In still other embodiments, R¹⁰ can be an unsubstituted C₁₋₈ alkyl. Examples of suitable C₁₋₈ alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl (straight and branched), hexyl (straight and branched), heptyl (straight and branched) and octyl (straight and branched). In yet still other embodiments, R¹⁰ can be an unsubstituted C₃₋₂₀ cycloalkyl. In even still other embodiments, R¹⁰ can be an unsubstituted C₃₋₈ cycloalkyl. In some embodiments, R¹⁰ can be an unsubstituted C₁₋₈ haloalkyl. Examples of suitable C₁₋₈ haloalkyls include, but are not limited to, CF₃, CHF₂, CH₂F, CH₂CF₃, CH₂CHF₂ and CH₂CH₂F.


[0166] In some embodiments of Formula (It), R¹¹ can be hydrogen. In other embodiments, R¹¹ can be deuterium. In still other embodiments, R¹¹ can be halogen, such as fluoro, chloro, bromo or iodo. In yet still other embodiments, R¹¹ can be an unsubstituted C₁₋₈ haloalkyl. Examples of suitable C₁₋₈ haloalkyls include, but are not limited to, CF₃, CHF₂, CH₂F, CH₂CF₃, CH₂CHF₂ and CH₂CH₂F.

[0167] When R¹¹ is an alkyl group, the number of carbon atoms of the alkyl group can vary. The alkyl group can be a long alkyl having 1 to 30 carbons, a medium alkyl having 1 to 12 carbon atoms or a lower alkyl having 1 to 6 carbon atoms. In some embodiments, R¹¹ can be an unsubstituted C₁₋₆ alkyl. In other embodiments, R¹¹ can be an unsubstituted C₂₋₆ alkyl. In still other embodiments, R¹¹ can be an unsubstituted C₁₋₈ alkyl. In still other embodiments, R¹¹ can be an unsubstituted C₁₋₁₂ alkyl. In yet still other embodiments, R¹¹ can be an unsubstituted C₁₋₃₀ alkyl. In some embodiments, R¹¹ can be an unsubstituted C₇₋₂₆ alkyl. In other embodiments, R¹¹ can be an unsubstituted C₁₂₋₂₆ alkyl. Examples of lower alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, a tert-butyl, pentyl (straight and branched) and hexyl (straight and branched). Examples of suitable C₁₋₈ alkyl

groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl (straight and branched), hexyl (straight and branched), heptyl (straight and branched) and octyl (straight and branched). Examples of unsubstituted C₁₋₃₀ alkyls include, but are not limited to, -(CH₂)₆CH₃, -(CH₂)₈CH₃, -(CH₂)₁₀CH₃, -(CH₂)₁₂CH₃, -(CH₂)₁₄CH₃, -(CH₂)₁₆CH₃, -(CH₂)₁₈CH₃, -(CH₂)₂₀CH₃, -(CH₂)₂₂CH₃ and -(CH₂)₂₄CH₃.

[0168] Similar to alkyls, alkenyls can be a long alkenyl having 2 to 30 carbons, a medium alkenyl having 2 to 12 carbon atoms or a lower alkenyl having 2 to 6 carbon atoms. In some embodiments, R¹¹ can be an unsubstituted C₂₋₃₀ alkenyl. In other embodiments, R¹¹ can be an unsubstituted C₂₋₁₆ alkenyl. In still other embodiments, R¹¹ can be an unsubstituted C₁₈ alkenyl. In yet still other embodiments, R¹¹ can be an unsubstituted C₂₀₋₃₀ alkenyl. Examples of unsubstituted C₂₋₃₀ alkenyls include, but are not limited to, -(CH₂)₇CH=CH(CH₂)₃CH₃, -(CH₂)₇CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₇CH=CH(CH₂)₇CH₃, -(CH₂)₇CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₇CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃, -(CH₂)₉CH=CH(CH₂)₅CH₃, -(CH₂)₃CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₁₁CH=CH(CH₂)₇CH₃, -(CH₂)₃CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃, -(CH₂)₄CH=CHCH(CH₃)₂ and -(CH₂)₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃.

[0169] In some embodiments, R¹¹ can be the aliphatic tail of a saturated or an unsaturated fatty acid. As an example, R¹¹ can be the aliphatic tail of caprylic acid (HOOC**(CH₂)₆CH₃**). In this example of caprylic acid, the aliphatic tail is bolded and italicized. When the saturated or an unsaturated fatty acid becomes part of a compound of Formula (It), the carbon of the carboxylic acid of the saturated or an unsaturated fatty acid becomes the carbon of **C**(=O)R¹¹ that is bold and underlined. For example, when R¹¹ is the aliphatic tail of caprylic acid, the compound of Formula (It) can have the following structure:



. A non-limiting list of suitable saturated or an unsaturated fatty acids are myristoleic acid, palmitoleic, sapienic acid, linoleic acid, oleic acid, linoleiaidic acid, elaidic acid, alpha-linolenic acid, vaccenic acid, arachidonic acid, erucic acid, eicosapentaenoic acid, (E)-8-methylnon-6-enoic acid, docosahexaenoic acid, caprylic acid,


capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, arachidic acid, behenic acid, lignoceric acid and cerotic acid.

[0170] In some embodiments of Formula (It), p17 can be 0. In other embodiments, each X^S can be deuterium, and p17 can be 6. In still other embodiments, each X^S can be deuterium, and p17 can be 5. In yet still other embodiments, each X^S can be deuterium, and p17 can be 4. In some embodiments, each X^S can be deuterium, and p17 can be 3. In other embodiments, each X^S can be deuterium, and p17 can be 2. In still other embodiments, each X^S can be deuterium, and p17 can be 1. In some embodiments, each X^S can be chloro, and p17 can be 6. In other embodiments, each X^S can be chloro, and p17 can be 5. In still other embodiments, each X^S can be chloro, and p17 can be 4. In yet still other embodiments, each X^S can be chloro, and p17 can be 3. In some embodiments, each X^S can be chloro, and p17 can be 2. In other embodiments, each X^S can be chloro, and p17 can be 1. In some embodiments, each X^S can be fluoro, and p17 can be 6. In other embodiments, each X^S can be fluoro, and p17 can be 5. In still other embodiments, each X^S can be fluoro, and p17 can be 4. In yet still other embodiments, each X^S can be fluoro, and p17 can be 3. In some embodiments, each X^S can be fluoro, and p17 can be 2. In other embodiments, each X^S can be fluoro, and p17 can be 1.

[0171] In some embodiments of Formula (It), when p17 is 0, then R¹⁰ cannot be hydrogen. In other embodiments, when p17 is 0, then R¹⁰ cannot be deuterium. In still other embodiments, when p17 is 0, then R¹⁰ cannot be halogen. In yet still other embodiments, when p17 is 0, then R¹⁰ cannot be hydroxy. In some embodiments, when p17 is 0, then R¹⁰ cannot be an unsubstituted C₁₋₈ alkyl. In other embodiments, when p17 is 0, then R¹⁰ cannot be an unsubstituted C₃₋₂₀ cycloalkyl, or an unsubstituted C₃₋₈ cycloalkyl. In still other embodiments, when p17 is 0, then R¹⁰ cannot be an unsubstituted C₁₋₈ haloalkyl. In some embodiments, when p17 is 0, then R¹¹ cannot be hydrogen. In other embodiments, when p17 is 0, then R¹¹ cannot be deuterium. In still other embodiments, when p17 is 0, then R¹¹ cannot be halogen. In some embodiments, when p17 is 0, then R¹¹ cannot be an unsubstituted C₁₋₃₀ alkyl, such as those described in this paragraph. In other embodiments, when p17 is 0, then R¹¹ cannot be an aliphatic tail of a saturated fatty acid. In some embodiments, when p17 is 0, then R¹¹ cannot be an unsubstituted C₂₋₃₀ alkenyl, such as those

described in this paragraph. In other embodiments, when p17 is 0, then R¹¹ cannot be an aliphatic tail of an unsaturated fatty acid. In still other embodiments, when p17 is 0, then R¹¹ cannot be an unsubstituted C₁₋₈ haloalkyl. In some embodiments, R¹⁰ cannot be hydrogen. In other embodiments, R¹⁰ cannot be an unsubstituted C₁₋₆ alkyl. In still other embodiments, R¹⁰ cannot be an unsubstituted C₃₋₁₀ cycloalkyl. In yet still other embodiments, R¹⁰ cannot be halogen. In even still other embodiments, R¹⁰ cannot be an unsubstituted C₁₋₈ haloalkyl. In some embodiments, R¹¹ cannot be an unsubstituted C₁₋₆ alkyl. In other embodiments, R¹¹ cannot be an unsubstituted C₁₋₈ haloalkyl. In still other embodiments, R¹¹ cannot be an unsubstituted C₁₇ alkenyl. In yet still other embodiments, R¹¹ cannot be an unsubstituted C₁₉ alkenyl. Examples of suitable halogen groups include fluoro, chloro, bromo and iodo. Examples of unsubstituted C₁₋₃₀ alkyls include, but are not limited to, -(CH₂)₆CH₃, -(CH₂)₈CH₃, -(CH₂)₁₀CH₃, -(CH₂)₁₂CH₃, -(CH₂)₁₄CH₃, -(CH₂)₁₆CH₃, -(CH₂)₁₈CH₃, -(CH₂)₂₀CH₃, -(CH₂)₂₂CH₃ and -(CH₂)₂₄CH₃. Examples of unsubstituted C₂₋₃₀ alkenyls include, but are not limited to, -(CH₂)₇CH=CH(CH₂)₃CH₃, -(CH₂)₇CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₇CH=CH(CH₂)₇CH₃, -(CH₂)₇CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₇CH=CH(CH₂)₇CH₃, -(CH₂)₇CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃, -(CH₂)₉CH=CH(CH₂)₅CH₃, -(CH₂)₃CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CH(CH₂)₄CH₃, -(CH₂)₁₁CH=CH(CH₂)₇CH₃, -(CH₂)₃CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃, -(CH₂)₄CH=CHCH(CH₃)₂ and -(CH₂)₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CHCH₂CH₃. Examples of suitable haloalkyl groups include, but are not limited to, chloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 1-chloro-2-fluoromethyl and 2-fluoroisobutyl. Examples of suitable C₃₋₂₀ cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl. Examples of suitable C₁₋₈ alkyl groups include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl (straight and branched), hexyl (straight and branched), heptyl (straight and branched) and octyl (straight and branched). Examples of suitable C₁₇ alkenyl groups include, but are not limited to, -(CH₂)₇CH=CH(CH₂)₇CH₃. Examples of suitable C₁₉ alkenyl groups include, but are not limited to, -(CH₂)₃CH=CHCH₂CH=CHCH₂CH=CHCH₂CH=CH(CH₂)₄CH₃. In some

embodiments, a compound of Formula (It), or a pharmaceutically acceptable salt thereof, cannot one or more of the following compounds,



acceptable salt of any of the foregoing.

[0172] In some embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), at least two of each of X^1 , X^2 , X^3 , X^4 , X^5 , X^6 , X^7 , X^{11} , X^{12} , X^{13} and X^{14} , respectively, cannot be fluoro. In other embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), three of each of X^1 , X^2 , X^3 , X^4 , X^5 , X^6 , X^7 , X^{11} , X^{12} , X^{13} and X^{14} , respectively, cannot be fluoro. In still other embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), at least two of each of X^1 , X^2 , X^3 , X^4 , X^5 , X^6 , X^7 , X^{11} , X^{12} , X^{13} and X^{14} , respectively, cannot be chloro. In yet still other embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), three of each of X^1 , X^2 , X^3 , X^4 , X^5 , X^6 , X^7 , X^{11} , X^{12} , X^{13} and X^{14} , respectively, cannot be chloro. In some embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), each of

$X^1, X^2, X^3, X^4, X^5, X^6, X^7, X^{11}, X^{12}, X^{13}$ and X^{14} , respectively, are independently deuterium, an unsubstituted C_{1-4} alkyl, fluoro or chloro, provided that at least two of each of $X^1, X^2, X^3, X^4, X^5, X^6, X^7, X^{11}, X^{12}, X^{13}$ and X^{14} , respectively, are fluoro or chloro. In other embodiments of Formulae (Ia), (Ib), (Id), (If), (Ig), (Ih), (Ik), (Ip), (Iq), (Ir) and (Is), each of $X^1, X^2, X^3, X^4, X^5, X^6, X^7, X^{11}, X^{12}, X^{13}$ and X^{14} , respectively, are independently hydrogen, deuterium, fluoro or chloro, provided that at least two of each of $X^1, X^2, X^3, X^4, X^5, X^6, X^7, X^{11}, X^{12}, X^{13}$ and X^{14} , respectively, are fluoro or chloro.

[0173] In some embodiments of Formulae (Ic), (Id), (Ie), (If), (Im), (In), (Io), (Ip), (Iq), (Ir) and (Is), each of $R^{5A}, R^{5B}, R^{5C}, R^{5D}, R^{5E}, R^{5F}, R^{5G}, R^{5H}, R^{5J}, R^{5K}$ and R^{5L} , respectively, cannot be hydrogen. In other embodiments of Formulae (Ic), (Id), (Ie), (If), (Im), (In), (Io), (Ip), (Iq), (Ir) and (Is), each of $R^{5A}, R^{5B}, R^{5C}, R^{5D}, R^{5E}, R^{5F}, R^{5G}, R^{5H}, R^{5J}, R^{5K}$ and R^{5L} , respectively, cannot be an unsubstituted C_{1-4} alkyl, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.


[0174] In some embodiments of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and (It), each of $p1, p2, p3, p4, p5, p6, p7, p8, p9, p10, p11, p12, p13, p14, p15, p16$ and $p17$, respectively, cannot be 0.


[0175] In some embodiments of Formula (Ie), when R^{4a} is $-C(X^{16})_3$, then at least two of X^{16} cannot be fluoro. In other embodiments of Formula (Ie), when R^{4a} is $-C(X^{16})_3$, then $-C(X^{16})_3$ cannot be $-CF_3$. In still other embodiments of Formula (Ie), when R^{4a} is $-C(X^{16})_3$, then at least two of X^{16} cannot be chloro. In yet still other embodiments of Formula (Ie), when R^{4a} is $-C(X^{16})_3$, then $-C(X^{16})_3$ cannot be $-CCl_3$. In some embodiments of Formula (Ie), when R^{4b} is $-C(X^{16})_3$, then at least two of X^{16} cannot be fluoro. In other embodiments of Formula (Ie), when R^{4b} is $-C(X^{16})_3$, then $-C(X^{16})_3$ cannot be $-CF_3$. In still other embodiments of Formula (Ie), when R^{4b} is $-C(X^{16})_3$, then at least two of X^{16} cannot be chloro. In yet still other embodiments of Formula (Ie), when R^{4b} is $-C(X^{16})_3$, then $-C(X^{16})_3$ cannot be $-CCl_3$.

[0176] In some embodiments of Formulae (Im), (In) and (Io), each X^8 , each X^9 and X^{10} , respectively, can be deuterium or chloro. In other embodiments of Formulae (Im), (In) and (Io), each X^8 , each X^9 and X^{10} , respectively, can be deuterium or fluoro. In still other embodiments, more than one of each X^8 , each X^9 and X^{10} , respectively, is fluoro. For

example, 2, 3, 4, or 5 of X^8 , X^9 and X^{10} , respectively, is fluoro. In yet still other embodiments, more than one of each X^8 , each X^9 and X^{10} , respectively, is chloro (such as 2, 3, 4, or 5 of X^8 , X^9 and X^{10} , respectively, is chloro).. In some embodiments, the rings to which X^8 and X^9 are respectively attached cannot be fully fluorinated. In other embodiments, the rings to which X^8 and X^9 are respectively attached cannot be fully chlorinated. In some embodiments, X^{10} cannot be fluoro. In other embodiments, X^{10} cannot be chloro.

[0177] In some embodiments, the compound cannot be selected from:





and , or a pharmaceutically acceptable salt of any of the foregoing.

Methods

[0178] The various compounds contemplated herein can be obtained from known starting materials by various routes known to those skilled in the art. Some suitable routes are illustrated in the Examples and following references: Radchenko et al., *Journal of Organic Chemistry* (2010) 75:5941-5952; U.S. Patent Publication No. 2008/0287468 (filed 11 October 2007); WO 2002/059083 (filed 23 October 2001); Nisato et al. *Journal of Heterocyclic Chemistry* (1985) 22:961-963; WO 2005/000810 (filed 22 June 2004); WO 2007/036733 (filed 29 September 2006); Lewin et al., *Journal of Medicinal Chemistry* (1998) 41:988-995; WO 2010/017047 (filed 27 July 2009); WO 2013/033059 (filed 28 August 2012). Salts can be formed using methods known to those skilled in the art and described herein, for example, reacting an amine with a suitable acid (such as HCl). In some embodiments, the pharmaceutically acceptable salt of a compound described herein (for example, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It)) can be the HCl salt.

Pharmaceutical Compositions

[0179] Some embodiments described herein relate to a pharmaceutical composition, that can include an effective amount of one or more compounds described herein (for example, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing) and a pharmaceutically acceptable carrier, diluent, excipient or combination thereof.

[0180] The term “pharmaceutical composition” refers to a mixture of one or more compounds disclosed herein with other chemical components, such as diluents or carriers. The pharmaceutical composition facilitates administration of the compound to an organism. Pharmaceutical compositions can also be obtained by reacting compounds with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, and salicylic acid. Pharmaceutical compositions will generally be tailored to the specific intended route of administration.

[0181] The term “physiologically acceptable” defines a carrier, diluent or excipient that does not abrogate the biological activity and properties of the compound nor cause appreciable damage or injury to an animal to which delivery of the composition is intended.

[0182] As used herein, a “carrier” refers to a compound that facilitates the incorporation of a compound into cells or tissues. For example, without limitation, dimethyl sulfoxide (DMSO) is a commonly utilized carrier that facilitates the uptake of many organic compounds into cells or tissues of a subject.

[0183] As used herein, a “diluent” refers to an ingredient in a pharmaceutical composition that lacks appreciable pharmacological activity but may be pharmaceutically necessary or desirable. For example, a diluent may be used to increase the bulk of a potent drug whose mass is too small for manufacture and/or administration. It may also be a liquid for the dissolution of a drug to be administered by injection, ingestion or inhalation. A common form of diluent in the art is a buffered aqueous solution such as, without limitation, phosphate buffered saline that mimics the pH and isotonicity of human blood.

[0184] As used herein, an “excipient” refers to an essentially inert substance that is added to a pharmaceutical composition to provide, without limitation, bulk, consistency, stability, binding ability, lubrication, disintegrating ability etc., to the composition. A “diluent” is a type of excipient.

[0185] The pharmaceutical compositions described herein can be administered to a human patient *per se*, or in pharmaceutical compositions where they are mixed with other active ingredients, as in combination therapy, or carriers, diluents, excipients or combinations

thereof. Proper formulation is dependent upon the route of administration chosen. Techniques for formulation and administration of the compounds described herein are known to those skilled in the art.

[0186] The pharmaceutical compositions disclosed herein may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or tableting processes. Additionally, the active ingredients are contained in an amount effective to achieve its intended purpose. Many of the compounds used in the pharmaceutical combinations disclosed herein may be provided as salts with pharmaceutically compatible counterions.

[0187] Multiple techniques of administering a compound exist in the art including, but not limited to, oral, rectal, pulmonary, topical, aerosol, injection, infusion and parenteral delivery, including intramuscular, subcutaneous, intravenous, intramedullary injections, intrathecal, direct intraventricular, intraperitoneal, intranasal and intraocular injections.

[0188] One may also administer the compound in a local rather than systemic manner, for example, via injection or implantation of the compound directly into the affected area, often in a depot or sustained release formulation. Furthermore, one may administer the compound in a targeted drug delivery system, for example, in a liposome coated with a tissue-specific antibody. The liposomes will be targeted to and taken up selectively by the organ. For example, intranasal or pulmonary delivery to target a respiratory infection may be desirable.

[0189] The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied with a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, may be the labeling

approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. Compositions that can include a compound described herein formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

Methods of Use

[0190] Some embodiments provided herein relate to a method of treating a disease or condition that can include administering to a subject an effective amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing. Other embodiments provided herein relate to a method of treating a disease or condition that can include contacting a cell in the central and/or peripheral nervous system of a subject with an effective amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing. In some embodiments, the subject can be at risk of developing a disease or condition that is responsive to acetaminophen and/or a NSAID. In some embodiments, the disease or condition can be one or more of the following: pain, fever, inflammation, ischemic injury (such as myocardial and/or cerebral) and/or neuronal injury. In some embodiments, the subject can be post-operative and has, or is believed to have or has actually developed post-operative pain. In some embodiments, the subject can be in need of treatment for acute pain and has, is believed to have or has actually developed acute pain. In some embodiments, the subject can be in need of treatment for chronic pain and has, is believed to have or has actually developed chronic pain. In some embodiments, the subject can be in need of treatment for neuropathic pain and has, is believed to have or has actually developed neuropathic pain. The basis for determining the need for treatment can be based on an underlying condition or conditions, from indication by the subject and/or on other bases known to practitioners. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided (such as administered) prophylactically, for example, prophylactically for pain (such as post-operative pain).

[0191] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can contact a cell in the central nervous system, for example, the brain and/or spinal cord, and thereby treat a disease or condition described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can contact a cell in the peripheral nervous system, for example, the ganglia and/or nervous system outside the brain and spinal cord, and thereby treat a disease or condition described herein.

[0192] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can contact a TRP (transient receptor potential) channels modulator (such as TRPV1 and/or TRPA1), and thereby treat a disease or condition described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can contact a cannabinoid receptors modulator (such as CB1 and/or CB2), and thereby treat a disease or condition described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can contact a serotonin receptor (for example, 5HT1, 5HT2, 5HT3, 5HT4, 5HT5, 5HT6 and/or 5HT7) and modulate its activity, and thereby treat a disease or condition described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can act as an anandamide reuptake inhibitor, and thereby treat a disease or condition described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be a substrate for the fatty acid amide hydrolase (FAAH), and thereby treat a disease or condition described herein.

[0193] Some embodiments described herein generally relate to a method of treating pain of any etiology, including acute pain and chronic and neuropathic pain, and any pain in which acetaminophen is prescribed. Examples of pain include post-surgical pain; post-operative pain (including dental pain); migraine; headache and trigeminal neuralgia; pain associated with burn, wound and/or kidney stone; pain associated with trauma (including traumatic head injury); neuropathic pain (e.g., central and peripheral pain); pain associated with musculo- skeletal disorders; strains; sprains; contusions; fractures; myalgia; nociceptive pain (for example, rheumatoid arthritis and osteoarthritis pain); cystitis; visceral pain (such as, pancreatitis, inflammatory bowel disease and internal organ pain); ankylosing spondylitis; sero-negative (non-rheumatoid) arthropathies; non-articular rheumatism and peri-articular disorders; and mixed pain. Central pain includes post-stroke pain, pain associated with multiple sclerosis, spinal cord injury, migraine and HIV-related neuropathic pain. Peripheral pain includes post-herpetic neuralgia and diabetic neuropathy. Mixed pain includes pain associated with cancer (including "break-through pain" and pain associated with terminal cancer), lower back and fibromyalgia. Examples of pain with an inflammatory component (in addition to some of those described above) include rheumatic pain, pain associated with mucositis and pain associated with dysmenorrhea. In some embodiments, a method and/or a composition described herein can be used for treating or preventing post-surgical pain. In some embodiments, a method and/or a composition described herein can be used for treating or preventing cancer pain or pain associated with a subject having cancer. In some embodiments, a method and/or a composition described herein can be used for treating or preventing osteoarthritis and/or rheumatoid arthritis pain. In some embodiments, a method and/or a composition described herein can be used for treating or preventing migraine pain. In some embodiments, a method and/or a composition described herein can be used for treating or preventing lower back pain and/or fibromyalgia pain. In some embodiments, a method and/or a composition described herein can be used for treating or preventing pain that is selected from pain associated with surgery, trauma, osteoarthritis, rheumatoid arthritis, lower back pain, fibromyalgia, postherpetic neuralgia, diabetic neuropathy, HIV-associated neuropathy and complex regional pain syndrome. Additionally information regarding pain is

provided in Melnikova, I., "Pain market" (2010) 9(8):589-590, which is hereby incorporated by reference in its entirety.

[0194] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be used for treating or preventing pain and/or a fever (for example, in adults, children and/or infants, and in animal health to treat animals such as the cat, dog, or horse). As used herein, an "infant" is a human that is 1 year old or younger, a "child" is a human >1 to 17 years old, and an "adult" is a human 18 years or older. Compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be used to treat a variety and varying degrees of pain. In some embodiments, the pain can be acute pain (for example, acute pain following surgery, such as orthopedic surgery of adults, children, and/or infants). In some embodiments, the pain can be chronic pain (for example, pain lasting days, weeks, months, or years, and optionally following an initial event, such as an injury, trauma, surgery, or onset of disease).

[0195] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be used for treating and/or preventing a fever, such as endotoxin-induced fever (for example, endotoxin-induced fever in adults, children, and/or infants). In some embodiments, the fever can be selected from low-grade fever, moderate fever, high-grade fever and hyperpyrexia fever. In some embodiments, the fever can be selected from Pel-Ebstein fever, continuous fever, intermittent fever and remittent fever.

[0196] As described herein, compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be used in various subjects. In some embodiments, the subject can be a child and/or an infant, for example, a child or infant with a fever. In other embodiments, the subject can be an adult. In other embodiments, the subject can be an animal such as a cat, dog, or horse. As described herein, compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or

pharmaceutically acceptable salts of any of the foregoing, can be administered by physicians and/or veterinarians as appropriate.

[0197] Some embodiments described herein relate to a method of delaying the onset of analgesia in a subject in need thereof, wherein the method can include administering to the subject an effective amount of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It) , or pharmaceutically acceptable salts of any of the foregoing, that delays drug action by greater than 5 minutes, or 10 minutes, or 15 minutes, or 30 minutes, or 1 hour, or 2, hours, or 3 hours, or 4 hours, or 6 hours, or 8 hours, or 10 hours, or 12 hours, or 18 hours or 24 hours. Other embodiments described herein relate to a method of delaying the onset of analgesia in a subject in need thereof, wherein the method can include contacting a cell in the central and/or peripheral nervous system of a subject with an effective amount of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, that delays drug action by greater than 5 minutes, or 10 minutes, or 15 minutes, or 30 minutes, or 1 hour, or 2, hours, or 3 hours, or 4 hours, or 6 hours, or 8 hours, or 10 hours, or 12 hours, or 18 hours or 24 hours.

[0198] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, may provide greater reduction or prevention of pain than acetaminophen in the early/acute phase (0-10 minutes). In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, may provide greater reduction or prevention of pain than acetaminophen in the late/tonic phase (10-35 minutes).

[0199] As described herein, there can be a variety of advantages of using a compound described herein. Examples of such advantages include, but are not limited to, improved effectiveness, tolerance by a subject's body, inflammatory analgesic properties, osteoarthritis analgesic properties, incisional analgesic properties and neuropathic analgesic properties. Compounds described herein can have a longer metabolic half-life, for example, a longer metabolic half-life compared to APAP when both APAP and compounds described

herein are administered by the same route and concentration. Additional advantage can include comparable or improved properties, such as those described herein, while minimizing and/or avoiding one or more side effects associated with NSAIDs, APAP and opioids. For example, a person of ordinary skill in the art would appreciate that the compounds described herein can avoid one or more undesirable side effects associated with inhibition of COX receptors and/or activation of opiate receptors, can have a lower likelihood for abuse or addiction compared to opioids, and can have a lower likelihood for loss of analgesic potency compared to opioids. Further advantages include lower dosages of compounds described herein, wherein the lower dosages achieves comparable or improved properties, such as inflammatory analgesic properties, osteoarthritis analgesic properties, incisional analgesic properties and/or neuropathic analgesic properties, compared to the same property achieved using a known NSAID, APAP or opioid. For example, a compound described herein can have comparable or improved osteoarthritis analgesic properties compared to a NSAID, such as celecoxib.

[0200] Furthermore, a person of ordinary skill in the art would appreciate that the compounds described would avoid many of the negative side effects associated with the NAPQI metabolite of acetaminophen. As described herein, acetaminophen can form the reactive metabolite, N-acetyl-p-benzoquinone imine (NAPQI), which is linked to liver toxicity. Acetaminophen is metabolically activated by cytochrome P450 enzymes to form NAPQI, and NAPQI depletes endogenous glutathione (GSH). The depletion of endogenous glutathione leaves cells vulnerable to oxidative damage. The formation of NAPQI is the result of the phenol or aniline ring of acetaminophen. Unlike acetaminophen, the compounds of the present application do not include a phenol or aniline ring and it is impossible to connect a substituent through a double bond (such as a carbonyl or imine group) at either end of bicyclo[1.1.1]pentane (i.e., at the 1 or 3 positions). As a result, one skilled in the art would not expect compounds described herein to form the reactive metabolite NAPQI.

[0201] As described herein, compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be administered by a variety of methods. In any of the methods described herein, administration can be by injection, infusion and/or intravenous

administration over the course of 1 minute, 5 minutes, 10 minutes, 30 minutes, 1 hour, 2 hours, 6 hours, 12 hours, 24 hours or longer, or any intermediate time. Such administration can, in some circumstances, substitute for or significantly reduce the need for administration of an opiate. Some methods described herein can include intravenous administration to a subject in need thereof, for example, to a subject to manage post-operative or other acute or chronic pain, in either a bolus dose or by infusion over minutes, hours or days. Other methods described herein can include oral, intravenous, subcutaneous and/or intraperitoneal administration to a subject in need thereof, for example, to a subject to manage post-operative or other acute pain or chronic pain.

[0202] Other embodiments described herein relate to a method for selecting a therapy for managing or treating pain in a subject in need thereof, that can include evaluating whether the subject is at risk for hepatic toxicity from pain therapy, and selecting a therapy that includes a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, to reduce or eliminate such risk. The method can further include providing the selected therapy that includes a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, to the subject. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be of significant benefit in pain management in hospitals or other care facilities (for example, a nursing home).

[0203] As used herein, the terms “prevent” and “preventing,” mean a subject does not experience and/or develop pain and/or fever. Examples of forms of prevention include prophylactic administration to a subject who is going to undergo surgery.

[0204] As used herein, the terms “treat,” “treating,” “treatment,” “therapeutic,” and “therapy” do not necessarily mean total cure or abolition of the disease or condition. Any alleviation of any undesired signs or symptoms of a disease or condition, to any extent can be considered treatment and/or therapy, for example the severity of the pain and/or fever is less compared to the severity of the pain and/or fever if the subject has not been

administered/received the compound. Furthermore, treatment may include acts that may worsen the subject's overall feeling of well-being or appearance.

[0205] The terms “therapeutically effective amount” and “effective amount” are used to indicate an amount of an active compound, or pharmaceutical agent, that elicits the biological or medicinal response indicated. For example, a therapeutically effective amount of compound can be the amount needed to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. This response may occur in a tissue, system, animal or human and includes alleviation of the signs or symptoms of the disease being treated. Determination of an effective amount is well within the capability of those skilled in the art, in view of the disclosure provided herein. The therapeutically effective amount of the compounds disclosed herein required as a dose will depend on the route of administration, the type of animal, including human, being treated, and the physical characteristics of the specific animal under consideration. The dose can be tailored to achieve a desired effect, but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

[0206] The amount of the compound, or an active salt or derivative thereof, required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician. In cases of administration of a pharmaceutically acceptable salt, dosages may be calculated as the free base. As will be understood by those of skill in the art, in certain situations it may be necessary to administer the compounds disclosed herein in amounts that exceed, or even far exceed, the above-stated, preferred dosage range in order to effectively and aggressively treat particularly aggressive diseases or conditions.

[0207] In general, however, a suitable dose will often be in the range of from about 0.15 mg/kg to about 100 mg/kg. For example, a suitable dose may be in the range from about 1 mg/kg to about 75 mg/kg of body weight per day, such as about 0.75 mg/kg to about 50 mg/kg of body weight of the recipient per day, about 1 mg/kg to 90 mg/kg of body weight of the recipient per day, or about 10 mg/kg to about 60 mg/kg of body weight of the recipient per day.

[0208] The compound may be administered in unit dosage form; for example, containing 1 to 2000 mg, 10 to 1000 mg or 5 to 500 mg of active ingredient per unit dosage form.

[0209] The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations.

[0210] As will be readily apparent to one skilled in the art, the useful *in vivo* dosage to be administered and the particular mode of administration will vary depending upon the age, weight, the severity of the affliction, and mammalian species treated, the particular compounds employed, and the specific use for which these compounds are employed. The determination of effective dosage levels, that is the dosage levels necessary to achieve the desired result, can be accomplished by one skilled in the art using routine methods, for example, human clinical trials, *in vivo* studies and *in vitro* studies. For example, useful dosages of compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts thereof, can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Such comparison can be done against an established analgesic drug, such as acetaminophen.

[0211] Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the modulating effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vivo* and/or *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations. Dosage intervals can also be determined using MEC value. Compositions should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

[0212] It should be noted that the attending physician would know how to and when to terminate, interrupt, or adjust administration due to toxicity or organ dysfunctions.

Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency, will also vary according to the age, body weight, and response of the individual patient. A program comparable to that discussed above may be used in animal health and veterinary medicine.

[0213] Compounds disclosed herein can be evaluated for efficacy and toxicity using known methods. For example, the toxicology of a particular compound, or of a subset of the compounds, sharing certain chemical moieties, may be established by determining *in vitro* toxicity towards a cell line, such as a mammalian, and preferably human, cell line. The results of such studies are often predictive of toxicity in animals, such as mammals, or more specifically, humans. Alternatively, the toxicity of particular compounds in an animal model, such as mice, rats, rabbits, dogs or monkeys, may be determined using known methods. The efficacy of a particular compound may be established using several recognized methods, such as *in vitro* methods, animal models, or human clinical trials. When selecting a model to determine efficacy, the skilled artisan can be guided by the state of the art to choose an appropriate model, dose, route of administration and/or regime.

Combination Drugs

[0214] One or more compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided alone or in combination with another drug(s). In some embodiments, the other drug(s) can be an opioid analgesic. Any of the known opioid analgesics can be combined with a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing. As non-limiting examples, such opioid analgesics include morphine, codeine, hydrocodone, oxycodone, fentanyl, pethidine, methadone, pentazocine, sufentanil, levorphanol, dihydrocodeine, nalbuphine, butorphanol, tramadol, meptazinol, buprenorphine,

dipipanone, alfentanil, remifentanil, oxymorphone, tapentadol, propoxyphene and hydromorphone.

[0215] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided in a dosage form (for example, an oral dosage form, an intravenous dosage form and/or an intraperitoneal dosage form), in combination with one of the following exemplary opioids: 1-20 mg hydrocodone (such as hydrocodone bitartrate), preferably 2.5 mg, 5 mg, 7.5 mg or 10 mg of hydrocodone or salt thereof; or 1-20 mg oxycodone, preferably 2.5 mg, 5 mg, 7.5 mg or 10 mg of oxycodone or salt thereof (such as the hydrochloride salt). In some embodiments, the amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be in the range of about 20 to about 2000 mg, for example 20 ± 0.5 to 2000 ± 5 mg.

[0216] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be combined with one or more non-steroidal anti-inflammatory drugs (NSAIDs). As non-limiting examples, such NSAIDs include celecoxib, ketorolac, ketoprofen, indomethacin, sulindac, etodolac, mefenamic acid, meclofenamic acid, meclofenamate sodium, flufenamic acid, tolmetin, diclofenac, diclofenac sodium, ibuprofen, naproxen, naproxen sodium, fenoprofen, flurbiprofen, oxaprozin, piroxicam, meloxicam, ampiroxicam, droxicam, lornoxicam, cinnoxicam, sudoxicam, and tenoxicam, and pharmaceutically acceptable salts of the foregoing. In some embodiments, an NSAID can be a COX-2 inhibitor.

[0217] In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided in a dosage form (for example, an oral dosage form, an intravenous dosage form and/or an intraperitoneal dosage form), in combination with one of the following exemplary NSAIDs: 10-1000 mg ibuprofen, for example 100 mg, 250 mg, 500 mg or 750 mg of ibuprofen or salt thereof; 100-1000 mg naproxen, for example 100 mg, 250 mg, 500 mg or 750 mg of naproxen or salt thereof (such

as the sodium salt); 100-1000 mg ketorolac, for example 100 mg, 250 mg, 500 mg or 750 mg of ketorolac or salt thereof; 100-1000 mg ketoprofen, for example 100 mg, 250 mg, 500 mg or 750 mg of ketoprofen or salt thereof; or 10-1000 mg celecoxib, for example 100 mg, 250 mg, 500 mg or 750 mg of celecoxib or salt thereof. In some embodiments, the amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be in the range of about 20 to about 2000 mg, for example 20 ± 0.5 to 2000 ± 5 mg.

[0218] Other combinations include combination of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, with butalbital, codeine, dihydrocodeine, and/or aspirin. The other drug(s) can be provided using routes known to those skilled in the art and/or described herein. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, and another drug(s) can be provided in the same dosage form. In other embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, and another drug(s) can be provided in the separate dosage forms. In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, and another drug(s) can be by the same route (for example, both intravenously) or by different routes (for example, one orally and the other intraperitoneally). In some embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided before another drug(s) (such as an opiate). In other embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided simultaneously with another drug(s) (such as an opiate). In still other embodiments, a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is)

and/or (It), or pharmaceutically acceptable salts of any of the foregoing, can be provided after another drug(s) (such as an opiate).

[0219] In some embodiments, a combination of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, and an opioid analgesic can synergistically relieve pain. In some embodiments, the synergistic relief of pain can reduce opioid use. Some embodiments disclosed herein relate to a method of managing, treating and/or reducing pain that can include administering an effective amount of a combination of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, and an opioid analgesic to a subject. Some embodiments disclosed herein relate to a method for reducing opioid use in pain management, that can include administering an amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, in combination with an amount of an opioid analgesic, wherein the amount of the opioid analgesic in the combination is less than the amount of opioid analgesic needed to achieve approximately the same level of pain management when the opioid analgesic is administered alone. Methods known for evaluating pain management is known to those skilled in the art, for example, pain assessment tools. Some embodiments disclosed herein relate to a method for decreasing the risk of opioid dependency that can include administering an amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, in combination with an amount of an opioid analgesic, wherein the amount of the opioid analgesic in the combination is less than the amount of opioid analgesic needed to achieve approximately the same level of pain management when the opioid analgesic is administered alone. Some embodiments disclosed herein relate to a method for treating pain and/or fever along with treating opioid dependency that can include administering an amount of a compound of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and/or (It), or pharmaceutically acceptable salts of any of the foregoing, in combination with an amount of an opioid analgesic.

EXAMPLES

[0220] Additional embodiments are disclosed in further detail in the following examples, which are not in any way intended to limit the scope of the claims.

COMPOUNDS


[0221] The compounds of Formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and (It), and pharmaceutically acceptable salts of any of the foregoing, are provided illustrated in Table 1 can be prepared in various ways, including those techniques described herein and/or known to those skilled in the art as guided by the details provided herein. For example, the compounds of Formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and (It), and pharmaceutically acceptable salts of any of the foregoing, illustrated in Table 1 can be prepared as described in the Examples below. Those skilled in the art will understand that a number of structures shown in Table 1 are not stereospecific and/or are depicted as having unfilled valencies, and thus are generic to isotopic and/or stereochemical variants, including racemates, diastereomers, enantiomers and/or deuterated versions, which can be prepared in accordance with the guidance provided herein.

TABLE 1

No.	Compound Structure	No.	Compound Structure
1		2	
3		4	
5		6	
7		8	

No.	Compound Structure	No.	Compound Structure
9		10	
11		12	
13		14	
15		16	
17		18	
19		20	
21		22	
23		24	
25		26	
27			

EXAMPLE 1

3-(4-(Trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (1)

[0222] 3-(Methoxycarbonyl)bicyclo-[1.1.1]pentane-1-carboxylic acid (70.0g, 411mmol) was added to a dry 1L RBF and dissolved in 400mL DCM under N₂. Oxalyl chloride (34.8mL, 411mmol) was added followed by 1 drop of dry DMF. The reaction was stirred overnight at room temperature (rt) where it was determined to be complete by NMR. The solvent was removed *in vacuo*. 100mL of dry toluene was added and removed *in vacuo*. Methyl 3-(chlorocarbonyl)bicyclo[1.1.1]pentane-1-carboxylate (78.0g, 411mmol) was dissolved in a total of 400mL DCM and transferred to an addition funnel via cannula. 200mL DCM was added under N₂ to a dry 2L RBF. The DCM was cooled to 0°C and NH₃ was bubbled through the solution for 10 minutes. The DCM solution of methyl 3-(chlorocarbonyl)bicyclo[1.1.1]pentane-1-carboxylate was added slowly to the DCM NH₃ solution. NH₃ was bubbled through the suspension during the addition and for 30 minutes after the addition was complete. The ice bath was removed and the reaction was stirred for 2h where it was determined to be complete by NMR. The solvent was removed and the solid was stirred in 600mL Et₂O for 1h and then filtered. The solid was dried to give 60.9g (88%) of methyl 3-carbamoylbicyclo[1.1.1]pentane-1-carboxylate **1-1** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.36 (s, 1H), 7.03 (s, 1H), 3.60 (s, 3H), 2.12 (s, 6H).

[0223] Compound **1-1** (35.0g, 207mmol) was suspended in 400mL THF. Phosphorus pentasulfide (22.9g, 103mmol) was added and the reaction was stirred for 12h where it was determined to be complete by crude NMR. The solid was removed by filtration. The precipitate was re-suspended in THF and filtered again. The filtrate was concentrated and the residue was absorbed onto Florisil® and purified by flash chromatography (SiO₂, EtOAc:Hexanes) to afford an off white solid. The solid was suspended in Et₂O with stirring, filtered and dried under high vacuum to afford 19g (49%) of carbamothioylbicyclo[1.1.1]pentane-1-carboxylate **1-2** as a white solid. LC/MS (APCI) *m/z* 186.0 [C₈H₁₁NO₂S+H]⁺.

[0224] Compound **1-2** (5.11g, 27.6mmol) and NaHCO₃ (6.95g, 83mmol) was suspended in 20mL EtOH. 3-Bromo-1,1,1-trifluoropropan-2-one (7.90g, 41.4mmol) in 20mL

EtOH was added and the reaction was stirred at 80°C for 2h where the reaction was determined to be complete by LCMS. The solvent was removed and the residue was partitioned between 75mL water and 75mL EtOAc. The layers were separated and the water layer was extracted with EtOAc (2x75mL). The combined organics were dried (Na₂SO₄) and concentrated to give methyl 3-(imino((3,3,3-trifluoro-2-oxopropyl)thio)methyl)bicyclo[1.1.1]pentane-1-carboxylate **1-3** as an off white solid.

[0225] Compound **1-3** (8.15g, 27.6mmol) was dissolved in 40mL DCM and Et₃N (7.69mL, 55.2mmol) was added under N₂. The flask was cooled to 0°C and TFAA (4.99mL, 35.9mmol) was added dropwise. The bath was removed and the reaction was stirred to rt over 2h where it was determined to be complete by LCMS. The reaction was concentrated and dissolved in 150mL EtOAc. The organic layer was washed with saturated NaHCO₃ (1x100mL), brine (1x100mL), and dried (Na₂SO₄). The solvent was removed to give 8g of an orange solid. The compound was recrystallized from 50mL of heptane to give 6.12g (80%) of methyl 3-(4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **1-4** as an off white solid. LC/MS (APCI) *m/z* 278.0 [C₁₁H₁₀F₃NO₂S+H]⁺.

[0226] Compound **1-4** (6.10g, 22.0mmol) was dissolved in 22mL THF followed by 22mL 2M NaOH. The reaction was stirred for 2h where it was determined to be complete by LCMS. The THF was removed and the solution was cooled to 0°C and the pH was adjusted to ~3 with 2M HCl. The mixture was allowed to stand in the refrigerator overnight and then filtered. The off-white solid was dissolved in EtOAc, dried (Na₂SO₄), and the solvent removed. The compound was dried overnight under high vacuum to give 5.19g (90%) of 3-(4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **1-5** as an off white solid. LC/MS (APCI) *m/z* 264.0 [C₁₀H₈F₃NO₂S+H]⁺.


[0227] Compound **1-5** (5.19g, 19.7mmol) was dissolved in 66mL of a 1:1 ratio of toluene:t-BuOH. Et₃N (5.50mL, 39.4mmol) and DPPA (4.67mL, 21.7mmol) were added under N₂. The reaction was stirred for 1h and 8g of 4Å molecular sieves were added. The reaction was stirred for 3h at rt and then stirred at 100°C overnight. The reaction was cooled to rt and filtered through Celite rinsing with EtOAc. The solvent was removed until the compound was an off-white solid. The compound was dissolved in 80mL EtOAc and washed with saturated NH₄Cl (50mL), saturated NaHCO₃ (50mL), brine (50mL) and dried

(Na₂SO₄). The solvent was removed to give 6.6g of a beige solid that was suspended in 90mL heptane and heated gently until all of the solid dissolved. The solution was carefully boiled down to reduce the solvent volume. Upon cooling, the solid was removed by filtration. The compound was dried under high vacuum overnight to give 5.89g (89%) of tert-butyl (3-(4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **1-6** as an off-white solid. LC/MS (APCI) *m/z* 335.1 [C₁₄H₁₇F₃N₂O₂S+H]⁺.

[0228] Compound **1-6** (5.52g, 16.5mmol) was dissolved in 25mL DCM. TFA (12.6mL, 165mmol) was added via syringe and the reaction was stirred for 1.5h where it was determined to be complete by LCMS. The reaction was concentrated and 12mL of 2M HCl in Et₂O was added. The mixture was stirred for 5 minutes and then the solvent was removed. The HCl addition was repeated two times. HCl (2N in Et₂O, 12mL) was added followed by 20mL Et₂O. The mixture was stirred vigorously and filtered. The solid was dried in the high vacuum oven overnight to give 4g of an off white solid. The compound was dissolved in 10 mL of EtOH and approximately 5 mL of EtOH was boiled away. The mixture was cooled to rt and the resulting crystals were filtered and rinsed with MTBE. The compound was dried overnight under high vacuum to give 3.41g (76%) of Compound **1** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.03 (s, 3H), 8.49-8.48 (s, 1H) 2.46 (s, 6H); LC/MS (APCI) *m/z* 235.0 [C₉H₉F₃N₂S+H]⁺.

EXAMPLE 2

3-(5-Methyl-4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (2)



[0229] A suspension of methyl 3-carbamothioylbicyclo[1.1.1]pentane-1-carboxylate (8.45g, 45.6mmol) and NaHCO₃ (5.75g, 68.4mmol) in anhydrous EtOH (45.6mL) was treated with 3-bromo-1,1,1-tributan-2-one (16.7mL, 137mmol). The resulting suspension was heated to 80°C until complete (~2h). The mixture was cooled to rt and concentrated to afford the crude intermediate as a semi-solid. The crude methyl 3-(imino((4,4,4-trifluoro-3-oxobutan-2-yl)thio)methyl)bicyclo[1.1.1]pentane-1-carboxylate was dissolved in anhydrous DCM (76mL), cooled to 0°C and treated with Et₃N (12.7mL, 91.0mmol) and TFAA (8.38mL, 59.3mmol). The resulting solution was stirred at rt. After

stirring overnight, additional Et₃N (6.36mL, 45.6mmol) and TFAA (6.44mL, 45.6mmol) was added and stirring was continued for 2.5h. 1N HCl (100mL) was added and the mixture was extracted with DCM (4x100mL). The combined organics were dried (Na₂SO₄) and concentrated to provide the crude product which was further purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 13.3g (97%) of methyl 3-(5-methyl-4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **2-1** as a yellow oil. LC/MS (APCI) *m/z* 292.0 [C₁₂H₁₂F₃NO₂S+H]⁺.

[0230] A solution of Compound **2-1** (13.3g, 45.6mmol) in THF (73.0mL) and water (18.2mL) was treated with LiOH·H₂O (4.21g, 100mmol) and allowed to stir at rt until complete (1h). The solution was concentrated to 1/3 of the original volume, diluted with water (50mL) and washed with Et₂O (50mL). The aqueous layer was acidified with 1N HCl then extracted with EtOAc (4x100 mL). The combined organic layers were dried (Na₂SO₄) and concentrated to provide 11.7g (92%) of 3-(5-methyl-4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **2-2** as a white solid. LC/MS (APCI) *m/z* 278.0 [C₁₁H₁₀F₃NO₂S+H]⁺.


[0231] A solution of Compound **2-2** (11.6g, 41.8mmol) in anhydrous toluene (105mL) and tert-BuOH (105mL) was treated with activated 3Å mol sieves followed by Et₃N (11.7mL, 84.0mmol) and DPPA (10.8mL, 50.2mmol). The mixture was stirred at rt for 4h followed by heating to 90°C overnight. The mixture was cooled to rt and filter through a pad of Celite. The pad was washed with EtOAc and the filtrate was concentrated to provide the crude product as a viscous amber oil which was further purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 12.7g (87%) of tert-butyl (3-(5-methyl-4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **2-3** as a white solid. LC/MS (APCI) *m/z* 349.1 [C₁₅H₁₉F₃N₂O₂S+H]⁺.

[0232] A solution of Compound **2-3** (12.7g, 36.5mmol) in anhydrous DCM (91mL) was cooled to 0°C and treated with TFA (91mL). The resulting solution was stirred at rt until complete (1h). The solution was concentrated to provide a pale-yellow oil that was re-dissolved in anhydrous Et₂O (50mL) and treated with 2N HCl in Et₂O (20mL) while stirring. A white precipitate formed immediately and stirring became difficult. The mixture was further diluted with Et₂O to allow for efficient stirring then the mixture was concentrated

to 1/4 of the original volume. The salt exchange process was repeated. The white suspension in Et₂O was filtered and the product was collected and washed with copious amounts of Et₂O to provide 9.97g (96%) of Compound **2** as a white powder. ¹H NMR (400 MHz, DMSO-d₆) δ 8.93 (br s, NH, 3H), 2.57-2.52 (m, 3H), 2.41 (s, 6H); LC/MS (APCI) *m/z* 249.1 [C₁₀H₁₁F₃N₂S+H]⁺.

EXAMPLE 3

3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (3)



[0233] A suspension of ground Mg turnings (0.475g, 19.6mmol) in Et₂O (35.6mL) was treated with 1,2-dibromoethane (0.046mL, 0.533mmol) followed by dropwise addition of 1-bromo-4-(trifluoromethyl)benzene (2.49mL, 17.8mmol). Once formation was complete (~1h) as determined by consumption of Mg metal, the Grignard reagent was divided across three separate vials each containing a solution of propellane (18.0mL, 5.90mmol) at rt. The resulting mixtures were sealed and heated to 50°C for 3 days after which the suspensions were cooled to 0°C. CO₂ (dried over anhydrous CaSO₄) was bubbled through the mixtures for 5 minutes. The mixtures were warmed to rt and stirred for an additional 10 minutes then diluted with EtOAc (50mL) and water (20mL). The mixtures were acidified with 1N HCl (20mL) and extracted with EtOAc (3x50mL). The combined organics were dried (Na₂SO₄) and concentrated to provide the crude product which was purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 0.217g (5%) of 3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid **3-1** as a semi-pure yellow solid which contains residual 4-(trifluoromethyl)benzoic acid. The product was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.58 (d, *J*=8.2 Hz, 2H), 7.33 (d, *J*=8.1 Hz, 2H), 2.41 (s, 6H).


[0234] Compound **3-1** (0.216g, 0.843mmol) was dissolved in anhydrous toluene (2.11mL) and t-BuOH (2.11mL). The mixture was treated with powdered 3Å mol sieves, Et₃N (0.235mL, 1.69mmol), and DPPA (0.218mL, 1.01mmol). The resulting mixture was stirred at rt for 4h then heated to 90°C and allowed to stir overnight. The mixture was cooled and filtered through Celite. The mixture was diluted with EtOAc (100mL) and washed with

10% citric acid (20mL) then brine (20mL). The combined organics were dried (Na_2SO_4) and concentrated to afford the crude product which was further purified by flash chromatography (SiO_2 , Hexanes/EtOAc) to provide 0.091g (33%) of tert-butyl (3-(4-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)carbamate **3-2** as a white solid. LC/MS (APCI) m/z 228.0 [$\text{C}_{17}\text{H}_{20}\text{F}_3\text{NO}_2-\text{C}_5\text{H}_8\text{O}_2+\text{H}$] $^+$.

[0235] A solution of Compound **3-2** (0.090g, 0.275mmol) in EtOAc (1.375mL) was treated with 2N HCl (1.38mL, 2.75mmol) in Et_2O . The resulting solution was stirred at rt until complete. The resulting suspension was concentrated to dryness and the resulting solid was triturated and washed with Et_2O to provide 48mg (67%) of Compound **3** as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.74 (br s, 3H), 7.70 (d, $J=8.2$ Hz, 2H), 7.50 (d, $J=8.1$ Hz, 2H), 2.31 (s, 6H); LC/MS (APCI) m/z 228.0 [$\text{C}_{12}\text{H}_{12}\text{F}_3\text{N}+\text{H}$] $^+$.

EXAMPLE 4

3-(4-Methoxyphenyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (4)



[0236] A solution of propellane (20.4mL, 6.95mmol) in Et_2O (0.34M) was treated with a 1M solution of (4-chlorophenyl)magnesium bromide (6.95mL, 6.95mmol). The resulting mixture was heated to 50°C in a sealed vial and allowed to stir for 4 days. The mixture was cooled to 0°C and CO_2 (dried over anhydrous CaSO_4) was bubbled through the mixture for 5 minutes. The mixture was warmed to rt and stirred for an additional 10 minutes, diluted with EtOAc (50mL) and acidified with 1N HCl. The mixture was further diluted with H_2O (50mL) and extracted with EtOAc (3x50mL). The combined organic layers were dried (Na_2SO_4) and concentrated to provide the crude product which was purified by flash chromatography (SiO_2 , Hexanes/EtOAc) to afford 0.626g (41%) of 3-(4-chlorophenyl)bicyclo[1.1.1]pentane-1-carboxylic acid **4-1** as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.28 (d, $J=7.4$ Hz, 2H), 7.14 (d, $J=7.4$ Hz, 2H), 2.35 (s, 6H).

[0237] Compound **4-1** (413mg, 1.86mmol) was dissolved in *t*-BuOH (9.27mL). Activated $3\text{\AA}$ MS (500mg) were added followed by Et_3N (517 μL , 3.71mmol) and DPPA (480 μL , 2.23mmol). The resulting solution was stirred at 30°C for 4h, and then heated to reflux overnight. The solution was cooled to rt and then concentrated under reduced


pressure. The residual oil was diluted with EtOAc (50mL) and H₂O (50mL), and extracted with EtOAc (3x20mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product that was purified by flash chromatography (SiO₂, Hexanes/EtOAc) followed by RPLC (RediSep Rf Gold ® C18Aq, H₂O:ACN) to provide tert-butyl (3-(4-chlorophenyl)bicyclo[1.1.1]pentan-1-yl)carbamate **4-2** (300mg, 55%) as a white solid. LC/MS (APCI) *m/z* 194.0 [C₁₆H₂₀ClNO₂ – C₅H₉O+H]⁺.

[0238] To an oven dried vial was added anhydrous Cs₂CO₃ (0.250g, 0.766mmol), *t*-BuBrettPhos (4.95mg, 10.2μmol), and Compound **4-2** (0.150g, 0.511mmol). The vial was flushed with N₂ then anhydrous MeOH (0.103mL, 2.55mmol) was added. In a separate oven dried vial, *t*-BuBrettPhos Pd G3 (8.72mg, 10.2μmol) was added followed by dioxane (1.02mL). The suspension was stirred for ~1 minute to obtain a homogenous solution. The solution containing the Pd catalyst was then transferred to the vial containing the arylchloride and the resulting suspension was stirred at 50°C overnight. The mixture was cooled to rt, concentrated and directly purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 0.102g (69%) of tert-butyl (3-(4-methoxyphenyl)bicyclo[1.1.1]pentan-1-yl)carbamate as a solid **4-3**. LC/MS (APCI) *m/z* 188.0 [C₁₇H₂₃NO₃-C₅H₈O₂+H]⁺.

[0239] A solution of Compound **4-3** (0.138g, 0.477mmol) in EtOAc (2.38mL) was treated with 2N HCl in Et₂O (2.38mL, 4.77mmol). The resulting solution was stirred at rt until complete. Once complete, the resulting suspension was concentrated to dryness and the resulting solid was triturated and washed with Et₂O to provide 99mg (92%) of Compound **4** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.74-8.59 (br s, NH, 3H), 7.17 (d, *J*=8.7 Hz, 2H), 6.88 (d, *J*=8.6 Hz, 2H), 3.73 (s, 3H), 2.21 (s, 6H); LC/MS (APCI) *m/z* 173.1 [C₁₂H₁₅NO – NH₃+H]⁺.

EXAMPLE 5

3-(3-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (5)




[0240] A suspension of ground Mg turnings (0.178g, 7.33mmol) in Et₂O (6.67mL) was treated with 1,2-dibromoethane (0.017mL, 0.200mmol) followed by dropwise addition of 1-bromo-3-(trifluoromethyl)benzene (0.932mL, 6.67mmol) in 5mL of anhydrous

Et₂O. Once formation was complete (~1h) as determined by consumption of Mg metal, the Grignard reagent was added to a solution of propellane (18.2mL, 6.00mmol) at rt. The resulting mixture was sealed and heated to 50°C for 4 days after which the suspension was cooled to 0°C. CO₂ (dried over anhydrous CaSO₄) was bubbled through the mixture for 5 minutes. The mixture was warmed to rt and stirred for an additional 10 minutes then diluted with EtOAc (50mL) and water (20mL). The mixture was acidified with 1N HCl (20mL) and extracted with EtOAc (3x50mL). The combined organics were dried (Na₂SO₄) and concentrated to provide the crude product which was purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 0.272g (16%) of 3-(3-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentane-1-carboxylic acid **5-1** as a beige solid. ¹H NMR (400 MHz, CDCl₃) δ 7.52-7.51 (m, 1H), 7.47-7.35 (m, 3H), 2.40 (s, 6H).


[0241] A solution of Compound **5-1** (0.272g, 1.06mmol) in a mixture of toluene (2.66mL) and *t*-BuOH (2.66mL) was treated with powdered 3Å mol sieves, Et₃N (0.296mL, 2.13mmol), and DPPA (0.275mL, 1.28mmol). The resulting mixture was stirred at rt for 4 h then heated to 90°C and allowed to stir overnight. The mixture was cooled and filtered through Celite. The mixture was diluted with H₂O (50mL) and extracted with EtOAc (4x50mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to afford 0.173g (50%) of tert-butyl (3-(3-(trifluoromethyl)phenyl)bicyclo[1.1.1]pentan-1-yl)carbamate **5-2** as a soft yellow solid. LC/MS (APCI) *m/z* 128.1 [C₁₇H₂₀F₃NO-C₅H₈O₂+H]⁺.

[0242] A solution of Compound **5-2** (0.173g, 0.529mmol) in EtOAc (2.64mL) was treated with 2N HCl in Et₂O (2.64mL, 5.29mmol) and allowed to stir at rt until complete. Once complete, the resulting suspension was concentrated to dryness and the resulting solid was triturated and washed with Et₂O to provide 0.110g (79%) of Compound **5** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (br s, NH, 3H), 7.66-7.59 (m, 4H), 2.34 (s, 6H); LC/MS (APCI) *m/z* 128.1 [C₁₂H₁₂F₃N+H]⁺.

EXAMPLE 6(3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)glycine hydrochloride (6)

[0243] A solution of tert-butyl (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate (7.00g, 27.9mmol) in anhydrous DMF (55.7mL) was cooled to 0°C and NaH (1.67g, 41.8mmol) was added. The resulting solution was stirred for 10 minutes followed by the addition of tert-butyl bromoacetate (4.97mL, 33.4mmol). The resulting suspension was warmed to rt while stirring. Once complete (2h), the mixture was carefully quenched with 10% citric acid (~50mL) and diluted with H₂O (~50mL). The reaction was extracted with EtOAc (3x150mL) and the combined organics dried (Na₂SO₄), concentrated and the crude product purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 9.44g (93%) of tert-butyl *N*-(tert-butoxycarbonyl)-*N*-(3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)glycinate **6-1** as a viscous colorless oil which solidified to a white solid upon standing. LC/MS (APCI) *m/z* 266.1 [C₁₇H₂₆F₃NO₃-C₅H₈O+H]⁺.

[0244] A solution of Compound **6-1** (9.43g, 25.8mmol) in DCM (34.4mL) was treated with TFA (17.2mL) and allowed to stir at rt for 36h at which point additional TFA (10mL) was added. Stirring was resumed for an additional 24h. The solution was concentrated and the crude residue was purified by RP-HPLC (C₁₈ SiO₂, H₂O/ACN buffered with 0.1% formic acid) to afford the desired product as a white semi-solid. The material was suspended in Et₂O and treated with 2N HCl in ether to generate the HCl salt, concentrated, treated again with 2N HCl in Et₂O, and then concentrated again. The resulting solid was washed with Et₂O and dried to afford 5.07g (80%) of Compound **6** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.39 (br s, 2H), 3.84 (s, 2H), 2.27 (s, 6H); LC/MS (APCI) *m/z* 210.1 [C₈H₁₀F₃NO+H]⁺.

EXAMPLE 72-Hydroxy-*N*-(3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)acetamide (7)

[0245] A solution of 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (12.0g, 64.0mmol) in DMF (128mL) at 0°C was treated with 2-tertiary-butoxyacetic acid (8.23mL, 64.0mmol) and *N,N*-diisopropylethylamine (22.3mL, 128mmol). HATU (26.8g, 70.4mmol) was added and the resulting solution was stirred at rt overnight. The mixture was concentrated to approximately 1/3 of the original volume, diluted with H₂O (200mL) and extracted with EtOAc (3x150mL). The combined organics were washed with 10% citric acid (100mL), water (100mL) then brine (100mL). The organic layer was dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 14.0g (82%) of 2-(tert-butoxy)-*N*-(3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)acetamide **7-1** as a white solid. LC/MS (APCI) *m/z* 210.1 [C₁₂H₁₈F₃NO₂-C₄H₈+H]⁺.

[0246] A solution of Compound **7-1** (14.0g, 52.6mmol) in anhydrous DCM (70.2mL) was treated with TFA (35.1mL) and allowed to stir at rt overnight. The mixture was concentrated to afford the crude product which was further purified by flash chromatography (C₁₈ SiO₂, H₂O/ACN buffered with 0.1% formic acid) to provide 9.67g (88%) of Compound **7** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.52 (br s, NH, 1H), 5.46 (br s, OH, 1H), 3.76 (s, 2H), 2.23 (s, 6H); LC/MS (APCI) *m/z* 210.1 [C₈H₁₀F₃NO₂+H]⁺.

EXAMPLE 8

N-Ethyl-3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (8)




[0247] A solution of tert-butyl (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate (0.200g, 0.796mmol) in anhydrous DMF (3.98mL) was cooled to 0°C and NaH (0.048g, 1.194mmol) was added as a 60% dispersion in mineral oil. The resulting solution was stirred for 10 minutes followed by the addition of iodoethane (0.077mL, 0.955mmol). The resulting suspension was allowed to warm to rt and stirring was continued for 1.5h. The mixture was carefully quenched with 10% citric acid, diluted with H₂O (20mL) and extracted with EtOAc (4x20mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 0.125g (56%) of tert-butyl ethyl(3-

(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate **8-1** the desired product as a soft solid. LC/MS (APCI) m/z 180.1 [$C_{13}H_{20}F_3NO_2 \cdot C_5H_8O_2 + H$] $^+$.

[0248] A solution of Compound **8-1** (0.125g, 0.448mmol) in EtOAc (2.24mL) was treated with a 2N HCl solution in Et₂O (2.24mL, 4.48mmol) and allowed to stir at rt until complete. The mixture was concentrated and the resulting white solid triturated with Et₂O to provide 84.6mg (88%) of Compound **8** as white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.84 (br s, NH, 2H), 2.92 (q, $J=7.3$ Hz, 2H), 2.31 (s, 6H), 1.20 (t, $J=7.2$ Hz, 3H); LC/MS (APCI) m/z 180.1 [$C_8H_{12}F_3N + H$] $^+$.


EXAMPLE 9

2-((3-(Trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)amino)ethan-1-ol hydrochloride (**9**)




[0249] A solution of tert-butyl (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate (0.200g, 0.796mmol) in anhydrous DMF (3.98mL) was cooled to 0°C and NaH (0.0480g, 1.19mmol) was added as 60% dispersion in mineral oil. The resulting solution was stirred for 10 minutes followed by the addition of 2-bromoethoxy-*t*-butyldimethylsilane (0.205mL, 0.955mmol). The resulting suspension was allowed to warm to rt and stirred for an additional 1h. The mixture was carefully quenched with 10% citric acid, diluted with H₂O (20mL) and extracted with EtOAc (4x20 mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 0.196g (60%) of *N*-(2-((tert-butyldimethylsilyl)oxyethyl)-3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine **9-1** as a clear, colorless oil. LC/MS (APCI) m/z 310.2 [$C_{19}H_{34}F_3NO_3 \cdot C_5H_8O_2 + H$] $^+$.

[0250] A solution of Compound **9-1** (0.196g, 0.479mmol) in DCM (2mL) was treated with TFA (2mL) and allowed to stir at rt overnight. The solution was concentrated and the resulting residue was dissolved in Et₂O (~1mL). A 2N solution of HCl in Et₂O (1mL) was added to produce a white precipitate which was triturated with Et₂O to provide 104mg (94%) of Compound **9** as a white solid. ¹H NMR (400 MHz, D₂O with DDS Na salt as internal standard) δ 3.85-3.81 (m, 2H), 3.21-3.17 (m, 2H), 2.42 (s, 6H); LC/MS (APCI) m/z 196.1 [$C_8H_{12}F_3NO + H$] $^+$.

EXAMPLE 103-(4-Chlorophenyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (10)

[0251] To a solution of tert-butyl (3-(4-chlorophenyl)bicyclo[1.1.1]pentan-1-yl)carbamate (300mg, 1.02mmol) in EtOAc (5.10mL) was added HCl (2N in Et₂O, 5.11mL, 10.2mmol). The resulting solution was stirred for 2 days at rt. The suspension was concentrated, and the residual solid was triturated with Et₂O (2x20mL). The precipitate was collected by filtration, and the filter cake was washed with Et₂O (30mL). The white solid was dried under vacuum to afford Compound **10** (227mg, 97%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.87 (br s, 3H), 7.41-7.37 (m, 2H), 7.31-7.27 (m, 2H), 2.26 (s, 6H). LC/MS (APCI) *m/z* 194.0 [C₁₁H₁₂ClN+H]⁺.

EXAMPLE 113-(2-(Trifluoromethyl)pyrimidin-4-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (11)

[0252] A solution of 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (5.00g, 29.4mmol) in a mixture of toluene (73.5mL) and *t*-BuOH (73.5mL) was treated with powdered 3Å mol sieves, Et₃N (8.19mL, 58.8mmol) and DPPA (7.61mL, 35.3mmol). The resulting mixture was stirred at rt for 4h then heated to 90°C and allowed to stir overnight. The mixture was cooled and filtered through a pad of Celite. The mixture was diluted with H₂O (50mL) and extracted with EtOAc (4x50mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to afford 4.56g (64%) of methyl 3-((tert-butoxycarbonyl)amino)bicyclo[1.1.1]pentane-1-carboxylate **11-1** as a white solid. LC/MS (APCI) *m/z* 142.1 [C₁₂H₁₉NO₄-C₅H₈O₂+H]⁺.

[0253] To a solution of Compound **11-1** (3.05g, 12.6mmol) in THF (21.1mL), MeOH (21.1mL), and H₂O (21.1mL) at 0°C was added LiOH·H₂O (1.59g, 37.9mmol). The reaction was warmed to rt and stirred overnight. The reaction was concentrated *in vacuo* to remove THF and MeOH and then diluted with water to 50mL. The aqueous layer was

washed with Et₂O and then acidified to pH 3 with 1N HCl. The aqueous layer was extracted with EtOAc (3X50 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo* to provide 3-((tert-butoxycarbonyl)amino)bicyclo[1.1.1]pentane-1-carboxylic acid **11-2** (2.56g, 89%) as a white solid. LC/MS (APCI) *m/z* 128.0 [C₁₁H₁₇NO₄-C₅H₉O₂+H]⁺.

[0254] A solution of Compound **11-2** (1.01g, 4.44mmol) in Et₂O (22.2mL) was cooled to -5°C and treated with MeMgBr (3.0M in Et₂O, 4.90mL, 14.6mmol). After 30 min, the reaction was warmed to rt. After 16h, the reaction was cooled to 0°C and quenched with sat. aqueous NH₄Cl solution (5mL). Upon warming to rt the reaction was diluted with DCM and H₂O. The combined organic layers were dried (Na₂SO₄) and concentrated under reduced pressure to afford tert-butyl (3-acetylbicyclo[1.1.1]pentan-1-yl)carbamate **11-3** (522mg, 52%) as a white solid. LC/MS (APCI) *m/z* 126.1 [C₁₂H₁₉NO₃-C₅H₉O₂+H]⁺.


[0255] A solution of Compound **11-3** (376mg, 1.67mmol) in DMF (3.33mL) was treated with 1,1-dimethoxy-N,N-dimethylmethanamine (266μL, 2.00mmol) and heated to 100°C. After 2h, additional 1,1-dimethoxy-N,N-dimethylmethanamine (266μL, 2.00mmol) was added and the reaction was again heated to 100°C. After 2h, the reaction was concentrated *in vacuo* and then dissolved in EtOH (8.3mL) and treated with 2,2,2-trifluoroacetamide (319μL, 3.51mmol) and NaOEt (20% in EtOH, 1.37mL, 3.51mmol). The reaction mixture was heated to 110°C using a Biotage Initiator microwave reactor for 3h, cooled to rt and then concentrated *in vacuo*. The crude reaction mixture was partitioned between DCM and H₂O and then the aqueous layer was extracted with DCM (3x30mL). The combined organic layers were dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude product was purified by column chromatography to provide a yellow oil which was further purified by reverse phase column chromatography to provide tert-butyl (3-(2-(trifluoromethyl)pyrimidin-4-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **11-4** (168mg, 31%) as a white solid. LC/MS (APCI) *m/z* 230.0 [C₁₅H₁₈N₃O₂-C₅H₉O₂+H]⁺.

[0256] To a solution of Compound **11-4** (161mg, 0.490mmol) in EtOAc (2.4mL) was added HCl (2N in Et₂O, 2.93mL, 5.87mmol). The resulting solution was stirred for 2 days at rt. The suspension was concentrated, and the residual solid was triturated with Et₂O (2x20mL). The precipitate was collected by filtration, and the filter cake was washed with

Et₂O (30mL). The white solid was dissolved in 1:1 ACN:H₂O and lyophilized to obtain Compound **11** (116mg, 89%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.04 (d, *J*=5.1 Hz, 1H), 8.82 (s, 3H), 7.85 (d, *J*=5.1 Hz, 1H), 2.41 (s, 6H); ¹⁹F NMR (376 MHz, DMSO-*d*₆, unreferenced) δ -69.06; LC/MS (APCI) *m/z* 230.0 [C₁₀H₁₀F₃N₃+H]⁺.

EXAMPLE 12

3-(4-(Trifluoromethyl)pyrimidin-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (**12**)



[0257] Representative Procedure (reaction was performed in 3 parallel batches using a total of 1.99g, 12.69mmol, 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid): A solution of 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (727mg, 4.27mmol), 4-(trifluoromethyl)pyrimidine (561μL, 5.12mmol), H₂SO₄ (273μL, 5.12mmol), AgNO₃ (218mg, 1.28mmol) in a mixture of 1:1 H₂O/ACN (14.2 mL) was treated with sodium persulfate (712mg, 2.99mmol) and heated to 80°C. After 2h, the reaction was cooled to rt and the combined reaction mixtures were concentrated *in vacuo*. MeOH (50mL) was added followed by 3 drops of conc. H₂SO₄. The solution was heated to 70°C for 1h and then cooled to rt and filtered over Celite. The filtrate was concentrated to ~5mL and the solution was slowly added to a cooled solution of sat. aq. NaHCO₃ (100mL). The aqueous layer was then extracted with EtOAc (3x50mL). The combined organic layers were dried (Na₂SO₄), filtered and concentrated *in vacuo*. The crude product was purified by normal followed by reverse phase column chromatography to provide methyl 3-(4-(trifluoromethyl)pyrimidin-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **12-1** (181mg, 4%) as a white solid. LC/MS (APCI) *m/z* 273.0 [C₁₂H₁₁F₃N₂O₂+H]⁺.

[0258] A solution of Compound **12-1** (261mg, 0.959mmol) in 1:1 ACN:H₂O (4.8mL) was treated with HCl (3N in H₂O, 640μL, 1.92mmol) and heated to 75°C. After 90 min, the reaction was cooled to rt and then lyophilized to provide 3-(4-(trifluoromethyl)pyrimidin-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **12-2** (239mg, 97%) as a white solid. LC/MS (APCI) *m/z* 259.0 [C₁₁H₉F₃N₂O₂+H]⁺.


[0259] Compound **12-2** (239mg, 0.925mmol) was dissolved in *t*-BuOH (4.63mL). Activated 3Å MS (400mg) were added followed by Et₃N (387μL, 2.78mmol) and DPPA

(259 μ L, 1.20mmol). The resulting solution was stirred at 35°C for 4h, and then heated to 80°C overnight. The solution was cooled to rt and then concentrated under reduced pressure. The residual oil was diluted with EtOAc (50mL) and H₂O (50mL), and extracted with EtOAc (3x20mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product that was purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide tert-butyl (3-(4-(trifluoromethyl)pyrimidin-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **12-3** (119mg, 39%) as a white solid. LC/MS (APCI) *m/z* 230.0 [C₁₅H₁₈F₃N₃O₂-C₅H₉O₂+H]⁺.

[0260] A solution of Compound **12-3** (119mg, 0.360mmol) in anhydrous DCM (1.81mL) was cooled to 0°C and treated with TFA (278 μ L, 3.61mmol) at rt. After 1h, additional TFA was added. After an additional 30 min, the reaction was concentrated *in vacuo* and Et₂O (8mL) was added causing a white solid to crash out of solution. The heterogeneous solution was treated with HCl (2N in Et₂O, 3.0mL, 6.00mmol) and the reaction mixture was stirred for 2 min. The reaction mixture was concentrated *in vacuo* and the process was repeated an additional 2 times. The suspension was concentrated, and the residual solid was triturated with Et₂O. The precipitate was collected by filtration, and the filter cake was washed with Et₂O. The white solid was dissolved in 1:1 ACN:H₂O and lyophilized to obtain Compound **12** (90.1mg, 94%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆, 60 °C) δ 9.15 (d, *J*=5.1 Hz, 1H), 8.86 (s, 3H), 7.91 (d, *J*=5.1 Hz, 1H), 2.44 (s, 6H); ¹⁹F NMR (376 MHz, DMSO-*d*₆, 60°C, unreferenced) δ -68.65; LC/MS (APCI) *m/z* 230.0 [C₁₀H₁₀F₃N₃+H]⁺.

EXAMPLE 13

3-(6-(Trifluoromethyl)pyridin-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (13)



[0261] Representative Procedure (reaction was performed in 8 parallel batches using a total of 8.00g, 47.0mmol, 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid): A solution of 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (1.00g, 5.88mmol), 2-(trifluoromethyl)pyridine (0.900mL, 7.80mmol), H₂SO₄ (0.430mL, 7.80mmol), and AgNO₃ (200mg, 1.18mmol) in a mixture of 1:1 H₂O:ACN (19.8mL) was heated to 80°C and treated with sodium persulfate (1.40g, 5.88mmol) in water (5mL) dropwise. After 1h,

the reaction was cooled to rt and the combined reaction mixtures were concentrated *in vacuo*. MeOH (50mL) was added followed by 5 drops of conc. H₂SO₄. The solution was heated to 70°C for 1h and then cooled to rt and filtered over Celite. The filtrate was concentrated to ~5mL and the solution was slowly added to a cooled solution of sat aq. NaHCO₃ (100mL). The aqueous layer was then extracted with EtOAc (3x50mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by normal followed by reverse phase column chromatography to provide methyl 3-(6-(trifluoromethyl)pyridin-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **13-1** (600mg, 5%) as a white solid. LC/MS (APCI) *m/z* 272.0 [C₁₃H₁₂F₃NO₂+H]⁺.

[0262] A solution of methyl Compound **13-1** (600mg, 2.12mmol) in 1:1 ACN:H₂O (11mL) was treated with HCl (2N in H₂O, 3.38mL, 6.64mmol) and heated to 80°C. After 1h, the reaction was cooled to rt and then lyophilized to provide 3-(6-(trifluoromethyl)pyridin-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **13-2** (500mg, 88%) as an off-white solid. LC/MS (APCI) *m/z* 258.0 [C₁₂H₁₀F₃NO₂+H]⁺.


[0263] A solution of Compound **13-2** (500mg, 1.94mmol) was dissolved in 1:1 toluene: tBuOH (9.8 mL) and treated with activated 3 Å MS (400mg), followed by Et₃N (699μL, 5.02mmol), and DPPA (503μL, 2.33mmol). The resulting solution was stirred at 35°C for 4h, and then heated to 80°C overnight. The solution was cooled to rt and concentrated to afford the crude product that was then purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide tert-butyl (3-(6-(trifluoromethyl)pyridin-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **13-3** (338mg, 53%) as a white solid. LC/MS (APCI) *m/z* 329.0 [C₁₆H₁₉ F₃N₂O₂+H]⁺.

[0264] A solution of Compound **13-3** (132mg, 0.402mmol) in anhydrous DCM (3.00mL) was cooled to 0°C and treated with TFA (310μL, 4.02mmol) at rt. After 1h, additional TFA was added. After an additional 30 min, the reaction was concentrated *in vacuo* and Et₂O (8mL) was added causing a white solid to crash out of solution. The heterogeneous solution was treated with HCl (2N in Et₂O, 2.0mL, 4.00mmol) and the reaction mixture was stirred for 2 min. The reaction mixture was concentrated *in vacuo* and the process was repeated once more. The suspension was concentrated, and the residual solid was triturated with Et₂O. The precipitate was collected by filtration, and the filter cake was

washed with Et₂O. The white solid was dissolved in 1:1 ACN:H₂O and lyophilized to obtain Compound **13** (87.9mg, 83%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.78 (s, 3H), 8.08 (t, *J*=7.8 Hz, 1H), 7.81 (d, *J*=7.7 Hz, 1H), 7.71 (d, *J*=7.8 Hz, 1H), 2.37 (s, 6H); ¹⁹F NMR (376 MHz, DMSO-*d*₆, unreference) δ -66.43; LC/MS (APCI) *m/z* 229.0 [C₁₁H₁₁F₃N₂+H]⁺.

EXAMPLE 14


3-(5-(trifluoromethyl)benzo[d]thiazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (14)



[0265] A solution of 2-amino-4-(trifluoromethyl)benzenethiol hydrochloride (0.707g, 3.08mmol) and 3-((tert-butoxycarbonyl)amino)bicyclo[1.1.1]pentane-1-carboxylic acid (0.700g, 3.08mmol) in 1N aqueous HCl (15.4mL) was microwaved at 160°C for 60 minutes. The mixture was concentrated and diluted with EtOAc (50mL) and water (20mL). The mixture was neutralized with saturated aqueous NaHCO₃ (30mL) and extracted with EtOAc (4x30mL). The combined organics were dried (Na₂SO₄) and concentrated to provide a yellow residue which was further purified by flash chromatography (SiO₂, DCM/0-10% MeOH containing 7N NH₃) to provide a pale-yellow oil that solidified upon standing. The material was re-dissolved in EtOAc/Et₂O and treated with 2N HCl in Et₂O to generate the HCl salt. The resulting precipitate was collected via filtration and washed with Et₂O to provide 100mg (10%) of Compound **14** as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.01 (br s, NH, 3H), 8.39-8.35 (m, 3H), 7.79 (dd, *J*=1.7, 8.5 Hz, 1H), 2.52 (s, 6H); LC/MS (APCI) *m/z* 285.0 [C₁₃H₁₁F₃N₂S+H]⁺.

EXAMPLE 15

3-(5-(Trifluoromethyl)benzo[d]oxazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (15)



[0266] A solution of 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (0.650g, 3.82mmol) in DCM (19.1mL) was cooled to 0°C and treated with 2 drops of DMF followed by oxalyl chloride (0.401mL, 4.58mmol). The mixture was stirred at rt for 1h then concentrated. The residue was taken up in DCM (19.1mL), cooled to 0°C and treated with

Hünig's base (2.00mL, 11.5mmol) followed by 2-amino-4-(trifluoromethyl)phenol (0.677g, 3.82mmol). The mixture was stirred at rt for 1.5h then concentrated to provide a crude residue which was further purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 0.819g (65%) of methyl 3-((2-hydroxy-5-(trifluoromethyl)phenyl)carbamoyl)bicyclo[1.1.1]pentane-1-carboxylate **15-1** as an off-white solid. LC/MS (APCI) *m/z* 330.0 [C₁₅H₁₄F₃NO₄+H]⁺.

[0267] A solution of Compound **15-1** (0.800g, 2.430mmol) in TFA (12.2mL) and AcOH (12.2mL) was heated at 100°C in a sealed microwave vial overnight. The mixture was cooled to rt and concentrated to provide crude methyl 3-(5-(trifluoromethyl)benzo[d]oxazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **15-2** that also contained 3-(5-(trifluoromethyl)benzo[d]oxazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **15-3**. The mixture was used without further purification. LC/MS (APCI) *m/z* 312.0 [C₁₅H₁₂F₃NO₃+H]⁺.

[0268] A crude mixture of Compound **15-2** and Compound **15-3** (0.756g, 2.43mmol) was dissolved in THF (12.1mL) and treated with 1M aqueous LiOH (5.34mL, 5.34mmol) and allowed to stir overnight. The mixture was acidified with 1N HCl and concentrated. The crude material was triturated with hexanes to provide 0.541g (75%) of Compound **15-3** as an off-white solid which was used without further purification. LC/MS (APCI) *m/z* 298.0 [C₁₄H₁₀F₃NO₃+H]⁺.


[0269] A solution of Compound **15-3** (0.540g, 1.82mmol) in a mixture of toluene (4.54mL) and tert-BuOH (4.54mL) was treated with powdered 3Å mol sieves, Et₃N (0.506mL, 3.63mmol) and DPPA (0.471mL, 2.18mmol). The resulting mixture was stirred at rt for 4h then heated to 90°C and allowed to stir overnight. The mixture was cooled and filtered through a pad of Celite. The mixture was diluted with EtOAc (100mL) and washed with 10% citric acid (20mL) then brine (20mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product which was further purified by flash chromatography (SiO₂, hexanes/EtOAc) to afford 0.299g (45%) of tert-butyl (3-(5-(trifluoromethyl)benzo[d]oxazol-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **15-4** as a white solid. LC/MS (APCI) *m/z* 282.1 [C₁₈H₁₉F₃N₂O₃-C₄H₈+H]⁺.

[0270] A solution of Compound **15-4** (0.299g, 0.812mmol) in EtOAc (4.06mL) was treated with 2N HCl in Et₂O (4.06mL, 8.12mmol). The resulting solution was stirred at

rt until complete. The mixture was concentrated and the resulting solid was triturated with Et₂O and collected via filtration to provide 0.215g (87%) of Compound **15** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.91 (br s, NH, 3H), 8.16 (s, 1H), 7.96 (d, *J*=8.6 Hz, 1H), 7.80 (dd, *J*=1.7, 8.6 Hz, 1H), 2.54 (s, 6H); LC/MS (APCI) *m/z* 269.0 [C₁₃H₁₁F₃N₂O+H]⁺.

EXAMPLE 16

3-(5-chlorobenzo[d]oxazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (16)



[0271] A solution of 3-(methoxycarbonyl)bicyclo[1.1.1]pentane-1-carboxylic acid (0.650g, 3.82mmol) in DCM (19.1mL) was cooled to 0°C and treated with 2 drops of DMF followed by oxalyl chloride (0.401mL, 4.58mmol). The mixture was stirred at rt for 1 h then concentrated. The residue was taken up in DCM (19.1mL), cooled to 0°C and treated with Hünig's base (2.00mL, 11.5mmol) followed by 2-amino-4-chlorophenol (0.548g, 3.82mmol). The mixture was stirred at rt for 1.5h then concentrated to provide a crude residue which was further purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 0.604g (54%) of methyl 3-((5-chloro-2-hydroxyphenyl)carbamoyl)bicyclo[1.1.1]pentane-1-carboxylate **16-1** as a brown solid. LC/MS (APCI) *m/z* 296.0 [C₁₄H₁₄ClN₂O₄+H]⁺.

[0272] A solution of Compound **16-1** (0.600g, 2.03mmol) in TFA (10.1mL) and AcOH (10.1mL) was heated at 100°C in a sealed microwave vial overnight. The mixture was cooled to rt and concentrated to provide crude methyl 3-(5-chlorobenzo[d]oxazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate which was re-dissolved in THF (10.1mL) and treated with 1M aqueous LiOH (4.46mL, 4.46mmol). The resulting mixture was allowed to stir at rt overnight. The mixture was concentrated and the resulting residue was triturated with hexanes to provide 0.450g (84%) of semi-pure 3-(5-chlorobenzo[d]oxazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **16-2** as a light maroon powder. LC/MS (APCI) *m/z* 263.9 [C₁₃H₁₀ClNO₃+H]⁺.


[0273] A solution of Compound **16-2** (0.450g, 1.71mmol) in a mixture of toluene (4.27mL) and *t*-BuOH (4.27mL) was treated with powdered 3 Å molecular sieves, Et₃N (0.476mL, 3.41mmol) and DPPA (0.442mL, 2.05mmol). The resulting mixture was stirred at rt for 4h then heated to 90°C and allowed to stir overnight. The mixture was cooled and

filtered through a pad of Celite. The mixture was diluted with EtOAc (100mL) and washed with 10% citric acid (20mL) then brine (20mL). The combined organics were dried (Na_2SO_4) and concentrated to afford the crude product which was further purified by flash chromatography (SiO_2 , hexanes/EtOAc) to afford 0.356g (62%) of tert-butyl (3-(5-chlorobenzo[d]oxazol-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **16-3** as a light orange solid. LC/MS (APCI) m/z 335.0 [$\text{C}_{17}\text{H}_{19}\text{ClN}_2\text{O}_3+\text{H}$] $^+$.

[0274] A solution of Compound **16-3** (0.356g, 1.06mmol) in EtOAc (5.32mL) was treated with 2N HCl in Et_2O (5.32mL, 10.6mmol). The resulting solution was stirred at rt for 2 days. The resulting suspension was concentrated, re-suspended in Et_2O and filtered. The collected solid was washed with Et_2O and dried to provide 0.230g (80%) of Compound **16** as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.97 (br s, NH, 3H), 7.85 (d, $J=2.1$ Hz, 1H), 7.77 (d, $J=8.7$ Hz, 1H), 7.47 (dd, $J=2.1, 8.7$ Hz, 1H), 2.52 (s, 6H); LC/MS (APCI) m/z 235.0 [$\text{C}_{12}\text{H}_{11}\text{ClN}_2\text{O}+\text{H}$] $^+$.


EXAMPLE 17

3-(5-fluoro-1H-benzo[d]imidazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (17)



[0275] A solution of 2-amino-4-fluoroaniline (0.278g, 2.20mmol) and 3-((tert-butoxycarbonyl)amino)bicyclo[1.1.1]pentane-1-carboxylic acid (0.500g, 2.20mmol) in a 1N aqueous HCl (11.0mL) was microwaved at 160°C for 50 minutes. The dark mixture was filtered through a plug of C_{18} silica gel and washed with ACN. The filtrate was concentrated to provide the crude product which was further purified by flash chromatography (C_{18} SiO_2 , $\text{H}_2\text{O}/\text{ACN}$ buffered with 0.1% formic acid) to afford the desired product. The material was re-dissolved in EtOAc/ Et_2O and treated with 2N HCl in ether to generate the HCl salt. The resulting precipitate was collected via filtration and washed with Et_2O to provide 0.250g (45%) of Compound **17** as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.17 (br s, NH, 3H), 7.76 (dd, $J=4.5, 8.9$ Hz, 1H), 7.60 (dd, $J=2.3, 8.7$ Hz, 1H), 7.36 (dt, $J=2.3, 9.4$ Hz, 1H), 2.60 (s, 6H); LC/MS (APCI) m/z 218.1 [$\text{C}_{12}\text{H}_{12}\text{FN}_3+\text{H}$] $^+$.

EXAMPLE 18

3-(3-Methoxyphenyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (18)

[0276] Mg turnings (0.260g, 10.7mmol) were added to an oven dried sealed tube fitted with a septa and N₂ balloon. Anhydrous Et₂O (20mL) was added followed by 1-bromo-3-methoxybenzene (1.35mL, 10.7mmol) and 46μL of 1,2-dibromoethane. The flask was sealed and stirred at reflux for 3h. After cooling to rt, the flask was fitted with septa and N₂ balloon. The flask was charged with propellane solution (31.3mL, 10.7mmol), sealed, and stirred at 60°C for 3 days. The flask was cooled to 0°C under N₂ and CO₂ was bubbled through the solution for 10 minutes. Et₂O (40mL) and 10mL 2N NaOH was added. The mixture was filtered through Celite. The layers were separated and the organic layers were washed with 2M NaOH (20 mL). The combined water layers were acidified to pH 1 with 2N HCl. The water layers were extracted with EtOAc (3x75mL). The combined organic layers were washed with brine and dried (MgSO₄) to give 950mg of crude residue. The material was absorbed onto Celite and purified by reverse phase chromatography to give 350mg (15%) of 3-(3-methoxyphenyl)bicyclo[1.1.1]pentane-1-carboxylic acid **18-1** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.43 (s, 1H), 7.24-7.21 (m, 1H), 6.82-6.78 (m, 2H), 6.76-6.75 (m, 1H), 3.74 (s, 3H), 2.20 (s, 6H).


[0277] Compound **18-1** (0.688g, 3.15mmol) was added to a 100mL RBF and sealed under N₂. Et₃N (0.879mL, 6.30mmol) and DPPA (0.800mL, 3.71mmol) were added. The reaction was stirred for 1h. 4 Å molecular sieves were added and the reaction was stirred for 3h. The reaction was then refluxed overnight. The solvent was evaporated and the residue was absorbed onto Florisil and purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 437mg (48%) of tert-butyl (3-(3-methoxyphenyl)bicyclo[1.1.1]pentan-1-yl)carbamate **18-2**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.57 (s, 1H), 7.22-7.18 (m, 1H), 6.79-6.71 (m, 2H), 6.74-6.73 (m, 1H), 3.73 (s, 3H), 2.13 (s, 6H), 1.39 (s, 9H).

[0278] Compound **18-2** (0.535g, 1.85mmol) was dissolved in 2mL DCM and 1mL TFA was added. The reaction was stirred at rt for 3h where it was determined to be

complete by LCMS. The reaction was concentrated and 5mL 1N HCl in Et₂O was added with stirring. The mixture was concentrated and the cycle was repeated three times. 1N HCl in Et₂O (5mL) and 10mL Et₂O was added with stirring. The white precipitate was filtered and rinsed with Et₂O. The solid was dried overnight under high vacuum to give 340mg (81%) of Compound **18** as a white powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.90 (br s, NH, 3H), 7.25-7.22 (m, 1H), 6.84-6.70 (m, 3H), 3.74 (s, 3H), 2.25 (s, 6H); LC/MS (APCI) *m/z* 190.1 [C₁₂H₁₅NO+H]⁺.

EXAMPLE 19

3-(1,1-Difluoroethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (19)



[0279] In an oven-dried round bottom flask fitted with a septum, a 0.305M solution of propellane (62.0mL, 18.9mmol) was injected followed by a 0.5M solution of *p*-tolylmagnesium bromide (37.8mL, 18.9mmol). The septum was replaced with a glass stopper wrapped with Teflon tape and the resulting cloudy mixture was stirred at rt for 4 days. The mixture was then cooled to 0°C and dry (CaSO₄) CO₂ gas was bubbled through the mixture for 10 minutes. The mixture was then acidified with 1N HCl, diluted with H₂O (40mL) and extracted with EtOAc (4x30 mL). The combined organics were dried (Na₂SO₄) and concentrated under reduced pressure to provide the crude product that was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to afford 1.96g (51%) of 3-(*p*-tolyl)bicyclo[1.1.1]pentane-1-carboxylic acid **19-1** as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.4 (br s, COOH, 1 H), 7.11 (s, 4 H), 2.27 (s, 3 H), 2.18 (s, 6 H).

[0280] Compound **19-1** (4.74g, 23.4mmol) was added to an oven dried 500mL flask that was fitted with a septa and N₂ balloon. The solid was suspended in 70mL dry Et₂O and the suspension was cooled to 0°C. MeLi (32.2mL, 51.6mmol) was added dropwise and the reaction was stirred from 0°C to rt overnight. The reaction was determined to be complete by TLC. The reaction was cooled to 0°C and acidified to pH 1 with 1N HCl. The mixture was extracted with EtOAc (3x50mL). The combined organic layers were washed with brine (1x100mL) and dried (Na₂SO₄). The compound was purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 2.74g (58%) of 1-(3-(*p*-

tolyl)bicyclo[1.1.1]pentan-1-yl)ethan-1-one **19-2** as a colorless oil. ^1H NMR (400 MHz, DMSO- d_6) δ 7.12 (s, 4H), 2.27 (s, 3H), 2.19 (s, 6H), 2.13 (s, 3H).

[0281] Compound **19-2** (2.50g, 12.5mmol) was dissolved in 2.7M DeoxoFluor (13.9mL, 37.4mmol) in toluene. The reaction was stirred for two days where it was determined to be complete by TLC. The reaction was slowly added to 100mL of a 0°C solution saturated aqueous NaHCO_3 . The mixture was extracted with EtOAc (3x50mL). The organic layers were washed with brine and dried (Na_2SO_4). The solvent was removed and the residue was purified by flash chromatography (SiO_2 , EtOAc/Hexanes) to afford 1.45g (52%) of 1-(1,1-difluoroethyl)-3-(p-tolyl)bicyclo[1.1.1]pentane **19-3** as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 7.12 (s, 4H), 2.27 (s, 3H), 2.04 (s, 6H), 1.61 (t, $J=18.6$ Hz, 3H).


[0282] Compound **19-3** (1.44g, 6.48mmol) was dissolved in 30mL ACN and 30mL water. NaIO_4 (13.9g, 64.8mmol) was added followed by $\text{RuCl}_3 \cdot x\text{H}_2\text{O}$ (0.134g, 0.648mmol). DCM (10mL) was added followed by 10mL water and 10mL ACN to facilitate stirring. The reaction was stirred overnight where it was determined to be complete by TLC. The solvents were removed *in vacuo*. 300mL of 10 % iPrOH/EtOAc was added. MgSO_4 was added with vigorous stirring. The mixture was filtered through Celite, and the solvent removed to give a dark orange oil. The residue was absorbed onto Florisil® and was purified by flash chromatography (SiO_2 , EtOAc/Hexanes) to give 755mg (60%) of 3-(1,1-difluoroethyl)bicyclo[1.1.1]pentane-1-carboxylic acid **19-4** as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 12.57 (s, 1H), 2.02 (s, 6H), 1.56 (t, $J=18.7$ Hz, 3H).

[0283] Compound **19-4** (0.735g, 4.17mmol) was dissolved in 12mL *t*-BuOH. Et_3N (1.16mL, 8.34mmol) was added by DPPA (0.989mL, 4.59mmol). The reaction was stirred for 1h and 4Å molecular sieves were added. The reaction was stirred for 3h at rt. The reaction was then fitted with a reflux condenser refluxed overnight. The 4Å molecular sieves were removed via filtration and the reaction was concentrated. The compound was absorbed onto Florisil® and purified by flash chromatography (SiO_2 , EtOAc:Hexanes) to give 175mg (17%) of tert-butyl (3-(1,1-difluoroethyl)bicyclo[1.1.1]pentan-1-yl)carbamate **19-5** as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 7.61 (s, 1H), 1.95 (s, 6H), 1.56 (t, $J=18.6$ Hz, 3H), 1.37 (s, 9H).

[0284] Compound **19-5** (0.161g, 0.651mmol) was dissolved in 2mL DCM and 1mL TFA was added. The reaction was determined to be complete in 2h by LCMS. The reaction was concentrated and 1N HCl in Et₂O (5mL) was added with stirring. The mixture was concentrated and the cycle was repeated three times. 1N HCl in Et₂O (5mL) and 10mL Et₂O was added with stirring. The white precipitate was filtered and rinsed with Et₂O. The solid was dried overnight under high vacuum to give 96mg (80%) of Compound **19** as a white powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.03 (s, 3H), 2.07 (s, 6H), 1.61 (t, *J*=18.8 Hz, 3H); LC/MS (APCI) *m/z* 148.1 [C₇H₁₁F₂N+H]⁺.

EXAMPLES 20 AND 21

3-(1,1-Difluoro-2-methylpropyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (**20**) and 3-(1-fluoro-2-methylprop-1-en-1-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (**21**)



[0285] 3-(Methoxycarbonyl)bicyclo-[1.1.1]pentane-1-carboxylic acid (5.00g, 29.4mmol) was dissolved in 30mL DCM under N₂. Oxalyl chloride was added and the reaction was stirred overnight. The solvent was removed and the reaction was monitored by ¹H NMR. The material was dissolved in 30mL THF and cooled to -78°C under N₂. 2.0M iPrMgBr in Et₂O (14.7mL, 29.4mmol) was added and the reaction was stirred for 3h and quenched at -78°C with saturated aqueous NH₄Cl. The mixture was extracted with EtOAc (3x75mL) and the combined organic layers were washed with brine (100mL). The solution was dried (Na₂SO₄) and concentrated. The crude mixture was purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 3.91g (68%) of methyl 3-isobutrylbicyclo[1.1.1]pentane-1-carboxylate **20-1** as a colorless oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 3.62 (s, 3H), 2.88 (sep, *J*=6.9 Hz, 1H), 2.25 (s, 6H), 0.97 (d, *J*=6.8 Hz, 6H).

[0286] Compound **20-1** (1.00g, 5.10mmol) was dissolved in neat Deoxo-Fluor® (3.76mL, 20.4mmol) and the reaction was stirred at rt overnight. The reaction was diluted with 15mL of EtOAc and that mixture was slowly added at 0°C to 75mL of saturated aqueous NaHCO₃ with stirring. The mixture was extracted with EtOAc (3x50 mL). The combined organic layers were washed with brine (100 mL) and dried (Na₂SO₄). The solvent was removed and the residue was purified by flash chromatography (SiO₂, EtOAc:Hexanes) to

afford 850mg of a colorless oil that is an inseparable mixture of methyl 3-(1,1-difluoro-2-methylpropyl)bicyclo[1.1.1]pentane-1-carboxylate **20-2** and methyl 3-(1-fluoro-2-methylprop-1-en-1-yl)bicyclo[1.1.1]pentane-1-carboxylate **21-2**. Compound **20-2**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.24 (s, 3H), 2.17 – 1.08 (m, 1H), 2.14 (s, 6H), 0.97 (d, *J*=7.2 Hz, 6H); Compound **21-2**: 2.13 (s, 3H), 2.25 (s, 6H), 1.61-1.57 (m, 6H).

[0287] A mixture of Compound **20-2** and Compound **21-2** (0.800g, 3.67mmol) was dissolved in 15mL of 4:1 THF/water. LiOH (0.769g, 18.3mmol) was added and the reaction was stirred for 3h at rt where it was determined to be complete by TLC. The THF was removed and the mixture was acidified to pH 1 with 1N HCl(aq.). The mixture was extracted with EtOAc (3x30mL). The combined organic layers were washed with brine (75mL) and dried (Na₂SO₄). The mixture was dried on the high vacuum to afford 572mg of a mixture of 3-(1,1-difluoro-2-methylpropyl)bicyclo[1.1.1]pentane-1-carboxylic acid **20-3** and 3-(1-fluoro-2-methylprop-1-en-1-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **21-3**. Compound **20-3**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.51 (s, 1H), 2.15 – 2.08 (m, 1H), 2.09 (s, 6H), 0.973 (d, *J*=6.8 Hz, 6H); Compound **21-3**: 12.51 (s, 1H), 2.19 (s, 6H), 1.61-1.57 (m, 6H).

[0288] A mixture of Compound **20-3** and Compound **21-3** (1.00g, 4.90mmol) was dissolved in 14mL dry t-BuOH. Et₃N (1.36mL, 9.79mmol) was added followed by DPPA (1.05mL, 4.90mmol). The reaction was stirred at rt for 1h and 4Å molecular sieves were added. The reaction was stirred for 3h and then refluxed overnight. The mixture was filtered and the solvent was removed *in vacuo*. The reaction was dissolved in 50mL EtOAc and washed with brine (50mL) and dried (Na₂SO₄). The residue was absorbed on Florisil® and purified by flash chromatography (SiO₂, EtOAc/Hexanes) to provide 430mg of tert-butyl (3-(1,1-difluoro-2-methylpropyl)bicyclo[1.1.1]pentan-1-yl)carbamate **20-4** as a white solid and 302mg of tert-butyl (3-(1-fluoro-2-methylprop-1-en-1-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **21-4** as a white solid. Compound **20-4**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.60 (s, 1H), 2.19-2.07 (m, 1H), 2.02 (s, 6H), 1.37 (s, 9H), 0.96 (d, *J*=6.8 Hz, 6H). Compound **21-4**: ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.57 (s, 1H), 2.11 (s, 6H), 1.59–1.56 (m, 6H), 1.37 (s, 9H).


[0289] Compound **20-4** (0.400g, 1.45mmol) was dissolved in 2mL DCM and 1mL TFA was added. The reaction was determined to be complete in 2h by LCMS. The

reaction was concentrated and 1N HCl in Et₂O (5mL) was added with stirring. The mixture was concentrated and the cycle was repeated three times. 1N HCl in Et₂O (5mL) and 10mL Et₂O was added with stirring. The white precipitate was filtered, rinsed with Et₂O and the solid was dried overnight under high vacuum to give 210mg (68%) of Compound **20** as a white powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.03 (s, 3H), 2.21–2.09 (m, 1H), 2.13 (s, 6H), 0.97 (d, *J*=6.9 Hz, 6H); LC/MS (APCI) *m/z* 176.1 [C₉H₁₅F₂N+H]⁺.

[0290] Compound **21-4** (0.250g, 0.979mmol) was dissolved in 2mL DCM and 1mL TFA was added. The mixture was stirred for 2h at rt at which point it was determined to be complete by LCMS. The reaction was concentrated and 5mL of 1N HCl in Et₂O was added with stirring. The mixture was concentrated and the cycle was repeated three times. 1N HCl in Et₂O (5mL) and 10mL Et₂O was added with stirring. The white precipitate was filtered, rinsed with Et₂O and the solid was dried overnight under high vacuum to give 121mg (64%) of Compound **21** as a white powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.96 (s, 3H), 2.22 (s, 6H), 1.60-1.59 (m, 6H); LC/MS (APCI) *m/z* 156.1 [C₉H₁₄FN+H]⁺.

EXAMPLE 22

3-Phenylbicyclo[1.1.1]pentan-1-amine hydrochloride (22)




[0291] To a solution of propellane (0.34M in Et₂O, 50.0mL, 17.0mmol) was added PhMgBr (3M in Et₂O, 5.67mL, 17.0mmol). The reaction flask was sealed and stirred at rt. After 4 days, the mixture was cooled to -78°C and dry (CaSO₄) CO₂ gas was bubbled through the mixture for 10 minutes. The mixture was warmed to 0°C for 10 min and then acidified with 2N HCl, diluted with H₂O and extracted with EtOAc. The combined organics were dried (Na₂SO₄) and concentrated under reduced pressure to provide the crude product that was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to afford 3-phenylbicyclo[1.1.1]pentane-1-carboxylic acid **22-1** (900mg, 28%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.35-7.20 (m, 5H), 2.37 (s, 6H).

[0292] Compound **22-1** (366mg, 1.95mmol) was dissolved in *t*-BuOH (9.73mL). Activated 3Å MS (500mg) were added followed by Et₃N (543μL, 3.89mmol) and DPPA (503μL, 2.34mmol). The resulting solution was stirred at 30°C for 4h, and then heated to reflux overnight. The solution was cooled to rt and then concentrated under reduced

pressure. The residual oil was diluted with EtOAc (50mL) and H₂O (50mL), and extracted with EtOAc (3x20mL). The combined organics were dried (Na₂SO₄) and concentrated to afford the crude product that was further purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide tert-butyl (3-phenylbicyclo[1.1.1]pentan-1-yl)carbamate **22** (310mg, 61%) as a clear colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.32-7.27 (m, 2H), 7.24-7.19 (m, 3H), 2.28 (s, 6H), 1.47 (s, 9H). LC/MS (APCI) *m/z* 160.1 [C₁₆H₂₁NO₂-C₅H₉O₂+H]⁺.

EXAMPLE 23

(S)-2-Amino-3-methyl-N-(3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)butanamide hydrochloride (23)



[0293] tert-Butyl (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate (0.500g, 0.198mmol) was treated with 4N HCl in dioxane (2mL). The mixture was stirred at rt for 4h then concentrated under reduced pressure. The resulting 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride **23-1** was used without further purification.


[0294] A suspension of Compound **23-1** (0.190g, 1.01mmol) in DCM was treated with Et₃N (0.413mL, 2.96mmol), (S)-2-tert-butoxycarbonylamino-3-methyl-butyric acid (0.330g, 1.50mmol) and HATU (0.760g, 2.00mmol) then allowed to stir at rt for 2h. The mixture was diluted with EtOAc and washed with saturated aqueous NH₄Cl. The organic layer was dried (Na₂SO₄) and concentrated under reduced pressure to provide the crude product which was further purified by flash chromatography (SiO₂, EtOAc/Hexanes) to afford 220mg (63%) of tert-butyl (S)-(3-methyl-1-oxo-1-((3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)amino)butan-2-yl)carbamate **23-2** which was used immediately in the next step.

[0295] Compound **23-2** (0.200g, 0.571mmol) was treated with 4N HCl in dioxane (2mL) and allowed to stir at rt for 4h. The mixture was concentrated under reduced pressure and the resulting residue was lyophilized from a mixture of H₂O/ACN to afford 150mg (91%) of Compound **23** as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.17 (s, 1H), 8.03 (s,

3H), 3.50-3.51 (m, 1H), 2.26 (s, 6H), 2.08-2.13 (m, 1H), 0.94-0.97 (m, 6H); LC/MS (ESI⁺) m/z 251.2 [C₁₁H₁₇F₃N₂O+H]⁺.

EXAMPLE 24


Ethyl (3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-yl)carbamate (24)



[0296] A suspension of 3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (0.190g, 1.01mmol) in DCM (1mL) was cooled to 0°C then treated with Et₃N (0.441mL, 3.16mmol) followed by ethyl chloroformate (0.106mL, 1.11mmol). The resulting solution was allowed to warm to rt while stirring. Once the solution reached rt, the solvent was removed under reduced pressure and the resulting residue was purified by flash chromatography (SiO₂, EtOAc/Hexanes) to provide 100mg (45%) of Compound **24** as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 5.10 (s, 1H), 4.11-4.13 (m, 2H), 2.24 (s, 6H), 1.23-1.27 (m, 3H); LC/MS (ESI⁺) m/z 246.1 [C₉H₁₂F₃NO₂+Na]⁺.

EXAMPLE 25

3-(4-(1,1-difluoroethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-amine hydrochloride (25)



[0297] Representative Procedure (reaction was performed in 5 parallel batches using total of 17.77g starting thioamide). Methyl 3-carbamothioylbicyclo[1.1.1]pentane-1-carboxylate (3.00g, 16.2mmol) was added to a 20mL MW tube. 1-Bromobutane-2,3-dione (1.74mL, 17.0mmol) was dissolved in 16mL MeOH and added to the vial. The reaction was heated at 100°C using a Biotage Initiator microwave reactor for 30 minutes and then combined with the other batches and concentrated *in vacuo*. The crude reaction mixture was partitioned between sat. aq. NaHCO₃ and EtOAc and then the water layer was extracted with EtOAc (3x150mL). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was adsorbed onto Celite[®] and then purified by flash chromatography (SiO₂, EtOAc/Hexanes) to provide a yellow solid. The product was further purified by triturating the yellow solid (2x) with hexanes to provide

19.4g (80%) of methyl 3-(4-acetylthiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **25-1** as a white solid. LC/MS (APCI) m/z 252.0 [$C_{12}H_{13}NO_3S + H$]⁺.

[0298] Representative Procedure (reaction was performed in 4 parallel batches using total of 18.1g of Compound **25-1**). Compound **25-1** (4.10g, 16.3mmol) was suspended in 9mL Deoxo-Fluor and heated to 60°C. The mixture became a clear solution in 10 minutes. The reaction was stirred overnight and then cooled to rt. The crude reactions were cautiously added to a solution of 800mL sat. aq. $NaHCO_3$ and 100mL EtOAc at 0°C by pipette. (Note: vigorous gas evolution was observed during the quenching of the reaction in the absence of EtOAc). The organic layer was separated and the aqueous layer was extracted with EtOAc (3x150mL). The combined organic layers were washed with brine, dried (Na_2SO_4), filtered and concentrated *in vacuo*. The crude product was adsorbed onto Celite® and then purified by flash chromatography (SiO_2 , EtOAc/Hexanes) to provide 15.2g (77%) of methyl 3-(4-(1,1-difluoroethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylate **25-2** as a white solid. LC/MS (APCI) m/z 274.0 [$C_{12}H_{13}F_2NO_2S + H$]⁺.

[0299] To a solution of Compound **25-2** (13.5g, 49.4mmol) in THF (99mL) and water (24.7mL) at 0°C was added $LiOH \cdot H_2O$ (4.15g, 99mmol). The reaction was warmed to rt and stirred for 2h. The THF was then removed *in vacuo* and the reaction was diluted with water to ~100mL. The aqueous layer was washed with Et_2O and then acidified to pH 3 with 1N HCl. The aqueous layer was then extracted with EtOAc (3x150mL). The combined organic layers were washed with brine, dried (Na_2SO_4), filtered and concentrated *in vacuo* to provide 12.1g (95%) of 3-(4-(1,1-difluoroethyl)thiazol-2-yl)bicyclo[1.1.1]pentane-1-carboxylic acid **25-3** as a white solid. LC/MS (APCI) m/z 260.0 [$C_{11}H_{11}F_2NO_2S + H$]⁺.


[0300] Compound **25-3** (13.6g, 52.5mmol) was dissolved in *t*-BuOH (131mL) and anhydrous toluene (131mL). Activated 3Å MS (13.0g) were added followed by Et_3N (14.6mL, 105mmol) and DPPA (13.6mL, 62.9mmol). The resulting solution was stirred at 30°C for 4h, and then heated to reflux overnight. The solution was cooled to rt and then filtered through a pad of Celite®. The pad was then washed with EtOAc (50mL) and the combined filtrates were concentrated *in vacuo*. The crude product was adsorbed onto Celite® and then purified by flash chromatography (SiO_2 , EtOAc/Hexanes) to provide 13.1g (75%) of

tert-butyl (3-(4-(1,1-difluoroethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-yl)carbamate **25-4** as a white solid. LC/MS (APCI) m/z 331.1 [$C_{15}H_{20}F_2N_2O_2S+H$] $^+$.

[0301] A solution of Compound **25-4** (13.0g, 39.3mmol) in anhydrous DCM (98mL) was cooled to 0°C and treated with TFA (98mL) at rt. After 90 min, the reaction was concentrated *in vacuo* and Et₂O (50mL) was added causing a white solid to crash out of solution. The heterogeneous solution was treated with HCl (4N in Et₂O, 20mL, 80.0mmol) and the reaction mixture was stirred for 2 min. The reaction mixture was concentrated *in vacuo* and the process was repeated an additional 2X using 20mL and 10mL 4N HCl in dioxane respectively. The suspension was concentrated, and the residual solid was triturated with Et₂O. The precipitate was collected by filtration, and the filter cake was washed with Et₂O. The white solid was dried under vacuum to afford 10.2g (97%) of Compound **25** as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆, 60 °C) δ 8.92 (s, 3H), 7.96 (t, *J*=1.1 Hz, 1H), 2.45 (s, 6H), 1.99 (t, *J*=18.8 Hz, 3H); ¹⁹F NMR (376 MHz, DMSO-*d*₆, 60 °C, unreferenced) δ -84.32; LC/MS (APCI) m/z 231.0 [$C_{10}H_{12}F_2N_2S+H$] $^+$.

EXAMPLE 26

(5Z,8Z,11Z,14Z)-N-(3-(4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-yl)icosa-5,8,11,14-tetraenamide (26)




[0302] The following operations were performed in a manner as to minimize exposure to light. A solution of (5Z,8Z,11Z,14Z)-icosa-5,8,11,14-tetraenoic acid (40mg, 0.131mmol) in DCM (0.9mL) and DMF (10μL) was cooled to 0°C and oxalyl chloride (23μL, 0.263mmol) was added dropwise. The reaction mixture was stirred for an additional 1h followed by the addition of a solution of Compound **1** (107mg, 0.394mmol) in pyridine (32μL, 0.394mmol). The mixture was warmed to rt and stirred for an additional 30 minutes. The mixture was diluted with DCM (5mL) and washed with 10% aqueous HCl and water. The organic layer was dried (Na₂SO₄), concentrated and purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 62mg (91%) of Compound **26** as a viscous pale yellow oil: ¹H NMR (400 MHz, CDCl₃) δ 7.65 (s, 1H), 5.86 (br s, 1H), 5.43-5.30 (m, 8 H), 2.86-

2.79 (m, 6H), 2.56 (s, 6 H), 2.18-2.03 (m, 6 H), 1.72 (quin, $J=7.4$ Hz, 2 H), 1.39-1.24 (m, 6 H), 0.89 (t, $J=6.8$ Hz, 3 H).

EXAMPLE 27


(5Z,8Z,11Z,14Z)-N-(3-(5-methyl-4-(trifluoromethyl)thiazol-2-yl)bicyclo[1.1.1]pentan-1-yl)icosa-5,8,11,14-tetraenamide (27)



[0303] The following operations were performed in a manner as to minimize exposure to light. A solution of (5Z,8Z,11Z,14Z)-icosa-5,8,11,14-tetraenoic acid (40mg, 0.131mmol) in DCM (0.9mL) and DMF (10μL) was cooled to 0°C and oxalyl chloride (23μL, 0.263mmol) was added dropwise. The reaction mixture was stirred for an additional 1h followed by the addition of a solution of Compound **2** (112mg, 0.394mmol) in pyridine (32μL, 0.394mmol). The mixture was warmed to rt and stirred for an additional 30 minutes. The mixture was diluted with DCM (5mL) and washed with 10% aqueous HCl and water. The organic layer was dried (Na₂SO₄), concentrated and purified by flash chromatography (SiO₂, Hexanes/EtOAc) to provide 71mg (>99%) of Compound **27** as a viscous pale yellow oil: ¹H NMR (400 MHz, CDCl₃) δ 5.84 (br s, 1H), 5.43-5.30 (m, 8 H), 2.85-2.79 (m, 6H), 2.54 (q, $J=1.8$ Hz, 3 H), 2.50 (s, 6 H), 2.17-2.03 (m, 6 H), 1.71 (quin, $J=7.4$ Hz, 2 H), 1.40-1.24 (m, 6 H), 0.89 (t, $J=6.8$ Hz, 3 H).

EXAMPLE 28

3-(trifluoromethyl)bicyclo[1.1.1]pentan-1-amine hydrochloride (28)



[0304] Compound **28** was obtained using a procedure described in U.S. Pub. No. 2016/0075654, which is herein incorporated by reference.

EXAMPLE A
FORMALIN PAW TEST

[0305] One test compound or the vehicle was administered to each rat or mouse in each test group (8 rats or mice per group). Non-fasted male Sprague-Dawley rats weighing 200-225 g or non-fasted male ICR mice weighing 23 ± 3 g were used. Test compounds were administered at a concentration of 30 mg/kg or 60 mg/kg; morphine was administered at a concentration of 5 mg/kg; and acetaminophen was administered at a concentration of ≥ 300 mg/kg. The control group received the vehicle (9% PEG400/ 0.5% PVP/0.5% Tween-80/ 90% of 0.5% CMC in water) for rats or the vehicle (5% DMSO/15% PEG400/ 10% HPbCD/0.9% Saline) for mice. After 30 minutes, a 2% formalin solution (0.05 mL) was injected into one hind paw (sub-plantar) of each rat. For mice, after 10 minutes a 2% formalin solution (0.02 mL) was injected into one hind paw (sub-plantar) of each mouse. Responses were measured every 5 minutes after the formalin injection for 35 minutes. The results in Table 2 are for oral administration in rat and the results in Table 3 are for intraperitoneal administration in mouse. In Tables 2 and 3, 'A' designates <70 seconds of paw licking, 'B' designates ≥ 70 and <165 seconds of paw licking, and 'C' designates ≥ 165 seconds of paw licking.

[0306] The results are provided in Tables 2 and 3. As shown in Tables 2 and 3, compounds of Formulae (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ik), (Im), (In), (Io), (Ip), (Iq), (Ir), (Is) and (It), or a pharmaceutically acceptable salt of any of the foregoing, significantly decreased the pain response in both the early/acute phase (0-10 minutes) and the late/tonic phase (10-35 minutes). The results indicate that compounds described herein have significant analgesic properties.

TABLE 2

Compound No.	Dosage (mg/kg)	Early Phase	Late Phase
1	30	A	B
2	30	A	A
3	30	A	A
4	30	A	A
5	30	A	B

Compound No.	Dosage (mg/kg)	Early Phase	Late Phase
6	30	A	B
7	30	A	B
8	30	A	C
9	30	A	C
10	30	A	A
11	30	A	C
12	30	A	C
13	30	A	A
14	30	A	C
15	30	A	C
16	30	A	C
17	30	A	C
18	30	A	B
19	30	A	B
20	30	A	C
21	30	A	C
22	30	A	B
25	30	A	A

TABLE 3

Compound No.	Dosage (mg/kg)	Early Phase	Late Phase
23	60	A	C
24	60	A	B

EXAMPLE BUVB INFLAMMATORY PAIN TEST

[0307] UVB irradiation to the surface of the hind paw was used to induce cutaneous inflammatory pain in non-fasted male Sprague-Dawley rats weighing 175-300 g. On Day -3, animals were restrained and the plantar surface of the left hind paw was exposed to 350 mJoules/cm² of UVB irradiation using a narrow band UVB light source (Daavlin®, Bryan, Ohio). Following UVB irradiation, animals were returned to their home cage. Thermal hyperalgesia was assessed prior to dosing and approximately 1, 2, and 4 hours after dosing on Day 0 using a radiant heat plantar test apparatus (IITC Life Sciences®; Woodland Hills, CA). Animals were placed in individual acrylic chambers on a glass platform warmed

to $30 \pm 2^{\circ}\text{C}$ and allowed to acclimate to their surroundings for a minimum of 15 minutes before testing. The stimulus was presented to the plantar surface and the timer was activated. The stimulus intensity was set such that pre-injury paw withdrawal latencies were approximately 12-18 seconds. Paw withdrawal latency values were recorded at the first observed nocifensive behavior (paw withdrawal, flinching, biting and/or licking). Three paw withdrawal latencies were measured for each hind paw per time point. The mean of the 3 values was taken as the paw withdrawal latency for that time point. A maximum cutoff latency of 30 seconds was used to prevent injury to the animal. The mean and standard error of the mean (SEM) were determined for each paw for each treatment group at each time point. Test compounds were administered orally at concentrations of 3, 10, 30 or 100 mg/kg; celecoxib was administered orally at a concentration of 30 mg/kg; and acetaminophen was administered orally at a concentration of 300 mg/kg. The control group received the vehicle (sterile water). In Table 4, 'A' designates ≥ 8 seconds of paw withdrawal latency, 'B' designates ≥ 6 and < 8 seconds of paw withdrawal latency, and 'C' designates < 6 seconds of paw withdrawal latency.

[0308] The results provided in Table 4 are for oral administration in rats. As shown in Table 4, compounds of Formulae (Ie), (Ih) and (Ik), or a pharmaceutically acceptable salt of any of the foregoing, significantly decreased the pain response at 1, 2 and 4 h. The results indicate that compounds described herein have at least comparable or improved inflammatory analgesic properties relative to those of acetaminophen and NSAIDs, such as celecoxib. Furthermore, while acetaminophen was dosed at 300 mg/kg, compounds of Formulae (Ie), (Ih) and (Ik), or a pharmaceutically acceptable salt of any of the foregoing, were dosed at 3 mg/kg, 10 mg/kg and 100 mg/kg. Thus, comparable or improved inflammatory analgesic properties were achieved at significantly lower doses of compounds described herein (such as approximately 100-fold, approximately 30-fold and approximately 3-fold) relative to that of acetaminophen. As a lower dose of compounds described herein are needed to achieve comparable or improved inflammatory analgesic properties, compounds described herein are expected to be better tolerated by a subject's body with compared to APAP.

TABLE 4

Compound No.	Dosage (mg/kg)	1 h	2 h	4 h
1	3	B	B	B
	10	B	B	B
	30	A	A	A
	100	A	A	A
2	3	A	A	A
	10	A	B	A
	30	A	A	B
	100	A	A	A
6	3	C	B	A
	10	B	B	B
	30	A	B	B
	100	A	A	B
7	3	B	B	B
	10	C	C	B
	30	B	B	B
	100	C	C	B
Celecoxib	30	A	A	A
Acetaminophen	300	A	A	A
Vehicle	N/A	C	C	C

EXAMPLE C
OSTEOARTHRITIS PAIN TEST

[0309] Osteoarthritis pain was induced in non-fasted male Sprague-Dawley rats weighing 150-250 g by injecting 2 mg of monosodium iodoacetate (MIA) into the left knee joint. Mechanical hyperalgesia was measured using a digital Randall-Selitto device (dRS; IITC Life Sciences®; Woodland Hills, CA). Animals were allowed to acclimate to the testing room for a minimum of 15 minutes before testing. Animals were placed in a restraint sling that suspended the animal, leaving the hind limbs available for testing. The stimulus was applied to the knee joint by a blunt tip and pressure was applied gradually over approximately 10 seconds. Joint compression threshold values were recorded at the first observed nocifensive behavior (vocalization, struggle, or withdrawal). One reading per joint was taken at each time point, and a maximum stimulus cutoff of 500 grams was used to prevent injury to the animal. Mechanical hyperalgesia was assessed prior to MIA injection, and again 14 days later. After 14 days, test or control articles were administered once daily for 7 days with

mechanical hyperalgesia assessed prior to dosing, 1, 2, and 4 hours post-dosing on days 0, 3, and 6. Test compounds were administered orally at concentrations of 3, 10, 30 or 100 mg/kg and celecoxib was administered orally at a concentration of 30 mg/kg. The control group received the vehicle (sterile water). After day 0, groups receiving the 100 mg/kg dose were reduced to a dose of 3 mg/kg. In Table 5, 'A' designates ≥ 375 grams of joint compression, 'B' designates ≥ 325 and < 375 grams of joint compression, and 'C' designates < 325 grams of joint compression.

[0310] The results provided in Table 5 are for oral administration in rats. As shown in Table 5, compounds of Formula (Ie), or a pharmaceutically acceptable salt of any of the foregoing, significantly decreased the pain response at 1, 2 and 4 h on days 0, 3 and 6. As shown by the data in Table 5, compounds described herein have comparable or improved osteoarthritis analgesic properties relative to those of NSAIDs, such as celecoxib. Furthermore, starting on day 3, comparable or improved osteoarthritis analgesic properties were achieved compared to celecoxib at all doses evaluated for the compounds described herein.

TABLE 5

Compound No.	Dosage (mg/kg)	Day 0			Day 3			Day 6		
		1 h	2 h	4 h	1 h	2 h	4 h	1 h	2 h	4 h
1	3	-	-	-	A	A	A	A	B	A
	10	B	C	C	B	B	B	B	B	B
	30	B	B	B	A	A	B	A	B	B
	100	A	A	A	-	-	-	-	-	-
2	3	-	-	-	A	A	A	B	B	B
	10	C	C	C	B	A	C	B	B	A
	30	B	B	B	B	B	B	B	B	A
	100	A	A	A	-	-	-	-	-	-
Celecoxib	30	B	A	B	C	A	A	B	A	A
Vehicle	N/A	C	C	C	C	C	C	C	C	C

EXAMPLE DPOST-INCISIONAL (BRENNAN) PAIN TEST

[0311] Incisional pain was induced in non-fasted male Sprague-Dawley rats weighing 180-300 g by making a 1-cm plantar incision in the left hind paw and is designed to mimic pain and sensitivity similar to what is reported by patients after surgery. Dosing occurred at the time of incision and spontaneous pain was measured 4 and 24 hours post-dosing using a cumulative pain score. The unrestrained animals were placed on a metal mesh platform and allowed to acclimate for about 15 min. Each animal was closely observed during a 1-min period repeated every 5 min for 1 h (12 times total). Depending on the position in which the foot was found during the scoring period, a score of 0, 1, or 2 was given. Full weight bearing of the foot (score = 0) was present if the skin on the plantar surface of the paw wound was blanched or distorted by the mesh. If the foot was completely off the mesh, or one or more instances of licking or flinching was observed, a score of 2 was recorded and the licking or flinching was noted but not counted. If the plantar surface of the paw touched the mesh without blanching or distorting, a score of 1 was given. The sum of the 12 scores obtained during the 1 h session was calculated (maximum score = 24) for each animal. Observations were made during hours 4 and 24. Test articles were administered orally at concentrations of 10, 30 or 100 mg/kg and ketoprofen was administered subcutaneously at a concentration of 10 mg/kg. The control group received the vehicle (sterile water). In Table 6, 'A' designates a cumulative pain score of ≤ 11 , 'B' designates >11 and ≤ 16 , and 'C' designates >16 .

[0312] The results provided in Table 6 are for oral administration in rats. As shown in Table 6, compounds of Formula (Ie), or a pharmaceutically acceptable salt of any of the foregoing, significantly decreased the pain response at 4 and 24 h. The results in Table 6 demonstrate that compounds described herein have at least comparable or improved incisional analgesic properties compared to ketoprofen, an NSAID.

TABLE 6

Compound No.	Dosage (mg/kg)	4 h	24 h
1	10	C	B
	30	A	A
	100	A	A
28	10	C	C
	30	B	B
	100	B	A
Ketoprofen	10	B	A
Vehicle	N/A	C	C

EXAMPLE ESPINAL NERVE LIGATION (SNL) NEUROPATHIC PAIN TEST

[0313] Neuropathy was induced in non-fasted male Sprague-Dawley rats weighing 75-100 g by surgically ligating the 4th and 5th lumbar spinal nerves (L4 and L5), a procedure also known as spinal nerve ligation (SNL). Animals were placed in individual acrylic chambers on a metal mesh surface and allowed to acclimate to their surroundings for a minimum of 15 minutes before testing. Mechanical sensitivity was assessed using 8 Semmes-Weinstein filaments (Stoelting®; Wood Dale, IL, USA) with varying stiffness (0.4, 0.6, 1.0, 2.0, 4.0, 6.0, 8.0, and 15 g) according to an up-down method to determine 50% response thresholds. Each filament was presented perpendicular to the plantar surface with sufficient force to cause slight buckling against the paw and held for approximately 6 seconds or until a positive response is noted (paw sharply withdrawn). Testing was initiated with the 2.0 g filament. In the absence of a paw withdrawal response, the next stronger stimulus was presented. In the event of paw withdrawal, the next weaker stimulus was used. This process was repeated until 4 responses after the initial change in response (no response to positive response or positive response to no response) were obtained. If the animal did not respond after reaching the strongest filament or if the animal responded after reaching the weakest filament, the testing was stopped for that time point. The 50% response threshold was calculated using the formula:

$$50\% \text{ response threshold (g)} = (10^{(X_f + k\delta)}) / 10,000;$$

X_f = value (in log units) of the final von Frey filament used;

k = tabular value for the pattern of positive/negative responses (*See* Chaplan, *et al.* "Quantitative Assessment of Tactile Allodynia in the Rat Paw" *J. Neurosci. Meth.* 53.1 (1994): 56-63);

δ = mean difference (in log units) between stimuli.

[0314] The mean and standard error of the mean (SEM) were determined for each paw for each treatment group at each time point. Thresholds were determined prior to surgery and again 14 days later. After 14 days, test or control compounds were administered and thresholds were again determined 1, 2, and 4 hours after compound administration. Median response thresholds in compound-treated animals were compared to those in vehicle treated animals to determine anti-allodynic efficacy of test and control compounds. Test compounds were administered orally at concentrations of 10, 30 or 100 mg/kg and gabapentin was administered orally at a concentration of 100 mg/kg. The control group received the vehicle (sterile water). In Table 7, 'A' designates a 50% threshold of ≥ 8 grams, 'B' designates a 50% threshold of ≥ 6 and < 8 grams, and 'C' designates a 50% threshold of < 6 grams.

[0315] The results provided in Table 7 are for oral administration in rats. As shown in Table 7, compounds of Formula (Ie), or a pharmaceutically acceptable salt of any of the foregoing, significantly decreased the pain response at 1, 2 and 4 h. The data in Table 7 indicates that compounds described herein have at least comparable or improved neuropathic analgesic properties compared to gabapentin.

TABLE 7

Compound No.	Dosage (mg/kg)	1 h	2 h	4 h
1	10	C	C	C
	30	B	B	C
	100	A	A	A
2	10	C	C	C
	30	C	C	C
	100	A	A	B
Gabapentin	100	B	A	A
Vehicle	N/A	C	C	C

EXAMPLE F
NSAIDS AND OPIOID RECEPTOR TEST

[0316] Compounds were screened as DMSO solutions at 10 μ M, 100 μ M, or 200 μ M and values represent percent inhibition of control-specific binding. Table 8 demonstrates the lack of significant *in vitro* activity of compounds of Formulae (Ie) and (Ir) or a pharmaceutically acceptable salt of any of the foregoing, against a panel of COX-1/COX-2 or opiate receptors. This data indicates that compounds described herein have a mechanism of action that is different from that of NSAIDs and opioids. As NSAID and opioid mechanisms of action are precluded, a person of ordinary skill in the art understands that undesirable side effects associated with inhibition of COX receptors in NSAIDs and activation of opiate receptors in opioids can be avoided by compounds described herein. Furthermore, opioids are known to lose their analgesic potency after to continued administration. (See Benyamin, et al. "Opioid Complications and Side Effects" *Pain Physician* 11 (2008): S105-S120). As the data indicates compounds described herein have a different mechanism action compared to opioids, a person of ordinary skill in the art would appreciate that compounds described herein would not be expected to lose their analgesic potency like opioids after continued administration.

TABLE 8

Assay Name	% Inhibition					
	APAP	Cmpd 1	Cmpd 2		Cmpd 13	Cmpd 28
	10 μ M	10 μ M	10 μ M	200 μ M	200 μ M	100 μ M
COX-1 (human)	-7	-	-	15	6	-
COX-2 (human)	3	-	-	-2	8	-
Opiate δ_1 (human)	2	-	-	-	-	-
Opiate δ_2 (human)	-	7	10	-	6	2
Opiate κ (human)	-1	-4	-9	-	33	1
Opiate μ (human)	-5	11	9	-	56	3

EXAMPLE G
CONDITIONED PLACE PREFERENCE (ABUSE) TEST

[0317] A conditioned place preference (CPP) model was used to assess the reward properties of compounds of formulae (Ie), or a pharmaceutically acceptable salt of any of the foregoing, in adult male C57BL/6 mice. The CPP model apparatus consisted of a Plexiglas® test unit having three (A, B, C) compartments (9cmx9cmx16cm) configured to allow access to all chambers. Compartments A and C had textually distinct flooring and wall marking. Mice were placed in the middle compartment with free access to the other compartments. On day 1 and 2, the “naïve chamber preference” was assessed by placing the animal in the middle chamber twice a day (AM and PM) for 30 min on two consecutive days, with the chamber showing the greatest residence time being called the “preferred chamber”. On “drug pairing days” (days 5, 6, 7 and 8) mice were injected IP with vehicle (50mM citric acid) in the AM in the preferred chamber and the drug/vehicle in the afternoon in the non-preferred chamber for four consecutive days. On the morning of day 9 (the day after the last drug pairing), the “drug chamber preference” was assessed by placing the mouse (absent any injection) in the middle chamber, with free access to all chambers for 30 minutes. This was repeated five days later (day fourteen), with no intervening drug treatment.

[0318] The time spent on days 9 and 14 (e.g. day 0 and day 5 respectively after the last drug treatment) in the drug associated chamber was the test datum and is represented in FIG. 1. Mice undergoing least-preferred chamber pairing with morphine for 4 consecutive days displayed a significant increase in the least preferred chamber residence time ($p < 0.0001$) on day 9 and on day 14 (1 and 5 days after the last drug treatment). In contrast, there was no increase in the non-preferred chamber residence time when the least-preferred chamber was paired with saline, APAP or test compounds 2 or 28 on either days 9 or 14, or days 0 or 5. These results indicate that compounds described herein are less likely to be abused or addictive relative to opioids, such as morphine.

EXAMPLE H
BIOAVAILABILITY TEST

[0319] Oral bioavailability of compounds of Formulae (Ie), (Ih), (Ik), and (Ir), or a pharmaceutically acceptable salt of any of the foregoing, were evaluated in fasted male Sprague-Dawley (SD) rats weighing 200-225g. Compounds were dosed at 3 mg/kg (IV) and 30 mg/kg (PO). Unless otherwise noted, compounds were dissolved in 0.9% saline for IV dosing and 100% sterile water for PO dosing. Pharmacokinetic parameters were determined using Phoenix WinNonlin (v7.0).

[0320] The results provided in Table 9 indicate that the compounds of Formulae (Ie), (Ih), (Ik), and (Ir), or a pharmaceutically acceptable salt of any of the foregoing, show favorable pharmacokinetic profiles, and a person of ordinary skill in the art would appreciate that the compounds of the present application may have improved metabolic stability relative to acetaminophen. For example, a person of ordinary skill in the art would appreciate that the compounds described herein demonstrate a longer metabolic half-life ($t_{1/2}$) than that of acetaminophen. (*See* Hirate et al., *Biopharmaceutics & Drug Disposition* (1990) 11:245-252). Therefore, compounds described can be dosed less often compared to a known NSAID, APAP and/or opioid.

TABLE 9

Parameters	Cmpd 1		Cmpd 2		Cmpd 7	
	IV	PO	IV	PO	IV	PO
AUC _{inf} ($\mu\text{M} \cdot \text{h}$)	19.2	277	4.0	29	9.0	119
CL ($\text{mL} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$)	11	--	52	--	27	--
V _{dss} (L/kg)	1.4	--	3.6	--	2.5	--
C _{max} (μM)	--	49	--	12	--	67
T _{max} (h)	--	0.3	--	1.0	--	0.19
t _{1/2} (h)	3.1	5.6	2.7	2.4	2.3	2.5
F (%)	--	115	--	79	--	138

Parameters	Cmpd 6		Cmpd 13		Cmpd 25 ^a	
	IV	PO	IV	PO	IV	PO
AUC _{inf} ($\mu\text{M} \cdot \text{h}$)	1.8	29	3.0	35	13	144 ^b
CL ($\text{mL} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$)	121	--	73	--	17	--
V _{dss} (L/kg)	0.79	--	20.2	--	3.2	--
C _{max} (μM)	--	32	--	11	--	24
T _{max} (h)	--	0.22	--	0.6	--	1.2
t _{1/2} (h)	0.15	1.6	9.5	6.8	3.7	ND
F (%)	--	134	--	16	--	115

^a IV: DMSO, PEG400, and 30% HP β CD (5/20/75); PO: PEG400, PVP, Tween 80, 0.5% CMC in water (9/0.5/0.5/90). ^b AUC_{24h}

[0321] Although the foregoing has been described in some detail by way of illustrations and examples for purposes of clarity and understanding, it will be understood by those of skill in the art that numerous and various modifications can be made without departing from the spirit of the present disclosure. Therefore, it should be clearly understood that the forms disclosed herein are illustrative only and are not intended to limit the scope of the present disclosure, but rather to also cover all modification and alternatives coming with the true scope and spirit of the invention.