
SUMMARY OF THE INVENTION

The present invention provides a compound of formula (I):



(I)

;

wherein:

L is –NHCO- or –CONH-; and

R^1 and R^2 are each independently alkyl, cycloalkyl, ketocycloalkyl, heterocyclyl, aryl or heteroaryl, which is optionally mono-, di-, or tri-substituted independently with alkyl, alkoxy, halogen, cyano, nitro, trifluoroalkyl, amino, alkylamino, dialkylamino, acyl, aryl, heteroaryl, heterocyclyl, heterocyclyl-R^3, -NHR^3, -N(alkyl)R^3, -$C(O)NHR^3$, -C(O)N(alkyl)R^3, -$NHC(O)R^3$, -N(alkyl)$C(O)R^3$, -OH or -OR^3, wherein:

R^3 is C_1-C_6alkyl or C_1-C_6cycloalkyl, which is optionally substituted with halogen, -CN, -NH_2, -NH(C_1-C_3alkyl), -N(C_1-C_3alkyl)$_2$, C_1-C_3alkylheterocyclyl, C_1-C_3alkylcarbamate, -C(O)NH(C_1-C_3alkyl), -C(O)N(C_1-C_3alkyl)$_2$, -NHC(O)-C_1-C_3alkyl, -N(C_1-C_3alkyl)-C(O)-C_1-C_3alkyl, OH, or –O-C_1-C_6alkyl; or

a pharmaceutically acceptable salt thereof.

The present invention also provides a pharmaceutical composition comprising at least one compound of the invention or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.

The present ~~invention~~disclosure also provides a method of treating a disease or disorder, the method comprises administering a therapeutically effective amount of at least one compound ~~of the present invention~~disclosed herein or a pharmaceutically acceptable salt thereof to a mammal in need thereof, wherein the disease or disorder is a central nervous system disease or disorder. In some embodiments of the method, a symptom of the disease or disorder is treated.

In one aspect, there is provided the use of a therapeutically effective amount of at least one compound as defined above or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating a central nervous system disease or disorder.

DETAILED DESCRIPTION OF THE INVENTION