

FORMULATIONS OF FACTOR VIII CHIMERIC PROTEINS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application Serial Nos. 63/214,245 filed June 23, 2021, 63/214,246 filed June 23, 2021, 63/214,752 filed June 24, 2021, and 63/231,909 filed August 11, 2021, each of which are incorporated herein by reference in their entirety.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[0002] The content of the electronically submitted sequence listing in ASCII text file (Name: SA9-480PC_ST25.txt; Size: 158 kb; Created: June 22, 2022) is incorporated herein by reference in its entirety.

BACKGROUND OF THE DISCLOSURE

[0003] Hemophilia A is a bleeding disorder caused by defects in the gene encoding coagulation factor VIII (FVIII) and affects 1-2 in 10,000 male births. Graw *et al.*, *Nat. Rev. Genet.* 6(6): 488-501 (2005). Patients affected with hemophilia A can be treated with infusions of purified plasma FVIII or recombinantly produced FVIII. Many commercially available FVIII products are known to have a half-life of about 8-12 hours, requiring frequent intravenous administration to the patients. See Weiner M.A. and Cairo, M.S., *Pediatric Hematology Secrets*, Lee, M.T., 12. Disorders of Coagulation, Elsevier Health Sciences, 2001; Lillicrap, D. *Thromb. Res.* 122 Suppl 4:S2-8 (2008). In addition, a number of approaches have been tried in order to extend the FVIII half-life. For example, the approaches in development to extend the half-life of clotting factors include pegylation, glycopegylation, and conjugation with albumin. See Dumont *et al.*, *Blood.* 119(13): 3024-3030 (2012). Consistent results have been demonstrated in humans, for example, an rFVIII-Fc fusion protein was reported to improve half-life up to ~1.7-fold compared with ADVATE® in hemophilia A patients. See Powell *et al.*, *Blood.* 119(13): 3031-3037 (2012). Therefore, the half-life increases, despite minor improvements, indicate the presence of other half-life limiting factors, such as clearance by VWF. Pipe *et al.*, *Blood.* 128(16):2007-2016 (2016)).

[0004] Efanesoctocog alfa (also known as Efa and BIVV001) is a fusion protein that is designed to uncouple recombinant clotting factor VIII from VWF in circulation. The chimeric protein comprises a single recombinant factor VIII protein fused to dimeric Fc, a D'D3 domain of VWF, and two ELNN Polypeptides. Chhabra *et al.* *Blood* 135(17): 1484-1496 (2020). In one early study

involving patients with severe hemophilia A, a single intravenous injection of efanesoctocog alfa resulted in high sustained factor VIII activity levels, with a half-life that was up to four times the half-life associated with recombinant factor VIII. See Konkle et al., *NEJM*. 383:1018-27 (2020).

[0005] However, there remains a need for improved pharmaceutical compositions.

BRIEF SUMMARY OF THE DISCLOSURE

[0006] The present disclosure is directed to, *inter alia*, pharmaceutical compositions comprising a Factor VIII ("FVIII") protein, kits comprising such pharmaceutical compositions, and therapeutic methods and uses of the pharmaceutical compositions.

[0007] In some embodiments, the pharmaceutical composition comprises a chimeric protein that comprises a first polypeptide chain which comprises a FVIII protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof. In some embodiments, the chimeric protein comprises (i) a FVIII protein comprising a FVIII polypeptide, an ELNN Polypeptide inserted within the B domain (e.g., replacing at least a portion of the B domain) of the FVIII polypeptide, and a first Fc region; and (ii) a VWF protein comprising a VWF fragment (e.g., a fragment comprising the D'D3 domains of VWF, which fragment may comprise mutations), a second ELNN Polypeptide, a thrombin-cleavable linker (such as an $\alpha 2$ linker), and a second Fc region. In some embodiments, the chimeric protein disclosed herein is a FVIII-ELNN-Fc/D'D3-ELNN-Fc heterodimer.

[0008] In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) sucrose; (c) histidine; (d) arginine; (e) calcium chloride; and (f) a polysorbate. In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) sucrose; (c) L-histidine; (d) L-arginine; (e) calcium chloride; and (f) a polysorbate. In some embodiments, the pharmaceutical composition comprising: (a) a Factor VIII ("FVIII") protein; (b) sucrose; (c) L-histidine; (d) L-arginine-HCl; (e) calcium chloride dihydrate; and (f) a polysorbate.

[0009] In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) about 1% (w/v) to about 4% (w/v) sucrose; (c) about 5 mM to about 15 mM histidine; (d) about 150 mM to about 300 mM arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) about 1% (w/v) to about 4% (w/v) sucrose; (c) about 5 mM to about 15 mM histidine; (d) about 200 mM to about 300 mM arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) about 1% (w/v) to about 4% (w/v) sucrose; (c) about 5 mM to about 15 mM L-histidine;

(d) about 200 mM to about 300 mM L-arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprises: (a) a Factor VIII ("FVIII") protein; (b) about 1% (w/v) to about 4% (w/v) sucrose; (c) about 5 mM to about 15 mM L-histidine; (d) about 200 mM to about 300 mM L-arginine-HCl; (e) about 2.5 mM to about 10 mM calcium chloride dihydrate; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate.

[0010] In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) about 5% (w/v) to about 7.5% (w/v) sucrose; (c) about 5 mM to about 15 mM histidine; (d) about 150 mM to about 300 mM arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprises: (a) a FVIII protein; (b) about 5% (w/v) to about 7.5% (w/v) sucrose; (c) about 5 mM to about 15 mM histidine; (d) about 200 mM to about 300 mM arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprising: (a) a Factor VIII ("FVIII") protein; (b) about 5% (w/v) to about 7.5% (w/v) sucrose; (c) about 5 mM to about 15 mM L-histidine; (d) about 200 mM to about 300 mM L-arginine; (e) about 2.5 mM to about 10 mM calcium chloride; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate. In some embodiments, the pharmaceutical composition comprising: (a) a Factor VIII ("FVIII") protein; (b) about 5% (w/v) to about 7.5% (w/v) sucrose; (c) about 5 mM to about 15 mM L-histidine; (d) about 200 mM to about 300 mM L-arginine; (e) about 2.5 mM to about 10 mM calcium chloride dihydrate; and (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate.

[0011] In some embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5% (w/v) to about 7.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM histidine. In some embodiments, the pharmaceutical composition comprises at least 150 mM arginine. In some embodiments, the pharmaceutical composition comprises about 150 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises at least 250 mM arginine. In some embodiments, the pharmaceutical composition comprises about 200 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises about 2.5 mM to about 10 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises polysorbate 20. In some embodiments, the pharmaceutical composition comprises polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0012] In some embodiments, the composition comprises at least about 250 mM L-arginine. In some embodiments, the composition comprises at least about 250 mM L-arginine-HCl. In some embodiments, the composition comprises about 250 mM L-arginine. In some embodiments, the composition comprises about 250 mM L-arginine-HCl. In some embodiments, the polysorbate is polysorbate 80. In some embodiments, the polysorbate is polysorbate 20.

[0013] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
- (b) sucrose;
- (c) histidine;
- (d) arginine;
- (e) calcium chloride; and
- (f) a polysorbate.

[0014] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
- (b) about 1% (w/v) to about 4% (w/v) sucrose;
- (c) about 5 mM to about 15 mM histidine;
- (d) about 150 mM to about 300 mM arginine;
- (e) about 2.5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate.

[0015] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 150 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0016] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second

polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;

- (b) about 1% (w/v) to about 4% (w/v) sucrose;
- (c) about 5 mM to about 15 mM histidine;
- (d) about 200 mM to about 300 mM arginine;
- (e) about 2.5 mM to about 10 mM calcium chloride; and
- (f) about 0.008% (w/v) to about 0.1% (w/v) of a polysorbate.

[0017] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 200 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0018] In some embodiments, the pharmaceutical composition comprises about 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein. In some embodiments, the first polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 1 and the second polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains. In some embodiments, the chimeric protein is efanesoctocog alfa.

[0019] In some embodiments, the pharmaceutical composition comprises about 1% (w/v) to about 4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM histidine. In some embodiments, the pharmaceutical composition comprises at least 150 mM arginine. In some embodiments, the pharmaceutical composition comprises about 150 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises at least 250 mM arginine. In some embodiments, the pharmaceutical composition comprises about 200 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises about 2.5 mM to about 10 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises polysorbate 20. In some embodiments, the pharmaceutical composition comprises polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0020] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) about 1% (w/v) to about 4% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;

- (c) about 150 mM to about 300 mM arginine;
 - (d) about 2.5 mM to about 10 mM calcium chloride; and
 - (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.
- [0021] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 1% (w/v) to about 4% (w/v) sucrose;
 - (b) about 5 mM to about 15 mM histidine;
 - (c) about 200 mM to about 300 mM arginine;
 - (d) about 2.5 mM to about 10 mM calcium chloride; and
 - (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.
- [0022] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 2% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 250 mM arginine;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 20 or polysorbate 80.
- [0023] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 2% (w/v) sucrose;
 - (b) about 10 mM L-histidine;
 - (c) about 250 mM L-arginine-HCl;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 80.
- [0024] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.4 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0025] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 22.45 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0026] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 22.45 mg/ml sucrose;

- (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.62 mg/ml calcium chloride; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0027] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0028] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 20 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 52.7 mg/mL L-arginine-HCl;
 - (d) about 0.7 mg/mL calcium chloride dihydrate; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0029] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 20 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 52.7 mg/mL L-arginine-HCl;
 - (d) about 0.6 mg/mL calcium chloride; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0030] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 20 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 43.6 mg/mL L-arginine;
 - (d) about 0.7 mg/mL calcium chloride dihydrate; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0031] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 20 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 43.6 mg/mL L-arginine;
 - (d) about 0.6 mg/mL calcium chloride; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0032] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0033] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.4 mg/mL to 0.8 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0034] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 60 mg/mL L-arginine;
 - (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0035] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 60 mg/mL L-arginine;
 - (d) 0.4 mg/mL to 0.7 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0036] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 22.45 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0037] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 22.45 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 48.88 mg/ml L-arginine;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.

[0038] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0039] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0040] In some embodiments, the pharmaceutical composition comprises about 5% (w/v) to about 7.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM histidine. In some embodiments, the pharmaceutical composition comprises at least 150 mM arginine. In some embodiments, the pharmaceutical composition comprises about 150 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises at least 250 mM arginine. In some embodiments, the pharmaceutical composition comprises about 200 mM to about 300 mM arginine. In some embodiments, the pharmaceutical composition comprises about 2.5 mM to about 10 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises polysorbate 20. In some embodiments, the pharmaceutical composition comprises polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0041] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 150 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0042] In some embodiments, disclosed herein is a pharmaceutical composition comprising:

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 200 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and

- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.
- [0043] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 5% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 250 mM arginine;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 20 or polysorbate 80.
- [0044] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 5% (w/v) sucrose;
 - (b) about 10 mM L-histidine;
 - (c) about 250 mM L-arginine-HCl;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 80.
- [0045] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.4 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0046] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0047] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.62 mg/ml calcium chloride; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0048] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;

- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0049] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 50 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 52.7 mg/mL L-arginine-HCl;
 - (d) about 0.7 mg/mL calcium chloride dihydrate; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0050] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 50 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 52.7 mg/mL L-arginine-HCl;
 - (d) about 0.6 mg/mL calcium chloride; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0051] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 50 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 43.6 mg/mL L-arginine;
 - (d) about 0.7 mg/mL calcium chloride dihydrate; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0052] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) about 50 mg/mL sucrose;
 - (b) about 1.6 mg/mL L-histidine;
 - (c) about 43.6 mg/mL L-arginine;
 - (d) about 0.6 mg/mL calcium chloride; and
 - (e) about 0.5 mg/mL polysorbate 80.
- [0053] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0054] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;

- (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.4 mg/mL to 0.8 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0055] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 60 mg/mL L-arginine;
 - (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0056] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 45 mg/mL to 60 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 60 mg/mL L-arginine;
 - (d) 0.4 mg/mL to 0.7 mg/mL calcium chloride dihydrate; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
- [0057] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0058] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 48.88 mg/ml L-arginine;
 - (d) 0.82 mg/ml calcium chloride dihydrate; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0059] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.62 mg/ml calcium chloride; and
 - (e) 0.56 mg/ml polysorbate 80.
- [0060] In some embodiments, disclosed herein is a pharmaceutical composition comprising:
- (a) 56.12 mg/ml sucrose;

- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0061] In some embodiments, disclosed herein is a pharmaceutical composition according to any of the disclosed embodiments, which further comprises a pH of about 6.5 to about 7.5. In some embodiments, the pharmaceutical composition disclosed herein has a pH of about 6.8 to about 7.3. In some embodiments, the pharmaceutical composition disclosed herein has a pH of about 7.0. In some embodiments, the pharmaceutical composition disclosed herein has a pH of about 6.8. In some embodiments the pharmaceutical composition disclosed herein does not comprise NaCl. In some embodiments, the pharmaceutical composition does not comprise NaOH. In some embodiments, the pharmaceutical composition does not comprise sodium ions. In some embodiments the pharmaceutical composition disclosed herein comprises less than 8.8 mg/mL sodium chloride (NaCl). In some embodiments, the pharmaceutical composition disclosed herein comprises L-histidine. In some embodiments, the pharmaceutical composition disclosed herein comprises L-arginine. In some embodiments, the pharmaceutical composition disclosed herein comprises arginine-HCl. In some embodiments, the pharmaceutical composition disclosed herein comprises L-arginine-HCl. In some embodiments, the pharmaceutical composition disclosed herein comprises calcium chloride dihydrate.

[0062] In some embodiments, disclosed herein is a pharmaceutical composition according to any of the disclosed embodiments, wherein the pharmaceutical composition has a chimeric protein concentration of about 0.8 to about 1.2 mg/mL. In some embodiments, the pharmaceutical composition disclosed herein comprises 75 IU/mL to 2,000 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition disclosed herein has an osmolality about 525 to about 725 mOsm/kg. In some embodiments, the pharmaceutical composition disclosed herein has an osmolality about 600 to about 650 mOsm/kg. In some embodiments, the pharmaceutical composition disclosed herein has a turbidity of less than about 7 Nephelometric Turbidity Units (NTU).

[0063] Also disclosed herein is a method of treating hemophilia A in a subject in need thereof, comprising administering to the subject an effective amount of the pharmaceutical composition according to any of the embodiments disclosed herein. In some embodiments, the pharmaceutical composition is self-administered. In some embodiments, the pharmaceutical composition is administered intravenously. In some embodiments, the pharmaceutical composition is administered intravenously at a dose of about 20 IU/kg to about 70 IU/kg. In some embodiments, the pharmaceutical composition is administered intravenously at a dose of about 50 IU/kg. In

some embodiments, the pharmaceutical composition is administered intravenously once every 7-10 days. In some embodiments, the pharmaceutical composition is administered intravenously once weekly.

[0064] Also disclosed herein is a pharmaceutical kit comprising (i) a first container comprising a lyophilized pharmaceutical composition comprising

(a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;

(b) sucrose;

(c) histidine;

(d) arginine;

(e) calcium chloride; and

(f) polysorbate 20 or polysorbate 80, and

(ii) a second container comprising sterile water.

[0065] In some embodiments, the chimeric protein comprises a first polypeptide chain comprising the amino acid sequence set forth as SEQ ID NO: 1 and a second polypeptide chain comprising the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.

[0066] Also disclosed herein is a pharmaceutical kit comprising (i) a first container comprising a lyophilized pharmaceutical composition comprising

(a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;

(b) about 30 mg to about 135 mg sucrose;

(c) about 2.5 mg to about 7.5 mg histidine;

(d) about 140 mg to about 200 mg arginine;

(e) about 1.5 mg to about 5 mg calcium chloride; and

(f) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80, and

(ii) a second container comprising sterile water.

In some embodiments, the chimeric protein comprises a first polypeptide chain comprising the amino acid sequence set forth as SEQ ID NO: 1 and a second polypeptide chain comprising the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the

second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.

[0067] In some embodiments, the pharmaceutical kit disclosed herein comprises a lyophilized pharmaceutical composition comprising:

- (a) about 30 mg to about 135 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg histidine;
- (c) about 140 mg to about 200 mg arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80.

[0068] In some embodiments the pharmaceutical kit comprises a lyophilized pharmaceutical composition that does not comprise NaCl. In some embodiments the lyophilized pharmaceutical composition comprises less than 8.8 mg/mL sodium chloride (NaCl). In some embodiments, the pharmaceutical composition does not comprise NaOH. In some embodiments, the pharmaceutical composition does not comprise sodium ions. In some embodiments, the lyophilized pharmaceutical composition comprises L-histidine. In some embodiments, the lyophilized pharmaceutical composition comprises L-arginine. In some embodiments, the lyophilized pharmaceutical composition comprises arginine-HCl. In some embodiments, the lyophilized pharmaceutical composition comprises L-arginine-HCl. In some embodiments, the lyophilized pharmaceutical composition disclosed herein comprises calcium chloride dihydrate.

[0069] In some embodiments, the pharmaceutical kit comprises a lyophilized pharmaceutical composition comprising:

- (a) about 67.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 177.3 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0070] In some embodiments, the pharmaceutical kit comprises a lyophilized pharmaceutical composition comprising:

- (a) about 67.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 146.6 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0071] In some embodiments, the pharmaceutical kit disclosed herein comprises a lyophilized pharmaceutical composition comprising:

- (a) about 160 mg to about 200 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg histidine;
- (c) about 140 mg to about 200 mg arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80.

[0072] In some embodiments the pharmaceutical kit comprises a lyophilized pharmaceutical composition that does not comprise NaCl. In some embodiments the lyophilized pharmaceutical composition comprises less than 8.8 mg/mL sodium chloride (NaCl). In some embodiments, the pharmaceutical composition does not comprise NaOH. In some embodiments, the pharmaceutical composition does not comprise sodium ions. In some embodiments, the lyophilized pharmaceutical composition comprises L-histidine. In some embodiments, the lyophilized pharmaceutical composition comprises L-arginine. In some embodiments, the lyophilized pharmaceutical composition comprises arginine-HCl. In some embodiments, the lyophilized pharmaceutical composition comprises L-arginine-HCl. In some embodiments, the lyophilized pharmaceutical composition disclosed herein comprises calcium chloride dihydrate.

[0073] In some embodiments, the pharmaceutical kit comprises a lyophilized pharmaceutical composition comprising:

- (a) about 168.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 177.3 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0074] In some embodiments, the pharmaceutical kit comprises a lyophilized pharmaceutical composition comprising:

- (a) about 168.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 146.6 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0075] In some embodiments, the pharmaceutical kit comprises a lyophilized pharmaceutical composition having a moisture content of less than 2%. In some embodiments, the lyophilized pharmaceutical composition has a moisture content of less than 1.8%. In some embodiments, the lyophilized pharmaceutical composition has a moisture content of less than 1.6%. In some embodiments, the lyophilized pharmaceutical composition is in a lyophilized cake. In some embodiments, the lyophilized cake is white. In some embodiments, the lyophilized cake is less

than Y4 in the European Pharmacopoeia color scale. In some embodiments, the lyophilized pharmaceutical composition is a powder.

[0076] In some embodiments, the pharmaceutical kit comprises a first container comprising 100 IU to 10,000 IU of the chimeric protein. In some embodiments, the first container comprises 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein.

[0077] In some embodiments, the pharmaceutical kit further comprises instructions for combining the lyophilized pharmaceutical composition and sterile water. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the lyophilized pharmaceutical composition is reconstituted within 7 to 12 seconds.

[0078] In some embodiments, the pharmaceutical composition comprises a second container comprising sterilized water at a volume sufficient to produce, when combined with the lyophilized powder of the first container, a solution for injection disclosed herein is obtained.

[0079] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 525 to about 725 mOsm/kg. In some embodiments, the osmolality of the resulting solution is about 600 to about 650 mOsm/kg. In some embodiments, the pH of the resulting solution is about 6.5 to about 7.5. In some embodiments, the pH of the resulting solution is about 7.0. In some embodiments, the pH of the resulting solution is about 6.8. In some embodiments, the the turbidity of the resulting solution is less than about 7 Nephelometric Turbidity Units (NTU). In some embodiments, the protein concentration of the resulting solution is about 0.8 to about 1.2 mg/mL. In some embodiments, less than 3% of the protein is aggregated.

[0080] In some embodiments, the pharmaceutical kit comprises a second container comprising sterilized water at a volume sufficient to produce, when combined with the lyophilized pharmaceutical composition of the first container, a solution comprising:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

[0081] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;

- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

[0082] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0083] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0084] In some embodiments, the pharmaceutical kit comprises a second container comprising sterilized water at a volume sufficient to produce, when combined with the lyophilized pharmaceutical composition of the first container, a solution comprising:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

[0085] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

[0086] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0087] In some embodiments of the pharmaceutical kit disclosed herein, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

[0088] In some embodiments, the pharmaceutical kit comprises a second container comprising about 2 mL to about 5 mL of sterile water. In some embodiments, the pharmaceutical kit comprises a second container comprising about 3 mL of sterile water. In some embodiments, the pharmaceutical kit comprises a second container comprising about 3.3 mL of sterile water.

[0089] In some embodiments, the pharmaceutical kit comprises a first container which is a glass vial comprising a rubber stopper. In some embodiments, the pharmaceutical kit comprises a second container which is a syringe body. In some embodiments, the sterile water is in the syringe body. In some embodiments, the syringe body is associated with a plunger. In some embodiments, the pharmaceutical kit further comprises an adaptor to connect the glass vial to the syringe body. In some embodiments, the pharmaceutical kit further comprises infusion tubing associated with a needle to be connected to the syringe body, suitable for intravenous infusion.

[0090] Also disclosed herein is a method of treating hemophilia A in a subject in need thereof, comprising combining the lyophilized pharmaceutical composition and the sterile water of the pharmaceutical kit according to any of the embodiments disclosed herein, and administering to the subject an effective amount of the resulting combination (i.e. solution). In some embodiments, the subject combines the lyophilized pharmaceutical composition and the sterile water of the kit. In some embodiments, the combination is self-administered by the subject.

BRIEF DESCRIPTION OF THE DRAWINGS/FIGURES

[0091] FIG. 1 is a schematic representation of efanesoctocog alfa, an exemplary FVIII-ELNN-Fc/D'D3-ELNN-Fc heterodimer. FVIII: factor VIII; VWF: von Willebrand Factor; A1, A2, A3, C1, C2: domains of FVIII; D'D3: domains of VWF; Fc: Fc region of immunoglobulin constant region.

[0092] FIG. 2 shows the pH of pharmaceutical compositions containing various concentrations of L-histidine (10 mM, 20 mM, 50 mM) at efanesoctocog alfa concentrations of 1 mg/mL, 0.67 mg/mL (4000 IU/mL), and 0.045 mg/mL (250 IU/mL). Controls are provided with compositions containing no DS, no excipients or compositions containing only excipients. Abbreviations: DP: drug product. BDP: bulk drug product. DS: drug substance. IU: International Units.

[0093] FIG. 3 shows the measured turbidity (NTU) for pharmaceutical compositions containing various concentrations of L-histidine (10 mM, 20 mM, 50 mM) at efanesoctocog alfa concentrations of 1 mg/mL, 0.67 mg/mL (4000 IU/mL), and 0.045 mg/mL (250 IU/mL). Results of compositions at 5, 25, and 40°C are depicted. Time points were T0 and 1 week.

[0094] FIG. 4 shows the aggregation levels (%HMWS) for pharmaceutical compositions containing various concentrations of L-histidine (10 mM, 20 mM, 50 mM) at efanesoctocog alfa concentrations of 1 mg/mL, 0.67 mg/mL (4000 IU/mL), and 0.045 mg/mL (250 IU/mL). Results of compositions at 5 and 25°C are depicted. Time points were T0 and 1 week.

[0095] FIG. 5 shows the efanesoctocog alfa protein concentration (µg/mL) measured for each of the 13 buffer compositions tested. Compositions were tested at both 2 – 8 °C and 30 °C. Time points measured are start of experiment (T0), one month (T1), and 3 months (T3), and 6 months (T6). Abbreviations: DP: drug product. DDS: diluted drug substance. DoE: design of experiment.

[0096] FIG. 6 shows the efanesoctocog alfa specific activity (IU/mg) measured for each of the 13 buffer compositions tested. Compositions were tested at both 2 – 8 °C and 30 °C. Time points measured are start of experiment (T0), one month (T1), and 3 months (T3). Abbreviations: DP: drug product. DDS: diluted drug substance. DoE: design of experiment.

[0097] FIG. 7 shows the efanesoctocog alfa aggregation levels (%HMWS) measured for each of the 13 buffer compositions tested. Compositions were tested at both 2 – 8 °C and 30 °C. Time points measured are start of experiment (T0), one month (T1), and 3 months (T3), and 6 months (T6). The 5% specification limit is noted on the graph. Abbreviations: DP: drug product. DDS: diluted drug substance. DoE: design of experiment.

[0098] FIG. 8 shows the glass transition temperature (Tg) measured using DSC for each of the 13 buffer compositions tested. Compositions were tested at both 2 – 8 °C and 30 °C. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), 6 months (T6), and 12 months (T12). Abbreviations: DP: drug product. DDS: diluted drug substance. DoE: design of experiment.

[0099] FIG. 9 shows the aggregation levels (%HMWS) over time (hours) of efanesoctocog alfa drug substance (DS) compositions (1 mg/mL) containing 5% sucrose (dashed line) or 1% sucrose (solid line) at room temperature (RT)/room light (RL) conditions.

[0100] FIG. 10 shows the aggregation levels (%HMWS) over time (hours) of efanesoctocog alfa drug substance (DS) compositions (1 mg/mL) containing 5% sucrose (dashed line) or 1% sucrose (solid line) at 2-8 °C.

[0101] FIG. 11 shows the aggregation levels (%HMWS) over time (hours) of 250 IU efanesoctocog alfa bulk drug product (BDP) compositions containing 0, 1, 2, and 5% w/v sucrose concentrations at room temperature (RT)/room light (RL) conditions.

[0102] FIG. 12 shows the aggregation levels (%HMWS) over time (hours) of 250 IU efanesoctocog alfa bulk drug product (BDP) compositions containing 0, 1, 2, and 5% w/v sucrose concentrations at 2-8 °C.

[0103] FIG. 13 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU at 5 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0104] FIG. 14 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU at 30 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0105] FIG. 15 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU at 40 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0106] FIG. 16 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 4000 IU at 5 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0107] FIG. 17 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 4000 IU at 30 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0108] FIG. 18 shows the aggregation levels (%HMWS) of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 4000 IU at 40 °C. Lyo DP compositions containing 0, 1, 2,

and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months.

[0109] FIG. 19 shows aggregation levels (%HMWS) over time (hours) of efanesoctocog alfa DS and 4000 IU bulk drug product (BDP) liquid compositions containing either 2% w/v sucrose (solid line) or 5% w/v sucrose (dashed line). Samples were held at room temperature or 5°C and tested after 0, 5, 7, 25, 43, and 55 hours.

[0110] FIG. 20 shows the glass transition temperature (T_g) measured using DSC for compositions of efanesoctocog alfa DP at 250 IU or 4000 IU at 5 °C. Compositions containing 0, 1, 2, and 5% w/v sucrose were tested at both strengths. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0111] FIG. 21 shows the glass transition temperature (T_g) measured using DSC for compositions of efanesoctocog alfa DP at 250 IU or 4000 IU at 30 °C. Compositions containing 0, 1, 2, and 5% w/v sucrose were tested at both strengths. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0112] FIG. 22 shows the glass transition temperature (T_g) measured using DSC for compositions of efanesoctocog alfa DP at 250 IU or 4000 IU at 40 °C. Compositions containing 0, 1, 2, and 5% w/v sucrose were tested at both strengths. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0113] FIG. 23 shows the glass transition temperature (T_g) using DSC of efanesoctocog alfa liquid BDP at 250 IU or 4000 IU. Compositions containing 0, 1, 2, and 5% w/v sucrose were tested at both strengths.

[0114] FIG. 24 shows the residual moisture content of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU or 4000 IU at 5 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0115] FIG. 25 shows the residual moisture content of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU or 4000 IU at 30 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0116] FIG. 26 shows the residual moisture content of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU or 4000 IU at 40 °C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Time points measured are start of experiment (T0), one month (T1), 3 months (T3), and 6 months (T6).

[0117] FIG. 27 is a graphical depiction of the percentage of high molecular weight (HMW) aggregates of efanesoctocog alfa (BIVV001) as analyzed by size exclusion chromatography

(SEC) with buffers of varying arginine concentrations (Experiment 1). Data was collected at 0 minutes, 40 minutes, 80 minutes, 120 minutes, and 160 minutes post-thaw.

[0118] FIG. 28 is a graphical depiction of the percentage of high molecular weight (HMW) aggregates of efanesoctocog alfa (BIVV001) as analyzed by size exclusion chromatography (SEC) with buffers of varying arginine concentrations (Experiment 2). Data was collected at 0 minutes, 40 minutes, 80 minutes, 120 minutes, and 160 minutes post-thaw.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0119] The present disclosure is directed to, *inter alia*, formulations (including aqueous and lyophilized formulations, as well as related kits) comprising FVIII proteins. In some embodiments, the FVIII protein is a chimeric FVIII protein such as efanesoctocog alfa comprising two polypeptides, i.e., a first polypeptide comprising a FVIII protein comprising a first ELNN Polypeptide sequence insert fused to a first Fc region, and a second polypeptide comprising a VWF protein fused to a second Ig constant region by a second ELNN Polypeptide sequence, wherein the first ELNN Polypeptide sequence contains about 288 amino acids and the second ELNN Polypeptide sequence contains about 144 amino acids, and the first Ig constant region and the second Ig constant region are covalently linked together by disulfide bonds.

[0120] The present disclosure provides formulations (including aqueous and lyophilized formulations, as well as related kits) for a chimeric protein comprising (i) a factor VIII (FVIII) polypeptide and (ii) a von Willebrand factor (VWF) fragment comprising a D' domain of VWF and a D3 domain of VWF. Included herein are compositions that may be lyophilized, as well as compositions formed upon reconstitution of lyophilized formulations with a diluent. Therapeutic methods and uses are also provided.

I. Definitions

[0121] It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, "a nucleotide sequence," is understood to represent one or more nucleotide sequences. As such, the terms "a" (or "an"), "one or more," and "at least one" can be used interchangeably herein.

[0122] Furthermore, "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term "and/or" as used in a phrase such as "A and/or B" herein is intended to include "A and B," "A or B," "A" (alone), and "B" (alone). Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following aspects: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[0123] It is understood that wherever aspects are described herein with the language "comprising," otherwise analogous aspects described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0124] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, may provide one of skill with a general dictionary of many of the terms used in this disclosure.

[0125] Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation. The headings provided herein are not limitations of the various aspects of the disclosure. Accordingly, the terms defined immediately below are more fully defined by reference to the specification in its entirety.

[0126] The term "about" is used herein to mean approximately, roughly, around, or in the regions of. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" can modify a numerical value above and below the stated value by a variance of, e.g., 10 percent, up or down (higher or lower). In some embodiments, the term indicates deviation from the indicated numerical value by $\pm 10\%$, $\pm 5\%$, $\pm 4\%$, $\pm 3\%$, $\pm 2\%$, $\pm 1\%$, $\pm 0.9\%$, $\pm 0.8\%$, $\pm 0.7\%$, $\pm 0.6\%$, $\pm 0.5\%$, $\pm 0.4\%$, $\pm 0.3\%$, $\pm 0.2\%$, $\pm 0.1\%$, $\pm 0.05\%$, or $\pm 0.01\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 10\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 5\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 4\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 3\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 2\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 1\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.9\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.8\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.7\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.6\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.5\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.4\%$. In some embodiments,

"about" indicates deviation from the indicated numerical value by $\pm 0.3\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.1\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.05\%$. In some embodiments, "about" indicates deviation from the indicated numerical value by $\pm 0.01\%$.

[0127] Depending on context, the term "polynucleotide" or "nucleotide" may encompass a singular nucleic acid as well as plural nucleic acids. In some embodiments, a polynucleotide is an isolated nucleic acid molecule or construct, e.g., messenger RNA (mRNA) or plasmid DNA (pDNA). In some embodiments, a polynucleotide comprises a conventional phosphodiester bond. In some embodiments, a polynucleotide comprises a non-conventional bond (e.g., an amide bond, such as found in peptide nucleic acids (PNA)). The term "nucleic acid" may refer to any one or more nucleic acid segments, e.g., DNA or RNA fragments, present in a polynucleotide. By "isolated" nucleic acid or polynucleotide is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, a recombinant polynucleotide encoding a Factor VIII polypeptide contained in a vector is considered isolated for the purposes of the present disclosure. Further examples of an isolated polynucleotide include recombinant polynucleotides maintained in heterologous host cells or purified (partially or substantially) from other polynucleotides in a solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of polynucleotides of the present disclosure. Isolated polynucleotides or nucleic acids according to the present disclosure further include such molecules produced synthetically. In addition, a polynucleotide or a nucleic acid can include regulatory elements such as promoters, enhancers, ribosome binding sites, or transcription termination signals.

[0128] Certain proteins secreted by mammalian cells are associated with a secretory signal peptide which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Those of ordinary skill in the art are aware that signal peptides are generally fused to the N-terminus of the polypeptide, and are cleaved from the complete or "full-length" polypeptide to produce a secreted or "mature" form of the polypeptide. In some embodiments, a native signal peptide or a functional derivative of that sequence that retains the ability to direct the secretion of the polypeptide that is operably associated with it. Alternatively, a heterologous mammalian signal peptide, e.g., a human tissue plasminogen activator (TPA) or mouse β -glucuronidase signal peptide, or a functional derivative thereof, can be used.

[0129] As used herein, the term "polypeptide" is intended to encompass a singular "polypeptide" as well as plural "polypeptides," and refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term "polypeptide" refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the

product. Thus, peptides, dipeptides, tripeptides, oligopeptides, "protein," "amino acid chain," or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of "polypeptide," and the term "polypeptide" can be used instead of, or interchangeably with any of these terms. The term "polypeptide" is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non-naturally occurring amino acids. A polypeptide can be derived from a natural biological source or produced recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It can be generated in any manner, including by chemical synthesis.

[0130] An "isolated" polypeptide or a fragment, variant, or derivative thereof refers to a polypeptide that is not in its natural milieu. No particular level of purification is required. For example, an isolated polypeptide can simply be removed from its native or natural environment. Recombinantly produced polypeptides and proteins expressed in host cells are considered isolated for the purpose of the disclosure, as are native or recombinant polypeptides which have been separated, fractionated, or partially or substantially purified by any suitable technique.

[0131] Also included in the present disclosure are fragments or variants of polypeptides, and any combination thereof. The term "fragment" or "variant" when referring to polypeptide binding domains or binding molecules of the present disclosure include any polypeptides which retain at least some of the properties (*e.g.*, FcRn binding affinity for an FcRn binding domain or Fc variant, coagulation activity for an FVIII variant, or FVIII binding activity for the VWF fragment) of the reference polypeptide. Fragments of polypeptides include proteolytic fragments, as well as deletion fragments, in addition to specific antibody fragments discussed elsewhere herein, but do not include the naturally occurring full-length polypeptide (or mature polypeptide). Variants of polypeptide binding domains or binding molecules of the present disclosure include fragments as described above, and also polypeptides with altered amino acid sequences due to amino acid substitutions, deletions, or insertions. Variants can be naturally or non-naturally occurring. Non-naturally occurring variants can be produced using art-known mutagenesis techniques. Variant polypeptides can comprise conservative or non-conservative amino acid substitutions, deletions or additions.

[0132] The term "VWF protein" as used herein means any VWF fragment that interacts with FVIII and retains at least one or more properties that are normally provided to FVIII by full-length VWF, *e.g.*, preventing premature activation to FVIIIa, preventing premature proteolysis, preventing clearance, preventing association with phospholipid membranes that could lead to premature clearance, preventing binding to FVIII clearance receptors that can bind naked FVIII

but not VWF-bound FVIII, and/or stabilizing the FVIII heavy chain and light chain interactions. A VWF fragment referred to herein is a VWF polypeptide that is less than the full-length VWF protein, wherein the VWF fragment retains the ability to interact with and/or bind to FVIII. In some embodiments, a VWF protein is a fragment (which may be mutated) of full-length VWF that binds to a FVIII protein such that the FVIII protein has reduced binding to, or does not bind, full length VWF (e.g., endogenous VWF in a subject).

[0133] A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, if an amino acid in a polypeptide is replaced with another amino acid from the same side chain family, the substitution is considered to be conservative. In some embodiments, a string of amino acids can be conservatively replaced with a structurally similar string that differs in order and/or composition of side chain family members.

[0134] As known in the art, "sequence identity" between two polypeptides is determined by comparing the amino acid sequence of one polypeptide to the sequence of a second polypeptide. Similarly, "sequence identity" between two polynucleotides is determined by comparing the nucleotide sequence of one polynucleotide to the sequence of a second polynucleotide. The terms "% identical", "% identity" or similar terms are intended to refer, in particular, to the percentage of nucleotides or amino acids (as applicable) which are identical in an optimal alignment between the sequences to be compared. Said percentage is purely statistical, and the differences between the two sequences may be but are not necessarily randomly distributed over the entire length of the sequences to be compared. Comparisons of two sequences are usually carried out by comparing the sequences, after optimal alignment, with respect to a segment or "window of comparison", in order to identify local regions of corresponding sequences. For example, the optimal alignment for a comparison may be carried out manually or with the aid of the local homology algorithm by Smith and Waterman, 1981, *Adv. App. Math.* 2, 482, with the aid of the local homology algorithm by Needleman and Wunsch, 1970, *J. Mol. Biol.* 48, 443, with the aid of the similarity search algorithm by Pearson and Lipman, 1988, *Proc. Natl Acad. Sci. USA* 88, 2444, or with the aid of computer programs using said algorithms (GAP, BESTFIT, FASTA, BLAST P, BLAST N and TFASTA in Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.). In some embodiments, percent identity of two sequences is

determined using the BLASTN or BLASTP algorithm, as available on the United States National Center for Biotechnology Information (NCBI) website (e.g., at blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch&BLAST_SPEC=blast2seq&LINK_LOC=align2seq). In some embodiments, the algorithm parameters used for BLASTN algorithm on the NCBI website include: (i) Expect Threshold set to 10; (ii) Word Size set to 28; (iii) Max matches in a query range set to 0; (iv) Match/Mismatch Scores set to 1, -2; (v) Gap Costs set to Linear; and (vi) the filter for low complexity regions being used. In some embodiments, the algorithm parameters used for BLASTP algorithm on the NCBI website include: (i) Expect Threshold set to 10; (ii) Word Size set to 3; (iii) Max matches in a query range set to 0; (iv) Matrix set to BLOSUM62; (v) Gap Costs set to Existence: 11 Extension: 1; and (vi) conditional compositional score matrix adjustment. When discussed herein, whether any particular polypeptide is at least about 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 99%, or 100% identical to another polypeptide can be determined using methods and computer programs/software known in the art such as, but not limited to, the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). BESTFIT uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981), to find the best segment of homology between two sequences. When using BESTFIT or any other sequence alignment program to determine whether a particular sequence is, for example, 95% identical to a reference sequence according to the present disclosure, the parameters are set, of course, such that the percentage of identity is calculated over the full-length of the reference polypeptide sequence and that gaps in homology of up to 5% of the total number of amino acids in the reference sequence are allowed.

[0135] As used herein, an "amino acid corresponding to" or an "equivalent amino acid" in a VWF sequence or a FVIII protein sequence is identified by alignment to maximize the identity or similarity between a first VWF or FVIII sequence and a second VWF or FVIII sequence. The number used to identify an equivalent amino acid in a second VWF or FVIII sequence is based on the number used to identify the corresponding amino acid in the first VWF or FVIII sequence.

[0136] As used herein, the term "insertion site" refers to a position in a FVIII polypeptide, or fragment, variant, or derivative thereof, which is immediately downstream of the position at which a half-life extending moiety or heterologous moiety can be inserted. An "insertion site" is specified as a number, the number being the number of the amino acid in mature native FVIII (SEQ ID NO: 8) to which the insertion site corresponds, which is immediately C-terminal to the position of the insertion. For example, the phrase "comprises an ELNN Polypeptide at an insertion site which corresponds to amino acid 1656 of SEQ ID NO: 8" indicates that the heterologous moiety is

located between two amino acids corresponding to amino acid 1656 and amino acid 1657 of SEQ ID NO: 8.

[0137] The phrase "immediately downstream of an amino acid" as used herein refers to position right next to the terminal carboxyl group of the amino acid. For example, an insertion site immediately downstream of amino acid 745 corresponding to the mature wild type FVIII protein (SEQ ID NO: 8) means that the insertion site is between amino acid 745 and amino acid 746 corresponding to the mature wild type FVIII protein. Similarly, the phrase "immediately upstream of an amino acid" refers to the position right next to the terminal amine group of the amino acid.

[0138] The phrase "between two amino acids of an insertion site" as used herein refers to a position in which an ELNN Polypeptide or any other polypeptide is inserted between two adjacent amino acids. Thus, the phrases "inserted immediately downstream of an amino acid" and "inserted between two amino acids of an insertion site" are used synonymously with "inserted at an insertion site."

[0139] The terms "inserted," "is inserted," "inserted into" or grammatically related terms, as used herein with respect to insertions of ELNN Polypeptide into FVIII refers to the position of an ELNN Polypeptide in a chimeric protein relative to the analogous position in native mature human FVIII. As used herein the terms refer to the characteristics of the recombinant FVIII polypeptide relative to native mature human FVIII, and do not indicate, imply or infer any methods or process by which the chimeric protein was made. For example, in reference to a chimeric protein provided herein, the phrase "an ELNN Polypeptide is inserted immediately downstream of residue 745 of the FVIII polypeptide" means that the chimeric protein comprises an ELNN Polypeptide immediately downstream of an amino acid which corresponds to amino acid 745 in native mature human FVIII, e.g., bounded by amino acids corresponding to amino acids 745 and 746 of native mature human FVIII (without requiring the presence of an amino acid corresponding to 746 of native mature human FVIII), and does not connote an order or method of production for which the chimeric protein was constructed.

[0140] As used herein, the terms "ELNN Polypeptide" and "ELNN" are synonymous, and refer to extended length polypeptides with non-naturally occurring, substantially non-repetitive sequences that are composed mainly of small hydrophilic amino acids, with the sequence having a low degree or no secondary or tertiary structure under physiologic conditions. ELNNs can confer certain desirable pharmacokinetic, physicochemical and pharmaceutical properties when linked to a VWF protein or a FVIII sequence of the disclosure to create a chimeric polypeptide. Such desirable properties include but are not limited to enhanced pharmacokinetic parameters and solubility characteristics. As used herein, the terms "ELNN Polypeptide" and "ELNN" specifically exclude antibodies or antibody fragments such as single-chain antibodies or Fc fragments of a

light chain or a heavy chain. ELNN polypeptides are known in the art, and non-limiting descriptions relating to and examples of ELNN polypeptides known as XTEN® polypeptides are available in Schellenberger et al., (2009) *Nat Biotechnol* 27(12):1186-90; Brandl et al., (2020) *Journal of Controlled Release* 327:186-197; and Radon et al., (2021) *Advanced Functional Materials* 31, 2101633 (pages 1-33), the entire contents of each of which are incorporated herein by reference.

[0141] A "fusion" or "chimeric" protein comprises a first amino acid sequence linked to a second amino acid sequence with which it is not naturally linked in nature. The amino acid sequences which normally exist in separate proteins can be brought together in the fusion polypeptide, or the amino acid sequences which normally exist in the same protein can be placed in a new arrangement in the fusion polypeptide, e.g., fusion of a Factor VIII domain of the disclosure with an Ig Fc domain. A fusion protein is created, for example, by chemical synthesis, or by creating and translating a polynucleotide in which the peptide regions are encoded in the desired relationship. A chimeric protein can further comprise a second amino acid sequence associated with the first amino acid sequence by a covalent, non-peptide bond or a non-covalent bond.

[0142] With respect to sequences, the term "linked" as used herein refers to a first amino acid sequence or nucleotide sequence covalently or non-covalently joined to a second amino acid sequence or nucleotide sequence, respectively. The first amino acid or nucleotide sequence can be directly joined or juxtaposed to the second amino acid or nucleotide sequence or alternatively an intervening sequence can covalently join the first sequence to the second sequence. Depending on context, the term "linked" means not only a fusion of a first amino acid sequence to a second amino acid sequence at the C-terminus or the N-terminus, but also includes insertion of the whole first amino acid sequence (or the second amino acid sequence) into any two amino acids in the second amino acid sequence (or the first amino acid sequence, respectively). In some embodiments, the first amino acid sequence can be linked to a second amino acid sequence by a peptide bond or a linker. The first nucleotide sequence can be linked to a second nucleotide sequence by a phosphodiester bond or a linker. The linker can be a peptide or a polypeptide (for polypeptide chains) or a nucleotide or a nucleotide chain (for nucleotide chains) or any chemical moiety (for both polypeptide and polynucleotide chains). The term "linked" may also be indicated by a hyphen (-).

[0143] With respect to two polypeptides, the term "associated with" refers to one or more covalent or non-covalent bonds formed between a first polypeptide and a second polypeptide. In some embodiments, the term "associated with" means a covalent, non-peptide bond or a non-covalent bond. This association can be indicated by a colon, i.e., (:). In some embodiments, it means a covalent bond except a peptide bond. For example, the amino acid cysteine comprises a thiol group that can form a disulfide bond or bridge with a thiol group on a second cysteine

residue. In most naturally occurring IgG molecules, the CH1 and CL regions are associated by a disulfide bond and the two heavy chains are associated by two disulfide bonds at positions corresponding to 239 and 242 using the Kabat numbering system (position 226 or 229, EU numbering system). Examples of covalent bonds include, but are not limited to, a peptide bond, a metal bond, a hydrogen bond, a disulfide bond, a sigma bond, a pi bond, a delta bond, a glycosidic bond, an agnostic bond, a bent bond, a dipolar bond, a Pi backbond, a double bond, a triple bond, a quadruple bond, a quintuple bond, a sextuple bond, conjugation, hyperconjugation, aromaticity, hapticity, or antibonding. Non-limiting examples of non-covalent bond include an ionic bond (e.g., cation-pi bond or salt bond), a metal bond, a hydrogen bond (e.g., dihydrogen bond, dihydrogen complex, low-barrier hydrogen bond, or symmetric hydrogen bond), van der Waals force, London dispersion force, a mechanical bond, a halogen bond, aurophilicity, intercalation, stacking, entropic force, or chemical polarity. In some embodiments, the one or more covalent bonds between the first amino acid chain and the second amino acid chain is two disulfide bonds. In some embodiments, the one or more covalent bonds between the first amino acid chain and the second amino acid chain is two disulfide bonds between a first Fc portion on the first amino acid chain and a second Fc portion on the second amino acid chain, wherein the two disulfide bonds occur in the hinge region of the two Fc portions.

[0144] In some embodiments, a polypeptide has an enzymatic cleavage site cleaved by an enzyme that is activated during the clotting cascade, such that cleavage of such sites occurs at the site of clot formation. Exemplary such sites include, e.g., those recognized by thrombin, Factor XIa or Factor Xa. Other enzymatic cleavage sites are known in the art and described in elsewhere herein. In constructs that include more than one processing or cleavage site, it will be understood that such sites can be the same or different. As used herein, the term "half-life" refers to a biological half-life of a particular polypeptide *in vivo*. Half-life can be represented by the time required for half the quantity administered to a subject to be cleared from the circulation and/or other tissues in the animal. In some embodiments, when a clearance curve of a given polypeptide is constructed as a function of time, the curve is usually biphasic with a rapid α -phase and longer β -phase. The α -phase typically represents an equilibration of the administered Fc polypeptide between the intra- and extra-vascular space and is, in part, determined by the size of the polypeptide. The β -phase typically represents the catabolism of the polypeptide in the intravascular space. In some embodiments, FVIII and chimeric proteins comprising FVIII are monophasic, and thus do not have an alpha phase, but just the single beta phase. Therefore, in some embodiments, the term half-life as used herein refers to the half-life of the polypeptide in the β -phase. The typical beta phase half-life of a human antibody in humans is 21 days. In some embodiments, the half-life is expressed as the half-life of the terminal phase.

[0145] In some embodiments, a subject has hemophilia A. In some embodiments, the hemophilia A is severe hemophilia A.

[0146] "Administer" or "administering," as used herein refers to delivering to a subject a composition described herein, e.g., a chimeric protein. The composition, e.g., the chimeric protein, can be administered to a subject using methods known in the art. In particular, the composition can be administered intravenously, subcutaneously, intramuscularly, intradermally, or via any mucosal surface, e.g., orally, sublingually, buccally, nasally, rectally, vaginally or via pulmonary route. In some embodiments, the administration is intravenous. In some embodiments, the administration is subcutaneous. In some embodiments, the administration is self-administration. In some embodiments, a parent administers the chimeric protein to a child. In some embodiments, the chimeric protein is administered to a subject by a healthcare practitioner such as a medical doctor, a medic, or a nurse.

[0147] As used herein, the term "dose" refers to a single administration of a composition to a subject. A single dose can be administered all at once, e.g., as a bolus, or over a period of time, e.g., via an intravenous infusion. The term "multiple doses" means more than one dose, e.g., more than one administration.

[0148] When referring to co-administration of more than one composition, a dose of composition A can be administered concurrently with a dose of composition B. Alternatively, a dose of composition A can be administered before or after a dose of composition B. In some embodiments, composition A and composition B are combined into a single formulation.

[0149] As used herein, the term "interval" or "dosing interval" refers to the amount of time that elapses between a first dose of composition A and a subsequent dose of the same composition administered to a subject. A dosing interval can refer to the time that elapses between a first dose and a second dose, or a dosing interval can refer to the amount of time that elapses between multiple doses.

[0150] The term "dosing frequency" as used herein refers to the number of doses administered per a specific dosing interval. For example, a dosing frequency can be written as once a week, once every two weeks, etc. Therefore, a dosing interval of 7 days can be also written as a dosing interval of once in 7 days or once every week, or once a week.

[0151] As used herein the term "prophylactic treatment" refers to the administration of a therapy for the treatment of hemophilia A, where such treatment is intended to prevent or reduce the severity of one or more symptoms of hemophilia A, e.g., bleeding episodes, e.g., one or more spontaneous bleeding episodes, and/or joint damage. See Jimenez-Yuste et al., Blood Transfus. 12(3):314-19 (2014). To prevent or reduce the severity of such symptoms, e.g., bleeding episodes and the progression of joint disease, hemophilia A patients may receive regular infusions of

clotting factor as part of a prophylactic treatment regimen. The basis of such prophylactic treatment is the observation that hemophilia patients with a clotting factor, e.g., FVIII, level of 1% or more rarely experience spontaneous bleeding episodes and have fewer hemophilia-related comorbidities as compared to patients with severe hemophilia. See, e.g., Coppola A. et al, *Semin. Thromb. Hemost.* 38(1): 79-94 (2012). Health care practitioners treating these hemophilia patients surmised that maintaining factor levels at around 1% with regular infusions could potentially reduce the risk of hemophilia symptoms, including bleeding episodes and joint damage. See *id.* Subsequent research has confirmed these benefits in pediatric hemophilia patients receiving prophylactic treatment with clotting factor, rendering prophylactic treatment the goal for people with severe hemophilia. See *id.*

[0152] A "prophylactic" treatment can also refer to the preemptive administration of the composition described herein, e.g., a protein (such as a chimeric protein), to a subject in order to control, manage, prevent, or reduce the occurrence or severity of one or more symptoms of hemophilia A, e.g., bleeding episodes. In some embodiments, prophylactic treatment with a clotting factor, e.g., FVIII, is used to treat subjects with severe hemophilia A. In some embodiments, prophylactic treatment refers to administering a composition disclosed herein to a subject in need thereof to reduce the occurrence of one or more symptom of hemophilia A. A prophylactic treatment can include administration of multiple doses. The multiple doses used in prophylactic treatment are typically administered at particular dosing intervals. In some embodiments, the annualized bleeding rate can be reduced to less than 10, less than 9, less than 8, less than 7, less than 6, less than 5, less than 4, less than 3, less than 2, or less than 1.

[0153] The term "on-demand treatment" or "episodic treatment" refers to the "as needed" administration of a chimeric molecule in response to symptoms of hemophilia A, e.g., a bleeding episode, or before an activity that can cause bleeding. In some aspects, the on-demand treatment can be given to a subject when bleeding starts, such as after an injury, or when bleeding is expected, such as before surgery. In some aspects, the on-demand treatment can be given prior to activities that increase the risk of bleeding, such as contact sports. In some embodiments, the on-demand treatment is given as a single dose. In some embodiments, the on-demand treatment is given as a first dose, followed by one or more additional doses. When the chimeric protein is administered on-demand, the one or more additional doses can be administered at least about 12 hours, at least about 24 hours, at least about 36 hours, at least about 48 hours, at least about 60 hours, at least about 72 hours, at least about 84 hours, at least about 96 hours, at least about 108 hours, or at least about 120 hours after the first dose. It should be noted, however, that the dosing interval associated with on-demand treatment is not the same as the dosing interval used for prophylactic treatment.

[0154] In some embodiments, the subject in need of a general hemostatic agent is undergoing, or is about to undergo, surgery. The chimeric protein of the disclosure can be administered prior to or after surgery. The chimeric protein of the disclosure can also be administered during or after surgery to control an acute bleeding episode. When the chimeric protein is administered prior to surgery, the administration can be at least about 1 hour, at least about 2 hours, at least about 4 hours, at least about 8 hours, at least about 12 hours, at least about 24 hours, at least about 36 hours, at least about 48 hours, or at least about 72 hours prior to surgery. When the chimeric protein is administered to after surgery, the administration can be at least about 1 hour, at least about 2 hours, at least about 4 hours, at least about 8 hours, at least about 12 hours, at least about 24 hours, at least about 36 hours, at least about 48 hours, or at least about 72 hours after surgery. The surgery can include, but is not limited to, liver transplantation, liver resection, dental procedures, or stem cell transplantation.

[0155] "Treat", "treatment", "treating", as used herein refers to, e.g., the reduction in severity of a disease or condition; the reduction in the duration of a disease course; the amelioration of one or more symptoms associated with a disease or condition; the provision of beneficial effects to a subject with a disease or condition, without necessarily curing the disease or condition, or the prophylaxis of one or more symptoms associated with a disease or condition. In some embodiments, treating or treatment comprises maintaining a FVIII trough level at least about 1 IU/dL, 2 IU/dL, 3 IU/dL, 4 IU/dL, 5 IU/dL, 6 IU/dL, 7 IU/dL, 8 IU/dL, 9 IU/dL, 10 IU/dL, 11 IU/dL, 12 IU/dL, 13 IU/dL, 14 IU/dL, 15 IU/dL, 16 IU/dL, 17 IU/dL, 18 IU/dL, 19 IU/dL, or 20 IU/dL in a subject by administering a chimeric protein of the disclosure. As used herein, a "trough level" in a hemophilia A patient is the measurement of the lowest concentration reached by a factor therapy, e.g., a FVIII therapy, before the next dose is administered. In some embodiments, treating or treatment means maintaining a FVIII trough level of at least about 1 IU/dL between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII trough level of at least about 3 IU/dL between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII trough level of at least about 5 IU/dL between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII trough level between about 1 and about 20 IU/dL, about 2 and about 20 IU/dL, about 3 and about 20 IU/dL, about 4 and about 20 IU/dL, about 5 and about 20 IU/dL, about 6 and about 20 IU/dL, about 7 and about 20 IU/dL, about 8 and about 20 IU/dL, about 9 and about 20 IU/dL, or about 10 and about 20 IU/dL during the dosing interval.

[0156] In some embodiments, treatment or treating of a disease or condition comprises maintaining FVIII activity in a subject at a level comparable to at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, or 20% of the FVIII

activity in a non-hemophiliac subject between the doses. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 1% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 2% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 3% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 4% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 5% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 6% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 7% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 8% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 9% between the dosing interval. In some embodiments, treating or treatment means maintaining a FVIII activity level of at least about 10% between the dosing interval. In some embodiments, the minimum trough level required for treatment can be measured by one or more known methods (for example, the activated partial thromboplastin time (aPTT) assays or chromogenic assays, which are well known in the art) and can be adjusted (increased or decreased) for each person. Non-limiting examples of assays for measuring trough level are disclosed in U.S. Application Publication No. 20190375822, which is hereby incorporated by reference in its entirety.

II. Chimeric Proteins

[0157] In an aspect, the present disclosure is directed to pharmaceutical compositions comprising a chimeric protein or protein which comprises a first polypeptide chain which comprises a Factor VIII ("FVIII") protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof. In some embodiments, the chimeric protein comprises (i) a FVIII protein comprising a FVIII polypeptide, an ELNN Polypeptide inserted within the B domain of the FVIII polypeptide, and a first Fc region; and (ii) a VWF protein comprising a VWF fragment, a second ELNN Polypeptide sequence, an a2 linker, and a second Fc region. In some embodiments, the chimeric protein disclosed herein is a FVIII-ELNN-Fc/D'D3-ELNN-Fc heterodimer. Non-limiting examples of chimeric proteins that may be used in various embodiments are described in U.S. Patent No. 10,138,291 and US Patent No. 11,192,936 B2, the entire contents of each of which are incorporated herein by reference.

[0158] In some embodiments, the chimeric protein is efanesoctocog alfa. Efanesoctocog alfa, also known as "BIVV001", "efanesoctocogum alfa" and "rFVIII-Fc-VWF-XTEN", is described in

Chhabra et al. *Blood* 135(17): 1484-1496 (2020), the entire contents of which are hereby incorporated by reference in its entirety. A schematic representation of efanesoctocog alfa, as an exemplary FVIII-ELNN-Fc/D'D3-ELNN-Fc heterodimer, is presented in FIG. 1.

[0159] Efanesoctocog alfa is a large protein (over 300 kDa) comprising multiple parts on each of two polypeptide chains that are associated by covalent bonds and non-covalent interactions. The protein has a tendency to aggregate under certain conditions, which can reduce the stability of formulations unless an excipient such as L-arginine is selected and present in an amount that is sufficient to reduce the aggregation. For example, the aggregation may be reduced by adding high levels of L-arginine (e.g., about 250 mM).

[0160] Additional information regarding efanesoctocog alfa can be found in the International Nonproprietary Names for Pharmaceutical Substances (INN) WHO Drug Information, 2019, Vol. 33, No. 4, p.828-30. In some embodiments, the chimeric protein is a FVIII-ELNN-Fc/D'D3-ELNN-Fc heterodimer comprising (i) a first polypeptide comprising the amino acid sequence of SEQ ID NO: 1 and (ii) a second polypeptide comprising the amino acid sequence of SEQ ID NO: 2. In some embodiments, the chimeric protein comprises (i) a first polypeptide and (ii) a second polypeptide that are covalently linked via one or more disulfide bonds (e.g., two disulfide bonds). In some embodiments, the chimeric protein comprises a FVIII protein encoded by the nucleic acid sequence of SEQ ID NO: 4. In some embodiments, the chimeric protein comprises a VWF protein encoded by the nucleic acid sequence of SEQ ID NO: 6. In some embodiments, the efanesoctocog alfa has an activity of at least 1600 IU/mg. In some embodiments, the efanesoctocog alfa has an activity of at least 1700 IU/mg. In some embodiments, the efanesoctocog alfa has an activity of at least 1800 IU/mg. In some embodiments, the efanesoctocog alfa has an activity of at least 1900 IU/mg. In some embodiments, the efanesoctocog alfa has an activity of 1600 IU/mg to 2000 IU/mg.

[0161] In some embodiments, the chimeric protein comprises a FVIII protein comprising the amino acid sequence of SEQ ID NO: 1. In some embodiments, the chimeric protein comprises a FVIII protein comprising one or more disulfide bridges at one or more of the following locations: residues 153-179, 248-329, 528-554, 630-711, 1220-1246, 1287-1291, 1409-1557, 1562-1714, 1761-1821, and/or 1867-1925 of SEQ ID NO: 1. In some embodiments, the chimeric protein comprises a FVIII protein comprising one or more disulfide bridges at each of the following locations: residues 153-179, 248-329, 528-554, 630-711, 1220-1246, 1287-1291, 1409-1557, 1562-1714, 1761-1821, and 1867-1925 of SEQ ID NO: 1. In some embodiments, the chimeric protein comprises a FVIII protein comprising one or more Cys-SH residues at residues 310, 692, and/or 1388 of SEQ ID NO: 1. In some embodiments, the chimeric protein comprises a FVIII protein comprising a Cys-SH residues at each of residues 310, 692, and/or 1388 of SEQ ID NO: 1.

[0162] In some embodiments, the chimeric protein comprises a FVIII protein comprises one or more N-glycosylation sites at residues N41, N239, N1198, N1506, and/or N1797 of SEQ ID NO: 1. In some embodiments, the chimeric protein comprises a FVIII protein comprises one or more O-glycosylation sites at residues 746-1036 of SEQ ID NO: 1 and/or the Ser and Thr residues in the linker peptides. In some embodiments, the chimeric protein comprises a FVIII protein comprises one or more Tyr-sulfation sites at residues 346, 718, 719, 723, 729, 1052, and/or 1068 of SEQ ID NO: 1.

[0163] In some embodiments, the chimeric protein comprises a VWF protein comprising the amino acid sequence of SEQ ID NO: 2. In some embodiments, the chimeric protein comprises a VWF protein comprising one or more disulfide bridges at one or more of the following locations: residues 4-45, 13-41, 25-36, 29-64, 47-58, 66-88, 83-100, 86-95, 104-233, 126-268, 135-230, 151-158, 283-326, 297-321, 308-348, 328-334, 338-363, 367-410, 386-406, 390-402, 394-433, 414-427, 436-464, 459-474, 462-471, 698-758, and/or 804-862 of SEQ ID NO: 2. In some embodiments, the chimeric protein comprises a VWF protein comprising one or more disulfide bridges at each of the following locations: residues 4-45, 13-41, 25-36, 29-64, 47-58, 66-88, 83-100, 86-95, 104-233, 126-268, 135-230, 151-158, 283-326, 297-321, 308-348, 328-334, 338-363, 367-410, 386-406, 390-402, 394-433, 414-427, 436-464, 459-474, 462-471, 698-758, and/or 804-862 of SEQ ID NO: 2.

[0164] In some embodiments, the chimeric protein comprises a VWF protein comprising one or more N-glycosylation sites at residues N94, N384, N734 of SEQ ID NO: 2. In some embodiments, the chimeric protein comprises a VWF protein comprising one or more O-glycosylation sites at residues 478-625 of SEQ ID NO: 2 and/or the Ser and Thr residues in the linker peptides. In some embodiments, the chimeric protein comprises a VWF protein comprising one or more Tyr-sulfation sites at residues 632, 633, 637, and/or 643 of SEQ ID NO: 2. In some embodiments, the VWF protein comprises a VWF fragment comprising a D1, D2, D', and/or D3 domain of VWF. In one embodiment, the VWF fragment comprises a D1D2 region of VWF comprising the amino acid sequence of SEQ ID NO: 20. In some embodiments, the VWF protein further comprises a VWF signal peptide sequence. In one embodiment, the VWF signal peptide comprises the amino acid sequence of SEQ ID NO: 19. In one specific embodiment, the VWF protein comprises a VWF signal peptide comprising the amino acid sequence of SEQ ID NO: 19, a D1D2 region of VWF comprising the amino acid sequence of SEQ ID NO: 20, a D' domain of VWF comprising the amino acid sequence of SEQ ID NO: 21, a D3 domain of VWF comprising the amino acid sequence of SEQ ID NO: 22, an ELNN Polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 14 (AE144_5A), an α 2 linker comprising the amino acid sequence of SEQ ID NO: 15, and/or a Fc region comprising the amino acid sequence of SEQ ID NO: 23.

[0165] In some embodiments, the chimeric protein of the present disclosure comprises: (i) a FVIII protein comprising a FVIII polypeptide, a first ELNN Polypeptide sequence, and a first Fc region; and (ii) a VWF fragment comprising a D' domain of VWF and a D3 domain of VWF, a second ELNN Polypeptide sequence, an a2 linker of FVIII, and a second Fc region; wherein: the FVIII protein has a deletion of amino acids 746 to 1648 corresponding to mature FVIII; the first ELNN Polypeptide sequence is inserted within the FVIII polypeptide immediately downstream of amino acid 745 corresponding to mature FVIII; the first ELNN Polypeptide sequence comprises an amino acid sequence having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% sequence identity to the amino acid sequence of AE288 (SEQ ID NO: 9); the first Fc region is fused to the C-terminus of the FVIII polypeptide; the second ELNN Polypeptide sequence is fused to the C-terminus of the VWF fragment; the second ELNN Polypeptide sequence comprises an amino acid sequence having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% sequence identity to the amino acid sequence of AE144_5A (SEQ ID NO: 14); the a2 linker is fused to the C-terminus of the ELNN Polypeptide; the a2 linker comprises an amino acid sequence having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or about 100% sequence identity to the amino acid sequence of SEQ ID NO: 15; the second Fc region is fused to the C-terminus of the a2 linker; and the first Fc region is covalently linked to the second Fc region by a disulfide bond(e.g., two disulfide bonds).

[0166] In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising an amino acid sequence at least about 80%, 90%, 95%, or 100% identical to the amino acid sequence set forth in SEQ ID NO: 1; and a second polypeptide sequence comprising a VWF fragment comprising a D' domain of VWF and a D3 domain of VWF and an Fc region. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising FVIII polypeptide and an Fc region; and a second polypeptide sequence comprising an amino acid sequence at least about 80%, 90%, 95%, or 100% identical to the amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising an amino acid sequence at least about 80%, 90%, 95%, or 100% identical to the amino acid sequence set forth in SEQ ID NO: 1 and a second polypeptide sequence comprising an amino acid sequence at least about 80%, 90%, 95%, or

100% identical to the amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 7 and a second polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 1 and a second polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 2, wherein the first polypeptide sequence and the second polypeptide sequence are linked to each other by a disulfide bond. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 1 and a second polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 2, wherein the first polypeptide sequence and the second polypeptide sequence are linked to each other by two disulfide bonds. In some embodiments, the chimeric protein of the disclosure comprises two polypeptide sequences, a first polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 1 and a second polypeptide sequence comprising the amino acid sequence set forth in SEQ ID NO: 2, wherein the first polypeptide sequence comprises a first Fc portion, wherein the second polypeptide sequence comprises a second Fc portion, wherein the first Fc portion and the second Fc portion are linked to each other by two disulfide bonds in the hinge region.

[0167] In some embodiments, the chimeric protein of the disclosure comprises a FVIII protein comprising an amino acid sequence at least about 80%, 90%, 95%, or 100% identical to SEQ ID NO: 7, SEQ ID NO: 3, or SEQ ID NO: 1; and a VWF protein comprising an amino acid sequence at least about 80%, 90%, 95%, or 100% identical to SEQ ID NO: 2 or SEQ ID NO: 5.

[0168] In some embodiments, the chimeric protein of the disclosure comprises: (i) a FVIII protein comprising a first FVIII polypeptide fragment comprising the amino acid sequence of SEQ ID NO: 17; a first ELNN Polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 9 (AE288); a second FVIII polypeptide fragment comprising the amino acid sequence of SEQ ID NO: 18; and a first Fc region comprising the amino acid sequence of SEQ ID NO: 23; and (ii) a VWF protein comprising: a D' domain of VWF comprising the amino acid sequence of SEQ ID NO: 21; a D3 domain of VWF comprising the amino acid sequence of SEQ ID NO: 22; a second ELNN Polypeptide sequence comprising the amino acid sequence of SEQ ID NO: 14 (AE144_5A); an a2 linker comprising the amino acid sequence of SEQ ID NO: 15; and a second Fc region comprising the amino acid sequence of SEQ ID NO: 23, and wherein the first Fc region is covalently linked to the second Fc region by a disulfide bond (e.g., two disulfide bonds).

[0169] In some embodiments, the chimeric protein of the disclosure comprises a FVIII protein comprising a FVIII polypeptide, a first ELNN Polypeptide sequence, a first Fc region, and a VWF protein comprising a D' domain of VWF, a D3 domain of VWF, a second ELNN Polypeptide sequence, an a2 linker of FVIII and a second Fc region, wherein the FVIII polypeptide comprises the amino acid sequence of SEQ ID NO: 17, the first ELNN Polypeptide sequence comprises the amino acid sequence of AE288 (SEQ ID NO: 9) and is fused to the C-terminus of SEQ ID NO: 17, the FVIII polypeptide further comprises the amino acid sequence of SEQ ID NO: 18, the first Fc region comprises the amino acid sequence of SEQ ID NO: 23 and is fused to the C-terminus of SEQ ID NO: 18; the D' domain of VWF comprises the amino acid sequence of SEQ ID NO: 21; the D3 domain of VWF comprises the amino acid sequence of SEQ ID NO: 21, the second ELNN Polypeptide sequence comprises the amino acid sequence of AE144_5A (SEQ ID NO: 14) and is fused to the C-terminus of the D3 domain of VWF; the a2 linker comprises the amino acid sequence of SEQ ID NO: 15 and is fused to the C-terminus of the second ELNN Polypeptide sequence; the second Fc region comprises the amino acid sequence of SEQ ID NO: 23 and is fused to the C-terminus of the a2 linker; and wherein the first Fc region is covalently linked to the second Fc region by a disulfide bond.

[0170] In some embodiments, the chimeric protein of the disclosure comprises a FVIII protein comprising a FVIII signal peptide comprising the amino acid sequence of SEQ ID NO: 16. In some embodiments, the chimeric protein comprises a VWF protein comprising a VWF signal peptide comprising the amino acid sequence of SEQ ID NO: 19. In some embodiments, the chimeric protein comprises a VWF protein comprising a D1D2 domain of VWF comprising the amino acid sequence of SEQ ID NO: 20.

[0171] In some embodiments, the chimeric protein comprises a first polypeptide comprising the amino acid sequence of SEQ ID NO: 3 and a second polypeptide comprising the amino acid sequence of SEQ ID NO: 5. In some embodiments, the chimeric protein comprises a first polypeptide comprising the amino acid sequence of SEQ ID NO: 7 and a second polypeptide comprising the amino acid sequence of SEQ ID NO: 2. In some embodiments, the chimeric protein comprises a first polypeptide comprising the amino acid sequence of SEQ ID NO: 1 and a second polypeptide comprising the amino acid sequence of SEQ ID NO: 2.

[0172] In some embodiments, the chimeric protein comprises one or more disulfide bridges between the first polypeptide and the second polypeptide. In some embodiments, the chimeric protein comprises two disulfide bridges between the first polypeptide and the second polypeptide. In some embodiments, the chimeric protein comprises a first polypeptide comprising the amino acid sequence of SEQ ID NO: 1 and a second polypeptide comprising the amino acid sequence of SEQ ID NO: 2, wherein the chimeric protein comprises a disulfide bridge between residue 1726

of SEQ ID NO: 1 and residue 663 of SEQ ID NO: 2, and a disulfide bridge between residue 1729 of SEQ ID NO: 1 and residue 666 of SEQ ID NO: 2.

IV. Pharmaceutical Compositions

[0173] In an aspect, the present disclosure is directed to pharmaceutical compositions of a chimeric protein which are formulated to improve protein stability. In some embodiments, the disclosed pharmaceutical compositions demonstrate increased stability based on analysis by visual inspection, protein concentration, pH stability, formation of high molecular weight species (HMWS), and/or change in turbidity. Analysis of these properties of stability can be made using conventional techniques, including size exclusion chromatography (SEC), reversed-phase high-performance liquid chromatography (RP-HPLC), and many others.

[0174] The pharmaceutical compositions disclosed herein comprise a specified amount of the chimeric protein. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 0.8 to about 1.2 mg/mL. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 0.8 mg/mL. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 0.9 mg/mL. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 1.0 mg/mL. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 1.1 mg/mL. In some embodiments, the pharmaceutical composition has a chimeric protein concentration of about 1.2 mg/mL.

[0175] In some embodiments, the pharmaceutical composition comprises about 75 IU/mL to about 2,000 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 75 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 100 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 150 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 200 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 250 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 300 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 350 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 400 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 450 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 500 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 550 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 600 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 650

IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 700 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 750 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 800 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 850 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 900 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 950 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1000 IU/mL of the chimeric protein.

[0176] In some embodiments, the pharmaceutical composition comprises about 1100 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1150 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1200 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1250 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1300 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1350 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1400 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1450 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1500 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1550 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1600 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1650 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1700 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1750 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1800 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1850 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1900 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 1950 IU/mL of the chimeric protein. In some embodiments, the pharmaceutical composition comprises about 2000 IU/mL of the chimeric protein.

[0177] Pharmaceutical compositions containing the chimeric protein of the present disclosure also contain a suitable pharmaceutically acceptable carrier. For example, they can contain

excipients and/or auxiliaries that provide enhanced stability of the chimeric protein or facilitate processing of the active compounds into preparations designed for delivery to the site of action.

[0178] In an aspect, disclosed herein are pharmaceutical compositions comprising a specified amount of a chimeric protein along with excipients as disclosed. The pharmaceutical compositions disclosed herein comprise various concentrations of these excipients as disclosed, and the concentrations can be expressed in various ways. For example, the concentration of a given excipient can be expressed as a molar concentration (e.g., M or mM), as a weight/volume percent, (e.g., grams per 100 ml diluent), or as milligrams per milliliter (mg/ml). Pharmaceutical compositions provided herein can contain specified amounts of the various excipients at a level of precision ranging from approximate, e.g., concentrations expressed only to one significant figure (e.g., about 0.1% (w/v)), or with more precision, e.g., out to 2, 3, 4, 5, or 6 significant figures (e.g., about 3.88 mg/ml, with precision out to three significant figures). The necessary level of precision can vary depending on, e.g., the requirements of a given regulatory agency, or the manufacturing process.

[0179] Pharmaceutical compositions disclosed herein may include a stabilizing agent.

[0180] In some embodiments, the pharmaceutical compositions disclosed herein include specified amounts or concentrations of sucrose.

[0181] In some embodiments, the pharmaceutical composition comprises 4.5% (w/v) to 8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5% (w/v) to about 7.5% (w/v) sucrose.

[0182] In some embodiments, the pharmaceutical composition comprises about 4.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 4.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 4.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 4.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 4.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.2% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 5.9% (w/v) sucrose.

sucrose. In some embodiments, the pharmaceutical composition comprises about 6.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.2% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 6.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.2% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 7.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 8% (w/v) sucrose.

[0183] In some embodiments, the pharmaceutical composition comprises 168.3 mg sucrose. In some embodiments, the pharmaceutical composition comprises about 168.3 mg sucrose. In some embodiments, the amount of sucrose can vary up to 10% of a specific amount. In some embodiments, the specific amount of sucrose is 168.3 mg. In some embodiments, the amount of sucrose can vary up to 5% of a specific amount. In some embodiments, the specific amount of sucrose is 168.3 mg. In some embodiments, the amount of sucrose can vary up to 1% of a specific amount. In some embodiments, the specific amount of sucrose is 168.3 mg.

[0184] In some embodiments, the pharmaceutical composition comprises 1% (w/v) to 4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.5% (w/v) to about 2.5% (w/v) sucrose.

[0185] In some embodiments, the pharmaceutical composition comprises about 1.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.2% (w/v)

sucrose. In some embodiments, the pharmaceutical composition comprises about 1.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 1.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.2% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 2.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.0% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.1% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.2% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.3% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.4% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.5% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.6% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.7% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.8% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 3.9% (w/v) sucrose. In some embodiments, the pharmaceutical composition comprises about 4.0% (w/v) sucrose.

[0186] In some embodiments, the pharmaceutical composition comprises 67.34 mg sucrose. In some embodiments, the pharmaceutical composition comprises about 67.34 mg sucrose. In some embodiments, the amount of sucrose can vary up to 10% of a specific amount. In some embodiments, the specific amount of sucrose is 67.34 mg. In some embodiments, the amount of sucrose can vary up to 5% of a specific amount. In some embodiments, the specific amount of

sucrose is 67.34 mg. In some embodiments, the amount of sucrose can vary up to 1% of a specific amount. In some embodiments, the specific amount of sucrose is 67.34 mg.

[0187] Pharmaceutical compositions disclosed herein may include a buffer. In some embodiments, the pharmaceutical compositions disclosed herein include specified amounts or concentrations of histidine. In some embodiments, the histidine included in the pharmaceutical composition is L-histidine. In some embodiments, the pharmaceutical composition comprises about 5 mM to about 15 mM histidine.

[0188] In some embodiments, the pharmaceutical composition comprises about 5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 5.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 6 mM histidine. In some embodiments, the pharmaceutical composition comprises about 6.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 7 mM histidine. In some embodiments, the pharmaceutical composition comprises about 7.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 8 mM histidine. In some embodiments, the pharmaceutical composition comprises about 8.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 9 mM histidine. In some embodiments, the pharmaceutical composition comprises about 9.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 10 mM histidine. In some embodiments, the pharmaceutical composition comprises about 10.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 11 mM histidine. In some embodiments, the pharmaceutical composition comprises about 11.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 12 mM histidine. In some embodiments, the pharmaceutical composition comprises about 12.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 13 mM histidine. In some embodiments, the pharmaceutical composition comprises about 13.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 14 mM histidine. In some embodiments, the pharmaceutical composition comprises about 14.5 mM histidine. In some embodiments, the pharmaceutical composition comprises about 15 mM histidine. In some embodiments, the histidine is L-histidine.

[0189] In some embodiments, the pharmaceutical compositions disclosed herein include specified amounts or concentrations of arginine. In some embodiments, the pharmaceutical composition comprises arginine hydrochloride (HCl). In some embodiments, the arginine is L-arginine. In some embodiments, the composition comprises L-arginine-HCl.

[0190] In some embodiments, the pharmaceutical composition comprises at least 150 mM arginine. In some embodiments, the pharmaceutical composition comprises at least 200 mM

arginine. In some embodiments, the pharmaceutical composition comprises at least 250 mM arginine. In some embodiments, the pharmaceutical composition comprises about 150 mM to about 300 mM arginine.

[0191] In some embodiments, the pharmaceutical composition comprises about 200 mM to about 300 mM arginine.

[0192] In some embodiments, the pharmaceutical composition comprises about 150 mM arginine. In some embodiments, the pharmaceutical composition comprises about 160 mM arginine. In some embodiments, the pharmaceutical composition comprises about 170 mM arginine. In some embodiments, the pharmaceutical composition comprises about 180 mM arginine. In some embodiments, the pharmaceutical composition comprises about 190 mM arginine. In some embodiments, the pharmaceutical composition comprises about 200 mM arginine. In some embodiments, the pharmaceutical composition comprises about 210 mM arginine. In some embodiments, the pharmaceutical composition comprises about 220 mM arginine. In some embodiments, the pharmaceutical composition comprises about 230 mM arginine. In some embodiments, the pharmaceutical composition comprises about 240 mM arginine. In some embodiments, the pharmaceutical composition comprises about 250 mM arginine. In some embodiments, the pharmaceutical composition comprises about 260 mM arginine. In some embodiments, the pharmaceutical composition comprises about 270 mM arginine. In some embodiments, the pharmaceutical composition comprises about 280 mM arginine. In some embodiments, the pharmaceutical composition comprises about 290 mM arginine. In some embodiments, the pharmaceutical composition comprises about 300 mM arginine. In some embodiments, the arginine is L-arginine. In some embodiments, the composition comprises L-arginine-HCl.

[0193] Pharmaceutical compositions disclosed herein may include a bulking agent. In some embodiments, the pharmaceutical compositions disclosed herein include specified amounts or concentrations of calcium chloride (CaCl_2). In some embodiments, the composition comprises $\text{CaCl}_2 \bullet 2\text{H}_2\text{O}$, CaCl_2 (anhydrous), $\text{CaCl}_2 \bullet 4\text{H}_2\text{O}$, or $\text{CaCl}_2 \bullet 6\text{H}_2\text{O}$. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition comprises about 2.5 mM to about 10 mM calcium chloride. In some embodiments, the composition comprises calcium chloride dihydrate.

[0194] In some embodiments, the pharmaceutical composition comprises about 2.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 3 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 3.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 4 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about

4.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 5.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 6 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 6.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 7 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 7.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 8 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 8.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 9 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 9.5 mM calcium chloride. In some embodiments, the pharmaceutical composition comprises about 10 mM calcium chloride. In some embodiments, the composition comprises calcium chloride dihydrate.

[0195] In some embodiments, the pharmaceutical compositions disclosed herein do not include bulking agents other than calcium chloride. In some embodiments, calcium chloride is the sole bulking agent. In some embodiments, the pharmaceutical composition comprises less than 8.8 mg/mL sodium chloride (NaCl). In some embodiments, the pharmaceutical composition is substantially free of sodium chloride. In some embodiments, the pharmaceutical composition is free of sodium chloride.

[0196] In some embodiments, the pharmaceutical compositions disclosed herein include specified amounts or concentrations of polysorbate 20 (PS20) or polysorbate 80 (PS80). In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) to about 0.1% (w/v) PS80 or PS20. In some embodiments, the pharmaceutical composition comprises at least about 0.03% PS20 or PS80. In some embodiments, the pharmaceutical composition comprises about 0.05% PS20 or PS80. In some embodiments, the pharmaceutical composition comprises PS20. In some embodiments, the pharmaceutical composition comprises PS80.

[0197] In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.01% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.02% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.03% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.04% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.05% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.06% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.07% (w/v) polysorbate 20. In some

embodiments, the pharmaceutical composition comprises about 0.08% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.09% (w/v) polysorbate 20. In some embodiments, the pharmaceutical composition comprises about 0.1% (w/v) polysorbate 20.

[0198] In some embodiments, the pharmaceutical composition comprises about 0.008% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.01% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.02% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.03% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.04% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.05% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.06% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.07% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.08% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.09% (w/v) polysorbate 80. In some embodiments, the pharmaceutical composition comprises about 0.1% (w/v) polysorbate 80.

[0199] In some embodiments, the pharmaceutical composition is a pre-lyophilization solution. In some embodiments, pre-lyophilization solution does not comprise NaCl. In some embodiments, the pre-lyophilization solution does not comprise NaOH. In some embodiments, the pre-lyophilization solution does not comprise sodium ions.

[0200] In some embodiments, the pharmaceutical composition comprises

- (a) about 1% (w/v) to about 4% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 150 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0201] In some embodiments, the pharmaceutical composition comprises

- (a) about 1% (w/v) to about 4% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 200 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the

composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0202] In some embodiments, the pharmaceutical composition comprises

- (a) 2% (w/v) to 3% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM histidine;
- (c) 225 mM to about 300 mM arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises 225 mM to about 300 mM L-arginine-HCl. In some embodiments, the composition comprises 5 mM to 6 mM calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0203] In some embodiments, the pharmaceutical composition comprises

- (a) 2% (w/v) to 3% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0204] In some embodiments, the pharmaceutical composition comprises

- (a) 2% (w/v) to 3% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine-HCl;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0205] In some embodiments, the pharmaceutical composition comprises

- (a) 2% (w/v) to 3% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine-HCl;
- (d) 5 mM to 6 mM calcium chloride dihydrate; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0206] In some embodiments, the pharmaceutical composition comprises:

- (a) about 2.25 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;

- (c) about 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and
- (e) 0.056 % (w/v) polysorbate 80.

[0207] In some embodiments, the pharmaceutical composition comprises:

- (a) about 2.25 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine;
- (d) about 5.61 mM calcium chloride; and
- (e) about 0.056 % (w/v) polysorbate 80.

[0208] In some embodiments, the pharmaceutical composition comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and
- (e) 0.056 % (w/v) polysorbate 80.

[0209] In some embodiments, the pharmaceutical composition comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0210] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 2.25 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and
- (e) 0.056 % (w/v) polysorbate 80.

[0211] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 2.25 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine;
- (d) about 5.61 mM calcium chloride; and
- (e) about 0.056 % (w/v) polysorbate 80.

[0212] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate (5.61 mM calcium chloride); and
- (e) 0.056 % (w/v) polysorbate 80.

[0213] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0214] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2% (w/v) to 3% (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0215] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0216] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0217] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0218] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 2.25 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) (w/v) polysorbate 80.

[0219] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise NaCl. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise NaOH. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise sodium ions.

[0220] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0221] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) to about 3% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and

(e) about 0.05% polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0222] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0223] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 200 mM to about 300 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0224] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0225] In some embodiments, the pharmaceutical composition comprises

- (a) about 2% (w/v) sucrose;
- (b) about 10 mM histidine;

(c) about 250 mM arginine;

(d) about 5 mM calcium chloride; and

[0226] (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0227] In some embodiments, the pharmaceutical composition comprises:

(a) about 20 mg/ml sucrose;

(b) about 1.552 mg/ml L-histidine;

(c) about 52.665 mg/ml L-arginine-HCl;

(d) about 0.735 mg/ml calcium chloride; and

(e) about 0.5 mg/ml polysorbate 80.

[0228] In some embodiments, the pharmaceutical composition comprises:

(a) about 20 mg/ml sucrose;

(b) about 1.552 mg/ml L-histidine;

(c) about 52.665 mg/ml L-arginine-HCl;

(d) about 0.735 mg/ml calcium chloride dihydrate; and

(e) about 0.5 mg/ml polysorbate 80.

[0229] In some embodiments, the pharmaceutical composition comprises:

(a) about 20 mg/ml sucrose;

(b) about 1.552 mg/ml L-histidine;

(c) about 52.665 mg/ml L-arginine-HCl;

(d) about 0.555 mg/ml calcium chloride; and

(e) about 0.5 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0230] In some embodiments, the pharmaceutical composition comprises:

(a) about 20 mg/ml sucrose;

(b) about 1.552 mg/ml L-histidine;

(c) about 43.550 mg/ml L-arginine;

(d) about 0.735 mg/ml calcium chloride dihydrate; and

(e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0231] In some embodiments, the pharmaceutical composition comprises:

- (a) about 20 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 43.550 mg/ml L-arginine;
- (d) about 0.735 mg/ml calcium chloride; and
- (e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0232] In some embodiments, the pharmaceutical composition comprises:

- (a) about 20 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 43.550 mg/ml L-arginine;
- (d) about 0.555 mg/ml calcium chloride; and
- (e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0233] In some embodiments, the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

[0234] In some embodiments, the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0235] In some embodiments, the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

[0236] In some embodiments, the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0237] In some embodiments, the pharmaceutical composition comprises

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 150 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

[0238] In some embodiments, the pharmaceutical composition comprises

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 200 mM to about 300 mM arginine;
- (d) about 2.5 mM to about 10 mM calcium chloride; and
- (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0239] In some embodiments, the pharmaceutical composition comprises

- (a) 5% (w/v) to 6% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM histidine;
- (c) 225 mM to about 300 mM arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises 225 mM to about 300 mM L-arginine-HCl. In some embodiments, the composition comprises 5 mM to 6 mM calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0240] In some embodiments, the pharmaceutical composition comprises

- (a) 5% (w/v) to 6% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0241] In some embodiments, the pharmaceutical composition comprises

- (a) 5% (w/v) to 6% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine-HCl;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0242] In some embodiments, the pharmaceutical composition comprises

- (a) 5% (w/v) to 6% (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine-HCl;
- (d) 5 mM to 6 mM calcium chloride dihydrate; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0243] In some embodiments, the pharmaceutical composition comprises:

- (a) about 5.61 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and
- (e) 0.056 % (w/v) polysorbate 80.

[0244] In some embodiments, the pharmaceutical composition comprises:

- (a) about 5.61 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine;
- (d) about 5.61 mM calcium chloride; and
- (e) about 0.056 % (w/v) polysorbate 80.

[0245] In some embodiments, the pharmaceutical composition comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and

(e) 0.056 % (w/v) polysorbate 80.

[0246] In some embodiments, the pharmaceutical composition comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0247] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 5.61 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate; and
- (e) 0.056 % (w/v) polysorbate 80.

[0248] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 5.61 % (w/v) sucrose;
- (b) about 11.2 mM L-histidine;
- (c) about 280.58 mM L-arginine;
- (d) about 5.61 mM calcium chloride; and
- (e) about 0.056 % (w/v) polysorbate 80.

[0249] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine-HCl;
- (d) about 5.61 mM calcium chloride dihydrate (5.61 mM calcium chloride); and
- (e) 0.056 % (w/v) polysorbate 80.

[0250] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0251] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5% (w/v) to 6% (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0252] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 7.5 mM to 12.5 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0253] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 225 mM to about 300 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0254] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5 mM to 6 mM calcium chloride; and
- (e) 0.056 % (w/v) polysorbate 80.

[0255] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 5.61 % (w/v) sucrose;
- (b) 11.2 mM L-histidine;
- (c) 280.58 mM L-arginine;
- (d) 5.61 mM calcium chloride; and
- (e) 0.01% (w/v) to about 0.075% (w/v) polysorbate 80.

[0256] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise NaCl. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise NaOH. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution does not comprise sodium ions.

[0257] In some embodiments, the pharmaceutical composition comprises

- (a) about 5% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0258] In some embodiments, the pharmaceutical composition comprises

- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 20 or polysorbate 80. In some embodiments, the composition comprises polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0259] In some embodiments, the pharmaceutical composition comprises

- (a) about 5% (w/v) sucrose;
- (b) about 5 mM to about 15 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

- [0260] In some embodiments, the pharmaceutical composition comprises
- (a) about 5% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 200 mM to about 300 mM arginine;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.
- [0261] In some embodiments, the pharmaceutical composition comprises
- (a) about 5% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 250 mM arginine;
 - (d) about 2.5 mM to about 10 mM calcium chloride; and
 - (e) about 0.05% polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.
- [0262] In some embodiments, the pharmaceutical composition comprises
- (a) about 5% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 250 mM arginine;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 80. In some embodiments, the histidine is L-histidine. In some embodiments, the arginine is L-arginine. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.
- [0263] In some embodiments, the pharmaceutical composition comprises:
- (a) about 50 mg/ml sucrose;
 - (b) about 1.552 mg/ml L-histidine;
 - (c) about 52.665 mg/ml L-arginine-HCl;
 - (d) about 0.735 mg/ml calcium chloride; and
 - (e) about 0.5 mg/ml polysorbate 80.
- [0264] In some embodiments, the pharmaceutical composition comprises:

- (a) about 50 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 52.665 mg/ml L-arginine-HCl;
- (d) about 0.735 mg/ml calcium chloride dihydrate; and
- (e) about 0.5 mg/ml polysorbate 80.

[0265] In some embodiments, the pharmaceutical composition comprises:

- (a) about 50 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 52.665 mg/ml L-arginine-HCl;
- (d) about 0.555 mg/ml calcium chloride; and
- (e) about 0.5 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0266] In some embodiments, the pharmaceutical composition comprises:

- (a) about 50 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 43.550 mg/ml L-arginine;
- (d) about 0.735 mg/ml calcium chloride dihydrate; and
- (e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0267] In some embodiments, the pharmaceutical composition comprises:

- (a) about 50 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 43.550 mg/ml L-arginine;
- (d) about 0.735 mg/ml calcium chloride; and
- (e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0268] In some embodiments, the pharmaceutical composition comprises:

- (a) about 50 mg/ml sucrose;
- (b) about 1.552 mg/ml L-histidine;
- (c) about 43.550 mg/ml L-arginine;
- (d) about 0.555 mg/ml calcium chloride; and

(e) about 0.5 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate. In some embodiments, the pharmaceutical composition is a pre-lyophilization solution.

[0269] In some embodiments, the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

[0270] In some embodiments, the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0271] In some embodiments, the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

[0272] In some embodiments, the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0273] In some embodiments, the pharmaceutical composition has a pH of about 6.5 to about 7.5. In some embodiments, the pharmaceutical composition has a pH of about 7.0. In some embodiments, the pharmaceutical composition has a pH of about 6.8.

[0274] In some embodiments, the pharmaceutical composition has a pH of about 6.5. In some embodiments, the pharmaceutical composition has a pH of about 6.6. In some embodiments, the pharmaceutical composition has a pH of about 6.7. In some embodiments, the pharmaceutical

composition has a pH of about 6.8. In some embodiments, the pharmaceutical composition has a pH of about 6.9. In some embodiments, the pharmaceutical composition has a pH of about 7.0. In some embodiments, the pharmaceutical composition has a pH of about 7.1. In some embodiments, the pharmaceutical composition has a pH of about 7.2. In some embodiments, the pharmaceutical composition has a pH of about 7.3. In some embodiments, the pharmaceutical composition has a pH of about 7.4. In some embodiments, the pharmaceutical composition has a pH of about 7.5.

[0275] In some embodiments, the pharmaceutical composition has a pH of 6.5. In some embodiments, the pharmaceutical composition has a pH of 6.6. In some embodiments, the pharmaceutical composition has a pH of 6.7. In some embodiments, the pharmaceutical composition has a pH of 6.8. In some embodiments, the pharmaceutical composition has a pH of 6.9. In some embodiments, the pharmaceutical composition has a pH of 7.0. In some embodiments, the pharmaceutical composition has a pH of 7.1. In some embodiments, the pharmaceutical composition has a pH of 7.2. In some embodiments, the pharmaceutical composition has a pH of 7.3. In some embodiments, the pharmaceutical composition has a pH of 7.4. In some embodiments, the pharmaceutical composition has a pH of 7.5.

[0276] In some embodiments, a volume of 3.367 mL of the pre-lyophilization solution is added to a container or vial. In some embodiments, the pre-lyophilized solution is subjected to lyophilization, resulting in a lyophilized pharmaceutical composition.

[0277] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 30 mg to about 135 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg histidine;
- (c) about 140 mg to about 200 mg arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80.

[0278] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 30 mg to about 135 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg L-histidine;
- (c) about 140 mg to about 200 mg L-arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0279] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 30 mg to 135 mg sucrose;

- (b) 2.5 mg to 7.5 mg histidine;
- (c) 140 mg to 200 mg arginine;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 20 or polysorbate 80.

[0280] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 30 mg to 135 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 140 mg to 200 mg L-arginine-HCl;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0281] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 30 mg to 135 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 140 mg to 200 mg L-arginine;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0282] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 67.34 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 177.3 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0283] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 67.34 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 146.6 mg L-arginine;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

[0284] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.2 mg L-histidine;
- (c) 177.3 mg L-arginine-HCl;
- (d) 2.5 mg calcium chloride; and

- (e) 1.7 mg polysorbate 80.
- [0285] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 67.34 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 146.6 mg L-arginine;
 - (d) 2.5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0286] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 67.34 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 2.47 mg calcium chloride dihydrate; and
- [0287] (e) 1.68 mg polysorbate 80. In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 67.34 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 1.87 mg calcium chloride; and
 - (e) 1.68 mg polysorbate 80.
- [0288] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 67.34 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 146.63 mg L-arginine;
 - (d) 2.47 mg calcium chloride dihydrate; and
 - (e) 1.68 mg polysorbate 80.
- [0289] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 67.34 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 146.63 mg L-arginine;
 - (d) 1.87 mg calcium chloride; and
 - (e) 1.68 mg polysorbate 80.
- [0290] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 30 mg to 135 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;

- (d) 2.5 mg calcium chloride dihydrate; and
 - (e) 1.7 mg polysorbate 80.
- [0291] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 30 mg to 135 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;
 - (d) 1.9 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0292] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 30 mg to 135 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 146.6 mg L-arginine;
 - (d) 2.5 mg calcium chloride dihydrate; and
 - (e) 1.7 mg polysorbate 80.
- [0293] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 30 mg to 135 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 146.6 mg L-arginine;
 - (d) 1.9 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0294] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 67.34 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;
 - (d) 2.5 mg calcium chloride dihydrate; and
 - (e) 1.7 mg polysorbate 80.
- [0295] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 67.34 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;
 - (d) 1.9 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0296] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 67.34 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;

- (c) 146.6 mg L-arginine;
- (d) 2.5 mg calcium chloride dihydrate; and
- (e) 1.7 mg polysorbate 80.

[0297] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 146.6 mg L-arginine;
- (d) 1.9 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0298] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 140 mg to 200 mg L-arginine;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80. In some embodiments, the composition comprises 140 mg to 200 mg L-arginine-HCl.

[0299] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0300] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 1.5 mg to 5 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0301] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1 mg to 5 mg polysorbate 80.

[0302] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 1.87 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80.

[0303] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1 mg to 5 mg polysorbate 80.

[0304] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 1.87 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80.

[0305] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 160 mg to about 200 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg histidine;
- (c) about 140 mg to about 200 mg arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80.

[0306] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) about 160 mg to about 200 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg L-histidine;
- (c) about 140 mg to about 200 mg L-arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0307] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 160 mg to 200 mg sucrose;
- (b) 2.5 mg to 7.5 mg histidine;
- (c) 140 mg to 200 mg arginine;

- (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1 mg to 5 mg polysorbate 20 or polysorbate 80.
- [0308] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 160 mg to 200 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;
 - (c) 140 mg to 200 mg L-arginine-HCl;
 - (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1 mg to 5 mg polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.
- [0309] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 160 mg to 200 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;
 - (c) 140 mg to 200 mg L-arginine;
 - (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1 mg to 5 mg polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.
- [0310] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) about 168.3 mg sucrose;
 - (b) about 5.2 mg L-histidine;
 - (c) about 177.3 mg L-arginine-HCl;
 - (d) about 2.5 mg calcium chloride; and
 - (e) about 1.7 mg polysorbate 20 or polysorbate 80.
- [0311] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) about 168.3 mg sucrose;
 - (b) about 5.2 mg L-histidine;
 - (c) about 146.6 mg L-arginine;
 - (d) about 2.5 mg calcium chloride; and
 - (e) about 1.7 mg polysorbate 20 or polysorbate 80.
- [0312] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 168.3 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;
 - (d) 2.5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0313] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.3 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 146.6 mg L-arginine;
 - (d) 2.5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0314] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 2.47 mg calcium chloride dihydrate; and
 - (e) 1.68 mg polysorbate 80.
- [0315] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 1.87 mg calcium chloride; and
 - (e) 1.68 mg polysorbate 80.
- [0316] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 146.63 mg L-arginine;
 - (d) 2.47 mg calcium chloride dihydrate; and
 - (e) 1.68 mg polysorbate 80.
- [0317] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 146.63 mg L-arginine;
 - (d) 1.87 mg calcium chloride; and
 - (e) 1.68 mg polysorbate 80.
- [0318] In some embodiments, the lyophilized pharmaceutical composition comprises:
- (a) 160 mg to 200 mg sucrose;
 - (b) 5.2 mg L-histidine;
 - (c) 177.3 mg L-arginine-HCl;
 - (d) 2.5 mg calcium chloride dihydrate; and
 - (e) 1.7 mg polysorbate 80.

[0319] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 160 mg to 200 mg sucrose;
- (b) 5.2 mg L-histidine;
- (c) 177.3 mg L-arginine-HCl;
- (d) 1.9 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0320] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 160 mg to 200 mg sucrose;
- (b) 5.2 mg L-histidine;
- (c) 146.6 mg L-arginine;
- (d) 2.5 mg calcium chloride dihydrate; and
- (e) 1.7 mg polysorbate 80.

[0321] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 160 mg to 200 mg sucrose;
- (b) 5.2 mg L-histidine;
- (c) 146.6 mg L-arginine;
- (d) 1.9 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0322] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.3 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 177.3 mg L-arginine-HCl;
- (d) 2.5 mg calcium chloride dihydrate; and
- (e) 1.7 mg polysorbate 80.

[0323] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.3 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 177.3 mg L-arginine-HCl;
- (d) 1.9 mg calcium chloride; and
- (e) 1.7 mg polysorbate 80.

[0324] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.3 mg sucrose;
- (b) 2.5 mg to 7.5 mg L-histidine;
- (c) 146.6 mg L-arginine;
- (d) 2.5 mg calcium chloride dihydrate; and

- (e) 1.7 mg polysorbate 80.
- [0325] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.3 mg sucrose;
 - (b) 2.5 mg to 7.5 mg L-histidine;
 - (c) 146.6 mg L-arginine;
 - (d) 1.9 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0326] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 140 mg to 200 mg L-arginine;
 - (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80. In some embodiments, the composition comprises 140 mg to 200 mg L-arginine-HCl.
- [0327] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0328] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 146.63 mg L-arginine;
 - (d) 1.5 mg to 5 mg calcium chloride; and
 - (e) 1.7 mg polysorbate 80.
- [0329] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;
 - (c) 177.32 mg L-arginine-HCl;
 - (d) 2.47 mg calcium chloride dihydrate; and
 - (e) 1 mg to 5 mg polysorbate 80.
- [0330] In some embodiments, the lyophilized pharmaceutical composition comprises:
 - (a) 168.35 mg sucrose;
 - (b) 5.23 mg L-histidine;

- (c) 177.32 mg L-arginine-HCl;
- (d) 1.87 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80.

[0331] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1 mg to 5 mg polysorbate 80.

[0332] In some embodiments, the lyophilized pharmaceutical composition comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 1.87 mg calcium chloride; and
- (e) 1 mg to 5 mg polysorbate 80.

[0333] In some embodiments, the lyophilized pharmaceutical composition has a moisture content of less than 2%. In some embodiments, the lyophilized pharmaceutical composition has a moisture content of less than 1.8%. In some embodiments, the lyophilized pharmaceutical composition has a moisture content of less than 1.6%.

[0334] In some embodiments, the lyophilized pharmaceutical composition is in a lyophilized cake. In some embodiments, the lyophilized cake is white. In some embodiments, the lyophilized cake is less than Y4 in the European Pharmacopoeia color scale. See Degree of Coloration of Liquids (Method 2.2.2), European Pharmacopoeia, 10th Ed. (2021).

[0335] In some embodiments, the lyophilized pharmaceutical composition and sterile water are combined to produce an injectable solution. In some embodiments, the lyophilized pharmaceutical composition is combined with about 2 mL to about 5 mL of sterile water. In some embodiments, the lyophilized pharmaceutical composition is combined with about 3 mL of sterile water. In some embodiments, the lyophilized pharmaceutical composition is combined with 3 mL of sterile water. In some embodiments, the sterile water is USP grade sterile water. In some embodiments, the sterile water is USP grade sterile water for injection. In some embodiments, the sterile water is pyrogen-free or nonpyrogenic. In some embodiments, the sterile water does not contain a bacteriostatic or antimicrobial agent. In some embodiments, the sterile water contains a bacteriostatic or antimicrobial agent. In some embodiments, the sterile water is sterilized using a filter. In some embodiments, the sterile water is sterilized using a 0.1 μ m filter. In some embodiments, the sterile water is distilled water. In some embodiments, the sterile water is sterile,

nonpyrogenic, distilled water, hypotonic, with an osmolarity of zero mOsmol/L, and does not contain a bacteriostatic or antimicrobial agent.

[0336] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine;
- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and

[0337] (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine;
- (d) 0.5 mg/mL to 0.8 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate

[0338] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0339] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0340] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0341] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0342] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 10 mg/mL to about 40 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0343] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 10 mg/mL to about 40 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0344] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 10 mg/mL to about 40 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 44.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80

[0345] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 10 mg/mL to about 40 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 44.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0346] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0347] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0348] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0349] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 48.88 mg/ml L-arginine;

- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0350] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 50 mg/mL to about 70 mg/mL L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0351] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 50 mg/mL to about 70 mg/mL L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0352] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 40 mg/mL to about 60 mg/mL L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0353] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 40 mg/mL to about 60 mg/mL L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0354] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.7 mg/mL to about 0.9 mg/mL calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0355] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.5 mg/mL to about 0.9 mg/mL calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0356] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.7 mg/mL to about 0.9 mg/mL calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl.

[0357] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.5 mg/mL to about 0.7 mg/mL calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0358] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and

(e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80.

[0359] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0360] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80.

[0361] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0362] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0363] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80.

[0364] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0365] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 22.45mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80.

[0366] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 67.34 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 177.32 mg L-arginine-HCl;
- (d) about 2.47 mg calcium chloride; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0367] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 67.34 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 146.63 mg L-arginine;
- (d) about 2.47 mg calcium chloride dihydrate; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0368] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 67.34 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 146.63 mg L-arginine;
- (d) about 1.87 mg calcium chloride; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0369] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0370] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 1.87 mg calcium chloride; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0371] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 67.34 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0372] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 67.34 mg sucrose;

- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 1.87 mg calcium chloride; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0373] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine;
- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and

[0374] (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine;
- (d) 0.5 mg/mL to 0.8 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate

[0375] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0376] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and

(e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0377] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0378] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0379] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 45 mg/mL to about 60 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0380] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 45 mg/mL to about 60 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0381] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 45 mg/mL to about 60 mg/ml sucrose;

- (b) about 1.74 mg/ml L-histidine;
- (c) about 44.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80

[0382] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 45 mg/mL to about 60 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 44.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0383] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0384] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0385] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0386] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.5 mg/mL to about 2.0 mg/mL L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0387] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 50 mg/mL to about 70 mg/mL L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0388] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 50 mg/mL to about 70 mg/mL L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0389] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 40 mg/mL to about 60 mg/mL L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0390] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 40 mg/mL to about 60 mg/mL L-arginine;

- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0391] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.7 mg/mL to about 0.9 mg/mL calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0392] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.5 mg/mL to about 0.9 mg/mL calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments, the composition comprises calcium chloride dihydrate.

[0393] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.7 mg/mL to about 0.9 mg/mL calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl.

[0394] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.5 mg/mL to about 0.7 mg/mL calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0395] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80.

[0396] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0397] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80.

[0398] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.4 mg/mL to about 0.7 mg/mL polysorbate 80. In some embodiments the composition comprises L-arginine-HCl. In some embodiments, the composition comprises calcium chloride dihydrate.

[0399] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;

- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0400] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 59.11 mg/ml L-arginine-HCl;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80.

[0401] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.82 mg/ml calcium chloride dihydrate; and
- (e) about 0.56 mg/ml polysorbate 80.

[0402] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 56.12 mg/ml sucrose;
- (b) about 1.74 mg/ml L-histidine;
- (c) about 48.88 mg/ml L-arginine;
- (d) about 0.62 mg/ml calcium chloride; and
- (e) about 0.56 mg/ml polysorbate 80.

[0403] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 168.4 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 177.32 mg L-arginine-HCl;
- (d) about 2.47 mg calcium chloride; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0404] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 168.4 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 146.63 mg L-arginine;
- (d) about 2.47 mg calcium chloride dihydrate; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0405] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) about 168.4 mg sucrose;
- (b) about 5.23 mg L-histidine;
- (c) about 146.63 mg L-arginine;
- (d) about 1.87 mg calcium chloride; and
- (e) about 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0406] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0407] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 177.32 mg L-arginine-HCl;
- (d) 1.87 mg calcium chloride; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0408] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;

- (d) 2.47 mg calcium chloride dihydrate; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0409] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 168.35 mg sucrose;
- (b) 5.23 mg L-histidine;
- (c) 146.63 mg L-arginine;
- (d) 1.87 mg calcium chloride; and
- (e) 1.68 mg polysorbate 80

in 3 mL of sterile water.

[0410] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the lyophilized pharmaceutical composition is reconstituted within 7 to 12 seconds.

[0411] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 525 to about 725 mOsm/kg. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 600 to about 650 mOsm/kg.

[0412] In some embodiments, the pharmaceutical composition has an osmolality of about 525 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 550 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 575 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 600 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 625 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 650 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 675 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 700 mOsm/kg. In some embodiments, the pharmaceutical composition has an osmolality of about 725 mOsm/kg.

[0413] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.5 to about 7.5. In some embodiments, the pharmaceutical composition has a pH of about 7.0. In some embodiments, the pharmaceutical composition has a pH of about 6.8. In some embodiments, the pharmaceutical composition has a pH of 6.8.

[0414] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.5. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.6. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.7. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.8. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.9. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.0. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.1. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.2. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.3. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.4. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.5.

[0415] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 0.8 to about 1.2 mg/mL.

[0416] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 0.8 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 0.9 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 1.0 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 1.1 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 1.2 mg/mL.

[0417] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is 0.8 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are

combined, then the protein concentration of the resulting solution is 0.9 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is 1.0 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is 1.1 mg/mL. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is 1.2 mg/mL.

[0418] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, then the turbidity of the resulting solution is less than about 7 Nephelometric Turbidity Units.

[0419] In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 1% of the chimeric protein is aggregated. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 2% of the chimeric protein is aggregated. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 2.5% of the chimeric protein is aggregated. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 3% of the chimeric protein is aggregated. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 3.5% of the chimeric protein is aggregated. In some embodiments, when the lyophilized pharmaceutical composition and the sterile water are combined, less than 4% of the chimeric protein is aggregated.

V. Pharmaceutical Kits

[0420] In some embodiments, the pharmaceutical composition is provided with a second container comprising sterile water. Disclosed herein are pharmaceutical kits which comprise a first container containing the pharmaceutical composition and a second container containing sterile water.

[0421] Disclosed herein is a pharmaceutical kit comprising:

- (i) a first container comprising a lyophilized pharmaceutical composition comprising
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) sucrose;
 - (c) histidine;
 - (d) arginine;

- (e) calcium chloride; and
 - (f) a polysorbate (such as polysorbate 20 or polysorbate 80), and
- (ii) a second container comprising sterile water.

[0422] Also disclosed herein is a pharmaceutical kit comprising:

- (i) a first container comprising a lyophilized pharmaceutical composition comprising
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) about 30 mg to about 135 mg sucrose;
 - (c) about 2.5 mg to about 7.5 mg histidine;
 - (d) about 140 mg to about 200 mg arginine;
 - (e) about 1.5 mg to about 5 mg calcium chloride; and
 - (f) about 1 mg to about 5 mg of a polysorbate (such as polysorbate 20 or polysorbate 80), and
- (ii) a second container comprising sterile water.

[0423] Also disclosed herein is a pharmaceutical kit comprising:

- (i) a first container comprising a lyophilized pharmaceutical composition comprising
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) about 160 mg to about 200 mg sucrose;
 - (c) about 2.5 mg to about 7.5 mg histidine;
 - (d) about 140 mg to about 200 mg arginine;
 - (e) about 1.5 mg to about 5 mg calcium chloride; and
 - (f) about 1 mg to about 5 mg of a polysorbate (such as polysorbate 20 or polysorbate 80), and
- (ii) a second container comprising sterile water.

[0424] In some embodiments, the first container comprises 100 IU to 10,000 IU of the chimeric protein.

[0425] In some embodiments, the first container comprises 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein. In some embodiments, the first container comprises 250 IU of the chimeric protein. In some embodiments, the first container comprises 500 IU of the chimeric protein. In some embodiments, the first container comprises 1000 IU of the chimeric

protein. In some embodiments, the first container comprises 2000 IU of the chimeric protein. In some embodiments, the first container comprises 3000 IU of the chimeric protein. In some embodiments, the first container comprises 4,000 IU of the chimeric protein.

[0426] In some embodiments, the second container comprises about 2 mL to about 5 mL of sterile water. In some embodiments, the second container comprises 2 mL to 5 mL of sterile water. In some embodiments, the second container comprises about 3 mL of sterile water. In some embodiments, the second container comprises 3 mL of sterile water.

[0427] In some embodiments, the second container comprises about 2 mL of sterile water. In some embodiments, the second container comprises about 2.1 mL of sterile water. In some embodiments, the second container comprises about 2.2 mL of sterile water. In some embodiments, the second container comprises about 2.3 mL of sterile water. In some embodiments, the second container comprises about 2.4 mL of sterile water. In some embodiments, the second container comprises about 2.5 mL of sterile water. In some embodiments, the second container comprises about 2.6 mL of sterile water. In some embodiments, the second container comprises about 2.7 mL of sterile water. In some embodiments, the second container comprises about 2.8 mL of sterile water. In some embodiments, the second container comprises about 2.9 mL of sterile water. In some embodiments, the second container comprises about 3 mL of sterile water. In some embodiments, the second container comprises about 3.1 mL of sterile water. In some embodiments, the second container comprises about 3.2 mL of sterile water. In some embodiments, the second container comprises about 3.3 mL of sterile water. In some embodiments, the second container comprises about 3.4 mL of sterile water. In some embodiments, the second container comprises about 3.5 mL of sterile water. In some embodiments, the second container comprises about 3.6 mL of sterile water. In some embodiments, the second container comprises about 3.7 mL of sterile water. In some embodiments, the second container comprises about 3.8 mL of sterile water. In some embodiments, the second container comprises about 3.9 mL of sterile water. In some embodiments, the second container comprises about 4 mL of sterile water. In some embodiments, the second container comprises about 4.1 mL of sterile water. In some embodiments, the second container comprises about 4.2 mL of sterile water. In some embodiments, the second container comprises about 4.3 mL of sterile water. In some embodiments, the second container comprises about 4.4 mL of sterile water. In some embodiments, the second container comprises about 4.5 mL of sterile water. In some embodiments, the second container comprises about 4.6 mL of sterile water. In some embodiments, the second container comprises about 4.7 mL of sterile water. In some embodiments, the second container comprises about 4.8 mL of sterile water. In some

embodiments, the second container comprises about 4.9 mL of sterile water. In some embodiments, the second container comprises about 5 mL of sterile water.

[0428] In some embodiments, the pharmaceutical kit further comprises instructions for combining the lyophilized pharmaceutical composition and the sterile water.

[0429] In some embodiments, the first container is a glass vial comprising a rubber stopper.

[0430] In some embodiments, the second container is a syringe body. In some embodiments, the syringe body is associated with a plunger. In some embodiments, the pharmaceutical kit further comprises an adaptor to connect the glass vial to the syringe body. In some embodiments, the pharmaceutical kit further comprises infusion tubing associated with a needle to be connected to the syringe body, suitable for intravenous infusion. In some embodiments, the second container is a pre-filled syringe.

VI. Methods and Uses of the Pharmaceutical Composition

[0431] Also disclosed herein is a use of the pharmaceutical composition or a method for treating hemophilia A in a subject in need thereof, comprising administering to the subject an effective amount of the pharmaceutical composition of the disclosure. In some embodiments, the treatment of hemophilia A comprises preventing a bleeding episode in a human subject in need thereof. In some embodiments, the treatment of hemophilia A comprises treating a bleeding episode in a human subject in need thereof. In some embodiments, the treatment of hemophilia A comprises controlling the incidence or frequency of a bleeding episode in a human subject in need thereof. In some embodiments, the treatment of hemophilia A comprises decreasing the incidence or frequency of a bleeding episode in a human subject in need thereof.

[0432] In some embodiments, the composition is used to treat a bleeding disease or condition in a subject in need thereof. The bleeding disease or condition is selected from the group consisting of a bleeding coagulation disorder, hemarthrosis, muscle bleed, oral bleed, hemorrhage, hemorrhage into muscles, oral hemorrhage, trauma, trauma capitis, gastrointestinal bleeding, intracranial hemorrhage, intra-abdominal hemorrhage, intrathoracic hemorrhage, bone fracture, central nervous system bleeding, bleeding in the retropharyngeal space, bleeding in the retroperitoneal space, bleeding in the iliopsoas sheath and any combinations thereof. In still other embodiments, the subject is scheduled to undergo a surgery. In some embodiments, the treatment is prophylactic or on-demand.

[0433] In some embodiments, the use or method comprises combining the lyophilized pharmaceutical composition and the sterile water of a kit of the disclosure, and administering to the subject an effective amount of the resulting combination. In some embodiments, the subject combines the lyophilized pharmaceutical composition and the sterile water of the kit. In some embodiments, the combination is self-administered by the subject.

[0434] In some embodiments, the multiple doses comprise at least two doses, at least three doses, at least four doses, at least five doses, at least six doses, at least seven doses, at least eight doses, at least nine doses, at least ten doses, at least eleven doses, at least twelve doses, at least thirteen doses, at least fourteen doses, at least fifteen doses, at least sixteen doses, at least seventeen doses, at least eighteen doses, at least nineteen doses, at least twenty doses, or more. In some embodiments, the multiple doses are administered for at least about 1 month, at least about 2 months, at least about 3 months, at least about 4 months, at least about 5 months, at least about 6 months, at least about 12 months, at least about 18 months, at least about 2 years, at least about 3 years, at least about 4 years, at least about 5 years, at least about 10 years, at least about 15 years, at least about 20 years, or for at least about 25 years.

[0435] A chimeric protein described herein can be administered by any means known in the art. In some embodiments, the chimeric protein is administered by a route selected from the group consisting of intravenous injection, intravenous infusion, subcutaneous administration, intramuscular administration, oral administration, nasal administration, and pulmonary administration. In some embodiments, the chimeric protein is administered intravenously. In some embodiments, the chimeric protein is administered subcutaneously.

[0436] Having now described the present disclosure in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the disclosure. All patents, publications, and articles referred to herein are expressly and specifically incorporated herein by reference.

EXAMPLES

Example 1: Evaluation of buffer selection and target pH on stability of chimeric protein

[0437] The experiments of this example showed favorable pH and buffer species from various buffers like acetate, succinate, histidine, potassium phosphate, and Tris within a wide pH range from 4.0 – 8.0. Eleven buffers were studied in combination with efanesoctocog alfa at a target concentration of 1 mg/mL. The buffers studied are shown in Table 1. No additional excipients were included in the pharmaceutical compositions in this study. Each pharmaceutical composition comprising efanesoctocog alfa and the specific buffer was evaluated by measuring actual protein concentration, pH, visual inspection, high molecular weight species (HMWS), and turbidity. Studied samples were stored for up to 4 weeks at 5 °C, 25 °C, and 40 °C.

Table 1:

Buffers Studied	
Abbreviation	Full Name
A4.0	10 mM sodium acetate pH 4.0

A5.0	10 mM sodium acetate pH 5.0
S5.5	10 mM sodium succinate pH 5.5
H5.5	10 mM histidine-HCl pH 5.5
H6.0	10 mM histidine-HCl pH 6.0
H6.5	10 mM histidine-HCl pH 6.5
H7.0	10 mM histidine-HCl pH 7.0
H7.5	10 mM histidine-HCl pH 7.5
KP6.5	10 mM potassium phosphate pH 6.5
KP7.0	10 mM potassium phosphate pH 7.0
T8.0	10 mM Tris pH 8.0

[0438] Results of initial visual appearance, measured protein concentration, and pH of preparations of 1 mg/mL efanesoctocog alfa in the respective buffers are shown in Table 2. For H7.0, H7.5, and T8.0, the measured pH was 6.88, 7.05, and 7.84 respectively, which was lower than the target (expected) pH.

Table 2:

Buffer	Visual Appearance	Measured Concentration (mg/mL)	Measured pH
A4.0	Precipitated	Not measured	Not measured
A5.0	Cloudy, White, No visible particles	Not measured	Not measured
S5.5	Clear, White, No visible particles	1.06	5.48
H5.5	Clear, White, No visible particles	1.06	5.46
H6.0	Clear, White, No visible particles	1.04	5.99
H6.5	Clear, White, No visible particles	1.05	6.40
H7.0	Clear, White, No visible particles	1.04	6.88
H7.5	Clear, White, No visible particles	1.06	7.05
KP6.5	Clear, White, No visible particles	1.03	6.49
KP7.0	Clear, White, No visible particles	1.07	6.94
T8.0	Cloudy, White, No visible particles	1.04	7.84

[0439] Pharmaceutical compositions A4.0 and A5.0 were eliminated from further study as they showed precipitation and poor visual appearance. The remaining pharmaceutical compositions were stored at 5°C and percentage of HMWS (%HMWS) were measured by size exclusion chromatography (SEC) at 0, 7, and 30 days. Results are shown in Table 3.

Table 3: Aggregation (%HMWS) by SEC at 5°C

Sample	0 Days	7 Days	30 Days
S5.5	88.14	86.50	95.15
H5.5	58.21	60.79	87.86
H6.0	68.38	68.50	69.73
H6.5	28.39	27.64	28.64
H7.0	7.01	6.31	4.32
H7.5	6.63	7.07	4.29
KP6.5	24.00	22.15	23.92

KP7.0	18.48	17.34	16.86
T8.0	27.25	49.11	65.52
standard	2.56	2.48	2.85

[0440] Pharmaceutical compositions containing histidine buffer at pH 6.88 (H7.0) and pH 7.05 (H7.5) showed the lowest %HMWS. Slight loss in HMWS was observed at 1 month (30 days). Pharmaceutical compositions containing histidine buffer at pH 6.5 (H6.5) or potassium phosphate buffer at pH 6.5 (KP6.5) and pH 7 (KP7.0) produced comparable %HMWS, but showed significantly higher %HMWS than pharmaceutical compositions containing histidine buffer at pH 6.88 (H7.0) and pH 7.05 (H7.5).

[0441] The pharmaceutical compositions were stored at 25°C and %HMWS was measured by size exclusion chromatography (SEC) at 0, 0.25, 4, and 7 days. Results are shown in Table 4. Pharmaceutical compositions containing histidine buffer at pH 6.88 (H7.0) and pH 7.05 (H7.5) showed the lowest %HMWS at the start of the experiment (0 days).

Table 4: Aggregation (%HMWS) by SEC at 25°C

Sample	0 Days	0.25 Day	4 Days	7 Days
S5.5	88.14	87.53	94.28	95.30
H5.5	58.21	87.23	93.53	93.63
H6.0	68.38	78.07	91.17	93.00
H6.5	28.39	28.39	39.27	65.32
H7.0	7.01	7.47	14.07	16.72
H7.5	6.63	5.23	39.41	46.55
KP6.5	24.00	22.99	24.77	25.26
KP7.0	18.48	19.43	19.61	20.22
T8.0	27.25	27.45	25.17	46.22
standard	2.56	2.36	2.85	2.27

[0442] The pharmaceutical compositions were stored at 40°C and %HMWS was measured by size exclusion chromatography (SEC) at 0, 0.25, 1, and 7 days. Results are shown in Table 5.

Table 5: Aggregation (%HMWS) by SEC at 40°C

Sample	0 Days	0.25 Day	1 Day	7 Days
S5.5	88.14	97.11	97.25	97.21
H5.5	58.21	98.85	97.44	98.35
H6.0	68.38	72.70	82.80	96.48
H6.5	28.39	73.51	85.18	93.39
H7.0	7.01	22.06	42.36	68.71
H7.5	6.63	7.79	14.71	46.99
KP6.5	24.00	23.14	25.82	35.79
KP7.0	18.48	18.69	20.00	28.70
T8.0	27.25	30.33	24.96	58.07

standard	2.56	2.28	2.79	2.79
----------	------	------	------	------

[0443] Pharmaceutical compositions containing histidine buffer at pH 6.88 (H7.0) and pH 7.05 (H7.5) showed the lowest %HMWS at the start. Significant degradation of the efanesoctocog alfa was observed at 40°C for all buffers tested.

[0444] Concentration stability was also assessed for the pharmaceutical compositions at 5°C, 25°C, and 40°C at various time points. Results are shown in Table 6. For all but one sample, there was a slight increase in concentration (mg/mL) at 1 month at 5°C, demonstrating possible instability or aggregation; only potassium phosphate at pH 6.5 (KP6.5) did not show increase in concentration. The samples did not show any significant change in concentration after 1 week at all temperatures.

Table 6: Concentration Stability Results

pH BUFFER STUDY: Concentration (mg/mL), Time (days)																
5°C	Sample	0	7	30	25°C	Sample	0	0.25	4	7	40°C	Sample	0	0.25	1	7
	S5.5	1.01	1.01	1.25		S5.5	1.01	1.06	0.95	1.04		S5.5	1.01	1.02	0.97	1.04
	H5.5	1.05	1.04	1.68		H5.5	1.05	0.99	1.04	1.07		H5.5	1.05	1.06	0.96	1.06
	H6.0	1.05	1.05	1.50		H6.0	1.05	1.07	1.04	1.09		H6.0	1.05	1.06	1.01	1.12
	H6.5	1.06	1.06	1.62		H6.5	1.06	0.97	1.07	1.04		H6.5	1.06	1.11	1.08	1.11
	H6.8	1.06	1.04	1.38		H6.8	1.06	0.96	1.01	1.04		H6.8	1.06	1.10	1.07	1.07
	H7.3	1.07	1.06	1.41		H7.3	1.07	1.01	1.05	1.06		H7.3	1.07	1.03	1.05	1.03
	KP6.5	1.06	1.06	1.06		KP6.5	1.06	1.08	0.99	1.00		KP6.5	1.06	1.11	0.92	1.04
	KP7.0	1.07	1.01	1.36		KP7.0	1.07	1.07	1.04	1.04		KP7.0	1.07	1.08	1.08	1.02
	T8.0	1.10	1.00	1.55		T8.0	1.10	1.05	1.07	1.08		T8.0	1.10	1.11	1.03	1.06

[0445] pH stability was also assessed for the pharmaceutical compositions at 5°C, 25°C, and 40°C at various time points. pH values for the samples did not change significantly when stored for one month at 5°C, one week at 25°C, or 1 week at 40°C (data not shown). Measured values were all within +/- 0.5 units from time zero value over all time points.

[0446] Turbidity was also assessed for the pharmaceutical compositions at 5°C, 25°C, and 40°C at various time points. There was a 2- to 3-fold increase in absorbance in compositions containing potassium phosphate 7.0 (KP7.0) and histidine 5.5 (H5.5) after 7 days at 25°C and 40°C, respectively. Overall, the absorbance values did not change significantly when stored for one month at 5°C, one week at 25°C, or 1 week at 40°C (data not shown).

[0447] Summary

[0448] Pharmaceutical compositions containing histidine and potassium phosphate within a pH range of 6.8 to 7.3 showed minimal change in concentration, pH, and turbidity, as well as low

aggregation (%HMWS). Efanesoctocog alfa appears to be most stable in a pharmaceutical composition having a pH range of 6.8 to 7.3.

[0449] The initial aggregation was significantly lower in histidine buffer than in potassium phosphate buffer. The smallest increase in aggregation was seen in phosphate buffer pH 6.5 and pH 7.0 over 7 days at stress temperatures of 25°C and 40°C.

Example 2: Evaluation of histidine molarity and protein concentration on stability of efanesoctocog alfa

[0450] These determined the effect of the molarity of histidine and the concentration of protein on achieving a target pH of 7.0. The studies were also conducted to determine the effect of buffer molarity on the stability of efanesoctocog alfa at protein concentrations of 1, 0.67 and 0.045 mg/mL.

[0451] Study Design: Pharmaceutical compositions were studied with each containing one of three different molarities of histidine (10 mM, 20 mM, and 50 mM) and one of three different concentrations of efanesoctocog alfa (1 mg/mL, 0.67 mg/mL (4000 IU/mL), and 0.045 mg/mL (250 IU/mL)) in the presence of other excipients at constant concentrations. In addition to histidine and efanesoctocog alfa, all pharmaceutical compositions were formulated to include the following excipients: 250 mM Arginine hydrochloride (ArgHCl), 5 mM CaCl₂, 5% sucrose, and 0.05% polysorbate 80 (PS80). Liquid stability was evaluated for one week at 5°C, 25°C and 40°C. Stability of each pharmaceutical composition was evaluated by visual inspection, turbidity, protein concentration (Lunatic, Unchained Labs, Pleasanton CA), pH, osmolality, %HMWS by SEC, and protein concentration by RP-HPLC.

[0452] The maximum pH for pharmaceutical compositions comprising histidine at 10 mM, 20 mM, and 50 mM at any protein concentration (1 mg/mL, 4000 IU/mL, or 250 IU/mL) was 6.8, 6.95, and 7.1, respectively. Target pH 7.0 was achieved at 20 mM and 50 mM histidine. As the molarity of histidine increased, the pH of the formulation increased, potentially due to the increased buffering capacity. (see FIG. 2). There was no significant effect of protein concentration on the pH and there was no significant change in pH over time and temperature (data not shown).

[0453] At the start of the experiment (T₀), higher osmolality was observed as the histidine molarity increased from 10 mM to 50mM. Visual inspection showed the nine formulations studied at T₀ and 1 week at 5°C, 25°C and 40°C were clear and colorless. However, the protein aggregated significantly at 40 °C after 1 week, and there was a significant increase in %HMWS at 1 week at 40 °C. There was no effect of molarity on concentration and osmolality over 1 week at the three different temperatures (data not shown).

[0454] Turbidity results are shown in **Table 7**. Turbidity (NTU) showed a direct relation with protein concentration. At 5°C and 25°C the samples were clear at T₀ and remained clear after 1

week. Although turbidity increased significantly at 40°C for 1mg/mL and 4000IU samples, 1mg/mL samples were clear after 1 week. Further results of turbidity testing are shown in FIG. 3.

Table 7.

SAMPLE	NTU			
	T0	1 week		
	T0	5°C	25°C	40°C
10 mM HIS (1 mg/mL)	0.7	1.0	1.0	2.3
20 mM HIS (1 mg/mL)	0.7	1.3	1.0	3.0
50 mM HIS (1 mg/mL)	0.8	0.9	1.0	2.6
10 mM HIS 4000 IU	0.7	0.7	0.7	1.6
20 mM HIS 4000 IU	0.7	0.6	0.7	1.7
50 mM HIS 4000 IU	0.7	0.7	0.7	1.9
10 mM HIS 250 IU	0.7	0.1	0.1	0.1
20 mM HIS 250 IU	0.7	0.1	0.2	0.2
50 mM HIS 250 IU	0.7	0.1	n/a	0.2

Results of the analysis of %HMWS by SEC for samples observed at 5°C and 25°C are shown in FIG. 4. Aggregation was significantly higher for all molarities and protein concentrations at 40°C (data not shown). SEC testing revealed that the greater the protein concentration, the greater the %HMWS at T0. It was also observed that the %HMWS of efanesoctocog alfawas comparable at 5 °C, 25 °C, and 40 °C across all concentrations of histidine tested (10, 20 or 50 mM). Thus, changes in molarity of histidine did not appear to have an effect on aggregation levels. Samples containing 1mg/mL of protein showed aggregation increases with temperature.

[0455] Samples containing 4000 IU/mL (0.67mg/mL) of protein showed loss of aggregates after 1 week at 25°C. Samples containing 250 IU/mL (0.045 mg/mL) showed loss of aggregates after 1 week at at both 25°C and 40°C. Generally, at 25 °C, there was some HMWS aggregation, but over time the HMWS appeared to be breaking down into smaller molecules and presented as LMW peaks after a short duration (approximately 25 minutes). In contrast to 25 °C, HMWS did not appear to break down into LMW species at 40 °C. The 250 IU/mL (0.045 mg/mL) sample was not stable at 40 °C.

[0456] Summary

[0457] A maximum pH of 6.8 was achieved for the pharmaceutical compositions containing efanesoctocog alfaat 1 mg/mL, 4000 IU/mL, or 250 IU/mL; 10 mM histidine; 250 mM ArgHCl; 5% sucrose; 5 mM CaCl₂; and 0.05% PS80. Increasing molarity of histidine buffer showed no significant effects on pH, osmolality, protein concentration, turbidity, %HMWS at T0, or %HMWS over 1 week stability at 5°C, 25°C, and 40°C. Efanesoctocog alfashowed significant aggregation

at a stress temperature of 40°C after 1 week for all protein concentrations studied. For pharmaceutical compositions containing 4000 IU/mL and 250 IU/mL protein, significant loss of aggregate peaks was observed at 25°C and 40°C over 1 week. There was a significant increase in turbidity at 1 week at 40°C for pharmaceutical compositions containing 1 mg/mL and 4000IU/mL of efanesoctocog alfa.

Example 3: Evaluation of polysorbate 80 (PS80) concentration on protein stability

[0458] These studies were determined the effect of the concentration of PS80 on the stability of efanesoctocog alfa formulations.

[0459] Two concentrations of efanesoctocog alfa (0.67 mg/mL (4000 IU/mL) and 0.045 mg/mL (250 IU/mL)) in 10 mM Histidine, 250 mM Arginine hydrochloride (ArgHCl), 5 mM CaCl₂, 5% sucrose, and polysorbate 80 (PS80) were studied. The range of concentrations of PS80 for the 250 IU/mL samples was 0.002% to 0.07%. The range of concentrations of PS80 for the 4000 IU/mL samples was 0.03% to 0.1%. Samples were subjected to stress by orbital agitation (250 rpm for 0, 6, and 24 hours) or freeze/thaw (-80 °C not less than 24hrs, thaw at room temperature) for 0, 1, 3, and 5 cycles. Analysis included aggregation by SEC, PS80 concentration by high performance liquid chromatography (HPLC) with charged aerosol detection (CAD), particle test by micro-flow imaging (MFI) and high accuracy liquid particle counter (HIAC), pH, turbidity, and protein concentration. Results of each analysis is shown in Table 8.

[0460] Summary

[0461] The results of this study indicated that a concentration of PS80 $\geq$ 0.03% stabilized both 250IU and 4000IU concentrations of efanesoctocog alfa. No changes in PS80 content were observed for samples after 24 hrs agitation or 5 freeze/thaw cycles. No trend in particulates was observed for for the whole study of PS80 range. No trends or changes in pH or turbidity were observed over the PS80 range. Therefore, a pharmaceutical composition containing a concentration of PS80 $\geq$ 0.03% may increase stability of efanesoctocog alfa at a minimum concentration of 250 IU and a maximum concentration of 4000 IU with minimal impact on other composition properties.

Table 8: Sub-Visible Particles by MFI and HIAC

	Strength	PS80 Con. (%)	Agitation Study						Freeze/Thaw study					
			T0		T6hrs		T24hrs		F/T 1X		F/T 3X		F/T 5X	
			≥10µm	≥25µm	≥10µm	≥25µm	≥10µm	≥25µm	≥10µm	≥25µm	≥10µm	≥25µm	≥10µm	≥25µm
Particle # by MFI	250 IU	0.002	44	11	14	5	23	2	14	7	7	2	28	5
		0.005	11	5	11	7	9	5	9	0	11	0	9	2
		0.010	14	2	11	5	28	5	14	0	11	5	28	9
		0.030	39	7	39	7	14	7	2	0	21	0	5	2
		0.050	18	0	9	5	41	11	50	7	57	0	32	9
		0.070	14	7	57	18	25	5	16	5	9	0	0	0
Particle # by HIAC	250 IU	0.030	7	2	9	2	257	64	18	5	5	0	21	5
		0.050	7	2	11	0	18	5	41	14	16	5	7	5
		0.070	9	0	7	0	156	48	41	5	32	7	9	0
		0.100	11	2	9	5	119	32	7	0	16	2	11	2
		0.002	60	8	18	0	15	0	100	25	2	0	7	0
		0.005	27	0	27	0	7	2	0	0	0	0	0	0
Particle # by HIAC	250 IU	0.010	20	0	17	0	27	2	2	0	3	2	0	0
		0.030	28	3	20	0	33	2	2	2	3	0	2	0
		0.050	22	2	10	2	23	2	7	0	0	0	0	0
		0.070	17	2	43	2	10	0	23	7	2	0	2	0
		0.030	25	0	17	0	18	0	0	0	5	2	2	0
		0.050	12	2	25	0	12	2	2	0	2	0	2	0
Particle # by HIAC	4000 IU	0.070	10	2	2	0	17	3	0	0	2	0	3	0
		0.100	32	0	12	0	13	2	2	0	52	22	2	0

Example 4: Evaluation of excipient concentrations on protein stability

[0462] The experiments described in this example assess the stability of pharmaceutical compositions comprising efanesoctocog alfa drug product (DP) at 1000 IU formulated with dilution buffers containing varied concentrations of excipients. Minimum and maximum excipient concentration ranges were intentionally designed to deviate from current manufacturing tolerances in order to challenge the formulation robustness. A series of 13 different dilution buffers were evaluated (see Table 9). The Design of Experiment (DoE) model was based on five parameters (i.e., five excipients) at two levels (i.e., $\pm 5\%$ of target values). This included a control center point with the target formulation excipient quantities (i.e., Buffer #13).

[0463] Efanesoctocog alfa drug substance (DS) was pre-formulated at a concentration of 1 mg/mL in 10 mM L-Histidine, 250 mM L-Arginine Hydrochloride (HCl), 5 mM Calcium Chloride, 5 % (w/v) Sucrose and 0.05 % (w/v) Polysorbate 80 (PS80) at pH 7.01. Efanesoctocog alfa DS was diluted down to one of several strengths using one of the 13 dilution buffers to generate diluted drug substance (DDS) of the efanesoctocog alfa.

Table 9: Dilution Buffers Tested

Buffer No.	L-Arginine HCl (g/L)	Sucrose (g/L)	L-Histidine (g/L)	Calcium Chloride Dihydrate (g/L)	Polysorbate-80 (g/L)
1	50.032	52.500	1.630	0.698	0.475
2	55.298	47.500	1.474	0.698	0.525
3	55.298	52.500	1.474	0.698	0.475
4	55.298	47.500	1.630	0.772	0.525
5	50.032	47.500	1.630	0.772	0.475
6	55.298	52.500	1.630	0.772	0.525
7	50.032	52.500	1.474	0.772	0.525
8	50.032	52.500	1.630	0.698	0.525
9	55.298	47.500	1.630	0.698	0.475
10	50.032	47.500	1.474	0.698	0.525
11	55.298	52.500	1.474	0.772	0.475
12	50.032	47.500	1.474	0.772	0.475
13	52.665	50.000	1.552	0.735	0.500

[0464] Efanesoctocog alfa DDS was then lyophilized to generate efanesoctocog alfa DP at the intended strength per vial of 1000 IU. The resulting pharmaceutical compositions containing

efanesoctocog alfa DP at 1000 IU were assessed at various stability conditions (2 – 8 °C and 30 °C) and stability timepoints (T0, one month (T1), 3 months (T3), 6 months (T6), and 12 months (T12)). Pharmaceutical composition samples were tested for specific activity, protein concentration, aggregation, visual inspection, reconstitution time, pH, residual moisture, product/excipient crystallization, and glass transition temperature.

[0465] Protein concentration was measured using a RP-HPLC method with fluorescence detection. A stability specification range of 75 – 125 % of initial value was targeted for protein concentration.

[0466] Table 10 and FIG. 11 display concentration stability data up to 6 months (T6). Concentration values were generally lower than the associated target value of 167 µg/mL but remained within the 75 – 125 % of initial DP concentration value, thus falling within the acceptable range of the specification.

Table 10: Protein concentration results (µg/mL)

Buffer No.	T0 DDS	T0 DP 2-8 °C	T1 DP 30 °C	T3 DP 2-8 °C	T3 DP 30 °C	T6 DP 2-8 °C	T6 DP 30 °C	DP % Decrease T0 → T6
1	159.7	156.1	151.6	160.6	148.1	138.5	138.6	11.21
2	165.2	158.2	148.9	154.5	149.5	139.7	142	10.24
3	151.5	158.5	144.7	152.5	151.7	138	139.2	12.18
4	160.4	160.6	144.7	160	154.5	139.6	139.9	12.89
5	155.6	161.8	150.1	159.7	153.2	144	140.3	13.29
6	159.6	156.0	152.8	162	156	140.2	138.9	10.96
7	165.8	163.8	154.7	153.6	NA	138.9	144.5	11.78
8	155.4	161.8	156.3	150.7	154.2	134.9	140.7	13.04
9	157	162.2	160.8	153.1	155.2	134.1	141.8	12.58
10	157.7	169.5	161.9	154	157.6	140.6	141.8	16.34
11	154.1	165.8	160.6	155.4	151.1	135.4	138.2	16.65
12	161.2	170.2	170.4	160.6	159.9	141.8	141.2	17.04
13	145.4	167.9	162.7	152.6	153.1	142.5	137.6	18.05
Average	157.58	162.49	155.40	156.10	156.62	139.09	140.36	N/A
Std. Dev	5.49	4.76	7.63	3.88	11.10	2.96	1.91	N/A

[0467] Specific activities of DP and DDS were calculated using concentration data from Figure 11. Results are shown in Table 11 and Figure 13. The specific activity specification is > 1600 IU/ mg for

DDS and > 1280 IU/mg for DP of all strengths. For DP, the specific activity of each tested sample exceeded the specification at each time point.

[0468] Testing of efanesoctocog alfa DDS found that pre-lyophilization activity values were within the 80 – 125 % acceptable range of the specification.

Table 11: Specific activity results (IU/mg)

Buffer No.	T0 DDS	DDS Spec	T0 DP (2-8 °C)	T1 DP (30 °C)	T3 DP (2-8 °C)	T3 DP (30°C)	DP Spec
1	1748.3	1600	2053.7	1528.5	N/A	1497.0	1280.0
2	1679.8	1600	2138.1	1409.6	N/A	1554.5	1280.0
3	2093.7	1600	2251.5	1361.5	N/A	1624.3	1280.0
4	1791.1	1600	2036.6	1563.5	N/A	1622.7	1280.0
5	1822.0	1600	2060.9	1416.6	N/A	1560.1	1280.0
6	1830.8	1600	1946.4	1550.0	N/A	1443.6	1280.0
7	1769.6	1600	2040.5	1473.7	N/A	1352.60	1280.0
8	1718.8	1600	2098.6	1542.0	N/A	1655.0	1280.0
9	1759.2	1600	1600.6	1496.3	N/A	1607.6	1280.0
10	1750.8	1600	1620.5	1423.6	N/A	N/A	1280.0
11	1835.8	1600	1591.8	1504.8	N/A	1653.9	1280.0
12	1719.6	1600	1608.5	1426.6	N/A	1627.9	1280.0
13	N/A	1600	1626.2	1432.4	N/A	1659.0	1280.0

[0469] DP and DDS aggregation testing was performed by SEC-HPLC and results are shown in FIG. 7. At T0, DP aggregation ranged between 1.8 - 1.9 %. At T3 and T6, 30 °C samples displayed reduced aggregation values as low as 1.1 % which may be due to column performance or system variation. When comparing T0 to T6 2-8 °C results, no significant change in aggregation was observed. All aggregation values obtained are below 5% for efanesoctocog alfa DP. T1 30 °C and T3 2-8 °C sample aggregation analyses are yet to be completed.

[0470] Residual moisture values did not show significant variation on stability, with all results remaining below 1.2 %, within the 3 % stability specification limit.

[0471] Upon visual inspection, efanesoctocog alfa DP lyophilate cakes were observed to be white to off-white in color. These cakes fully dissolved upon reconstitution within ~ 60 seconds. Reconstitution values at T0 and T1 timepoints were slightly lower (within 25 seconds). All efanesoctocog alfa DP samples were reconstituted within 70 seconds (well below the proposed 3.5-minute specification), and no unexpected appearance observations were noted.

[0472] Samples were tested using powder X-ray diffraction. Samples appeared as amorphous content at T1 to T6 timepoints as confirmed by the amorphous halo pattern of the diffractogram. No evidence of product or excipient crystallisation such as crystallisation peaks was observed.

[0473] The glass transition temperature (T_g) was measured by differential scanning calorimetry (DSC) in duplicate for each sample; averaged results are shown in FIG. 16. The highest and lowest values for T_g were 71.75 °C (T6 30°C Buffer #10) and 61.95 °C (Buffer #3 T6 2-8 °C) respectively. The highest T_g value (Buffer #10 T6 30°C) coincided with the lowest moisture content (0.9 %) observed across the samples at that timepoint. The lowest T_g value (Buffer #3 T6 2-8 °C) coincided with the higher relative moisture content (1.11 %) observed for Buffer #3 at this timepoint. Samples did not show any obvious trend toward increasing or decreasing T_g values at different storage conditions on stability. Enthalpy values measured between 0.19 and 0.46 J/g°C.

[0474] All stability sample results shown in Figure 16 displayed T_g values above 60 °C. This is favorable toward stable long-term storage of the efanesoctocog alfa at a temperature of 2-8 °C, with an inclusive six month storage at room temperature.

Example 5: Evaluation of Arginine HCl concentration on protein stability

[0475] The experiments described in this example are used to assess the stability of efanesoctocog alfa in pharmaceutical compositions formulated with different concentrations of Arginine HCl.

[0476] Two pharmaceutical compositions are evaluated where the only difference is concentration of L-Arginine-HCl. The first composition contains 1.0 mg/mL efanesoctocog alfa in 10 mM Histidine, pH 7.0, 250 mM L-Arginine-HCl, 5 mM CaCl₂, 5% sucrose, and 0.05% Polysorbate-80. The second composition contains 1.0 mg/mL efanesoctocog alfa in 10 mM Histidine, pH 7.0, 175 mM L-Arginine-HCl, 5 mM CaCl₂, 5% sucrose, and 0.05% Polysorbate-80. Each composition is diluted approximately 60, 120 and 240 minutes after thawing at room temperature in preparation for size exclusion chromatography (SEC) analysis. Duplicate SEC injections of each sample are made and the % HMWS is determined for each injection.

[0477] Lyophilized efanesoctocog alfa drug product is reconstituted in 3.0 mL of water for injection (WFI). An aliquot of the reconstituted solution is transferred to an HPLC vial for analysis.

[0478] The SEC assay uses an HPLC instrument equipped with a pump, a temperature controlled autosampler, a column heater, and Fluorescence detector. The sample solutions are analyzed using the following instrumentation and method parameters:

- Mobile Phase: Dulbecco's Phosphate- Buffered Saline (D-PBS) containing Calcium and Magnesium adding 0.36 M Sodium Chloride (0.9 mM Calcium Chloride, 0.5 mM Magnesium Chloride, 2.7 mM Potassium Chloride, 1.5 mM Potassium Phosphate Mono basic, 496 mM Sodium Chloride, 8.1 mM Sodium Phosphate Dibasic, pH 7. 0 ± 0.1)
- Column Heater: 26°C
- Run Time: 40 minutes
- FL Detector: Ex/Em = 280/350 nm, PMT Gain = 5
- Injection Mass Load: 2 µg
- Autosampler Temperature: 5°C
- Flow Rate: 0.5 mL/min, Isocratic

[0479] Pharmaceutical compositions of efanesoctocog alfa containing 250 mM L-Arginine-HCl are found to be more stable than compositions containing 175 mM L-Arginine-HCl.

Example 6: Further evaluation of L-Arginine-HCl concentration on protein stability

[0480] The effect of arginine concentration over a broader range was also examined. Eight concentrations of L-arginine-HCl (50, 100, 125, 150, 175, 200, 250, and 300 mM) were added to a base formulation of 10 mM histidine, 5 mM calcium chloride, 5% sucrose, and 0.05% polysorbate 80. A summary of the study design is provided in Table 12. Upon thawing at room temperature (25°C), the formulations were prepared, placed in an HPLC vial and stored in the autosampler at 5°C ± 3°. HPLC injections for each time point were taken from the same vial. The % aggregation for each formulation were obtained. Aggregation was measured by SEC (%HMWS) for each sample.

[0481] The SEC assay uses an HPLC instrument equipped with a pump, a temperature controlled autosampler, a column heater, and Fluorescence detector. The sample solutions are analyzed using the following instrumentation and method parameters:

[0482] Mobile Phase: Dulbecco's Phosphate- Buffered Saline (D-PBS) containing Calcium and Magnesium adding 0.36 M Sodium Chloride (0.9 mM Calcium Chloride, 0.5 mM Magnesium Chloride, 2.7 mM Potassium Chloride, 1.5 mM Potassium Phosphate Mono basic, 496 mM Sodium Chloride, 8.1 mM Sodium Phosphate Dibasic, pH 7. 0 ± 0.1)

- Column Heater: 26°C
- Run Time: 40 minutes
- FL Detector: Ex/Em = 280/350 nm, PMT Gain = 5

- Injection Mass Load: 2 µg
- Autosampler Temperature: 5°C
- Flow Rate: 0.5 mL/min, Isocratic

Table 12. Arginine Concentration Study Overview

Study	efanesoctocog alfa concentration	Arginine Levels	Study conditions	Study timepoints	Quality Attributes
Selection of L-Arginine HCl as a stabilizing excipient	1 mg/mL	50, 100, 125, 150, 175, 200, 250, and 300 mM	Freezing at -80°C and ambient room temperature thaw	40-, 80-, 120-, and 160-minutes post thaw	% Aggregates by SE-HPLC

[0483] Two experiments (each run in duplicate) were run using the conditions outlined in Table 12. The results from Experiment 1 are shown in Table 13 and **FIG. 27**. In Experiment 1, High Molecular Weight (HMW) aggregates were found at a higher percentage in formulations with lower concentrations of arginine. Higher concentrations of arginine (200 mM to 300 mM) exhibited a consistently low percentage of HMW aggregates out to 160 minutes post-thaw.

Table 13. High Molecular Weight Aggregate Percentage – Experiment 1

Arginine	%HMW AREA				
	0 min	40 min	80 min	120 min	160 min
0 mM	4.24	4.22	4.25	4.24	4.26
50 mM	8.38	8.27	8.16	8.05	7.93
100 mM	7.36	7.13	6.94	6.83	6.60
125 mM	7.35	6.92	6.56	6.12	5.85
150 mM	4.79	4.42	4.32	4.19	4.01
175 mM	5.73	5.13	4.46	4.26	3.97
200 mM	4.10	3.77	3.78	3.54	3.63
250 mM	3.72	3.71	3.70	3.71	3.70
300 mM	3.60	3.60	3.60	3.59	3.61

[0484] The results from Experiment 2 can be seen in Table 14 and **FIG. 28**. In Experiment 2, HMW aggregates were found at a higher percentage in formulations with lower concentrations of arginine. Consistent with the observations of Experiment 1, Experiment 2 also showed that, higher

concentrations of arginine (200 mM to 300 mM) exhibited a consistently low percentage of HMW aggregates out to 160 minutes post-thaw.

Table 14. High Molecular Weight Aggregate Percentage – Experiment 2

Arginine	%HMW AREA				
	0 min	40 min	80 min	120 min	160 min
0 mM	4.24	4.22	4.25	4.24	4.26
50 mM	8.38	8.27	8.16	8.05	7.93
100 mM	7.36	7.13	6.94	6.83	6.60
125 mM	7.35	6.92	6.56	6.12	5.85
150 mM	4.79	4.42	4.32	4.19	4.01
175 mM	5.73	5.13	4.46	4.26	3.97
200 mM	4.10	3.77	3.78	3.54	3.63
250 mM	3.72	3.71	3.70	3.71	3.70
300 mM	3.60	3.60	3.60	3.59	3.61

[0485] The osmolality of samples across the range of arginine concentrations was also studied in two separate experiments (each run in duplicate). The osmolality (mOsm/kg) of the formulations was shown to increase with increased concentration of arginine. In Experiment 1, the first set of replicates was assayed on the osmometer after having been out at room temperature for 24 hours. The second set of replicates is from the retains that were stored at -80°C. The results from Experiment 1 are displayed in **Table 15**. In Experiment #2, the single set of replicate samples is from the retains that were stored at -80°C. The results from Experiment 2 are displayed in **Table 16**.

Table 15. Osmolality of formulations with increasing arginine concentrations – Experiment 1

	Experiment 1	
	Rep. 1 (RT)	Rep. 2 (-80°C)
Sample ID	mOsm/kg	mOsm/kg
290 reference	295	288
0 mM	218	n/a
50 mM Arg	327	331
100 mM Arg	371	420
125 mM Arg	408	425
150 mM Arg	495	484
175 mM Arg	530	524

200 mM Arg	534	537
250 mM Arg	655	667
300 mM Arg	758	699

Table 16. Osmolality of formulations with increasing arginine concentrations – Experiment 2

	Experiment 2 (-80°C)
Sample ID	mOsm/kg
290 reference	288
0 mM	184
50 mM Arg	302
100 mM Arg	356
125 mM Arg	398
150 mM Arg	476
175 mM Arg	513
200 mM Arg	538
250 mM Arg	n/a
300 mM Arg	701

[0486] The pH was not adjusted for the samples tested, and the pH of formulations across the range of arginine concentrations was subsequently assessed. The results are shown in Table 17.

[0487] Table 17. pH of formulations with increasing arginine concentrations

Sample	Desired Arg concentration	replicate 1	replicate 2	replicate 3
		pH as measured by pH meter 1	pH as measured by pH meter 2	pH as measured by pH meter 3
Formulation 1	0	7.00	6.99	6.95
Formulation 2	50	6.74	6.74	6.76
Formulation 3	100	6.74	6.70	6.75
Formulation 4	125	6.74	6.69	6.70
Formulation 5	150	6.69	6.65	6.67
Formulation 6	175	6.69	6.62	6.67
Formulation 7	200	6.67	6.60	6.70
Formulation 8	250	6.54	6.54	6.65
Formulation 9	300	6.50	6.51	6.62

[0488] Conclusion: In these experiments, surprisingly the formulations comprising efanesoctocog alfa having at least 250 mM arginine showed the lowest %HMW at time 0 (T0) and subsequent timepoints tested (see FIG. 27 and FIG. 28), while lower levels of arginine increased aggregation at T0. At the beginning of time zero, the control, 0% arginine, had a T0 of ~ 4% HMW area, and the 50 mM, 100 mM, 125 mM and 150 mM started with a substantially greater %HMW, showing that the arginine at low concentrations from 50 mM – 200 mM began at higher aggregation levels that were greater than the control, 0 mM arginine. There was a decrease in %HMW area over time at the 50 mM, 100 mM, 125 mM and 150 mM concentrations.

[0489] Thus, the use of at least 250 mM arginine in the formulation appears to stabilize efanesoctocog alfa over time with the least amount of aggregation.

Example 7: Evaluation of low sucrose on protein stability

[0490] The experiments described in this example assess the stability of efanesoctocog alfa in pharmaceutical compositions formulated with different concentrations of sucrose. In addition to sucrose, each composition also contained 10 mM L-histidine, 250 mM L-arginine-HCl, 5 mM CaCl₂, and 0.05% (w/v) Polysorbate 80, at a pH of 6.8.

[0491] Efanesoctocog alfa drug substance (DS) comprises 1 mg/ml efanesoctocog alfa, 10 mM L-histidine, 250 mM L-arginine-HCl, 5 mM CaCl₂, and 0.05% (w/v) polysorbate 80, and sucrose, at a pH of 6.8. Efanesoctocog alfa drug product (DP) is DS diluted to a desired activity in international units (IU), such as IU per vial (e.g., IU per 3.367 mL when a vial is filled with 3.367 mL of drug product). DP may be lyophilized (e.g., in a vial) resulting in lyophilized DP that may then be reconstituted prior to injection.

[0492] Aggregation of efanesoctocog alfa drug substance (DS) compositions (1 mg/mL) containing 5% or 1% sucrose were tested at either room temperature (RT)/room light (RL) conditions or at 2-8 °C. As shown in FIG. 9 and FIG. 10, DS compositions containing 5% sucrose showed less aggregation (%HMWS) over time than DS compositions containing 1% sucrose.

[0493] To further analyze the impact on sucrose concentration on efanesoctocog alfa compositions, pharmaceutical compositions of efanesoctocog alfa at 0, 1, 2, and 5% w/v sucrose were subjected to different stability conditions and stability was assessed using several methods. Compositions were tested at the following product stages: drug substance (DS, which was had not

been lyophilized or adjusted for any particular IU strength) frozen storage stability, DS freeze/thaw (FT) stability, liquid DS post-thaw stability, diluted DS and drug product (DP) stability, DP lyophilization stability, and DP-post reconstitution. Drug substance was tested at freezing temperatures (-80°C or -30°C). Drug product samples were tested at room temperature (RT), 5°C, 30°C, and 40°C. Lyophilized DP samples were tested at two different DP concentrations (4000 IU and 250 IU).

[0494] Aggregation of efanesoctocog alfa bulk drug product (BDP) compositions at 250 IU were tested at either room temperature (RT)/room light (RL) conditions or at 2-8 °C. BDP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Results at RT/RL are shown in FIG. 11 and results at 2-8 °C are shown in FIG. 12. BDP compositions containing 0% sucrose showed the highest aggregation levels for both conditions. However, none of the BDP compositions at 250 IU showed significant increases in %HMWS over the 24 hour time period tested. For all compositions and both conditions, it was observed that aggregation levels began to decrease over time after about 7 days (data not shown).

[0495] Aggregation of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU or 4000 IU were tested at 5°C, 30°C, and 40°C. Aggregation was measured by SEC (%HMWS) for each sample. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months. Results of Lyo DP compositions at 250 IU are shown in FIG. 13 (5°C), FIG. 14 (30°C), and FIG. 15 (40°C). Results of Lyo DP compositions at 4000 IU are shown in FIG. 16 (5°C), FIG. 17 (30°C), and FIG. 18 (40°C). Generally, all sucrose concentrations showed no significant increase in %HMWS over 6 months at 5°C, 30°C and 40°C. None of the compositions tested showed significant aggregation trends.

[0496] Aggregation of efanesoctocog alfa DS and 4000 IU bulk drug product (BDP) liquid compositions containing either 2% or 5% sucrose were tested by SEC. Samples were held at room temperature or 5°C and tested after 0, 5, 7, 25, 43, and 55 hours. Samples were also tested after lyophilization. Results are shown in FIG. 19. Less aggregation over time was observed for compositions containing 5% sucrose. However, compositions containing 2% sucrose did not show significant increases in aggregation over the time period tested.

[0497] A summary of the results of the aggregation testing is provided in Table 18. An increase 1.0 or more %HMWS from a baseline value set forth as the corresponding value for 5% sucrose was deemed to be unacceptable. 0% sucrose was found to be unacceptable.

Table 18: Sucrose screening stability results (%HMWS).

Sucrose conc, %	Frozen Storage, % (-80°C) (1 mg/mL)		Liquid DS Rate %/h (1 mg/mL)	Liquid BDP Rate %/h (4000 IU)		Cum. %/h (4000 IU)	Lyo DP (Δ6M) (4000 IU)			Lyo DP (Δ6M) (250 IU)		
	Δ6M	Δ5XFT		5°C	RT		5°C	30°C	40°C	5°C	30°C	40°C
5	0.3	1.0	0.03	0.01	0.02	0.01	0.3	0.2	0.4	0.0	-0.1	0.0
2	0.3	1.2	0.02	0.02	0.02	0.02	0.5	0.7	0.9	0.0	0.2	0.0
1	0.3	1.3	0.08	0.03	0.04	0.03	0.7	0.9	0.3	0.1	0.0	0.0
0	0.6	2.2	ND	ND	ND	ND	0.2	0.0	-0.2	0.2	0.1	0.1

ND = Not Determined

[0498] As shown in Table 18, stability results indicated that freeze/thaw and frozen storage stability data was favorable for 1% to 5% w/v sucrose at -80°C. Freeze/thaw and frozen storage stability was unfavorable at all concentrations at -30 °C (data not shown). DP samples showed a minor but acceptable increase in aggregation through 6 months at 40 °C for the 1% and 2% formulations.

[0499] The glass transition temperature (T_g) of compositions of efanesoctocog alfa DP at 250 IU or 4000 IU were tested at 5°C, 30°C, and 40°C. T_g was measured by modulated differential scanning calorimetry (DSC). DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T₀, 1 month, 2 months, 3 months, and 6 months. Results are shown in FIG. 20 (5°C), FIG. 21 (30°C), and FIG. 22 (40°C). Generally, T_g was observed to correlate with sucrose concentration, with the higher sucrose concentration showing lower T_g. No significant changes in T_g were observed at 5°C, 30°C and 40°C through 6 months for all compositions. No major changes in T_g were observed at the different DP concentrations.

[0500] The glass transition temperature (T_g) of efanesoctocog alfa drug product (DP) containing 0, 1, 2, and 5% w/v sucrose was measured by differential scanning calorimetry (DSC). DP samples were tested at two different DP concentrations (4000 IU and 250 IU). Results are shown in FIG. 23.

[0501] The DSC protocol for T_g determination includes:

- (i) equilibration at 15°C,
- (ii) modulate at 1°C every 60 seconds,
- (iii) isothermal for 5 minutes

(iv) ramp at 3°C/min to 130°C.

[0502] Residual moisture content of compositions of efanesoctocog alfa lyophilized drug product (Lyo DP) at 250 IU or 4000 IU were tested at 5°C, 30°C, and 40°C. Lyo DP compositions containing 0, 1, 2, and 5% w/v sucrose were tested. Samples were tested at T0, 1 month, 2 months, 3 months, and 6 months. Results are shown in FIG. 24 (5°C), FIG. 25 (30°C), and FIG. 26 (40°C). No significant change in residual moisture content was observed at 5°C, 30°C and 40°C through 6 months for all compositions. All residual moisture values fell below the spec of 3%.

[0503] Upon visual inspection, concentration of sucrose had no effect on cake appearance. over 6 months at 2-8°C, 30°C and 40°C (data not shown).

[0504] Based on the results of these experiments, sucrose concentrations of 1% to 2% (w/v) appear to be the acceptable concentrations for pharmaceutical compositions of efanesoctocog alfa.

[0505] The foregoing description of the specific embodiments will so fully reveal the general nature of the disclosure that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present disclosure. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[0506] Other embodiments of the disclosure will be apparent to those skilled in the art from consideration of the specification and practice of the disclosure disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the disclosure being indicated by the following claims.

[0507] All patents and publications cited herein are incorporated by reference herein in their entirety.

SEQUENCES

Table 19: Exemplary Chimeric protein Sequences

A-FVIII(ELNN)-Fc: SEQ ID NO: 1						
ATRRYYLGAV	ELSWDMQSD	LGELPVDARF	PPRVPKSFPP	NTSVVYKCTL	FVEFTDHLFN	60
IAKPRPPWVG	LLGPTIQAEV	YDTVVITLKN	MASHPVSLHA	VGVSYWKASE	GAEYDDQTSQ	120
REKEDDKVFP	GGSHTYVWQV	LKENGPMASD	PLCLTYSYLS	HVDLVKDLNS	GLIGALLVCR	180
EGSLAKEKTQ	TLHKFILLFA	VFDEGKSWHS	ETKNSLMQDR	DAASARAWPK	MHTVNGYVNR	240
SLPGLIGCHR	KSVYWHVIGM	GTTPEVHSIF	LEGHTFLVRN	HRQASLEISP	ITFLTAQTLL	300
MDLGQFLLC	HISSHQHDMG	EAYVKVDSCP	EEPQLRMKN	EEAEDYDDL	TDSEMDVVR	360
DDDNPSFIIQ	IRSVAKKHPK	TWVHYIAAEE	EDWDYAPLVL	APDDRSYKSQ	YLNNGPQRIG	420
RKYKKVREMA	YTDETFKTR	AIQHESGILG	PLLYGEVGDT	LLIIFKNQAS	RPYNIYPHGT	480
TDVRPLYSR	LPGVKHLKLD	FPILPGEIFK	YKWTVTVEDG	PTKSDPRCLT	RYSSSFVNME	540
RDLASGLIGP	LLICYKESVD	QRGNQIMSDK	RNVILFSVFD	ENRSWYLTEN	TQRFLEPNPAG	600
VQLEDPEFQA	SNIMHSINGY	VFDSLQLSVC	LHEVAYWYIL	SIGAQTDFLS	VFFSGYTFKH	660
KMVYEDTLTL	FPFSGETVFM	SMENPGLWIL	GCHNSDFRNR	GMTALLKVSS	CDKNTGDYEE	720
DSYEDISAYL	LSKNNATIEPR	SFSQNGTSES	ATPESGPGSE	PATSGSETPG	TSESATPESG	780
PGSEPATSGS	ETPGTSESAT	PESGPGTSTE	PSEGSAPGSP	AGSPTSTEEG	TSESATPESG	840
PGSEPATSGS	ETPGTSESAT	PESGPGSPAG	SPTSTEEGSP	AGSPTSTEEG	TSTEPSGESA	900
PGTSESATPE	SGPGTSESAT	PESGPGTSTE	ATPESGPGSE	PATSGSETPG	SEPATSGSET	960
PGSPAGSPTS	TEEGTSTEPS	EGSAPGTSTE	PSEGSAPGSE	PATSGSETPG	TSESATPESG	1020
PGTSTEPS	SAPASSITR	TTLQSDQEEI	DYDDTISVEM	KKEDFDIYDE	DENQSPRSFQ	1080
KKTRHYFTIA	VERLWDYGMS	SSPHVLNRNA	QSGSVPQFKK	VVFQFTDGS	FTQPLYRGEL	1140
NEHLGLLGPY	IRAEVEDNIM	VTFRNQASRP	YSFYSSLISY	EEDQRQGAEP	RKNFVKPNET	1200
KTYFWKVQHH	MAPTKDEFDC	KAWAYFSDVD	LEKDVHSGLI	GPLLVCHTNT	LNPAGHRQVT	1260
VQEFALFFETI	FDETKSWYFT	ENMERNCRAP	CNIQMEDPTF	KENYRFHAIN	GYIMDTLPGL	1320
VMAQDQRIRW	YLLSMGNSNEN	IHSIHFSGHV	FTVRKKEEYK	MALYNLYPGV	FETVEMLPK	1380
AGTWRVECLI	GEHLHAGMST	LFLVYSNKCQ	TPLGMASGHI	RDFQITASGQ	YGQWAPKLAR	1440
LHYSGSINAW	STKEPFSWIK	VDLLAPMIIH	GIKTQGARQK	FSSLYISQFI	IMYSLDGKKW	1500
QTYRGNSTGT	LMVFFGNVDS	SGIKHNIENP	PIIARYIRLH	PTHYSIRSTL	RMELMGCDLN	1560
SCSMPLGMES	KAISDAQITA	SSYFTNMFAT	WSPSKARLHL	QGRSNAWRPQ	VNNPKEWLQV	1620
DFQKTMKVTG	VTTQGVKSLL	TSMYVKEFLI	SSSQDGHQWT	LFFQNGKVKV	FQGNQDSFT	1680
VVNSLDPPLL	TRYLRHPQS	WVHQIALRME	VLGCEAQDLY	DKTHTCPPCP	APPELLGGPSV	1740
FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	GVEVHNAKTK	PREEQYNSTY	1800
RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	GQPREPQVYT	LPPSRDELTK	1860
NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTTPVLD	DGSFFLYSKL	TVDKSRWQQG	1920
NVFCSCVMHE	ALHNHYTQKS	LSLSPG				
VWF(D'D3)- ELNN-a2 Linker-Fc: SEQ ID NO: 2						
SLSCRPPMVK	LVCADNLR	EGLECTKTCQ	NYDLECMSMG	CVSGCLCPPG	MVRHENRCVA	60
LERCPCHFQG	KEYAPGETVK	IGCNTCVCRD	RKWNCTDHVC	DATCSTIGMA	HYLTFDGLKY	120
LFPGEQCYVL	VQDYCGSNPG	TFRILVGNKG	CSHPSVKCKK	RVTILVEGGE	IELFDGEVNV	180
KRPMKDETHF	EVVSESGRYII	LLLGKALSIV	WDRHLSISVV	LKQTYQEKVC	GLCGNFDGIQ	240
NNDLTSSNLQ	VEEDPVDGFG	SWKVSSQCAD	TRKVPLDSSP	ATCHNNIMKQ	TMVDSSCRIL	300
TSDVFQDCNK	LVDPEPYLDV	CIYDTCSCES	IGDCAAFCDT	TAAYAHVCAQ	HGKVVTWRTA	360
TLCPQSCER	NLRENGYEA	WRYNSCAPAC	QVTCQHPEPL	ACPVCVEGC	HAHCPPGKIL	420
DELLQTCVDP	EDCPVCEVAG	RRFASGKKVT	LNPSPDEHCQ	ICHCDVNNLT	CEACQEPGTS	480
ESATPESGPG	SEPATSGSET	PGTSESATPE	SGPGSEPATS	GSETPGTSES	ATPESGPGTS	540
TEPSEGSAPG	SPAGSPTSTE	EGTSESATPE	SGPGSEPATS	GSETPGTSES	ATPESGPGSP	600
AGSPTSTEEG	SPAGSPTSTE	EGASSDKNTG	DYYEDSYEDI	SAYLLSKNNA	IEPRSFSDKT	660
HTCPPCPAPE	LLGGPSVFLF	PPKPKDTLMI	SRTPEVTCV	VDVSHEDPEV	KFNWYVDGVE	720
VHNAKTKPRE	EQYNSTYRVV	SVLTVLHQDW	LNGKEYKCKV	SNKALPAIE	KTISKAKGQP	780
REPQVYTLPP	SRELTKNQV	SLTCLVKGYF	PSDIAVEWES	NGQPENNYKT	TPPVLDSDGS	840
FFLYSKLTVD	KSRWQQGNVF	SCSVMHEALH	NHYTQKSLSL	SPG		883

Amino acid sequence of FVIII protein (312A) – includes signal peptide: SEQ ID NO: 3

1	MQIELSTCFF	LCLLRFCFSA	TRRYYLGAVE	LSWDYMQSDL	GELPVDARFP
51	PRVPKSFPPN	TSVVYKKTLF	VEFTDHLFNI	AKPRPPWMGL	LGPTIQAEVY
101	DTVVITLKNM	ASHPVSLHAV	GVSYWKASEG	AEYDDQTSQR	EKEDDKVFPG
151	GSHTYVWQVL	KENGPMASDP	LCLTYSYLSH	VDLVKDLNSG	LIGALLVCRE
201	GSLAKEKTQT	LHKFILLFAV	FDEGKSWHSE	TKNSLMQDRD	AASARAWPKM
251	HTVNGYVNRN	LPGLIGCHRK	SVYWHVIGMG	TTPEVHSIFL	EGHTFLVRNH
301	RQASLEISPI	TFLTAQTLLM	DLGQFLLFCH	ISSHQHDGME	AYVKVDSCPE
351	EPQLRMKNNE	EAEDYDDDLT	DSEMDVVREF	DDNSPSFIQI	RSVAKKHPKT
401	WVHYIAAEEE	DWDYAPLVLA	PDDRSYKSQY	LNNGPQRIGR	KYKKVRFMAY
451	TDETFKTREA	IQHESGILGP	LLYGEVGDIT	LIIIFKNQASR	PYNIYPHGIT
501	DVRPLYSRRL	PKGKHLKDF	PILPGEIFKY	KWTVTVEDGP	TKSDPRCLTR
551	YYSSFVNMER	DLASGLIGPL	LICYKESVDQ	RGNQIMSDKR	NVILFSVFDE
601	NRSWYLTENI	QRFLPNPAGV	QLEDPEFQAS	NIMHSINGYV	FDSLQLSVCL
651	HEVAYWYILS	IGAQTDFLSV	FFSGYTFFKH	MVYEDTLTLF	PFSGETVFMS
701	MENPGLWILG	CHNSDFRNRG	MTALLKVSSC	DKNTGDYIED	SYEDISAYLL
751	SKNNAIEPRS	FSQNGTSESA	TPESGPGSEP	ATSGSETPGT	SESATPESGP
801	GSEPATSGSE	TPGTSESATP	ESGPGTSTEP	SEGSAPGSPA	GSPTSTEEGT
851	SESATPESGP	GSEPATSGSE	TPGTSESATP	ESGPGSPAGS	PTSTEEGSPA
901	GSPTSTEEGT	STEPSEGSAP	GTSESATPES	GPGTSESATP	ESGPGTSESA
951	TPESGPGSEP	ATSGSETPGS	EPATSGSETP	GSPAGSPTST	EEGTSTEPSE
1001	GSAPGTSTEP	SEGSAPGSEP	ATSGSETPGT	SESATPESGP	GTSTEPSEGS
1051	APASSEITRT	TLQSDQEEID	YDDTISVEMK	KEDFDIYDED	ENQSPRSFQK
1101	KTRHYFIAAV	ERLWDYGMSS	SPHVLNRNAQ	SGSVPPQFKKV	VFQEFDTGGSF
1151	TQPLYRGELN	EHLGLLGPYI	RAEVEDNIMV	TFRNQASRPY	SFYSSLISYE
1201	EDQRQGAEP	KNFVKPNETK	TYFWKVQHMH	APTKDEFDCK	AWAYFSDVDL
1251	EKDVSGLIG	PLLVCHTNTL	NPAHGRQVTV	QEFALFFTIF	DETKSWYFTE
1301	NMERNCRAPC	NIQMEDPTFK	ENYRFHAING	YIMDTLPGLV	MAQDQIRIRWY
1351	LLSMGSNENI	HSIHFSGHVF	TVRKKEEYKM	ALYNLYPGVF	ETVEMLPSCA
1401	GIWRVECLIG	EHLHAGMSTL	FLVYSNKCQT	PLGMASGHIR	DFQITASGQY
1451	GQWAPKLARL	HYSGSINAWS	TKEPFSWIKV	DLLAPMIIHG	IKTQGARQKF
1501	SSLYISQFII	MYSLDGKKWQ	TYRGNSTGTL	MVFFGNVDSS	GIKHNIFFNPP
1551	IIARYIRLHP	THYSIRSTLR	MELMGCDLNS	CSMPLGMESK	AISDAQITAS
1601	SYFTNMFATW	SPSKARLHLQ	GRSNAWRPQV	NNPKEWLQVD	FQKTMKVTGV
1651	TTQGVKSLLT	SMYVKEFLIS	SSQDGHQWTL	FFQNGKVVKV	QCNQDSFTPV
1701	VNSLDPPLLT	RYLRIHPQSW	VHQIALRMEV	LGCEAQDLYD	KHTCPCPCA
1751	PELLGGPSVF	LFPPKPKDTL	MISRTPEVTC	VVVDVSHEDP	EVKFNWYVDG
1801	VEVHNAKTKP	REEQYNSTYR	VVSVLTVLHQ	DWLNGKEYKC	KVSNKALPAP
1851	IEKTISKAKG	QPREPQVYTL	PPSRDELTKN	QVSLTCLVKG	FYPDSIAVEW
1901	ESNGQPENNY	KTTTPVLDS	GSFFLYSKLT	VDKSRWQQGN	VFSCSVMHEA
1951	LHNHYTQKSL	SLSPG*			

Nucleotide sequence encoding FVIII 312A: SEQ ID NO: 4

1	ATGCAAATAG	AGCTCTCCAC	CTGCTTCTTT	CTGTGCCTTT	TGCGATTCTG
51	CTTTAGTGCC	ACCAGAAGAT	ACTACCTGGG	TGCAGTGGAA	CTGTCATGGG

101	ACTATATGCA	AAGTGATCTC	GGTGAGCTGC	CTGTGGACGC	AAGATTTTCCT
151	CCTAGAGTGC	CAAAATCTTT	TCCATTCAAC	ACCTCAGTCG	TGTACAAAAA
201	GACTCTGTTT	GTAGAATTCA	CGGATCACCT	TTTCAACATC	GCTAAGCCAA
251	GGCCACCCTG	GATGGGTCTG	CTAGGTCCTA	CCATCCAGGC	TGAGGTTTAT
301	GATACAGTGG	TCATTACACT	TAAGAACATG	GCTTCCCATC	CTGTCAGTCT
351	TCATGCTGTT	GGTGTATCCT	ACTGGAAAGC	TTCTGAGGGA	GCTGAATATG
401	ATGATCAGAC	CAGTCAAAGG	GAGAAAGAAG	ATGATAAAGT	CTTCCCTGGT
451	GGAAGCCATA	CATATGTCTG	GCAGGTCCTG	AAAGAGAATG	GTCCAATGGC
501	CTCTGACCCA	CTGTGCCTTA	CCTACTCATA	TCTTTCTCAT	GTGGACCTGG
551	TAAAAGACTT	GAATTCAGGC	CTCATTGGAG	CCCTACTAGT	ATGTAGAGAA
601	GGGAGTCTGG	CCAAGGAAAA	GACACAGACC	TTGCACAAAT	TTATACTACT
651	TTTTGCTGTA	TTTGATGAAG	GGAAAAGTTG	GCACTCAGAA	ACAAAGAAGT
701	CCTTGATGCA	GGATAGGGAT	GCTGCATCTG	CTCGGGCCTG	GCCTAAAAATG
751	CACACAGTCA	ATGGTTATGT	AAACAGGTCT	CTGCCAGGTC	TGATTGGATG
801	CCACAGGAAA	TCAGTCTATT	GGCATGTGAT	TGGAATGGGC	ACCACCTCTG
851	AAGTGCACTC	AATATTCTCT	GAAGGTCACA	CATTTCTTGT	GAGGAACCAT
901	CGCCAGGCGT	CCTTGGAAT	CTCGCCAATA	ACTTTCTCTA	CTGCTCAAAC
951	ACTCTTGATG	GACCTTGGAC	AGTTTCTIAC	GTTTTGTCTA	ATCTCTTCCC
1001	ACCAACATGA	TGGCATGGAA	GCTTATGTCA	AAGTAGACAG	CTGTCCAGAG
1051	GAACCCCAAC	TACGAATGAA	AAATAATGAA	GAAGCGGAAG	ACTATGATGA
1101	TGATCTTACT	GATTCTGAAA	TGGATGTGGT	CAGGTTTGAT	GATGACAACT
1151	CTCCTTCCTT	TATCCAAATT	CGCTCAGTTG	CCAAGAAGCA	TCCTAAAACT
1201	TGGGTACATT	ACATTGCTGC	TGAAGAGGAG	GACTGGGACT	ATGCTCCCTT
1251	AGTCCTCGCC	CCCGATGACA	GAAGTTATAA	AAGTCAATAT	TTGAACAATG
1301	GCCCTCAGCG	GATTGGTAGG	AAGTACAAAA	AAGTCCGATT	TATGGCATACT
1351	ACAGATGAAA	CCTTTAAGAC	TCGTGAAGCT	ATTCAGCATG	AATCAGGAAT
1401	CTTGGGACCT	TTACTTTATG	GGGAAGTTGG	AGACACACTG	TTGATTATAT
1451	TTAAGAATCA	AGCAAGCAGA	CCATATAACA	TCTACCCTCA	CGGAATCACT
1501	GATGTCCGTC	CTTTGTATTC	AAGGAGATTA	CCAAAAGGTG	TAAAACATTT
1551	GAAGGATTTT	CCAATTCTGC	CAGGAGAAAT	ATTCAAATAT	AAATGGACAG
1601	TGACTGTAGA	AGATGGGCCA	ACTAAATCAG	ATCCTCGGTG	CCTGACCCGC
1651	TATTACTCTA	GTTTCGTTAA	TATGGAGAGA	GATCTAGCTT	CAGGACTCAT
1701	TGGCCCTCTC	CTCATCTGCT	ACAAAGAATC	TGTAGATCAA	AGAGGAAACC
1751	AGATAATGTC	AGACAAGAGG	AATGTCATCC	TGTTTTCTGT	ATTTGATGAG
1801	AACCGAAGCT	GGTACCTCAC	AGAGAATATA	CAACGCTTTC	TCCCCAATCC
1851	AGCTGGAGTG	CAGCTTGAGG	ATCCAGAGTT	CCAAGCCTCC	AACATCATGC
1901	ACAGCATCAA	TGGCTATGTT	TTTGATAGTT	TGCAGTTGTC	AGTTTGTGTTG
1951	CATGAGGTGG	CATACTGGTA	CATTCTAAGC	ATTGGAGCAC	AGACTGACTT
2001	CCTTTCTGTC	TTCTTCTCTG	GATATACCTT	CAAACACAAA	ATGGTCTATG
2051	AAGACACACT	CACCCATATC	CCATTCTCAG	GAGAAACTGT	CTTCATGTCG
2101	ATGGAAAACC	CAGGTCTATG	GATTCTGGGG	TGCCACAAC	CAGACTTTTCG
2151	GAACAGAGGC	ATGACCGCCT	TACTGAAGGT	TTCTAGTTGT	GACAAGAACA
2201	CTGGTGATTA	TTACGAGGAC	AGTTATGAAG	ATATTTTCAGC	ATACTTGCTG
2251	AGTAAAAACA	ATGCCATTGA	ACCAAGAAGC	TTCTCTCAAA	ACGGTACCTC
2301	AGAGTCTGCT	ACCCCCGAGT	CAGGGCCAGG	ATCAGAGCCA	GCCACCTCCG
2351	GGTCTGAGAC	ACCCGGGACT	TCCGAGAGTG	CCACCCCTGA	GTCCGGACCC
2401	GGGTCCGAGC	CCGCCACTTC	CGGCTCCGAA	ACTCCCGGCA	CAAGCGAGAG
2451	CGCTACCCCA	GAGTCAGGAC	CAGGAACATC	TACAGAGCCC	TCTGAAGGCT
2501	CCGCTCCAGG	GTCCCCAGCC	GGCAGTCCCA	CTAGCACCAG	GGAGGGAACC
2551	TCTGAAAGCG	CCACACCCGA	ATCAGGGCCA	GGGTCTGAGC	CTGCTACCAG
2601	CGGCAGCGAG	ACACCAGGCA	CCTCTGAGTC	CGCCACACCA	GAGTCCGGAC
2651	CCGGATCTCC	CGCTGGGAGC	CCCACCTCCA	CTGAGGAGGG	ATCTCCTGCT

2701	GGCTCTCCAA	CATCTACTGA	GGAAGGAACC	TCAACCGAGC	CATCCGAGGG
2751	ATCAGCTCCC	GGCACCTCAG	AGTCGGCAAC	CCCGGAGTCT	GGACCCGGAA
2801	CTTCCGAAAG	TGCCACACCA	GAGTCCGGTC	CCGGGACTTC	AGAATCAGCA
2851	ACACCCGAGT	CCGGCCCTGG	GTCTGAACCC	GCCACAAGTG	GTAGTGAGAC
2901	ACCAGGATCA	GAACCTGCTA	CCTCAGGGTC	AGAGACACCC	GGATCTCCGG
2951	CAGGCTCACC	AACCTCCACT	GAGGAGGGCA	CCAGCACAGA	ACCAAGCGAG
3001	GGCTCCGCAC	CCGGAACAAG	CACTGAACCC	AGTGAGGGTT	CAGCACCCGG
3051	CTCTGAGCCG	GCCACAAGTG	GCAGTGAGAC	ACCCGGCACT	TCAGAGAGTG
3101	CCACCCCCGA	GAGTGGCCCA	GGCACTAGTA	CCGAGCCCTC	TGAAGGCAGT
3151	GCGCCAGCCT	CGAGCGAAAT	AACCTCGTACT	ACTCTTCAGT	CAGATCAAGA
3201	GGAAATTGAC	TATGATGATA	CCATATCAGT	TGAAATGAAG	AAGGAAGATT
3251	TTGACATTTA	TGATGAGGAT	GAAAATCAGA	GCCCCCGCAG	CTTCAAAAG
3301	AAAACACGAC	ACTATTTTAT	TGCTGCAGTG	GAGAGGCTCT	GGGATTATGG
3351	GATGAGTAGC	TCCCCACATG	TTCTAAGAAA	CAGGGCTCAG	AGTGGCAGTG
3401	TCCCTCAGTT	CAAGAAAGTT	GTTTTCCAGG	AATTTACTGA	TGGCTCCTTT
3451	ACTCAGCCCT	TATACCGTGG	AGAACTAAAT	GAACATTTGG	GACTCCTGGG
3501	GCCATATATA	AGAGCAGAAG	TTGAAGATAA	TATCATGGTA	ACTTTCAGAA
3551	ATCAGGCCCT	TCGTCCCTAT	TCCTTCTIAT	CTAGCCTTAT	TTCTTATGAG
3601	GAAGATCAGA	GGCAAGGAGC	AGAACCTAGA	AAAAACTTTG	TCAAGCCTAA
3651	TGAAACCAAA	ACTTACTTTT	GGAAAGTGCA	ACATCATATG	GCACCCACTA
3701	AAGATGAGTT	TGACTGCAAA	GCCTGGGCTT	ATTTCTCTGA	TGTTGACCTG
3751	GAAAAAGATG	TGCACTCAGG	CCTGATTGGA	CCCCTTCTGG	TCTGCCACAC
3801	TAACACACTG	AACCCTGCTC	ATGGGAGACA	AGTGACAGTA	CAGGAATTTG
3851	CTCTGTTTTT	CACCATCTTT	GATGAGACCA	AAAGCTGGTA	CTTCACTGAA
3901	AATATGGAAA	GAACTGCAG	GGCTCCCTGC	AATATCCAGA	TGGAAGATCC
3951	CACTTTTTAA	GAGAATTATC	GCTTCCATGC	AATCAATGGC	TACATAATGG
4001	ATACACTACC	TGGCTTAGTA	ATGGCTCAGG	ATCAAAGGAT	TCGATGGTAT
4051	CTGCTCAGCA	TGGGCAGCAA	TGAAAACATC	CATTCTATTG	ATTTCAGTGG
4101	ACATGTGTTC	ACTGTACGAA	AAAAAGAGGA	GTATAAAATG	GCCTGTACAC
4151	ATCTCTATCC	AGGTGTTTTT	GAGACAGTGG	AAATGTTACC	ATCCAAAGCT
4201	GGAAATTTGG	GGGTGGAATG	CCTTATTGGC	GAGCATCTAC	ATGCTGGGAT
4251	GAGCACACTT	TTTCTGGTGT	ACAGCAATAA	GTGTCAGACT	CCCCTGGGAA
4301	TGGCTTCTGG	ACACATTAGA	GATTTTCAGA	TTACAGCTTC	AGGACAATAT
4351	GGACAGTGGG	CCCCAAAGCT	GGCCAGACTT	CATTATTCCG	GATCAATCAA
4401	TGCCTGGAGC	ACCAAGGAGC	CCTTTTCTTG	GATCAAGGTG	GATCTGTTGG
4451	CACCAATGAT	TATTCACGGC	ATCAAGACCC	AGGGTGCCCG	TCAGAAGTTC
4501	TCCAGCCTCT	ACATCTCTCA	GTTTATCATC	ATGTATAGTC	TTGATGGGAA
4551	GAAGTGGCAG	ACTTATCGAG	GAAATTCCAC	TGGAACCTTA	ATGGTCTTCT
4601	TTGGCAATGT	GGATTCATCT	GGGATAAAAC	ACAATATTTT	TAACCCTCCA
4651	ATTATTGCTC	GATACATCCG	TTTGCACCCA	ACTCATTATA	GCATTGCGAG
4701	CACTCTTTCG	ATGGAGTTGA	TGGGCTGTGA	TTTAAATAGT	TGCAGCATGC
4751	CATTGGGAAT	GGAGAGTAAA	GCAATATCAG	ATGCACAGAT	TACTGCTTCA
4801	TCCTACTTTA	CCAATATGTT	TGCCACCTGG	TCTCCTTCAA	AAGCTCGACT
4851	TCACCTCCAA	GGGAGGAGTA	ATGCCTGGAG	ACCTCAGGTG	AATAATCCAA
4901	AAGAGTGGCT	GCAAGTGGAC	TTCCAGAAGA	CAATGAAAGT	CACAGGAGTA
4951	ACTACTCAGG	GAGTAAAATC	TCTGCTTACC	AGCATGTATG	TGAAGGAGTT
5001	CCTCATCTCC	AGCAGTCAAG	ATGGCCATCA	GTGGACTCTC	TTTTTTCAGA
5051	ATGGCAAAGT	AAAGGTTTTT	CAGGGAAATC	AAGACTCCTT	CACACCTGTG
5101	GTGAACTCTC	TAGACCCACC	GTTACTGACT	CGCTACCTTC	GAATTCACCC
5151	CCAGAGTTGG	GTGCACCAGA	TTGCCCTGAG	GATGGAGGTT	CTGGGCTGCG
5201	AGGCACAGGA	CCTCTACGAC	AAAACCTACA	CATGCCCACC	GTGCCCAGCT
5251	CCAGAACTCC	TGGGCGGACC	GTCAGTCTTC	CTCTTCCCCC	CAAAACCCAA

5301	GGACACCCTC	ATGATCTCCC	GGACCCCTGA	GGTCACATGC	GTGGTGGTGG
5351	ACGTGAGCCA	CGAAGACCCT	GAGGTCAAGT	TCAACTGGTA	TGTGGACGGC
5401	GTGGAAGTGC	ATAATGCCAA	GACAAAGCCG	CGGGAGGAGC	AGTACAACAG
5451	CACGTACCGT	GTGGTCAGCG	TCCTCACCGT	CCTGCACCAA	GACTGGCTGA
5501	ATGGCAAGGA	GTACAAGTGC	AAGGTCTCCA	ACAAAGCCCT	CCCAGCCCCC
5551	ATCGAGAAAA	CCATCTCCAA	AGCCAAAGGG	CAGCCCCGAG	AACCACAGGT
5601	GTACACCCTG	CCCCATCCC	GGGATGAGCT	GACCAAGAAC	CAAGTTAGCC
5651	TGACCTGCCT	GGTCAAAGGC	TTCTATCCCA	GCGACATCGC	CGTGGAGTGG
5701	GAGAGCAATG	GGCAGCCGGA	GAACAACACT	AAGACCACGC	CTCCCGTGT
5751	GGACTCCGAC	GGCTCCTTCT	TCCTCTACTC	CAAGCTCACC	GTGGACAAGA
5801	GCAGGTGGCA	GCAGGGGAAC	GTCTTCTCAT	GCTCCGTGAT	GCATGAGGCT
5851	CTGCACAACC	ACTACACGCA	GAAGAGCCTC	TCCCTGTCTC	CGGGTTGA

**Amino acid sequence of VWF059A protein – includes signal peptide and D1D2 region:
SEQ ID NO: 5**

1	MIPARFAGVL	LALALILPGT	LCAEGTRGRS	STARCSLFGS	DFVNTFDGSM
51	YSFAGYCSYL	LAGGCQKRSE	SIIGDFQNGK	RVSLSVYLGE	FFDIHLFVNG
101	TVTQGDQQRVS	MPYASKGLYL	ETEAGYYKLS	GEAYGFVARI	DGSGNFQVLL
151	SDRYFNKTCG	LCGNFNIFAE	DDFMTQEGTL	TSDPYDFANS	WALSSGEQWC
201	ERASPPSSSC	NISSGEMQKG	LWEQCQLLKS	TSVFARCHPL	VDPEPFVALC
251	EKTLCECAGG	LECACPALLE	YARTCAQEGM	VLYGWDHSA	CSPVCPAGME
301	YRQCVSPCAR	TCQSLHINEM	CQERCVDGCS	CPEGQLLDEG	LCVESTECPC
351	VHSGKRYPPG	TSLSRDCNTC	ICRNSQWICS	NEECPGECVL	TGQSHFKSFD
401	NRYFTFSGIC	QYLLARDCQD	HSFSIVIVETV	QCADDRDAVC	TRSVTVRLPG
451	LHNSLVKLKH	GAGVAMDQGD	IQLPLLKGD	RIQHTVTASV	RLSYGEDLQM
501	DWDGRGRLLV	KLSPVYAGKT	CGLCGNYNGN	QGDDFLTPSG	LAEPREDFG
551	NAWKHLGDCQ	DLQKQHSDDC	ALNPRMTRES	EEACAVLTSP	TFEACHRAVS
601	PLPYLRNCRY	DVCSCSDGRE	CLCGALASYA	AACAGRGVRV	AWREPGRCEL
651	NCPKGQVYLQ	CGTPCNLTCT	SLSYPDEECN	EACLEGCFCP	PGLYMDERGD
701	CVPKACQPCY	YDGEIFQPED	IFSDHHTMCY	CEDGFMHCTM	SGVPGSLLPD
751	AVLSSPLSHR	SKRSLSCRPP	MVKLVCPADN	LRAEGLECTK	TCQNYDLECM
801	SMGCVSGCLC	PPGMVRHENR	CVALERCPCF	HQKEYAPGE	TVKIGCNTCV
851	CRDRKWNCTD	HVCDATCSTI	GMAHYLTFDG	LKYLFPGEQ	YVLVQDYCGS
901	NPGETFRILVG	NKGCSHPSVK	CKKRVITLVE	GGEIELFDGE	VNVKRPMDKE
951	THFEVVESEGR	YIILLGKAL	SVVWDRHLSI	SVVLKQTYQE	KVCGLCGNFD
1001	GIQNNDLTSS	NLQVEEDPVD	FGNSWKVSSQ	CADTRKVPLD	SSPATCHNNI
1051	MKQTMVDSSC	RILTSDFVQD	CNKLVDPPEY	LDVCIYDTC	CESIGDCAAF
1101	CDTIAAYAHV	CAQHGVVVTW	RTATLCPQSC	EERNLRENGY	EAEWRYNCSA
1151	PACQVTCQHP	EPLACPVCV	EGCHAHCPPG	KILDELLQTC	VPEDPCVCE
1201	VAGRRFASGK	KVTILNPSDPE	HCQICHCDVV	NLTCEACQEP	GTSESATPES
1251	GPSEPATSG	SETPGTSESA	TPESGPGSEP	ATSGSETPGT	SESATPESGP
1301	GTSTEPSEGS	APGSPAGSPT	STEEGTSESA	TPESGPGSEP	ATSGSETPGT
1351	SESATPESGP	GSPAGSPTST	EEGSPAGSPT	STEEGASSDK	NTGDYEDSY
1401	EDISAYLLSK	NNAIEPRSF	DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT
1451	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	GVEVHNAKTK	PREEQYNSTY
1501	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	GQPREPQVYT
1551	LPPSRDELTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTTPVLDS
1601	DGSFFLYSKL	TVDKSRWQQG	NVFSCSVMHE	ALHNHYTQKS	LSLSPG*

Nucleotide sequence encoding VWF059A protein: SEQ ID NO: 6

1	ATGATTCTG	CCAGATTTC	CGGGGTGCTG	CTTGCTCTGG	CCCTCATTTT
51	GCCAGGGACC	CTTTGTGCAG	AAGGAAGTCTG	CGGCAGGTCA	TCCACGGGCC
101	GATGCAGCCT	TTTCGGAAGT	GACTTCGTCA	ACACCTTTGA	TGGGAGCATG
151	TACAGCTTTG	CGGGATACTG	CAGTTACCTC	CTGGCAGGGG	GCTGCCAGAA
201	ACGCTCCTTC	TCGATTATTG	GGGACTTCCA	GAATGGCAAG	AGAGTGAGCC
251	TCTCCGTGTA	TCTTGGGGAA	TTTTTTGACA	TCCATTTGTT	TGTCAATGGT
301	ACCGTGACAC	AGGGGGACCA	AAGAGTCTCC	ATGCCCTATG	CCTCCAAAGG
351	GCTGTATCTA	GAAACTGAGG	CTGGGTACTA	CAAGCTGTCC	GGTGAGGCCT
401	ATGGCTTTGT	GGCCAGGATC	GATGGCAGCG	GCAACTTTCA	AGTCTGTCTG
451	TCAGACAGAT	ACTTCAACAA	GACCTGCGGG	CTGTGTGGCA	ACTTTAACAT
501	CTTTGCTGAA	GATGACTTTA	TGACCCAAGA	AGGGACCTTG	ACCTCGGACC
551	CTTATGACTT	TGCCAACTCA	TGGGCTCTGA	GCAGTGGAGA	ACAGTGGTGT
601	GAACGGGCAT	CTCCTCCAG	CAGCTCATGC	AACATCTCCT	CTGGGGAAAT
651	GCAGAAGGGC	CTGTGGGAGC	AGTGCCAGCT	TCTGAAGAGC	ACCTCGGTGT
701	TTGCCCCTG	CCACCTCTG	GTGGACCCCG	AGCCTTTTGT	GGCCCTGTGT
751	GAGAAGACTT	TGTGTGAGTG	TGCTGGGGGG	CTGGAGTGCG	CCTGCCCTGC
801	CCTCCTGGAG	TACGCCCGGA	CCTGTGCCCA	GGAGGGAATG	GTGCTGTACG
851	GCTGGACCGA	CCACAGCGCG	TGCAGCCAG	TGTGCCCTGC	TGGTATGGAG
901	TATAGGCAGT	GTGTGTCCCC	TTGCGCCAGG	ACCTGCCAGA	GCCTGCACAT
951	CAATGAAATG	TGTCAGGAGC	GATGCGTGGA	TGGCTGCAGC	TGCCCTGAGG
1001	GACAGCTCCT	GGATGAAGGC	CTCTGCGTGG	AGAGCACCGA	GTGTCCCTGC
1051	GTGCATTCCG	GAAAGCGCTA	CCCTCCCGGC	ACCTCCCTCT	CTCGAGACTG
1101	CAACACCTGC	ATTTGCCGAA	ACAGCCAGTG	GATCTGCAGC	AATGAAGAAT
1151	GTCCAGGGGA	GTGCCTTGTC	ACTGGTCAAT	CCCACTTCAA	GAGCTTTGAC
1201	AACAGATACT	TCACCTTCAG	TGGGATCTGC	CAGTACCTGC	TGGCCCGGGA
1251	TTGCCAGGAC	CACTCCTTCT	CCATTGTGAT	TGAGACTGTC	CAGTGTGCTG
1301	ATGACCGCGA	CGCTGTGTGC	ACCCGCTCCG	TCACCGTCCG	GCTGCCTGGC
1351	CTGCACAACA	GCCTTGTGAA	ACTGAAGCAT	GGGGCAGGAG	TGCCATAGGA
1401	TGGCCAGGAC	ATCCAGCTCC	CCCTCCTGAA	AGGTGACCTC	CGCATCCAGC
1451	ATACAGTGAC	GGCCTCCGTG	CGCCTCAGCT	ACGGGGAGGA	CCTGCAGATG
1501	GACTGGGATG	GCCGCGGGAG	GCTGCTGGTG	AAGCTGTCCC	CCGTCTATGC
1551	CGGGAAGACC	TGCGGCCTGT	GTGGGAATTA	CAATGGCAAC	CAGGGCGACG
1601	ACTTCCTTAC	CCCCTCTGGG	CTGGCGGAGC	CCCGGGTGGA	GGACTTCGGG
1651	AACGCCTGGA	AGCTGCACGG	GGACTGCCAG	GACCTGCAGA	AGCAGCACAG
1701	CGATCCCTGC	GCCCTCAACC	CGCGCATGAC	CAGGTTCTCC	GAGGAGGCGT
1751	GCGCGGTCT	GACGTCCCCC	ACATTGAGAG	CCTGCCATCG	TGCCGTACAG
1801	CCGCTGCCCT	ACCTGCGGAA	CTGCCGCTAC	GACGTGTGCT	CCTGCTCGGA
1851	CGGCCGCGAG	TGCCTGTGCG	GCGCCCTGGC	CAGCTATGCC	GCGGCCTGCG
1901	CGGGGAGAGG	CGTGCGCGTC	GCGTGCGCGC	AGCCAGGCCG	CTGTGAGCTG
1951	AACTGCCCGA	AAGGCCAGGT	GTACCTGCAG	TGCGGGACCC	CCTGCAACCT
2001	GACCTGCCGC	TCTCTCTCTT	ACCCGGATGA	GGAATGCAAT	GAGGCCTGCC
2051	TGGAGGGCTG	CTTCTGCCCC	CCAGGGCTCT	ACATGGATGA	GAGGGGGGAC
2101	TGCGTGCCCA	AGGCCCAAGT	CCCCTGTTAC	TATGACGGTG	AGATCTTCCA
2151	GCCAGAAGAC	ATCTTCTCAG	ACCATCACAC	CATGTGCTAC	TGTGAGGATG
2201	GCTTCATGCA	CTGTACCATG	AGTGAGTCC	CCGGAAGCTT	GCTGCCTGAC
2251	GCTGTCTCTA	GCAGTCCCCT	GTCTCATCGC	AGCAAAAGGA	GCCTATCCTG
2301	TCGGCCCCCC	ATGGTCAAGC	TGGTGTGTCC	CGCTGACAAAC	CTGCGGGCTG
2351	AAGGGCTCGA	GTGTACCAAA	ACGTGCCAGA	ACTATGACCT	GGAGTGCATG
2401	AGCATGGGCT	GTGTCTCTGG	CTGCCTCTGC	CCCCCGGGCA	TGGTCCGGCA

2451	TGAGAACAGA	TGTGTGGCCC	TGGAAAGGTG	TCCCTGCTTC	CATCAGGGCA
2501	AGGAGTATGC	CCCTGGAGAA	ACAGTGAAGA	TTGGCTGCAA	CACTTGTGTC
2551	TGTCGGGACC	GGAAGTGGAA	CTGCACAGAC	CATGTGTGTG	ATGCCACGTG
2601	CTCCACGATC	GGCATGGCCC	ACTACCTCAC	CTTCGACGGG	CTCAAATACC
2651	TGTTCCCCGG	GGAGTGCCAG	TACGTTCTGG	TGCAGGATTA	CTGCGGCAGT
2701	AACCCTGGGA	CCTTTCGGAT	CCTAGTGGGG	AATAAGGGAT	GCAGCCACCC
2751	CTCAGTGAAG	TGCAAGAAAC	GGGTACCAT	CCTGGTGGAG	GGAGGAGAGA
2801	TTGAGCTGTT	TGACGGGGAG	GTGAATGIGA	AGAGGCCCAT	GAAGGATGAG
2851	ACTCACTTTG	AGGTGGTGGA	GTCTGGCCGG	TACATCATTC	TGCTGCTGGG
2901	CAAAGCCCTC	TCCGTGGTCT	GGGACCGCCA	CCTGAGCATC	TCCGTGGTCC
2951	TGAAGCAGAC	ATACCAGGAG	AAAGTGTGTG	GCCTGTGTGG	GAATTTTGAT
3001	GGCATCCAGA	ACAATGACCT	CACCAGCAGC	AACCTCCAAG	TGGAGGAAGA
3051	CCCTGTGGAC	TTTGGGAACT	CCTGGAAAGT	GAGCTCGCAG	TGTGCTGACA
3101	CCAGAAAAGT	GCCTCTGGAC	TCATCCCCTG	CCACCTGCCA	TAACAACATC
3151	ATGAAGCAGA	CGATGGTGGA	TTCCTCCTGT	AGAATCCTTA	CCAGTGACGT
3201	CTTCCAGGAC	TGCAACAAGC	TGGTGGACCC	CGAGCCATAT	CTGGATGICT
3251	GCATTTACGA	CACCTGCTCC	TGTGAGTCCA	TTGGGGACTG	CGCCGCATTC
3301	TGCGACACCA	TTGCTGCCTA	TGCCCACGTG	TGTGCCACGC	ATGGCAAGGT
3351	GGTGACCTGG	AGGACGGCCA	CATTGTGCCC	CCAGAGCTGC	GAGGAGAGGA
3401	ATCTCCGGGA	GAACGGGTAT	GAGGCTGAGT	GGCGCTATAA	CAGCTGTGCA
3451	CCTGCCTGTC	AAGTCACGTG	TCAGCACCTT	GAGCCACTGG	CCTGCCCTGT
3501	GCAGTGTGTG	GAGGGCTGCC	ATGCCCACTG	CCCTCCAGGG	AAAATCCTGG
3551	ATGAGCTTTT	GCAGACCTGC	GTTGACCCTG	AAGACTGTCC	AGTGTGTGAG
3601	GTGGCTGGCC	GGCGTTTTGC	CTCAGGAAAG	AAAGTCACCT	TGAATCCCAG
3651	TGACCCTGAG	CACCTGCCAG	TTTGCCACTG	TGATGTTGTC	AACCTCACCT
3701	GTGAAGCCTG	CCAGGAGCCG	GGTACATCAG	AGAGCGCCAC	CCCTGAAAGT
3751	GGTCCCGGGA	GCGAGCCAGC	CACATCTGGG	TCGGAACGCG	CAGGCACATC
3801	CGAGTCTGCA	ACTCCCGAGT	CCGGACCTGG	CTCCGAGCCT	GCCACTAGCG
3851	GCTCCGAGAC	TCCGGGAACT	TCCGAGAGCG	CTACACCAGA	AAGCGGACCC
3901	GGAAACCAGTA	CCGAACCTAG	CGAGGGCTCT	GCTCCGGGCA	GCCCAGCCGG
3951	CTCTCCTACA	TCCACGGAGG	AGGGCACTTC	CGAATCCGCC	ACCCCGGAGT
4001	CAGGGCCAGG	ATCTGAACCC	GCTACCTCAG	GCAGTGAGAC	GCCAGGAACG
4051	AGCGAGTCCG	CTACACCGGA	GAGTGGGCCA	GGGAGCCCTG	CTGGATCTCC
4101	TACGTCCACT	GAGGAAGGGT	CACCAGCGGG	CTCGCCACCC	AGCACTGAAG
4151	AAGGTGCCTC	GTCTGACAAG	AACACTGGTG	ATTATTACGA	GGACAGTTAT
4201	GAAGATATTT	CAGCATACTT	GCTGAGTAAA	AACAATGCCA	TTGAACCAAG
4251	AAGCTTCTCT	GACAAAACCT	ACACATGCCC	ACCGTGCCCA	GCTCCAGAAC
4301	TCCTGGGCGG	ACCGTCAGTC	TTCCTCTTCC	CCCCAAAACC	CAAGGACACC
4351	CTCATGATCT	CCCGGACCCC	TGAGGTCACT	TGCGTGGTGG	TGGACGTGAG
4401	CCACGAAGAC	CCTGAGGTCA	AGTTCAACTG	GTATGTGGAC	GGCGTGGAAG
4451	TGCATAATGC	CAAGACAAAG	CCGCGGGAGG	AGCAGTACAA	CAGCACGTAC
4501	CGTGTGGTCA	GCGTCTCAC	CGTCTGTCAC	CAAGACTGGC	TGAATGGCAA
4551	GGAGTACAAG	TGCAAGGTCT	CCAACAAAGC	CCTCCAGGCC	CCCATCGAGA
4601	AAACCATCTC	CAAAGCCAAA	GGGCAGCCCC	GAGAACCACA	GGTGTACACC
4651	CTGCCCCCAT	CCCGGGATGA	GCTGACCAAG	AACCAAGTTA	GCCTGACCTG
4701	CCTGGTCAAA	GGCTTCTATC	CCAGCGACAT	CGCCGTGGAG	TGGGAGAGCA
4751	ATGGGCAGCC	GGAGAACAAC	TACAAGACCA	CGCCTCCCGT	GTTGGACTCC
4801	GACGGCTCCT	TCTTCCTCTA	CTCCAAGCTC	ACCGTGGACA	AGAGCAGGTG
4851	GCAGCAGGGG	AACGTCTTCT	CATGCTCCGT	GATGCATGAG	GCTCTGCACA
4901	ACCACTACAC	GCAGAAGAGC	CTCTCCCTGT	CTCCGGGTTG	A

FVIII(ELNN)-Fc: SEQ ID NO: 7

TRRYYLGAWE	LSWDYMQSDL	GELPVDARFP	PRVPKSFPPN	TSVVYKKTLE	VEFTDHLFNI	60
AKPRPPWML	LGPTIQAEVY	DTVITLKNM	ASHPVSLHAV	GVSYWKASEG	AEYDDQTSQR	120
EKEDDKVFP	GSHTYVWQVL	KENGPMASDP	LCLTYSYLSH	VDLVKDLNSG	LIGALLVCRE	180
GSLAKEKTQT	LHKFILLFAV	FDEGKSWHSE	TKNSLMQDRD	AASARAWPKM	HTVNGYVNRS	240
LPGLIGCHRK	SVYWHVIGMG	TTPEVHSIFL	EGHTFLVRNH	RQASLEISPI	TFLTAQTLLM	300
DLGQFLFLFCH	ISSHQHDGME	AYVKVDSCPE	EPQLRMKNNE	EAEDYDDDLT	DSEMDVVRFD	360
DDNSPSFTIQI	RSVAKKHPKT	WVHYIAAEEE	DWDYAPLVLA	PDDRSYKSQY	LNNGPQRIGR	420
KYKKVRFEMAY	TDETFKTREA	IQHESGILGP	LLYGEVGDTL	LIIFKNQASR	PYNIYPHGIT	480
DVRPLYSRRL	PKGVKHLKDF	PILPGEIFKY	KWTVTVEDGP	TKSDPRCLTR	YYSSTFVNMER	540
DLASGLIGPL	LICYKESVDQ	RGNQIMSDKR	NVILFSVFDE	NRSWYL TENI	QRFLPNPAGV	600
QLEDPEFQAS	NIMHSINGYV	FDSLQLSVCL	HEVAYWYILS	IGAQTDFLSV	FFSGYTFFKHK	660
MVYEDTLTLF	PFSGETVEMS	MENPGLWILG	CHNSDFRNRG	MTALLKVSSC	DKNTGDYVED	720
SYEDISAYLL	SKNNAIEPRS	FSQNGTSESA	TPESGPGSEP	ATSGSETPGT	SESATPESGP	780
GSEPATSGSE	TPGTSESATP	ESGPGTSTEP	SEGSAPGSPA	GSPTSTEEGT	SESATPESGP	840
GSEPATSGSE	TPGTSESATP	ESGPGSPAGS	PTSTEEGSPA	GSPTSTEEGT	STEPSEGSAP	900
GTSESATPES	GPGTSESATP	ESGPGTSESA	TPESGPGSEP	ATSGSETPGS	EPATSGSETP	960
GSPAGSPSTST	EEGTSTEPSE	GSAPGTSTEP	SEGSAPGSEP	ATSGSETPGT	SESATPESGP	1020
GTSTEPSEGS	APASSEITRT	TLQSDQEEID	YDDTISVEMK	KEDFDIYDED	ENQSPRSFQK	1080
KTRHYFIAAV	ERLWDYGMSS	SPHVLRNRAQ	SGSVPQFKKV	VFQEFDTGSF	TQPLYRGELN	1140
EHLGLLGPYI	RAEVEDNIMV	TFRNQASRPY	SEYSSLI SYE	EDQRQGAEP	RNFVKPNETK	1200
TYFWKVQHHM	APTKDEFDCK	AWAYFSDVDL	EKDVHSGLIG	PLLVCHTNTL	NPAHGRQVTV	1260
QEFALFFTFIF	DETKSWYFTE	NMERNCRAPC	NIQMEDPTFK	ENYRFHAING	YIMDTLPGLV	1320
MAQDQRIRWY	LLSMGNSNENI	HSIHFSGHVF	TVRKKEEYKM	ALYNLYPGVF	ETVEMLPSPA	1380
GIWRVECLIG	EHLHAGMSTL	FLVYSNKCQT	PLGMASGHIR	DFQITASGQY	GQWAPKLARL	1440
HYSGSINAWS	TKEPFSWIKV	DLLAPMI IHG	IKTQGARQKF	SSLYISQFII	MYSLDGKKWQ	1500
TYRGNSTGTL	MVFFGNVDSS	GIKHNI FNPP	IIARYIRLHP	THYSIRSTLR	MELMGCDLNS	1560
CSMPLGMESK	AISDAQITAS	SYFTNM FATW	SPSKARLHLQ	GRSNAWRPQV	NNPKEWLQVD	1620
FQKTMKVTVG	TTQGVKSLLT	SMYVKEFLIS	SSQDGHQWTL	FFQNGKVKVF	QGNQDSFTPV	1680
VNSLDPLPLT	RYLRHPPQSW	VHQIALRMEV	LGCEAQDLYD	KTHTCPPCPA	PELLGGPSVF	1740
LFPPKPKDTL	MISRTPEVTC	VVDVSHEDP	EVKFNWYVDG	VEVHNAKTKP	REEQYNSTYR	1800
VVSVLTVLHQ	DWLNGKEYKC	KVSNKALPAP	IEKTISKAKG	QPREPQVYTL	PPSRDELTKN	1860
QVSLTCLVKG	FYPSDIAVEW	ESNGQPENNY	KTPPPVLDS	GSFFLYSKLT	VDKSRWQQGN	1920
VFSCSVMHEA	LHNHYTQKSL	SLSPG				1945

Table 20. Additional chimeric protein sequences

Description / SEQ ID NO.	Sequence
Full length FVIII	ATTRYYLGAV ELSDYMQSD LGELPVDARF PPRVPKSFPP NTSVVYKKTLE FVEFTDHLFN 60
	IAKPRPPWML LLGPTIQAEV YDTVITLKN MASHPVSLHA VGVSYWKASE GAEDDDQTSQ 120
	REKEDDKVFP GGSHTYVWQV LKENGPMASD PLCLTYSYLS HVDLVKDLNS GLIGALLVCR 180
SEQ ID NO: 8	EGSLAKEKTQ TLHKFILLFA VFDEGKSWHS ETKNLSMQDR DAASARAWPK MHTVNGYVNR 240
	SLPGLIGCHR KSVYWHVIGM GTTPEVHSIF LEGHTFLVRN HRQASLEISP ITFLTAQTLL 300
	MDLGQFLFLFC HISSHQHDGM EAYVKVDSCP EEPQLRMKNN EEAEDYDDDL TDSEMDVVRF 360
	DDNSPSFTIQI IRSVAKKHPK TWVHYIAAEE EDWDYAPLVL APDDRSYKSQ YLNNGPQRIG 420
	RKYKKVRFMA YTDETFKTRE AIQHESGILG PLYGEVGD TLIIIFKNQAS RPYNIYPHGI 480
	TDVRPLYSRRL LPKGVKHLKD FPILPGEIFK YKWTVTVEDG PTKSDPRCLT RYYSSTFVNME 540
	RDLASGLIGP LLICYKESVD QRGNQIMSDK RNVLFSVFD ENRSWYL TENI QRFLPNPAG 600
	VQLEDPEFQA SNIMHSINGY VFDLQLSVCL LHEVAYWYIL SIGAQTDFLS VFFSGYTFFKH 660
	KMVYEDTLTL FPFSGETVFM SMENPGLWIL GCHNSDFRNR GMTALLKVSS CDKNTGDYVE 720
	DSYEDISAYL LSKNNAIEPR SFSQNSRHPS TRQKQFNATT IPENDIEKTD PWFHRTFMP 780
	KIQNVSSSDL LMLLRQSPTP HGLSLSLDQE AKYETFSDDP SPGAIDSNN LSEMTFRFPQ 840
	LHSGDMVFT PESGLQLRLN EKLGTAAATE LKKLDFKVSS TSNNLISTIP SDNLAAGTDN 900
	TSSLGPPSMP VHYDSQLDTT LFGKKSSPLT ESGGPLSLSE ENNDSKLLS GLMNSQESSW 960
	GKNVSSSTESG RLFGKGRAGH PALLTKDNAL FKVSISLLKT NKTSNNSATN RKTHIDGP SL 1020

	LIENSPSPVWQ NILESSTFEK KVTPLIHDRM LMDKNATALR LNHSMSNKTS SKNMEMVQQK 1080
	KEGPIPPDAQ NPDMSFFKML FLPEASARWIQ RTHGKNLSNS GQGPSPKQIV SLGPEKSVEG 1140
	QNFLSEKNKV VVGKGFTKD VGLKEMVFPs SRNLFLTNLD NLHENNTHNQ EKKIQEEIEK 1200
	KETLIQENVV LPQIHTVTGT KNFMKNLFLL STRQNVESY DGAYAPVLQD FRSLNDSTNR 1260
	TKKHTAHFSK KGEEENLEGL GNQTKQIVEK YACTTRISP N TSQQNFVTQR SKRALKQFRL 1320
	PLEETELEKR IIVDDTSTQW SKNMKHLTPS TLTQIDYNEK EKGAITQSPL SDCLTRSHSI 1380
	PQANRSPLPI AKVSSFPSIR PIYLTRVLQF DNSSHLPAAS YRKKDSGVQE SSHFLQGAKK 1440
	NNLSLAILTL EMTGDQREVG SLGTSATNSV TYKKVENTVL PKPDLPKTSG KVELLPKVHI 1500
	YQKDLFPET SNGSPGHLDL VEGSLLQGT GAIKWNEANR PGKVPFLRVA TESSAKTPSK 1560
	LLDPLAWDNH YGTQIPKEEW KSQEKSPKT AFKKKDTILS LNACESNHAI AAINEGQNK 1620
	EIEVWAKQG RTERLCSQNP PVLKRHRQEI TRTTLQSDQE EIDYDDTISV EMKKEDFDIY 1680
	DEDENQSPRS FQKTRHYFI AAVEERLWDYG MSSSPHVLNR RAQSGSVQPF KKVVFQEFDT 1740
	GSFTQPLYRG ELNEHLGLLG PYIRAEVEDN IMVTFRNQAS RPYSFYSSLI SYEEDQRQGA 1800
	EPKRNFKVKN ETPTYFWKVQ HHMAPTKDEF DCKAWYFSD VDLEKDVHSG LIGPLLVCHT 1860
	NTLNPAHGRQ VTVQEFALFF TIFDETKSWY FTEMNERNCR APCNIQMEDP TFKENYRFHA 1920
	INGYIMDTLP GLVMAQDQRI RWYLLSMGSN ENIHSIHFSG HVFTVRKKEE YKMALYNLYP 1980
	GVFETVEMLP SKAGIWRVEC LIGELHLAGM STLFLVYSNK CQTPLGMASG HIRDFQITAS 2040
	GQYQWAPKL ARLHYSGSIN AWSTKEPFSW IKVDLLAPMI IHGIKTQGAR QKFSSLYISQ 2100
	FIIMYSLDGK KWQTYRGNT GTLMVFFGNV DSSGIKHNI NPPIIARYIR LHPHYHSIR 2160
	TLRMELMGCD LNSCSMPLGM ESKATSDAQI TASSYFTNMF ATWSPSKARL HLQGRSNAWR 2220
	PQVNNPKEWL QVDFQTKMV TGVTTQGVKS LLTSMYVKEF LISSSQDGHQ WTLFFQNGKV 2280
	KVFQGNQDSF TPVNSLDPP LLTRYLRHP QSWVHQIALR MEVLGCEAQD LY 2332
AE288	GTSESATPES GPGSEPATSG SETPGTSESA TPESGPGSEP ATSGSETPGT SESATPESGP 60
SEQ ID NO: 9	GTSTEPSEGS APGSPAGSPT STEEGTSESA TPESGPGSEP ATSGSETPGT SESATPESGP 120
	GSPAGSPTST EEGSPAGSPT STEEGTSTEP SEGSA PGTS SATPESGPAT SESATPESGP 180
	GTSESATPES GPGSEPATSG SETPGSEPAT SGSETPGSPA GSPTSTEEGT STEPSEGSAP 240
	GTSTEPSEGS APGSEPATSG SETPGTSESA TPESGPGTST EPSEGSAP 288
pSYN VWF059	TSTEEGASIS DKNTGDYYED SYEDISAYLL SKNNAIEPRS FSDKTH
SEQ ID NO: 10	
pSYN VWF059A	TSTEEGASSD KNTGDYYEDS YEDISAYLLS KNNNAIEPRSF SDKTH
SEQ ID NO: 11	
pSYN FVIII 312	ATRRYYLGAV ELSWDYMQSD LGELPVDARF PPRVPKSFPE NTSVVYKKT L FVEFTDHLFN 60
SEQ ID NO: 12	IAKPRPPWVG LLGPTIQAEV YDTVVITLKN MASHPVSLHA VGVSYWKASE GA EYDDQTSQ 120
	REKEDDKVFP GGSHTYVWQV LKENGPMASD PLCLTYSYLS HVDLVKDLNS GLIGALLVCR 180
	EGSLAKEKTO TLHKFILLFA VFDEGKSWHS ETKNLSMQDR DAASARAWPK MHTVNGVNR 240
	SLPGLIGCHR KSVYWHVIGM GTTPEVHSIF LEGHTFLVRN HRQASLEISP ITFLTAQTL 300
	MDLGQFLLFC HISSHQHDGM EAYVKVDSCP EEPQLRMKN EEAEDYDDDL TDSEMDVVR 360
	DDNSPSFIQ IRSVAKKHPK TWVHYIAAEE EDWDYAPLVL APDDRSYKSQ YLNNGPQRIG 420
	RKYKKVRFMA YTDETFKTRE AIQHESGILG PLYGEVGD LLIIFKNQAS RPYNIYPHGI 480
	TDVRPLYRR LPKGVKHLKD FPILPGEIFK YKWTVTVEDG PTKSDPRCLT RYSSSFVNME 540
	RDLASGLIGP LLICYKESVD QRGNQIMSDK RNVLFSVFD ENRSWYLTEN IQRFPLNPAG 600
	VQLEDPEFQA SNIMHSINGY VFDSLQLSVC LHEVAYWYIL SIGAQTDFLS VFFSGYTFKH 660
	KMVEDTLTL FPFSGETVFM SMENPGLWIL GCHNSDFRNR GMTALLKVSS CDKNTGDYYE 720
	DSYEDISAYL LSKNNAIEPR SFSQNGTSES ATPESGPGSE PATSGSETPG TSESATPESG 780
	PGSEPATSGS ETPGTSESAT PESGPGTSTE PSEGSAPGSP AGSPTSTEEG TSESATPESG 840
	PGSEPATSGS ETPGTSESAT PESGPGSPAG SPTSTEEGSP AGSPTSTEEG TSTEPSEGS 900
	PGTSESATPE SGPGTSESAT PESGPGTSES ATPESGPGSE PATSGSETPG SEPATSGSET 960
	PGSPAGSPTS TEEGTSTEPS EGSAPGTSTE PSEGSAPGSE PATSGSETPG TSESATPESG 1020
	PGTSTEPSEG SAPASSEITR TTLQSDQEEI DYDDTISVEM KKEDFDIYDE DENQSPRSFQ 1080
	KKTRHYFIAA VERLWDYGMS SSPHVLNRRA QSGSVQFQK VVFQEFDTGS RQPLVYRGEL 1140
	NEHLGLLGPY IRAEVEDNIM VTFRNQASRP YSFYSSLI SYEEDQRQGAEP RKNFVKPNET 1200
	KTYFWKVQHH MAPTKDEFDC KAWAYFSDVD LEKDVHSGLI GPLLVCHTNT LNPAGHGRQVT 1260
	VQEFALFFT I FDETKSWYFT ENMERNCRAP CNIQMEDPTF KENYRFHAIN GYIMDTLPGL 1320
	VMAQDQRIRW YLLSMGSNEN IHSIHFSGHV FTVRKKEEYK MALYNLYPGV FETVEMLP 1380

	AGIWRVECLI GEHLHAGMST LFLVYSNKCQ TPLGMASGHI RQFQITASGQ YGQWAPKLAR 1440 LHYSGSINAW STKEPFSWIK VDLLAPMIH GIKTQGARQK FSSLYISQFI IMYSLDGKKW 1500 QTYRGNSTGT LMVFFGNVDS SGIKHNIFNP PIARYIRLH PTHYSIRSTL RMELMGCDLN 1560 SCSMPLGMES KAISDAQITA SSYFTNMFAT WSPSKARLHL QGRSNAWRPQ VNNPKEWLQV 1620 DFQKTMKVTG VTTQGVKSLL TSMYVKEFLI SSSQDGHQWT LFFQNGKVKV FQGNQDSFTP 1680 VVNSLDPPLL TRYLRHPQS WVHQIALRME VLGCEAQDLY DKHTCPCPCP APELLGGPSV 1740 FLFPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PREEQYNSTY 1800 RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPFSRDELTK 1860 NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD SGSFFLYSKL TVDKSRWQQG 1920 NVFSCSVME ALHNHYTQKS LSLSPGK
ELNN_AE288_2 SEQ ID NO: 13	GSPAGSPTST EEGTSESATP ESGPGSEPAT SGSETPGTSE SATPESGPGT STEPSEGSAP 60 GTSTEPSEGS APGTSTEPSE GSAPGTSTEP SEGSAPTST EPSEGSAPGT STEPSEGSAP 120 GSPAGSPTST EEGTSTEPSE GSAPGTSESA TPESGPGSEP ATSGSETPGT SESATPESGP 180 GSEPATSGSE TPGTSESATP ESGPGTSTEP SEGSAPTSE SATPESGPGS PAGSPTSTEE 240 GSPAGSPTST EEGSPAGSPT STEEGTSESA TPESGPGTST EPSEGSAP 288
ELNN AE144_5A SEQ ID NO: 14	TSESATPESG PGSEPATSGS ETPGTSESAT PESGPGSEPA TSGSETPGTS ESATPESGPG 60 TSTEPSEGS SA PGSPAGSPTS TEEGTSESAT PESGPGSEPA TSGSETPGTS ESATPESGPG 120 SPAGSPTSTE EGSPAGSPTS TEEG
a2 Linker of chimeric protein SEQ ID NO: 15	DKNTGDYIED SYEDISAYLL SKNNAIEPRS FS 32
Signal Peptide of FVIII SEQ ID NO: 16	MQIELSTCFFLCLLRFCFS
FVIII fragment 1 of chimeric protein SEQ ID NO: 17	ATRRYYLGAV ELSWDYMQSD LGELPVDARF PPRVPKSFPP NTSVVYKKTLL FVEFTDHLFN 60 IAKPRPPWMG LLGPTTQAEV YDTVVITLKN MASHPVSLHA VGVSYWKASE GAIEDDQTSQ 120 REKEDDKVFP GGSHTYVWQV LKENGPMASD PLCLTYSYLS HVDLVKDLNS GLIGALLVCR 180 EGSLAKEKTQ TLHKFILLFA VFDEGKSWHS ETKNLSMQDR DAASARAWPK MHTVNGYVNR 240 SLPGLIGCHR KSVYWHVIGM GTTPEVHSIF LEGHTFLVRN HRQASLEISP ITFLTAQTLL 300 MDLGQFLLFC HISSHQHDMG EAYVKVDSCEP EEPQLRMKN EEAEDYDDDL TDSEMDVVRV 360 DDDNPSFSIQ IRSVAKKHPK TWVHYIAAEE EDWDYAPLVL APDDRSYKSK YLNNGPQRIG 420 RKYKKVRFMA YTDETFKTRE AIQHESGILG PLYGEVGDV LLIIFKNQAS RPYNIYPHGI 480 TDVVRPLYSR LKPGVKHLKD FPILPGEIFK YKWTVTVEDG PTKSDPRCLT RYYSFVNME 540 RDLASGLIGP LLICYKESVD QRGNQIMSDK RNVILFSVFD ENRSWYLTEN IQRFLPNPAG 600 VQLEDPEFQA SNIMHSINGY VFDSLQLSVC LHEVAYWYIL SIGAQTDFLS VFFSGYTFKH 660 KMYEDTLTL PFSGETVFM SMENPGLWIL GCHNSDFRNR GMTALLKVSS CDKNTGDYIE 720 DSYEDISAYL LSKNNAIEPR SFSQN
FVIII fragment 2 of chimeric protein SEQ ID NO: 18	EITRTTLQSD QEEIDYDDTI SVEMKKEDFD IYDEDENQSP RSFQKKTRHY FIAAVERLWD 60 YGMSSSPHVL RNRAQSGSVP QFKKVVVFQEF TDGSFTQPLY RGELNEHLGL LGPYIRAEVE 120 DNIMVTFRNQ ASRPYSFYSS LISYEEDQRO GAEPKRNFKV PNETKTYFWK VQHMMAPT KD 180 EFDCKAWAYF SDVDLEKDVH SGLIGPLLVLC HTNTLNPAHG RQVTVQEFAL FFTIFDETKS 240 WYFTENMERN CRAPCNIQME DPTFKENYRF HAINGYIMDT LPGLVMAQDQ RIRWYLLSMG 300 SNENIHSIH FSGHVFTVRKK EBYKMALYNL YPGVFETVEM LPSKAGIWRV ECLIGEHLHA 360 GMSTLFLVYS NKQCTPLGMA SGHIRDFQIT ASGQYGQWAP KLARLHYSGS INAWSTKEPF 420 SWIKVDLLAP MIHGIKTQG ARQKFSSLYI SQFIIMYSLD GKKWQTYRGN STGILMVFFG 480 NVDSSGIKHN IFNPPIARY IRLHPTHYSI RSTLRMELMG CDLNSCSMPL GMESKAISDA 540 QITASSYFTN MFATWSPSKA RLHLQGRSNA WRPQVNNPKE WLQVDFQKTM KVTGVTTQGV 600 KSLLTSMYVK EFLISSSQDG HQWTLFFQNG KVKVFQGNQD SFTPVVNSLD PPLTRYLRI 660 HPQSWVHQIA LRMEVIGCEA QDLY

VWF Signal Peptide	MIPARFAGVL LALALILPGT LC						
SEQ ID NO: 19							
VWF D1D2 domain of chimeric protein	AEGTRGRSST	ARCSLFGSDF	VNTFDGSMYS	FAGYCSYLLA	GGCQKRSFSI	IGDFQNGKRV	60
	SLSVYLGEFF	DIHLFVNGTV	TQGDQRVSM	YASKGLYLET	EAGYYKLSGE	AYGFVARIDG	120
	SGNFQVLLSD	RYFNKTCGLC	GNFNIFAEDD	FMTQEGTLTS	DPYDFANSWA	LSSGEQWCER	180
SEQ ID NO: 20	ASPPSSSCNI	SSGEMQKGLW	EQCQLLKSTS	VFARCHPLVD	PEPFFVALCEK	TLCECAGGLE	240
	CACPALLEYA	RTCAQEGMVL	YGWTDHSACS	PVCPAGMEYR	QCVSPCARTC	QSLHINEMCQ	300
	ERCVDGCSCP	EGQLLDEGLC	VESTECPCVH	SGKRYPPGTS	LSRDCNTCIC	RNSQWICSNE	360
	ECPGECCLVTG	QSHFKSFDNR	YFTFSGICQY	LLARDCQDHS	FSIVIETVQC	ADDRDAVCTR	420
	SVTVRLPGLH	NSLVKLKHGA	GVAMDGQDIQ	LPPLKGDRLI	QHTVTASVRL	SYGEDLQMDW	480
	DGRGRLLVKL	SPVYAGKTCG	LCGNYNNGQG	DDFLTPSGLA	EPRVEDFGNA	WKLHGDCQDL	540
	QKQHSDPICAL	NPRMTRFSEE	ACAVLTSPTF	EACHRAVSPL	PYLRNCRYDV	CSCSDGRECL	600
	CGALASYAAA	CAGRGVRVAW	REPGRCELNC	PKGQVYLQCG	TPCNLTCSRSL	SYPDEECNEA	660
	CLEGCFCPPG	LYMDERGDV	PKAQCPYYD	GEIFQPEDIF	SDHHTMCYCE	DGMHCTMSG	720
	VPGSLLPDAV	LSSPLSHRSK	R				
VWF D' domain of chimeric protein	SLSCRPPMVK	LVCPADNLRA	EGLECTKTICQ	NYDLECMSMG	CVSGCLCPPG	MVRHENRCVA	60
	LERPCPFHQG	KEYAPGETVK	TGCNTCVCRD	RKWNCTDHVC	DAT		103
SEQ ID NO: 21							
VWF D3 domain of chimeric protein	CSTIGMAHYL	TFDGLKYLFP	GECQYVLVQD	YCGSNPGTFR	ILVGNKGCSH	PSVKCKKRV	60
	ILVEGGEIEL	FDGEVNVKRP	MKDETHFEVV	ESGRYIILL	GKALSVVWDR	HLSISVVLKQ	120
	TYQEKVCGLC	GNFDGIQNN	LTSSNLQVEE	DPVDFGNSWK	VSSQCADTRK	VPLDSSPATC	180
SEQ ID NO: 22	HNNIMQTMV	DSSCRILTSD	VFQDCNKLVD	PEPYLDVCIY	DTCSCEISGD	CAAFCDTIAA	240
	YAHVCAQH GK	VVTWRTATLC	PQSCEERNLR	ENGYEA EWRY	NSCAPACQVT	CQHPPELACP	300
	VQCVEGCHAH	CPPGKILDEL	LQTCVDPEDC	PVCEVAGRRF	ASGKKVTLNP	SDPEHCQICH	360
	CDVNNLTCEA	CQEP					374
Fc region	DKTHTCPPCP	APPELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
	GVEVHNAKTK	PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
SEQ ID NO: 23	GQPREPQVYT	LPPSRDELTK	NQVSLTCLVK	GFYPSTDAVE	WESNGQPENN	YKTTTPVLDS	180
	DGSFFLYSKL	TVDKSRWQQG	NVFSCSVMHE	ALHNYHTQKS	LSLSPG		
ELNN AE288_3	GTSESATPES	GGPSEPATSG	SETPGTSESA	TPESGPGSEP	ATSGSETPGT	SESATPESGP	60
	GTSTEPSEGS	APGSPAGSPT	STEEGTSESA	TPESGPGSEP	ATSGSETPGT	SESATPESGP	120
SEQ ID NO: 24	GSPAGSPTST	EEGSPAGSPT	STEEGTSTEP	SEGSAPGTSE	SATPESGPGT	SESATPESGP	180
	GTSESATPES	GGPSEPATSG	SETPGSEPAT	SGSETPGSPA	GSPTSTEEGT	STEPSEGSAP	240
	GTSTEPSEGS	APGSEPATSG	SETPGTSESA	TPESGPGTST	EPSEGSAPAS	S	291

CLAIMS

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising:
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) sucrose;
 - (c) histidine;
 - (d) arginine;
 - (e) calcium chloride; and
 - (f) a polysorbate.
2. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises about 5% (w/v) to about 7.5% (w/v) sucrose.
3. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises about 1% (w/v) to about 4% (w/v) sucrose.
4. The pharmaceutical composition of any one of claims 1 to 3, wherein the pharmaceutical composition comprises about 5 mM to about 15 mM histidine.
5. The pharmaceutical composition of any one of claims 1-4, wherein the pharmaceutical composition comprises about 200 mM to about 300 mM arginine.
6. The pharmaceutical composition of any one of claims 1-5, wherein the pharmaceutical composition comprises about 2.5 mM to about 10 mM calcium chloride.
7. The pharmaceutical composition of any one of claims 1-6, wherein the polysorbate is polysorbate 20.

8. The pharmaceutical composition of any one of claims 1- 7, wherein the polysorbate is polysorbate 80.
9. The pharmaceutical composition of any one of claims 1-8, wherein the pharmaceutical composition comprises about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.
10. A pharmaceutical composition comprising:
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) about 1% (w/v) to about 4% (w/v) sucrose;
 - (c) about 5 mM to about 15 mM histidine;
 - (d) about 200 mM to about 300 mM arginine;
 - (e) about 2.5 mM to about 10 mM calcium chloride; and
 - (f) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.
11. The pharmaceutical composition of claim 10, wherein the polysorbate is polysorbate 20.
12. The pharmaceutical composition of claim 10 or claim 11, wherein the polysorbate is polysorbate 80.
13. The pharmaceutical composition of any one of claims 10-12, wherein the pharmaceutical composition comprises:
 - (a) about 2% (w/v) sucrose;
 - (b) about 10 mM histidine;
 - (c) about 250 mM arginine;
 - (d) about 5 mM calcium chloride; and
 - (e) about 0.05% polysorbate 20 or polysorbate 80.

14. The pharmaceutical composition of any one of claims 10-13, wherein the pharmaceutical composition comprises:

- (a) about 2% (w/v) sucrose;
- (b) about 10 mM L-histidine;
- (c) about 250 mM L-arginine-HCl;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 80.

15. The pharmaceutical composition of any one of claims 10-12, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.4 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

16. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

17. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

18. The pharmaceutical composition of any one of claims 15-17, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

19. The pharmaceutical composition of any one of claims 15-18, wherein the pharmaceutical composition comprises:

- (a) about 20 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 52.7 mg/mL L-arginine-HCl;
- (d) about 0.7 mg/mL calcium chloride dihydrate; and
- (e) about 0.5 mg/mL polysorbate 80.

20. The pharmaceutical composition of any one of claims 15-19, wherein the pharmaceutical composition comprises:

- (a) about 20 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 52.7 mg/mL L-arginine-HCl;
- (d) about 0.6 mg/mL calcium chloride; and
- (e) about 0.5 mg/mL polysorbate 80.

21. The pharmaceutical composition of any one of claims 15-20, wherein the pharmaceutical composition comprises:

- (a) about 20 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 43.6 mg/mL L-arginine;
- (d) about 0.7 mg/mL calcium chloride dihydrate; and

- (e) about 0.5 mg/mL polysorbate 80.

22. The pharmaceutical composition of any one of claims 15-21, wherein the pharmaceutical composition comprises:

- (a) about 20 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 43.6 mg/mL L-arginine;
- (d) about 0.6 mg/mL calcium chloride; and
- (e) about 0.5 mg/mL polysorbate 80.

23. The pharmaceutical composition of any one of claims 15-22, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

24. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.4 mg/mL to 0.8 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

25. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 60 mg/mL L-arginine;

- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

26. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 10 mg/mL to 40 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 60 mg/mL L-arginine;
- (d) 0.4 mg/mL to 0.7 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

27. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

28. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

29. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;

- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

30. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition comprises:

- (a) 22.45 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

31. The pharmaceutical composition of any one of claims 1 to 30, wherein the pharmaceutical composition has a pH of about 6.5 to about 7.5.

32. The pharmaceutical composition of any one of claims 1 to 31, wherein the pharmaceutical composition has a pH of about 7.0.

33. The pharmaceutical composition of any one of claims 1 to 31, wherein the pharmaceutical composition has a pH of about 6.8.

34. The pharmaceutical composition of any one of claims 1 to 33, wherein the pharmaceutical composition comprises less than 8.8 mg/mL sodium chloride (NaCl).

35. The pharmaceutical composition of any one of claims 1 to 34, wherein the pharmaceutical composition does not comprise NaCl.

36. The pharmaceutical composition of any one of claims 1 to 13, wherein the histidine is L-histidine.

37. The pharmaceutical composition of any one of claims 1 to 13, wherein the arginine is L-arginine.

38. The pharmaceutical composition of any one of claims 1 to 13, which comprises arginine-HCl.
39. The pharmaceutical composition of any one of claims 1 to 13, which comprises L-arginine-HCl.
40. The pharmaceutical composition of any one of claims 1 to 15, which comprises calcium chloride dihydrate.
41. The pharmaceutical composition of any one of claims 1 to 40, wherein the pharmaceutical composition has a chimeric protein concentration of about 0.8 to about 1.2 mg/mL.
42. The pharmaceutical composition of any one of claims 1 to 41, wherein the pharmaceutical composition comprises 75 IU/mL to 2,000 IU/mL of the chimeric protein.
43. The pharmaceutical composition of any one of claims 1 to 42, wherein the pharmaceutical composition has an osmolality about 525 to about 725 mOsm/kg.
44. The pharmaceutical composition of any one of claims 1 to 43, wherein the pharmaceutical composition comprises has an osmolality about 600 to about 650 mOsm/kg.
45. The pharmaceutical composition of any one of claims 1 to 44, wherein the pharmaceutical composition comprises has a turbidity of less than about 7 Nephelometric Turbidity Units (NTU).
46. The pharmaceutical composition of any one of claims 1 to 45, wherein the first polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 1 and the second polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.
47. The pharmaceutical composition of any one of claims 1 to 46, wherein the chimeric protein is efanesoctocog alfa.
48. The pharmaceutical composition of any one of claims 1 to 41 or 43 to 47, which comprises about 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein.

49. A method of treating hemophilia A in a subject in need thereof, comprising administering to the subject an effective amount of the pharmaceutical composition of any one of claims 1 to 48.
50. The method of claim 49, wherein the pharmaceutical composition is self-administered.
51. The method of claim 49 or 50, wherein the pharmaceutical composition is administered intravenously.
52. The method of any one of claims 49 to 51, wherein the pharmaceutical composition is administered intravenously at a dose of 20 IU/kg to 70 IU/kg.
53. The method of any one of claims 49 to 52, wherein the pharmaceutical composition is administered intravenously at a dose of 50 IU/kg.
54. The method any one of claims 49 to 53, wherein the pharmaceutical composition is administered intravenously once every 7-10 days.
55. The method of claim any one of claims 49 to 54, wherein the pharmaceutical composition is administered intravenously once weekly.
56. A pharmaceutical kit comprising:
- (i) a first container comprising a lyophilized pharmaceutical composition comprising
 - (a) a chimeric protein comprising a first polypeptide chain which comprises a Factor VIII ("FVIII") protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;
 - (b) about 30 mg to about 135 mg sucrose;
 - (c) about 2.5 mg to about 7.5 mg histidine;
 - (d) about 140 mg to about 200 mg arginine;
 - (e) about 1.5 mg to about 5 mg calcium chloride; and
 - (f) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80, and
 - (ii) a second container comprising sterile water.

57. The pharmaceutical kit of claim 56, which does not comprise NaCl.
58. The pharmaceutical kit of claim 56 or 57, wherein the histidine is L-histidine.
59. The pharmaceutical kit any one of claims 56 to 58, wherein the arginine is L-arginine.
60. The pharmaceutical kit of claim any one of claims 56 to 59, which comprises arginine-HCl.
61. The pharmaceutical kit of claim any one of claims 56 to 60, which comprises L-arginine-HCl.
62. The pharmaceutical kit of claim any one of claims 56 to 61, which comprises calcium chloride dihydrate.
63. The pharmaceutical kit of claim any one of claims 56 to 61, wherein the lyophilized pharmaceutical composition comprises:
- (a) about 67.3 mg sucrose;
 - (b) about 5.2 mg L-histidine;
 - (c) about 177.3 mg L-arginine-HCl;
 - (d) about 2.5 mg calcium chloride; and
 - (e) about 1.7 mg polysorbate 20 or polysorbate 80.
64. The pharmaceutical kit of any one of claims 56 to 61, wherein the lyophilized pharmaceutical composition comprises:
- (a) about 67.3 mg sucrose;
 - (b) about 5.2 mg L-histidine;
 - (c) about 146.6 mg L-arginine-HCl;
 - (d) about 2.5 mg calcium chloride; and
 - (e) about 1.7 mg polysorbate 20 or polysorbate 80.
65. The pharmaceutical kit of any one of claims 56 to 64, wherein the lyophilized pharmaceutical composition has a moisture content of less than 2%.
66. The pharmaceutical kit of any one of claims 56 to 65, wherein the lyophilized pharmaceutical composition has a moisture content of less than 1.8%.

67. The pharmaceutical kit of any one of claims 56 to 66, wherein the lyophilized pharmaceutical composition has a moisture content of less than 1.6%.
68. The pharmaceutical kit of any one of claims 56 to 67, wherein the lyophilized pharmaceutical composition is in a lyophilized cake.
69. The pharmaceutical kit of claim 68, wherein the lyophilized cake is white.
70. The pharmaceutical kit of claim 68 or 69, wherein the lyophilized cake is less than Y4 in the European Pharmacopoeia color scale.
71. The pharmaceutical kit of claim any one of claims 56 to 70, wherein the first container comprises 100 IU to 10,000 IU of the chimeric protein.
72. The pharmaceutical kit of any one of claims 56 to 71, wherein the first container comprises 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein.
73. The pharmaceutical kit of any one of claims 56 to 72, further comprising instructions for combining the lyophilized pharmaceutical composition and the sterile water.
74. The pharmaceutical kit of any one of claims 56 to 73, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the lyophilized pharmaceutical composition is reconstituted within 7 to 12 seconds.
75. The pharmaceutical kit of any one of claims 56 to 74, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 525 to about 725 mOsm/kg.
76. The pharmaceutical kit of any one of claims 56 to 75, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 600 to about 650 mOsm/kg.
77. The pharmaceutical kit of claim any one of claims 56 to 76, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.5 to about 7.5.

78. The pharmaceutical kit of any one of claims 56 to 77, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.0.
79. The pharmaceutical kit of any one of claims 56 to 78, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 0.8 to about 1.2 mg/mL.
80. The pharmaceutical kit of claim 79, wherein less than 3% of the protein is aggregated.
81. The pharmaceutical kit of any one of claims 56 to 80, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the turbidity of the resulting solution is less than about 7 Nephelometric Turbidity Units (NTU).
82. The pharmaceutical kit of any one of claims 56 to 81, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
 - (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.
83. The pharmaceutical kit of any one of claims 56 to 81, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:
- (a) 10 mg/mL to 40 mg/mL sucrose;
 - (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
 - (c) 40 mg/mL to 60 mg/mL L-arginine;
 - (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
 - (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

84. The pharmaceutical kit of any one of claims 56 to 81, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:
- (a) 22.45 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 59.11 mg/ml L-arginine-HCl;
 - (d) 0.62 mg/ml calcium chloride; and
 - (e) 0.56 mg/ml polysorbate 80.
85. The pharmaceutical kit of any one of claims 56 to 81, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:
- (a) 22.45 mg/ml sucrose;
 - (b) 1.74 mg/ml L-histidine;
 - (c) 48.88 mg/ml L-arginine;
 - (d) 0.62 mg/ml calcium chloride; and
 - (e) 0.56 mg/ml polysorbate 80.
86. The pharmaceutical kit of any one of claims 56 to 85, wherein the second container comprises about 2 mL to about 5 mL of the sterile water.
87. The pharmaceutical kit of any one of claims 56 to 86, wherein the second container comprises about 3 mL of the sterile water.
88. The pharmaceutical kit of any one of claims 56 to 86, wherein the second container comprises about 3.3 mL of the sterile water.
89. The pharmaceutical kit of any one of claims 56 to 88, wherein the first container is a glass vial comprising a rubber stopper.
90. The pharmaceutical kit of claim any one of claims 56 to 89, wherein the second container is a syringe body.

91. The pharmaceutical kit of claim 90, wherein the sterile water is in the syringe body.
92. The pharmaceutical kit of claim 90 or 91, wherein the syringe body is associated with a plunger.
93. The pharmaceutical kit of any one of claims 89 to 92, further comprising an adaptor to connect the glass vial to the syringe body.
94. The pharmaceutical kit of any one of claims 89 to 93, further comprising infusion tubing associated with a needle to be connected to the syringe body, suitable for intravenous infusion.
95. The pharmaceutical kit of any one of claims 56 to 94, wherein the first polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 1 and the second polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.
96. A method of treating hemophilia A in a subject in need thereof, comprising combining the lyophilized pharmaceutical composition and the sterile water of the kit of any one of claims 56 to 95, and administering to the subject an effective amount of the resulting combination.
97. The method of claim 96, wherein the subject combines the lyophilized pharmaceutical composition and the sterile water of the kit.
98. The method of claim 96 or 97, wherein the combination is self-administered by the subject.
99. The pharmaceutical composition of any one of claims 1-9, wherein the pharmaceutical composition comprises:
- (a) about 5% (w/v) to about 7.5% (w/v) sucrose;
 - (b) about 5 mM to about 15 mM histidine;
 - (c) about 200 mM to about 300 mM arginine;
 - (d) about 2.5 mM to about 10 mM calcium chloride; and
 - (e) about 0.008% (w/v) to about 0.1% (w/v) polysorbate 20 or polysorbate 80.

100. The pharmaceutical composition of any one of claims 1-9 or 99, wherein the pharmaceutical composition comprises:

- (a) about 5% (w/v) sucrose;
- (b) about 10 mM histidine;
- (c) about 250 mM arginine;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 20 or polysorbate 80.

101. The pharmaceutical composition of any one of claims 1-9 or 99-100, wherein the pharmaceutical composition comprises:

- (a) about 5% (w/v) sucrose;
- (b) about 10 mM L-histidine;
- (c) about 250 mM L-arginine-HCl;
- (d) about 5 mM calcium chloride; and
- (e) about 0.05% polysorbate 80.

102. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.4 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

103. The pharmaceutical composition of claim 102, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

104. The pharmaceutical composition of any one of claims 102-103, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

105. The pharmaceutical composition of any one of claims 102-104, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

106. The pharmaceutical composition of any one of claims 102-105, wherein the pharmaceutical composition comprises:

- (a) about 50 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 52.7 mg/mL L-arginine-HCl;
- (d) about 0.7 mg/mL calcium chloride dihydrate; and
- (e) about 0.5 mg/mL polysorbate 80.

107. The pharmaceutical composition of any one of claims 102-106, wherein the pharmaceutical composition comprises:

- (a) about 50 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 52.7 mg/mL L-arginine-HCl;
- (d) about 0.6 mg/mL calcium chloride; and

- (e) about 0.5 mg/mL polysorbate 80.

108. The pharmaceutical composition of any one of claims 102-107, wherein the pharmaceutical composition comprises:

- (a) about 50 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 43.6 mg/mL L-arginine;
- (d) about 0.7 mg/mL calcium chloride dihydrate; and
- (e) about 0.5 mg/mL polysorbate 80.

109. The pharmaceutical composition of any one of claims 102-108, wherein the pharmaceutical composition comprises:

- (a) about 50 mg/mL sucrose;
- (b) about 1.6 mg/mL L-histidine;
- (c) about 43.6 mg/mL L-arginine;
- (d) about 0.6 mg/mL calcium chloride; and
- (e) about 0.5 mg/mL polysorbate 80.

110. The pharmaceutical composition of any one of claims 102-109, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

111. The pharmaceutical composition of any one of claims 102-110, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 50 mg/mL to 70 mg/mL L-arginine-HCl;

- (d) 0.4 mg/mL to 0.8 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

112. The pharmaceutical composition of any one of claims 102-111, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 60 mg/mL L-arginine;
- (d) 0.7 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

113. The pharmaceutical composition of any one of claims 102-112, wherein the pharmaceutical composition comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 60 mg/mL L-arginine;
- (d) 0.4 mg/mL to 0.7 mg/mL calcium chloride dihydrate; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

114. The pharmaceutical composition of any one of claims 102-113, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

115. The pharmaceutical composition of any one of claims 102-114, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;

- (c) 48.88 mg/ml L-arginine;
- (d) 0.82 mg/ml calcium chloride dihydrate; and
- (e) 0.56 mg/ml polysorbate 80.

116. The pharmaceutical composition of any one of claims 102-115, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

117. The pharmaceutical composition of any one of claims 102-116, wherein the pharmaceutical composition comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

118. The pharmaceutical composition of any one of claims 99-117, wherein the pharmaceutical composition has a pH of about 6.5 to about 7.5.

119. The pharmaceutical composition of claim 118, wherein the pharmaceutical composition has a pH of about 7.0.

120. The pharmaceutical composition of claim 118, wherein the pharmaceutical composition has a pH of about 6.8.

121. The pharmaceutical composition of any one of claims 99-120, wherein the pharmaceutical composition comprises less than 8.8 mg/mL sodium chloride (NaCl).

122. The pharmaceutical composition any one of claims 99-121, which does not comprise NaCl.
123. The pharmaceutical composition of any one of claims 1-11, wherein the histidine is L-histidine.
124. The pharmaceutical composition of any one of claims 99-100, wherein the arginine is L-arginine.
125. The pharmaceutical composition of any one of claims 99-100, which comprises arginine-HCl.
126. The pharmaceutical composition of any one of claims 99-100, which comprises L-arginine-HCl.
127. The pharmaceutical composition of any one of claims 99-102, which comprises calcium chloride dihydrate.
128. The pharmaceutical composition of any one of claims 99-127, wherein the pharmaceutical composition has a chimeric protein concentration of about 0.8 to about 1.2 mg/mL.
129. The pharmaceutical composition of any one of claims 99-127, wherein the pharmaceutical composition comprises 75 IU/mL to 2,000 IU/mL of the chimeric protein.
130. The pharmaceutical composition of any one of claims 99-129, wherein the pharmaceutical composition has an osmolality about 525 to about 725 mOsm/kg.
131. The pharmaceutical composition of claim 130, wherein the pharmaceutical composition comprises has an osmolality about 600 to about 650 mOsm/kg.
132. The pharmaceutical composition of any one of claims 99-131, wherein the pharmaceutical composition comprises has a turbidity of less than about 7 Nephelometric Turbidity Units.
133. The pharmaceutical composition of any one of claims 99-132, wherein the first polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 1 and the second polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide

chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.

134. The pharmaceutical composition of any one of claims 99-133, wherein the chimeric protein is efanesoctocog alfa.

135. The pharmaceutical composition of any one of claims 99-127 or 130-134, which comprises about 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein.

136. A method of treating hemophilia A in a subject in need thereof, comprising administering to the subject an effective amount of the pharmaceutical composition of any one of claims 99 to 135.

137. The method of claim 136, wherein the pharmaceutical composition is self-administered.

138. The method of claim 136, wherein the pharmaceutical composition is administered intravenously.

139. The method of claim 137, wherein the pharmaceutical composition is administered intravenously.

140. The method of any one of claims 136-139, wherein the pharmaceutical composition is administered intravenously at a dose of 20 IU/kg to 70 IU/kg.

141. The method of any one of claims 136-139, wherein the pharmaceutical composition is administered intravenously at a dose of 50 IU/kg.

142. The method of any one of claims 136-140, wherein the pharmaceutical composition is administered intravenously once every 7-10 days.

143. The method of any one of claims 136-140, wherein the pharmaceutical composition is administered intravenously once weekly.

144. A pharmaceutical kit comprising:

- (i) a first container comprising a lyophilized pharmaceutical composition comprising
 - (a) a chimeric protein comprising

a first polypeptide chain which comprises a Factor VIII ("FVIII") protein or a portion thereof and a first immunoglobulin ("Ig") constant region or a portion thereof, and
a second polypeptide chain which comprises a von Willebrand Factor ("VWF") protein and a second Ig constant region or a portion thereof;

- (b) sucrose;
 - (c) histidine;
 - (d) arginine;
 - (e) calcium chloride; and
 - (f) polysorbate 20 or polysorbate 80, and
- (ii) a second container comprising sterile water.

145. The pharmaceutical kit of claim 144, wherein the lyophilized pharmaceutical composition comprises:

- (a) about 160 mg to about 200 mg sucrose;
- (b) about 2.5 mg to about 7.5 mg histidine;
- (c) about 140 mg to about 200 mg arginine;
- (d) about 1.5 mg to about 5 mg calcium chloride; and
- (e) about 1 mg to about 5 mg polysorbate 20 or polysorbate 80.

146. The pharmaceutical kit of any one of claims 144-145, which does not comprise NaCl.

147. The pharmaceutical kit of any one of claims 144-146, wherein the histidine is L-histidine.

148. The pharmaceutical kit of any one of claims 144-147, wherein the arginine is L-arginine.

149. The pharmaceutical kit of any one of claims 144-147, which comprises arginine-HCl.

150. The pharmaceutical kit of any one of claims 144-147, which comprises L-arginine-HCl.

151. The pharmaceutical kit of any one of claims 144-150, which comprises calcium chloride dihydrate.

152. The pharmaceutical kit of any one of claims 144-151, wherein the lyophilized pharmaceutical composition comprises:

- (a) about 168.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 177.3 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

153. The pharmaceutical kit of any one of claims 144-151, wherein the lyophilized pharmaceutical composition comprises:

- (a) about 168.3 mg sucrose;
- (b) about 5.2 mg L-histidine;
- (c) about 146.6 mg L-arginine-HCl;
- (d) about 2.5 mg calcium chloride; and
- (e) about 1.7 mg polysorbate 20 or polysorbate 80.

154. The pharmaceutical kit of any one of claims 144-153, wherein the lyophilized pharmaceutical composition has a moisture content of less than 2%.

155. The pharmaceutical kit of any one of claims 144-154, wherein the lyophilized pharmaceutical composition has a moisture content of less than 1.8%.

156. The pharmaceutical kit of any one of claims 144-155, wherein the lyophilized pharmaceutical composition has a moisture content of less than 1.6%.

157. The pharmaceutical kit of of any one of claims 144-156, wherein the lyophilized pharmaceutical composition is in a lyophilized cake.

158. The pharmaceutical kit of claim 157, wherein the lyophilized cake is white.

159. The pharmaceutical kit of claim 158, wherein the lyophilized cake is less than Y4 in the European Pharmacopoeia color scale.

160. The pharmaceutical kit of of any one of claims 144-159, wherein the first container comprises 100 IU to 10,000 IU of the chimeric protein.

161. The pharmaceutical kit of any one of claims 144-160, wherein the first container comprises 250 IU, 500 IU, 1000 IU, 2000 IU, 3000 IU, or 4,000 IU of the chimeric protein.
162. The pharmaceutical kit of any one of claims 144-161, further comprising instructions for combining the lyophilized pharmaceutical composition and the sterile water.
163. The pharmaceutical kit of any one of claims 144-162, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the lyophilized pharmaceutical composition is reconstituted within 7 to 12 seconds.
164. The pharmaceutical kit of any one of claims 144-163, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 525 to about 725 mOsm/kg.
165. The pharmaceutical kit of any one of claims 144-164, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the osmolality of the resulting solution is about 600 to about 650 mOsm/kg.
166. The pharmaceutical kit of any one of claims 144-165, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 6.5 to about 7.5.
167. The pharmaceutical kit of any one of claims 144-166, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the pH of the resulting solution is about 7.0.
168. The pharmaceutical kit of any one of claims 144-167, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the protein concentration of the resulting solution is about 0.8 to about 1.2 mg/mL.
169. The pharmaceutical kit of claim 168, wherein less than 3% of the protein is aggregated.
170. The pharmaceutical kit of any one of claims 144-169, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the turbidity of the resulting solution is less than about 7 Nephelometric Turbidity Units.

171. The pharmaceutical kit of claim 144, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 70 mg/mL L-arginine-HCl;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

172. The pharmaceutical kit of claim 171, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 45 mg/mL to 60 mg/mL sucrose;
- (b) 1.5 mg/mL to 2.0 mg/mL L-histidine;
- (c) 40 mg/mL to 60 mg/mL L-arginine;
- (d) 0.5 mg/mL to 0.9 mg/mL calcium chloride; and
- (e) 0.4 mg/mL to 0.7 mg/mL polysorbate 80.

173. The pharmaceutical kit of any one of claims 144 or 171-172, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 59.11 mg/ml L-arginine-HCl;
- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

174. The pharmaceutical kit of any one of claims 144 or 171-173, wherein when the lyophilized pharmaceutical composition and the sterile water are combined, then the resulting solution comprises:

- (a) 56.12 mg/ml sucrose;
- (b) 1.74 mg/ml L-histidine;
- (c) 48.88 mg/ml L-arginine;

- (d) 0.62 mg/ml calcium chloride; and
- (e) 0.56 mg/ml polysorbate 80.

175. The pharmaceutical kit of any one of claims 144-174, wherein the second container comprises about 2 mL to about 5 mL of the sterile water.

176. The pharmaceutical kit of any one of claims 144-175, wherein the second container comprises about 3 mL of the sterile water.

177. The pharmaceutical kit of any one of claims 144-175, wherein the second container comprises about 3.3 mL of the sterile water.

178. The pharmaceutical kit of any one of claims 144-177, wherein the first container is a glass vial comprising a rubber stopper.

179. The pharmaceutical kit of any one of claims 144-178, wherein the second container is a syringe body.

180. The pharmaceutical kit of claim 179, wherein the sterile water is in the syringe body.

181. The pharmaceutical kit of claim 179 or 180, wherein the syringe body is associated with a plunger.

182. The pharmaceutical kit of any one of claims 179-181, further comprising an adaptor to connect the glass vial to the syringe body.

183. The pharmaceutical kit of any one of claims 179-182, further comprising infusion tubing associated with a needle to be connected to the syringe body, suitable for intravenous infusion.

184. The pharmaceutical kit of any one of claims 144 to 183, wherein the first polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 1 and the second polypeptide chain comprises the amino acid sequence set forth as SEQ ID NO: 2, wherein the first polypeptide chain and the second polypeptide chain are covalently linked by two disulfide bonds between Fc domains in the first and second polypeptide chains.

185. A method of treating hemophilia A in a subject in need thereof, comprising combining the lyophilized pharmaceutical composition and the sterile water of the kit of any one of claims 144 to 184, and administering to the subject an effective amount of the resulting combination.
186. The method of claim 185, wherein the subject combines the lyophilized pharmaceutical composition and the sterile water of the kit.
187. The method of claim 185 or 186, wherein the combination is self-administered by the subject.
188. The pharmaceutical composition of any one of claims 1 to 34, wherein the pharmaceutical composition does not comprise NaOH.
189. The pharmaceutical composition of any one of claims 1 to 34, wherein the pharmaceutical composition does not comprise sodium ions.
190. The pharmaceutical kit of claim 56, which does not comprise NaOH.
191. The pharmaceutical kit of claim 56, which does not comprise sodium ions.
192. The pharmaceutical composition any one of claims 99-121, which does not comprise NaOH.
193. The pharmaceutical composition any one of claims 99-121, which does not comprise sodium ions.
194. The pharmaceutical kit of any one of claims 144-145, which does not comprise NaOH.
195. The pharmaceutical kit of any one of claims 144-145, which does not comprise sodium ions.