

PHOSPHO-INDOLES AS HIV INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application Nos. 60/611,061, filed September 16, 2005, 60/711,445 filed August 25, 2005 and 60/711,565, filed August 26, 2005, all entitled "Phospho-Indoles as HIV Inhibitors."

FIELD OF THE INVENTION

The present invention provides new human immunodeficiency virus (HIV) reverse transcriptase inhibitory compounds and their pharmaceutically acceptable salts, prodrugs, analogs and derivatives. Also included are methods of using these compounds for the prophylaxis and treatment of HIV infection and AIDS, and pharmaceutical compositions that contain the compounds.

BACKGROUND OF THE INVENTION


Numerous compounds have been synthesized to combat the human immunodeficiency virus (HIV) since it was discovered to be the etiological cause of the acquired immunodeficiency syndrome (AIDS) in 1983. A focal point of AIDS research efforts has been and continues to be the development of inhibitors of human immunodeficiency virus (HIV-1) reverse transcriptase, the enzyme responsible for the reverse transcription of the retroviral RNA to proviral DNA (W.C. Greene, New England Journal of Medicine (1991), 324:308-17; Mitsuya et al., Science (1990), 249:1533-44; E.J. DeClercq, Retrovirus (1992), 8:119-34). Inhibitors include non-nucleoside reverse transcriptase inhibitors or NNRTIs that bind to a specific allosteric site of the HIV reverse transcriptase near the polymerase site and interfere with reverse transcription by either altering the conformation or the mobility of the reverse transcriptase, thus leading to noncompetitive inhibition of the enzyme (Kohlstaedt et al., Science (1992), 256:1783-90).

Several classes of compounds have been identified as N1
Examples of these include:

- 1) 1-[(2-hydroxyethoxy)methyl]-6-(phenylthio)thymines (HEPT) (Tanaka et al., J. Med. Chem. (1991), 34:349-57; Pontikis et al., J. Med. Chem. (1997), 40:1845-54; Danel et al., J. Med. Chem. (1996), 39:2427-31; Baba et al., Antiviral Res. (1992), 17:245-64);
- 2) bis(heteroaryl)piperazines (BHAP) (Romero et al., J. Med. Chem. (1993), 36:1505-8);
- 3) dihydroalkoxybenzyloxypyrimidine (DABO) (Danel et al., Acta Chemica Scandinavica (1997), 51:426-30; Mai et al., J. Med. Chem. (1997), 40:1447-54);
- 4) 2', 5'-bis-O-(tertbutyldimethylsilyl)-3'-spiro-5''-(4''-amino-1'', 2''-oxathiole-2'', 2''-dioxide)pyrimidines (TSAO) (Balzarini et al., PNAS USA (1992), 89:4392-96);
- 5) phenylethylthiazolylthiourea (PETT) derivatives (Bell et al., J. Med. Chem. (1995), 38:4929-36; Cantrell et al., J. Med. Chem. (1996), 39:4261-74);
- 6) tetrahydro-imidazo[4,5,1-jk][1,4]-benzodiazepine-2(1H)-one and -thione (TIBO) derivatives (Pauwels et al., Nature (1990), 343:470-4);
- 7) phosphorus-substituted imidazole derivatives (PCT Publication No. WO 03/091264 A2 to Gilead Sciences, Inc.);
- 8) alpha-anilinophenylacetamide (alpha-APA) derivatives (Pauwels et al., PNAS USA (1993), 90:1711-15); and
- 9) indole derivatives (U.S. Patent No. 5,527,819 to Merck & Co. and counterpart PCT Publication No. WO 94/19321).

Indole derivatives described in U.S. Patent No. 5,527,819 assigned to Merck & Co. have been shown to be inhibitors of HIV reverse transcriptase. Some of these compounds exhibited IC₅₀ values against HIV reverse transcriptase at concentrations of from 3-35 nM. A process for synthesizing optionally substituted indoles by a palladium-catalyzed annulation between a ketone and an iodoaniline was also developed at Merck & Co. (Chen et al., J. Org. Chem. (1997), 62(9):2676-77).

The compounds disclosed in the '819 patent are generally represented by the following broad structural Formula (III):



in which the variables X, Y, Z, R and R⁶ are broadly defined.

U.S. Patent No. 5,124,327 to Merck & Co. discloses a class of optionally substituted sulfonylphenyl indole compounds. The patent reports that the compounds are active as reverse transcriptase inhibitors and could be useful in the treatment of HIV infection and AIDS.

U.S. Patent No. 6,710,068 to Idenix Pharmaceuticals, Ltd., discloses a class of phenylindoles substituted with at least two moieties other than hydrogen on either or both rings. See also PCT Publication No. WO 02/083126.

PCT Publication No. WO 2004/014364 to Idenix Pharmaceuticals discloses another class of phenylindoles that displays enhanced anti-HIV activity. These compounds are also substituted with at least two moieties other than hydrogen on either or both rings. In addition, these compounds incorporate a number of different substituents with a carboxamide functionality at position-2 on the indole, the position shown in formula (II) above as "Z". Typical placement of substituents is at the 3' and 5' positions on the phenyl ring, and at the 4' and 5', 5' and 6', or 5' and 7' positions on the benzo ring of the indole moiety.

Bristol Myers Squibb disclose various optionally substituted indoles, azaindoles, piperazines, and pyrrolidines for the treatment of HIV and/or AIDS in several U.S. patents and U.S. and PCT publications. See U.S. Publication Nos. 2004/0006090; 2004/0063746; 2003/0096825; 2003/0236277; and WO 03/068221.

WO 01/02388 to SmithKline Beecham S.P.A discloses optionally substituted phenylindoles with a carbamyl substituent that are alleged to have utility in the treatment of HIV, AIDS, osteoporosis, cancers, and Alzheimer's disease.

Warner-Lambert Company discloses various indole-thiazepinone diazepinones, benzothiophenes, benzofurans, and indole-2-carboxamides for the treatment of HIV in U.S. 5,424,329; U.S. 5,565,446; U.S. 5,703,069; and WO 96/29077.

Shinogi & Co. report optionally substituted indole derivatives that are viral integrase inhibitors useful as anti-HIV drugs in U.S. Publication No. 2002/0019434 and U.S. Patent Nos. 6,716,605 and 6,506,787.

U.S. Patent No. 5,945,440 to Kleinschroth et al. discloses a class of indolocarbazole amides for the treatment of a variety of diseases including cancer, viral diseases (including HIV), cardiac and vascular diseases, bronchopulmonary diseases, inflammatory disorders, degenerative diseases of the central nervous system, and other diseases.

U.S. Patent No. 4,866,084 to Gunasekera et al. teaches certain bisindole alkaloid compounds that have antiviral and antitumor activity, including HSV (herpes simplex virus). U.S. Patent 5,935,982 to Dykstra et al. reports a different class of bisindoles that have utility versus retroviral infections and especially HIV.

U.S. Patent No. 5,852,011 to Matsunaga et al. discloses a class of indole derivatives substituted by a heteroaryl function and an amide function. The compounds are said to possess antitumor, antiviral, and antimicrobial properties.

U.S. Patent No. 5,935,982 to Dykstra et al. discloses a class of bis-indoles and specifically propose their use for treating retroviral infections, and especially infection by HIV.

U.S. Patent No. 5,929,114 to Domagala et al. discloses a class of arylthio and bithiobisarylamide compounds, including indole derivative, that reportedly have antibacterial and antiviral activity.

U.S. Patent No. 5,830,894 to Pevear et al. discloses a class of triazinoindole derivatives that reportedly have anti-pestivirus activity, most notably BVDV activity.

Indoles have been used in the treatment of diseases other than HIV. U.S. Patent No. 5,981,525 to Farina et al. discloses a complex array of indoles for use in the treatment of osteoporosis based on their ability to inhibit osteoclast H^+ -ATPase and thus reduce bone resorption. U.S. Patent No. 6,025,390, also to Farina et al., teaches another

group of indole derivatives, termed heteroaromatic pentadienoic acid derivatives, for the treatment of osteoporosis. U.S. Patent No. 5,489,685 to Houpis et al. discloses a series of compounds that are furo(2,3-b) pyridine carboxylic acid esters, allegedly useful in the treatment of HIV.

It is known that over a period of time, antiviral agents that are active against HIV induce mutations in the virus that reduce the efficacy of the drug. This was apparently the problem exhibited by the Merck indoles in U.S. Patent No. 5,527,819 (Williams *et al.*, J. Med. Chem., 1993, 36(9), 1291-94). Drug resistance most typically occurs by mutation of a gene that encodes an enzyme used in viral replication, and most typically in the case of HIV, reverse transcriptase, protease, or DNA integrase. It has been demonstrated that the efficacy of a drug against HIV infection can be prolonged, augmented, or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation from that caused by the principal drug. Alternatively, the pharmacokinetics, biodistribution, or other parameters of a drug can be altered by such combination or alternation therapy. In general, combination therapy is typical rather than alternation therapy since combination therapy induces multiple simultaneous pressures on the virus. However, one cannot predict which mutations will be induced in the HIV-1 genome by a given drug, whether the mutations are permanent or transient, or how an infected cell with a mutated HIV-1 sequence will respond to therapy with other agents in combination or alternation. These factors are exacerbated by the fact that there is a paucity of data on the kinetics of drug resistance in long-term cell cultures treated with modern antiretroviral agents.

Therefore, there is a need to provide new compounds and methods for the treatment of HIV.


It is thus an object of the present invention to provide new compounds, compositions, methods and uses for the treatment of patients infected with HIV.

It is yet another object of the present invention to provide new compositions and methods for the treatment of patients infected with HIV that exhibit activity against drug-resistant forms of the virus.

SUMMARY OF THE INVENTION

3-phosphoindole compounds display antiviral activity against HIV, in particular against strains of HIV that have developed cross resistance to other anti-HIV drugs. Compounds, compositions and methods for treatment of HIV infection are disclosed that include the 3-phosphoindole compounds. The 3-phosphoindoles can be in the form of a wide variety of moieties, including, but not limited to, phosphates, phosphonates, a phosphorothioate, including thiophosphates, thiophosphonates, phosphate, and a phosphoramidate, including iminiophosphates and iminophosphonates.

In one embodiment, the compounds that can have anti-HIV activity are of Formula (A) below. These compounds include an phosphorus-linked substituent at position-3 and particular substituents at position-2 on the indole, and a monosubstitution at position R^{5'} or a disubstitution at positions R^{4'} and R^{5'}, R^{5'} and R^{6'}, or R^{5'} and R^{7'} on the benzo ring.



In one particular embodiment, substituent "X" represents a phenyl ring that is unsubstituted or is substituted by one or more halogens or lower alkyl groups such as methyl or ethyl. Particular substituents at position 2 on the indole moiety include, for example, hydrogen, hydroxy, halogen, alkyl, aryl, heteroaryl, and especially substituents having a carboxamide or carboxamide moiety shown as "Z" in Formula (A). Substituents for the benzo ring of the indole moiety include but are not limited to chlorine, fluorine, bromine, iodine, CF₃, CN, NO₂, and methoxy.

The active compound may be a salt or prodrug that, upon administration, provides directly or indirectly the parent compound or that itself exhibits a desired activity. In another embodiment, the compound of Formula A includes a charged heteroatom, and in a particular embodiment, the compound includes an N-oxide group. Modifications affecting the biological activity of the compounds of the present invention

also are included here, as are any changes that produce increased activity of the parent compounds.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an illustration of three (3) general structures of the present invention given as Formula (A), Formula (B), and Formula (C).

Figure 2 is an illustration of phosphorylated compounds (I)-(X) of the general Formulae (A) and (B).

Figure 3 is an illustration of 9- and 10-membered bicyclic phosphorylated compounds (XI)-(XX) of the general Formula (C).

DETAILED DESCRIPTION OF THE INVENTION

Provided are a composition of matter, method of use and a pharmaceutical composition for the treatment of retroviral infections in mammals and, in particular, HIV in humans. Included within the present invention are the following features:

- 3-phosphoindoles and pharmaceutically acceptable salts and prodrugs thereof as described herein, optionally substantially free of other chemical entities;
- 3-phosphoindoles of Formulas A-C and pharmaceutically acceptable salts and prodrugs thereof as described herein, optionally substantially free of other chemical entities;
- 3-phosphoindoles and pharmaceutically acceptable salts and prodrugs thereof as described herein that are effective against HIV in a host;
- 3-phosphoindoles and pharmaceutically acceptable salts and prodrugs thereof as described herein that are effective against drug-resistant strains of HIV in a host, and, in certain embodiments, where the drug-resistant strains of HIV is due to a reverse transcriptase mutation, such as lysine 103 → asparagine and/or tyrosine 181 → cysteine;
- 3-phosphoindoles and pharmaceutically acceptable salts and prodrugs thereof as described herein for use in the treatment or prophylaxis of an HIV infection in a host, or in the manufacture of a medicament for the treatment or prophylaxis of

an HIV infection in a host, especially in individuals diagnosed as having an HIV infection or being at risk for such infection;

- 3-phosphoindoles and pharmaceutically acceptable salts and prodrugs thereof as described herein for use in the treatment or prophylaxis of an HIV infection, or in the manufacture of a medicament for the treatment or prophylaxis of an HIV infection, which is resistant to one or more reverse transcriptase inhibitors, in a host;
- 3-phosphoindoles and their pharmaceutically acceptable salts and prodrugs thereof as described herein for use in the treatment or prophylaxis of an HIV infection as a form of salvage therapy in a host, or in the manufacture of a medicament for the treatment or prophylaxis of an HIV infection as a form of salvage therapy in a host, especially in individuals diagnosed as having an HIV infection or being at risk for such infection;
- 3-phosphoindoles and their pharmaceutically acceptable salts and prodrugs thereof as described herein for use in the treatment or prophylaxis, or in the manufacture of a medicament for the treatment or prophylaxis, of an HIV infection that is resistant to one or more reverse transcriptase inhibitors due to a reverse transcriptase mutation, such as lysine 103 → asparagine and/or tyrosine 181 → cysteine, in a host, especially in individuals diagnosed as having an HIV infection or being at risk for such infection;
- processes for the preparation of 3-phosphoindoles, optionally substantially isolated from other chemical entities;
- pharmaceutical compositions comprising an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug thereof, together with a pharmaceutically acceptable carrier or diluent;
- pharmaceutical compositions comprising an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug thereof, in combination with one or more other anti-HIV agents, optionally with a pharmaceutically acceptable carrier or diluent;

- pharmaceutical compositions for the treatment or prophylaxis of an HIV infection that is resistant to one or more reverse transcriptase inhibitors, in a host, comprising an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination with at least one other anti-HIV agent, optionally with a pharmaceutically acceptable carrier or diluent;
- pharmaceutical compositions for the treatment or prophylaxis of an HIV infection in a host as a form of salvage therapy comprising an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent;
- pharmaceutical compositions for the treatment or prophylaxis of an HIV infection that is resistant to one or more reverse transcriptase inhibitors due to a reverse transcriptase mutation, such as lysine 103 → asparagine and/or tyrosine 181 → cysteine, in a host comprising an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent;
- methods for the treatment or prophylaxis of an HIV infection in a host, wherein the HIV can be resistant to one or more reverse transcriptase inhibitors, comprising administering to said host an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination or alternation with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent;
- methods for the treatment or prophylaxis of an HIV infection in a host as a form of salvage therapy comprising administering to said host an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination or alternation with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent;
- methods for the treatment or prophylaxis of an HIV infection, which is resistant to one or more reverse transcriptase inhibitors due to a reverse transcriptase

mutation, such as lysine 103 → asparagine and/or tyrosine 181


host comprising administering to said host an effective anti-HIV treatment amount of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination or alternation with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent;

- use of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination or alternation with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent, for the treatment or prophylaxis of an HIV infection in a host;
- use of a 3-phosphoindole or its pharmaceutically acceptable salt or prodrug, optionally in combination or alternation with at least one other anti-HIV agent, and optionally with a pharmaceutically acceptable carrier or diluent, for the treatment or prophylaxis of an HIV infection, or in the manufacture of a medicament for the treatment or prophylaxis of an HIV infection, which is resistant to one or more reverse transcriptase inhibitors, which can be due to a reverse transcriptase mutation, such as lysine 103 → asparagine and/or tyrosine 181 → cysteine, in a host. This use can be a form of salvage therapy; and
- any or all of the foregoing in which the host is a human.

I. Active Compounds of the Present Invention

In a general embodiment of the invention, a 3-phosphoindole or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof is provided. The 3-phosphoindole can be in the form of a phosphate, phosphonate, thiophosphate, thiophosphonate, iminiophosphate or iminophosphonate.

In a first embodiment of the present invention, the compound is represented generally by the following chemical Formula (A):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

each X and Y independently is

- a) H;
- b) halogen (F, Cl, Br, or I), typically F;
- c) R³;
- d) CF₃;
- e) C₁₋₆ alkyl;
- f) C₂₋₆ alkenyl;
- g) C₂₋₆ alkynyl;
- h) alkylheterocycle;
- i) 3-14 membered carbocycle, aryl, heterocycle, any of which may comprise a monocyclic, bicyclic, tricyclic or spiro structure;

- j) OH;
- k) OR^2 ;
- l) O-alkyl;
- m) O-alkenyl;
- n) O-alkynyl;
- o) O-alkylaryl;
- p) O-aryl;
- q) O-heterocycle;
- r) O-aralkyl;
- s) O-carbocycle;
- t) SH;
- u) SR^2 ;
- v) S-alkyl;
- w) S-alkenyl;
- x) S-alkynyl;
- y) S-alkylaryl;
- z) S-aryl;
- aa) S-heterocycle;
- bb) S-aralkyl;
- cc) S-carbocycle;
- dd) NH_2 ;
- ee) NHR^2 ;
- ff) NR^2R^2 ;
- gg) NH-alkyl;

- hh) N-dialkyl;
- ii) NH-aryl;
- jj) N-alkaryl;
- kk) N-aralkyl;
- ll) NH-heterocycle;
- mm) N-alkyl-heterocycle;
- nn) N-alkenyl-heterocycle;
- oo) N-alkynyl-heterocycle; or


alternatively, X and Y may come together to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle wherein each ring comprises 3-7 members;

Z is:

- a) H;
- b) CN;
- c) NO₂;
- d) C₁₋₆ alkyl;
- e) C₂₋₆ alkenyl;
- f) C₂₋₆ alkynyl;
- g) alkaryl;
- h) aralkyl;
- i) heterocycle;
- j) alkyl-heterocycle;
- k) aryl;
- l) alkoxy;
- m) OR²;

- n) SR^2 ;
- o) $S(O)_nR^2$;
- p) $S(O)_n-NR^2R^3$;
- q) $N(R^2)(R^3)$;
- r) carboxamido;
- s) amido;
- t) acyl;
- u) $C(=W)-R^3$;
- v) $C(=W)NH-C(R^3)(R^3)-C(=W)-N(R^2)(R^2)$;
- w) $C(=W)NH-P(=W)(R^3)-A-R^3$;
- x) $C(=W)NH-A-S(O)_n-NR^2$;
- y) $C(=W)NH-CR^3R^3-S(O)_n-NR^2R^2$;
- z) $C(=W)-NH-A-C(=W)-N(R^2)(R^2)$;
- aa) $C(=W)-N(R^2)(R^2)$;
- bb) $C(=W)-NH-A-R^3$;
- cc) $C(=W)-NH-NH-R^3$;
- dd) $C(=W)-NH-C(R^3)(R^3)-C(=W)NH-C(R^3)(R^3)-C(=W)-N(R^2)(R^2)$;
- ee) $C(=W)-NH-R^2$;
- ff) $C(=W)-NH-A-C(=W)-NH-A-C(=W)-NH_2$;
- gg) $C(R^2)(R^3)(R^3)$;
- hh) $C(R^2)(R^3)-NH-R^2$;
- ii) $A-S(O)_n-R^3$;
- jj) $C(=W)-A-C(=W)-A-C(=W)R^3$;
- kk) $A-R^3$;

- ll) $C(=W)-(O)R^2$;
- mm) $C(=W)-A-C(=W)-NH_2$;
- nn) an amino acid residue;
- oo) $C(=W)-N(R^2)-A-(\text{amino acid residue})$;
- pp) $C(=W)-N(R^2)-A-(\text{amino acid residue})-C(=W)-R^3$;
- qq) $C(=W)-\text{amino acid residue}$;
- rr) $C(=W)-N(R^2)-A-(\text{amino acid residue})-A-C(=W)-R^3$;
- ss) $C(=W)-OR^2$;
- tt) $C(=W)-S(R^2)$;
- uu) $C(=W)-NH-NH-R^2$;
- vv) $C(=W)-NH-N(R^2)-A-C(=W)R^3$;
- ww) $C(=W)-N(R^2)-C(=W)-R^3$;
- xx) $C(=W)-A-NH-C(=W)R^3$;
- yy) $C(=W)-A-NH-C(=W)OR^2$;
- zz) $C(=W)-A-R^3$;
- aaa) $C(=W)-NH-NH-CH_2-C(=W)R^3$;
- bbb) $P(=W)(R^3)(R^3)$; or
- ccc) $A-P(=W)(R^3)(R^3)$.
- ddd) $C(=W)-NH-C_{1-10}\text{alkyl-heteroaryl}$
- eee) $C(=W)-NH-C_{1-4}\text{alkyl-heteroaryl}$
- fff) $C(=W)-NH-CH_2\text{-heteroaryl}$



ggg)

wherein in embodiments (ddd), (eee) and (fff), the heteroaryl can optionally include a charged heteroatom, and in particular can include an N-oxide

wherein each X, Y and Z independently may be unsubstituted or substituted by one or more of C₁₋₆ alkyl; alkoxy; OH; oxo; halo (F, Cl, Br, or I); NR²R²; optionally substituted aryl; optionally substituted heterocycle; O-C(=W)-alkyl; C(=W)-OR²; CN; NO₂; NH-C(=W)-alkyl; NH-S(O)_n-alkyl; NH-S(O)_n-NR²R²; or C₃₋₆ cycloalkyl;

each W is independently:

- a) O;
- b) S;
- c) NH;
- d) N-N(R²)(R²);
- e) N(R²);
- f) N-OH;
- g) N-O-alkyl; or
- h) N-O-R²;

R¹ is:

- a) H;
- b) -R²;
- c) C(=W)-R³;
- d) C(=W)-O(R²);
- e) C(=W)-S(R²);
- f) C(=W)-NH-R²;
- g) C(=W)-N(R²)(R²);
- h) C(=W)-NH-A-(amino acid residue);
- i) A-(amino acid residue)-R³;

j) $S(O)_n-R^3$; or

k) $S(O)_2-N(R^2)(R^2)$;

any of which optionally may be substituted by one or more of C_{1-6} alkyl; OH; alkoxy; aryl; halo; CN; NO_2 ; or $N(R^2)(R^2)$;

each R^2 is independently:

a) H;

b) CF_3 ;

c) CN;

d) optionally substituted, branched or unbranched, alkyl, such as a C_{1-6} alkyl;

e) optionally substituted, branched or unbranched, alkenyl, such as a C_{2-6} alkenyl;

f) optionally substituted, branched or unbranched, alkynyl, such as a C_{2-6} alkynyl;

g) 3-14 membered carbocycle;

h) optionally substituted aryl;

i) optionally substituted aralkyl;

j) optionally substituted alkylaryl;

k) optionally substituted heterocycle;

l) optionally substituted alkylheterocycle;

m) optionally substituted heterocycle-alkyl;

n) A-heterocycle;

o) acyl;

p) alkoxy;

q) $CH_2-S(O)_nR^3$;

r) $C(alkyl)_2-S(O)_nalkyl$;

- s) $\text{CH(alkyl)-S(O)}_n\text{alkyl}$;
- t) CH_2NH_2 ;
- u) $\text{CH}_2\text{NH(alkyl)}$;
- v) $\text{CH}_2\text{N(alkyl)}_2$;
- w) CH(alkyl)-NH_2 ;
- x) $\text{CH(alkyl)-NH(alkyl)}$;
- y) $\text{CH(alkyl)-N(alkyl)}_2$;
- z) $\text{C(alkyl)}_2\text{-NH}_2$;
- aa) $\text{C(alkyl)}_2\text{-NH(alkyl)}$;
- bb) $\text{C(alkyl)}_2\text{-N(alkyl)}_2$;
- cc) $\text{CH}_2\text{-C(=W)H}$;
- dd) $\text{CH}_2\text{-C(=W)alkyl}$;
- ee) A-alkyl;
- ff) $\text{C(alkyl)}_2\text{-C(=W)alkyl}$;
- gg) $\text{CH}_2\text{-C(=W)H}$;
- hh) $\text{CH}_2\text{-C(=W)alkenyl}$;
- ii) $\text{CH(alkenyl)-C(=W)H}$;
- jj) A-S(O)alkyl;
- kk) $\text{CH(NH)-S(O)}_n\text{alkyl}$; or
- ll) A-N(NH)alkyl;
- mm) $\text{C(R}^3\text{)(R}^3\text{)-S(O)}_n\text{NH}_2$;
- nn) $\text{C(R}^3\text{)(R}^3\text{)-S(O)}_n\text{CF}_3$;
- oo) $\text{C(R}^3\text{)(R}^3\text{)-NH}_2$;

each R^3 is independently:

- a) H;
- b) OH;
- c) halogen (F, Cl, Br, or I);
- d) CF₃;
- e) CN;
- f) optionally substituted, branched or unbranched, alkyl, such as a C₁₋₆ alkyl;
- g) optionally substituted, branched or unbranched, alkenyl, such as a C₂₋₆ alkenyl;
- h) optionally substituted, branched or unbranched, alkynyl, such as a C₂₋₆ alkynyl;
- i) 3-14 membered carbocycle;
- j) optionally substituted aryl;
- k) optionally substituted aralkyl;
- l) optionally substituted alkylaryl;
- m) optionally substituted heterocycle;
- n) optionally substituted alkylheterocycle;
- o) optionally substituted heterocycle-alkyl;
- p) A-heterocycle;
- q) acyl;
- r) carboxamido;
- s) carbamoyl;
- t) alkoxy;
- u) OH
- v) OR²;

- w) O-alkyl;
- x) O-alkenyl;
- y) O-alkynyl;
- z) O-alkaryl;
- aa) O-aralkyl;
- bb) O-carbocycle;
- cc) O-heterocycle;
- dd) O-aryl;
- ee) SH
- ff) SR^2 ;
- gg) S-alkyl;
- hh) S-alkenyl;
- ii) S-alkynyl;
- jj) S-alkaryl;
- kk) S-aralkyl;
- ll) S-carbocycle;
- mm) S-heterocycle;
- nn) S-aryl;
- oo) $S(O)_nR^2$;
- pp) amino;
- qq) NH^2 ;
- rr) NHR^2 ;
- ss) $N(R^2)(R^2)$;
- tt) $NH-S(O)_nR^2$;

- uu) NHC(=W)-aryl ;
- vv) NHC(=W)-alkyl ;
- ww) $\text{NH-C(=W)-heterocycle}$;
- xx) $\text{CH}_2\text{-S(O)}_n\text{R}^2$;
- yy) C(=W)R^2 ;
- zz) $\text{C(=W)-N(R}^2\text{)-R}^2$;
- aaa) $\text{C(alkyl)}_2\text{-S(O)}_n\text{R}^2$;
- bbb) $\text{CH(alkyl)-S(O)}_n\text{R}^2$;
- ccc) $\text{C(alkyl)}_2\text{-NH}_2$;
- ddd) $\text{CH(alkyl)-N(alkyl)R}^2$;
- eee) $\text{C(R}^2\text{)(R}^2\text{)-NR}^2\text{R}^2$;
- fff) $\text{CH}_2\text{N(alkyl)R}^2$;
- ggg) CH(alkyl)-NHR^2 ;
- hhh) $\text{C(alkyl)}_2\text{-NHR}^2$;
- iii) $\text{C(alkyl)}_2\text{-N(alkyl)R}^2$;
- jjj) $\text{CH}_2\text{-C(=W)H}$;
- kkk) $\text{CH}_2\text{-C(=W)alkyl}$;
- lll) $\text{CR}^2\text{R}^2\text{-C(=W)R}^2$;
- mmm) A-R^2 ;
- nnn) $\text{C(R}^2\text{)}_2\text{-C(=W)R}^2$;
- ooo) $\text{CH}_2\text{-C(=W)H}$;
- ppp) $\text{CH}_2\text{-C(=W)alkenyl}$;
- qqq) $\text{CH(alkenyl)-C(=W)H}$;
- rrr) A-S(O)R^2 ;

sss) $\text{CH}(\text{NH})\text{-S}(\text{O})_n\text{R}^2$; or

ttt) $\text{A-N}(\text{NH})\text{R}^2$;

uuu) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{NH}_2$;

vvv) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{CF}_3$;

www) $\text{C}(\text{R}^2)(\text{R}^2)\text{-NH}_2$;

wherein the optional substitution comprises one or more of

a) a substituted or unsubstituted heterocycle;

b) $\text{C}(=\text{W})\text{O-aryl}$;

c) $\text{C}(=\text{W})\text{O-alkyl}$;

d) $\text{C}(=\text{W})\text{NH}_2$;

e) $\text{C}(=\text{W})\text{NH-alkyl}$;

f) $\text{C}(=\text{W})\text{NH-aryl}$;

g) $\text{C}(=\text{W})\text{N-di-alkyl}$;

h) $\text{C}(=\text{W})\text{N(alkyl)-aryl}$;

i) α -amino acid;

j) α -amino ester;

k) α -amino-carboxamide;

l) β -amino acid;

m) β -amino ester; or

n) β -amino-carboxamide;

wherein, when the optional substitution includes a substituted heterocycle, then the heterocycle substitution is selected from the group consisting of:

a) $\text{C}(=\text{W})\text{O-aryl}$;

b) $\text{C}(=\text{W})\text{O-alkyl}$;

- c) $C(=W)NH_2$;
 - d) $C(=W)NH$ -aryl;
 - e) $C(=W)NH$ -alkyl;
 - f) $C(=W)N$ -di-alkyl;
 - g) $C(=W)N$ (alkyl)-aryl;
 - h) α -amino acid;
 - i) α -amino ester;
 - j) α -amino-carboxamide;
 - k) β -amino acid;
 - l) β -amino ester; and
 - m) β -amino-carboxamide;
 - n) halo; or
 - o) cyano,
- alone or in any combination;

n independently is 0, 1 or 2;

each A is independently a disubstituted spacer selected from the group consisting of:

- a) C_{1-6} alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- b) C_{2-12} alkenylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- c) C_{2-12} alkynylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- d) optionally substituted arylene;

- e) O-alkylene, branched or unbranched, and optionally having heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- f) aralkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- g) optionally substituted cycloalkyl; and
- h) optionally substituted heterocycle;

wherein "A" may be joined by any desired linkage such as, for example, an ether, thioether, amino, carboxamido, ester or carbon-carbon linkage, or any combination thereof;

each $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ independently is:

- a) H;
- b) halogen (F, Cl, Br, I);
- c) NO_2 ;
- d) CN;
- e) CF_3 ;
- f) OH
- g) OR^2 ;
- h) SH
- i) SR^2 ;
- j) NR^2R^2 ;
- k) $NHS(O)_nR^2$;
- l) $NHCO-C_{1-3}$ alkyl;
- m) $S(O)_nR^2$;
- n) aryl;

- o) heterocycle;
- p) C₁₋₆ alkyl;
- q) C₂₋₆ alkenyl;
- r) C₂₋₆ alkynyl;
- s) C(=W)-S(O)_nR²;
- t) C(=W)-S(O)_n-NR²R²;
- u) C(=W)-aryl;
- v) C(=W)-alkyl;
- w) C(=W)-heterocycle; or
- x) C(=W)-NR²R²;

each of which optionally may be substituted with one or more of :

- a) OR²;
- b) S(O)_nR²;
- c) C(=W)-S(O)_nR²;
- d) C(=W)-S(O)_n-NR²R²;
- e) C(=W)-aryl;
- f) C(=W)-alkyl;
- g) C(=W)-heterocycle;
- h) C(=W)NR²R²;
- i) NO₂;
- j) CN;
- k) CF₃;
- l) halogen (F, Cl, Br, I);
- m) NHS(O)_nR²;

- n) NHCO-C_{1-3} alkyl;
- o) aryl;
- p) heterocycle;
- q) C_{1-6} alkyl;
- r) C_{2-6} alkenyl;
- s) C_{2-6} alkynyl; or
- t) NR^2R^2 .

In one embodiment of Formula (A), X is an optionally substituted phenyl; Y is any of its definitions; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, and $\text{R}^{7'}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is a carboxamide moiety.

In an alternative embodiment, R^1 is acyl, alkyl, aryl, alkaryl, or aralkyl.

In yet another alternative embodiment, $\text{R}^{4'}$ is fluoro, nitro or cyano, W is oxygen, and Y is O-alkyl.


In a second embodiment of Formula (A), X is an optionally substituted phenyl; Y is any of its definitions; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, and $\text{R}^{7'}$ are all hydrogen; $\text{R}^{5'}$ is chlorine; and Z is a carboxamide or carboxamido-heterocyclyl moiety.

In yet another embodiment of Formula (A), X is tolyl, thiazolyl or pyridyl; Y is H, OH, or O-alkyl; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, and $\text{R}^{7'}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamido, acyl, an alkyl-sulphonyl group or a carboxylic acid derivative.

In an alternative to the preceding embodiment, X, Y, R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$ and $\text{R}^{5'}$ all are as defined above, and Z is a carboxamido-alkylene-heterocycle, typically carboxamido-alkylene-pyridyl; a thioamido-pyridyl wherein the pyridyl is unsubstituted or substituted by OH, OMe or lower alkyl; an imino-nitrile; or an alkylsulphonyl-aryl group.

In yet another embodiment, Z is a carboxamido-alkylene-heterocycle wherein the heterocycle includes at least one N-oxide group.

In a second embodiment, the invention provides a phenylindole for use in the treatment of HIV represented by the following general Formula (B):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

each W, Y, Z, R¹, R², R³, A, n, R⁴', R⁵', R⁶' and R⁷' is as defined above for Formula (A);

and

each R²'', R³'', R⁴'', R⁵'', and R⁶'' independently is:

- a) H;
- b) halogen;
- c) NO₂;
- d) CN;
- e) OR²;
- f) SR²;
- g) NH₂;
- h) NR²R³;
- i) N(R²)-C(=W)-C₁₋₄ alkyl;
- j) N(R²)-SO₂-C₁₋₄ alkyl;
- k) C₁₋₆ alkyl;
- l) C₂₋₆ alkenyl;

- m) C_{2-6} alkynyl;
- n) aryl;
- o) CF_3 ;
- p) $CR^2R^2-S(O)_n-R^3$;
- q) $CR^2R^2NR^2R^3$;
- r) C-OH;
- s) $CR^2R^2-C(=W)R^2$;
- t) acyl;
- u) $C(=W)R^2$;
- v) $C(=W)OR^2$;
- w) $C(=W)SR^2$;
- x) $C(=W)-NR^2R^3$;
- y) $C(=W)NH(CH_2)_p-(\text{amino acid residue})$;
- z) amino residue; or
- aa) A-(amino acid residue);

wherein any of the above optionally may be substituted; or

alternatively, $R^{2''}$ or $R^{6''}$ may be joined to Y to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle wherein each ring comprises 3-14 members.

The following are non-limiting examples of embodiments of Formula (B):


- a) Y is alkyl; W is O, $S(O)_n$, or NH; R^1 , $R^{4'}$, $R^{6'}$, $R^{7'}$, $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ are all hydrogen; $R^{5'}$ is halogen; and Z is carboxamide;
- b) Y is aryl; W is O, $S(O)_n$, or NH; R^1 , $R^{4'}$, $R^{6'}$, $R^{7'}$, $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ are all hydrogen; $R^{5'}$ is amino-alkyl, thioamino-alkyl, or aminocarbonyl-alkyl; and Z is carboxamide;

- c) Y is $-\text{OH}$ or $-\text{SR}^2$; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{NH}_2$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{4''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl; and Z is carboxamide;
- d) Y is $-\text{OH}$; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{4''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{4'}$ and $\text{R}^{5'}$ are halogen; $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl; and Z is carboxamide;
- e) Y is $-\text{OH}$; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{NR}^2\text{R}^2$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide;
- f) Y is alkenyl; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide or carboxamide-heterocycle;
- g) Y is alkynyl or $-\text{NR}^2\text{R}^3$; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{O-alkyl}$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide;
- h) Y is alkenylene; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide;
- i) Y is ethyl; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{OH}$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is chloro; and Z is carboxamide;
- j) Y is $-\text{O-methyl}$; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide;
- k) Y is $-\text{O-ethyl}$; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{NH}_2$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is carboxamide-heterocycle;
- l) Y is $-\text{O-ethyl}$; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is chloro; and Z is carboxamide;
- m) Y is $-\text{O-H}$; W is O, $\text{S}(\text{O})_n$, or NH ; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is a carboxamide alkyl group that is unsubstituted or substituted by one or more $-\text{NO}_2$, $-\text{NH}-\text{C}(=\text{O})\text{-alkyl}$, or $-\text{NH}-\text{S}(\text{O})_n\text{-alkyl}$;
- n) Y is $-\text{O-H}$; W is O, $\text{S}(\text{O})_n$, or $\text{N}-\text{NH}_2$; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$, $\text{R}^{7'}$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are all hydrogen; $\text{R}^{5'}$ is halogen; and Z is a carboxamido-alkylene-

heterocyclyl group wherein said heterocycle imidazole, thiazole, pyridyl or furan, and wherein the heterocycle is unsubstituted or further substituted by one or more halo, oxo, -OH, -NO₂, -MeOH, -NH-C(=O)-alkyl, or -NH-S(O)_n-alkyl;

- o) Y is -O-H; W is O, S(O)_n, or NH; R¹, R^{4'}, R^{6'}, R^{7'}, R^{2''}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all hydrogen; R^{5'} is halogen; and Z is carboxamide-cyclopropyl or carboxamide-cyclobutyl;
- p) Y is -O-methyl; W is O, S(O)_n, or NH; R¹, R^{4'}, R^{6'}, R^{7'}, R^{2''}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all hydrogen; R^{5'} is halogen; and Z is a carboxamide-ethyl, carboxamide -ethanol, or carboxamide -ethyl-methoxy;
- q) Y is -OH or -NR²R³; W is O, S(O)_n, or N-O-alkyl; R¹, R^{4'}, R^{6'}, R^{7'}, R^{2''}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all hydrogen; R^{5'} is halogen; and Z is a carboxamide alkyl-phenyl group wherein the phenyl is further substituted by one or more halo, oxo, -OH, -OCH₃, -NO₂, -MeOH, or -NH-C(=O)-alkyl groups;
- r) Y is -OH or -SR²; W is O, S(O)_n, or N-NR²R²; R¹, R^{4'}, R^{6'}, R^{7'}, R^{2''}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all hydrogen; R^{5'} is halogen; and Z is an isopropanol carboxamide moiety; and
- s) Y is -OH; W is O, S(O)_n, or N-OH; R¹, R^{4'}, R^{6'}, R^{7'}, R^{2''}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all hydrogen; R^{5'} is halogen; and Z is a thioacetamide group.

In a third embodiment, the present invention provides an optionally substituted 9-11 membered bicyclic 3-phosphoindole for use in the treatment of HIV represented by the following general Formula (C):



or a pharmaceutically acceptable salt, prodrug, *N*—oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

each W, Z, R¹, R², R³, A, n, R^{4'}, R^{5'}, R^{6'} and R^{7'} is as defined above for Formula (A);

each R^{3''}, R^{4''}, R^{5''}, and R^{6''} is as defined above for Formula (B);

----- indicates the presence of a single or double bond, wherein:

when ----- indicates the presence of a double bond, each Y and T independently is:

- a) CR³;
- b) N; or
- c) S(=W);

such that at least one of Y and T is CR³; and

when ----- indicates the presence of a single bond, then each Y and T independently is:

- a) CHR³;
- b) C(R³)(R³);
- c) O;
- d) S; or
- e) NR²;

such that at least one of Y and T is C(R³)(R³); and

m is 1 or 2, with the proviso that m can only be 2 for T or Y = CR².

The following are non-limiting examples of embodiments of Formula (C):

- a) W is O, Y is CR², T is (CH₂)₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is carboxamide;


- b) W is S, Y is O, T is CH₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is carboxamido-heterocycle wherein said heterocycle is optionally substituted furan, imidazole, thiazole, or pyridyl;
- c) W is S, Y is NR², T is CH₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is carboxamide;
- d) W is O, Y is SR², T is CH₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is alkyl-carboxamide;
- e) W is S, Y is SR², T is CH₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is carboxamide;
- f) W is O, Y is CR², T is C-C(=W)R³, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is methoxymethyl-carboxamide;
- g) W is S, Y is O, T is C-C(=W)R³, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is halogen, and Z is carboxamide-cyclobutyl;
- h) W is O, Y is SR², T is C-C(=W)R³, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is chloro, and Z is alkyl carboxamide wherein alkyl is optionally substituted by one or more halo, oxo, -OH, -NO₂, -MeOH, -NH-C(=O)alkyl, or -NH-S(O)_n-alkyl;
- i) W is S, Y is NR², T is CC(=W)R³, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is chloro, and Z is carboxamide;
- j) W is S, Y is N, T is C-C(=W)R³, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is fluoro, and Z is carboxamide;
- k) W is NH, Y is CR², T is NR₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is amino-alkyl, and Z is carboxamide;
- l) W is NR², Y is O, T is (CH₂)₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is chloro, and Z is carboxamide;
- m) W is N-OH, Y is O, T is CH₂, R¹, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}, and R^{6''} are all H; R^{5'} is thioamino-alkyl, and Z is alkyl carboxamide ;

- n) W is S, Y is SR^2 , T is CH_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is a carboxamide-heterocyclyl group wherein the heterocycle is an optionally substituted pyridine, thiazole, imidazole or furan;
- o) W is N-O-alkyl, Y is NR^2 , T is CH_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is carboxamide-alkyl group wherein the alkyl is optionally substituted by one or more $-NO_2$, $-NH_2$, $-NH-C(=W)alkyl$, or $-NH-S(O)_n-alkyl$;
- p) W is NH, Y is SR^2 , T is $(CH_2)_2$, $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is carboxamide-alkyl-phenyl wherein phenyl is optionally substituted by one or more halo, oxo, OH, NO_2 , MeOH, $-NH-C(=O)alkyl$ or $-NH-S(O)_n-alkyl$ groups;
- q) W is NR^2 , Y is $(CH_2)_2$, T is NR_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is a carboxamide-cyclopropyl group;
- r) W is N-OH, Y is CH, T is O, $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is a carboxamide-cyclopropyl group;
- s) W is N-O-alkyl, Y is O, T is $(CH_2)_2$, $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is chloro, and Z is carboxamide-methoxyethyl;
- t) W is O, Y is N, T is CH_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is carboxamide-ethanol;
- u) W is N- NR^2R^2 , Y is NR^2 , T is CH_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is halogen, and Z is carboxamide; and
- v) W is O, Y is CR^2 , T is CH_2 , $R^1, R^{4'}, R^{6'}, R^{7'}, R^{3''}, R^{4''}, R^{5''}$, and $R^{6''}$ are all H; $R^{5'}$ is aminocarbonyl-alkyl, and Z is carboxamide.


In one set of embodiments of Formula A, B or C, Z is $C(=W)-R^3$; $C(=W)-NH-A-C(=W)-N(R^2)(R^2)$; $C(=W)-NH-A-R^3$; $C(=W)-NH-R^2$; or $C(=W)-A-R^3$.

In one embodiment of Formula A, B or C, the compound includes a charged heteroatom. In particular, a charged nitrogen, such as through an N-oxide, is part of the compound. The charged heteroatom can be on a heteroaromatic ring that is attached to the indole, for example through $C(O)NH-alkyl$, or specifically through $C(O)NH-CH_2$.

In a particular embodiment of Formula A, B or C, Z is $C(=O)-NH-R^2$ wherein R^2 is an optionally substituted alkylheterocycle, wherein the heterocycle is of the formula




, wherein each Rx is independently CH or N^+-O^- . In a particular embodiment, one of Rx is N^+-O^- . The alkyl group can be C_{1-10} alkyl, or C_{1-4} alkyl, or can specifically be a methylene or ethylene. In a specific embodiment, R^2 is of the formula




where n is 0, 1 or 2.


In particular embodiments of Formula A, B or C, the compound is:




B;




C




D;











E;




F;



G;




Additional examples of an N-oxide embodiment of a compound of Formula A, B or C are:



. Either substituent on the phenyl ring may be substituted by CN, Me, halo, alkyl, alkenyl, alkynyl, alkyl-CN or alkenyl-CN, as the most commonly synthesized and tested N-oxides in the series to date.

II. Particular Sub-Embodiments of the Present Invention

In the first embodiment of the invention, the compounds are represented generally by the Formula (A):



or a pharmaceutically acceptable salt, prodrug, N—oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

X and Y, each independently, is:

- a) 3-14 membered carbocycle, aryl, heterocycle, any of which may comprise a monocyclic, bicyclic, tricyclic or spiro structure, or optionally may be substituted;
- b) H;
- c) OH;
- d) Cl, Br, I, F;
- e) CF₃;
- f) C₁₋₆ alkyl;
- g) C₂₋₆ alkenyl;
- h) C₂₋₆ alkynyl;
- i) alkylheterocycle;
- j) NH₂;
- k) NH-alkyl;
- l) N-dialkyl;
- m) NH-aryl;
- n) N-alkaryl;
- o) N-aralkyl;
- p) NH-heterocycle;
- q) N-alkyl-heterocycle;
- r) N-alkenyl-heterocycle;
- s) N-alkynyl-heterocycle;
- t) O-alkyl;
- u) O-alkenyl;
- v) O-alkynyl;

- w) O-alkylaryl;
- x) O-aryl;
- y) O-heterocycle;
- z) O-aralkyl;
- aa) O-carbocycle;
- bb) SR^2 ; or
- cc) NR^2R^3 ;

alternatively, X and Y may be joined to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle wherein each ring comprises 3-7 members;

Z is:

- a) H;
- b) alkoxy;
- c) NO_2 ;
- d) $N(R^2)(R^3)$;
- e) OR^2 ;
- f) carboxamido;
- g) amido;
- h) acyl;
- i) $S(O)_nR^2$;
- j) $S(O)_n-NR^2R^3$;
- k) C_{1-6} alkyl;
- l) C_{2-6} alkenyl;
- m) C_{2-6} alkynyl;
- n) alkaryl;

- o) aralkyl;
- p) heterocycle;
- q) alkyl-heterocycle;
- r) aryl;
- s) CN;
- t) $C(=W)-R^2$;
- u) $C(=W)NH-C(R^2)(R^2)-C(=W)-N(R^2)(R^2)$;
- v) $C(=W)NH-P(=W)(R^2)-A-R^2$;
- w) $C(=W)NH-A-S(O)_n-NR^2$;
- x) $C(=W)NH-CR^2R^3-S(O)_n-NR^2R^3$;
- y) $C(=W)-NH-A-C(=W)-N(R^2)(R^3)$;
- z) $C(=W)-N(R^2)(R^3)$;
- aa) $C(=W)-NH-A-R^2$;
- bb) $C(=W)-NH-NH-R^2$;
- cc) $C(=W)-NH-C(R^2)(R^2)-C(=W)NH-C(R^2)(R^3)C(=W)-N(R^2)(R^3)$;
- dd) $C(=W)-NH-R^2$;
- ee) $C(=W)-NH-A-C(=W)-NH-A-C(=W)-NH_2$;
- ff) $C(R^2)(R^2)(R^3)$;
- gg) $C(R^2)(R^2)-NH-R^2$;
- hh) $A-S(O)_n-R^2$;
- ii) $C(=W)-A-C(=W)-A-C(=W)R^3$;
- jj) $A-R^2$;
- kk) $C(=W)-(O)R^2$;
- ll) $C(=W)-A-C(=W)-NH_2$;

- mm) an amino acid residue;
- nn) $C(=W)-N(R^2)-A-(\text{amino acid residue})$;
- oo) $C(=W)-N(R^2)-A-(\text{amino acid residue})-C(=W)-R^2$;
- pp) $C(=W)-\text{amino acid residue}$;
- qq) $C(=W)-N(R^2)-A-(\text{amino acid residue})-A-C(=W)-R^2$;
- rr) $C(=W)-OR^3$;
- ss) $C(=W)-S(R^2)$;
- tt) $C(=W)-NH-NH-R^2$;
- uu) $C(=W)-NH-N(R^2)-A-C(=W)R^2$;
- vv) $C(=W)-N(R^2)-C(=W)-R^3$;
- ww) $C(=W)-A-NH-C(=W)R^2$;
- xx) $C(=W)-A-NH-C(=W)OR^3$;
- yy) $C(=W)-A-R^3$;
- zz) $C(=W)-NH-NH-CH_2-C(=W)R^2$;
- aaa) $P(=W)(R^2)(R^2)$; or
- bbb) $A-P(=W)(R^2)(R^2)$;

wherein each of the foregoing X, Y and Z independently may be unsubstituted or substituted by one or more of:

- a) H;
- b) C_{1-6} alkyl;
- c) alkoxy;
- d) OH;
- e) oxo;
- f) halo;

- g) NR^2R^2 ;
- h) optionally substituted aryl;
- i) optionally substituted heterocyclyl;
- j) $\text{O}-\text{C}(=\text{W})\text{-alkyl}$;
- k) $\text{C}(=\text{W})\text{-OR}^2$;
- l) CN ;
- m) NO_2 ;
- n) $\text{NH}-\text{C}(=\text{W})\text{alkyl}$;
- o) $\text{NH}-\text{S}(\text{O})_n\text{-alkyl}$;
- p) $\text{NH}-\text{S}(\text{O})_n\text{-NR}^2\text{R}^2$; or
- q) C_{3-6} cycloalkyl;

W is:

- a) O ;
- b) S ;
- c) NH ;
- d) $\text{N}-\text{N}(\text{R}^2)(\text{R}^2)$;
- e) $\text{N}(\text{R}^2)$;
- f) $\text{N}-\text{OH}$; or
- g) $\text{N}-\text{O-alkyl}$;

R^1 is:

- a) H ;
- b) R^2 ;
- c) $\text{C}(=\text{W})\text{-R}^2$;
- d) $\text{C}(=\text{W})\text{-O}(\text{R}^2)$;

- e) $C(=W)-S(R^2)$;
- f) $C(=W)-NH-R^2$;
- g) $C(=W)-N(R^2)(R^2)$;
- h) $C(=W)-NH-A-(\text{amino acid residue})$;
- i) $A-(\text{amino acid residue})-R^2$;
- j) $S(O)_n-R^3$; or
- k) $S(O)_2-N(R^2)(R^2)$;

any of which optionally may be substituted by one or more:

- a) C_{1-6} alkyl;
- b) OH;
- c) alkoxy;
- d) aryl;
- e) halo;
- f) CN;
- g) NO_2 ; or
- h) $N(R^2)(R^2)$;

R^2 is:

- a) H;
- b) OH;
- c) halogen;
- d) optionally substituted, branched or unbranched alkyl;
- e) optionally substituted, branched or unbranched alkenyl;
- f) optionally substituted, branched or unbranched alkynyl;
- g) 3-14 membered carbocycle;

- h) alkylheterocycle;
- i) acyl;
- j) carboxamido;
- k) carbamoyl;
- l) alkoxy;
- m) optionally substituted aryl;
- n) optionally substituted aralkyl;
- o) optionally substituted alkylaryl;
- p) O-alkyl;
- q) O-alkenyl;
- r) O-alkynyl;
- s) O-alkaryl;
- t) O-aralkyl;
- u) O-carbocycle;
- v) O-heterocycle;
- w) O-aryl;
- x) CF_3 ;
- y) CN;
- z) $\text{S(O)}_n\text{-R}^3$;
- aa) $\text{N(R}^3\text{)(R}^3\text{)}$;
- bb) $\text{NH-S(O)}_n\text{-R}^3$;
- cc) NHC(=W)-aryl ;
- dd) NHC(=W)-alkyl ;
- ee) $\text{NHC(=W)-heterocycle}$;

- ff) $\text{CH}_2\text{-S(O)}_n\text{R}^3$;
- gg) C(=W)R^3 ;
- hh) $\text{C(=W)NR}^3\text{R}^3$;
- ii) $\text{C(alkyl)}_2\text{-S(O)}_n\text{R}^3$;
- jj) $\text{CH(alkyl)-S(O)}_n\text{R}^3$;
- kk) $\text{C(alkyl)}_2\text{-NH}_2$;
- ll) $\text{CH(alkyl)-N(alkyl)R}^3$;
- mm) $\text{CR}^3\text{R}^3\text{-NR}^3\text{R}^3$;
- nn) $\text{CH}_2\text{N(alkyl)R}^3$;
- oo) CH(alkyl)-NHR^3 ;
- pp) $\text{C(alkyl)}_2\text{-NHR}^3$;
- qq) $\text{C(alkyl)}_2\text{-N(alkyl)R}^3$;
- rr) $\text{CH}_2\text{-C(=W)H}$;
- ss) $\text{CH}_2\text{-C(=W)alkyl}$;
- tt) $\text{CR}^3\text{R}^3\text{-C(=W)R}^3$;
- uu) A-R^3 ;
- vv) $\text{C(R}^3)_2\text{-C(=W)R}^3$;
- ww) $\text{CH}_2\text{-C(=W)H}$;
- xx) $\text{CH}_2\text{-C(=W)alkenyl}$;
- yy) $\text{CH(alkenyl)-C(=W)H}$;
- zz) A-S(O)R^3 ;
- aaa) $\text{CH(NH)-S(O)}_n\text{R}^3$; or
- bbb) A-N(NH)R^3 ;

wherein said optional substitution comprises one or more of:

- a) a substituted or unsubstituted heterocycle;
- b) $C(=W)O$ -aryl;
- c) $C(=W)O$ -alkyl;
- d) $C(=W)NH_2$;
- e) $C(=W)NH$ -alkyl;
- f) $C(=W)NH$ -aryl;
- g) $C(=W)N$ -di-alkyl;
- h) $C(=W)N$ (alkyl)-aryl;
- i) α -amino acid;
- j) α -amino ester;
- k) α -amino-carboxamide;
- l) β -amino acid;
- m) β -amino ester; or
- n) β -amino carboxamide;

R^3 is:

- a) H;
- b) OH;
- c) C_{1-6} alkyl;
- d) C_{2-6} alkenyl;
- e) C_{2-6} alkynyl;
- f) alkoxy;
- g) CF_3 ;
- h) CN;
- i) amino;

- j) NR^2R^2 ;
- k) O-alkyl;
- l) O-alkenyl;
- m) O-alkynyl;
- n) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{NH}_2$;
- o) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{CF}_3$;
- p) $\text{C}(\text{R}^2)(\text{R}^2)\text{-NH}_2$;
- q) A-heterocycle;
- r) $\text{C}(\text{R}^2)(\text{R}^2)\text{-NR}^2\text{R}^2$;
- s) $\text{C}(\text{R}^2)(\text{R}^2)\text{-C}(=\text{W})\text{R}^2$;
- t) aryl;
- u) carbocycle;
- v) heterocycle;
- w) cycloalkyl;
- x) alkaryl;
- y) alkylheterocycle;
- z) aralkyl; or
- aa) heterocycle-alkyl;

any of which may be unsubstituted or substituted with one or more of the following, taken in any combination:

- a) halo;
- b) OH;
- c) OR^2 ;
- d) SR^2 ;

- e) COOH ;
- f) carboxylic acid ester;
- g) $\text{C}(=\text{W})\text{R}^2$;
- h) $\text{C}(=\text{W})\text{OR}^2$;
- i) $\text{C}(=\text{W})\text{OR}^3$;
- j) $\text{C}(=\text{W})\text{SR}^2$;
- k) $\text{A}-\text{C}(=\text{W})\text{NH}_2$;
- l) $\text{C}(=\text{W})\text{NR}^2\text{R}^3$;
- m) NR^2R^2 ;
- n) NR^2R^3 ;
- o) $\text{NR}^2-\text{S}(\text{O})_n\text{R}^3$;
- p) $\text{NR}^2-\text{C}(=\text{W})-\text{C}_{1-6}\text{alkyl}$;
- q) $\text{S}(\text{O})_n\text{R}^3$;
- r) $\text{C}_{1-6}\text{alkoxy}$;
- s) $\text{C}_{1-6}\text{thioether}$;
- t) amino acid residue;
- u) $\text{NH}-\text{A}-(\text{amino acid residue})$;
- v) $\text{C}(=\text{W})\text{NH}-\text{A}-(\text{amino acid residue})$; and

wherein when said optional substitution comprises a substituted heterocycle, then substitution is selected from the group consisting of:

- a) $\text{C}(=\text{W})\text{O-aryl}$;
- b) $\text{C}(=\text{W})\text{O-alkyl}$;
- c) $\text{C}(=\text{W})\text{NH}_2$;
- d) $\text{C}(=\text{W})\text{NH-aryl}$;

- e) $C(=W)NH$ -alkyl;
- f) $C(=W)N$ -di-alkyl;
- g) $C(=W)N$ (alkyl)-aryl;
- h) α -amino acid;
- i) α -amino ester;
- j) α -amino-carboxamide;
- k) β -amino acid;
- l) β -amino ester; or
- m) β -amino-carboxamide;
- n) halo; or
- o) cyano,

taken alone or in any combination;

n independently is 0, 1 or 2;

each A is independently a disubstituted spacer selected from the group consisting of:

- a) C_{1-6} alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- b) C_{2-12} alkenylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- c) C_{2-12} alkynylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- d) optionally substituted arylene;

- e) O-alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- f) aralkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- g) optionally substituted cycloalkyl; and
- h) optionally substituted heterocycle;

wherein "A" may be joined by any desired linkage such as, for example, an ether, thioether, amino, carboxamido, ester or carbon-carbon linkage, or any combination thereof;

Each $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ independently is:

- a) H;
- b) halogen;
- c) NO_2 ;
- d) CN;
- e) CF_3 ;
- f) OR^2 ;
- g) NR^2R^2 ;
- h) $NHS(O)_nR^2$;
- i) $NHCO-C_{1-3}$ alkyl;
- j) $S(O)_nR^2$;
- k) aryl;
- l) heterocycle;
- m) C_{1-6} alkyl;
- n) C_{2-6} alkenyl;

- o) C_{2-6} alkynyl;
- p) $C(=W)-S(O)_nR^2$;
- q) $C(=W)-S(O)_n-NR^2R^2$;
- r) $C(=W)$ -aryl;
- s) $C(=W)$ -alkyl;
- t) $C(=W)$ -heterocycle; or
- u) $C(=W)-NR^2R^2$;

each of which optionally may be substituted with one or more of :

- a) OR^2 ;
- b) $S(O)_nR^2$;
- c) $C(=W)-S(O)_nR^2$;
- d) $C(=W)-S(O)_n-NR^2R^2$;
- e) $C(=W)$ -aryl;
- f) $C(=W)$ -alkyl;
- g) $C(=W)$ -heterocycle;
- h) $C(=W)NR^2R^2$;
- i) H;
- j) NO_2 ;
- k) CN;
- l) CF_3 ;
- m) halogen;
- n) $NHS(O)_nR^2$;
- o) $NHCO-C_{1-3}$ alkyl;
- p) aryl;

- q) heterocycle;
- r) C₁₋₆ alkyl;
- s) C₂₋₆ alkenyl;
- t) C₂₋₆ alkynyl; or
- u) NR²R².

In one embodiment of Formula (A), X is an optionally substituted phenyl; Y is any of its definitions; R¹, R^{4'}, R^{6'}, and R^{7'} are all hydrogen; R^{5'} is halogen; and Z is a carboxamide moiety.

In an alternative embodiment, R¹ is acyl, alkyl, aryl, alkaryl, or aralkyl.

In yet another alternative embodiment, R^{4'} is fluoro, nitro or cyano, W is oxygen, and Y is O-alkyl.

In a second embodiment of Formula (A), X is an optionally substituted phenyl; Y is any of its definitions; R¹, R^{4'}, R^{6'}, and R^{7'} are all hydrogen; R^{5'} is chlorine; and Z is a carboxamide or carboxamido-heterocyclyl moiety.

In yet another embodiment of Formula (A), X is tolyl, thiazolyl or pyridyl; Y is H, OH, or O-alkyl; R¹, R^{4'}, R^{6'}, and R^{7'} are all hydrogen; R^{5'} is halogen; and Z is carboxamido, acyl, an alkyl-sulphonyl group or a carboxylic acid derivative.

In an alternative to the preceding embodiment, X, Y, R¹, R^{4'}, R^{6'}, R^{7'} and R^{5'} all are as defined above, and Z is a carboxamido-alkylene-heterocyclyl wherein the heterocycle typically is imidazole, furan, pyridine, pyrimidine, or thiazole that optionally may be substituted; a thioamido-pyridyl wherein the pyridyl is unsubstituted or substituted by OH, OMe or lower alkyl; an imino-nitrile, or an alkylsulphonyl-aryl group.

A first series of subembodiments of the present invention is given where the Formula (A) is as provided above, and W, X and Y are defined as:

- a) W is O, X is alkyl, and Y is -O-alkyl;
- b) W is O, X is -O-aryl, and Y is alkyl;
- c) W is O, X is -O-aryl, and Y is $-NR^2R^3$;
- d) W is O, X is -O-alkyl, and Y is alkyl;
- e) W is O, X is -O-alkyl, and Y is halo;
- f) W is O, X is -O-heterocycle, and Y is alkyl;
- g) W is O, X is aryl, and Y is -O-alkyl;
- h) W is O, X is heterocyclyl, and Y is -O-aryl;
- i) W is O, X is alkyl, and Y is -O-heterocyclyl;
- j) W is $-NR^2R^2$, X is heterocyclyl, and Y is -O-aryl;
- k) W is $-NR^2R^2$, X is alkyl, and Y is halo;
- l) W is S, X is alkyl, and Y is -O-alkyl;
- m) W is S, X is alkyl, and Y is $-NR^2R^3$;
- n) W is S, X is -O-aryl, and Y is alkyl;
- o) W is S, X is -O-aryl, and Y is C-halo;
- p) W is S, X is -O-alkyl, and Y is alkyl;
- q) W is S, X is -O-heterocycle, and Y is alkyl;
- r) W is S, X is aryl, and Y is -O-alkyl;
- s) W is S, X is heterocyclyl, and Y is -O-aryl;
- t) W is S, X is alkyl, and Y is -O-heterocyclyl;
- u) W is O, X is aryl, and Y is -O-aryl;
- v) W is $-NR^2$, X is -O-alkyl, and Y is $-NR^2R^3$;
- w) W is O, X is -O-aryl, and Y is -O-aryl;
- x) W is O, X is alkyl, and Y is alkyl; and

- y) W is $-\text{NR}^2$, X is $-\text{O-alkyl}$, and Y is alkyl.

A second series of subembodiments of the present invention is given where Formula (A) is as given, R^1 is H, alkyl, acyl, aryl, aralkyl, or alkaryl; and Z is defined as:

- a) $-\text{C}(=\text{W})\text{NR}^2\text{R}^3$, R^2 is H, and R^3 is NR^2R^2 ;
- b) $-\text{C}(=\text{W})\text{NR}^2\text{R}^3$, R^2 is H, and R^3 is NR^2R^2 , or R^2 is C_{1-5} alkyl optionally substituted with OH and R^3 is $-\text{NH}_2$;
- c) $-\text{C}(=\text{W})\text{NR}^2\text{R}^3$, R^2 is H, and R^3 is $(\text{CH}_2)_m\text{C}(=\text{W})\text{NR}^2\text{R}^2$;
- d) $-\text{C}(=\text{W})\text{NR}^2\text{R}^3$, R^2 is H, and R^3 is alkyl substituted by optionally substituted aryl or heterocycle;
- e) $-\text{C}(=\text{W})\text{R}^3$, R^3 is an amino acid residue or $-\text{NH}(\text{CH}_2)_p\text{-(amino acid residue)}$;
- f) $-\text{C}(=\text{W})\text{NHNHC}_2\text{H}_5\text{OH}$;
- g) $-\text{C}(=\text{W})\text{NHCH}_2\text{C}(=\text{W})\text{NH}_2$;
- h) $-\text{C}(=\text{W})\text{NHCH}_2\text{CONHNH}_2$;
- i) $-\text{C}(=\text{W})\text{NHCH}_2\text{CH}_2\text{-(2-NO}_2\text{, 5-methyl imidazole)}$;
- j) $-\text{C}(=\text{W})\text{NHCH}_2\text{NHCH}(\text{CH}_3)\text{C}(=\text{W})\text{OH}$;
- k) $-\text{C}(=\text{W})\text{NHCH}=\text{CHC}(=\text{W})\text{NH}_2$;
- l) $-\text{C}(=\text{W})\text{NR}^2\text{R}^5\text{NR}^2\text{R}^3$, R^5 is $(=\text{O})$, R^2 and R^3 are as defined above;
- m) $-\text{C}(=\text{W})\text{NR}^2\text{NR}^2\text{-C}(=\text{W})\text{R}^3$, R^2 is H or alkyl, and R^3 is aryl;
- n) $-\text{C}(=\text{W})\text{N}(-\text{NR}^2\text{R}^3)\text{-N}(-\text{NR}^2\text{R}^3)\text{R}^3$, R^2 is H, R^3 is R^2 or alkoxy;
- o) $-\text{C}(=\text{W})\text{NHR}^2\text{C}(=\text{W})\text{-Q}$, Q is heterocycle, and R^2 is as defined above;
- p) $-\text{C}(=\text{W})\text{NR}^2\text{R}^3$, R^2 is as defined above, and R^3 is $-\text{OH}$;
- q) $-\text{COR}^2\text{R}^3$, R^2 is amino and R^3 is a heterocycle;
- r) $-\text{C}(=\text{W})\text{NHNHC}(=\text{W})\text{R}^2$ and R^2 is NH_2 ;
- s) $-\text{C}(=\text{W})\text{-R}^2\text{-CH-A-C}(=\text{W})\text{NH}_2$, and R^2 is NH;

- t) $-\text{C}(=\text{W})-\text{R}^2-\text{CH}-\text{A}-\text{C}(=\text{W})\text{H}$, and R^2 is NH ;
- u) $-\text{C}(=\text{W})-\text{R}^2-\text{CH}-\text{A}-\text{C}(=\text{W})\text{OH}$, and R^2 is NH ;
- v) $-\text{C}(=\text{W})-\text{R}^2-\text{CH}-\text{A}-\text{R}^3$, R^2 is NH , and R^3 is CH_3 ;
- w) $-\text{C}(=\text{W})\text{NHR}^2\text{C}(=\text{W})\text{NH}_2$, and R^2 is optionally substituted, branched chain alkyl;
- x) $-\text{C}(=\text{W})\text{R}^2\text{R}^3$, R^2 is NH or alkyl, and R^3 is NH_2 ;
- y) $-\text{C}(=\text{W})\text{R}^2-\text{C}(=\text{W})\text{OR}^3$, and R^2 and R^3 are as defined as above;
- z) $-\text{C}(=\text{W})\text{R}^2-\text{NH}-\text{C}(=\text{W})\text{C}_{1-4}$ alkoxy, and R^2 is as defined above;
- aa) $-\text{C}(=\text{W})\text{R}^2\text{C}(=\text{W})\text{C}_{1-4}$ alkoxy, and R^2 is as defined above;
- bb) $-\text{C}(=\text{W})\text{R}^2$, and R^2 is NH_2 ;
- cc) $-\text{C}(=\text{W})\text{R}^2-\text{NH}-\text{C}(=\text{W})\text{OR}^3$, and R^2 and R^3 are as defined as above;
- dd) $-\text{C}(=\text{W})\text{R}^2-\text{C}(=\text{W})\text{R}^2$, and R^2 is as defined above;
- ee) $-\text{C}(=\text{W})\text{NHR}^2$ where R^2 is optionally substituted aryl, cycloalkyl or a heterocyclyl ring;
- ff) $-\text{C}(=\text{W})\text{R}^2-\text{W}-\text{R}^3$, where R^2 and R^3 are as defined as above;
- gg) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(\text{R}^2)-\text{C}(=\text{W})-\text{NH}_2$, and R^2 is as defined above;
- hh) $-\text{C}(=\text{W})-\text{NH}-\text{NH}_2$;
- ii) $-\text{C}(=\text{W})-\text{NH}-\text{NH}(\text{R}^2)$, and R^2 is as defined above;
- jj) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(\text{C}[=\text{W}]\text{NH}_2)(\text{CH}_2-\text{C}[=\text{W}]-\text{O}-\text{aryl})$;
- kk) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(-[\text{CH}_2]_4-\text{NH}-\text{C}[=\text{W}]-\text{t-BuO})(-\text{C}[=\text{W}]-\text{NH}_2)$;
- ll) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(-\text{CH}_2-\text{CH}_2-\text{C}[=\text{W}]-\text{t-BuO})(\text{C}[=\text{W}]-\text{NH}_2)$;
- mm) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(\text{R}^3)(-\text{C}[=\text{W}]-\text{NH}_2)$, and R^3 is as defined above;
- nn) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(-\text{CH}_2-\text{R}^3)(-\text{C}[=\text{W}]-\text{NH}_2)$, and R^3 is as defined above;
- oo) $-\text{C}(=\text{W})-\text{NH}-\text{CH}(-\text{CH}_2\text{OH})(-\text{C}[=\text{W}]-\text{NH}_2)$;

- pp) $-C(=W)-NH-CH(C[=W]-NH_2)(C[=W]-NH_2)$;
- qq) $-C(=W)-NHR^2-C[=W]NH_2$, and R^2 is as defined above;
- rr) $-C(=W)-NH-CH(-[CH_2]_4-NH-C[=W]-O-CH_2-R^3)(-C[=W]-NH_2)$, and R^3 is as defined above;
- ss) $-C(=W)-NH-CH(-CH_2-C[=W]-NH_2)(-C[=W]-NH_2)$;
- tt) $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined above;
- uu) $-C(=W)-NH-CH(-[CH_2]_4-NH_2)(-C[=W]-NH_2)$;
- vv) $-C(=W)-NH-CH(-CH[R^2][OH])(-C[=W]-NH_2)$, and R^2 is as defined above;
- ww) $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, and R^2 is as defined above;
- xx) $-C(=W)-NH-CH(-R^2-C[=W]-NH_2)(-C[=W]-NH_2)$, and R^2 is as defined above;
- yy) $-C(=W)-NH-CH(-R^2-SCH_3)(-C[=W]-NH_2)$, and R^2 is as defined above;
- zz) $-C(=W)-NH-CH(-C[=N]-NH_2)(-C[=W]-NH_2)$;
- aaa) $-C(=W)-NH-CH(-R^3)(-C[=W]-NH_2)$, and R^3 is as defined above;
- bbb) $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, and R^3 is as defined above;
- ccc) $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, and R^2 is as defined above;
- ddd) $-C(=W)-NH-R^2-C[=W]R^3$, and R^2 is alkylene and R^3 is aryl or heteroaryl;
- eee) $-C(=W)-NH-R^2-R^3-C[=W]-NH_2$, where R^2 is alkylene and R^3 is aryl or heteroaryl;
- fff) $-C(=W)-NH-NH-R^2-R^3-C(=W)NH_2$, where R^2 is alkylene and R^3 is aryl or heteroaryl;
- ggg) $-C(=W)-NH-NH-CH(R^3)-C(=W)R^2$, and R^2 is NH_2 and R^3 is optionally substituted aryl or heteroaryl;
- hhh) $-C(=W)NHR^2(-R^3)-C(=W)NH_2$, where R^2 is alkylene and R^3 is optionally substituted alkyl, aryl or heteroaryl;
- iii) $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)OH$, where R^2 is alkylene and R^3 is optionally substituted alkyl, aryl or heteroaryl;

- jjj) $-\text{C}(=\text{W})\text{NHR}^2(-\text{R}^3)-\text{C}(=\text{W})\text{NH}-\text{R}^2-\text{NH}_2$, where R^2 is alkylene and R^3 is optionally substituted alkyl, aryl or heteroaryl;
- kkk) $-\text{C}(=\text{W})\text{NHR}^2(-\text{R}^3)-\text{C}(=\text{W})\text{NH}-\text{R}^3$, where R^2 is alkylene and R^3 is optionally substituted alkyl, aryl or heteroaryl;
- lll) $-\text{C}(=\text{W})-\text{R}^2-(\text{CH}_2)_p-\text{A}-\text{C}(=\text{W})-\text{NH}_2$, where R^2 is $-\text{NH}$, p is 0-10, A is a divalent linker or an optionally substituted aryl or heteroaryl, and W is O or S ;
- mmm) $-\text{C}(=\text{W})\text{NH}-\text{R}^3$, where R^3 is an optionally substituted heterocycle;
- nnn) $-\text{C}(=\text{W})-\text{NH}-\text{R}^2-\text{R}^5-\text{R}^3$, where W is O or S , R^2 is alkylene, alkenylene or alkynylene, R^5 is $-\text{SO}_2$, and R^3 is $-\text{NH}_2$;
- ooo) $-\text{C}(=\text{W})-\text{NH}-\text{NH}-\text{R}^2(\text{R}^3)-\text{R}^5-\text{NH}_2$, where W is O or S , R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene, or heteroaryl, and R^5 is $-\text{SO}_2$;
- ppp) $-\text{C}(=\text{W})-\text{NH}-\text{R}^3(\text{R}^5-\text{NH}_2)$ where W is O or S , R^3 is aryl, arylene or heteroaryl, and R^5 is SO_2 ;
- qqq) $-\text{C}(=\text{W})-\text{NH}-\text{R}^2-\text{R}^3(\text{R}^5-\text{NH}_2)$ where W is O or S , R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heteroaryl, and R^5 is SO_2 ;
- rrr) $-\text{C}(=\text{W})-\text{NH}-\text{R}^3(\text{R}^2\text{R}^5-\text{NH}_2)$ where W is O or S , R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heteroaryl, and R^5 is SO_2 ;
- sss) $-\text{C}(=\text{W})-\text{NHR}^2(-\text{R}^3)-\text{C}(=\text{W})\text{NH}-\text{R}^2-\text{C}(=\text{W})\text{OH}$, where R^2 is an optionally substituted alkylene, alkenylene, or alkynylene, and R^3 is any of its definitions as provided above;
- ttt) $-\text{C}(=\text{W})-\text{NHR}^2(-\text{R}^3)-\text{C}(=\text{W})\text{NH}-\text{R}^2-\text{C}(=\text{W})\text{NH}_2$, where R^2 is an optionally substituted alkylene, alkenylene, or alkynylene, and R^3 is any of its definitions as provided above;
- uuu) $-\text{C}(=\text{W})-\text{NHR}^2-\text{C}(=\text{W})\text{NH}-\text{R}^2-\text{C}(=\text{W})\text{OH}$, where R^2 is an optionally substituted alkylene, alkenylene, or alkynylene, and R^3 is any of its definitions as provided above; or

vvv) $-C(=W)-NHR^2-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is an optionally substituted alkylene, alkenylene, or alkynylene, and R^3 is any of its definitions as provided above; and

wherein R^2 , R^3 and A each is as defined previously.

A third series of subembodiments of the present invention is given for the Formula (A) as provided above where $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are defined as:

- a) $R^{6'}$ and $R^{7'}$ are both hydrogen, and $R^{4'}$ and $R^{5'}$ independently are halo; $-NO_2$; $-N$; $-OR^2$; $-NR^2R^2$; $-NH-R^5-C_{1-3}$ alkyl; $-NHCO-C_{1-3}$ alkyl; oxime; hydrazine; $-H(SO_2)C_{1-3}$ alkyl; $-NH-O-C_{1-3}$ alkyl; $-NHOH$; or C_{1-3} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-C(=W)OH$, halogen, $-NR^2R^2$, C_{1-3} alkoxy, or C_{1-3} thioether;
- b) $R^{4'}$ and $R^{7'}$ are both hydrogen, and $R^{5'}$ and $R^{6'}$ independently are halo; $-NO_2$; $-N$; $-OR^2$; $-NR^2R^2$; $-NHSO_2-C_{1-6}$ alkyl; $-NHCO-C_{1-6}$ alkyl; oxime; hydrazine; $-H-O-C_{1-6}$ alkyl; $-NH-OH$; or C_{1-3} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-(=W)OH$, halogen, $-NR^2R^2$, C_{1-3} alkoxy, or C_{1-3} thioether;
- c) $R^{4'}$ and $R^{6'}$ are both hydrogen, and $R^{5'}$ and $R^{7'}$ independently are halo; $-NO_2$; $-N$; $-OR^2$; $-NR^2R^2$; $-NHSO_2-C_{1-6}$ alkyl; $-NHCO-C_{1-6}$ alkyl; oxime; hydrazine; $-H-O-C_{1-6}$ alkyl; $-NH-OH$; or C_{1-6} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-(=W)OH$, halogen, $-NR^2R^2$, C_{1-3} alkoxy, or C_{1-3} thioether;
- d) $R^{4'}$ and $R^{7'}$ are both hydrogen, and $R^{5'}$ and $R^{6'}$ independently are halo; $-NO_2$; $-N$; $-OR^2$; $-NR^2R^2$; $-NH-O-C_{1-6}$ alkyl; $-NHOH$; or C_{1-6} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-C(=W)OH$, halogen or $-NH_2$;
- e) $R^{4'}$ and $R^{6'}$ are both hydrogen, and $R^{5'}$ and $R^{7'}$ independently are halo; $-NO_2$; $-N$; $-OR^2$; $-NR^2R^2$; $-NH-O-C_{1-3}$ alkyl; $-NHOH$; or C_{1-6} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-C(=W)OH$, halogen or $-NH_2$;

- f) $R^{6'}$ and $R^{7'}$ are both hydrogen, and $R^{4'}$ and $R^{5'}$ independently are halo; $-C_{1-6}$ alkyl, alkenyl or alkynyl optionally substituted with one or more halo;
- g) $R^{4'}$ and $R^{7'}$ are both hydrogen, and $R^{5'}$ and $R^{6'}$ independently are halo; $-C_{1-6}$ alkyl, alkenyl or alkynyl optionally substituted with one or more halo;
- h) $R^{4'}$ and $R^{6'}$ are both hydrogen, and $R^{5'}$ and $R^{7'}$ independently are halo, $-C_{1-6}$ alkyl, alkenyl or alkynyl optionally substituted with one or more halo;
- i) $R^{6'}$ and $R^{7'}$ are both hydrogen, and $R^{4'}$ and $R^{5'}$ independently are Cl, F, Br, I, methyl, ethyl or CF_3 ;
- j) $R^{4'}$ and $R^{7'}$ are both hydrogen, and $R^{5'}$ and $R^{6'}$ independently are Cl, F, Br, I, methyl, ethyl or CF_3 ;
- k) $R^{4'}$ and $R^{6'}$ are both hydrogen, and $R^{5'}$ and $R^{7'}$ independently are Cl, F, Br, I, methyl, ethyl or CF_3 ; and

wherein A is as defined above.

A fourth series of subembodiments for Formula (A) is defined where W, X and Y are as defined in the first series of subembodiments, and Z is as defined in the second series of subembodiments.

A fifth series of subembodiments for Formula (A) is defined where W, X and Y are as defined in the first series of subembodiments, and $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are as defined in the third series of subembodiments

A sixth series of subembodiments for Formula (A) is defined where Z is as defined in the second series of subembodiments, and $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are as defined in the third series of subembodiments.

Non-limiting species of the first embodiment as given by Formula (A) above are defined when:

- 1) Z is $-C(=W)NHNHC_2H_5OH$, $R^{4'}$, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is Cl, X is H, and Y is H;
- 2) Z is $-C(=W)NHCH_2C(=W)NH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is OH;

- 3) Z is $-C(=W)NHCH_2CONHNH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -alkyl;
- 4) Z is $-C(=W)NHCH_2CH_2-(2-NO_2, 5-Me-imidazole)$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -alkenyl;
- 5) Z is $-C(=W)NHCH_2NHCH(CH_3)C(=W)OH$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -alkynyl;
- 6) Z is $-C(=W)CH=CHC(=W)NH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -alkaryl;
- 7) Z is $-C(=W)NHNHCH_2C(=W)NH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -aryl;
- 8) Z is $-C(=W)NHCH_2C(=W)R^2$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is $-O$ -heterocycle;
- 9) Z is $-C(=W)NHCH_2-A-C(=W)NH_2$, where A is a divalent spacer, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is C_{1-6} alkyl;
- 10) Z is $-C(=W)R^2CH-A-C(=W)H$, where A is a divalent spacer, R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is C_{2-6} alkynyl;
- 11) Z is $-C(=W)R^2CH-A-C(=W)OH$, where A is a divalent spacer, R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is C_{2-6} alkynyl;
- 12) Z is $-C(=W)R^2-CH-A-R^3$, where A is a divalent spacer, R^2 is NH , R^3 is CH_3 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is aryl;
- 13) Z is $-C(=W)NHR^2-C(=W)R^2$, where R^2 is an optionally substituted, branched chain alkylene or NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is H, and Y is heterocycle;
- 14) Z is $-C(=W)R^2$, where R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is OH, and Y is H;
- 15) Z is $-C(=W)R^2R^3$ -heterocycle, where R^2 is NH , R^3 is CH_2 , heterocycle is optionally substituted morpholine, imidazole or pyrrole, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is C_{1-6} alkyl, and Y is H;

16) Z is $-C(=W)R^2C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyloxy, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -alkyl, and Y is H;

17) Z is $-C(=W)R^2-NH-C(=W)-C_{1-4}$ alkoxy, where R^2 is an optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -alkenyl, and Y is H;

18) Z is $-C(=W)R^3-C(=W)R^2$, where R^2 is C_{1-4} alkoxy, R^3 is optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -alkynyl, and Y is H;

19) Z is $-C(=W)R^2R^3$, where R^2 is optionally substituted alkyl, R^3 is optionally substituted phenyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -alkaryl, and Y is H;

20) Z is $-C(=W)R^2-NH-C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -aralkyl, and Y is H;

21) Z is $-C(=W)R^2C(=W)-NH_2$, where R^2 is optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -aryl, and Y is H;

22) Z is $-C(=W)R^2-W-R^3$, where R^2 and R^3 are optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-O$ -heterocycle, and Y is H;

23) Z is $-C(=W)R^2C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyloxy, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-C_{2-6}$ -alkenyl, and Y is H;

24) Z is $-C(=W)-NH-CH(R^2)-C(=W)-NH_2$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is $-C_{2-6}$ -alkynyl, and Y is H;

25) Z is $-C(=W)-NH-NH_2$, where $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is aryl, and Y is H;

26) Z is $-C(=W)-NH-NH(R^2)$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, X is heterocycle, and Y is H;

27) Z is $-C(=W)-NH-CH(C[=W]NH_2)(CH_2-C[=W]-O-CH_2-aryl)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is carbocycle, and Y is H;

28) Z is $-C(=W)-NH-CH(-[CH_2]_4-NH-C[=W]-t-BuO)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkaryl, and Y is C_{1-6} alkyl;

- 29) Z is $-C(=W)-NH-CH(-CH_2-CH_2-C[=W]-t-BuO)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -aryl, and Y is $-O$ -alkyl;
- 30) Z is $-C(=W)-NH-CH(-CH_2R^3)(-C[=W]-NH_2)$, where R^3 is CF_3 , $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{2-6}$ alkenyl, and Y is $-OH$;
- 31) Z is $-C(=W)-NH-CH(-CH_2R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{2-6}$ alkynyl, and Y is H;
- 32) Z is $-C(=W)-NH-CH(-CH_2OH)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkylaryl, and Y is $-O$ -alkyl;
- 33) Z is $-C(=W)-NH-CH(-C[=W]-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-heterocycle$, and Y is $-OH$;
- 34) Z is $-C(=W)-NH-CH(-R^3)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkenyl, and Y is $-O$ -alkyl;
- 35) Z is $-C(=W)-NH-CH(-[CH_2]_4-NH-C[=W]-O-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O$ -alkylene;
- 36) Z is $-C(=W)-NH-CH(-CH_2-C[=W]-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-carbocycle$, and Y is $-O$ -alkyl;
- 37) Z is $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkyl, and Y is $-heterocycle$;
- 38) Z is $-C(=W)-NH-CH(-[CH_2]_4-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkenyl, and Y is $-OH$;
- 39) Z is $-C(=W)-NH-CH(-CH[R^2][OH])(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkynyl, and Y is $-O$ -alkyl;
- 40) Z is $-C(=W)-NH-CH(R^2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O$ -alkylene;

- 41) Z is $-C(=W)-NH-CH(-R^2-C[=W]-NH_2)(-C[=W]-NH_2)$, where R^2 is NH_2 , $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{2-6}$ alkylene, and Y is $-OH$;
- 42) Z is $-C(=W)-NH-CH(C[=NH]-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-carbocycle$, and Y is $-O-alkyl$;
- 43) Z is $-C(=W)-NH-CH(-C[=NH]-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O-alkylene$;
- 44) Z is $-C(=W)-NH-CH(-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O-aryl$, and Y is $-C_{1-6}$ alkyl;
- 45) Z is $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O-alkylene$;
- 46) Z is $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, where R^2 is NH_2 , $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{1-6}$ alkynyl, and Y is $-O-aryl$;
- 47) Z is $-C(=W)-NH-NH-CH(R^3)-C[=W]R^2$, where R^2 is $-NH_2$, R^3 is optionally substituted aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O-alkylaryl$, and Y is $-OH$;
- 48) Z is $-C(=W)-NHR^2(-R^3)-C[=W]NH_2$, where R^2 is $-alkylene$, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O-alkynyl$, and Y is $-OH$;
- 49) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)OH$, where R^2 is $alkylene$, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O-alkenyl$, and Y is $-O-alkyl$;
- 50) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-NH_2$, where R^2 is $alkylene$, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is H, and Y is $-O-alkyl$;
- 51) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^3-C(=W)OH$, where R^2 is $alkylene$, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O-alkylene$;

- 52) Z is $-C(=W)R^2(CH_2)_p-A-C(=W)NH_2$, where R^2 is $-NH$, p is 0-10, A is divalent, optionally substituted aryl or heterocycle, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{1-6}$ alkyl, and Y is $-O$ -alkenyl;
- 53) Z is $-C(=W)NH-R^3$, where R^3 is optionally substituted heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkyl, and Y is $-OH$;
- 54) Z is $-C(=W)NH-R^2-R^5-R^3$, where R^2 is alkylene, alkenylene or alkynylene, R^5 is $-SO_2$, R^3 is $-NH_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -aryl, and Y is $-O$ -alkyl;
- 55) Z is $-C(=W)NH-NH-R^2(R^3)-R^5-NH_2$, where R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heterocyclyl, R^5 is $-SO_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-O$ -alkynyl, and Y is $-OH$;
- 56) Z is $-C(=W)NH-R^3(R^5-NH_2)$, where R^3 is aryl, arylene or heterocyclyl, R^5 is $-SO_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-heterocyclyl$, and Y is $-O$ -alkyl;
- 57) Z is $-C(=W)NH-R^2R^3(R^5-NH_2)$, where R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heterocyclyl, R^5 is $-SO_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-aryl$, and Y is $-O$ -alkenyl;
- 58) Z is $-C(=W)NH-R^3(R^2R^5-NH_2)$, where R^2 is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heterocyclyl, R^5 is $-SO_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-OH$, and Y is $-O$ -alkylaryl;
- 59) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)OH$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, R^3 is any of the definitions provided in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{2-6}$ alkynyl, and Y is $-O$ -alkenyl;
- 60) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, R^3 is any of the definitions provided in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{2-6}$ alkenyl, and Y is $-H$;


61) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)OH$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-C_{1-6}$ alkyl, and Y is $-H$;

62) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is $-H$, and Y is $-O$ -alkyl;

63) Z is $-C(=W)NH-R^3$ where W is O, and R^3 is a heterocycle, and in particular pyridyl, pyrimidyl or imidazo that is optionally substituted by one or more halo, cyano, alkyl, alkenyl, alkynyl, or cyanoalkyl groups; $R^{4'}$ and $R^{5'}$ independently are H, F or Cl, $R^{6'}$ and $R^{7'}$ are H, X is optionally substituted phenyl, and Y is $-O$ -alkyl; and

wherein W is as defined for Formula (A).

In a second embodiment, the invention provides a phenylindole for use in the treatment of HIV represented by the following general Formula (B):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

each W, Y, Z, R^1 , R^2 , R^3 , A, n, $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ is as defined above for Formula (A); and

each $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ independently is:

- a) H;
- b) halogen;

- c) NO_2 ;
- d) CN ;
- e) OR^2 ;
- f) SR^2 ;
- g) NH_2 ;
- h) NR^2R^3 ;
- i) $\text{N(R}^2\text{)-C(=W)-C}_{1-4}\text{ alkyl}$;
- j) $\text{N(R}^2\text{)-SO}_2\text{-C}_{1-4}\text{ alkyl}$;
- k) $\text{C}_{1-6}\text{ alkyl}$;
- l) $\text{C}_{2-6}\text{ alkenyl}$;
- m) $\text{C}_{2-6}\text{ alkynyl}$;
- n) aryl ;
- o) CF_3 ;
- p) $\text{CR}^2\text{R}^2\text{-S(O)}_n\text{-R}^3$;
- q) $\text{CR}^2\text{R}^2\text{NR}^2\text{R}^3$;
- r) C-OH ;
- s) $\text{CR}^2\text{R}^2\text{-C(=W)R}^2$;
- t) acyl ;
- u) C(=W)R^2 ;
- v) C(=W)OR^2 ;
- w) C(=W)SR^2 ;
- x) $\text{C(=W)-NR}^2\text{R}^3$;
- y) $\text{C(=W)NH(CH}_2\text{)}_p\text{-(amino acid residue)}$;
- z) amino residue; or

- aa) A-(amino acid residue);
- bb) cyanoalkyl;
- cc) cyanoalkenyl; or
- dd) cyanoalkynyl,

wherein any of the above optionally may be substituted; or

alternatively, $R^{2''}$ or $R^{6''}$ may be joined to Y to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle comprising 4 -14 members.

A first series of subembodiments of the Formula (B) include all the subembodiments provided for Formula (A) above.

A second series of subembodiments of the present invention is given for the embodiment of the Formula (B) as provided above where $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ are defined as:

- a) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ each independently is halogen; -NO₂; -CN; -OR²; -NH-R⁵-C₁₋₆ alkyl; -NHCO-C₁₋₆ alkyl; oxime; hydrazine; -N(OH)C₁₋₆ alkyl; C₁₋₆ alkoxy; -OH; -NR²R²; or -C₁₋₆ alkyl, alkenyl or alkynyl optionally substituted with one or more of -OH, -SR, -CN, -halo, -C(=W)H, -C(=W)OH, halogen, NR²R², -C₁₋₆ thioether, or -C₁₋₆alkoxy;
- b) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ each independently is halogen, or -C₁₋₆ alkyl, alkenyl, alkynyl optionally substituted with one or more halogen;
- c) $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$ and $R^{6''}$ are hydrogen;
- d) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are methyl;
- e) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are chloro;
- f) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are fluoro;
- g) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, $R^{3''}$ is iodo and $R^{5''}$ is bromo;
- h) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, $R^{3''}$ is methyl, and $R^{5''}$ is chloro; and

- i) $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, $R^{3''}$ is chloro, and $R^{5''}$ is methyl.

A third series of subembodiments of the present invention is given for the embodiment of the Formula (B) as provided above where W, Y, $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ are defined as:

- a) W is O, Y is -O-alkyl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ each independently is halogen; -NO₂; -CN; -OR²; -NH-R⁵-C₁₋₆ alkyl; -NHCO-C₁₋₆ alkyl; oxime; hydrazine; -N(OH)C₁₋₆ alkyl; C₁₋₆ alkoxy; -OH; -NR²R²; or -C₁₋₆ alkyl, alkenyl or alkynyl optionally substituted with one or more of -OH, -SR, -C(=W)H, -C(=W)OH, halogen, NR²R², -C₁₋₆ thioether, or -C₁₋₆alkoxy;
- b) W is S, Y is OH, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ each independently is halogen, or -C₁₋₆ alkyl, alkenyl, alkynyl optionally substituted with one or more halogen;
- c) W is O, Y is C₁₋₆ alkyl, $R^{2''}$, $R^{3''}$, $R^{4''}$, $R^{5''}$ and $R^{6''}$ are hydrogen;
- d) W is S, Y is C₁₋₆ alkylene, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are methyl;
- e) W is NH, Y is -O-alkyl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and
- f) $R^{5''}$ are chloro;
- g) W is S, Y is -O-alkenyl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and
- h) $R^{5''}$ are fluoro;
- i) W is O, Y is aryl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are fluoro;
- j) W is NH, Y is -O-alkynyl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are fluoro;
- k) W is S, Y is S, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, and $R^{3''}$ and $R^{5''}$ are fluoro;
- l) W is O, Y is -O-alkyl, $R^{2''}$, $R^{4''}$, and $R^{6''}$ are hydrogen, $R^{3''}$ is iodo and $R^{5''}$ is bromo;
- m) W is O, Y is -O-alkaryl, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is -OH;

- n) W is S, Y is $-\text{NR}^2\text{R}^3$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{4''}$ is $-\text{NH}_2$;
- o) W is S, Y is $-\text{SR}^2$, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{4''}$ is $-\text{NO}_2$;
and
- p) W is O, Y is $-\text{O-aralkyl}$, $\text{R}^{2''}$, $\text{R}^{4''}$, and $\text{R}^{6''}$ are hydrogen, $\text{R}^{3''}$ is chloro, and $\text{R}^{5''}$ is methyl.

A fourth series of subembodiments are defined when Z is as defined in any one of the second series of subembodiments for Formula (A), $\text{R}^{4'}$, $\text{R}^{5'}$, $\text{R}^{6'}$ and $\text{R}^{7'}$ are as defined in the third series of subembodiments for Formula (A), and $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ all are defined as in the second series of subembodiments for Formula (B).

A fifth series of subembodiments are defined where W and Y are as defined in the first series of subembodiments for Formula (A), Z is as defined in any one of the second series of subembodiments for Formula (A), $\text{R}^{4'}$, $\text{R}^{5'}$, $\text{R}^{6'}$ and $\text{R}^{7'}$ are as defined in the third series of subembodiments for Formula (A), and $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ all are defined as in the third series of subembodiments for Formula (B).

A sixth series of subembodiments are defined when Z is as defined in any one of the second series of subembodiments for Formula (A), $\text{R}^{4'}$, $\text{R}^{5'}$, $\text{R}^{6'}$ and $\text{R}^{7'}$ are as defined in the third series of subembodiments for Formula (A), and W, Y, $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ all are defined as in the third series of subembodiments for Formula (B).

Non-limiting species of the second embodiment as given by Formula (B) above are defined when:

- A) Z is $-\text{C}(=\text{W})\text{NHNHC}_2\text{H}_5\text{OH}$, $\text{R}^{4'}$, $\text{R}^{6'}$ and $\text{R}^{7'}$ are H, $\text{R}^{5'}$ is Cl, Y is H, $\text{R}^{2''}$, $\text{R}^{4''}$ and $\text{R}^{6''}$ are H, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl;
- B) Z is $-\text{C}(=\text{W})\text{NHCH}_2\text{C}(=\text{W})\text{NH}_2$, $\text{R}^{4'}$ is F or Cl, $\text{R}^{6'}$ and $\text{R}^{7'}$ are H, $\text{R}^{5'}$ is F or Cl, Y is OH, $\text{R}^{2''}$, $\text{R}^{4''}$ and $\text{R}^{6''}$ are H, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl;
- C) Z is $-\text{C}(=\text{W})\text{NHCH}_2\text{CONHNH}_2$, $\text{R}^{4'}$ is F or Cl, $\text{R}^{6'}$ and $\text{R}^{7'}$ are H, $\text{R}^{5'}$ is F or Cl, Y is $-\text{O-alkyl}$, $\text{R}^{2''}$, $\text{R}^{4''}$ and $\text{R}^{6''}$ are H, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl;

- D) Z is $-C(=W)NHCH_2CH_2-(2-NO_2, 5-Me-imidazole)$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is $-O-alkenyl$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- E) Z is $-C(=W)NHCH_2NHC(=W)OH$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is $-O-alkynyl$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- F) Z is $-C(=W)CH=CHC(=W)NH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is $-O-alkaryl$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- G) Z is $-C(=W)NHNHCH_2C(=W)NH_2$, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is $-O-aryl$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- H) Z is $-C(=W)NHCH_2C(=W)R^2$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is $-O-heterocycle$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- I) Z is $-C(=W)NHCH_2-A-C(=W)NH_2$, where A is a divalent spacer, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is C_{1-6} alkyl, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NH_2$;
- J) Z is $-C(=W)R^2CH-A-C(=W)H$, where A is a divalent spacer, R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is C_{2-6} alkynyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- K) Z is $-C(=W)R^2CH-A-C(=W)OH$, where A is a divalent spacer, R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is C_{2-6} alkynyl, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NO_2$;
- L) Z is $-C(=W)R^2-CH-A-R^3$, where A is a divalent spacer, R^2 is NH , R^3 is CH_3 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is aryl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- M) Z is $-C(=W)NHR^2-C(=W)R^2$, where R^2 is an optionally substituted, branched chain alkylene or NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is heterocycle, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;
- N) Z is $-C(=W)R^2$, where R^2 is NH , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-OH$;

O) Z is $-C(=W)R^2R^3$ -heterocycle, where R^2 is NH, R^3 is CH_2 , heterocycle is optionally substituted morpholine, imidazole or pyrrole, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

P) Z is $-C(=W)R^2NH-C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyloxy, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

Q) Z is $-C(=W)R^2-NH-C(=W)-C_{1-4}$ alkoxy, where R^2 is an optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

R) Z is $-C(=W)R^3-C(=W)R^2$, where R^2 is C_{1-4} alkoxy, R^3 is optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

S) Z is $-C(=W)R^2R^3$, where R^2 is optionally substituted alkyl, R^3 is optionally substituted phenyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

T) Z is $-C(=W)R^2-NH-C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

U) Z is $-C(=W)R^2C(=W)-NH_2$, where R^2 is optionally substituted alkyl, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

V) Z is $-C(=W)-A-R^3$, where R^3 is an optionally substituted alkyl, A is an alkylene linker, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

X) Z is $-C(=W)R^2C(=W)-O-R^3$, where R^2 is optionally substituted alkyl, R^3 is benzyloxy, $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

Y) Z is $-C(=W)-NH-CH(R^2)-C(=W)-NH_2$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

Z) Z is $-C(=W)-NH-NH_2$, where $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

AA) Z is $-C(=W)-NH-NH(R^2)$, where R^2 is NH_2 , $R^{4'}$ is F or Cl, $R^{6'}$ and $R^{7'}$ are H, $R^{5'}$ is F or Cl, Y is H, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

BB) Z is $-C(=W)-NH-CH(C[=W]NH_2)(CH_2-C[=W]-O-NR^2R^2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, R^2 is alkyl or aryl, Y is H, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NH_2$;

CC) Z is $-C(=W)-NH-CH(-[CH_2]_4-NH-C[=W]-t-BuO)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is C_{1-6} alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

DD) Z is $-C(=W)-NH-CH(-CH_2-CH_2-C[=W]-t-BuO)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

EE) Z is $-C(=W)-NH-CH(-CH_2R^3)(-C[=W]-NH_2)$, where R^3 is CF_3 , $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-OH$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

FF) Z is $-C(=W)-NH-CH(-CH_2R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is H, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NO_2$;

GG) Z is $-C(=W)-NH-CH(-acyl)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

HH) Z is $-C(=W)-NH-CH(-C[=W]-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-OH$, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-OH$;

II) Z is $-C(=W)-NH-CH(-R^3)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

JJ) Z is $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -heterocycle, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

KK) Z is $-C(=W)-NH-CH(-[CH_2]_4-NH_2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -OH, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

LL) Z is $-C(=W)-NH-CH(-CR^2R^2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -O-alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

MM) Z is $-C(=W)-NH-CH(R^2)(-C[=W]-NH_2)$, where $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -O-alkylene, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

NN) Z is $-C(=W)-NH-CH(-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-C_{1-6}$ alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

OO) Z is $-C(=W)-NH-CH(-CH_2-R^3)(-C[=W]-NH_2)$, where R^3 is as defined in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -O-alkylene, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

PP) Z is $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, where R^2 is NH_2 , $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -O-aryl, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NH_2$;

UU) Z is $-C(=W)-NH-NH-CH(R^3)-C[=W]R^2$, where R^2 is $-NH_2$, R^3 is optionally substituted aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -OH, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

VV) Z is $-C(=W)-NHR^2(-R^3)-C[=W]NH_2$, where R^2 is -alkylene, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is -OH, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

WW) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)OH$, where R^2 is alkylene, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

XX) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-NH_2$, where R^2 is alkylene, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

YY) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^3-C(=W)OH$, where R^2 is alkylene, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkylene, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

ZZ) Z is $-C(=W)R^2(CH_2)_p-A-C(=W)NH_2$, where R^2 is $-NH$, p is 0-10, A is divalent, optionally substituted aryl or heterocycle, R^3 is optionally substituted alkyl, aryl or heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkenyl, $R^{2''}$, $R^{3''}$, $R^{5''}$ and $R^{6''}$ are H, and $R^{4''}$ is $-NO_2$;

AAA) Z is $-C(=W)NH-R^3$, where R^3 is optionally substituted heterocycle, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-OH$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

BBB) Z is $-C(=W)NH-A-S(O)_n-R^2$, where A is alkylene, alkenylene or alkynylene, R^2 is $-NH_2$, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

CCC) Z is $-C(=W)NH-NH-R^2(R^3)-A-C(=W)NH_2$, where R^2 is alkyl, A is alkylene, alkenylene or alkynylene, R^3 is aryl, arylene or heterocyclyl, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-OH$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

DDD) Z is $-C(=W)NH-A-[S(O)_2-NH_2]$, where A is alkylene, alkenylene or alkynylene, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkenyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

EEE) Z is $-C(=W)NH-A-C(=W)NH-R^2-C(=W)OH$, where R^2 and R^3 are any of the definitions provided in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O$ -alkenyl, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;


FFF) Z is $-C(=W)NHR^2(-R^3)-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, R^3 is any of the definitions provided in the first general embodiment, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-H$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

GGG) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)OH$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-H$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl;

HHH) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is optionally substituted alkylene, alkenylene or alkynylene, $R^{4'}$ and $R^{5'}$ independently are F or Cl, $R^{6'}$ and $R^{7'}$ are H, Y is $-O\text{-alkyl}$, $R^{2''}$, $R^{4''}$ and $R^{6''}$ are H, and $R^{3''}$ and $R^{5''}$ are methyl; and

wherein W is as defined for Formula (A).

In a third embodiment, the present invention provides an optionally substituted 9-11 membered bicyclic 3-phosphoindole for use in the treatment of HIV represented by the following general Formula (C):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

each W, Z, R^1 , R^2 , R^3 , A, n, $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ is as defined above for Formula (A);

each $R^{3''}$, $R^{4''}$, $R^{5''}$, and $R^{6''}$ is as defined above for Formula (B);

----- indicates the presence of a single or double bond, where:

when ----- indicates the presence of a double bond, each Y and T independently is:

- a) CR^2 ;
- b) N; or
- c) SR^2 ;

with the caveat that one of Y and T must be CR^2 ;

when ----- indicates the presence of a single bond, then each Y and T independently is:

- a) CR^2 ;
- b) O;
- c) NR^2 ; or
- d) SR^2 ;

with the caveat that one of Y and T must be CR^2 ; and

m is 1 or 2, with the proviso that m can only be 2 for T or $\text{Y} = \text{CR}^2$.

A first series of subembodiments of the Formula (C) include all the subembodiments provided for Formula (A) above in which substituent "X" is aryl, heterocyclyl, O-aryl, and O-heterocyclyl.

A second series of subembodiments of the present invention is given for the embodiment of Formula (C) as provided above where Y and T are defined as:

- j) T is $-\text{CH}_2$ and Y is $-\text{O}-$;
- k) T is $-\text{C}-\text{C}(=\text{O})-\text{OCH}_3$ and Y is $-\text{O}-$;
- l) T is $-\text{O}$ and Y is $-\text{CR}^2$;
- m) T is $(-\text{CH}_2)_2$ and Y is $-\text{NR}^2$;
- n) T is $-\text{CH}_2$ and Y is $-\text{SR}^2$;
- o) T is $-\text{O}$ and Y is $-\text{C-aryl}$;
- p) T is $-\text{NR}^2$ and Y is $-\text{C-halo}$;

- q) T is $-\text{SR}_2$ and Y is $-\text{C-heterocycle}$;
- r) T is $-\text{C-C}_{1-6}$ alkyl and Y is $-\text{O-}$;
- s) T is $-\text{C-C(=S)CH}_2$ and Y is $-\text{CH}_2-$;
- t) T is $-\text{CH-OCH}_3$ and Y is $-\text{SR}^2$;
- u) T is $-\text{C-OH}$ and Y is $-\text{CH}_2-$;
- v) T is $-\text{C-O-C}_{1-6}$ alkyl and Y is $-\text{CH}_2-$;
- w) T is $-\text{C-NH}_2$ and Y is $-\text{CH}_2-$;
- x) T is $-\text{C-NH-C}_{1-6}$ alkyl and Y is $-\text{CH}_2-$; and
- y) T is $(-\text{CH}_2)_2$ and Y is $(-\text{CH}_2)_2$.

A third series of subembodiments of the present invention is given for the embodiment of the Formula (C) as provided above where $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ are defined as:

- a) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{3''}$ and $\text{R}^{5''}$ each independently is halogen; $-\text{NO}_2$; $-\text{CN}$; $-\text{OR}^2$; $-\text{NH-R}^5-\text{C}_{1-6}$ alkyl; $-\text{NHCO-C}_{1-6}$ alkyl; hydrazine; $-\text{N(OH)C}_{1-6}$ alkyl; C_{1-6} alkoxy; $-\text{OH}$; $-\text{NR}^2\text{R}^2$; or $-\text{C}_{1-6}$ alkyl, alkenyl or alkynyl optionally substituted with one or more of $-\text{OH}$, $-\text{SR}$, $-\text{C(=W)H}$, $-\text{C(=W)OH}$, halogen, NR^2R^2 , $-\text{C}_{1-6}$ thioether, or $-\text{C}_{1-6}$ alkoxy;
- b) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{3''}$ and $\text{R}^{5''}$ each independently is halogen, or $-\text{C}_{1-6}$ alkyl, alkenyl, alkynyl optionally substituted with one or more halogen;
- c) $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$ and $\text{R}^{6''}$ are hydrogen;
- d) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are methyl;
- e) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are chloro;
- f) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{3''}$ and $\text{R}^{5''}$ are fluoro;
- g) $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, $\text{R}^{3''}$ is iodo and $\text{R}^{5''}$ is bromo;
- h) $\text{R}^{3''}$, $\text{R}^{4''}$ and $\text{R}^{6''}$ are hydrogen, and $\text{R}^{5''}$ is chloro;

- i) $R^{4''}$ and $R^{6''}$ are hydrogen, $R^{3''}$ is chloro, and $R^{5''}$ is methyl.
- i) $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is halogen; $-\text{NO}_2$; $-\text{CN}$; $-\text{OR}^2$; $-\text{NH}-\text{R}^5$ - C_{1-6} alkyl; $-\text{NHCO}-\text{C}_{1-6}$ alkyl; hydrazine; $-\text{N}(\text{OH})\text{C}_{1-6}$ alkyl; C_{1-6} alkoxy; $-\text{OH}$; $-\text{NR}^2\text{R}^2$; or $-\text{C}_{1-6}$ alkyl, alkenyl or alkynyl optionally substituted with one or more of $-\text{OH}$, $-\text{SR}$, $-\text{C}(=\text{W})\text{H}$, $-\text{C}(=\text{W})\text{OH}$, halogen, NR^2R^2 , $-\text{C}_{1-6}$ thioether, or $-\text{C}_{1-6}\text{alkoxy}$;
- j) $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is $-\text{NO}_2$;
- k) $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is $-\text{OR}^2$;
- l) $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is halogen; and
- m) $R^{3''}$, $R^{5''}$ and $R^{6''}$ are hydrogen, and $R^{4''}$ is $-\text{NH}_2$.

A fourth series of subembodiments for Formula (C) are defined when Z is as defined in any one of the second series of subembodiments for Formula (A), $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are as defined in the third series of subembodiments for Formula (A), and $R^{3''}$, $R^{4''}$, $R^{5''}$ and $R^{6''}$ all are defined as in the third series of subembodiments for Formula (C).

A fifth series of subembodiments are defined when Z is as defined in any one of the second series of subembodiments for Formula (A), $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are as defined in the third series of subembodiments for Formula (A), and Y and T are as defined as in the second series of subembodiments for Formula (C).

A sixth series of subembodiments for Formula (C) are defined when Z is as defined in any one of the second series of subembodiments for Formula (A), $R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$ are as defined in the third series of subembodiments for Formula (A), and $R^{3''}$, $R^{4''}$, $R^{5''}$ and $R^{6''}$ all are defined as in the second series of subembodiments for Formula (B).

Species of the third embodiment as given by Formula (C) above are defined when:

- a) Z is $-\text{C}(=\text{W})\text{N}(\text{R}^2\text{R}^3)\text{C}(=\text{W})\text{NH}_2$, where R^2 is NH and R^3 is alkyl; R^1 , $\text{R}^{4'}$, $\text{R}^{6'}$ and $\text{R}^{7'}$ are H ; $\text{R}^{5'}$ is halo or $-\text{NO}_2$; $\text{R}^{4''}$ and $\text{R}^{6''}$ are H ; $\text{R}^{3''}$ and $\text{R}^{5''}$ independently are $-\text{OH}$ or $-\text{NR}^2\text{R}^2$; Y is O ; and T is CH_2 ;

- b) Z is $-C(=W)-NH-CH(-C[=W]NH_2)(-CH_2-C[=W]-O-CH_3-aryl)$; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or $-CN$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are $-C_{1-5}$ alkoxy or $-C_{1-5}$ alkyl optionally substituted with $-C_{1-5}$ alkoxy; Y is SR^2 ; and T is CH_2 ;
- c) Z is $-C(=W)NHNH_2$; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or $-NR^2R^2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are $-NO_2$ or $-CN$; Y is CH_2 ; and $T-R^2-R^3$ is CH_2 ;
- d) Z is $-C(=W)NH-CH_2-C(=W)NHNH_2$; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or $-NH-R^5-C_{1-6}$ alkyl, where R^5 is $-C(O)$ or $-S(O)_n$ and n is 0, 1 or 2; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are $-OR^2$ or $-CN$; Y is $C-C_{1-6}$ alkyl; and T is N;
- e) Z is $-C(=W)NH-CH_2-C(=W)NHNH_2$; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or $-NH-R^5-C_{1-6}$ alkyl, where R^5 is $-C(O)$ or $-S(O)_n$ and n is 0, 1 or 2; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are $-OR^2$ or $-CN$; Y is $C-C_{1-6}$ alkyl; and T is SR^2 and R^2 is as defined for Formula (A);
- f) Z is $-C(=W)NH-CH(R^2)-C(=W)NH_2$, where R^2 is $-NH_2$ or alkyl; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or $-NHCO-C_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are $-NHOH$; Y is $C-alkenyl$; and T is $(CH_2)_2$;
- g) Z is $-C(=W)NR^2-C(=W)R^3$, where R^2 is optionally substituted alkyl, alkenyl or alkynyl; R^3 is NH_2 ; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or oxime; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NH-R^5-C_{1-6}$ alkyl, where R^5 is $-C(O)$ or $-S(O)_n$ and n is 0, 1 or 2; Y is $C-alkynyl$; and T is $-O$;
- h) Z is $-C(=W)NH-R^2-SR^2$ where R^2 is optionally substituted alkyl, alkenyl or alkynyl; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or hydrazine; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NHCO-C_{1-6}$ alkyl; Y is $C-carbocycle$; and T is NR^2 where R^2 is as defined for Formula (A);
- i) Z is $-C(=W)-NH-N(R^3)(R^2)$, where R^2 is hydroxy or alkoxy, and R^3 is H or alkyl; $R^1, R^{4'}, R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or C_{1-6} alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-C(=W)OH$, halogen, $-NR^2R^2$, C_{1-3} alkoxy, or C_{1-3} thioether; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-CN$; Y is $C-aryl$; and T is N;

- j) Z is $-C(=W)NH-CH_2NH-CH(CH_3)C(=W)OH$; R^1 , $R^{4'}$, $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or CF_3 ; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NH-O-C_{1-3}$ alkyl; Y is O; and T is CH_2 ;
- k) Z is $-C(=W)-NH-N(R^2)-CH(R^2)-C(=W)R^2$, where R^2 is H or NH_2 ; R^1 , $R^{4'}$, $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or H; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or halogen; Y is $-NR^2$ and T is CR^2 where R^2 is as defined in the first general embodiment;
- l) Z is $-C(=W)-N(R^2)-C(=W)R^3$, where R^2 is NH and R^3 is CH_3 ; R^1 , $R^{4'}$, $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ is halo or H; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or halogen; Y is $-O$; and T is (C-O-alkyl);
- m) Z is $-C(=W)-NH-CH=CH-C(=W)R^2$, where R^2 is NH_2 ; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or H; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-OR^2$ where R^2 is as defined in the first general embodiment; Y is $-O$; and T is (C- CH_2 - CH_3);
- n) Z is $-C(=W)-NH-CH=CH-C(=W)R^2$, where R^2 is $-NH_2$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-NO_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NHOH$; Y is $-O$ and T is C- CH_3 ;
- o) Z is $-C(=W)R^2(CH_2)_2-C(=W)R^2$, where R^2 is NH or NH_2 ; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-OR^2$ where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-3}$ alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)OH$, halogen, or $-NH_2$; Y is $-SR^2$; and T is CH_2 ;
- p) Z is $-C(=W)-R^2-CH_2-A-C(=W)R^2$, where R^2 is as defined in the first general embodiment, and A is a divalent linker as defined in the first general embodiment; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-CN$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-3}$ alkoxy; Y is $-NR^2$ where R^2 is as defined in the first general embodiment; and T is $(CH_2)_2$;
- q) Z is $-C(=W)-R^2-CH_2-A-C(=W)R^2$, where R^2 is alkyl or $-NH_2$, and A is a divalent linker; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-OH$; Y is $-CH_2$ and T is $(CH_2)_2$;

- r) Z is $-C(=W)-A-R^2-C(=W)OR^3$, where R^2 is $-NH$ and R^3 is $-H$ or alkyl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-NHR^5-C_{1-3}$ alkyl, where R^5 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; Y is $-C-C_{1-6}$ alkyl; and T is CH_2 ;
- s) Z is $-C(=W)-NH-C(=W)OR^3$, wherein R^3 is as defined for Formula (A); R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-NHCO-C_{1-3}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are halo or H; Y is $-C$ -alkenyl; and T is O;
- t) Z is $-C(=W)R^3-NH-C(=W)-R^2$, where R^2 is $-NH_2$ and R^3 is $-NH$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or oxime; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-3}$ alkyl or alkenyl optionally substituted with one or more halogen; Y is $-C$ -alkynyl; and T is $(CH_2)_2$;
- u) Z is $-C(=W)-N(C=O)-N(R^2)-R^3$, where R^2 is H or alkyl, R^3 is $-NH_2$, R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or hydrazine; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H; Y is $-C$ -aryl; and T is $(CH_2)_2$;
- v) Z is $-C(=W)-N(R^2)-N(R^2)-C(=W)R^3$, where R^2 is H or alkyl, and R^3 is NH_2 ; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-NHS(O)_2-C_{1-3}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-methyl$; Y is $-C$ -heterocycle; and T is O;
- w) Z is $-C(=W)-N(-N[R^2][R^3])-R^3$, where R^2 is H or alkyl, and R^3 is NH_2 ; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-C_{1-6}$ alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)H$, $-C(=W)OH$, halogen, NR^2R^2 , C_{1-3} alkoxy, or C_{1-3} thioether, where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; Y is $-C$ -carbocycle; and T is NR^2 where R^2 is as provided above;
- x) Z is $-C(=W)R^2-C(=W)NH_2$, where R^2 is alkyl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-H$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or chlorine; Y is $-SR^2$ and T is CH_2 ;
- y) Z is $-C(=W)R^2-SR^2$, where R^2 is $-NH$ or alkyl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are halo or $-CF_3$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or fluorine; Y is $-O$; and T is $(CH_2)_2$;

z) Z is $-C(=W)-CH(R^2)-C(=NH)R^2$, where R^2 is H or NH_2 ; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are halo or $-H$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-CF_3$; Y is $-NR^2$; and T is CR^2 ; where R^2 is as defined for Formula (A);

aa) Z is $-C(=W)-NH-A-NH-C(=W)-A-C(=W)-R^2$ where R^2 is NH_2 and A is a divalent linker; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are $-CN$ or $-NO_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NO_2$; Y is $-CH_2$; and T is NR^2 , where R^2 is as defined for Formula (A);

bb) Z is $-C(=W)-R^2-CH(-A-C[=W]R^2)(-C[=W]-NH_2)$, where R^2 is H, alkyl, or NH ; A is a divalent spacer linkage as defined for Formula (A); R^1 , $R^{4'}$, and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-NHCO-C_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or hydrazinyl; Y is $(CH_2)_2$; and T is N;

cc) Z is $-C(=W)-NH-A(-C[=W]-NH_2)$, where A is an alkylene or arylene divalent spacer linkage; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-NH(SO_2)C_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NHOH$; Y is $-C$ -alkenyl; and T is N;

dd) Z is $-C(=W)-NH-CH(-A-R^3)(-C[=W]-NH_2)$, where A is an alkylene or alkenylene spacer; R^3 is OH; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or CF_3 ; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NR^2R^2$ where R^2 is as defined for Formula (A); Y is $-C$ -carbocycle; and T is SR^2 ;

ee) Z is $-C(=W)-NH-CH(R^2)(R^3)$, where R^2 and R^3 each is $C(=W)NH_2$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{4'}$ are H or halo; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-5}$ alkoxy; Y is $-O$; and T is CH_2 ;

ff) Z is $-C(=W)-A-CH(-R^2-C[=W]-NH_2)(-C[=W]-NH_2)$, where A is a spacer linkage as defined for Formula (A); R^2 is NH ; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{4'}$ are H or $-NO_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-OH$; Y is $-N$; and T is CH_2 ;

gg) Z is $-C(=W)-NH-CH(-CH-R^2-OH)(-C[=W]-NH_2)$, where R^2 is NH or alkylene; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{4'}$ are H or $-halogen$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-halogen$; Y is $-SR^2$; and T is $CH-OCH_3$;

hh) Z is $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, where R^2 is alkyl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{4'}$ are H or $-NO_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-CN$; Y is $-C$ -aryl; and T is N;

ii) Z is $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, where R^2 is alkyl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{4'}$ are H or $-OH$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-CN$; Y is $-C$ -heterocycle; and T is $(CH_2)_2$;

jj) Z is $-C(=W)-NH-CH(-R^2-C[=W]-NH_2)(-C[=W]-NH_2)$, where R^2 is alkylene; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-NO_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NHOH$; Y is $-O$ and T is CH_2 ;

kk) Z is $-C(=W)-NH-CH(-R^2)(-C[=W]-NH_2)$, where R^2 is $-S(O)R^3$ and R^3 is as defined for Formula (A); R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-OR^2$, $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-3}$ alkoxy; Y is $-NR^2$, where R^2 is as defined in the first general embodiment, and T is CH_2 ;

ll) Z is $-C(=W)-NH-CH(-C[=NH]-NH_2)(-C[=W]-NH_2)$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-OH$; Y is $-SR^2$, and T is $(CH_2)_2$;

mm) Z is $-C(=W)-NH-NH-CH(R^3)-C[=W]R^2$, where R^2 is $-NH_2$ and R^3 is optionally substituted aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-CN$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; Y is $-C-C_{1-6}$ alkyl; and T is N;

nn) Z is $-C(=W)-NHR^2(-R^3)-C[=W]NH_2$, where R^2 is alkyl, and R^3 is optionally substituted alkyl, aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-NH-R^{5'-C_{1-3}}$ alkyl, where R^5 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-6}$ alkyl or alkenyl optionally substituted with one or more $-OH$, $-SR$, $-C(=W)OH$, halogen, or NH_2 ; Y is $-C$ -alkenyl; and T is SR^2 where R^2 is as defined for Formula (A);

oo) Z is $-C(=W)-NHR^2(-R^3)-C[=W]NH-R^2-C(=W)OH$, where R^2 is alkyl, and R^3 is optionally substituted alkyl, aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-NHCO-C_{1-3}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-C_{1-6}$ alkyl or alkenyl optionally substituted with one or more halogen; Y is $-C$ -alkynyl; and T is CH_2 ;

pp) Z is $-C(=W)-NHR^2(-R^3)-C[=W]NH-R^2-NH_2$, where R^2 is alkyl, and R^3 is optionally substituted alkyl, aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or

–oxime; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or –halogen; Y is $-\text{CH}_2$; and T-NR^2 where R^2 is as defined for Formula (A);

qq) Z is $-\text{C}(=\text{W})-\text{NHR}^2(-\text{R}^3)-\text{C}(=\text{W})\text{NH-R}^3$, where R^2 is alkyl, and R^3 is optionally substituted alkyl, aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or –hydrazine; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H; Y is $-\text{NR}^2$ where R^2 is as defined in the first general embodiment; and T is $(\text{CH}_2)_2$;

rr) Z is $-\text{C}(=\text{W})-\text{R}^2-\text{A}-\text{C}(=\text{W})-\text{NH}_2$, where R^2 is $-\text{NH}$, and A is a divalent spacer linkage that is optionally substituted aryl or heteroaryl; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-\text{NH}(\text{SO}_2)\text{C}_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or methyl; Y is $-\text{C-C}_{1-6}$ alkyl; and T is N;

ss) Z is $-\text{C}(=\text{W})\text{NH-R}^3$, where R^3 is an optionally heterocycle; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-\text{C}_{1-6}$ alkyl or alkenyl optionally substituted with one or more $-\text{OH}$, $-\text{SR}$, $\text{C}(=\text{W})\text{H}$, $\text{C}(=\text{W})\text{OH}$, halogen, NR^2R^2 , C_{1-3} alkoxy, or C_{1-3} thioether; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or chloro; Y is $-\text{C-alkenyl}$; and T is NR^2 where R^2 is as defined for Formula (A);

tt) Z is $-\text{C}(=\text{W})\text{NH-A-S(O)}_2-\text{R}^3$, where A is an alkylene, alkenylene or alkynylene divalent spacer linkage, R^3 is $-\text{NH}_2$, and R^5 is $-\text{SO}_2$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or fluoro; Y is $-(\text{CH}_2)_2$; and T is $(\text{CH}_2)_2$;

uu) Z is $-\text{C}(=\text{W})\text{NH-NH-A-(R}^3)-\text{S(O)}_2-\text{NH}_2$, where A is an alkylene, alkenylene or alkynylene divalent spacer linkage, R^3 is aryl, arylene or heteroaryl, and R^5 is $-\text{SO}_2$; R^1 , $R^{6'}$ and $R^{7'}$ are H; $R^{4'}$ and $R^{5'}$ are H or $-\text{CF}_3$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or CF_3 ; Y is $-\text{O}$; and T is CH_2 ;

vv) Z is $-\text{C}(=\text{W})\text{NH-R}^3(\text{SO}_2-\text{NH}_2)$, where R^3 is aryl, arylene or heteroaryl; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or –halo; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-\text{NO}_2$; Y is $-\text{O}$; and T is CR^2 where R^2 is as defined for Formula (A);

ww) Z is $-\text{C}(=\text{W})\text{NH-A-(NH}_2)$, where A is an alkylene, alkenylene or alkynylene divalent spacer linkage; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-\text{NO}_2$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or –halo; Y is $-\text{O}$; and T is C-carbocycle;

xx) Z is $-C(=W)NH-R^3(-A-SO_2-NH_2)$, where A is an alkylene, alkenylene or alkynylene divalent spacer linkage; R^3 is aryl, arylene or heteroaryl; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-CN$; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NH-R^5-C_{1-3}$ alkyl, where R^5 is as defined in the first general embodiment; Y is $-C$ -carbocycle; and T is NR^2 where R^2 is as defined for Formula (A);

yy) Z is $-C(=W)NH(-R^3)-C(=W)NH-A-C(=W)OH$, where A is an optionally substituted alkylene, alkenylene or alkynylene divalent spacer linkage, R^3 is as defined in the first general embodiment; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-OR^2$ where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-CN$; Y is $-C$ -aryl; and T is CH_2 ;

zz) Z is $-C(=W)NH(-R^3)-C(=W)NH-A-C(=W)NH_2$, where A is an optionally substituted divalent spacer linkage as defined for Formula (A); R^3 is as defined in the first general embodiment; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-NR^2R^2$, where R^2 is as defined in the first general embodiment; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-NH-R^5-C_{1-6}$ alkyl, where R^5 is as defined in the first general embodiment; Y is $-C$ -heterocycle; and T is SR^2 ;

aaa) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)OH$, where R^2 is as defined for Formula (A); A is an optionally substituted divalent spacer linkage; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-NHSO_2-C_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or hydrazine; Y is $-O$ and T is $(CH_2)_2$; and

bbb) Z is $-C(=W)NHR^2-C(=W)NH-R^2-C(=W)NH_2$, where R^2 is as defined for Formula (A); A is an optionally substituted divalent spacer linkage; R^1 , $R^{4'}$ and $R^{7'}$ are H; $R^{5'}$ and $R^{6'}$ are H or $-NHCO-C_{1-6}$ alkyl; $R^{4''}$ and $R^{6''}$ are H; $R^{3''}$ and $R^{5''}$ independently are H or $-OH$; Y is $-NR^2$ where R^2 is as defined in the first general embodiment, and T is CH_2 ;

wherein W is as defined above for the first general embodiment of Formula (A).

For all embodiments, subembodiments are defined when:

1) Z is $-C(=W)-NH-R^2$; $-C(=W)-NR^2R^3$; $-C-R^2R^3$; $-CR^2-C(=W)R^3$; $-R^2-C(=W)R^3$; $-R^2-C(=W)R^2$; $-R^2R^3$; R^3 ; or $C(=W)-NH-CR^2R^2-C(=W)-NH-CR^2R^3-C(=W)-NR^2R^3$;

2) $R^{4'}$, $R^{6'}$ and $R^{7'}$ are H, and $R^{5'}$ is i) halogen, and chlorine in particular; ii) hydrazine; iii) C_{1-6} alkyl, alkenyl, alkynyl, amino-alkyl, thioamino-alkyl, or aminocarbonyl-alkyl, each of which optionally may be substituted with one or more $-OH$, $-C(=W)H$, $-C(=W)OH$, $-NR^2R^3$, $-C_{1-3}$ alkoxy, or $-C_{1-3}$ thioether; or one of the following combinations:

- a) $R^{5'}$, $R^{6'}$ and $R^{7'}$ are hydrogen, and $R^{4'}$ is halogen;
- b) $R^{4'}$, $R^{5'}$ and $R^{7'}$ are hydrogen, and $R^{6'}$ is halogen;
- c) $R^{4'}$, $R^{5'}$ and $R^{6'}$ are hydrogen, and $R^{7'}$ is halogen;
- d) $R^{5'}$, $R^{6'}$ and $R^{7'}$ are hydrogen, and $R^{4'}$ is CF_3 ;
- e) $R^{4'}$, $R^{5'}$ and $R^{7'}$ are hydrogen, and $R^{6'}$ is CF_3 ; and
- f) $R^{4'}$, $R^{5'}$ and $R^{6'}$ are hydrogen, and $R^{7'}$ is CF_3 .

III. Definitions

The following definitions and term construction are intended, unless otherwise indicated.

Ranges, specific values, and typical values listed for radicals, substituents and derivatives are for illustration only, and do not exclude other defined values or values within defined ranges for the radicals, substituents and derivatives. Whenever a range is described herein, the range independently includes each member of the range. As an illustrative example, when C_{1-6} -alkyl are referred to, this listing independently includes C_1 -alkyl, C_2 -alkyl, C_3 -alkyl, C_4 -alkyl, C_5 -alkyl and C_6 -alkyl.

“Halo” is fluoro, chloro, bromo or iodo.

“Alkyl”, “alkoxy”, “alkenyl”, “alkynyl”, etc., includes both straight chain and branched groups. However, reference to an individual radical such as “propyl” embraces only that straight-chain radical, whereas a branched chain isomer such as “isopropyl” is specifically termed such.

“Alkyl” as used herein and unless otherwise specified, is a saturated, straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of, for example, C₁₋₁₀, and specifically includes methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, *t*-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. When the context of this document allows alkyl to be substituted, the moieties with which the alkyl group may be substituted include but not limited to hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, aryl, heterocyclyl, halo, carboxy, acyl, acyloxy, amido, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either protected or unprotected as needed, as known to those skilled in the art and as taught, for example, in Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Third Ed., 1999.

The term “lower alkyl” as used herein and unless otherwise specified, includes a C₁₋₄ saturated, straight, branched, or if appropriate, cyclic (for example, cyclopropyl) alkyl group, including both substituted and unsubstituted forms. Unless otherwise specifically stated in this application, when alkyl is a suitable moiety, lower alkyl is typical. Similarly, when alkyl or lower alkyl is a suitable moiety, unsubstituted alkyl or lower alkyl is typical.

The terms “alkenyl” and “alkynyl” refer to alkyl moieties, including both substituted and unsubstituted forms wherein at least one saturated C-C bond is replaced by a double or triple bond. Thus, C₂₋₆ alkenyl may be vinyl, allyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, or 5-hexenyl. Similarly, C₂₋₆ alkynyl may be ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 1-pentyne, 2-pentyne, 3-pentyne, 4-pentyne, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, or 5-hexynyl.

The term "alkylene" refers to a saturated, straight chain, divalent alkyl radical of the formula $-(CH_2)_n-$, wherein "n" may be any whole integer from 1 to 10.

As used herein with exceptions as noted, "aryl" is intended to mean any stable monocyclic, bicyclic or tricyclic carbon ring of up to 8 members in each ring, wherein at least one ring is aromatic as defined by the Huckel $4n+2$ rule. Examples of aryl ring systems include phenyl, naphthyl, tetrahydronaphthyl, and biphenyl. The aryl group may be substituted with one or more moieties including but not limited to hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, alkyl, heterocyclyl, halo, carboxy, acyl, acyloxy, amido, nitro, cyano, sulfonamido, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either protected or unprotected as needed, as known to those skilled in the art and as taught, for example, in Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Third Ed., 1999.

The term "heterocycle" or "heterocyclic" as used herein except where noted, refers to a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring which is either saturated or unsaturated, including heteroaryl, and which consists of carbon atom(s) and from one to four heteroatoms including but not limited to O, S, N and P; and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and/or the nitrogen heteroatom quaternized, and including any bicyclic group in which any of the above-identified heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. The heteroaromatic ring may be partially or totally hydrogenated, as desired. For example, dihydropyridine may be used in place of pyridine. Functional oxygen and nitrogen groups on a heteroaryl may be protected as necessary or desired. Suitable protecting groups for oxygen or nitrogen include trimethylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, trityl, substituted trityl, alkyl, methanesulfonyl, p-toluenesulfonyl, or acyl groups such as acetyl and propionyl.

Non-limiting examples of heteroaryl and heterocyclic groups include furyl, pyridyl, pyrimidyl, pyridazinyl, pyrazinyl, piperidinyl, piperazinyl, thienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, tetrazolyl, triazolyl, triazinyl, thiazinyl, oxazolyl, purinyl, carbazolyl, quinolinyl, pyrazolyl, morpholinyl, benzimidazolyl, and the like. Any of the heteroaromatic and heterocyclic moieties may be optionally substituted as

described above for aryl, including substitution(s) with one or more hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, alkyl, heterocyclyl, halo, carboxy, acyl, acyloxy, amido, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either protected or unprotected as needed, as known to those skilled in the art and as taught, for example, in Greene et al., Protective Groups in Organic Synthesis, John Wiley and Sons, Third Ed., 1999.

The term “acyl” refers to a compound of the formula “RC(O)-”, wherein R is substituted or unsubstituted alkyl or aryl as defined herein.

The term “carboxyl” refers to a compound of the formula “RCOOH”, wherein R is substituted or unsubstituted alkyl or aryl as defined herein.

The term “aralkyl” as used herein unless otherwise specified, refers to an aryl group as defined above linked to the molecule through an alkyl group as defined above.

The term “alkaryl” as used herein unless otherwise specified, refers to an alkyl group as defined above linked to the molecule through an aryl group as defined above.

The term “alkoxy” as used herein unless otherwise specified, refers to a moiety of the structure “-O-alkyl”, where alkyl is as defined above.

The term “amino” as used herein unless otherwise specified, refers to a moiety represented by the structure “-NR₂”, and includes primary amines, and secondary and tertiary amines optionally substituted by alkyl, aryl, heterocyclyl, and/or sulfonyl groups. Thus, R₂ may represent two hydrogens, two alkyl moieties, or one hydrogen and one alkyl moiety.

The term “amido” as used herein unless otherwise specified, refers to a moiety represented by the structure “-C(O)NR₂”, wherein R₂ is an H, alkyl, aryl, acyl, heterocyclyl and/or a sulfonyl group.

As used herein, an “amino acid” or an “amino acid residue” is a natural amino acid or some portion thereof (i.e., Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Hyl, Hyp, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val) in D or L form, or an unnatural amino acid having one or more open valences such as, for example, *t*-butylglycine, ornithine, hippuric acid and phosphothreonine. Other unnatural amino acids are those represented by the formula “NH₂(CH₂)_yCOOH”, wherein y is 2-12, and includes

aminoalkanoic acids such as ϵ -amino caproic acid ($\text{H}_2\text{N}-(\text{CH}_2)_5\text{-COOH}$). The term also comprises natural and unnatural amino acids bearing amino-protecting groups such as acyl, trifluoroacetyl and benzyloxycarbonyl, as well as natural and unnatural aminot acids protected at carboxy moieties by protecting groups such as C_{1-6} alkyl, phenyl or benzyl ester and amide, and protecting groups known to those of skill in the art. In all instances where natural and unnatural amino acids contain one or more chiral centers, all possible stereochemical configurations, including both "D" and "L" forms and mixtures thereof, including racemic mixtures, are contained herein.

The term "quaternary amine" as used herein includes quaternary ammonium salts that have a positively charged nitrogen. They are formed by the reaction between a basic nitrogen in the compound of interest and an appropriate quaternizing agent such as, for example, methyl iodide or benzyl iodide. Appropriate counterions accompanying a quaternary amine include acetate, trifluoroacetate, chloro, bromo and iodo ions.

As used herein, the term "*N*-oxides" denotes a state of the compounds of the present invention in which one or more nitrogen atoms are oxidized with an oxygen atom.

As used herein, a "retrovirus" includes any virus that expresses reverse transcriptase. Examples of a retrovirus include but are not limited to, HIV-1, HIV-2, HTLV-I, HTLV-II, FeLV, FIV, SIV, AMV, MMTV, and MoMuLV.

As used herein, "reverse transcriptase" or "RT" refers to an enzyme having a non-nucleoside inhibitory binding site similar to that of HIV-1 RT, and to which ligands, which bind the composite binding pocket of the compounds of the present invention, also will bind. One measure of RT activity is viral replication. A measure of HIV-1 viral replication is the automated assay that utilizes MTT, as described earlier in this specification. Another measure is the p24 core antigen enzyme immunoassay, such as, for example, the assay commercially available from Coulter Corporation/Immunotech, Inc.[®] (Westbrook, MI). Another means for measuring RT activity is by assaying recombinant HIV-1 reverse transcriptase activity, such as, for example, by using the Quan-T-RT[™] assay system commercially available from Amersham[®] (Arlington Heights, IL) and as described by Bosworth et al., *Nature*, **1989**, 341:167-168.

As used herein, a compound that “inhibits replication of human immunodeficiency virus (HIV)” means a compound that, when contacted with HIV-1, for example, via HIV-infected cells, effects a reduction in the amount of HIV-1 as compared with an untreated control. Inhibition of replication of HIV-1 may be measured by any means known to those skilled in the art, such as, for example, by the p24 assay disclosed above.

The reagent denoted “mCPBA” in the synthesis schemes is meta-chloroperoxybenzoic acid.

The term “salvage therapy” as used herein means a compound that can be taken with any regimen after a patient’s initial treatment regimen has failed.

As used herein, the term “host” refers to a multicellular or unicellular organism in which the virus can replicate. Thus, “host” includes a cell line, an mammal and, preferably, a human. Alternatively, a host can be carrying a part of the HIV genome whose replication or function may be altered by the compounds of the present invention. The term host specifically refers to infected cells, cells transfected with all or part of the HIV genome, and mammals, especially primates including chimpanzees and humans. In most mammal applications of the present invention, the host is a human patient. Veterinary applications, however, are clearly anticipated by the present invention, such as, for example, in chimpanzees.

IV. Pharmaceutically Acceptable Salts, Prodrugs, Stereoisomers and Tautomers

An active compound may be administered as a salt or prodrug that, upon administration to the recipient, is capable of providing directly or indirectly the parent compound, or that exhibits activity itself. Nonlimiting examples include a pharmaceutically-acceptable salt, alternatively referred to as a “physiologically-acceptable salt”. In addition, modifications made to a compound can affect its biologic activity, in some cases increasing the activity over the parent compound. This activity can be assessed by preparing a salt or prodrug form of the compound, and testing its antiviral activity by using methods described herein or other methods known to those of skill in the art of NNRTIs.

The phrase “pharmaceutically acceptable salt or prodrug” is used throughout the specification to describe any pharmaceutically acceptable form (such as an ester, amide, salt of an ester, salt of an amide or related group) of a compound that, upon administration to a patient, provides an active compound of the invention. The terms “stereoisomer” and “tautomer” as used herein include all possible stereoisomeric and tautomeric forms of the compounds of the present invention, as well as their quaternary amine, salt, solvate, prodrug, derivative, and *N*-oxide forms. Where the compounds of the general formulae (I) and (II) contain one or more chiral centers, all possible enantiomeric and diastereomeric forms are included.

The term “pharmaceutically acceptable salt” refers to the state of a compound in which the compound carries a counterion that is pharmaceutically acceptable, and wherein the salt retains the desired biological activity of the herein-identified compounds while exhibiting minimal undesired toxicological effects. Such salts are non-toxic, therapeutically useful forms of the compounds of the present invention. Any salt that retains the desired biological activity of the compounds contained herein and that exhibits minimal or no undesired or toxicological effects is intended for inclusion here. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable organic or inorganic acids and bases. Non-pharmaceutically acceptable acids and bases also find use herein, as for example, in the synthesis and/or purification of the compounds of interest. Thus, all “salts” are intended for inclusion here.

Non-limiting examples of suitable salts include those derived from inorganic acids, such as, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, bicarbonic acid, carbonic acid; and salts formed with organic acids, such as, for example, formic acid, acetic acid, oxalic acid, tartaric acid, succinic acid, malic acid, malonic acid, ascorbic acid, citric acid, benzoic acid, tannic acid, palmoic acid, alginic acid, polyglutamic acid, tosic acid, methanesulfonic acid, naphthalenesulfonic acid, naphthalenedisulfonic acid, α -ketoglutaric acid, α -glycerophosphoric acid and polygalacturonic acid. Suitable salts include those derived from alkali metals such as lithium, potassium and sodium, from alkaline earth metals such as calcium and magnesium, as well as from other bases well known to those of skill in the pharmaceutical art. Other suitable salts include those derived from metal cations such as zinc, bismuth, barium, or aluminum, or with a cation formed from an amine,

such as ammonia, NN-dibenzylethylene-diamine, D-glucosamine, tetraethylammonium, or ethylenediamine. Moreover, suitable salts include those derived from a combination of acids and bases, such as, for example, a zinc tannate salt.

A pharmaceutically acceptable prodrug refers to a compound that is metabolized (i.e., hydrolyzed or oxidized, for example) in the host to form a compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, and/or dephosphorylated to produce the active compound.

The compounds of the present invention either possess antiviral activity against retroviruses and HIV in particular, or are metabolized to a compound that exhibits such activity.

Any of the oxo-pyrimidine compounds described herein may be administered as a prodrug to increase the activity, bioavailability, stability, or otherwise alter the properties of the oxo-pyrimidine. A number of prodrug ligands also are known. In general, acylation, alkylation or other lipophilic modifications of a heteroatom of the oxo-pyrimidine will increase the stability of the compound. Examples of substituent groups that can replace one or more hydrogens on a heterocycle include, but are not limited to, alkyl, aryl, steroidal, carbohydrate including sugars, 1,2-diacylglycerol, phospholipid, phosphatidylcholine, phosphocholine, and/or alcohol. Any of these may be used in combination with the disclosed oxo-pyrimidine compound to achieve a desired effect.

V. Methods of Treatment

In one embodiment, a method of treatment or prophylaxis of an HIV infection in a host is provided, comprising administering a 3-phosphoindole compound to a host in need thereof. In one particular embodiment, the compound is administered orally, parenterally, enterally, intravenously, intradermally, subcutaneously, percutaneously, transdermally, intranasally, topically or by inhalation therapy. The compound may be administered to a host that has been diagnosed with an HIV infection by measurement of a viral load in the host blood or tissue. In other embodiments the host can have been diagnosed by measurement of an anti-HIV antibody titer in blood. In another

embodiment, the compounds are administered to reduce or prevent symptoms of AIDS (acquired immune deficiency syndrome) in a host. In yet another embodiment the compounds of the invention are administered to a host at risk of infection with HIV.

In another embodiment, the active compound exhibits activity against drug-resistant forms of HIV, and thus exhibits decreased cross-resistance against currently approved antiviral therapies. The phrase “activity against a drug-resistant form of HIV” means that a compound (or its prodrug or pharmaceutically acceptable salt) is active against the mutant strain with an EC_{50} of less than approximately 50, 25, 10 or 1 micromolar concentration. In one embodiment, the non-nucleoside reverse transcriptase inhibitor (NNRTI) displays an EC_{50} (in molar concentration) against a mutant HIV strain of less than approximately 5, 2.5, 1 or 0.1 micromolar. In one non-limiting embodiment, the HIV mutant strain has a reverse transcriptase mutation at lysine 103 → asparagine and/or tyrosine 181 → cysteine.

The 3-phosphoindoles can be assessed for their ability to inhibit reverse transcriptase activity *in vitro* according to standard screening methods. The spectrum of activity exhibited by any particular compound is determined by evaluating the compound in assays described in this specification or with other confirmatory assays known to those skilled in the art of anti-HIV compounds. Compounds typically exhibit an EC_{50} of less than 10-15 μ M.

In one embodiment, the efficacy of the 3-phosphoindoles is measured by the HIV-specific enzyme-linked immunosorbent assay, p24 ELISA. Drug efficacy is expressed as percent inhibition of the HIV p24 antigen in this rapid and sensitive assay. In a related embodiment useful for specific experiments, the efficacy of the anti-HIV compound is determined by a “plaque reduction assay” which measures the concentration of compound necessary to reduce the plaque number of the virus *in vitro*, according to the methods set forth more particularly herein, by 50% (i.e., the EC_{50} of the compound). In some embodiments the compound exhibits an EC_{50} of less than 15, or less than 10 micromolar to nanomolar amounts *in vitro*.

VI. Combination or Alternation Therapy

In a certain embodiments, the 3-phosphoindole compound is administered in combination and/or alternation with one or more other anti-retroviral or anti-HIV agent.

In one embodiment, the effect of administering two or more such agents in combination and/or alternation produces a synergistic effect in inhibiting HIV replication. In another embodiment, the effect of administering two or more such agents in combination and/or alternation produces an additive effect in inhibiting HIV replication.

In combination therapy, effective dosages of two or more agents are administered together, whereas during alternation therapy an effective dosage of each agent is administered serially. The dosages will depend upon absorption, inactivation, and excretion rates of the drugs as well as other factors known to those of skill in the art. Dosage values also will vary with the severity of the condition to be alleviated. For any particular individual, specific dosage regimens and schedules should be adjusted over time to meet the needs of the individual and the professional judgment of the person administering or supervising the administration of the compositions.

Drug resistance most typically occurs by mutation of a gene that encodes for an enzyme used in the viral replication cycle, and most typically in the case of HIV, in either the reverse transcriptase or protease genes. It has been demonstrated that the efficacy of an anti-HIV drug can be prolonged, augmented or restored by administering the compound in combination or alternation with a second, and perhaps third, antiviral compound that induces a different mutation(s) from that selected for by the principle drug. Such drug combinations simultaneously reduce the possibility of resistance to any single drug and any associated toxic effects. Alternatively, the pharmacokinetics, biodistribution, or other parameters of the drug can be altered by such combination or alternation therapy. For example, the use of a combination of drugs may permit an individual drug within that combination to be given at a dosage lower than what would be required when the drug is administered as a monotherapeutic. Likewise, when drugs that target different stages of the viral life cycle are combined, there exists the possibility for potentiating their effects. Moreover, use of combinations of drugs could lower or eliminate undesirable side-effects from a single drug while still producing anti-viral activity. In general, combination therapy is typical over alternation therapy because it places multiple, simultaneous pressures on the virus.

The second antiviral agent for the treatment of HIV can be, for example, a protease inhibitor, an HIV-integrase inhibitor, a chemokine inhibitor, or a reverse transcriptase inhibitor ("RTI"), the latter of which can either be a synthetic nucleoside

reverse transcriptase inhibitor ("NRTI") or a non-nucleoside reverse transcriptase inhibitor ("NNRTI"). In other embodiments, a second or third compound may be a pyrophosphate analog or a fusion-binding inhibitor. A list compiling resistance data collected *in vitro* and *in vivo* for certain antiviral compounds is found in Schinazi et al., Mutations in retroviral genes associated with drug resistance, *International Antiviral News*, 1997, 5(8).

In certain embodiments, the indole compound is administered in combination and/or alternation with FTC (2',3'-dideoxy-3'-thia-5-fluorocytidine); 141W94 (amprenavir, Glaxo Wellcome, Inc.); Viramune (nevirapine); Rescriptor (delavirdine); DMP-266 (efavirenz); DDI (2',3'-dideoxyinosine); 3TC (3'-thia-2',3'-dideoxycytidine); DDC (2',3'-dideoxycytidine), abacavir (1592U89), which is (1S,4R)-4-[(2-amino-6-cyclopropyl-amino)-9H-purin-9-yl]-2-cyclopentene-1-methanol succinate, **Tenofovir DF** (Viread), D4T, or AZT.

Other examples of antiviral agents that can be used in combination and/or alternation with the compounds disclosed herein include, but are not limited to, foscarnet; carbovir; acyclovir; interferon; fusion inhibitors such as **enfuvirtide**; and β -D-dioxolane nucleosides such as β -D-dioxolanylguanine (DXG), β -D-dioxolanyl-2,6-diaminopurine (DAPD), and β -D-dioxolanyl-6-chloropurine (ACP). Interferons that may be used include Schering-Plough's alpha interferon-2b products, Intron® A and PEG-Intron™; and Hoffman La Roche's Co-Pegasus and PEGASYS (pegylated interferon alfa-2a). Combinations with which the 3-phosphoindoles can be administered include **Epzicom** (ABC+3TC), **Trizivir** (ABC + 3TC + AZT), **Truvada** (FTC +Viread) and **Combivir** (AZT+3TC).

Examples of protease inhibitors that can be used in combination and/or alternation with the compounds disclosed herein include, but are not limited to indinavir ({1(1S,2R),5(S)}-2,3,5-trideoxy-N-(2,3-dihydro-2-hydroxy-1H-inden-1-yl)-5-[2-[(1,1-dimethylethyl)amino]carbonyl]-4-(3-pyridinylmethyl)-1-piperazinyl]-2-(phenylmethyl)-D-erythro-pentoamide sulfate; Merck & Co., Inc.); nelfinavir (Agouron); ritonavir (Abbott Labs), saquinavir (Roche); **Amprenavir**; **Atazanavir**; **Fosamprenavir**; **Kaletra**; and DMP-450 {[4R-4(r-a,5-a,6-b,7-6)-hexahydro-5,6-bis(hydroxy)-1,3-bis(3-amino)-phenyl]methyl-4,7-bis(phenylmethyl)-2H-1,3-diazepin-2-one}-bismesylate (Triangle Pharmaceuticals, Inc.).

Other compounds that can be administered in combination or alternation with the phenylindole to augment its anti-viral properties include (1S, 4R)-4-[2-amino-6-cyclopropyl-amino-9H-purin-9-yl]-2-cyclopentene-1-methanol succinate (1592U89, a carbovir analog, from GlaxoSmithKline); BILA 1906 (N-{1S-[[[3-[2S-{(1,1-dimethylethyl)amino]carbonyl}-4R-]3-pyrindinylmethyl)thio]-1-piperidinyl]-2R-hydroxy-1S-phenylmethyl)propyl]-amino]carbonyl]-2-methylpropyl}-2-quinolinecarboxamide) (Bio Mega/Boehringer Ingelheim); BILA 2185 (N-(1,1-dimethylethyl)-1-[2S-[[[2-2,6-dimethyl-phenoxy]-1-oxoethyl]amino]-2R-hydroxy-4-phenylbutyl]4R-pyridinylthio-2-piperidine-carboxamide) (Bio Mega/Boehringer Ingelheim); BM+51.0836 (triazoloiso-indolinone derivative) and BMS 186,318 (aminodiol derivative HIV-1 protease inhibitor) (Bristol-Myers Squibb); d4API (9-[2,5-dihydro-5-(phosphonomethoxy)-2-furanyl]-adenine) (Gilead); HBY097 (S-4-isopropoxycarbonyl-6-methoxy-3-[methylthio-methyl]-3,4-dihydroquinoxalin-2(1H)-thione); HEPT (1-[(2-hydroxy-ethoxy)methyl]6-[phenylthio]-thymine); KNI-272 ((2S,3S)-3-amino-2-hydroxy-4-phenylbutyric acid-containing tripeptide); L-697,593 (5-ethyl-6-methyl-3-(2-phthalimido-ethyl)pyridin-2(1H)-one); L-732,524 (hydroxy-aminopentane amide HIV-1, protease inhibitor) (Merck & Co.); L-697,661 (3-{[(4,7-dichloro-1,3-benzoxazol-2-yl)methyl]amino}-5-ethyl-6-methyl-pyridin-2(1H)-one); L-FDDC ((-)-β-L-5-fluoro-2',3'-dideoxycytidine); L-FDOC ((-)-β-L-5-fluoro-dioxolane cytosine); PFA (phosphonoformate; "foscarnet"; Astra); PMEA (9-(2-phosphonylmethoxyethyl)adenine) (Gilead); PMPA ((R)-9-(2-phosphonylmethoxypropyl)-adenine) (Gilead); Ro 31-8959 (hydroxyethylamine derivative HIV-1 protease inhibitor) (Roche); RPI-3121 (peptidyl protease inhibitor, 1-[(3S)-3-(n-alpha-benzyloxy-carbonyl)-1-asparginyl]-amino-2-hydroxy-4-phenylbutyryl]-n-tert-butyl-1-proline amide); 2720 (6-chloro-3,3-dimethyl-4-(isopropenyloxycarbonyl)-3,4-dihydro-quinoxalin-2(1H)thione); SC-52151 (hydroxyethylurea isostere protease inhibitor) (G.D. Searle); SC-55389A (hydroxyethyl-urea isostere protease inhibitor) (G.D. Searle); TIBO R82150 ((+)-(5S)-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)-imidazo-[4,5,1-jk]-[1,4]-benzodiazepin-2(1H)-thione) (Janssen Pharmaceuticals); TIBO 82913 ((+)-(5S)-4,5,6,7-tetrahydro-9-chloro-5-methyl-6-(3-methyl-2-butenyl)imidazo[4,5,1-jk]-[1,4]-benzo-diazepin-2(1H)-thione) (Janssen Pharmaceuticals); TSAO-m3T ([2',5'-bis-O-(tert-butyldimethylsilyl)-3'-spiro-5'-(4'-amino-1',2'-oxathiole-2',2'-dioxide)]-β-D-pentofuranosyl-N3-methyl-thymine); U90152 (1-[3-[(1-methylethyl-amino]2-pyridinyl]-

4-[[5—[(methylsulphonyl)-amino]-1H-indol-2-yl]-carbonyl]-piperazine); UC (thio-carboxanilide derivatives) (Uniroyal); UC-781 (N-[4-chloro-3-(3-methyl-2-butenyloxy)phenyl]-2-methyl-3-furancarbothioamide); UC-82 (N-[4-chloro-3-(3-methyl-2-butenyloxy)phenyl]-2-methyl-3-thiophenecarbothioamide); VB 11,328 (hydroxyethylsulphonamide protease inhibitor) (Vertex/Glaxo Wellcome); XM 323 (cyclic urea protease inhibitor) (Dupont Merck); and penciclovir. In yet another embodiment, the indole compound of the invention is administered in combination with the protease inhibitor LG 1350.

The following drugs have been approved by the FDA or are currently or have been in clinical trials for use in the treatment of HIV infection, and therefore in one embodiment, can be used in combination and/or alternation with the compounds of the present invention.

Drug Name	Manufacturer
3TC, Epivir® brand lamivudine	<u>GlaxoSmithKline</u>
abacavir generic Ziagen®, ABC, or 1592U89	<u>GlaxoSmithKline</u>
ABC, Ziagen® brand abacavir, or 1592U89	<u>GlaxoSmithKline</u>
ABT-378/r, or Kaletra® brand lopinavir/ritonavir	<u>Abbott Laboratories</u>
<i>AG-1549, S-1153, or capravirine (CPV)</i>	<u>Pfizer</u>
<i>AG1661, Remune® brand HIV-1 Immunogen, or Salk vaccine</i>	<u>Immune Response Corp.</u>
Agenerase® brand amprenavir (APV), 141W94, or VX-478	<u>GlaxoSmithKline</u>
<i>aldesleukin generic Proleukin®, or Interleukin-2 (IL-2)</i>	<u>Chiron Corporation</u>
<i>amdoxovir, or DAPD</i>	<u>Gilead Sciences</u>
amprenavir generic Agenerase®, APV, 141W94, or VX-478	<u>GlaxoSmithKline</u>
APV, Agenerase® brand amprenavir, 141W94, or VX-478	<u>GlaxoSmithKline</u>
atazanavir generic Reyataz™, or BMS-232632	<u>Bristol-Myers Squibb</u>
AZT, Retrovir® brand zidovudine (ZDV)	<u>GlaxoSmithKline</u>
Bis(POC) PMPA, Viread® brand tenofovir DF	<u>Gilead Sciences</u>

Drug Name	Manufacturer
<u>BMS-232632</u> , or Reyataz™ brand atazanavir	<u>Bristol-Myers Squibb</u>
<u>BMS-56190</u> , or <u>DPC-083</u>	<u>Bristol-Myers Squibb</u>
<u>calanolide A</u>	<u>Sarawak Medichem</u>
<u>capravirine</u> (CPV), AG-1549, or S-1153	<u>Pfizer</u>
Combivir® brand zidovudine + lamivudine, or AZT + 3TC	<u>GlaxoSmithKline</u>
<u>CPV</u> (capravirine), AG-1549, or S-1153	<u>Pfizer</u>
<u>Crixivan</u> ® brand indinavir (IDV), or MK-639	<u>Merck & Co.</u>
d4T, Zerit® brand stavudine, or BMY-27857	<u>Bristol-Myers Squibb</u>
<u>DAPD</u> , or amdoxovir	<u>Gilead Sciences</u>
ddC, or Hivid® brand zalcitabine	<u>Hoffmann-La Roche</u>
ddI, Videx® brand didanosine, or BMY-40900	<u>Bristol-Myers Squibb</u>
<u>delavirdine</u> generic Rescriptor®, DLV, or U-90152S/T	<u>Pfizer</u>
<u>didanosine</u> generic Videx®, ddI, or BMY-40900	<u>Bristol-Myers Squibb</u>
<u>DLV</u> , Rescriptor® brand delavirdine, or U-90152S/T	<u>Pfizer</u>
<u>DPC-083</u> , or <u>BMS-56190</u>	<u>Bristol-Myers Squibb</u>
<u>Droxia</u> ® brand hydroxyurea (HU)	<u>Bristol-Myers Squibb</u>
<u>efavirenz</u> generic Sustiva®, or EFV	<u>Bristol-Myers Squibb</u>
<u>EFV</u> , Sustiva® brand efavirenz	<u>Bristol-Myers Squibb</u>
<u>emtricitabine</u> generic <u>Emtriva</u> ™, or <u>FTC</u>	<u>Gilead Sciences</u>
<u>Emtriva</u> ® brand <u>emtricitabine</u> , or <u>FTC</u>	<u>Gilead Sciences</u>
<u>enfuvirtide</u> generic Fuzeon™, or T-20	<u>Trimeris and Hoffmann-La Roche</u>
<u>Epivir</u> ® brand lamivudine, or 3TC	<u>GlaxoSmithKline</u>
<u>epoetin alfa</u> (erythropoietin) generic Procrit®	<u>Ortho Biotech</u>

Drug Name	Manufacturer
<u>erythropoietin</u> (epoetin alfa) generic Procrit®	<u>Ortho Biotech</u>
<u>Fortovase®</u> brand saquinavir (Soft Gel Cap), or SQV (SGC)	<u>Hoffmann-La Roche</u>
<u>fosamprenavir</u> , or <u>GW-433908</u> , or <u>VX-175</u>	<u>GlaxoSmithKline</u>
<u>FTC</u> , or <u>Emtriva®</u> brand emtricitabine	<u>Gilead Sciences</u>
<u>Fuzeon™</u> brand enfuvirtide, or T-20	<u>Trimeris</u> and <u>Hoffmann-La Roche</u>
<u>GW-433908</u> , or <u>fosamprenavir</u> , or <u>VX-175</u>	<u>GlaxoSmithKline</u>
<u>HE2000</u> , or <u>alpha-epibromide</u>	<u>HollisEden Pharmaceuticals</u>
<u>HIV-1 Immunogen</u> generic <u>Remune®</u> , <u>Salk vaccine</u> , or <u>AG1661</u>	<u>Immune Response Corp.</u>
<u>Hivid®</u> brand zalcitabine, or ddC	<u>Hoffmann-La Roche</u>
<u>HU</u> , or <u>Droxia®</u> brand hydroxyurea	<u>Bristol-Myers Squibb</u>
<u>hydroxyurea</u> generic <u>Droxia®</u> , or <u>HU</u>	<u>Bristol-Myers Squibb</u>
<u>IDV</u> , <u>Crixivan®</u> brand indinavir, or <u>MK-639</u>	<u>Merck & Co.</u>
<u>IL-2</u> (<u>Interleukin-2</u>), or <u>Proleukin®</u> brand <u>aldesleukin</u>	<u>Chiron Corporation</u>
<u>indinavir</u> generic <u>Crixivan®</u> , <u>IDV</u> , or <u>MK-639</u>	<u>Merck & Co.</u>
<u>Interleukin-2 (IL-2)</u> , or <u>Proleukin®</u> brand <u>aldesleukin</u>	<u>Chiron Corporation</u>
<u>Invirase®</u> brand saquinavir (Hard Gel Cap), <u>SQV (HGC)</u> , or <u>Ro-31-8959</u>	<u>Hoffmann-La Roche</u>
<u>Kaletra®</u> brand lopinavir/ritonavir, or <u>ABT-378/r</u>	<u>Abbott Laboratories</u>
<u>lamivudine</u> generic <u>Epivir®</u> , or <u>3TC</u>	<u>GlaxoSmithKline</u>
<u>lopinavir/ritonavir</u> generic <u>Kaletra®</u> , or <u>ABT-378/r</u>	<u>Abbott Laboratories</u>
<u>MK-639</u> , <u>Crixivan®</u> brand indinavir (<u>IDV</u>)	<u>Merck & Co.</u>
<u>nelfinavir</u> generic <u>Viracept®</u> , <u>NFV</u> , or <u>AG-1343</u>	<u>Pfizer</u>
<u>nevirapine</u> generic <u>Viramune®</u> , <u>NVP</u> , or <u>BI-RG-587</u>	<u>Boehringer Ingelheim</u>

Drug Name	Manufacturer
NFV, Viracept® brand nelfinavir, or AG-1343	<u>Pfizer</u>
Norvir® brand ritonavir (RTV), or ABT-538	<u>Abbott Laboratories</u>
NVP, Viramune® brand nevirapine, or BI-RG-587	<u>Boehringer Ingelheim</u>
<i>PNU-140690, or tipranavir</i>	<u>Boehringer Ingelheim</u>
<i>PRO-542</i>	<u>Progenics Pharmaceuticals</u>
Procrit® brand epoetin alfa (erythropoietin)	<u>Ortho Biotech</u>
<i>Proleukin® brand aldesleukin, or Interleukin-2 (IL-2)</i>	<u>Chiron Corporation</u>
<i>Remune® brand HIV-1 Immunogen, or Salk vaccine</i>	<u>Immune Response Corp.</u>
Rescriptor® brand delavirdine (DLV), or U-90152S/T	<u>Pfizer</u>
Retrovir® brand zidovudine (ZDV), or AZT	<u>GlaxoSmithKline</u>
Reyataz™ brand atazanavir, or BMS-232632	<u>Bristol-Myers Squibb</u>
ritonavir generic Norvir®, RTV, or ABT-538	<u>Abbott Laboratories</u>
RTV, Norvir® brand ritonavir, or ABT-538	<u>Abbott Laboratories</u>
<i>Salk vaccine, Remune® brand HIV-1 Immunogen, or AG1661</i>	<u>Immune Response Corp.</u>
<u>saquinavir (Hard Gel Cap) generic Invirase®, SQV (HGC), or Ro-31-8959</u>	<u>Hoffmann-La Roche</u>
<u>saquinavir (Soft Gel Cap) generic Fortovase®, or SQV (SGC)</u>	<u>Hoffmann-La Roche</u>
<u>SCH-C</u>	<u>Schering-Plough</u>
Serostim® brand somatropin	<u>Serono Laboratories</u>
somatropin generic Serostim®	<u>Serono Laboratories</u>
<u>SQV (HGC), Invirase® brand saquinavir (Hard Gel Cap), or Ro-31-8959</u>	<u>Hoffmann-La Roche</u>
<u>SQV (SGC), or Fortovase® brand saquinavir (Soft Gel Cap)</u>	<u>Hoffmann-La Roche</u>
<u>stavudine generic Zerit®, d4T, or BMY-27857</u>	<u>Bristol-Myers Squibb</u>

Drug Name	Manufacturer
<u>Sustiva®</u> brand efavirenz (EFV)	<u>Bristol-Myers Squibb</u>
<u>T-1249</u>	<u>Trimeris and Hoffmann-La Roche</u>
<u>T-20</u> , or Fuzeon™ brand enfuvirtide	<u>Trimeris and Hoffmann-La Roche</u>
<u>TDF</u> , tenofovir DF generic Viread™, or Bis(POC) PMPA	<u>Gilead Sciences</u>
tenofovir DF (TDF) generic Viread®, Bis(POC) PMPA	<u>Gilead Sciences</u>
<u>tipranavir</u> , or PNU-140690	<u>Boehringer Ingelheim</u>
<u>TMC-114</u>	<u>Tibotec-Virco Group</u>
<u>TMC-125</u>	<u>Tibotec-Virco Group</u>
<u>Trizivir®</u> brand abacavir + zidovudine + lamivudine (ABC + AZT + 3TC)	<u>GlaxoSmithKline</u>
<u>Videx®</u> brand didanosine, ddI, or BMY-40900	<u>Bristol-Myers Squibb</u>
<u>Videx® EC</u> brand didanosine (ddI): delayed-release capsules	<u>Bristol-Myers Squibb</u>
<u>Viracept®</u> brand nelfinavir (NFV), or AG-1343	<u>Pfizer</u>
<u>Viramune®</u> brand nevirapine (NVP), or BI-RG-587	<u>Boehringer Ingelheim</u>
<u>Viread®</u> brand tenofovir DF, or Bis(POC) PMPA	<u>Gilead Sciences</u>
<u>VX-175</u> , or fosamprenavir, or GW-433908	<u>GlaxoSmithKline</u>
<u>zalcitabine</u> generic Hivid®, or ddC	<u>Hoffmann-La Roche</u>
<u>ZDV</u> , Retrovir® brand zidovudine, or AZT	<u>GlaxoSmithKline</u>
<u>Zerit®</u> brand stavudine, d4T, or BMY-27857	<u>Bristol-Myers Squibb</u>
<u>Ziagen®</u> brand abacavir (ABC), or 1592U89	<u>GlaxoSmithKline</u>
<u>zidovudine</u> generic Retrovir®, AZT, or ZDV	<u>GlaxoSmithKline</u>

Additional drugs in clinical trials that can be used in combination and/or alternation with the 3-phosphoindoles include:

PHASE I	PHASE II	PHASE III
GW5634 (GSK)	MIV-150 (Medivir/Chiron)	Tipranavir (B-I)
RO033-4649 (Roche)	TMC125 (Tibotec)	
GW640385 (GSK/Vertex)	TMC114 (Tibotec)	
Elvucitabine (Achillion Ph.)	Alovudine (FLT) (B-I)	
MIV-210 (GSK/Medivir)	Racivir (Pharmasset)	
SPD754 (Shire Pharm.)	Reverset (Incyte Corp.)	
FP21399 (Fuji Pharm.)	AMD070 (AnorMed)	
GW873140 (GSK)	BMS-488043 (BMS)	
Schering C/D (417690)	PRO 542 (Progenics Pharm)	
	TAK-220 (Takeda)	
	TNX-355 (Tanox)	
	UK-427,857 (Pfizer)	

The following drugs have been approved by the FDA for use in the treatment of complications of HIV infection and AIDS, which can be used in combination and/or alternation with the compounds of the present invention.

Brand Name	Generic Name	Use	Manufacturer Name
Abelcet, Ambisome	Amphotericin B, ABLC	antifungal for aspergillosis	various
Bactrim, Septra	sulfamethoxazole and trimethoprim	antiprotozoal antibiotic for Pneumocystis carinii pneumonia treatment and prevention	various
Biaxin, Klacid	clarithromycin	antibiotic for Mycobacterium avium prevention and treatment	Abbott Laboratories
Cytovene	ganciclovir, DHPG	antiviral for CMV retinitis	Roche
DaunoXome	daunorubicin-liposomal	chemotherapy for Kaposi's sarcoma	Gilead
Diflucan	fluconazole	antifungal for candidiasis,	Pfizer

Brand Name	Generic Name	Use	Manufacturer Name
		cryptococcal meningitis	
Doxil	doxorubicin hydrochloride-liposomal	chemotherapy for Kaposi's sarcoma	Ortho Biotech
Famvir	famciclovir	antiviral for herpes	Novartis
Foscarnet	foscavir	antiviral for herpes, CMV retinitis	Astra Pharmaceuticals
Gamimmune N	immune globulin, gamma globulin, IGIV	immune booster to prevent bacterial infections in children	Bayer Biologicals
Intron A	interferon alfa-2b	Karposi's sarcoma, hepatitis C	Schering
Marinol	dronabinol	treat appetite loss	Roxane Laboratories
Megace	megestrol acetate	treat appetite, weight loss	Bristol Myers-Squibb
Mepron	atovaquone	antiprotozoal antibiotic for Pneumocystis carinii pneumonia treatment and prevention	GlaxoSmithKline
Mycobutin, Ansamycin	rifabutin	antimycobacterial antibiotic for Mycobacterium avium prevention	Adria Pharmaceuticals
NebuPent	pentamidine	antiprotozoal antibiotic for Pneumocystis carinii pneumonia prevention	Fujisawa
Neutrexin	trimetrexate glucuronate and leucovorin	antiprotozoal antibiotic for Pneumocystis carinii pneumonia treatment	MedImmune
Panretin gel	alitretinoin gel 0.1%	AIDS-related Karposi's sarcoma	Ligand Pharmaceuticals
Procrit, Epogen	erythropoetin, EPO	treat anemia related to AZT therapy	Amgen
Roferon A	interferon alfa-2a	Karposi's sarcoma and hepatitis C	Roche
Serostim	somatropin rDNA	treat weight loss	Serono
Sporanox	itraconazole	antifungal for blastomycosis,	Janssen Pharmaceuticals

Brand Name	Generic Name	Use	Manufacturer Name
		histoplasmosis, aspergillosis, and candidiasis	
Taxol	paclitaxel	Karposi's sarcoma	Bristol Myers-Squibb
Valcyte	valganciclovir	antiviral for CMV retinitis	Roche
Vistide	cidofovir, HPMPC	antiviral for CMV retinitis	Gilead
Vitrasert implant	ganciclovir insert	antiviral for CMV retinitis	Bausch & Lomb
Vitravene intravitreal injectable	fomivirsen sodium injection	antiviral for CMV retinitis	Isis Pharmaceuticals
Zithromax	azithromycin	antibiotic for Mycobacterium avium	Pfizer

Several products have been allowed to proceed as Investigational New Drugs (IND) by the FDA for the treatment of complications of HIV infection and AIDS. Therefore, the following drugs can be used in combination and/or alternation with the compounds of the present invention.

- Trimetrexate glucuronate for the treatment of Pneumocystis carinii pneumonia in AIDS patients who cannot tolerate standard forms of treatment.
- Ganciclovir for the treatment of cytomegalovirus retinitis in AIDS patients.
- Aerosolized pentamidine for the prevention of Pneumocystis carinii pneumonia in AIDS patients.
- Erythropoietin for the treatment of zidovudine-related anemia.
- Atovaquone for the treatment of AIDS patients with Pneumocystis carinii pneumonia who are intolerant or unresponsive to trimethoprim-sulfamethoxazole.
- Rifabutin for prophylaxis against Mycobacterium avium complex bacteremia in AIDS patients.

- Vistide intravenous cidofovir for HIV-infected persons with relapsing cytomegalovirus (CMV) retinitis that has progressed despite treatment (Hoffmann-La Roche).
- Serostim, a mammalian derived recombinant human growth hormone, for the treatment of AIDS-related wasting (Serono Laboratories).

In general, during alternation therapy, an effective dosage of each agent is administered serially. During combination therapy, effective dosages of two or more agents are administered together. Dosages administered depend upon factors such as absorption, biodistribution, metabolism and excretion rates for each drug as well as other factors known to those skilled in the art. It is to be noted that dosage amounts will vary with the severity of the condition to be alleviated, the age, weight, and general physical condition of the subject who receives the drug. It is to be understood further that for any particular subject, specific dosage regimens and schedules should be adjusted over time according to the response of the subject to the drug, the needs of the subject, and the professional judgment of the person administering or supervising the administration of the compositions. Examples of suitable dosage ranges for anti-HIV compounds, including nucleoside derivatives such as, for example, D4T, DDI and 3TC, or protease inhibitors like nelfinavir and indinavir, are to be found in the scientific literature and Physicians' Desk Reference. Suggested ranges for effective dosages of the compounds of the present invention are guidelines only, and are not intended to limit the scope or use of the invention.

The disclosed combination and alternation regimen are useful in the treatment and prevention of retroviral infections and other related conditions, such as, for example, AIDS-related complex (ARC), persistent generalized lymphadenopathy (PGL), AIDS-related neurological conditions, anti-HIV antibody positive and HIV-positive conditions, Kaposi's sarcoma, thrombocytopenia purpura, and opportunistic infections. In addition, these compounds or formulations can be used prophylactically to prevent or retard the progression of clinical illness in individuals who are anti-HIV antibody or HIV-antigen positive, or who have been exposed to HIV.

VII. Pharmaceutical Compositions

The indole compounds of the present invention can be administered to a subject in need thereof, optionally in combination or alternation with another anti-HIV or anti-retroviral agent, and/or with a pharmaceutically acceptable carrier, diluent or excipient. In one embodiment, a subject infected with HIV may be treated by administering to that subject an effective amount of an indole derivative, a salt, prodrug, stereoisomer or tautomer thereof, in the presence of a pharmaceutically acceptable carrier or diluent. For subjects with multiple drug resistance, the oxo-pyrimidine compound is administered either alone or in combination with one or more other anti-retroviral agents or anti-HIV agents. The active compounds may be administered by any appropriate route, for example, orally, parenterally, enterally, intravenously, intradermally, subcutaneously, percutaneously, transdermally, intranasally, topically or by inhalation therapy, and may be in solid, liquid or vapor form.

The active compound(s) are included within the pharmaceutically acceptable carrier, diluent or excipient in an amount sufficient to deliver to a patient a therapeutically effective amount of the active compound in order to inhibit viral replication *in vivo*, especially HIV replication, without causing serious toxic effects in a treated subject. By an "inhibitory amount" is meant an amount of active ingredient sufficient to halt viral replication as measured by, for example, an assay such as the ones referred to herein.

One dose of the indole compound for all the conditions mentioned is in the range of from about 0.1 to 100 mg/kg of body weight per day, or from about 1 to 75 mg/kg of body weight per day, and even more typically from about 1 to 20 mg/kg of body weight per day. The effective dosage range of the pharmaceutically acceptable derivatives is calculated based on the weight of the parent indole derivative compound to be delivered. If the derivative compound itself exhibits activity, then the effective dosage can be estimated as above using the weight of the derivative, or by other means known to those of skill in the art.

The compounds are conveniently administered in units of any suitable dosage form, including but not limited to one containing from about 7 to 3000 mg, or from about 70 to 1400 mg, and even more typically from about 25 to 1000 mg of active ingredient

per unit dosage form. For example, an oral dosage of from about 50 to 1000 mg is usually convenient.

Ideally, the active ingredient is administered to achieve peak plasma concentrations of the active compound of from about 0.02 to 70 μM , and typically of from about 0.5 to 10 μM . For example, this can be achieved by intravenous injection of a 0.1 to 25% solution of active ingredient, optionally in saline, or administered as a bolus of active ingredient. It is to be understood that for any particular subject, specific dosage regimens should be adjusted over time to meet individual needs. The concentrations set forth here are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered all at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

One mode of administration of the active compound is oral. Oral compositions usually include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules, compressed into tablets, or delivered in liquid form. For oral therapeutic administration, the active compound may be incorporated with excipients or formulated as solid dispersions or solid solutions, and used in the form of tablets, troches, or capsules. By a "solid dispersion" is meant a solid state comprising at least two components where one component is dispersed more or less evenly throughout the other component. By "solid solution" is meant a solid state comprising at least two components that are chemically and physically integrated to produce a homogeneous product. A solid solution is typical over a solid dispersion because it more easily forms a liquid solution upon contact with an appropriate liquid medium, thereby increasing the bioavailability of a drug. Pharmaceutically compatible binding agents and/or adjuvant materials also may be included as part of this composition.

The tablets, pills, capsules, troches and the like may contain any of the following ingredients or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose; a disintegrating agent such as alginic acid, Primogel, or cornstarch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent like sucrose or saccharin; and a flavoring agent, such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it may contain a liquid carrier such as a fatty oil in addition to any material of the kinds given above. In

addition, dosage unit forms may contain various other materials that modify the physical form of the dosage unit, such as, for example, coatings of sugar, shellac, or other enteric agents.

The indole compounds may be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain sucrose as a sweetening agent, preservatives, dyes, colorings, and flavorings in addition to the active compounds.

The active compounds or their pharmaceutically acceptable salts or prodrugs can be mixed with other active materials that do not impair the desired action, or with materials that supplement the desired action, such as antibiotics, antifungals, anti-inflammatories, protease inhibitors, or other nucleoside or non-nucleoside antiviral agents. Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates, or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. A parenteral preparation normally will include sterile water and may be enclosed in ampoules, disposable syringes, or multiple dose vials made of glass or plastic.

If administered intravenously, typical carriers are physiological saline, phosphate buffered saline (PBS), a glucose solution, or a mixed solution comprising glucose and saline. If administration is percutaneous, such as, for example, through the use of a patch or ointment, the associated carrier may comprise a penetration-enhancing agent and/or a suitable wetting agent which are not harmful to the skin. If inhalation or insufflation is the desired route of administration, then the composition of the present invention includes the compound in the form of a solution, suspension or dry powder that can be delivered through the oral and/or nasal orifices.


Liposomal suspensions, which include liposomes targeted to infected cells with monoclonal antibodies to viral antigens, also are typical as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art,

for example, as described in U.S. Patent No. 4,522,811, which is incorporated herein by reference in its entirety. For example, liposomal formulations may be prepared by dissolving appropriate lipid(s) such as stearyl phosphatidyl ethanolamine, stearyl phosphatidyl choline, arachadoyl phosphatidyl choline, and cholesterol, in an inorganic solvent that later is evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the active compound, or a salt or prodrug thereof, is then introduced into the container. The container is swirled to free lipid material from its sides and to disperse lipid aggregates, thereby forming the liposomal suspension.


VIII. Processes for Preparing the Active Compounds

General Schemes


Scheme 1:




Scheme 2:



Scheme 3:




Scheme 4:







5

Scheme 5:




Scheme 6:

Scheme 7:Scheme 8:Scheme 9:




Scheme 10:




Particularized Schemes

Scheme 11: N-branched Heterocycles




5


Scheme 12: Phosphinate Heterocycles




Scheme 13: Alkyl/Alkenyl/Alkynyl Phosphinates




Scheme 14: Alkyl Phosphinates



Scheme 15: Alkyl Methyl Carboxamides



Scheme 16: Phosphinic Acids



Methods

The following are descriptions of the methodologies used in the foregoing schemes. Numbers in bold print correspond to numbered compounds found in Schemes 1-16. Methods A, B, C, D, F, L, N, T, U and AA are known to those skilled in the art of making this general class of compounds, but have been adapted for preparing the specific compounds of the present invention. This also is true for Methods H, I, J, M, P, Q, Y, Z, AB and AC. Methods E, G, K, O, R, S, W, V and X are new processes for preparing the compounds of the invention.

METHOD A

Bromobenzene **1** (1 eq) was stirred with diethylphosphite (1.1 eq.) and triethylamine (1.1 eq.) in a pressure tube and the mixture was degased with N₂ for about 15 minutes. Then palladium tetrakis (0.05 eq.) was added and the mixture was stirred under pressure at about 85°C overnight. The next morning, Et₂O was added to the reaction and the mixture was filtered. The filtrate was evaporated and the crude was purified by chromatography on silica gel (eluent: petroleum ether/EtOAc from 8/2 to 6/4). Alternatively, the reaction can be performed using 10 volumes of 1,2-diethoxyethane under the same conditions.

METHOD B

All the reagents (1 eq. of diethylphosphonate **2** or dimethylphosphonate **19**, 6 eq. of sodium hydroxide and ethanol or methanol (3ml/mmol)) were stirred at room temperature for about 5 hours. Then, ethanol (or methanol) was evaporated in vacuo and the mixture acidified with HCl 2.5N to reach pH = 1. Then the mixture was saturated with NaCl and extracted with ethyl acetate. Combined organic layers were dried over Na₂SO₄, filtered and concentrated to give the desired products **3** or **20**.

METHOD C

The compound **20** or **3** (1.2 eq.) was stirred at room temperature under N₂ with thionyl chloride (3.6 eq.), dichloromethane (5ml/mmol) and a few drops of dimethylformamide for about 5 hours. An aliquot was taken and put in dry methanol and Et₃N to monitor the reaction. When the reaction was complete, dichloromethane and

thionyl chloride, co-evaporated with toluene, were evaporated in vacuo to give an oil which was stored under N₂.

METHOD D

To a solution of arylphosphonyl dichloride **21** (1 mL, 6.35 mmol) in anhydrous dichloromethane (25 mL) ethanol or methanol (1,12 mL, 19.04 mmol) was added dropwise at about 0°C followed by the addition of triethylamine (2,65 mL, 19.04 mmol). The reaction mixture was stirred at room temperature for about 2 hours, and then was washed with a solution of HCl 1N (50 mL). The aqueous layer was extracted with dichloromethane. Combined organic phases were dried and concentrated under reduced pressure. The crude oil was purified by chromatography on silica gel (eluent: C₆H₁₂/EtOAc 6/4) to afford dimethylarylphosphonate **19** or diethylarylphenylphosphonate **2**.

METHOD E

The diethylarylphosphonate **2** (1 eq.) was stirred under N₂ at room temperature with bromotrimethylsilane (10 eq.) and dichloromethane (5ml/mmol) for about 5 hours. Then the bromotrimethylsilane and dichloromethane were evaporated to give a yellow oil. Next, oxalyl chloride, dimethylformamide and dichloromethane were added under N₂ and the mixture was stirred at room temperature overnight, following which the oxalyl chloride and solvents were evaporated. The oil in dichloromethane solution was stirred and allowed to cool to about 0°C, and ethanol or methanol (1,12 mL, 19.04 mmol) was added under N₂, followed by the dropwise addition of triethylamine. The mixture then was warmed to room temperature. The solvent was evaporated and the resulting product was purified by chromatography on silica gel (eluent: CH₂Cl₂/EtOAc : 8/2) to afford the compound **14**.

METHOD F

To a stirred solution of diethylarylphosphonate **2** (1 eq.) in dichloromethane (5ml/mmol) was added bromotrimethylsilane (5 eq.) at room temperature under N₂. After about 3 hours of stirring at room temperature, the solvent was evaporated and dichloromethane (5ml/mmol), a few drops of dimethylformamide and oxalyl chloride (2,5eq) were added. This mixture was stirred at room temperature overnight, and the solvent was evaporated to give an oil. A solution of the oil (1.2 eq.) in diethyl ether

(3ml/mmol) was stirred and cooled to about -17°C, after which anhydrous methanol (1.2 eq.) was added dropwise to remove the double addition components and triethylamine. The resulting mixture was warmed to room temperature, stirred for about 1 hour, and then filtered on autocup under N₂ to remove triethylamine salts. The solvent was evaporated to give the compound **15**.

METHOD G

n-BuLi (2.5M in hexane, 1.2 eq.) was added dropwise to a stirred and cooled (to about -90 °C) solution of bromoindole **11** (1 eq.) in anhydrous THF (10ml/mmol) under N₂. After keeping the solution at about -90°C for about 5 minutes, an appropriate chlorophosphorus reagent **15**, **4** or diphenylphosphonic chloride (1.2 eq.) was added dropwise to the solution at the same temperature. The reaction was allowed to warm slowly to about -40°C (TLC monitoring, eluent CH₂Cl₂/ ETOAc 9/1). Water then was added. Extraction with ethyl acetate, drying and evaporation led to a crude oil that was purified by chromatography on silica gel to give the compound **16**, **5** or **26**.

METHOD H

A stirred and cooled (to about 0°C) solution of compound **5**, **16**, **26**, **28** or **30** in methanol in a pressure tube was saturated with NH₃ gas for about 10 minutes. Then the mixture was stirred at about 50° C overnight, and after TLC monitoring, excess ammonia and methanol were evaporated in vacuo and the crude purified by chromatography on silica gel to give the carboxamide **6**, **7**, **27**, **29** or **31**.

METHOD I

Lithium hydroxide (14 eq.) was added to a stirred solution of compound **16** or **5** in tetrahydrofuran (20ml/mmol) and water (20ml/mmol). This mixture next was stirred at room temperature and monitored by TLC. If necessary, equivalents of lithium hydroxide were added until the reaction was complete. Then the THF was evaporated and HCL (1N) was added to reach pH 1. The aqueous layer was extracted with ethyl acetate and combined organic phases were dried, filtered and concentrated under reduced pressure to give the compound **17** or **8**.

METHOD J

The compound **17** or **8** (1 eq.) was stirred with dichloromethane (20ml/mmol) or DMF, and 1-hydroxybenzotriazole (1 eq.) was added followed by the addition of EDCI (1 eq.) followed by the amine (1eq) were added. This mixture was stirred at room temperature overnight. The next morning the mixture was washed with water (to a pH of 5-6), and organic layers were dried, filtered and concentrated under reduced pressure. The resulting oil was purified by chromatography on silica gel (eluent: CH₂Cl₂/EtOAc) to give a powder **18** or **9**.

METHOD K

In a microwave tube, the compound **5** or **6** (1eq.) was stirred with DMF (5ml/mmol), and TMSBr (5 eq.) was added. The tube was heated under microwave irradiations under pressure at about 60 °C (maximum power input 100W, CEM discover apparatus) for about 50 minutes. DMF was evaporated in vacuo, and the mixture was put in a pressure tube. Trimethylphosphite (4ml/mmol) was added and the mixture was stirred and heated at about 90 °C overnight. The mixture then was cooled in an ice water bath and HCl (1N) was added dropwise. The mixture was extracted with ethyl acetate, and combined organic layers were washed with HCl (1N) until no HP(OMe)₂ remained. Next it was dried, filtered and concentrated under reduce pressure to provide an oil, and the resulting oil was purified by chromatography on silica gel to give the compound **16** or **7**.

METHOD L

To a stirred and cooled (to about 0°C) solution of ethyl indole-2-carboxylate **10** (1 eq.) in DMF (2ml/mmol) under N₂, was added NaH (60% in oil, 1.2 eq.) portionwise. When gas evolution stopped, benzenesulfonyl chloride (1.2 eq.) was added. The reaction mixture was stirred for about 1 hour (TLC monitoring, eluent dichloromethane); a small amount of water then was added carefully and the DMF was evaporated. The crude residue was dissolved in ethyl acetate and washed with water and brine. After drying and evaporation of the solvents, the compound was purified by chromatography on silica gel (eluent: C₆H₁₂/EtOAc 9/1 to 7/3) to give the ethyl 1-phenylsulfonylindole-2-carboxylate.

To a stirred solution of ethyl 1-phenylsulfonylindole-2-carboxylate (1 eq.) in DMF (2.5ml/mmol) under N₂, was added a solution of bromine (4 eq.) in DMF

(0.5ml/mmol). This reaction mixture was stirred at room temperature for about 4 hours, following which water was added and the mixture was extracted with dichloromethane (x3). The organic layer was washed with a saturated solution of Na₂SO₅, dried and evaporated to give a crude yellow oil. Purification by chromatography on silica gel (eluent: C₆H₁₂/EtOAc 9/1) afforded 3-brominated indole **11**.

METHOD M

A mixture of **5** or **16** (1 eq.), acrylonitrile (10 eq.), palladium acetate (20% mol.), triethylamine (1 eq.) and tri-*ortho*-tolylphosphine (1 eq.) in degassed acetonitrile (30mL/mmol) was stirred and heated under microwave irradiation in a pressure tube for about 45 minutes. Next water was added and the aqueous layer was extracted with dichloromethane. The combined organic layers were dried and concentrated and purified by chromatography on silica gel (eluent: Petroleum ether/EtOAc 8/2) to give the compound **32** as a mixture of enantiomers E and Z (which are separated by preparative HPLC).

METHOD N

Bromobenzene **1** (1 eq.), dimethylformamide (1ml/mmol), triethylamine (3 eq.) and anilinium salt* (1.25 eq.) were put in a pressure tube and degassed with N₂ for about 15 minutes. Then palladium tetrakis was added and this mixture was stirred at about 85°C overnight, following which the solvent was evaporated and water was added to achieve a pH of about 5-6. The mixture was basified with NaHCO₃ until a pH 8 was reached, and then extracted with diethyl ether. The aqueous layer was acidified with HCl (1N) to reach a pH = 1, and extracted with ethyl acetate. Combined organic layers were dried, filtered and concentrated under reduced pressure to give the compound **14**.

*Anilinium salt was synthesised according to the procedure of Montchamp et al (*J. Am. Chem. Soc.*, **2001**, *123*, 510-511).

METHOD O

Tetramethylorthosilicate (1.2 eq.) was added to a stirred solution of compound **14** (1.2 eq.) in toluene (4ml/mmol) under N₂. This mixture was heated at reflux for about 1.5 hours, and then cooled to room temperature. Next the mixture was degassed with N₂ before adding triethylamine (3.3 eq.), bromoindole **11** (1 eq.), and palladium tetrakis (0.05 eq.). The mixture was stirred at about 100°C over a week-end (about 48 hours),

after which the reaction was cooled to room temperature and water was added to achieve a pH of about 8-9. The mixture next was extracted with ethyl acetate and the organic phases were washed with KHSO_4 (1N) saturated with NaCl. It then was dried, filtered and concentrated under reduced pressure to give an oil, which was purified by chromatography on silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to give the compound **12**.

METHOD P

To a stirred solution of compound **12** (1 eq.) in methanol (25ml/mmol) was added trimethylsilyldiazomethane (13 eq.) in a several fractions at room temperature under N_2 . This mixture was stirred overnight, after which water was added and the methanol was evaporated under reduce pressure. Then NaHCO_3 was added to reach pH 8, and the mixture was extracted with ethyl acetate. The organic phases were dried, filtered and concentrated under reduced pressure. The crude was purified by chromatography on silica gel (eluent: $\text{C}_6\text{H}_{12}/\text{EtOAc}$) to give the compound **16**.

METHOD Q

The compound **18** was dissolved in chloroform (or in CH_2Cl_2) under stirring at room temperature; m-chloroperoxybenzoic acid was added and the reaction allowed to stir overnight (about 15 hours). Then the mixture was diluted with dichloromethane and extracted with a mixture of saturated K_2CO_3 / H_2O (1/3). The aqueous layer was extracted three times with dichloromethane. The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure (TLC: dichloromethane/methanol = 9/1). The crude product was purified by chromatography on silica gel to give the compound **22**.

METHOD R

In a microwaves sealed tube, the compounds **11** (1 eq.) and **23** (1.1 eq.) were added to toluene (8ml/mmol) and triethylamine (3.3 eq.) and degased with N_2 for about 10 minutes. Then $\text{Pd}(\text{PPh}_3)_2$ was added and the tube was heated under microwave irradiation under pressure at about 120°C (maximum power input 200W, CEM discover apparatus) for about 30 minutes. The reaction was monitored by TLC and the tube was heated for about 30 minutes if necessary. The solvent was evaporated and the crude product was purified by chromatography on silica gel (eluent: $\text{C}_6\text{H}_{12}/\text{EtOAc}$: 8/2) to give the compound **5**.

METHOD S

In a microwave sealed tube, the compounds **11** (1 eq.) and **24** (2 eq.) were added to toluene (8ml/mmol) and then degased with N₂ over about a 10 minute time period. Next, Pd(OAc)₂ in about 20% solution was added, and the tube was under microwave irradiation under pressure at about 150 °C (maximum power input 200W, CEM discover apparatus) for about 45 minutes. The reaction was monitored by TLC and if any starting material was present, the tube was heated for about another 45 minutes at approximately 170°C. Next HCl (1N) (8ml/mmol) was added and the solution was extracted with ethyl acetate. The combined organic layers were dried, filtered and concentrated under reduced pressure. The crude product was purified by chromatography on silica gel (eluent: petroleum ether/EtOAc : 1/1) to give the compound **5** or **16**.

Alternatively the reaction is carried out at about 150°C in xylene for about 5 hours.

METHOD T

Pyridine (1 eq.) was carefully added to a vigorously stirred solution of alkyl chloroformate (1 eq.) and arylphosphinic acid (1 eq.) in dichloromethane (2ml/mmol) at room temperature. Once effervescence stopped, the solution was refluxed for about 15 minutes and then allowed to cool to room temperature. The solution was poured into 0.1M hydrochloric acid (1ml/mmol) and the organic layer was separated. After washing with water and drying over Na₂SO₄, the solvent was removed in vacuo to give the compound **23**.

METHOD U

Alkyl or aryl bromide (0.15 mol) was added dropwise to a mixture of magnesium (3.6 g) and dry tetrahydrofuran (40 mL) under N₂ atmosphere at about 50°C. After this addition, the reaction mixture was stirred at about 50°C. for an additional 1–2 hours to allow the reaction to run to completion. Then the mixture was added dropwise to a solution of triethyl phosphite (0.1 mol) and THF (25 mL) at between 40 °C. to 50 °C. in N₂ atmosphere, and stirred for about 3 hours at about 50°C. After removal of the solvent under reduced pressure, the crude product was distilled from the semisolid residue *in vacuo* to give the compound **24**.

METHOD V

The compound **6** or **7** (1 eq.) and Lawesson's reagent (4 eq.) were heated in toluene (10ml/mmol) at about 90 °C. under N₂ in a pressure tube. The reaction was monitored by TLC and heating continued until no starting material remained (about 5.5 hours). The crude solution was filtered and the filtrate was evaporated to dryness and purified by chromatography on silica gel to give the compound **25** and the compound **25'**.

METHOD W

To a stirred and cooled (to about -90 °C.) solution of bromoindole **11** (1 eq.) was added dropwise n-butyllithium (1.2 eq.) under N₂. After about 10 minutes, benzenephosphonyl dichloride **21** (1.1 eq.) in tetrahydrofuran (15ml/mmol) was added dropwise at a temperature of about -70 °C., and the temperature then was raised to about -90 °C. and maintained for about 15 minutes. This was followed by the addition of methylmagnesium bromide (1.1 eq.), and the mixture was allowed to warm to about -40°C for about 1 hour, after which it was quenched with water and extracted with ethyl acetate then dichloromethane. The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by chromatography on silica gel (eluent: CH₂Cl₂/AcOEt : 9/1 to 7/3) to give the compound **28**.

METHOD X

Dimethylamine (1 eq.) was added dropwise to a stirred and cooled (to about -55 °C.) solution of phenyl phosphonic dichloride **21** (1 eq.) in diethyl ether (1.5ml/mmol). Then triethylamine was added (1 eq.) and the mixture was allowed to warm to room temperature. The mixture next was filtered and the filtrate was evaporated to give an oil/mixture. To rid the oil/mixture of phenyl phosphonic dichloride, the oil/mixture was dissolved in EtOAc and washed twice with an HCl solution of pH 4-5. The organic phases were dried with Na₂SO₄, filtered and concentrated under reduced pressure to give an oil. Then, n-butyllithium (1.2 eq.) was added dropwise to a stirred and cooled (to about -80 °C.) solution of bromoindole **11** (1 eq.) in tetrahydrofuran (5ml/mmol) under N₂. At the end of the addition, the mixture was warmed to about -60 °C. and tetrahydrofuran (1.2 eq.; 3ml/mmol) was added dropwise to the oil. Then the mixture was warmed slowly to about -10 °C, and the reaction was quenched with water (8ml/mmol). HCl (1N) was added to achieve a pH of about 5, and the solvent was

evaporated *in vacuo*. The aqueous phase was extracted with ethyl acetate and the combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by chromatography on silica gel (eluent: C₆H₁₂/EtOAc : 9/1 to 0/10) to give the compound **30**.

METHOD Y Suzuki cross-coupling

Pd(PPh₃)₄ (10% mol.), a solution of Na₂CO₃ (6 eq.) in H₂O (2M) and a solution of aryl, alkyl or heteroaryl boronic acid (2 eq.) in EtOH (0.3M) were added to a stirred solution of **5** or **16** (bearing an halogeno or triflate substituent) in degassed toluene under N₂. The reaction tube was then heated under microwave irradiation at about 110 °C. until no starting material remained. Water was added and the reaction media was extracted with EtOAc, dried and concentrated. The crude residue was purified by column chromatography on silica gel (PE/EtOAc 8/2) to afford **37**.

METHOD Z

Under a nitrogen atmosphere, a dry reaction tube was charged with cuprous oxide (10%mol), a ligand (20%mol), a nucleophile (1.5 eq.), cesium carbonate (2 eq.) and the aryl halide **5** or **16** (bearing an halogeno or triflate substituent) (1 eq.), followed by the addition of anhydrous and degassed acetonitrile (0.6 mL per mmol of aryl halide). The tube was sealed and stirring was applied at about 80 °C. until the reaction ran to completion. The reaction mixture then was cooled to room temperature, diluted with *tert*-butylmethyl ether and filtered through a plug of celite, the filter cake being further washed with butylmethyl ether. The filtrate was concentrated *in vacuo* to remove the acetonitrile and was redissolved in *tert*-butylmethyl ether. This organic layer filtrate was washed twice with water and once with brine before being dried on Na₂SO₄ and filtered. The solvent was removed *in vacuo* to yield the crude product, which was purified by chromatography on silica gel to give the compound **42**.

METHOD AA

Alkyl halide was heated at about 90 °C. overnight in trimethylphosphite (10mL/mmol). The reaction media was cooled to about 0°C in an ice bath, and a solution of HCl (1N) was added carefully. The aqueous layer was extracted with EtOAc. The

combined organic layers were washed with HCl (1N) and with water, and then dried and concentrated to give compound **53** as a colorless oil.

METHOD AB

In a microwave sealed tube, the compounds **17** (1 eq.) , formaldehyde (37% wt in water, 1 eq.), and morpholine (1 eq.) were added to *t*-butyl alcohol (4ml/mmol). The tube was heated under microwave irradiation under pressure at about 170 °C (maximum power input 200W, CEM discover apparatus) for about 60 minutes. The reaction was monitored by TLC and if any starting material remained, the tube was heated for about 45 minutes at approximately 170 °C. The solvents were evaporated *in vacuo*, and the crude residue was purified by chromatography on silica gel(MeOH/EtOAc : 2/98) to give the compound **18**.


METHOD AC







In a microwave tube, the compound **5** or **6** (1 eq.) was stirred with DMF (5ml/mmol), and TMSBr (5 eq.) was added. The tube was heated under microwave irradiation under pressure at about 60 °C. (maximum power input 100W, CEM discover apparatus) for about 50 minutes. After cooling, water was added and compound **60** was collected by filtration.








IX. Representative Examples of the Active Compounds








Tables 1 and 2 contain a non-limiting list of representative compounds that may be prepared by the methods and according to Schemes 1-16 provided above.









Table 1









Compound	structure	Description
2a		Colorless oil, ¹ H NMR (CDCl ₃ , 250 MHz) δ 1.33 (t, <i>J</i> = 7.1 Hz, 6H), 4.05-4.25 (m, 4H), 7.46-7.57 (m, 3H), 7.78-7.87 (m, 2H), ³¹ P NMR (CDCl ₃ , 101.256 MHz) δ 19.3.








Compound	structure	Description
2c		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.26 (t, $J = 7.5$ Hz, 3H), 1.33 (t, $J = 7.05$ Hz, 6H), 2.7 (q, $J = 7.5$ Hz, 2H), 4.04-4.19 (m, 4H), 7.37-7.4 (m, 2H), 7.58-7.69 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.53.
2d		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.35 (t, $J = 7.05$ Hz, 6H), 4.05-4.26 (m, 4H), 7.62 (td, $J = 3.9$ and 7.8 Hz, 1H), 7.8-7.83 (m, 1H), 7.97-8.1 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 16.4, ^{19}F NMR (CDCl_3 , 282.4 MHz) δ 9.27 (s, 3F), MS (ESI, EI^+) $m/z = 283$ (MH^+).
2e		White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, $J = 7.05$ Hz, 6H), 4.09-4.25 (m, 4H), 7.62 (td, $J = 3.9$ and 7.8 Hz, 1H), 7.83-7.85 (m, 1H), 8.02-8.12 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.12, MS (ESI, EI^+) $m/z = 240$ (MH^+).
2f		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.34 (t, $J = 7.05$ Hz, 6H), 4.05-4.21 (m, 4H), 7.32-7.38 (m, 1H), 7.66-7.71 (m, 1H), 7.73-7.78 (m, 1H), 7.92-7.97 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 16.37, MS (ES^+) $m/z = 292.9/294.8$ (MH).
2g		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.35 (t, $J = 7.05$ Hz, 6H), 4.04-4.23 (m, 4H), 7.53 (t, $J = 1.8$ Hz, 1H), 7.65 (d, $J = 1.8$ Hz, 1H), 7.7 (d, $J = 1.8$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 14.74.
2h		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.34 (t, $J = 7.05$ Hz, 6H), 4.06-4.22 (m, 4H), 5.44 (d, $J = 47.4$ Hz, 2H), 7.51-7.6 (m, 2H), 7.78-7.85 (m, 2H), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -209.67 (t, $J = 47.4$ Hz, 1F), MS (ES^+) $m/z = 247$ (MH).








Compound	structure	Description
2i		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.34 (t, $J = 7.05$ Hz, 6H), 4.04-4.2 (m, 8H), 5.86 (s, 1H), 7.48-7.54 (m, 1H), 7.68-7.7 (m, 1H), 7.8-7.87 (m, 1H), 7.93-7.98 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 18.45, MS (ES^+) $m/z = 287$ (MH).
2j		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.35 (t, $J = 7.05$ Hz, 6H), 2.39 (s, 3H), 4.05-4.21 (m, 4H), 7.34-7.36 (m, 1H), 7.5-7.6 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 17.18, MS (ES^+) $m/z = 263.2$ (MH).
2k		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.37 (t, $J = 7.05$ Hz, 6H), 4.07-4.27 (m, 4H), 7.37-7.65 (m, 6H), 7.78-7.85 (m, 2H), 8.04-8.09 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 18.84, MS (ES^+) $m/z = 291.17$ (MH).
2l		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 6H), 2.15 (s, 3H), 4.06-4.22 (m, 4H), 5.44 (s, 2H), 7.38-7.60 (m, 3H), 7.95-8.03 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 17.99, MS (ESI, EI^+) $m/z = 287$ (MH^+).
2m		Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 6.9$ Hz, 3H), 1.323 (t, $J = 6.9$ Hz, 3H), 3.84 (s, 3H), 4.03-4.16 (m, 4H), 7.06-7.1 (m, 1H), 7.3-7.4 (m, 3H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.21, MS (ESI, EI^+) $m/z = 245.13$ (MH^+).
2q		Slight yellow oil, ^1H NMR (CDCl_3 , 400 MHz) δ 1.37 (t, $J = 7.05$ Hz, 6H), 4.11-4.28 (m, 4H), 8.046-8.048 (m, 1H), 8.237-8.24 (m, 1H), 8.27-8.273 (m, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 14.36, MS (ESI, EI^+) $m/z = 351$ (MH^+).
2r		Slight yellow oil, ^1H NMR (CDCl_3 , 400 MHz) δ 1.36 (t, $J = 7.07$ Hz, 6H), 4.08-4.26 (m, 4H), 7.49-7.52 (m, 1H), 7.67-7.73 (m, 1H), 7.85-7.89 (m, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 14.73








Compound	structure	Description
		(d, $J = 7.8$ Hz, 1P), MS (ESI, EI^+) $m/z = 301$ (MH^+).
2t		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.24 (d, $J = 6.9$ Hz, 6H), 1.25 (t, $J = 7.2$ Hz, 6H), 3.07 (heptuplet, $J = 6.9$ Hz, 1H), 4.02-4.11 (m, 4H), 7.84-7.94 (m, 2H), 8.04 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 15.12, MS (ESI, EI^+) $m/z = 282.3$ (MH^+).
2u		Yellow pale oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.35 (t, $J = 7.05$ Hz, 6H), 4.07-4.22 (m, 4H), 7.25-7.29 (m, 1H), 7.37-7.45 (m, 1H), 7.57-7.61 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 14.82 (d, $J = 9.1$ Hz, 1P), MS (ES^+) $m/z = 267.11$ (MH^+).
2v		Colourless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 0.94 (t, $J = 7.5$ Hz, 3H), 1.32 (t, $J = 7.2$ Hz, 6H), 1.66 (sextuplet, $J = 7.5$ Hz, 2H), 2.63 (t, $J = 7.5$ Hz, 2H), 4-4.21 (m, 4H), 7.34-7.41 (m, 2H), 7.58-7.67 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.53, MS (ES^+) $m/z = 257.37$ (MH^+).
2w		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.67-0.73 (m, 2H), 0.97-1.03 (m, 2H), 1.23 (t, $J = 7.05$ Hz, 6H), 1.97-2.06 (m, 1H), 3.93-4.06 (m, 4H), 7.27-7.31 (m, 1H), 7.38-7.51 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.35, MS (ES^+) $m/z = 255.06$ (MH^+).
2x		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.23 (t, $J = 7.05$ Hz, 6H), 2.37 (s, 3H), 3.97-4.09 (m, 4H), 7.49-7.57 (m, 2H), 7.69 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 15.74, MS (ES^+) $m/z = 309.24$ (MH^+).
2y		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.23 (t, $J = 7.05$ Hz, 6H), 2.34 (s, 3H), 2.63 (q, $J = 7.05$ Hz, 2H), 3.94-4.05 (m, 4H), 7.29-7.31 (m, 2H), 7.44-7.45 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.71, MS (ES^+) $m/z = 257.4$ (MH^+).
2z		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.88 (t, $J = 7.2$ Hz, 3H), 1.22 (t, $J = 7.05$ Hz, 6H), 1.54-1.62 (m, 2H), 2.34 (s, 3H), 2.55-2.6 (m, 2H), 3.94-4.05 (m, 4H), 7.27-7.36 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.73, MS (ES^+) $m/z = 271.36$ (MH^+).

Compound	structure	Description
2aa		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.28 (d, $J = 7.0$ Hz, 6H), 1.34 (t, $J = 7.2$ Hz, 6H), 2.97 (m, 1H), 4.13 (m, 4H), 7.36-7.45 (m, 2H), 7.58-7.75 (m, 2H), MS (ESI, EI^+) $m/z = 257$ (MH^+).
2ab		Brown oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.22 (t, $J = 7.2$ Hz, 6H), 2.37 (s, 3H), 3.92-4.05 (m, 4H), 7.33-7.37 (m, 2H), 7.57-7.64 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 18.62.
2ae		Yellow oil, ^1H NMR (d_6 -DMSO, 400 MHz) δ 1.22 (t, $J = 7.3$ Hz, 6H), 2.28 (s, 6H), 3.93-4.03 (m, 4H), 7.29-7.32 (m, 1H), 7.4-7.49 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 19.86, MS (ESI, EI^+) $m/z = 243$ (MH^+).
2af		Yellow oil, MS (ESI, EI^+) $m/z = 243$ (MH^+).
2ag		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.24 (t, $J = 7.05$ Hz, 6H), 2.28 (s, 3H), 2.42 (s, 3H), 3.96-4.06 (m, 4H), 7.22 (td, $J = 4.2$ and 7.5 Hz, 1H), 7.41 (d, $J = 7.5$ Hz, 1H), 7.62 (dd, $J = 7.5$ and 13.8 Hz, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 20.29.
2ah		Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 7.05$ Hz, 6H), 4.04-4.19 (m, 4H), 7.12-7.19 (m, 2H), 7.78-7.87 (m, 2H), ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -109.18, ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 18.33.
2ai		Pale yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, $J = 6.9$ Hz, 6H), 4.08-4.23 (m, 4H), 6.96-7.04 (m, 1H), 7.30-7.38 (m, 2H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 14.77-15.3 (m, 1P), ^{19}F NMR (CDCl_3 , 235.36 MHz) δ -107.56 (m, 2F), MS (ESI, EI^+) $m/z = 251$ (MH^+).
2ao		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 7.5$ Hz, 3H), 1.24 (t, $J = 7.05$ Hz, 6H), 2.89 (q, $J = 7.5$ Hz, 2H), 3.97-4.07 (m, 4H), 7.29-7.41 (m, 2H), 7.53-7.58 (m, 1H), 7.74 (ddd, $J = 1.2$ and 7.5 and 13.8 Hz, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.71, MS (ESI, EI^+) $m/z = 243$ (MH^+).

Compound	structure	Description
2ap		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.23 (t, $J = 7.05$ Hz, 6H), 3.81 (s, 3H), 3.96-4.06 (m, 4H), 7.15-7.32 (m, 3H), 7.44-7.51 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 17.71; MS (ESI, EI^+) $m/z = 245$ (MH^+)
2aq		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.22 (t, $J = 7.05$ Hz, 6H), 3.94-4.07 (m, 4H), 4.56 (s, 2H), 4.61 (s, 2H), 7.28-7.37 (m, 5H), 7.5-7.72 (m, 4H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 17.97; MS (ES^+) $m/z = 335$ (MH^+)
2ar		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.22 (t, $J = 7.2$ Hz, 6H), 3.97-4.03 (m, 4H), 7.43-7.56 (m, 2H), 7.65-7.7 (m, 2H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.6; MS (ES^+) $m/z = 271.2$ (MH^+)
2as		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.24 (t, $J = 7.05$ Hz, 6H), 2.42 (s, 3H), 3.99-4.1 (m, 4H), 7.82-7.94 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 15.09, MS (ES^+) $m/z = 254$ (MH^+).
2at		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.25 (t, $J = 7.2$ Hz, 6H), 4.01-4.13 (m, 4H), 7.49-7.57 (m, 1H), 7.65-7.7 (m, 1H), 7.88-7.92 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 13.5 (d, $J = 8.7$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 282.4 MHz) δ -108.43 (m, 1F).
2au		Orange oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.25 (t, $J = 7.2$ Hz, 6H), 4.04-4.13 (m, 4H), 7.82-7.90 (m, 1H), 7.96-8.01 (m, 1H), 8.15-8.18 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 12.83 (d, $J = 8.02$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 282.4 MHz) -108.7 (q, $J = 8.2$ Hz, 1F), MS (ES^+) $m/z = 258.2$ (MH^+).
2aw		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.25 (t, $J = 7.05$ Hz, 6H), 3.97-4.13 (m, 4H), 7.46-7.53 (m, 1H), 7.57-7.66 (m, 2H), 7.84-7.96 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 13.64, MS (ESI, EI^+) $m/z = 249.05$ (MH^+).
3a		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 7.3$ Hz, 3H), 4.08 (q, $J = 7.3$ Hz, 2H), 7.42-7.56 (m, 3H), 7.79-7.86 (m, 2H), 10.67 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 21.3.

Compound	structure	Description
3b		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.31 (t, $J = 7.05$ Hz, 3H), 2.32 (s, 6H), 4.03-4.15 (m, 2H), 7.15 (s, 1H), 7.42 (d, $J = 13.8$ Hz, 2H), 9.64 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 22.36, MS (ESI, EI^+) $m/z = 215$ (MH^+).
3c		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.3 (t, $J = 7.05$ Hz, 3H), 4.02-4.11 (m, 2H), 7.09-7.16 (m, 2H), 7.76-7.85 (m, 2H), 8.37 (brs, 1H), ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -106 (m, 1F), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 20, MS (ESI, EI^+) $m/z = (\text{MH}^+)$.
3d		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 7.05$ Hz, 3H), 4.11 (quintuplet, $J = 7.05$ Hz, 2H), 7.55-7.61 (m, 1H), 7.78-7.8 (m, 1H), 7.95-8.13 (m, 3H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 18.09, ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -66.03, MS (ESI, EI^+) $m/z = 255$ (MH^+).
3e		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 3H), 4.04 (quintuplet, $J = 7.05$ Hz, 2H), 7.27-8.09 (m, 5H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 16.59, MS (ESI, EI^+) $m/z = 212$ (MH^+).
3f		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 3H), 4.1 (quintuplet, $J = 7.05$ Hz, 2H), 7.51 (t, $J = 1.8$ Hz, 1H), 7.62 (d, $J = 1.8$ Hz, 1H), 7.67 (d, $J = 1.8$ Hz, 1H), 11.63 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.74, MS (ESI, EI^+) $m/z = 255$ (MH^+).
3g		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.26 (t, $J = 7.5$ Hz, 3H), 1.33 (t, $J = 7.2$ Hz, 3H), 2.69 (q, $J = 7.5$ Hz, 2H), 4.1 (quintuplet, $J = 7.2$ Hz, 2H), 7.36-7.39 (m, 2H), 7.61-7.69 (m, 2H), 10.83 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21.63, MS (ESI, EI^+) $m/z = 215$ (MH^+).
3h		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.25 (d, $J = 6.9$ Hz, 6H), 1.31 (t, $J = 7.05$ Hz, 3H), 2.88-2.98 (m, 1H), 4.03-4.13 (m, 2H), 7.35-7.41 (m, 2H), 7.60-7.69 (m, 2H), 11.26 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21.85, MS (ESI, EI^+) $m/z = 229$ (MH^+).

Compound	structure	Description
3i		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (d, $J = 7.0$ Hz, 6H), 2.35 (s, 3H), 4.04-4.14 (m, 2H), 7.32 (s, 1H), 7.45-7.59 (m, 2H), 8.28 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.16.
3j		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.31 (d, $J = 7.0$ Hz, 6H), 2.35 (s, 3H), 4.05-4.17 (m, 2H), 7.32 (s, 1H), 7.36-7.83 (m, 8H), 8.03 (d, $J = 14.3$ Hz, 1H), 10.16 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21.13, MS (ESI, EI^+) $m/z = 263$ (MH^+).
3k		Colourless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 7.0$ Hz, 3H), 3.83 (s, 3H), 4.08 (quintuplet, $J = 7.0$ Hz, 2H), 7.05-7.09 (m, 1H), 7.3-7.44 (m, 3H), 10.25 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 20.36.
3n		Light orange oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, $J = 7.2$ Hz, 3H), 4.16 (quintuplet, $J = 7.2$ Hz, 2H), 8.06 (brs, 1H), 8.22 (brs, 1H), 8.27 (brs, 1H), 10.95 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.19, MS (ESI, EI^+) $m/z = 322.99$ (MH^+).
3q		Light orange oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.34 (t, $J = 7.05$ Hz, 3H), 2.4 (s, 3H), 4.11 (quintuplet, $J = 7.05$ Hz, 2H), 7.04-7.08 (m, 1H), 7.27-7.35 (m, 1H), 7.4-7.43 (m, 1H), 11.37 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.37 (d, $J = 9.6$ Hz, 1P), MS (ESI, EI^+) $m/z = 219$ (MH^+).
3r		Yellowish oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.18 (t, $J = 7.1$ Hz, 3H), 3.87-3.97 (m, 2H), 7.55-7.67 (m, 3H), 7.98-8.18 (m, 3H), 8.53 (d, $J = 8.4$ Hz, 1H), ^{31}P NMR (d_6 -DMSO, 300 MHz) δ 14.76, MS (ESI, EI^+) $m/z = 237$ (MH^+).
3s		Yellowish oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.25 (d, $J = 6.9$ Hz, 6H), 1.31 (t, $J = 6.9$ Hz, 3H), 2.70-3.00 (m, 1H), 4.00-4.20 (m, 2H), 7.89 (d, $J = 5.4$ Hz, 1H), 7.25-7.36 (m, 1H), 7.47 (d, $J = 13.5$ Hz, 1H), 12.22 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 17.93, ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -112.35 (m, 1F).

Compound	structure	Description
3t		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.27 (d, $J = 6.9$ Hz, 6H), 1.33 (t, $J = 6.9$ Hz, 3H), 2.94-3.03 (m, 1H), 4.07-4.16 (m, 2H), 5.15 (brs, 1H), 7.66 (brs, 1H), 7.83-7.95 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 17.61, MS (ESI, EI^+) $m/z = 254.3$ (MH^+).
3u		Colourless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 3H), 4.1 (quintuplet, $J = 7.05$ Hz, 2H), 7.24-7.28 (m, 1H), 7.35-7.44 (m, 1H), 7.55-7.60 (m, 1H), 12.16 (s, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.63 (d, $J = 9.20$ Hz, 1P).
3v		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.17 (t, $J = 7.2$ Hz, 3H), 1.17 (t, $J = 7.5$ Hz, 3H), 2.32 (s, 3H), 2.61 (q, $J = 7.5$ Hz, 2H), 3.81-3.91 (m, 2H), 7.21 (brs, 1H), 7.29 (brs, 1H), 7.34 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 15.62, MS (ES^+) $m/z = 229.3$ (MH^+).
3x		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.18 (t, $J = 6.9$ Hz, 3H), 2.36 (s, 3H), 3.87 (quintuplet, $J = 7.2$ Hz, 2H), 7.33-7.42 (m, 2H), 7.44-7.55 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 16.41.
3z		Yellow pale oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 6.9$ Hz, 3H), 2.30 (brs, 3H), 2.45 (brs, 3H), 3.87 (quintuplet, $J = 7.5$ Hz, 2H), 7.18-7.27 (m, 2H), 7.57 (d, $J = 14.4$ Hz, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 15.79.
3aa		Yellow pale oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.17 (t, $J = 7.05$ Hz, 3H), 2.27 (brs, 6H), 3.85 (quintuplet, $J = 7.2$ Hz, 2H), 7.23-7.27 (m, 1H), 7.37-7.47 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 16.92, MS (ESI, EI^+) $m/z = 215$ (MH^+).
3ad		Pale yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.17 (t, $J = 7.1$ Hz, 6H), 3.84-3.94 (m, 2H), 7.51-7.69 (m, 3H), ^{19}F NMR (d_6 -DMSO, 300 MHz) δ -141.87 (m, 1F), -137.91 (m, 1F), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 12.80 (d, $J = 6.5$ Hz, 1P), MS (ESI, EI^+) $m/z = 223$ (MH^+).

Compound	structure	Description
3ae		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.18 (t, $J = 7.5$ Hz, 3H), 1.19 (t, $J = 7.05$ Hz, 3H), 2.91 (q, $J = 7.5$ Hz, 2H), 3.89 (quintuplet, $J = 7.05$ Hz, 2H), 7.24-7.36 (m, 2H), 7.46-7.52 (m, 1H), 7.74 (ddd, $J = 1.2$ and 7.5 and 13.8 Hz, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 15.66, MS (ESI, EI^+) $m/z = 215$ (MH^+).
3af		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.18 (t, $J = 6.9$ Hz, 3H), 2.4 (s, 3H), 3.85-3.97 (m, 2H), 7.78-7.81 (m, 1H), 7.82-7.84 (m, 1H), 7.84-7.87 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 11.67, MS (ES^+) $m/z = 226.2$ (MH^+).
3ag		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.17 (t, $J = 7.1$ Hz, 3H), 1.3 (s, 9H), 3.82-3.91 (m, 2H), 7.41-7.53 (m, 2H), 7.59-7.61 (m, 1H), 7.66-7.71 (m, 1H), MS (ES^+) $m/z = 243$ (MH^+).
3ai		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 6.9$ Hz, 3H), 3.87-3.97 (m, 2H), 7.4-7.49 (m, 1H), 7.6-7.64 (m, 1H), 7.78-7.82 (m, 1H), ^{19}F NMR (d_6 -DMSO, 282.4 MHz) δ -109.3 (q, $J = 8.2$ Hz, 1F), MS (ES^+) $m/z = 283/285$ (MH^+).
3aj		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 7.05$ Hz, 3H), 3.87-3.97 (m, 2H), 7.55-7.65 (m, 1H), 7.85-7.91 (m, 1H), 8.03-8.09 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 11.94 (d, $J = 8.02$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 282.4 MHz) -112.04 (m, 1F), MS (ES^+) $m/z = 228.3$ (MH^+).
3ak		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.20 (t, $J = 7.05$ Hz, 3H), 3.87-3.97 (m, 2H), 7.4-7.47 (m, 1H), 7.51-7.59 (m, 2H), 7.83-7.91 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 11.26, MS (ESI, EI^+) $m/z = 221.03$ (MH^+).
3al		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 7.2$ Hz, 3H), 3.9-4 (m, 2H), 7.31-7.53 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 8.05 (d, $J = 5.5$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 282.4 MHz) -110.71 (d, $J = 19.4$ Hz, 1F), -118.33 (dd, $J = 5.9$ and 19.2 Hz, 1F), MS (ES^+) $m/z = 223.03$ (MH^+).

Table 2


Compound	structure	Description
5a		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1,27 (t, $J = 7.1$ Hz, 3H), 1,36 (t, $J = 7.1$ Hz, 3H), 4.03 (m, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 7.51-7.83 (m, 11H), 8.05-8.11 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 23.3, MS (ESI, EI^+) $m/z = 532$ (MH^+).
5b		Thick yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.30 (t, $J = 7.2$ Hz, 3H), 4.22-4.34 (m, 2H), 5-5.12 (m, 2H), 7.26-7.33 (m, 6H), 7.42-7.54 (m, 5H), 7.60-7.66 (m, 1H), 7.77-7.78 (m, 1H), 7.85-7.92 (m, 3H), 8.07-8.10 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 25.24, MS (ESI, EI^+) $m/z = 594$ (MH^+).
5c		Slight yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1,27 (t, $J = 7.1$ Hz, 3H), 1,36 (t, $J = 7.1$ Hz, 3H), 2.30 (s, 6H), 3.94-4.06 (m, 2H), 4.44 (q, $J = 7.1$ Hz, 2H), 7.25 (s, 1H), 7.39 (s, 1H), 7.42 (s, 1H), 7.53 (dd, $J = 2.1$ and 9.0 Hz, 1H), 7.65-7.71 (m, 2H), 7.77-7.82 (m, 2H), 8.05-8.11 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 23.6, MS (ESI, EI^+) $m/z = 560$ (MH^+).

Compound

structure


Description

5d




Thick yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1,37 (t, $J = 7.2$ Hz, 3H), 1,44 (t, $J = 7.2$ Hz, 3H), 4.05-4.19 (m, 2H), 4.52 (q, $J = 7.2$ Hz, 2H), 7.11-7.17 (m, 2H), 7.32-7.36 (m, 1H), 7.48-7.54 (m, 2H), 7.60-7.63 (m, 1H), 7.84-7.93 (m, 4H), 8.07-8.09 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 23.6, ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -105.36 (m, 1F), MS (ESI, EI^+) $m/z = 550$ (MH^+).

5e




Thick yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1,38 (t, $J = 7.05$ Hz, 3H), 1,46 (t, $J = 7.2$ Hz, 3H), 4.09-4.2 (m, 2H), 4.53 (q, $J = 7.2$ Hz, 2H), 7.35 (dd, $J = 2.1$ and 9 Hz, 1H), 7.37-7.43 (m, 1H), 7.48-7.54 (m, 3H), 7.6-7.95 (m, 5H), 8.07-8.1 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 22.76, MS (ESI, EI^+) $m/z = 566$ (MH^+).

5f



Thick yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.23 (t, $J = 7.5$ Hz, 3H), 1,37 (t, $J = 7.05$ Hz, 3H), 1,44 (t, $J = 7.2$ Hz, 3H), 2.68 (q, $J = 7.5$ Hz, 2H), 4.06-4.18 (m, 2H), 4.5 (q, $J = 7.2$ Hz, 2H), 7.3-7.39 (m, 3H), 7.47-7.52 (m, 2H), 7.59-7.76 (m, 3H), 7.87 (d, $J = 1.8$ Hz, 1H), 7.91 (dd, $J = 1.5$ and 9 Hz, 1H), 8.07-8.1 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21, MS (ESI, EI^+) $m/z = 560$ (MH^+).

5g




Yellow solid, ^1H NMR (CDCl_3 , 400 MHz) δ 1.39 (t, $J = 7$ Hz, 3H), 1,44 (t, $J = 7.2$ Hz, 3H), 4.09-4.23 (m, 2H), 4.53 (qd, $J = 2$ and 7.2 Hz, 2H), 7.35 (dd, $J = 2$ and 9.2 Hz, 1H), 7.48-7.53 (m, 2H), 7.57-7.64 (m, 2H), 7.68-7.79 (m, 1H), 7.88 (d, $J = 2$ Hz, 1H), 7.94 (dd, $J = 1.6$ and 8.8 Hz, 1H), 8.04-8.1 (m, 3H), 8.16-8.2 (m, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 22.62, ^{19}F NMR (CDCl_3 , 235.36 MHz) δ -63.2, MS (ESI, EI^+) m/z

Compound

structure


Description

5h

= 600 (MH⁺).


Yellow solid, ¹H NMR (CDCl₃, 300 MHz) δ 1.4 (t, *J* = 7.05 Hz, 3H), 1.46 (t, *J* = 7.2 Hz, 3H), 4.04-4.24 (m, 2H), 4.55 (q, *J* = 7.2 Hz, 2H), 7.37 (dd, *J* = 2.1 and 9 Hz, 1H), 7.51-7.67 (m, 4H), 7.80 (dd, *J* = 1.5 and *J* = 7.8 Hz, 1H), 7.84 (d, *J* = 2.1 Hz, 1H), 7.95 (dd, *J* = 1.5 and 8.7 Hz, 1H), 8.07-8.17 (m, 4H), ³¹P NMR (CDCl₃, 101.256 MHz) δ 21.64, MS (ESI, EI⁺) *m/z* = 557 (MH⁺).

5i




White solid, ¹H NMR (CDCl₃, 300 MHz) δ 1.38 (t, *J* = 7.05 Hz, 3H), 1.46 (t, *J* = 7.35 Hz, 3H), 4.10-4.19 (m, 2H), 4.53 (q, *J* = 7.2 Hz, 2H), 7.3-7.37 (m, 2H), 7.49-7.55 (m, 2H), 7.6-7.67 (m, 2H), 7.77-7.84 (m, 1H), 7.87 (d, *J* = 2.1 Hz, 1H), 7.93 (dd, *J* = 1.8 and 9 Hz, 1H), 8-8.06 (m, 1H), 8.07-8.11 (m, 2H), ³¹P NMR (CDCl₃, 121.49 MHz) δ 22.11, MS (ES⁺) *m/z* = 609.73/611.77 (MH⁺).

5j



Thick yellow oil, ¹H NMR (d₆-DMSO, 300 MHz) δ 1.16 (d, *J* = 6.9 Hz, 3H), 1.17 (d, *J* = 6.9 Hz, 3H), 1.27 (t, *J* = 7.05 Hz, 3H), 1.38 (t, *J* = 7.2 Hz, 3H), 2.88-2.97 (m, 1H), 4.03 (q, *J* = 7.2 Hz, 2H), 4.45 (q, *J* = 7.05 Hz, 2H), 7.45-7.69 (m, 7H), 7.75-7.79 (m, 2H), 8.05-8.1 (m, 3H), MS (ES⁺) *m/z* = 573.8 (MH⁺).

5k




Yellow solid; ¹H NMR (d₆-DMSO, 300 MHz) δ 1.30 (t, *J* = 7.0 Hz, 3H), 1.35 (t, *J* = 7.0 Hz, 3H), 4.05-4.12 (m, 2H), 4.46 (q, *J* = 7.0 Hz, 2H), 5.56 (d, *J* = 12 Hz, 1H), 7.44-8.10 (m, 18H), ³¹P NMR (d₆-DMSO, 121.49 MHz) δ 22.07, MS (ES⁺) *m/z* = 608 (MH⁺).

Compound

structure


Description

5l




Thick yellow oil, ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 22.73, MS (ESI $^+$) m/z = 580 (MH $^+$).

5n




White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.37 (t, J = 7.2 Hz, 3H), 1.45 (t, J = 7.2 Hz, 3H), 3.81 (s, 3H), 4.05-4.18 (m, 2H), 4.53 (q, J = 7.2 Hz, 2H), 7.07 (dt, J = 1.2 Hz and 8.1 Hz, 1H), 7.28-7.63 (m, 7H), 7.87 (d, J = 2.1, 1H), 7.90 (dd, J = 1.5 Hz and 9 Hz, 1H), 8.08 (dd, J = 1.5 Hz and 7.5 Hz, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 24.00

5p



Yellowish powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.38 (t, J = 7.2 Hz, 3H), 1.47 (t, J = 7.2 Hz, 3H), 3.85 (s, 3H), 4.13 (q, J = 7.2 Hz, 2H), 4.53 (q, J = 7.2 Hz, 2H), 6.97 (dd, J = 2.7 Hz and 8.7 Hz, 2H), 7.34 (dd, J = 2.1 Hz and 9.3 Hz, 1H), 7.49-7.55 (m, 2H), 7.6-7.66 (m, 1H), 7.8-7.94 (m, 4H), 8.08-8.11 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 24.36, MS (ESI, EI $^+$) m/z = 562 (MH $^+$).

5q



Yellowish powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, J = 7.2 Hz, H), 1.38 (t, J = 7.05 Hz, 3H), 3.55 (s, 3H), 4.05-4.23 (m, 2H), 4.42 (q, J = 7.2 Hz, 2H), 6.82-6.87 (m, 1H), 7.05-7.11 (m, 1H), 7.35 (dd, J = 2.1 Hz and 9 Hz, 1H), 7.89-7.53 (m, 3H), 7.6-7.66 (m, 1H), 7.9-7.98 (m, 2H), 8.03 (d, J = 2.1 Hz, 1H), 8.09-8.12 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21.9, MS (ESI, EI $^+$) m/z = 562 (MH $^+$).


Compound	structure	Description
5r		Yellowish powder, ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 20.6, ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -63.3, MS (ESI, EI^+) m/z = 668 (MH^+).
5s		Yellowish powder, ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 20.86 (d, J = 7.8 Hz, 1P), ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -63.2 (s, 3F), -108.7 (q, J = 7.3 Hz, 1F), MS (ESI, EI^+) m/z = 618 (MH^+).
5u		Yellowish powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (quintuplet, J = 7.2 Hz, 6H), 4.09-4.3 (m, 2H), 4.42 (q, J = 6.9 Hz, 2H), 7.31 (dd, J = 2.1 Hz and 9 Hz, 1H), 7.47-7.64 (m, 6H), 7.71 (d, J = 1.8 Hz, 1H), 7.87-7.95 (m, 2H), 8.04-8.09 (m, 3H), 8.16 (qd, J = 1.2 Hz and 7.2 Hz, 1H), 8.69-8.72 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 24.31, MS (ESI, EI^+) m/z = 582 (MH^+).
5v		Yellowish powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.23 (d, J = 7.2 Hz, 6H), 1.38 (t, J = 7.2 Hz, 3H), 1.46 (t, J = 7.2 Hz, 3H), 2.83-3.06 (m, 1H), 4.05-4.25 (m, 2H), 4.55 (q, J = 7.0 Hz, 2H), 7.05-7.10 (m, 1H), 7.33 (dd, J = 2.1 Hz and 9.0 Hz, 1H), 7.31-7.41 (m, 1H), 7.45-7.64 (m, 4H), 7.86 (d, J = 1.8 Hz, 1H), 7.94 (dd, J = 8.7 Hz and 1.8 Hz, 1H), 8.06-8.11 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 22.83 (d, J = 7.65 Hz), MS (ES^+) m/z = 592 (MH^+).

Compound

structure


Description

5w




Yellow resin, ^1H NMR (CDCl_3 , 300 MHz) δ 1.42 (t, $J = 7.05$ Hz, 3H), 1.49 (t, $J = 7.2$ Hz, 3H), 4.11-4.26 (m, 2H), 4.57 (q, $J = 7.2$ Hz, 2H), 7.23-7.27 (m, 1H), 7.38 (dd, $J = 1.8$ and 9 Hz, 1H), 7.5-7.57 (m, 3H), 7.62-7.72 (m, 2H), 7.87-7.88 (m, 1H), 7.94-7.98 (m, 1H), 8.09-8.12 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 6.78, MS (ES^+) $m/z = 584.26$ (MH^+).

5x




Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.27 (t, $J = 6.9$ Hz, 3H), 1.37 (t, $J = 7.2$ Hz, 3H), 2.34 (s, 3H), 3.97-4.02 (m, 2H), 4.44 (q, $J = 7.2$ Hz, 2H), 7.34-7.38 (m, 2H), 7.51-7.55 (dd, $J = 2.1$ and 9 Hz, 1H), 7.65-7.72 (m, 4H), 7.75-7.83 (m, 2H) ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 23.57, MS (ESI, EI^+) $m/z = 546$ (MH^+).

5aa



Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.27 (t, $J = 7.05$ Hz, 3H), 1.39 (t, $J = 7.05$ Hz, 3H), 2.25 (2s, 6H), 3.96-4.04 (m, 2H), 4.44 (q, $J = 7.05$ Hz, 2H), 7.3-7.34 (m, 1H), 7.48-7.58 (m, 3H), 7.66-7.72 (m, 2H), 7.77-7.83 (m, 2H), 8.09-8.11 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 23.64, MS (ESI, EI^+) $m/z = 560$ (MH^+).

5ad




Dark yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.29 (t, $J = 6.9$ Hz, 3H), 1.37 (t, $J = 7.2$ Hz, 3H), 4.03-4.13 (m, 2H), 4.46 (q, $J = 7.1$ Hz, 2H), 7.44-7.63 (m, 4H), 7.65-7.71 (m, 2H), 7.78-7.83 (m, 1H), 7.88 (d, $J = 2.4$ Hz, 1H), 8.06-8.11 (m, 3H), ^{19}F NMR (d_6 -DMSO) δ -109.98 (m, 1F) + impureté, ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 20.41 (t, $J = 8.4$ Hz, 1P), MS (ESI, EI^+) $m/z = 569$ ($\text{M}+2\text{H}^+$).

Compound

structure


Description

5ae




Yellow oil; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.22-1.30 (m, 6H), 4.95-4.13 (m, 2H), 4.31 (d, J = 7.2 Hz, 2H), 7.25-7.45 (m, 3H), 7.55 (dd, J = 9.3 and 2.0 Hz, 1H), 7.66-7.85 (m, 6H), 8.06-8.10 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 13.91, ^{19}F NMR (d_6 -DMSO) δ -107.59 (m, 1F), MS (ESI, EI^+) m/z = 550 (MH^+).

5ah




Yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.77 (t, J = 7.5 Hz, 3H), 1.23 (t, J = 6.9 Hz, 3H), 1.26 (t, J = 7.2 Hz, 3H), 2.76 (q, J = 7.5 Hz, 2H), 3.83-4.13 (m, 2H), 4.24-4.35 (m, 2H), 7.3-7.42 (m, 2H), 7.48-7.6 (m, 3H), 7.65-7.8 (m, 4H), 8.06-8.1 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 22.65, MS (ESI, EI^+) m/z = 560 ($\text{M}+\text{H}^+$).

5ai



Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.25 (s, 9H), 1.25 (t, J = 7.2 Hz, 3H), 1.39 (t, J = 7.2 Hz, 3H), 3.99 (m, 2H), 4.46 (q, J = 7.2 Hz, 2H), 7.47-7.55 (m, 2H), 7.6-7.7 (m, 4H), 7.75-7.83 (m, 3H), 8.05-8.1 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 22.6, MS (ES^+) m/z = 588.17 (MH^+).

5aj



Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.4 (t, J = 7.05 Hz, 3H), 1.47 (t, J = 7.2 Hz, 3H), 2.44 (s, 3H), 4-4.25 (m, 2H), 4.55 (q, J = 7.2 Hz, 2H), 7.37 (dd, J = 2.1 and 9 Hz, 1H), 7.52-7.65 (m, 4H), 7.85-7.97 (m, 4H), 8.08-8.11 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 21.58, MS (ES^+) m/z = 571.4 (MH^+).


Compound	structure	Description
5ak		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.21 (t, J = 7.2 Hz, 3H), 1.26 (t, J = 6.9 Hz, 3H), 4.03 (q, J = 7.2 Hz, 2H), 4.18-4.26 (m, 2H), 7.52-7.63 (m, 3H), 7.66-7.71 (m, 4H), 7.78-7.83 (m, 1H), 7.92-8 (m, 1H), 8.05-8.09 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 18.06, MS (ES^+) m/z = 565.93 (MH^+).
5al		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.3 (t, J = 7.05 Hz, 3H), 1.37 (t, J = 7.05 Hz, 3H), 4-4.16 (m, 2H), 4.46 (q, J = 7.05 Hz, 2H), 7.56 (dd, J = 2.1 and 9 Hz, 1H), 7.66-7.71 (m, 2H), 7.79-7.84 (m, 1H), 7.91-7.99 (m, 2H), 8.08-8.18 (m, 5H), ^{19}F NMR (d_6 -DMSO, 282.40 MHz) -112.68 (q, J = 8.2 Hz, 1F), MS (ES^+) m/z = 575 (MH^+).
6a		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.34 (t, J = 7.1 Hz, 3H), 4.05 (m, 1H), 4.20 (m, 1H), 7.32 (dd, J = 2.1 and 8.7 Hz, 1H), 7.49-7.61 (m, 5H), 7.68-7.75 (m, 2H), 8.02 (brs, 1H), 10.27 (brs, 1H), 12.77 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 31.1, MS (ESI, EI^+) m/z = 363 (MH^+).
6b		White solid; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.32 (t, J = 7.0 Hz, 3H), 2.26 (s, 6H), 3.90-4.03 (m, 1H), 4.09-4.22 (m, 1H), 7.21 (s, 1H), 7.29-7.33 (m, 3H), 7.57 (dd, J = 1.8 and 9.0 Hz, 1H), 7.60 (d, J = 1.8 Hz, 1H), 7.99 (brs, 1H), 10.3 (brs, 1H), 12.7 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 31.3, MS (ESI, EI^+) m/z = 391 (MH^+).
6c		White Solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.34 (t, J = 6.9 Hz, 3H), 4-4.04 (m, 1H), 4.18-4.21 (m, 1H), 7.31-7.4 (m, 3H), 7.57-7.62 (m, 2H), 7.74-7.83 (m, 2H), 8.02 (brs, 1H), 10.18 (brs, 1H), 12.79 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 29.29, ^{19}F NMR (d_6 -DMSO, 282.4 MHz)

Compound

structure

Description


6d



δ -106.3, MS (ESI, EI^+) m/z = 381 (MH^+).


White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.35 (t, J = 7.05 Hz, 3H), 4-4.09 (m, 1H), 4.17-4.25 (m, 1H), 7.3-7.35 (m, 1H), 7.55-7.74 (m, 6H), 8.05 (brs, 1H), 10.07 (brs, 1H), 12.84 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 28.24, MS (ESI, EI^+) m/z = 397 (MH^+).

6e




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.13 (t, J = 7.5 Hz, 3H), 1.33 (t, J = 6.9 Hz, 3H), 2.62 (q, J = 7.5 Hz, 2H), 3.95-4.06 (m, 1H), 4.14-4.22 (m, 1H), 7.3 (dd, J = 1.65 and 8.85 Hz, 1H), 7.43-7.5 (m, 3H), 7.56-7.61 (m, 3H), 8 (brs, 1H), 10.31 (brs, 1H), 12.74 (brs, 1H), MS (ESI, EI^+) m/z = 391 (MH^+).

6f




White solid, ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 29.02, ^{19}F NMR (CDCl_3 , 235.36 MHz) δ -61.07, MS (ESI, EI^+) m/z = 431 (MH^+).

6g



White solid, ^1H NMR (d_6 -DMSO, 400 MHz) δ 1.36 (t, J = 7.05 Hz, 3H), 4-4.12 (m, 1H), 4.17-4.27 (m, 1H), 7.33 (dd, J = 2 and 8.76 Hz, 1H), 7.59 (dd, J = 1.56 and 8.78 Hz, 1H), 7.66 (d, J = 1.99 Hz, 1H), 7.73 (td, J = 3.3 and 7.78 Hz, 1H), 7.97-8.08 (m, 3H), 8.17-8.21 (m, 1H), 9.98 (brs, 1H), 12.83 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 28.62, MS (ESI, EI^+) m/z = 388 (MH^+).

6i




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.16 (d, J = 6.9 Hz, 3H), 1.17 (d, J = 6.9 Hz, 3H), 1.34 (t, J = 7.05 Hz, 3H), 2.88-2.97 (m, 1H), 3.97-4.06 (m, 1H), 4.15-4.23 (m, 1H), 7.31 (dd, J = 1.8 and 8.7 Hz, 1H), 7.42-7.66 (m, 6H), 8.01 (brs, 1H), 10.31 (brs, 1H), 12.74 (brs, 1H), ^{31}P NMR

Compound

structure

Description


6j



(*d*₆-DMSO, 121.49 MHz) δ 30.22, MS (ES⁺) *m/z* = 404.8 (MH⁺).


White solid; ¹H NMR (*d*₆-DMSO, 300 MHz) δ 1.37 (t, *J* = 7.0 Hz, 3H), 3.99-4.09 (m, 2H), 4.25 (q, *J* = 7.0 Hz, 2H), 5.56 (d, *J* = 12 Hz, 1H), 7.35-8.04 (m, 14H), 10.20 (brs, 1H), 12.78 (brs, 1H), ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 22.07, MS (ES⁺) *m/z* = 439 (MH⁺).

6k




Yellow solid; ¹H NMR (*d*₆-DMSO, 300 MHz) δ 1.35 (t, *J* = 7.0 Hz, 3H), 4.01-4.12 (m, 2H), 4.17-4.25 (m, 2H), 7.35 (dd, *J* = 8.8 and 1.8 Hz), 7.50-7.71 (m, 4H), 7.83-7.87 (m, 1H), 8.04 (brs, 1H), 9.87 (brs, 1H), 12.86 (brs, 1H), ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 26.47 (d, *J* = 7.4 Hz), ¹⁹F NMR (*d*₆-DMSO, 282.4 MHz) δ -111.96 (q, *J* = 7.9 Hz, MS (ES⁺) *m/z* = 459/460 (MH⁺).

6l



White solid; ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 28.35, MS (ES⁺) *m/z* = 411 (MH⁺).

6m




White powder, ¹H NMR (*d*₆-DMSO, 300 MHz) δ 1.39 (t, *J* = 7.04 Hz, 3H), 3.77 (s, 3H), 4.25-3.95 (two m, 2H), 7.29-7.14 (m, 4H), 7.33 (dd, *J* = 8.8 and 1.98 Hz, 1H), 7.49-7.42 (m, 1H), 7.59 (dd, *J* = 8.8 and *J* < 1.5 Hz, 1H), 7.66 (d, *J* = 1.93 Hz, 1H), 8.00 (brs, 1H), 10.20 (brs, 1H), 12.75 (brs, 1H), ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 30.85, MS (ESI, EI⁺) *m/z* = 393 (MH⁺), 785 (2M+H⁺).

Compound

structure


Description

6q




White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.35 (t, $J = 6.9$ Hz, 3H), 4.05-4.14 (m, 1H), 4.18-4.26 (m, 1H), 7.33 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.59 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.74 (d, $J = 1.8$ Hz, 1H), 7.83-7.9 (m, 2H), 7.97-8.03 (m, 2H), 9.74 (brs, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 300 MHz) δ 27.25 (d, $J = 6.57$ Hz, 1P), MS (ESI, EI^+) $m/z = 449$ (MH^+)

6r




White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.35 (t, $J = 7.2$ Hz, 3H), 4.08-4.29 (m, 2H), 7.35 (dd, $J = 2.1$ Hz and 8.7 Hz, 1H), 7.6 (dd, $J = 1.5$ Hz and 8.7 Hz, 1H), 7.8 (d, $J = 1.8$ Hz, 1H), 8.08 (brs, 1H), 8.31 (brs, 1H), 8.36 (brs, 1H), 8.41 (brs, 1H), 9.6 (brs, 1H), 12.88 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.48 MHz) δ 26.63, MS (ESI, EI^+) $m/z = 499$ (MH^+).

6t











White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.35 (t, $J = 6.9$ Hz, 3H), 3.95-4.08 (m, 1H), 4.26-4.37 (m, 1H), 7.27 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.38 (d, $J = 1.8$ Hz, 1H), 7.56-7.65 (m, 4H), 7.86-7.94 (m, 1H), 8.02-8.06 (m, 1H), 8.12 (brs, 1H), 8.19 (d, $J = 8.1$ Hz, 1H), 8.54-8.57 (m, 1H), 10.45 (brs, 1H), 12.83 (brs, 1H), MS (ESI, EI^+) $m/z = 413$ (MH^+).

6u



Yellowish powder, ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 29.27 (d, $J = 8.0$ Hz, 1P), MS (ES^+) $m/z = 423$ (MH^+).

Compound	structure	Description
6v		Colourless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.48 (t, $J = 7.05$ Hz, 3H), 4.05-4.18 (m, 1H), 4.28-4.41 (m, 1H), 6.18 (brs, 1H), 7.22-7.26 (m, 1H), 7.31-7.4 (m, 2H), 7.53-7.55 (m, 1H), 7.57-7.58 (m, 1H), 7.67-7.68 (m, 1H), 10.90 (brs, 1H), 11.13 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 27.7 (d, $J = 7.8$ Hz, 1P), MS (ES^+) $m/z = 415.28$ (MH^+).
6w		Off white powder, ^1H NMR (d_6 -DMSO, 400 MHz) δ 1.31 (t, $J = 7$ Hz, 3H), 2.33 (s, 3H), 3.94-4.04 (m, 1H), 4.11-4.22 (m, 1H), 7.30-7.34 (m, 3H), 7.57-7.62 (m, 4H), 7.98 (brs, 1H), 10.3 (brs, 1H), 12.72 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 31.46, MS (ESI , EI^+) $m/z = 377$ (MH^+).
6z		Yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.29 (t, $J = 7.05$ Hz, 3H), 2.22 (s, 3H), 2.37 (s, 3H), 3.81-3.94 (m, 1H), 4.11-4.24 (m, 1H), 7.21-7.26 (m, 1H), 7.27 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.32 (d, $J = 1.8$ Hz, 1H), 7.4 (d, $J = 7.5$ Hz, 1H), 7.48-7.6 (m, 2H), 8.01 (brs, 1H), 10.42 (brs, 1H), 12.75 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 31.55, MS (ESI , EI^+) $m/z = 391$ (MH^+).
6ab		Off white powder; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.33 (t, $J = 6.9$ Hz, 3H), 2.32 (s, 3H), 3.95-4.04 (m, 1H), 4.12-4.24 (m, 1H), 7.31 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.41-7.61 (m, 6H), 8.00 (brs, 1H), 10.28 (brs, 1H), 12.74 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 31.28, MS (ESI , EI^+) $m/z = 377$ (MH^+).


Compound	structure	Description
6ac		Pale yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1,35 (t, $J = 6.9$ Hz, 3H), 3.99-4.13 (m, 1H), 4.17-4.28 (m, 1H), 7.32-7.43 (m, 3H), 7.50-7.61 (m, 2H), 7.68 (d, $J = 1.8$ Hz, 1H), 8.04 (brs, 1H), 9.91 (brs, 1H), 12.9 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 28.0 (t, $J = 8.1$ Hz, 1P), ^{19}F NMR (d_6 -DMSO) δ -106.92 (m, 2F), MS (ESI, EI^+) $m/z = 399$ (MH^+).
6ad		Pale yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1,35 (t, $J = 6.9$ Hz, 3H), 3.97-4.11 (m, 1H), 4.16-4.29 (m, 1H), 7.22-7.42 (m, 3H), 7.56-7.59 (m, 2H), 7.61-7.75 (m, 1H), 7.77-7.88 (m, 1H), 8.01 (brs, 1H), 10.08 (brs, 1H), 12.78 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 25.2 (d, $J = 11.3$ Hz, 1P), ^{19}F NMR (d_6 -DMSO) δ -104.85 (m, 1F), MS (ESI, EI^+) $m/z = 381$ (MH^+).
6ah		Yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.94 (t, $J = 7.5$ Hz, 3H), 1.3 (t, $J = 7.05$ Hz, 3H), 2.73-2.98 (m, 2H), 3.83-3.96 (m, 1H), 4.1-4.25 (m, 1H), 7.28 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.32-7.38 (m, 3H), 7.5-7.6 (m, 2H), 7.66-7.73 (m, 1H), 8.02 (brs, 1H), 10.42 (brs, 1H), 12.76 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.69 MHz) δ 29.63, MS (ESI, EI^+) $m/z = 391$ (MH^+).
6ai		Off white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.31 (t, $J = 7.05$ Hz, 3H), 3.92-4.05 (m, 1H), 4.15-4.28 (m, 1H), 7.25 (dd, $J = 1.95$ and 8.85 Hz, 1H), 7.35 (d, $J = 2.1$ Hz, 1H), 7.47-7.61 (m, 4H), 8-8.1 (m, 2H), 10.17 (brs, 1H), 12.71 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 25.31, MS (ESI, EI^+) $m/z = 397$ (MH^+).

Compound

structure


Description

6aj




Orange oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.48 (t, $J = 7.05$ Hz, 3H), 4.05-4.23 (m, 1H), 4.29-4.4 (m, 1H), 6.33 (brs, 1H), 7.33 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.4-7.52 (m, 1H), 7.57-7.62 (m, 2H), 7.64-7.72 (m, 1H), 7.81-7.85 (m, 1H), 10.81 (brs, 1H), 11.42 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 26.56 (d, $J = 3.1$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) -107 (q, $J = 7.34$ Hz, 1F), MS (ES^+) $m/z = 406.3$ (MH^+).

6ak




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.35 (t, $J = 7.0$ Hz, 3H), 4.02-4.25 (m, 2H), 7.34 (d, $J = 8.5$ Hz, 2H), 7.59 (d, $J = 8.5$ Hz, 1H), 7.74 (d, $J = 8.5$ Hz, 1H), 8.04 (m, 2H), 8.16 (m, 1H), 8.29 (brs, 1H), 9.70 (brs, 1H), 14.73 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 26.87, MS (ES^+) $m/z = 575$ (MH^+), MS (ES^+) $m/z = 422$ (MH^+).

7a








Slight yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.27 (s, 6H), 3.73 (d, $J = 11.7$ Hz, 3H), 7.23 (brs, 1H), 7.29-7.33 (m, 3H), 7.56-7.59 (m, 2H), 8 (brs, 1H), 10.27 (brs, 1H), 12.75 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 33.27, MS (ESI, EI^+) $m/z = 377$ (MH^+).

7b



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.79 (d, $J = 11.7$ Hz, 3H), 7.35 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.6 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.67 (d, $J = 1.8$ Hz, 1H), 7.7 (t, $J = 1.8$ Hz, 2H), 7.89 (t, $J = 1.8$ Hz, 1H), 8.06 (brs, 1H), 9.78 (brs, 1H), 12.89 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 29.44, MS (ESI, EI^+) $m/z = 417$ (MH^+).

Compound	structure	Description
7c		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.77 (d, $J = 11.7$ Hz, 3H), 7.33 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.54-7.74 (m, 6H), 8.05 (brs, 1H), 10.01 (brs, 1H), 12.84 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 30.23, MS (ESI, EI^+) $m/z = 383$ (MH^+).
7d		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.14 (t, $J = 7.5$ Hz, 3H), 2.63 (q, $J = 7.5$ Hz, 2H), 3.75 (d, $J = 11.4$ Hz, 3H), 7.32 (dd, $J = 1.95$ and 8.85 Hz, 1H), 7.41-7.52 (m, 3H), 7.57-7.63 (m, 3H), 8.02 (brs, 1H), 10.27 (brs, 1H), 12.78 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 32.2; MS (ESI, EI^+) $m/z = 377$ (MH^+).
7e		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.8 (d, $J = 11.7$ Hz, 3H), 7.34 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.6 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.64 (d, $J = 1.8$ Hz, 1H), 7.76 (td, $J = 3.3$ and 7.8 Hz, 1H), 7.94-8.07 (m, 4H), 9.97 (brs, 1H), 12.7 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 29.98, ^{19}F NMR (CDCl_3 , 282.4 MHz) δ 118.06 (s, 3F), MS (ESI, EI^+) $m/z = 417$ (MH^+).
7f		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.79 (d, $J = 11.7$ Hz, 3H), 7.33 (dd, $J = 1.95$ and 8.7 Hz, 1H), 7.59 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.65 (d, $J = 1.8$ Hz, 1H), 7.69-7.75 (m, 1H), 7.96-8.08 (m, 3H), 8.17-8.21 (m, 1H), 9.9 (brs, 1H), 12.88 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 29.65, MS (ES^+) $m/z = 373.86$ (MH^+).
7g		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.16 (d, $J = 6.9$ Hz, 3H), 1.17 (d, $J = 6.9$ Hz, 3H), 2.88-2.98 (m, 1H), 3.76 (d, $J = 11.4$ Hz, 3H), 7.32 (dd, $J = 1.95$ and 8.85 Hz, 1H), 7.41-7.68 (m, 6H), 8.03 (brs, 1H), 10.28 (brs, 1H), 12.77 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.16, MS


Compound	structure	Description
7h		(ES ⁺) m/z = 390.9 (MH ⁺). White solid, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 3.77 (d, J = 11.7 Hz, 3H), 5.47 (d, J = 47.4 Hz, 2H), 7.33 (dd, J = 2.1 and 8.7 Hz, 1H), 7.55-7.81 (m, 6H), 8.04 (brs, 1H), 10.19 (brs, 1H), 12.78 (brs, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 31.62, ¹⁹ F NMR (<i>d</i> ₆ -DMSO, 282.40 MHz) δ -207.99 (t, J = 47.4 Hz, 1F), MS (ES ⁺) m/z = 380.83 (MH ⁺).
7i		White solid; ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 3.81 (d, J = 11.7 Hz, 3H), 7.33 (dd, J = 8.7 and 2.1 Hz, 1H), 7.41-7.72 (m, 9H), 7.87-8.01 (m, 2H), 8.05 (brs, 1H), 10.21 (brs, 1H), 12.79 (brs, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 31.74, MS (ES ⁺) m/z = 425 (MH ⁺).
7j		White solid; ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 3.79 (d, J = 11.7 Hz, 3H), 7.35 (dd, J = 8.7 and 2.1 Hz, 1H), 7.51-7.72 (m, 9H), 7.83-7.87 (m, 2H), 8.06 (brs, 1H), 9.83 (brs, 1H), 12.88 (brs, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 28.46 (d, J = 7.2 Hz), ¹⁹ F NMR (<i>d</i> ₆ -DMSO, 282.4 MHz) δ -110.60 (m), MS (ES ⁺) m/z = 445/447 (MH ⁺).
7k		White solid; ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 2.32 (s, 3H), 3.77 (d, J = 11.4 Hz, 3H), 7.34 (dd, J = 9.0 and 2.1 Hz, 1H), 7.47-7.59 (m, 5H), 8.04 (brs, 1H), 10.04 (brs, 1H), 12.88 (brs, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 30.31, MS (ES ⁺) m/z = 397 (MH ⁺).
7l		White powder, ¹ H NMR (CDCl ₃ , 300 MHz) δ 3.81 (s, 3H), 3.87 (d, J = 11.4 Hz, 3H), 6.02 (brs, 1H), 7.06-7.08 (m, 1H), 7.28-7.37 (m, 4H), 7.49-7.52 (m, 1H), 7.73 (brs, 1H), 10.67 (brs, 1H), 11.01 (brs, 1H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 32.67, MS (ESI, EI ⁺) m/z = 379 (MH ⁺).

Compound

structure


Description

7q




White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.92 (d, $J = 11.7$ Hz, 3H), 6.12 (brs, 1H), 7.36 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.48-7.51 (m, 1H), 7.55-7.65 (m, 3H), 7.9 (d, $J = 12.9$ Hz, 1H), 10.81 (brs, 1H), 10.97 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 29.27 (d, $J = 6.8$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -62.89 (s, 3F), -107.86 (q, $J = 7.6$ Hz, 1F), MS (ESI, EI^+) $m/z = 435$ (MH^+).

7s




White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.94 (d, $J = 11.7$ Hz, 3H), 6.12 (brs, 1H), 7.37 (dd, $J = 2.1$ Hz and 9 Hz, 1H), 7.57 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.63-7.637 (m, 1H), 8.04 (brs, 1H), 8.21 (brs, 1H), 8.26 (brs, 1H), 10.81 (brs, 1H), 10.94 (brs, 1H), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -62.96, MS (ESI, EI^+) $m/z = 485$ (MH^+).

7t



White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.92 (d, $J = 11.7$ Hz, 3H), 6.04 (brs, 1H), 7.3 (dd, $J = 2.1$ Hz and 9 Hz, 1H), 7.42-7.65 (m, 5H), 7.77 (qd, $J = 1.2$ Hz and 7.2 Hz, 1H), 7.9-7.93 (m, 1H), 8.03-8.06 (m, 1H), 8.68 (d, $J = 8.4$ Hz, 1H), 10.58 (brs, 1H), 11.19 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.98, MS (ESI, EI^+) $m/z = 399$ (MH^+).

7u




Off-white powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.24 (d, $J = 6.9$ Hz, 3H), 1.23 (d, $J = 6.9$ Hz, 3H), 2.96 (hept., $J = 6.9$ Hz, 1H), 3.83 (d, $J = 11.7$ Hz, 3H), 5.99 (brs, 1H), 7.07-7.13 (m, 1H), 7.20-7.29 (m, 1H), 7.34 (dd, $J = 2.1$ Hz and 9 Hz, 1H), 7.49-7.53 (m, 2H), 7.68 (d, $J = 2.1$ Hz, 1H), 7.95 (dd, $J = 9.6$ Hz and 1.8 Hz, 1H), 8.08-8.14 (m, 2H), 10.54 (brs, 1H), 10.95 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.35 (d, $J = 8.4$ Hz, 1P), MS (ES^+) $m/z = 409$ (MH^+).

Compound

structure


Description

7v




Off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 3.9 (d, $J = 11.7$ Hz, 3H), 6.09 (brs, 1H), 7.24-7.28 (m, 1H), 7.32-7.40 (m, 2H), 7.53-7.59 (m, 2H), 7.65-7.66 (m, 1H), 10.81 (brs, 1H), 10.87 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 29.77 (d, $J = 7.78$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -111.22 (m, 1F), MS (ES^+) $m/z = 401.29$ (MH^+).

7w




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.12 (t, $J = 7.5$ Hz, 3H), 2.28 (s, 3H), 2.58 (q, $J = 7.5$ Hz, 2H), 3.74 (d, $J = 11.4$ Hz, 3H), 7.26 (brs, 1H), 7.31 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.34-7.40 (m, 2H), 7.56-7.60 (m, 2H), 8 (brs, 1H), 10.29 (brs, 1H), 12.75 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.21, MS (ES^+) $m/z = 391.3$ (MH^+).

7x



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.83 (t, $J = 7.2$ Hz, 3H), 1.54 (sextuplet, $J = 7.2$ Hz, 2H), 2.58 (t, $J = 7.2$ Hz, 2H), 3.75 (d, $J = 11.4$ Hz, 3H), 7.31 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.4-7.51 (m, 3H), 7.55-7.60 (m, 3H), 8 (brs, 1H), 10.25 (brs, 1H), 12.75 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.17, MS (ES^+) $m/z = 391.26$ (MH^+).

7y




White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 2.40 (s, 3H), 3.88 (d, $J = 11.7$ Hz, 3H), 5.89 (d, $J = 16.5$ Hz, 1H), 5.97 (brs, 1H), 7.33-7.67 (m, 7H), 10.46 (s, 1H), 10.89 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.54, MS (ES^+) $m/z = 414$ (MH^+).

Compound

structure


Description

7z




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.69 (d, $J = 11.4$ Hz, 3H), 7.20 (brs, 1H), 7.25 (brs, 1H), 7.29 (brs, 1H), 7.38 (dd, $J = 6.3$ and 8.4 Hz, 1H), 7.44 (dd, $J = 1.5$ and 8.7 Hz, 1H) 8.02 (brs, 1H), 10.77 (brs, 1H), 12.96 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 31.74, MS (ES^+) $m/z = 394.9$ (MH^+).

7aa




Off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 0.69-0.71 (m, 2H), 0.98-1.02 (m, 2H), 1.85-1.95 (m, 1H), 3.86 (d, $J = 11.7$ Hz, 3H), 5.98 (brs, 1H), 7.2-7.22 (m, 1H), 7.28-7.35 (m, 3H), 7.47-7.58 (m, 3H), 7.71-7.72 (m, 1H), 10.50 (brs, 1H), 11.04 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.04, MS (ES^+) $m/z = 389.08$ (MH^+).

7ab




Yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.33 (s, 3H), 3.74 (d, $J = 11.7$ Hz, 3H), 7.31-7.33 (m, 3H), 7.57-7.60 (m, 4H), 8 (brs, 1H), 10.26 (brs, 1H), 12.76 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.48, MS (ESI, EI^+) $m/z = 363$ (MH^+).

7ae



Yellow pale powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.23 (s, 6H), 3.73 (d, $J = 11.4$ Hz, 3H), 7.25-7.33 (m, 2H), 7.36-7.43 (m, 1H), 7.49 (d, $J = 12.9$ Hz, 1H), 7.56-7.60 (m, 2H), 7.99 (brs, 1H), 10.28 (brs, 1H), 12.74 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 33.45, MS (ESI, EI^+) $m/z = 377$ (MH^+).

7ah




Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.85 (d, $J = 11.4$ Hz, 3H), 6.08 (brs, 1H), 7.30 (dd, $J = 2.0$ and 9.0 Hz, 1H), 7.36-7.56 (m, 4H), 7.68 (d, $J = 1.8$ Hz, 1H), 7.73-7.81 (m, 2H), 10.78 (brs, 1H), 10.03 (brs, 1H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 33.3, MS (ESI, EI^+) $m/z = 349$ (MH^+).

Compound

structure


Description

7ai




Pale orange powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.84 (d, $J = 11.4$ Hz, 3H), 5.97 (brs, 1H), 7.18-7.23 (m, 1H), 7.32-7.56 (m, 5H), 7.69-7.83 (m, 3H), 10.44 (brs, 1H), 11.04 (brs, 1H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 33.8, MS (ESI, EI^+) $m/z = 315$ (MH^+).

7aj




Orange powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.79 (d, $J = 11.4$ Hz, 3H), 7.32-7.44 (m, 3H), 7.51-7.61 (m, 2H), 7.67 (d, $J = 1.8$ Hz, 1H), 8.05 (brs, 1H), 9.86 (brs, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 29.99 (t, $J = 8.3$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 300 MHz) δ -106.93 (m, 2F), MS (ESI, EI^+) $m/z = 385$ (MH^+).

7ak




Pale yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.75 (d, $J = 11.4$ Hz, 3H), 7.31-7.38 (m, 3H), 7.54-7.62 (m, 2H), 7.71-7.83 (m, 2H), 8.01 (brs, 1H), 10.10 (brs, 1H), 12.76 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 32.30, ^{19}F NMR (d_6 -DMSO) δ -106.35 (m, 1F), MS (ESI, EI^+) $m/z = 367$ (MH^+).

7an



Off white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.92 (t, $J = 7.5$ Hz, 3H), 2.78-2.92 (m, 2H), 3.7 (d, $J = 11.7$ Hz, 3H), 7.25-7.38 (m, 4H), 7.52-7.59 (m, 2H), 7.67-7.75 (m, 1H), 8.02 (brs, 1H), 10.36 (brs, 1H), 12.78 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 32.41, MS (ESI, EI^+) $m/z = 377$ (MH^+).

7ao



White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.75 (d, $J = 11.7$ Hz, 3H), 7.27 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.34 (d, $J = 1.5$ Hz, 1H), 7.49-7.65 (m, 4H), 8.04-8.11 (m, 2H), 10.15 (brs, 1H), 12.8 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 27.33, MS (ESI, EI^+) $m/z = 383$ (MH^+).


Compound	structure	Description
7ap		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.75 (d, J = 11.7 Hz, 3H), 4.47 (s, 2H), 4.54 (s, 2H), 7.23-7.37 (m, 6H), 7.5-7.73 (m, 6H), 8.01 (brs, 1H), 10.21 (brs, 1H), 12.8 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.13, MS (ES^+) m/z = 468.86 (MH^+).
7aq		Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.88 (d, J = 11.7 Hz, 3H), 4.73 (s, 2H), 5.98 (brs, 1H), 7.3-7.35 (m, 1H), 7.42-7.49 (m, 2H), 7.56-7.59 (m, 1H), 7.66-7.73 (m, 2H), 7.8-7.85 (m, 1H), 10.34 (brs, 1H), 10.99 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 32.85, MS (ES^+) m/z = 378.86 (MH^+).
7ar		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.24 (s, 9H), 3.76 (d, J = 11.4 Hz, 3H), 7.3-7.34 (m, 1H), 7.4-7.65 (m, 5H), 7.76-7.81 (m, 1H), 8.02 (brs, 1H), 10.27 (brs, 1H), 12.75 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.29, MS (ES^+) m/z = 405.11 (MH^+).
7as		White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.4 (s, 3H), 3.9 (d, J = 11.7 Hz, 3H), 6.1 (brs, 1H), 7.34 (dd, J = 1.8 and 8.7 Hz, 1H), 7.53-7.6 (m, 3H), 7.76-7.85 (m, 2H), 10.84 (brs, 1H), 10.92 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 7.27, MS (ES^+) m/z = 388.3 (MH^+).
7au		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.78 (d, J = 11.4 Hz, 3H), 6.60 (d, J = 8.4 Hz, 1H), 7.34 (d, J = 1.8 Hz, 1H), 7.74 (dd, J = 1.7 and 8.4 Hz, 1H), 8.03-8.18 (m, 3H), 8.29 (brs, 1H), 9.67 (brs, 1H), 12.89 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 27.88, MS (ES^-) m/z = 406 (MH^+).

Compound

structure


Description

7av




White powder, ^1H NMR ($\text{d}_6\text{-DMSO}$, 300 MHz) δ 3.78 (d, $J = 11.4$ Hz, 3H), 7.34 (dd, $J = 1.8$ and 8.4 Hz, 1H), 7.42-7.56 (m, 1H), 7.58-7.73 (m, 3H), 7.80-7.89 (m, 2H), 8.05 (brs, 1H), 10.01 (brs, 1H), 12.86 (brs, 1H), ^{31}P NMR ($\text{d}_6\text{-DMSO}$, 121.49 MHz) δ 30.0, MS (ES $^-$) $m/z = 427/429$ (MH^+).


7aw




7ax



7ay



7az




Compound


structure

Description


7aaa




7aab




7aac



7aad



7aae




Compound


structure

Description

7aaf




8a




Off white solid; ^1H NMR (CDCl_3 , 250 MHz) δ 1.43 (t, J = 7.0 Hz, 3H), 2.33 (s, 6H), 4.03-4.10 (m, 1H), 4.28-4.35 (m, 1H), 7.20 (s, 1H), 7.35 (dd, J = 2.0 and 8.7 Hz, 1H), 7.41 (s, 1H), 7.46 (s, 1H), 7.56-7.60 (m, 1H), 10.7 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 33.9, MS (ESI, EI^+) m/z = 392 (MH^+).

9a




Pale yellow solid, ^1H NMR (CDCl_3 , 300 MHz) δ 1.45 (t, J = 7.2 Hz, 3H), 2.29 (s, 6H), 4-4.13 (m, 1H), 4.22-4.33 (m, 1H), 4.65 (dd, J = 5.6 and 16.1 Hz, 1H), 4.80 (dd, J = 6.1 and 16.1 Hz, 1H), 7.15 (brs, 1H), 7.24-7.34 (m, 6H), 7.75 (d, J = 2.1 Hz, 1H), 8.54-8.56 (m, 2H), 10.60 (brs, 1H), 11.81 (t, J = 5.7 Hz, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 32.8, MS (ESI, EI^+) m/z = 482 (MH^+).

11a




Off white solid, ^1H NMR (d_6 -DMSO) δ 1.37 (t, J = 7.2 Hz, 3H), 4.48 (q, J = 7.2 Hz, 2H), 7.59-7.68 (m, 4H), 7.77 (m, 1H), 7.96-8.09 (m, 3H), MS (ESI, EI^+) m/z = 442-444 (MH^+).

11b




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 4 (s, 3H), 7.66-7.72 (m, 3H), 7.78-7.83 (m, 1H), 7.93 (d, J = 9 Hz, 1H), 7.99-8.02 (m, 2H), MS (ES^+) m/z = 468.18/470.15 (MH^+).

11c




Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.46 (t, J = 7.2 Hz, 3H), 4.53 (q, J = 7.1 Hz, 2H), 7.31-7.36 (m, 1H), 7.42-7.59 (m, 5H), 7.95-8.04 (m, 3H), MS (ESI, EI^+) m/z = 408 (MH^+).

14a




Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.37 (s, 3H), 7.45 (d, J = 544 Hz, 1H), 7.35 (dd, J = 2.85 and 7.95 Hz, 2H), 7.57 (d, J = 13.2 Hz, 1H), 7.6 (d, J = 13.5 Hz, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 23.8.

16a




Yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.37 (t, J = 7.2 Hz, 3H), 3.75 (d, J = 11.7 Hz, 3H), 4.46 (q, J = 7.2 Hz, 2H), 7.54-7.58 (m, 1H), 7.67-7.9 (m, 5H), 8.07-8.24 (m, 6H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 22.98, MS (ES^+) m/z = 542.7 (MH^+).

16b




Thick yellow oil; ^1H NMR (CDCl_3 , 300 MHz) δ 1.46 (t, J = 7.05 Hz, 3H), 3.8 (d, J = 11.7 Hz, 3H), 4.54 (q, J = 7.05 Hz, 2H), 7.32-7.38 (m, 2H), 7.5-8.1 (m, 10H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 24.54, MS (ES^+) m/z = 595.74/597.72 (MH^+).

16c




Thick yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.36 (t, J = 7.2 Hz, 3H), 3.7 (d, J = 11.7 Hz, 3H), 4.44 (q, J = 7.2 Hz, 2H), 5.49 (d, J = 47.4 Hz, 2H), 7.54 (dd, J = 2.1 and 9 Hz, 1H), 7.6-7.88 (m, 8H), 8.05-8.11 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 23.82, MS (ES^+) m/z = 550 (MH^+).

16d




Yellowish powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.26 (d, J = 6.9 Hz, 6H), 1.28 (t, J = 6.9 Hz, 3H), 1.48 (t, J = 6.9 Hz, 3H), 2.96 (quint., J = 6.9 Hz, 1H), 3.83 (d, J = 11.4 Hz, 3H), 4.56 (q, J = 6.9 Hz, 2H), 7.07-7.13 (m, 1H), 7.37 (dd, J = 2.1 Hz and 9.0 Hz, 1H), 7.31-7.40 (m, 1H), 7.49-7.68 (m, 4H), 7.85 (d, J = 1.8 Hz, 1H), 7.95 (dd, J = 9.6 Hz and 1.8 Hz, 1H), 8.08-8.14 (m, 2H), MS (ES^+) m/z =

16e

578 (MH⁺).


Yellow resin, ¹H NMR (*d*₆-DMSO, 300 MHz) δ 0.83 (t, *J* = 7.5 Hz, 3H), 1.38 (t, *J* = 7.2 Hz, 3H), 1.55 (sextuplet, *J* = 7.5 Hz, 2H), 2.59 (t, *J* = 7.5 Hz, 2H), 3.69 (d, *J* = 11.7 Hz, 3H), 4.45 (q, *J* = 7.2 Hz, 2H), 7.44-7.8 (m, 9H), 8.05-8.12 (m, 3H), ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 24.36, MS (ES⁺) *m/z* = 560.38 (MH⁺).

16f




Thick yellow oil, ¹H NMR (*d*₆-DMSO, 300 MHz) δ 2.29 (s, 6H), 3.64 (d, *J* = 11.7 Hz, 3H), 4.04 (s, 3H), 7.25-7.34 (m, 3H), 7.6-7.75 (m, 3H), 7.81-7.84 (m, 1H), 7.9-7.94 (m, 1H), 8.12-8.16 (m, 2H), MS (ES⁺) *m/z* = 550 (MH⁺).

16g




Off-white solid, ¹H NMR (CDCl₃, 300 MHz) δ 0.66-0.71 (m, 2H), 0.91-0.99 (m, 2H), 1.81-1.94 (m, 1H), 3.69 (s, 3H), 3.79 (d, *J* = 11.4 Hz, 3H), 7.13-7.17 (m, 1H), 7.28-7.4 (m, 3H), 7.51-7.64 (m, 2H), 8.39-8.4 (m, 1H), 9.96 (brs, 1H), ³¹P NMR (CDCl₃, 121.49 MHz) δ 31.29.

16h




Thick yellow oil, ¹H NMR (*d*₆-DMSO, 300 MHz) δ 1.15 (t, *J* = 7.5 Hz, 3H), 1.39 (t, *J* = 7.2 Hz, 3H), 2.32 (s, 3H), 3.69 (d, *J* = 11.7 Hz, 3H), 4.46 (q, *J* = 7.5 Hz, 2H), 7.29 (brs, 1H), 7.4-7.46 (m, 2H), 7.53-7.56 (m, 1H), 7.63-7.81 (m, 4H), 8.03-8.12 (m, 3H), ³¹P NMR (*d*₆-DMSO, 121.49 MHz) δ 24.57, MS (ES⁺) *m/z* = 560.5 (MH⁺).

16i




Yellow resin, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.39 (t, J = 7.2 Hz, 3H), 2.35 (s, 3H), 1.38 (t, J = 7.2 Hz, 3H), 3.71 (d, J = 11.5 Hz, 3H), 4.46 (q, J = 7.2 Hz, 2H), 7.53-7.85 (m, 8H), 8.05-8.12 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 22.40, MS (ES^+) m/z = 610/612 (MH^+).

16j




Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.45 (t, J = 7.2 Hz, 3H), 3.80 (d, J = 11.4 Hz, 3H), 4.54 (q, J = 7.2 Hz, 2H), 7.36 (dd, J = 2.1 and 9.0 Hz, 1H), 7.47-7.67 (m, 6H), 7.84-7.96 (m, 4H), 8.09-8.12 (m, 2H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 26.7, MS (ESI, EI^+) m/z = 518 (MH^+).

16k




Pale yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.36 (t, J = 7.5 Hz, 3H), 3.69 (d, J = 11.7 Hz, 3H), 4.45 (q, J = 7.1 Hz, 2H), 7.41 (td, J = 8.9 and 2.5 Hz, 2H), 7.54 (dd, J = 9.0 and 2.1 Hz, 1H), 7.66-7.71 (m, 2H), 7.78-7.90 (m, 4H), 8.05-8.11 (m, 3H), ^{19}F NMR (d_6 -DMSO) δ -105.46 (m, 1F), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 24.32, MS (ESI, EI^+) m/z = 558 ($\text{M}+\text{Na}^+$).

16m




Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.38 (t, J = 7.05 Hz, 3H), 3.7 (d, J = 11.7 Hz, 3H), 3.78 (s, 3H), 4.46 (q, J = 7.05 Hz, 2H), 7.19-7.4 (m, 3H), 7.47-7.56 (m, 2H), 7.66-7.71 (m, 2H), 7.79-7.81 (m, 2H), 8.06-8.12 (m, 3H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 24.91.

16n




Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.37 (t, $J = 7.1$ Hz, 3H), 2.29 (s, 6H), 3.66 (d, $J = 11.4$ Hz, 3H), 4.44 (q, $J = 7.1$ Hz, 2H), 7.25 (s, 1H), 7.37 (s, 1H), 7.42 (s, 1H), 7.53 (dd, $J = 2.1$ and 9.0 Hz, 1H), 7.65-7.71 (m, 2H), 7.78-7.83 (m, 2H), 8.05-8.11 (m, 3H), MS (ESI, EI^+) $m/z = 546$ (MH^+).

16p



Brown oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 3H), 3.69 (d, $J = 11.4$ Hz, 3H), 4.42 (q, $J = 7.05$ Hz, 2H), 4.52 (s, 2H), 4.57 (s, 2H), 7.28-7.37 (m, 6H), 7.52-7.78 (m, 9H), 8.05-8.1 (m, 2H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 24.18, MS (ES^+) $m/z = 638$ (MH^+).

16q




Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.37 (t, $J = 7.2$ Hz, 3H), 2.4 (s, 3H), 3.73 (d, $J = 11.4$ Hz, 3H), 4.45 (q, $J = 7.2$ Hz, 2H), 7.55 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.66-7.71 (m, 2H), 7.78-7.82 (m, 1H), 7.89-7.9 (m, 2H), 7.94-8.12 (m, 5H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 22.16, MS (ES^+) $m/z = 557.4$ (MH^+).

Compound

structure

Description

17a



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.33 (s, 3H), 3.73 (d, $J = 11.7$ Hz, 3H), 7.36-7.43 (m, 3H), 7.57-7.66 (m, 3H), 7.81-7.82 (m, 1H), 13.01 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.55, MS (ES^+) $m/z = 362$ (M-H^+).

Compound	structure	Description
17b		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.15 (t, $J = 7.5$ Hz, 3H), 2.3 (s, 3H), 2.61 (q, $J = 7.5$ Hz, 2H), 3.73 (d, $J = 11.7$ Hz, 3H), 7.28 (brs, 1H), 7.36-7.40 (m, 1H), 7.39-7.5 (m, 2H), 7.57-7.61 (m, 1H), 7.78 (brs, 1H), 13.01 (brs, 1H), 14.82 (brs, s, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.89, MS (ES^+) $m/z = 392$ (MH^+).
17c		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.29 (s, 6H), 3.78 (d, $J = 11.7$ Hz, 3H), 7.28 (brs, 1H), 7.37 (brs, 1H), 7.42 (brs, 1H), 7.46-7.48 (m, 2H), 13.35 (s, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 35.15, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -115.84 (s, 1F) + MS (ES^+) $m/z = 395.9$ (MH^+).
17d		Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.88 (d, $J = 12.0$ Hz, 3H), 7.37 (dd, $J = 2.0$ and 9.0 Hz, 1H), 7.47-7.63 (m, 5H), 7.81-7.89 (m, 2H), 10.33 (brs, 1H), MS (ESI , EI^+) $m/z = 350$ (MH^+).
17e		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.71 (d, $J = 11.7$ Hz, 3H), 3.76 (s, 3H), 7.13-7.16 (m, 1H), 7.29-7.45 (m, 4H), 7.56 (dd, $J = 1.8$ and 9.0 Hz, 1H), 7.91 (d, $J = 1.8$ Hz, 1H), 12.86 (brs, 1H), 14.51 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 31.68, MS (ESI , EI^+) $m/z = 380$ (MH^+).
17f		Pale brown oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.72 (d, $J = 11.7$ Hz, 3H), 7.25-7.64 (m, 6H), 8.06 (d, $J = 2.1$ Hz, 1H), 12.99 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 29.23 (d, $J = 7.3$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 300 MHz) δ -112.01 (m, 1F), MS (ESI , EI^+) $m/z = 368$ (MH^+).


Compound	structure	Description
17g		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.28 (s, 6H), 3.71 (d, J = 11.7 Hz, 3H), 7.2-7.24 (m, 1H), 7.36-7.45 (m, 3H), 7.57-7.6 (m, 1H), 7.78-7.79 (m, 1H), 13.05 (brs, 1H); 14.8 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.87, MS (ES^+) m/z = 378 (MH^+).
17h		Off-white solid, MS (ES^+) m/z = 389 (MH^+).
18a		Pink solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.3 (s, 3H), 3.76 (d, J = 11.7 Hz, 3H), 4.57-4.75 (m, 2H), 7.31-7.61 (m, 9H), 8.52-8.54 (m, 2H), 11.37 (t, J = 5.7 Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.81, MS (ES^+) m/z = 453.9 (MH^+).
18b		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.3 (s, 3H), 3.74 (d, J = 11.4 Hz, 3H), 4.56-4.65 (m, 2H), 6.38-6.45 (m, 2H), 7.31-7.65 (m, 8H), 11.23-11.27 (m, 1H), 12.84 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.72, MS (ES^+) m/z = 442.9 (MH^+).
18c		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.78 (d, J = 11.7 Hz, 3H), 4.58-4.74 (m, 2H), 5.46 (d, J = 47.4 Hz, 2H), 7.32-7.38 (m, 3H), 7.55-7.82 (m, 6H), 8.51-8.53 (m, 2H), 11.3 (t, J = 5.7 Hz, 1H), 12.92 (d, J = 2.1 Hz, 1H), ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -211.09 (t, J = 47.4 Hz, 1F), MS (ES^+) m/z = 472 (MH^+).

Compound

structure


Description

18d




Off-white powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.35 (s, 3H), 3.88 (d, $J = 11.7$ Hz, 3H), 4.77 (qd, $J = 6$ Hz and 15.9 Hz, 2H), 7.04-7.35 (m, 7H), 7.68 (brs, 1H), 8.59-9.61 (m, 2H), 11.22 (brs, 1H), 11.71 (t, $J = 5.7$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 32.55 (d, $J = 8.2$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -111.55 (q, $J = 8.2$ Hz, 1F), MS (ES^+) $m/z = 471.92$ (MH^+).

18e




Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.88 (d, $J = 11.7$ Hz, 3H), 4.7-4.77 (m, 2H), 6.97 (tt, $J = 2.4$ Hz and 8.7 Hz, 1H), 7.2-7.33 (m, 6H), 7.64 (brs, 1H), 8.52-8.6 (m, 2H), 11.55-11.58 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.11 (t, $J = 8.5$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -106.21 (m, 2F), MS (ES^+) $m/z = 476.17$ (MH^+).

18f



White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.28 (s, 6H), 3.82 (d, $J = 11.7$ Hz, 3H), 4.69 (dd, $J = 6$ Hz and 15 Hz, 1H), 4.78 (dd, $J = 6$ Hz and 15 Hz, 1H), 7.13 (brs, 1H), 7.24-7.28 (m, 5H), 7.29-7.31 (m, 1H), 7.38-7.39 (m, 1H), 7.68-7.69 (m, 1H), 7.76-7.8 (m, 1H), 8.56 (dd, $J = 1.5$ Hz and 4.5 Hz, 1H), 8.7 (d, $J = 2.1$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.18, MS (ES^+) $m/z = 468.33$ (MH^+).

18g




White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.3 (s, 6H), 3.03 (t, $J = 7.2$ Hz, 2H), 3.76-3.88 (m, 2H), 3.79 (d, $J = 11.7$ Hz, 3H), 7.16-7.3 (m, 6H), 7.47-7.5 (m, 1H), 7.62-7.63 (m, 1H), 8.43-8.45 (m, 2H), 10.83 (brs, 1H), 11.36-11.38 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.38, MS (ES^+) $m/z = 482.38$ (MH^+).

Compound

structure


Description

18h




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.75 (d, $J = 11.4$ Hz, 3H), 4.66 (dd, $J = 5.4$ and 15.9 Hz, 1H), 4.78 (dd, $J = 5.7$ and 15.9 Hz, 1H), 7.24 (brs, 1H), 7.3-7.35 (m, 3H), 7.45-7.49 (m, 1H), 7.6 (dd, $J = 1.8$ and 9 Hz, 1H), 7.62 (d, $J = 1.8$ Hz, 1H), 8.4 (dd, $J = 1.2$ and 4.8 Hz, 1H), 8.58 (d, $J = 1.5$ Hz, 1H), 11.41 (t, $J = 5.4$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.81, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -132.93 (d, $J = 5.9$ Hz, 1F), MS (ES^+) $m/z = 486.3$ (MH^+).

18i




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.75 (d, $J = 11.4$ Hz, 3H), 4.7 (dd, $J = 5.1$ and 15.9 Hz, 1H), 4.84 (dd, $J = 6$ and 15.9 Hz, 1H), 7.22 (brs, 1H), 7.33 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.32-7.36 (m, 2H), 7.59 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.63 (d, $J = 1.5$ Hz, 1H), 8.6 (dd, $J = 2.55$ and 14.4 Hz, 1H), 8.62 (d, $J = 2.4$ Hz, 1H), 8.72 (d, $J = 1.5$ Hz, 1H), 11.46 (t, $J = 5.7$ Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.69, MS (ES^+) $m/z = 469.4$ (MH^+).

18k



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (s, 6H), 3.74 (d, $J = 11.7$ Hz, 3H), 4.51 (dd, $J = 5.25$ and 15 Hz, 1H), 4.67 (dd, $J = 6.0$ and 15 Hz, 1H), 7.22-7.38 (m, 6H), 7.53-7.64 (m, 4H), 11.34 (t, $J = 5.7$ Hz, 1H), 12.66 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.79, MS (ES^+) $m/z = 545.33/547.31$ (MH^+).

18l




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.28 (s, 6H), 3.77 (d, $J = 11.4$ Hz, 3H), 4.64-4.82 (m, 2H), 7.24 (brs, 1H), 7.34 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.36 (brs, 1H), 7.4 (brs, 1H), 7.52 (dd, $J = 1.5$ and 5.4 Hz, 1H), 7.58-7.63 (m, 2H), 8.76 (d, $J = 5.1$ Hz, 1H), 9.14 (d, $J = 1.5$ Hz, 1H), 11.48 (t, $J = 6.0$ Hz, 1H), 12.85 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.68, MS (ES^+) $m/z = 469.4$ (MH^+).

Compound

structure


Description

18m




Off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 1.19 (t, $J = 7.6$ Hz, 3H), 2.60 (q, $J = 7.6$ Hz, 2H), 2.31 (s, 3H), 3.84 (d, $J = 11.7$ Hz, 3H), 4.76 (m, 2H), 7.18-7.43 (m, 6H), 7.70 (m, 1H), 7.81 (m, 1H), 8.57 (d, $J = 4.8$ Hz, 1H), 8.72 (s, 1H), 10.62 (brs, 1H), 11.78 (t, $J = 7.6$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.08.

18n




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (s, 6H), 3.74 (d, $J = 11.5$ Hz, 3H), 4.39 (dd, $J = 5.7$ and 15.6 Hz, 1H), 4.53 (dd, $J = 5.7$ and 15.6 Hz, 1H), 5.62 (brs, 2H), 6.52 (t, $J = 7.2$ Hz, 1H), 6.63-6.70 (m, 1H), 6.97-7.35 (m, 6H), 7.58-7.65 (m, 2H), 11.34 (t, $J = 5.7$ Hz, 1H), 12.66 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 8.51, MS (ES^+) $m/z = 482$ (MH^+).

18p







Off-white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.12 (t, $J = 7.5$ Hz, 3H), 2.28 (s, 3H), 2.58 (q, $J = 7.5$ Hz, 2H), 3.77 (d, $J = 11.7$ Hz, 3H), 4.68 (dd, $J = 5.4$ and 17.1 Hz, 1H), 4.78 (dd, $J = 5.7$ and 16.8 Hz, 1H), 7.27 (brs, 1H), 7.33 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.36-7.46 (m, 2H), 7.52 (dd, $J = 1.2$ and 5.1 Hz, 1H), 7.60 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.63 (d, $J = 1.5$ Hz, 1H), 8.76 (d, $J = 5.4$ Hz, 1H), 9.14 (d, $J = 1.2$ Hz, 1H), 11.47 (t, $J = 5.4$ Hz, 1H), 12.82 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.62, MS (ES^+) $m/z = 483$ (MH^+).

18q




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.78 (d, $J = 11.7$ Hz, 3H), 4.61-4.78 (m, 2H), 7.35 (dd, $J = 2.1$ and 8.7 Hz, 2H), 7.38-7.41 (m, 1H), 7.5-7.55 (m, 1H), 7.57-7.6 (m, 3H), 7.73 (d, $J = 2.1$ Hz, 1H), 7.82-7.85 (m, 2H), 10.87-10.91 (m, 1H), 12.96 (brs, 1H), MS (ES^+) $m/z = 498.35$ (MH^+).


Compound	structure	Description
18r		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.27 (s, 6H), 3.72 (d, $J = 11.7$ Hz, 3H), 4.65-4.68 (m, 2H), 7.22 (brs, 1H), 7.29 (brs, 1H), 7.33 (brs, 1H), 7.38-7.48 (m, 4H), 8.53-8.55 (m, 2H), 11.86 (t, $J = 5.4$ Hz, 1H), 13.08 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.16, MS (ES^+) $m/z = 485.9$ (MH^+).
18s		
18t		
18u		

Compound	structure	Description
----------	-----------	-------------


18v




18w







18x







18y



Compound	structure	Description
----------	-----------	-------------




Compound	structure	Description
----------	-----------	-------------




Compound	structure	Description
----------	-----------	-------------


18ah




18ai



18aj




18ak




Compound	structure	Description
----------	-----------	-------------


18al




18am







18an



18ao



Compound	structure	Description
18ap		Yellow pale powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.12 (t, $J = 7.5$ Hz, 3H), 2.61 (q, $J = 7.5$ Hz, 2H), 3.77 (d, $J = 11.7$ Hz, 3H), 4.61 (dd, $J = 5.4$ and 15.9 Hz, 1H), 4.72 (dd, $J = 6.0$ and 15.9 Hz, 1H), 7.31-7.61 (m, 9H), 8.51-8.53 (m, 2H), 11.4 (t, $J = 5.4$ Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.74, MS (ES^+) $m/z = 467.89$ (MH^+).
18aq		Yellow pale powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.15 (d, $J = 6.9$ Hz, 6H), 2.9 (quintuplet, $J = 6.9$ Hz, 1H), 3.77 (d, $J = 11.4$ Hz, 3H), 4.61 (dd, $J = 5.4$ and 16.5 Hz, 1H), 4.72 (dd, $J = 6$ and 16.2 Hz, 1H), 7.33 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.37-7.39 (m, 2H), 7.41-7.54 (m, 3H), 7.58-7.66 (m, 3H), 8.52-8.54 (m, 2H), 11.42 (t, $J = 5.7$ Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 33.64, MS (ES^+) $m/z = 481.94$ (MH^+).
18ar		Pale yellow solid, ^1H NMR (CDCl_3 , 400 MHz) δ 3.86 (d, $J = 11.8$, 3H), 4.67 (dd, $J = 5.7$ and 15.9 Hz, 1H), 4.79 (dd, $J = 6.0$ and 15.9 Hz, 1H), 7.25-7.28 (dd, $J = 1.9$ and 8.4 Hz, 1H), 7.30-7.34 (m, 3H), 7.41-7.45 (m, 2H), 7.52-7.56 (m, 1H), 7.69-7.75 (m, 3H), 8.55-8.57 (m, 2H), 10.76 (s, 1H), 11.72 (t, $J = 5.8$ Hz, 1H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 34.2, MS (ESI, EI^+) $m/z = 440$ (MH^+).
18as		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.62 (s, 3H), 3.74 (d, $J = 11.7$ Hz, 3H), 4.53-4.67 (m, 2H), 6.9 (brs, 1H), 7.32 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.46-7.65 (m, 8H), 11.09 (t, $J = 5.1$ Hz, 1H), 12.85 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 33.71, MS (ESI, EI^+) $m/z = 443$ (MH^+).


Compound	structure	Description
18at		Off white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.76 (d, $J = 11.4$ Hz, 3H), 4.56 (dd, $J = 4.8$ and 15.6 Hz, 1H), 4.66 (dd, $J = 5.7$ and 15.6 Hz, 1H), 6.38-6.45 (m, 2H), 7.33 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.48-7.7 (m, 8H), 11.21 (t, $J = 5.25$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 33.59, MS (ESI, EI^+) $m/z = 429$ (MH^+).
18au		Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.76 (s, 3H), 3.84 (d, $J = 11.4$ Hz, 3H), 4.7 (dd, $J = 5.1$ and 15.6 Hz, 1H), 4.83 (dd, $J = 5.5$ and 15.75 Hz, 1H), 6.33-6.36 (m, 2H), 7-7.04 (m, 1H), 7.23-7.32 (m, 4H), 7.38-7.4 (m, 1H), 7.48 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.7 (d, $J = 2.1$ Hz, 1H), 11.61-11.64 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.64, MS (ESI, EI^+) $m/z = 459$ (MH^+).
18av		Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.76 (s, 3H), 3.86 (d, $J = 11.7$ Hz, 3H), 4.69 (dd, $J = 5.7$ and 15.9 Hz, 1H), 4.83 (dd, $J = 5.7$ and 15.9 Hz, 1H), 7.03-7.07 (m, 1H), 7.2-7.37 (m, 7H), 7.71 (brs, 1H), 8.55-8.58 (m, 2H), 11.48 (brs, 1H), 11.8 (t, $J = 5.7$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.93, MS (ESI, EI^+) $m/z = 470$ (MH^+).
18aw		Pale yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.78 (d, $J = 11.7$ Hz, 3H), 4.65 (m, 2H), 7.33-7.65 (m, 9H), 8.52 (d, $J = 6$ Hz, 2H), 11.13 (t, $J = 5.5$ Hz, 1H), 12.94 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 30.76 (d, $J = 7.4$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 300 MHz) δ -111.0 (m, 1F), MS (ESI, EI^+) $m/z = 458$ (MH^+).

Compound

structure


Description

18ax




Yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.25 (s, 6H), 3.75 (d, $J = 11.4$ Hz, 3H), 4.6 (dd, $J = 5.4$ and 16.2 Hz, 1H), 4.72 (dd, $J = 5.7$ and 16.2 Hz, 1H), 7.24-7.39 (m, 6H), 7.59-7.63 (m, 2H), 8.52-8.54 (m, 2H), 11.41 (t, $J = 5.7$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.88, MS (ES^+) $m/z = 467.9$ (MH^+).

18ay




White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.3 (s, 6H), 2.7-2.75 (m, 4H), 3.73-3.77 (m, 4H), 3.86 (d, $J = 11.7$ Hz, 3H), 4.43 (dd, $J = 5.7$ and 12.9 Hz, 1H), 4.52 (dd, $J = 6$ and 12.9 Hz, 1H), 7.17 (brs, 1H), 7.32 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.36 (brs, 1H), 7.41 (brs, 1H), 7.49 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.71-7.72 (m, 1H), 10.5 (brs, 1H), 11.46 (t, $J = 5.7$ Hz, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 34.52, MS (ES^+) $m/z = 475.87$ (MH^+).

18az



Pale yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (s, 6H), 3.73 (d, $J = 11.7$ Hz, 3H), 4.56 (dd, $J = 5.4$ and 15.3 Hz, 1H), 4.73 (dd, $J = 5.4$ and 15.3 Hz, 1H), 7.15-7.4 (m, 7H), 7.47 (td, $J = 1.8$ and 7.5 Hz, 1H), 7.59 (dd, $J = 1.8$ and 8.1 Hz, 1H), 7.63 (d, $J = 2.1$ Hz, 1H), 11.32 (t, $J = 5.4$ Hz, 1H), 12.83 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.76, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -118.4 (m, 1F), MS (ES^+) $m/z = 485$ (MH^+).

18aaa



White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.25 (s, 6H), 3.66 (s, 3H), 3.72 (d, $J = 11.7$ Hz, 3H), 4.52-4.7 (m, 2H), 6.9 (brs, 1H), 7.22 (brs, 2H), 7.26 (brs, 1H), 7.33 (dd, $J = 2.1$ and 9.3 Hz, 1H), 7.59-7.61 (m, 3H), 11.17 (t, $J = 5.1$ Hz, 1H), 12.83 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.93, MS (ES^+) $m/z = 471.36$ (MH^+).


Compound	structure	Description
18aab		Off-white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.75 (d, $J = 11.7$ Hz, 3H), 4.58 (dd, $J = 5.1$ and 15.3 Hz, 1H), 4.7 (dd, $J = 5.7$ and 15.3 Hz, 1H), 7.17-7.28 (m, 2H), 7.32-7.52 (m, 5H), 7.57-7.7 (m, 5H), 11.24 (t, $J = 5.4$ Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.64, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -118.41 (m, 1F), MS (ES^+) $m/z = 456.8$ (MH^+).
18aac		Off white powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.67-2.72 (m, 4H), 3.71-3.75 (m, 4H), 3.86 (d, $J = 11.7$ Hz, 3H), 4.39-4.5 (m, 2H), 7.3 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.41-7.54 (m, 4H), 7.66-7.67 (m, 1H), 7.73-7.81 (m, 2H), 10.49 (brs, 1H), 11.42 (t, $J = 5.7$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.56, MS (ES^+) $m/z = 447.92$ (MH^+).
18aad		Pale yellow powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.78 (d, $J = 11.7$ Hz, 3H), 4.54-4.74 (m, 2H), 7.16-7.28 (m, 2H), 7.28-7.65 (m, 7H), 7.74-7.75 (m, 1H), 10.84 (brs, 1H), 12.95 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 29.36, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -138 (m, 1F), -126.5 (m, 2F), MS (ES^+) $m/z = 493$ (MH^+).
18aae		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.64 (s, 3H), 3.77 (d, $J = 11.4$ Hz, 3H), 4.54-4.67 (m, 2H), 6.91 (brs, 1H), 7.33-7.39 (m, 3H), 7.5-7.62 (m, 3H), 7.67-7.68 (m, 1H), 10.68-10.72 (m, 1H), 12.95 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 30.92 (t, $J = 8.5$ Hz, 1P), ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -106.77 (m, 2F), MS (ES^+) $m/z = 479$ (MH^+).

Compound

structure


Description

18aaf




Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.39 (s, 3H), 3.88 (d, $J = 11.7$ Hz, 3H), 4.76 (d, $J = 5.4$ Hz, 2H), 7.25-7.33 (m, 4H), 7.57-7.61 (m, 2H), 7.71-7.8 (m, 2H), 8.58-8.6 (m, 2H), 11.27 (brs, 1H), 11.59 (t, $J = 5.4$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.35, MS (ES^+) $m/z = 479.35$ (MH^+).

18aag




Pale yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.39 (s, 3H), 3.72 (s, 3H), 3.84 (d, $J = 11.4$ Hz, 3H), 4.71 (d, $J = 5.4$ Hz, 2H), 7.12 (s, 1H), 7.32 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.47-7.59 (m, 4H), 7.66-7.74 (m, 2H), 11.21 (brs, 1H), 11.38 (t, $J = 5.4$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.21, MS (ES^+) $m/z = 482.38$ (MH^+).

18aaj







Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (s, 6H), 3.72 (d, $J = 11.7$ Hz, 3H), 4.53 (dd, $J = 5.1$ and 15.3 Hz, 1H), 4.68 (dd, $J = 5.4$ and 15 Hz, 1H), 7.06-7.12 (m, 1H), 7.21-7.34 (m, 5H), 7.48-7.53 (m, 1H), 7.58 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.63 (d, $J = 1.8$ Hz, 1H), 11.29 (t, $J = 5.1$ Hz, 1H), 12.82 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.79, MS (ES^+) $m/z = 503$ (MH^+).









18aak











Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.74 (d, $J = 11.4$ Hz, 3H), 4.60 (dd, $J = 5.4$ and 15.3 Hz, 1H), 4.77 (dd, $J = 5.4$ and 15.3 Hz, 1H), 7.22 (brs, 1H), 7.29 (brs, 1H), 7.31-7.38 (m, 4H), 7.49-7.52 (m, 2H), 7.6 (dd, $J = 1.5$ and 9 Hz, 1H), 7.64 (d, $J = 1.8$ Hz, 1H), 11.35 (t, $J = 5.4$ Hz, 1H), 12.84 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.72, MS (ES^+) $m/z = 501$ (MH^+).








Compound	structure	Description
18aal		Light yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (s, 6H), 3.73 (d, $J = 11.7$ Hz, 3H), 4.71 (dd, $J = 4.8$ and 15.3 Hz, 1H), 4.85 (dd, $J = 4.8$ and 15.3 Hz, 1H), 7.22-7.31 (m, 3H), 7.32 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.5-7.56 (m, 1H), 7.6 (dd, $J = 1.8$ and 9 Hz, 1H), 7.64 (dd, $J = 1.8$ and 10.5 Hz, 1H), 7.67 (d, $J = 1.2$ Hz, 1H), 7.76-7.79 (m, 1H), 11.38 (t, $J = 5.4$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.74, MS (ES^+) $m/z = 535$ (MH^+)
18aam		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.75 (d, $J = 11.7$ Hz, 3H), 4.64 (dd, $J = 5.4$ and 15.9 Hz, 1H), 4.78 (dd, $J = 6$ and 15.9 Hz, 1H), 7.23 (brs, 1H), 7.29-7.42 (m, 5H), 7.6 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.65 (d, $J = 1.8$ Hz, 1H), 7.78 (td, $J = 1.8$ and 7.5 Hz, 1H), 8.54-8.57 (m, 1H), 11.39 (t, $J = 5.7$ Hz, 1H), 12.82 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.70, MS (ES^+) $m/z = 468$ (MH^+)
18aan		White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.69 (d, $J = 11.7$ Hz, 3H), 4.65 (ddd, $J = 5.7$ and 15.3 and 21.6 Hz, 2H), 7.21 (brs, 1H), 7.25 (brs, 1H), 7.3 (brs, 1H), 7.37-7.45 (m, 2H), 7.46 (dd, $J = 1.5$ and 8.7 Hz, 1H), 7.82 (td, $J = 1.8$ and 7.8 Hz, 1H), 8.49 (dd, $J = 1.95$ and 4.65 Hz, 1H), 8.64 (d, $J = 1.8$ Hz, 1H), 11.81 (t, $J = 5.7$ Hz, 1H), 13.06 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.10, MS (ES^+) $m/z = 486$ (MH^+)
18aao		Light yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.75 (d, $J = 11.4$ Hz, 3H), 5.67 (d, $J = 7.5$ Hz, 1H), 7.2 (brs, 1H), 7.85 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.85-7.95 (m, 3H), 8.02-8.16 (m, 6H), 8.22 (d, $J = 1.8$ Hz, 1H), 8.28 (brs, 1H), 8.46-8.53 (m, 2H), 11.47 (d, $J = 7.5$ Hz, 1H), 12.76 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 31.79, MS (ES^+) $m/z = 482$ (MH^+)

Compound	structure	Description
18aap		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.80 (d, $J = 11.4$ Hz, 3H), 5.63 (d, $J = 7.5$ Hz, 1H), 7.2 (brs, 1H), 7.33 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.33-7.45 (m, 5H), 7.52-7.61 (m, 6H), 7.69 (d, $J = 2.1$ Hz, 1H), 7.77 (brs, 1H), 11.32 (d, $J = 7.5$ Hz, 1H), 12.8 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.12.
18aaq		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.77 (d, $J = 11.4$ Hz, 3H), 5.34 (d, $J = 11.1$ Hz, 1H), 5.86 (d, $J = 17.7$ Hz, 1H), 6.79 (dd, $J = 11.1$ and 17.7 Hz, 1H), 7.32 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.48-7.61 (m, 4H), 7.73 (dd, $J = 1.5$ and 7.8 Hz, 1H), 7.78-7.83 (m, 1H), 8.03 (brs, 1H), 10.2 (brs, 1H), 12.8 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 31.86, MS (ES^+) $m/z = 374.83$ (MH^+)
18aar		Sticky off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 3.84 (d, $J = 11.4$ Hz, 3H), 4.70 (dd, $J = 5.4$ and 15 Hz, 1H), 4.85 (dd, $J = 6$ and 15 Hz, 1H), 7.16 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.22-7.51 (m, 9H), 7.65-7.73 (m, 3H), 11.63 (brs, 1H), 11.68 (t, $J = 5.4$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.98, MS (ES^+) $m/z = 438.87$ (MH^+)
18aas		
18aat		

Compound	structure	Description
19a		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.76 (d, J = 11.1 Hz, 3H), 7.44-7.58 (m, 3H), 7.76-7.84 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 22.2.
19b		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.81 (d, J = 11.1 Hz, 6H), 7.59-7.66 (m, 1H), 7.84-7.87 (m, 1H), 8-8.1 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 17.94, MS (ES^+) m/z = 212 (MH).
19c		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.27 (t, J = 7.5 Hz, 3H), 2.72 (q, J = 7.5 Hz, 2H), 3.78 (d, J = 11.1 Hz, 6H), 7.4-7.43 (m, 2H), 7.59-7.69 (m, 2H), MS (ES^+) m/z = 215 (MH).
19d		Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.77 (d, J = 11.1 Hz, 6H), 7.28-7.39 (m, 2H), 7.57-7.66 (m, 2H), MS (ES^+) m/z = 201 (MH).
19e		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.78 (d, J = 11.4 Hz, 6H), 7.33-7.4 (m, 1H), 7.69-7.77 (m, 2H), 7.91-7.96 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.2, MS (ES^+) m/z = 265/267 (MH).
19f		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.79 (d, J = 11.1 Hz, 6H), 5.44 (d, J = 47.4 Hz, 2H), 7.5-7.63 (m, 2H), 7.78-7.85 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 22, ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -209.94 (t, J = 47.4 Hz, 1F), MS (ES^+) m/z = 219 (MH).
19g		Yellowish oil, ^1H NMR (CDCl_3 , 300 MHz) δ 2.41 (s, 3H), 3.77 (d, J = 11.1 Hz, 6H), 7.06-7.1 (m, 1H), 7.24-7.32 (m, 1H), 7.4-7.44 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 20.13 (d, J = 9.7 Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -112.5 (m, 1F).
19h		Colourless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 0.89 (t, J = 7.5 Hz, 3H), 1.6 (sextuplet, J = 7.5 Hz, 2H), 2.63 (t, J = 7.5 Hz, 2H), 3.65 (d, J = 11.4 Hz, 3H), 3.654 (d, J = 11.1 Hz, 3H), 7.47-7.50 (m, 3H), 7.54-7.55 (m, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 21.04, MS (ES^+) m/z = 229.3

Compound	structure	Description
19i		(MH ⁺). Colorless oil, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 1.18 (t, <i>J</i> = 7.5 Hz, 3H), 2.35 (s, 3H), 2.63 (q, <i>J</i> = 7.5 Hz, 2H), 3.64 (d, <i>J</i> = 11.1 Hz, 3H), 3.65 (d, <i>J</i> = 11.1 Hz, 3H), 7.31 (brs, 2H), 7.45-7.48 (m, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 21.40, MS (ES ⁺) <i>m/z</i> = 229.3 (MH ⁺).
19j		Colorless oil, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 0.68-0.74 (m, 2H), 0.97-1.03 (m, 2H), 2-2.07 (m, 1H), 3.65 (d, <i>J</i> = 11.1 Hz, 3H), 7.29-7.33 (m, 1H), 7.4-7.49 (m, 3H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 21.05, MS (ES ⁺) <i>m/z</i> = 227 (MH ⁺).
19k		Colorless oil, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 2.37 (s, 3H), 3.67 (d, <i>J</i> = 11.1 Hz, 3H), 7.51-7.72 (m, 3H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 18.44, MS (ES ⁺) <i>m/z</i> = 279/281 (MH ⁺).
19l		Yellow oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.27 (d, <i>J</i> = 6.9 Hz, 6H), 2.95 (quintuplet, <i>J</i> = 6.9 Hz, 1H), 3.77 (d, <i>J</i> = 11.1 Hz, 6H), 7.39-7.46 (m, 2H), 7.56-7.7 (m, 2H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 22.41.
19m		Colorless oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 3.8 (d, <i>J</i> = 11.1 Hz, 6H), 6.98-7.05 (m, 1H), 7.27-7.36 (m, 2H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 17.87 (t, <i>J</i> = 9.96 Hz, 1P), ¹⁹ F NMR (CDCl ₃ , 282.40 MHz) δ -107.27 (m, 2F).
19n		Yellow oil, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 3.65 (d, <i>J</i> = 11.1 Hz, 6H), 3.8 (s, 3H), 7.14-7.31 (m, 3H), 7.45-7.52 (m, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 20.45, MS (ESI, EI ⁺) <i>m/z</i> = 217.2 (MH ⁺)
19p		Oil, ¹ H NMR (<i>d</i> ₆ -DMSO, 300 MHz) δ 2.33 (s, 6H), 3.63 (d, <i>J</i> = 11.1 Hz, 6H), 7.28-7.29 (m, 2H), 7.32-7.34 (m, 1H), ³¹ P NMR (<i>d</i> ₆ -DMSO, 121.49 MHz) δ 21.34, MS (ESI, EI ⁺) <i>m/z</i> = 215 (MH ⁺)

Compound	structure	Description
19q		Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.79 (d, $J = 11.1$ Hz, 6H), 7.25-7.3 (m, 1H), 7.43-7.64 (m, 3H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 20 (d, $J = 8.66$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.4 MHz) δ -130.9 (m, 1F), MS (ESI, EI^+) $m/z = 205$ (MH^+).
19r		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.65 (d, $J = 11.1$ Hz, 6H), 4.56 (s, 2H), 4.61 (s, 2H), 7.28-7.41 (m, 4H), 7.51-7.71 (m, 5H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 20.68; MS (ES^+) $m/z = 307$ (MH^+)
20a		Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 3.71 and 3.74 (2s, 3H), 7.41-7.54 (m, 3H), 7.77-7.84 (m, 2H), 10.70 (brs, 1H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 22.3.
20b		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.76 (d, $J = 11.4$ Hz, 3H), 7.57-7.64 (m, 1H), 7.82-7.85 (m, 1H), 7.99-8.1 (m, 2H), 8.9 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 18.13, MS (ES^+) $m/z = 197.9$ (MH).
20c		Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 2.37 (s, 3H), 3.71 (d, $J = 11.4$ Hz, 3H), 7.32-7.35 (m, 2H), 7.58-7.64 (m, 2H), 10.73 (brs, 1H), ^3MS (ES^+) $m/z = 187$ (MH).
20d		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.71 (d, $J = 11.1$ Hz, 3H), 7.26-7.32 (m, 1H), 7.65-7.95 (m, 4H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.07, MS (ES^+) $m/z = 251/253$ (MH^+).
20e		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.75 (d, $J = 11.4$ Hz, 3H), 5.42 (d, $J = 47.4$ Hz, 2H), 7.4-7.6 (m, 3H), 7.8-7.85 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 21.42, ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -209.80 (t, $J = 47.4$ Hz, 1F), MS (ES^+) $m/z = 205$ (MH).
20f		Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 0.93 (t, $J = 7.5$ Hz, 3H), 1.64 (sextuplet, $J = 7.5$ Hz, 2H), 2.6 (t, $J = 7.5$ Hz, 2H), 3.71 (d, $J = 11.4$ Hz, 3H), 7.34-7.37 (m, 2H), 7.59-7.66 (m, 2H), 9.78 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 22.76, MS (ES^+) $m/z = 215.2$ (MH^+).

Compound	structure	Description
20g		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.36 (s, 3H), 3.53 (d, J = 10.0 Hz, 3H), 7.47-7.64 (m, 3H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 14.79, MS (ES^+) m/z = 263/265 (MH^+).
20h		Yellow pale oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.25 (d, J = 6.9 Hz, 6H), 2.89-2.98 (m, 1H), 3.72 (d, J = 11.4 Hz, 3H), 6.59 (brs, 1H), 7.34-7.43 (m, 2H), 7.59-7.69 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 23.21.
20i		Yellow pale oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.24 (t, J = 7.5 Hz, 3H), 2.67 (q, J = 7.5 Hz, 2H), 3.72 (d, J = 11.4 Hz, 3H), 7.35-7.39 (m, 2H), 7.59-7.67 (m, 3H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 23.07, MS (ES^+) m/z = 201 (MH^+).
20j		Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 3.66 (d, J = 11.4 Hz, 3H), 6.92-6.98 (m, 1H), 7.24-7.31 (m, 2H), 7.64 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 16.92.
20k		Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 2.37 (s, 3H), 3.71 (d, J = 11.4 Hz, 3H), 7.02-7.05 (m, 1H), 7.24-7.32 (m, 1H), 7.36-7.41 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 19.66 (d, J = 9.6 Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -112.7 (q, J = 9.32 Hz, 1F).
20l		Yellow oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.51 (d, J = 11.1 Hz, 3H), 3.79 (s, 3H), 7.12-7.28 (m, 3H), 7.39-7.46 (m, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 17.03, MS (ESI, EI^+) m/z = 203.2 (MH^+).
20m		Pale pink oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.53 (d, J = 11.1 Hz, 3H), 7.37-7.62 (m, 4H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 15.11 (d, J = 8.77 Hz, 1P), ^{19}F NMR (d_6 -DMSO, 235.36 MHz) δ -111.8 (m, 1F), MS (ESI, EI^+) m/z = 191 (MH^+).

Compound	structure	Description
20n		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.31 (s, 6H), 3.50 (d, $J = 11.1$ Hz, 3H), 7.20 (s, 1H), 7.26 (s, 1H), 7.31 (s, 1H), 12.03 (brs, 1H), MS (ESI, EI^+) $m/z = 201$ (MH^+)
20p		Oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.51 (d, $J = 11.4$ Hz, 3H), 4.56 (s, 2H), 4.59 (s, 2H), 7.3-7.71 (m, 9H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 16.22, MS (ES^+) $m/z = 293$ (MH^+).


Compound	structure	Description
22a		White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.87 (d, $J = 11.7$ Hz, 3H), 4.55-4.74 (m, 2H), 6.97 (tt, $J = 2.4$ Hz and 8.7 Hz, 1H), 7.18-7.26 (m, 2H), 7.28-7.3 (m, 1H), 7.32-7.35 (m, 2H), 7.57 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.61 (d, $J = 2.1$ Hz, 1H), 8.33 (d, $J = 6.9$ Hz, 2H), 11.51 (t, $J = 5.7$ Hz, 1H), 12.15 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.42 (t, $J = 8.5$ Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ POSITIF, MS (ES^+) $m/z = 492.28$ (MH^+).
22b		Light orange powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.31 (s, 6H), 3.83 (d, $J = 11.7$ Hz, 3H), 4.6-4.77 (m, 2H), 7.16-7.37 (m, 6H), 7.46-7.49 (m, 1H), 7.66-7.67 (m, 1H), 8.16-8.18 (m, 1H), 8.48 (brs, 1H), 11.31 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.5, MS (ES^+) $m/z = 484.43$ (MH^+).
22c		White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.33 (s, 6H), 3.02 (t, $J = 7.2$ Hz, 2H), 3.77-3.82 (m, 2H), 3.81 (d, $J = 11.7$ Hz, 3H), 7.19-7.21 (m, 3H), 7.26-7.33 (m, 3H), 7.47-7.51 (m, 1H), 7.63-7.64 (m, 1H), 8.04-8.07 (m, 2H), 10.59 (brs, 1H), 11.32-11.35 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.58, MS (ES^+) $m/z = 498.42$ (MH^+).

Compound

structure


Description

22d




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.74 (d, $J = 11.7$ Hz, 3H), 4.57 (dd, $J = 5.1$ and 15.6 Hz, 1H), 4.67 (dd, $J = 5.7$ and 15.6 Hz, 1H), 7.22-7.34 (m, 4H), 7.44-7.49 (m, 1H), 7.57-7.60 (m, 2H), 8.11-8.14 (m, 1H), 8.55-8.57 (m, 1H), 11.37 (t, $J = 5.4$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.78, ^{19}F NMR (d_6 -DMSO, 282.40 MHz) δ -126.51 (m, 1F), MS (ES^+) $m/z = 502.4$ (MH^+).

22e




White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 2.34 (s, 6H), 3.86 (d, $J = 11.4$ Hz, 3H), 4.82-4.85 (m, 2H), 7.18 (brs, 1H), 7.33 (dd, $J = 1.95$ and 8.85 Hz, 1H), 7.36 (brs, 1H), 7.41 (brs, 1H), 7.47 (dd, $J = 1.8$ and 8.85 Hz, 1H), 7.7 (d, $J = 1.8$ Hz, 1H), 8-8.01 (m, 1H), 8.23 (brs, 1H), 8.41-8.42 (m, 1H), 10.45 (brs, 1H), 11.89-11.93 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.03, MS (ES^+) $m/z = 485.4$ (MH^+).

22f



White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 2.33 (s, 6H), 3.87 (d, $J = 11.4$ Hz, 3H), 4.92 (d, $J = 5.4$ Hz, 1H), 7.18 (brs, 1H), 7.32 (dd, $J = 1.8$ and 9 Hz, 1H), 7.38 (brs, 1H), 7.43 (brs, 1H), 7.45 (dd, $J = 2.1$ and 8.4 Hz, 1H), 7.74 (d, $J = 1.8$ Hz, 1H), 8.17 (d, $J = 3.9$ Hz, 1H), 8.4 (d, $J = 3.9$ Hz, 1H), 8.62 (brs, 1H), 10.39 (brs, 1H), 11.85 (t, $J = 5.7$ Hz, 1H), MS (ES^+) $m/z = 485.4$ (MH^+).

22g



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.26 (s, 6H), 3.74 (d, $J = 11.4$ Hz, 3H), 4.59 (dd, $J = 5.1$ and 15.9 Hz, 1H), 4.69 (dd, $J = 5.40$ and 15.9 Hz, 1H), 7.22 (brs, 1H), 7.31-7.34 (m, 3H), 7.59 (dd, $J = 1.8$ and 9 Hz, 2H), 8.34 (d, $J = 1.8$ Hz, 1H), 8.64 (t, $J = 1.8$ Hz, 1H), 9.01 (d, $J = 2.1$ Hz, 1H), 11.34 (t, $J = 5.7$ Hz, 1H), 12.87 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.70, MS (ES^+) $m/z = 485.3$ (MH^+).


Compound	structure	Description
22h		White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 1.20 (t, J = 7.5 Hz, 3H), 2.33 (s, 3H), 2.62 (q, J = 7.5 Hz, 2H), 3.84 (d, J = 11.7 Hz, 3H), 4.65 (dd, J = 5.7 and 15.5 Hz, 1H), 4.73 (dd, J = 5.7 and 15.6 Hz, 1H), 7.19-7.49 (m, 6H), 7.68 (m, 1H), 7.81 (m, 1H), 8.15 (d, J = 6.6 Hz, 1H), 8.42 (s, 1H), 10.70 (brs, 1H), 11.86 (t, J = 5.7 Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.10, MS (ES^+) m/z = 498 (MH^+).
22i		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.27 (s, 6H), 3.76 (d, J = 11.7 Hz, 3H), 4.59-4.74 (m, 2H), 7.24 (brs, 1H), 7.34 (dd, J = 2.1 and 8.7 Hz, 1H), 7.33 (brs, 1H), 7.37 (brs, 1H), 7.53-7.55 (m, 1H), 7.58-7.62 (m, 2H), 8.56 (dd, J = 2.1 and 6.9 Hz, 1H), 9.03-9.04 (m, 1H), 11.45 (t, J = 5.4 Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.67, MS (ES^+) m/z = 485.35 (MH^+).
22j		Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.27 (s, 6H), 3.70 (d, J = 11.7 Hz, 3H), 4.59-4.60 (m, 2H), 7.22 (brs, 1H), 7.28 (brs, 1H), 7.33 (brs, 1H), 7.35-7.47 (m, 4H), 8.14 (d, J = 5.1 Hz, 1H), 8.29 (brs, 1H), 11.73 (t, J = 5.1 Hz, 1H), 13.07 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.03, MS (ES^+) m/z = 502.45 (MH^+).
22k		Off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 2.19 (s, 6H), 3.88 (d, J = 11.4 Hz, 3H), 4.96 (m, 2H), 7.24 (brs, 1H), 7.20-7.49 (m, 6H), 7.75 (m, 1H), 8.15 (d, J = 5.1 Hz, 1H), 9.40 (s, 1H), 10.23 (brs, 1H), 11.79 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 34.2, MS (ES^+) m/z = 485.35 (MH^+).

Compound

structure


Description

22l




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.29 (s, 6H), 3.64 (d, $J = 12$ Hz, 3H), 4.32 (s, 3H), 4.95 (d, $J = 5.4$ Hz, 2H), 7.26 (brs, 1H), 7.35 (dd, $J = 2.1$ and 9 Hz, 1H), 7.38 (brs, 1H), 7.42 (brs, 1H), 7.57 (d, $J = 1.8$ Hz, 1H), 7.62 (dd, $J = 1.8$ and 9 Hz, 1H), 8.08 (d, $J = 6.6$ Hz, 2H), 8.93 (d, $J = 6.6$ Hz, 2H), 11.6 (t, $J = 5.4$ Hz, 1H), 12.88 (d, $J = 3.6$ Hz, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.64, MS (ES^+) $m/z = 482.43$ (M^+).

22m




Off-white solid, ^1H NMR (CDCl_3 , 300 MHz) δ 2.32 (s, 6H), 3.84 (d, $J = 12$ Hz, 3H), 4.68 (t, $J = 5.1$ Hz, 2H), 7.17 (brs, 1H), 7.28-7.38 (m, 6H), 8.26-8.28 (m, 2H), 11.39 (brs, 1H), 12.44 (t, $J = 5.1$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 33.73, MS (ES^+) $m/z = 502$ (MH^+).

22n



Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.25 (s, 6H), 3.74 (d, $J = 11.7$ Hz, 3H), 4.55 (dd, $J = 5.4$ and 15.9 Hz, 1H), 4.65 (dd, $J = 5.4$ and 15.9 Hz, 1H), 7.23 (brs, 1H), 7.28 (brs, 1H), 7.31-7.35 (m, 2H), 7.38-7.4 (m, 2H), 7.59 (dd, $J = 1.8$ and 6.9 Hz, 2H), 8.19 (d, $J = 6.9$ Hz, 1H), 11.36 (t, $J = 5.4$ Hz, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.83, MS (ES^+) $m/z = 484.19$ (MH^+).

22p




Off-white solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.28 (s, 6H), 3.77 (d, $J = 11.7$ Hz, 3H), 4.64-4.74 (m, 2H), 7.24-7.47 (m, 7H), 7.57-7.64 (m, 2H), 8.34-8.36 (m, 1H), 11.32-11.37 (m, 1H), 12.86 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 32.67, MS (ES^+) $m/z = 484.5$ (MH^+).

Compound


structure

Description


22q




22r







22s







22t



Compound	structure	Description
----------	-----------	-------------




Compound	structure	Description
----------	-----------	-------------




Compound	structure	Description
----------	-----------	-------------


22ac




22ad



22ae




22af




Compound	structure	Description
----------	-----------	-------------


22ag




22ah



22ai



22aj




Compound

structure


Description

22ak




Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.4 (s, 3H), 3.86 (d, $J = 12$ Hz, 3H), 4.58 (dd, $J = 5.4$ and 15.6 Hz, 1H), 4.73 (dd, $J = 5.4$ and 15.6 Hz, 1H), 7.33-7.35 (m, 3H), 7.53-7.60 (m, 3H), 7.7-7.8 (m, 2H), 8.27-8.3 (m, 2H), 11.55 (t, $J = 5.4$ Hz, 1H), 11.61 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.62, MS (ES^+) $m/z = 495.4$ (MH^+).


22al



22am




23a








Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 5.03-5.2 (m, 2H), 7.3-7.47 (m, 5H), 7.48-7.63 (m, 3H), 7.64 (d, $J = 565.8$ Hz, 1H), 7.75-7.83 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 25.55, MS (ESI, EI^+) $m/z = 233$ (MH^+).

23b



Colorless oil, ^1H NMR (CDCl_3 , 250 MHz) δ 1.59 (s, 9H), 7.76 (d, $J = 552$ Hz, 1H), 7.5-7.82 (m, 5H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.27, MS (ESI, EI^+) $m/z = 198$ (MH^+).

Compound	structure	Description
23c		Colorless oil, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.38 (s, 3H), 3.68 (d, J = 12 Hz, 3H), 7.49 (d, J = 566 Hz, 1H), 7.4 (dd, J = 3 and 7.8 Hz, 2H), 7.62 (d, J = 13.8 Hz, 1H), 7.64 (d, J = 13.5 Hz, 1H)


Compound	structure	Description
25a		White oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.46 (t, J = 7.2 Hz, 3H), 3.74 (d, J = 14.4 Hz, 3H), 4.46-4.57 (m, 2H), 7.33 (dd, J = 1.95 Hz and 8.7 Hz, 1H), 7.43-7.56 (m, 5H), 7.62-7.67 (m, 1H), 7.93-8.01 (m, 4H), 8.09-8.13 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 73.28, MS (ESI, EI^+) m/z = 534 (MH^+).
25b		White powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.80 (d, J = 14.4 Hz, 3H), 5.92 (brs, 1H), 7.32 (dd, J = 1.95 Hz and 9 Hz, 1H), 7.42-7.58 (m, 4H), 7.8-7.9 (m, 3H), 9.24 (brs, 1H), 10.15 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 80.48, MS (ESI, EI^+) m/z = 365 (MH^+).
25'a		Yellow powder, ^1H NMR (CDCl_3 , 400 MHz) δ 3.86 (d, J = 11.6 Hz, 3H), 7.31 (dd, J = 1.95 Hz and 8.8 Hz, 1H), 7.42-7.47 (m, 3H), 7.52-7.56 (m, 1H), 7.69-7.78 (m, 3H), 8.05 (brs, 1H), 10.63 (brs, 1H), 12.5 (brs, 1H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 34.54, MS (ESI, EI^+) m/z = 365 (MH^+).
25'b		Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 1.45 (t, J = 7.2 Hz, 3H), 4.03-4.13 (m, 1H), 4.25-4.36 (m, 1H), 7.3 (dd, J = 1.95 Hz and 8.85 Hz, 1H), 7.41-7.47 (m, 3H), 7.51-7.56 (m, 1H), 7.72-7.79 (m, 3H), 8.01 (brs, 1H), 10.59 (brs, 1H), 12.59 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 32.32, MS (ESI, EI^+) m/z = 379 (MH^+).

Compound

structure


Description

26a




White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 1.18 (t, $J = 7.2$ Hz, 3H), 3.78 (q, $J = 7.2$ Hz, 2H), 7.33 (dd, $J = 2.1$ and 9 Hz, 1H), 7.45-7.75 (m, 14H), 7.93-7.97 (m, 1H), 8-8.04 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 22.23, MS (ESI, EI^+) $m/z = 564$ (MH^+).

27a




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 6.13 (d, $J = 1.8$ Hz, 1H), 7.22 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.53-7.72 (m, 11H), 7.92 (brs, 1H), 10.37 (brs, 1H), 12.84 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 26.61, MS (ESI, EI^+) $m/z = 395$ (MH^+).

28a




Yellow thick oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, $J = 7.2$ Hz, 3H), 4.2-4.41 (m, 2H), 7.32-7.35 (dd, $J = 2.1$ Hz and 9 Hz, 1H), 7.47-7.65 (m, 6H), 7.74-7.81 (m, 2H), 7.88-7.93 (m, 2H), 8-8.03 (m, 2H), ^{31}P NMR (CDCl_3 , 101.256 MHz) δ 25.04, MS (ESI, EI^+) $m/z = 502$ (MH^+).

29a



White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.24 (d, $J = 13.5$ Hz, 3H), 7.1 (m, 1H), 7.25 (dd, $J = 1.2$ and 8.7 Hz, 1H), 7.51-7.64 (m, 4H), 7.74-7.8 (m, 2H), 7.83-7.86 (m, 1H), 10.53 (brs, 1H), 12.62 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 30.62, MS (ESI, EI^+) $m/z = 333$ (MH^+).

30a




Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.44 (t, $J = 6.5$ Hz, 3H), 2.66 (d, $J = 11.1$ Hz, 6H), 4.48 (q, $J = 6.7$ Hz, 2H), 7.34 (d, $J = 9.9$ Hz, 1H), 7.45-7.52 (m, 5H), 7.60 (m, 1H), 7.85-7.95 (m, 4H), 8.07-8.09 (m, 2H), ^{31}P NMR (CDCl_3 , 101 MHz) δ 25.11, MS (ESI, EI^+) $m/z = 531$ (MH^+).

Compound

structure


Description

31a




Pale orange powder, ^1H NMR ($\text{d}_6\text{-DMSO}$, 300 MHz) δ 2.64 (d, $J = 11.7$ Hz, 6H), 7.32 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.5-7.59 (m, 4H), 7.72-7.79 (m, 3H), 7.86 (brs, 1H), 10.85 (brs, 1H), 12.61 (brs, 1H), MS (ESI, EI^+) $m/z = 384$ ($\text{M}+\text{Na}$).

32a




Slight yellow solid; ^1H NMR (CDCl_3 , 300 MHz) δ 1.38-1.49 (m, 6H), 4.1-4.24 (m, 2H), 4.55 (q, $J = 7.2$ Hz, 2H), 5.97 (d, $J = 16.8$ Hz, 1H), 7.16-7.68 (m, 7H), 7.86-8.12 (m, 6H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 22.82, MS (ES^+) $m/z = 582.88$ (MH^+).

32b




Slight yellow solid; ^1H NMR (CDCl_3 , 300 MHz) δ 1.38-1.49 (m, 6H), 4.1-4.24 (m, 2H), 4.55 (q, $J = 7.2$ Hz, 2H), 5.56 (d, $J = 12$ Hz, 1H), 7.16-7.68 (m, 7H), 7.86-8.12 (m, 6H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 22.82, MS (ES^+) $m/z = 582.88$ (MH^+).

33a



White solid, ^1H NMR (CDCl_3 , 300 MHz) δ 3.90 (d, $J = 11.7$ Hz, 3H), 5.80 (brs, 1H), 5.93 (d, $J = 16.5$ Hz, 1H), 7.39-7.92 (m, 5.93, 8H), 10.16 (brs, 1H), 10.85 (brs, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 31.53, MS (ES^+) $m/z = 400$ (MH^+).

38a




Compound


structure

Description


38b




38c




38d



38e



38f




Compound


structure

Description


38g




38h




38i









38j



38k



Compound	structure	Description
43a		
43b		
43c		
47a		Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 6.9$ Hz, 6H), 4.20-4.10 (m, 4H), 9.01 (d, $J = 6.6$ Hz, 2H), 9.31 (d, $J = 3.3$ Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 12.37, MS (ESI, EI^+) $m/z = 217$ (MH^+).
47b		Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.2 (t, $J = 7.2$ Hz, 6H), 3.95-4.1 (m, 4H), 7.25-7.3 (m, 1H), 7.93-8 (m, 1H), 8.62-8.64 (m, 1H), 8.83-8.85 (m, 1H) ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 15.72.
47c		Yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.36 (t, $J = 7.2$ Hz, 6H), 4.16-4.31 (m, 4H), 7.41-7.46 (m, 1H), 7.77-7.85 (m, 1H), 7.96-8.01 (m, 1H), 8.8-8.82 (m, 1H) ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 7.16, MS (ESI, EI^+) $m/z = 216$ (MH^+).
47d		Brown oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.27 (t, $J = 6.9$ Hz, 6H), 4.01-4.16 (m, 4H), 7.56-7.62 (m, 2H), 8.67-8.71 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 14.63, MS


Compound	structure	Description
		(ESI, EI ⁺) m/z = 216 (MH ⁺).
47e		Oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.33 (t, J = 7.2 Hz, 6H), 4.04-4.19 (m, 4H), 7.32-7.35 (m, 1H), 7.41-7.45 (m, 1H), 7.97-8.01 (m, 1H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 13.35, MS (ESI, EI ⁺) m/z = 221.11 (MH ⁺).
47f		Yellow pale oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.34 (t, J = 7.05 Hz, 6H), 4.06-4.22 (m, 4H), 7.16-7.20 (m, 1H), 7.65-7.72 (m, 2H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 12.05, MS (ESI, EI ⁺) m/z = 221.11 (MH ⁺).
48a		Yellow oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.27 (t, J = 7.05 Hz, 3H), 4.01-4.07 (m, 2H), 7.31-7.4 (m, 2H), 7.91-7.95 (m, 1H), 12.73 (s, 1H), ³¹ P NMR (CDCl ₃ , 101.256 MHz) δ 14.67.
48b		Yellow oil, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.31 (t, J = 7.05 Hz, 3H), 4.04-4.14 (m, 2H), 7.12-7.16 (m, 1H), 7.61-7.68 (m, 2H), 9.12 (brs, 1H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 14.7.
50a		Yellowish powder, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.41 (t, J = 7.05 Hz, 3H), 1.49 (t, J = 7.05 Hz, 3H), 4.13-4.28 (m, 2H), 4.57 (qd, J = 1.8 Hz and 7.05 Hz, 2H), 7.16-7.19 (m, 1H), 7.36 (dd, J = 2.4 Hz and 9 Hz, 1H), 7.50-7.56 (m, 2H), 7.61-7.73 (m, 3H), 7.9-7.96 (m, 2H), 8.1-8.14 (m, 2H), MS (ESI, EI ⁺) m/z = 537 (MH ⁺).
50b		Yellowish powder, ¹ H NMR (CDCl ₃ , 300 MHz) δ 1.38 (t, J = 7.05 Hz, 3H), 1.42 (t, J = 7.05 Hz, 3H), 4.08-4.26 (m, 2H), 4.52 (q, J = 7.05 Hz, 2H), 7.34 (dd, J = 2.1 Hz and 9 Hz, 1H), 7.35-7.41 (m, 1H), 7.47-7.53 (m, 2H), 7.58-7.64 (m, 1H), 7.77-7.84 (m, 1H), 7.88-7.92 (m, 1H), 8.03-8.15 (m, 4H), 8.72-8.74 (m, 1H), ³¹ P NMR (CDCl ₃ , 121.49 MHz) δ 19.72.

Compound

structure


Description

50c




Oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.37 (t, $J = 7.05$ Hz, 3H), 1.45 (t, $J = 7.2$ Hz, 3H), 4.05-4.22 (m, 2H), 4.52 (q, $J = 7.2$ Hz, 2H), 7.34 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.39 (dd, $J = 2.1$ and 3.3 Hz, 2H), 7.48-7.54 (m, 2H), 7.6-7.65 (m, 1H), 7.88 (d, $J = 1.8$ Hz, 1H), 7.92 (dd, $J = 1.8$ and 9 Hz, 1H), 8.02-8.1 (m, 3H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 18.36.

51




White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.33 (t, $J = 7.05$ Hz, 3H), 4.03-4.09 (m, 1H), 4.15-4.24 (m, 1H), 7.21-7.23 (m, 1H), 7.34 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.55-7.61 (m, 2H), 7.7 (d, $J = 2.1$ Hz, 1H), 8-8.03 (m, 2H), 10.12 (brs, 1H), 12.76 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 23.46, MS (ES^+) $m/z = 368.8$ (MH^+).

52a




Beige powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.83 (d, $J = 12$ Hz, 3H), 5.93 (brs, 1H), 7.3-7.34 (m, 2H), 7.39-7.42 (m, 1H), 7.47 (dd, $J = 1.8$ Hz and 8.7 Hz, 1H), 7.7 (d, $J = 1.8$ Hz, 1H), 7.91 (ddd, $J = 1.2$ Hz and 2.7 Hz and 8.1 Hz, 1H), 10.44 (brs, 1H), 10.95 (brs, 1H), MS (ES^-) $m/z = 353.06$ (MH^-).

52b



White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.77 (d, $J = 12$ Hz, 3H), 7.22-7.24 (m, 1H), 7.34 (dd, $J = 1.8$ and 8.7 Hz, 1H), 7.56-7.61 (m, 2H), 7.66 (d, $J = 1.8$ Hz, 1H), 8.01-8.05 (m, 2H), 10.1 (brs, 1H), 12.81 (brs, 1H), MS (ES^+) $m/z = 354.7$ (MH^+).

53a



Colorless oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.66 (d, $J = 3.9$ Hz, 3H), 1.76 (d, $J = 5.4$ Hz, 3H), 2.57 (dd, $J = 7.65$ and 21.75 Hz, 2H), 3.74 (d, $J = 10.5$ Hz, 6H), 5.12-5.21 (m, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 7.32, MS (ESI , EI^+) $m/z = 179$ (MH^+).


Compound	structure	Description
56a		Thick yellow oil, ^1H NMR (CDCl_3 , 300 MHz) δ 1.24 (d, J = 3.9 Hz, 3H), 1.5 (t, J = 7.2 Hz, 3H), 1.54 (d, J = 6 Hz, 3H), 2.78 (dd, J = 8.1 and 19.5 Hz, 2H), 3.7 (d, J = 11.1 Hz, 3H), 4.57 (q, J = 7.2 Hz, 2H), 5.07-5.1 (m, 1H), 7.37 (dd, J = 2.1 and 8.7 Hz, 1H), 7.5-7.55 (m, 2H), 7.63-7.67 (m, 1H), 7.87 (d, J = 1.8 Hz, 1H), 7.96 (dd, J = 1.2 and 9 Hz, 1H), 8.1-8.13 (m, 2H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 8.68, MS (ESI, EI^+) m/z = 511 (MH^+).
57a		White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.03-1.38 (m, 4H), 1.49-1.75 (m, 4H), 1.87-1.98 (m, 2H), 3.59 (d, J = 11.1 Hz, 3H), 7.33 (dd, J = 1.95 and 8.7 Hz, 1H), 7.59 (d, J = 8.7 Hz, 1H), 7.67 (brs, 1H), 7.87 (brs, 1H), 10.23 (brs, 1H), 12.69 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 48.93, MS (ESI, EI^+) m/z = 355 (MH^+).
57b		Pale yellow solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.17 (d, J = 3.6 Hz, 3H), 1.55 (d, J = 5.7 Hz, 3H), 2.81 (m, 2H), 3.62 (d, J = 11.1 Hz, 3H), 5.0 (m, 1H), 7.33 (dd, J = 2.0 and 8.7 Hz, 1H), 7.57 (d, J = 8.7 Hz, 1H), 7.68 (d, J = 2.0 Hz, 1H), 7.85 (brs, 1H), 7.87 (brs, 1H), 10.10 (brs, 1H), 12.64 (brs, 1H), MS (ESI, EI^+) m/z = 341 (MH^+).
59a		Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 3.88 (d, J = 12 Hz, 3H), 4.4 (d, J = 5.4 Hz, 2H), 6.98 (tt, J = 2.4 Hz and 8.7 Hz, 1H), 7.26-7.31 (m, 2H), 7.38 (dd, J = 1.8 Hz and 8.7 Hz, 1H), 7.59 (d, J = 1.5 Hz, 1H), 7.46 (dd, J = 1.8 Hz and 8.7 Hz, 1H), 11.24 (brs, 1H), 11.79 (t, J = 5.4 Hz, 1H), ^{31}P NMR (CDCl_3 , 121.49 MHz) δ 30.82 (t, J = 8.6 Hz, 1P), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -106.02 (m, 2F), MS (ES^+) m/z = 424.04 (MH^+).

Compound

structure


Description

59b




Yellow powder, ^1H NMR (CDCl_3 , 300 MHz) δ 2.8 (t, J = 6.9 Hz, 2H), 3.79-3.85 (m, 2H), 3.87 (d, J = 11.7 Hz, 3H), 6.96 (tt, J = 2.4 Hz and 8.7 Hz, 1H), 7.25-7.33 (m, 2H), 7.36 (d, J = 1.8 Hz, 1H), 7.53 (dd, J = 1.8 Hz and 8.7 Hz, 1H), 7.60 (d, J = 1.8 Hz, 1H), 11.13 (brs, 1H), 11.46 (t, J = 5.4 Hz, 1H), ^{19}F NMR (CDCl_3 , 282.40 MHz) δ -106.25 (m, 2F), MS (ES^+) m/z = 437.96 (MH^+).

60a




White solid, ^1H NMR (d_6 -DMSO, 300 MHz) δ 7.14-7.17 (m, 1H), 7.32-7.73 (m, 7H), 8.16 (brs, 1H), 11.78 (brs, 1H), 12 (brs, 1H), MS (ESI , EI^+) m/z 335 (MH^+).

60b




White powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 3.72 (s, 3H), 6.9-6.93 (m, 2H), 7.19 (dd, J = 1.5 Hz and 8.7 Hz, 1H), 7.43-7.46 (m, 1H), 7.58-7.69 (m, 3H), 8-8.03 (m, 1H), 11.25 (brs, 1H), 12.15 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 121.49 MHz) δ 31.44, MS (ESI , EI^+) m/z = 365 (MH^+).

60d




Off white powder, ^1H NMR (d_6 -DMSO, 400 MHz) δ 2.3 (s, 3H), 7.26 (dd, J = 2 and 8.8 Hz, 1H), 7.3-7.37 (m, 2H), 7.46-7.54 (m, 3H), 7.82 (brs, 1H), 7.88 (brs, 1H), 10.69 (brs, 1H), 12.49 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101.256 MHz) δ 23.81, MS (ESI , EI^+) m/z = 349 (MH^+).

60f




Off white powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 2.21 (s, 6H), 7.2-7.28 (m, 2H), 7.37-7.53 (m, 3H), 7.79 (brs, 1H), 7.87 (brs, 1H), 10.68 (brs, 1H), 12.47 (brs, 1H), MS (ESI , EI^+) m/z = 363 (MH^+).

Compound	structure	Description
60g	 <chem>Nc1cc(O=P(c2cc(Cl)ccc2n1)c3cc(F)cc(F)c3)cc(F)c3cc(F)cc(F)c3</chem>	Orange powder, ^1H NMR (d_6 -DMSO, 300 MHz) δ 7.27 (dd, $J = 1.95$ and 8.87 Hz, 1H), 7.48-7.54 (m, 3H), 7.61-7.71 (m, 1H), 7.85 (d, $J = 1.8$ Hz, 1H), 7.9 (brs, 1H), 10.54 (brs, 1H), 12.53 (brs, 1H), ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 20.39, ^{19}F NMR (d_6 -DMSO, 282.4 MHz) δ -133.8 (m, 1F), -137.3 (m, 1F), MS (ESI, EI^+) $m/z = 371$ (MH^+).

5 The following examples are provided to illustrate the present invention, and are in no way intended to limit the scope of the invention.

EXAMPLES

X. Compound synthesis

General Synthetic Method

Compound	FW	Equivalent
CHCl ₃ (or CH ₂ Cl ₂)		75ml/mmol
mCPBA (70% in water)	172.57	2.5


Experimental :

The pyridine precursor was dissolved in chloroform (or in CH₂Cl₂) under stirring at room temperature; m-chloroperoxybenzoic acid was added and the reaction allowed to stir overnight (~15H).


The mixture was diluted with dichloromethane and extracted with a mixture of saturated K₂CO₃ / H₂O (1/3).

The aqueous layer was extracted three times with dichloromethane. The combined organic layer was dried over sodium sulphate, filtered and concentrated under reduced pressure. (TLC: dichloromethane/methanol = 9/1). The crude product was then purified by chromatography.


Compounds synthesised:




Compound A




Compound B




Compound C




Compound D




Compound E




Compound F




Compound G




Compound H




Compound I




Compound J




Compound K




Compound L



Compound M



Compound N




Compound P

Compound O	
------------	--


Example 1. Ethyl 3-bromo-5-chloro-1-(phenylsulfonyl)-1H-indole-2-carboxylate

a) Synthesis of ethyl 5-chloro-1-(phenylsulfonyl)-1H-indole-2-carboxylate as starting material (based on work of Silvestri R., De Martino G., La Regina G., Artico M., Massa S., Vargiu L., Mura M., Loi A.-G., Marceddu T., La Colla P. *J. Med. Chem.* **2003**, 46:2482-2493):



To a stirred and cooled (0°C) solution of ethyl-5-chloroindole-2-carboxylate (1.052g, 4.70 mmol) in DMF (25 mL) under N₂, was added NaH (60% in oil, 230 mg, 5.64 mmol) portionwise. After the end of gas evolution, phenylsulfonyl chloride (0.72 mL, 5.64 mmol) was added. The reaction mixture was stirred for 1 h (TLC monitoring, eluant dichloromethane). A little amount of water was added carefully and DMF was evaporated. Crude residue was dissolved in EtOAc and washed with water and brine. After drying and evaporation of solvents the compound was purified by chromatography on silica gel (eluant: cyclohexane/EtOAc 9/1 to 7/3) to afford protected indole (1.547 g, 90% yield). Off-white solid; ¹H NMR (*d*₆-DMSO) δ 1.30 (t, *J* = 7.2 Hz, 3H), 4.35 (q, *J* = 7.2 Hz, 2H), 7.37 (s, 1H), 7.53 (dd, *J* = 2.2 and 9.1 Hz), 7.62-7.77 (m, 3H), 7.80 (d, *J* = 2.2 Hz, 1H), 7.99 (m, 2H), 8.06 (d, *J* = 9.1 Hz); MS (ESI, EI⁺) *m/z* = 364 (MH⁺).

b) Synthesis of ethyl 3-bromo-5-chloro-1-(phenylsulfonyl)-1H-indole-2-carboxylate intermediate:




To a stirred solution of ethyl 5-chloro-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate (4.83 g, 13.27 mmol) in DMF (40 mL) under N₂, was added a solution of bromine (1.3 mL, 26.54 mmol) in DMF (10 mL). Reaction media was stirred at RT 4H, water was added (150 mL) and was extracted with dichloromethane (3×100 mL). Organic layer was washed with a saturated solution of Na₂SO₅, dried and evaporated to give a crude yellow oil. Purification by chromatography on silica gel (eluant: cyclohexane/EtOAc 9/1) afforded 3-brominated indole (5.548 g, 93% yield). Off white solid; ¹H NMR (*d*₆-DMSO) δ 1.37 (t, *J* = 7.2 Hz, 3H), 4.48 (q, *J* = 7.2 Hz, 2H), 7.59-7.68 (m, 4H), 7.77 (m, 1H), 7.96-8.09 (m, 3H); MS (ESI, EI⁺) *m/z* = 442-444 (MH⁺).

Method A: Typical procedure for the synthesis of Ethyl 5-chloro-3-(dialkoxyphosphoryl)-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate and Ethyl 5-chloro-3-[alkoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate

To a stirred and cooled (-90°C) solution of ethyl 3-bromo-5-chloro-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate (0.50 mmol) in anhydrous THF (2.5 mL) under N₂, was added *n*-BuLi (2.5M in hexanes, 0.24 mL, 0.60 mmol) dropwise. After 5 min at -90°C, appropriate chlorophosphorus reagent (0.60 mmol) was added dropwise at the same temperature. The reaction was allowed to warm up to RT over 3 H (TLC monitoring, eluant dichloromethane/ EtOAc 9/1). Water was then added (5 mL). Extraction with EtOAc (3× 20 mL) drying and evaporation led to a crude oil that was purified by chromatography on silica gel.

Example 2. Ethyl 5-chloro-3-(diethoxyphosphoryl)-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate




Method A: Purification by chromatography on silica gel (eluant: dichloromethane/EtOAc 9/1 to 8/2) afforded desired indole (176 mg, 71% yield). White solid; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.19 (t, $J = 7.1$ Hz, 6H), 1.39 (t, $J = 7.1$ Hz, 3H), 3.99-4.09 (m, 4H), 4.46 (q, $J = 7.1$ Hz, 2H), 7.77 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.67-7.82 (m, 4H), 8.07-8.12 (m, 2H); ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 9.7; MS (ESI, EI^+) $m/z = 500$ (MH^+).

Example 3. Ethyl 5-chloro-3-[ethoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1H-indole-2-carboxylate


a) Synthesis of ethyl hydrogen phenylphosphonochloridate intermediate was synthesized according to Smith A.B. III, Ducry L., Corbett, R.M., Hirschmann R. *Org. Lett.* **2000**, 2:3887-3890:

i) Synthesis of diethyl phenylphosphonate:




Colorless oil; ^1H NMR (CDCl_3 , 250 MHz) δ 1.33 (t, $J = 7.1$ Hz, 6H), 4.05-4.25 (m, 4H), 7.46-7.57 (m, 3H), 7.78-7.87 (m, 2H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 19.3.

ii) Synthesis of ethyl hydrogen phenylphosphonate:




Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 1.32 (t, $J = 7.3$ Hz, 3H), 4.08 (q, $J = 7.3$ Hz, 2H), 7.42-7.56 (m, 3H), 7.79-7.86 (m, 2H), 10.67 (brs, 1H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 21.3;

iii) Synthesis of ethyl hydrogen phenylphosphonochloridate having the following physical characteristics:



^{31}P NMR (CDCl_3 , 101 MHz) δ 10.20 and 10.24.


b) Synthesis of final product Ethyl 5-chloro-3-[ethoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate:



Method A: Purification by chromatography on silica gel (eluant: dichloromethane/EtOAc 9/1) afforded debrominated indole (318 mg) and then desired indole (326 mg, 41% yield). Colorless oil; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.27 (t, $J = 7.1$ Hz, 3H), 1.36 (t, $J = 7.1$ Hz, 3H), 4.03 (m, 2H), 4.38 (q, $J = 7.1$ Hz, 2H), 7.51-7.83 (m, 11H), 8.05-8.11 (m, 3H); ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 23.3; MS (ESI, EI^+) $m/z = 532$ (MH^+).

Example 4. Ethyl 5-chloro-3-[ethoxy(3,5-dimethylphenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate


a) Synthesis of diethyl 3,5-dimethylphenylphosphonate intermediate according to method described by Hirao T., Masunaga T., Oshiro Y., Agawa T. *Synthesis* **1981**, 56-57.



Purification by chromatography on silica gel (eluant: cyclohexane/EtOAc 6/4) afforded product (1.625 g, 61% yield) as a colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 1.33 (t, $J = 7.0$ Hz, 6H), 2.35 (s, 6H), 4.02-4.18 (m, 4H), 7.18 (s, 1H), 7.40 (s, 1H), 7.45 (s, 1H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 20.3.

Other intermediates are synthesized according to Example 3 (i, ii and iii).


b) Synthesis of Ethyl 5-chloro-3-[ethoxy(3,5-dimethylphenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate:



Method A: Purification by chromatography on silica gel (eluant: dichloromethane/EtOAc 95/5) afforded product (750 mg, 56% yield) as a light yellow solid. ^1H NMR (d_6 -DMSO, 300 MHz) δ 1,27 (t, $J = 7.1$ Hz, 3H), 1,36 (t, $J = 7.1$ Hz, 3H), 2,30 (s, 6H), 3,94-4,06 (m, 2H), 4,44 (q, $J = 7.1$ Hz, 2H), 7,25 (s, 1H), 7,39 (s, 1H), 7,42 (s, 1H), 7,53 (dd, $J = 2.1$ and 9.0 Hz, 1H), 7,65-7,71 (m, 2H), 7,77-7,82 (m, 2H), 8,05-8,11 (m, 3H); ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 23.6; MS (ESI, EI^+) $m/z = 560$ (MH^+).

Example 5. Ethyl 5-chloro-3-[methoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate

a) synthesis of dimethyl phenylphosphonate:




To a solution of benzenephosphonyl dichloride (1 mL, 6.35 mmol) in anhydrous dichloromethane (25 mL) was added dropwise at 0°C ethanol (1,12 mL, 19.04 mmol) following by triethylamine (2,65 mL, 19.04 mmol). The reaction mixture was stirred at RT 2h. The reaction mixture was washed with a solution of HCl 1N (50 mL). Aqueous layer was extracted with dichloromethane. Combined organic phases were dried and concentrated under reduce pressure. The crude oil was purified by chromatography on silica gel (Eluant: cyclohexane/EtOAc 6/4) to afford dimethyl phenylphosphonate as

colorless oil (1,110g, 82% yield). ^1H NMR (CDCl_3 , 300 MHz) δ 3.76 (d, $J = 11.1$ Hz, 3H), 7.44-7.58 (m, 3H), 7.76-7.84 (m, 2H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 22.2.


Other intermediates are synthesised according to example 3 (i, ii and iii).

b) synthesis of final product Ethyl 5-chloro-3-[methoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate:



Method A: Colorless oil; ^1H NMR (CDCl_3 , 300 MHz) δ 1.45 (t, $J = 7.2$ Hz, 3H), 3.80 (d, $J = 11.4$ Hz, 3H), 4.54 (q, $J = 7.2$ Hz, 2H), 7.36 (dd, $J = 2.1$ and 9.0 Hz, 1H), 7.47-7.67 (m, 6H), 7.84-7.96 (m, 4H), 8.09-8.12 (m, 2H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 26.7; MS (ESI, EI^+) $m/z = 518$ (MH^+).


Example 6. Ethyl 2-(aminocarbonyl)-5-chloro-1*H*-indol-3-yl-(phenyl)phosphinate



Ethyl-5-chloro-3-[ethoxy(phenyl)phosphoryl]-1-(phenylsulfonyl)-1*H*-indole-2-carboxylate (268 mg, 0.50 mmol) was dissolved in a saturated solution of ammonia in methanol (5 mL) in a pressure tube. The tube was heated under microwave irradiation under pressure at 65°C (Maximum power input 100W, CEM discover apparatus) for 2 H. After evaporation of solvents, purification by chromatography on silica gel (eluant: dichloromethane/MeOH 95/5 to 9/1) afforded desired carboxamide indole (107 mg, 81%


yield). White solid; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.34 (t, $J = 7.1$ Hz, 3H), 4.05 (m, 1H), 4.20 (m, 1H), 7.32 (dd, $J = 2.1$ and 8.7 Hz, 1H), 7.49-7.61 (m, 5H), 7.68-7.75 (m, 2H), 8.02 (brs, 1H), 10.27 (brs, 1H), 12.77 (brs, 1H); ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 31.1; MS (ESI, EI^+) $m/z = 363$ (MH^+).

Example 7. Ethyl 2-(aminocarbonyl)-5-chloro-1H-indol-3-yl-(3,5-dimethylphenyl)phosphonate



Same procedure as described in example 5. white solid; ^1H NMR (d_6 -DMSO, 300 MHz) δ 1.32 (t, $J = 7.0$ Hz, 3H), 2.26 (s, 6H), 3.90-4.03 (m, 1H), 4.09-4.22 (m, 1H), 7.21 (s, 1H), 7.29-7.33 (m, 3H), 7.57 (dd, $J = 1.8$ and 9.0 Hz, 1H), 7.60 (dd, $J = 1.8$ Hz, 1H), 7.99 (brs, 1H), 10.3 (brs, 1H), 12.7 (brs, 1H); ^{31}P NMR (d_6 -DMSO, 101 MHz) δ 31.3; MS (ESI, EI^+) $m/z = 391$ (MH^+).

Example 8. Methyl 2-(aminocarbonyl)-5-chloro-1H-indol-3-yl-(phenyl)phosphinate



Same procedure as described in example 5. Pale yellow powder; ^1H NMR (CDCl_3 , 300 MHz) δ 3.85 (d, $J = 11.4$ Hz, 3H), 6.08 (broad s, 1H), 7.30 (dd, $J = 2.0$ and 9.0 Hz, 1H), 7.36-7.56 (m, 4H), 7.68 (d, $J = 1.8$ Hz, 1H), 7.73-7.81 (m, 2H), 10.78 (broad s, 1H), 10.03 (broad s, 1H); ^{31}P NMR (CDCl_3 , 101 MHz) δ 33.3; MS (ESI, EI^+) $m/z = 349$ (MH^+).

Example 9: Biological Activity Against Drug Resistant Strains of HIV

In one embodiment, the efficacy of an anti-HIV compound is measured *in vitro* by a rapid, sensitive, and automated assay that involves the reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT). An HIV-transformed cell line that is highly permissive and selective for HIV infection, such as, for example, the T-4 cell line, MT-4, is chosen as the target cell line (Koyanagi et al., *Int. J. Cancer*, **1985**, 36:445-451). *In situ* reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) as assessed spectrophotometrically is the standard by which the viability of both mock-infected cells and HIV-infected cells is measured. Inhibition of the HIV-induced cytopathic effect serves as the end-point. A 50% cytotoxic concentration (CC₅₀ in μM) is defined as the concentration of compound that reduces the absorbance of the mock-infected control sample by 50%. The percent efficacy of an anti-HIV compound is calculated by the formula (expressed as a %):

$$(\text{OD}_{\text{HIV test compound}}) - (\text{OD}_{\text{control}}) / (\text{OD}_{\text{mock infected cells}}) - (\text{OD}_{\text{control}})$$

Here, (OD_{HIV test compound}) is the optical density measured for a specific amount of a test compound in HIV-infected cells; (OD_{control}) is the optical density measured for untreated HIV-infected, control cells; and (OD_{mock infected cells}) is the optical density measured for control, mock-infected cells that are untreated. Optical density values typically are assessed at 540 nm. The dosage of an anti-HIV test compound that provides 50% protection according to the preceding formula is defined as the 50% inhibitory concentration (IC₅₀ in μM). The selectivity index (SI) is defined as the ratio of the CC₅₀ to the IC₅₀.

In another embodiment, the p24 ELISA assay is used to determine the efficacy of an anti-HIV compound. This viral replication immunoassay measures the amount of p24 viral capsid (core) antigen present, and is available commercially from sources such as, for example, Coulter Corporation/Immunotech, Inc.[®] (Westbrook, MI).

Still other embodiments include a reverse transcriptase assay in which the amount of viral replication is measured by utilizing a homopolymer poly rA:oligo dT template primer system that quantifies the incorporation into cells of tritiated thymidine monophosphate by scintillation counting methods (Southern Research Institute, University of Alabama, Birmingham, AL); a syncytial inhibition assay that employs

CEM-SS, HeLa-CD4, or HeLa-CD4-LTR-b-galactosidase cells having an immunofluorescent, chemiluminescent, or colorimetric endpoint; and an attachment-and fusion-inhibition assay that utilizes indicator cell lines and quantitation by chemiluminescent, colorimetric or microscopic evaluation (Southern Research Institute, University of Alabama, Birmingham, AL).

In one embodiment the indole compounds of the present invention do not exhibit cross resistance with other non-nucleoside reverse transcriptase inhibitors (NNRTIs), in that the compounds of the present invention display an EC_{50} (in molar concentration) in a mutant HIV strain of less than approximately 50, 25, 10 or 1 μ M concentration. In a typical embodiment, the NNRTIs display an EC_{50} in a mutant HIV strain of less than approximately 5, 2.5, 1 or 0.1 μ M concentration. The degree of cross-resistance against a drug resistant strain of HIV is measured by assessing the EC_{50} of the desired oxopyrimidine compound in the target mutated, i.e., drug resistant, virus.

Therefore, in another important embodiment of this invention, a method for treating a patient with a cross-resistant HIV is provided that includes administering an effective HIV-treatment amount of an indole compound, a salt, prodrug, stereoisomer or tautomer thereof.

Biological Activity Against Drug Resistant Strains of HIV

In one embodiment the phenylindoles of the present invention do not exhibit cross resistance with other non-nucleosides reverse transcriptase inhibitors (NNRTI), in that it displays an EC_{50} (in molar concentrations) in a mutant HIV strain of less than approximately 50, 25, 10 or 1 micromolar concentration. In a typical embodiment, the non-nucleosides reverse transcriptase inhibitors (NNRTI) displays an EC_{50} (in molar concentrations) in a mutant HIV strain of less than approximately 5, 2.5, 1 or 0.1 micromolar concentration. The degree of cross-resistance against a drug resistant strain of HIV can easily be measured by assessing the EC_{50} of the desired indole in the target mutated i.e., drug resistant, virus.

Therefore, in another important embodiment of this invention, a method for treating a patient with a cross-resistant HIV is provided that includes administering an effective HIV-treatment amount of a phenylindole or its prodrug or salt.




TABLE 3 - ENZYME ASSAY DATA

Compound Number	Indole Substituent Position							IC ₅₀ WT (μM)	IC ₅₀ Y181C (μM)	IC ₅₀ K103N (μM)	IC ₅₀ K103N/ Y181C (μM)
	1	2	3	4	5	6	7				
7ah	H	CONH2	P(O)(Ph)OMe	H	Cl	H	H	0.309	0.349	0.389	6.75
6a	H	CONH2	P(O)(Ph)OEt	H	Cl	H	H	0.722	1.968	1.634	>5
6b	H	CONH2	P(O)(Ph:3,5-di-Me)Oet	H	Cl	H	H	1.641	4.311	2.607	>5
18ar	H	CONHCH2(4-pyridine)	P(O)(Ph)OMe	H	H	H	H	0.156	0.489	0.93	1.662

9a	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3,5-di-Me)OEt	H	Cl	H	H	3,413	4.489	3.356	>5
27a	H	CONH ₂	P(O)Ph ₂	H	Cl	H	H	>5	>5	>5	>5
7ai	H	CONH ₂	P(O)(Ph)OMe	H	Cl	H	H	0,393	5.606	6.587	>5
60a	H	CONH ₂	P(O)(Ph)OH	H	Cl	H	H	>5	>5	>5	>5
29a	H	CONH ₂	P(O)(Ph)CH ₃	H	Cl	H	H	0,233	2.524	>5	>5
31a	H	CONH ₂	P(O)(Ph)N(Me) ₂	H	Cl	H	H	1,401	>5	>5	>5
6c	H	CONH ₂	P(O)(Ph:4-F)OEt	H	Cl	H	H	3,62	>5	>5	>5
6ac	H	CONH ₂	P(O)(Ph:3,5-di-F)OEt	H	Cl	H	H	0.726	3.5	12.72	>5
6ad	H	CONH ₂	P(O)(Ph:2-F)OEt	H	Cl	H	H	0,433	4.346	3.293	>5

TABLE 4 - CELL-BASED ASSAY DATA USING BH-10 CELLS

Compound No.	Indole Substituent Position							EC ₅₀	EC ₅₀	EC ₅₀
	1	2	3	4	5	6	7	WT//B (μM)	Y181C (μM)	K103N/Y181C (μM)
	1									
7ah	H	CONH2	P(O)(Ph)OMe	H	Cl	H	H	0,0004	0,0049	0,0034
6a	H	CONH2	P(O)(Ph)OEt	H	Cl	H	H	0,0012	0,0137	0,0164
6b	H	CONH2	P(O)(Ph:3,5-di-Me)OEt	H	Cl	H	H	0,003	0,028	0,0517
18ar	H	CONHCH2(4-pyridine)	P(O)(Ph)OMe	H	Cl	H	H	0,001	0,0056	0,0039
9a	H	CONHCH2(4-pyridine)	P(O)(Ph:3,5-di-Me)OEt	H	Cl	H	H	0,011	0,0577	0,008
27a	H	CONH2	P(O)Ph2	H	Cl	H	H	>1,25	>1,25	>1,25
7ai	H	CONH2	P(O)(Ph)OMe	H	H	H	H	0,001	1,1091	0,7056
60a	H	CONH2	P(O)(Ph)OH	H	Cl	H	H	>1,25	>1,25	>1,25
29a	H	CONH2	P(O)(Ph)CH3	H	Cl	H	H	0,0008	0,2664	1,3828
31a	H	CONH2	P(O)(Ph)N(Me)2	H	Cl	H	H	0,0139	0,4668	>1,25

6c	H	CONH2		P(O)(Ph:4-F)OEt	H	Cl	H	H	H	0,0345	>1,25	>1,25
6ac	H	CONH2		P(O)(Ph:3,5-di-F)OEt	H	Cl	H	H	H	0,0037	0,9862	>1,25
6ad	H	CONH2		P(O)(Ph:2-F)OEt	H	Cl	H	H	H	0,0009	0,0583	0,017
6ae	H	CONH2		P(O)(Ph:3-F)OEt	H	Cl	H	H	H	0,0005	0,0616	0,011
7ak	H	CONH2		P(O)(Ph:4-F)OMe	H	Cl	H	H	H	0,0055	0,0345	0,468
7aj	H	CONH2		P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	H	0,0006	0,0177	0,0091
6ab	H	CONH2		P(O)(Ph:3-Me)OEt	H	Cl	H	H	H	0,0005	0,0348	0,0196
6m	H	CONH2		P(O)(Ph:3-OMe)OEt	H	Cl	H	H	H	0,0023		0,0664
6w	H	CONH2		P(O)(Ph:4-Me)OEt	H	Cl	H	H	H	0,0072		>1,25
6f	H	CONH2		P(O)(Ph:3-CF3)OEt	H	Cl	H	H	H	0,0067		0,498
6g	H	CONH2		P(O)(Ph:3-C≡N)OEt	H	Cl	H	H	H	0,0095		0,1951
6x	H	CONH2		P(O)(Ph:2-Me)OEt	H	Cl	H	H	H	0,0213		0,4483
7l	H	CONH2		P(O)(Ph:3-OMe)OMe	H	Cl	H	H	H	0,0006	0,0385	0,0218
7ac	H	CONH2		P(O)(Ph:3-Me)OMe	H	Cl	H	H	H	0,0013	0,0096	0,0055
6s	H	CONH2		P(O)(Ph:3-F,5-Me)OEt	H	Cl	H	H	H	0,0014		0,0207
57a	H	CONH2		P(O)(CH(CH2)5)OMe	H	Cl	H	H	H	0,0017		0,1615

6y	H	CONH2	P(O)(Ph:3,4-di-Me)OEt	H	Cl	H	H	0,0026		>1,25
7r	H	CONH2	P(O)(Ph:3-F,5-Me)OMe	H	Cl	H	H	0,0004	0,0005	0,0004
25b	H	CONH2	P(S)(Ph)OMe	H	Cl	H	H	0,0004	0,0414	0,0899
7s	H	CONH2	P(O)(Ph:3,5-di-CF3)OMe	H	Cl	H	H	0,0128	0,0683	0,0479
7q	H	CONH2	P(O)(Ph:3-F,5-CF3)OMe	H	Cl	H	H	0,0011	0,0166	0,0029
6d	H	CONH2	P(O)(Ph:3-Cl)OEt	H	Cl	H	H	0,0014	0,0166	0,0158
7n	H	CONH2	P(O)(Ph:2-OMe)OMe	H	Cl	H	H	0,0017	0,066	0,0015
6t	H	CONH2	P(O)(1-naphthalene)OEt	H	Cl	H	H	0,0325	>1,25	>1,25
6e	H	CONH2	P(O)(Ph:3-Et)OEt	H	Cl	H	H	0,002	0,0341	0,0012
7ad	H	CONH2	P(O)(Ph:2-Me)OMe	H	Cl	H	H	0,0013	0,032	0,0047
7ae	H	CONH2	P(O)(Ph:3,4-di-Me)OMe	H	Cl	H	H	0,0005	0,0111	0,0011
6ai	H	CONH2	P(O)(Ph:2-Cl)OEt	H	Cl	H	H	0,0012	0,1456	0,0173
7am	H	CONH2	P(O)(Ph:2,5-di-F)OMe	H	Cl	H	H	0,0003	0,0149	0,0012
7al	H	CONH2	P(O)(Ph:2-F)OMe	H	Cl	H	H	0,0005	0,0139	0,0026
7c	H	CONH2	P(O)(Ph:3-Cl)OMe	H	Cl	H	H	0,0004	0,0087	0,0026
7d	H	CONH2	P(O)(Ph:3-Et)OMe	H	Cl	H	H	0,0006	0,0065	0,0014

7e	H	CONH2		P(O)(Ph:3-CF3)OMe	H	Cl	H	H	0,0009	0,0032	0,0012
7ao	H	CONH2		P(O)(Ph:2-Cl)OMe	H	Cl	H	H	0,0004	0,002	0,0014
7t	H	CONH2		P(O)(1-naphthalene)OMe	H	Cl	H	H	0,0006	0,0057	0,0026
7a	H	CONH2		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0002	0,0012	0,001
6ah	H	CONH2		P(O)(Ph:2-Et)OEt	H	Cl	H	H	0,0162	0,211	0,3072
18at	H	CONHCH2(2-furan)		P(O)(Ph)OMe	H	Cl	H	H	0,0018	0,0015	0,0005
7an	H	CONH2		P(O)(Ph:2-Et)OMe	H	Cl	H	H	0,0029	0,0505	0,0013
6aa	H	CONH2		P(O)(Ph:2,5-di-Me)OEt	H	Cl	H	H	0,0002	0,0019	0,0016
18as	H	CONHCH2(5-imidazole:1-Me)		P(O)(Ph)OMe	H	Cl	H	H	0,0003	0,0099	0,0006
7ag	H	CONH2		P(O)(Ph:2,5-di-Me)OMe	H	Cl	H	H	0,0003	0,0407	0,001
7-Aza	H	CONH2		P(O)(Ph)OMe	H	Cl	H	N7	0,0147	>1,25	0,1929
7b	H	CONH2		P(O)(Ph:3,5-di-Cl)OMe	H	Cl	H	H	0,0012	0,0053	0,0168
57b	H	CONH2		P(O)(CH2CH=CMe2)OMe	H	Cl	H	H	0,0008	0,0131	0,1743
18a	H	CONHCH2(4-pyridine)		P(O)(Ph:3-Me)OMe	H	Cl	H	H	0,001	0,0101	0,0031
18b	H	CONHCH2(2-furan)		P(O)(Ph:3-Me)OMe	H	Cl	H	H	0,0036	0,0101	0,004
18av	H	CONHCH2(4-pyridine)		P(O)(Ph:3-F)OMe	H	Cl	H	H	0,0007	0,0013	0,0024

18av	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3-OMe)OMe	H	Cl	H	H	0,0008	0,0033	0,0014
18au	H	CONHCH ₂ (2-furan)	P(O)(Ph:3-OMe)OMe	H	Cl	H	H	0,0008	0,0033	0,006
6i	H	CONH ₂	P(O)(Ph:3-iPr)OEt	H	Cl	H	H	0,0012	0,0091	0,0035
7g	H	CONH ₂	P(O)(Ph:3-iPr)OMe	H	Cl	H	H	0,0025	0,0088	0,0077
51	H	CONH ₂	P(O)(2-thiophene)OEt	H	Cl	H	H	0,0022	0,0529	0,0656
52b	H	CONH ₂	P(O)(2-thiophene)OMe	H	Cl	H	H	0,0005	0,0087	0,0025
7af	H	CONH ₂	P(O)(Ph:2,3-di-Me)OMe	H	Cl	H	H	0,0022	0,1156	0,033
18aab	H	CONHCH ₂ (Ph:2-F)	P(O)(Ph)OMe	H	Cl	H	H	0,0038	0,0335	0,0102
18ax	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,003	0,0072	0,0023
18aan	H	CONHCH(Ph)CONH ₂ (S)	P(O)(Ph)OMe ("R")	H	Cl	H	H	0,015	0,3044	0,6004
18aao	H	CONHCH(Ph)CONH ₂ (S)	P(O)(Ph)OMe ("S")	H	Cl	H	H	0,0381	0,287	0,1177
7z	H	CONH ₂	P(O)(Ph:3,5-di-Me)OMe	F	Cl	H	H	0,0021	0,0085	0,0022
7h	H	CONH ₂	P(O)(Ph:3-CH ₂ F)OMe	H	Cl	H	H	0,0009	0,0061	0,0025
18c	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3-CH ₂ F)OMe	H	Cl	H	H	0,002	0,0029	0,0047
18ap	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3-Et)OMe	H	Cl	H	H	0,0058	0,0037	0,0058
18aac	H	CONHCH ₂ N(CH ₂ CH ₂) ₂ O	P(O)(Ph)OMe	H	Cl	H	H	0,0005	0,0101	0,014

7ap	H	CONH2	P(O)(Ph:3-CH2OCH2Ph)OMe	H	Cl	H	H	0,1528	1,1352	>1,25
18r	H	CONHCH2(4-pyridine)	P(O)(Ph:3,5-di-Me)OMe	F	Cl	H	H	0,0038	0,0165	0,0036
7f	H	CONH2	P(O)(Ph:3-C≡N)OMe	H	Cl	H	H	0,0009	0,0105	0,0153
18ay	H	CONHCH2N(CH2CH2)2O	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0035	0,0198	0,0093
7aq	H	CONH2	P(O)(Ph:3-CH2OH)OMe	H	Cl	H	H	0,0025	0,0831	0,1265
18d	H	CONHCH2(4-pyridine)	P(O)(Ph:3-F,5-Me)OMe	H	Cl	H	H	0,0023	0,0094	0,0037
18aq	H	CONHCH2(4-pyridine)	P(O)(Ph:3-iPr)OMe	H	Cl	H	H	0,012	0,0317	0,0186
18aaq	H	CONHCH2(Ph)	P(O)(Ph)OMe	H	Cl	H	H	0,0072	0,0184	0,0276
52a	H	CONH2	P(O)(3-thiophene)OMe	H	Cl	H	H	0,0005	0,0015	0,0063
22m	H	CONHCH2(4-pyridine:4-O)	P(O)(Ph:3,5-di-Me)OMe	F	Cl	H	H	0,0021	0,0026	0,0019
7aa	H	CONH2	P(O)(Ph:3-cPr)OMe	H	Cl	H	H	0,0022	0,007	0,0037
18aad	H	CONHCH2(Ph:2-F)	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,011	0,0382	0,0125
7ar	H	CONH2	P(O)(Ph:3-tBu)OMe	H	Cl	H	H	0,0029	0,0193	0,0306
59a	H	CONHCH2C≡N	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,0014	0,0473	0,0084
59b	H	CONH(CH2)2C≡N	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,0035	0,0693	0,0215
18e	H	CONHCH2(4-pyridine)	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,0035	0,0148	0,012

22a	H	CONHCH ₂ (4-pyridine-4-O)	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,0031	0,0524	0,0113
18q	H	CONHCH ₂ (Ph:4-C≡N)	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,003	0,0357	0,0542
22n	H	CONHCH ₂ (4-pyridine-4-O)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,002	0,0088	0,0045
7i	H	CONH ₂	P(O)(Ph:3-Ph)OMe	H	Cl	H	H	0,0099	0,3983	0,3801
18aae	H	CONHCH ₂ (5-imidazole:1-Me)	P(O)(Ph:3,5-di-F)OMe	H	Cl	H	H	0,0026	0,0105	0,0027
18az	H	CONHCH ₂ (Ph:2-F)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,003	0,0167	0,0085
18aaa	H	CONHCH ₂ (5-imidazole:1-Me)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0032	0,0064	0,0033
22l	H	CONHCH ₂ (4-pyridine:4-Me.I)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0061	0,2693	0,0774
18f	H	CONHCH ₂ (3-pyridine)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0023	0,0059	0,0034
18g	H	CONH(CH ₂) ₂ (4-pyridine)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0116	0,0306	0,0098
22c	H	CONH(CH ₂) ₂ (4-pyridine:4-O)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0028	0,0101	0,0268
18aah	H	CONHCH ₂ (Ph:2,6-di-F)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,007	0,0228	0,0169
7v	H	CONH ₂	P(O)(Ph:3-F,5-Cl)OMe	H	Cl	H	H	0,0008	0,0149	0,0023
18aai	H	CONHCH ₂ (Ph:2,5-di-F)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0036	0,0156	0,0183
22b	H	CONHCH ₂ (3-pyridine:3-O)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0035	0,0187	0,0032
18n	H	CONHCH ₂ (Ph:2-NH ₂)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0195	0,0878	0,0231

18k	H	CONHCH ₂ (Ph:4-Br)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0393	0,1023	0,0504
18aaj	H	CONHCH ₂ (Ph:2,4-di-F)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0316	0,1089	0,0416
7as	H	CONH ₂	P(O)(Ph:3-Me,5-C≡N)OMe	H	Cl	H	H	0,0014	0,0048	0,0029
7x	H	CONH ₂	P(O)(Ph:3-nPr)OMe	H	Cl	H	H	0,003	0,0138	0,0087
18aaj	H	CONHCH ₂ (Ph:2-Cl)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0155	0,0214	0,0258
18aak	H	CONHCH ₂ (Ph:2-CF ₃)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	>1,25	>1,25	0,9067
18aaf	H	CONHCH ₂ (4-pyridine)	P(O)(Ph:3-Me,5-C≡N)OMe	H	Cl	H	H	0,0018	0,0039	0,0011
18aag	H	CONHCH ₂ (5-imidazole:1-Me)	P(O)(Ph:3-Me,5-C≡N)OMe	H	Cl	H	H	0,0031	0,0188	0,0029
	H	CONHCH ₂ (Ph:2,3,6-tri-F)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0179	0,0171	0,0228
22ak	H	CONHCH ₂ (4-pyridine:4-O)	P(O)(Ph:3-Me,5-C≡N)OMe	H	Cl	H	H	0,007	0,0067	0,011
18aal	H	CONHCH ₂ (2-pyridine)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0301	0,0197	0,0218
18aam	H	CONHCH ₂ (3-pyridine)	P(O)(Ph:3,5-di-Me)OMe	F	Cl	H	H	0,0173	0,0068	0,0225
22j	H	CONHCH ₂ (3-pyridine:3-O)	P(O)(Ph:3,5-di-Me)OMe	F	Cl	H	H	0,0068	0,0099	0,006
22p	H	CONHCH ₂ (2-pyridine:2-O)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0019	0,0513	0,0179
7at	H	CONH ₂	P(O)(Ph:3-F,5-C≡N)OMe	H	Cl	H	H	0,0057	0,0221	0,0103
18l	H	CONHCH ₂ (4-pyrimidine)	P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0017	0,0025	0,0022

7w	H	CONH2		P(O)(Ph:3-Me,5-Et)OMe	H	Cl	H	H	0,0004	0,0055	0,0012
22i	H	CONHCH2(4-pyrimidine:1-O)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0014	0,0062	0,0079
22h	H	CONHCH2(3-pyridine:3-O)		P(O)(Ph:3-Me,5-Et)OMe	H	Cl	H	H	0,0016	0,0058	0,0036
18h	H	CONHCH2(4-pyridine:3-F)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0017	0,0143	0,0043
22d	H	CONHCH2(4-pyridine:3-F,4-O)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0029	0,0015	0,0067
18i	H	CONHCH2(2-pyrazine)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,003	0,0168	0,0018
18j	H	CONHCH2(5-pyrimidine)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0036	0,0051	0,0019
22g	H	CONHCH2(5-pyrimidine:1-O)		P(O)(Ph:3,5-di-Me)OMe	H	Cl	H	H	0,0021	0,0107	0,003
33a	H	CONH2		P(O)(Ph:3-CH=CHC≡N)OMe	H	Cl	H	H	0,0032	0,002	0,0034
7j	H	CONH2		P(O)(Ph:3-F,5-Br)OMe	H	Cl	H	H	0,0018	0,003	0,0041

Ph is phenyl; Me is methyl; and Et is ethyl. Relative solubilities (at OD 600nm) for the compounds ranged from about

75 $\mu\text{M}/\text{mL}$ for Compound No. 5 to $>1000 \mu\text{M}/\text{mL}$ for Compound Nos. 7-11. A cytochrome-P450 binding assay completed for Compound No. 3 indicated binding of 15.87 μM , and MDRC4 and CNDO cell-based assays for this same compound versus other mutant forms such as, for example, K103N/M184V/NAMS and K103N/Y181C/M184V/NAMS, gave EC_{50} values of from 0.0013-0.4466 μM .

TOXICOLOGY

1. pH dependent aqueous solubility

The aqueous solubility of each compound was determined by the conventional shake-flask method at a saturated concentration of 1mM. The flask (vial) was shaken for 3 hrs at ambient temperature, and then centrifuged. The supernatant was analyzed by HPLC with UV detection for solubility determination. In general, a higher aqueous solubility is more desirable for drug candidates.

2. Human plasma protein binding

The human plasma protein binding was determined using the Equilibrium Dialysis method. Dialysis was conducted with pooled human plasma at 37°C for approximately 6 hours at a drug concentration of 1 μ M. At the end of the dialysis, buffer samples from the buffer-side of the dialysis cell were collected and analyzed by LC/MS/MS for free drug concentration. For NNRTIs, a lower protein binding is more desirable.

3. Bi-directional CACO-2 permeability

The objective of this assay is to determine the bi-directional permeability classification and efflux-limited absorption potential of a test compound in a Caco-2 cell monolayer system. Typically, the assay involves the measurement of non-specific binding to the Transwell apparatus in a pH7.4 assay buffer, the bi-directional permeability assessment of a test compound across Caco-2 cell monolayers, apical-to-basolateral transport assessment, basolateral-to-apical transport assessment, and monolayer integrity. High permeability and no efflux suggest that intestinal permeability is not expected to be a limiting factor for oral absorption in humans.

4. CYP450 inhibition

In vitro CYP450 inhibition screening allows for the prediction of potential drug-drug interactions. To determine whether a test compound inhibits a particular P450 enzyme activity, changes in the metabolism of a P450-specific substrate by human liver microsomes are monitored with varying concentrations of the test compound. Potency and rank order of the inhibition can be assessed by determination of the IC₅₀ values for a particular isozyme. For NNRTIs, higher IC₅₀ values would suggest less inhibition, and thus less potential for drug-drug interaction in patients.

CYP3A4 inhibition was screened using the CYP3A4/BFC high throughput inhibitor screening kit (BD Biosciences), CYP2D6 inhibition was screened using the CYP2D6/AMMC high throughput inhibitor screening kit (BD Biosciences), and CYP2C9 inhibition was screened using the P450-Glo™ Assay kit (Promega).

5. In vitro metabolic stability in liver microsomes

The metabolic stability assay is to evaluate the stability of test compound in biological matrices. Data are useful in terms of understanding and predicting the elimination mechanisms of the test compound. CYP450-dependent metabolism of drugs can also vary widely from one species to another. Evaluating in vitro metabolism of a drug by liver microsomes from multiple species allows the comparison of metabolism in animal species to that of humans. This can help identify the most relevant animal model for PK and toxicology studies. The metabolic stability of a test compound was evaluated in vitro in liver microsomes of the rat, dog, monkey, and human. 10 μM of the test compound was pre-incubated for 5 min at 37°C with 1mg/mL liver microsomes in 0.1M Tris buffer, pH 7.4 containing 5mM MgCl₂ and 0.1mM EDTA. Following pre-incubation, NADPH (final concentration of 3 mM) was added to start the reaction and samples incubated for 0 and 1 or 2 hours. After terminating the reaction, the supernatant was analyzed by HPLC-UV or LC/MS/MS for the disappearance of the parent and formation of metabolite(s). The %parent remaining was the peak area ratio of the 1 or 2 hr sample to the time 0 sample. In general, less metabolism (higher %parent values) is more desirable.

6. In vitro metabolism in liver microsomes – metabolic pathway

Phase I biotransformation of a test compound is evaluated in this assay. Samples from the metabolic stability experiment were analyzed by LC/MS/MS for metabolite profiling and identification. Structures of metabolites were elucidated based on multiple MS/MS experiments such as full scan, neutral loss scan, and product ion scan. Metabolic pathways were then postulated based on the structures of major metabolites. For compounds from the phosphinate series, N-oxidation of the terminal pyridine ring is the predominant pathway, followed by the oxidation of the dimethyl-phenyl group, where the methyl group(s) is hydroxylated, further oxidized to an aldehyde, and finally to a

carboxylic acid. The elucidation of metabolic pathways is important in terms of understanding the elimination mechanism of a test compound and assisting the design of new molecules with improved DM-PK profiles.

Cat.	Cmpd	Aqueous solubility (uM)				Protein binding (human plasma)		CYP450 inhibition, human (IC50, uM)			CYP3A4 induction, human hepatocytes		Metabolic stability in liver microsomes %Parent after 2hrs			
		pH1	pH3	pH7	pH9	%Free	CYP3A4	CYP2D6	CYP2C9	(mRNA)	Human	Rat	Dog	Monke		
Parent N		>1000	231	2.1	1.7	0.2%	0.66	2.04		weak to moderate	22%*	1%*	*	1%*		
N-Oxide D		52	74.2	50.6	66.8	1.6%	1.07	>10	>10		76%	1%	82%	24%		
Parent N		820	590	9.1	10.1	0.9%	0.81	1.35	5.35		15%	2%	29%	1%		
N-Oxide B		190	170	200	190	3.8%	1.04	>10	>10		94%	1%	85%	63%		
Parent P		>1000	724.3	24	33.8	<0.5%	0.77	2.01	1.68		3%	0%	20%	3%		
N-Oxide G						1.6%	1.14	>10	8.85		85%	24%	77%	41%		

*%Parent after 1hr
incubation

Cat.	Cmpd	Metabolic pathway (in vitro)
Parent	N	N-oxidation, hydroxylation to CH ₃ OH- to CHO- to COOH-
N-		
Oxide	D	Hydroxylation to CH ₃ OH- to CHO- to COOH-
Parent	O	N-oxidation, hydroxylation to CH ₃ OH- to CHO- to COOH-
N-		
Oxide	B	Hydroxylation to CH ₃ OH- to CHO- to COOH-
Parent	P	N-oxidation, hydroxylation to CH ₃ OH- to CHO- to COOH-
N-		
Oxide	G	Hydroxylation to CH ₃ OH- to CHO- to COOH-


7. PK and oral bioavailability in the rat and dog.

The pharmacokinetics of a test compound was evaluated in Sprague-Dawley rats and Beagle dogs. A typical PK study involved dosing of 2 to 3 animals via a single IV bolus injection at 1 mg/kg and another 3 animals via a single oral gavage at 5 mg/kg. Blood samples were collected at various time points over a 24-hr period. Plasma was separated and analyzed for test compound and its metabolite(s) by LC/MS/MS. PK parameters were calculated from the plasma concentration – time profile using a noncompartmental method. Oral bioavailability (F) was calculated based on dose-normalized AUC values from oral and IV administration. Higher oral bioavailability values are better.

MEAN PK PARAMETERS (N = 3 OR 2)												
Cmpd	Cat.	Species	Route	Dose	Vehicle	Cmax	Tmax	T1/2	AUClast	AUCinf	AUCinf	F
				(mg/kg)		(ng/mL)	(hr)	(hr)	(hr*ng/mL)	(hr*ng/mL)	(mL/kg/hr)	(%)
D	N-Oxide	Rat	IV	1.0	PEG400	165.7		0.9	174.3	188.3	5417.8	6368.6
D	N-Oxide	Rat	PO	5.0	PEG400	45.0	0.5	2.2	132.0	150.9	30.2	16.0%
N	Parent	Rat	IV	1.0	PEG400	118.8		0.9	108.3	125.9	7992.2	9587.9
N	Parent	Rat	PO	5.0	PEG400	NC	NC	NC	NC	NC	NC	NC
N	Parent	Dog	IV	1.0	PEG400	407.0		4.2	923.5	970.1	1192.3	4079.3
N	Parent	Dog	PO	5.0	PEG400	60.9	0.7	2.8	116.4	138.6	27.7	2.9%
D	N-Oxide	Dog	IV	1.0	PEG400	429.7		3.2	1007.0	1220.5	821.8	3715.7
D	N-Oxide	Dog	PO	5.0	0.5%MC	116.2	2.7	2.6	696.1	746.7	149.3	12.4%

WE CLAIM:

1. An optionally substituted 3-phosphoindole compound.
2. The compound of claim 1 wherein the 3-phosphoindoles is in the form of a phosphate, a phosphonate, a thiophosphate, a thiophosphonate, an iminiophosphate or an iminophosphonate.
3. The compound of claim 1 wherein the compound is a phosphate or a phosphonate.
4. The compound of claim 1 wherein the compound is further substituted at the 2 position.
5. The compound of claim 1 wherein the compound is substituted at the nitrogen of the indole.
6. The compound of claim 1 wherein the compound is of Formula (A):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

X and Y, each independently, is

- b) 3-14 membered carbocycle, aryl, heterocycle, any of which may comprise a monocyclic, bicyclic, tricyclic or spiro structure, or optionally may be substituted;
- b) H;
- dd) OH;
- ee) Cl, Br, I, F;

- ff) CF_3 ;
- gg) C_{1-6} alkyl;
- hh) C_{2-6} alkenyl;
- ii) C_{2-6} alkynyl;
- jj) alkylheterocycle;
- kk) NH_2 ;
- ll) NH-alkyl;
- mm) N-dialkyl;
- nn) NH-aryl;
- oo) N-alkaryl;
- pp) N-aralkyl;
- qq) NH-heterocycle;
- rr) N-alkyl-heterocycle;
- ss) N-alkenyl-heterocycle;
- tt) N-alkynyl-heterocycle;
- uu) O-alkyl;
- vv) O-alkenyl;
- ww) O-alkynyl;
- xx) O-alkylaryl;
- yy) O-aryl;
- zz) O-heterocycle;
- aaa) O-aralkyl;
- bbb) O-carbocycle;
- ccc) SR^2 ; or

ddd) NR^2R^3 ;

alternatively, X and Y may be joined to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle wherein each ring comprises 3-7 members;

Z is:

mm) H;

nn) alkoxy;

oo) NO_2 ;

pp) $\text{N}(\text{R}^2)(\text{R}^3)$;

qq) OR^2 ;

rr) carboxamido;

ss) amido;

tt) acyl;

uu) $\text{S}(\text{O})_n\text{R}^2$;

vv) $\text{S}(\text{O})_n\text{-NR}^2\text{R}^3$;

ww) C_{1-6} alkyl;

xx) C_{2-6} alkenyl;

yy) C_{2-6} alkynyl;

zz) alkaryl;

aaa) aralkyl;

bbb) heterocycle;

ccc) alkyl-heterocycle;


ddd) aryl;

eee) CN;

fff) $\text{C}(=\text{W})\text{-R}^2$;

- ggg) $C(=W)NH-C(R^2)(R^2)-C(=W)-N(R^2)(R^2)$;
- hhh) $C(=W)NH-P(=W)(R^2)-A-R^2$;
- iii) $C(=W)NH-A-S(O)_n-NR^2$;
- jjj) $C(=W)NH-CR^2R^3-S(O)_n-NR^2R^3$;
- kkk) $C(=W)-NH-A-C(=W)-N(R^2)(R^3)$;
- lll) $C(=W)-N(R^2)(R^3)$;
- mmm) $C(=W)-NH-A-R^2$;
- nnn) $C(=W)-NH-NH-R^2$;
- ooo) $C(=W)-NH-C(R^2)(R^2)-C(=W)NH-C(R^2)(R^3)C(=W)-N(R^2)(R^3)$;
- ppp) $C(=W)-NH-R^2$;
- qqq) $C(=W)-NH-A-C(=W)-NH-A-C(=W)-NH_2$;
- rrr) $C(R^2)(R^2)(R^3)$;
- sss) $C(R^2)(R^2)-NH-R^2$;
- ttt) $A-S(O)_n-R^2$;
- uuu) $C(=W)-A-C(=W)-A-C(=W)R^3$;
- vvv) $A-R^2$;
- www) $C(=W)-(O)R^2$;
- xxx) $C(=W)-A-C(=W)-NH_2$;
- mm)an amino acid residue;
- aaa) $C(=W)-N(R^2)-A-(\text{amino acid residue})$;
- bbb) $C(=W)-N(R^2)-A-(\text{amino acid residue})-C(=W)-R^2$;
- ccc) $C(=W)-\text{amino acid residue}$;
- ddd) $C(=W)-N(R^2)-A-(\text{amino acid residue})-A-C(=W)-R^2$;

- eee) $C(=W)-OR^3$;
- fff) $C(=W)-S(R^2)$;
- ggg) $C(=W)-NH-NH-R^2$;
- hhh) $C(=W)-NH-N(R^2)-A-C(=W)R^2$;
- iii) $C(=W)-N(R^2)-C(=W)-R^3$;
- jjj) $C(=W)-A-NH-C(=W)R^2$;
- kkk) $C(=W)-A-NH-C(=W)OR^3$;
- lll) $C(=W)-A-R^3$;
- mmm) $C(=W)-NH-NH-CH_2-C(=W)R^2$;
- aaa) $P(=W)(R^2)(R^2)$; or
- bbb) $A-P(=W)(R^2)(R^2)$;
- hhh) $A-P(=W)(R^3)(R^3)$;
- iii) $C(=W)-NH-C_{1-10}alkyl-heteroaryl$;
- jjj) $C(=W)-NH-C_{1-4}alkyl-heteroaryl$;
- kkk) $C(=W)-NH-CH_2-heteroaryl$; and



wherein in embodiments (ddd), (eee) and (fff), the heteroaryl can optionally include a charged heteroatom, and in particular can include an N-oxide

wherein each of the foregoing X, Y and Z independently may be unsubstituted or substituted by one or more of:

- b) H;
- ii) C_{1-6} alkyl;
- iii) alkoxy;

- iv) OH;
- v) oxo;
- vi) halo;
- vii) NR^2R^2 ;
- viii) optionally substituted aryl;
- ix) optionally substituted heterocyclyl;
- x) O-C(=W)-alkyl ;
- xi) C(=W)-OR^2 ;
- x) CN;
- xi) NO_2 ;
- xii) NH-C(=W)-alkyl ;
- xiii) $\text{NH-S(O)}_n\text{-alkyl}$;
- xiv) $\text{NH-S(O)}_n\text{-NR}^2\text{R}^2$; or
- xv) C_{3-6} cycloalkyl;

W is:

- h) O;
- i) S;
- j) NH;
- k) $\text{N-N(R}^2\text{)(R}^2\text{)}$;
- l) $\text{N(R}^2\text{)}$;
- m) N-OH; or
- n) N-O-alkyl;

R^1 is:

- l) H;

- m) R^2 ;
- n) $C(=W)-R^2$;
- o) $C(=W)-O(R^2)$;
- p) $C(=W)-S(R^2)$;
- q) $C(=W)-NH-R^2$;
- r) $C(=W)-N(R^2)(R^2)$;
- s) $C(=W)-NH-A-(\text{amino acid residue})$;
- t) $A-(\text{amino acid residue})-R^2$;
- u) $S(O)_n-R^3$; or
- v) $S(O)_2-N(R^2)(R^2)$;

any of which optionally may be substituted by one or more:

- i. C_{1-6} alkyl;
- ii. OH;
- iii. alkoxy;
- iv. aryl;
- v. halo;
- vi. CN;
- vii. NO_2 ; or
- viii. $N(R^2)(R^2)$;

Each R^2 is independently selected from:

- aaa) H;
- bbb) OH;
- ccc) halogen;
- ddd) optionally substituted, branched or unbranched alkyl;

- eee) optionally substituted, branched or unbranched alkenyl;
- fff) optionally substituted, branched or unbranched alkynyl;
- ggg) 3-14 membered carbocycle;
- hhh) alkylheterocycle;
- iii) acyl;
- jjj) carboxamido;
- kkk) carbamoyl;
- lll) alkoxy;
- mmm) optionally substituted aryl;
- nnn) optionally substituted aralkyl;
- ooo) optionally substituted alkylaryl;
- ppp) O-alkyl;
- qqq) O-alkenyl;
- rrr) O-alkynyl;
- sss) O-alkaryl;
- ttt) O-aralkyl;
- uuu) O-carbocycle;
- vvv) O-heterocycle;
- www) O-aryl;
- xxx) CF_3 ;
- yyy) CN;
- zzz) $\text{S(O)}_n\text{-R}^3$;
- aaaa) $\text{N(R}^3\text{)(R}^3\text{)}$;
- bbbb) $\text{NH-S(O)}_n\text{-R}^3$;

cccc)	NHC(=W)-aryl;
dddd)	NHC(=W)-alkyl;
eeee)	$\text{NHC(=W)-heterocycle;}$
ffff)	$\text{CH}_2\text{-S(O)}_n\text{R}^3;$
gggg)	$\text{C(=W)R}^3;$
hhhh)	$\text{C(=W)NR}^3\text{R}^3;$
iiii)	$\text{C(alkyl)}_2\text{-S(O)}_n\text{R}^3;$
jjjj)	$\text{CH(alkyl)-S(O)}_n\text{R}^3;$
kkkk)	$\text{C(alkyl)}_2\text{-NH}_2;$
llll)	$\text{CH(alkyl)-N(alkyl)R}^3;$
mmmm)	$\text{CR}^3\text{R}^2\text{-NR}^3\text{R}^3;$
nnnn)	$\text{CH}_2\text{N(alkyl)R}^3;$
oooo)	$\text{CH(alkyl)-NHR}^3;$
pppp)	$\text{C(alkyl)}_2\text{-NHR}^3;$
qqqq)	$\text{C(alkyl)}_2\text{-N(alkyl)R}^3;$
rrrr)	$\text{CH}_2\text{-C(=W)H;}$
ssss)	$\text{CH}_2\text{-C(=W)alkyl;}$
tttt)	$\text{CR}^3\text{R}^3\text{-C(=W)R}^3;$
uuuu)	$\text{A-R}^3;$
vvvv)	$\text{C(R}^3)_2\text{-C(=W)R}^3;$
wwww)	$\text{CH}_2\text{-C(=W)H;}$
xxxx)	$\text{CH}_2\text{-C(=W)alkenyl;}$
yyyy)	$\text{CH(alkenyl)-C(=W)H;}$
zzzz)	$\text{A-S(O)R}^3;$

aaa) $\text{CH}(\text{NH})\text{-S}(\text{O})_n\text{R}^3$; or

bbb) $\text{A-N}(\text{NH})\text{R}^3$;

wherein said optional substitution comprises one or more of:

A) a substituted or unsubstituted heterocycle;

B) $\text{C}(=\text{W})\text{O-aryl}$;

C) $\text{C}(=\text{W})\text{O-alkyl}$;

D) $\text{C}(=\text{W})\text{NH}_2$;

E) $\text{C}(=\text{W})\text{NH-alkyl}$;

F) $\text{C}(=\text{W})\text{NH-aryl}$;

G) $\text{C}(=\text{W})\text{N-di-alkyl}$;

H) $\text{C}(=\text{W})\text{N(alkyl)-aryl}$;

I) α -amino acid;

J) α -amino ester;

K) α -amino-carboxamide;

L) β -amino acid;

M) β -amino ester; or

N) β -amino-carboxamide;

Each R^3 is:

bb) H;

cc) OH;

dd) C_{1-6} alkyl;

ee) C_{2-6} alkenyl;

ff) C_{2-6} alkynyl;

gg) alkoxy;

- hh) CF_3 ;
- ii) CN ;
- jj) amino;
- kk) NR^2R^2 ;
- ll) O-alkyl;
- mm) O-alkenyl;
- nn) O-alkynyl;
- oo) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{NH}_2$;
- pp) $\text{C}(\text{R}^2)(\text{R}^2)\text{-S}(\text{O})_n\text{CF}_3$;
- qq) $\text{C}(\text{R}^2)(\text{R}^2)\text{-NH}_2$;
- rr) A-heterocycle;
- ss) $\text{C}(\text{R}^2)(\text{R}^2)\text{-NR}^2\text{R}^2$;
- tt) $\text{C}(\text{R}^2)(\text{R}^2)\text{-C}(=\text{W})\text{R}^2$;
- uu) aryl;
- vv) carbocycle;
- ww) heterocycle;
- xx) cycloalkyl;
- yy) alkaryl;
- zz) alkylheterocycle;
- aaa) aralkyl; or
- bbb) heterocycle-alkyl;

any of which may be unsubstituted or substituted with one or more of the following taken in any combination:

- i. halo;

- xxiv. OH;
- xxv. OR^2 ;
- xxvi. SR^2 ;
- xxvii. COOH;
- xxviii. carboxylic acid ester;
- xxix. $C(=W)R^2$;
- xxx. $C(=W)OR^2$;
- xxxi. $C(=W)OR^3$;
- xxxii. $C(=W)SR^2$;
- xxxiii. $A-C(=W)NH_2$;
- xxxiv. $C(=W)NR^2R^3$;
- xxxv. NR^2R^2 ;
- xxxvi. NR^2R^3 ;
- xxxvii. $NR^2-S(O)_nR^3$;
- xxxviii. $NR^2-C(=W)-C_{1-6}alkyl$;
- xxxix. $S(O)_nR^3$;
- xl. $C_{1-6}alkoxy$;
- xli. $C_{1-6} thioether$;
- xl.ii. amino acid residue;
- xl.iii. $NH-A-(amino\ acid\ residue)$;
- xl. iv. $C(=W)NH-A-(amino\ acid\ residue)$; and
- xl. v. wherein when said optional substitution comprises a substituted heterocycle, then substitution is selected from the group consisting of:

- b. $C(=W)O$ -aryl;
 - c. $C(=W)O$ -alkyl;
 - d. $C(=W)NH_2$;
 - e. $C(=W)NH$ -aryl;
 - f. $C(=W)NH$ -alkyl;
 - g. $C(=W)N$ -di-alkyl;
 - h. $C(=W)N(alkyl)$ -aryl;
 - i. α -amino acid;
 - j. α -amino ester;
 - k. α -amino-carboxamide;
 - k. β -amino acid;
 - l. β -amino ester; or
 - m. β -amino-carboxamide;
 - n. halo; or
 - o. cyano,
- taken alone or in any combination;

n independently is 0, 1 or 2;

A is a disubstituted spacer selected from the group consisting of:

- j. C_{1-6} alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- k. C_{2-12} alkenylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;

- l. C₂₋₁₂ alkynylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- m. optionally substituted arylene;
- n. O-alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- o. aralkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- p. optionally substituted cycloalkyl; and
- q. optionally substituted heterocycle;

wherein "A" may be joined by any desired linkage or any combination of desired linkages.

Each R^{4'}, R^{5'}, R^{6'} and R^{7'} independently is:

- a. H;
- b. halogen;
- c. NO₂;
- d. CN;
- e. CF₃;
- f. OR²
- g. NR²R²;
- h. NHS(O)_nR²;
- i. NHCO-C₁₋₃ alkyl;
- j. S(O)_nR²;
- k. aryl;


- l. heterocycle;
- m. C₁₋₆ alkyl;
- n. C₂₋₆ alkenyl;
- o. C₂₋₆ alkynyl;
- p. C(=W)-S(O)_nR²;
- q. C(=W)-S(O)_n-NR²R²;
- r. C(=W)-aryl;
- s. C(=W)-alkyl;
- t. C(=W)-heterocycle; or
- u. C(=W)-NR²R²;


each of which optionally may be substituted with one or more of :






- r. OR²;
- ii. S(O)_nR²;
- iii. C(=W)-S(O)_nR²;
- iv. C(=W)-S(O)_n-NR²R²;
- xxi. C(=W)-aryl;
- xxii. C(=W)-alkyl;
- xxiii. C(=W)-heterocycle;
- xxiv. C(=W)NR²R²;
- xxv. H;
- xxvi. NO₂;
- xxvii. CN;
- xxviii. CF₃;
- xxix. halogen;

- xxx. $\text{NHS}(\text{O})_n\text{R}^2$;
- xxxi. NHCO-C_{1-3} alkyl;
- xxxii. aryl;
- xxxiii. heterocycle;
- xxxiv. C_{1-6} alkyl;
- xxxv. C_{2-6} alkenyl;
- xxxvi. C_{2-6} alkynyl; or
- xxxvii. NR^2R^2 .

- 7. The compound of claim 6 wherein $\text{R}^{5'}$ is halogen, CH_3 , CN , CF_3 or NO_2 .
- 8. The compound of claim 7 wherein $\text{R}^{5'}$ is Cl .
- 9. The compound of claim 6 wherein W is O or S .
- 10. The compound of claim 6 wherein Z is optionally substituted amide or carboxamide.
- 11. The compound of claim 6 wherein X is substituted or unsubstituted phenyl.
- 12. The compound of claim 6 wherein Y is $-\text{O-C}_{1-6}$ alkyl.
- 13. The compound of claim 6 wherein the compound is selected from the group consisting of:








5

14. The compound of claim 1 wherein the compound is of Formula (B):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine, stereochemical isomer or tautomer thereof, wherein:

Y is:

- a. 3-14 membered carbocycle, aryl, heterocycle, any of which may comprise a monocyclic, bicyclic, tricyclic or spiro structure, or any of which optionally may be substituted;
- b. H;
- c. OH;
- d. halogen;
- e. CF₃;
- f. C₁₋₆ alkyl;
- g. C₂₋₆ alkenyl;
- h. C₂₋₆ alkynyl;
- i. alkylheterocycle;
- j. NH₂;
- k. NH-alkyl;
- l. N-dialkyl;
- m. NH-aryl;
- n. N-alkaryl;

- o. N-aralkyl;
- p. NH-heterocycle;
- q. N-alkyl-heterocycle;
- r. N-alkenyl-heterocycle;
- s. N-alkynyl-heterocycle;
- t. O-alkyl;
- u. O-alkenyl;
- v. O-alkynyl;
- w. O-alkylaryl;
- x. O-aryl;
- y. O-heterocycle;
- z. O-aralkyl;
- aa. O-carbocycle;
- bb. SR^2 ; or
- cc. NR^2R^3 ;

Z is:

- aaa) H;
- bbb) alkoxy;
- ccc) NO_2 ;
- ddd) $N(R^2)(R^3)$;
- eee) OR^2 ;
- fff) carboxamido;
- ggg) amido;
- hhh) acyl;

- iii) $S(O)_nR^2$;
- jjj) $S(O)_n-NR^2R^3$;
- kkk) C_{1-6} alkyl;
- lll) C_{2-6} alkenyl;
- mmm) C_{2-6} alkynyl;
- nnn) alkaryl;
- ooo) aralkyl;
- ppp) heterocycle;
- qqq) alkyl-heterocycle;
- rrr) aryl;
- sss) CN;
- ttt) $C(=W)-R^2$;
- uuu) $C(=W)NH-C(R^2)(R^2)-C(=W)-N(R^2)(R^2)$;
- vvv) $C(=W)NH-P(=W)(R^2)-A-R^2$;
- www) $C(=W)NH-A-S(O)_n-NR^2$;
- xxx) $C(=W)NH-CR^2R^3-S(O)_n-NR^2R^3$;
- yyy) $C(=W)-NH-A-C(=W)-N(R^2)(R^3)$;
- zzz) $C(=W)-N(R^2)(R^3)$;
- aaaa) $C(=W)-NH-A-R^2$;
- bbbb) $C(=W)-NH-NH-R^2$;
- cccc) $C(=W)-NH-C(R^2)(R^2)-C(=W)NH-C(R^2)(R^3)C(=W)-N(R^2)(R^3)$;
- dddd) $C(=W)-NH-R^2$;
- eeee) $C(=W)-NH-A-C(=W)-NH-A-C(=W)-NH_2$;

- ffff) $C(R^2)(R^2)(R^3)$;
- gggg) $C(R^2)(R^2)-NH-R^2$;
- hhhh) $A-S(O)_n-R^2$;
- iiii) $C(=W)-A-C(=W)-A-C(=W)R^3$;
- jjjj) $A-R^2$;
- kkkk) $C(=W)-(O)R^2$;
- llll) $C(=W)-A-C(=W)-NH_2$;
- mmmm) an amino acid residue;
- nnnn) $C(=W)-N(R^2)-A-(\text{amino acid residue})$;
- oooo) $C(=W)-N(R^2)-A-(\text{amino acid residue})-C(=W)-R^2$;
- pppp) $C(=W)-\text{amino acid residue}$;
- qqqq) $C(=W)-N(R^2)-A-(\text{amino acid residue})-A-C(=W)-R^2$;
- rrrr) $C(=W)-OR^3$;
- ssss) $C(=W)-S(R^2)$;
- tttt) $C(=W)-NH-NH-R^2$;
- uuuu) $C(=W)-NH-N(R^2)-A-C(=W)R^2$;
- vvvv) $C(=W)-N(R^2)-C(=W)-R^3$;
- wwww) $C(=W)-A-NH-C(=W)R^2$;
- xxxx) $C(=W)-A-NH-C(=W)OR^3$;
- yyyy) $C(=W)-A-R^3$;
- zzzz) $C(=W)-NH-NH-CH_2-C(=W)R^2$;
- aaa) $P(=W)(R^2)(R^2)$; or
- bbb) $A-P(=W)(R^2)(R^2)$;

wherein each of the foregoing Y and Z independently may be unsubstituted or substituted by one or more of:

- r. H;
- s. C₁₋₆ alkyl;
- t. alkoxy;
- u. OH;
- v. oxo;
- w. halo;
- x. NR²R²;
- y. optionally substituted aryl;
- z. optionally substituted heterocyclyl;
- aa. O-C(=W)-alkyl;
- bb. C(=W)-OR²;
- cc. CN;
- dd. NO₂;
- ee. NH-C(=W)-alkyl;
- ff. NH-S(O)_n-alkyl;
- gg. NH-S(O)_n-NR²R²; or
- hh. C₃₋₆ cycloalkyl;

W is:

- h) O;
- i) S;
- j) NH;
- k) N-N(R²)(R²);

- l) $N(R^2)$;
- m) $N-OH$; or
- n) $N-O-alkyl$;

R^1 is:

- cc) H ;
- dd) R^2 ;
- ee) $C(=W)-R^2$;
- ff) $C(=W)-O(R^2)$;
- gg) $C(=W)-S(R^2)$;
- hh) $C(=W)-NH-R^2$;
- ii) $C(=W)-N(R^2)(R^2)$;
- jj) $C(=W)-NH-A-(\text{amino acid residue})$;
- kk) $A-(\text{amino acid residue})-R^2$;
- ll) $S(O)_n-R^3$; or
- mm) $S(O)_2-N(R^2)(R^2)$;

any of which optionally may be substituted by one or more:

- i. C_{1-6} alkyl;
- ix. OH ;
- x. alkoxy;
- xi. aryl;
- xii. halo;
- xiii. CN ;
- xiv. NO_2 ; or
- xv. $N(R^2)(R^2)$;

Each R^2 is:

- a) H;
- b) OH;
- c) halogen;
- ccc) optionally substituted, branched or unbranched alkyl;
- ddd) optionally substituted, branched or unbranched alkenyl;
- eee) optionally substituted, branched or unbranched alkynyl;
- fff) 3-14 membered carbocycle;
- ggg) alkylheterocycle;
- hhh) acyl;
- iii) carboxamido;
- jjj) carbamoyl;
- kkk) alkoxy;
- lll) optionally substituted aryl;
- mmm) optionally substituted aralkyl;
- nnn) optionally substituted alkaryl;
- ooo) O-alkyl;
- ppp) O-alkenyl;
- qqq) O-alkynyl;
- rrr) O-alkaryl;
- sss) O-aralkyl;
- ttt) O-carbocycle;
- uuu) O-heterocycle;
- vvv) O-aryl;

- www) CF_3 ;
- xxx) CN ;
- yyy) $\text{S(O)}_n\text{-R}^3$;
- zzz) $\text{N(R}^3\text{)(R}^3\text{)}$;
- aaaa) $\text{NH-S(O)}_n\text{-R}^3$;
- bbbb) NHC(=W)-aryl ;
- cccc) NHC(=W)-alkyl ;
- dddd) $\text{NHC(=W)-heterocycle}$;
- eeee) $\text{CH}_2\text{-S(O)}_n\text{R}^3$;
- ffff) C(=W)R^3 ;
- gggg) $\text{C(=W)NR}^3\text{R}^3$;
- hhhh) $\text{C(alkyl)}_2\text{-S(O)}_n\text{R}^3$;
- iiii) $\text{CH(alkyl)-S(O)}_n\text{R}^3$;
- jjjj) $\text{C(alkyl)}_2\text{-NH}_2$;
- kkkk) $\text{CH(alkyl)-N(alkyl)R}^3$;
- llll) $\text{CR}^3\text{R}^2\text{-NR}^3\text{R}^3$;
- mmmm) $\text{CH}_2\text{N(alkyl)R}^3$;
- nnnn) CH(alkyl)-NHR^3 ;
- oooo) $\text{C(alkyl)}_2\text{-NHR}^3$;
- pppp) $\text{C(alkyl)}_2\text{-N(alkyl)R}^3$;
- qqqq) $\text{CH}_2\text{-C(=W)H}$;
- rrrr) $\text{CH}_2\text{-C(=W)alkyl}$;
- ssss) $\text{CR}^3\text{R}^3\text{-C(=W)R}^3$;
- tttt) A-R^3 ;

uuuu) $C(R^3)_2-C(=W)R^3$;

vvvv) $CH_2-C(=W)H$;

www) $CH_2-C(=W)$ alkenyl;

xxxx) $CH(alkenyl)-C(=W)H$;

yyyy) $A-S(O)R^3$;

zzzz) $CH(NH)-S(O)_nR^3$; or

aaaaa) $A-N(NH)R^3$;

wherein said optional substitution comprises one or more of:

I. a substituted or unsubstituted heterocycle;

II. $C(=W)O$ -aryl;

III. $C(=W)O$ -alkyl;

IV. $C(=W)NH_2$;

V. $C(=W)NH$ -alkyl;

VI. $C(=W)NH$ -aryl;

VII. $C(=W)N$ -di-alkyl;

VIII. $C(=W)N(alkyl)$ -aryl;

IX. α -amino acid;

X. α -amino ester;

XI. α -amino-carboxamide;

XII. β -amino acid;

XIII. β -amino ester; or

XIV. β -amino-carboxamide;

XV. halo; or

XVI. cyano,

Each R^3 is:

- bb) H;
- cc) OH;
- dd) C_{1-6} alkyl;
- ee) C_{2-6} alkenyl;
- ff) C_{2-6} alkynyl;
- gg) alkoxy;
- hh) CF_3 ;
- ii) CN;
- jj) amino;
- kk) NR^2R^2 ;
- ll) O-alkyl;
- mm) O-alkenyl;
- nn) O-alkynyl;
- oo) $C(R^2)(R^2)-S(O)_nNH_2$;
- pp) $C(R^2)(R^2)-S(O)_nCF_3$;
- qq) $C(R^2)(R^2)-NH_2$;
- rr) A-heterocycle;
- ss) $C(R^2)(R^2)-NR^2R^2$;
- tt) $C(R^2)(R^2)-C(=W)R^2$;
- uu) aryl;
- vv) carbocycle;
- ww) heterocycle;
- xx) cycloalkyl;

yy) alkaryl;

zz) alkylheterocycle;

aaa) aralkyl; or

bbb) heterocycle-alkyl;

any of which may be unsubstituted or substituted with one or more of the following taken in any combination:

- I. halo;
- II. OH;
- III. OR^2 ;
- IV. SR^2 ;
- V. COOH;
- VI. carboxylic acid ester;
- VII. $C(=W)R^2$;
- VIII. $C(=W)OR^2$;
- IX. $C(=W)OR^3$;
- X. $C(=W)SR^2$;
- XI. $A-C(=W)NH_2$;
- XII. $C(=W)NR^2R^3$;
- XIII. NR^2R^2 ;
- XIV. NR^2R^3 ;
- XV. $NR^2-S(O)_nR^3$;
- XVI. $NR^2-C(=W)-C_{1-6}alkyl$;
- XVII. $S(O)_nR^3$;
- XVIII. $C_{1-6}alkoxy$;

- XIX. C₁₋₆ thioether;
- XX. amino acid residue;
- XXI. NH-A-(amino acid residue);
- XXII. C(=W)NH-A-(amino acid residue); and
- XXIII. wherein when said optional substitution comprises a substituted heterocycle, then substitution is selected from the group consisting of:

- i. C(=W)O-aryl;
 - ii. C(=W)O-alkyl;
 - iii. C(=W)NH₂;
 - iv. C(=W)NH-aryl;
 - v. C(=W)NH-alkyl;
 - vi. C(=W)N-di-alkyl;
 - vii. C(=W)N(alkyl)-aryl;
 - viii. α -amino acid;
 - ix. α -amino ester;
 - x. α -amino-carboxamide;
 - xi. β -amino acid;
 - xii. β -amino ester; or
 - xiii. β -amino-carboxamide;
- taken alone or in any combination;

n independently is 0, 1 or 2;

A is a disubstituted spacer selected from the group consisting of:

- i) C₁₋₆ alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- j) C₂₋₁₂ alkenylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- k) C₂₋₁₂ alkynylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- l) optionally substituted arylene;
- m) O-alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- n) aralkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- o) optionally substituted cycloalkyl; and
- p) optionally substituted heterocycle;

wherein "A" may be joined by any desired linkage or any combination of desired linkages.

Each R^{4'}, R^{5'}, R^{6'} and R^{7'} independently is:

- a. H;
- b. halogen;
- c. NO₂;
- d. CN;
- e. CF₃;
- f. OR²

- g. NR^2R^2 ;
- h. $\text{NHS(O)}_n\text{R}^2$;
- i. NHCO-C_{1-3} alkyl;
- j. $\text{S(O)}_n\text{R}^2$;
- k. aryl;
- l. heterocycle;
- m. C_{1-6} alkyl;
- n. C_{2-6} alkenyl;
- o. C_{2-6} alkynyl;
- p. $\text{C(=W)-S(O)}_n\text{R}^2$;
- q. $\text{C(=W)-S(O)}_n\text{-NR}^2\text{R}^2$;
- r. C(=W)-aryl ;
- s. C(=W)-alkyl ;
- t. C(=W)-heterocycle ; or
- u. $\text{C(=W)-NR}^2\text{R}^2$;

each of which optionally may be substituted with one or more of :

- I. OR^2 ;
- II. $\text{S(O)}_n\text{R}^2$;
- III. $\text{C(=W)-S(O)}_n\text{R}^2$;
- IV. $\text{C(=W)-S(O)}_n\text{-NR}^2\text{R}^2$;
- V. C(=W)-aryl ;
- VI. C(=W)-alkyl ;
- VII. C(=W)-heterocycle ;
- VIII. $\text{C(=W)NR}^2\text{R}^2$;

- IX. H;
- X. NO_2 ;
- XI. CN;
- XII. CF_3 ;
- XIII. halogen;
- XIV. $\text{NHS(O)}_n\text{R}^2$;
- XV. NHCO-C_{1-3} alkyl;
- XVI. aryl;
- XVII. heterocycle;
- XVIII. C_{1-6} alkyl;
- XIX. C_{2-6} alkenyl;
- XX. C_{2-6} alkynyl; or
- XXI. NR^2R^2 ; and

Each $\text{R}^{2''}$, $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ independently is:

- bb) H;
- cc) halogen;
- dd) NO_2 ;
- ee) CN;
- ff) OR^2 ;
- gg) SR^2 ;
- hh) NH_2 ;
- ii) NR^2R^3 ;
- jj) $\text{N(R}^2\text{)-C(=W)-C}_{1-4}$ alkyl;
- kk) $\text{N(R}^2\text{)-SO}_2\text{-C}_{1-4}$ alkyl;


- ll) C₁₋₆ alkyl;
- mm) C₂₋₆ alkenyl;
- nn) C₂₋₆ alkynyl;
- oo) aryl;
- pp) CF₃;
- qq) CR²R²-S(O)_n-R³;
- rr) CR²R²NR²R³;
- ss) C-OH;
- tt) CR²R²-C(=W)R²;
- uu) acyl;
- vv) C(=W)R²;
- ww) C(=W)OR²;
- xx) C(=W)SR²;
- yy) C(=W)-NR²R³;
- zz) C(=W)NH(CH₂)_p-(amino acid residue);
- aaa) amino residue; or
- bbb) A-(amino acid residue);

wherein any of the above optionally may be substituted; or

alternatively, R^{2''} or R^{6''} may be joined to Y to form an optionally substituted bicyclic or tricyclic phosphorylated heterocycle comprising 4-14 members.

- 15. The compound of claim 14 wherein R^{5'} is halogen, CH₃, CN, CF₃ or NO₂.
- ¹⁶ ~~22~~. The compound of claim 15 wherein R^{5'} is Cl.
- ¹⁷ ~~23~~. The compound of claim 14 wherein W is O or S.

- ¹⁸~~24~~. The compound of claim 14 wherein Z is optionally substituted amido or carboxamide.
- ¹⁹~~25~~. The compound of claim 14 wherein X is substituted or unsubstituted phenyl.
- ²⁰~~26~~. The compound of claim 14 wherein Y is $-O-C_{1-6}$ alkyl.
- ²¹~~27~~. The compound of claim 1 wherein the compound is of Formula (C):



or a pharmaceutically acceptable salt, prodrug, *N*-oxide, quaternary amine,

stereochemical isomer or tautomer thereof, wherein:

Y is:

- a. 3-14 membered carbocycle, aryl, heterocycle, any of which may comprise a monocyclic, bicyclic, tricyclic or spiro structure;
- b. H;
- c. OH;
- d. halogen;
- e. CF_3 ;
- f. C_{1-6} alkyl;
- g. C_{2-6} alkenyl;
- h. C_{2-6} alkynyl;
- i. alkylheterocycle;
- j. NH_2 ;

- k. NH-alkyl;
- l. N-dialkyl;
- m. NH-aryl;
- n. N-alkaryl;
- o. N-aralkyl;
- p. NH-heterocycle;
- q. N-alkyl-heterocycle;
- r. N-alkenyl-heterocycle;
- s. N-alkynyl-heterocycle;
- t. O-alkyl;
- u. O-alkenyl;
- v. O-alkynyl;
- w. O-alkylaryl;
- x. O-aryl;
- y. O-heterocycle;
- z. O-aralkyl;
- aa. O-carbocycle;
- bb. SR^2 ; or
- cc. NR^2R^3 ;

Z is:

- aaa) H;
- bbb) alkoxy;
- ccc) NO_2 ;
- ddd) $N(R^2)(R^3)$;

- eee) OR^2 ;
- fff) carboxamido;
- ggg) amido;
- hhh) acyl;
- iii) $\text{S(O)}_n\text{R}^2$;
- jjj) $\text{S(O)}_n\text{-NR}^2\text{R}^3$;
- kkk) C_{1-6} alkyl;
- lll) C_{2-6} alkenyl;
- mmm) C_{2-6} alkynyl;
- nnn) alkaryl;
- ooo) aralkyl;
- ppp) heterocycle;
- qqq) alkyl-heterocycle;
- rrr) aryl;
- sss) CN;
- ttt) C(=W)-R^2 ;
- uuu) $\text{C(=W)NH-C(R}^2\text{)(R}^2\text{)-C(=W)-N(R}^2\text{)(R}^2\text{)}$;
- vvv) $\text{C(=W)NH-P(=W)(R}^2\text{)-A-R}^2$;
- www) $\text{C(=W)NH-A-S(O)}_n\text{-NR}^2$;
- xxx) $\text{C(=W)NH-CR}^2\text{R}^3\text{-S(O)}_n\text{-NR}^2\text{R}^3$;
- yyy) $\text{C(=W)-NH-A-C(=W)-N(R}^2\text{)(R}^3\text{)}$;
- zzz) $\text{C(=W)-N(R}^2\text{)(R}^3\text{)}$;
- aaaa) C(=W)-NH-A-R^2 ;
- bbbb) C(=W)-NH-NH-R^2 ;

- cccc) $C(=W)-NH-C(R^2)(R^2)-C(=W)NH-C(R^2)(R^3)C(=W)-N(R^2)(R^3);$
- dddd) $C(=W)-NH-R^2;$
- eeee) $C(=W)-NH-A-C(=W)-NH-A-C(=W)-NH_2;$
- ffff) $C(R^2)(R^2)(R^3);$
- gggg) $C(R^2)(R^2)-NH-R^2;$
- hhhh) $A-S(O)_n-R^2;$
- iiii) $C(=W)-A-C(=W)-A-C(=W)R^3;$
- jjjj) $A-R^2;$
- kkkk) $C(=W)-(O)R^2;$
- llll) $C(=W)-A-C(=W)-NH_2;$
- mmmm) an amino acid residue;
- nnnn) $C(=W)-N(R^2)-A-(\text{amino acid residue});$
- oooo) $C(=W)-N(R^2)-A-(\text{amino acid residue})-C(=W)-R^2;$
- pppp) $C(=W)-\text{amino acid residue};$
- qqqq) $C(=W)-N(R^2)-A-(\text{amino acid residue})-A-C(=W)-R^2;$
- rrrr) $C(=W)-OR^3;$
- ssss) $C(=W)-S(R^2);$
- tttt) $C(=W)-NH-NH-R^2;$
- uuuu) $C(=W)-NH-N(R^2)-A-C(=W)R^2;$
- vvvv) $C(=W)-N(R^2)-C(=W)-R^3;$
- wwww) $C(=W)-A-NH-C(=W)R^2;$
- xxxx) $C(=W)-A-NH-C(=W)OR^3;$
- yyyy) $C(=W)-A-R^3;$

zzzz) $C(=W)-NH-NH-CH_2-C(=W)R^2$;

aaa) $P(=W)(R^2)(R^2)$; or

bbb) $A-P(=W)(R^2)(R^2)$;

wherein each of the foregoing Y and Z independently may be unsubstituted or substituted by one or more of:

nn) H;

oo) C_{1-6} alkyl;

pp) alkoxy;

qq) OH;

rr) oxo;

ss) halo;

tt) NR^2R^2 ;

uu) optionally substituted aryl;

vv) optionally substituted heterocyclyl;

ww) $O-C(=W)$ -alkyl;

xx) $C(=W)-OR^2$;

yy) CN;

zz) NO_2 ;

aaa) $NH-C(=W)$ -alkyl;

bbb) $NH-S(O)_n$ -alkyl;

ccc) $NH-S(O)_n-NR^2R^2$; or

ddd) C_{3-6} cycloalkyl;

W is:

h) O;

- i) S;
- j) NH;
- k) N-N(R²)(R²);
- l) N(R²);
- m) N-OH; or
- n) N-O-alkyl;

R¹ is:

- l) H;
- m) R²;
- n) C(=W)-R²;
- o) C(=W)-O(R²);
- p) C(=W)-S(R²);
- q) C(=W)-NH-R²;
- r) C(=W)-N(R²)(R²);
- s) C(=W)-NH-A-(amino acid residue);
- t) A-(amino acid residue)-R²;
- u) S(O)_n-R³; or
- v) S(O)₂-N(R²)(R²);

any of which optionally may be substituted by one or more:

- i. C₁₋₆ alkyl;
- ii. OH;
- iii. alkoxy;
- iv. aryl;
- v. halo;

- vi. CN;
- vii. NO₂; or
- viii. N(R²)(R²);

Each R² is:

- ccc) H;
- ddd) OH;
- eee) halogen;
- fff) optionally substituted, branched or unbranched alkyl;
- ggg) optionally substituted, branched or unbranched alkenyl;
- hhh) optionally substituted, branched or unbranched alkynyl;
- iii) 3-14 membered carbocycle;
- jjj) alkylheterocycle;
- kkk) acyl;
- lll) carboxamido;
- mmm) carbamoyl;
- nnn) alkoxy;
- ooo) optionally substituted aryl;
- ppp) optionally substituted aralkyl;
- qqq) optionally substituted alkylaryl;
- rrr) O-alkyl;
- sss) O-alkenyl;
- ttt) O-alkynyl;
- uuu) O-alkaryl;
- vvv) O-aralkyl;

www)	O-carbocycle;
xxx)	O-heterocycle;
yyy)	O-aryl;
zzz)	CF ₃ ;
aaaa)	CN;
bbbb)	S(O) _n -R ³ ;
cccc)	N(R ³)(R ³);
dddd)	NH-S(O) _n -R ³ ;
eeee)	NHC(=W)-aryl;
fff)	NHC(=W)alkyl;
gggg)	NHC(=W)-heterocycle;
hhhh)	CH ₂ -S(O) _n R ³ ;
iiii)	C(=W)R ³ ;
jjjj)	C(=W)NR ³ R ³ ;
kkkk)	C(alkyl) ₂ -S(O) _n R ³ ;
llll)	CH(alkyl)-S(O) _n R ³ ;
mmmm)	C(alkyl) ₂ -NH ₂ ;
nnnn)	CH(alkyl)-N(alkyl)R ³ ;
oooo)	CR ³ R ² -NR ³ R ³ ;
pppp)	CH ₂ N(alkyl)R ³ ;
qqqq)	CH(alkyl)-NHR ³ ;
rrrr)	C(alkyl) ₂ -NHR ³ ;
ssss)	C(alkyl) ₂ -N(alkyl)R ³ ;
tttt)	CH ₂ -C(=W)H;

- uuuu) $\text{CH}_2\text{-C(=W)alkyl}$;
- vvvv) $\text{CR}^3\text{R}^3\text{-C(=W)R}^3$;
- www) A-R^3 ;
- xxxx) $\text{C(R}^3)_2\text{-C(=W)R}^3$;
- yyyy) $\text{CH}_2\text{-C(=W)H}$;
- zzzz) $\text{CH}_2\text{-C(=W)alkenyl}$;
- aaaa) $\text{CH(alkenyl)-C(=W)H}$;
- bbbb) A-S(O)R^3 ;
- cccc) $\text{CH(NH)-S(O)}_n\text{R}^3$; or
- dddd) A-N(NH)R^3 ;

wherein said optional substitution comprises one or more of:

- i. a substituted or unsubstituted heterocycle;
- ii. C(=W)O-aryl ;
- iii. C(=W)O-alkyl ;
- iv. C(=W)NH_2 ;
- v. C(=W)NH-alkyl ;
- vi. C(=W)NH-aryl ;
- vii. C(=W)N-di-alkyl ;
- viii. $\text{C(=W)N(alkyl)-aryl}$;
- ix. α -amino acid;
- x. α -amino ester;
- xi. α -amino-carboxamide;
- xii. β -amino acid;
- xiii. β -amino ester; or

xiv. β -amino-carboxamide;

Each R^3 is:

- bb) H;
- cc) OH;
- dd) C_{1-6} alkyl;
- ee) C_{2-6} alkenyl;
- ff) C_{2-6} alkynyl;
- gg) alkoxy;
- hh) CF_3 ;
- ii) CN;
- jj) amino;
- kk) NR^2R^2 ;
- ll) O-alkyl;
- mm) O-alkenyl;
- nn) O-alkynyl;
- oo) $C(R^2)(R^2)-S(O)_nNH_2$;
- pp) $C(R^2)(R^2)-S(O)_nCF_3$;
- qq) $C(R^2)(R^2)-NH_2$;
- rr) A-heterocycle;
- ss) $C(R^2)(R^2)-NR^2R^2$;
- tt) $C(R^2)(R^2)-C(=W)R^2$;
- uu) aryl;
- vv) carbocycle;
- ww) heterocycle;

xx) cycloalkyl;

yy) alkaryl;

zz) alkylheterocycle;

aaa) aralkyl; or

bbb) heterocycle-alkyl;

any of which may be unsubstituted or substituted with one or more of the following taken in any combination:

- i) halo;
- ii) OH;
- iii) OR^2 ;
- iv) SR^2 ;
- v) COOH;
- vi) carboxylic acid ester;
- vii) $C(=W)R^2$;
- viii) $C(=W)OR^2$;
- ix) $C(=W)OR^3$;
- x) $C(=W)SR^2$;
- xi) $A-C(=W)NH_2$;
- xii) $C(=W)NR^2R^3$;
- xiii) NR^2R^2 ;
- xiv) NR^2R^3 ;
- xv) $NR^2-S(O)_nR^3$;
- xvi) $NR^2-C(=W)-C_{1-6}alkyl$;
- xvii) $S(O)_nR^3$;

- xviii) C₁₋₆alkoxy;
- xix) C₁₋₆ thioether;
- xx) amino acid residue;
- xxi) NH-A-(amino acid residue);
- xxii) C(=W)NH-A-(amino acid residue); and
- xxiii) wherein when said optional substitution comprises a substituted heterocycle, then substitution is selected from the group consisting of:

- I. C(=W)O-aryl;
- II. C(=W)O-alkyl;
- III. C(=W)NH₂;
- IV. C(=W)NH-aryl;
- V. C(=W)NH-alkyl;
- VI. C(=W)N-di-alkyl;
- VII. C(=W)N(alkyl)-aryl;
- VIII. α -amino acid;
- IX. α -amino ester;
- X. α -amino-carboxamide;
- XI. β -amino acid;
- XII. β -amino ester; or
- XIII. β -amino-carboxamide;

taken alone or in any combination;

n independently is 0, 1 or 2;

A is a disubstituted spacer selected from the group consisting of:

- a. C₁₋₆ alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- b. C₂₋₁₂ alkenylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- c. C₂₋₁₂ alkynylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- d. optionally substituted arylene;
- e. O-alkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- f. aralkylene, branched or unbranched, and optionally having one or more heteroatoms, aryl, cycloalkyl or heterocyclyl functions in or attached to the chain;
- g. optionally substituted cycloalkyl; and
- h. optionally substituted heterocycle;

wherein "A" may be joined by any desired linkage or any combination of desired linkages.

Each R^{4'}, R^{5'}, R^{6'} and R^{7'} independently is:

- a. H;
- b. halogen;
- c. NO₂;
- d. CN;
- e. CF₃;
- f. OR²

- g. NR^2R^2 ;
- h. $\text{NHS(O)}_n\text{R}^2$;
- i. NHCO-C_{1-3} alkyl;
- j. $\text{S(O)}_n\text{R}^2$;
- k. aryl;
- l. heterocycle;
- m. C_{1-6} alkyl;
- n. C_{2-6} alkenyl;
- o. C_{2-6} alkynyl;
- p. $\text{C(=W)-S(O)}_n\text{R}^2$;
- q. $\text{C(=W)-S(O)}_n\text{-NR}^2\text{R}^2$;
- r. C(=W)-aryl ;
- s. C(=W)-alkyl ;
- t. C(=W)-heterocycle ; or
- u. $\text{C(=W)-NR}^2\text{R}^2$;

each of which optionally may be substituted with one or more of :

- i. OR^2 ;
- ii. $\text{S(O)}_n\text{R}^2$;
- iii. $\text{C(=W)-S(O)}_n\text{R}^2$;
- iv. $\text{C(=W)-S(O)}_n\text{-NR}^2\text{R}^2$;
- v. C(=W)-aryl ;
- vi. C(=W)-alkyl ;
- vii. C(=W)-heterocycle ;
- viii. $\text{C(=W)NR}^2\text{R}^2$;

- ix. H;
- x. NO_2 ;
- xi. CN;
- xii. CF_3 ;
- xiii. halogen;
- xiv. $\text{NHS}(\text{O})_n\text{R}^2$;
- xv. NHCO-C_{1-3} alkyl;
- xvi. aryl;
- xvii. heterocycle;
- xviii. C_{1-6} alkyl;
- xix. C_{2-6} alkenyl;
- xx. C_{2-6} alkynyl; or
- xxi. NR^2R^2 .

Each $\text{R}^{3''}$, $\text{R}^{4''}$, $\text{R}^{5''}$, and $\text{R}^{6''}$ independently is:

- a. H;
- b. halogen;
- c. NO_2 ;
- d. CN;
- e. OR^2 ;
- f. SR^2 ;
- g. NH_2 ;
- h. NR^2R^3 ;
- i. $\text{N}(\text{R}^2)\text{-C(=W)-C}_{1-4}$ alkyl;
- j. $\text{N}(\text{R}^2)\text{-SO}_2\text{-C}_{1-4}$ alkyl;

- k. C_{1-6} alkyl;
- l. C_{2-6} alkenyl;
- m. C_{2-6} alkynyl;
- n. aryl;
- o. CF_3 ;
- p. $CR^2R^2-S(O)_n-R^3$;
- q. $CR^2R^2NR^2R^3$;
- r. $C-OH$;
- s. $CR^2R^2-C(=W)R^2$;
- t. acyl;
- u. $C(=W)R^2$;
- v. $C(=W)OR^2$;
- w. $C(=W)SR^2$;
- x. $C(=W)-NR^2R^3$;
- y. $C(=W)NH(CH_2)_p$ -(amino acid residue);
- z. amino residue; or
- aa. A-(amino acid residue);

wherein any of the above optionally may be substituted;

wherein ----- indicates the presence of a single or double bond, where:

when ----- indicates the presence of a double bond, each Y and T independently is:

- j. CR^2 ;
- k. N; or
- l. SR^2 ;

with the caveat that one of Y and T must be CR^2 ;

when ----- indicates the presence of a single bond, then each Y and T independently is:

- m. CR^2 ;
- n. O;
- o. NR^2 ; or
- p. SR^2 ;

with the caveat that one of Y and T must be CR^2 ; and

m is 1 or 2, with the proviso that m can only be 2 for T or $\text{Y} = \text{CR}^2$.

22. The compound of claim 21 wherein $\text{R}^{5'}$ is halogen, CH_3 , CN, CF_3 or NO_2 .
23. The compound of claim 22 wherein $\text{R}^{5'}$ is Cl.
24. The compound of claim 21 wherein Z is optionally substituted amido or carboxamide.
25. The compound of claim 21 wherein W is O or S.
26. The compound of claim 21 wherein Y is $-\text{O}-\text{C}_{1-6}$ alkyl.
27. A pharmaceutical composition comprising an HIV-treatment effective amount of a 3-phosphoindole compound or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-oxide or quaternary amine, with a pharmaceutically acceptable carrier or diluent
28. The composition of claim 27 wherein the 3-phosphoindoles is in the form of a phosphate, a phosphonate, a phosphorothioate, or a phosphoramidate.
29. The composition of claim 27 wherein the compound is a phosphate or a phosphonate.
30. The composition of claim 27 wherein the compound is further substituted at the 2 position.

31. The composition of claim 27 wherein the compound is substituted at the nitrogen of the indole.
32. The composition of claim 27 comprising an effective anti-HIV treatment amount of a compound of claims 1, 6, 14, or 21 or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-oxide or quaternary amine, with a pharmaceutically acceptable carrier or diluent.
33. The composition of claim 27 wherein the compound is in combination with at least one other anti-HIV agent, optionally with a pharmaceutically acceptable carrier or diluent.
34. The pharmaceutical composition of claim 33 wherein the other anti-HIV agent is a reverse transcriptase inhibitor.
35. The pharmaceutical composition of claim 33 wherein the other anti-HIV agent is a reverse transcriptase inhibitor.
36. The pharmaceutical composition of claim 35 wherein the reverse transcriptase inhibitor induces a mutation lysine 103 → asparagine and/or tyrosine 181 → cysteine in HIV reverse transcriptase.
37. Use of an effective amount of a 3-phosphoindole compound or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-oxide or quaternary amine, with a pharmaceutically acceptable carrier or diluent in the manufacture of a medicament for the treatment or prophylaxis of an HIV infection in a host.
38. The use of claim 37 wherein the 3-phosphoindoles is in the form of a phosphate, a phosphonate, a phosphorothioate, or a phosphoramidate.
39. The use of claim 37 wherein the compound is a phosphate or a phosphonate.
40. The use of claim 37 wherein the compound is further substituted at the 2 position.
41. The use of claim 37 wherein the compound is substituted at the nitrogen of the indole.
42. The use of claim 37 wherein the compound is a compound of one of claims 1, 9, or 16, or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-

oxide or quaternary amine, optionally with a pharmaceutically acceptable carrier or diluent.

43. The use of claim 37 wherein the host is a human.
44. The use of claim 37 wherein the compound is administered in combination and/or alternation with at least one other anti-HIV agent, optionally with a pharmaceutically acceptable carrier or diluent.
45. The use of claim 44 wherein the other anti-HIV agent is a reverse transcriptase inhibitor.
46. The use of claim 45 wherein the reverse transcriptase inhibitor induces a mutation lysine 103 → asparagine and/or tyrosine 181 → cysteine in HIV reverse transcriptase.
47. The use of claim 37, wherein the HIV has a mutation at lysine 103 → asparagine and/or tyrosine 181 → cysteine in HIV reverse transcriptase.
48. The use of claim 37 wherein the HIV is resistant to one or more reverse transcriptase inhibitor(s).
49. The use of claim 38 wherein the compound is administered in combination and/or alternation with at least one other anti-HIV agent, optionally with a pharmaceutically acceptable carrier or diluent.
50. Use of an anti-HIV effective treatment amount of a 3-phosphoindole compound or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-oxide or quaternary amine, optionally with a pharmaceutically acceptable carrier or diluent in the manufacture of a medicament for for salvage therapy in the treatment or prophylaxis of an HIV infection in a host..
51. The use of claim 50 wherein the compound is a compound of one of claims 1, 9, or 16, or its pharmaceutically acceptable salt, prodrug, stereoisomer, tautomer, *N*-oxide or quaternary amine.
52. The use of claim 50 wherein the compound is administered in combination and/or alternation with at least one other anti-HIV agent, optionally with a pharmaceutically acceptable carrier or diluent.