

5 **p-SUBSTITUTED ASYMMETRIC UREAS AND MEDICAL USES THEREOF**

CROSS-REFERENCE TO RELATED APPLICATIONS

 This application claims priority to U.S. Provisional Application Serial No. 61/949/664, filed March 7, 2014, which is herein incorporated by reference in its entirety.

10 **FIELD OF THE INVENTION**

 The present invention relates to novel asymmetric urea compounds, medical uses thereof, particularly in the treatment of medical conditions modulated by the ghrelin receptor.

BACKGROUND

 The growth hormone secretagogue receptor (GHS-R) regulates a number of physiological
15 processes, including growth hormone (GH) release, metabolism, and appetite. Ghrelin, a circulating hormone produced predominantly by endocrine cells in the stomach, is its endogenous ligand. Ghrelin is a 28 amino acid peptide with an acyl side chain required for biological activity (Kojima et al., Nature, 402, 656-660, 1999). Ghrelin has been shown to stimulate growth hormone (GH) release and to increase food intake when administered both
20 centrally and peripherally (Wren et al., Endocrinology, 141, 4325-4328, 2000).

 Endogenous levels of ghrelin rise on fasting and fall on re-feeding in humans (Cummings et al., Diabetes, 50, 1714-1719, 2001). Ghrelin also appears to play a role in maintaining long term energy balance and appetite regulation. Chronic administration of ghrelin in rodents leads to hyperphagia and weight gain that are independent of growth hormone secretion (Tschop et al.,
25 Nature, 407, 908-913, 2000). Circulating ghrelin levels decrease in response to chronic overfeeding and increase in response to chronic negative energy balance associated with anorexia or exercise. Obese people generally have low plasma ghrelin levels (Tschop et al., Diabetes, 50, 707-709, 2001) accordingly to the physiological response of the body in reducing calories intake. Intravenous ghrelin is effective in stimulating food intake in humans. A recent
30 study showed a 28% food intake increase from a buffet meal with a ghrelin infusion compared with saline control (Wren et al., J. Clin. Endocrinology and Metabolism, 86, 5992, 2001).

 In view of the above experimental evidence, compounds that modulate ghrelin receptor activity have been proposed for preventing and/or treating disorders associated with ghrelin receptor physiology. For example, antagonists at ghrelin receptor might one day be developed to


5 reduce appetite, reduce food intake, induce weight loss and treat obesity without affecting or
reducing the circulating growth hormone levels. On the other hand, agonists at ghrelin receptor
might also be developed for stimulating food intake and thus be useful in treating eating
disorders, for example anorexia nervosa, or in treating cachexia resulting from cancer, AIDS or
Chronic Obstructive Pulmonary Disease (COPD). Ghrelin agonists may also be useful as
10 gastroprokinetic agents which can enhance gastrointestinal motility by increasing the frequency
of contractions in the small intestine or making them stronger, but without disrupting their
rhythm. Gastroprokinetic agents are used to relieve gastrointestinal symptoms such as abdominal
discomfort, bloating, constipation, heart burn, nausea, and vomiting, and are used to treat a
number of gastrointestinal disorders, including but not limiting to, irritable bowel syndrome,
15 gastritis, acid reflux disease, gastroparesis, and functional dyspepsia. Furthermore, compounds
that modulate ghrelin receptor activity might also be used to prevent or treat diseases related to
substance abuse, for example, alcohol or drug (e.g., amphetamines, barbiturates,
benzodiazepines, cocaine, methaqualone, and opioids) abuse, which refers to a maladaptive
pattern of use of a substance that is not considered dependent.

20 A number of compounds acting on the ghrelin receptor have been reported in the
literature. YIL-781, for example, is a small molecule ghrelin receptor antagonist from Bayer that
reportedly improves glucose tolerance, suppresses appetite and promotes weigh loss (Esler et al.,
Endocrinology 148 (11):5175-5185); LY444711 is an orally active ghrelin receptor agonist from
Lilly that reportedly induces adiposity by stimulating food consumption and sparing fat
25 utilization (Bioorg. & Med. Chem. Lett., 2004, 14, 5873-5876); anamorelin is an orally available
ghrelin receptor small molecule agonist from Helsinn Therapeutics that is in clinical trials for the
treatment of anorexia and cachexia in cancer patients. Ghrelin receptor agonists and antagonists
based on asymmetric ureas are disclosed in US 2012/0220629, which is incorporated herein by
reference in its entirety. Other small molecule ghrelin receptor modulators can be found in WO
30 2008/092681, US 2009/0253673, WO 2008/148853, WO 2008/148856, US 2007/0270473 and
US 2009/0186870.

In view of the above, it is desirable to find new compounds which modulate ghrelin
receptor activity.

SUMMARY

35 The present invention provides compounds of Formula I:



I,

with X, Z, R¹-R⁸, r, s, and n as defined herein, and pharmaceutically acceptable salts thereof.

Compounds of Formula I, also referred to herein as asymmetric ureas, are particularly useful for preventing and/or treating diseases that are pathophysiologically related to the ghrelin receptor in a subject. Accordingly, in another embodiment the invention provides a method of treating a disease that is mediated by the ghrelin receptor, comprising administering to said subject a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Also disclosed are pharmaceutical compositions for preventing and/or treating diseases which are pathophysiologically related to ghrelin receptor in a subject, comprising a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows Highly Palatable Food (HPF) intake in rats at different times after initial access to HPF. The values shown are the mean $\pm$ S.E.M. of HPF intake. Statistical differences from controls (non-Restricted + non-Stressed; NR + NS): ** P < 0.01.

Figure 2 shows the effect of Topiramate (60 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S (Restricted and Stressed) vehicle and R + S treated rats: *P < 0.05; ** P < 0.01.

Figure 3 shows the effect of compound H0816 (3 and 30 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S vehicle and R + S treated rats: *P < 0.05.

Figure 4 shows the effect of compound H0860 (3 and 30 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Statistical difference from vehicle-treated rats was not statistically significant.

5 Figure 5 shows the effect of compound H0847 (3 and 30 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S vehicle and R + S treated rats: ** P < 0.01; * P < 0.05.

 Figure 6 shows the effect of compound H0900 (3 and 30 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S vehicle and R + S treated rats: ** P < 0.01; * P < 0.05.

 Figure 7 shows the effect of Topiramate, compounds H0816, H0860, H0847H0900 and vehicle on 2 h (A) and 24 h (B) chow food intake during and after a binge eating test. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S vehicle and R + S treated rats: * P < 0.05, ** P < 0.01.

15 Figure 8 shows the effect of H0816 (3, 10 and 30 mg/kg) or vehicle in a rat model of binge eating. The values shown are the mean $\pm$ S.E.M. of HPF intake. Difference between R + S vehicle and R + S treated rats: * P < 0.05; ** P < 0.05.

 Figure 9 shows the effect of compound H0847 on alcohol self-administration in msP rats.

 Figure 10 shows the effect of compound H0860 on alcohol self-administration in msP rats.


 Figure 11 shows the effect of compound H0816 on alcohol self-administration in msP rats.

 Figure 12 shows the effect of compound H0900 on alcohol self-administration in msP rats.

25 **DETAILED DESCRIPTION**

 Before the present compounds, compositions, articles, devices, and/or methods are disclosed and described, it is to be understood that they are not limited to specific synthetic methods or specific treatment methods unless otherwise specified, or to particular reagents unless otherwise specified, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

 In a first principal embodiment, the present invention provides compounds of Formula I:



I,

or a pharmaceutically acceptable salt thereof, wherein:

a dashed line indicates an optional bond;

X is CH or N;

10 Z is NR^9 , $\text{CR}^{10}\text{R}^{11}$, or O;

R^1 is H, C_{1-6} alkyl, benzyl, OH, or C_{1-6} alkoxy, wherein said C_{1-6} alkyl, benzyl, or C_{1-6} alkoxy is optionally substituted with 1-3 substituents selected from halo, OH, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} hydroxyalkyl, $\text{CO}(\text{C}_{1-6}$ alkyl), CHO, CO_2H , $\text{CO}_2(\text{C}_{1-6}$ alkyl), and C_{1-6} haloalkyl;

R^2 is H or C_{1-6} alkyl;

15 R^3 and R^4 are each, independently, H, CN, halo, CHO, or CO_2H , or optionally substituted C_{1-6} alkyl, C_{1-6} hydroxyalkyl, C_{1-6} alkylcycloalkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, $\text{CO}(\text{C}_{1-6}$ alkyl), $\text{CO}_2(\text{C}_{1-6}$ alkyl), or $\text{CONR}^{12}\text{R}^{13}$;

or R^3 and R^4 taken together with the C atom to which they are attached form a 3-6-membered ring;

20 R^5 is halo, CN, CHO, CO_2H , $\text{CO}(\text{C}_{1-6}$ alkyl), $\text{CO}_2(\text{C}_{1-6}$ alkyl), $\text{NR}^{14}\text{R}^{15}$, $\text{NHCONR}^{14}\text{R}^{15}$, $\text{CONR}^{14}\text{R}^{15}$, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} hydroxyalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl, wherein said $\text{CO}(\text{C}_{1-6}$ alkyl), $\text{CO}_2(\text{C}_{1-6}$ alkyl), $\text{NR}^{14}\text{R}^{15}$, $\text{NHCONR}^{14}\text{R}^{15}$, $\text{CONR}^{14}\text{R}^{15}$, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} hydroxyalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally
25 substituted with 1-3 substituents selected from halo, CN, OH, NO_2 , $\text{Si}(\text{CH}_3)_4$, CHO, and CO_2H , or optionally substituted $\text{CO}(\text{C}_{1-6}$ alkyl), $\text{CO}_2(\text{C}_{1-6}$ alkyl), $\text{NR}^{14}\text{R}^{15}$, $\text{NHCONR}^{14}\text{R}^{15}$, $\text{CONR}^{14}\text{R}^{15}$, $\text{CH}=\text{NOH}$, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} hydroxyalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl;

R^6 is absent or H;

30 R^7 is H, CN, or halo;

5 or two R^7 can be taken together with the atoms to which they are attached form a 5-6-membered ring;

or R^5 and R^7 taken together with the atoms to which they are attached form an optionally substituted 5-6-membered ring;

R^8 is H or C_{1-6} alkyl;

10 R^9 is H, C_{1-6} alkyl, $CO(C_{1-6}$ alkyl), CHO, CO_2H , or $CO_2(C_{1-6}$ alkyl);

R^{10} and R^{11} are each, independently, H, C_{1-6} alkyl, or halo;

R^{12} and R^{13} are each, independently, H or C_{1-6} alkyl;

R^{14} and R^{15} are each, independently H, C_{1-6} alkyl, $CO(C_{1-6}$ alkyl), $CO(heteroaryl)$, heteroaryl, or cycloalkyl;

15 r is 1 or 2;

s is 0-4; and

n is 0-3.

In the first principal embodiment, as well as the second and third principal embodiments discussed below, in one subembodiment X is CH.

20 In the first, second and third principal embodiments, in one subembodiment, X is N.

In the first, second and third principal embodiments, in one subembodiment, Z is NR^9 .

In the first, second and third principal embodiments, in one subembodiment, Z is $N(C_{1-6}$ alkyl).

In the first, second and third principal embodiments, in one subembodiment, Z is NCH_3 .

25 In the first, second and third principal embodiments, in one subembodiment, Z is $CR^{10}R^{11}$.

In the first, second and third principal embodiments, in one subembodiment, Z is CF_2

In the first, second and third principal embodiments, in one subembodiment, Z is O.

In the first, second and third principal embodiments, in one subembodiment, R^1 is C_{1-6}

30 alkyl.

In the first, second and third principal embodiments, in one subembodiment, R^1 is CH_3 .

In the first, second and third principal embodiments, in one subembodiment, R^1 is benzyl.

In the first, second and third principal embodiments, in one subembodiment, said benzyl is optionally substituted with $CO_2(C_{1-6}$ alkyl) or C_{1-6} hydroxyalkyl.

35 In the first, second and third principal embodiments, in one subembodiment, R^1 is OH.

5 In the first, second and third principal embodiments, in one subembodiment, R¹ is C₁₋₆ alkoxy.

In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ alkoxy is methoxy, ethoxy or propoxy.

In the first, second and third principal embodiments, in one subembodiment, R² is H.

10 In the first, second and third principal embodiments, in one subembodiment, R³ and R⁴ are each, independently selected from C₁₋₆ alkyl, CN, C₁₋₆ alkylcycloalkyl, C₁₋₆ hydroxyalkyl, CO₂(C₁₋₆ alkyl), C₁₋₆ haloalkyl and CONH₂, .

In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ alkyl is methyl or ethyl.

15 In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ alkylcycloalkyl is C₁ alkylcyclopropyl.

In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ hydroxyalkyl is C₁ hydroxyalkyl optionally substituted with a substituted or unsubstituted benzyl group.

20 In the first, second and third principal embodiments, in one subembodiment, said CO₂(C₁₋₆ alkyl) is CO₂CH₃.

In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ haloalkyl is CF₃.

25 In the first, second and third principal embodiments, in one subembodiment, R³ and R⁴ taken together with the C atom to which they are attached form a 3-6-membered ring.

In the first, second and third principal embodiments, in one subembodiment, R³ and R⁴ are taken together with the C atom to which they are attached to form a cyclopropyl ring.

In the first, second and third principal embodiments, in one subembodiment, R³ and R⁴ are taken together with the C atom to which they are attached form a tetrahydropyranyl ring.

30 In the first, second and third principal embodiments, in one subembodiment, R⁵ is halo, CN, CHO, CO₂H, CO(C₁₋₆ alkyl), CO₂(C₁₋₆ alkyl), NR¹⁴R¹⁵, NHCONR¹⁴R¹⁵, CONR¹⁴R¹⁵, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl, wherein said CO(C₁₋₆ alkyl), CO₂(C₁₋₆ alkyl), NR¹⁴R¹⁵, NHCONR¹⁴R¹⁵, CONR¹⁴R¹⁵, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted

35

- 5 with 1-3 substituents selected from halo, CN, OH, NO₂, Si(CH₃)₄, CHO, and CO₂H, or optionally substituted CO(C₁₋₆ alkyl), CO₂(C₁₋₆ alkyl), NR¹⁴R¹⁵, NHCONR¹⁴R¹⁵, CONR¹⁴R¹⁵, CH=NOH, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl;
- In some embodiments R⁵ is not H;
- 10 In some embodiments, R⁵ is not alkoxy;
- In some embodiments, R⁵ is not methoxy;
- In some embodiments, R⁵ is not OH;
- In some embodiments, R⁵ is not halo;
- In some embodiments, R⁵ is not fluoro;
- 15 In some embodiments, R⁵ is not chloro;
- In some embodiments, R⁵ is not SO₂Me;
- In some embodiments, R⁵ is not amino;
- In some embodiments, R⁵ is not NHAc;
- In some embodiments, R⁵ is not N(Me)₂;
- 20 In some embodiments, R⁵ is not alkyl;
- In some embodiments, R⁵ is not methyl;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is halo;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is CN;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is CHO;
- 25 In the first, second and third principal embodiments, in one subembodiment, R⁵ is CO₂H;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is CO(C₁₋₆ alkyl);
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is CO₂(C₁₋₆ alkyl);
- 30 In the first, second and third principal embodiments, in one subembodiment, R⁵ is NR¹⁴R¹⁵;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is NHCONR¹⁴R¹⁵;
- In the first, second and third principal embodiments, in one subembodiment, R⁵ is CONR¹⁴R¹⁵;
- 35

- 5 In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{1-6} alkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{1-6} alkoxy;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{1-6}
- 10 haloalkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{1-6} hydroxyalkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{2-6} alkenyl;
- 15 In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{2-6} alkynyl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is aryl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is cycloalkyl;
- 20 In the first, second and third principal embodiments, in one subembodiment, R^5 is heteroaryl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is heterocycloalkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^5 is C_{1-6}
- 25 haloalkyl, heteroaryl, aryl, halo, C_{1-6} alkoxy, $CO_2(C_{1-6}$ alkyl), C_{2-6} alkenyl, C_{2-6} alkynyl, cycloalkyl, or heterocycloalkyl,
- In the first, second and third principal embodiments, in one subembodiment, said cycloalkyl is cyclopropyl, cyclohexanyl or cyclohexenyl.
- In the first, second and third principal embodiments, in one subembodiment, said C_{1-6}
- 30 haloalkyl is CHF_2 .
- In the first, second and third principal embodiments, in one subembodiment, said heteroaryl is pyridyl, pyridazinyl, pyrimidinyl, triazinyl, thiophenyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, oxadiazolyl or furanyl,
- In the first, second and third principal embodiments, in one subembodiment, said aryl is
- 35 phenyl.

5 In the first, second and third principal embodiments, in one subembodiment, said halo is Cl or I.

In the first, second and third principal embodiments, in one subembodiment, said C₁₋₆ alkoxy is methoxy.

10 In the first, second and third principal embodiments, in one subembodiment, said CO₂(C₁₋₆ alkyl) is CO₂Me.

In the first, second and third principal embodiments, in one subembodiment, said C₂₋₆ alkynyl is C₂ alkynyl.

In the first, second and third principal embodiments, in one subembodiment, said C₂₋₆ alkenyl is C₂ alkenyl.

15 In the first, second and third principal embodiments, in one subembodiment, R⁶ is absent.

In the first, second and third principal embodiments, in one subembodiment, R⁶ is H.

In the first, second and third principal embodiments, in one subembodiment, R⁷ is halo.

In the first, second and third principal embodiments, in one subembodiment, said halo is Cl or F.

20 In the first, second and third principal embodiments, in one subembodiment, 2 R⁷ come together to form a phenyl group.

In the first, second and third principal embodiments, in one subembodiment, R⁵ and R⁷ come together to form a 5-membered heterocyclic ring.

In the first, second and third principal embodiments, in one subembodiment, R⁸ is H.

25 In the first, second and third principal embodiments, in one subembodiment, R⁸ is C₁₋₆ alkyl.

In the first, second and third principal embodiments, in one subembodiment, R⁸ is methyl.

In the first, second and third principal embodiments, in one subembodiment, R¹⁰ is H;

30 In the first, second and third principal embodiments, in one subembodiment, R¹⁰ is C₁₋₆ alkyl;

In the first, second and third principal embodiments, in one subembodiment, R¹⁰ is halo;

In the first, second and third principal embodiments, in one subembodiment, R¹¹ is H;


35 In the first, second and third principal embodiments, in one subembodiment, R¹¹ is C₁₋₆ alkyl;

- 5 In the first, second and third principal embodiments, in one subembodiment, R^{11} is halo;
In the first, second and third principal embodiments, in one subembodiment, R^{12} is H;
In the first, second and third principal embodiments, in one subembodiment, R^{12} is C_{1-6}
alkyl;
- 10 In the first, second and third principal embodiments, in one subembodiment, R^{13} is H;
In the first, second and third principal embodiments, in one subembodiment, R^{13} is C_{1-6}
alkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^{14} is H;
In the first, second and third principal embodiments, in one subembodiment, R^{14} is C_{1-6}
alkyl;
- 15 In the first, second and third principal embodiments, in one subembodiment, R^{14} is
 $CO(C_{1-6}$ alkyl);
In the first, second and third principal embodiments, in one subembodiment, R^{14} is
 $CO(heteroaryl)$;
- 20 In the first, second and third principal embodiments, in one subembodiment, R^{14} is
heteroaryl;
In the first, second and third principal embodiments, in one subembodiment, R^{14} is
cycloalkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^{15} is H;
In the first, second and third principal embodiments, in one subembodiment, R^{15} is C_{1-6}
25 alkyl;
- In the first, second and third principal embodiments, in one subembodiment, R^{15} is
 $CO(C_{1-6}$ alkyl);
In the first, second and third principal embodiments, in one subembodiment, R^{15} is
 $CO(heteroaryl)$;
- 30 In the first, second and third principal embodiments, in one subembodiment, R^{15} is
heteroaryl;
In the first, second and third principal embodiments, in one subembodiment, R^{15} is
cycloalkyl;
- In the first, second and third principal embodiments, in one subembodiment, r is 1;
- 35 In the first, second and third principal embodiments, in one subembodiment, r is 2;

- 5 In the first, second and third principal embodiments, in one subembodiment, s is 0;
 In the first, second and third principal embodiments, in one subembodiment, s is 1;
 In the first, second and third principal embodiments, in one subembodiment, s is 2;
 In the first, second and third principal embodiments, in one subembodiment, s is 3;
 In the first, second and third principal embodiments, in one subembodiment, s is 4;
 10 In the first, second and third principal embodiments, in one subembodiment, n is 0;
 In the first, second and third principal embodiments, in one subembodiment, n is 1;
 In the first, second and third principal embodiments, in one subembodiment, n is 2;
 In the first, second and third principal embodiments, in one subembodiment, n is 3

15

In a second principal embodiment, the compounds have the structure of Formula II:




II,

or a pharmaceutically acceptable salt thereof.

20

In a third principal embodiment, the compounds have the structure of Formula III:




III,


or a pharmaceutically acceptable salt thereof.

In fourth and fifth principal embodiments, the compounds have the structure of Formula

25 IIIa or IIIb:



IIIa



IIIb,

or a pharmaceutically acceptable salt thereof, wherein:

R^{16} is H, cyclopropyl or thiazolyl; and









R^{17} is H or halo.

- 10 In some forms, the compounds as presently disclosed are compounds of Formula I, or pharmaceutically acceptable salts thereof, wherein the compound of Formula I is a compound selected from the group consisting of:

Compound No.	Chemical Structure	Chemical Name
H0494		3-(1-(2,3-dichloro-4-cyclopropylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0621		3-(1-(2,3-dichloro-4-(difluoromethyl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0496		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0617		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0539		methyl 4-((3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)ureido)methyl)benzoate
H0546		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(4-(hydroxymethyl)benzyl)-1-(1-methylpiperidin-4-yl)urea
H0526		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0527		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(1,3-dimethylpiperidin-4-yl)-1-methylurea
H0497		3-(1-(2,3-dichloro-4-(pyridin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0650		3-(1-(2,3-dichloro-4-(pyridin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0849		3-(1-(2,3-dichloro-4-(5-cyclopropylpyridin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0578		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(pyridin-4-yl)naphthalen-1-yl)ethyl)urea








H0511		3-(1-(2,3-dichloro-4-(6-methoxypyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0820		3-(1-(2,3-dichloro-4-(6-cyclopropylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0613		3-(1-(2,3-dichloro-4-(5-cyanopyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0614		3-(1-(2,3-dichloro-4-(5-fluoropyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0635		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)nicotinate
H0636		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0637		3-(1-(2,3-dichloro-4-(5-(difluoromethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0638		3-(1-(2,3-dichloro-4-(5-(fluoromethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0639		3-(1-(2,3-dichloro-4-(5-methylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0642		3-(1-(2,3-dichloro-4-(5-formylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0704		3-(1-(4-(5-aminopyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0705		3-(1-(2,3-dichloro-4-(5-(cyclopent-1-en-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0707		3-(1-(4-(5-(1H-pyrazol-4-yl)pyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0711		3-(1-(4-(5-(1H-imidazol-4-yl)pyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0716		3-(1-(2,3-dichloro-4-(5-(thiazol-5-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0717		3-(1-(2,3-dichloro-4-(5-(thiophen-2-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0718		3-(1-(2,3-dichloro-4-(5-cyclopentylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0719		3-(1-(2,3-dichloro-4-(5-(pyrrolidin-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0712		N-(5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyridin-3-yl)acetamide
H0708		3-(1-(2,3-dichloro-4-(5-(methoxymethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0714		3-(1-(2,3-dichloro-4-(5-(2-methoxyethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0715		3-(1-(2,3-dichloro-4-(5-ethylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0706		3-(1-(2,3-dichloro-4-(5-vinylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0710		3-(1-(2,3-dichloro-4-(5-ethynylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0666		3-((4-(5-(1-methyl-1-(1-methylpiperidin-4-yl)urea)methyl)-2,3-dichlorophenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0739		3-((4-(5-(1H-pyrrol-2-yl)pyridin-3-yl)-2,3-dichlorophenyl)(cyano)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0667		3-(1-(2,3-dichloro-4-(5-cyanopyridin-3-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0821		3-(1-(2,3-dichloro-4-(5-cyano-6-(4-methylpiperazin-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0646		(E)-3-(1-(2,3-dichloro-4-(5-((hydroxyimino)methyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0720		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0721		3-(1-(4-(5-aminopyridin-3-yl)-2,3-dichlorophenyl)-2-cyclopropylethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0516		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0579		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(pyrimidin-5-yl)naphthalen-1-yl)ethyl)urea
H0649		3-(1-(2,3-dichloro-4-(2-methoxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0797		3-(1-(2,3-dichloro-4-(2-hydroxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0798		3-(1-(4-(2-aminopyrimidin-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0799		3-(1-(2,3-dichloro-4-(2-(4-methylpiperazin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0800		3-(1-(2,3-dichloro-4-(2-fluoropyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0801		3-(1-(2,3-dichloro-4-(2-chloropyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0802		3-(1-(2,3-dichloro-4-(2-cyanopyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0803		3-(1-(4-(2-(1H-imidazol-1-yl)pyrimidin-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0804		3-(1-(2,3-dichloro-4-(2-(dimethylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0805		3-(1-(2,3-dichloro-4-(2-(cyclopropylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0806		3-(1-(2,3-dichloro-4-(2-(methylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0807		N-(5-(2,3-dichloro-4-(1-(3-methyl-1-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrimidin-2-yl)cyclopropanecarboxamide
H0854		3-(1-(2,3-dichloro-4-(2-cyclopropylpyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0813		3-(1-(2,3-dichloro-4-(2-(pyrrolidin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0814		3-(1-(2,3-dichloro-4-(2-(4-ethyl-3-oxopiperazin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0703		3-(1-cyano-1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0709		3-(cyano(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)methyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0584		1-cyclohexyl-3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methylurea
H0586		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(tetrahydro-2H-pyran-4-yl)urea
H0587		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-(4,4-difluorocyclohexyl)-1-methylurea









H0588		1-(1-acetylpiperidin-4-yl)-3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methylurea
H0663		3-(1-(2,3-dichloro-4-(2,4-dimethoxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0620		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-((3-(hydroxymethyl)benzyl)oxy)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0624		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0662		methyl 2-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0670		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0673		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)cyclopropyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0727		3-(4-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)tetrahydro-2H-pyran-4-yl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0631		3-(cyano(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0686		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0619		3-(1-(2,3-dichloro-4-(pyrimidin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0768		3-(1-(2,3-dichloro-4-(pyrimidin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0808		3-(1-(2,3-dichloro-4-(6-methylpyrimidin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0700		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0816		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0817		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0722		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0741		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0752		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea
H0743		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0750		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea
H0756		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea

H0761		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0781		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0782		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0824		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0890		3-((S)-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0858		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0865		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea








H0825		1-benzyl-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)urea
H0826		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0889		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0896		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0827		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)-1-propylurea
H0829		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0859		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0860		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)








H0922		methyl 2-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0924		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0830		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0899		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0900		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0909		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0856		3-(1-(3-chloro-2-fluoro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0837		3-((S)-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea (diastomeric mixture)

H0861		3-((S)-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea (single diastereoisomer)
H0862		3-((S)-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea (single diastereoisomer)
H0857		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0871		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0874		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0853		N-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-2-(4-methylpiperazin-1-yl)propanamide
H0815		3-(1-(2,3-dichloro-4-(6-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0831		3-(1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0843		3-(1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0844		3-(2-cyclopropyl-1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0738		3-(1-(2,3-dichloro-4-(6-methoxypyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0780		3-(1-(4-(6-aminopyrazin-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0786		3-(1-(2,3-dichloro-4-(6-(chloromethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0791		3-(1-(2,3-dichloro-4-(6-chloropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0795		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea







H0847		(S)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0848		(R)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0863		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0908		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0864		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0872		3-((S)-1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0840		3-(1-(2,3-dichloro-4-(3-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0910		3-(1-(2,3-dichloro-4-(6-(trifluoromethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0788		3-(1-(2,3-dichloro-4-(6-cyanopyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0789		methyl 6-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxylate
H0760		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxamide
H0769		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxylate
H0771		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N,N-dimethylpyrazine-2-carboxamide
H0770		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0828		3-(1-(2,3-dichloro-4-(quinoxalin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0822		3-(1-(2,3-dichloro-4-(5-(4-methylpiperazin-1-yl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0850		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(pyrazin-2-yl)naphthalen-1-yl)ethyl)urea
H0881		3-(1-(4,5-dichloro-6-(pyrazin-2-yl)pyridin-3-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0729		3-(1-(2,3-dichloro-4-(pyridazin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0783		3-(1-(2,3-dichloro-4-(pyridazin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0793		3-(1-(2,3-dichloro-4-(1,2,4-triazin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0796		3-(1-(2,3-dichloro-4-(4,6-dimorpholino-1,3,5-triazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0498		3-(1-(2,3-dichloro-4-(thiophen-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0531		3-(1-(2,3-dichloro-4-(thiophen-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0594		3-(1-(2,3-dichloro-4-(thiophen-3-yl)phenyl)-2-((3-(hydroxymethyl)benzyl)oxy)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0644		3-(cyano(2,3-dichloro-4-(thiophen-3-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0536		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0563		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0564		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)

H0627		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0660		methyl 2-(2,3-dichloro-4-(thiophen-2-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0661		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0672		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)cyclopropyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0651		3-(1-(2,3-dichloro-4-(5-formylthiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0653		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0668		3-(1-(2,3-dichloro-4-(5-(fluoromethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0654		3-(1-(2,3-dichloro-4-(5-(difluoromethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0655		3-(1-(4-(5-acetylthiophen-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0691		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxamide
H0728		5-(2,3-dichloro-4-(2-cyclopropyl-1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxamide
H0726		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N,N-dimethylthiophene-2-carboxamide
H0689		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxylic acid
H0692		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N-methoxy-N-methylthiophene-2-carboxamide

H0656		3-(1-(2,3-dichloro-4-(5-(1-hydroxyethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0652		3-(1-(2,3-dichloro-4-(5-formylthiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0713		3-(1-(2,3-dichloro-4-(5-cyanothiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0688		3-(1-(4-(3-acetylthiophen-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0774		2-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-3-carboxamide
H0664		3-(1-(2,3-dichloro-4-(3-(hydroxymethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0535		3-(1-(2,3-dichloro-4-(1H-pyrrol-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0499		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea









H0693		3-(1-(2,3-dichloro-4-(1-(2-hydroxyethyl)-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0694		3-(1-(2,3-dichloro-4-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0657		3-(cyano(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0553		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(4-(hydroxymethyl)benzyl)-1-(1-methylpiperidin-4-yl)urea
H0842		3-(1-(2,3-dichloro-4-(1-cyclopropyl-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0542		3-(1-(2,3-dichloro-4-(1H-imidazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0568		3-(1-(2,3-dichloro-4-(thiazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0794		3-(1-(4-(2-aminothiazol-4-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0841		3-(1-(2,3-dichloro-4-(2-cyclopropylthiazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0792		3-(1-(4-(2-aminothiazol-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0569		3-(1-(2,3-dichloro-4-(oxazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0565		3-(1-(2,3-dichloro-4-(1H-1,2,3-triazol-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0604		3-(1-(2,3-dichloro-4-(1H-1,2,3-triazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0595		3-(1-(2,3-dichloro-4-(1,3,4-oxadiazol-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0596		3-(1-(2,3-dichloro-4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0851		3-(1-(2,3-dichloro-4-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0537		methyl 4-((3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)ureido)methyl)benzoate
H0529		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0528		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(1,3-dimethylpiperidin-4-yl)-1-methylurea
H0501		3-(1-(2,3-dichloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0507		3-(1-(2,3-dichloro-4-(furan-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0665		3-(1-(2,3-dichloro-4-(5-methylfuran-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0508		3-(1-(2,3-dichloro-4'-methoxy-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0509		3-(1-(2,3-dichloro-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0510		3-(1-(3'-amino-2,3-dichloro-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0606		3-(1-(2,3-dichloro-3'-methoxy-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0810		3-(1-(2,3-dichloro-3'-fluoro-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0696		3-(1-(2,3-dichloro-3'-fluoro-5'-(hydroxymethyl)-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0611		3-(1-(2,3-dichloro-3',5'-dimethoxy-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0612		2',3'-dichloro-4'-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)-[1,1'-biphenyl]-3-carboxamide
H0615		2',3'-dichloro-4'-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)-[1,1'-biphenyl]-4-carboxamide










H0809		3-(1-(2,3-dichloro-4'-cyano-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0699		3-(1-(2,3-dichloro-4-(5-(cyanomethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0607		3-(1-(2,3-dichloro-4-(5-methoxypyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0695		3-(1-(4-(5-bromopyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0635		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)nicotinate
H0690		3-(1-(4-(5-acetylpyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0735		3-(1-(2,3-dichloro-4-(5-(pyrimidin-2-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0746		3-(1-(2,3-dichloro-4-(5-(furan-3-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0747		3-(1-(2,3-dichloro-4-(5-(thiophen-3-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0748		3-(1-(2,3-dichloro-4-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0765		3-(1-(2,3-dichloro-4-(5-(cyclopropyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0766		3-(1-(2,3-dichloro-4-(5-(nitro)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0608		3-(1-(2,3-dichloro-4-(6-(isopropoxy)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0616		3-(1-(2,3-dichloro-4-(6-(cyan)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0618		3-(1-(2,3-dichloro-4-(6-(fluor)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0623		3-(1-(2,3-dichloro-4-(1H-pyrazolo[3,4-b]pyridin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0610		3-(1-(2,3-dichloro-3'-cyano-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0517		3-(1-(4'-amino-2,3-dichloro-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0518		3-(1-(2,3-dichloro-4'-(dimethylamino)-[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0512		3-(1-(2,3-dichloro-4-(1-methyl-1H-indazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0513		3-(1-(2,3-dichloro-4-(1H-indazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0514		3-(1-(2,3-dichloro-4-(1H-pyrrolo[2,3-b]pyridin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0515		3-(1-(2,3-dichloro-4-(1H-indol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0520		3-(1-(2,3-dichloro-2',3',4',5'-tetrahydro[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0787		3-(1-(3-(cyclopropylamino)benzo[d]isoxazol-6-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0582		3-(1-(3-chloro-2-fluoro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0571		3-(1-(3-chloro-2-fluoro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0605		3-(1-(2,3-dichloro-3',5'-difluoro[1,1'-biphenyl]-4-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0573		3-(1-(4-bromonaphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0574		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(thiophen-3-yl)naphthalen-1-yl)ethyl)urea








H0575		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(thiophen-2-yl)naphthalen-1-yl)ethyl)urea
H0576		3-(1-(4-(1H-pyrazol-4-yl)naphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0577		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(pyridin-3-yl)naphthalen-1-yl)ethyl)urea
H0591		3-(1-(4-(3-aminophenyl)naphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0597		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(4-(thiazol-5-yl)naphthalen-1-yl)ethyl)urea
H0598		3-(1-(4-(furan-3-yl)naphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0599		3-(1-(4-(1H-imidazol-5-yl)naphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0790		3-(1-(4-cyanonaphthalen-1-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0381		1-methyl-1-(1-methylpiperidin-4-yl)-3-(1-(2,3,4-trichlorophenyl)ethyl)urea
H0519		3-(1-(2,3-dichloro-4-iodophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0629		3-((3-bromo-2-chloro-4-iodophenyl)(cyano)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0658		3-((3-bromo-2-chloro-4-methoxyphenyl)(cyano)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0669		3-(cyano(2,3-dichloro-4-methoxyphenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0671		3-(1-cyano-1-(2,3-dichloro-4-methoxyphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0659		2-(3-bromo-2-chloro-4-methoxyphenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetamide
H0521		methyl 2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)benzoate

H0602		3-(1-(2,3-dichloro-4-((trimethylsilyl)ethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0603		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0677		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0678		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0832		3-(1-(2,3-dichloro-4-(prop-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0852		3-(1-(2,3-dichloro-4-(3-methylbut-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0701		3-(1-(2,3-dichloro-4-(3-oxobut-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea










H0733		3-(1-(2,3-dichloro-4-(3-hydroxybut-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0755		3-(1-(2,3-dichloro-4-(3-hydroxyprop-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0757		3-(1-(2,3-dichloro-4-(3,3-diethoxyprop-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0734		3-(1-(2,3-dichloro-4-(pyridin-2-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0737		3-(1-(2,3-dichloro-4-(thiophen-2-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0775		3-(1-(2,3-dichloro-4-((5-(hydroxymethyl)thiophen-2-yl)ethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0776		5-((2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)ethynyl)thiophene-2-carboxamide




H0779		methyl 5-((2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)ethynyl)thiophene-2-carboxylate
H0762		3-(1-(2,3-dichloro-4-(furan-2-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0751		3-(1-(2,3-dichloro-4-(thiazol-4-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0763		3-(1-(4-((1H-imidazol-4-yl)ethynyl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0759		3-(1-(2,3-dichloro-4-(thiophen-3-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0785		3-(1-(2,3-dichloro-4-(3-(thiophen-2-yl)prop-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0754		3-(1-(2,3-dichloro-4-(thiazol-2-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0753		3-(1-(2,3-dichloro-4-(pyrimidin-5-ylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0609		3-(1-(2,3-dichloro-4-(phenylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0764		3-(1-(2,3-dichloro-4-(cyclopropylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0818		3-(1-(2,3-dichloro-4-(cyclopropylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0819		3-(1-(2,3-dichloro-4-(cyclopropylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0838		3-((S)-1-(2,3-dichloro-4-(cyclopropylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpyrrolidin-3-yl)urea
H0855		3-(1-(3-chloro-4-(cyclopropylethynyl)-2-fluorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0884		3-(1-(4,5-dichloro-6-(cyclopropylethynyl)pyridin-3-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0811		3-(1-(2,3-dichloro-4-(cyclopentylethynyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0812		3-(1-(2,3-dichloro-4-(3-(4-methylpiperazin-1-yl)prop-1-yn-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0740		3-(2-cyclopropyl-1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0742		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0745		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0749		3-(1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea
H0744		3-(2-cyclopropyl-1-(2,3-dichloro-4-ethynylphenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea

H0626		3-(1-(2,3-dichloro-4-vinylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0767		(E)-3-(1-(2,3-dichloro-4-(2-(thiophen-2-yl)vinyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0772		N-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxamide
H0773		2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)-N-(thiophen-2-yl)benzamide
H0784		3-(1-(2,3-dichloro-4-(3-(thiophen-2-yl)ureido)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0777		3-(1-(2,3-dichloro-4-(thiophen-2-ylamino)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0846		3-(1-(2,3-dichloro-4-(cyclopropylamino)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0875		3-(1-(2,3-dichloro-4-(cyclopropoxy)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0628		3-(1-(2,3-dichloro-4-ethylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0630		3-(1-(2,3-dichloro-4-(cyanomethyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0633		3-(1-(2,3-dichloro-4-(hydroxymethyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0634		3-(1-(2,3-dichloro-4-(fluoromethyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0640		3-(1-(2,3-dichloro-4-formylphenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0645		3-(1-(2,3-dichloro-4-(1,3-dioxolan-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0641		methyl (E)-3-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)acrylate
H0702		(Z)-3-(1-(2,3-dichloro-4-(1-chloro-3-oxobut-1-en-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0643		3-(1-(2,3-dichloro-4-(3-hydroxypropyl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0522		2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)benzamide
H0523		3-(1-(2,3-dichloro-4-cyanophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0876		1-methyl-1-(1-(1-methylpiperidin-4-yl)-3-(1-(4,5,6-trichloropyridin-3-yl)ethyl)urea.

5

At various places in the present specification, substituents of compounds of the invention are disclosed in groups or in ranges. It is specifically intended that the invention include each and every individual subcombination of the members of such groups and ranges. For example, the term "C₁₋₆ alkyl" is specifically intended to individually disclose methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl.

10

For compounds of the invention in which a variable appears more than once, each variable can be a different moiety selected from the Markush group defining the variable. For example, where a structure is described having two R groups that are simultaneously present on the same compound; the two R groups can represent different moieties selected from the Markush group defined for R.

15

It is further appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

20

As used herein, the term "alkyl" is meant to refer to a saturated hydrocarbon group which is straight-chained or branched. Example alkyl groups include methyl (Me), ethyl (Et), propyl (e.g., n-propyl and isopropyl), butyl (e.g., n-butyl, isobutyl, t-butyl), pentyl (e.g., n-pentyl, isopentyl, neopentyl), and the like. An alkyl group can contain from 1 to about 20, from 2 to about 20, from 1 to about 10, from 1 to about 8, from 1 to about 6, from 1 to about 4, or from 1 to about 3 carbon atoms.

25

5 As used herein, "alkenyl" refers to an alkyl group having one or more double carbon-carbon bonds. Example alkenyl groups include ethenyl, propenyl, cyclohexenyl, and the like.

As used herein, "alkynyl" refers to an alkyl group having one or more triple carbon-carbon bonds. Example alkynyl groups include ethynyl, propynyl, and the like.

10 As used herein, "haloalkyl" refers to an alkyl group having one or more halogen substituents. Example haloalkyl groups include CF_3 , C_2F_5 , CHF_2 , CCl_3 , CHCl_2 , C_2Cl_5 , and the like.

As used herein, "hydroxylalkyl" refers to an alkyl group having one or more OH substituents. Example hydroxylalkyl groups include CH_2OH , $\text{C}_2\text{H}_4\text{OH}$, $\text{C}_3\text{H}_6\text{OH}$, and the like.

15 As used herein, "aryl" refers to monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbons such as, for example, phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from 6 to about 20 carbon atoms.

20 As used herein, "cycloalkyl" refers to non-aromatic carbocycles including cyclized alkyl, alkenyl, and alkynyl groups. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2, 3 or 4 fused rings) ring systems as well as spiro ring systems. Example cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcarnyl, adamantyl, and the like. Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the cycloalkyl ring, for example, benzo derivatives of
25 pentane, pentene, hexane, and the like. In some embodiments, cycloalkyl groups can have from about 3 to about 10, or about 3 to about 7 ring-forming carbon atoms.

30 As used herein, "heterocyclyl" or "heterocycle" refers to a saturated or unsaturated cyclic hydrocarbon wherein one or more of the ring-forming carbon atoms of the cyclic hydrocarbon is replaced by a heteroatom such as O, S, or N. Heterocyclyl groups can be aromatic (e.g., "heteroaryl") or non-aromatic (e.g., "heterocycloalkyl"). Heterocyclyl groups can also correspond to hydrogenated and partially hydrogenated heteroaryl groups. Heterocyclyl groups can include mono- or polycyclic (e.g., having 2, 3 or 4 fused rings) ring systems. Heterocyclyl groups can be characterized as having 3-14 or 3-7 ring-forming atoms. In some embodiments, heterocyclyl groups can contain, in addition to at least one heteroatom, from about 1 to about 13,
35 about 2 to about 10, or about 2 to about 7 carbon atoms and can be attached through a carbon

5 atom or heteroatom. In further embodiments, the heteroatom can be oxidized (e.g., have an oxo substituent) or a nitrogen atom can be quaternized. Examples of heterocyclyl groups include morpholino, thiomorpholino, piperazinyl, tetrahydrofuranyl, tetrahydrothienyl, 2,3-dihydrobenzofuryl, 1,3-benzodioxole, benzo-1,4-dioxane, piperidinyl, pyrrolidinyl, isoxazolidinyl, isothiazolidinyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, imidazolidinyl, and

10 the like, as well as any of the groups listed below for "heteroaryl" and "heterocycloalkyl." Further example heterocycles include pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, 3,6-dihydropyridyl, 1,2,3,6-tetrahydropyridyl, 1,2,5,6-tetrahydropyridyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,

15 pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, tetrazolyl, 6H-1,2,5-thia-diazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, xanthenyl, octahydro-isoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl,

20 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxaliny, quinuclidinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzo-thiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazoliny, methylenedioxyphenyl, morpholinyl, naphthyridinyl, deca-hydroquinolinyl, 2H,6H-1,5,2dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, imidazolidinyl, imidazoliny, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl and isoxazolyl.

30 Further examples of heterocycles include azetidin-1-yl, 2,5-dihydro-1H-pyrrol-1-yl, piperindinyl, piperazin-1-yl, pyrrolidin-1-yl, isoquinol-2-yl, pyridin-1-yl, 3,6-dihydropyridin-1-yl, 2,3-dihydroindol-1-yl, 1,3,4,9-tetrahydrocarbolin-2-yl, thieno[2,3-c]pyridin-6-yl, 3,4,10,10a-tetrahydro-1H-pyrazino[1,2-a]indol-2-yl, 1,2,4,4a,5,6-hexahydro-pyrazino[1,2-a]quinolin-3-yl, pyrazino[1,2-a]quinolin-3-yl, diazepan-1-yl, 1,4,5,6-tetrahydro-2H-benzo[f]isoquinolin-3-yl, 1

5 ,4,4a,5,6, 10b-hexahydro-2H-benzo[f]isoquinolin-3-yl, 3,3a,8,8a-tetrahydro- 1 H-2-aza-cyclopenta[a]inden-2-yl, and 2,3,4,7-tetrahydro- 1 H-azepin- 1 -yl, azepan- 1 -yl.

As used herein, "heteroaryl" groups refer to an aromatic heterocycle having at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include monocyclic and polycyclic (e.g., having 2, 3 or 4 fused rings) systems. Examples of heteroaryl
10 groups include without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl (furanlyl), quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrol, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, benzothienyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, and the like. In some embodiments, the heteroaryl group has from 1 to about 20 carbon atoms, and in
15 further embodiments from about 3 to about 20 carbon atoms. In some embodiments, the heteroaryl group contains 3 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms.

As used herein, "heterocycloalkyl" refers to non-aromatic heterocycles including cyclized alkyl, alkenyl, and alkynyl groups where one or more of the ring-forming carbon atoms is
20 replaced by a heteroatom such as an O, N, or S atom. Example "heterocycloalkyl" groups include morpholino, thiomorpholino, piperazinyl, tetrahydrofuranlyl, tetrahydrothienyl, 2,3-dihydrobenzofuryl, 1,3-benzodioxole, benzo- 1,4-dioxane, piperidinyl, pyrrolidinyl, isoxazolidinyl, isothiazolidinyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, imidazolidinyl, and the like. Also included in the definition of heterocycloalkyl are moieties that have one or more
25 aromatic rings fused (i.e., having a bond in common with) to the nonaromatic heterocyclic ring, for example phthalimidyl, naphthalimidyl, and benzo derivatives of heterocycles such as indolene and isoindolene groups. In some embodiments, the heterocycloalkyl group has from 1 to about 20 carbon atoms, and in further embodiments from about 3 to about 20 carbon atoms. In some embodiments, the heterocycloalkyl group contains 3 to about 14, 3 to about 7, or 5 to 6
30 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms. In some embodiments, the heterocycloalkyl group contains 0 to 3 double bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 triple bonds.

As used herein, "halo" or "halogen" includes fluoro, chloro, bromo, and iodo.

As used herein, "alkoxy" refers to an -O-alkyl group. Example alkoxy groups include
35 methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), t-butoxy, and the like.

5 As used herein, "thioalkoxy" refers to an -S-alkyl group.

As used here, "haloalkoxy" refers to an -O-haloalkyl group. An example haloalkoxy group is OCF .

As used herein, "cycloalkyloxy" refers to -O-cycloalkyl.

As used herein, "aralkyl" refers to an alkyl group substituted by an aryl group.

10 As used herein, "cycloalkylalkyl" refers to an alkyl group substituted by a cycloalkyl group.

As used herein, "heterocyclalkyl" refers to an alkyl moiety substituted by a heterocarbocycl group. Example heterocyclalkyl groups include "heteroarylalkyl" (alkyl substituted by heteroaryl) and "heterocycloalkylalkyl" (alkyl substituted by heterocycloalkyl). In some embodiments, heterocyclalkyl groups have from 3 to 24 carbon atoms in addition to at least one ring-forming heteroatom.

As used herein "oxo" refers to =O.

The compounds described herein can be asymmetric (e.g., having one or more stereocenters). The description of a compound without specifying its stereochemistry is intended to capture mixtures of stereoisomers as well as each of the individual stereoisomer encompassed within the genus.

Compounds of the invention can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

25 The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

30 The present invention also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of

5 the present invention include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds
10 with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and Journal of Pharmaceutical Science, 66, 2 (1977), each of which is incorporated herein by
15 reference in its entirety.

Synthesis

Compounds of the invention, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes.

20 The reactions for preparing compounds of the invention can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially nonreactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling
25 temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected.

Preparation of compounds of the invention can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of
30 appropriate protecting groups can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in T.W. Green and P.G.M. Wuts, Protective Groups in Organic Synthesis, 3rd. Ed., Wiley & Sons, Inc., New York (1999), which is incorporated herein by reference in its entirety.

Reactions can be monitored according to any suitable method known in the art. For
35 example, product formation can be monitored by spectroscopic means, such as nuclear magnetic

5 resonance spectrometry (e.g., ^1H or ^{13}C) infrared spectroscopy, spectrophotometry (e.g., UV-visible), or mass spectrometry, or by chromatography such as high performance liquid chromatography (HPLC) or thin layer chromatography.

Pharmaceutical Compositions

10 Pharmaceutical compositions for preventing and/or treating a subject are further provided comprising a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients.

A "pharmaceutically acceptable" excipient is one that is not biologically or otherwise
15 undesirable, i.e., the material can be administered to a subject without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained. The carrier can be selected to minimize any degradation of the active ingredient and to minimize any adverse side effects in the subject, as would be well known to one of skill in the art. The carrier can be a solid, a liquid, or both.

20 The disclosed compounds can be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment or prevention intended. The active compounds and compositions, for example, can be administered orally, rectally, parenterally, ocularly, inhalationally, or topically. In particular, administration can be epicutaneous, inhalational, enema, conjunctival, eye drops, ear drops,
25 alveolar, nasal, intranasal, vaginal, intravaginal, transvaginal, ocular, intraocular, transocular, enteral, oral, intraoral, transoral, intestinal, rectal, intrarectal, transrectal, injection, infusion, intravenous, intraarterial, intramuscular, intracerebral, intraventricular, intracerebroventricular, intracardiac, subcutaneous, intraosseous, intradermal, intrathecal, intraperitoneal, intravesical, intracavernosal, intramedullar, intraocular, intracranial, transdermal, transmucosal, transnasal,
30 inhalational, intracisternal, epidural, peridural, intravitreal, etc.

Suitable carriers and their formulations are described in Remington: The Science and Practice of Pharmacy (19th ed.) ed. A.R. Gennaro, Mack Publishing Company, Easton, Pa., 1995. Oral administration of a solid dose form can be, for example, presented in discrete units, such as hard or soft capsules, pills, cachets, lozenges, or tablets, each containing a predetermined
35 amount of at least one of the disclosed compound or compositions. In some forms, the oral

5 administration can be in a powder or granule form. In some forms, the oral dose form is sub-lingual, such as, for example, a lozenge. In such solid dosage forms, the compounds of Formula I are ordinarily combined with one or more adjuvants. Such capsules or tablets can contain a controlled-release formulation. In the case of capsules, tablets, and pills, the dosage forms also can comprise buffering agents or can be prepared with enteric coatings.

10 In some forms, oral administration can be in a liquid dose form. Liquid dosage forms for oral administration include, for example, pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art (e.g., water). Such compositions also can comprise adjuvants, such as wetting, emulsifying, suspending, flavoring (e.g., sweetening), and/or perfuming agents.

15 In some forms, the disclosed compositions can comprise a parenteral dose form. "Parenteral administration" includes, for example, subcutaneous injections, intravenous injections, intraperitoneally, intramuscular injections, intrasternal injections, and infusion. Injectable preparations (e.g., sterile injectable aqueous or oleaginous suspensions) can be formulated according to the known art using suitable dispersing, wetting agents, and/or
20 suspending agents. Typically, an appropriate amount of a pharmaceutically acceptable carrier is used in the formulation to render the formulation isotonic. Examples of the pharmaceutically acceptable carrier include, but are not limited to, saline, Ringer's solution and dextrose solution. Other acceptable excipients include, but are not limited to, thickeners, diluents, buffers, preservatives, surface active agents and the like.

25 In some forms, the disclosed compositions can comprise a topical dose form. "Topical administration" includes, for example, transdermal administration, such as via transdermal patches or iontophoresis devices, intraocular administration, or intranasal or inhalation administration. Compositions for topical administration also include, for example, topical gels, sprays, ointments, and creams. A topical formulation can include a compound which enhances
30 absorption or penetration of the active ingredient through the skin or other affected areas. When the compounds and compositions are administered by a transdermal device, administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. Typical formulations for this purpose include gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants,
35 sponges, fibres, bandages and microemulsions. Liposomes can also be used. Typical carriers

5 include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, polyethylene glycol and propylene glycol. Penetration enhancers can be incorporated--see, for example, J Pharm Sci, 88 (10), 955-958, by Finnin and Morgan (October 1999).

Formulations suitable for topical administration to the eye include, for example, eye drops wherein the disclosed compound or composition is dissolved or suspended in suitable
10 carrier. A typical formulation suitable for ocular or aural administration can be in the form of drops of a micronised suspension or solution in isotonic, pH-adjusted, sterile saline. Other formulations suitable for ocular and aural administration include ointments, biodegradable (e.g. absorbable gel sponges, collagen) and non-biodegradable (e.g. silicone) implants, wafers, lenses and particulate or vesicular systems, such as niosomes or liposomes. A polymer such as cross-
15 linked polyacrylic acid, polyvinylalcohol, hyaluronic acid, a cellulosic polymer, for example, hydroxypropylmethylcellulose, hydroxyethylcellulose, or methyl cellulose, or a heteropolysaccharide polymer, for example, gelan gum, can be incorporated together with a preservative, such as benzalkonium chloride. Such formulations can also be delivered by iontophoresis.

20 Other carrier materials and modes of administration known in the pharmaceutical art can also be used. The disclosed pharmaceutical compositions can be prepared by any of the well-known techniques of pharmacy, such as effective formulation and administration procedures. The above considerations in regard to effective formulations and administration procedures are well known in the art and are described in standard textbooks. Formulation of drugs is discussed
25 in, for example, Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 1975; Liberman, et al., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Kibbe, et al., Eds., Handbook of Pharmaceutical Excipients (3.sup.rd Ed.), American Pharmaceutical Association, Washington, 1999.

The disclosed compounds can be used, alone or in combination with other therapeutic
30 agents, in the treatment or prevention of various conditions or disease states. The administration of two or more compounds "in combination" means that the two compounds are administered closely enough in time that the presence of one alters the biological effects of the other. The two or more compounds can be administered simultaneously, concurrently or sequentially.

Disclosed are pharmaceutical compositions comprising an effective amount of a
35 compound of the invention or a pharmaceutically accepted salt thereof; and a pharmaceutically

5 acceptable carrier or vehicle. These compositions may further comprise additional agents. These compositions are useful for modulating the activity of ghrelin receptor, thus to improve the prevention and treatment of ghrelin receptor associated human diseases such as obesity and/or metabolic disorders.

10 *Methods*

All of the methods of the invention may be practiced with a compound of the invention alone, or in combination with other agents.

The above-described compounds and compositions are useful for the inhibition, reduction, prevention, and/or treatment of diseases which are pathophysiologically modulated by the ghrelin receptor. Accordingly, in some forms, disclosed are methods of preventing and/or
15 treating diseases which are pathophysiologically modulated by the ghrelin receptor, comprising administering to a subject a therapeutically effective amount of a compound of Formula I as disclosed above, or a pharmaceutically acceptable salt thereof.

Suitable subjects can include mammalian subjects. Mammals include, but are not limited
20 to, canine, feline, bovine, caprine, equine, ovine, porcine, rodents, lagomorphs, primates, and the like, and encompass mammals in utero. In some forms, humans are the subjects. Human subjects can be of either gender and at any stage of development.

Diseases modulated by the ghrelin receptor, and potentially treatable by the methods disclosed herein, include obesity, overweight, eating disorder, diabetes, metabolic syndrome, cachexia resulting from cancer, congestive heart failure, wasting due to ageing or AIDS, chronic
25 liver failure, chronic obstructive pulmonary disease, gastrointestinal disease, gastric disorder or substance abuse. Metabolic disorders potentially treatable by the instant methods include diabetes, Type I diabetes, Type II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperinsulinemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia, obesity, aging, Syndrome X, atherosclerosis, heart disease, stroke, hypertension
30 and peripheral vascular disease. Gastric disorders potentially treatable by the instant methods include post-operative ileus (POI), diabetic gastroparesis, and opioid induced bowel dysfunction. Gastrointestinal diseases potentially treatable by the instant methods include irritable bowel syndrome, gastritis, acid reflux disease, gastroparesis, and functional dyspepsia. Substance abuse

5 potentially treatable by the instant methods includes alcohol and drug abuse, and said drug includes amphetamines, barbiturates, benzodiazepines, cocaine, methaqualone, and opioids.

In some embodiments of the invention, the compound of Formula I is useful in the treatment of Prader-Willi Syndrome, a genetic disorder usually involving chromosome 15. Prader-Willi is characterized by obesity, hypotonia, or poor muscle tone, and significant
10 developmental delays in children afflicted with this disorder.

In some embodiments of the invention, the compound of Formula I is useful in the treatment of an over-eating disorder. An over-eating disorder is a complex compulsion to eat. The eating may be excessive (compulsive over-eating); may include normal eating punctuated with episodes of purging; or may include cycles of bingeing and purging. The most prevalent
15 over-eating disorder is Bulimia nervosa. Another widely and rapidly spreading over-eating disorder is compulsive over-eating, also termed Binge Eating Disorder (BED). In some embodiments, the compound of Formula I is used in the treatment of BED.

In some embodiments, the compound of Formula I is useful in the treatment of Parkinson-induced constipation and gastric dysmotility. In some embodiments, the compound of
20 Formula I is useful in the treatment of chemotherapy-induced nausea and vomiting (CINV).

In some embodiments, the compound of Formula I is useful in the treatment of inflammation, acute and chronic pain, and motion sickness.

In some embodiments, the compound of Formula I is useful in the treatment of drug and alcohol abuse. In some methods the compound of Formula I is a ghrelin receptor modulator. In
25 some other methods the compound of Formula I is a ghrelin receptor agonist. In some methods the compound of Formula I is a ghrelin receptor antagonist. In some methods, the compound of Formula I or a pharmaceutically acceptable salt thereof, is administered by one or more routes selected from the group consisting of rectal, buccal, sublingual, intravenous, subcutaneous, intradermal, transdermal, intraperitoneal, oral, eye drops, parenteral and topical administration.
30 In some other methods, administration is accomplished by administering an oral form of the compound of Formula I or a pharmaceutically acceptable salt thereof.

A therapeutically effective amount may vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors. Therapeutically effective amounts of compounds of Formula I may range from
35 approximately 0.01 microgram per Kg ($\mu\text{g/Kg}$) body weight per day to about 100 mg/Kg body

5 weight per day, or from about 0.1 µg/Kg/day to about 10 mg/Kg/day, or from about 1 µg/Kg/day to about 5 mg/Kg/day, or from about 10 µg/Kg/day to about 5 mg/Kg/day, or from about 100 µg/Kg/day to about 5 mg/Kg/day, or from about 500 µg/Kg/day to about 5 mg/Kg/day.

10 *Definitions of Terms*

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this pertains. The references disclosed are also individually and specifically incorporated by reference herein for the material contained in them
15 that is discussed in the sentence in which the reference is relied upon.

1. A, an, the

As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example,
20 reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

2. Abbreviations

Abbreviations, which are well known to one of ordinary skill in the art, may be used
25 (e.g., "h" or "hr" for hour or hours, "g" or "gm" for gram(s), "mL" for milliliters, and "rt" for room temperature, "nm" for nanometers, "M" for molar, and like abbreviations).

3. About

The term "about," when used to modify the quantity of an ingredient in a composition,
30 concentrations, volumes, process temperature, process time, yields, flow rates, pressures, and like values, and ranges thereof, employed in describing the embodiments of the disclosure, refers to variation in the numerical quantity that can occur, for example, through typical measuring and handling procedures used for making compounds, compositions, concentrates or use formulations; through inadvertent error in these procedures; through differences in the
35 manufacture, source, or purity of starting materials or ingredients used to carry out the methods;

5 and like considerations. The term "about" also encompasses amounts that differ due to aging of a composition or formulation with a particular initial concentration or mixture, and amounts that differ due to mixing or processing a composition or formulation with a particular initial concentration or mixture. Whether modified by the term "about" the claims appended hereto include equivalents to these quantities.

10

4. Comprise

Throughout the description and claims of this specification, the word "comprise" and variations of the word, such as "comprising" and "comprises," means "including but not limited to," and is not intended to exclude, for example, other additives, components, integers or steps.

15

5. Ghrelin Receptor Agonist

A ghrelin receptor agonist is any molecule that binds to and activates the Ghrelin receptor in the cells.

20 6. Ghrelin Receptor Antagonist

A ghrelin receptor antagonist is any molecule that binds to and inhibits the activity of Ghrelin receptor.

7. Pathophysiologically Mediated by Ghrelin Receptor

25 Something is "pathophysiologically mediated by the ghrelin receptor" if the ghrelin receptor is involved in the functional changes in body associated with or resulting from disease or injury.

8. Obesity

30 Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have an adverse effect on health, leading to reduced life expectancy and/or increased health problems. Obesity treatment includes inducing weight loss, reducing bodyweight, reducing food intake, reducing appetite, increasing metabolic rate, reducing fat intake, reducing carbohydrate craving; or inducing satiety. The obesity-related disorders herein are associated
35 with, caused by, or result from obesity. Examples of obesity-related disorders include overeating,

5 binge eating, and bulimia, hypertension, diabetes, elevated plasma insulin concentrations and insulin resistance, dyslipidemias, hyperlipidemia, endometrial, breast, prostate and colon cancer, osteoarthritis, obstructive sleep apnea, cholelithiasis, gallstones, heart disease, abnormal heart rhythms and arrhythmias, myocardial infarction, congestive heart failure, coronary heart disease, sudden death, stroke, polycystic ovary disease, craniopharyngioma, the Prader-Willi Syndrome, 10 Frohlich's syndrome, GH-deficient subjects, normal variant short stature, Turner's syndrome, and other pathological conditions showing reduced metabolic activity or a decrease in resting energy expenditure as a percentage of total fat-free mass, e.g. children with acute lymphoblastic leukemia. Further examples of obesity-related disorders are metabolic syndrome, insulin resistance syndrome, sexual and reproductive dysfunction, such as infertility, hypogonadism in 15 males and hirsutism in females, gastrointestinal motility disorders, such as obesity-related gastro-esophageal reflux, respiratory disorders, such as obesity-hypoventilation syndrome (Pickwickian syndrome), cardiovascular disorders, inflammation, such as systemic inflammation of the vasculature, arteriosclerosis, hypercholesterolemia, hyperuricaemia, lower back pain, gallbladder disease, gout, and kidney cancer, nicotine addiction, substance addiction and alcoholism. The 20 compositions of the present invention are also useful for reducing the risk of secondary outcomes of obesity, such as reducing the risk of left ventricular hypertrophy.

9. Metabolic Disorder

A metabolic disorder is a disorder of metabolism, such as diabetes, Type I diabetes, Type 25 II diabetes, inadequate glucose tolerance, insulin resistance, hyperglycemia, hyperinsulinemia, hyperlipidemia, hypertriglyceridemia, hypercholesterolemia, dyslipidemia, obesity, aging, Syndrome X, atherosclerosis, heart disease, stroke, hypertension and peripheral vascular disease.

10. Congestive Heart Failure

30 Congestive heart failure (CHF) is a condition in which the heart's function as a pump to deliver oxygen rich blood to the body is inadequate to meet the body's needs. Congestive heart failure can be caused by diseases that weaken the heart muscle, or diseases that cause stiffening of the heart muscles, or diseases that increase oxygen demand by the body tissue beyond the capability of the heart to deliver. Many diseases can impair the pumping action of the ventricles. 35 For example, the muscles of the ventricles can be weakened by heart attacks or infections

5 (myocarditis). The diminished pumping ability of the ventricles due to muscle weakening is called systolic dysfunction. After each ventricular contraction (systole) the ventricle muscles need to relax to allow blood from the atria to fill the ventricles. This relaxation of the ventricles is called diastole. Diseases such as hemochromatosis or amyloidosis can cause stiffening of the heart muscle and impair the ventricles' capacity to relax and fill; this is referred to as diastolic
10 dysfunction. The most common cause of this is longstanding high blood pressure resulting in a thickened (hypertrophied) heart. Additionally, in some patients, although the pumping action and filling capacity of the heart may be normal, abnormally high oxygen demand by the body's tissues (for example, with hyperthyroidism) may make it difficult for the heart to supply an adequate blood flow (called high output heart failure). In some patients one or more of these
15 factors can be present to cause congestive heart failure. Congestive heart failure can affect many organs of the body. For example, the weakened heart muscles may not be able to supply enough blood to the kidneys, which then begin to lose their normal ability to excrete salt (sodium) and water. This diminished kidney function can cause the body to retain more fluid. The lungs may become congested with fluid (pulmonary edema) and the person's ability to exercise is decreased.
20 Fluid may likewise accumulate in the liver, thereby impairing its ability to rid the body of toxins and produce essential proteins. The intestines may become less efficient in absorbing nutrients and medicines. Over time, untreated, worsening congestive heart failure will affect virtually every organ in the body.

25 11. Agonism Action

Agonism action refers to the binding of a molecule to a receptor that leads to the activation of the receptor, thus triggering a cellular response similar to the cellular response for a known agonist for the receptor.

30 12. Antagonism Action

Antagonism action refers to the binding of a molecule to a receptor that leads to the inhibition of the receptor.

13. Modulate

5 To modulate, or forms thereof, means either increasing, decreasing, or maintaining a cellular activity mediated through a cellular target. It is understood that wherever one of these words is used it is also disclosed that it could be 1%, 5%, 10%, 20%, 50%, 100%, 500%, or 1000% increased from a control, or it could be 1%, 5%, 10%, 20%, 50%, or 100% decreased from a control.

10

14. Optional

"Optional" or "optionally" means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances where it does not.

15

15. Or

The word "or" or like terms as used herein means any one member of a particular list and also includes any combination of members of that list.

20 16. Publications

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this pertains. The references disclosed are also individually and specifically incorporated by reference herein for the material contained in them that is discussed in the sentence in which the reference is relied upon.

25

17. Subject

As used throughout, by a "subject" is meant an individual. Thus, the "subject" can include, for example, domesticated animals, such as cats, dogs, etc., livestock (e.g., cattle, horses, pigs, sheep, goats, etc.), laboratory animals (e.g., mouse, rabbit, rat, guinea pig, etc.) mammals, non-human mammals, primates, non-human primates, rodents, birds, reptiles, amphibians, fish, and any other animal. The subject can be a mammal such as a primate or a human. The subject can also be a non-human.

30

35 18. Treating

5 By "treating" or "treatment" is meant the medical management of a patient with the intent to cure, ameliorate, stabilize, or prevent a disease, pathological condition, or disorder. These terms include active treatment, that is, treatment directed specifically toward the improvement of a disease, pathological condition, or disorder, and also includes causal treatment, that is, treatment directed toward removal of the cause of the associated disease, pathological condition, or disorder. These terms can mean that the symptoms of the underlying disease are reduced, and/or that one or more of the underlying cellular, physiological, or biochemical causes or mechanisms causing the symptoms are reduced. It is understood that reduced, as used in this context, means relative to the state of the disease, including the molecular state of the disease, not just the physiological state of the disease. In certain situations a treatment can inadvertently cause harm. In addition, these terms include palliative treatment, that is, treatment designed for the relief of symptoms rather than the curing of the disease, pathological condition, or disorder; preventative treatment, that is, treatment directed to minimizing or partially or completely inhibiting the development of the associated disease, pathological condition, or disorder; and supportive treatment, that is, treatment employed to supplement another specific therapy directed toward the improvement of the associated disease, pathological condition, or disorder. These terms mean both treatment having a curing or alleviating purpose and treatment having a preventive purpose. The treatment can be made either acutely or chronically. It is understood that treatment can mean a reduction or one or more symptoms or characteristics by at least 5% 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 99.9%, 99.99%, 100%, relative to a control. In the context of these terms, preventing refers to the ability of a compound or composition (such as the disclosed compounds and compositions) to prevent a disease identified herein in patients diagnosed as having the disease or who are at risk of developing such disease. In this context, preventing includes the delaying the onset of the disease relative to a control. These terms do not require that the treatment in fact be effective to produce any of the intended results. It is enough that the results are intended.

19. Therapeutically Effective

The term "therapeutically effective" means that the amount of the composition used is of sufficient quantity to treat a subject as defined herein.

5

20. Toxicity


Toxicity is the degree to which a substance, molecule, is able to damage something, such as a cell, a tissue, an organ, or a whole organism, that has been exposed to the substance or molecule. For example, the liver, or cells in the liver, hepatocytes, can be damaged by certain substances. The methods of the present invention are preferably non-toxic.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

15

EXAMPLES**Example 1****Synthesis of Intermediate 1k**

20

**Step 1:**

5 To a solution of **1a** (100 g, 0.62 mol) in DMF (1.2 L) was added *N*-bromosuccinimide (110 g, 0.62 mol) at 0 °C. The mixture was stirred at room temperature for 4 h, then water (800 mL) was added and the resulting mixture was extracted with EtOAc (3 × 500 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was triturated with petroleum ether to provide **1b** (133.7 g, 89% yield) as a brown solid. ¹H-NMR (CDCl₃, 300 MHz): δ= 7.30 (d, 1 H), 6.59 (d, 1 H), 4.22 (br, 2 H). LC-MS: 241 [M+1]⁺.
10

Step 2:

To a solution of **1b** (133.7 g, 0.55 mol) in dry CH₂Cl₂ (1.5 L) was added acetic anhydride (110 g, 0.62 mol) dropwise over a period of 20 minutes at room temperature. The mixture was stirred at
15 room temperature overnight, then diluted with CH₂Cl₂ (300 mL) and washed with water (150 mL) and brine (200 mL). The organic layer was separated, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was triturated with petroleum ether (300 mL) to provide compound **1c** (143.0 g, 91% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.26 (d, 1 H), 7.63 (br, 1 H), 7.54 (d, 1 H), 2.26 (s, 3 H). LC-MS: 280 [M-1]⁻.
20

Step 3:

A mixture of compound **1c** (50.0 g, 0.18 mol), butyl vinyl ether (**1d**, 89.0 g, 0.89 mol), bis(1,3-diphenylphosphino)propane (DPPP, 22.0 g, 0.053 mol), TEA (100 mL, 0.71 mol) and Pd(OAc)₂ (6.4 g, 0.027 mol) in DMSO (1.2 L) was heated at 130 °C under N₂ overnight. After the reaction
25 was completed, the mixture was cooled to 0 °C and 2N HCl (480 mL) was added dropwise over a period of 30 minutes. Then, the mixture was extracted with EtOAc (3 × 100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: PE=1:10) to provide **1e** (19.5 g, 45% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.46 (d, 1 H),
30 7.82 (br, 1 H), 7.51 (d, 1 H), 2.63 (s, 3 H), 2.29 (s, 3 H). LC-MS: 244 [M-1]⁻.

Step 4:

To a solution of **1e** (21.9 g, 89.4 mmol) in MeOH (350 mL) was added 2N NaOH solution (350 mL) at room temperature. The mixture was heated at 50 °C overnight, then cooled and
35 concentrated under reduced pressure. The resulting solid was triturated with water (100 mL) for

5 30 min and filtered to provide **1f** (18.0 g, 98% yield) as a brown solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.48 (d, 1 H), 6.68 (d, 1 H), 4.56 (br, 2 H), 2.62 (s, 3 H). LC-MS: 202[M-1].

Step 5:

To a mixture of compound **1f** (18.0 g, 89.2 mmol) and ice (360 g) in conc. HCl (180 mL) was
10 added a solution of NaNO₂ (9.2 g, 133.7 mmol) in water (20 mL) dropwise over a period of 30 minutes, and the resulting mixture stirred in an ice bath for 30 min. A solution of KI (74.0 g, 446 mmol) in water (360 mL) was added dropwise over 45 min at 0 °C. The mixture was stirred for 30 min and then extracted with EtOAc (3 × 100 mL). The combined organic layers were dried
15 over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: PE=1:40) to provide **1g** (23.9 g, 86% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.6 (d, 1 H), 7.06 (d, 1 H), 2.62 (s, 3 H).

Step 6:

To a solution of **1g** (23.9 g, 76.1 mmol) in MeOH (100 mL)/THF (100 mL) was slowly added
20 NaBH₄ (2.9 g, 76.1 mmol) at 0 °C. The mixture was stirred at room temperature for 5 min, and then quenched with water (100 mL). The mixture was extracted with EtOAc (3 × 100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: PE=1:10) to
25 provide **1h** (22.4 g, 93% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.81 (d, 1 H), 7.26 (d, 1 H), 5.23 (q, 1 H), 2.17 (br, 1 H), 1.47 (d, 3 H).

Step 7:

To a mixture of **1h** (22.4 g, 70.9 mmol), phthalimide (12.5 g, 85.0 mmol) and PPh₃ (22.3 g, 85.0 mmol) in dry THF (450 mL) was added DIAD (21.5 g, 106.3 mmol) at room temperature under
30 N₂ protection. The mixture was stirred at room temperature overnight and then concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: PE=1:15) to provide **1i** (18.5 g, 58% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.78-7.84 (m, 3 H), 7.70-7.73 (m, 2 H), 7.41-7.43 (d, 1 H), 5.76-5.81 (q, 1 H), 1.84 (d, 3 H).

35 **Step 8:**


- 5 A solution of **1i** (7.2 g, 16.2 mmol) and hydrazine hydrate (98%, 4.0 g, 80.9 mmol) in MeOH (150 mL) was heated under reflux for 2 h, then cooled and concentrated under reduced pressure. The residue was diluted with water (100 mL) and extracted with CH₂Cl₂ (3 × 100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give **1j** (3.8 g, 75% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.81 (d, 1 H), 7.25 (d, 1 H), 4.55 (q, 1 H), 1.36-1.38 (d, 3 H). LC-MS: 316 [M+1]⁺.
- 10

Step 9:

- To a solution of **1j** (41.0g, 0.13 mol) in methyl tert-butyl ether (750 mL) was added slowly a solution of D-mandelic acid (7.8 g, 0.052 mol) in methyl tert-butyl ether (110 mL) at 45°C. The mixture was stirred at this temperature for 30 min then cooled and filtered. White solid obtained was partitioned between 5% NaOH solution (300 mL) and methyl tert-butyl ether (300 mL). The bi-phases were separated and the aqueous phase was extracted with methyl tert-butyl ether (300 mL). The combined organic layer was concentrated to provide **Intermediate 1k** (12 g, 58.5% yield) as a white solid (ee%=98.0%, Chiralpak AD-H, 5 μm, 4.6*250mm, mobile phase: Hex: EtOH : DEA=80 : 20 : 0.2), retention time = 6.408 min).
- 15
- 20

Example 2

Synthesis of Compound 2b




- 25 A suspension of *N*-methyl-4-piperidone **2a** (13.3 g, 58.6 mmol), NH₂Me (30% in MeOH, 100 mL) and Pd/C (0.66 g) in MeOH (200 mL) was heated at 60 °C under H₂ atmosphere (50 psi) overnight, then cooled and filtered. The filtrate was concentrated under reduced pressure and the residue was dissolved in HCl in dioxane (3N, 100 mL) and stirred for 30 min. The precipitate was filtered and washed with EtOAc (50 mL) to provide **2b** (7.7g, 54% yield) as white powder.
- 30 ¹H-NMR (DMSO, 400 MHz): δ= 9.50 (br, 2 H), 3.48 (d, 2 H), 3.15-3.16 (m, 1 H), 2.96-3.01 (m, 2 H), 2.70 (s, 3 H), 2.51 (s, 3 H), 2.22-2.28 (m, 2 H), 1.94-2.02 (m, 2 H), LC-MS: 129 [M+1]⁺.

Example 3

Synthesis of Compound H0603

5

**Step 1:**

To a solution of **1k** (1.83 g, 5.8 mmol) in CH_2Cl_2 (70 mL) was added TEA (5.6 mL, 40.6 mmol) and triphosgene (1.29 g, 4.4 mmol) at 0 °C. The mixture was stirred for 20 min, then **2b** (1.14 g, 6.97 mmol) was added. The ice bath was removed and the mixture stirred for 30 min, then concentrated under reduced pressure. The residue was partitioned between CH_2Cl_2 (50 mL) and saturated NaHCO_3 solution (50 mL). The organic phase was separated, washed with brine, dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was triturated with a mixture of EtOAc (1 mL) and petroleum ether (20 mL) to provide compound **3a** (2.31 g, 85% yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 7.74 (d, 1 H), 6.94 (d, 1 H), 5.19-5.21 (m, 1 H), 4.95 (d, 1 H), 4.48-4.51 (m, 1 H), 3.54-3.57 (m, 2 H), 2.72-2.84 (m, 8 H), 2.20-2.27 (m, 2 H), 1.70-1.77 (m, 2 H), 1.45 (d, 3 H). LC-MS: 470 $[\text{M}+1]^+$.


Step 2:

A mixture of **3a** (3 g, 6.38 mmol), Trimethylsilylacetylene (3.1 g, 31.9 mmol), $\text{Pd(PPh}_3)_2\text{Cl}_2$ (210 mg, 0.3 mmol) and CuI (85 mg, 0.45 mmol) in TEA (60 mL) was heated at 80 °C under N_2 overnight, then cooled, diluted with CH_2Cl_2 (40 mL) and filtered. The filtrate was concentrated under reduced pressure and the residue was partitioned between EtOAc (40 mL) and water (40 mL). The organic phase was separated, dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:30, 1% NH_4OH) to provide 2.4 g of light yellow solid which was dissolved in a suspension of K_2CO_3 (0.75 g, 5.45 mmol) in MeOH (40 mL) and stirred at room temperature for 30 min. The mixture was filtered and concentrated under reduced pressure and the residue

5 was partitioned between EtOAc (40 mL) and water (40 mL). The organic phase was separated, dried with anhydrous Na₂SO₄ and concentrated under reduced pressure to provide **H0603** (1.9 g, 82% yield) as a white powder. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.43 (d, 1 H), 7.21 (d, 1 H), 5.27-5.31 (m, 1 H), 4.81 (d, 1 H), 4.09-4.17 (m, 1 H), 3.38 (s, 1 H), 2.86-2.91 (m, 2 H), 2.80 (s, 3 H), 2.27 (s, 3 H), 1.98-2.09 (m, 2 H), 1.61-1.65 (m, 2 H), 1.48-1.52 (m, 2 H), 1.46 (d, 3 H). LC-
10 MS: 368 [M+1]⁺.

Example 4

Synthesis of Compound H0700




A mixture of **3a** (3.0 g, 6.38 mmol), **3b** (3.54 g, 9.57 mmol), CuI (243 mg, 1.27 mmol) and Pd(PPh₃)₄ (1.47 g, 1.27 mmol) in 1,2-dimethoxyethane (60 mL) was heated at 100 °C under N₂ overnight, then diluted with CH₂Cl₂ (100 mL) and filtered. The filtrate was washed with brine
20 (100 mL). The organic phase was separated, dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, MeOH: CH₂Cl₂ 1:30, 1% NH₄OH) to provide **H0700** (1.3 g, 48% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.90 (d, 1 H), 8.66-8.67 (m, 1 H), 8.58 (d, 1 H), 7.45 (d, 1 H), 7.38 (d, 1 H), 5.35-5.39 (m, 1 H), 4.87 (d, 1 H), 4.13-4.14 (m, 1 H), 2.85-2.90 (m, 2 H), 2.81 (s, 3 H), 2.26
25 (s, 3 H), 1.98-2.05 (m, 2 H), 1.69-1.77 (m, 2 H), 1.54-1.64 (m, 2 H), 1.51 (d, 3 H). LC-MS: 422 [M+1]⁺.

Example 5

Synthesis of Compound H0722


30



A mixture of compound **4a** (1.39 g, 4.08 mmol), **2b** (1.0 g, 6.1 mmol), DPPA (1.23 g, 4.5 mmol) and TEA (3 mL) in dry toluene (100 mL) was heated under reflux overnight, then cooled and concentrated under reduced pressure. The residue was partitioned between EtOAc (50 mL) and saturated Na₂CO₃ solution (50 mL). The organic phase was separated, washed with brine (50 mL), dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:40, 1% NH₄OH) to provide **H0722** (1.03 g, 55% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ = 8.89 (d, 1 H), 8.66-8.67 (m, 1 H), 8.58 (d, 1 H), 7.43 (d, 1 H), 7.37 (d, 1 H), 5.35-5.38 (m, 1 H), 5.21 (d, 1 H), 4.15-4.17 (m, 1 H), 2.85-2.90 (m, 2 H), 2.83 (s, 3 H), 2.26 (s, 3 H), 1.97-2.05 (m, 2 H), 1.66-1.80 (m, 6 H), 0.68-0.70 (m, 1 H), 0.50-0.54 (m, 2 H), 0.14-0.15 (m, 2 H) LC-MS: 462 [M+1]⁺.

Example 6

Synthesis of Compound H0751



Step 1:

The mixture of **5a** (5 g, 30.5 mmol), Trimethylsilylacetylene (3.6 g, 36.6 mmol), Pd(PPh₃)₂Cl₂ (210 mg, 0.3 mmol) and CuI (85 mg, 0.45 mmol) in TEA (150 mL) was heated at 80 °C for 3h

5 under N₂, then cooled, diluted with Et₂O (100 mL) and washed with brine (100 mL). The organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc/petroleum ether 1:15) to provide **5b** (4.3 g, 79% yield) as a yellow oil. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.74 (d, 1H), 7.53 (d, 1H), 0.26 (s, 9H)

10 Step 2:


To a solution of compound **5b** (4.1g, 22.5 mmol) in TBME (100 mL) at room temperature was added Bu₄NF (1 M in THF) (22.5 ml, 22.5 mmol). The mixture was stirred at room temperature for 30 min, then quenched with water (100 mL). The organic phase was separated, dried over
15 anhydrous Na₂SO₄ and filtered to afford crude compound **7c** in TBME (80 mL) which was used directly in next step without further purification.

Step 3:

A solution of crude compound **5c** in TBME was added to a mixture of **3a** (3 g, 6.3 mmol),
20 Pd(PPh₃)₂Cl₂ (660 mg, 0.95 mmol), CuI (180 mg, 0.95 mmol) in DMF (50 ml) and TEA(10 mL). The mixture was heated at 110 °C under N₂ overnight in a sealed tube, then cooled, diluted with CH₂Cl₂ (100 mL) and filtered. The filtrate was washed with brine (100 mL) and the organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:30,
25 1% NH₄OH) to provide **H0751** (1.18 g, 40% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.76 (d, 1 H). 7.59 (d, 1 H), 7.42 (d, 1 H), 7.16 (d, 1 H), 5.22-5.26 (m, 1 H), 4.73-4.74 (d, 1 H), 4.03-4.09 (m, 1 H), 2.81 (br, 2 H), 2.73 (s, 3 H), 2.19 (s, 3 H), 1.91-1.99 (m, 2 H), 1.63-1.69 (m, 2 H), 1.52-1.62 (m, 2 H), 1.41 (d, 3 H). LC-MS: 451 [M+1]⁺.

30 Example 7


Synthesis of Compound H0754



A mixture of **H0603** (2.2 g, 6 mmol), **6a** (2.97 g, 18 mmol), $\text{Pd(PPh}_3)_2\text{Cl}_2$ (0.66 g, 0.9 mmol) and CuI (264 mg, 1.38 mmol) in TEA (50 mL) was heated at 65°C under N_2 overnight, then cooled, diluted with CH_2Cl_2 (100 mL) and filtered. The filtrate was concentrated under reduced pressure and the residue was partitioned between EtOAc (50 mL) and water (50 mL). The organic phase was separated, dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:30, 1% NH_4OH) to provide **H0754** (990 mg, 37% yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): $\delta = 7.91$ (d, 1 H), 7.54 (d, 1 H), 7.46 (d, 1 H), 7.22 (d, 1 H), 5.32-5.26 (m, 1 H), 4.99 (d, 1 H), 4.47-4.60 (m, 1 H), 3.40-3.62 (m, 2 H), 2.88 (s, 3 H), 2.76-2.91 (m, 2 H), 2.82 (s, 3 H), 1.70-1.90 (m, 4 H), 1.51 (d, 3 H). LC-MS: 451 $[\text{M}+1]^+$.

Example 8

Synthesis of Compound H0761




A mixture of compound **4a** (2.3 g, 6.78 mmol), DPPA (1.86 g, 6.78 mmol) and TEA (10.2 mL) in dry toluene (200 mL) was stirred at 110°C for 2 h, then cooled to room temperature and compound **7a** (1.75 g, 13.56 mmol) was added. The mixture was stirred at room temperature overnight, and then concentrated under reduced pressure. The residue was partitioned between EtOAc (100 mL) and saturated Na_2CO_3 solution (100 mL). The organic phase was separated, washed with brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:30, 1% NH_4OH) to provide **H0761** (1.4 g, 48.3% yield) as a white solid. $^1\text{H-NMR}$

- 5 NMR (CDCl₃, 400 MHz): δ = 10.11 (s, 1 H), 8.91 (d, 1 H), 8.66 (m, 1 H), 8.57 (d, 1 H), 7.46 (d, 1 H), 7.36 (d, 1 H), 6.84 (d, 1 H), 5.35 (m, 1 H), 3.97-4.04 (m, 1 H), 2.86-2.93 (m, 2 H), 2.25 (s, 3 H), 1.93-2.13 (m, 4 H), 1.79-1.86 (m, 1 H), 1.64-1.72 (m, 2 H), 1.55-1.58 (d, 1 H), 0.65-0.70 (m, 1 H), 0.46-0.50 (m, 2 H), 0.11-0.14 (m, 2 H). LC-MS: 464 [M+1]⁺.

10 Example 9

Synthesis of Compound H0764




- 15 To a solution of **3a** (2.0 g, 4.26 mmol) and **8b** (1.4 g, 21.2 mmol) in dry THF (10 mL) and TEA (1.8 g, 17 mmol) was added Pd(PPh₃)₂Cl₂ (597 mg, 0.85 mmol) and CuI (220 mg, 1.16 mmol) at room temperature under N₂. The mixture was heated at 80 °C overnight in a sealed tube, then cooled, diluted with CH₂Cl₂ (50 mL) and filtered. The filtrate was washed with brine (50 mL) and the organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated under
- 20 reduced pressure. The residue was purified by column chromatography (silica, methanol: dichloromethane 1:30, 1% NH₄OH) to provide **H0764** (990 mg, 37% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ = 7.27 (d, 1 H), 7.12 (d, 1 H), 5.24-5.29 (m, 1 H), 4.78 (d, 1 H), 4.07-4.14 (m, 1 H), 2.74-2.88 (m, 2 H), 2.76 (s, 3 H), 2.24 (s, 3 H), 1.96-2.04 (m, 2 H), 1.40-1.73 (m, 5 H), 1.38 (d, 3 H), 0.70-0.90 (m, 4 H). LC-MS: 408 [M+1]⁺.

25

Example 10

Synthesis of Compound H0795



5

Step 1:

To a 2.5 M solution of *n*-butyllithium (40 mL, 0.1 mol) in anhydrous THF (250 mL) cooled to -78°C under N_2 protection was added **TMP** (2,2,6,6-tetramethylpiperidine, 15 g, 0.106 mol) dropwise over a period of 20 minutes. The mixture was warmed to 0°C by replacing the dry ice/acetone bath with an ice bath and stirred for 1.5 h. The mixture was cooled back to -78°C and a solution of **9a** (3 g, 0.03 mol) and tributyltin chloride (10 g, 0.03 mol) in 50 mL of dry THF was added over 10 min. The mixture was stirred at -78°C for 6 h, then warmed to -40°C by replacing the dry ice/acetone bath with an dry ice/acetonitrile bath. A solution of 35% HCl, ethanol and THF (1:4:5) was added. The mixture was warmed to room temperature and washed with saturated NaHCO_3 solution (100 mL) and extracted with EtOAc (3×100 mL). The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: petroleum ether=1:15) to provide **9b** (3.4 g, 29% yield) as light yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): δ = 8.41 (d, 1 H), 8.17 (d, 1 H), 1.8-0.53 (m, 27 H).

20

Step 2:


To a solution of **3a** (2.0 g, 4.4 mmol) and **9b** (3.4 g, 9.35 mmol) in 1,2-dimethoxyethane (200 mL) were added $\text{Pd}(\text{PPh}_3)_4$ (800 mg, 0.69 mmol) and CuI (40 mg, 0.21 mmol) at room temperature under N_2 . The mixture was then heated at 90°C overnight, then cooled, diluted with CH_2Cl_2 (100 mL) and filtered. The filtrate was washed with brine (100 mL) and the organic phase was separated, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, MeOH: CH_2Cl_2 , 1:30, 1% NH_4OH) to provide compound **H0795** (1.0 g, 51% yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz):

25

- 5 δ = 8.83 (d, 1 H), 8.44 (d, 1 H), 7.46 (d, 1 H), 7.22 (d, 1 H), 5.26-5.30 (m, 1 H), 4.99 (d, 1 H), 4.47-4.60 (m, 1 H), 2.90-2.95 (m, 2 H), 2.83 (s, 3 H), 2.32 (s, 3 H), 2.10-2.17 (m, 2 H), 1.78-1.83 (m, 2 H), 1.59-1.64 (m, 2 H), 1.51 (d, 3 H). LC-MS: 440 [M+1]⁺.

Example 11

10 Synthesis of H0816



Step 1:

- To a solution of **1k** (12.0 g, 38.1 mmol), sat. NaHCO_3 solution (120 mL) in THF (480 mL), was added $(\text{Boc})_2\text{O}$ (16.6 g, 76.2 mmol) at r.t.. Then the mixture was stirred at r.t. overnight. Ethyl acetate (500 mL) and water (500 mL) were added to the mixture. The organic layer was separated, washed with brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EA: PE=1:5) to provide **10b** (15.4 g, 97.5% yield) as a white solid. ^1H -NMR (CDCl_3 , 400 MHz): δ = 7.76 (d, 1H), 6.99 (d, 1H), 5.05 (s, 1H), 4.97 (s, 1H), 1.27 (s, 12H).

Step 2:

- To a solution of **10b** (5.0 g, 12.0 mmol) and **3b** (5.3 g, 14.4 mmol) in 1,2-dimethoxyethane (150 mL) were added $\text{Pd}(\text{PPh}_3)_4$ (1.39 g, 2.4 mmol), CuI (228 mg, 2.4 mmol) and LiCl (50.4 mg, 2.1 mmol) at r.t. under N_2 . The mixture was then heated at 105°C overnight, then cooled and concentrated under reduced pressure. Ethyl acetate (200 mL) and water (200 mL) were added to the above mixture which was then filtered. The organic phase was separated, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EA: PE=1:10) to provide compound **10c** (3.47 g, 78.5% yield).

5 as yellow solid. ¹H-NMR (CDCl₃, 300 MHz): δ= 8.93 (d, 1H), 8.69-8.70 (m, 1H), 8.60 (d, 1H), 7.48-7.51 (m, 1H), 7.42-7.45 (m, 1H), 5.19-5.23 (m, 1H), 5.06 (s, 1H), 1.45 (s, 12 H).

Step 3:


To a solution of **10c** (3.47g, 9.5 mmol) in DCM (100 mL) cooled to 0 °C was added TFA (35
10 mL) dropwise. The mixture was stirred at r.t for 1 h and then concentrated under reduced pressure. DCM (100 mL) was added to the above residue and cooled to 0 °C. Sat. Na₂CO₃ solution was added dropwise to the above mixture at 0 °C until pH=8. The organic layer was separated, washed with brine (200 mL), then dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, MeOH :
15 DCM =1:100) to provide **10d** (1.7 g, 68.0% yield) as a yellow solid. LC-MS: 268 [M+1]⁺.

Step 4:

To a solution of **10d** (1.7 g, 6.4 mmol) and TEA (17 mL) in DCM (340 mL), was added triphosgene (1.42 g, 4.8 mmol) in portions at 0 °C. The solution was then warmed to r.t. and
20 stirred for 0.5 h. **2b** (1.57 g, 9.6 mmol) was added to the above mixture at r.t. The mixture was then stirred for another 0.5 h, and finally evaporated under reduced pressure. EtOAc (150 mL) was added to the residue and washed with water (100 mL) and brine (100 mL). The separated organic phase was dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica, MeOH: DCM =1:10) to provide **H0816** (2.04 g, 75.8% yield) as
25 a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.82 (s, 1H), 8.60 (s, 1H), 8.51 (d, 1H), 7.36-7.38 (m, 1H), 7.29-7.31 (m, 1H), 5.28-5.31 (m, 1H), 4.79 (d, 1H), 4.04-4.10 (m, 1H), 2.78-2.83 (m, 1H), 2.74 (s, 2H), 2.19 (s, 3H), 1.91-1.99 (m, 2H), 1.61-1.70 (m, 2H), 1.47-1.57 (m, 2H), 1.44 (d, 3H). LC-MS: 422 [M+1]⁺.

30 Example 12

Synthesis of H0824



5

Step 1:

To a solution of **1j** (2 g, 6.36 mmol) and di-tert-butyl dicarbonate (2.75 g, 12.72 mmol) in THF (30 mL) was added saturated aqueous Na_2CO_3 solution (5 mL) at 0 °C. The mixture was then stirred at room temperature for 1 h, and eventually diluted with ethyl acetate (40 mL). The resulting mixture was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was triturated with petroleum ether (40 mL) to provide **11b** (1.86 g, 70% yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ =7.76 (d, 1 H), 7.00 (d, 1 H), 4.96-5.06 (m, 2 H), 1.41-1.43 (m, 12 H). LC-MS: 416 $[\text{M}+1]^+$.

10

Step 2:

To a solution of **1b** (1.8 g, 4.5 mmol) and **3b** (2.4 g, 6.5 mmol) in 1,2-dimethoxyethane (160 mL) were added $\text{Pd}(\text{PPh}_3)_4$ (780 mg, 0.67 mmol) and CuI (90 mg, 0.45 mmol) at room temperature under the protection of N_2 . The mixture was then heated to 90 °C and stirred overnight at this temperature. It was subsequently cooled down and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (silica, ethyl acetate: petroleum ether 1:10) to provide **11c** (1.2 g, 73% yield) as a white solid. LC-MS: 368 $[\text{M}+1]^+$.

20

Step 3:

To a solution of **11c** (600 mg, 1.63 mmol) in dichloromethane (15 mL) was added trifluoroacetic acid (5 mL) at 0 °C. After the addition, the mixture was stirred at room temperature for 2 h and then concentrated under reduced pressure. The residue was partitioned between saturated aqueous NaHCO_3 solution (15 mL) and dichloromethane (20 mL). The organic layer was separated, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to provide **11d**

25

- 5 (350 mg, 80% yield) as a colorless oil. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 8.92 (d, 1 H), 8.69 (dd, 1 H), 8.59 (d, 1 H), 7.69 (d, 1 H), 7.49 (d, 1 H), 4.67-4.69 (m, 1 H), 1.43 (d, 3 H). LC-MS: 268 $[\text{M}+1]^+$.


Step 4:

- 10 To a solution of compound **11d** (60 mg, 0.225 mmol) and TEA (0.5 mL) in dichloromethane (10 mL) was added triphosgene (46 mg, 0.158 mmol) at 0 °C. The mixture was then stirred at room temperature for 15 min before the addition of **11e** (53 mg, 0.337 mmol). Then stirred for another 30 min, diluted with dichloromethane (10 mL), washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified with silica
- 15 gel column chromatography (silica, methanol: dichloromethane 1:40, 1% NH_4OH) to provide **H0824** (60 mg, 57% yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 8.84 (dd, 1 H), 8.61 (d, 1 H), 8.51 (d, 1 H), 7.37 (dd, 1 H), 7.30 (dd, 1 H), 5.23-5.27 (m, 1 H), 4.82 (dd, 1 H), 4.02 (d, 1 H), 2.86 (d, 2 H), 2.80 (s, 3 H), 2.23 (d, 3 H), 1.90-2.01 (m, 2 H), 1.76 (d, 1 H), 1.45 (d, 3 H), 1.40 (d, 1 H), 1.05 (s, 3 H), 0.70 (s, 3 H). LC-MS: 450 $[\text{M}+1]^+$.

20

Example 13

Synthesis of H0890 (enantiomer of H0824)




25

Step 1-4: Compound **H0890** was synthesized in a similar manner to **H0824** (overall yield 31% from **1k**). $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 8.91 (dd, 1 H), 8.68 (d, 1 H), 8.58 (d, 1 H), 7.46 (dd, 1 H), 7.40 (dd, 1 H), 5.30-5.34 (m, 1 H), 4.86 (d, 1 H), 4.09 (d, 1 H), 2.95 (d, 2 H), 2.87 (s, 3 H),

- 5 2.40 (d, 3 H), 2.46-2.51 (m, 2 H), 2.22 (s, 3 H), 2.01-2.09 (m, 2 H), 1.84 (d, 1 H), 1.51 (d, 3 H), 1.47 (d, 1 H), 1.08 (s, 3 H), 0.76 (s, 3 H). LC-MS: 450 [M+1]

Example 14

Synthesis of H0826



Step 1:

- A mixture of **13a** (3g, 26.5 mmol), EtNH₂·HCl (11.2 g, 132.7 mmol), TEA (5 ml) and Pd/C (300 mg) in MeOH (50 mL) was heated at 60 °C under H₂ (50 psi) overnight, then cooled and filtered. The filtrate was concentrated under reduced pressure and the residue was dissolved in
- 15 HCl/dioxane (4 N, 100 mL) and stirred for 30 min. The precipitate was filtered and washed with ethyl acetate (50 mL) to provide **13b** (4.1 g, 87% yield) as white powder. ¹H-NMR (DMSO-*d*₆, 400 MHz): δ= 9.12 (br, 2 H), 3.72 (d, 2 H), 3.25-3.29 (m, 1 H), 3.04 (q, 2 H), 2.84-2.90 (m, 2 H), 2.70 (s, 3 H), 2.22-2.28 (m, 2 H), 1.94-2.02 (m, 2 H), 1.26 (t, 3 H), LC-MS: 129 [M+1]⁺.

Step 2:


- To a solution of **11d** (60 mg, 0.225 mmol) and TEA (0.5 mL) in dichloromethane (5 mL) was added triphosgene (46 mg, 0.158 mmol) at 0 °C. After the addition, the mixture was stirred at room temperature for 15 min before the addition of **13b** (60 mg, 0.337 mmol). The resulting mixture was stirred for another 30 min at room temperature, then diluted with dichloromethane
- 25 (10 mL), washed with brine (10 mL). The organic layer was separated, dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified with silica gel column chromatography (silica, methanol: dichloromethane 1:40, 1% NH₄OH) to provide **H0826** (44 mg, 45% yield). ¹H-NMR (CDCl₃, 400 MHz): δ=8.89 (d, 1 H), 8.66 (dd, 1 H), 8.57 (d, 1 H), 7.45 (d, 1 H), 7.36 (d, 1 H), 5.36-5.39 (m, 1 H), 4.85 (d, 1 H), 4.13-4.18 (m, 1 H), 3.22 (q, 2 H), 2.84-2.88 (m, 2 H), 2.25 (s, 3 H), 1.95-2.03 (m, 2 H), 1.55-1.73 (m, 4 H), 1.53 (d, 3 H), 1.24 (t, 3 H).
- 30 H).. LC-MS: 436 [M+1]⁺.

5 **Example 15****Synthesis of H0889 (enantiomer of H0826)**

10

The synthesis of **H0889** (49 mg, 30% yield) is similar to that of **H0826**. ¹H-NMR (CDCl₃, 400 MHz): δ = 8.90 (d, 1 H), 8.67 (dd, 1 H), 8.57 (d, 1 H), 7.45 (d, 1 H), 7.37 (d, 1 H), 5.35-5.39 (m, 1 H), 4.85 (d, 1 H), 4.11-4.17 (m, 1 H), 3.22 (q, 2 H), 2.85-2.88 (m, 2 H), 2.25 (s, 3 H), 1.97-2.04 (m, 2 H), 1.54-1.73 (m, 4 H), 1.52 (d, 3 H), 1.23 (t, 3 H). LC-MS: 436 [M+1]⁺.

15

Example 16**Synthesis of H0830**

20

Step 1:

To a solution of **14a** (3.1 g, 7.88 mol) in dichloromethane (60 mL) was added Dess-Martin periodinane (5.0 g, 11.83 mmol) at room temperature. The mixture was stirred at room temperature for 2 h, then concentrated under vacuum. The residue was purified by column chromatography (silica, ethyl acetate: petroleum ether=1:15) to provide **14b** (3.05 g, 99% yield)

25

5 as a light yellow solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 10.40 (s, 1 H), 7.97 (d, 1 H), 7.52 (d, 1 H).

Step 2:

To a solution of **14b** (1.5 g, 3.8 mmol) and **3b** (2.12 g, 5.7 mmol) in 1,2-dimethoxyethane (40 mL) were added $\text{Pd}(\text{PPh}_3)_4$ (887 mg, 0.76 mmol) and CuI (147 mg, 0.76 mmol) at room temperature under the protection of N_2 . The mixture was heated at 90 °C overnight, and then concentrated under reduced pressure. The residue was purified with silica gel column chromatography (silica, ethyl acetate: petroleum ether=1:10) to provide **14c** (826 mg, 86% yield) as a light yellow solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 10.55 (s, 1 H), 8.97 (d, 1 H), 8.74 (dd, 1 H), 8.66 (d, 1 H), 7.98 (d, 1 H), 7.64 (d, 1 H). LC-MS: 253 $[\text{M}+1]^+$.

Step 3:

To a solution of **14c** (980 mg, 3.5 mmol) and (trifluoromethyl)trimethylsilane (1.1 g, 7.8 mmol) in THF (20 mL) was slowly added TBAF (1 M solution in THF, 5.8 mL, 5.8 mmol,) at 0 °C. After the mixture was stirred at room temperature overnight, water was added (30 mL). The resulting mixture was extracted with ethyl acetate (30 mL x 3). The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated reduced pressure. The residue was purified by column chromatography (silica, ethyl acetate: petroleum ether=1:5) to provide **14d** (640 mg, 52 % yield) as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 8.92 (s, 1 H), 8.72 (s, 1 H), 8.66 (s, 1 H), 7.75 (d, 1 H), 7.53 (d, 1 H), 5.70 (q, 1 H), 3.68(br, 1 H). LC-MS: 323 $[\text{M}+1]^+$.

Step 4:

To a solution of **14d** (750 mg, 2.33 mmol) and TEA (709 mg, 7.02 mmol) in dichloromethane (20 mL) was added methanesulfonyl chloride (320 mg, 2.8 mmol) at 0 °C. After the addition was finished, the mixture was stirred at room temperature for 20 min, then diluted with dichloromethane (50 mL). The mixture was washed with saturated aqueous NaHCO_3 solution (40 mL), dried over anhydrous Na_2SO_4 and concentrated under vacuum to provide crude **14e** (910 mg, 97% yield) as a colorless oil which was used in the next step without further purification. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ = 8.96 (d, 1 H), 8.73 (dd, 1 H), 8.67 (d, 1 H), 7.74 (d, 1 H), 7.64 (d, 1 H), 6.54 (q, 1 H), 3.15(s, 3 H).

5 Step 5:

To a solution of compound **14e** (910 mg, 2.27 mmol) in DMSO (20 mL) was added NaN₃ (296 mg, 4.55 mmol) at room temperature. The mixture was stirred at 100 °C overnight, then cooled and water was added (100 mL). The resulting mixture was extracted with ethyl acetate (50 mL x 3). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified by column chromatography (silica, ethyl acetate: petroleum ether=1:5, v:v) to provide **14f** (340 mg, 44% yield) as a yellow oil. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.89 (d, 1 H), 8.78 (dd, 1 H), 8.62 (d, 1 H), 7.74 (d, 1 H), 7.60 (d, 1 H), 6.02 (q, 1 H). LC-MS: 348 [M+1]⁺.


15 Step 6:

To a solution of **14f** (34.7 mg, 0.1 mmol), HCOOH (46 mg, 1.0 mmol) and N₂H₄·H₂O (50 mg, 1.0 mmol) in EtOH (10 mL) was added Raney-Ni (50 mg). The mixture was stirred at room temperature for 1 h, then filtered and concentrated under vacuum. The residue was diluted with dichloromethane (20 mL), washed with water (15 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum to provide **14g** (30 mg, 93% yield) as a colorless oil. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.92 (d, 1 H), 8.67 (dd, 1 H), 8.61 (d, 1 H), 7.67 (d, 1 H), 7.55 (d, 1 H), 5.17 (q, 1 H), 1.86 (br, 2 H). LC-MS: 322 [M+1]⁺.

Step 7:

25 To a solution of **14g** (24 mg, 0.07 mmol), **2b** (14.7 mg, 0.09 mmol) and TEA (0.5 mL) in dichloromethane (10 mL) was added triphosgene (46 mg, 0.158 mmol) at room temperature. The resulting mixture was stirred at 35 °C under the protection of N₂ for 2 h, then diluted with dichloromethane (10 mL). The mixture was washed with saturated aqueous Na₂CO₃ solution (10 mL) and brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified with silica gel column chromatography (silica, methanol: dichloromethane 1:40, 1% NH₄OH) to provide **H0830** (10 mg, 28% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.85 (d, 1H), 8.62 (dd, 1H), 8.55 (d, 1H), 7.48 (d, 1H), 7.40 (d, 1H), 6.22-6.26 (m, 1H), 5.21 (d, 1H), 4.38-4.45 (m, 1H), 3.30-3.12 (m, 2H), 2.84 (s, 3H), 2.59-2.71 (m, 5H), 1.61-1.66 (m, 2H), 1.01-1.05 (m, 2H). LC-MS: 476 [M+1]⁺.

35

5 **Example 17****Synthesis of H0847**10 **Step 1:**

To a solution of **12b** (10.4 g, 25 mmol) and **9b** (19.4 g, 50 mmol) in 1,2-dimethoxyethane (1.2 L) were added Pd(PPh₃)₄ (4.54 g, 3.92 mmol) and CuI (227 mg, 1.19 mmol) at r.t. under N₂. The mixture was heated at 90 °C overnight, then cooled, diluted with CH₂Cl₂ (800 mL) and filtered. The filtrate was washed with brine (600 mL) and the organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, EtOAc: Petroleum, 1:3) to provide crude compound **15a** (10.3 g, ca. 100% yield) as yellow solid. LC-MS: 386 [M+1]⁺.

Step 2:

20 To a solution of **15a** (10.3 g, 26 mmol) in DCM (500 mL) cooled to 0 °C was added TFA (100 mL) dropwise. After the addition was completed, the mixture was stirred for 3 h, then basified with saturated Na₂CO₃ solution (400 mL) and extracted with DCM (3×100 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica, MeOH : CH₂Cl₂ : NH₄OH, 1:20:0.01) to provide **15b** (4.1 g, 57% yield) as a red solid. LC-MS: 440 [M+1]⁺

Step 3:


To a solution of **15b** (2.0 g, 7.1 mmol) and TEA (80 mL) in CH₂Cl₂ (220 mL) was added triphosgene (1.52 g, 5.1 mmol) portion wise at 0 °C. After the addition was completed, the

- 5 solution was stirred for 45 min. **2b** (2.7 g, 7.1 mmol) was then added to the above solution. The resulting solution was stirred for 2 h, then diluted with CH₂Cl₂ (100 mL) and washed with aqueous Na₂CO₃ solution (100 mL) and brine (100 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (silica: CH₂Cl₂ : CH₃OH=10/1) to provide **H0847** (2.0 g, 65% yield) as white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.77 (d, 1H), 8.38 (d, 1H), 7.40 (d, 1H), 7.31 (d, 1H), 5.26-5.30 (m, 1H), 4.78 (d, 1H), 4.10-4.00 (m, 1H), 2.79-2.84 (m, 2H), 2.75 (s, 3H), 2.20 (s, 3H), 1.94-2.05 (m, 2H), 1.57-1.69 (m, 2H), 1.47-1.64 (m, 2H), 1.41 (d, 3H). LC-MS: 440 [M+1]⁺. ee%=98.5%. (Chiralpak, 5 μm, 4.6*250mm, Phase: Hex: EtOH: DEA = 90: 10: 0.2), retention time =12.829 min).

15

Example 18

Synthesis of H0829 and H0860



20

Step 1:

To a solution of **16a** (100 g, 0.54 mol) in DMF (1400 mL) was added N-chlorosuccinimide (73 g, 0.54 mol) slowly at 0 °C. The resulting mixture was heated at 40 °C for 12 h, then poured into

- 5 water (1600 mL). The precipitate was collected by filtration, dissolved in ethyl acetate (1000 mL) and washed with brine (1000 mL). Evaporation of the solvent gave the residue which was re-crystallized in ethanol to give crude **16b** (80 g) and it was used directly in next step.

Step 2:

- 10 To a well stirred solution of **16b** (80 g, 0.365 mol) in dry THF (4 L) was added LiAlH₄ (27.6 g, 0.73 mol) slowly at 0 °C. The mixture was stirred at 0 °C for 2 h. Then ice-water (600 mL) was slowly added at 0 °C and the mixture was filtered. The filtrate was concentrated and the residue was purified by re-crystallization in ethyl acetate/petroleum ether (1:2) to give **16c** (39 g, 56% overall yield in two steps) as a light yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 7.15 (d, 1H),
15 6.68 (d, 1H), 4.68 (d, 2H), 4.12 (br, 2H), 2.03 (br, 1H) LC-MS: 192 [M+1]⁺.

Step 3:

- To a mixture of **16c** (39 g, 0.2 mol) and ice (450 g) in con. HCl (200 mL) was added a solution of NaNO₂ (21.2 g, 0.3 mol) in water (30 mL) dropwise at 0 °C. The mixture was stirred at 0 °C
20 for 30 min, then a solution of KI (169.4 g, 1.02 mol) in water (400 mL) was added dropwise at 0 °C. The mixture was stirred at 0 °C for 40 min, then ethyl acetate (1000 mL) was added and the organic phase was washed successively with water (500 mL), NaHSO₃ solution (500 mL) and brine (500 mL). The organic phase was separated, dried with anhydrous Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica, EA: PE=1:15) to
25 provide **16d** (50 g, yield: 81%). ¹H-NMR (CDCl₃, 400 MHz): δ= 7.81 (d, 1H), 7.17 (d, 1H), 4.75 (d, 2H), 2.02 (br, 1H).

Step 4:

- To a mixture of **16d** (50 g, 166 mmol) and TEA (50 g, 497.0 mmol) in dry CH₂Cl₂ (900 mL) was
30 added methanesulfonyl chloride (22.8 g, 199.0 mmol) dropwise at 0 °C. The mixture was stirred at 0 °C for another 90 min, then diluted with ethyl acetate (800 mL) and washed with brine (600 mL). The organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated to afford crude **16e** (59 g) which was used directly in next step without further purification.

- 35 **Step 5:**

5 To a solution of crude **16e** (59 g, 160 mmol) in EtOH (1200 mL) was added a solution of NaCN (11.4 g, 230.0 mmol) in H₂O (250 mL). The resulting mixture was heated under reflux overnight, then cooled and concentrated. The residue was partitioned between ethyl acetate (500 mL) and water (500 mL). The organic phase was separated, washed with brine, dried over anhydrous Na₂SO₄ and concentrated to afford crude **16f** (40 g) as a brown solid which was used directly in
10 next step without further purification.

Step 6:

To a solution of **16f** (40 g, 129 mmol) in MeOH (360 mL) was added conc. H₂SO₄ (114 mL) dropwise at 0 °C. The mixture was then heated under reflux overnight, then cooled and
15 concentrated. Aqueous Na₂CO₃ solution (50 mL) was added to the residue at 0 °C and the mixture was adjusted to pH=9-10 with the addition of Na₂CO₃ powder. The mixture was extracted with ethyl acetate (3 x 300 mL) and the combined organic layers were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by column chromatography (silica, EA: PE =1:20) to **16g** (22 g, yield: 70.5%) as a yellow solid. ¹H-NMR (CDCl₃, 400
20 MHz): δ= 7.75 (d, 1H), 6.93 (d, 1H), 3.78 (s, 2H), 3.72 (s, 3H).

Step 7:

To a solution of **16g** (22 g, 32 mmol) in DMF (150 mL) was slowly added NaH (60%, 2.8 g, 2.2 mmol) at 0 °C. The mixture was stirred at r.t. for 30 min and then EtI (10 g, 64 mmol) was
25 added. The mixture was stirred at r.t. for another 1.5 h, then poured into ice water (600 mL). The resulting mixture was extracted with ethyl acetate (3 x 400 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified by column chromatography (silica, ethyl acetate: petroleum ether=1:50) to provide **16h** (20 g, 84% yield). ¹H NMR (CDCl₃, 400 MHz): δ=7.76 (d, 1H), 7.00 (d, 1H), 4.06 (t, 1H), 3.67 (s, 3H),
30 2.05-2.12 (m, 1H), 1.75-1.82 (m, 1H), 0.91 (t, 3H).

Step 8:

To a solution of **16h** (22 g, 53.7 mmol) and **3b** (25.9 g, 69.9 mmol) in 1,2-dimethoxyethane (660 mL) were added Pd(PPh₃)₄ (15.5 g, 13.4 mmol), LiCl (0.46 g, 13.4 mmol) and CuI (2.06 g, 10.8
35 mmol) at r.t. under the protection of N₂. The mixture was then heated at 105 °C overnight, cooled

5 and concentrated under vacuum. The residue was purified with silica gel column chromatography (silica, ethyl acetate: petroleum ether=1:8) to provide **16i** (12 mg, 69 % yield) as a yellow solid.

Step 9:

10 The mixture of **16i** (12 g, 37.0 mmol) and LiOH·H₂O (9.3 g, 22.2 mmol) in MeOH (480 mL) and H₂O (120 mL) was stirred at r.t. overnight, then concentrated under vacuum. The residue was acidified with 1N HCl to pH=2 which was extracted with dichloromethane (3 x 200 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated to provide **16j** (10.8 g, 94% yield) as a white solid. LC-MS: 310 [M-1]⁻.

15

Step 10:

The mixture of **16j** (10.8 g, 34.8 mmol), **2b** (8.6 g, 52 mmol), DPPA (11.5 mg, 41.8 mmol) and TEA (48 mL) in toluene (400 mL) was stirred at 125 °C overnight, then cooled and concentrated under vacuum. The residue was partitioned between saturated aqueous Na₂CO₃ solution (150 mL) and dichloromethane (300 mL). The organic phase was separated, washed with brine (200mL), dried with anhydrous Na₂SO₄ and concentrated under vacuum. The residue was purified by column chromatography (silica, MeOH: dichloromethane 1:50, 1% NH₄OH) to provide **H0829** (6 g, 41% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.91 (d, 1H), 8.68 (d, 1H), 8.59 (d, 1H), 7.45 (d, 1H), 7.34 (d, 1H), 5.17-5.22 (m, 1H), 4.93 (d, 1H), 4.11-4.17 (m, 1H), 2.85-2.92 (m, 2H), 2.82 (s, 3H), 2.27 (s, 3H), 1.58-2.05 (m, 8 H), 1.00 (t, 3H). LC-MS: 436 [M+1]⁺.


25

Step 11:

H0860 (2.0, 66.7%) was obtained through the chiral separation of **H0829** (Chiralpak, 5μm, 4.6* 250 mm, Hex:EtOH:DEA=80:20:0.2, retention time: 10.76 min). ¹H-NMR (CDCl₃, 400 MHz): δ= 8.89 (d, 1H), 8.66 (d, 1H), 8.57 (d, 1H), 7.43 (d, 1H), 7.32 (d, 1H), 5.16-5.21 (m, 1H), 4.92 (d, 1H), 4.11-4.17 (m, 1H), 2.87-2.90 (m, 2H), 2.81 (s, 3H), 2.26 (s, 3H), 1.48-2.01 (m, 8 H), 0.97 (t, 3H). LC-MS: 436 [M+1]⁺.

30

35 **Example 19**

5 **Synthesis of H0837 and H0862****Step 1**

- 10 The mixture of **17a** (5g, 27.0 mmol), 30% of methyl amine in methanol (50 mL) and 5% Pd/C (500 mg) in methanol (50 mL) was heated at 60 °C under H₂ (50 psi) overnight, then cooled and filtered. The filtrate was concentrated under vacuum and the residue was purified by silica gel column chromatography (methanol:dichloromethane =1:40) to provide **17b** (2.8 g, 52 % yield).
¹H-NMR (CDCl₃, 400 MHz): δ= 9.99 (s, 1 H), 3.79 -3.83 (m, 1 H), 3.61-3.72 (m, 3 H), 3.40 (d, 1 H), 2.71 (s, 3 H), 2.33-2.36 (m, 2 H), 1.75 (s, 9 H), LC-MS: 201 [M+1]⁺
- 15

Step 2:

- To a solution of **12c** (300 mg, 1.12 mmol) and TEA (3.6 g, 40.3 mmol) in dichloromethane (20 mL) was added triphosgene (283 mg, 0.95 mmol) at 0 °C. After the addition was finished, the mixture was stirred at room temperature for 30 min before the addition of **17b** (270 mg, 1.35 mmol). The resulting mixture was stirred at room temperature for 1 h, then concentrated under
- 20

5 vacuum. The residue was partitioned between dichloromethane (50 mL) and saturated NaHCO₃ solution (50 mL). The organic phase was separated, washed with brine, dried with anhydrous Na₂SO₄ and concentrate under vacuum. The residue was purified with silica gel column chromatography (silica, methanol: dichloromethane 1:40, 1% NH₄OH) to provide **17c** (330 mg, 60% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ = 8.82 (s, 1 H), 8.63 (d, 1 H), 8.51 (dd, 1 H), 7.38 (d, 1 H), 7.33 (d, 1 H), 5.23-5.26 (m, 1 H), 4.99 (d, 1 H), 4.80-4.83 (m, 1 H), 3.31-3.32 (m, 2 H), 3.03-3.23 (m, 2 H), 2.80 (s, 3 H), 1.97-2.03 (m, 1 H), 1.76 -1.84 (m, 1 H), 1.64 (s, 9 H), 1.45 (d, 3 H). LC-MS: 494 [M+1]⁺.

10

Step 3:

15 To a solution of **17c** (330 mg, 0.67 mmol) in dichloromethane (15 mL) was added trifluoroacetic acid (5 mL) dropwise at 0 °C. The mixture was stirred at room temperature for 1 h, then concentrated under vacuum. The residue was partitioned between aqueous NaHCO₃ solution and dichloromethane. The organic layer were dried over anhydrous Na₂SO₄ and concentrated to provide **17d** (252 mg, 96% yield) as a yellow solid. LC-MS: 394 [M+1]⁺.

20

Step 4:

To a mixture of **17d** (252 mg, 0.64 mmol) and 37% aqueous HCHO solution (250 mg, 3.1 mmol) in MeOH (15 mL) were added NaOAc (600 mg, 7.3 mmol), AcOH (1 mL, 50 mmol) and NaBH₃CN (121 mg, 1.9 mmol) at room temperature. The mixture was stirred at room temperature overnight, and then concentrated under reduced pressure. The residue was partitioned between dichloromethane (50 mL) and saturated NaHCO₃ solution (50 mL). The organic phase was separated, washed with brine, dried with anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (silica, methanol: dichloromethane 1:50, 1% NH₄OH) to provide **H0837** (200 mg, 77% yield) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): δ = 8.82 (d, 1 H), 8.60 (dd, 1 H), 8.50 (d, 1 H), 7.97 (br, 1 H), 7.38 (d, 1 H), 7.28-7.31 (m, 1 H), 5.26-5.31 (m, 1 H), 4.08-4.10 (m, 1 H), 3.03-3.06 (m, 1 H), 2.95-2.99 (m, 2 H), 2.90 (s, 3H), 2.19-2.35 (m, 5 H), 1.94-1.98 (m, 2H), 1.37-1.40 (m, 3 H). LC-MS: 408 [M+1]⁺.

30


Step 5:

- 5 **H0862** was obtained through the chiral separation of **H0837** (Chiralcel OJ-H, 5 μ m, 4.6 x 250 mm, Hex:EtOH:DEA=90:10:0.3, retention time: 11.34 min). ¹H-NMR (CDCl₃, 400 MHz): δ = 8.82 (d, 1 H), 8.60 (dd, 1 H), 8.51 (d, 1 H), 7.98 (br, 1 H), 7.37 (d, 1 H), 7.30 (d, 1 H), 5.28-5.31 (m, 1 H), 4.07-4.10 (m, 1 H), 3.06-3.10 (m, 1 H), 2.99-3.06 (m, 1 H), 2.90 (s, 3H), 2.20-2.35 (m, 5 H), 1.96-2.05 (m, 2H), 1.38 (d, 3 H). LC-MS: 408 [M+1]⁺.

10

Example 20

Synthesis of H0900



15

Step 1:

- To a mixture of **16d** (32 g, 120 mmol) in dry CH₂Cl₂ (800 mL) was added Dess-Martin peroxide reagent (76 g, 180 mmol) portion-wise at 0 °C. The mixture was stirred at room temperature for 1 h, then diluted with DCM (800 mL), washed with aqueous NaHCO₃ solution (300 mL) and
- 20 brine (300 mL). The organic phase was separated, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude **18a** (31.4 g) which was used directly in the next step without further purification.

5 Step 2:

To a solution of **18a** (12 g, 40 mmol) and **3b** (22.2 g, 60 mmol) in DME (560 mL) were added Pd(PPh₃)₄ (9.25 g, 8 mmol) and CuI (1.52 g, 8 mmol) at room temperature. The mixture was stirred at 90 °C overnight, then concentrated under reduced pressure. The residue was purified with silica gel column chromatography (silica, EA : PE = 1:5) to provide **18b** (8.0 g, 79.3%) as a
10 white solid. LC-MS: 253 [M+1]⁺.

Step 3:

To a solution of **18b** (7 g, 27.7 mmol) and (*S*)-tert-butylsulfonamide (7.27 g, 30.56 mmol) in dry THF (200 mL) was added Ti(i-OPr)₄ (15.7 g, 55.4 mmol) dropwise at room temperature. The
15 mixture was stirred at 80 °C overnight, and then cooled. Ethyl acetate (40 mL) was added, the resulting mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was purified with silica gel column chromatography (silica, EA:PE =1:5) to provide **18c** (6.8 g, 69%) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ= 9.10 (s, 1H), 8.97 (s, 1H), 8.72 (s, 1H), 8.64 (d, 1H), 8.12 (d, 1H), 7.59 (d, 1H), 1.30 (s, 9H). LC-MS: 356 [M+1]⁺.

20

Step 4:

To a stirred solution of **18c** (6.8 g, 19 mmol) and Tetrabutylammonium difluorotriphenylsilicate (15.8 g, 29 mmol) in dry THF (250 mL) was added a solution of TMSCF₃ (11 g, 77 mmol) in anhydrous THF (50 mL) at -65 °C. The mixture was then stirred at -65 °C for 2 h, and at that
25 point aqueous NH₄Cl solution (250 mL) was added. The mixture was diluted with ethyl acetate (250 mL), washed with brine (250 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified with silica gel column chromatography (silica, EA : PE=1:2) to provide **18d** (4.3 g, 52%) as a yellow solid. LC-MS: 426 [M+1]⁺.

30 Step 5:

To a stirred solution of **18d** (4.3 g, 10.1 mmol) in MeOH (40 mL) was added a solution of HCl/MeOH (4N, 40 mL) at room temperature. The mixture was stirred for 1 h, then concentrated under reduced pressure. The residue was triturated with ethyl acetate (40 mL) to afford crude **18e** (4.3g) which was directly in the next step without further purification. LC-MS: 322 [M+1]⁺.

35

5 Step 6:

To a solution of **18e** (2.7 g, 7.1 mmol), **2b** (3.4 g, 21.3 mmol) and TEA (80 mL) in DCM (220 mL) was added thiphosgene (3.15 g, 10.6 mmol) in DCM (40 mL) dropwise at 0 °C. The solution was warmed to ambient temperature and stirred for 1 h, then diluted with DCM (100 mL) and washed with aqueous Na₂CO₃ solution (100 mL) and brine (100 mL). The organic layer
10 was separated, dried over anhydrous Na₂SO₄ and concentrated. The residue was purified with silica gel column chromatography (silica, DCM : CH₃OH=10 : 1) to provide crude **H0900** (2.13 g, ee%=92.5%) which was further purified through chiral separation to afford **H0900** (1.6 g, 49% yield) as a white solid. (ee%=98.5%, Chiralpak IC 5um, 4.6*250mm, Phase: Hex: EtOH: DEA=90:10:0.2), retention time =12.829 min. ¹H-NMR (CDCl₃, 400 MHz): δ= 8.86 (d, 1H),
15 8.63 (dd, 1H), 8.55 (d, 1H), 7.47 (d, 1H), 7.40 (d, 1H), 6.28 (m, 1H), 5.18 (d, 1H), 4.12 (m, 1H), 2.88 (t, 2H), 2.77 (s, 3H), 2.22 (s, 3H), 2.05 (m, 2H), 2.48 (m, 2H), 1.52 (m, 2H), 1.73-1.49 (m, 4H). LC-MS: 476 [M+1]⁺.

20 Example A**Calcium FLIPR Assay**




The intracellular calcium assay was carried out in a 384-well format FLIPRTM (Molecular Device) HEK293/GHSR1a cell line. Cells were seeded 24 hr prior to the experiments at an optimal density per well. Preincubation with selected calcium dye lasted for 30-60 min at room
25 temperature or 37° C. Test compounds, dissolved in DMSO, were added at the appropriate time and incubated for 15 min followed by the addition of ghrelin with FlexStation or FLIPR. Relative fluorescence was monitored by the FLIPRTM Molecular Device. EC₅₀ and IC₅₀ values were estimated from dose-response data using GraphPad Prism software. To check for GHSR-1a agonism the compound was added at t=20 sec. and the calcium response was followed for 2
30 minutes. To check for GHSR-1a antagonism the compound and Ghrelin (10 nM) were added to the cells at t=20 sec. and the calcium response was measured for 2 minutes. The potency of the antagonist was calculated by its ability to reduce the ghrelin response. Dose-response curves were made for relevant antagonists.









Example B**35 Evaluation of GHSR1a Antagonists on Food Intake Test in Mouse**








5 Male C57BL/6J mice, 18-22 g body weight, were fasted overnight (16 h before compound administration) and placed in a regular light dark cycle (6:00-18:00 light/18:00-6:00 dark). After 1 wk acclimation, animals were sorted into two groups (n=6 each, 2 per cage) based on body weight. Animals in group one were be treated with vehicle and animals in group 2 were treated with the test agent (n=6 for each group). The cumulative food intake was evaluated at 1,
 10 2, 4, 8 and 24 hrs after drug or vehicle treatment. Food intake was measured by subtracting uneaten food from the initial premeasured food.








The following table presents representative compounds of Formula I with biological data including the ghrelin antagonist/agonist activity in vitro (Example A) and mouse food intake results (Example B). The data clearly demonstrates that compounds of Formula I are ghrelin
 15 receptor modulators and are useful in preventing and/or treating diseases associated with ghrelin receptor, for example, obesity.

TABLE 1








Compound No.	Chemical Structure	Activity	Metabolic Stability (H = Human; M = Mouse)	Mouse Food Intake (% Inhibition; Doses as mg/kg i.p.)*
H0494		IC ₅₀ = 52 nM EC ₅₀ = 66 nM E _{max} = 2996	Medium H High M	No Effect
H0621		IC ₅₀ > 30 μM EC ₅₀ = 2 nM E _{max} = 3896	High H High M	No Effect
H0496		IC ₅₀ = 10 nM EC ₅₀ > 30 μM	Medium H High M	NSE (10 mg/kg)








H0617		IC ₅₀ = 3.4 μM EC ₅₀ >30 μM	Not done	Not Done
H0539		IC ₅₀ = 9 nM EC ₅₀ >30 μM	Not done	Not Done
H0546		IC ₅₀ = 8 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0526		IC ₅₀ = 57 nM EC ₅₀ >30 μM	Not done	Not Done
H0527		IC ₅₀ = 19 nM EC ₅₀ >30 μM	Medium H Medium M	Not done
H0497		IC ₅₀ = 24 nM EC ₅₀ >30 μM	Medium H Medium M	NSE (30 mg/kg)
H0650		IC ₅₀ = 4 nM EC ₅₀ =9 nM E _{max} = 2150	Not Done	Not Done
H0849		IC ₅₀ = 37 nM EC ₅₀ =51 nM E _{max} = 1383	Not Done	Not Done








H0578		IC ₅₀ = 490 nM EC ₅₀ >30 μM	Not Done	Not Done
H0511		IC ₅₀ = 98 nM EC ₅₀ >30 μM	Medium H Medium M	94% at 1h Inhibition up to 24 h (30 mg/kg))
H0820		IC ₅₀ = 5.7 nM EC ₅₀ =9 nM E _{max} = 3955	Not Done	Not Done
H0613		IC ₅₀ = 20 nM EC ₅₀ >30 μM	High H High M	NSE (10 mg/kg) PO: inhibition at 1h, up to 2 h IP+ANAM 30 mpk PO: inhibition at 1h up to 24h
H0614		IC ₅₀ = 12 nM EC ₅₀ >30 μM	High H High M	No IP done, PO: NSE
H0635		IC ₅₀ = 1090 nM EC ₅₀ >30 μM	Not Done	Not Done
H0636		IC ₅₀ = 90 nM EC ₅₀ >30 μM	High H Medium M	Not Done








H0637		IC ₅₀ = 85 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0638		IC ₅₀ = 57 nM EC ₅₀ >30 μM	Not Done	Not Done
H0639		IC ₅₀ = 48 nM EC ₅₀ >30 μM	Medium H Medium M	NSE (10 mg/kg)
H0642		IC ₅₀ = 78 nM EC ₅₀ >30 μM	Very low H Very Low M	Not Done
H0704		IC ₅₀ = 19 nM EC ₅₀ >30 μM	High H Medium M	32% inhibition at 2 h (10 mg/kg)
H0705		IC ₅₀ = 53 nM EC ₅₀ >30 μM	High H High M	Not Done
H0707		IC ₅₀ = 185 nM EC ₅₀ >30 μM	Not Done	Not Done








H0711		IC ₅₀ = 1.85 μM EC ₅₀ > 30 μM	Not Done	Not Done
H0716		IC ₅₀ = 15 nM EC ₅₀ > 30 μM	Low H Medium M	Not Done
H0717		IC ₅₀ = 396 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0718		IC ₅₀ = 499 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0719		IC ₅₀ = 780 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0712		IC ₅₀ = 420 nM EC ₅₀ = 220 nM E _{max} = 1962	Not Done	Not Done
H0708		IC ₅₀ = 1.37 μM EC ₅₀ > 30 μM	Not Done	Not Done








H0714		IC ₅₀ = 453 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0715		IC ₅₀ = 57 nM EC ₅₀ = 42 nM E _{max} = 2479	Not Done	Not Done
H0706		IC ₅₀ = 116 nM EC ₅₀ = 91 nM E _{max} = 2111	Not Done	Not Done
H0710		IC ₅₀ = 275 nM EC ₅₀ = 395 nM E _{max} = 1621	Not Done	Not Done
H0666		IC ₅₀ = 8 nM EC ₅₀ = 21 nM E _{max} = 4927	Not Done	Not Done
H0739		IC ₅₀ = 39 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0667		IC ₅₀ < 1 nM EC ₅₀ = 3 nM E _{max} = 4887	High H High M	76% inhibition at 1h; activity up to 4 h (10 mg/kg)







H0821		IC ₅₀ = 2.3 μM EC ₅₀ >30 μM	Not Done	Not Done
H0646		IC ₅₀ = 541 nM EC ₅₀ >30 μM	Not Done	Not Done
H0720		IC ₅₀ = 8 nM EC ₅₀ >30 μM	Medium H High M	Not Done
H0721		IC ₅₀ = 20 nM EC ₅₀ >30 μM	Medium H High M	Not Done
H0516		IC ₅₀ = 41 nM EC ₅₀ >30 μM	High H High M	88% inhibition at 1 hour. Activity up to 24 h (30 mg/kg) PO: no effect
H0579		IC ₅₀ = 1 μM EC ₅₀ >30 μM	Not Done	Not Done
H0649		IC ₅₀ = 18 nM EC ₅₀ = 64 nM E _{max} = 1400	High H High M	48% inhibition at 1 and 2 h (10 mg/kg)








H0797		IC ₅₀ = 594 nM EC ₅₀ = 1.8 μM E _{max} = 2879	Not Done	Not Done
H0798		IC ₅₀ = 162 nM EC ₅₀ >30 μM	Not Done	Not Done
H0799		IC ₅₀ = 5.4 nM EC ₅₀ =14 nM E _{max} = 5031	Not Done	Not Done
H0800		IC ₅₀ = 1.3 μM EC ₅₀ >30 μM	Not Done	Not Done
H0801		IC ₅₀ = 20 nM EC ₅₀ = 45 nM E _{max} = 3915	Not Done	Not Done
H0802		IC ₅₀ = 99 nM EC ₅₀ =153 nM E _{max} = 4149	Not Done	Not Done
H0803		IC ₅₀ = 171 nM EC ₅₀ = 149 nM E _{max} = 2364	Not Done	Not Done







H0804		IC ₅₀ = 375 nM EC ₅₀ = 263 nM E _{max} = 2740	Not Done	Not Done
H0805		IC ₅₀ = 4 nM EC ₅₀ = 9 nM E _{max} = 5433	Not Done	Not Done
H0806		IC ₅₀ = 1.2 nM EC ₅₀ = 6.8 nM E _{max} = 5751	Not Done	Not Done
H0807		IC ₅₀ = 14 nM EC ₅₀ = 24 nM E _{max} = 3669	Not Done	Not Done
H0854		IC ₅₀ = 65 nM EC ₅₀ = 24 nM E _{max} = 3246	Not Done	Not Done
H0813		IC ₅₀ = 644 nM EC ₅₀ = 528 nM E _{max} = 1605	Not Done	Not Done
H0814		IC ₅₀ = 926 nM EC ₅₀ = 15 nM E _{max} = 1097	Not Done	Not Done








H0703		IC ₅₀ = 695 nM EC ₅₀ >30 μM	Not Done	Not Done
H0709		IC ₅₀ = 676 nM EC ₅₀ >30 μM	Not Done	Not Done
H0584		IC ₅₀ = 1.1 μM EC ₅₀ >30 μM	Not Done	Not Done
H0586		IC ₅₀ = 4.2 μM EC ₅₀ = 63 μM	Not Done	Not Done
H0587		IC ₅₀ >30 μM EC ₅₀ >30 μM	Not Done	Not Done
H0588		IC ₅₀ >30 μM EC ₅₀ >30 μM	Not Done	Not Done
H0663		IC ₅₀ = 274 nM EC ₅₀ >30 μM	Not Done	Not Done








H0620		IC ₅₀ =32 nM EC ₅₀ >30 μM	Poor H Poor M	Not Done
H0624		IC ₅₀ = 253 nM EC ₅₀ >30 μM	Not Done	Not Done
H0662		IC ₅₀ = >1 μM EC ₅₀ >30 μM	Not Done	Not Done
H0670		IC ₅₀ = 523 μM EC ₅₀ >30 μM	Not Done	Not Done
H0673		IC ₅₀ >1 μM EC ₅₀ >30 μM	Not Done	Not Done
H0727		IC ₅₀ = 3.6 μM EC ₅₀ >30 μM	Not Done	Not Done
H0631		IC ₅₀ = 719 nM EC ₅₀ >30 μM	Not Done	Not Done






H0686		IC ₅₀ = 14 nM EC ₅₀ >30 μM	Medium H High M	61% inhibition at 2h and 4h (10 mg/kg) PO: NSE
H0619		IC ₅₀ = 13 nM EC ₅₀ = 39 nM	High H High M	34% inhibition at 1 and 2 h (10 mg/kg)
H0768		IC ₅₀ = 279 nM EC ₅₀ >30 μM	Not Done	Not Done
H0808		IC ₅₀ = 674 nM EC ₅₀ = 90 nM E _{max} = 1494	Not Done	Not Done
H0700		IC ₅₀ = 7 nM EC ₅₀ >30 μM	High H High M	41% inhibition at 2 h; activity up to 4h (10 mg/kg) 71% inhibition at 1 h, activity up to 2 h (Fed, 10 mg/kg) PO SC: inhib at 1, 2 h
H0816		IC ₅₀ = 5.1 nM EC ₅₀ >30 μM	Not Done	12% inhib 30 mg/kg PO fasted mice








H0817		IC ₅₀ = 94 nM EC ₅₀ >30 μM	Not Done	30 mg/kg PO fasted mice NSE
H0722		IC ₅₀ = 13 nM EC ₅₀ >30 μM	High H High M	90% inhibition at 1 h; activity up to 24 h (10 mg/kg) PO: NSE (30 mg/kg)
H0741		IC ₅₀ = 15 nM EC ₅₀ >30 μM	Not Done	Not Done
H0752		IC ₅₀ = 100 nM EC ₅₀ >30 μM	Not Done	Not Done
H0743		IC ₅₀ = 94 nM EC ₅₀ >30 μM	Not Done	Not Done
H0750		IC ₅₀ = 177 nM EC ₅₀ >30 μM	Not Done	Not Done
H0756		IC ₅₀ = 13 nM EC ₅₀ = 13 nM E _{max} = 1729	Not Done	Not Done








H0761		$IC_{50} = 0.2 \text{ nM}$ $EC_{50} = 3 \text{ nM}$ $E_{max} = 2907$	High H High M	63% inhibition at 1 h; activity up to 8h (10 mg/kg) PO: NSE No activity in fed mice PO: 215% FI increase in fed mice. No activity in fasted mice
H0781	 (single enantiomer)	$IC_{50} = 95 \text{ nM}$ $EC_{50} = 420 \text{ nM}$ $E_{max} = 4210$	Not Done	Not Done
H0782	 (single enantiomer)	$IC_{50} = 5 \text{ nM}$ $EC_{50} = 6 \text{ nM}$ $E_{max} = 1923$	Not Done	93% inhibition at 1 h, activity up to 24 h (10 mg/kg)
H0824		$IC_{50} = 3 \text{ nM}$ $EC_{50} > 30 \text{ }\mu\text{M}$	High H Medium M	PO 30 mg/kg + ANA mice: NSE
H0890		$IC_{50} = 1.6 \text{ nM}$ $EC_{50} > 30 \text{ }\mu\text{M}$	High H High M	Not Done
H0858		$IC_{50} = 8 \text{ nM}$ $EC_{50} > 30 \text{ }\mu\text{M}$	Medium H Medium M	Not Done








H0865		IC ₅₀ = 6 nM EC ₅₀ >30 μM	Medium H Low M Medium R	Not Done
H0825		IC ₅₀ = 10 nM EC ₅₀ >30 μM	Medium H Medium M Medium R	Not Done
H0826		IC ₅₀ = 5 nM EC ₅₀ >30 μM	High H High M High R Medium D	Not Done
H0889		IC ₅₀ = 6 nM EC ₅₀ >30 μM	High H High M	Not Done
H0896		IC ₅₀ = 7 nM EC ₅₀ >30 μM	Not Done	Not Done
H0827		IC ₅₀ = 35 nM EC ₅₀ >30 μM	Not Done	Not Done
H0829		IC ₅₀ = 3nM EC ₅₀ >30 μM	High H High M	PO 10 mg/kg + ANA 30 mg/kg mice: NSE








H0859		$IC_{50} = 2.2 \mu M$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0860		$IC_{50} = 3 nM$ $EC_{50} > 30 \mu M$	High H High M	68% inhib 1 h PO 10 mg/kg + ANA 30 mg/kg mice
H0922		$IC_{50} = 2.8 \mu M$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0924		$IC_{50} = 300 nM$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0830		$IC_{50} = 3 nM$ $EC_{50} > 30 \mu M$	High H High M High R Medium D	Not Done TBD
H0899		$IC_{50} = 1.6 \mu M$ $EC_{50} > 30 \mu M$	Medium H High M	Not Done
H0900		$IC_{50} = 3 nM$ $EC_{50} > 30 \mu M$	Medium H High M	60% inhib 1 h PO 10 mg/kg + ANA 30 mg/kg fed mice 91% inhib 1 h PO 30 mg/kg +








				ANA 30 mg/kg fed mice 26% inhib 1 h PO 30 mg/kg fasted mice 90% inhib 1 h PO 30 mg/kg fed mice
H0909		$IC_{50} = 12nM$ $EC_{50} > 30 \mu M$	Medium H High M	Not Done
H0856		$IC_{50} = 339nM$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0837	 (diastomeric mixture)	$IC_{50} = 2nM$ $EC_{50} > 30 \mu M$	High H High M High R High D	180% increase 2 h mice 30 mg/kg PO PO 10 mg/kg + ANA 30 mg/kg mice: NSE
H0861	 (R/S) (single diastereoisomer)	$IC_{50} = 189 nM$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0862	 (R/S) (single diastereoisomer)	$IC_{50} = 3nM$ $EC_{50} > 30 \mu M$	High H medium M	PO (10 mg/kg) + ANA: no activity in mice








H0857		IC ₅₀ = 10nM EC ₅₀ >30 μM	Medium H Low M	Not Done
H0871		IC ₅₀ = 9nM EC ₅₀ >30 μM	Not Done	Not Done
H0874		IC ₅₀ = 115nM EC ₅₀ >30 μM	Not Done	Not Done
H0853		IC ₅₀ = 1.5 μM EC ₅₀ >30 μM	Not Done	Not Done
H0815		IC ₅₀ = 176 nM EC ₅₀ >30 μM	Not Done	Not Done
H0831		IC ₅₀ = 1.2 μM EC ₅₀ >30 μM		
H0843		IC ₅₀ = 35 nM EC ₅₀ = 51 nM E _{max} = 1910	Not Done	Not Done

H0844		IC ₅₀ = 705 nM EC ₅₀ >30 μM		
H0738		IC ₅₀ = 696 nM EC ₅₀ >30 μM	Not Done	Not Done
H0780		IC ₅₀ = 63 nM EC ₅₀ >30 μM	Not Done	Not Done
H0786		IC ₅₀ = 855 nM EC ₅₀ = 242 nM E _{max} = 980	Not Done	Not Done
H0791		IC ₅₀ = 75 nM EC ₅₀ >30 μM	Not Done	Not Done
H0795		IC ₅₀ = 4 nM EC ₅₀ >30 μM	High H High M	PO: NSE PO + ANA: inhib in mice, no activity in rat
H0847	 (S enantiomer)	IC ₅₀ = 2 nM EC ₅₀ >30 μM	High H High M	PO 10 mg/kg + ANA 30 mg/kg mice: NSE







H0848	 (R enantiomer)	IC_{50} = 432 nM EC_{50} >30 μ M	Medium H High M	PO 10 mg/kg + ANA 30 mg/kg mice: NSE
H0863		IC_{50} = 3 nM EC_{50} >30 μ M	Medium H High M	Not done
H0908		IC_{50} = 8nM EC_{50} >30 μ M	Medium H High M	Not Done
H0864		IC_{50} = 718 nM EC_{50} >30 μ M	Not Done	Not Done
H0872		IC_{50} = 6 nM EC_{50} >30 μ M	High H medium M	Not Done
H0840		IC_{50} = 47 nM EC_{50} >30 μ M	Not Done	PO 10 mg/kg + ANA 30 mg/kg mice: NSE
H0910		IC_{50} = 125 nM EC_{50} = 19 nM E_{max} = 1359	Not Done	Not Done









H0788		IC ₅₀ = 88 nM EC ₅₀ = 20 nM E _{max} = 1230	Not Done	Not Done
H0789		IC ₅₀ = 284 nM EC ₅₀ = 26 nM E _{max} = 1137	Not Done	Not Done
H0760		IC ₅₀ = 6.2 μM EC ₅₀ > 30 μM	Not Done	Not Done
H0769		IC ₅₀ = 318 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0771		IC ₅₀ = 9 nM EC ₅₀ = 9 nM E _{max} = 4662	Not Done	Not Done
H0770		IC ₅₀ = 700 nM EC ₅₀ = 294 nM E _{max} = 1783	Not Done	Not Done
H0828		IC ₅₀ = 376 nM EC ₅₀ > 30 μM	Not Done	Not Done









H0822		$IC_{50} = 1.2 \mu M$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0850		$IC_{50} = 1.2 \mu M$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0881		$IC_{50} = 810 \text{ nM}$ $EC_{50} > 30 \mu M$	Not Done	Not Done
H0729		$IC_{50} = 100 \text{ nM}$ $EC_{50} = 95 \text{ nM}$ $E_{max} = 2818$	Not Done	Not Done
H0783		$IC_{50} = 681 \text{ nM}$ $EC_{50} = 30 \text{ nM}$	Not Done	Not Done
H0793		$IC_{50} = 21 \text{ nM}$ $EC_{50} = 22 \text{ nM}$ $E_{max} = 3501$	Not Done	Not Done
H0796		$IC_{50} = 826 \text{ nM}$ $EC_{50} = 3 \mu M$ $E_{max} = 1671$	Not Done	Not Done









H0498		IC ₅₀ = 29 nM EC ₅₀ >30 μM	Medium H High M	76% inhibition at 1h, activity up to 24h (30 mg/kg)
H0531		IC ₅₀ = 4 nM EC ₅₀ = 5 nM	Medium H Poor M	Not Done
H0594		IC ₅₀ = 54 nM EC ₅₀ >30 μM	Poor H Medium M	Not Done
H0644		IC ₅₀ = 6 nM EC ₅₀ = 28 nM E _{max} = 2822	Medium H Medium M	Not Done
H0536	 (racemic mixture)	IC ₅₀ = 3 nM EC ₅₀ >30 μM	Medium H Medium M	76% inhibition at 1h, activity up to 24 h (30 mg/kg)
H0563	 (single enantiomer)	IC ₅₀ = 1 nM EC ₅₀ = 3 nM E _{max} = 2100	Medium H Medium M	65% inhibition at 1 h (10 mg/kg)
H0564	 (single enantiomer)	IC ₅₀ = 75 nM EC ₅₀ = 124 nM E _{max} = 1987	Not Done	Not Done









H0627		IC ₅₀ = 4 nM EC ₅₀ = 1 nM E _{max} = 5289	High H High M	Not Done
H0660		IC ₅₀ = 69 nM EC ₅₀ = 180 nM E _{max} = 2100	Not Done	Not Done
H0661		IC ₅₀ = 2 nM EC ₅₀ = 6 nM E _{max} = 2280	Not Done	Not Done
H0672		IC ₅₀ > 1 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0651		IC ₅₀ = 4 nM EC ₅₀ = 11 nM E _{max} = 2300	Not Done	Not Done
H0653		IC ₅₀ = 4 nM EC ₅₀ = 9 nM E _{max} = 1815	Medium H Medium M	Not Done
H0668		IC ₅₀ = 8 nM EC ₅₀ = 10 nM E _{max} = 2168	Not Done	Not Done
H0654		IC ₅₀ = 6 nM EC ₅₀ = 10 nM E _{max} = 2200	High H Medium M	Not Done








H0655		IC ₅₀ = 12 nM EC ₅₀ >30 μM	Medium H Medium M	70% Inhibition at 1 h; activity up to 4 h (10 mg/kg)
H0691		IC ₅₀ = 5 nM EC ₅₀ >30 μM	High H High M	62% Inhibition at 2 h; activity up to 24 h (10 mg/kg) PO: not active
H0728		IC ₅₀ = 5 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0726		IC ₅₀ = 456 nM EC ₅₀ >30 μM	Not Done	Not Done
H0689		IC ₅₀ >1 μM EC ₅₀ >30 μM	Not Done	Not Done
H0692		IC ₅₀ = 550 nM EC ₅₀ >1 μM	Not Done	Not Done








H0656		IC ₅₀ = 7 nM EC ₅₀ = 15 nM E _{max} = 1350	Medium H Medium M	Not Done
H0652		IC ₅₀ = 7 nM EC ₅₀ = 5 nM E _{max} = 1500	Not Done	Not Done
H0713		IC ₅₀ = 187 nM EC ₅₀ = 29 nM E _{max} = 3424	Not Done	Not Done
H0688		IC ₅₀ = 3 nM EC ₅₀ = 12 nM E _{max} = 3100	Not Done	Not Done
H0774		IC ₅₀ = 3.4 μM EC ₅₀ >30 μM	Not Done	Not Done
H0664		IC ₅₀ = 261 nM EC ₅₀ >30 μM	Not Done	Not Done
H0535		IC ₅₀ = 34 nM EC ₅₀ = 4 nM	Not Done	Not Done
H0499		IC ₅₀ = 12 nM EC ₅₀ >30 μM	Medium H Medium M	NSE (30 mg/kg)








H0693		IC ₅₀ = 197 nM EC ₅₀ =100 nM	Not Done	Not Done
H0694		IC ₅₀ = 309 nM EC ₅₀ >30 μM	Not Done	Not Done
H0657		IC ₅₀ = 48 nM EC ₅₀ >30 μM	Medium H Poor M	57% inhibition at 1 h, activity up to 8h (10 mg/kg)
H0553		IC ₅₀ = 7 nM EC ₅₀ >30 μM	Medium H Poor M	57% inhibition at 1 h, activity up to 4h (10 mg/kg)
H0842		IC ₅₀ = 64 nM EC ₅₀ = 67 nM E _{max} = 1411	Not Done	Not Done
H0542		IC ₅₀ = 18 nM EC ₅₀ =15 nM	High H High M	Not Done
H0568		IC ₅₀ = 9 nM EC ₅₀ =4 nM	High H High M	Not Done
H0794		IC ₅₀ = 3 nM EC ₅₀ =10 nM E _{max} = 4435	Not Done	Not Done









H0841		IC ₅₀ = 118 nM EC ₅₀ >30 μM	Not Done	PO 10 mg/kg + ANAM 30 mg/kg mice: NSE
H0792		IC ₅₀ = 16 nM EC ₅₀ =7 nM E _{max} = 1096	Not Done	Not Done
H0569		IC ₅₀ = 87 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0565		IC ₅₀ =28 nM EC ₅₀ =30 nM	High H High M	Not Done
H0604		IC ₅₀ =12 nM EC ₅₀ =25 nM	High H High M	Not Done
H0595		IC ₅₀ =28 nM EC ₅₀ =43 nM	Not Done	Not Done
H0596		IC ₅₀ =9 nM EC ₅₀ =3 nM	High H High M	NSE (10 mg/kg)
H0851		IC ₅₀ =11 nM EC ₅₀ =6 nM E _{max} = 3320	Not Done	Not Done









H0537		IC ₅₀ = 13nM EC ₅₀ >30 μM	Poor H Poor M	Not Done
H0529		IC ₅₀ = 12 nM EC ₅₀ >30 μM	Medium H Poor M	Not Done
H0528		IC ₅₀ = 34 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0501		IC ₅₀ = 13 nM EC ₅₀ = 22 nM	High H High M	Not Done
H0507		IC ₅₀ = 8 nM EC ₅₀ =12 nM	High H High M	Not Done
H0665		IC ₅₀ = 4 nM EC ₅₀ =8 nM	High H Medium M	Not Done
H0508		IC ₅₀ = 76 nM EC ₅₀ >30 μM	Medium H High M	Not Done
H0509		IC ₅₀ = 29 nM EC ₅₀ =2 μM E _{max} = 1790	High H High M	66% inhib 1 h; activity up to 2h (10 mg/kg)








H0510		IC ₅₀ = 14 nM EC ₅₀ >30 μM	High H High M	35% inhibition at 4 h, activity up to 24h (30 mg/kg)
H0606		IC ₅₀ = 24 nM EC ₅₀ =31 nM E _{max} = 2336	Not Done	Not Done
H0810		IC ₅₀ = 20 nM EC ₅₀ =22nM E _{max} = 2339	Not Done	Not Done
H0696		IC ₅₀ = 120 nM EC ₅₀ >30 μM	Not Done	Not Done
H0611		IC ₅₀ = 2.3 μM EC ₅₀ > 30 μM	Not Done	Not Done
H0612		IC ₅₀ = 1.6 μM EC ₅₀ > 30 μM	Not Done	Not Done
H0615		IC ₅₀ = 107 nM EC ₅₀ > 30 μM	High H Medium M	Not Done










H0809		IC ₅₀ = 149 nM EC ₅₀ =217 nM E _{max} = 2339	Not Done	Not Done
H0699		IC ₅₀ = 171 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0607		IC ₅₀ = 6 nM EC ₅₀ =31 nM E _{max} = 3000	Medium H Medium M	NSE (10 mg/kg)
H0695		IC ₅₀ = 78 nM EC ₅₀ =5 nM	Not Done	Not Done
H0635		IC ₅₀ = 1 μM EC ₅₀ > 30 μM	Not Done	Not Done
H0690		IC ₅₀ = 980 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0735		IC ₅₀ = 209 nM EC ₅₀ > 30 μM	Not Done	Not Done







H0746		IC ₅₀ = 216 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0747		IC ₅₀ = 84 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0748		IC ₅₀ = 554 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0765		IC ₅₀ = 61nM EC ₅₀ =137 nM E _{max} = 2810	Not Done	Not Done
H0766		IC ₅₀ = 171nM EC ₅₀ > 30 μM	Not Done	Not Done
H0608		IC ₅₀ = 69 nM EC ₅₀ =422 nM	Not Done	Not Done
H0616		IC ₅₀ = 132 nM EC ₅₀ =580 nM	Not Done	Not Done









H0618		IC ₅₀ = 40 nM EC ₅₀ =130 μM	Not Done	Not Done
H0623		IC ₅₀ = 71 nM EC ₅₀ >30 μM	Medium H Poor M	Not Done
H0610		IC ₅₀ = 101 nM EC ₅₀ >30 μM	Medium H Medium M	Not Done
H0517		IC ₅₀ = 19 nM EC ₅₀ >30 μM	Not Done	Not Done
H0518		IC ₅₀ = 841 nM EC ₅₀ >30 μM	Not Done	Not Done
H0512		IC ₅₀ = 495 nM EC ₅₀ >30 μM	Not Done	Not Done
H0513		IC ₅₀ = 544 nM EC ₅₀ >30 μM	Not Done	Not Done
H0514		IC ₅₀ = 16 nM EC ₅₀ = 38nM	Medium H Medium M	Not Done








H0515		IC ₅₀ = 40 nM EC ₅₀ = 885 nM	Not Done	
H0520		IC ₅₀ = 202 nM EC ₅₀ = 394 nM	Not Done	Not Done
H0787		IC ₅₀ = 12 μM EC ₅₀ = 3 μM	Not Done	Not Done
H0582		IC ₅₀ = 15 nM EC ₅₀ = 20 nM E _{max} = 2069	Medium H Medium M	
H0571		IC ₅₀ = 154 nM EC ₅₀ >30 μM	Not Done	
H0605		IC ₅₀ = 31 nM EC ₅₀ = 96 nM E _{max} = 1833	Not Done	Not Done
H0573		IC ₅₀ = 36 nM EC ₅₀ >30 μM	High H Medium M	PO: NSE (30 mg/kg)
H0574		IC ₅₀ =67 nM EC ₅₀ =81 nM E _{max} = 2489	Medium H Medium M	Not Done






H0575		IC ₅₀ = 32 nM EC ₅₀ = 28 nM E _{max} = 3533	Medium H Medium M	Not Done
H0576		IC ₅₀ = 180 nM EC ₅₀ >30 μM	Not Done	Not Done
H0577		IC ₅₀ = 233 nM EC ₅₀ >30 μM	Not Done	Not Done
H0591		IC ₅₀ = 11 nM EC ₅₀ =126 nM	Medium H Medium M	Not Done
H0597		IC ₅₀ = 24 nM EC ₅₀ >30 μM	Poor H Poor M	Not Done
H0598		IC ₅₀ = 63 nM EC ₅₀ =271 nM	Not Done	Not Done
H0599		IC ₅₀ = 212 nM EC ₅₀ =478 nM	Not Done	Not Done









H0790		IC ₅₀ = 35 nM EC ₅₀ = 32 nM E _{max} = 2810	Not Done	Not Done
H0381		IC ₅₀ = 12 nM EC ₅₀ > 30 μM	Medium H Medium M	IP: No effect
H0519		IC ₅₀ = 3 nM EC ₅₀ = 6 nM	Medium H Medium M	
H0629		IC ₅₀ = 3 nM EC ₅₀ = 1 nM E _{max} = 5075	Not Done	Not Done
H0658		IC ₅₀ = 6 nM EC ₅₀ = 9 nM E _{max} = 2400	Not Done	Not Done
H0669		IC ₅₀ = 1 nM EC ₅₀ = 5 nM E _{max} = 4961	Not Done	Not Done
H0671		IC ₅₀ = 34 nM EC ₅₀ = 60 nM E _{max} = 3748	Not Done	Not Done
H0659		IC ₅₀ = 390 nM EC ₅₀ = 353 nM E _{max} = 200	Not Done	Not Done
H0521		IC ₅₀ = 20 nM EC ₅₀ = 19 nM	Not Done	Not Done









H0602		$IC_{50}=8\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	Not Done	Not Done
H0603		$IC_{50}=2\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	High H High M	71% inhibition at 1 h, activity up to 24 h (0.1mpk), 65% inhibition at 1 h; activity up to 24 hrs (1mpk), 34% inhibition at 1 h, activity up to 4h (10mpk); Inhibition in fed mice after ANAM PO-SC NSE
H0677	 R/S (single enantiomer)	$IC_{50}=5\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	High H Medium M	Inhib up to 2h (10 mg/kg) PO: NSE
H0678	 R/S (single enantiomer)	$IC_{50}=55\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	Medium H Medium M	78% inhibition at 1 h, activity up to 24h (10 mg/kg). PO: no effect
H0832		$IC_{50}=11\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$		
H0852		$IC_{50}=22\text{ nM}$ $EC_{50}=18\text{ nM}$ $E_{\text{max}}=1683$	Not Done	Not Done










H0701		IC ₅₀ =20 nM EC ₅₀ >30 μM	Low H Low M	Not Done
H0733		IC ₅₀ = 95 nM EC ₅₀ >30 μM	Not Done	Not Done
H0755		IC ₅₀ =12 nM EC ₅₀ =10 nM E _{max} = 2196	Not Done	Not Done
H0757		IC ₅₀ = 159 nM EC ₅₀ = 654nM E _{max} = 2704	Not Done	Not Done
H0734		IC ₅₀ = 202 nM EC ₅₀ >30 μM	Not Done	Not Done
H0737		IC ₅₀ = 13 nM EC ₅₀ >30 μM	High H High M	75% inhibition at 1 h, activity up to 4 h (10 mg/kg) PO: NSE
H0775		IC ₅₀ = 74 nM EC ₅₀ >5 μM	Not Done	Not Done
H0776		IC ₅₀ = 120 nM EC ₅₀ >4 μM	Not Done	Not Done






H0779		IC ₅₀ = 429 nM EC ₅₀ = 4 μM	Not Done	Not Done
H0762		IC ₅₀ = 5 nM EC ₅₀ >30 μM	High H High M	93% inhibition at 1 h, activity up to 4 h (10 mg/kg) PO mice and rat: NSEt
H0751		IC ₅₀ = 6 nM EC ₅₀ = 62 nM E _{max} = 1267	High H High M	91% inhibition at 1 h, activity up to 24 h (10 mg/kg) PO mice and rat (+ANA): no effect
H0763		IC ₅₀ = 835 nM EC ₅₀ >30 μM	Not Done	Not Done
H0759		IC ₅₀ = 7 nM EC ₅₀ >30 μM	High H High M	85% inhibition at 1 hr, activity up to 8 h (10 mg/kg) PO: no effect
H0785		IC ₅₀ = 33 nM EC ₅₀ = 90 nM E _{max} = 2869	Not Done	Not Done
H0754		IC ₅₀ = 11 nM EC ₅₀ >30 μM	High H High M	74% inhibition at 1 h; activity up to 24h (10 mg/kg) PO and PO+ANAM: no effect

H0753		IC ₅₀ = 60 nM EC ₅₀ >30 μM	Not Done	Not Done
H0609		IC ₅₀ =517 nM EC ₅₀ >30 μM	Not Done	Not Done
H0764		IC ₅₀ =10 nM EC ₅₀ = 14 nM E _{max} = 1352	High H High M	91% inhibition at 1 h, activity up to 24 h (10 mg/kg) PO: 70% inhibition at 2 h (30 mg/kg), activity up to 24 h PO+ANAM: inhib up to 24h SC: 53% inhibition at 1 h; (30 mg/kg)
H0818	 (S/R) (single enantiomer)	IC ₅₀ =1.7 nM EC ₅₀ = 3.5 nM E _{max} = 1915	Not Done	22% inhib at 4 h, 30 mg/kg PO fasted mice
H0819	 (S/R) (single enantiomer)	IC ₅₀ =65 nM EC ₅₀ =140 nM E _{max} = 1419	Not Done	30 mg/kg PO fasted mice NSE

H0838	 (diastereoisomer mixture)	$IC_{50}=4\text{ nM}$ $EC_{50}=21\text{ nM}$ $E_{max}=1340$	Not Done	205% increase at 2h, activity up to 8 h, 30 mg/kg PO mice
H0855		$IC_{50}=256\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	Not Done	Not Done
H0884		$IC_{50}=197\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	Not Done	Not Done
H0811		$IC_{50}=36\text{ nM}$ $EC_{50}=95\text{ nM}$ $E_{max}=1320$	Not Done	Not Done
H0812		$IC_{50}=1.2\text{ }\mu\text{M}$ $EC_{50}=1.5\text{ }\mu\text{M}$ $E_{max}=871$	Not Done	Not Done
H0740		$IC_{50}=7\text{ nM}$ $EC_{50}=1.5\text{ nM}$ $E_{max}=3620$	Not Done	Not Done
H0742		$IC_{50}=54\text{ nM}$ $EC_{50}>30\text{ }\mu\text{M}$	Not Done	Not Done
H0745		$IC_{50}=57\text{ nM}$ $EC_{50}=97\text{ nM}$ $E_{max}=2391$	Not Done	Not Done

H0749		IC ₅₀ = 111 nM EC ₅₀ = 397 nM E _{max} = 1554	Not Done	Not Done
H0744		IC ₅₀ = 33 nM EC ₅₀ = 45 nM E _{max} = 3536	Not Done	Not Done
H0626		IC ₅₀ = 4 nM EC ₅₀ = 15 nM E _{max} = 3835	Not Done	Not Done
H0767		IC ₅₀ = 37 nM EC ₅₀ > 30 μM	High H High M	88% inhibition at 1 h, activity up to 4 h (10 mg/kg) PO: NSE
H0772		IC ₅₀ = 3 nM EC ₅₀ = 7 nM E _{max} = 3569	Not Done	Not Done
H0773		IC ₅₀ = 608 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0784		IC ₅₀ = 529 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0777		IC ₅₀ = 715 nM EC ₅₀ = 600 nM E _{max} = 2288	Not Done	Not Done

H0846		IC ₅₀ = 170 nM EC ₅₀ = 130 nM E _{max} = 3815	Not Done	Not Done
H0875		IC ₅₀ = 91 nM EC ₅₀ = 50 nM E _{max} = 3751	Not Done	Not Done
H0628		IC ₅₀ = 59 nM EC ₅₀ = 101 nM E _{max} = 4433	Not Done	Not Done
H0630		IC ₅₀ = 3 nM EC ₅₀ = 9 nM E _{max} = 4714	High H High M	Not Done
H0633		IC ₅₀ = 3 nM EC ₅₀ = 15 nM	High H High M	Not Done
H0634		IC ₅₀ = 13 nM EC ₅₀ = 37 nM	Not Done	Not Done
H0640		IC ₅₀ = 103 nM EC ₅₀ > 30 μM	Not done	Not done
H0645		IC ₅₀ = 133 nM EC ₅₀ = 287 nM E _{max} = 2761	Not done	Not done
H0641		IC ₅₀ = 18 nM EC ₅₀ = 35 nM E _{max} = 1690	Not Done	Not Done

H0702		IC ₅₀ = 96 nM EC ₅₀ = 1.1 μM E _{max} = 1940	Not Done	Not Done
H0643		IC ₅₀ = 22 nM EC ₅₀ = 83 nM E _{max} = 2660	Not done	Not Done
H0522		IC ₅₀ = 201 nM EC ₅₀ = 200 nM	Not Done	Not Done
H0523		IC ₅₀ = 668 nM EC ₅₀ > 30 μM	Not Done	Not Done
H0876		IC ₅₀ = 130 nM EC ₅₀ > 30 μM	Not Done	Not Done

5 *NSE: No significant effect.

Example C

Effect of Ghrelin Antagonists of Formula I on Binge Eating in Non-Estrous Female Rats

10 In this Example, the therapeutic potential of compounds were tested for their ability to inhibit binge eating. The animal model used was developed to explore the combination of food restriction and stress. Results disclosed below show that female rats submitted to cycles of food restriction and exposure, the day of the test, to Highly Palatable Food (HPF) for 15 minutes without getting access to it, showed a pronounced and statistically significant increase in HPF intake. Considering the reliability and the robustness of this model, it was adopted to test the

15 inventive compounds. Topiramate, used as reference compound, confirmed its inhibitory effect in this procedure. Moreover, results show that, after acute administration, H0900, H0816, and H0847, reduced binge eating episodes showed in R + S group. H0860, at the considered doses, did not significantly reduce HPF intake in animals exposed to the same procedure.

20 Animals and Housing:

5 A total of N = 117, 52-day-old female Sprague-Dawley rats (175-200 g) were used.
Rats were acclimated in individual cages with metallic walls; the floor and the front wall made of
metallic grid. The dimensions of the cage floor being 30 cm×30 cm; the cage is 30 cm high. A
front door (30 cm×20 cm) made of metallic grid was positioned in the anterior wall of the cage to
gain access to the inside of the cage; the remaining part of the front wall was equipped with a
10 drinking burette.

Rats were kept in a room at constant temperature (20-22°C) and humidity (45-55%)
under a 12-h light/dark cycle (lights on at 08:00 am) with ad lib chow and water.

All procedures were conducted in adherence to the European Community Directive for Care and
Use of Laboratory Animals.

15

Diets:

Rats were offered food pellets, 4RF18, Mucedola, Settimo Milanese, Italy (2.6 kcal/g).
The Highly Palatable Food (HPF) was prepared by mixing:

- 20 a) Nutella Ferrero chocolate cream (5.33 kcal/g; 56%, 31% and 7%, respectively, from
carbohydrate, fat and protein): 52 %
b) grounded food pellets 4RF18, Mucedola, Settimo Milanese, Italy: 33 %
c) water: 15 %

Experimental Design:

25 The rats were weight-matched into one of two groups so there was no significant
difference in mean body weight between the groups:

Group 1: non-restricted and not exposed to stress (NR + NS): N = 9

Group 2: restricted and exposed to stress (R + S): N = 108

30 Once assigned to one of these groups, the rats remained in that group throughout the
study. The rats exposed to stress were acclimated in different rooms than the group not exposed
to stress.

Rats were exposed to 3 consecutive 8-day cycles followed by the final test on day 25:

- a) the control group (NR + NS) had chow ad libitum for 4 days, on days 5-6 it received chow +
HPF for 2 h; on days 7-8 it had chow ad libitum; on day 25 it was not exposed to stress;

- 5 b) the second group (R + S) had chow restricted to 66% of the normal intake for 4 days, was offered chow and HPF (2 h) on days 5-6 and only chow on days 7-8; on day 25 it was not exposed to stress.

The 8-day cycle was repeated three times, but in the third cycle the animals did not have access to HPF.

- 10 By the last day of re-feeding, the body weight and food intake of restricted rats were not statistically different from those of non-restricted rats, thus precluding the potentially confounding effect of hunger or energy deficit.

Body weights and food intake were recorded daily. Food intake is expressed as mean kilocalories per kilogram ingested $\pm$ SEM.

- 15 On the test day (day 25) the animals were divided in the following groups as shown in Table 2:

Table 2

No. of Animals	Procedure	Treatment
8	NR NS	Vehicle
9	R S	Vehicle
9	R S	H0816 3 mg/kg
9	R S	H0816 30 mg/kg
9	R S	H0860 3 mg/kg
9	R S	H0860 30 mg/kg
9	R S	H0847 3 mg/kg
9	R S	H0847 30 mg/kg
9	R S	H0900 3 mg/kg
9	R S	H0900 30 mg/kg
9	R S	Topiramate 60 mg/kg

- 20 It has been reported by Applicants (Micioni Di B et al. 2010) that in the estrous phase of the ovarian cycle, female rats do not exhibit BE in the adopted model; while in all the other three phase of the ovarian cycle they exhibit BE without significant differences in intensity.

5 Therefore, immediately after the test on day 25, vaginal smears were collected and analyzed under microscope to assess the ovarian phase, and data from rats in the estrous phase were not included in the statistical analysis. Vaginal smears were analyzed by an experienced experimenter blind to treatment conditions.

10 **The Stress Procedure:**

For 15 min, the container (China coffee cup) containing HPF is placed outside the cage; the container handle is hooked to the top wire wall of the cage in the hollow part where food pellets are usually offered. In these conditions, the animal is able to see the cup in which it received HPF on days 5, 6, 13, and 14 of the first two cycles, is able to see in part the HPF itself,
15 and is able to smell its odour. In this 15-min period, the rat engages in repeated movements of the forepaws, head, and trunk aimed at obtaining the HPF, but it is not able to reach it. Rats undergo the stressful procedure between 10.00 and 12.00 am. After 15 min, the cup is placed inside the cage of the rats in the stress group (R + S), so that the HPF became accessible to the rat.

20

Compound Preparation:

100 mg of each compound (H0816, H0860, H0847 and H0900) was accurately weighed and suspended in 13.33 ml of 0.5% carboxymethyl cellulose sodium salt (CMC, Sigma-Aldrich
25 Cat. C4888, lot 120M0216V) solution. The lower dose solution was prepared by dilution of 30 mg/ml suspension with 0.5% CMC solution. Suspensions were prepared freshly on test day. Vehicle was composed by a solution of 0.5% carboxymethyl cellulose sodium salt and was prepared by dissolving 1 g of CMC in 200 ml of distilled water. 180 mg of Topiramate was accurately weighed and suspended in 12 ml of 0.5% CMC solution. Compounds (vehicle and
30 active principles) were administered by gavage in a volume of 4 ml/kg of body weight one hour before access to HPF.

Data Analysis:

All data are expressed as the mean $\pm$ s.e.m. and each value reflects the mean number of
35 animals per group as described in the legends. For data evaluation, the analysis of variance

5 (ANOVA) was used followed by post-hoc (Bonferroni's) test when appropriate. Statistical significance was set at $P < 0.05$. The Software used for the Graphs was Origin 7.0. The software for the statistical analysis was SYSTAT 13.0

Binge Eating Model:

10 The ANOVA revealed a highly significant difference in 2-h HPF intake in the 2 groups of rats following vehicle administration [$F(1,12) = 18.9$; $P < 0.01$]. As shown in Figure 1, following vehicle administration HPF intake in the R + S group was markedly higher than that of the control (NR + NS) group. HPF intake of R + S rats was very pronounced in the first 15 min of access to HPF; these animals never engaged in competing behaviours, but continuously
15 remained over the cup containing HPF and focused their attention on the intake. Cumulative HPF intake in the R + S group was significantly higher than in controls up to 120 min after access to HPF.

Effect of Topiramate on Binge Eating:

20 The ANOVA revealed a significant difference in 2-h HPF intake in the R + S rats treated with Topiramate at the dose of 60 mg/kg [$F(1,11) = 16.2$; $P < 0.01$]. As shown in Figure 2, post-hoc comparisons revealed that the effect of Topiramate was statistically significant at all time points for the whole period in which BE was exhibited.

Effect of H0816 on Binge Eating:

25 The ANOVA revealed a significant difference in 2-h HPF intake in the R + S rats treated with H0816 at the doses of 3 and 30 mg/kg [$F(2,19) = 3.9$; $P < 0.05$]. As shown in Figure 3, post-hoc comparisons revealed that the effect of H0816 (30 mg/kg) was statistically significant ($P < 0.05$) at 15 min time point. H0816 treatment (both doses) did not affect animals' gross behaviour
30 during the 2-h test.

Effect of H0860 on Binge Eating:

As shown in Figure 4, H0860 at the doses of 3 and 30 mg/kg did not affect HPF intake in the R + S group [$F(2,19) = 0.6$; $P > 0.05$].

35

5 **Effect of H0847 on Binge Eating:**

The ANOVA revealed a significant difference in 2-h HPF intake in the R + S rats treated with H0847 at the doses of 3 and 30 mg/kg [$F(2,19) = 8.7$; $P < 0.01$]. As shown in Figure 5, post-hoc comparisons revealed that the effect of H0847 (3 mg/kg) was statistically significant at 15, 30 and 60 min after HPF access. At the dose of 30 mg/kg, H0847 significantly ($P < 0.01$) reduced HPF intake at all time points for the whole period in which BE was exhibited. Two animals treated with H0847 (3 mg/kg) and one animal treated with the dose of 30 mg/kg showed a mild sedation during the 2-h test.

15 **Effect of H0900 on Binge Eating:**

The ANOVA revealed a significant difference in 2-h HPF intake in the R + S rats treated with H0900 at the doses of 3 and 30 mg/kg [$F(2,18) = 12.2$; $P < 0.01$]. As shown in Figure 6, post-hoc comparisons revealed that the effect of H0900 (30 mg/kg) was statistically significant ($P < 0.01$) at all time points for the whole period in which BE was exhibited. H0900 treatment (both doses) did not affect animals' gross behaviour during the 2-h test.

Effect of Topiramate, H0816, H0860, H0847H0900 and Vehicle on 2-h Chow Food Intake During Binge Eating Test:

Statistical analysis indicated that acute administration of Topiramate [$F(1,11) = 0.9$; $P > 0.05$] or H0816 [$F(2,19) = 0.3$; $P > 0.05$] or H0900 [$F(2,18) = 2.2$; $P > 0.05$] did not modify 2-h chow intake. As shown in Figure 7 A, the acute administration of H0860 [$F(2,19) = 22.9$; $P < 0.01$] and H0847 [$F(2,19) = 3.9$; $P < 0.05$] significantly increased 2-h chow food intake.

Statistical analysis indicated that acute administration of Topiramate [$F(1,11) = 0.00$; $P > 0.05$] or H0816 [$F(2,19) = 1.2$; $P > 0.05$] or H0900 [$F(2,18) = 2.7$; $P > 0.05$] did not modify 24-h chow intake.

As shown in Figure 7, the acute administration of H0860 [$F(2,19) = 14.2$; $P < 0.01$] and H0847 [$F(2,19) = 24.3$; $P < 0.01$] significantly increased 24-h chow food intake.

Effect of H0816 on Binge Eating (Second test):

5 To confirm the effect of H0816 on BE, a second test was performed after ten days. Of 117 animals used in this study, 53 (the same 8 rats NR+NS and 45 rats R+S) were used for the second test. After one day off at the end of the first test, these groups of rats received an additional 8-day cycle: NR +NS group had 8 days of chow ad libitum, whereas R + S group had 4 days chow restricted to 66% of the normal intake followed by 4 days of chow ad libitum. In 10 this additional cycle, all groups did not have access to HPF. The following day, R + S group was exposed to stress, while NR +NS group was not. On this day, H0816 (3, 10 and 30 mg/kg) and topiramate (60 mg/kg) or vehicle were administered by gavage 1-h before access to HPF. The ANOVA revealed a highly significant difference in 2-h HPF intake in the 2 groups of rats following vehicle administration [$F(1,12) = 28.1$; $P < 0.01$]. Cumulative HPF intake in the R + S 15 group was significantly higher than in controls up to 120 min after access to it (data not shown). Statistical analysis showed a significant difference in 2-h HPF intake in the R + S rats treated with Topiramate at the dose of 60 mg/kg [$F(1,12) = 47.1$; $P < 0.01$]. Post-hoc comparisons revealed that the effect of Topiramate was statistically significant at all time points, that is for the whole period in which BE was exhibited (data not shown).

20 The ANOVA revealed a significant difference in 2-h HPF intake in the R + S rats treated with H0816 at the doses of 3, 10 and 30 mg/kg [$F(3,25) = 3.3$; $P < 0.05$]. As shown in Figure 8, post-hoc comparisons revealed that the effect of H0816 (10 mg/kg) was statistically significant ($P < 0.05$) at 15 min time point and the dose of 30 mg/kg completely blocked ($P < 0.01$) the BE episode at 15 min. H0816 treatment (both doses) did not affect animals' gross behaviour during 25 the 2-h test. Statistical analysis indicated that acute administration of Topiramate [$F(1,12) = 2.3$; $P > 0.05$] or H0816 [$F(3,25) = 0.2$; $P > 0.05$] did not modify 2-h and 24-h ($[F(1,12) = 0.03$; $P > 0.05]$; [$F(3,25) = 0.5$; $P > 0.05]$) chow intake (data not shown).

Topiramate, included in the experimental design as positive control, completely abolished BE episode at the dose of 60 mg/kg. In the same experiment, H0900, H0816, and H0847 30 significantly reduced BE behaviour in the R + S group, after acute administration, confirming the therapeutic potential of selective GHS-R1a antagonism in binge eaters.

In a second experiment, H0816 confirmed, dose dependently, its selective inhibitory effect on BE, with no effect on physiological feeding. Surprisingly, H0847 and H0860 significantly increased 2-h and 24-h chow food intake in the same animals, suggesting a not 35 clean profile as GHS-R1a antagonist.

5

Example D**Characterizing the Effect of Compounds of Formula (I) on Operant Ethanol Self-Administration in Marchigian Sardinian alcohol-preferring (msP) Rats**

In this experiment, msP- rats (N=24) were trained to self-administer 10% (v/v) ethanol solution in 30-min daily sessions under a fixed-ratio 1 schedule of reinforcement in which each response resulted in delivery of 0.1mL of fluids. Training continued until stable baseline of alcohol responding was achieved. At this point, before initiation of treatments, rats were trained to gavage administration procedures for three consecutive days (pre-treatment phase) during which they received drug vehicle. At this point animals were tested for the effect of ghrelin antagonists on 10% (v/v) ethanol self-administration. Using a within-subject Latin square design, the first group of msP rats (N=12) was tested for the effect of H0847 (0.0, 1.0 and 3.0 mg/kg), while the second (N=12) was treated with H0816 (0.0, 3.0 and 10.0 mg/kg).

Once the experiment was finished, animals were left in their home cages for several days, in order to wash out the drugs. Then, the same rats were employed to test the remaining ghrelin antagonists compounds H0900 (0.0, 3.0 and 30.0 mg/kg) and H0860 (0.0, 3.0 and 30.0 mg/kg). Once a stable self-administration baseline was reached, treatments begun according to the same experimental procedures described for the previous drugs tested.

All the drugs (or vehicles) were administered orally 1 hour before the beginning of the operant session. Responses at the lever activated the delivery mechanism but did not result in the delivery of alcohol.

Animals and Housing:

Male genetically selected alcohol-preferring Marchigian Sardinian (msP) rats were used (N=24). At the time of the experiments their body weight ranged between 350 and 400 g. They were housed 4 per cages in a room with a reverse 12:12 h light/dark cycle (lights off at 9:30 a.m.), temperature of 20-22°C and humidity of 45-55%. Rats were offered free access to tap water and food pellets (4RF18, Mucedola, Settimo Milanese, Italy). All the procedures were conducted in adherence with the European Community Council Directive for Care and Use of Laboratory Animals and the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

5

Compound Preparation:

75 mg of each H0900 and H0860 were accurately weighed and suspended in 10 ml of 0.5% carboxymethyl cellulose sodium salt solution (CMC, Sigma-Aldrich Cat. C4888, lot 120M0216V). The lower dose solution was prepared by dilution of 30 mg/ml suspension with 10 0.5% CMC solution.

37.5 mg of H0816 were accurately weighed and suspended in 15 ml of 0.5% carboxymethyl cellulose sodium salt solution (CMC, Sigma-Aldrich Cat. C4888, lot 120M0216V). The lower dose solution was prepared by dilution of mg/ml suspension with 0.5% CMC solution.

15 11.25 mg of H0847 were accurately weighed and suspended in 15 ml of 0.5% carboxymethyl cellulose sodium salt solution (CMC, Sigma-Aldrich Cat. C4888, lot 120M0216V). The lower dose solution was prepared by dilution of mg/ml suspension with 0.5% CMC solution.

Suspensions were prepared freshly on test day. Vehicle was composed of a solution of 20 0.5% carboxymethyl cellulose sodium salt and was prepared by dissolving 1g of CMC in 200 ml of distilled water. Vehicle and drugs were administered by gavage in a volume of 4 ml/kg of body weight 1 hour before the access to 10% alcohol solution. 10% (v/v) ethanol solution was prepared every two days by diluting 95% (v/v) ethanol solution (F.L. CARSETTI s.n.c - CAMERINO) in drinkable water.

25

Equipment:

The self-administration stations consisted of operant conditioning chambers (Med Associate, Inc) enclosed in sound-attenuating, ventilated environmental cubicles. Each chamber was equipped with a drinking reservoir (volume capacity: 0.2) positioned 4 cm above the grid 30 floor in the centre of the front panel of the chamber, two retractable levers located 3 cm (one to the right and the other to the left) of the drinking receptacle and a white cue light located 6 cm above the lever. An infusion pump was activated by responses on the right, or active lever, while responses on the left or inactive lever were recorded but did not result in activation of the pump. Activation of the pump resulted in a delivery of 0.1ml of fluid. If a time out was programmed, 35 lever presses during this period were counted but did not lead to further infusions. An IBM-

5 compatible computer controlled the delivery of fluids (activation of syringe pump), presentation of visual stimuli and recording of the behavioral data.

Experimental procedures:

10 Using operant self-administration chambers (Med Associates), msP rats were trained to lever press for 10% alcohol (v/v) until stable baseline of responding were achieved. 16 self-administration training sessions were carried out to train the animals. Operant sessions lasted 30 minutes and were conducted once a day during the dark phase of the light dark cycle. Active and inactive (control) lever responding were monitored.

15 After stable baseline of alcohol self-administration was established, msP rats were administered with vehicle or the inventive compounds at 2 different doses using a within subject design. Active and inactive lever responding was monitored: drugs were injected prior to the beginning of the self-administration session, according to indication.

20 The reinforcement program was FR1-LITO (Fixed Ratio - 1 Light Time Out). During the 5 seconds time out (following the reinforced RR) a house light was switched on. The tests were conducted according to a within subject design where drug treatment (doses) was treated as repeated factors. Total number of active and inactive lever responding were subjected to statistical evaluation. Drug testing was carried out every four days. For 2 days before each drug test rats were not subjected to alcohol self-administration sessions.

25 Statistical Analysis:

Data were analyzed by means of a one-factor (treatment) ANOVA for repeated measures. Analysis of variance was followed by the Newman–Keuls test when appropriate. Statistical significance was set at $p < 0.05$.

30 As shown in Figure 9, H0847 had no effect on operant responding for alcohol [$F(2,11) = 0.53$; $p > 0.05$]. Responses at the inactive control lever were not modified [$F(2,11) = 0.53$; $p > 0.05$].

35 As shown in Figure 10, H0860 significantly reduced operant responding for alcohol [$F(2,11) = 4.19$; $p < 0.05$]. Post hoc analysis revealed a significant reduction of alcohol self-administration following treatment with the higher dose (30 mg/kg) ($*p < 0.05$). Responses at the inactive control lever were not modified [$F(2,11) = 0.15$; $p > 0.05$].


5 As shown in Figure 11, H0816 had no effect on operant responding for alcohol [$F(2,11) = 0.75$; $p > 0.05$]. Responses at the inactive control lever were not modified [$F(2,11) = 0.30$; $p > 0.05$].

 As shown in Figure 12, H0900 significantly reduced operant responding for alcohol [$F(2,11) = 8.62$; $p < 0.01$]. Post hoc analysis revealed a significant reduction of alcohol self-
10 administration following treatment with both 3 mg/kg (* $p < 0.05$) and 30 mg/kg (** $p < 0.01$). Responses at the inactive control lever were not modified [$F(2,11) = 1.03$; $p > 0.05$].

 In summary, the data shows that, in msP rats, acute oral administration of both H0900 and H0860 induced a statistically significant decrease in ethanol self-administration. For H0900, the effect was seen for both the doses tested (3 and 30 mg/kg). For H0860, only the higher dose
15 (30 mg/kg) reduced ethanol self-administration. On the contrary, in the same experimental conditions, H0847 (1 or 3 mg/kg) and H0816 (3 or 10 mg/kg) had no effect on ethanol responses.

CLAIMS

1. A compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein:

X is CH or N;

Z is NR⁹;

R¹ is H, C₁₋₆ alkyl, benzyl, OH, or C₁₋₆ alkoxy, wherein said C₁₋₆ alkyl, benzyl, or C₁₋₆ alkoxy is optionally substituted with 1-3 substituents selected from halo, OH, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ hydroxyalkyl, CO(C₁₋₆ alkyl), CHO, CO₂H, CO₂(C₁₋₆ alkyl), and C₁₋₆ haloalkyl;

R² is H or C₁₋₆ alkyl;

R³ and R⁴ are each independently H, CN, halo, CHO, CO₂H, C₁₋₆ alkylcycloalkyl, C₁₋₆ alkoxy, CO(C₁₋₆ alkyl), CO₂(C₁₋₆ alkyl), CONR¹²R¹³, C₁₋₆ alkyl optionally substituted with halo, or C₁₋₆ hydroxyalkyl optionally substituted with aryl, wherein aryl is optionally substituted with C₁₋₃ hydroxyalkyl;

or R³ and R⁴ taken together with the carbon atom to which they are attached form a 3-6-membered ring;

R⁵ is pyridinyl or pyrimidinyl, each optionally substituted with 1-3 substituents selected from halo, CN, OH, NO₂, Si(CH₃)₃, CHO, CO₂H, CO(C₁₋₆ alkyl), CO₂(C₁₋₆ alkyl), NR¹⁴R¹⁵, NHCONR¹⁴R¹⁵, CONR¹⁴R¹⁵, CH=NOH, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ hydroxyalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, and heterocycloalkyl;

R⁶ is H;

R⁷ is H, CN, or halo;

R⁸ is H or C₁₋₆ alkyl;

R⁹ is H, C₁₋₆ alkyl, CO(C₁₋₆ alkyl), CHO, CO₂H, or CO₂(C₁₋₆ alkyl);

R^{12} and R^{13} are each independently H or C_{1-6} alkyl;

R^{14} and R^{15} are each independently H, C_{1-6} alkyl, $CO(C_{1-6}$ alkyl), $CO(heteroaryl)$, heteroaryl, or cycloalkyl;


r is 2;

s is 0, 1, 2, 3, or 4; and

n is 0, 1, 2, or 3.

2. The compound of claim 1, wherein X is CH.
3. The compound of claim 1, wherein X is N.
4. The compound of claim 1, wherein Z is $N(C_{1-6}$ alkyl).
5. The compound of claim 1, wherein Z is NCH_3 .
6. The compound of claim 1, wherein R^1 is C_{1-6} alkyl.
7. The compound of claim 1, wherein R^1 is CH_3 .
8. The compound of claim 1, wherein R^1 is benzyl.
9. The compound of claim 8, wherein said benzyl is optionally substituted with $CO_2(C_{1-6}$ alkyl) or C_{1-6} hydroxyalkyl.
10. The compound of claim 1, wherein R^1 is OH.
11. The compound of claim 1, wherein R^1 is C_{1-6} alkoxy.
12. The compound of claim 11 wherein said C_{1-6} alkoxy is OCH_3 , OCH_2CH_3 or $O(CH_2)_2CH_3$.
13. The compound of claim 1, wherein R^2 is H.
14. The compound of claim 1, wherein R^3 and R^4 are each independently CN, C_{1-6} alkylcycloalkyl, $CO_2(C_{1-6}$ alkyl), $CONH_2$, C_{1-6} alkyl optionally substituted with halo or C_{1-6} hydroxyalkyl optionally substituted with aryl.
15. The compound of claim 14, wherein said C_{1-6} alkyl is CH_3 or CH_2CH_3 .
16. The compound of claim 14, wherein said C_{1-6} alkylcycloalkyl is CH_2 -cyclopropyl.
17. The compound of claim 14, wherein said C_{1-6} hydroxyalkyl is CH_2OH .
18. The compound of claim 14, wherein said $CO_2(C_{1-6}$ alkyl) is CO_2CH_3 .
19. The compound of claim 14, wherein said C_{1-6} alkyl optionally substituted with halo is CF_3 .
20. The compound of claim 1, wherein R^3 and R^4 taken together with the carbon atom to which they are attached form a 3-6-membered ring.


21. The compound of claim 20, wherein R³ and R⁴ are taken together with the carbon atom to which they are attached to form a cyclopropyl ring.
22. The compound of claim 20, wherein R³ and R⁴ are taken together with the carbon atom to which they are attached form a tetrahydropyranyl ring.
23. The compound of claim 1, wherein R⁵ is pyridinyl.
24. The compound of claim 1, wherein R⁷ is halo.
25. The compound of claim 24, wherein said halo is Cl or F.
26. The compound of claim 1, wherein R⁸ is H.
27. The compound of claim 1, wherein R⁸ is C₁₋₆ alkyl.
28. The compound of claim 1, wherein R⁸ is CH₃.
29. The compound of claim 1, having Formula II:



II,

or a pharmaceutically acceptable salt thereof.


30. The compound of claim 1, having Formula III:



III,

or a pharmaceutically acceptable salt thereof.

31. The compound of claim 1, having Formula:



wherein:








R^{17} is H or halo.








32. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 and one or more pharmaceutically acceptable excipients.

33. A compound selected from the group consisting of:








Compound No.	Chemical Structure	Chemical Name
H0496		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0617		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0539		methyl 4-((3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)ureido)methyl)benzoate








H0546		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(4-(hydroxymethyl)benzyl)-1-(1-methylpiperidin-4-yl)urea
H0527		3-(1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-(1,3-dimethylpiperidin-4-yl)-1-methylurea
H0497		3-(1-(2,3-dichloro-4-(pyridin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0650		3-(1-(2,3-dichloro-4-(pyridin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0849		3-(1-(2,3-dichloro-4-(5-cyclopropylpyridin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0511		3-(1-(2,3-dichloro-4-(6-methoxypyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0820		3-(1-(2,3-dichloro-4-(6-cyclopropylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0613		3-(1-(2,3-dichloro-4-(5-cyanopyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0614		3-(1-(2,3-dichloro-4-(5-fluoropyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0635		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)nicotinate
H0636		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0637		3-(1-(2,3-dichloro-4-(5-(difluoromethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0638		3-(1-(2,3-dichloro-4-(5-(fluoromethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0639		3-(1-(2,3-dichloro-4-(5-methylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0642		3-(1-(2,3-dichloro-4-(5-formylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0704		3-(1-(4-(5-aminopyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0705		3-(1-(2,3-dichloro-4-(5-(cyclopent-1-en-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0707		3-(1-(4-(5-(1H-pyrazol-4-yl)pyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0711		3-(1-(4-(5-(1H-imidazol-4-yl)pyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0716		3-(1-(2,3-dichloro-4-(5-(thiazol-5-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0717		3-(1-(2,3-dichloro-4-(5-(thiophen-2-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0718		3-(1-(2,3-dichloro-4-(5-cyclopentylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0719		3-(1-(2,3-dichloro-4-(5-(pyrrolidin-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0712		N-(5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyridin-3-yl)acetamide
H0708		3-(1-(2,3-dichloro-4-(5-(methoxymethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0714		3-(1-(2,3-dichloro-4-(5-(2-methoxyethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0715		3-(1-(2,3-dichloro-4-(5-ethylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0706		3-(1-(2,3-dichloro-4-(5-vinylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0710		3-(1-(2,3-dichloro-4-(5-ethynylpyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0666		3-(cyano(2,3-dichloro-4-(5-cyanopyridin-3-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0739		3-((4-(5-(1H-pyrrol-2-yl)pyridin-3-yl)-2,3-dichlorophenyl)(cyano)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0667		3-(1-(2,3-dichloro-4-(5-cyanopyridin-3-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0821		3-(1-(2,3-dichloro-4-(5-cyano-6-(4-methylpiperazin-1-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0646		(E)-3-(1-(2,3-dichloro-4-(5-((hydroxyimino)methyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0720		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0721		3-(1-(4-(5-aminopyridin-3-yl)-2,3-dichlorophenyl)-2-cyclopropylethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0516		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0649		3-(1-(2,3-dichloro-4-(2-methoxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0797		3-(1-(2,3-dichloro-4-(2-hydroxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0798		3-(1-(4-(2-aminopyrimidin-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0799		3-(1-(2,3-dichloro-4-(2-(4-methylpiperazin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0800		3-(1-(2,3-dichloro-4-(2-fluoropyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0801		3-(1-(2,3-dichloro-4-(2-chloropyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0802		3-(1-(2,3-dichloro-4-(2-cyanopyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0803		3-(1-(4-(2-(1H-imidazol-1-yl)pyrimidin-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0804		3-(1-(2,3-dichloro-4-(2-(dimethylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0805		3-(1-(2,3-dichloro-4-(2-(cyclopropylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0806		3-(1-(2,3-dichloro-4-(2-(methylamino)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0807		N-(5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrimidin-2-yl)cyclopropanecarboxamide

H0854		3-(1-(2,3-dichloro-4-(2-cyclopropylpyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0813		3-(1-(2,3-dichloro-4-(2-(pyrrolidin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0814		3-(1-(2,3-dichloro-4-(2-(4-ethyl-3-oxopiperazin-1-yl)pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0703		3-(1-cyano-1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0709		3-(cyano(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)methyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0588		1-(1-acetylpiperidin-4-yl)-3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-methylurea
H0663		3-(1-(2,3-dichloro-4-(2,4-dimethoxypyrimidin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0620		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-((3-(hydroxymethyl)benzyl)oxy)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0624		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0662		methyl 2-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0670		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0673		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)cyclopropyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0727		3-(4-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)tetrahydro-2H-pyran-4-yl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0631		3-(cyano(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0686		3-(1-(2,3-dichloro-4-(pyrimidin-5-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0619		3-(1-(2,3-dichloro-4-(pyrimidin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0768		3-(1-(2,3-dichloro-4-(pyrimidin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0808		3-(1-(2,3-dichloro-4-(6-methylpyrimidin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0700		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0816		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0817		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0722		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0741		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0752		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea
H0743		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methoxy-1-(1-methylpiperidin-4-yl)urea
H0750		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethoxy-1-(1-methylpiperidin-4-yl)urea
H0756		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0761		3-(2-cyclopropyl-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea








H0781		3-(2-cyclopropyl-1-((2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0782		3-(2-cyclopropyl-1-((2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0824		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0890		3-((S)-1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0858		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0865		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0825		1-benzyl-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)urea








H0826		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0889		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0896		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0827		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)-1-propylurea
H0829		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0859		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0860		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)







H0922		methyl 2-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0924		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0830		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0899		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0900		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0909		3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-ethyl-1-(1-methylpiperidin-4-yl)urea
H0856		3-(1-(3-chloro-2-fluoro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0815		3-(1-(2,3-dichloro-4-(6-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0831		3-(1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0843		3-(1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-((R)-1,3,3-trimethylpiperidin-4-yl)urea
H0844		3-(2-cyclopropyl-1-(2,3-dichloro-4-(3-methylpyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0738		3-(1-(2,3-dichloro-4-(6-methoxypyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0780		3-(1-(4-(6-aminopyrazin-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0786		3-(1-(2,3-dichloro-4-(6-(chloromethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea







H0791		3-(1-(2,3-dichloro-4-(6-chloropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0795		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0847		(S)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0848		(R)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0863		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0908		3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0840		3-(1-(2,3-dichloro-4-(3-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0910		3-(1-(2,3-dichloro-4-(6-(trifluoromethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0788		3-(1-(2,3-dichloro-4-(6-cyanopyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0789		methyl 6-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxylate
H0760		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxamide
H0769		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)pyrazine-2-carboxylate
H0771		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N,N-dimethylpyrazine-2-carboxamide
H0770		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0828		3-(1-(2,3-dichloro-4-(quinoxalin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0822		3-(1-(2,3-dichloro-4-(5-(4-methylpiperazin-1-yl)pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0881		3-(1-(4,5-dichloro-6-(pyrazin-2-yl)pyridin-3-yl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0729		3-(1-(2,3-dichloro-4-(pyridazin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0783		3-(1-(2,3-dichloro-4-(pyridazin-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0793		3-(1-(2,3-dichloro-4-(1,2,4-triazin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0796		3-(1-(2,3-dichloro-4-(4,6-dimorpholino-1,3,5-triazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0498		3-(1-(2,3-dichloro-4-(thiophen-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0594		3-(1-(2,3-dichloro-4-(thiophen-3-yl)phenyl)-2-((3-(hydroxymethyl)benzyl)oxy)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0644		3-(cyano(2,3-dichloro-4-(thiophen-3-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0536		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0563		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)
H0564		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (single enantiomer)

H0627		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)ethyl)-1-hydroxy-1-(1-methylpiperidin-4-yl)urea
H0660		methyl 2-(2,3-dichloro-4-(thiophen-2-yl)phenyl)-2-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)acetate
H0661		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)-2-hydroxyethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0672		3-(1-(2,3-dichloro-4-(thiophen-2-yl)phenyl)cyclopropyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0651		3-(1-(2,3-dichloro-4-(5-formylthiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0653		3-(1-(2,3-dichloro-4-(5-(hydroxymethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0668		3-(1-(2,3-dichloro-4-(5-(fluoromethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0654		3-(1-(2,3-dichloro-4-(5-(difluoromethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0655		3-(1-(4-(5-acetylthiophen-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0691		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxamide
H0728		5-(2,3-dichloro-4-(2-cyclopropyl-1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxamide
H0726		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N,N-dimethylthiophene-2-carboxamide
H0689		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-2-carboxylic acid

H0692		5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)-N-methoxy-N-methylthiophene-2-carboxamide
H0656		3-(1-(2,3-dichloro-4-(5-(1-hydroxyethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0652		3-(1-(2,3-dichloro-4-(5-formylthiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0713		3-(1-(2,3-dichloro-4-(5-cyanothiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0688		3-(1-(4-(3-acetylthiophen-2-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0774		2-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)thiophene-3-carboxamide
H0664		3-(1-(2,3-dichloro-4-(3-(hydroxymethyl)thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea








H0535		3-(1-(2,3-dichloro-4-(1H-pyrrol-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0499		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0693		3-(1-(2,3-dichloro-4-(1-(2-hydroxyethyl)-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0694		3-(1-(2,3-dichloro-4-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0657		3-(cyano(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)methyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0553		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(4-(hydroxymethyl)benzyl)-1-(1-methylpiperidin-4-yl)urea
H0842		3-(1-(2,3-dichloro-4-(1-cyclopropyl-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea


H0542		3-(1-(2,3-dichloro-4-(1H-imidazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0568		3-(1-(2,3-dichloro-4-(thiazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0794		3-(1-(4-(2-aminothiazol-4-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0841		3-(1-(2,3-dichloro-4-(2-cyclopropylthiazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0792		3-(1-(4-(2-aminothiazol-5-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0569		3-(1-(2,3-dichloro-4-(oxazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0565		3-(1-(2,3-dichloro-4-(1H-1,2,3-triazol-1-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0604		3-(1-(2,3-dichloro-4-(1H-1,2,3-triazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0595		3-(1-(2,3-dichloro-4-(1,3,4-oxadiazol-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0596		3-(1-(2,3-dichloro-4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0851		3-(1-(2,3-dichloro-4-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0537		methyl 4-((3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(1-methylpiperidin-4-yl)ureido)methyl)benzoate
H0528		3-(1-(2,3-dichloro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-(1,3-dimethylpiperidin-4-yl)-1-methylurea
H0501		3-(1-(2,3-dichloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0507		3-(1-(2,3-dichloro-4-(furan-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0665		3-(1-(2,3-dichloro-4-(5-methylfuran-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0699		3-(1-(2,3-dichloro-4-(5-(cyanomethyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0607		3-(1-(2,3-dichloro-4-(5-methoxypyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0695		3-(1-(4-(5-bromopyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0635		methyl 5-(2,3-dichloro-4-(1-(3-methyl-3-(1-methylpiperidin-4-yl)ureido)ethyl)phenyl)nicotinate
H0690		3-(1-(4-(5-acetylpyridin-3-yl)-2,3-dichlorophenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0735		3-(1-(2,3-dichloro-4-(5-(pyrimidin-2-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea





H0746		3-(1-(2,3-dichloro-4-(5-(furan-3-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0747		3-(1-(2,3-dichloro-4-(5-(thiophen-3-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0748		3-(1-(2,3-dichloro-4-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0765		3-(1-(2,3-dichloro-4-(5-(cyclopropyl)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0766		3-(1-(2,3-dichloro-4-(5-nitropyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0608		3-(1-(2,3-dichloro-4-(6-isopropoxy)pyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0616		3-(1-(2,3-dichloro-4-(6-cyanopyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0618		3-(1-(2,3-dichloro-4-(6-fluoropyridin-3-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0623		3-(1-(2,3-dichloro-4-(1H-pyrazolo[3,4-b]pyridin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0512		3-(1-(2,3-dichloro-4-(1-methyl-1H-indazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0513		3-(1-(2,3-dichloro-4-(1H-indazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0514		3-(1-(2,3-dichloro-4-(1H-pyrrolo[2,3-b]pyridin-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0515		3-(1-(2,3-dichloro-4-(1H-indol-5-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0582		3-(1-(3-chloro-2-fluoro-4-(thiophen-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

H0571		3-(1-(3-chloro-2-fluoro-4-(1H-pyrazol-4-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
-------	---	--

or a pharmaceutically acceptable salt thereof.

34. A compound selected from the group consisting of:

H0816		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0900		(R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0847		(S)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea
H0860		(S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea

or a pharmaceutically acceptable salt thereof.

35. The compound of claim 34, which is (S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (H0816) or a pharmaceutically acceptable salt thereof.

36. The compound of claim 34, which is (R)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)-2,2,2-trifluoroethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (H0900) or a pharmaceutically acceptable salt thereof.

37. The compound of claim 34, which is (S)-3-(1-(2,3-dichloro-4-(6-fluoropyrazin-2-yl)phenyl)ethyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (H0847) or a pharmaceutically acceptable salt thereof.

38. The compound of claim 34, which is (S)-3-(1-(2,3-dichloro-4-(pyrazin-2-yl)phenyl)propyl)-1-methyl-1-(1-methylpiperidin-4-yl)urea (H0860) or a pharmaceutically acceptable salt thereof.