SUMMARY

[0005] Although many of these methods are able to reduce or combat the errors introduced by the sample preparation and sequencing processes for all molecules that are converted and sequenced, these methods are not able to infer the counts of molecules that were converted but not sequenced. Since this count of converted by unsequenced molecules can be highly variable from genomic region to region, these counts can dramatically and adversely affect the sensitivity that can be achieved.

[0006] To address this issue, input double-stranded deoxyribonucleic acid (DNA) can be converted by a process that tags both halves of the individual double-stranded molecule, in some cases differently. This can be performed using a variety of techniques, including ligation of hairpin, bubble, or forked adapters or other adaptors having double-stranded and single stranded segments (the unhybridized portion of a bubble, forked or hairpin adapter are deemed single-stranded herein). If tagged correctly, each original Watson and Crick (i.e., strand) side of the input double-stranded DNA molecule can be differently tagged and identified by the sequencer and subsequent bioinformatics. For all molecules in a particular region, counts of molecules where both Watson and Crick sides were recovered ("Pairs") versus those where only one half was recovered ("Singlets") can be recorded. The number of unseen molecules can be estimated based on the number of Pairs and Singlets detected.

[0007] An aspect of the present disclosure provides a method for detecting and/or quantifying rare deoxyribonucleic acid (DNA) in a heterogeneous population of original DNA fragments, comprising tagging the original DNA fragments in a single reaction using a library of a plurality of different tags such that greater than 30% of the fragments are tagged at both ends, wherein each of the tags comprises a molecular barcode. The single reaction can be in a single reaction vessel. Greater than 50% of the fragments can be tagged at both ends. The plurality of different tags can be no more than any of 100, 500, 1000, 10,000 or 100,000 different tags.

[0007a] Another aspect provides a method, comprising (a) providing a sample comprising a set of double-stranded polynucleotide molecules, each double-stranded polynucleotide molecule including first and second complementary strands; (b) tagging said double-stranded polynucleotide molecules with a set of duplex tags, wherein each duplex tag differently tags said first and second complementary strands of a double-stranded polynucleotide molecule in said set; (c) sequencing at least some of said tagged strands to produce a set of sequence reads; (d) reducing and/or tracking redundancy in said set of sequence reads; (e) sorting

sequence reads into paired reads and unpaired reads, wherein (i) each paired read corresponds to sequence reads generated from a first tagged strand and a second differently tagged complementary strand derived from a double-stranded polynucleotide molecule in said set, and (ii) each unpaired read represents a first tagged strand having no second differently tagged complementary strand derived from a double-stranded polynucleotide molecule represented among said sequence reads in said set of sequence reads; (f) determining quantitative measures of (i) said paired reads and (ii) said unpaired reads that map to each of one or more genetic loci; and (g) estimating with a programmed computer processor a quantitative measure of total double-stranded polynucleotide molecules in said set that map to each of said one or more genetic loci based on said quantitative measure of paired reads and unpaired reads mapping to each locus.

[0008] Another aspect provides a set of library adaptors that can be used to tag the molecules of interest (e.g., by ligation, hybridization, etc.). The set of library adaptors can comprise plurality of polynucleotide molecules with molecular barcodes, wherein the plurality of polynucleotide molecules are less than or equal to 80 nucleotide bases in length, wherein the molecular barcodes are at least 4 nucleotide bases in length, and wherein (a) the molecular barcodes are different from one another and have an edit distance of at least 1 between one another; (b) the molecular barcodes are located at least one nucleotide base