

DESCRIPTION

TEST DEVICE, TEST KIT, AND TEST METHOD

Technical Field

[0001]

5 The present invention relates to a test device, a test kit, and a test method capable of antibody test.

Background Art

[0002]

10 A conventionally known test device for performing an immunoassay is a test device employing immunochromatography.

[0003]

15 Immunochromatography is an immunoassay that employs a property (capillary action) in which a specimen slowly flows on a porous test piece such as a cellulose membrane while dissolving a reagent therein. By using this property in an immunochromatographic test device, antigens (or antibodies) in the specimen migrate on a cellulose membrane while forming immune complexes with antibodies (or antigens) (labeled antibodies (or antigens)) labeled with metal colloid or the like previously prepared at a dropping portion on which the specimen is dropped. Subsequently, the immune complexes are trapped on capture antibodies (or antigens) previously prepared on the cellulose membrane, and colored, and finally the colored state is visually observed for determination.

25 Immunochromatographic test devices have been adopted for

diagnosis of pregnancy, influenza, and the like.

[0004]

A known example of an immunochromatographic test device is a test device or the like in which a test piece is enclosed in a rectangular case. The test piece contains previously immobilized labeled antibodies and capture antibodies, and the case includes a dropping window on which a specimen is dropped and a detection window through which the state of the test piece is visually observed (for example, see Patent Literature 1).

[0005]

Currently, a method for detecting IgG antibodies or IgM antibodies that are types of immunoglobulin in blood and that specifically react with a nucleocapsid protein (hereinafter referred to as "N antigen") and a spike protein (hereinafter referred to as "S antigen") of virus (SARS-CoV-2) causing novel coronavirus infection (COVID-19) is established.

[0006]

As a simple and quick test device for determining the presence or absence of a history of novel coronavirus infection, a kit capable of measuring IgG antibodies and/or IgM antibodies in human whole blood, serum, or plasma by immunochromatography is also increasingly used.

[0007]

For the antibody test devices, the labeled antigen in the

test piece is the S antigen alone, the N antigen alone, or the S antigen and the N antigen together or the like. To trap an immune complex formed by binding the target antibody (for example, IgG antibody) in the specimen to the labeled antigen, the test piece includes a line (capture antibody line) to which the antibody (capture antibody, or anti-IgG antibody) is applied.

[0008]

The S antigen and the N antigen are proteins of the SARS-CoV-2 virus (hereinafter referred to as novel coronavirus). Therefore, when it is possible to detect (trap) an immune complex formed by binding the antibody in the specimen to the S antigen or the N antigen on the capture antibody line, the presence or absence of a history of infection with the virus can be determined.

Citation List

Patent Literature

[0009]

Patent Literature 1: Japanese Patent No. 6217141

Summary of Invention

Technical Problem

[0010]

However, the conventional test device by immunochromatography (hereinafter referred to as "immunochromatographic test device") has difficulty in

performing fractionated determination.

[0011]

Specifically, the conventional immunochromatographic test device has a problem in which it cannot determine whether an antibody (neutralizing antibody, for example, IgG antibody) having a neutralization effect of reducing the infectivity of a predetermined virus (for example, novel coronavirus) is produced by natural infection or by vaccination.

[0012]

It is known that an antibody that neutralizes the novel coronavirus targets a spike protein (S antigen) necessary for entrance into human cells and prevents infection by the virus. Thus, research and development of vaccine capable of producing an antibody (IgG antibody, and the like) to target (capable of binding to) the S antigen have been advanced.

[0013]

When the use of a vaccine expands, it is desirable to determine the efficacy of the vaccination quickly and simply. Here, the conventional immunochromatographic test device can also detect the presence of IgG antibody (functioning as a neutralizing antibody) capable of binding to the S antigen which can be used as the labeled antigen in the test piece.

[0014]

However, the IgG antibody having such a neutralization effect is produced not only by vaccination but also by viral

infection (natural infection). Specifically, the immunochromatographic test device has a problem in which the test piece containing the S antigen used as the labeled antigen can detect the neutralizing antibody (for example, IgG antibody) in the specimen, but cannot determine whether the antibody is produced by vaccination or natural infection.

[0015]

In a case of using both the S antigen and the N antigen as the labeled antigen in the test piece of the conventional immunochromatographic examination device, the same problem occurs. In other words, an immune complex (one type of antibody) that binds to the S antigen or the N antigen is trapped on the capture antibody line as described above. In this case, when the capture antibody line is colored, the immunochromatographic test device can only determine whether the specimen contains at least the antibody (IgG antibody functioning as the neutralizing antibody) to trap the S antigen or the antibody (for example, IgG antibody) to trap the N antigen. Thus, the immunochromatographic test device in this case also has the problem in which whether the neutralizing antibody is produced by vaccination cannot be determined.

[0016]

In such a situation, a test device and a test method for quickly and simply determining the efficacy of vaccination

against a predetermined virus, for example, a novel coronavirus, or determining the history of vaccination inoculation (the presence or absence of the neutralizing antibody produced by natural infection) are not necessarily provided at a practical level, and practical application of the test device and the test method is desired.

[0017]

In view of the aforementioned circumstances, it is an object of the present invention to provide a test device and a test method by immunochromatography that can quickly and simply determine the efficacy of vaccination against a predetermined virus, for example, a novel coronavirus, or determine the history of vaccination inoculation.

Solution to Problem

[0018]

The present invention involves a test device by immunochromatography including: a first test piece; a second test piece separate from the first test piece; and a case configured to integrally accommodate the first test piece and the second test piece. The case includes a dropping window on which a specimen is dropped, a first detection window through which a part of the first test piece can be visually observed from the outside, and a second detection window through which a part of the second test piece can be visually observed from the outside. The first test piece can detect the presence of a

first antibody in the specimen, and the second test piece can detect the presence of a second antibody different from the first antibody in the specimen. A result from the first test piece visually observed through the first detection window and
5 a result from the second test piece visually observed through the second detection window can be determined in parallel and at the same time.

[0019]

The present invention also involves a test kit including
10 the above-described test device, and a determination guide that allows for determining a state of a specimen including a possibility that an antibody is produced by vaccination on the basis of a test result from the first test piece and a test result from the second test piece.

15 [0020]

The present invention also involves a test method using the above-described test device, the method including: a first test step of dropping the specimen on the first test piece to detect the presence or absence of the first antibody in the
20 specimen; a second test step of dropping the specimen on the second test piece different from the first test piece to detect the presence or absence of the second antibody in the specimen; and a determination step of combining results from the first and second test steps to determine the state of the
25 specimen.

[0021]

The present invention also involves a test method for determining the presence or absence of a first antibody and a second antibody in a specimen by immunochromatography, where
5 the first antibody is an antibody produced by vaccination or infection with a predetermined virus, and the second antibody is an antibody produced by infection with the virus. The method includes: dropping the specimen on a first test piece to determine the presence or absence of the first antibody in
10 the specimen; dropping the specimen on a second test piece different from the first test piece to determine the presence or absence of the second antibody in the specimen; and combining a test result from the first test piece and a test result from the second test piece to determine whether the
15 first antibody is produced by the vaccination in the specimen.

Advantageous Effects of Invention

[0022]

According to the present invention, an object is to provide a test device, a test kit, and a test method by
20 immunochromatography that can quickly and simply determining the efficacy of vaccine against a predetermined virus, for example, a novel coronavirus. There can be provided.

Brief Description of Drawings

[0023]

25 FIG. 1(A) is a plan view of a test device according to an

embodiment of the present invention, FIG. 1(B) is a plan view of a test piece, and FIG. 1(C) is a cross-sectional side view of the test piece.

FIG. 2 is a schematic view illustrating the principles of test by immunochromatography.

FIG. 3 is a flowchart illustrating an example of a test method according to the present embodiment.

FIG. 4 is a schematic view illustrating an example of the test method according to the present embodiment.

FIG. 5 is a schematic view illustrating an example of a test kit according to the present embodiment.

FIG. 6 is a schematic view illustrating an example of the test kit according to the present embodiment.

Description of Embodiments

[0024]

Hereinafter, an embodiment of the present invention will be described with reference to the drawings. FIG. 1 to FIG. 4 are views illustrating examples of the present embodiment of the present invention. Portions assigned with the same reference numeral in the drawings indicate the same member. It is noted that in the drawings, some of the structures are appropriately omitted for the simplification of the drawings. The size, shape, thickness, and the like of members are appropriately exaggerated.

[0025]

<Test device>

FIG. 1 includes views illustrating a test device 10 according to the present embodiment. FIG. 1(A) is a plan view, FIG. 1(B) is an extracted plan view of an internal of the test device 10, and FIG. 1(C) is a cross-sectional side view of FIG. 1(B).

[0026]

The test device 10 is a device for antibody test by immunochromatography. The test device 10 performs a reaction between an antibody in a specimen and a labeled body containing an antigen, causes the produced immune complex to migrate by capillary action, traps the immune complex with a previously immobilized antibody, and determines the presence or absence of a target antibody.

[0027]

Specifically, as illustrated in FIG. 1(A), the test device 10 according to the present embodiment includes a first test piece 11, a second test piece 12, and a case 13.

[0028]

The case 13 is, for example, an elongated rectangular box in plan view, and includes a dropping window 130 on which a specimen is dropped, a first detection window 131, and a second detection window 132. The first test piece 11 and the second test piece 12 are both band-like porous members (for example, cellulose membrane), and are integrally accommodated

in the case 13 although they are provided as separate pieces.
The first test piece 11 and the second test piece 12 are
aligned so that a band lengthwise direction conforms to the
lengthwise direction of the case 13 (in FIG. 1, a left-right
5 direction). In this example, the first test piece 11 and the
second test piece 12 are accommodated in the case 13 so that
they are aligned (disposed in parallel) along the short-side
direction (in FIG. 1, vertical direction) of the case 13.

[0029]

10 The first detection window 131 is provided so that a part
of the first test piece 11 can be visually observed from the
outside. The second detection window 132 is provided so that a
part of the second test piece 12 can be visually observed from
the outside.

15 [0030]

The first test piece 11 can detect the presence of a
first antibody in the specimen to be dropped through the
dropping window 130. The second test piece 12 can detect the
presence of a second antibody in the same specimen as the
specimen dropped on the first test piece 11 or in a specimen
20 that is a sample of the same specimen and is dropped
approximately simultaneously with dropping on the first test
piece 11.

[0031]

25 The test device 10 is configured to determine a detection

result from the first test piece 11 visually observed through the first detection window 131 and a result from the second test piece 12 visually observed through the second detection window 132 in parallel and at the same time. More

5 specifically, the test device 10 is configured such that the results from the first test piece 11 and from the second test piece 12 are visually observed in parallel and approximately at the same time, producing various types of determinations. Herein, the expressions "in parallel and at the same time" and
10 "in parallel" mean that the test pieces 11 and 12 and the detection results thereof can be recognized collectively (at a glance without largely moving a line of sight), and are not limited to a case where the detection results of the test pieces 11 and 12 are obtained at an exactly simultaneous time,
15 and to a case where the results of the test pieces 11 and 12 are disposed (aligned) in parallel.

[0032]

The first antibody and the second antibody to be detected in the present embodiment are antibodies that are of the same
20 type but trap different antigens. Specifically, the "first antibody" in the present embodiment is an immunoglobulin present only in human blood, and for example, is an antibody that binds to a part of a spike protein of a predetermined virus (for example, a SARS-CoV-2 virus). More specifically,
25 the spike protein includes an S1 unit and an S2 unit, and the

first antibody in the present embodiment is an antibody (for example, IgG antibody) that binds to (trap) a receptor-binding domain (RBD) in the S1 unit. Hereinafter, the spike protein is referred to as an "S antigen", a receptor-binding domain in the S1 unit (RBD portion extracted from the S1 unit) is referred to as an "S1/RBD antigen", and an IgG antibody (first antibody) that binds to the S1/RBD antigen is referred to as an "S1/RBD-IgG antibody".

[0033]

The "second antibody" in the present embodiment is an immunoglobulin present only in human blood, and for example, is an antibody (for example, IgG antibody) that binds to (trap) a nucleocapsid protein of the predetermined virus (for example, a SARS-CoV-2 virus). Hereinafter, the nucleocapsid protein is referred to as an "NP antigen", and an IgG antibody (second antibody) that binds to the NP antigen is referred to as an "NP-IgG antibody".

[0034]

As described above, the first antibody (S1/RBD-IgG antibody) and the second antibody (NP-IgG antibody) in the present embodiment are the same type of antibody (for example, IgG antibody) present in human blood, but trap different antigens. In this respect, the first antibody and the second antibody are different types of IgG antibodies.

[0035]

Referring to FIGs. 1(B) and 1(C), the first test piece 11 is a member that includes a sample pad 111 to serve as a dropping portion on which the specimen is dropped, a first labeled body-containing portion 112 containing a first labeled body, a first detection portion 113, and an absorption pad 114 configured to absorb the dropped specimen. These constituents are connected so that their end portions superimpose each other in the band lengthwise direction. The specimen dropped on the sample pad 111 through the dropping window 130 migrates in the arrow direction toward the absorption pad 114 (in FIG. 1, from the left upstream to the right downstream). The "specimen" in the present embodiment is whole blood, serum, or plasma of human (subject), and may be a specimen liquid in which a developer is added thereto.

[0036]

More specifically, the first labeled body-containing portion 112 is a conjugate pad containing an antigen (labeled antigen) and an antibody (labeled antibody) each labeled with colored particles such as gold nanoparticles. In this example, the labeled antigen in the first labeled body-containing portion 112 is an S1/RBD antigen capable of binding to the S1/RBD-IgG antibody (capable of trapping the S1/RBD-IgG antibody). The "first labeled body" in the present embodiment is a labeled antigen capable of binding to the S1/RBD-IgG antibody. In other words, this labeled antigen is an S1/RBD

labeled antigen. The first labeled body binds to the S1/RBD-IgG antibody to form an immune complex (hereinafter referred to as a "first immune complex IC1") and migrates from the upstream side (in FIG. 1, left side) to the downstream side (in FIG. 1, right side). The labeled antibody in the first labeled body-containing portion 112 is a labeled antibody (labeled antibody LB1) of a non-human animal.

[0037]

The first detection portion 113 is a membrane filter that is provided downstream of the first labeled body-containing portion 112. The first detection portion 113 includes a capture antibody line TL1 and a control antibody line CL1. In these antibody lines TL1 and CL1, the respective antibodies, a capture antibody T1 and a control antibody C1, for example, are linearly applied and immobilized in a widthwise direction of a band. The capture antibody T1 is an antibody derived from an animal and capable of trapping the first immune complex IC1. The control antibody C1 is an antibody capable of trapping a labeled antibody LB1 (an antibody not involved in the first immune complex IC1). The absorption pad 114 is provided downstream of the first detection portion 113.

[0038]

The second test piece 12 is a member that includes the sample pad 111 to serve as a dropping portion on which the specimen is dropped, a second labeled body-containing portion

122 containing a second labeled body, a second detection portion 123, and the absorption pad 114 to absorb the dropped specimen. These constituents are connected so that their end portions superimpose each other in the band lengthwise

5 direction. The second test piece 12 is separate from the first test piece 11, and is disposed in parallel but separately from the first test piece 11. In this example, the sample pad 111 and the absorption pad 114 are used both for the first test piece 11 and the second test piece 12. Specifically, the

10 sample pad 111 is provided to be superimposed on the end portion (in this example, a left end portion) of each of both the first test piece 11 and the second test piece 12 in the band lengthwise direction, and the absorption pad 114 is provided to be superimposed on the other end portion (in this
15 example, a right end portion) of each of both the first test piece 11 and the second test piece 12 in the band lengthwise direction. The specimen dropped on the sample pad 111 migrates in the arrow direction toward the absorption pad 114 (in FIG. 1, from the left upstream to the right downstream).

20 [0039]

More specifically, the second labeled body-containing portion 122 is a conjugate pad containing an antigen (labeled antigen) and an antibody (labeled antibody) each labeled with colored particles such as gold nanoparticles. In this example,
25 the labeled antigen in the second labeled body-containing

portion 122 is an NP antigen capable of binding to the NP-IgG antibody (capable of trapping the NP-IgG antibody). The "second labeled body" in the present embodiment is a labeled antigen capable of binding to the NP-IgG antibody. In other words, this labeled antigen is an NP labeled antigen. The second labeled body binds to the NP-IgG antibody to form an immune complex (hereinafter referred to as a "second immune complex IC2") and migrates from the upstream side (in FIG. 1, left side) to the downstream side (in FIG. 1, right side). The labeled antibody in the second labeled body-containing portion 122 is a labeled antibody (labeled antibody LB2) of a non-human animal.

[0040]

The second detection portion 123 is a membrane filter that is provided downstream of the second labeled body-containing portion 122. The second detection portion 123 includes a capture antibody line TL2 and a control antibody line CL2. In these antibody lines TL2 and CL2, the respective antibodies, a capture antibody T2 and a control antibody C2, for example, are linearly applied and immobilized in the widthwise direction of the band. The capture antibody T2 is an antibody derived from an animal and capable of trapping the second immune complex IC2. The capture antibody T2 may be the same antibody as the capture antibody T1 in the first detection portion 113.

[0041]

The control antibody C2 is an antibody capable of trapping a labeled antibody LB2 (an antibody not involved in the second immune complex IC2).

5 [0042]

The first test piece 11 is not limited to a laminated structure in which the sample pad 111, the conjugate pad (the first labeled body-containing portion) 112, the membrane filter (the first detection portion) 113, and the absorption pad 114 each partly superimpose each other as illustrated in FIG. 1(C). For example, the first test piece 11 may have a structure in which the sample pad 111, the conjugate pad 112, the membrane filter 113, and the absorption pad 114 are continuously disposed along the band lengthwise direction of a single band-like porous member so that their respective corresponding regions are adjacent to each other. The first test piece 11 may also have a structure in which only some of these constituents are laminated, or the like. In the following description, the first test piece 11 having the laminated structure as illustrated in FIG. 1(C) will be described as an example. However, when the first test piece 11 has a shape of a single band (or a band in which some of the constituents superimpose each other), the constituents described as the sample pad 111, the conjugate pad 112, the membrane filter 113, and the absorption pad 114 should be

interpreted as regions corresponding to the respective constituents. For example, when "the sample pad 111" is described, it should be interpreted as "a region corresponding to the sample pad 111". The same is applied to the second test piece 12.

[0043]

<Detection principle of test device>

Referring to FIG. 2, the detection principle of the test device 10 will be described. In the detection principle, the first test piece 11 is comparable with the second test piece 12, and thus antigens and antibodies of the first test piece 11 and the second test piece 12 are collectively illustrated in FIG. 2. Specifically, in FIG. 2, the labeled antigen (the S1/RBD labeled antigen and the NP labeled antigen) are collectively referred to as LG, the labeled antibodies (the labeled antibodies LB1 and LB2) are collectively referred to as LB, the antibodies (the S1/SBD-IgG antibody and the NP-IgG antibody) to be detected in the specimen are collectively referred to as AB, the first immune complex IC1 and the second immune complex IC2) are collectively referred to as IC, the capture antibodies T1 and T2 are collectively referred to as T, and the control antibodies C1 and C2 are collectively referred to as C.

[0044]

When a specimen S is dropped through the dropping window

130 (not illustrated in FIG. 2), the specimen S permeates the first labeled body-containing portion 112 of the first test piece 11 and the second labeled body-containing portion 122 of the second test piece 12 through the sample pad 111

5 respectively (at approximately the same time) through the sample pad 111. Furthermore, the specimen S migrates to the downstream side through two routes via the first test piece 11 and the second test piece 12.

[0045]

10 In the first test piece 11, the specimen migrates to the first detection portion 113 while dissolving the labeled antigen (the S1/RBD labeled antigen, and in FIG. 2, the labeled antigen LG) in the first labeled body-containing portion 112. When the specimen S contains the S1/RBD-IgG
15 antibody to be tested (in FIG. 2, the antibody AB), the S1/RBD-IgG antibody (the antibody AB) binds to the S1/RBD labeled antigen (the labeled antigen LG) to form the first immune complex IC1 (in FIG. 2, the immune complex IC) and migrates downstream. The first immune complex IC1 (the immune
20 complex IC) binds to the capture antibody T1 (in FIG. 2, the complementary antibody T) in the capture antibody line TL1 and is trapped on the capture antibody line TL1. The trapping produces a state in which colored particles derived from the labeled antigen are concentrated on the capture antibody line
25 TL1, and the capture antibody line TL1 is colored. The first

detection window 131 is opened so that this coloring can be visually observed. When the colored state is visually observed, the presence or absence of the SI/RBD-IgG antibody in the specimen can be determined. In other words, when the capture antibody line TL1 is colored, it is determined that the specimen contains the SI/RBD-IgG antibody (positive).

[0046]

When the specimen migrates, the labeled antibody LB1 (in FIG. 2, the labeled antibody LB) in the first labeled body-containing portion 112 also migrates downstream. The labeled antibody LB1 (the labeled antibody LB) does not bind to the SI/RBD labeled antigen (the labeled antigen LG) (without participation in the first immune complex IC1) and is not trapped on the capture antibody line TL1. However, the labeled antibody LB1 is trapped by the control antibody C1 (in FIG. 2, the control antibody C) immobilized downstream. This trapping produces a state in which colored particles derived from the labeled antibody LB1 are concentrated on the control antibody line CL1, and the control antibody line CL1 is colored. The first detection window 131 is opened so that this coloring can be visually observed. When the colored state is visually observed, it is confirmed that the specimen passed through the capture antibody line TL1 and reached the control antibody line CL1. In other words, when the capture antibody line TL1 is not colored and the control antibody line CL1 alone is

colored, it is determined that the S1/RBD-IgG antibody is not present in the specimen (negative).

[0047]

The second test piece 12 determines the presence or
5 absence of the NP-IgG antibody in the same specimen as the
specimen S that is tested by the first test piece 11. Since
this principle is the same as that in the first test piece 11,
a description will be partially omitted. The specimen migrates
to the second detection portion 123 while dissolving the
10 labeled antigen (the NP labeled antigen and the labeled
antigen LG) in the second labeled body-containing portion 122.
When the specimen contains the NP-IgG antibody to be tested
(the antibody AB), the NP-IgG antibody binds to the NP labeled
antigen (the labeled antigen LG) to form the second immune
15 complex IC2 (the immune complex IC), and the second immune
complex IC2 is trapped on the capture antibody line TL2. By
this trapping, the capture antibody line TL2 is colored. The
colored state can be visually observed through the second
detection window 132. When the capture antibody line TL2 is
20 colored, it is determined that the specimen contains the NP-
IgG antibody (positive). In contrast, when the capture
antibody line TL2 is not colored and the control antibody line
CL2 alone is colored, it is determined that the NP-IgG
antibody is not present (negative).

[0048]

In the present embodiment, the specimen dropped through one dropping window 130 in the test device 10 is absorbed by the sample pad 111 of the first test piece 11 and the second test piece 12 approximately at the same time, and then
5 migrates downstream through two routes via the test pieces 11 and 12. The first test piece 11 can determine the presence or absence of the S1/RBD-IgG antibody in the specimen. The second test piece 12 can determine the presence or absence of the NP-IgG antibody in the specimen.

10 [0049]

Thus, the test device 10 according to the present embodiment can detect the presence or absence of two types of antibodies (the S1/RBD-IgG antibody and the NP-IgG antibody in the specimen) in parallel and at the same time by one test,
15 where these antibodies are present in the same specimen and of the same type (for example, IgG antibody), and trap different antigens. This configuration can determine the state of the specimen by combining detection results of the two types of antibodies that trap different antigens contains in the same
20 antibody. Thus, the test device 10 according to the present embodiment can be applied to various types of determinations including determining the efficacy of predetermined vaccination.

[0050]

25 <Test Method>

Hereinafter, a method for testing an antibody according to the present embodiment will be described. A test method using the aforementioned test device 10 will be described. The test method includes a first test step of dropping the specimen on the first test piece 11 in the test device 10 to detect the presence or absence of the first antibody in the specimen, a second test step of dropping the specimen on the second test piece 12 in the test device 10 to detect the presence or absence of the second antibody in the specimen, and a determination step of combining results from the first test step and the second test step, that is, a detection state by the first detection portion 113 that can be visually observed through the first detection window 131 and a detection state by the second detection portion 123 that can be visually observed through the second detection window 132 to determine the state of the specimen. At the determination step, results of the presence or absence of the S1/RBD-IgG antibody and the presence or absence of the NP-IgG antibody in the specimen are combined to determine the final determination. Thus, one determination is derived from a plurality of determination candidates including a determination on a possibility that the antibody is produced by vaccination.

[0051]

As described in the detection principle of the test

device, when the specimen is dropped through the dropping window 130 in the test device 10, the first detection portion in the first test piece 11 detects the presence or absence of the S1/RBD-IgG antibody in the specimen. The S1/RBD-IgG antibody is an antibody (neutralizing antibody) to target a spike protein (S antigen) of the novel coronavirus. The second detection portion in the second test piece 12 detects the presence or absence of the NP-IgG antibody in the same specimen. In the present embodiment, the state of the specimen, specifically whether natural infection with the coronavirus has occurred and whether the neutralizing antibody is produced, is determined on the basis of the results.

[0052]

However, when the presence or absence of the neutralizing antibody in the specimen can be detected by the test device 10, whether the neutralizing antibody is produced by viral infection or vaccination cannot be determined. The test method according to the present embodiment makes it possible to determine whether the detected S1/RBD-IgG antibody (neutralizing antibody) may have been produced by vaccination by combining the result from the first test piece 11 (the presence or absence of the S1/RBD-IgG antibody in the specimen) and the result from the second test piece 12 (the presence or absence of the NP-IgG antibody in the specimen).

[0053]

Referring to FIG. 3 and FIG. 4, a description will be given of the test method according to the present embodiment, and a determination on whether the neutralizing antibody is acquired or the efficacy of vaccination (for example, a determination on a possibility that the neutralizing antibody has been produced by vaccination (a possibility that the detected S1/RBD-IgG antibody has been produced by vaccination)). FIG. 3 is a flowchart illustrating an example of the test method according to the present embodiment. FIG. 4 includes plan views illustrating an outline of the test results of the first test piece 11 and the second test piece 12.

[0054]

<<First case>>

A determination in a first case using the test method according to the present embodiment will be described. The specimen is dropped on the test device 10 (Step S01), and the first detection window 131 in the first test piece 11 is visually observed to check the test result. When the capture antibody line TL1 is colored, that is, when the result is positive (Yes in Step S03), it is determined that the S1/RBD-IgG antibody (the neutralizing antibody) is present in the specimen (Step S05).

[0055]

Subsequently, the second detection window 132 in the

second test piece 12 detected in parallel and at the same time is visually observed to check the test result. When the capture antibody line TL2 is colored, that is, when the result is positive (Yes in Step S07), this means that the NP-IgG antibody is present (see FIG. 4(A)).

[0056]

When the detection results of the first test piece 11 and the second test piece 12 are combined, the determination in the first case is that "subject has a history of natural viral infection, and neutralizing antibody is produced (acquired): determination a" (Step S09).

[0057]

In the first case, the neutralizing antibody is acquired, but whether the neutralizing antibody is acquired by vaccination cannot be specified. However, when it is established that a subject is vaccinated before the test, a possibility that the neutralizing antibody is acquired by vaccination is also considered.

[0058]

<<Second case>>

Next, a determination in a second case will be described. When the capture antibody line TL1 of the first test piece 11 is colored, that is, when the result is positive (Yes in Step S03), it is determined that the S1/RBD-IgG antibody (the neutralizing antibody) is present (acquired). In this stage,

however, it is not obvious whether the neutralizing antibody is produced (acquired) by vaccination or viral infection (natural infection).

[0059]

5 When the capture antibody line TL2 is not colored, that is, when the result is negative (No in Step S07), it is determined that the NP-IgG antibody is not present (see FIG. 4(B)).

[0060]

10 In this case, from only the result that the NP-IgG antibody is not present, the possibility of natural viral infection can be determined to be low. Therefore, when it is not established that the subject is vaccinated, it is determined from the positive result in the first test piece 11
15 that the neutralizing antibody has been produced by natural infection. When it is established that the subject is vaccinated, the determination is that the neutralizing antibody has been produced by vaccination.

[0061]

20 Like this, when the detection results of the first test piece 11 and the second test piece 12 are combined, the determination in the second case is that "neutralizing antibody is produced (acquired), and neutralizing antibody may be acquired by vaccination: determination b" (Step S11).

25 Again, in this case, particularly, when it is established that

the subject is vaccinated before the test, the possibility that the neutralizing antibody in the specimen is acquired by vaccination is higher, as described above.

[0062]

5 <<Third case>>

Next, a determination in a third case will be described. When the capture antibody line TL1 of the first test piece 11 is not colored, that is, when the result is negative (No in Step S03), it is determined that the S1/RBD-IgG antibody (neutralizing antibody) is not present (is not acquired) (Step S13). When the capture antibody line TL2 of the second test piece 12 is colored, that is, when the result is positive (Yes in Step S15), it is determined that the NP-IgG antibody is present (see FIG. 4(C)).

15 [0063]

In this case, when the detection results of the first test piece 11 and the second test piece 12 are combined, the determination in the third case is that "subject has a history of natural viral infection, and neutralizing antibody is not produced: determination c" (Step S17). In the third case, for example, when it is established that the subject is vaccinated before the test, the determination is that the vaccination is not effective (has not been still effective).

[0064]

25 <<Fourth case>>

Next, a determination in a fourth case will be described. When the capture antibody line TL1 of the first test piece 11 is not colored, that is, when the result is negative (No in Step S03), it is determined that the S1/RBD-IgG antibody (neutralizing antibody) is not present (is not acquired). When the capture antibody line TL2 of the second test piece 12 is not colored, that is, when the result is negative (No in Step S15), it is determined that the NP-IgG antibody is not present (see FIG. 4(D)).

10 [0065]

In this case, when the detection results of the first test piece 11 and the second test piece 12 are combined, the determination in the fourth case is that "subject has no history of natural viral infection, and neutralizing antibody is not produced: determination d" (Step S19). In the fourth case, for example, when it is established that the subject is vaccinated before the test, the determination is that the vaccination is not effective (has not been still effective).

[0066]

20 Thus, the test device 10 and the test method according to the present invention can derive one determination from a plurality of determination candidates including the efficacy of a predetermined vaccine (the determination a illustrated in Step S09 in FIG. 3 and in FIG. 4(E), the determination b illustrated in Step S11 in FIG. 3 and in FIG. 4(F), the

25

determination c illustrated in Step S17 in FIG. 3 and in FIG. 4(G), and the determination d illustrated in Step S19 in FIG. 3 and in FIG. 4(G)). Therefore, a simple and quick method can test and determine the efficacy of a vaccine against a predetermined virus (for example, novel coronavirus), for example, a fact where the neutralizing antibody is not acquired although it is established that the subject is vaccinated (the third case and the fourth case), or when the neutralizing antibody is present in the specimen, whether the neutralizing antibody is produced by natural viral infection or vaccination (the second case) or the like.

[0067]

The test method according to the present embodiment makes various determinations of the antibodies by the following processes. In this case, the test method is not limited to a method using the test device 10.

[0068]

Specifically, the method is a test method for determining the presence or absence of a first antibody and a second antibody in the specimen by immunochromatography where the first antibody is an antibody produced by vaccination or infection with a predetermined virus, and the second antibody is an antibody produced by infection with the virus. Then, the method includes: dropping the specimen on the first test piece 11 to determine the presence or absence of the first antibody

in the specimen, dropping the same specimen on the second test piece 12 different from the first test piece 11 to determine the presence or absence of the second antibody in the specimen, and combining a test result from the first test piece 11 and a test result from the second test piece 12 to derive a determination from determination candidates including the possibility that the first antibody is produced in the specimen by vaccination.

[0069]

In this case, particularly, when the test result from the first test piece 11 is positive and the test result from the second test piece 12 is negative, it is determined that there is a possibility that the first antibody may have been produced in the specimen by the vaccination.

[0070]

<Test kit>

Next, a test kit 30 according to the present embodiment will be described with reference to FIG. 5. The test kit 30 according to the present embodiment includes the test device 10 and a determination guide 20. The determination guide 20 allows the user to determine the state of the specimen including a possibility that the antibody is produced by predetermined vaccination on the basis of the test result from the first test piece 11 and the test result from the second test piece 12.

[0071]

The determination guide 20 may be a printed matter separate from the test device 10, such as paper, as illustrated in FIG. 5(A), or may be a guide that is displayed (printed) directly on the test device 10 as illustrated in FIG. 5(B). For example, when the determination guide 20 is paper, a flowchart as illustrated in FIG. 3 may be printed on paper. Sets of the test results of the first test piece 11 and the second test piece 12 as illustrated in FIG. 4, that is, the colored state (schematic view) (FIGs. 4(A) to 4(D)), and determinations based on the colored states (FIGs. 4(E) to 4(F), the determinations a to d) may be printed.

[0072]

The determination guide 20 may be provided, for example, in a form of an application of a smartphone, a personal computer, or the like although the illustration thereof is omitted. In this case, when the application is initiated, for example, questions as illustrated in the flowchart in FIG. 3 are successively displayed. When answers are input in accordance with the display, any one of the determinations a to d may be displayed. Furthermore, when the application is initiated, a set of the colored states (schematic views) of the first test piece 11 and the second test piece 12 and the determination based on the colored states (any of the determinations a to d) may be displayed. Alternatively, the

application may be configured so that when the application is initiated, the application is shifted to a photographing mode or the like. When the actual test results of the first test piece 11 and the second test piece 12 are photographed and sent, any one of the determinations a to d illustrated in FIGs. 4(E) to 4(H) may be displayed.

[0073]

The determination guide 20 is not limited to the aforementioned examples and may be in any form as long as an examiner can uniquely acquire the state of the specimen (specifically, any of the determinations a to d illustrated in FIGs. 4(E) to 4(H)) including a possibility that the antibody is produced by predetermined vaccination based on the test results of the first test piece 11 and the second test piece 12.

[0074]

FIG. 6 is a view illustrating another example of the determination guide 20. The aforementioned determinations a to d are determinations in a stage where the presence or absence of vaccination in the subject is not (cannot be) specified. However, the determination guide 20 in this example illustrated in FIG. 6 makes a more detailed determinations possible on the basis of the aforementioned determinations a to d and the presence or absence of vaccination in the subject (a fact that the subject is vaccinated).

[0075]

Specifically, when the determination a is given and when it is established that the subject is vaccinated, the determination is that "neutralizing antibody has been produced by vaccination or natural infection". To a user of the test kit 30, for example, "Neutralizing antibody has been produced by vaccination or natural infection (determination a-1)" or the like is displayed.

[0076]

When the determination a is given and when it is not established that the subject is vaccinated, the determination is that "neutralizing antibody has been produced by natural infection". To the user, for example, "Neutralizing antibody may be produced by natural infection (Determination a-2)" or the like is displayed.

[0077]

When the determination b is given and when it is established that the subject is vaccinated, it is determined that "there is a high possibility that neutralizing antibody is produced by vaccination (since the result from the second test piece 12 is negative)". To the user, for example, "There is a high possibility that neutralizing antibody is produced by vaccination (Determination b-1)" or the like is displayed.

[0078]

When the determination b is given and when it is

established that the subject is vaccinated, it is determined that "there is a possibility that neutralizing antibody may be produced by natural infection (since the result from the first test piece 11 is positive)". To the user, for example, "there is a possibility that neutralizing antibody may be produced by natural infection (Determination b-2)" or the like is displayed.

[0079]

When the determination c is given and when it is established that the subject is vaccinated, it is determined that "there is a high possibility that neutralizing antibody is not produced although subject is vaccinated. Although subject has a history of natural infection, neutralizing antibody is not produced by the natural infection". To the user, for example, "It seems that neutralizing antibody has not been produced (by vaccination), but subject has a history of natural infection (Determination c-1)" or the like is displayed.

[0080]

When the determination c is given and when it is not established that the subject is vaccinated, it is determined that "subject has a history of natural infection but neutralizing antibody is not produced (by natural infection)". To the user, for example, "Subject has a history of natural infection, but neutralizing antibody is not produced

(Determination c-2)" or the like is displayed.

[0081]

When the determination d is given and when it is established that the subject is vaccinated, it is determined
5 that "There is a high possibility that neutralizing antibody is not produced although the subject is vaccinated. Subject has no history of natural infection". To the user, for example, "It seems that neutralizing antibody has not been produced by vaccination. Subject has no history of natural
10 infection (Determination d-1)" or the like is displayed.

[0082]

When the determination d is given and when it is not established that the subject is vaccinated, it is determined that "subject has no history of natural infection, and
15 neutralizing antibody is not produced". To the user, for example, "subject has no history of natural infection and neutralizing antibody is not produced (Determination d-2)" or the like is displayed. For example, to a subject who does not acquire the neutralizing antibody, a recommendation such as
20 that "Vaccination recommended" or the like may be displayed.

[0083]

The flow of determination illustrated in FIG. 6 may be displayed, for example, after the flowchart (Steps S09, S11, S17 and S19) illustrated in FIG. 3 that is the determination
25 guide 20. For example, following the results of determinations

illustrated in FIG. 4, which is the determination guide 20,
the user may be allowed to select the presence or absence of
vaccination to derive the final determinations a-1 to d-2
illustrated in FIG. 6. The final determinations a-1 to d-2
5 illustrated in FIG. 6 may be derived by the application or the
like.

[0084]

When the determination guide 20 as illustrated in FIG. 6
is used, the user obtains the final determinations a-1 to d-2.
10 Thus, the state of the subject can be simply and uniquely
ascertained.

[0085]

An example of the present embodiment of the present
invention have been described above, but the present invention
15 is not limited to the example.

For example, one test device 10 may include a plurality
of dropping windows 130. Specifically, the dropping window 130
may be provided in each of both the first test piece 11 and
the second test piece 12, and the same specimen may be dropped
20 through each of the dropping windows 130.

[0086]

An example in which the first test piece 11 and the
second test piece 12 are disposed in parallel along the short-
side direction of the case 13 has been described, and the
25 present invention is not limited to this example. The first

test piece 11 and the second test piece 12 may be aligned linearly (separate from each other) along the lengthwise direction of the case 13. However, when the test pieces 11 and 12 are aligned linearly, it is possible that the colored states of the test pieces may be more difficult to be visually observed. Furthermore, it is necessary to drop the specimen twice. When the test pieces 11 and 12 are disposed in parallel, the colored states of the test pieces are likely to be collectively recognized as a pattern, for example, as illustrated in FIG. 4, and easily compared with the determination guide 20 and the like. The sample pad 111 can be shared by the first test piece 11 and the second test piece 12 as illustrated in FIG. 1, and the specimen can be supplied to the first test piece 11 and the second test piece 12 by one procedural step of dropping the specimen.

[0087]

In the present embodiment, a case where the antibody to be detected is an IgG antibody has been described. However, the antibody to be detected may be an IgM antibody or other immunoglobulins.

[0088]

A configuration of the sample pad 111 and the absorption pad 114 is not limited to the configuration in which the sample pad 111 and the absorption pad 114 are used for both the first test piece 11 and the second test piece 12. The

sample pad 111 and the absorption pad 114 may be provided separately in each of both the first test piece 11 and the second test piece 12.

[0089]

5 The labeled antibody LB1 in the first labeled body-containing portion 112, the labeled antibody LB2 in the second labeled body-containing portion 122, and the control antibodies C1 and C2 may be different antibodies, or at least one set of the antibodies among these may be the same
10 antibodies.

[0090]

 It should be noted that the test device 10 of the present invention is not limited to the above-described embodiment and can be modified in various manner without departing from the
15 scope of the present invention.

Reference Signs List

[0091]

10	test device
11	first test piece
20	12 second test piece
	20 determination guide
	30 test kit
	111 sample pad
	112 first labeled body-containing portion (conjugate
25	pad)

113 first detection portion (membrane filter)
114 absorption pad
122 second labeled body-containing portion (conjugate
pad)

5 123 second detection portion (membrane filter)

130 dropping window

131 first detection window

132 second detection window

C1, C2 control antibody

10 CL1 control antibody line

CL2 control antibody line

T1, T2 capture antibody

TL1 capture antibody line

TL2 capture antibody line

15

CLAIMS

1. A test device by immunochromatography comprising:
a first test piece;
a second test piece separate from the first test piece;

5 and

a case configured to integrally accommodate the first
test piece and the second test piece, wherein:

the case includes

a dropping window on which a specimen is dropped,

10 a first detection window through which a part of
the first test piece can be visually observed from outside,
and

a second detection window through which a part of
the second test piece can be visually observed from outside;

15 the first test piece can detect presence of a first
antibody in the specimen;

the second test piece can detect presence of a second
antibody different from the first antibody in the specimen;
and

20 a result from the first test piece visually observed
through the first detection window and a result from the
second test piece visually observed through the second
detection window can be determined in parallel and at the same
time.

25 2. The test device according to claim 1, wherein the first

antibody and the second antibody are antibodies that are of the same type and trap different antigens.

3. The test device according to claim 1 or 2, wherein the first test piece and the second test piece are disposed in parallel.

4. The test device according to claim 1 or 2, wherein:

the first test piece includes at least a first labeled body-containing portion containing a first labeled body capable of trapping the first antibody, and a first detection portion capable of trapping a complex of the first antibody and the first labelled body; and

the second test piece includes at least a second labeled body-containing portion containing a second labeled body capable of trapping the second antibody, and a second detection portion capable of trapping a complex of the second antibody and the second labelled body.

5. The test device according to claim 3, wherein:

the first labeled body includes, as an antigen, a part of a spike protein of a predetermined virus; and

the second labeled body includes, as an antigen, a nucleocapsid protein of the virus.

6. The test device according to claim 4, wherein:

the first antibody is an antibody produced by vaccination or infection with a predetermined virus; and

the second antibody is an antibody produced by infection

with the virus.

7. The test device according to claim 4 or 5, wherein the virus is a SARS-CoV-2 virus.

8. A test kit comprising:

5 the test device according to any one of claims 1 to 6;
and

a determination guide that allows for determining a state of a specimen including a possibility that an antibody is produced by vaccination on a basis of a test result from the first test piece and a test result from the second test piece.

9. A test method using the test device according to any one of claims 1 to 6, the method comprising:

a first test step of dropping the specimen on the first test piece to detect presence or absence of the first antibody in the specimen;

a second test step of dropping the specimen on the second test piece different from the first test piece to detect presence or absence of the second antibody in the specimen;
and

20 a determination step of combining results from the first and second test steps to determine a state of the specimen.

10. The test method according to claim 8, wherein the determination step derives a determination from a plurality of determination candidates including a determination on a possibility that antibodies are produced by vaccination.

11. A test method for determining presence or absence of a first antibody and a second antibody in a specimen by immunochromatography, where the first antibody is an antibody produced by vaccination or infection with a predetermined virus, and the second antibody is an antibody produced by infection with the virus,

the method comprising:

dropping the specimen on a first test piece to determine presence or absence of the first antibody in the specimen;

dropping the specimen on a second test piece different from the first test piece to determine presence or absence of the second antibody in the specimen; and

combining a test result from the first test piece and a test result from the second test piece to derive a determination from determination candidates including a possibility that the first antibody is produced by the vaccination in the specimen.

12. The test method according to claim 10, wherein when the test result from the first test piece is positive and the test result from the second test piece is negative, it is determined that the first antibody may have been produced in the specimen by the vaccination.

13. The test method according to claim 10, wherein the virus is a SARS-CoV-2 virus.