


Claims

1. An extended-release conjugate of a GLP-1 agonist comprising an insoluble matrix with a multiplicity of covalently attached linker-agonist peptide, wherein the extended-release conjugate has the formula M-(L-E)_a, wherein:

M is an insoluble matrix comprising a crosslinked polyethylene glycol hydrogel having crosslinkers cleavable by beta elimination;

L is a cleavable linker of formula (3)



wherein R¹ is CN or R³SO₂, wherein R³ is optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, or (R⁹)₂N, wherein each R⁹ is independently H or optionally substituted alkyl;

R² is H;

one of R⁵ is (CH₂)_yZ*, (CH₂CH₂O)_xCH₂CH₂Z* or (CH₂)_yNH-CO-(CH₂CH₂O)_xCH₂CH₂Z*, wherein x is 1-100, y = 1-6, and the other R⁵ is H or unsubstituted alkyl, and Z* comprises the residue of a functional group coupled to M, wherein the functional group comprises N₃, NH₂, NH-CO₂^tBu, SH, S^tBu, maleimide, CO₂H, CO₂^tBu, 1,3-diene, cyclopentadiene, furan, alkyne, cyclooctyne, acrylate, aminooxy, keto or acrylamide;


E is an agonist peptide of SEQ ID NO: 2; and

a is an integer representing the number of L-E units required to provide a concentration of 1-1000 mg peptide in one ml of the volume occupied by the matrix.

2. The extended-release conjugate of claim 1 wherein R¹ is CN or CH₃SO₂; R² is H; one R⁵ is (CH₂)_yZ* wherein y is 5; and the other R⁵ is H.

3. The extended-release conjugate of claim 1 or 2 wherein the hydrogel is in the form of microspheres.

4. The extended-release conjugate of any one of claims 1-3 wherein said crosslinkers contained in the hydrogel are of formula (1)



wherein: m is 0 or 1; and

wherein X and one of R¹, R² and R⁵ is coupled to a polymer contained in the hydrogel, and wherein X comprises the residue of a functional group coupled to a polymer contained in the hydrogel, wherein the functional group comprises N₃, NH₂, NH-CO₂^tBu, SH, S^tBu, maleimide, CO₂H, CO₂^tBu, 1,3-diene, cyclopentadiene, furan, alkyne, cyclooctyne, acrylate, aminooxy, keto or acrylamide,

with the proviso that at least one of R¹ and R² is CN; NO₂;

optionally substituted aryl;

optionally substituted heteroaryl;

optionally substituted alkenyl;

optionally substituted alkynyl;

COR³ or SOR³ or SO₂R³ wherein

R³ is H or optionally substituted alkyl;

aryl or arylalkyl, each optionally substituted;

heteroaryl or heteroarylalkyl, each optionally substituted; or

OR⁹ or NR⁹₂ wherein each R⁹ is independently H or optionally substituted alkyl, or both R⁹ groups taken together with the nitrogen to which they are attached form a heterocyclic ring;

SR⁴ wherein

R⁴ is optionally substituted alkyl;

aryl or arylalkyl, each optionally substituted; or


heteroaryl or heteroarylalkyl, each optionally substituted;

wherein R¹ and R² may be joined to form a 3-8 membered ring; and

wherein any remaining R¹ and R² is H or is alkyl, arylalkyl or heteroarylalkyl, each optionally substituted; and

any remaining R⁵ is independently H or is alkyl, alkenylalkyl, alkynylalkyl, (OCH₂CH₂)_p O-alkyl wherein p=1-1000, aryl, arylalkyl, heteroaryl or heteroarylalkyl, each optionally substituted; or

said crosslinkers when contained in the hydrogel are of formula (2)



wherein two of R¹, R² and R⁵ are coupled to a polymer contained in the hydrogel;



m is 0-1;

n is 1-1000;

s is 0-2;

t is 2,4, 8, 16 or 32;

Q is a core group having the valency t;



wherein R⁶ is H, optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted arylalkyl, or optionally substituted heteroarylalkyl; and

with the proviso that at least one of R¹ and R² is CN; NO₂;

optionally substituted aryl;

optionally substituted heteroaryl;

optionally substituted alkenyl;

optionally substituted alkynyl;

COR³ or SOR³ or SO₂R³ wherein

R³ is H or optionally substituted alkyl;

aryl or arylalkyl, each optionally substituted;

heteroaryl or heteroarylalkyl, each optionally substituted; or

OR⁹ or NR⁹₂ wherein each R⁹ is independently H or optionally substituted alkyl, or both R⁹ groups taken together with the nitrogen to which they are attached form a heterocyclic ring;

SR⁴ wherein

R⁴ is optionally substituted alkyl;

aryl or arylalkyl, each optionally substituted; or

heteroaryl or heteroarylalkyl, each optionally substituted;


wherein R¹ and R² may be joined to form a 3-8 membered ring; and

wherein any remaining R¹ and R² is H or is alkyl, arylalkyl or heteroarylalkyl, each optionally substituted; and

any remaining R⁵ is independently H or is alkyl, alkenylalkyl, alkynylalkyl, (OCH₂CH₂)_p O-alkyl wherein p=1-1000, aryl, arylalkyl, heteroaryl or heteroarylalkyl, each optionally substituted.

5. The extended-release conjugate of claim 1 wherein the linker-agonist peptide is covalently coupled to the crosslinkers.

6. A compound of formula (4)



wherein:

E is of SEQ ID NO: 2;

R¹ is CN or R³SO₂, wherein R³ is optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, or (R⁹)₂N, wherein each R⁹ is independently H or optionally substituted alkyl;

R² is H; and

one of R^5 is $(CH_2)_yZ$, $(CH_2CH_2O)_xCH_2CH_2Z$ or $(CH_2)_yNH-CO-(CH_2CH_2O)_xCH_2CH_2Z$, wherein x is 1-100, $y = 1-6$, and the other R^5 is H or unsubstituted alkyl, and Z is a functional group for coupling to an insoluble matrix, wherein said functional group comprises N_3 , NH_2 , $NH-CO_2^tBu$, SH , S^tBu , maleimide, CO_2H , CO_2^tBu , 1,3-diene, cyclopentadiene, furan, alkyne, cyclooctyne, acrylate, aminooxy, keto or acrylamide.

7. The compound of claim 6 wherein R^1 is CN or CH_3SO_2 ; R^2 is H; one R^5 is $(CH_2)_yZ$ wherein y is 5; and the other R^5 is H.

8. Use of the conjugate of any one of claims 1-5 in the manufacture of a medicament to be administered to a subject for the treatment of a condition benefited by a GLP-1 agonist, wherein the medicament is to be administered at one dosage per one to three months.