

ANTI-SIRP-BETA1 ANTIBODIES AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of priority of US Provisional Application No. 62/691,913, filed June 29, 2018, which is incorporated by reference herein in its entirety for any purpose.

FIELD OF THE PRESENT DISCLOSURE

[0002] The present disclosure relates to anti-SIRP β 1 antibodies and therapeutic uses of such antibodies.

BACKGROUND OF THE PRESENT DISCLOSURE

[0001] Signal regulatory protein beta (SIRP β) belongs to the SIRP family of transmembrane receptors, which are expressed within the myeloid cell lineage (including monocytes, macrophages, granulocytes, and dendritic cells) and in neuronal cells. The SIRP family of proteins is characterized by an extracellular region containing 2 membrane-proximal IgC domains and a distal IgV domain. Unlike SIRP α receptors, SIRP β proteins contain short cytoplasmic domains that lack cytoplasmic sequence motifs capable of recruiting protein tyrosine phosphatase SHP-2 and SHP-1. SIRP β 1 isoform 1 associates with the adaptor protein DAP12, which contains a single cytoplasmic immunoreceptor tyrosine-based activating (ITAM) motif. *See, e.g.*, Dietrich et al. 2000, J Immunol 164:9-12; Liu et al. 2005 J Biol Chem 280:36132-36140. SIRP β has been shown to be a microglial modulator of phagocytosis in Alzheimer's disease. *See, e.g.*, Gaikwad et al. 2009 Am J Pathol 175:2528-2539.

[0002] Accordingly, there is a need for therapeutic anti-SIRP β 1 antibodies to treat disease, disorders, and conditions associated with SIRP β 1 activity.

SUMMARY OF THE PRESENT DISCLOSURE

[0003] In one example, the present disclosure refers to an isolated antibody that binds to human SIRP β 1, wherein the antibody has at least one, at least two, at least three, at least four, or at least five properties selected from:

- a) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α ;
- b) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP γ ;
- c) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP β 1 isoform 3;
- d) binds to human SIRP β 1 isoform 1, but does not bind to mouse SIRP β 1;
- e) binds to human SIRP β 1 isoform 1, but does not bind to cynomolgus monkey SIRP β 1 isoform 1;
- f) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
- g) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
- h) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;

- i) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
- j) induces neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
- k) increases neutrophil-mediated tumor cell clearance *in vivo*;
- l) increases TREM2 expression on macrophages *in vitro* and/or *in vivo*;
- m) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
- n) increases viability of dendritic cells *in vitro* and/or *in vivo*.

[0003a] In one aspect, the present disclosure refers to an isolated antibody that binds to human SIRP β 1, wherein the antibody binds to human SIRP β 1 isoform 1, but does not bind to human SIRP β 1 isoform 3, and optionally wherein the antibody has at least one, at least two, at least three, or at least four properties selected from:

- a) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α ;
- b) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP γ ;
- c) binds to human SIRP β 1 isoform 1, but does not bind to mouse SIRP β 1;
- d) binds to human SIRP β 1 isoform 1, but does not bind to cynomolgus monkey SIRP β 1 isoform 1;
- e) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
- f) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
- g) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
- h) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
- i) induces neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
- j) increases neutrophil-mediated tumor cell clearance *in vivo*;
- k) increases TREM2 expression on macrophages *in vitro* and/or *in vivo*;
- l) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
- m) increases viability of dendritic cells *in vitro* and/or *in vivo*.

[0004] In some examples, the antibody binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α or human SIRP β 1 isoform 3. In some examples, the antibody binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α , SIRP γ , or human SIRP β 1 isoform 3. In some examples, the antibody has at least one, at least two, or at least three properties selected from:

- a) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
- b) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
- c) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
- d) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
- e) induces neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
- o) increases neutrophil-mediated tumor cell clearance *in vivo*;
- p) increases TREM2 expression on macrophages *in vitro* and/or *in vivo*;
- q) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
- f) increases viability of dendritic cells *in vitro* and/or *in vivo*.

[0005] In one example, the present disclosure refers to an isolated antibody that binds to human SIRP β 1, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises (a) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (b) HVR-H2 comprising an amino acid sequence of an HVR-H2 shown in Table 4 and/or Table 9; (c) HVR-H3 comprising an amino acid sequence of an HVR-H3 shown in Table 4 and/or Table 9; (d) HVR-L1 comprising an amino acid sequence of an HVR-L1 shown in Table 3 and/or Table 8; (e) HVR-L2 comprising an amino acid sequence of an HVR-L2 shown in Table 3 and/or Table 8; and (f) HVR-L3 comprising an amino acid sequence of an HVR-L3 shown in Table 3 and/or Table 8. In some examples, the heavy chain variable region comprises one, two, three or four framework regions selected from VH FR1, VH FR2, VH FR3, and

VH FR4 shown in Table 6. In some examples, the light chain variable region comprises one, two, three or four framework regions selected from VL FR1, VL FR2, VL FR3, and VL FR4 shown in Table 5.

[0005a] In one aspect, the present disclosure refers to an isolated antibody that binds to human SIRPβ1, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises (a) HVR-H1 comprising an amino acid sequence of SEQ ID NO 80; (b) HVR-H2 comprising an amino acid sequence of SEQ ID NO: 101; (c) HVR-H3 comprising an amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising an amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising an amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising an amino acid sequence of SEQ ID NO: 31.

[0006] In some examples, the antibody comprises HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9). In some examples, the antibody comprises a heavy chain variable region comprising an amino acid sequence at least 90%, at least 95%, at least 97%, or at least 99% identical to a sequence selected from SEQ ID NOs: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382. In some examples, the antibody comprises a heavy chain variable region selected from SEQ ID NOs: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382. In some examples, the antibody comprises a light chain variable region comprising an amino acid sequence at least 90%, at least 95%, at least 97%, or at least 99% identical to a sequence selected from SEQ ID NOs: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370. In some examples, the antibody comprises a light chain variable region selected from SEQ ID NOs: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the heavy chain variable region, and a light chain variable region 90%, at least 95%, at least 97%, or at least 99% identical to the light chain variable region, of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-

1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region and a light chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21.

[0007] In some examples, the antibody comprises HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9). In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the heavy chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the light chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a light chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the heavy chain variable region, and a light chain variable region 90%, at least 95%, at least 97%, or at least 99% identical to the light chain variable region, of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-

40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region and a light chain variable region of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21.

[0008] In some examples, the antibody comprises HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 8 and 9). In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the heavy chain variable region of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the light chain variable region of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a light chain variable region of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the heavy chain variable region, and a light chain variable region 90%, at least 95%, at least 97%, or at least 99% identical to the light chain variable region, of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some examples, the antibody comprises a heavy chain variable region and a light chain variable region of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21.

[0009] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 80, 228, or 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 101, 239, 239, 240, or 241; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31. In some examples, the antibody comprises a heavy chain variable

region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 366, 367, 368, or 369. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 267. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 366, 367, 368, or 369. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 267.

[0010] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 81, 99, 230, 231, or 232; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 102, 242, 243, 244, or 245; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126 or 253; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 371, 372, 373, 374, or 375. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 269 or 370. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 371, 372, 373, 374, or 375. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 269 or 370. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 371, 372, or 373 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 370; or a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 374 or 375 and a light chain variable region comprising the amino acid sequence of SEQ ID NO: 269.

[0011] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 85, 233, or 234; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 107, 246, 247, or 248; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131 or 254; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 376, 377, or 378. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 280. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 376, 377, or 378. In

some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 280.

[0012] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 97, 235, 236, or 237; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 121, 249, 250, 251, or 252; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 163; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 2; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 22; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 69. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 379, 380, 381, or 382. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 344. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 379, 380, 381, or 382. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 344.

[0013] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 239; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 367. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 267. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 367. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 267.

[0014] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 231; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 243; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 372. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 370. In

some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 372. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 270.

[0015] In some examples, the antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 246; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37. In some examples, the antibody comprises a heavy chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 376. In some examples, the antibody comprises a light chain variable region that is at least 90%, at least 95%, at least 97%, or at least 99% identical to the amino acid sequence of SEQ ID NO: 280. In some examples, the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO: 376. In some examples, the antibody comprises a light chain variable region comprising the amino acid sequence of SEQ ID NO: 280.

[0016] In some examples, the antibody is a monoclonal antibody. In some examples, the antibody is of the IgG class, the IgM class, or the IgA class. In some examples, the antibody is of the IgG class and has an IgG1, IgG2, or IgG4 isotype. In some examples, the antibody has an IgG1 isotype. In some examples, the antibody comprises a E430G substitution and a P331S substitution according to EU numbering.

[0017] In some examples, the antibody is an antibody fragment. In some examples, the fragment is a Fab, Fab', Fab'-SH, F(ab')₂, Fv or scFv fragment.

[0018] In some examples, the antibody has an affinity (K_D) for human SIRPβ1 isoform 1 of 0.1 nM to 50 nM, or 0.5 nM to 10 nM, or 0.5 nM to 5 nM. In some examples, affinity is measured using a ForteBio Octet[®] system.

[0019] In some examples, the antibody recognizes a first and a second antigen, wherein the first antigen is SIRPβ1 and the second antigen is:

- (a) an antigen facilitating transport across the blood-brain-barrier;
- (b) an antigen facilitating transport across the blood-brain-barrier selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), and diphtheria toxin receptor;
- (c) a disease-causing agent selected from the group consisting of disease-causing peptides or proteins or disease-causing nucleic acids, wherein the disease-causing nucleic acids are antisense GGCCCC (G2C4) repeat-expansion RNA, the disease-causing proteins are selected from the group consisting of amyloid beta, oligomeric

amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides;

- (d) ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from the group consisting of CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA-4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, BTLA, KIR, GAL9, TIM3, A2AR, LAG-3, and phosphatidylserine; and
- (e) a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells.

[0020] In some examples, an isolated antibody that binds to human SIRPβ1 is provided, wherein the antibody competes with one or more antibodies provided herein for binding to human SIRPβ1 isoform 1. In some examples, an isolated antibody that binds to human SIRPβ1 is provided, wherein the antibody binds essentially the same or overlapping SIRPβ1 isoform 1 epitope as one or more antibodies provided herein. In one example, the present disclosure refers to an isolated antibody that binds to human SIRPβ1, wherein the antibody competes with one or more antibodies disclosed herein for binding to human SIRPβ1 isoform 1, and/or wherein the antibody binds essentially the same or overlapping SIRPβ1 isoform 1 epitope as one or more antibodies as disclosed herein.

[0021] In one aspect, the present disclosure refers to an isolated nucleic acid comprising a nucleic acid sequence encoding the antibody as disclosed herein. In some examples, an isolated nucleic acid is provided, comprising a nucleic acid sequence encoding an anti-SIRPβ1 antibody provided herein. In one aspect, the present disclosure refers to a vector comprising the nucleic acid as disclosed herein. In some examples, an isolated host cell is provided, comprising the vector. In some examples, an isolated host cell is provided that expresses an anti-SIRPβ1 antibody provided herein. In one aspect, the present disclosure refers to an isolated host cell comprising the vector as disclosed herein, wherein the host cell expresses the antibody as disclosed herein. In some examples, methods of producing an antibody that binds to human SIRPβ1 are provided, comprising culturing a host cell that expresses an anti-SIRPβ1 antibody provided herein so that the antibody is produced. In some examples, the method further comprises recovering the antibody produced by the cell. In one aspect, the present disclosure refers to a method of producing an antibody that binds to human SIRPβ1, comprising culturing the cell as disclosed herein so that the antibody is produced, optionally further comprising recovering the antibody produced by the cell.

[0022] In one aspect, the present disclosure refers to a pharmaceutical composition comprising the antibody as disclosed herein and a pharmaceutically acceptable carrier. In some examples, a pharmaceutical composition is provided, comprising an anti-SIRPβ1 antibody provided herein and a pharmaceutically acceptable carrier.

[0023] In some examples, a method of treating cancer is provided, comprising administering to an individual in need thereof a therapeutically effective amount of an anti-SIRPβ1 antibody provided herein. In some examples, the cancer is selected from selected from sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, small cell lung cancer, melanoma, lymphoma, pancreatic

cancer, prostate cancer, ovarian cancer, and fibrosarcoma, glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma, diffuse large B-cell lymphoma, lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, indolent B cell lymphoma, aggressive B cell lymphoma, T cell lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, myelodysplastic syndromes, myeloproliferative neoplasms, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma.

[0024] In some examples, a method of treatment further comprises administering a therapeutic agent that inhibits or agonizes PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73. In some examples, a method of treatment further comprises administering to the individual at least one antibody that specifically binds to an inhibitory checkpoint molecule, and/or one or more standard or investigational anti-cancer therapies.

[0025] In some examples, a method of treatment further comprises administering at least one antibody that specifically binds to an inhibitory checkpoint molecule in combination with the anti-SIRPA antibody. In some examples, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is selected from an anti-PD-L1 antibody, an anti-CTLA4 antibody, an anti-PD-L2 antibody, an anti-PD-1 antibody, an anti-B7-H3 antibody, an anti-B7-H4 antibody, and anti-HVEM antibody, an anti- B- and T-lymphocyte attenuator (BTLA) antibody, an anti-Killer inhibitory receptor (KIR) antibody, an anti-GAL9 antibody, an anti-TIM-1 antibody, an anti-TIM3 antibody, an anti-TIM-4 antibody, an anti-A2AR antibody, an anti-CD39 antibody, an anti-CD73 antibody, an anti-LAG-3 antibody, an anti-phosphatidylserine antibody, an anti-CD27 antibody, an anti-CD30 antibody, an anti-TNF α antibody, an anti-CD33 antibody, an anti-Siglec-5 antibody, an anti-Siglec-7 antibody, an anti-Siglec-9 antibody, an anti-Siglec-11 antibody, an antagonistic anti-TREM1 antibody, an

antagonistic anti-TREM2 antibody, an anti-TIGIT antibody, an anti-VISTA antibody, an anti-CD2 antibody, an anti-CD5 antibody, and any combination thereof.

[0026] In some examples, a method of treatment further comprises administering one or more standard or investigational anti-cancer therapies selected from radiotherapy, cytotoxic chemotherapy, targeted therapy, imatinib therapy, trastuzumab therapy, etanercept therapy, adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and cytokine therapy.

[0027] In some examples, a method of treatment further comprises administering to the individual at least one antibody that specifically binds to an inhibitory cytokine. In some examples, the at least one antibody that specifically binds to an inhibitory cytokine is selected from an anti-CCL2 antibody, an anti-CSF-1 antibody, an anti-IL-2 antibody, and any combination thereof.

[0028] In some examples, a method of treatment further comprises administering to the individual at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein. In some examples, the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is selected from an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof.

[0029] In some examples, a method of treatment further comprises administering to the individual at least one stimulatory cytokine. In some examples, the at least one stimulatory cytokine is selected from IFN- α 4, IFN- β , IL-1 β , TNF- α , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- γ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.

[0030] In some examples, use of an anti-SIRP β 1 antibody provided herein for the preparation of a medicament is provided. In some examples, the medicament is for treating cancer. In some examples, an anti-SIRP β 1 antibody provided herein is provided for treating cancer. In one aspect, the present disclosure refers to use of an antibody as disclosed herein in the preparation of a medicament for treating cancer in an individual, optionally wherein the cancer is selected from sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, small cell lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma, glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma, diffuse large B-cell lymphoma, lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, indolent B cell lymphoma, aggressive B cell lymphoma, T cell lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, myelodysplastic syndromes, myeloproliferative neoplasms, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma.

[0031] In some examples, a method of treating a neurodegenerative disease or disorder is provided, comprising administering to an individual in need thereof a therapeutically effective amount of an anti-SIRP β 1 antibody provided herein. In some examples, the neurodegenerative disease or disorder is selected from dementia, frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, Nasu-Hakola disease, cognitive

deficit, memory loss, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis.

[0032] In some examples, use of an anti-SIRP β 1 antibody provided herein for the preparation of a medicament for treating a neurodegenerative disease or disorder is provided. In some examples, the neurodegenerative disease or disorder is selected from dementia, frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, Nasu-Hakola disease, cognitive deficit, memory loss, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis. In some examples, an anti-SIRP β 1 antibody provided herein for use in treating a neurodegenerative disease or disorder is provided. In some examples, the neurodegenerative disease or disorder is selected from dementia, frontotemporal dementia, progressive supranuclear palsy, Alzheimer's disease, vascular dementia, Nasu-Hakola disease, cognitive deficit, memory loss, seizures, retinal dystrophy, amyotrophic lateral sclerosis, traumatic brain injury, a spinal cord injury, stroke, Parkinson's disease, acute disseminated encephalomyelitis, retinal degeneration, age related macular degeneration, glaucoma, multiple sclerosis.

[0033] It is to be understood that one, some, or all of the properties of the various examples described herein may be combined to form other examples of the present disclosure. These and other aspects of the disclosure will become apparent to one of skill in the art. These and other examples of the disclosure are further described by the detailed description that follows.

BRIEF DESCRIPTION OF THE DRAWINGS

[0034] **FIG. 1A** shows an alignment of the amino acid sequences of SIRP β 1 isoform 1 (SEQ ID NO: 1; Uniprot Accession No. O00241) and SIRP β 1 isoform 3 (SEQ ID NO: 384; Uniprot Accession No. Q5TFQ8). **FIG. 1B** shows an alignment of the amino acid sequences of SIRP β 1 isoform 1 (SEQ ID NO: 1) with SIRP α (SEQ ID NO: 385; Uniprot Accession No. P78324), which shows high homology within the extracellular domains.

[0035] **FIG. 2** shows an alignment of the amino acid sequences of human SIRP β 1 isoform 1 (SEQ ID NO: 1) and mouse SIRP β 1 (SEQ ID NO: 386; Uniprot Accession No. Q8BFX8).

[0036] **FIG. 3A-3B** shows induction of human SIRP β 1-dependent luciferase expression in a cell-based reporter assay. Cells were stimulated with plate-bound anti-SIRP β 1 antibodies (3A) or anti-SIRP α (SA-9C2) (3B) or human IgG1 isotype control. Results are expressed as fold over background. The background level is set to 1 on y-axis.

[0037] **FIG. 4A** shows the expression pattern of two DAP12-associated receptors, SIRP β 1 and TREM1, on human monocytes. Shaded histograms represent background fluorescence from isotype

stained cells. Black outlined histograms represent receptor expression level with target-specific antibody stained cells. **FIG. 4B** shows the expression of SIRP β 1 on M1 and M2 polarized macrophages from 2 healthy donors. Shaded histograms represent background fluorescence from isotype stained cells. Black outlined histograms represent SIRP β 1 expression level on M2 macrophages, whereas dashed line histograms represent SIRP β 1 expression level on M1 macrophages. **FIG. 4C** shows the expression of SIRP β 1 and TREM1 on human monocyte-derived dendritic cells (DCs) with or without LPS stimulation. Shaded histograms represent background fluorescence from isotype stained cells. Black outlined histograms represent receptor expression level with target-specific antibody stained cells.

[0038] **FIG. 5A** profiles the RNA expression pattern of SIRP β 1 isoform 1 in various tissues from human donors. **FIG. 5B** profiles the RNA expression pattern of SIRP β 1 isoform 3 in various tissues from human donors. Peak expression for SIRP β 1 isoform 3 occurs in the brain with limited expression in peripheral tissues.

[0039] **FIG. 6A** shows SIRP β 1-mediated respiratory burst from primary human neutrophils obtained from two different donors. **FIG. 6B** shows SIRP β 1-mediated respiratory burst (top panel) or IL-8 release (bottom panel) from primary human monocytes. **FIG. 6C** shows SIRP β 1-mediated TNF α cytokine release from primary human macrophages or dendritic cells (DCs).

[0040] **FIG. 7** shows the anti-tumor activity of primary human neutrophils stimulated with plate-bound anti-SIRP β 1 antibodies. Luminescence values are presented on a relative scale with the signal of Raji-Luc co-cultured with neutrophils in the absence of opsonizing antibody set to 1 on the y-axis.

[0041] **FIG. 8** shows induction of human SIRP β 1-dependent luciferase expression in a cell-based reporter assay by affinity matured antibodies. Results presented are raw luminescence values. The background level is set to 10,000 on y-axis.

[0042] **FIG. 9** shows SIRP β 1-mediated TNF α cytokine release from primary human dendritic cells (DCs).

[0043] **FIG. 10A-10B** shows the cross-reactivity of anti-SIRP β 1 antibodies to multiple SIRP β 1 antigens. Soluble, human SIRP β 1-Fc isoform 1 (O00241) or isoform 3 (Q5TFQ8) or cynomolgus SIRP β 1-Fc (XM005568541) were coated onto plates and incubated with increasing concentrations of parental and affinity matured forms of anti-SIRP β 1 antibodies SB-1, SB-1-2, SB-2, SB-2-7, SB-8, SB-8-13, SB-40, and SB-40-21. All anti-SIRP β 1 antibodies bound only to human SIRP β 1-Fc isoform 1 antigen.

[0044] **FIG. 11A** shows the upregulation of TREM2 expression on monocyte-derived human macrophages from three different healthy donors by plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3, SB-2-8, SB-8-13, SB-8-15, and SB-40-20. Baseline TREM2 expression was determined from cells incubated on human IgG1 isotype control antibody. TREM2 was

detected with an anti-TREM2 antibody (clone ADI-22) conjugated with DyLight650 fluorophore. **FIG. 11B** shows the viability of monocyte-derived human macrophages cultured on plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3, SB-2-8, SB-8-13, SB-8-15, and SB-40-20. Viable cells were quantified by measuring luminescence values following addition of Cell Titer Glo substrate. Cells cultured on human IgG1 isotype control or in the absence of plate-bound antibody (No Ab) established baseline viability of cells.

[0045] **FIG. 12** shows the viability of monocyte-derived macrophages from two different healthy donors cultured on increasing concentrations of plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3 and SB-2-8 or human IgG1 isotype control. Viable cells were quantified by measuring luminescence values following addition of Cell Titer Glo substrate.

[0046] **FIG. 13** shows anti-SIRP β 1 antibodies enhance the agonistic activity of anti-TREM2 antibody in a macrophage viability assay. Monocyte-derived macrophages were cultured on plate-bound, full-length SB-1-3, SB-2-8, or huIgG1 isotype control in the presence or absence of soluble anti-TREM2 antibody. Viable cells were quantified by measuring luminescence values following addition of Cell Titer Glo substrate.

[0047] **FIG. 14A** shows the viability of bone marrow-derived macrophages obtained from human SIRP β 1 BAC transgenic mice cultured on plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3, SB-2-8, and SB-8-13 or human IgG1 isotype control. **FIG. 14B** shows the viability of bone marrow-derived dendritic cells obtained from human SIRP β 1 BAC transgenic mice cultured on plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3, SB-2-8, and SB-8-13 or human IgG1 isotype control. Viable cells were quantified by measuring luminescence values following addition of Cell Titer Glo substrate. Cells cultured on human IgG1 isotype control or in the absence of plate-bound antibody (No Ab) established baseline viability of cells.

[0048] **FIG. 15A-15C** show cross-reactivity of anti-SIRP β 1 antibodies to different receptors of the SIRP family. **FIG. 15A** and **FIG. 15B** show titration curves of various anti-SIRP receptor antibodies binding BW5147.G.1.4 cells overexpressing recombinant human SIRP β 1 or recombinant human SIRP α , respectively. **FIG. 15C** shows titration curves of antibodies binding Jurkat cells, an immortalized human T cell line that endogenously expresses SIRP γ . Positive control antibodies were anti-SIRP α/β clone 18D5, anti-SIRP $\alpha/\beta/\gamma$ clone KWAR23, and anti-SIRP γ clone LSB2.20.

DETAILED DESCRIPTION

[0049] The present disclosure relates to anti-SIRP β 1 antibodies (*e.g.*, monoclonal antibodies); methods of making and using such antibodies; pharmaceutical compositions comprising such antibodies; nucleic acids encoding such antibodies; and host cells comprising nucleic acids encoding such antibodies.

[0050] The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized methodologies such as those described in Sambrook *et al. Molecular Cloning: A Laboratory Manual* 3d edition (2001) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *Current Protocols in Molecular Biology* (F.M. Ausubel, *et al.* eds., (2003); *Monoclonal Antibodies: A Practical Approach* (P. Shepherd and C. Dean, eds., Oxford University Press, 2000).

[0051] All references cited herein, including patent applications and publications, are hereby incorporated by reference in their entirety.

I. Definitions

[0052] The terms “Signal-Regulatory Protein $\beta 1$,” “SIRP $\beta 1$,” and “SIRP $\beta 1$ polypeptide” are used interchangeably herein refer herein to any native SIRP $\beta 1$ from any vertebrate source, including mammals such as primates (*e.g.*, humans and cynos) and rodents (*e.g.*, mice and rats), unless otherwise indicated. In some embodiments, the term encompasses both wild-type sequences and naturally occurring variant sequences, *e.g.*, splice variants or allelic variants. In some embodiments, the term encompasses “full-length,” unprocessed SIRP $\beta 1$ as well as any form of SIRP $\beta 1$ that results from processing in the cell. In some embodiments, the SIRP $\beta 1$ is human SIRP $\beta 1$. In some embodiments, the amino acid sequence of an exemplary SIRP $\beta 1$ isoform 1 is Uniprot Accession No. O00241 as of February 28, 2018. In some embodiments, the amino acid sequence of an exemplary human SIRP $\beta 1$ is SEQ ID NO: 1. In some embodiments, the amino acid sequence of an exemplary SIRP $\beta 1$ isoform 3 is Uniprot Accession No. Q5TFQ8 as of January 31, 2018. In some embodiments, the amino acid sequence of an exemplary human SIRP $\beta 1$ isoform 3 is SEQ ID NO: 384. Unless specifically indicated otherwise, “SIRP $\beta 1$ ” as used herein refers to SIRP $\beta 1$ isoform 1.

[0053] The terms “anti- SIRP $\beta 1$ antibody,” an “antibody that binds to SIRP $\beta 1$,” and “antibody that specifically binds SIRP $\beta 1$ ” refer to an antibody that is capable of binding SIRP $\beta 1$ with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting SIRP $\beta 1$. In one embodiment, the extent of binding of an anti- SIRP $\beta 1$ antibody to an unrelated, non- SIRP $\beta 1$ polypeptide is less than about 10% of the binding of the antibody to SIRP $\beta 1$ as measured, *e.g.*, by a radioimmunoassay (RIA). In certain embodiments, an antibody that binds to SIRP $\beta 1$ has a dissociation constant (KD) of $< 1 \mu\text{M}$, $< 100 \text{ nM}$, $< 10 \text{ nM}$, $< 1 \text{ nM}$, $< 0.1 \text{ nM}$, $< 0.01 \text{ nM}$, or $< 0.001 \text{ nM}$ (*e.g.*, 10^{-8} M or less, *e.g.* from 10^{-8} M to 10^{-13} M , *e.g.*, from 10^{-9} M to 10^{-13} M). In certain embodiments, an anti-SIRP $\beta 1$ antibody binds to an epitope of SIRP $\beta 1$ that is conserved among SIRP $\beta 1$ polypeptides from different species.

[0054] With regard to the binding of an antibody to a target molecule, the term “specific binding” or “specifically binds” or is “specific for” a particular polypeptide or an epitope on a particular polypeptide target means binding that is measurably different from a non-specific interaction.

Specific binding can be measured, for example, by determining binding of a molecule compared to binding of a control molecule. For example, specific binding can be determined by competition with a control molecule that is similar to the target, for example, an excess of non-labeled target. In this case, specific binding is indicated if the binding of the labeled target to a probe is competitively inhibited by excess unlabeled target. The term “specific binding” or “specifically binds to” or is “specific for” a particular polypeptide or an epitope on a particular polypeptide target as used herein can be exhibited, for example, by a molecule having a KD for the target of about any of 10^{-4} M or lower, 10^{-5} M or lower, 10^{-6} M or lower, 10^{-7} M or lower, 10^{-8} M or lower, 10^{-9} M or lower, 10^{-10} M or lower, 10^{-11} M or lower, 10^{-12} M or lower or a KD in the range of 10^{-4} M to 10^{-6} M or 10^{-6} M to 10^{-10} M or 10^{-7} M to 10^{-9} M. As will be appreciated by the skilled artisan, affinity and KD values are inversely related. A high affinity for an antigen is measured by a low KD value. In one embodiment, the term “specific binding” refers to binding where a molecule binds to a particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

[0055] The term “immunoglobulin” (Ig) is used interchangeably with “antibody” herein. The term “antibody” herein is used in the broadest sense and specially covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific antibodies) including those formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

[0056] “Native antibodies” are usually heterotetrameric glycoproteins of about 150,000 Daltons, composed of two identical Light (“L”) chains and two identical heavy (“H”) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intra-chain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

[0057] For the structure and properties of the different classes of antibodies, *see, e.g., Basic and Clinical Immunology*, 8th Ed., Daniel P. Stites, Abba I. Terr and Tristram G. Parslow (eds.), Appleton & Lange, Norwalk, CT, 1994, page 71 and Chapter 6.

[0058] The light chain from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (“ κ ”) and lambda (“ λ ”), based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains (CH), immunoglobulins can be assigned to different classes or isotypes. There are five classes of

immunoglobulins: IgA, IgD, IgE, IgG, and IgM, having heavy chains designated alpha ("α"), delta ("δ"), epsilon ("ε"), gamma ("γ"), and mu ("μ"), respectively. The γ and α classes are further divided into subclasses (isotypes) on the basis of relatively minor differences in the CH sequence and function, *e.g.*, humans express the following subclasses: IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known and described generally in, for example, Abbas *et al.*, Cellular and Molecular Immunology, 4th ed. (W.B. Saunders Co., 2000).

[0059] The "variable region" or "variable domain" of an antibody, such as an anti-SIRPβ1 antibody of the present disclosure, refers to the amino-terminal domains of the heavy or light chain of the antibody. The variable domains of the heavy chain and light chain may be referred to as "V_H" and "V_L", respectively. These domains are generally the most variable parts of the antibody (relative to other antibodies of the same class) and contain the antigen binding sites.

[0060] The term "variable" refers to the fact that certain segments of the variable domains differ extensively in sequence among antibodies, such as anti-SIRPβ1 antibodies of the present disclosure. The variable domain mediates antigen binding and defines the specificity of a particular antibody for its particular antigen. However, the variability is not evenly distributed across the entire span of the variable domains. Instead, it is concentrated in three segments called hypervariable regions (HVRs) both in the light-chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three HVRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The HVRs in each chain are held together in close proximity by the FR regions and, with the HVRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (*see* Kabat *et al.*, *Sequences of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in the binding of antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent-cellular toxicity.

[0061] The term "monoclonal antibody" as used herein refers to an antibody, such as a monoclonal anti-SIRPβ1 antibody of the present disclosure, obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations and/or post-translation modifications (*e.g.*, isomerizations, amidations, etc.) that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. In contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized

by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including, for example, the hybridoma method, recombinant DNA methods, and technologies for producing human or human-like antibodies in animals that have parts or all of the human immunoglobulin loci or genes encoding human immunoglobulin sequences.

[0062] The terms “full-length antibody,” “intact antibody” or “whole antibody” are used interchangeably to refer to an antibody, such as an anti- SIRPβ1 antibody of the present disclosure, in its substantially intact form, as opposed to an antibody fragment. Specifically, whole antibodies include those with heavy and light chains including an Fc region. The constant domains may be native sequence constant domains (*e.g.*, human native sequence constant domains) or amino acid sequence variants thereof. In some cases, the intact antibody may have one or more effector functions.

[0063] An “antibody fragment” refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include Fab, Fab', F(ab')₂ and Fv fragments; diabodies; linear antibodies (*see* U.S. Patent 5641870, Example 2; Zapata *et al.*, *Protein Eng.* 8(10):1057-1062 (1995)); single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[0064] Papain digestion of antibodies, such as anti-SIRPβ1 antibodies of the present disclosure, produces two identical antigen-binding fragments, called “Fab” fragments, and a residual “Fc” fragment, a designation reflecting the ability to crystallize readily. The Fab fragment consists of an entire light chain along with the variable region domain of the heavy chain (V_H), and the first constant domain of one heavy chain (C_{H1}). Each Fab fragment is monovalent with respect to antigen binding, *i.e.*, it has a single antigen-binding site. Pepsin treatment of an antibody yields a single large F(ab')₂ fragment which roughly corresponds to two disulfide linked Fab fragments having different antigen-binding activity and is still capable of cross-linking antigen. Fab' fragments differ from Fab fragments by having a few additional residues at the carboxy terminus of the C_{H1} domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0065] The Fc fragment comprises the carboxy-terminal portions of both heavy chains held together by disulfides. The effector functions of antibodies are determined by sequences in the Fc region, the region which is also recognized by Fc receptors (FcR) found on certain types of cells.

[0066] “Functional fragments” of antibodies, such as anti-SIRP β 1 antibodies of the present disclosure, comprise a portion of an intact antibody, generally including the antigen binding or variable region of the intact antibody or the Fc region of an antibody which retains or has modified FcR binding capability. Examples of antibody fragments include linear antibody, single-chain antibody molecules and multispecific antibodies formed from antibody fragments.

[0067] The term “diabodies” refers to small antibody fragments prepared by constructing sFv fragments (see preceding paragraph) with short linkers (about 5-10) residues) between the V_H and V_L domains such that inter-chain but not intra-chain pairing of the variable domains is achieved, thereby resulting in a bivalent fragment, *i.e.*, a fragment having two antigen-binding sites. Bispecific diabodies are heterodimers of two “crossover” sFv fragments in which the V_H and V_L domains of the two antibodies are present on different polypeptide chains.

[0068] As used herein, a “chimeric antibody” refers to an antibody (immunoglobulin), such as a chimeric anti-SIRP β 1 antibody of the present disclosure, in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is(are) identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity. Chimeric antibodies of interest herein include PRIMATIZED[®] antibodies wherein the antigen-binding region of the antibody is derived from an antibody produced by, *e.g.*, immunizing macaque monkeys with an antigen of interest. As used herein, “humanized antibody” is used a subset of “chimeric antibodies.”

[0069] “Humanized” forms of non-human (*e.g.*, murine) antibodies, such as humanized forms of anti-SIRP β 1 antibodies of the present disclosure, are chimeric antibodies comprising amino acid residues from non-human HVRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the HVRs (*e.g.*, CDRs) correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized antibody optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, *e.g.*, a non-human antibody, refers to an antibody that has undergone humanization.

[0070] A “human antibody” is one that possesses an amino-acid sequence corresponding to that of an antibody, such as an anti-SIRP β 1 antibody of the present disclosure, produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries and yeast-based platform technologies. Human antibodies can

be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, immunized xenomice as well as generated via a human B-cell hybridoma technology.

[0071] The term “hypervariable region” or “HVR,” when used herein refers to the regions of an antibody-variable domain, such as that of an anti-SIRPβ1 antibody of the present disclosure, that are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six HVRs; three in the V_H (H1, H2, H3), and three in the V_L (L1, L2, L3). In native antibodies, H3 and L3 display the most diversity of the six HVRs, and H3 in particular is believed to play a unique role in conferring fine specificity to antibodies. Naturally occurring camelid antibodies consisting of a heavy chain only are functional and stable in the absence of light chain.

[0072] A number of HVR delineations are in use and are encompassed herein. In some embodiments, the HVRs may be Kabat complementarity-determining regions (CDRs) based on sequence variability and are the most commonly used (Kabat *et al.*, *supra*). In some embodiments, the HVRs may be Chothia CDRs. Chothia refers instead to the location of the structural loops (Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). In some embodiments, the HVRs may be AbM HVRs. The AbM HVRs represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody-modeling software. In some embodiments, the HVRs may be “contact” HVRs. The “contact” HVRs are based on an analysis of the available complex crystal structures. The residues from each of these HVRs are noted below.

Loop	Kabat	AbM	Chothia	Contact
L1	L24-L34	L24-L34	L26-L32	L30-L36
L2	L50-L56	L50-L56	L50-L52	L46-L55
L3	L89-L97	L89-L97	L91-L96	L89-L96
H1	H31-H35B	H26-H35B	H26-H32	H30-H35B (Kabat numbering)
H1	H31-H35	H26-H35	H26-H32	H30-H35 (Chothia numbering)
H2	H50-H65	H50-H58	H53-H55	H47-H58
H3	H95-H102	H95-H102	H96-H101	H93-H101

[0073] HVRs may comprise “extended HVRs” as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2), and 89-97 or 89-96 (L3) in the V_L, and 26-35 (H1), 50-65 or 49-65 (a preferred embodiment) (H2), and 93-102, 94-102, or 95-102 (H3) in the V_H. The variable-domain residues are numbered according to Kabat *et al.*, *supra*, for each of these extended-HVR definitions.

[0074] “Framework” or “FR” residues are those variable-domain residues other than the HVR residues as herein defined.

[0075] An “acceptor human framework” as used herein is a framework comprising the amino acid sequence of a V_L or V_H framework derived from a human immunoglobulin framework or a human consensus framework. An acceptor human framework “derived from” a human immunoglobulin

framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may comprise pre-existing amino acid sequence changes. In some embodiments, the number of pre-existing amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. Where pre-existing amino acid changes are present in a V_H, preferable those changes occur at only three, two, or one of positions 71H, 73H and 78H; for instance, the amino acid residues at those positions may be 71A, 73T and/or 78A. In one embodiment, the V_L acceptor human framework is identical in sequence to the V_L human immunoglobulin framework sequence or human consensus framework sequence.

[0076] A “human consensus framework” is a framework that represents the most commonly occurring amino acid residues in a selection of human immunoglobulin V_L or V_H framework sequences. Generally, the selection of human immunoglobulin V_L or V_H sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). Examples include for the V_L, the subgroup may be subgroup kappa I, kappa II, kappa III or kappa IV as in Kabat *et al.*, *supra*. Additionally, for the V_H, the subgroup may be subgroup I, subgroup II, or subgroup III as in Kabat *et al.*, *supra*.

[0077] An “amino-acid modification” at a specified position, *e.g.*, of an anti-SIRPβ1 antibody of the present disclosure, refers to the substitution or deletion of the specified residue, or the insertion of at least one amino acid residue adjacent the specified residue. Insertion “adjacent” to a specified residue means insertion within one to two residues thereof. The insertion may be N-terminal or C-terminal to the specified residue. The preferred amino acid modification herein is a substitution.

[0078] An “affinity-matured” antibody, such as an affinity matured anti-SIRPβ1 antibody of the present disclosure, is one with one or more alterations in one or more HVRs thereof that result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody that does not possess those alteration(s). In one embodiment, an affinity-matured antibody has nanomolar or even picomolar affinities for the target antigen. Affinity-matured antibodies are produced by procedures known in the art. For example, Marks *et al.* *Bio/Technology* 10:779-783 (1992) describes affinity maturation by V_H- and V_L-domain shuffling. Random mutagenesis of HVR and/or framework residues is described by, for example: Barbas *et al.* *Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al.* *Gene* 169:147-155 (1995); Yelton *et al.* *J. Immunol.* 155: 1994-2004 (1995); Jackson *et al.* *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.* *J. Mol. Biol.* 226:889-896 (1992).

[0079] “Fv” is the minimum antibody fragment which comprises a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (3 loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody. However, even a single

variable domain (or half of an Fv comprising only three HVRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0080] “Single-chain Fv” also abbreviated as “sFv” or “scFv” are antibody fragments that comprise the V_H and V_L antibody domains connected into a single polypeptide chain. Preferably, the sFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains, which enables the sFv to form the desired structure for antigen binding.

[0081] Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody, and vary with the antibody isotype.

[0082] The term “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including native-sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. Suitable native-sequence Fc regions for use in the antibodies of the present disclosure include human IgG1, IgG2, IgG3 and IgG4.

[0083] A “native sequence Fc region” comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof.

[0084] A “variant Fc region” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

[0085] “Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors, FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (“ITAM”) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (“ITIM”) in its cytoplasmic domain. Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. FcRs can also increase the serum half-life of antibodies.

[0086] As used herein, “percent (%) amino acid sequence identity” and “homology” with respect to a peptide, polypeptide or antibody sequence refers to the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific peptide or polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or MEGALIGN™ (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms known in the art needed to achieve maximal alignment over the full-length of the sequences being compared.

[0087] The term “compete” when used in the context of antibodies (*e.g.*, neutralizing antibodies) that compete for the same epitope means competition between antibody as determined by an assay in which the antibody being tested prevents or inhibits (*e.g.*, reduces) specific binding of a reference molecule (*e.g.*, a ligand, or a reference antibody) to a common antigen (*e.g.*, SIRPβ1 or a fragment thereof). Numerous types of competitive binding assays can be used to determine if antibody competes with another, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (*see, e.g.*, Stahli *et al.*, 1983, *Methods in Enzymology* 9:242-253); solid phase direct biotin-avidin EIA (*see, e.g.*, Kirkland *et al.*, 1986, *J. Immunol.* 137:3614-3619) solid phase direct labeled assay, solid phase direct labeled sandwich assay (*see, e.g.*, Harlow and Lane, 1988, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using 1-125 label (*see, e.g.*, Morel *et al.*, 1988, *Molec. Immunol.* 25:7-15); solid phase direct biotin-avidin EIA (*see, e.g.*, Cheung, *et al.*, 1990, *Virology* 176:546-552); and direct labeled RIA (Moldenhauer *et al.*, 1990, *Scand. J. Immunol.* 32:77-82). Typically, such an assay involves the use of purified antigen bound to a solid surface or

cells bearing either of these, an unlabelled test antibody and a labeled reference antibody.

Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test antibody. Usually the test antibody is present in excess. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur. Additional details regarding methods for determining competitive binding are provided in the examples herein.

Usually, when a competing antibody is present in excess, it will inhibit (*e.g.*, reduce) specific binding of a reference antibody to a common antigen by at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97.5%, and/or near 100%.

[0088] As used herein, an “interaction” between a SIRPβ1 polypeptide and a second polypeptide encompasses, without limitation, protein-protein interaction, a physical interaction, a chemical interaction, binding, covalent binding, and ionic binding. As used herein, an antibody “inhibits interaction” between two polypeptides when the antibody disrupts, reduces, or completely eliminates an interaction between the two polypeptides. An antibody of the present disclosure, thereof, “inhibits interaction” between two polypeptides when the antibody thereof binds to one of the two polypeptides. In some embodiments, the interaction can be inhibited by at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97.5%, and/or near 100%.

[0089] The term “epitope” includes any determinant capable of being bound by an antibody. An epitope is a region of an antigen that is bound by an antibody that targets that antigen, and when the antigen is a polypeptide, includes specific amino acids that directly contact the antibody. Most often, epitopes reside on polypeptides, but in some instances, can reside on other kinds of molecules, such as nucleic acids. Epitope determinants can include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl groups, and can have specific three-dimensional structural characteristics, and/or specific charge characteristics. Generally, antibodies specific for a particular target antigen will preferentially recognize an epitope on the target antigen in a complex mixture of polypeptides and/or macromolecules.

[0090] An “agonist” antibody or an “activating” antibody is an antibody that induces (*e.g.*, increases) one or more activities or functions of the antigen after the antibody binds the antigen.

[0091] An “antagonist” antibody or a “blocking” antibody or an “inhibitory” antibody is an antibody that reduces, inhibits, and/or eliminates (*e.g.*, decreases) antigen binding to one or more ligand after the antibody binds the antigen, and/or that reduces, inhibits, and/or eliminates (*e.g.*, decreases) one or more activities or functions of the antigen after the antibody binds the antigen. In some embodiments, antagonist antibodies, or blocking antibodies, or inhibitory antibodies substantially or completely inhibit antigen binding to one or more ligand and/or one or more activities or functions of the antigen.

[0092] An “isolated” antibody, such as an isolated anti-SIRPβ1 antibody of the present disclosure, is one that has been identified, separated and/or recovered from a component of its production environment (*e.g.*, naturally or recombinantly). Preferably, the isolated antibody is free of association with all other contaminant components from its production environment. Contaminant components from its production environment, such as those resulting from recombinant transfected cells, are materials that would typically interfere with research, diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody will be purified: (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments, to greater than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant T-cells since at least one component of the antibody’s natural environment will not be present. Ordinarily, however, an isolated polypeptide or antibody will be prepared by at least one purification step.

[0093] An “isolated” nucleic acid molecule encoding an antibody, such as an anti-SIRPβ1 antibody of the present disclosure, is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the environment in which it was produced. Preferably, the isolated nucleic acid is free of association with all components associated with the production environment. The isolated nucleic acid molecules encoding the polypeptides and antibodies herein is in a form other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from nucleic acid encoding the polypeptides and antibodies herein existing naturally in cells.

[0094] The term “vector,” as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid,” which refers to a circular double stranded DNA into which additional DNA segments may be ligated. Another type of vector is a phage vector. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors,” or simply, “expression vectors.” In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present

specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector.

[0095] “Polynucleotide,” or “nucleic acid,” as used interchangeably herein, refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase or by a synthetic reaction.

[0096] A “host cell” includes an individual cell or cell culture that can be or has been a recipient for vector(s) for incorporation of polynucleotide inserts. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in genomic DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation. A host cell includes cells transfected *in vivo* with a polynucleotide(s) of this invention.

[0097] “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed.

[0098] As used herein, the term “preventing” includes providing prophylaxis with respect to occurrence or recurrence of a particular disease, disorder, or condition in an individual. An individual may be predisposed to, susceptible to a particular disease, disorder, or condition, or at risk of developing such a disease, disorder, or condition, but has not yet been diagnosed with the disease, disorder, or condition.

[0099] As used herein, an individual “at risk” of developing a particular disease, disorder, or condition may or may not have detectable disease or symptoms of disease, and may or may not have displayed detectable disease or symptoms of disease prior to the treatment methods described herein. “At risk” denotes that an individual has one or more risk factors, which are measurable parameters that correlate with development of a particular disease, disorder, or condition, as known in the art. An individual having one or more of these risk factors has a higher probability of developing a particular disease, disorder, or condition than an individual without one or more of these risk factors.

[0100] As used herein, the term “treatment” refers to clinical intervention designed to alter the natural course of the individual being treated during the course of clinical pathology. Desirable effects of treatment include decreasing the rate of progression, ameliorating or palliating the pathological state, and remission or improved prognosis of a particular disease, disorder, or condition. An individual is successfully “treated”, for example, if one or more symptoms associated with a particular disease, disorder, or condition are mitigated or eliminated.

[0101] An “effective amount” refers to at least an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result. An effective amount can be provided in one or more administrations. An effective amount herein may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the treatment to

elicit a desired response in the individual. An effective amount is also one in which any toxic or detrimental effects of the treatment are outweighed by the therapeutically beneficial effects. For prophylactic use, beneficial or desired results include results such as eliminating or reducing the risk, lessening the severity, or delaying the onset of the disease, including biochemical, histological and/or behavioral symptoms of the disease, its complications and intermediate pathological phenotypes presenting during development of the disease. For therapeutic use, beneficial or desired results include clinical results such as decreasing one or more symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications required to treat the disease, enhancing effect of another medication such as via targeting, delaying the progression of the disease, and/or prolonging survival. An effective amount of drug, compound, or pharmaceutical composition is an amount sufficient to accomplish prophylactic or therapeutic treatment either directly or indirectly. As is understood in the clinical context, an effective amount of a drug, compound, or pharmaceutical composition may or may not be achieved in conjunction with another drug, compound, or pharmaceutical composition. Thus, an “effective amount” may be considered in the context of administering one or more therapeutic agents, and a single agent may be considered to be given in an effective amount if, in conjunction with one or more other agents, a desirable result may be or is achieved.

[00102] An “individual” for purposes of treatment, prevention, or reduction of risk refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sport, or pet animals, such as dogs, horses, rabbits, cattle, pigs, hamsters, gerbils, mice, ferrets, rats, cats, and the like. In some embodiments, the individual is human.

[00103] As used herein, administration “in conjunction” with another compound or composition includes simultaneous administration and/or administration at different times. Administration in conjunction also encompasses administration as a co-formulation or administration as separate compositions, including at different dosing frequencies or intervals, and using the same route of administration or different routes of administration. In some embodiments, administration in conjunction is administration as a part of the same treatment regimen.

[00104] The term “about” as used herein refers to the usual error range for the respective value readily known to the skilled person in this technical field. Reference to “*about*” a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter *per se*.

[00105] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly indicates otherwise. For example, reference to an “antibody” is a reference to from one to many antibodies, such as molar amounts, and includes equivalents thereof known to those skilled in the art, and so forth.

[00106] It is understood that aspect and embodiments of the present disclosure described herein include “comprising,” “consisting,” and “consisting essentially of” aspects and embodiments.

I. Anti-SIRPβ1 Antibodies

[00107] Provided herein are anti-SIRPβ1 antibodies. Antibodies provided are useful, *e.g.*, for the diagnosis or treatment of the SIRPβ1 mediated disorders.

[00108] In some embodiments, an anti-SIRPβ1 antibody of the present disclosure is an agonist of SIRPβ1 activity. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1 but does not bind to human SIRPα. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1 but does not bind to human SIRPγ. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1 but does not bind to human SIRPβ1 isoform 3. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1 but does not bind to mouse SIRPβ1 and/or cynomolgus monkey SIRPβ1. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure agonizes SIRPβ1 activity on CD14-positive monocytes. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces respiratory burst in immune cells, such as neutrophils, monocytes, and/or macrophages. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces or increases IL-8 expression in monocytes. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces or increases TNFα expression in macrophages and/or dendritic cells. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces neutrophil-mediated phagocytosis, for example, of tumor cells. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure increases neutrophil-mediated tumor cell clearance. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure increases or enhances anti-tumor properties or activity of neutrophils.

[00109] In some embodiments, an anti-SIRPβ1 antibody of the present disclosure recruits immune cells. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces syk phosphorylation. In some embodiments, an anti-SIRPβ1 antibody of the present disclosure induces syk phosphorylation when clustered by adjacent cells expressing Fc gamma receptors.

[00110] In some embodiments, SIRPβ1 antibody of the present disclosure downregulates SIRPβ1 expression on the surface of a cell. In some embodiments, SIRPβ1 antibody of the present disclosure does not downregulate SIRPβ1 expression on the surface of a cell. In some embodiments, SIRPβ1 antibody of the present disclosure blocks binding of a ligand to SIRPβ1. In some embodiments, SIRPβ1 antibody of the present disclosure does not block binding of a ligand to SIRPβ1.

[00111] In some embodiments, an anti-SIRPβ1 antibody is provided that has one or more properties selected from:

- a) binds to human SIRPβ1 isoform 1, but does not bind to human SIRPα;
- b) binds to human SIRPβ1 isoform 1, but does not bind to human SIRPγ;
- c) binds to human SIRPβ1 isoform 1, but does not bind to human SIRPβ1 isoform 3;

- d) binds to human SIRP β 1 isoform 1, but does not bind to mouse SIRP β 1;
- e) binds to human SIRP β 1 isoform 1, but does not bind to cynomolgus monkey SIRP β 1;
- f) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
- g) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
- h) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
- i) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
- j) induces or increases neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
- k) induces or increases neutrophil-mediated tumor cell clearance *in vivo*;
- l) upregulates TREM2 expression on macrophages *in vitro* and/or *in vivo*;
- m) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
- n) increases viability of dendritic cells *in vitro* and/or *in vivo*.

[00112] In some embodiments, an anti-SIRP β 1 antibody is provided that has the following properties:

- a) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α ;
- b) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP γ ;
- c) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP β 1 isoform 3;
- d) binds to human SIRP β 1 isoform 1, but does not bind to mouse SIRP β 1;
- e) binds to human SIRP β 1 isoform 1, but does not bind to cynomolgus monkey SIRP β 1;
- f) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
- g) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
- h) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
- i) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
- j) induces or increases neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
- o) induces or increases neutrophil-mediated tumor cell clearance *in vivo*;
- p) upregulates TREM2 expression on macrophages *in vitro* and/or *in vivo*;
- q) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
- k) increases viability of dendritic cells *in vitro* and/or *in vivo*.

A. Exemplary Antibodies and Certain Other Antibody Embodiments

[00113] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (b) HVR-H2 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (c) HVR-H3 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (d) HVR-L1 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8; (e) HVR-L2 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8; and (f) HVR-L3 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8.

[00114] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising HVR-H1, HVR-H2, HVR-H3, HVR-L1, HVR-L2, and HVR-L3, wherein: (a) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (b) HVR-H2 comprising an amino acid sequence of an HVR-H2 shown in Table 4 and/or Table 9; (c) HVR-H3 comprising an amino acid sequence of an HVR-H3 shown in Table 4 and/or Table 9; (d) HVR-L1 comprising an amino acid sequence of an HVR-L1 shown in Table 3 and/or Table 8; (e) HVR-L2 comprising an amino acid sequence of an HVR-L2 shown in Table 3 and/or Table 8; and (f) HVR-L3 comprising an amino acid sequence of an HVR-L3 shown in Table 3 and/or Table 8.

[00115] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising six HVRs of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9).

[00116] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising six HVRs of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9).

[00117] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising six HVRs of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 8 and 9).

[00118] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising six HVRs of an antibody selected from SB-1-3, SB-2-8, and SB-8-13 (shown in Tables 8 and 9).

[00119] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 228; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 238; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[00120] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 239; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[00121] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 240; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[00122] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 241; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[00123] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 230; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 242; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00124] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 231; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 243; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00125] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 232; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 244; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00126] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 99; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 245; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00127] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 230; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 242; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 253; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00128] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 246; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00129] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 234; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 247; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00130] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 248; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00131] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 246; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 254; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00132] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 235; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 249; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 163; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 2; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 22; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 69.

[00133] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 236; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 250; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 163; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 2; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 22; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 69.

[00134] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 237; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 251; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 163; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 2; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 22; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 69.

[00135] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 236; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 252; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 163; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 2; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 22; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 69.

[00136] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising at least one, at least two, or all three V_H HVR sequences selected from (a) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (b) HVR-H2 comprising an amino acid sequence of an HVR-H2 shown in Table 4 and/or Table 9; (c) HVR-H3 comprising an amino acid sequence of an HVR-H3 shown in Table 4 and/or Table 9. In some embodiments, an anti-SIRP β 1 antibody comprises HVR-H1, HVR, H2, and HVR-H3 of an antibody selected from SB-1, SB-2,

SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 4 and 9). In some embodiments, an anti-SIRP β 1 antibody comprises HVR-H1, HVR, H2, and HVR-H3 of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 4 and 9). In some embodiments, an anti-SIRP β 1 antibody comprises HVR-H1, HVR, H2, and HVR-H3 of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Table 9). In some embodiments, an anti-SIRP β 1 antibody comprises HVR-H1, HVR, H2, and HVR-H3 of an antibody selected from SB-1-3, SB-2-8, and SB-8-13 (shown in Table 9). In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 239; and (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125. In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 231; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 243; and (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126. In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 246; and (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131.

[00137] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising at least one, at least two, or all three V_L HVR sequences selected from (a) HVR-L1 comprising an amino acid sequence of an HVR-L1 shown in Table 3 and/or Table 8; (b) HVR-L2 comprising an amino acid sequence of an HVR-L2 shown in Table 3 and/or Table 8; (c) HVR-L3 comprising an amino acid sequence of an HVR-L3 shown in Table 3 and/or Table 8. In some embodiments, an anti-SIRP β 1 antibody comprises HVR-L1, HVR, L2, and HVR-L3 of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3 and 8). In some embodiments, an anti-SIRP β 1 antibody

comprises HVR-L1, HVR, L2, and HVR-L3 of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3 and 8). In some embodiments, an anti-SIRP β 1 antibody comprises HVR-L1, HVR, L2, and HVR-L3 of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Table 8). In some embodiments, an anti-SIRP β 1 antibody comprises HVR-L1, HVR, L2, and HVR-L3 of an antibody selected from SB-1-3, SB-2-8, and SB-8-13 (shown in Table 8). In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (b) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31. In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (b) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32. In some embodiments, an anti-SIRP β 1 antibody comprises (a) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (b) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00138] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) a V_H domain comprising at least one, at least two, or all three V_H HVR sequences selected from (i) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9, (ii) HVR-H2 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9, and (iii) HVR-H3 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; and (b) a V_L domain comprising at least one, at least two, or all three V_L HVR sequences selected from (i) HVR-L1 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8, (ii) HVR-L2 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8, and (iii) HVR-L3 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8.

[00139] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) a V_H domain comprising (i) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9, (ii) HVR-H2 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9, and (iii) HVR-H3 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; and (b) a V_L domain comprising (i) HVR-L1 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8, (ii) HVR-L2 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8, and (iii) HVR-L3 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8.

[00140] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising a V_H domain comprising HVR-H1, HVR-H2, and HVR-H3, and a V_L domain comprising HVR-L1, HVR-L2, and

HVR-L3, wherein HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 are the HVRs of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9).

[00141] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising a V_H domain comprising HVR-H1, HVR-H2, and HVR-H2, and a V_L domain comprising HVR-L1, HVR-L2, and HVR-L3, wherein HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 are the HVRs of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 3, 4, 8, and 9).

[00142] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising a V_H domain comprising HVR-H1, HVR-H2, and HVR-H2, and a V_L domain comprising HVR-L1, HVR-L2, and HVR-L3, wherein HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 are the HVRs of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21 (shown in Tables 8 and 9).

[00143] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising a V_H domain comprising HVR-H1, HVR-H2, and HVR-H2, and a V_L domain comprising HVR-L1, HVR-L2, and HVR-L3, wherein HVR-H1, HVR-H2, HVR-H2, HVR-L1, HVR-L2, and HVR-L3 are the HVRs of an antibody selected from SB-1-3, SB-2-8, and SB-8-13 (shown in Tables 8 and 9).

[00144] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) a V_H domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 229; (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 239; (iii) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 125; and (b) a V_L domain comprising (iv) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (v) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (vi) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 31.

[00145] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) a V_H domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 231; (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 243; (iii) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 126; and (b) a V_L domain comprising (iv) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (v) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 16; and (vi) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 32.

[00146] In some embodiments, provided herein are anti-SIRP β 1 antibodies comprising (a) a V_H domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 233; (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 246; (iii) HVR-H3 comprising the amino acid sequence of SEQ ID NO: 131; and (b) a V_L domain comprising (iv) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 383; (v) HVR-L2 comprising the amino acid sequence of SEQ ID NO: 19; and (vi) HVR-L3 comprising the amino acid sequence of SEQ ID NO: 37.

[00147] In another aspect, an anti-SIRP β 1 antibody comprises a heavy chain variable domain (V_H) sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382. In certain embodiments, a V_H sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence of SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, or 365 contains substitutions (*e.g.*, conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-SIRP β 1 antibody comprising that sequence retains the ability to bind to SIRP β 1. In certain embodiments, a total of 1 to 10 amino acids have been substituted, inserted, and/or deleted in SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382. In certain embodiments, substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FRs). Optionally, the anti-SIRP β 1 antibody comprises the V_H sequence of SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, or 382, including post-translational modifications of that sequence. In a particular embodiment, the V_H comprises one, two or three HVRs selected from: (a) HVR-H1 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (b)

HVR-H2 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9; (c) HVR-H3 comprising an amino acid sequence of an HVR-H1 shown in Table 4 and/or Table 9.

[00148] In another aspect, an anti-SIRP β 1 antibody is provided, wherein the antibody comprises a light chain variable domain (V_L) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370. In certain embodiments, a V_L sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence of SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370 contains substitutions (*e.g.*, conservative substitutions), insertions, or deletions relative to the reference sequence, but an anti-SIRP β 1 antibody comprising that sequence retains the ability to bind to SIRP β 1. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370. In certain embodiments, a total of 1 to 5 amino acids have been substituted, inserted and/or deleted in SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370. In certain embodiments, the substitutions, insertions, or deletions occur in regions outside the HVRs (*i.e.*, in the FRs). Optionally, the anti-SIRP β 1 antibody comprises the V_L sequence of SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, or 370, including post-translational modifications of that sequence. In a particular embodiment, the V_L comprises one, two or three HVRs selected from (a) HVR-L1 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8; (b) HVR-L2 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8; and (c) HVR-L3 comprising an amino acid sequence of an HVR-H1 shown in Table 3 and/or Table 8.

[00149] In some embodiments, the anti-SIRP β 1 antibody is provided, wherein the antibody comprises a V_H as in any of the embodiments provided above, and a V_L as in any of the embodiments provided above. In some embodiments, provided herein are anti-SIRP β 1 antibodies, wherein the antibody comprises a V_H as in any of the embodiments provided above, and a V_L as in any of the embodiments provided above. In one embodiment, the antibody comprises a V_H sequence

selected from SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, and 382; and V_L sequence selected from SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, and 370, including post-translational modifications of those sequences. In some embodiments, the antibody comprises the V_H sequence and V_L sequence of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, the antibody comprises the V_H sequence and V_L sequence of an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, the antibody comprises the V_H sequence and V_L sequence of an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, the antibody comprises the V_H sequence and V_L sequence of an antibody selected from SB-1-3, SB-2-8, and SB-8-13. In some embodiments, the antibody comprises the V_H sequence of SEQ ID NO: 367 and V_L sequence of SEQ ID NO: 267. In some embodiments, the antibody comprises the V_H sequence of SEQ ID NO: 372 and V_L sequence of SEQ ID NO: 370. In some embodiments, the antibody comprises the V_H sequence of SEQ ID NO: 376 and V_L sequence of SEQ ID NO: 280.

[00150] In some embodiments, an anti-SIRPβ1 antibody competes for binding with an antibody comprising a V_H sequence selected from SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, and 382, and a V_L sequence selected from SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, and 370. In some embodiments, an anti-SIRPβ1 antibody competes for binding with an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14,

SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody selected from SB-1-3, SB-2-8, and SB-8-13. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody comprising the V_H sequence of SEQ ID NO: 367 and V_L sequence of SEQ ID NO: 267. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody comprising the V_H sequence of SEQ ID NO: 372 and V_L sequence of SEQ ID NO: 370. In some embodiments, an anti-SIRP β 1 antibody competes for binding with an antibody comprising the V_H sequence of SEQ ID NO: 376 and V_L sequence of SEQ ID NO: 280.

[00151] In some embodiments, the antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody comprising a V_H sequence selected from SEQ ID NO: 268, 270, 272, 274, 276, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345, 347, 349, 351, 353, 355, 357, 359, 361, 363, 365, 366, 367, 368, 369, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, and 382, and a V_L sequence selected from SEQ ID NO: 267, 269, 271, 273, 275, 277, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346, 348, 350, 352, 354, 356, 358, 360, 362, 364, and 370. In some embodiments, the antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody selected from SB-1, SB-2, SB-3, SB-4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-10, SB-11, SB-12, SB-13, SB-14, SB-15, SB-16, SB-17, SB-18, SB-19, SB-20, SB-21, SB-22, SB-23, SB-24, SB-25, SB-26, SB-27, SB-28, SB-29, SB-30, SB-31, SB-32, SB-33, SB-34, SB-35, SB-36, SB-37, SB-38, SB-39, SB-40, SB-41, SB-42, SB-43, SB-44, SB-45, SB-46, SB-47, SB-48, SB-49, SB-50, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21.

[00152] In some embodiments, the antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody selected from SB-1, SB-2, SB-3, SB-

4, SB-5, SB-6, SB-7, SB-8, SB-9, SB-14, SB-28, SB-39, SB-40, SB-49, SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, the antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody selected from SB-1-2, SB-1-3, SB-1-4, SB-1-5, SB-2-7, SB-2-8, SB-2-9, SB-2-10, SB-2-11, SB-8-13, SB-8-14, SB-8-15, SB-8-16, SB-40-18, SB-40-19, SB-40-20, and SB-40-21. In some embodiments, the antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody selected from SB-1-3, SB-2-8, and SB-8-13. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody comprising the V_H sequence of SEQ ID NO: 367 and V_L sequence of SEQ ID NO: 267. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody comprising the V_H sequence of SEQ ID NO: 372 and V_L sequence of SEQ ID NO: 370. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope of human SIRP β 1 that is the same as or overlaps with the epitope bound by an anti-SIRP β 1 antibody comprising the V_H sequence of SEQ ID NO: 376 and V_L sequence of SEQ ID NO: 280. In some embodiments, the epitope of human SIRP β 1 is the same epitope as bound by an anti-SIRP β 1 antibody.

[00153] In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 30 to 148 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 30 to 136 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 30 to 80 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 40 to 90 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 50 to 100 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 60 to 110 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 70 to 120 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 80 to 130 of SEQ ID NO: 1. In some embodiments, an anti-SIRP β 1 antibody binds to an epitope within amino acids 90 to 140 of SEQ ID NO: 1.

[00154] In some embodiments, the anti-SIRP β 1 antibody according to any of the above embodiments is a monoclonal antibody, including a humanized and/or human antibody. In some embodiments, the anti-SIRP β 1 antibody is an antibody fragment, *e.g.*, a Fv, Fab, Fab', scFv, diabody, or F(ab')₂ fragment. In some embodiments, the anti-SIRP β 1 antibody is a substantially full-length antibody, *e.g.*, an IgG1 antibody, IgG2a antibody or other antibody class or isotype as defined herein.

[00155] In some embodiments, an anti-SIRP β 1 antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described in Sections 1-7 below:

(1) *Anti-SIRPβ1 antibody binding affinity*

[00156] In some embodiments of any of the antibodies provided herein, the antibody has a dissociation constant (K_D) of $< 1 \mu\text{M}$, $< 100 \text{ nM}$, $< 10 \text{ nM}$, $< 1 \text{ nM}$, $< 0.1 \text{ nM}$, $< 0.01 \text{ nM}$, or $< 0.001 \text{ nM}$ (*e.g.*, 10^{-8} M or less, *e.g.*, from 10^{-8} M to 10^{-13} M , *e.g.*, from 10^{-9} M to 10^{-13} M). Dissociation constants may be determined through any analytical technique, including any biochemical or biophysical technique such as ELISA, surface plasmon resonance (SPR), bio-layer interferometry (*see, e.g.*, Octet System by ForteBio), isothermal titration calorimetry (ITC), differential scanning calorimetry (DSC), circular dichroism (CD), stopped-flow analysis, and colorimetric or fluorescent protein melting analyses. In one embodiment, K_D is measured by a radiolabeled antigen binding assay (RIA). In some embodiments, an RIA is performed with the Fab version of an antibody of interest and its antigen, for example as described in Chen *et al. J. Mol. Biol.* 293:865-881(1999)). In some embodiments, K_D is measured using a BIACORE surface plasmon resonance assay, for example, an assay using a BIACORE -2000 or a BIACORE -3000 (BIAcore, Inc., Piscataway, NJ) is performed at 25°C with immobilized antigen CM5 chips at ~ 10 response units (RU). In some embodiments, K_D is measured using a ForteBio Octet[®] Red384 system (ForteBio, Menlo Park, CA), for example, as discussed in the examples herein.

[00157] In some embodiments, the K_D is determined using a monovalent antibody (*e.g.*, a Fab) or a full-length antibody. In some embodiments, the K_D is determined using a full-length antibody in a monovalent form.

(2) *Antibody fragments*

[00158] In some embodiments of any of the antibodies provided herein, the antibody is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, *see* Hudson *et al. Nat. Med.* 9:129-134 (2003). For a review of scFv fragments, *see, e.g.*, WO 93/16185; and U.S. Patent Nos. 5571894 and 5587458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased *in vivo* half-life, *see* U.S. Patent No. 5869046.

[00159] Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. *See, for example*, EP404097; WO 1993/01161; Hudson *et al. Nat. Med.* 9:129-134 (2003). Triabodies and tetrabodies are also described in Hudson *et al. Nat. Med.* 9:129-134 (2003). Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (*see, e.g.*, U.S. Patent No. 6248516).

[00160] Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (*e.g.*, *E. coli* or phage), as described herein.

(3) *Chimeric and Humanized antibodies*

[00161] In some embodiments of any of the antibodies provided herein, the antibody is a chimeric antibody. Certain chimeric antibodies are described, *e.g.*, in U.S. Patent No. 4816567. In one example, a chimeric antibody comprises a non-human variable region (*e.g.*, a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a “class switched” antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-binding fragments thereof.

[00162] In some embodiments of any of the antibodies provided herein, the antibody is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. In certain embodiments, a humanized antibody is substantially non-immunogenic in humans. In certain embodiments, a humanized antibody has substantially the same affinity for a target as an antibody from another species from which the humanized antibody is derived. *See, e.g.*, U.S. Pat. No. 5530101, 5693761; 5693762; and 5585089. In certain embodiments, amino acids of an antibody variable domain that can be modified without diminishing the native affinity of the antigen binding domain while reducing its immunogenicity are identified. *See, e.g.*, U.S. Pat. Nos. 5766886 and 5869619. Generally, a humanized antibody comprises one or more variable domains in which HVRs (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (*e.g.*, the antibody from which the HVR residues are derived), for example, to restore or improve antibody specificity or affinity.

[00163] Humanized antibodies and methods of making them are reviewed, for example, in Almagro *et al. Front. Biosci.* 13:1619-1633 (2008), and are further described, *e.g.*, in US Patent Nos. 5821337, 7527791, 6982321, and 7087409. Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the “best-fit” method (*see, e.g.*, Sims *et al. J. Immunol.* 151:2296 (1993)); framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (*see, e.g.*, Carter *et al. Proc. Natl. Acad. Sci. USA* 89:4285 (1992); and Presta *et al., J. Immunol.* 151:2623 (1993)); human mature (somatically mutated) framework regions or human germline framework regions (*see, e.g.*, Almagro and Fransson *Front. Biosci.* 13:1619-1633 (2008)); and

framework regions derived from screening FR libraries (*see, e.g., Baca et al. J. Biol. Chem.* 272:10678-10684 (1997) and Rosok *et al. J. Biol. Chem.* 271:22611-22618 (1996)).

(4) *Human Antibodies*

[00164] In some embodiments of any of the antibodies provided herein, the antibody is a human antibody. Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk *et al. Curr. Opin. Pharmacol.* 5:368-74 (2001) and Lonberg *Curr. Opin. Immunol.* 20:450-459 (2008).

[00165] Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. One can engineer mouse strains deficient in mouse antibody production with large fragments of the human Ig loci in anticipation that such mice would produce human antibodies in the absence of mouse antibodies. Large human Ig fragments can preserve the large variable gene diversity as well as the proper regulation of antibody production and expression. By exploiting the mouse machinery for antibody diversification and selection and the lack of immunological tolerance to human proteins, the reproduced human antibody repertoire in these mouse strains can yield high affinity fully human antibodies against any antigen of interest, including human antigens. Using the hybridoma technology, antigen-specific human MAbs with the desired specificity can be produced and selected. Certain exemplary methods are described in U.S. Pat. No. 5545807, EP 546073, and EP 546073. *See also*, for example, U.S. Patent Nos. 6075181 and 6150584 describing XENOMOUSE™ technology; U.S. Patent No. 5770429 describing HUMAB® technology; U.S. Patent No. 7041870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VELOCIMOUSE® technology. Human variable regions from intact antibodies generated by such animals may be further modified, *e.g.*, by combining with a different human constant region.

[00166] Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described. (*See, e.g., Kozbor J. Immunol.* 133:3001 (1984) and Boerner *et al. J. Immunol.* 147:86 (1991)). Human antibodies generated via human B-cell hybridoma technology are also described in Li *et al. Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Patent No. 7189826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines). Human hybridoma technology (Trioma technology) is also described in Vollmers *et al. Histology and Histopathology* 20(3):927-937 (2005) and Vollmers *et al. Methods and Findings in Experimental and Clinical Pharmacology* 27(3):185-91 (2005). Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain

sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

[0100] In some embodiments of any of the antibodies provided herein, the antibody is a human antibody isolated by *in vitro* methods and/or screening combinatorial libraries for antibodies with the desired activity or activities. Suitable examples include but are not limited to phage display (CAT, Morphosys, Dyax, Biosite/Medarex, Xoma, Symphogen, Alexion (formerly Proliferon), Affimed) ribosome display (CAT), yeast-based platform technology (Adimab), and the like. In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter *et al. Ann. Rev. Immunol.* 12: 433-455 (1994). For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. *See also* Sidhu *et al. J. Mol. Biol.* 338(2): 299-310, 2004; Lee *et al. J. Mol. Biol.* 340(5): 1073-1093, 2004; Fellouse *Proc. Natl. Acad. Sci. USA* 101(34):12467-12472 (2004); and Lee *et al. J. Immunol. Methods* 284(-2):1 19-132 (2004). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (*e.g.*, from human) to provide a single source of antibodies to a wide range of non-self and also self-antigens without any immunization as described by Griffiths *et al. EMBO J.* 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers comprising random sequence to encode the highly variable HVR3 regions and to accomplish rearrangement *in vitro*, as described by Hoogenboom *et al. J. Mol. Biol.*, 227: 381-388, 1992. Patent publications describing human antibody phage libraries include, for example: US Patent No. 5750373, and US Patent Publication Nos. 2007/0292936 and 2009/0002360. Antibodies isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

(5) *Constant Regions including Fc regions*

[0101] In some embodiments of any of the antibodies provided herein, the antibody comprises an Fc. In some embodiments, the Fc is a human IgG1, IgG2, IgG3, and/or IgG4 isotype. In some embodiments, the antibody is of the IgG class, the IgM class, or the IgA class.

[0102] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG2 isotype. In some embodiments, the antibody contains a human IgG2 constant region. In some embodiments, the human IgG2 constant region includes an Fc region. In some embodiments, the antibody induces the one or more SIRPβ1 activities or independently of binding to an Fc receptor. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0103] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG1 isotype. In some embodiments, the antibody contains a mouse IgG1 constant region. In some embodiments, the antibody contains a human IgG1 constant region. In some embodiments, the human IgG1 constant region includes an Fc region. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0104] In certain embodiments of any of the antibodies provided herein, the antibody has an IgG4 isotype. In some embodiments, the antibody contains a human IgG4 constant region. In some embodiments, the human IgG4 constant region includes an Fc region. In some embodiments, the antibody binds an inhibitory Fc receptor. In certain embodiments, the inhibitory Fc receptor is inhibitory Fc-gamma receptor IIB (FcγIIB).

[0105] In certain embodiments of any of the antibodies provided herein, the antibody has a hybrid IgG2/4 isotype. In some embodiments, the antibody includes an amino acid sequence comprising amino acids 118 to 260 according to EU numbering of human IgG2 and amino acids 261-447 according to EU numbering of human IgG4 (WO 1997/11971; WO 2007/106585).

[0106] In some embodiments, the Fc region increases clustering without activating complement as compared to a corresponding antibody comprising an Fc region that does not comprise the amino acid substitutions. In some embodiments, the antibody induces one or more activities of a target specifically bound by the antibody. In some embodiments, the antibody binds to SIRPβ1.

[0107] It may also be desirable to modify an anti-SIRPβ1 antibody of the present disclosure to modify effector function and/or to increase serum half-life of the antibody. For example, the Fc receptor binding site on the constant region may be modified or mutated to remove or reduce binding affinity to certain Fc receptors, such as FcγRI, FcγRII, and/or FcγRIII to reduce Antibody-dependent cell-mediated cytotoxicity. In some embodiments, the effector function is impaired by removing N-glycosylation of the Fc region (*e.g.*, in the CH2 domain of IgG) of the antibody. In some embodiments, the effector function is impaired by modifying regions such as 233-236, 297, and/or 327-331 of human IgG as described in WO 99/58572 and Armour *et al. Molecular Immunology* 40: 585-593 (2003); Reddy *et al. J. Immunology* 164:1925-1933 (2000). In other embodiments, it may also be desirable to modify an anti-SIRPβ1 antibody of the present disclosure to modify effector function to increase finding selectivity toward the ITIM-containing FcγRIIb (CD32b) to increase clustering of SIRPβ1 antibodies on adjacent cells without activating humoral responses including Antibody-dependent cell-mediated cytotoxicity and antibody-dependent cellular phagocytosis.

[0108] To increase the serum half-life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5739277, for example. As used herein, the term “salvage receptor binding epitope” refers to an

epitope of the Fc region of an IgG molecule (*e.g.*, IgG₁, IgG₂, IgG₃, or IgG₄) that is responsible for increasing the *in vivo* serum half-life of the IgG molecule. Other amino acid sequence modifications.

(6) *Multispecific Antibodies*

[0109] Multispecific are antibodies that have binding specificities for at least two different epitopes, including those on the same or another polypeptide (*e.g.*, one or more SIRPβ1 polypeptides of the present disclosure). In some embodiments, the multispecific antibody is a bispecific antibody. In some embodiments, the multispecific antibody is a trispecific antibody. In some embodiments, the multispecific antibody is a tetraspecific antibody. Such antibodies can be derived from full-length antibodies or antibody fragments (*e.g.*, F(ab')₂ bispecific antibodies). In some embodiments, the multispecific antibody comprises a first antigen binding region which binds to first site on SIRPβ1 and comprises a second antigen binding region which binds to a second site on SIRPβ1. In some embodiment, the multispecific antibodies comprises a first antigen binding region which binds to SIRPβ1 and a second antigen binding region that binds to a second polypeptide.

[0110] Provided herein are multispecific antibodies comprises a first antigen binding region, wherein the first antigen binding region comprises the six HVRs of an antibody described herein, which binds to SIRPβ1 and a second antigen binding region that binds to a second polypeptide. In some embodiments, the first antigen binding region comprises the V_H or V_L of an antibody described herein.

[0111] In some embodiments of any of the multispecific antibodies, the second polypeptide is an antigen facilitating transport across the blood-brain-barrier. In some embodiments, and antibody herein is conjugated to a peptide that facilitates transport across the blood-brain barrier. Numerous antigens and peptides are known in the art that facilitate transport across the blood-brain barrier (*see, e.g.*, Gabathuler R. *Neurobiol. Dis.* 37:48-57 (2010)). Such second antigens and peptides include, without limitation, transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), diphtheria toxin receptor, including CRM197 (a non-toxic mutant of diphtheria toxin), TMEM 30(A) (Flippase), protein transduction domains such as TAT, Syn-B, or penetratin, poly-arginine or generally positively charged peptides, Angiopep peptides such as ANG1005 (*see, e.g.*, Gabathuler, 2010), and other cell surface proteins that are enriched on blood-brain barrier endothelial cells (*see, e.g.*, Daneman *et al.* *PLoS One* 5(10):e13741 (2010)). In some embodiments, the second polypeptide is transferrin.

[0112] In some embodiments of any of the multispecific antibodies, the second polypeptide is a disease-causing protein selected from amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10,

Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides. In some embodiments, the second polypeptide is Tau. In some embodiments, the second polypeptide is A β . In some embodiments, the second polypeptide is TREM2. In some embodiments, the second polypeptide is α -synuclein.

[0113] In some embodiments of any of the multispecific antibodies, the second polypeptide is a ligand and/or protein expressed on immune cells, wherein the ligand and/or protein is selected from CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA-4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, BTLA, KIR, GAL9, TIM3, A2AR, LAG-3, and phosphatidylserine.

[0114] In some embodiments of any of the multispecific antibodies, the second polypeptide is a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells and any combination thereof.

[0115] In some embodiments of any of the multispecific antibodies, the second polypeptide is an immunoglobulin-like receptor, such as TREM2. In some embodiments of any of the multispecific antibodies, the second polypeptide is an immunoglobulin-like receptor expressed on myeloid lineage cells.

[0116] The multivalent antibody contains at least one polypeptide chain (and preferably two polypeptide chains), wherein the polypeptide chain or chains comprise two or more variable domains. For instance, the polypeptide chain or chains may comprise VD1-(X1)_n-VD2-(X2)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, Fc is one polypeptide chain of an Fc region, X1 and X2 represent an amino acid or polypeptide, and n is 0 or 1. Similarly, the polypeptide chain or chains may comprise V_H-C_{H1}-flexible linker-V_H-C_{H1}-Fc region chain; or V_H-C_{H1}-V_H-C_{H1}-Fc region chain. The multivalent antibody herein preferably further comprises at least two (and preferably four) light chain variable domain polypeptides. The multivalent antibody herein may, for instance, comprise from about two to about eight light chain variable domain polypeptides. The light chain variable domain polypeptides contemplated here comprise a light chain variable domain and, optionally, further comprise a CL domain.

[0117] Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain- light chain pairs having different specificities (see Milstein and Cuello *Nature* 305: 537 (1983), WO 93/08829, and Traunecker *et al. EMBO J.* 10:3655 (1991)), and “knob-in-hole” engineering (see, e.g., U.S. Patent No. 5731168). See also WO 2013/026833 (CrossMab). Multi-specific antibodies may also be made by engineering electrostatic

steering effects for making antibody Fc- heterodimeric molecules (WO 2009/089004A1); cross-linking two or more antibodies (*see, e.g.*, US Patent No. 4676980); using leucine; using “diabody” technology for making bispecific antibody fragments (*see, e.g.*, Hollinger *et al. Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993)); and using single-chain Fv (scFv) dimers (*see, e.g.*, Gruber *et al. J. Immunol.* 152:5368 (1994)); and preparing trispecific antibodies as described, *e.g.*, in Tutt *et al. J. Immunol.* 147: 60 (1991).

[0118] Engineered antibodies with three or more functional antigen binding sites, including “Octopus antibodies,” are also included herein (*see, e.g.*, US 2006/0025576). The antibody herein also includes a “Dual Acting FAb” or “DAF” comprising an antigen binding site that binds to multiple SIRPβ1 (*see*, US 2008/0069820, for example).

(7) *Antibody Variants*

[0119] In some embodiments of any of the antibodies provided herein, amino acid sequence variants of the antibodies are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody.

(i) *Substitution, Insertion, and Deletion Variants*

[0120] In some embodiments of any of the antibodies provided herein, antibody variants having one or more amino acid substitutions are provided. Amino acid sequence variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within the amino acid sequences of the antibody.

TABLE 1: Amino Acid Substitutions

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu

Original Residue	Exemplary Substitutions	Preferred Substitutions
Phe (F)	Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

[0121] Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;
- (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;
- (3) acidic: Asp, Glu;
- (4) basic: His, Lys, Arg;
- (5) residues that influence chain orientation: Gly, Pro; and
- (6) aromatic: Trp, Tyr, Phe.

[0122] For example, non-conservative substitutions can involve the exchange of a member of one of these classes for a member from another class. Such substituted residues can be introduced, for example, into regions of a human antibody that are homologous with non-human antibodies, or into the non-homologous regions of the molecule.

[0123] In making changes to the polypeptide or antibody described herein, according to certain embodiments, the hydropathic index of amino acids can be considered. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (−0.4); threonine (−0.7); serine (−0.8); tryptophan (−0.9); tyrosine (−1.3); proline (−1.6); histidine (−3.2); glutamate (−3.5); glutamine (−3.5); aspartate (−3.5); asparagine (−3.5); lysine (−3.9); and arginine (−4.5).

[0124] The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is understood in the art. Kyte *et al. J. Mol. Biol.*, 157:105-131 (1982). It is known that certain amino acids can be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, in certain embodiments, the substitution of amino acids whose hydropathic

indices are within ± 2 is included. In certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included.

[0125] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, *i.e.*, with a biological property of the protein.

[0126] The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0 $\pm$ 1); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (−0.4); proline (−0.5 $\pm$ 1); alanine (−0.5); histidine (−0.5); cysteine (−1.0); methionine (−1.3); valine (−1.5); leucine (−1.8); isoleucine (−1.8); tyrosine (−2.3); phenylalanine (−2.5) and tryptophan (−3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in certain embodiments, those which are within ± 1 are included, and in certain embodiments, those within ± 0.5 are included. One can also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as “epitopic core regions”.

[0127] In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (*e.g.*, conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in HVRs. Such alterations may, for example, be outside of antigen contacting residues in the HVRs. In certain embodiments of the variant VH and VL sequences provided above, each HVR either is unaltered, or contains no more than one, two or three amino acid substitutions.

[0128] Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides comprising a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (*e.g.*, for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

[0129] Any cysteine residue not involved in maintaining the proper conformation of the antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment, such as an Fv fragment).

(ii) *Glycosylation variants*

[0130] In some embodiments of any of the antibodies provided herein, the antibody is altered to increase or decrease the extent to which the antibody is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

[0131] Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0132] Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

[0133] Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 according to Kabat numbering of the CH2 domain of the Fc region. The oligosaccharide may include various carbohydrates, for example, mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the “stem” of the biantennary oligosaccharide structure. In some embodiments, modifications of the oligosaccharide in an antibody of the invention may be made in order to create antibody variants with certain improved properties.

[0134] In one embodiment, antibody variants are provided having a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. *See, e.g.*, US Patent Publication Nos. 2003/0157108 and 2004/0093621. Examples of publications related to “defucosylated” or “fucose-deficient” antibody variants include: US 2003/0157108; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; Okazaki *et al.* *J. Mol. Biol.* 336:1239-1249 (2004); Yamane-Ohnuki *et al.* *Biotech. Bioeng.* 87:614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Led 3 CHO cells deficient in protein fucosylation (Ripka *et al.* *Arch. Biochem. Biophys.* 249:533-545 (1986); US 2003/0157108), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (*see, e.g.*, Yamane-Ohnuki *et al.* *Biotech. Bioeng.* 87: 614 (2004) and Kanda *et al.* *Biotechnol. Bioeng.* 94(4):680-688 (2006)).

(iii) *Modified Constant regions*

[0135] In some embodiments of any of the antibodies provided herein, the antibody Fc is an antibody Fc isotype and/or modification. In some embodiments, the antibody Fc isotype and/or modification is capable of binding to Fc gamma receptor.

[0136] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG1 modified Fc. In some embodiments, the IgG1 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG1 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments, the one or more amino acid substitutions are selected from N297A (Bolt S *et al.* (1993) *Eur J Immunol* 23:403-411), D265A (Shields *et al.* (2001) *R. J. Biol. Chem.* 276, 6591–6604), L234A, L235A (Hutchins *et al.* (1995) *Proc Natl Acad Sci USA*, 92:11980-11984; Alegre *et al.*, (1994) *Transplantation* 57:1537-1543. 31; Xu *et al.*, (2000) *Cell Immunol*, 200:16-26), G237A (Alegre *et al.* (1994) *Transplantation* 57:1537-1543. 31; Xu *et al.* (2000) *Cell Immunol*, 200:16-26), C226S, C229S, E233P, L234V, L234F, L235E (McEarchern *et al.*, (2007) *Blood*, 109:1185-1192), P331S (Sazinsky *et al.*, (2008) *Proc Natl Acad Sci USA* 2008, 105:20167-20172), S267E, L328F, A330L, M252Y, S254T, and/or T256E, where the amino acid position is according to the EU numbering convention.

[0137] In some embodiments of any of the IgG1 modified Fc, the Fc comprises N297A mutation according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises D265A and N297A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises D270A mutations according to EU numbering. In some embodiments, the IgG1 modified Fc comprises L234A and L235A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234A and G237A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234A, L235A and G237A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises one or more (including all) of P238D, L328E, E233, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises one or more of S267E/L328F mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, L328E, E233D, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, L328E, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, S267E, L328E, E233D, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises P238D, S267E, L328E, G237D, H268D, P271G and A330R mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises C226S, C229S,

E233P, L234V, and L235A mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises L234F, L235E, and P331S mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises S267E and L328F mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises S267E mutations according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the Fc comprises a substitute of the constant heavy 1 (CH1) and hinge region of IgG1 with CH1 and hinge region of IgG2 (amino acids 118-230 of IgG2 according to EU numbering) with a Kappa light chain.

[0138] In some embodiments of any of the IgG1 modified Fc, the Fc includes two or more amino acid substitutions that increase antibody clustering without activating complement as compared to a corresponding antibody having an Fc region that does not include the two or more amino acid substitutions. Accordingly, in some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc is an antibody comprising an Fc region, where the antibody comprises an amino acid substitution at position E430G and one or more amino acid substitutions in the Fc region at a residue position selected from: L234F, L235A, L235E, S267E, K322A, L328F, A330S, P331S, and any combination thereof according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, A330S, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, A330S, and P331S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, and A330S according to EU numbering. In some embodiments, the IgG1 modified Fc comprises an amino acid substitution at positions E430G, K322A, and P331S according to EU numbering.

[0139] In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise herein may be combined with an A330L mutation (Lazar *et al. Proc Natl Acad Sci USA*, 103:4005-4010 (2006)), or one or more of L234F, L235E, and/or P331S mutations (Sazinsky *et al. Proc Natl Acad Sci USA*, 105:20167-20172 (2008)), according to the EU numbering convention, to eliminate complement activation. In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more of A330L, A330S, L234F, L235E, and/or P331S according to EU numbering. In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (*e.g.*, one or more (including all) of M252Y, S254T, and T256E mutations according to the

EU numbering convention). In some embodiments of any of the IgG1 modified Fc, the IgG1 modified Fc may further comprise one or more of E430G, E430S, E430F, E430T, E345K, E345Q, E345R, E345Y, S440Y, and/or S440W according to EU numbering.

[0140] Other aspects of the present disclosure relate to antibodies having modified constant regions (*i.e.*, Fc regions). An antibody dependent on binding to Fc γ R receptor to activate targeted receptors may lose its agonist activity if engineered to eliminate Fc γ R binding (*see, e.g.*, Wilson *et al. Cancer Cell* 19:101-113 (2011); Armour *et al. Immunology* 40:585-593 (2003); and White *et al. Cancer Cell* 27:138-148 (2015)). As such, it is thought that an anti- SIRP β 1 antibody of the present disclosure with the correct epitope specificity can activate the target antigen, with minimal adverse effects, when the antibody has an Fc domain from a human IgG2 isotype (CH1 and hinge region) or another type of Fc domain that is capable of preferentially binding the inhibitory Fc γ RIIB receptors, or a variation thereof.

[0141] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG2 modified Fc. In some embodiments, the IgG2 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG2 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments of any of the IgG2 modified Fc, the one or more amino acid substitutions are selected from V234A (Alegre *et al. Transplantation* 57:1537-1543 (1994); Xu *et al. Cell Immunol*, 200:16-26 (2000)); G237A (Cole *et al. Transplantation*, 68:563-571 (1999)); H268Q, V309L, A330S, P331S (US 2007/0148167; Armour *et al. Eur J Immunol* 29: 2613-2624 (1999); Armour *et al. The Haematology Journal* 1(Suppl.1):27 (2000); Armour *et al. The Haematology Journal* 1(Suppl.1):27 (2000)), C219S, and/or C220S (White *et al. Cancer Cell* 27, 138-148 (2015)); S267E, L328F (Chu *et al. Mol Immunol*, 45:3926-3933 (2008)); and M252Y, S254T, and/or T256E according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions V234A and G237A according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions C219S or C220S according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions A330S and P331S according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering.

[0142] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C127S amino acid substitution according to the EU numbering convention (White *et al.*, (2015) *Cancer Cell* 27, 138-148; Lightle *et al. Protein Sci.* 19:753-762 (2010); and WO 2008/079246). In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention

(White *et al. Cancer Cell* 27:138-148 (2015); Lightle *et al. Protein Sci.* 19:753-762 (2010); and WO 2008/079246).

[0143] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C220S amino acid substitution according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention.

[0144] In some embodiments of any of the IgG2 modified Fc, the Fc comprises a C219S amino acid substitution according to the EU numbering convention. In some embodiments of any of the IgG2 modified Fc, the antibody has an IgG2 isotype with a Kappa light chain constant domain that comprises a C214S amino acid substitution according to the EU numbering convention.

[0145] In some embodiments of any of the IgG2 modified Fc, the Fc includes an IgG2 isotype heavy chain constant domain 1(CH1) and hinge region (White *et al. Cancer Cell* 27:138-148 (2015)). In certain embodiments of any of the IgG2 modified Fc, the IgG2 isotype CH1 and hinge region comprise the amino acid sequence of 118-230 according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the antibody Fc region comprises a S267E amino acid substitution, a L328F amino acid substitution, or both, and/or a N297A or N297Q amino acid substitution according to the EU numbering convention.

[0146] In some embodiments of any of the IgG2 modified Fc, the Fc further comprises one or more amino acid substitution at positions E430G, E430S, E430F, E430T, E345K, E345Q, E345R, E345Y, S440Y, and S440W according to EU numbering. In some embodiments of any of the IgG2 modified Fc, the Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (*e.g.*, one or more (including all) of M252Y, S254T, and T256E mutations according to the EU numbering convention). In some embodiments of any of the IgG2 modified Fc, the Fc may further comprise A330S and P331S.

[0147] In some embodiments of any of the IgG2 modified Fc, the Fc is an IgG2/4 hybrid Fc. In some embodiments, the IgG2/4 hybrid Fc comprises IgG2 aa 118 to 260 and IgG4 aa 261 to 447. In some embodiments of any IgG2 modified Fc, the Fc comprises one or more amino acid substitutions at positions H268Q, V309L, A330S, and P331S according to EU numbering.

[0148] In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises one or more additional amino acid substitutions selected from A330L, L234F, L235E, or P331S according to EU numbering; and any combination thereof.

[0149] In certain embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises one or more amino acid substitutions at a residue position selected from C127S, L234A, L234F, L235A, L235E, S267E, K322A, L328F, A330S, P331S, E345R, E430G, S440Y, and any combination thereof according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S

according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, A330S, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, A330S, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, and A330S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E430G, K322A, and P331S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at position C127S according to EU numbering. In some embodiments of any of the IgG1 and/or IgG2 modified Fc, the Fc comprises an amino acid substitution at positions E345R, E430G and S440Y according to EU numbering.

[0150] In some embodiments of any of the antibodies provided herein, the modified antibody Fc is an IgG4 modified Fc. In some embodiments, the IgG4 modified Fc comprises one or more modifications. For example, in some embodiments, the IgG4 modified Fc comprises one or more amino acid substitutions (*e.g.*, relative to a wild-type Fc region of the same isotype). In some embodiments of any of the IgG4 modified Fc, the one or more amino acid substitutions are selected from L235A, G237A, S229P, L236E (Reddy *et al. J Immunol* 164:1925-1933(2000)), S267E, E318A, L328F, M252Y, S254T, and/or T256E according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the Fc may further comprise L235A, G237A, and E318A according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the Fc may further comprise S228P and L235E according to the EU numbering convention. In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc may further comprise S267E and L328F according to the EU numbering convention.

[0151] In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc comprises may be combined with an S228P mutation according to the EU numbering convention (Angal *et al. Mol Immunol.* 30:105-108 (1993)) and/or with one or more mutations described in (Peters *et al. J Biol Chem.* 287(29):24525-33 (2012)) to enhance antibody stabilization.

[0152] In some embodiments of any of the IgG4 modified Fc, the IgG4 modified Fc may further comprise one or more mutations to enhance the antibody half-life in human serum (*e.g.*, one or more

(including all) of M252Y, S254T, and T256E mutations according to the EU numbering convention).

[0153] In some embodiments of any of the IgG4 modified Fc, the Fc comprises L235E according to EU numbering. In certain embodiments of any of the IgG4 modified Fc, the Fc comprises one or more amino acid substitutions at a residue position selected from C127S, F234A, L235A, L235E, S267E, K322A, L328F, E345R, E430G, S440Y, and any combination thereof, according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G, L243A, L235A, and P331S according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G and P331S according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at position E430 according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc region comprises an amino acid substitution at positions E430G and K322A according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions S267E and L328F according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at position C127S according to EU numbering. In some embodiments of any of the IgG4 modified Fc, the Fc comprises an amino acid substitution at positions E345R, E430G and S440Y according to EU numbering.

(8) *Other antibody modifications*

[0154] In some embodiments of any of the antibodies, the antibody is a derivative. The term “derivative” refers to a molecule that includes a chemical modification other than an insertion, deletion, or substitution of amino acids (or nucleic acids). In certain embodiments, derivatives comprise covalent modifications, including, but not limited to, chemical bonding with polymers, lipids, or other organic or inorganic moieties. In certain embodiments, a chemically modified antigen binding protein can have a greater circulating half-life than an antigen binding protein that is not chemically modified. In certain embodiments, a chemically modified antigen binding protein can have improved targeting capacity for desired cells, tissues, and/or organs. In some embodiments, a derivative antigen binding protein is covalently modified to include one or more water soluble polymer attachments, including, but not limited to, polyethylene glycol, polyoxyethylene glycol, or polypropylene glycol. *See, e.g.*, U.S. Pat. Nos. 4640835, 4496689, 4301144, 4670417, 4791192 and 4179337. In certain embodiments, a derivative antigen binding protein comprises one or more polymer, including, but not limited to, monomethoxy-polyethylene glycol, dextran, cellulose, , copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids

(either homopolymers or random copolymers), poly-(N-vinyl pyrrolidone)-polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (*e.g.*, glycerol) and polyvinyl alcohol, as well as mixtures of such polymers.

[0155] In certain embodiments, a derivative is covalently modified with polyethylene glycol (PEG) subunits. In certain embodiments, one or more water-soluble polymer is bonded at one or more specific position, for example at the amino terminus, of a derivative. In certain embodiments, one or more water-soluble polymer is randomly attached to one or more side chains of a derivative. In certain embodiments, PEG is used to improve the therapeutic capacity for an antigen binding protein. In certain embodiments, PEG is used to improve the therapeutic capacity for a humanized antibody. Certain such methods are discussed, for example, in U.S. Pat. No. 6133426, which is hereby incorporated by reference for any purpose.

[0156] Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed “peptide mimetics” or “peptidomimetics.” Fauchere, *J. Adv. Drug Res.*, 15:29 (1986); and Evans *et al. J. Med. Chem.*, 30:1229 (1987), which are incorporated herein by reference for any purpose. Such compounds are often developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides can be used to produce a similar therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (*i.e.*, a polypeptide that has a biochemical property or pharmacological activity), such as human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from: -CH₂NH-, -CH₂S-, -CH₂-CH₂-, -CH=CH-(*cis* and *trans*), -COCH₂-, -CH(OH)CH₂-, and -CH₂SO-, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) can be used in certain embodiments to generate more stable peptides. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation can be generated by methods known in the art (Rizo and Gierasch *Ann. Rev. Biochem.*, 61:387 (1992), incorporated herein by reference for any purpose); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0157] Drug conjugation involves coupling of a biological active cytotoxic (anticancer) payload or drug to an antibody that specifically targets a certain tumor marker (*e.g.* a polypeptide that, ideally, is only to be found in or on tumor cells). Antibodies track these proteins down in the body and attach themselves to the surface of cancer cells. The biochemical reaction between the antibody and the target protein (antigen) triggers a signal in the tumor cell, which then absorbs or internalizes the antibody together with the cytotoxin. After the ADC is internalized, the cytotoxic drug is released

and kills the cancer. Due to this targeting, ideally the drug has lower side effects and gives a wider therapeutic window than other chemotherapeutic agents. Techniques to conjugate antibodies are disclosed are known in the art (*see, e.g.*, Jane de Lartigue OncLive July 5, 2012; ADC Review on antibody-drug conjugates; and Ducry *et al. Bioconjugate Chemistry* 21 (1):5-13 (2010).

II. *Antibody activities*

[0158] In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to human SIRPα. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to human SIRPγ. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to human SIRPβ1 isoform 3. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to mouse SIRPβ1. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to cynomolgus monkey SIRPβ1. In some embodiments, an anti-SIRPβ1 antibody is provided that binds to human SIRPβ1 isoform 1, but does not bind to cynomolgus monkey SIRPβ1 isoform 1.

[0159] In some embodiments, binding of an anti-SIRPβ1 antibody to an antigen may be determined using a ForteBio Octet® Red384 system (ForteBio, Menlo Park, CA), for example, at described in Example 1. Antibody binding may be determined using, for example, a Fab fragment of the antibody (for monovalent binding) or a full-length antibody such as an IgG (for bivalent binding, or avidity). An exemplary binding assay for full-length antibodies using a ForteBio Octet® Red384 system is as follows. Antibodies are loaded onto AHQ sensors. The loaded antibodies are then exposed to antigen, and the off-rate measured in assay buffer at various time intervals (such as 3 minutes). Kinetics data may then be fit using a binding model in the data analysis software provided with the system. For antibody Fab fragment affinity measurements, in some embodiments, an antigen-Fc may be loaded onto AHQ sensors and then exposed to the antibody Fab fragment. The off-rate is measured as above, and the data analyzed using the software provided with the system.

[0160] In some embodiments, an anti-SIRPβ1 antibody is provided that agonizes SIRPβ1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRPβ1 antibody is provided that agonizes SIRPβ1 activity on CD14-positive monocytes *in vitro*. In some embodiments, an anti-SIRPβ1 antibody is provided that agonizes SIRPβ1 activity on CD14-positive monocytes *in vivo*. In some embodiments, the anti-SIRPβ1 antibody agonizes SIRPβ1 activity on CD14-positive monocytes *in vitro* in an assay in which the anti-SIRPβ1 antibody is immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell. In some embodiments, an anti-SIRPβ1 antibody is provided that agonizes SIRPβ1 activity on CD14-positive monocytes *in vivo*. A nonlimiting exemplary assay for determining whether an anti-SIRPβ1 antibody agonizes SIRPβ1 activity on CD14-positive monocytes *in vitro* is described in Examples 3 and 4. For example,

isolated monocytes are starved for a period of time, such as 4 hours, and then incubated with anti-SIRP β 1 antibody, e.g., in the presence of anti-IgG antibody. Following incubation, cells are lysed and phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is measured, for example, using an anti-phosphotyrosine and/or anti-phosphoserine antibody. An increase in phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is indicative of an increase in SIRP β 1 activity (i.e., agonist of SIRP β 1 activity). As another example, monocytes may be incubated with anti-SRP β 1 antibody in the presence of accessory cells that express Fc gamma receptors, such as B cells. Following incubation, cells are lysed and phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is determined as above. An increase in phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is indicative of an increase in SIRP β 1 activity (i.e., that the anti-SIRP β 1 antibody is an agonist of SIRP β 1 activity). A nonlimiting exemplary assay to determine whether an anti-SIRP β 1 antibody agonizes SIRP β 1 activity on CD14-positive monocytes *in vivo* comprises administering the anti-SIRP β 1 antibody to a human SIRP β 1 BAC transgenic C57BL/6 mouse, and isolating CD14-positive monocytes, e.g., by FACS. The monocytes are then lysed and phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is measured as described above. An increase in phosphorylation of one or more of Syk, ERK, AKT, SIRP β 1, and/or DAP12 is indicative of an increase in SIRP β 1 activity (i.e., agonist of SIRP β 1 activity).

[0161] In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro*. In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vivo*. Nonlimiting exemplary *in vitro* assays for determining whether an anti-SIRP β 1 antibody induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes, are described in Examples 5 and 6. For example, primary neutrophils are contacted with anti-SIRP β 1 antibody and the production of reactive oxygen species (ROS) is detected using a general oxidative stress indicator, such as fluorescent dye CM-H₂DCFDA. Alternatively, the assay may be carried out using anti-SIRP β 1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell.

[0162] In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases IL-8 expression in monocytes *in vitro*. In some embodiments, an anti-SIRP β 1 antibody is provided that induces or increases IL-8 expression in monocytes *in vivo*. A nonlimiting exemplary *in vitro* assay for determining whether an anti-SIRP β 1 antibody induces or

increases IL-8 expression in monocytes is described in Example 5. For example, primary monocytes may be stimulated with anti-SIRPβ1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell, for example, overnight. The supernatant may be collected to assay for IL-8 release. A nonlimiting exemplary assay to determine whether an anti-SIRPβ1 antibody induces or increases IL-8 expression in monocytes *in vivo* comprises administering the anti-SIRPβ1 antibody to human SIRPβ1 BAC transgenic C57BL/6 mice and obtaining one or more blood samples. Serum concentration of IL-8 may be determined using a commercial assay, such as the DuoSet ELISA kit (R&D Systems).

[0163] In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases TNFα expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases TNFα expression in macrophages and/or dendritic cells *in vitro*. In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases TNFα expression in macrophages and/or dendritic cells *in vivo*. A nonlimiting exemplary *in vitro* assay for determining whether an anti-SIRPβ1 antibody induces or increases TNFα expression is described in Example 6. For example, monocyte-derived macrophages or dendritic cells may be stimulated with LPS in the presence of anti-SIRPβ1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell, for example, overnight. The supernatant may be collected to assay for TNFα release. A nonlimiting exemplary assay to determine whether an anti-SIRPβ1 antibody induces or increases TNFα expression in macrophages and/or dendritic cells *in vivo* comprises administering the anti-SIRPβ1 antibody to human SIRPβ1 BAC transgenic C57BL/6 mice along with LPS (such as 5 μg of LPS/mouse). Mice are then sacrificed, e.g., by CO₂ asphyxiation and peritoneal fluid is recovered and clarified by centrifugation. TNF-α concentration may be determined using a commercial assay, such as the DuoSet ELISA kit (R&D Systems).

[0164] In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro*. In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases neutrophil-mediated phagocytosis, for example, of tumor cells *in vivo*. In some embodiments, an anti-SIRPβ1 antibody is provided that induces or increases neutrophil-mediated tumor cell clearance *in vivo*. A nonlimiting exemplary *in vitro* assay for determining whether an anti-SIRPβ1 antibody induces or increases neutrophil-mediated phagocytosis is described in Example 7. For example, primary neutrophils are contacted with anti-SIRPβ1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell. Cancer cells, such as Raji

B cell lymphoma cells engineered to express luciferase, are co-cultured with the neutrophils and immobilized anti-SIRP β 1 antibody in the presence of opsonizing antibody, such as anti-CD20 antibody. Viable Raji cells are quantified by measuring luciferase activity. A reduction in viable Raji cells in the presence of anti-SIRP β 1 antibody compared to an IgG control antibody indicates that the anti-SIRP β 1 antibody induces or increases neutrophil-mediated phagocytosis. A nonlimiting exemplary assay to determine whether an anti-SIRP β 1 antibody induces or increases neutrophil-mediated tumor cell clearance *in vivo* comprises injecting intravenously GFP-expressing B16F10 melanoma cells into human SIRP β 1 BAC transgenic mice. Mice are then treated with the anti-SIRP β 1 antibody with an opsonizing anti-gp75 antibody. Peripheral blood is collected, red blood cells are lysed, and white blood cells resuspended in FACS buffer. Neutrophils are stained with anti-CD11b and anti-Ly6G antibodies and analyzed by flow cytometry for the acquisition of green fluorescence signal from ingested tumor cells. An increase in green fluorescence signal in the neutrophils compared to the same experiment with an isotype-matched control antibody indicates that anti-SIRP β 1 antibody induces or increases neutrophil-mediated phagocytosis.

[0165] A further nonlimiting exemplary assay to determine whether an anti-SIRP β 1 antibody induces or increases neutrophil-mediated tumor cell clearance *in vivo* uses a lung metastasis model, e.g., as follows. Fc receptor γ -chain deficient C57BL/6 mice, which lack expression of activating Fc γ Rs, are injected intravenously with B16F10 melanoma cells. Bone marrow neutrophils isolated from human SIRP β 1 BAC transgenic mice are then injected intravenously with the anti-SIRP β 1 antibody in combination with an opsonizing anti-gp75 antibody. After a period of time, mice are euthanized and lungs are harvested and fixed. The number of metastatic tumor nodules with dark pigmentation are visually counted. A reduction in the number of metastatic tumor nodules compared to the same experiment with an isotype-matched control antibody indicates that the anti-SIRP β 1 antibody induces or increases neutrophil-mediated tumor cell clearance *in vivo*.

[0166] In some embodiments, an anti-SIRP β 1 antibody is provided that increases TREM2 expression on macrophages *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRP β 1 antibody is provided that increases TREM2 expression on macrophages *in vitro*. In some embodiments, an anti-SIRP β 1 antibody is provided that increases TREM2 expression on macrophages *in vivo*. A nonlimiting exemplary *in vitro* assay for determining whether an anti-SIRP β 1 antibody increases TREM2 expression on macrophages is described in Example 16. For example, monocyte-derived macrophages differentiated in culture with M-CSF are incubated with anti-SIRP β 1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell. TREM2 expression on the macrophages is analyzed, e.g., by FACS analysis. A nonlimiting exemplary assay to determine whether an anti-SIRP β 1 antibody increases TREM2 expression on macrophages *in vivo* is as follows. Human SIRP β 1 BAC transgenic C57BL/6 mice are challenged with aged sterile thioglycolate broth injected

into the peritoneum. Following challenge, mice are treated with the anti-SIRP β 1 antibody. Mice are then sacrificed, and peritoneal macrophages are harvested after euthanasia, e.g., by rinsing the peritoneal cavity with PBS. Macrophages are defined as CD11b⁺ Ly6C⁻ F4/80⁺ cells, and may be analyzed by FACS to determine TREM2 expression levels.

[0167] In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody. In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of macrophages *in vitro*, alone and/or in combination with an agonist anti-TREM2 antibody. In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of macrophages *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody. In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of dendritic cells *in vitro* and/or *in vivo*. In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of dendritic cells *in vitro*. In some embodiments, an anti-SIRP β 1 antibody is provided that increases viability of dendritic cells *in vivo*. Nonlimiting exemplary *in vitro* assays for determining whether an anti-SIRP β 1 antibody increases viability of macrophages *in vitro*, alone or in combination with an agonist anti-TREM2 antibody, are described in Examples 17, 18, and 19. For example, monocyte-derived macrophages differentiated in culture with M-CSF are incubated with anti-SIRP β 1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell, and with or without anti-TREM2 antibody. Cell viability is measured, for example, using Cell Titer Glo kit (Promega), a reagent that produces a luminescence signal relative to ATP concentration in the sample. A similar assay may be used to determine whether an anti-SIRP β 1 antibody increases the viability of bone marrow-derived macrophages or dendritic cells *in vitro*. For this assay, bone marrow cells are cultured with M-CSF to differentiate macrophages, or with GM-CSF to differentiate dendritic cells. The macrophages or dendritic cells are incubated with anti-SIRP β 1 antibody immobilized on a solid support, such as a culture plate, or bound by a secondary anti-IgG antibody, or bound by Fc gamma receptor on an accessory cell, and with or without anti-TREM2 antibody. As above, cell viability is measured, for example, using Cell Titer Glo kit (Promega), a reagent that produces a luminescence signal relative to ATP concentration in the sample. A nonlimiting exemplary assay to determine whether an anti-SIRP β 1 antibody increases viability of macrophages *in vivo* is as follows. Prolonged treatment with a blocking antibody to CSF1R depletes peritoneal and tissue resident F4/80⁺ macrophages due to the abrogation of M-CSF-mediated survival signal. For this assay, human SIRP β 1 BAC transgenic C57BL/6 mice are administered anti-CSF1R antibody along with the anti-SIRP β 1 antibody. Following treatment, mice are sacrificed, e.g., by CO₂ asphyxiation, and peripheral blood is collected by cardiac puncture. Peritoneal macrophages are collected by lavage with PBS.

Macrophage populations are counted by FACS by gating on CD11b+ Ly6C- F4/80+ cells in blood and peritoneum.

III. Nucleic acids, vectors, and host cells

[0168] Anti-SIRP β 1 antibodies of the present disclosure may be produced using recombinant methods and compositions, *e.g.*, as described in U.S. Patent No. 4816567. In some embodiments, isolated nucleic acids having a nucleotide sequence encoding any of the anti-SIRP β 1 antibodies of the present disclosure are provided. Such nucleic acids may encode an amino acid sequence comprising the V_L and/or an amino acid sequence comprising the V_H of the anti-SIRP β 1 antibody (*e.g.*, the light and/or heavy chains of the antibody). In some embodiments, one or more vectors (*e.g.*, expression vectors) comprising such nucleic acids are provided. In some embodiments, a host cell comprising such nucleic acid is also provided. In some embodiments, the host cell comprises (*e.g.*, has been transduced with): (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_L of the antibody and an amino acid sequence comprising the V_H of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_L of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence comprising the V_H of the antibody. In some embodiments, the host cell is eukaryotic, *e.g.*, a Chinese Hamster Ovary (CHO) cell or lymphoid cell (*e.g.*, Y0, NS0, Sp20 cell). Host cells of the present disclosure also include, without limitation, isolated cells, *in vitro* cultured cells, and *ex vivo* cultured cells.

[0169] Methods of making an anti-SIRP β 1 antibody of the present disclosure are provided. In some embodiments, the method includes culturing a host cell of the present disclosure comprising a nucleic acid encoding the anti-SIRP β 1 antibody, under conditions suitable for expression of the antibody. In some embodiments, the antibody is subsequently recovered from the host cell (or host cell culture medium).

[0170] For recombinant production of an anti-SIRP β 1 antibody of the present disclosure, a nucleic acid encoding the anti-SIRP β 1 antibody is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody).

[0171] Suitable vectors comprising a nucleic acid sequence encoding any of the anti-SIRP β 1 antibodies of the present disclosure, or cell-surface expressed fragments or polypeptides thereof polypeptides (including antibodies) described herein include, without limitation, cloning vectors and expression vectors. Suitable cloning vectors can be constructed according to standard techniques, or may be selected from a large number of cloning vectors available in the art. While the cloning vector selected may vary according to the host cell intended to be used, useful cloning vectors generally have the ability to self-replicate, may possess a single target for a particular restriction endonuclease,

and/or may carry genes for a marker that can be used in selecting clones comprising the vector. Suitable examples include plasmids and bacterial viruses, *e.g.*, pUC18, pUC19, Bluescript (*e.g.*, pBS SK+) and its derivatives, mpl8, mpl9, pBR322, pMB9, ColE1, pCR1, RP4, phage DNAs, and shuttle vectors such as pSA3 and pAT28. These and many other cloning vectors are available from commercial vendors such as BioRad, Strategene, and Invitrogen.

[0172] Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells. For example, anti-SIRP β 1 antibodies of the present disclosure may be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria (*e.g.*, U.S. Patent Nos. 5648237, 5789199, and 5840523). After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

[0173] In addition to prokaryotes, eukaryotic microorganisms, such as filamentous fungi or yeast, are also suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been “humanized,” resulting in the production of an antibody with a partially or fully human glycosylation pattern (*e.g.*, Gerngross *Nat. Biotech.* 22:1409-1414 (2004); and Li *et al. Nat. Biotech.* 24:210-215 (2006)).

[0174] Suitable host cells for the expression of glycosylated antibody can also be derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures can also be utilized as hosts (*e.g.*, U.S. Patent Nos. 5959177, 6040498, 6420548, 7125978, and 6417429, describing PLANTIBODIES™ technology for producing antibodies in transgenic plants).

[0175] Vertebrate cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293 cells as described, *e.g.*, in Graham *et al. J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as described, *e.g.*, in Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK; buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep G2); mouse mammary tumor (MMT 060562); TRI cells, as described, *e.g.*, in Mather *et al. Annals N.Y. Acad. Sci.* 383:44-68 (1982); MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR- CHO cells (Urlaub *et al. Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); and myeloma cell lines such as Y0, NS0 and Sp2/0. For a review of certain mammalian host cell lines

suitable for antibody production, *see, e.g.*, Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003).

IV. Pharmaceutical compositions/formulations

[0176] Provided herein are pharmaceutical compositions and/or pharmaceutical formulations comprising the anti-SIRPβ1 antibodies of the present disclosure and a pharmaceutically acceptable carrier.

[0177] In some embodiments, pharmaceutically acceptable carrier preferably are nontoxic to recipients at the dosages and concentrations employed. The antibodies described herein may be formulated into preparations in solid, semi-solid, liquid or gaseous forms. Examples of such formulations include, without limitation, tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. Pharmaceutically acceptable carriers can include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers of diluents, which are vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. In certain embodiments, the pharmaceutical composition can comprise formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition.

[0178] In certain embodiments, pharmaceutically acceptable carriers include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. Further examples of formulations that are suitable for various types of administration can be found in *Remington: The Science and Practice of Pharmacy*,

Pharmaceutical Press 22nd ed. (2013). For a brief review of methods for drug delivery, *see*, Langer, *Science* 249:1527-1533 (1990).

[0179] Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can comprise antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives.

[0180] Formulations may be optimized for retention and stabilization in the brain or central nervous system. When the agent is administered into the cranial compartment, it is desirable for the agent to be retained in the compartment, and not to diffuse or otherwise cross the blood brain barrier. Stabilization techniques include cross-linking, multimerizing, or linking to groups such as polyethylene glycol, polyacrylamide, neutral protein carriers, *etc.* in order to achieve an increase in molecular weight.

[0181] Other strategies for increasing retention include the entrapment of the antibody, such as an anti-SIRP β 1 antibody of the present disclosure, in a biodegradable or bioerodible implant. The rate of release of the therapeutically active agent is controlled by the rate of transport through the polymeric matrix, and the biodegradation of the implant. Implants may be particles, sheets, patches, plaques, fibers, microcapsules and the like and may be of any size or shape compatible with the selected site of insertion. Biodegradable polymeric compositions which may be employed may be organic esters or ethers, which when degraded result in physiologically acceptable degradation products, including the monomers. Anhydrides, amides, orthoesters or the like, by themselves or in combination with other monomers, may find use. The polymers will be condensation polymers. The polymers may be cross-linked or non-cross-linked. Of particular interest are polymers of hydroxylaliphatic carboxylic acids, either homo- or copolymers, and polysaccharides. Included among the polyesters of interest are polymers of D-lactic acid, L-lactic acid, racemic lactic acid, glycolic acid, polycaprolactone, and combinations thereof. Among the polysaccharides of interest are calcium alginate, and functionalized celluloses, particularly carboxymethylcellulose esters characterized by being water insoluble, a molecular weight of about 5 kD to 500 kD, *etc.* Biodegradable hydrogels may also be employed in the implants of the subject invention. Hydrogels are typically a copolymer material, characterized by the ability to imbibe a liquid.

V. Therapeutic uses

[0182] As disclosed herein, anti-SIRP β 1 antibodies of the present disclosure may be used for preventing, reducing risk, or treating diseases and disorders. In various embodiments, the anti-SIRP β 1 antibodies provided herein have one or more activities selected from: agonizing SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*; inducing respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*; inducing IL-8 expression in

monocytes *in vitro* and/or *in vivo*; inducing TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*; inducing neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*; increasing neutrophil-mediated tumor cell clearance *in vivo*; upregulating TREM2 expression on macrophages *in vitro* and/or *in vivo*; increasing viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and increasing viability of dendritic cells *in vitro* and/or *in vivo*.

[0183] In various embodiments, the anti-SIRP β 1 antibodies provided herein may be used for preventing, reducing risk, or treating dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, Taupathy disease, Nasu-Hakola disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, wound healing, Crohn's disease, inflammatory bowel disease, ulcerative colitis, obesity, Malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, seizures, spinal cord injury, traumatic brain injury, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, lupus, arthritis, multiple sclerosis, low bone density, osteoporosis, osteogenesis, osteopetrotic disease, Paget's disease of bone, and/or cancer. In some such embodiments, the anti-SIRP β 1 antibody is an agonist antibody.

[0184] In some embodiments, provided herein are methods of preventing, reducing risk, or treating an individual having dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, Taupathy disease, Nasu-Hakola disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, wound healing, Crohn's disease, inflammatory bowel disease, ulcerative colitis, obesity, Malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, Sarcoidosis, diseases of aging, seizures, spinal cord injury, traumatic brain injury, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, lupus, arthritis, multiple sclerosis, low bone density, osteoporosis, osteogenesis, osteopetrotic disease, Paget's disease of bone, and/or cancer, by administering to the individual a therapeutically effective amount of an anti-SIRP β 1 antibody of the present disclosure. In some such embodiments, the anti-SIRP β 1 antibody is an agonist of (*e.g.*, induces or increases) SIRP β 1 activity.

[0185] As disclosed herein, anti-SIRP β 1 antibodies of the present disclosure may also be used for inducing and/or promoting innate immune cell survival. In some embodiments, the present disclosure provides methods of inducing or promoting innate immune cell survival in an individual in need thereof, by administering to the individual a therapeutically effective amount of an agonist anti-SIRP β 1 antibody of the present disclosure.

[0186] As disclosed herein, anti-SIRP β 1 antibodies of the present disclosure may also be used for inducing and/or promoting wound healing, such as after injury. In some embodiments, the wound healing may be colonic wound repair following injury. In some embodiments, the present disclosure provides methods of inducing or promoting wound healing in an individual in need thereof, by administering to the individual a therapeutically effective amount of an agonist anti-SIRP β 1 antibody of the present disclosure.

[0187] In some embodiments, a subject or individual is a mammal. Mammals include, without limitation, domesticated animals (*e.g.*, cows, sheep, cats, dogs, and horses), primates (*e.g.*, humans and non-human primates such as monkeys), rabbits, and rodents (*e.g.*, mice and rats). In some embodiments, the subject or individual is a human.

[0188] An antibody provided herein (and any additional therapeutic agent) can be administered by any suitable means, including parenteral, intrapulmonary, intranasal, intralesional administration, intracerebrospinal, intracranial, intraspinal, intrasynovial, intrathecal, oral, topical, or inhalation routes. Parenteral infusions include intramuscular, intravenous administration as a bolus or by continuous infusion over a period of time, intraarterial, intra-articular, intraperitoneal, or subcutaneous administration. In some embodiments, the administration is intravenous administration. In some embodiments, the administration is subcutaneous. Dosing can be by any suitable route, *e.g.* by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

[0189] Antibodies provided herein would be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as described herein, or

about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

[0190] For the prevention or treatment of disease, the appropriate dosage of an antibody of the invention (when used alone or in combination with one or more other additional therapeutic agents) will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

[0191] Depending on the type and severity of the disease, about 1 µg/kg to 15 mg/kg (*e.g.*, 0.1 mg/kg-10 mg/kg) of antibody can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of disease symptoms occurs. One exemplary dosage of the antibody would be in the range from about 0.05 mg/kg to about 10 mg/kg. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 4.0 mg/kg, or 10 mg/kg (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, *e.g.*, every week or every three weeks (*e.g.*, such that the patient receives from about two to about twenty, or *e.g.*, about six doses of the antibody). In certain embodiments, dosing frequency is three times per day, twice per day, once per day, once every other day, once weekly, once every two weeks, once every four weeks, once every five weeks, once every six weeks, once every seven weeks, once every eight weeks, once every nine weeks, once every ten weeks, or once monthly, once every two months, once every three months, or longer. An initial higher loading dose, followed by one or more lower doses may be administered. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

Dementia

[0192] Dementia is a non-specific syndrome (*i.e.*, a set of signs and symptoms) that presents as a serious loss of global cognitive ability in a previously unimpaired person, beyond what might be expected from normal ageing. Dementia may be static as the result of a unique global brain injury. Alternatively, dementia may be progressive, resulting in long-term decline due to damage or disease in the body. While dementia is much more common in the geriatric population, it can also occur before the age of 65. Cognitive areas affected by dementia include, without limitation, memory, attention span, language, and problem solving. Generally, symptoms must be present for at least six months to before an individual is diagnosed with dementia.

[0193] Exemplary forms of dementia include, without limitation, frontotemporal dementia, Alzheimer's disease, vascular dementia, semantic dementia, and dementia with Lewy bodies.

[0194] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat dementia. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having dementia.

Frontotemporal dementia

[0195] Frontotemporal dementia (FTD) is a condition resulting from the progressive deterioration of the frontal lobe of the brain. Over time, the degeneration may advance to the temporal lobe. Second only to Alzheimer's disease (AD) in prevalence, FTD accounts for 20% of pre-senile dementia cases. The clinical features of FTD include memory deficits, behavioral abnormalities, personality changes, and language impairments (Cruts, M. & Van Broeckhoven, C., Trends Genet. 24:186-194 (2008); Neary, D., *et al.*, Neurology 51:1546-1554 (1998); Ratnavalli, E., Brayne, C., Dawson, K. & Hodges, J. R., Neurology 58:1615-1621 (2002)).

[0196] A substantial portion of FTD cases are inherited in an autosomal dominant fashion, but even in one family, symptoms can span a spectrum from FTD with behavioral disturbances, to Primary Progressive Aphasia, to Cortico-Basal Ganglionic Degeneration. FTD, like most neurodegenerative diseases, can be characterized by the pathological presence of specific protein aggregates in the diseased brain. Historically, the first descriptions of FTD recognized the presence of intraneuronal accumulations of hyperphosphorylated Tau protein in neurofibrillary tangles or Pick bodies. A causal role for the microtubule associated protein Tau was supported by the identification of mutations in the gene encoding the Tau protein in several families (Hutton, M., *et al.*, Nature 393:702-705 (1998). However, the majority of FTD brains show no accumulation of hyperphosphorylated Tau but do exhibit immunoreactivity to ubiquitin (Ub) and TAR DNA binding protein (TDP43) (Neumann, M., *et al.*, Arch. Neurol. 64:1388-1394 (2007)). A majority of those FTD cases with Ub inclusions (FTD-U) were shown to carry mutations in the progranulin gene.

[0197] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat FTD. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having FTD.

Alzheimer's disease

[0198] Alzheimer's disease (AD) is the most common form of dementia. There is no cure for the disease, which worsens as it progresses, and eventually leads to death. Most often, AD is diagnosed in people over 65 years of age. However, the less-prevalent early-onset *alzheimer's* can occur much earlier.

[0199] Common symptoms of Alzheimer's disease include, behavioral symptoms, such as difficulty in remembering recent events; cognitive symptoms, confusion, irritability and aggression, mood swings, trouble with language, and long-term memory loss. As the disease progresses bodily

functions are lost, ultimately leading to death. Alzheimer's disease develops for an unknown and variable amount of time before becoming fully apparent, and it can progress undiagnosed for years.

[0200] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat Alzheimer's disease. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having Alzheimer's disease.

Nasu-Hakola disease

[0201] Nasu-Hakola disease (NHD), which may alternatively be referred to as polycystic lipomembranous osteodysplasia with sclerosing leukoencephalopathy (PLOS), is a rare inherited leukodystrophy characterized by progressive presenile dementia associated with recurrent bone fractures due to polycystic osseous lesions of the lower and upper extremities. NHD disease course is generally divided into four stages: latent, osseous, early neurologic, and late neurologic. After a normal development during childhood (latent stage), NHD starts manifesting during adolescence or young adulthood (typical age of onset 20-30 years) with pain in the hands, wrists, ankles, and feet. Patients then start suffering from recurrent bone fractures due to polycystic osseous and osteoporotic lesions in the limb bones (osseous stage). During the third or fourth decade of life (early neurologic stage), patients present with pronounced personality changes (*e.g.*, euphoria, lack of concentration, loss of judgment, and social inhibitions) characteristic of a frontal lobe syndrome. Patients also typically suffer from progressive memory disturbances. Epileptic seizures are also frequently observed. Finally (late neurologic stage), patients progress to a profound dementia, are unable to speak and move, and usually die by the age of 50.

[0202] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat Nasu-Hakola disease (NHD). In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having NHD.

Parkinson's disease

[0203] Parkinson's disease, which may be referred to as idiopathic or primary parkinsonism, hypokinetic rigid syndrome (HRS), or paralysis agitans, is a neurodegenerative brain disorder that affects motor system control. The progressive death of dopamine-producing cells in the brain leads to the major symptoms of Parkinson's. Most often, Parkinson's disease is diagnosed in people over 50 years of age. Parkinson's disease is idiopathic (having no known cause) in most people. However, genetic factors also play a role in the disease.

[0204] Symptoms of Parkinson's disease include, without limitation, tremors of the hands, arms, legs, jaw, and face, muscle rigidity in the limbs and trunk, slowness of movement (bradykinesia), postural instability, difficulty walking, neuropsychiatric problems, changes in speech or behavior, depression, anxiety, pain, psychosis, dementia, hallucinations, and sleep problems.

[0205] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat Parkinson's disease. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having Parkinson's disease.

Amyotrophic lateral sclerosis

[0206] As used herein, amyotrophic lateral sclerosis (ALS) or motor neuron disease or Lou Gehrig's disease are used interchangeably and refer to a debilitating disease with varied etiology characterized by rapidly progressive weakness, muscle atrophy and fasciculations, muscle spasticity, difficulty speaking (dysarthria), difficulty swallowing (dysphagia), and difficulty breathing (dyspnea).

[0207] It has been shown that progranulin play a role in ALS (Schymick, JC *et al.*, (2007) J Neurol Neurosurg Psychiatry.;78:754-6) and protects against the damage caused by ALS causing proteins such as TDP-43 (Laird, AS *et al.*, (2010). PLoS ONE 5: e13368). It was also demonstrated that pro-NGF induces p75 mediated death of oligodendrocytes and corticospinal neurons following spinal cord injury (Beatty *et al.*, Neuron (2002),36, pp. 375-386; Giehl *et al.*, Proc. Natl. Acad. Sci USA (2004), 101, pp 6226-30).

[0208] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat ALS. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having ALS.

Huntington's disease

[0209] Huntington's disease (HD) is an inherited neurodegenerative disease caused by an autosomal dominant mutation in the Huntingtin gene (HTT). Expansion of a cytosine-adenine-guanine (CAG) triplet repeat within the Huntingtin gene results in production of a mutant form of the Huntingtin protein (Htt) encoded by the gene. This mutant Huntingtin protein (mHtt) is toxic and contributes to neuronal death. Symptoms of Huntington's disease most commonly appear between the ages of 35 and 44, although they can appear at any age.

[0210] Symptoms of Huntington's disease, include, without limitation, motor control problems, jerky, random movements (chorea), abnormal eye movements, impaired balance, seizures, difficulty chewing, difficulty swallowing, cognitive problems, altered speech, memory deficits, thinking difficulties, insomnia, fatigue, dementia, changes in personality, depression, anxiety, and compulsive behavior.

[0211] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat Huntington's disease (HD). In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having HD.

Taupathy disease

[0212] Taupathy diseases, or Tauopathies, are a class of neurodegenerative disease caused by aggregation of the microtubule-associated protein tau within the brain. Alzheimer's disease (AD) is the most well-known Taupathy disease, and involves an accumulation of tau protein within neurons in the form of insoluble neurofibrillary tangles (NFTs). Other Taupathy diseases and disorders include progressive supranuclear palsy, dementia pugilistica (chronic traumatic encephalopathy), Frontotemporal dementia and parkinsonism linked to chromosome 17, Lytico-Bodig disease (Parkinson-dementia complex of Guam), Tangle-predominant dementia, Ganglioglioma and gangliocytoma, Meningioangiomas, Subacute sclerosing panencephalitis, lead encephalopathy, tuberous sclerosis, Hallervorden-Spatz disease, lipofuscinosis, Pick's disease, corticobasal degeneration, Argyrophilic grain disease (AGD), Huntington's disease, frontotemporal dementia, and frontotemporal lobar degeneration.

[0213] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat Taupathy disease. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having Taupathy disease.

Multiple sclerosis

[0214] Multiple sclerosis (MS) can also be referred to as disseminated sclerosis or encephalomyelitis disseminata. MS is an inflammatory disease in which the fatty myelin sheaths around the axons of the brain and spinal cord are damaged, leading to demyelination and scarring as well as a broad spectrum of signs and symptoms. MS affects the ability of nerve cells in the brain and spinal cord to communicate with each other effectively. Nerve cells communicate by sending electrical signals called action potentials down long fibers called axons, which are contained within an insulating substance called myelin. In MS, the body's own immune system attacks and damages the myelin. When myelin is lost, the axons can no longer effectively conduct signals. MS onset usually occurs in young adults, and is more common in women.

[0215] Symptoms of MS include, without limitation, changes in sensation, such as loss of sensitivity or tingling; pricking or numbness, such as hypoesthesia and paresthesia; muscle weakness; clonus; muscle spasms; difficulty in moving; difficulties with coordination and balance, such as ataxia; problems in speech, such as dysarthria, or in swallowing, such as dysphagia; visual problems, such as nystagmus, optic neuritis including phosphenes, and diplopia; fatigue; acute or chronic pain; and bladder and bowel difficulties; cognitive impairment of varying degrees; emotional symptoms of depression or unstable mood; Uhthoff's phenomenon, which is an exacerbation of extant symptoms due to an exposure to higher than usual ambient temperatures; and Lhermitte's sign, which is an electrical sensation that runs down the back when bending the neck.

[0216] In some embodiments, administering an anti-SIRP β 1 antibody of the present disclosure can prevent, reduce the risk, and/or treat multiple sclerosis. In some embodiments, administering an anti-SIRP β 1 antibody may induce one or more SIRP β 1 activities in an individual having multiple sclerosis.

Cancer

[0217] Yet further aspects of the present disclosure provide methods for preventing, reducing risk, or treating an individual having cancer, comprising administering to the individual a therapeutically effective amount of an isolated anti-SIRP β 1 antibody of the present disclosure. Any of the isolated antibodies of the present disclosure may be used in these methods. In some embodiments, the isolated antibody is an agonist antibody of the present disclosure.

[0218] The tumor microenvironment is known to contain a heterogeneous immune infiltrate, which includes T lymphocytes, macrophages and cells of myeloid/granulocytic lineage. In particular, the presence of M2-macrophages in tumors is associated with poor prognosis. Therapies that reduce the number of these cells in the tumor, such as CSF1R blocking agents, are showing beneficial effects in preclinical models and early stage clinical studies. A seminal preclinical study has also shown synergies between drugs that target tumor-associated macrophages (*e.g.*, CSF1/CSF1R blocking antibodies) and checkpoint blocking antibodies that target T cells, indicating that manipulating both cell types shows efficacy in tumor models where individual therapies are poorly effective (Zhu Y; *Cancer Res.* 2014 Sep 15; 74(18):5057-69). Without wishing to be bound by any particular theory, it is thought that inducing SIRP β 1 signaling in tumor associated macrophages and/or neutrophils may inhibit suppression of the immune response in the tumor microenvironment, resulting in a therapeutic anti-tumor immune response.

[0219] In certain embodiments, a cancer to be prevented or treated by the methods of the present disclosure includes, without limitation, squamous cell carcinoma (*e.g.*, epithelial squamous cell carcinoma), lung cancer, small-cell lung cancer, non-small cell lung cancer, squamous non-small cell lung cancer (NSCLC), adenocarcinoma of the lung, squamous carcinoma of the lung, non-squamous NSCLC, glioma, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, gastric or stomach cancer including gastrointestinal cancer and gastrointestinal stromal cancer, renal cancer (*e.g.* clear cell carcinoma), ovarian cancer, liver cancer, colorectal cancer, endometrial cancer, kidney cancer (*e.g.*, renal cell carcinoma (RCC)), prostate cancer (*e.g.* hormone refractory prostate adenocarcinoma), thyroid cancer, neuroblastoma, pancreatic cancer, glioblastoma (glioblastoma multiforme), cervical cancer, stomach cancer, bladder cancer, hepatoma, breast cancer, colon carcinoma, and head and neck cancer (or carcinoma), gastric cancer, germ cell tumor, pediatric sarcoma, sinonasal natural killer, melanoma (*e.g.*, metastatic malignant melanoma, such as cutaneous or intraocular malignant melanoma), bone cancer, skin cancer, uterine cancer, cancer of the anal region, testicular cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium,

carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, cancer of the ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, myelodysplastic syndromes, colorectal neoplasms, solid tumors, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally-induced cancers including those induced by asbestos, virus-related cancers (*e.g.*, human papilloma virus (HPV)-related tumor), and hematologic malignancies derived from either of the two major blood cell lineages, *i.e.*, the myeloid cell line (which produces granulocytes, erythrocytes, thrombocytes, macrophages and mast cells) or lymphoid cell line (which produces B, T, NK and plasma cells), such as all types of leukemias, lymphomas, and myelomas, *e.g.*, acute, chronic, lymphocytic and/or myelogenous leukemias, such as acute leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), and chronic myelogenous leukemia (CML), undifferentiated AML (M0), myeloblastic leukemia (M1), myeloblastic leukemia (M2; with cell maturation), promyelocytic leukemia (M3 or M3 variant [M3V]), myelomonocytic leukemia (M4 or M4 variant with eosinophilia [M4E]), monocytic leukemia (M5), erythroleukemia (M6), megakaryoblastic leukemia (M7), isolated granulocytic sarcoma, and chloroma; lymphomas, such as Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), indolent lymphoma, large B cell diffuse lymphoma, B cell hematologic malignancy, *e.g.*, B-cell lymphomas, T-cell lymphomas, lymphoplasmacytoid lymphoma, monocytoid B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma, anaplastic (*e.g.*, Ki 1+) large-cell lymphoma, adult T-cell lymphoma/leukemia, mantle cell lymphoma, angio immunoblastic T-cell lymphoma, angiocentric lymphoma, intestinal T-cell lymphoma, primary mediastinal B-cell lymphoma, precursor T-lymphoblastic lymphoma, T-lymphoblastic; and lymphoma/leukemia (T-Lbly/T-ALL), peripheral T-cell lymphoma, lymphoblastic lymphoma, post-transplantation lymphoproliferative disorder, true histiocytic lymphoma, primary central nervous system lymphoma, primary effusion lymphoma, lymphoblastic lymphoma (LBL), hematopoietic tumors of lymphoid lineage, acute lymphoblastic leukemia, diffuse large B-cell lymphoma, Burkitt's lymphoma, follicular lymphoma, diffuse histiocytic lymphoma (DHL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, cutaneous T-cell lymphoma (CTLC) (also called mycosis fungoides or Sezary syndrome), and lymphoplasmacytic lymphoma (LPL) with Waldenstrom's macroglobulinemia; myelomas, such as IgG myeloma, light chain myeloma, nonsecretory myeloma, smoldering myeloma (also called indolent myeloma), solitary plasmacytoma, and multiple myelomas, chronic lymphocytic leukemia (CLL), hairy cell lymphoma; hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; seminoma, teratocarcinoma, tumors of the central and

peripheral nervous, including astrocytoma, schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xeroderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma, hematopoietic tumors of lymphoid lineage, for example T-cell and B-cell tumors, including but not limited to T-cell disorders such as T-prolymphocytic leukemia (T-PLL), including of the small cell and cerebriform cell type; large granular lymphocyte leukemia (LGL) preferably of the T-cell type; a/d T-NHL heptasyllabic lymphoma; peripheral/post-thymic T cell lymphoma (pleomorphic and immunoblastic subtypes); angiocentric (nasal) T-cell lymphoma; cancer of the head or neck, renal cancer, rectal cancer, cancer of the thyroid gland; acute myeloid lymphoma, as well as any combinations of said cancers. An anti-SIRPβ1 antibody of the present disclosure may also be used to treat metastatic cancer.

[0220] In some embodiments, the cancer is selected from sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, small cell lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma. In some embodiments, the cancer is triple-negative breast carcinoma. In some embodiments, the cancer may be an early stage cancer or a late stage cancer. In some embodiments, the cancer may be a primary tumor. In some embodiments, the cancer may be a metastatic tumor at a second site derived from any of the above types of cancer.

[0221] In some embodiments, the cancer is selected from glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma; diffuse large B-cell lymphoma; lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, indolent B cell lymphoma, aggressive B cell lymphoma, T cell lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, myelodysplastic syndromes, myeloproliferative neoplasms, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma.

[0222] In some embodiments, the present disclosure provides methods of preventing, reducing risk, or treating an individual having cancer, by administering to the individual a therapeutically effective amount of an anti-SIRPβ1 antibody of the present disclosure.

[0223] In some embodiments, the present disclosure provides methods of preventing, reducing risk, or treating an individual having cancer, wherein the individual is refractory to checkpoint inhibitor therapy, by administering to the individual a therapeutically effective amount of an anti-SIRP β 1 antibody of the present disclosure.

[0224] In some embodiments, an anti-SIRP β 1 antibody of the present disclosure may be administered in conjunction with a therapeutic agent that acts as a checkpoint inhibitor. In some embodiments, the method further includes administering to the individual at least one antibody that specifically binds to an inhibitory immune checkpoint molecule, and/or another standard or investigational anti-cancer therapy. In some embodiments, the inhibitory checkpoint molecule is selected from PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, and CD73. In typical embodiments, the therapeutic agent is an antibody to a checkpoint inhibitor selected from D1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73. In some embodiments, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is administered in combination with the anti-SIRP β 1 antibody of the present disclosure. In some embodiments, a combination of antibodies directed against checkpoint inhibitors is administered in conjunction with an anti-SIRP β 1 antibody of the present invention.

[0225] In some embodiments, an anti-SIRP β 1 antibody of the present disclosure may be administered in conjunction with at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein, *e.g.*, an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof.

[0226] In some embodiments, the method further includes administering to the individual at least one antibody that specifically binds to an inhibitory immune checkpoint molecule, and/or another standard or investigational anti-cancer therapy. In some embodiments, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is administered in combination with the anti-SIRP β 1 antibody of the present disclosure. In some embodiments, the at least one antibody that specifically binds to an inhibitory checkpoint molecule is selected from an anti-PD-L1 antibody, an anti-CTLA4 antibody, an anti-PD-L2 antibody, an anti-PD-1 antibody, an anti-B7-H3 antibody, an anti-B7-H4 antibody, and anti-HVEM antibody, an anti-B- and T-lymphocyte attenuator (BTLA) antibody, an anti-Killer inhibitory receptor (KIR) antibody, an anti-GAL9 antibody, an anti-TIM3

antibody, an anti-A2AR antibody, an anti-LAG-3 antibody, an anti-phosphatidylserine antibody, an anti-CD27 antibody, and any combination thereof. In some embodiments, the standard or investigational anti-cancer therapy is one or more therapies selected from radiotherapy, cytotoxic chemotherapy, targeted therapy, imatinib (Gleevec®), trastuzumab (Herceptin®), adoptive cell transfer (ACT), chimeric antigen receptor T cell transfer (CAR-T), vaccine therapy, and cytokine therapy. In some embodiments, the method further includes administering to the individual at least one antibody that specifically binds to an inhibitory cytokine. In some embodiments, the at least one antibody that specifically binds to an inhibitory cytokine is administered in combination with the anti-SIRPβ1 antibody of the present disclosure. In some embodiments, the at least one antibody that specifically binds to an inhibitory cytokine is selected from an anti-CCL2 antibody, an anti-CSF-1 antibody, an anti-IL-2 antibody, and any combination thereof.

[0227] In some embodiments, the method further includes administering to the individual at least one agonistic antibody that specifically binds to a stimulatory immune checkpoint protein. In some embodiments, the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is administered in combination with the anti-SIRPβ1 antibody of the present disclosure. In some embodiments, the at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein is selected from an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, and any combination thereof.

[0228] In some embodiments, an anti-SIRPβ1 antibody of the present invention is administered in combination with radiation therapy and/or a chemotherapeutic agent. Chemotherapeutic agents include, for example, the following groups: anti-metabolites/anti-cancer agents, such as pyrimidine analogs (5-fluorouracil, floxuridine, capecitabine, gemcitabine and cytarabine) and purine analogs, folate antagonists and related inhibitors (methotrexate, pemetrexed, mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones, eribulin and navelbine; epidipodophyllotoxins (etoposide, teniposide); DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil, cisplatin, cyclophosphamide, Cytosan, dactinomycin, daunorubicin, doxorubicin, epirubicin, hexamethylmelamineoxaliplatin, iphosphamide, melphalan, merchlorhtamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, temozolamide, teniposide, triethylenethiophosphoramide and etoposide (VP 16)); DNA methyltransferase inhibitors (azacytidine); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin (adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin) and

mitomycin; enzymes (L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepa), alkylsulfonates (busulfan), nitrosoureas (carmustine (BCNU) and analogs, streptozocin), triazenes (dacarbazine (DTIC)); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (TNP470, genistein, pomalidomide) and growth factor inhibitors (vascular endothelial growth factor (VEGF) inhibitors, such as ziv-aflibercept; fibroblast growth factor (FGF) inhibitors); inhibitors of apoptosis protein (IAP) antagonists (birinapant); histone deacetylase (HDAC) inhibitors (vorinostat, romidepsin, chidamide, panobinostat, mocetinostat, abexinostat, belinostat, entinostat, resminostat, givinostat, quisinostat, SB939); proteasome inhibitors (ixazomib); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab, panitumumab, pertuzumab, cetuximab, adalimumab, golimumab, infliximab, rituximab, ocrelizumab, ofatumumab, obinutuzumab, alemtuzumab, abciximab, atlizumab, daclizumab, denosumab, efalizumab, elotuzumab, rovelizumab, ruplizumab, ustekinumab, visilizumab, gemtuzumab ozogamicin, brentuximab vedotin); chimeric antigen receptors; cell cycle inhibitors (flavopiridol, roscovitine, bryostatin-1) and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin, irinotecan (CPT-11) and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); PARP inhibitors (niraparib, olaparib); focal adhesion kinase (FAK) inhibitors (defactinib (VS-6063), VS-4718, VS-6062, GSK2256098); growth factor signal transduction kinase inhibitors (cediranib, galunisertib, rociletinib, vandetanib, afatinib, EGF816, AZD4547); c-Met inhibitors (capmatinib, INC280); ALK inhibitors (ceritinib, crizotinib); mitochondrial dysfunction inducers, toxins such as Cholera toxin, ricin, Pseudomonas exotoxin, Bordetella pertussis adenylate cyclase toxin, or diphtheria toxin, and caspase activators; and chromatin disruptors. In some embodiments, a chemotherapeutic agent is a B-Raf inhibitor, a MEK inhibitor, a VEGF inhibitor, a VEGFR inhibitor, a tyrosine kinase inhibitor, an anti-mitotic agent, or any combination thereof.

[0229] In some embodiments, an anti-SIRP β 1 antibody of the present disclosure is administered in combination with adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and/or cytokine therapy.

[0230] In some embodiments, an anti-SIRP β 1 antibody of the present disclosure is administered in combination with at least one antibody that specifically binds to an inhibitory cytokine, *e.g.*, an inhibitory cytokine such as an anti-CCL2 antibody, an anti-CSF-1 antibody, or an anti-IL-2 antibody.

[0231] In some embodiments, an anti-SIRP β 1 antibody of the present disclosure is administered in combination with at least one stimulatory cytokine. In some embodiments that may be combined with any of the preceding embodiments, the at least one stimulatory cytokine is selected from IFN- α 4, IFN- β , IL-1 β , TNF- α , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- γ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.

VI. Diagnostic uses

[0232] In some embodiments of any of the antibodies, any of the anti-SIRP β 1 antibodies provided herein is useful for detecting the presence of SIRP β 1 in a sample or an individual. The term “detecting” as used herein encompasses quantitative or qualitative detection. Provided herein are methods of using the antibodies of this disclosure for diagnostic purposes, such as the detection of SIRP β 1 in an individual or in tissue samples derived from an individual. In some embodiments, the individual is a human.

[0233] The detection method may involve quantification of the antigen-bound antibody. Antibody detection in biological samples may occur with any method known in the art, including immunofluorescence microscopy, immunocytochemistry, immunohistochemistry, ELISA, FACS analysis, immunoprecipitation, or micro-positron emission tomography. In certain embodiments, the antibody is radiolabeled, for example with ^{18}F and subsequently detected utilizing micro-positron emission tomography analysis. Antibody-binding may also be quantified in a patient by non-invasive techniques such as positron emission tomography (PET), X-ray computed tomography, single-photon emission computed tomography (SPECT), computed tomography (CT), and computed axial tomography (CAT).

VII. Articles of Manufacture

[0234] Provided herein are articles of manufacture (*e.g.*, kit) comprising an anti-SIRP β 1 antibody described herein. Article of manufacture may include one or more containers comprising an antibody described herein. Containers may be any suitable packaging including, but is not limited to, vials, bottles, jars, flexible packaging (*e.g.*, sealed Mylar or plastic bags), and the like. The containers may be unit doses, bulk packages (*e.g.*, multi-dose packages) or sub-unit doses.

[0235] In some embodiments, the kits may further include a second agent. In some embodiments, the second agent is a pharmaceutically-acceptable buffer or diluting agent including, but not limited to, such as bacteriostatic water for injection (BWFI), phosphate- buffered saline, Ringer's solution and dextrose solution. In some embodiments, the second agent is a pharmaceutically active agent. In some embodiments, the second agent is a pharmaceutically active agent described herein.

[0236] In some embodiments of any of the articles of manufacture, the article of manufactures further include instructions for use in accordance with the methods of this disclosure. The instructions generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. In some embodiments, these instructions comprise a description of administration of the isolated antibody of the present disclosure (*e.g.*, an anti-SIRP β 1 antibody described herein) to prevent, reduce risk, or treat an individual having a disease, disorder, or injury selected from dementia, frontotemporal dementia, Alzheimer's disease, vascular dementia, mixed dementia, Creutzfeldt-Jakob disease, normal pressure hydrocephalus, amyotrophic lateral sclerosis, Huntington's disease, taupathy disease, Nasu-Hakola disease, stroke, acute trauma, chronic trauma, lupus, acute and chronic colitis, rheumatoid arthritis, wound healing, Crohn's disease, inflammatory bowel disease, ulcerative colitis, obesity, malaria, essential tremor, central nervous system lupus, Behcet's disease, Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, Shy-Drager syndrome, progressive supranuclear palsy, cortical basal ganglionic degeneration, acute disseminated encephalomyelitis, granulomatous disorders, sarcoidosis, diseases of aging, seizures, spinal cord injury, traumatic brain injury, age related macular degeneration, glaucoma, retinitis pigmentosa, retinal degeneration, respiratory tract infection, sepsis, eye infection, systemic infection, lupus, arthritis, multiple sclerosis, low bone density, osteoporosis, osteogenesis, osteopetrotic disease, Paget's disease of bone, and cancer, according to any methods of this disclosure. In some embodiments, the instructions include instructions for use of the anti-SIRP β 1 antibody and the second agent (*e.g.*, second pharmaceutically active agent).

[0237] The present disclosure will be more fully understood by reference to the following Examples. They should not, however, be construed as limiting the scope of the present disclosure. All citations throughout the disclosure are hereby expressly incorporated by reference.

EXAMPLES

[0238] The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of non-critical parameters that could be changed or modified to yield essentially similar results.

Example 1: Production, identification, and characterization of agonist anti-SIRP β 1 antibodies

[0239] The amino acid sequence of the human SIRP β 1 isoform 1 protein (SIRP β 1) is set forth below in SEQ ID NO: 1. Human SIRP β 1 contains a signal peptide located at amino residues 1-29 of SEQ ID NO: 1; an extracellular immunoglobulin-like variable-type (IgV) domain located at amino

residues 30-136 of SEQ ID NO: 1; two immunoglobulin-like constant (IgC) domains located at amino acids 147-246 and 253-347 SEQ ID NO: 1; a transmembrane domain located at amino residues 372-392 of SEQ ID NO: 1; and an intracellular domain located at amino residues 393-398 of SEQ ID NO: 1.

Human SIRP β 1 amino acid sequence (SEQ ID NO: 1):

10	20	30	40	50
MPVPASWPHL	PSPFLLMTLL	LGRLTGVAGE	DELQVIQPEK	SVSVAAGESA
60	70	80	90	100
TLRCAMTSLI	PVGPIMWFRG	AGAGRELIYN	QKEGHFPRVT	TVSELTKRNN
110	120	130	140	150
LDFSISISNI	TPADAGTYYC	VKFRKGSPDD	VEFKSGAGTE	LSVRAKPSAP
160	170	180	190	200
VVSGPAVRAT	PEHTVSFTCE	SHGFSPRDI	LKWFKNGNEL	SDFQTNVDPA
210	220	230	240	250
GDSVSYSIHS	TARVVLTRGD	VHSQVICEIA	HITLQGDPLR	GTANLSEAIR
260	270	280	290	300
VPPTLEVTQQ	PMRAENQANV	TCQVSNFYPR	GLQLTWLENG	NVSRTEAST
310	320	330	340	350
LIENKDGTYN	WMSWLLVNTC	AHRDDVLTCT	QVEHDGQQAV	SKSYALEISA
360	370	380	390	
HQKEHGSDIT	HEAALAPTAP	LLVALLGPK	LLLVGVSAI	YICWKQKA

[0240] The human SIRP β 1 amino acid sequence comprises a lysine residue (aa380) within the transmembrane domain that interacts with an aspartic acid in DAP12, a key adaptor protein that transduces signaling from SIRP β 1, as well as TREM1, TREM2, and other related IgV family members. A BLAST analysis of human SIRP β 1 identified multiple isoforms possibly derived from alternative splicing. Among these transcript variants, SIRP β 1 isoform 3 shares the most sequence identity to SIRP β 1 isoform 1 (**FIG. 1A**). Human SIRP β 1 is also related to human SIRP α . An alignment of the amino acid sequences of human SIRP β 1 and human SIRP α was generated by 2-way blast (**FIG. 1B**). The extracellular region of both receptors share 85% sequence identity. In contrast, human SIRP β 1 displays poorer homology to mouse SIRP β 1 (**FIG. 2**) suggesting that the SIRP gene loci in both species experience divergent selection pressure.

[0241] Antibodies that bind the extracellular domain of human SIRP β 1, particularly the extracellular IgV domain (amino acid residues 30-136 of SEQ ID NO: 1) can be generated using mouse hybridoma technology, phage display technology, and yeast-based platform technology. Antibodies are screened for their ability to bind cells that express SIRP β 1 and for their ability to activate SIRP β 1 signaling, activity, and function in cells and in a whole animal *in vivo* as described

in Examples 2-20 below. For example, agonistic anti-SIRP β 1 antibodies can be produced that target the IgV domain (amino acid residues 30-136). By analogy to SIRP α , the IgV domain of SIRP β 1 is predicted bind to an unknown ligand(s), and through multimerization of receptor, lead to activation.

Production of His-tagged and Fc-conjugated human SIRP α and human SIRP β 1 IgV domains

[0242] For mammalian expression of human SIRP α and human SIRP β 1 IgV domain (also referred to as “domain 1”) antigens (SEQ ID NOs: 387 and 388, respectively), synthetic genes based on cDNA were cloned into mammalian expression vectors, followed by transient transfection and expression in HEK293/T cells. Constructs included a heterologous signal peptide and C-terminal human IgG1 Fc and/or or His tag. Briefly, expression vectors containing the antigen of interest were transfected by complexing with a transfection reagent followed by exposure to HEK293/T cells for one hour followed by dilution of culture media to a final density of 4 million cells per mL. The cells were then cultured for 7 days with fresh feed media every 48 hours. After 7 days, the supernatant was collected following centrifugation and purification was performed using Ni-sepharose and, in some cases, an SEC column purification to reach >95% non-aggregated monomer content. SIRP α and SIRP β 1 monomer antigens were prepared by fragmenting a SIRP α / β 1 Fc fusion antigen with modified hinge region (Lynaugh *et al.*, MAb. 2013 Oct;5(5):641-45) with FabRICATOR (IdeS) protease (Genovis, Cat # A2-FR2-1000), followed by Protein A affinity purification to remove undigested Fc fusion protein and SEC to remove aggregated monomer.

Library screening for anti-SIRP β 1 antibodies

[0243] Eight naïve (pre-immune) human synthetic yeast libraries each of $\sim 10^9$ diversity were designed, generated, and propagated as described previously (*see, e.g.*: Xu *et al.*, 2013; WO2009036379; WO2010105256; WO2012009568; Xu *et al.*, Protein Eng Des Sel. 2013 Oct;26(10):663-70). Ten parallel selections were performed, using the eight naïve libraries for human SIRP β 1-Fc fusion antigen selections and two pools of the eight libraries for human SIRP β 1 monomer selections. For the first two rounds of selection, a magnetic bead sorting technique utilizing the Miltenyi MACs system was performed, essentially as described (Siegel *et al.*, J Immunol Methods. 2004 Mar;286(1-2): 141-53). Briefly, yeast cells ($\sim 10^9$ cells/library, total density $\sim 10^{10}$) were incubated with 3 ml of 10 nM biotinylated SIRP β 1-Fc fusion antigen or 100 nM biotinylated SIRP β 1 monomer antigen for 15 min at room temperature in FACS wash buffer PBS with 0.1% BSA. After washing once with 50 ml ice-cold wash buffer, the cell pellet was resuspended in 40 mL wash buffer, and 500 μ L Streptavidin MicroBeads (Miltenyi Biotec, Bergisch Gladbach, Germany. Cat # 130-048-101) were added to the yeast and incubated for 15 min at 4°C. Next, yeast cells were pelleted, resuspended in 5 mL wash buffer, and loaded onto a MACS LS column (Miltenyi Biotec, Bergisch Gladbach, Germany. Cat.#130-042-401). After the 5 mL was loaded, the column was washed 3 times with 3 ml FACS wash buffer. The column was then

removed from the magnetic field, and the yeast were eluted with 5 mL of growth media and then grown overnight. The following four rounds of sorting were performed using flow cytometry.

[0244] For the first FACS selection round, approximately 1×10^8 yeast were pelleted, washed three times with wash buffer, and incubated with 10 nM biotinylated SIRP β 1-Fc fusion antigen or 100 nM biotinylated SIRP β 1 monomer antigen for 10 min at room temperature. Yeast were then washed twice and stained with goat anti-human F(ab')₂ kappa-FITC diluted 1:100 (Southern Biotech, Birmingham, Alabama, Cat# 2062-02) and either streptavidin-Alexa Fluor 633 (Life Technologies, Grand Island, NY, Cat # S21375) diluted 1:500, or Extravidin-phycoerythrin (Sigma-Aldrich, St Louis, Cat # E4011) diluted 1:50, secondary reagents for 15 min at 4°C. After washing twice with ice-cold wash buffer, the cell pellets were resuspended in 0.4 mL wash buffer and transferred to strainer-capped sort tubes. Sorting was performed using a FACS ARIA sorter (BD Biosciences) and sort gates were determined to select only SIRP β 1 binding clones. In the following selection round approximately 2×10^7 yeast were prepared as above but incubation was with a polyspecific reactivity reagent to conduct a negative sort to decrease polyspecific binders (Xu *et al.*, PEDS. 2013 Oct;26(10):663-70), and binders to control protein, HIS-tagged human SIRP α monomer. The next round utilized labeling with 10 nM human SIRP β 1-Fc fusion antigens and 100 nM human SIRP β 1 monomer antigen. After the final round of sorting, yeast cells were plated and individual colonies were picked for characterization. The final round utilized labeling with 100 nM and 10 nM human SIRP β 1 monomer antigen.

[0245] Heavy chains from the second and fourth FACS sorting selection round outputs were used to prepare light chain diversification libraries used for additional selections. For these selections, the first selection round utilized Miltenyi MACs beads and labeling with 10 nM human SIRP β 1-Fc fusion antigen. Four rounds of FACS sorting followed. The first round used 100 nM human SIRP β 1 monomer antigen. The second FACS round was a negative sort to decrease binding to reagent binders, polyspecific binders, and binders to control protein human SIRP α HIS tagged monomer. The last two rounds utilized human SIRP β 1 monomer titration (100 nM, 10 nM, and 1 nM) to select highest affinity binders, 100 nM human SIRP α monomer, and competition with control AM4-5 antibody, which binds to SIRP β 1 IgV domain, to assess competitor representation in the enriched population. (See United States Patent Application Publication No. US2014/0242095.) After the final round of sorting, yeast cells were plated and individual colonies were picked for characterization.

Antibody IgG and Fab production and purification

[0246] Yeast clones were grown to saturation and then induced for 48 h at 30°C with shaking. After induction, yeast cells were pelleted and the supernatants were harvested for purification. IgGs were purified using a Protein A column and eluted with acetic acid, pH 2.0. Fab fragments were generated by papain digestion and purified over CaptureSelect IgG-CH1 affinity matrix

(LifeTechnologies, Cat # 1943200250).

Octet® binding experiments

[0247] The affinities of the anti-SIRPβ1 antibodies were determined by measuring their dissociation constants (K_D) using a ForteBio Octet® Red384 system (ForteBio, Menlo Park, CA), performed generally as previously described (Estep *et al.*, MAb. 2013 Mar-Apr;5(2):270-8). Briefly, Octet affinity measurements were performed by loading IgGs on-line onto AHQ sensors. Sensors were equilibrated off-line in assay buffer for 30 min and then monitored on-line for 60 seconds for baseline establishment. For avid binding measurement, sensors with loaded IgGs were exposed to 100 nM antigen (human SIRPα domain 1 (D1) Fc fusion or SIRPβ1 domain 1 (D1) Fc fusion) for 3 min, and then transferred to assay buffer for 3 min for off-rate measurement. Additional avid binding was determined by loading biotinylated SIRPβ1 monomer on SA sensors and exposing to 100 nM IgG in solution. Monovalent binding measurements were obtained by loading human SIRPα- or SIRPβ1-Fc fusion antigens onto AHQ sensor, followed by exposure to 100 nM anti-SIRPβ1 antibody Fab. Additional monovalent measurements were made by loading biotinylated human SIRPα or SIRPβ1 monomer to SA sensor followed by exposure to 100 nM Fab in solution. Kinetics data were fit using a 1:1 binding model in the data analysis software provided by ForteBio (ForteBio Data Analysis Software 7.0).

Epitope binning

[0248] Epitope binning of the anti-SIRPβ1 antibodies was performed on a ForteBio Octet® Red384 system (ForteBio, Menlo Park, CA) using a standard sandwich format binning assay. Control anti-target IgG was loaded onto AHQ sensors and unoccupied Fc-binding sites on the sensor were blocked with a non-relevant human IgG1 antibody. The sensors were then exposed to 100 nM target antigen followed by a second anti-target antibody. Data was processed using ForteBio Data Analysis Software 7.0. Additional binding by the second antibody after antigen association indicates an unoccupied epitope (non-competitor), while no binding indicates epitope blocking (competitor). This process was iterated for two reference antibodies that bind the IgV domain of SIRPβ1: (i) AM4-5, which defines bin 1; and (ii) SB-17, which defines bin 2.

[0249] The final set of anti-SIRPβ1 antibodies were selected based on antigen binding affinities. Antibodies that were positive for binding to human SIRPβ1 were tested for cross-reactivity with human SIRPα. The bin category of each of the antibodies are listed below in **Table 2**. In Table 2, “ND” refers to antibodies for which the Bin category was not determined; “NB” refers to antibodies for which there was no binding to indicated antigen; “PF” refers to antibodies for which antigen binding kinetics showed poor fit to 1:1 binding model. No detectable binding to mouse SIRPβ1 D1-Fc, human SIRPα D1-Fc, or mouse SIRPα D1-Fc was observed for any of the anti-SIRPβ1 antibodies.

Table 2: Biochemical Characterization of anti-SIRPβ1 antibodies

Clone Index (SB-#)	Bin Code (Human SIRPβ1)	Fab K_D Human SIRPβ1-Fc (M) Monovalent	IgG K_D Human SIRPβ1 D1-Fc (M) Avid
1*	2	9.79E-08	9.66E-10
2*	2	2.66E-07	1.97E-09
3	2	P.F.	P.F.
4	2	P.F.	P.F.
5	2	P.F.	P.F.
6	2	1.31E-07	1.30E-09
7	2	P.F.	P.F.
8*	2	1.26E-07	2.72E-09
9	2	P.F.	3.66E-09
10	2	P.F.	P.F.
11	2	6.78E-07	3.69E-09
12	1	N.B.	4.94E-09
13	1	P.F.	P.F.
14	2	P.F.	P.F.
15	2	4.07E-07	3.45E-09
16	2	N.B.	P.F.
17	2	2.46E-07	3.32E-09
18	2	P.F.	P.F.
19	2	N.B.	7.41E-09
20	2	P.F.	P.F.
21	1	N.B.	5.86E-09
22	2	N.B.	P.F.
23	2	P.F.	P.F.
24	1	N.B.	P.F.
25	2	N.B.	P.F.
26	2	N.B.	6.17E-09
27	2	P.F.	P.F.
28	2	P.F.	5.59E-09
29	2	N.B.	1.07E-08
30	2	N.B.	7.87E-09
31	2	P.F.	3.75E-09
32	1	N.B.	7.13E-09
33	2	N.B.	9.23E-09
34	1	N.B.	P.F.
35	2	N.B.	7.47E-09
36	2	N.B.	P.F.
37	2	N.B.	7.50E-09
38	2	N.B.	7.33E-09
39	1	N.B.	2.14E-08

40*	1,2	4.27E-07	9.64E-09
41	1,2	N.B.	1.03E-08
42	1	N.B.	1.23E-08
43	1	N.B.	1.23E-08
44	1,2	N.B.	1.89E-08
45	1	N.B.	1.39E-08
46	2	N.B.	2.00E-08
47	1,2	N.B.	1.44E-08
48	1,2	N.B.	2.77E-08
49	2	1.56E-07	4.39E-09
50	2	P.F.	P.F.

Antibody heavy chain and light chain variable domain sequences

[0250] Using standard techniques, the amino acid sequences encoding the light chain variable domains and the heavy chain variable domains of the antibodies were determined. The EU or Kabat sequences of the antibodies are set forth in **Tables 3-6**, as follows. The EU or Kabat light chain HVR sequences of the antibodies are set forth in **Table 3**. The EU or Kabat heavy chain HVR sequences of the antibodies are set forth in **Table 4**. The EU or Kabat light chain framework (FR) sequences of the antibodies are set forth in **Table 5**. The EU or Kabat heavy chain framework (FR) sequences of the antibodies are set forth in **Table 6**.

Table 3: EU or Kabat light chain HVR sequences of anti-SIRPβ1 antibodies

Ab ID	HVR L1	SEQ ID NO	HVR L2	SEQ ID NO	HVR L3	SEQ ID NO
SB-1	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-2	RASQSVSSSYLA	383	GASSRAT	16	QQSSSHPFT	32
SB-3	RASQSVSSSYLA	2	DASNRAT	17	QQRLFHPPT	33
SB-4	RASQSVSSSYLA	383	GASSRAT	16	QQYADAPIT	34
SB-5	RASQSVSSSYLA	2	DASNRAT	17	QQRLFHPPT	33
SB-6	RASQSVSSSYLA	383	GASSRAT	16	QQSGHLPIT	35
SB-7	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQHYIAPFT	36
SB-8	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-9	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQYHSVPPIT	38
SB-10	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQAIESPLT	39
SB-11	RSSQSLLYSNGYNYLD	5	LGSNRAS	20	VQALQTPLT	40
SB-12	RASQSVSSNLA	6	SASTRAT	21	QQLDNLPYT	41
SB-13	RSSQSLLYSNGYNYLD	5	LGSNRAS	20	MQALRSPIT	42
SB-14	RASQSVSSSYLA	2	DSSNRAT	22	QQFSYYPIT	43
SB-15	RASQSISSYLN	7	AASSLQS	23	QQAYSHPFT	44
SB-16	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQLFSTPFT	45

Ab ID	HVR L1	SEQ ID NO	HVR L2	SEQ ID NO	HVR L3	SEQ ID NO
SB-17	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQYYDDPYT	46
SB-18	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	LQALQTPIT	47
SB-19	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQAIGVPPT	48
SB-20	KSSQSVLYSSNNKNYLA	8	WASTRES	18	QQYYLSPFT	49
SB-21	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQTLRIPPT	50
SB-22	RASQGISSWLA	9	AASNLSQS	24	QQGNSYPIT	51
SB-23	RASQSISSYLN	7	AASSLQS	23	QQAYPYPLT	52
SB-24	RASQSVSSNLA	6	GASTRAT	25	QQLNIHPWT	53
SB-25	RASQGISSWLA	9	AASSLQS	23	QQVNSFPWT	54
SB-26	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQARGLPT	55
SB-27	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQAVSDPPT	56
SB-28	RASQSVSSSYLA	383	GASSRAT	16	QQDGNFPLT	57
SB-29	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQARGSPIT	58
SB-30	RASQSVSSFLA	10	GASSRAT	16	QQFLSSPWT	59
SB-31	RASQGISSWLA	9	AASSLQS	23	QQAVSHPFT	60
SB-32	KSSQSVLYSSNNKNYLA	8	WASTRES	18	QQDFLTPT	61
SB-33	QASQDISNYLN	11	DASNLET	26	QQFAFLPLT	62
SB-34	RASQSVSSNLA	6	GASTRAT	25	QQDNTFPYT	63
SB-35	RSSQSLLHSNGYNYLD	4	LGSNRAS	20	MQTLQVPLT	64
SB-36	RASQGISSWLA	9	AASSLQS	23	QQAFSHRT	65
SB-37	RASQSVSSSYLA	2	DASNRAT	17	QQRHTYPLT	66
SB-38	RASQGISSWLA	9	AASSLQS	23	QQAVSYPIT	67
SB-39	RASQSISSYLN	7	GASSLQS	27	QQSYDFPLT	68
SB-40	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69
SB-41	QASQDITNYLN	12	DASNLET	26	QQADNFPYT	70
SB-42	RASQSISSYLN	7	SASSLQS	28	QQGDSFPIT	71
SB-43	RASQSVSSSYLA	2	DASKRAT	29	QQRFDLPIT	72
SB-44	RASQSIGSWLA	13	KASSLES	30	QEYGSYRT	73
SB-45	RASQSVSSFLA	10	GASSRAT	16	QQVVSVPPT	74
SB-46	RASQSVRSSSYLA	14	GASSRAT	16	QQLYSSPYT	75
SB-47	QASQDISNYLN	11	DASNLET	26	QQADYFPIT	76
SB-48	RASQGIDSWLA	15	AASSLQS	23	QQASNFPIT	77
SB-49	QASQDITNYLN	12	DASNLET	26	QQYFHPPLT	78
SB-50	KSSQSVLFSSNNKNYLA	3	WASTRES	18	QQFLHTPRT	79

Table 4: EU or Kabat heavy chain HVR sequences of anti-SIRPβ1 antibodies

Ab ID	HVR H1	SEQ ID NO	HVR H2	SEQ ID NO	HVR H3	SEQ ID NO
SB-1	SYAMS	80	TISGSGGSTYYADSVKG	101	DFTEVVGWLGMDV	125
SB-2	SYGMN	81	VIWYDGSNKYYADSVKG	102	DQTAAAIWGMMDV	126
SB-3	GYMH	82	WINPSSGGTNYAQKFQG	103	EGIAATDAYFDL	127
SB-4	SYGIS	83	WISAYNGNTNYAQKLQG	104	SGTHFGTYSYSNWFD	128
SB-5	GYMH	82	WINPNSGGTNYAQKFQG	105	EGDEDWFD	129
SB-6	SYAMS	80	TISGSGGSTYYADSVKG	101	DFTEVVGWLGMDV	125
SB-7	SYAIS	84	GIPIFGTASYAQKFQG	106	ETRQDSAHYYGMDV	130
SB-8	SGYYWG	85	SIYHSGSTYYNPSLKS	107	GGAMTPAGMDV	131
SB-9	SYGIH	86	WISAYNGNTNYAQKLQG	104	DGLHYGDYIVYYGMDV	132
SB-10	SYAIS	84	GIPIFGTANYAQKFQG	108	GVPRGDLGMDV	133
SB-11	NYAIS	87	GIPIFGTANYAQKFQG	108	PVDSSSYSLGYYGMDV	134
SB-12	GYMH	82	WINPNSGGTNYAQKFQG	109	DTYAYSYGMDV	135
SB-13	YYYWS	88	SIYYSGSTNYNPSLKS	110	GDTSGGAYFDL	136
SB-14	SYAIS	84	GIPIFGTASYAQKFQG	106	DRGGVGFY	137
SB-15	SNSYYWG	89	SIYYSGSTYYNPSLKS	111	EVGAPPSYFPI	138
SB-16	SYAIS	84	SIPIFGTANYAQKFQG	112	ANYDSSGYSLDL	139
SB-17	SYGIS	83	WISAYNGNTNYAQKLQG	104	GPLYGDYHVRYGMDV	140
SB-18	SYAIS	84	GIPIFGTANYAQKFQG	108	AKPRGDYGMMDV	141
SB-19	SYAIS	84	GIPIFGTANYAQKFQG	108	DGGGGYAYEYFQH	142
SB-20	SYAIS	84	SIPIFGTANYAQKFQG	112	DGREYGGHYGMDV	143
SB-21	SNGIS	90	WISAYNGNTNYAQKLQG	104	VGNMDQEYFDL	144
SB-22	SNYMS	91	VIYSDGSTYYADSVKG	113	PTRYGYDRLGMDV	145
SB-23	SYAIS	84	GIPIFGTANYAQKFQG	114	TTYRDYYMDV	146
SB-24	SGYYWA	92	SIYHSGSTYYNPSLKS	107	DRSRGYPVYGMDV	147
SB-25	SLAIS	93	GIPIFGTANYAQKFQG	108	SGGDYSGYDYASGMDV	148
SB-26	SYAIS	84	GIPIFGTANYAQKFQG	108	DGSAGRQEHGMDV	149
SB-27	SYAIS	84	GIPIFGTANYAQKFQG	108	QDLGSSSHWHFDL	150
SB-28	SSSYYWG	94	SISYSGSTYYNPSLKS	115	DPRDYSSGSSGGGWGYF DL	151
SB-29	SYAIS	84	SIPIFGTANYAQKFQG	112	APYGSSSGYGYFDL	152
SB-30	SGYYWS	95	YIYYSGSTVYNPSLKS	116	EGPGPSYFDP	153
SB-31	SYMH	96	IINPGGGSTSYAQKFQG	117	EGLYSSGWYIDV	154
SB-32	YYYWS	88	YIYSSGSTNYNPSLKS	118	GDSSSGGLDL	155
SB-33	SNYMS	91	VIYSGGSTYYADSVKG	119	GQYTGSLDV	156
SB-34	GYMH	82	WINPNSGGTKYAQKFQG	120	DTYYTPYGMDV	157
SB-35	SYAIS	84	GIPIFGTANYAQKFQG	108	GRPQSESYLLDY	158
SB-36	SYMH	96	IINPSGGSTSYAQKFQG	121	EGPEQLWYLDY	159
SB-37	NYAIS	87	GIPIFGTANYAQKFQG	108	SRWGASGYYYMDV	160
SB-38	SYMH	96	IINPSGGSTSYAQKFQG	121	ESGTDFTISY	161
SB-39	SGYYWA	92	SIYHSGSTYYNPSLKS	107	GGSNYGDYGRFDY	162
SB-40	SYMS	97	IINPSGGSTSYAQKFQG	121	DTGEYSYSPHGMDV	163
SB-41	SSSYYWG	94	SIYYSGSTYYNPSLKS	111	VGQYPIYGMDV	164
SB-42	SYGIS	83	WISAYNGNTNYAQKLQG	104	GPGHYVAGMDV	165
SB-43	SGYYWA	92	SIYHSGSTYYNPSLKS	107	DAPGYPMLGMDV	166
SB-44	SNYMS	91	VIYSGDITYADSVKG	122	EGSSFWSGSAVSYYGM DV	167
SB-45	SGYYWA	92	SIYHSGSTYYNPSLKS	107	DLSRGYAVSGMDV	168
SB-46	SYAMS	80	AISGSGGSTYYADSVKG	123	ASPWELDV	169
SB-47	SSSYAWG	98	SIYYSGSTYYNPSLKS	111	DLGHYDYWSGSRDYYY GMDV	170
SB-48	SYGMH	99	VISYDGSNKYYADSVKG	124	DGTIAAAGWPPEYFQH	171
SB-49	SSDYYWG	100	SIYYSGSTYYNPSLKS	111	GPTGYKDKWRYYYGM DV	172
SB-50	SYAIS	84	GIPIFGTANYAQKFQG	108	EGGGHASYYHYGMDV	173

Table 5: EU or Kabat light chain Framework sequences of anti-SIRPβ1 antibodies

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
SB-1	EIVMTQSP GTLSLSPG ERATLSC	174	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-2	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-3	EIVMTQSP ATLSLSPG ERATLSC	176	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISRLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-4	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-5	EIVLTQSPA TSLSPGER ATITC	177	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISRLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-6	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-7	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-8	EIVMTQSP GTLSLSPG ERATLSC	174	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-9	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-10	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGVYYC	195	FGGGTKVEIK	200
SB-11	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGVYYC	195	FGGGTKVEIK	200
SB-12	EIVMTQSP ATLSVSPG ERATLSC	180	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TEFTLTISLQS EDFAVYYC	196	FGGGTKVEIK	200
SB-13	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQVLIY	189	GVPDRFSGSGS GTDFTLKISRVE AEDVGVYYC	195	FGGGTKVEIK	200
SB-14	EIVMTQSP ATLSLSPG ERATLSC	176	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISRLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-15	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-16	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-17	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-18	EIVLTQSPA TSLSPGER ATLSC	182	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGVYYC	195	FGGGTKVEIK	200
SB-19	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGVYYC	195	FGGGTKVEIK	200

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
SB-20	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISSLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-21	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGYYC	195	FGGGTKVEIK	200
SB-22	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-23	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-24	EIVMTQSP ATLSVSPG ERATLSC	180	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TEFTLTISSLQS EDFAVYYC	196	FGGGTKVEIK	200
SB-25	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-26	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGYYC	195	FGGGTKVEIK	200
SB-27	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISSLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-28	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-29	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIF	191	GVPDRFSGSGS GTDFTLKISRVE AEDVGYYC	195	FGGGTKVEIK	200
SB-30	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-31	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-32	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISSLQ AEDVAVYYC	194	FGGGTKVEIK	200
SB-33	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTFTISSLQ PEDIATYYC	198	FGGGTKVEIK	200
SB-34	EIVMTQSP ATLSVSPG ERATLSC	180	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TEFTLTISSLQS EDFAVYYC	196	FGGGTKVEIK	200
SB-35	DIVMTQSP LSPVTPG EPASISC	179	WYLQKPGQ SPQLLIY	188	GVPDRFSGSGS GTDFTLKISRVE AEDVGYYC	195	FGGGTKVEIK	200
SB-36	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-37	EIVLTQSPA TSLSPGER ATLSC	182	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-38	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISSLQ PEDFATYYC	197	FGGGTKVEIK	200

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
SB-39	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-40	EIVLTQSPA TSLSPGER ATLSC	182	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISLLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-41	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTFTISLQ PEDIATYYC	198	FGGGTKVEIK	200
SB-42	EIVMTQSP ATLSVSPG ERATITC	184	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-43	EIVLTQSPA TSLSPGER ATLSC	182	WYQQKPGQ APRLLIY	186	GIPARFSGSGSG TDFTLTISLLEP EDFAVYYC	193	FGGGTKVEIK	200
SB-44	DIQMTQSP STLSASVG DRVITIC	185	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTEFTLTISLQ PDDFATYYC	199	FGGGTKVEIK	200
SB-45	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-46	EIVLTQSPG TSLSPGER ATLSC	175	WYQQKPGQ APRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKVEIK	200
SB-47	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTFTISLQ PEDIATYYC	198	FGGGTKVEIK	200
SB-48	DIQMTQSP SSVSASVG DRVITIC	183	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTLTISLQ PEDFATYYC	197	FGGGTKVEIK	200
SB-49	DIQMTQSP SSLSASVG DRVITIC	181	WYQQKPGK APKLLIY	190	GVPSRFSGSGS GTDFTFTISLQ PEDIATYYC	198	FGGGTKVEIK	200
SB-50	DIVMTQSP DSLAVSLG ERATINC	178	WYQQKPGQ PPKLLIY	187	GVPDRFSGSGS GTDFTLTISLQ AEDVAVYYC	194	FGGGTKVEIK	200

Table 6: EU or Kabat heavy chain Framework sequences of anti-SIRPβ1 antibodies

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-1	EVQLLESGGGL VQPGGSLRLSC AASGFTFS	201	WVRQAPGKG LEWVS	210	RFTISRDNKNT LYLQMNSLRAE DTAVYYCAK	215	WGQGTTV TVSS	222
SB-2	QVQLVESGGG VVQPGRSLRLS CAASGFTFS	202	WVRQAPGKG LEWVA	211	RFTISRDNKNT LYLQMNSLRAE DTAVYYCAR	216	WGQGTTV TVSS	222
SB-3	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTRDTSIST AYMELSLRSD DTAVYYCAR	217	WGRGTLV TVSS	223
SB-4	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTSTST AYMELSLRSD DTAVYYCAR	218	WGQGTLV TVSS	224
SB-5	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTRDTSIST AYMELSLRSD DTAVYYCAR	217	WGQGTLV TVSS	224

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-6	EVQLLES GGGL VQPGSLRLSC AASGFTFS	201	WVRQAPGKG LEWVS	210	RFTISRDN SKNT LYLQMNSLRAE DTAVYYCAK	215	WGQGTTV TVSS	222
SB-7	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-8	QVQLQESGPG L VKPSETLSLTC AVSGYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTTV TVSS	222
SB-9	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTSTST AYMELRSLRSD DTAVYYCAR	218	WGQGTTV TVSS	222
SB-10	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-11	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGKGTTV TVSS	225
SB-12	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTRDTSIST AYMELRSLRSD DTAVYYCAR	217	WGQGTTV TVSS	222
SB-13	QVQLQESGPG L VKPSETLSLTC TVSGGIS	206	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGRGTLV TVSS	223
SB-14	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTLV TVSS	224
SB-15	QLQLQESGPG L VKPSETLSLTC TVSGGIS	207	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTMT TVSS	226
SB-16	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGRGTLV TVSS	223
SB-17	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTSTST AYMELRSLRSD DTAVYYCAR	218	WGQGTTV TVSS	222
SB-18	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-19	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTLV TVSS	224
SB-20	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-21	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTSTST AYMELRSLRSD DTAVYYCAR	218	WGRGTLV TVSS	223
SB-22	EVQLVESGGGL VQPGSLRLSC AASGFTVS	208	WVRQAPGKG LEWVS	210	RFTISRDN SKNT LYLQMNSLRAE DTAVYYCAR	216	WGQGTTV TVSS	222
SB-23	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGKGTTV TVSS	225
SB-24	QVQLQESGPG L VKPSETLSLTC AVSGYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTTV TVSS	222

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-25	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-26	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTTV TVSS	222
SB-27	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGRGTLV TVSS	223
SB-28	QLQLQESGPG L VKPSETLSLTC TVSGGSIS	207	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGRGTLV TVSS	223
SB-29	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGRGTLV TVSS	223
SB-30	QVQLQESGPG L VKPSQTLSTC TVSGGSIS	209	WIRQHPGKG LEWIG	214	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTLLV TVSS	224
SB-31	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTRDTSTST VYMELSSLRSED TAVYYCAR	221	WGQGTLLV TVSS	224
SB-32	QVQLQESGPG L VKPSETLSLTC TVSGGSIS	206	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGRGTLV TVSS	223
SB-33	EVQLVESGGGL VQPGGSLRLSC AASGFTVS	208	WVRQAPGKG LEWVS	210	RFTISRDNKNT LYLQMNSLRAE DTAVYYCAR	216	WGQGTMT VTVSS	226
SB-34	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTRDTSIST AYMELSLRSD DTAVYYCAR	217	WGQGTTV TVSS	222
SB-35	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTLLV TVSS	224
SB-36	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTRDTSTST VYMELSSLRSED TAVYYCAR	221	WGQGTLLV TVSS	224
SB-37	QVQLVQSGAE VKKPGSSVKVS CKASGGTFS	204	WVRQAPGQG LEWMG	212	RVTITADESTST AYMELSSLRSED TAVYYCAR	219	WGQGTMT VTVSS	226
SB-38	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTRDTSTST VYMELSSLRSED TAVYYCAR	221	WGQGTLLV TVSS	224
SB-39	QVQLQESGPG L VKPSETLSLTC AVSGYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTLLV TVSS	224
SB-40	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTRDTSTST VYMELSSLRSED TAVYYCAR	221	WGQGTTV TVSS	222
SB-41	QLQLQESGPG L VKPSETLSLTC TVSGGSIS	207	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTTV TVSS	222
SB-42	QVQLVQSGAE VKKPGASVKV SCKASGYTFT	203	WVRQAPGQG LEWMG	212	RVTIMTDTSTST AYMELSLRSD DTAVYYCAR	218	WGQGTTV TVSS	222
SB-43	QVQLQESGPG L VKPSETLSLTC AVSGYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKNQ FSLKLSSVTAAD TAVYYCAR	220	WGQGTTV SVSS	227

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-44	EVQLVESGGGLVQPGGSLRLSCAASGFTVS	208	WVRQAPGKGLEWVS	210	RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR	216	WGQGTTVTVSS	222
SB-45	QVQLQESGPGLVKPSETLSLTCAVSGYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKNQFSLKLSSVTAADTAVYYCAR	220	WGQGTTVTVSS	222
SB-46	EVQLLES GGGLVQPGGSLRLSCAASGFTFS	201	WVRQAPGKGLEWVS	210	RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR	216	WGQGTMTVTVSS	226
SB-47	QLQLQESGPGLVKPSETLSLTCTVSGGSIS	207	WIRQPPGKGL EWIG	213	RVTISVDTSKNQFSLKLSSVTAADTAVYYCAR	220	WGQGTTVTVSS	222
SB-48	QVQLVESGGGVVQPGRLRLSCAASGFTFS	202	WVRQAPGKGLEWVA	211	RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR	216	WGQGTLLVTVSS	224
SB-49	QLQLQESGPGLVKPSETLSLTCTVSGGSIS	207	WIRQPPGKGL EWIG	213	RVTISVDTSKNQFSLKLSSVTAADTAVYYCAR	220	WGQGTTVTVSS	222
SB-50	QVQLVQSGAEVKKPGSSVKVSCKASGGTFS	204	WVRQAPGQGLEWMG	212	RVTITADESTSTAYMELSSLRSEDTAVYYCAR	219	WGQGTTVTVSS	222

Characterization of SIRPβ1 antibody binding

[0251] Initial characterization of anti-SIRPβ1 antibodies involved determining their ability to bind cell lines expressing human or mouse SIRPβ1. Cells were harvested, plated at 10⁶/ml in a 96-well plate, washed, and incubated in 100 μl FACS buffer containing 1 μg/ml anti-SIRPβ1 antibody for 0.5 hour on ice. Cells were then washed twice and incubated in 100 μl FACS buffer containing 0.5 μg/ml PE-conjugated secondary antibody for 30 minutes on ice. Cells were washed twice in cold FACS buffer and acquired on a BD FACS Canto. Data analysis and calculation of mean fluorescence intensity (MFI) values or % positive cells was performed with FlowJo (TreeStar) software version 10.0.7.

[0252] **Table 7** shows the mean fluorescence intensity (MFI) values of anti-SIRPβ1 antibodies binding to a Chinese hamster ovary (CHO) cell line expressing low levels of recombinant human SIRPβ1. The human IgG1 isotype control established the background fluorescence signal set to 1. Of the 50 anti-SIRPβ1 antibody clones tested, 32 clones bound to cells with an MFI ≥ 2-fold over background. As a negative control, the anti-SIRPβ1 antibodies were also screened for surface binding to CHO cells overexpressing recombinant mouse SIRPβ1. As expected, none of the test antibodies bound to mouse SIRPβ1. Given the high sequence similarity between receptors of the SIRP family, anti-SIRPβ1 antibodies were also screened for cross-reactivity to human and mouse SIRPα. In cell binding assays, none of the anti-SIRPβ1 antibodies bound cells overexpressing human or mouse SIRPα.

Table 7: Cell Binding Characterization of anti-SIRP β 1 antibodies

Clone Index (SB-#)	Cell Binding Human SIRP β FOB (Fold Over Background)	Cell Binding Human SIRP α FOB (Fold Over Background)	Cell Binding Mouse SIRP β FOB (Fold Over Background)	Cell Binding Mouse SIRP α FOB (Fold Over Background)
1*	5.7	2.3	1.1	1.4
2*	8.7	3.0	1.3	1.1
3	2.5	1.8	1.3	1.3
4	1.7	2.0	1.2	0.9
5	1.6	2.0	1.3	1.5
6	4.3	1.2	1.2	1.0
7	3.7	2.0	1.5	1.0
8*	4.1	2.1	1.7	1.2
9	1.8	2.1	1.0	0.8
10	3.8	2.4	1.4	1.0
11	6.3	2.3	1.2	0.9
12	1.6	2.0	1.3	1.2
13	1.9	1.9	1.2	0.9
14	4.5	2.5	1.3	1.2
15	8.0	2.2	1.3	1.4
16	2.0	2.0	1.5	1.4
17	2.8	2.5	1.3	1.6
18	2.8	2.4	1.2	0.9
19	1.8	2.2	1.3	0.9
20	2.6	2.2	1.3	1.2
21	1.9	2.0	1.3	1.2
22	1.4	1.9	1.4	0.9
23	2.4	1.9	1.5	1.4
24	3.7	2.5	1.3	1.1
25	2.1	2.5	1.2	1.3
26	2.0	2.1	1.3	1.2
27	3.9	2.4	1.3	1.1
28	4.1	2.1	1.2	1.0
29	1.8	1.9	1.2	0.9
30	1.7	2.1	1.4	1.5
31	3.1	1.8	1.4	1.0
32	1.5	2.2	1.2	1.2
33	1.4	2.2	1.3	1.4
34	1.8	2.0	1.2	1.3
35	2.1	2.6	1.1	1.0
36	1.7	2.1	1.1	1.2
37	3.8	2.2	1.2	1.0
38	1.5	2.1	1.3	1.1
39	5.0	2.0	1.4	1.1
40*	7.0	2.5	1.1	1.2

41	1.7	2.2	1.3	1.7
42	3.3	2.7	1.2	1.2
43	2.8	2.3	1.1	1.6
44	2.8	2.9	1.5	1.2
45	8.5	2.5	1.4	1.2
46	5.3	2.1	1.3	1.4
47	1.6	2.1	1.3	1.7
48	1.7	2.1	1.1	2.4
49	3.4	2.7	1.2	1.7
50	2.8	2.5	1.2	1.7

[0253] Additionally, anti-SIRP β 1 antibodies were also screened for antigen specificity by using a reporter cell line expressing the luciferase gene under the control of an NFAT (nuclear factor of activated T-cells) promoter. The cell line BW5147.G.1.4 (ATCC® TIB48™), derived from mouse thymus lymphoma T lymphocytes, was infected with Cignal Lenti NFAT-luciferase virus (Qiagen), resulting in BWZ/NFAT-luciferase reporter cells. Subsequently, cells were transduced with either a lentivirus expressing human SIRP α -DAP12 chimera, in which the intracellular ITIM motif of SIRP α was substituted with the intracellular ITAM motif of DAP12, or with two lentiviruses expressing human SIRP β 1 and human DAP12. Test antibodies, as well as the human IgG1 isotype control, were adsorbed onto a 96-well plate at 10 μ g/mL. After washing, 10⁵ NFAT-luciferase reporter cells expressing the huSIRP α /DAP12 chimera (BWZ-huSIRP α) or co-expressing huSIRP β 1 and DAP12 (BWZ-huSIRP β 1) were seeded onto plates and incubated overnight at 37°C. Luciferase activity was measured by adding OneGlo Reagent (Promega) to each well and incubating the samples for 3 min at room temperature on a plate shaker. The luminescence signal was quantified using a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software. As shown in **FIG. 3A**, 48 out of 50 anti-SIRP β 1 clones induced luciferase expression >5-fold over background in BWZ-huSIRP β 1 reporter cells. In contrast, anti-SIRP β 1 antibodies failed to induce luciferase expression when BWZ-huSIRP α reporter cells were added onto antibody-coated wells (**FIG. 3B**). In summary, only reporter cells expressing human SIRP β 1 induced luciferase expression in the presence of immobilized anti-SIRP β 1 antibodies, as measured by luminescence signal. These results establish that most anti-SIRP β 1 antibodies capable of binding membrane-bound antigen demonstrate specificity towards the target antigen without cross-reacting to other SIRP receptors.

Example 2: SIRP β 1 expression profile on myeloid cells

[0254] The SIRP family comprises several transmembrane glycoproteins primarily expressed within the myeloid cell compartment. The expression pattern of SIRP β 1 was verified on primary human cells isolated from healthy human donors. Human primary monocytes were isolated from

heparinized human peripheral blood obtained from two healthy donors (Blood Centers of the Pacific) using RosetteSep Human Monocyte Enrichment Cocktail (STEMCELL Technologies), according to the manufacturer's protocol. Both SIRP β 1 and TREM1 are preferentially expressed on CD14-high monocytes. Monocytes were seeded in RPMI (Invitrogen) containing 10% Fetal Calf Serum (Hyclone) and 50 ng/ml M-CSF or GM-CSF (Peprotech) to induce differentiation of M2-like or M1-like macrophages, respectively. After 5-6 days, macrophages were harvested by scraping cells attached to plastic. Alternatively, monocytes were seeded in RPMI medium containing 10% Fetal Calf Serum (Hyclone) and 20 ng/ml IL-4 and GM-CSF (Peprotech) to induce differentiation of immature dendritic cells. After 6-7 days, dendritic cells were harvested by scraping cells attached to plastic.

[0255] As shown in **FIG. 4A**, CD14 expression defines a subset of primary human monocytes. Classical and intermediate monocytes express high levels of CD14, whereas, non-classical monocytes lack CD14 expression. Like TREM1, SIRP β 1 is abundantly expressed in CD14+ monocytes. Monocytes isolated from peripheral blood as described above were cultured for 5 days with M-CSF or GM-CSF to generate M2 or M1 macrophages, respectively. SIRP β 1 expression decreases upon differentiation of primary monocytes into macrophages with M1-like macrophages expressing higher levels of the receptor relative to M2-like macrophages (**FIG. 4B**). Similarly, SIRP β 1 expression decreases upon differentiation of primary monocytes into immature dendritic cells. LPS-induced maturation of dendritic cells further downregulates SIRP β 1 expression, in contrast to the increased expression observed for TREM1 (**FIG. 4C**). Unlike TREM1, LPS treatment downregulates SIRP β 1 expression on DCs. Peak expression for SIRP β 1 occurs in blood, spleen, and lung. Surveying SIRP β 1 expression pattern from RNA-seq data sets from multiple tissue samples gathered from healthy donors confirms that SIRP β 1 transcript levels predominate in the blood, consistent with high expression of the receptor in circulating monocytes and neutrophils (**FIG. 5A**). However, RNA-seq data reveals that the expression pattern of SIRP β 1 isoform 3 transcript differs from that of SIRP β 1 isoform 1; SIRP β 1 isoform 3 is mostly expressed in the brain and not peripheral tissues (**FIG. 5B**). (Data were obtained from Genotype-Tissue Expression (GTEx) Consortium database.)

Example 3: SIRP β 1 antibodies induce Syk phosphorylation

[0256] Spleen tyrosine kinase (Syk) is an intracellular signaling molecule that functions downstream of SIRP β 1 by phosphorylating several substrates, thereby facilitating the formation of a signaling complex leading to cellular activation and inflammatory processes. The ability of agonist SIRP β 1 antibodies to induce Syk activation is determined by culturing mouse monocytes and measuring the phosphorylation state of Syk protein in cell extracts. In these experiments, a secondary antibody is used to cross-link anti-SIRP β 1 antibodies on cells to induce intracellular signaling.

[0257] Bone marrow-derived monocytes (BMDM) from wild-type (WT) mice, from human SIRP β 1 BAC transgenic mice, and from mice that lack expression of functional Fc receptor common gamma chain gene (Fc γ R KO; REF: Takai T 1994. Cell 76(3):519-29) are starved for 4 hours in 1% serum RPMI and then removed from tissue culture dishes with PBS-EDTA, washed with PBS, and counted. The cells are coated with full-length SIRP β 1 antibodies or with huIgG1 isotype control for 15 minutes on ice. After washing with cold PBS, cells are incubated at 37°C for the indicated period of time in the presence of goat anti-human IgG. After stimulation, cells are lysed with lysis buffer (1% v/v NP-40%, 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM EDTA, 1.5 mM MgCl₂, 10% glycerol, plus protease and phosphatase inhibitors) followed by centrifugation at 16,000 g for 10 min at 4°C to remove insoluble materials. Lysates are then immunoprecipitated with anti-Syk antibody (N-19 for BMDM or 4D10 for human DCs, Santa Cruz Biotechnology). Precipitated proteins are fractionated by SDS-PAGE, transferred to PVDF membranes and probed with anti-phosphotyrosine antibody (4G10, Millipore). To confirm that all substrates are adequately immunoprecipitated, immunoblots are reprobed with anti-Syk antibody (Abcam, for BMDM) or anti-Syk (Novus Biological, for human DCs). Visualization is performed with the enhanced chemiluminescence (ECL) system (GE healthcare), as described (*e.g.*, Peng *et al.*, (2010) Sci Signal., 3(122): ra38). These studies provide support that anti-SIRP β 1 antibodies of the present invention induce syk phosphorylation.

Example 4: SIRP β 1 antibodies induce Syk phosphorylation when clustered by adjacent cells that expresses Fc gamma receptors.

[0258] Activation of spleen tyrosine kinase (Syk) is facilitated by crosslinking two or more SIRP β 1 receptors with antibodies, thereby facilitating the formation of a signaling complex leading to cellular activation and inflammatory processes. *In vivo* cross-linking is mediated by adjacent cells that express Fc receptors (FcR), such as B cells and other leukocytes (White AL Cancer Immunol Immunother (2013) 62:941–948; Wilson NS 2011, Cancer Cell 19, 101–113; Bartholomaeus P J Immunol 2014; 192:2091-2098). In these experiments, accessory cells expressing Fc gamma receptors (*i.e.*, B cells) are used to cross-link anti-SIRP β 1 antibodies to induce intracellular signaling.

[0259] The ability of Fc receptors to induce activation of Syk through antibody clustering is determined by culturing mouse monocytes in the presence of cells expressing Fc receptors and measuring the phosphorylation state of Syk protein in cell extracts. Bone marrow-derived monocytes (BMDM) from wild-type (WT) mice and human SIRP β 1 BAC transgenic mice are starved for 4 hours in 1% serum RPMI and then removed from tissue culture dishes with PBS-EDTA, washed with PBS, and counted. The cells are coated with full-length SIRP β 1 antibodies, or huIgG1 isotype control for 15 minutes on ice. After washing with cold PBS, cells are incubated for 5 minutes at

37°C with glutaraldehyde-fixed cells that express Fc receptors and that are previously prepared as follows. Briefly, Fc receptor expressing cells are either B cells isolated from mouse spleens using MACS microbeads (CD19⁺ B-cell isolation kit Miltenyi Biotec) according to the manufacturer's protocol, or alternatively, the P815 cell line that overexpresses FcR2b and FcR3. 2×10^6 cells/ml cells are fixed with 0.05% glutaraldehyde for 1 minute at room temperature, the reaction is stopped with 1 μM Glycine and cells are then washed extensively with PBS. After stimulation, cells are lysed with lysis buffer (1% v/v NP-40%, 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM EDTA, 1.5 mM MgCl₂, 10% glycerol, plus protease and phosphatase inhibitors) followed by centrifugation at 16,000 g for 10 min at 4°C to remove insoluble materials. Lysates are then immunoprecipitated with anti-Syk antibody (N-19 for BMDM or 4D10 for human DCs, Santa Cruz Biotechnology). Precipitated proteins are fractionated by SDS-PAGE, transferred to PVDF membranes and probed with anti-phosphotyrosine antibody (4G10, Millipore). To confirm that all substrates are adequately immunoprecipitated, immunoblots are reprobed with anti-Syk antibody (Abcam, for BMDM) or anti-Syk (Novus Biological, for human DCs). Visualization is performed with the enhanced chemiluminescence (ECL) system (GE healthcare), as described (*e.g.*, Peng *et al.*, (2010) *Sci Signal.*, 3(122): ra38). These studies provide support that anti-SIRPβ1 antibodies of the present invention induce syk phosphorylation when clustered by adjacent cells expressing Fc gamma receptors.

Example 5: SIRPβ1 antibodies induce respiratory burst in immune cells

[0260] The agonistic function of SIRPβ1 antibodies of the present disclosure was evaluated in primary human innate immune cells (*e.g.*, monocytes and neutrophils).

[0261] Anti-SIRPβ1 and isotype control antibodies were diluted in serum-free RPMI media and mixed with 100,000 neutrophils on 96-well plates at a concentration of 10 μg/ml. Primary neutrophils were isolated with EasySep™ Direct Human Neutrophil Isolation Kit (STEMCELL) according to the manufacturer's instructions from peripheral blood obtained the same day. To detect the production of reactive oxygen species (ROS), cells were labeled with 2 μM of the fluorescent dye, CM-H₂DCFDA. Cells were stimulated with soluble, full-length human IgG1 isotype control or the anti-SIRPβ1 antibodies SB-1, -2, -3, -4, -5, -6, -7, -8, -9, -11, -14, -15, -17, -27, -28, -31, -39, -40, -41, -45, -46, and -49. Following 1 hour of antibody-mediated stimulation in the presence of CM-H₂DCFDA at 37°C, the relative fluorescence units in cells were measured at excitation wavelength 495 nm and emission wavelength 530 nm. Specific fluorescence index of stimulated cells was obtained by subtraction of background fluorescence of labeled cells incubated in medium alone and/or with isotype control antibody (huIgG1). Plates were read with a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software.

[0262] In **FIG. 6A**, primary human neutrophils from 2 healthy donors were stimulated with 10 $\mu\text{g/mL}$ of soluble full-length anti-SIRP β 1 antibodies or human IgG1 isotype control and labeled with 2 μM CM-H₂DCFDA for 1 hour at 37°C to monitor SIRP β 1-mediated ROS production. The SIRP β 1 antibodies SB-1, -2, -3, and -5 proved highly agonistic in solution. In contrast, the SIRP β 1 antibodies SB-15, -17, -27, -28, -31, -39, -40, -41, -45, -46, and -49 weakly activated SIRP β 1-mediated respiratory burst in solution.

[0263] Cells were left untreated or stimulated with soluble, full-length human IgG1 isotype control or the anti-SIRP β 1 antibodies SB-1, -2, -3, -4, -5, -6, -7, -8, -9, -11, -12, -13, -14, -15, -17, -23, -26, -27, -28, -31, -33, -34, -35, -36, -37, -39, -40, -41, -45, -46, and -49. In **FIG. 6B**, primary human monocytes were stimulated with 20 $\mu\text{g/mL}$ of soluble full-length anti-SIRP β 1 antibodies or human IgG1 isotype control and labeled with 2 μM CM-H₂DCFDA for 1 hour at 37°C to monitor SIRP β 1-mediated ROS production and IL-8 production. The SIRP β 1 antibodies SB-2, -3, -4, -5, -6, -7, -28, and -49 proved highly agonistic in solution. In contrast, the SIRP β 1 antibodies SB-14, -15, -17, -27, -41, -45, and -46 weakly activated SIRP β 1-mediated respiratory burst while in solution. Additionally, primary monocytes were stimulated overnight at 37°C with anti-SIRP β 1 antibodies adsorbed onto 96-well plates at 10 $\mu\text{g/mL}$. The supernatant fraction was subsequently collected and assayed for IL-8 release. The SIRP β 1 antibodies SB-1, -2, -3, -7, -8, -9, -14, 28, and -49 proved highly agonistic as plate-bound antibodies. In contrast, the SIRP β 1 antibodies SB-4, -12, -23, -26, -33, -34, -35, -36, and -37 weakly activated SIRP β 1-mediated cytokine release.

Example 6: SIRP β 1 increases secretion of inflammatory cytokines from macrophages.

[0264] Published literature describe bone marrow-derived macrophages (BMDM) or primary peritoneal macrophage possessing altered TLR responses when deficient in DAP12. To determine whether SIRP β 1 antibodies of the present disclosure induce changes in inflammatory cytokine production, primary human monocyte-derived macrophages and dendritic cells are cultured with plate-bound test antibodies in combination with non-saturating levels of TLR stimulators and the level of cytokines are measured after 24h. To generate monocyte-derived macrophages and dendritic cells, human primary monocytes were isolated from heparinized human blood (Blood Centers of the Pacific) using RosetteSep Human Monocyte Enrichment Cocktail (STEMCELL Technologies), according to the manufacturer's protocol. Monocytes were seeded in RPMI (Invitrogen) containing 10% Fetal Calf Serum (Hyclone) and 50 ng/ml M-CSF to induce macrophage differentiation. After 5-6 days, macrophages were harvested by scraping cells attached to plastic. Alternatively, monocytes were seeded in RPMI medium containing 10% Fetal Calf Serum (Hyclone) and 20 ng/ml IL-4 and GM-CSF (Peprotech) to induce differentiation of immature dendritic cells. After 6-7 days, dendritic cells were harvested by scraping cells attached to plastic. Macrophages or dendritic cells are plated on 96-well plates coated with indicated antibody at

10^5 cells/well and incubated for 24h at 37°C. Cells are co-stimulated with TLR4 agonist, LPS (*Salmonella abortus equi*).

[0265] As shown in **FIG. 6C**, monocyte-derived macrophages and dendritic cells were stimulated overnight at 37°C with 0.5 ng/mL LPS in the presence of anti-SIRPβ1 antibodies (SB-1, -2, -3, -4, -5, -6, -7, -8, -9, -11, -12, -14, -15, -16, -17, -18, -19, -20, -21, -28, -32, -40, and -49) adsorbed onto 96-well plates at 10 μg/mL. The supernatant fraction was subsequently collected and assayed for TNFα release. As a negative control, DCs were stimulated with LPS in the presence of plate-bound anti-SIRPα antibody, SA-56-90, which suppresses TNFα release. In all experiments measuring respiratory burst, production of reactive oxygen species (ROS) was monitored by labeling cells with 2 μM of the fluorescent indicator, CM-H₂DCFDA. The SIRPβ1 antibodies SB-1, -2, -6, -8, and -49 proved highly agonistic as plate-bound antibodies. In contrast, most remaining SIRPβ1 antibodies weakly activated SIRPβ1-mediated cytokine release. Similarly, plate-bound SB-1 and SB-40, but not SB-32, enhanced LPS-induced TNFα release from monocyte-derived dendritic cells. In contrast, LPS-induced TNFα release was decreased in dendritic cells stimulated with plate-bound anti-SIRPα antibody, SA-90, confirming that agonistic anti-SIRPβ1 antibodies activate cells, whereas, agonistic SIRPα antibodies inhibit cellular activity.

Example 7: Anti-tumor activity of SIRPβ1-stimulated neutrophils

[0266] Though neutrophils are efficient phagocytes for antibody- or complement-opsonized target cells, in the context of the tumor microenvironment, neutrophils generally contribute towards tumor progression, invasion, and angiogenesis. However, recent publications suggest that tumor-associated neutrophils, like other myeloid cells, retain the potential to polarize towards an anti-tumor phenotype. Since neutrophils express high levels of SIRPβ1, anti-SIRPβ1 antibodies were evaluated for their ability to induce neutrophil-mediated tumor cell clearance *in vitro*. Primary neutrophils were isolated with EasySep™ Direct Human Neutrophil Isolation Kit (STEMCELL) according to the manufacturer's instructions from peripheral blood of healthy donors obtained the same day. Isolated human neutrophils were then added onto 96-well plates previously coated with 10 μg/mL anti-SIRPβ1 antibodies or isotype control. Subsequently, Raji B cell lymphoma cells engineered to stably express luciferase were mixed with neutrophils in a 1:1 ratio with or without opsonizing antibody (anti-CD20 human IgG1). Co-cultured cells were incubated overnight at 37°C, and viable Raji cells were quantified by measuring luciferase activity following the addition of OneGlo reagent (Promega) and incubating samples at room temperature for 3 min on a plate shaker. The luminescence signal was detected with a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software.

[0267] As shown in **FIG. 7**, human neutrophils failed to eliminate Raji cells when co-cultured in the presence of immobilized isotype control antibody. The addition of anti-CD20 huIgG1 did not

restore tumor cell clearance with unstimulated neutrophils. Similarly, neutrophils stimulated with immobilized anti-SIRP β 1 antibodies failed to significantly eliminate Raji cells. However, SIRP β 1-stimulated neutrophils cleared anti-CD20 opsonized Raji cells. The agonistic anti-SIRP β 1 antibodies SB-1, SB-2, SB-3, and SB-4 reduced luminescence signal from viable Raji cells ~50% compared to isotype control-treated neutrophils. The anti-SIRP β 1 antibodies SB-28 and SB-49 demonstrated weak activity against opsonized Raji cells. These results suggest that agonistic anti-SIRP β 1 antibodies enhance the anti-tumor properties of neutrophils.

Example 8: Affinity maturation of anti-SIRP β 1 antibodies

[0268] Five anti-SIRP β 1 antibodies, SB-1, SB-2, SB-8, and SB-40 (termed “parent” antibodies), with various physical and functional attributes were affinity-matured. Briefly, diversified antibody libraries were created in yeast for each of the starting parent antibodies. The diversity was created by utilizing standard molecular cloning techniques to combine the parental heavy chain CDR-H3 and light chain (LC) with pre-existing genetic diversity in the CDR-H1 and CDR-H2 regions of the heavy chain (HC) (termed “H1/H2” optimization). This resulted in six libraries of roughly 10^5 clones in size; 1:755-768 for selection of antibodies with improved affinity. Selection pressures used for screening the libraries included human SIRP α and SIRP β 1 antigen equilibrium titration, parental antibody Fab competition kinetics, and the use of polyspecificity reagent deselection (as described, for example, in WO 2014/179363; Xu *et al.*, *Protein Eng Des Sel*, 26(10): 663-670). FACS flow cytometry was then employed to visualize and select antibodies, using standard techniques (see, *e.g.*, Chao *et al. Nature Protocols*, 2006;1:755-768). The desired population was then carried forward into additional selection rounds. After 6 rounds of enrichment, yeast cells were plated out in order to obtain single antibody isolates, which were then produced and characterized as described in Example 1. Seventeen affinity-improved antibodies from four of the five starting parental antibodies were thus obtained.

Antibody IgG and Fab production and purification

[0269] Yeast clones were grown to saturation and then induced for 48 h at 30°C with shaking. After induction, yeast cells were pelleted and the supernatants were harvested for purification.

[0270] Immunoglobulins were purified using a Protein A column and eluted with acetic acid, pH 2.0. Fab fragments were generated by papain digestion and purified over CaptureSelect IgG-CHI affinity matrix (LifeTechnologies).

Affinity determination

[0271] The affinities of the anti-SIRP β 1 antibodies were determined by measuring K_D values by ForteBio Octet[®] and Meso Scale Discovery (MSD) instrument. Octet[®] affinity measurements were performed at room temperature, generally as previously described (Estep *et al. MAb*s. 2013 Mar-Apr;5(2):270-8). Briefly, Octet[®] affinity measurements were performed by loading IgGs on-line

onto AHQ sensors. Sensors were equilibrated off-line in assay buffer for 30 min and then monitored on-line for 60 seconds for baseline establishment. For avid binding measurement, sensors with loaded IgGs were exposed to 100 nM antigen (human SIRP α or SIRP β 1 Fc fusion) for 3 min, and then transferred to assay buffer for 3 min for off-rate measurement. Additional avid binding was determined by loading biotinylated SIRP β 1 monomer on SA sensors and exposing to 100 nM IgG in solution. Monovalent binding measurements were obtained by loading human SIRP β 1 Fc fusion antigens to AHQ sensor, followed by exposure to 100 nM anti-SIRP β 1 antibody Fab. Additional monovalent measurements were made by loading biotinylated human SIRP β 1 monomer to SA sensor followed by exposure to 100 nM Fab in solution. Kinetics data were fit using a 1:1 binding model in the data analysis software provided by ForteBio.

[0272] For MSD-SET K_D measurements, solution equilibrium titrations (SET) were performed in PBS + 0.1% IgG-Free BSA (PBSF) with recombinant human SIRP β 1, held constant at 100 pM and incubated with 3-to 5-fold serial dilutions of antibody starting at around 50 nM. Antibodies (20 nM in PBS) were coated onto standard bind MSD-ECL plates overnight at 4°C or at room temperature for 30 min. Plates were then blocked with 1% BSA for 30 min with shaking at 700 rpm, followed by three washes with wash buffer (PBSF + 0.05% Tween 20). SET samples were applied and incubated on the plates for 150s with shaking at 700 rpm followed by one wash. Antigen captured on a plate was detected with 250 ng/ml sulfotag-labeled streptavidin in PBSF by incubation on the plate for 3 min. The plates were washed three times with wash buffer and then read on the MSD Sector Imager 2400 instrument using 1x Read Buffer T with surfactant. The percent free antigen was plotted as a function of titrated antibody in Prism and fit to a quadratic equation to extract the K_D . To improve throughput, liquid handling robots were used throughout MSD-SET experiments, including SET sample preparation.

[0273] Cell binding affinity measurements were performed at 4°C using BWZ reporter cells either expressing either human SIRP α or SIRP β 1. Briefly, cells were harvested, washed in PBS and incubated with increasing concentration of anti-SIRP β 1 antibodies or isotype control. Antibodies were diluted in FACS buffer (PBS + 2% FBS). After incubation on ice for 30 min, cells were washed two times in FACS buffer and incubated with anti-human PE conjugated secondary antibody (BD Biosciences) for 30 min on ice. Cells were then washed twice in 200ul FACS buffer, and subsequently analyzed on a FACS Canto screening instrument (BD). Apparent K_D values were determined by non-linear curve fitting (modified OneSiteTotal, Graph Pad Prism).

Anti-SIRP β 1 antibody selection

[0274] Affinity-matured anti-SIRP β 1 antibody clones, which showed improved affinity compared to the respective parental antibody, were characterized further. After initial screening of all affinity-matured antibody clones, clones for each parental antibody were selected for further analysis.

Antibody heavy chain and light chain variable domain sequences

[0275] Using standard techniques, the amino acid sequences encoding the light chain variable and the heavy chain variable domains of the affinity matured antibodies were determined. The Kabat light chain HVR sequences of the affinity matured antibodies are set forth in Table 8. The Kabat heavy chain HVR sequences of the antibodies are set forth in Table 9. The Kabat heavy chain framework (FR) sequences of the antibodies are set forth in Table 10. The Kabat light chain framework (FR) sequences of the antibodies are set forth in Table 11.

Table 8: Kabat light chain HVR sequences of affinity matured anti-SIRPβ1 antibodies

Ab ID	HVR L1	SEQ ID NO	HVR L2	SEQ ID NO	HVR L3	SEQ ID NO
SB-1 (p)	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-1-2	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-1-3	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-1-4	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-1-5	RASQSVSSSYLA	383	GASSRAT	16	QLLGSSPRT	31
SB-2 (p)	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-2-7	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-2-8	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-2-9	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-2-10	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-2-11	RASQSVSSSYLA	383	GASSRAT	16	QQSSHPFT	32
SB-8 (p)	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-8-13	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-8-14	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-8-15	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-8-16	RASQSVSSSYLA	383	GASNRAT	19	QQVYSSPYT	37
SB-40 (p)	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69
SB-40-18	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69
SB-40-19	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69
SB-40-20	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69
SB-40-21	RASQSVSSSYLA	2	DSSNRAT	22	QQRDEHPPWT	69

(p) denotes parental antibody sequence

Table 9: Kabat heavy chain HVR sequences of affinity matured anti-SIRPβ1 antibodies

Ab ID	HVR H1	SEQ ID NO	HVR H2	SEQ ID NO	HVR H3	SEQ ID NO
SB-1 (p)	SYAMS	80	TISGSGGSTYYADSVKG	101	DFTEVVGWLGMDV	125
SB-1-2	SFGMN	228	AITASGGSTFYADSVKG	238	DFTEVVGWLGMDV	125
SB-1-3	AYGMN	229	AITSSGRSTYYADSVKG	239	DFTEVVGWLGMDV	125
SB-1-4	AYGMN	229	AIRASGGATYYADSVKG	240	DFTEVVGWLGMDV	125
SB-1-5	AYGMN	229	AISASGRSTFYADSVKG	241	DFTEVVGWLGMDV	125
SB-2 (p)	SYGMN	81	VIWYDGSNKYYADSVKG	102	DQTAAAIWGMMDV	126
SB-2-7	RYGMH	230	AISGLAGPT-YADSVKG	242	DQTAAAIWGMMDV	126
SB-2-8	DYGMH	231	AISAFAGST-YADSVKG	243	DQTAAAIWGMMDV	126
SB-2-9	TYGMH	232	HIWYEGSNKVYADSVKG	244	DQTAAAIWGMMDV	126
SB-2-10	SYGMH	99	AISGLAGQT-YADSVKG	245	DQTAAAIWGMMDV	126
SB-2-11	RYGMH	230	AISGLAGPT-YADSVKG	242	DQTAAWGIWGMMDV	253
SB-8 (p)	SGYYWG	85	SIYHSGSTYYNPSLKS	107	GGAMTPAGMDV	131
SB-8-13	AHYYWG	233	SIFHSGHTYYNPSLKS	246	GGAMTPAGMDV	131
SB-8-14	PHYYWG	234	SIYHSGHTYYNPSLKS	247	GGAMTPAGMDV	131
SB-8-15	AHYYWG	233	SIYQSGHTYYNPSLKS	248	GGAMTPAGMDV	131
SB-8-16	AHYYWG	233	SIFHSGHTYYNPSLKS	246	AGAMTPAGMDV	254
SB-40 (p)	SYYMS	97	IINPSGGSTSYAQKFQG	121	DTGEYSYSPHGMDV	163
SB-40-18	SYYMA	235	WINPAVGATIYSQKFQG	249	DTGEYSYSPHGMDV	163
SB-40-19	SYYMV	236	IINPSSGATNYAQKFQG	250	DTGEYSYSPHGMDV	163
SB-40-20	SFYIS	237	IINPSSGHTNYAQKLQG	251	DTGEYSYSPHGMDV	163
SB-40-21	SYYMV	236	IINPSSGDTNYAQKFQG	252	DTGEYSYSPHGMDV	163

(p) denotes parental antibody sequence

Table 10: Kabat light chain FR sequences of affinity matured anti-SIRPβ1 antibodies

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
SB-1 (p)	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-1-2	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-1-3	EIVMT QSPGTL	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
	SLSPGE RATLSC							
SB-1-4	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-1-5	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2 (p)	EIVLTQ SPGTL LSPGER ATLSC	175	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2-7	EVVLT QSPGTL SLSPGE RATLSC	255	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2-8	EVVLT QSPGTL SLSPGE RATLSC	255	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2-9	EVVLT QSPGTL SLSPGE RATLSC	255	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2-10	EIVLTQ SPGTL LSPGER ATLSC	175	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-2-11	EIVLTQ SPGTL LSPGER ATLSC	175	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-8 (p)	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-8-13	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-8-14	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-8-15	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200
SB-8-16	EIVMT QSPGTL SLSPGE RATLSC	174	WYQQKPGQA PRLLIY	186	GIPDRFSGSGSG TDFTLTISRLEP EDFAVYYC	192	FGGGTKV EIK	200

Ab ID	VL FR1	SEQ ID NO	VL FR2	SEQ ID NO	VL F3	SEQ ID NO	VL FR4	SEQ ID NO
SB-40 (p)	EIVLTQ SPATLS LSPGER ATLSC	182	WYQQKPGQA PRLLIY	186	GIPARFSGSGSG TDFTLTISSELP EDFAVYYC	193	FGGGTKV EIK	200
SB-40-18	EIVLTQ SPATLS LSPGER ATLSC	182	WYQQKPGQA PRLLIY	186	GIPARFSGSGSG TDFTLTISSELP EDFAVYYC	193	FGGGTKV EIK	200
SB-40-19	EIVLTQ SPATLS LSPGER ATLSC	182	WYQQKPGQA PRLLIY	186	GIPARFSGSGSG TDFTLTISSELP EDFAVYYC	193	FGGGTKV EIK	200
SB-40-20	EIVLTQ SPATLS LSPGER ATLSC	182	WYQQKPGQA PRLLIY	186	GIPARFSGSGSG TDFTLTISSELP EDFAVYYC	193	FGGGTKV EIK	200
SB-40-21	EIVLTQ SPATLS LSPGER ATLSC	182	WYQQKPGQA PRLLIY	186	GIPARFSGSGSG TDFTLTISSELP EDFAVYYC	193	FGGGTKV EIK	200

(p) denotes parental antibody sequence

Table 11: Kabat heavy chain FR sequences of affinity matured anti-SIRPβ1 antibodies

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-1 (p)	EVQLLE SGGGLV QPGGSL RLSCAA SGFTFS	201	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA K	215	WGQGT TVSS	222
SB-1-2	QVQLVE SGGGVV QPGRSL RLSCAA SGFTFG	256	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA K	215	WGQGT TVSS	222
SB-1-3	EVQLVE SGGGLV KPGGSL RLSCAA SGFTFS	257	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA K	215	WGQGT TVSS	222
SB-1-4	QVQLVE SGGGVV QPGGSL RLSCAA SGFTFS	258	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA K	215	WGQGT TVSS	222
SB-1-5	QVQLVE SGGGVV QPGRSL RLSCAA SGFTFS	202	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA K	215	WGQGT TVSS	222
SB-2 (p)	QVQLVE SGGGVV QPGRSL RLSCAA SGFTFS	202	WVRQAPGKG LEWVA	211	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
SB-2-7	EVQLLE SGGGLV QPGGSL RLSCAA SGFTFA	259	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222
SB-2-8	QVQLVE SGGGVV QPGRSL RLSCAA SGFTFL	260	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222
SB-2-9	QVQLVE SGGGVV QPGRSL RLSCAA SGFTFK	261	WVRQAPGKG LEWVA	211	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222
SB-2-10	EVQLLE SGGGLV QPGGSL RLSCAA SGFTFG	262	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222
SB-2-11	QVQLVE SGGGLV QPGGSL RLSCAA SGFTFA	263	WVRQAPGKG LEWVS	210	RFTISRDNSKN TLYLQMNSLR AEDTAVYYCA R	216	WGQGT TVSS	222
SB-8 (p)	QVQLQE SGPGLV KPSETLS LTCAVS GYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKN QFSLKLSSVTA ADTAVYYCAR	220	WGQGT TVSS	222
SB-8-13	QVQLQE SGPGLV KPSETLS LTCAVS GYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKN QFSLKLSSVTA ADTAVYYCAR	220	WGQGT TVSS	222
SB-8-14	QVQLQE SGPGLV KPSETLS LTCAVS GYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKN QFSLKLSSVTA ADTAVYYCAR	220	WGQGT TVSS	222
SB-8-15	QVQLQE SGPGLV KPSETLS LTCAVS GYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKN QFSLKLSSVTA ADTAVYYCAR	220	WGQGT TVSS	222
SB-8-16	QVQLQE SGPGLV KPSETLS LTCAVS GYSIS	205	WIRQPPGKGL EWIG	213	RVTISVDTSKN QFSLKLSSVTA ADTAVYYCAR	220	WGQGT TVSS	222
SB-40 (p)	QVQLVQ SGAEVK KPGASV KVSCKA SGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTRDTST STVYMELSSL RSEDVAVYYC AR	221	WGQGT TVSS	222
SB-40-18	QVQLVQ SGAEVK	203	WVRQAPGQR LEWMG	264	RVTITRDTAS TAYMELSSLR	265	WGQGT TVSS	222

Ab ID	VH FR1	SEQ ID NO	VH FR2	SEQ ID NO	VH F3	SEQ ID NO	VH FR4	SEQ ID NO
	KPGASV KVSCKA SGYTFT				SED TAVYYCA R			
SB-40-19	QVQLVQ SGAEVK KPGASV KVSCKA SGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTST STAYMELRSL RSDDTAVYYC AR	218	WGQGTT VTVSS	222
SB-40-20	QVQLVQ SGAEVK KPGASV KVSCKA SGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTTDTST STAYMELSSL RSED TAVYYC AR	266	WGQGTT VTVSS	222
SB-40-21	QVQLVQ SGAEVK KPGASV KVSCKA SGYTFT	203	WVRQAPGQG LEWMG	212	RVTMTRDTSIS TAYMELSRLR SDDTAVYYCA R	217	WGQGTT VTVSS	222

(p) denotes parental antibody sequence

Characterization of affinity-matured anti-SIRPβ1 antibody binding

[0276] A final set of affinity matured anti-SIRPβ1 antibodies were selected based on antigen binding affinities. Antibodies that were positive for binding to human SIRPβ1 were tested for cross-reactivity to human SIRPα. The biochemical characteristics of each antibody are listed below in Table 12. In Table 12, “N.B.” refers to antibodies for which there is no binding to the indicated antigen; “P.F.” refers to antibodies for which antigen binding kinetics show poor fit to 1:1 binding model; “N.M.” refers to not measurable.

Table 12: Biochemical Characterization of affinity matured anti-SIRPβ1 antibodies

Clone Index	Optimization Method	Fab K _D Human SIRPβ Fc (M) Monovalent	IgG K _D Mouse SIRPβ Fc (M) Avid	IgG K _D Human SIRPα Fc (M) Avid	MSD Fab K _D Human SIRPβ HIS (M) Monovalent
SB-1	Parent	1.72E-07	N.B.	N.B.	N.M.
SB-1-2	H1H2	3.78E-08	N.B.	N.B.	N.M.
SB-1-3	H1H2	4.54E-08	N.B.	N.B.	N.M.
SB-1-4	H1H2	4.33E-08	N.B.	N.B.	N.M.
SB-1-5	H1H2	8.24E-08	N.B.	N.B.	N.M.
SB-2	Parent	9.77E-07	N.B.	N.B.	N.M.
SB-2-7	H1H2	6.21E-09	N.B.	N.B.	N.M.
SB-2-8	H1H2	5.02E-09	N.B.	N.B.	N.M.
SB-2-9	H1H2	1.92E-08	N.B.	N.B.	N.M.
SB-2-10	H1H2	1.23E-08	N.B.	N.B.	N.M.
SB-2-11	H3	4.40E-09	N.B.	N.B.	N.M.
SB-8	Parent	1.51E-07	N.B.	N.B.	N.M.
SB-8-13	H1H2	2.85E-09	N.B.	N.B.	6.3E-11

SB-8-14	H1H2	3.25E-09	N.B.	1.83E-08	5.3E-11
SB-8-15	H1H2	3.73E-09	N.B.	N.B.	4.8E-11
SB-8-16	H3	2.32E-09	N.B.	N.B.	5.5E-11
SB-40	Parent	9.18E-07	N.B.	N.B.	N.M.
SB-40-18	H1H2	1.81E-08	N.B.	N.B.	N.M.
SB-40-19	H1H2	4.48E-08	N.B.	N.B.	N.M.
SB-40-20	H1H2	9.41E-09	N.B.	N.B.	N.M.
SB-40-21	H1H2	4.04E-08	N.B.	N.B.	N.M.

[0277] Subsequent characterization of affinity matured anti-SIRPβ1 antibodies involved determining their ability to bind cell lines expressing human or mouse SIRPβ1. Cells were harvested, plated at 10⁶/ml in a 96-well plate, washed, and incubated in 100μl FACS buffer containing 1 μg/ml anti-SIRPβ1 antibody for 0.5 hour on ice. Cells were then washed twice and incubated in 100ul FACS buffer containing 0.5μg/ml PE-conjugated secondary antibody for 30 minutes on ice. Cells were washed twice in cold FACS buffer and acquired on a BD FACS Canto. Data analysis and calculation of mean fluorescence intensity (MFI) values or % positive cells was performed with FlowJo (TreeStar) software version 10.0.7.

[0278] **Table 13** shows the mean fluorescence intensity (MFI) values of affinity matured anti-SIRPβ1 antibodies binding to the Chinese hamster ovary (CHO) cell line expressing low levels of human SIRPβ1. The human IgG1 isotype control established the background fluorescence signal set to 1. Of the 22 anti-SIRPβ1 antibody clones tested, 17 clones bound to cells with an MFI ≥ 3-fold over background. As a negative control, the anti-SIRPβ1 antibodies were also screened for surface binding to CHO cells overexpressing mouse SIRPβ1. As expected, none of the test antibodies showed significant binding to mouse SIRPβ1, as clones were originally selected for binding the human antigen. Given the high sequence similarity between receptors of the SIRP family, anti-SIRPβ1 antibodies were also screened for cross-reactivity to human SIRPα. In cell binding assays, none of the anti-SIRPβ1 antibodies bound cells overexpressing human SIRPα.

Table 13: Cell Binding Characterization of affinity matured anti-SIRPβ1 antibodies

Clone Index	Cell Binding Human SIRPβ FOB (Fold Over Background)	Cell Binding Human SIRPα FOB (Fold Over Background)	Cell Binding Mouse SIRPβ FOB (Fold Over Background)	EC50 (nM)
SB-1	2	1	1	0.072
SB-1-2	4	1	1	0.059
SB-1-3	4	0	2	0.062
SB-1-4	3	1	1	0.073
SB-1-5	3	1	2	0.038
SB-2	2	1	1	0.235
SB-2-7	5	1	1	0.132
SB-2-8	5	1	1	0.174

SB-2-9	4	1	2	0.469
SB-2-10	4	1	2	0.234
SB-2-11	6	0	2	0.220
SB-8	2	1	2	0.441
SB-8-13	7	1	2	0.253
SB-8-14	7	1	1	0.304
SB-8-15	7	1	1	0.737
SB-8-16	9	1	2	0.147
SB-40	2	1	1	0.707
SB-40-18	4	1	2	0.320
SB-40-19	3	1	2	0.713
SB-40-20	3	1	2	0.405
SB-40-21	5	1	2	0.715

[0279] Though the original four parental anti-SIRP β 1 antibodies did not demonstrate SIRP α cross-reactivity, Octet[®] analysis revealed that one of the progeny antibodies (SB-8-14) acquired avid binding to SIRP α despite the application of negative selection pressure during library screening. See Table 12 above. To determine if avid binding triggers SIRP α -dependent signaling, anti-SIRP β 1 antibodies were assessed for the ability to induce gene expression in human SIRP α and SIRP β 1 reporter cells. As described previously, test antibodies or two anti-SIRP α antibodies (clones SA-90, SA-94) were adsorbed onto 96-well plates at 10 μ g/mL. After washing, 10⁵ BWZ-huSIRP α or BWZ-huSIRP β 1 NFAT-luciferase reporter cells were seeded onto wells and incubated overnight at 37°C. Luciferase activity was quantified by adding OneGlo reagent (Promega) to each well and incubating samples at room temperature for 3 min on a plate shaker. The luminescence signal was quantified using a BioTek Synergy[™] Microplate Reader using GEN5[™] 2.04 software. As shown in **FIG. 8**, both the parental anti-SIRP β 1 antibodies and their affinity improved progeny retained the ability to induce luciferase expression in the BWZ-huSIRP β 1 reporter cells. All affinity matured clones, including SB-8-14, failed to significantly induce luciferase expression when BWZ-huSIRP α reporter cells were added onto antibody-coated wells (**FIG. 8**). In contrast, both anti-SIRP α antibody clones induced luciferase expression in BWZ-huSIRP α and not in BWZ-huSIRP β 1 reporter cells (**FIG. 8**). These results establish that despite avid binding to soluble SIRP α of one of the affinity matured progeny antibodies disclosed herein, all anti-SIRP β 1 antibodies tested failed to functionally engage membrane-bound SIRP α . Instead, the anti-SIRP β 1 antibodies demonstrated functional specificity towards membrane-bound human SIRP β 1.

Example 9: Affinity matured anti-SIRP β 1 antibodies increase secretion of inflammatory cytokines from dendritic cells.

[0280] To determine whether affinity matured anti-SIRP β 1 antibodies retain the ability to induce changes in inflammatory cytokine production, primary human monocyte-derived dendritic cells are

cultured with plate-bound test antibodies in combination with non-saturating levels of TLR stimulators and the level of cytokines are measured after 24 hours. To generate monocyte-derived dendritic cells, human primary monocytes were isolated from heparinized human blood (Blood Centers of the Pacific) using RosetteSep Human Monocyte Enrichment Cocktail (STEMCELL Technologies), according to the manufacturer's protocol. Monocytes were seeded in RPMI (Invitrogen) containing 10% Fetal Calf Serum (Hyclone) and 20 ng/ml IL-4 and GM-CSF (Peprotech) to induce differentiation of immature dendritic cells. After 6-7 days, dendritic cells were harvested by scraping cells attached to plastic. Dendritic cells are plated on 96-well plates coated with indicated antibody at 10^5 cells/well and incubated for 24h at 37°C. Cells are co-stimulated with TLR4 agonist, LPS (*Salmonella abortus equi*).

[0281] In FIG. 9, monocyte-derived dendritic cells from 3 healthy donors were stimulated overnight at 37°C with 0.5 ng/mL LPS in the presence of indicated affinity matured anti-SIRPβ1 antibodies adsorbed onto 96-well plates at 2 µg/mL. The supernatant fraction was subsequently collected and assayed for TNFα release. The SIRPβ1 antibodies SB-1-3, SB-2-8, and SB-8-13 proved highly agonistic as plate-bound antibodies. In contrast, remaining SIRPβ1 antibodies, SB-2-8, SB-8-15, and SB-40-20, weakly activated SIRPβ1-mediated cytokine release.

Example 10: Anti-SIRPβ1 antibodies are specific to human SIRPβ1 isoform 1

[0282] Among the multiple SIRPβ1 isoforms catalogued, isoform 1 is the full-length version of the receptor primarily expressed in the periphery. Recombinant antigen based on the sequence of SIRPβ1 isoform 1 was used for selecting anti-SIRPβ1 antibodies from yeast library pools. To ascertain if the selected anti-SIRPβ1 antibodies cross-reacted with other isoforms of human SIRPβ1 or with cynomolgus SIRPβ1, recombinant antigens based on human SIRPβ1 isoform 3 sequence and cynomolgus SIRPβ1 isoform 1 sequence were produced by transient transfection in HEK293 cells. Purified, Fc-tagged recombinant proteins were adsorbed onto 96-well plates at 1 µg/mL overnight at 4°C. Plates were subsequently washed with PBS + 0.05% Tween-20 and blocked with 1% BSA + PBS for 2 hours at room temperature. Antibodies diluted in blocking buffer at 1 µg/mL were added to wells, serially diluted, and allowed to bind antigen overnight at 4°C. The following day, the primary antibodies were removed and anti-human kappa light chain antibody conjugated to horseradish peroxidase diluted 1:10,000 in blocking buffer were added to wells and incubated for 1 hour at room temperature. Plates were washed with PBS + 0.05% Tween-20 and 100 µL of TMB substrate solution was added to wells to generate a colorimetric signal proportional to amount of anti-SIRPβ1 antibody bound to antigen. Once the reaction reached an appropriate color intensity, 100 µL of Stop solution was added to quench the enzyme. Plates were read with a BioTek Synergy™ Microplate Reader using GEN5™ 2.04 software.

[0283] **FIG. 10A-10B** shows binding curves of anti-SIRPβ1 antibodies bound to indicated SIRPβ1 antigens based on ELISA data. Eight anti-SIRPβ1 antibodies (four parental antibodies and four corresponding affinity matured antibodies) were selected for analysis. EC50 values calculated from these binding curves demonstrated that all eight anti-SIRPβ1 antibodies recognized human SIRPβ1 isoform 1 with high affinity. For example, the EC50 values for SB-1 and SB-1-2 are 0.034 nM and 0.045 nM, respectively. The EC50 values for SB-2 and SB-2-7 are 0.032 nM and 0.029 nM, respectively. The EC50 values for SB-8 and SB-8-13 are 0.022 nM and 0.019 nM, respectively. The EC50 values for SB-40 and SB-40-21 are 0.040 nM and 0.025 nM, respectively. However, the anti-SIRPβ1 antibodies tested did not cross-react to either human SIRPβ1 isoform 3 or to cynomolgus SIRPβ1. These results demonstrate that the screening and selection platform yielded highly specific antibodies to human SIRPβ1 isoform 1 with no evidence of cross-reactivity towards human SIRPα, human SIRPβ1 isoform 3, murine SIRPβ1, or cynomolgus SIRPβ1.

Example 11: Analysis of the effect of anti-SIRPβ1 antibodies in increasing recruitment of immune cells *in vivo*

[0284] The ability of anti-SIRPβ1 antibodies to modulate the recruitment of inflammatory cells (neutrophil granulocytes, monocytes, and macrophages) in the peritoneal cavity (PEC) of human SIRPβ1 BAC-transgenic mice after intraperitoneal (IP) administration of either antibody alone or in combination with LPS is evaluated as follows. Briefly, mice receive first an IP injection of 40 mg/kg anti-SIRPβ1 antibody or isotype control antibody mIgG1 (clone MOPC-21, Bioxcell). 14 hours later, mice receive an IP injection of 4 mg/kg LPS, or PBS as a control. 6 hours after LPS or PBS injection, cells are harvested from the PEC as described (*see, e.g., Gawish R et al, 2014 FASEB J*) and analyzed by FACS. For FACS analysis, PEC cells are incubated with anti-CD11b-Pacific Blue, anti-CD11c PeCy7, anti-MCH-II- APCCy7, anti-Gr1-FITC, anti-Ly6G-PE and a viability dye (Life Technologies, Cat# L34957) for 1 hour on ice, then washed twice with cold FACS buffer. 4% PFA-fixed samples are then acquired. Data are acquired on a BD FACS CANTO II cytometer (Becton Dickinson) and analyzed with FlowJo software. These studies provide support that anti-SIRPβ1 antibodies of the present disclosure increase recruitment of immune cells *in vivo*.

Example 12: SIRPβ1 expression in the tumor microenvironment

[0285] Groups of 3 human SIRPβ1 BAC-transgenic mice (females, 8 weeks old) are challenged subcutaneously with 1×10^6 MC38 or CT26 colon carcinoma cells, or EMT-6 murine mammary carcinoma cells, suspended in 100μl PBS. Animals are anesthetized with isoflurane prior to implant. When the tumors reach a size of 700-1000 mm³, tumors are explanted to analyze SIRPβ1 expression in the tumor microenvironment by FACS. As a comparison, the spleen of the tumor bearing mice or control spleen of naïve mice is also analyzed. For expression analysis by FACS, tumor and spleens

are incubated in PBS containing 1mg/ml collagenase and then processed through a cell strained to obtain a single cell suspension. Cells are then incubated with anti-CD45-PerCp-Cy7, anti-CD11b-PerCP-Cy5.5, anti-CD3-PC, anti-Gr1-FITC, anti-NK1.1-PE, anti-SIRP β 1-APC antibodies and a viability dye (Life Technologies, Cat# L34957) for 30min on ice, then washed twice with cold FACS buffer. 4% PFA-fixed samples are then acquired. Data are acquired on a BD FACS CANTO II cytometer (Becton Dickinson) and analyzed with FlowJo software. These studies provide support that myeloid cells or cells of myeloid lineage express SIRP β 1 in the tumor environment.

Example 13: Analysis of tumor growth in human SIRP β 1 BAC transgenic mice

[0286] Groups of wild-type (WT, n=11) and human SIRP β 1 BAC transgenic mice (SIRP β 1tg, n=14) mice (sex and age-matched littermates, 10 weeks old (+/- 2 weeks)) are challenged subcutaneously with 1×10^6 MC38 colon carcinoma tumor cells suspended in 100 μ l PBS. Mice are then anesthetized with isoflurane prior to tumor implant. Following tumor implant, the mice are administered various concentrations of an anti-SIRP β 1 antibody of the present invention. Tumor growth is monitored with a caliper biweekly starting at day 5. The endpoint of the experiment is a tumor volume of 2000 mm³ or 60 days. The effect of anti-SIRP β 1 antibody administration on reducing tumor growth or tumor size (expressed as volume, mm³), as compared to that observed in control animals, is determined. These studies provide support that anti-SIRP β 1 antibodies of the present invention reduce tumor growth *in vivo*.

Example 14: Anti-SIRP β 1 antibodies induce the expression of CD83 and CD86 on human dendritic cells (DCs)

[0287] To evaluate the ability of anti-SIRP β 1 antibodies to modify expression of CD83 and CD86, both plate-bound and soluble antibodies are incubated with dendritic cells (DCs), and the expression of CD83, CD86, CCR7, and phosphorylated ERK are measured. Antibodies are plated overnight at 4°C in 12 well plates at 2 or 10 μ g/ml in PBS. Wells are washed 3X with PBS the next day. Primary human monocytes isolated from peripheral blood of healthy donors are added to antibody coated wells in RPMI media supplemented with 10% FBS and 20 ng/mL IL-4 and GM-CSF and incubated at 37°C, 5% CO₂ for 5 days. On day 5, immature human DCs are harvested and analyzed by FACS for CD86, CD83, CD1a, and HLA-DR, on a BD FACS Canto. Data analysis is performed with FlowJo (TreeStar) software version 10.0.7. Levels of CD83 and CD86 are evaluated for CD1a+/HLA-/DR+ cell populations. For intracellular ERK phosphorylation, cells are fixed with 1% formaldehyde, permeabilized with cytofix/cytoperm kit (BD), and intracellular Erk phosphorylation is determined with flow cytometry after staining with PE-ERK antibody (BD). These studies provide support that anti-SIRP β 1 antibodies of the present invention increase expression of CD83, CD86 and CCR7 in human dendritic cells. Additionally, these studies provide support that anti-

SIRP β 1 antibodies of the present invention increase or induce ERK phosphorylation in human dendritic cells.

Example 15: Screening for anti-SIRP β 1 and/or anti-SIRP β 1 bispecific antibodies that induce phosphorylation of SIRP β 1, DAP12, SYK, ERK, and AKT, which indicate activation of the PI3K pathway

[0288] Primary human myeloid cells or murine myeloid cell lines engineered to express human SIRP β 1 (J774, RAW 264.7, BMM cells, or osteoclasts) are removed from tissue culture dishes with PBS-EDTA, washed with PBS, and counted. J774 (40×10^6) or RAW 264.7 cells (10×10^6 BMM or osteoclasts) are incubated with an anti-SIRP β 1 and/or an anti-SIRP β 1 bispecific antibody (such as an anti-SIRP β 1/TREM2 bispecific antibody) or with an isotype-matched control antibody at $1 \mu\text{g}/10^6$ cells for 20 min on ice or under other conditions. Cells are lysed in ice-cold radioimmunoprecipitation assay (RIPA) buffer for 20 min followed by centrifugation at 16,000 g for 10 min at 4°C to remove insoluble materials. The resulting supernatant is subjected to immunoprecipitation reactions with the indicated antibodies (DAP12, ERK, or AKT) and protein A- or protein G-agarose (Sigma). The beads are extensively washed with RIPA buffer and the proteins are separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). The proteins are then transferred to nitrocellulose membranes by Western blotting, incubated with the appropriate antibodies (antibodies that specifically recognize the phosphorylated form of DAP12, ERK, or AKT), and visualized with the enhanced chemiluminescence (ECL) system (Pierce), as described (*e.g.*, Peng *et al.*, (2010) *Sci Signal.*, 3(122): ra38). These studies provide support that anti-SIRP β 1 antibodies of the present invention induce phosphorylation of SIRP β 1, DAP12, SYK, ERK, and AKT. Additionally, these studies provide support that anti-SIRP β 1 antibodies of the present invention are effective at activating the PI3K pathway.

Example 16: Anti-SIRP β 1 antibodies upregulate TREM2 expression on human macrophages

[0289] SIRP β 1 and TREM2 are DAP12-associated receptors expressed on myeloid cells. To evaluate if SIRP β 1 modifies the expression of TREM2, macrophages were cultured on plate-bound antibodies and analyzed for cell surface levels of TREM2 by flow cytometry.

[0290] Antibodies (SB-1-3, SB-2-8, SB-8-13, SB-8-15, SB-40-20, and huIgG1 control) were adsorbed onto 96-well plates at 37°C for 4 hours at $2 \mu\text{g}/\text{mL}$ in PBS. Wells were washed twice with PBS. Monocyte-derived human macrophages, differentiating in culture with M-CSF for 5-6 days, were harvested and plated at 100,000 cells per well. Macrophages were incubated overnight at 37°C, 5% CO₂. FACS analysis of TREM2 expression was performed on a BD FACS Canto II and data analysis was performed with FlowJo (TreeStar) software.

[0291] As shown in **FIG. 11A**, Plate-bound anti-SIRP β 1 antibodies SB-2-8, SB-8-15, and SB-40-20 increased TREM2 expression approximately 2-fold relative to huIgG1 isotype control treated macrophages obtained from 3 healthy donors. In 2 out of 3 healthy donors (624 and 626), however, plate-bound anti-SIRP β 1 antibodies SB-1-3 and SB-8-13 only partially increased TREM2 expression relative to huIgG1 isotype control treated macrophages. In macrophages from a third healthy donor (625), plate-bound SB-1-3 and SB-8-13 increased TREM2 expression approximately 2-fold relative to huIgG1 isotype control. Based on these results, anti-SIRP β 1 antibodies SB-2-8, SB-8-15, and SB-40-20 function as agonists to induce cell surface TREM2 expression.

Example 17: Anti-SIRP β 1 antibodies increase viability of human macrophages

[0292] Evidence in the literature suggests that TREM2 promotes macrophage/microglia viability. Since SIRP β 1 associates with DAP12 adaptor molecule and may influence TREM2 expression, anti-SIRP β 1 antibodies were evaluated for the ability to increase human macrophage viability.

[0293] Antibodies (SB-1-3, SB-2-8, SB-8-13, SB-8-15, SB-40-20, and huIgG1 control) were adsorbed onto 96-well plates at 37°C for 4 hours at 2 μ g/mL in PBS. Wells were washed twice with PBS. Monocyte-derived human macrophages, differentiating in culture with M-CSF for 5-6 days, were harvested and washed to remove residual M-CSF. Macrophages were resuspended in RPMI growth media supplemented with 10% FBS and 1% Penn/Strep at 500,000 cell per mL. Cells were diluted with equal volume of PBS and 100 μ L containing 25,000 cells were added to each well. Macrophages were incubated at 37°C for 2 days. Analysis of viability was performed using Cell Titer Glo kit (Promega), a reagent that produces a luminescence signal relative to ATP concentration in the sample. Plates were read with a Biotek Synergy Microplate Reader using GEN5 2.04 software.

[0294] As shown in **FIG. 11B**, baseline luminescence was established with macrophages cultured on plate-bound, full-length huIgG1 isotype control. In comparison to macrophages seeded in the absence of antibody (No Ab), there is no significant change in macrophage viability with the isotype control. However, macrophages cultured on plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3, SB-8-13, and SB-8-15 demonstrated a significant increase in viability relative to isotype control treated cells. Plate-bound, full-length anti-SIRP β 1 antibodies SB-2-8 and SB-40-20 failed to increase macrophage viability relative to the isotype control treated cells.

[0295] Given the differences in activity initially observed between the anti-SIRP β 1 antibodies, subsequent assays determined if antibody activity depended on concentration of adsorbed protein. Anti-SIRP β 1 antibodies SB-1-3 and SB-2-8 or the human IgG1 isotype control were coated onto 96-well plates at 37°C for 4 hours starting at 10 μ g/mL in PBS and serially diluted 3-fold. As described previously, macrophages were diluted with equal volume of PBS and 100 μ L containing 25,000 cells were added to each well. Macrophages were incubated at 37°C for 2 days. Analysis of viability was

performed using Cell Titer Glo kit (Promega) and plates were read with a Biotek Synergy Microplate Reader using GEN5 2.04 software.

[0296] As shown in **FIG. 12**, plate-bound, full-length anti-SIRP β 1 antibody SB-1-3 increased macrophage viability in a dose-dependent manner relative to isotype control treated cells. In contrast, plate-bound, full-length anti-SIRP β 1 antibody SB-2-8 failed to increase macrophage viability even at high concentrations of coated protein. Thus, anti-SIRP β 1 antibodies SB-1-3, SB-8-13, and SB-8-15 demonstrated agonistic activity in this assay.

Example 18: Additive and synergistic effect of combination treatment of macrophages with anti-SIRP β 1 and anti-TREM2 antibodies

[0297] Agonistic anti-TREM2 antibodies have been shown to increase macrophage viability when added to cells in a soluble format. Viability assays were performed to determine if co-stimulating macrophages with anti-SIRP β 1 antibodies further enhances the agonist activity anti-TREM2 antibody.

[0298] Anti-SIRP β 1 antibodies SB-1-3 and SB-2-8 or the human IgG1 isotype control were coated onto 96-well plates at 37°C for 4 hours at 10 μ g/mL in PBS. As described previously, macrophages were diluted with equal volume of PBS and 100 μ L containing 25,000 cells were added to each well. Where indicated, macrophages were also treated with 50 μ g/mL of anti-TREM2 antibody. Macrophages were incubated at 37°C for 2 days. Analysis of viability was performed using Cell Titer Glo kit (Promega) and plates were read with a Biotek Synergy Microplate Reader using GEN5 2.04 software.

[0299] As shown in **FIG. 13**, adding agonist anti-TREM2 antibody to cells cultured on human IgG1 isotype control significantly increased macrophage viability relative to cells cultured in the absence of agonist anti-TREM2 antibody, establishing the reproducibility of this assay.

Furthermore, culturing macrophages on plate-bound, full-length anti-SIRP β 1 antibody SB-1-3 significantly increased viability relative to isotype treated cells, whereas anti-SIRP β 1 antibody SB-2-8 did not show that activity (**FIG. 12**). Co-stimulating macrophages with plate-bound anti-SIRP β 1 antibody SB-1-3 and soluble agonist anti-TREM2 antibody shows additivity by further enhancing macrophage viability. Interestingly, co-stimulating macrophages with plate-bound anti-SIRP β 1 antibody SB-2-8 and soluble anti-TREM2 antibody shows synergistic activity by increasing macrophage viability. Thus, anti-SIRP β 1 antibodies enhance the activity of agonistic anti-TREM2 antibodies.

Example 19: Anti-SIRP β 1 antibodies increase the viability of bone marrow-derived macrophages

[0300] Human SIRP β 1 BAC transgenic mice that recapitulate the expression of the human antigen in mouse myeloid cells were generated. Murine bone marrow cells from human SIRP β 1 transgenic

mice were obtained by flushing tibia and femurs with PBS. Bone marrow cells were cultured in RPMI media supplemented with 50 ng/mL M-CSF to differentiate macrophages or 10 ng/mL GM-CSF to differentiate dendritic cells. Anti-SIRP β 1 antibodies were evaluated for their ability to increase murine myeloid cell viability.

[0301] Anti-SIRP β 1 antibodies SB-1-3, SB-2-8, and SB-8-13 or the human IgG1 isotype control were coated onto 96-well plates at 37°C for 4 hours at 10 μ g/mL in PBS. Macrophages and dendritic cells were diluted with equal volume of PBS and 100 μ L containing 25,000 cells were added to each well. Cells were incubated at 37°C for 2 days. Analysis of viability was performed using Cell Titer Glo kit (Promega) and plates were read with a Biotek Synergy Microplate Reader using GEN5 2.04 software. As shown in **FIG. 14A**, consistent with human macrophages, culturing bone marrow-derived macrophages on plate-bound, full-length anti-SIRP β 1 antibodies SB-1-3 and SB-8-13 enhanced macrophage viability relative to isotype control treated cells. In contrast, anti-SIRP β 1 antibody SB-2-8 failed to increase macrophage viability. As shown in **FIG. 14B**, when dendritic cells (DC) were cultured on plate-bound antibodies, all anti-SIRP β 1 antibodies increased DC viability relative to isotype control treated cells. However, only anti-SIRP β 1 antibody SB-8-13 treated DC showed increased viability relative to untreated dendritic cells (No Ab.). Thus, anti-SIRP β 1 antibodies show agonistic activity with murine myeloid cells expressing human SIRP β 1.

Example 20: Cross-reactivity of affinity matured anti-SIRP β 1 antibodies to related SIRP receptors

[0302] A characteristic feature of proteins of the SIRP family is their extensive amino acid sequence conservation in the extracellular domain. For example, the extracellular region of SIRP β 1 shares ~90% sequence identity with SIRP α and ~77% sequence identity with SIRP γ . Antibodies developed against one SIRP protein family member often cross-react with multiple SIRP receptor family members.

[0303] Despite the homology in sequence/structure, the SIRP receptors exhibit different functional properties and expression patterns. For example, SIRP α contains a cytoplasmic ITIM motif to mediate immune suppression upon binding to CD47. In contrast, SIRP β 1, which lacks intracellular signaling motifs and does not bind CD47, associates with the ITAM-containing DAP12 adaptor protein to propagate activation signals. SIRP γ , unlike other SIRP receptors that are primarily expressed on myeloid cells, is expressed on lymphocytes and NK cells and serves to stabilize cell-cell adhesion during antigen presentation to T cells.

[0304] To determine antigen specificity of anti-SIRP β 1 antibodies of the present disclosure, cell-based affinity measurements were performed to ascertain the apparent affinities of affinity-matured anti-SIRP β 1 antibodies to cell surface-expressed antigen. BW5147.G.1.4 cells, an immortalized mouse T-cell line, were engineered to overexpress human SIRP β 1 (BWZ-huSIRP β 1) or human

SIRP α (BWZ-huSIRP α). Jurkat cells, an immortalized human T cell line, endogenously express SIRP γ .

[0305] In these experiments, two antibodies previously described were used as positive controls: antibody 18D5 (an anti-SIRP antibody that cross-reacts with SIRP α and SIRP β) and antibody KWAR23 (an anti-SIRP antibody that cross-reacts with SIRP α , SIRP β , and SIRP γ). These antibodies were recombinantly produced on a human IgG4 backbone. A commercially available anti-SIRP γ antibody (clone LSB2.20; Biolegend, San Diego) conjugated with phycoerythrin (PE) fluorophore was used to verify SIRP γ expression on Jurkat cells. Serial dilutions of each of the anti-SIRP β 1 monoclonal antibodies and control antibodies were added to 10^5 cells and allowed to achieve binding equilibrium at 4°C. After addition of fluorescently-labeled secondary antibody and brief washing steps, MFI values as a function of titrated antibody concentration were recorded via FACS analysis.

[0306] In **Table 14** below, EC50 values assign relative affinity by adding increasing concentrations of antibody variants to cells overexpressing SIRP β 1. Receptor bound antibodies were detected by staining cells with anti-human IgG PE secondary antibody (Southern Biotech). Curves were fit using nonlinear regression analysis with Graphpad Prism 6 software. As shown in **Fig. 15A**, anti-SIRP β 1 antibodies bound BWZ-huSIRP β 1 cells with distinct EC50 profiles. Anti-SIRP β 1 antibody SB1-3 showed the best apparent affinity relative to other anti-SIRP β 1 antibodies, as well as compared to that of the positive control antibody, 18D5. Anti-SIRP β 1 antibodies failed to bind cells expressing SIRP α (**Fig. 15B**) or SIRP γ (**Fig. 15C**) demonstrating that these antibodies are antigen-specific.

Table 14: EC50 values of antibodies binding BWZ-human SIRP β 1 cells

	SB1-3	SB2-8	SB8-13	SB8-15	SB40-20	18D5
EC ₅₀ (nM)	0.02837	0.2025	0.1993	0.5962	8.704	0.4769

CERTAIN SEQUENCES*SB-1: Light Chain Variable Region*

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQLLGSSPRTFSGGGTKVEIK (SEQ ID NO: 267)

SB-1: Heavy Chain Variable Region

EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSTISGSGGSTYYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGT TVTVSS (SEQ ID NO:
268)

SB-2: Light Chain Variable Region

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSSHPFTFSGGGTKVEIK (SEQ ID NO: 269)

SB-2: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMNWRQAPGKGLEWVAVIWDGSKNYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAAIWGMVWGQGT TVTVSS (SEQ ID NO:
270)

SB-3: Light Chain Variable Region

EIVMTQSPATLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGS
GTDFTLTISLLEPEDFAVYYCQQLRFHPPTFSGGGTKVEIK (SEQ ID NO: 271)

SB-3: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTGYMHWRQAPGGLEWMGWINPSSGGTNYAQKFQGR
VTMTRDTSISTAYMELSLRLSDDTAVYYCAREGIAATDAYFDLWGRGTLTVTVSS (SEQ ID NO:
272)

SB-4: Light Chain Variable Region

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQYADAPITFSGGGTKVEIK (SEQ ID NO: 273)

SB-4: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWRQAPGGLEWMGWISAYNGNTNYAQKLQGR
VTMTTDTSTSTAYMELSLRLSDDTAVYYCARSGTHFGTYSYSNWFDPWGQGT LTVTVSS (SEQ ID
NO: 274)

SB-5: Light Chain Variable Region

EIVLTQSPATLSSLSPGERATITCRASQSVSSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGS
GTDFTLTISLLEPEDFAVYYCQQLRFHPPTFSGGGTKVEIK (SEQ ID NO: 275)

SB-5: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTGYYMHWVRQAPGQGLEWMGWINPNSGGTNYAQKFQGR
VTMTRDTSISTAYMELSRRLRSDDTAVYYCAREGDEDWFDPPWGQGTLLTVTVSS (SEQ ID NO: 276)

SB-6: Light Chain Variable Region

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSGHLPITFGGGTKVEIK (SEQ ID NO: 277)

SB-6: Heavy Chain Variable Region

EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSTISGSGGSTYYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGTITVTVSS (SEQ ID NO:
268)

SB-7: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQHYIAPFTFGGGTKVEIK (SEQ ID NO: 278)

SB-7: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPIFGTASYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARETRQDSAHHYGMVWGQGTITVTVSS (SEQ ID NO:
279)

SB-8: Light Chain Variable Region

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASNRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQVYSSPYTFGGGKVEIK (SEQ ID NO: 280)

SB-8: Heavy Chain Variable Region

QVQLQESGPGLVKPSETLSLTCAVSGYSISSGYWGWIRQPPGKGLEWIGSIYHSGSTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARGGAMTPAGMDVWGQGTITVTVSS (SEQ ID NO:
281)

SB-9: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQYHSVPPITFGGGTKVEIK (SEQ ID NO: 282)

SB-9: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGIHWVRQAPGQGLEWMGWISAYNGNTNYAQKLQGR
VTMTTDTSTSTAYMELRSLRSDDTAVYYCARDGLHYGDYIVYYGMVWGQGTITVTVSS (SEQ ID
NO: 283)

SB-10: Light Chain Variable Region

DIVMTQSPSLPVPVTPGEPASISCRSSQSLLHSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDFR
SGSGSGTDFTLKISRVEAEDVGVYYCMQAIESPLTFGGGTKVEIK (SEQ ID NO: 284)

SB-10: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARGVPRGDLGMDVWGQGTTVTVSS (SEQ ID NO:
285)

SB-11: Light Chain Variable Region

DIVMTQSPSLPVPVTPGEPASISCRSSQSLLYSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDFR
SGSGSGTDFTLKISRVEAEDVGVYYCVQALQTPITFGGGGTKVEIK (SEQ ID NO: 286)

SB-11: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSNYAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARPVDSSSYSLGYYYGMDVWGKGTTVTVSS (SEQ ID
NO: 287)

SB-12: Light Chain Variable Region

EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIYSASTRATGIPARFSGSGS
GTEFTLTISLQSEDFAVYYCQQLDNLPYTFGGGTKVEIK (SEQ ID NO: 288)

SB-12: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTGYMHVVRQAPGGGLEWMGWINPNSGGTSYAQKFQGR
VTMTRDTSISTAYMELSRRLRSDDTAVYYCARDTYAYSYGMDVWGQGTTVTVSS (SEQ ID NO:
289)

SB-13: Light Chain Variable Region

DIVMTQSPSLPVPVTPGEPASISCRSSQSLLYSNGYNYLDWYLQKPGQSPQVLIYLGSNRASGVPDFR
SGSGSGTDFTLKISRVEAEDVGVYYCMQALRSPITFGGGGTKVEIK (SEQ ID NO: 290)

SB-13: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQPPGKGLEWIGSIYYSGSTNYPNLSKSRV
TISVDTSKNQFSLKLSSVTAADTAVYYCARGDTSGGAYFDLWGRGTLTVTVSS (SEQ ID NO:
291)

SB-14: Light Chain Variable Region

EIVMTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
GTDFTLTISLLEPEDFAVYYCQQFSYYPITFGGGGTKVEIK (SEQ ID NO: 292)

SB-14: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPGGGLEWMGGIIPIFGTASYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARDRGVGFYWGQGTTLTVTVSS (SEQ ID NO: 293)

SB-15: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQAYSHPFTFGGGTKVEIK (SEQ ID NO: 294)

SB-15: Heavy Chain Variable Region

QLQLQESGPGLVKPSSETLSLTCTVSGSISNSYYWGWIRQPPGKGLEWIGSIYYSGSTYYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCAREVGAPPSYPFDIWGQGTMTVTVSS (SEQ ID NO:
295)

SB-16: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQLFSTPFTFGGGTKVEIK (SEQ ID NO: 296)

SB-16: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPGGGLEWMGSIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARANYDSSGYSGLDLWGRGTLVTVSS (SEQ ID NO:
297)

SB-17: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQYYDDPYTFGGGKVEIK (SEQ ID NO: 298)

SB-17: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSYGISWVRQAPGGGLEWMGWISAYNGNTNYAQKLQGR
VTMTTDTSTSTAYMELRSLRSDDTAVYYCARGPLLYGDYHVRYGMDVWGQGTTVTVSS (SEQ ID
NO: 299)

SB-18: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRSSQSLHNSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDRF
SGSGSGTDFTLKISRVEAEDVGVYYCLQALQTPITFGGGTKVEIK (SEQ ID NO: 300)

SB-18: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARAKPRGDYGMVWGQGTTVTVSS (SEQ ID NO:
301)

SB-19: Light Chain Variable Region

DIVMTQSPSLPVTTPGEPASISCRSSQSLHNSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDRF
SGSGSGTDFTLKISRVEAEDVGVYYCMQAIGVPPTFGGGTKVEIK (SEQ ID NO: 302)

SB-19: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPQGQGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARDGGGGYAYEYFQHWGQGTTLVTVSS (SEQ ID NO:
303)

SB-20: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISSLQAEDVAVYYCQQYYLSPFTFGGGTKVEIK (SEQ ID NO: 304)

SB-20: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPQGQGLEWMGSIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARDGREYGGHYGMDVWGQGTTVTVSS (SEQ ID NO:
305)

SB-21: Light Chain Variable Region

DIVMTQSPSLPVTTPGEPASISCRSSQSLLSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDRF
SGSGSGTDFTLKISRVEAEDVGVYYCMQTLRIPPTFGGGTKVEIK (SEQ ID NO: 306)

SB-21: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSNGISWVRQAPQGQLEWMGWISAYNGNTNYAQKLQGR
VTMTTDTSTSTAYMELRSLRSDDTAVYYCARVGNMDQEYFDLWGRGTLVTVSS (SEQ ID NO:
307)

SB-22: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVTTITCRASQGISSWLAWYQQKPGKAPKLLIYAASNLSGVPSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQGNSYPITFGGGTKVEIK (SEQ ID NO: 308)

SB-22: Heavy Chain Variable Region

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSDGSTYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARPTRYGYDRLGMDVWGQGTTVTVSS (SEQ ID NO:
309)

SB-23: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVTTITCRASQSISSYLNWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQAYPYPLTFGGGTKVEIK (SEQ ID NO: 310)

SB-23: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPQGQGLEWMGGIAPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARTTYRDYYMDVWGKGTTVTVSS (SEQ ID NO:
311)

SB-24: Light Chain Variable Region

EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIYGASTRATGIPARFSGSGS
GTEFTLTISLQSEDFAVYYCQQLNIHPWTFGGGTKVEIK (SEQ ID NO: 312)

SB-24: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWAWIRQPPGKGLEWIGSIYHSGSTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARDRSRGYPVYGMDVWGQGTTVTVSS (SEQ ID NO:
313)

SB-25: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVITITCRASQGISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQVNSFPWTFGGGTKVEIK (SEQ ID NO: 314)

SB-25: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSLAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARSGGDYSYDYASGMDVWGQGTTVTVSS (SEQ ID
NO: 315)

SB-26: Light Chain Variable Region

DIVMTQSPSLPVTTPGEPASISCRSSQSLHNSNGYNYLDWYLQKPGQSPQLLIYLGSNRASGVPDFR
SGSGSGTDFTLKISRVEAEDVGVYYCMQARGLPTFGGGTKVEIK (SEQ ID NO: 316)

SB-26: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARDGSAGRQEHGMDVWGQGTTVTVSS (SEQ ID NO:
317)

SB-27: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQAQVSDPPTFGGGTKVEIK (SEQ ID NO: 318)

SB-27: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGGGLEWMGGIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARQDLGSSHHWFDLWGRGTLVTVSS (SEQ ID NO:
319)

SB-28: Light Chain Variable Region

EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQDGNFPLTFGGGTKVEIK (SEQ ID NO: 320)

SB-28: Heavy Chain Variable Region

QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYWGWIRQPPGKGLEWIGSISYSGSTYYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCARDPRDYSSGSSGGWGYFDLWGRGTLVTVSS (SEQ
ID NO: 321)

SB-29: Light Chain Variable Region

DIVMTQSPSLPVTTPGEPASISCRSSQSLLSNGYNYLDWYLQKPGQSPQLLI FLGSNRASGVDPDRF
SGSGSGTDFTLKISRVEAEDVGVYYCMQARGSPITFGGGTKVEIK (SEQ ID NO: 322)

SB-29: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGGGLEWMGSIIPIFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSED TAVYYCARAPYGSSSGYGYFDLWGRGTLVTVSS (SEQ ID NO:
323)

SB-30: Light Chain Variable Region

EIVLTQSPGTLSPGERATLSCRASQSVSSSFLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQFLSSPWTFGGGTKVENQ (SEQ ID NO: 324)

SB-30: Heavy Chain Variable Region

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWIGYIYSGSTVYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCAREGPGYPSYFDPWGQGT L VTVSS (SEQ ID NO:
325)

SB-31: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVITITCRASQGISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQAVSHPFTFGGGTKVEIK (SEQ ID NO: 326)

SB-31: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLEWMGIINPGGGSTSYAQKFQGR
VTMTRDTSTSTVYMELSSLRSED TAVYYCAREGLYSSGWYIDVWGQGT L VTVSS (SEQ ID NO:
327)

SB-32: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVDPDR
FSGSGSGTDFTLTISLQAEDVAVYYCQQDFLTPITFGGGTKVEIK (SEQ ID NO: 328)

SB-32: Heavy Chain Variable Region

QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQPPGKGLEWIGYIYSSGSTNYNPSLKS RV
TISVDTSKNQFSLKLSSVTAADTAVYYCARGDSSSGGLDLWGRGTLVTVSS (SEQ ID NO: 329)

SB-33: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVTTITCQASQDISNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFSGSGS
GTDFTFTISSLQPEDATYYCQQFAFLPLTFGGGKVENQ (SEQ ID NO: 330)

SB-33: Heavy Chain Variable Region

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGSTYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARGQYTGSLDVWGQGTMTVTVSS (SEQ ID NO: 331)

SB-34: Light Chain Variable Region

EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIYGASTRATGIPARFSGSGS
GTEFTLTISSLQSEDFAVYYCQQDNTFPYTFGGGKVEIK (SEQ ID NO: 332)

SB-34: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTGYMHWRQAPGQGLEWMGWINPNSGGTKYAQKFQGR
VTMTRDTSISTAYMELSLRLSDDTAVYYCARDTYTPYGMDVWGQGTITVTVSS (SEQ ID NO:
333)

SB-35: Light Chain Variable Region

DIVMTQSPSLPVTTPGEPASISCRSSQSLHNSGYNLDWYLQKPGQSPQLLIYLGSNRASGVPDFR
SGSGSGTDFTLKISRVEAEDVGVYYCMQTLQVPLTFGGGKVEIK (SEQ ID NO: 334)

SB-35: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARGRPQSESYLLDYWGQGTITVTVSS (SEQ ID NO:
335)

SB-36: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVTTITCRASQGISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQAFSHRTFGGGKVEIK (SEQ ID NO: 336)

SB-36: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWRQAPGQGLEWMGIINPSGGSTSYAQKFQGR
VTMTRDTSSTVYMELSSLRSEDVAVYYCAREGPEQLWYLDYWGQGTITVTVSS (SEQ ID NO:
337)

SB-37: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRATGIPARFSGSGS
GTDFTLTISSLEPEDFAVYYCQQRHTYPLTFGGGKVEIK (SEQ ID NO: 338)

SB-37: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSNYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCARSRWGASGYYYYMDVWGQGTMTVTVSS (SEQ ID NO:
339)

SB-38: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVITITCRASQGISSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQAVSYPIITFGGGTKVEIK (SEQ ID NO: 340)

SB-38: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHVWRQAPGGGLEWMGIINPSGGSTSYAQKFQGR
VTMTRDTSTSTVYMELSSLRSEDATVYYCARESGTDFGTISYWGQGTLLTVSS (SEQ ID NO:
341)

SB-39: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVITITCRASQSISSYLNWYQQKPGKAPKLLIYGASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQSYDFPLITFGGGTKVEIK (SEQ ID NO: 342)

SB-39: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYWAWIRQPPGKGLEWIGSIYHSGSTYYNPSLKS
VTISVDTSKNQFSLKLSSVTAADTAVYYCARGGSNYGDYGRFDYWGQGTLLTVSS (SEQ ID NO:
343)

SB-40: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
GTDFTLTISLQPEDFAVYYCQQRDEHPPWTITFGGGTKVEIK (SEQ ID NO: 344)

SB-40: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSYMSWVRQAPGGGLEWMGIINPSGGSTSYAQKFQGR
VTMTRDTSTSTVYMELSSLRSEDATVYYCARDTGEYSYSPHGMVWGQGTITVTVSS (SEQ ID NO:
345)

SB-41: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVITITCQASQDITNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFSGSGS
GTDFTFTISLQPEDIATYYCQQADNFPYITFGGGTKVEIK (SEQ ID NO: 346)

SB-41: Heavy Chain Variable Region

QLQLQESGPGLVKPSSETLSLTCTVSGGSISSSSYWGWIRQPPGKGLEWIGSIYYSGSTYYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCARVGQYPIYGMDVWGQGTITVTVSS (SEQ ID NO:
347)

SB-42: Light Chain Variable Region

EIVMTQSPATLSVSPGERATITCRASQSISSYLNWYQQKPGKAPKLLIYSASSLQSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQGDSFPITFGGGTKVEIK (SEQ ID NO: 348)

SB-42: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQAPGGLEWMGWISAYNGNTNYAQKLQGR
VTMTTDTSTSTAYMELRSLRSDDTAVYYCARGPGHYVAGMDVWGQGTTVTVSS (SEQ ID NO:
349)

SB-43: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASKRATGIPARFSGSGS
GTDFTLTISLLEPEDFAVYYCQQRFDFPITFGGGTKVEIK (SEQ ID NO: 350)

SB-43: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWAWIRQPPGKGLEWIGSIYHSGSTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARDAPGYPMLGMDVWGQGTTVSVSS (SEQ ID NO:
351)

SB-44: Light Chain Variable Region

DIQMTQSPSTLSASVGRVTITCRASQSIGSWLAWYQQKPGKAPKLLIYKASSLESGVPSRFSGSGS
GTEFTLTISLQPDDEFATYYCQEYGSYRTFGGGTKVEIK (SEQ ID NO: 352)

SB-44: Heavy Chain Variable Region

EVQLVESGGGLVQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLEWVSVIYSGGDTYYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCAREGSSFWSGSAVSYYGMDVWGQGTTVTVSS (SEQ ID
NO: 353)

SB-45: Light Chain Variable Region

VLTQSPGTLTLSPGERATLSCRASQSVSSSFLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSG
TDFTLTISRLEPEDFAVYYCQQVSVPTFGGGTKVEIK (SEQ ID NO: 354)

SB-45: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISSGYYWAWIRQPPGKGLEWIGSIYHSGSTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARDLSRGYAVSGMDVWGQGTTVTVSS (SEQ ID NO:
355)

SB-46: Light Chain Variable Region

EIVLTQSPGTLTLSPGERATLSCRASQSVRSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQLYSSPYTFGGGTKVEIK (SEQ ID NO: 356)

SB-46: Heavy Chain Variable Region

EVQLLES GGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCARASPWELDVWGQGTMTVTVSS (SEQ ID NO: 357)

SB-47: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFSGSGS
GTDFTFTISSSLQPEDIATYYCQQADYFPITFGGGTKVEIK (SEQ ID NO: 358)

SB-47: Heavy Chain Variable Region

QLQLQESGPGGLVKPSETLSLTCTVSGGSISSSSYAWGWIRQPPGKGLEWIGSIYYSGSTYYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCARDLGHYDYWSGSRDYYYGMDVWGQGTITVTVSS
(SEQ ID NO: 359)

SB-48: Light Chain Variable Region

DIQMTQSPSSVSASVGDRVTITCRASQGIDSWLAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGS
GTDFTLTISSSLQPEDFATYYCQQASNFPITFGGGTKVEIK (SEQ ID NO: 360)

SB-48: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVISYDGSNKYYADSVKGR
FTISRDN SKNTLYLQMNSLRAEDTAVYYCARDGTIAAAGWPPEYFQHWGQGTITVTVSS (SEQ ID
NO: 361)

SB-49: Light Chain Variable Region

DIQMTQSPSSLSASVGDRVTITCQASQDITNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFSGSGS
GTDFTFTISSSLQPEDATYYCQQYFHPPLTFGGGKVEIK (SEQ ID NO: 362)

SB-49: Heavy Chain Variable Region

QLQLQESGPGGLVKPSETLSLTCTVSGGSISSSDYYWGWIRQPPGKGLEWIGSIYYSGSTYYNPSLKS
RVTISVDTSKNQFSLKLSSVTAADTAVYYCARGPTGYKDKWRYYYGMDVWGQGTITVTVSS (SEQ
ID NO: 363)

SB-50: Light Chain Variable Region

DIVMTQSPDSLAVSLGERATINCKSSQSVLFSSNNKNYLAWYQQKPGQPPKLLIYWASTRESGVPDR
FSGSGSGTDFTLTISSSLQAEDVAVYYCQQFLHTPRTFGGGKVEIK (SEQ ID NO: 364)

SB-50: Heavy Chain Variable Region

QVQLVQSGAEVKKPGSSVKVSCASGGTFSSYAISWVRQAPGQGLEWMGGIIPFGTANYAQKFQGR
VTITADESTSTAYMELSSLRSEDVAVYYCAREGGGHASYHYGYMDVWGQGTITVTVSS (SEQ ID
NO: 365)

SB-1-2: Light Chain Variable Region

EIVMTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQLLGSSPRTFGGGKVEIK (SEQ ID NO: 267)

SB-1-2: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFGSFGMNWVRQAPGKGLEWVSAITASGGSTFYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGTTVTVSS (SEQ ID NO:
366)

SB-1-3: Light Chain Variable Region

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQLLGSSPRTFGGGTKVEIK (SEQ ID NO: 267)

SB-1-3: Heavy Chain Variable Region

EVQLVESGGGLVKPGGSLRLSCAASGFTFSAYGMNWRQAPGKGLEWVSAITSSGRSTYYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGTTVTVSS (SEQ ID NO:
367)

SB-1-4: Light Chain Variable Region

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQLLGSSPRTFGGGTKVEIK (SEQ ID NO: 267)

SB-1-4: Heavy Chain Variable Region

QVQLVESGGGVVQPGGSLRLSCAASGFTFSAYGMNWRQAPGKGLEWVS AIRASGGATYYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGTTVTVSS (SEQ ID NO:
368)

SB-1-5: Light Chain Variable Region

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQLLGSSPRTFGGGTKVEIK (SEQ ID NO: 267)

SB-1-5: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFSAYGMNWRQAPGKGLEWVSAISASGRSTFYADSVKGR
FTISRDNSKNTLYLQMNSLRAEDTAVYYCAKDFTEVVGWLGMDVWGQGTTVTVSS (SEQ ID NO:
369)

SB-2-7: Light Chain Variable Region

EVVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQSSSHPFTFGGGTKVEIK (SEQ ID NO: 370)

SB-2-7: Heavy Chain Variable Region

EVQLLES GGGLVQPGGSLRLSCAASGFTFARYGMHWVRQAPGKGLEWVSAISGLAGPTYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAAIWGM DVWGQGTTVTVSS (SEQ ID NO:
371)

SB-2-8: Light Chain Variable Region

EVVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSSSHPFTFGGGTKVEIK (SEQ ID NO: 370)

SB-2-8: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFLDYGMHWVRQAPGKGLEWVSAISAFAGSTYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAAIWGM DVWGQGT TTVTVSS (SEQ ID NO:
372)

SB-2-9: Light Chain Variable Region

EVVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSSSHPFTFGGGTKVEIK (SEQ ID NO: 370)

SB-2-9: Heavy Chain Variable Region

QVQLVESGGGVVQPGRSLRLSCAASGFTFKTYGMHWVRQAPGKGLEWVAHIWYEGSNKVYADSVKGRF
FTISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAAIWGM DVWGQGT TTVTVSS (SEQ ID NO:
373)

SB-2-10: Light Chain Variable Region

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSSSHPFTFGGGTKVEIK (SEQ ID NO: 269)

SB-2-10: Heavy Chain Variable Region

EVQLLES GGGLVQPGGSLRLSCAASGFTFGSYGMHWVRQAPGKGLEWVSAISGLAGQTYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAAIWGM DVWGQGT TTVTVSS (SEQ ID NO:
374)

SB-2-11: Light Chain Variable Region

EIVLTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQSSSHPFTFGGGTKVEIK (SEQ ID NO: 269)

SB-2-11: Heavy Chain Variable Region

QVQLVES GGGLVQPGGSLRLSCAASGFTFARYGMHWVRQAPGKGLEWVSAISGLAGPTYADSVKGRF
TISRDN SKNTLYLQMNSLRAEDTAVYYCARDQTAAWGIWGM DVWGQGT TTVTVSS (SEQ ID NO:
375)

SB-8-13: Light Chain Variable Region

EIVMTQSPGTLSSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASN RATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQVYSSPYTFGGG TKVEIK (SEQ ID NO: 280)

SB-8-13: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISAHYYWGWIRQPPGKGLEWIGSIFHSGHTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARGGAMTPAGMDVWGQGTTVTVSS (SEQ ID NO:
376)

SB-8-14: Light Chain Variable Region

EIVMTQSPGTLTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASNRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQVYSSPYTFGGGTKVEIK (SEQ ID NO: 280)

SB-8-14: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISPHYYWGWIRQPPGKGLEWIGSIYHSGHTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARGGAMTPAGMDVWGQGTTVTVSS (SEQ ID NO:
377)

SB-8-15: Light Chain Variable Region

EIVMTQSPGTLTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASNRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQVYSSPYTFGGGTKVEIK (SEQ ID NO: 280)

SB-8-15: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISPHYYWGWIRQPPGKGLEWIGSIYHSGHTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARGGAMTPAGMDVWGQGTTVTVSS (SEQ ID NO:
377)

SB-8-16: Light Chain Variable Region

EIVMTQSPGTLTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASNRATGIPDRFSGSG
SGTDFTLTISRLEPEDFAVYYCQQVYSSPYTFGGGTKVEIK (SEQ ID NO: 280)

SB-8-16: Heavy Chain Variable Region

QVQLQESGPGLVKPSSETLSLTCAVSGYSISAHYYWGWIRQPPGKGLEWIGSIFHSGHTYYNPSLKSR
VTISVDTSKNQFSLKLSSVTAADTAVYYCARAGAMTPAGMDVWGQGTTVTVSS (SEQ ID NO:
378)

SB-40-18: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
GTDFTLTISSELEPEDFAVYYCQQRDEHPPWTFGGGTKVEIK (SEQ ID NO: 344)

SB-40-18: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMAWVRQAPGQRLEWMGWINPAVGATIYSQKFQGR
VTITRDTASTAYMELSSLRSEDTAVYYCARDTGEYSYSPHGMDVWGQGTTVTVSS (SEQ ID NO:
379)

SB-40-19: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
 GTDFTLTISLLEPEDFAVYYCQQRDEHPPWTFGGGTKVEIK (SEQ ID NO: 344)

SB-40-19: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSYVMVWRQAPGGLEWMGIINPSSGATNYAQKFQGR
 VTMTTDTSTSTAYMELRSLRSDDTAVYYCARDTGEYSYSPHGMDVWGQGTTVTVSS (SEQ ID NO:
 380)

SB-40-20: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
 GTDFTLTISLLEPEDFAVYYCQQRDEHPPWTFGGGTKVEIK (SEQ ID NO: 344)

SB-40-20: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSFYISWVRQAPGGLEWMGIINPSSGHTNYAQKLQGR
 VTMTTDTSTSTAYMELSSLRSED TAVYYCARDTGEYSYSPHGMDVWGQGTTVTVSS (SEQ ID NO:
 381)

SB-40-21: Light Chain Variable Region

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDSSNRATGIPARFSGSGS
 GTDFTLTISLLEPEDFAVYYCQQRDEHPPWTFGGGTKVEIK (SEQ ID NO: 344)

SB-40-21: Heavy Chain Variable Region

QVQLVQSGAEVKKPGASVKVSCASGYTFTSYVMVWRQAPGGLEWMGIINPSSGDTNYAQKFQGR
 VTMTTRDTSISTAYMELSRSLRSDDTAVYYCARDTGEYSYSPHGMDVWGQGTTVTVSS (SEQ ID NO:
 382)

SIRP α domain 1 (IgV domain)

EELQVIQPKSVLVAAGETATLRCTATSLIPVGPIQWFRGAGPGRELIYNQKEGHFPRVTTVSDLTK
 RNNMDFSIRIGNITPADAGTYVCVKFRKGSPDDVEFKSGAGTELSVRKPS (SEQ ID NO: 387)

SIRP β 1 domain 1 (IgV domain)

EDELQVIQPEKSVSVAAGESATLRCAMTSLIPVGPIMWFRGAGAGRELIYNQKEGHFPRVTTVSELT
 KRNNDFFSISISNITPADAGTYVCVKFRKGSPDDVEFKSGAGTELSVRKPS (SEQ ID NO:
 388)

CLAIMS

What is claimed is:

1. An isolated antibody that binds to human SIRP β 1, wherein the antibody binds to human SIRP β 1 isoform 1, but does not bind to human SIRP β 1 isoform 3, and optionally wherein the antibody has at least one, at least two, at least three, or at least four properties selected from:
 - a) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP α ;
 - b) binds to human SIRP β 1 isoform 1, but does not bind to human SIRP γ ;
 - c) binds to human SIRP β 1 isoform 1, but does not bind to mouse SIRP β 1;
 - d) binds to human SIRP β 1 isoform 1, but does not bind to cynomolgus monkey SIRP β 1 isoform 1;
 - e) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
 - f) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
 - g) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
 - h) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;
 - i) induces neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
 - j) increases neutrophil-mediated tumor cell clearance *in vivo*;
 - k) increases TREM2 expression on macrophages *in vitro* and/or *in vivo*;
 - l) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
 - m) increases viability of dendritic cells *in vitro* and/or *in vivo*.
2. The isolated antibody of claim 1, wherein the antibody does not bind to human SIRP α or SIRP γ .
3. The isolated antibody of claim 1 or claim 2, wherein the antibody has at least one, at least two, or at least three properties selected from:
 - a) agonizes SIRP β 1 activity on CD14-positive monocytes *in vitro* and/or *in vivo*;
 - b) induces or increases respiratory burst in immune cells, such as neutrophils and/or monocytes *in vitro* and/or *in vivo*;
 - c) induces or increases IL-8 expression in monocytes *in vitro* and/or *in vivo*;
 - d) induces or increases TNF α expression in macrophages and/or dendritic cells *in vitro* and/or *in vivo*;

- e) induces neutrophil-mediated phagocytosis, for example, of tumor cells *in vitro* and/or *in vivo*;
 - f) increases neutrophil-mediated tumor cell clearance *in vivo*;
 - g) increases TREM2 expression on macrophages *in vitro* and/or *in vivo*;
 - h) increases viability of macrophages *in vitro* and/or *in vivo*, alone and/or in combination with an agonist anti-TREM2 antibody; and
 - i) increases viability of dendritic cells *in vitro* and/or *in vivo*.
4. An isolated antibody that binds to human SIRP β 1, wherein the antibody comprises a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises (a) HVR-H1 comprising an amino acid sequence of SEQ ID NO 80; (b) HVR-H2 comprising an amino acid sequence of SEQ ID NO: 101; (c) HVR-H3 comprising an amino acid sequence of SEQ ID NO: 125; (d) HVR-L1 comprising an amino acid sequence of SEQ ID NO: 383; (e) HVR-L2 comprising an amino acid sequence of SEQ ID NO: 16; and (f) HVR-L3 comprising an amino acid sequence of SEQ ID NO: 31.
5. The antibody of claim 4, wherein the heavy chain variable region comprises one, two, three or four framework regions selected from VH FR1, VH FR2, VH FR3, and VH FR4 of SEQ ID NOs: 210, 210, 215, and 222, and/or wherein the light chain variable region comprises one, two, three or four framework regions selected from VL FR1, VL FR2, VL FR3, and VL FR4 of SEQ ID NOs: 174, 186, 192, and 200.
6. The antibody of any one of claims 1-5, wherein the antibody comprises a heavy chain variable region comprising an amino acid sequence at least 90%, at least 95%, at least 97%, or at least 99% identical to a sequence of SEQ ID NO: 268; and/or wherein the antibody comprises a light chain variable region comprising an amino acid sequence at least 90%, at least 95%, at least 97%, or at least 99% identical to a sequence of SEQ ID NO: 267.
7. The antibody of any one of claims 1-6, wherein the antibody comprises a heavy chain variable region of SEQ ID NO: 268; and/or wherein the antibody comprises a light chain variable region of SEQ ID NO: 267.
8. The antibody of any one of claims 1-6, wherein the antibody is a monoclonal antibody, and/or wherein the antibody is of the IgG class, the IgM class, or the IgA class.
9. The antibody of claim 8, wherein the antibody is of the IgG class and has an IgG1, IgG2, or IgG4 isotype.
10. The antibody of claim 9, wherein the antibody has an IgG1 isotype, optionally comprising a E430G substitution and a P331S substitution according to EU numbering.

11. The antibody of any one of claims 1-10, wherein the antibody is an antibody fragment, optionally comprising a Fab, Fab', Fab'-SH, F(ab')₂, Fv or scFv fragment.
12. The antibody of any one of claims 1-11, wherein the antibody has an affinity (K_D) for human SIRP β 1 isoform 1 of 0.1 nM to 50 nM, or 0.5 nM to 10 nM, or 0.5 nM to 5 nM, optionally wherein affinity is measured using a ForteBio Octet[®] system.
13. The antibody of any one of claims 1-12, wherein the antibody recognizes a first and a second antigen, wherein the first antigen is SIRP β 1 and the second antigen is:
- (a) an antigen facilitating transport across the blood-brain-barrier;
 - (b) an antigen facilitating transport across the blood-brain-barrier selected from the group consisting of transferrin receptor (TR), insulin receptor (HIR), insulin-like growth factor receptor (IGFR), low-density lipoprotein receptor related proteins 1 and 2 (LPR-1 and 2), and diphtheria toxin receptor;
 - (c) a disease-causing agent selected from the group consisting of disease-causing peptides or proteins or disease-causing nucleic acids, wherein the disease-causing nucleic acids are antisense GGCCCC (G2C4) repeat-expansion RNA, the disease-causing proteins are selected from the group consisting of amyloid beta, oligomeric amyloid beta, amyloid beta plaques, amyloid precursor protein or fragments thereof, Tau, IAPP, alpha-synuclein, TDP-43, FUS protein, C9orf72 (chromosome 9 open reading frame 72), c9RAN protein, prion protein, PrPSc, huntingtin, calcitonin, superoxide dismutase, ataxin, ataxin 1, ataxin 2, ataxin 3, ataxin 7, ataxin 8, ataxin 10, Lewy body, atrial natriuretic factor, islet amyloid polypeptide, insulin, apolipoprotein AI, serum amyloid A, medin, prolactin, transthyretin, lysozyme, beta 2 microglobulin, gelsolin, keratoepithelin, cystatin, immunoglobulin light chain AL, S-IBM protein, Repeat-associated non-ATG (RAN) translation products, DiPeptide repeat (DPR) peptides, glycine-alanine (GA) repeat peptides, glycine-proline (GP) repeat peptides, glycine-arginine (GR) repeat peptides, proline-alanine (PA) repeat peptides, ubiquitin, and proline-arginine (PR) repeat peptides;
 - (d) ligands and/or proteins expressed on immune cells, wherein the ligands and/or proteins selected from the group consisting of CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, PD-L1, CTLA-4, PD-L2, PD-1, B7-H3, B7-H4, HVEM, BTLA, KIR, GAL9, TIM3, A2AR, LAG-3, and phosphatidylserine; and
 - (e) a protein, lipid, polysaccharide, or glycolipid expressed on one or more tumor cells.

14. An isolated nucleic acid comprising a nucleic acid sequence encoding the antibody of any one of the preceding claims.
15. A vector comprising the nucleic acid of claim 14.
16. An isolated host cell comprising the vector of claim 15, wherein the host cell expresses the antibody of any one of claims 1-13.
17. A method of producing an antibody that binds to human SIRP β 1, comprising culturing the cell of claim 16 so that the antibody is produced, optionally further comprising recovering the antibody produced by the cell.
18. A pharmaceutical composition comprising the antibody of any one of claims 1-13 and a pharmaceutically acceptable carrier.
19. Use of an antibody of any one of claims 1-13 in the preparation of a medicament for treating cancer in an individual, optionally wherein the cancer is selected from sarcoma, bladder cancer, brain cancer, breast cancer, colon cancer, rectal cancer, endometrial cancer, kidney cancer, renal pelvis cancer, leukemia, lung cancer, small cell lung cancer, melanoma, lymphoma, pancreatic cancer, prostate cancer, ovarian cancer, and fibrosarcoma, glioblastoma multiforme; renal clear cell carcinoma; adrenocortical carcinoma; bladder urothelial carcinoma, diffuse large B-cell lymphoma, lung adenocarcinoma; pancreatic adenocarcinoma, renal cell cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, indolent B cell lymphoma, aggressive B cell lymphoma, T cell lymphoma, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), multiple myeloma, myelodysplastic syndromes, myeloproliferative neoplasms, breast invasive carcinoma, cervical squamous cell carcinoma, endocervical adenocarcinoma, cholangiocarcinoma, colon adenocarcinoma, diffuse large B-cell lymphoma, esophageal carcinoma, head and neck squamous cell carcinoma, kidney chromophobe, renal papillary cell carcinoma, lower grade glioma, hepatocellular carcinoma, lung squamous cell carcinoma, mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectal adenocarcinoma, cutaneous melanoma, stomach adenocarcinoma, testicular germ cell tumors, thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, and uveal melanoma.
20. The use of claim 19, wherein the individual is to be administered (a) a therapeutic agent that inhibits or agonizes PD1, PDL1, CD40, OX40, ICOS, CD28, CD137/4-1BB, CD27, GITR, CTLA4, PD-L2, B7-H3, B7-H4, HVEM, LIGHT, BTLA, CD30, TIGIT, VISTA, KIR, GAL9, TIM1, TIM3, TIM4, A2AR, LAG3, DR-5, CD2, CD5, CD39, or CD73; (b) at least one antibody that specifically binds to an inhibitory checkpoint molecule, and/or (c) one or more standard or

investigational anti-cancer therapies optionally wherein the at least one antibody that specifically binds to an inhibitory checkpoint molecule is selected from an anti-PD-L1 antibody, an anti-CTLA4 antibody, an anti-PD-L2 antibody, an anti-PD-1 antibody, an anti-B7-H3 antibody, an anti-B7-H4 antibody, and anti-HVEM antibody, an anti- B- and T-lymphocyte attenuator (BTLA) antibody, an anti-Killer inhibitory receptor (KIR) antibody, an anti-GAL9 antibody, an anti-TIM-1 antibody, an anti-TIM3 antibody, an anti-TIM-4 antibody, an anti-A2AR antibody, an anti-CD39 antibody, an anti-CD73 antibody, an anti-LAG-3 antibody, an anti-phosphatidylserine antibody, an anti-CD27 antibody, an anti-CD30 antibody, an anti-TNF α antibody, an anti-CD33 antibody, an anti-Siglec-5 antibody, an anti-Siglec-7 antibody, an anti-Siglec-9 antibody, an anti-Siglec-11 antibody, an antagonistic anti-TREM1 antibody, an antagonistic anti-TREM2 antibody, an anti-TIGIT antibody, an anti-VISTA antibody, an anti-CD2 antibody, an anti-CD5 antibody, and any combination thereof; and wherein the one or more standard or investigational anti-cancer therapies are selected from radiotherapy, cytotoxic chemotherapy, targeted therapy, imatinib therapy, trastuzumab therapy, etanercept therapy, adoptive cell transfer (ACT) therapy, chimeric antigen receptor T cell transfer (CAR-T) therapy, vaccine therapy, and cytokine therapy.

21. The use of claim 19 or claim 20, wherein the individual is to be administered at least one antibody that specifically binds to an inhibitory cytokine optionally selected from an anti-CCL2 antibody, an anti-CSF-1 antibody, an anti-IL-2 antibody, and any combination thereof.

22. The use of any one of claims 19-21, wherein the individual is to be administered at least one agonistic antibody that specifically binds to a stimulatory checkpoint protein optionally selected from an agonist anti-CD40 antibody, an agonist anti-OX40 antibody, an agonist anti-ICOS antibody, an agonist anti-CD28 antibody, an agonistic anti-TREM1 antibody, an agonistic anti-TREM2 antibody, an agonist anti-CD137/4-1BB antibody, an agonist anti-CD27 antibody, an agonist anti-glucocorticoid-induced TNFR-related protein GITR antibody, an agonist anti-CD30 antibody, an agonist anti-BTLA antibody, an agonist anti-HVEM antibody, an agonist anti-CD2 antibody, an agonist anti-CD5 antibody, and any combination thereof.

23. The use of any one of claims 19-22, wherein the individual is to be further administered at least one stimulatory cytokine optionally selected from IFN- α 4, IFN- β , IL-1 β , TNF- α , IL-6, IL-8, CRP, IL-20 family members, LIF, IFN- γ , OSM, CNTF, GM-CSF, IL-11, IL-12, IL-15, IL-17, IL-18, IL-23, CXCL10, IL-33, MCP-1, MIP-1-beta, and any combination thereof.