


Figure 1

Mean Induration Size by Treatment Group and Study Day




Treatment:

BCG = 5×10^5 CFU BCG (range $2 - 8 \times 10^5$),
 Group 1 = 5×10^3 CFU VPM1002 (range $2 - 8 \times 10^3$),
 Group 2 = 5×10^4 CFU VPM1002 (range $2 - 8 \times 10^4$),
 Group 3 = 5×10^5 CFU VPM1002 (range $2 - 8 \times 10^5$).

Figure 2

Mean Erythema Size by Treatment Group and Study Day



Treatment:

BCG = 5×10^5 CFU BCG (range $2 - 8 \times 10^5$),
 Group 1 = 5×10^3 CFU VPM1002 (range $2 - 8 \times 10^3$),
 Group 2 = 5×10^4 CFU VPM1002 (range $2 - 8 \times 10^4$),
 Group 3 = 5×10^5 CFU VPM1002 (range $2 - 8 \times 10^5$).

Figure 3


Mean Changes from Baseline for IFN- γ -Response after Stimulation with Ag85B in Naïve Subjects

Figure 4

**Mean Changes from Baseline for IFN- γ -Response after Stimulation with Ag85B
in Pre-Immunized Subjects**




Figure 5

Change from Baseline of Single Multifunctional CD4⁺ T cells in Naïve Subjects