

STAPLE-CONTAINING POLYPEPTIDES AND APPLICATION THEREOF

[0001] The present disclosure claims the priority of:

CN2021114005918, filed on November 19, 2021;

CN2022106482638, filed on June 8, 2022.

TECHNICAL FIELD

[0002] The present disclosure relates to a series of staple-containing polypeptides and use thereof. Specifically, the present disclosure discloses the polypeptides with sequences represented by formulas (I-1) to (I-5).

BACKGROUND






[0003] Overweight and obesity are serious health problems facing all mankind. They are often accompanied by other diseases such as coronary artery disease, hypertension, type 2 diabetes, nonalcoholic fatty liver disease, kidney disease and certain cancers. The World Health Organization (WHO) defines obesity as one of the top ten chronic diseases. Obesity, along with hypertension, hyperlipidemia and hyperglycemia are known as the “Quartet of Death” and may become the number one killer in the 21st century. Data from WHO show that the prevalence of obesity in China was approximately 6.2% in 2016. In addition, the Lancet published a survey report of worldwide adult body weight in 2016. The survey found that the population of worldwide obese adults has exceeded that of healthy adults, and China has surpassed the United States to become the country with the largest obese population in the world. The number of people suffering from diabetes, hypertension, cardiovascular disease and other diseases caused by overweight and obesity is increasing year by year, and ages of these people are getting younger.

[0004] Glucagon (GCG) is a hormone secreted by pancreas and binding to the glucagon receptor (GCGR) to produce physiological functions. Glucagon promotes the rise of blood sugar by increasing gluconeogenesis and glycogenolysis. In addition, GCG can also reduce the synthesis of fatty acid in liver adipose tissue and promote fat decomposition. Glucagon-like peptide 1 (GLP-1) is a hormone secreted by intestinal L cells. It can reduce body weight by


suppressing appetite and reducing food intake, as well as increasing energy consumption and promoting the thermogenesis of brown adipose tissue. On the basis of maintaining the efficacy of a GLP-1 agonist, the introduction of GCG activity will have the medicinal effects of: helping to further promote the secretion of insulin from pancreatic β cells; promoting the metabolism of brown adipose tissue; enhancing the β -oxidation of the fatty acids in liver and reducing the generation of lipids and cholesterol; improving the survival rate of cardiomyocytes; accelerating the lipid metabolism of white adipose tissue and reducing the fat content. The GLP-1/GCG dual-target synergy is likely to have better effect in improving blood sugar and weight loss than a single-target action. Therefore, the research on the GLP-1/GCG dual-target drugs in the treatment of obesity and related diseases is of great significance.



SUMMARY

[0005] The present disclosure provides a polypeptide with a sequence represented by the following formulas,






[0006] wherein

[0007] the structure of Aib is  ;

[0008] X₂ is selected from  and  ;

[0009] ¹Lys represents a modified lysine, and the modification is that the amino groups on the two lysine side chains are connected to  ;

[0010] X is selected from  and  , wherein “*” indicates the position connected to X₀;

[0011] X₀ is  ;

[0012] m is selected from 2 and 3;


[0013] n is selected from 8, 9 and 10;


[0014] p is selected from 1 and 2.

[0015] In some embodiments of the present disclosure, the above-mentioned m is 2, and other variables are as defined in the present disclosure.

[0016] In some embodiments of the present disclosure, the above-mentioned n is 9, and other variables are as defined in the present disclosure.


[0017] In some embodiments of the present disclosure, the above-mentioned p is 1, and other variables are as defined in the present disclosure.

[0018] In some embodiments of the present disclosure, the above-mentioned X₂ is  , and other variables are as defined in the present disclosure.

[0019] In some embodiments of the present disclosure, the above-mentioned X₀ is  , and other variables are as defined in the present disclosure.

is , and other variables are
as defined in the present disclosure.

[0022] The present disclosure also provides a polypeptide represented by the following formulas,



4

disease.

[0024] In some embodiments of the present disclosure, the GLP-1R/GCGR-related disease is selected from obesity and nonalcoholic steatohepatitis (NASH).

Technical effect

[0025] The compounds of the present disclosure have strong agonistic activity on GLP-1R/GCGR; the compounds of the present disclosure exhibit excellent weight-loss efficacy in DIO mice; the compounds of the present disclosure exhibit excellent NASH-improving efficacy in STZ-NASH mice; the compounds of the present disclosure have extremely high plasma protein binding and excellent plasma stability; the compounds of the present disclosure have excellent pharmacokinetic properties.

Definition and term

[0026] Unless otherwise specified, the following terms and phrases used herein are intended to have the following meanings. A specific term or phrase should not be considered indefinite or unclear in the absence of a particular definition, but should be understood in the conventional sense. When a trade name appears herein, it is intended to refer to its corresponding commodity or active ingredient thereof.


[0027] The term “pharmaceutically acceptable” is used herein in terms of those compounds, materials, compositions, and/or dosage forms, which are suitable for use in contact with human and animal tissues within the scope of reliable medical judgment, with no excessive toxicity, irritation, allergic reaction or other problems or complications, commensurate with a reasonable benefit/risk ratio.


[0028] The term “pharmaceutically acceptable salt” means a salt of compounds disclosed herein that is prepared by reacting the compound having a specific substituent disclosed herein with a relatively non-toxic acid or base. When compounds disclosed herein contain a relatively acidic functional group, a base addition salt can be obtained by bringing the compound into contact with a sufficient amount of base in a pure solution or a suitable inert solvent. When compounds disclosed herein contain a relatively basic functional group, an acid addition salt can be obtained by bringing the compound into contact with a sufficient amount of acid in a


pure solution or a suitable inert solvent. Some specific compounds disclosed herein contain both basic and acidic functional groups and can be converted to any base or acid addition salt.


[0029] The pharmaceutically acceptable salt disclosed herein can be prepared from the parent compound that contains an acidic or basic moiety by conventional chemical methods. Generally, such salt can be prepared by reacting the free acid or base form of the compound with a stoichiometric amount of an appropriate base or acid in water or an organic solvent or a mixture thereof.



[0030] “Amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that perform a similar function to the naturally occurring amino acids. The naturally occurring amino acids are those encoded by a genetic code, as well as those that are later modified, such as hydroxyproline, γ -carboxyglutamic acid, and O-phosphoserine. The amino acid analog refers to a compound that has the same basic chemical structure (e.g., an α carbon bonded to a hydrogen, a carboxyl group, an amino group, or an R group) as a naturally occurring amino acid, such as homoserine, norleucine, methionine sulfoxide, and methionine methylsulfonium. Such analog may have a modified R group (e.g., norleucine) or a modified peptide backbone, but retains the same basic chemical structure as the naturally occurring amino acid. The amino acid mimetic refers to a chemical compound that has a structure different from the general chemical structure of an amino acid but performs a similar function to a naturally occurring amino acid.


[0031] A or Ala disclosed herein represents alanine, with a structure of  ; R or


Arg represents arginine, with a structure of  ; N or Asn represents



asparagine, with a structure of  ; D or Asp represents aspartic acid, with a


structure of  ; C or Cys represents cysteine, with a structure of



 ; Q or Gln represents glutamine, with a structure of
 
 ; E or


Glu represents glutamic acid, with a structure of
 
 ; G or Gly represents



glycine, with a structure of
 
 ; H or His represents histidine, with a structure of



 ; I or Ile represents isoleucine, with a structure of
 
 ; L or


Leu represents leucine, with a structure of
 
 ; K or Lys represents lysine, with a


structure of
 
 ; M or Met represents methionine, with a structure of



 ; F or Phe represents phenylalanine, with a structure of


 ; P or Pro represents proline, with a structure of
 
 ; S or Ser

represents serine, with a structure of
 
 ; T or Thr represents threonine, with a

structure of
 
 ; W or Trp represents tryptophan, with a structure of


 ; Y or Tyr represents tyrosine, with a structure of



[0032] The term “treatment” includes inhibiting, slowing, stopping or reversing the existing symptoms or the progression or severity of a patient’s condition.

[0033] Unless otherwise specified, the term “isomer” is intended to include geometric isomers, cis- or trans- isomers, stereoisomers, enantiomers, optical isomers, diastereomers, and tautomers.

[0034] Compounds disclosed herein may be present in a specific geometric or stereoisomeric form. The present disclosure contemplates all such compounds, including cis and trans isomers, (-)- and (+)-enantiomers, (*R*)- and (*S*)-enantiomers, diastereoisomers, (*D*)-isomers, (*L*)-isomers, and a racemic mixture and other mixtures, for example, a mixture enriched in enantiomer or diastereoisomer, all of which are encompassed within the scope disclosed herein. The substituent such as alkyl may have an additional asymmetric carbon atom. All these isomers and mixtures thereof are encompassed within the scope disclosed herein.

[0035] Unless otherwise specified, the term “enantiomer” or “optical isomer” means stereoisomers that are in a mirrored relationship with each other.

[0036] Unless otherwise specified, the term “cis-trans isomer” or “geometric isomer” is produced by the inability of a double bond or a single bond between ring-forming carbon atoms to rotate freely.

[0037] Unless otherwise specified, the term “diastereomer” means a stereoisomer in which two or more chiral centers of are contained in a molecule and is in a non-mirrored relationship between molecules.

[0038] Unless otherwise specified, “(+)” means dextroisomer, “(-)” means levoisomer, and “(±)” means racemate.

[0039] Unless otherwise specified, a wedged solid bond () and a wedged dashed bond () indicate the absolute configuration of a stereocenter; a straight solid bond () and a straight dashed bond () indicate the relative configuration of a stereocenter; a wavy line () indicates a wedged solid bond () or a wedged dashed bond (); or a wavy line () indicates a straight solid bond () or a straight dashed bond ().

[0040] Unless otherwise specified, the term “enriched in one isomer”, “isomer enriched”, “enriched in one enantiomer” or “enantiomeric enriched” means that the content of one isomer or enantiomer is less than 100%, and the content of the isomer or enantiomer is 60% or more, or 70% or more, or 80% or more, or 90% or more, or 95% or more, or 96% or more, or 97% or more, or 98% or more, or 99% or more, or 99.5% or more, or 99.6% or more, or 99.7% or more, or 99.8% or more, or 99.9% or more.


[0041] Unless otherwise specified, the term “isomer excess” or “enantiomeric excess” means the difference between the relative percentages of two isomers or two enantiomers. For example, if one isomer or enantiomer is present in an amount of 90% and the other isomer or enantiomer is present in an amount of 10%, the isomer or enantiomeric excess (ee value) is 80%.


[0042] Optically active (*R*)- and (*S*)-isomer, or *D* and *L* isomer can be prepared using chiral synthesis or chiral reagents or other conventional techniques. If one kind of enantiomer of certain compound disclosed herein is to be obtained, the pure desired enantiomer can be obtained by asymmetric synthesis or derivative action of chiral auxiliary followed by separating the resulting diastereomeric mixture and cleaving the auxiliary group. Alternatively, when the molecule contains a basic functional group (such as amino) or an acidic functional group (such as carboxyl), the compound reacts with an appropriate optically active acid or base to form a salt of the diastereomeric isomer which is then subjected to diastereomeric resolution through the conventional method in the art to afford the pure enantiomer. In addition, the enantiomer and the diastereoisomer are generally isolated through chromatography which uses a chiral stationary phase and optionally combines with a chemical derivative method (for example, carbamate generated from amine).


[0043] Compounds disclosed herein may contain an unnatural proportion of atomic isotopes at one or more of the atoms that make up the compounds. For example, a compound may be labeled with a radioisotope such as tritium (³H), iodine-125 (¹²⁵I) or C-14(¹⁴C). For another example, hydrogen can be replaced by heavy hydrogen to form a deuterated drug. The bond between deuterium and carbon is stronger than that between ordinary hydrogen and carbon. Compared with undeuterated drugs, deuterated drugs have advantages of reduced toxic side effects, increased drug stability, enhanced efficacy, and prolonged biological half-life of drugs.

All changes in the isotopic composition of compounds disclosed herein, regardless of radioactivity, are included within the scope of the present disclosure.


[0044] When an enumerated linking group does not indicate its linking direction, its linking

direction is arbitrary. For example, when the linking group L in  is -M-W-, the -M-W- can be linked to the ring A and the ring B in the same direction as the reading order

from left to right to constitute , or can be linked to the ring A and the ring B in the reverse direction as the reading order from left to right to constitute

. A combination of the linking groups, substituents and/or variants thereof is allowed only when such combination can result in a stable compound.

[0045] Unless otherwise specified, when a group has one or more connectable sites, any one or more sites of the group can be connected to other groups through chemical bonds. When the connection position of the chemical bond is variable, and there is H atom(s) at a connectable site(s), with the connection of the chemical bond, the number of H atom(s) at this site will correspondingly decrease as the number of the connected chemical bond increases, and the group will become a group of corresponding valence. The chemical bond between the site and other groups can be represented by a straight solid bond () , a straight dashed bond () , or a wavy line () . For example, the straight solid bond in -OCH₃ indicates that the group is connected to other groups through the oxygen atom in the group; the straight dashed bond in  indicates that the group is connected to other groups through two ends of the nitrogen

atom in the group; the wavy line in  indicates that the group is connected to other groups through the 1- and 2-carbon atoms in the phenyl group.

[0046] Unless otherwise specified, the term “C₁₋₃ alkyl” is used to indicate a linear or branched saturated hydrocarbon group consisting of 1 to 3 carbon atoms. The C₁₋₃ alkyl group

includes C₁₋₂ and C₂₋₃ alkyl groups and the like. It may be monovalent (e.g., methyl), divalent (e.g., methylene) or multivalent (e.g., methenyl). Examples of C₁₋₃ alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (including *n*-propyl and isopropyl), and the like.

[0047] The structures of compounds disclosed herein can be confirmed by conventional methods well known to those skilled in the art. If the present disclosure relates to an absolute configuration of a compound, the absolute configuration can be confirmed by conventional techniques in the art, such as single crystal X-Ray diffraction (SXRD). In a single crystal X-Ray diffraction (SXRD), the diffraction intensity data of a cultivated single crystal are collected using a Bruker D8 venture diffractometer with a light source of CuK α radiation in a scanning mode of ϕ/ω scan; after collecting the relevant data, the crystal structure is further analyzed by the direct method (Shelxs97) to confirm the absolute configuration.

[0048] Compounds disclosed herein can be prepared by a variety of synthetic methods well known to those skilled in the art, including the following enumerated embodiment, the embodiment formed by the following enumerated embodiment in combination with other chemical synthesis methods, and equivalents well known to those skilled in the art. Alternative embodiments include, but are not limited to examples disclosed herein.

[0049] Solvents used in the present disclosure are commercially available.

[0050] The following abbreviations are used in the present disclosure: aq represents aqueous; eq represents equivalent or equivalence; DCM represents dichloromethane; PE represents petroleum ether; DMSO represents dimethyl sulfoxide; MeOH represents methanol; Boc represents tert-butoxycarbonyl, which is an amine protecting group; Dde represents (4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl, which is a side chain protecting group for amino acids; r.t. represents room temperature; O/N represents overnight; THF represents tetrahydrofuran; Boc₂O represents di-tert-butyl dicarbonate; TFA represents trifluoroacetic acid; DIEA represents diisopropylethylamine; DMF represents N,N-dimethylformamide; HBTU represents O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate; HOBT represents 1-hydroxybenzotriazole; HOAT represents 1-hydroxy-7-azabenzotriazole; DIC represents N,N'-diisopropylcarbodiimide; DBU represents 1,8-diazabicyclo[5.4.0]undec-7-ene; PhSiH₃ represents phenylsilane; Pd(PPh₃)₄ represents


tetrakis(triphenylphosphine)palladium; AEEA represents 2-(2-(2-aminoethoxy)ethoxy)acetic acid; DIEA represents diisopropylethylamine.

[0051] Compounds are named according to general naming principles in the art or by ChemDraw® software, and commercially available compounds are named with their vendor directory names.

DETAILED DESCRIPTION


[0052] The present disclosure is described in detail below by means of examples. However, it is not intended that these examples have any disadvantageous limitations to the present disclosure. The present disclosure has been described in detail herein, and embodiments are also disclosed herein. It will be apparent to those skilled in the art that various changes and modifications may be made to the embodiments disclosed herein without departing from the spirit and scope disclosed herein.

Example 1



WX001

[0053] Synthesis of intermediate 1



Intermediate 1

[0054] 1. Attachment of resin to intermediate 1

[0055] 1.1 5.0 g of 2-chlorotriyl chloride resin (2-CTC Resin, degree of substitution S=1.00 mmol/g) and 1.93 g of Fmoc-AEEA-OH were weighed and added to a reaction column, and DCM (40 mL) was further added. Then, DIEA (3.5 mL) was added to the reaction column and the system was blown with nitrogen for 2 h. Then, MeOH (5 mL) was added to the reaction column and the system was further blown with nitrogen for 30 min. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 minute each

time, and the reaction column was drained until no liquid flowed out.

[0056] 1.2 20% piperidine/DMF (100 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (100 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0057] 2. Coupling of amino acids

[0058] 2.1 Coupling of Fmoc-AEEA-OH

[0059] 1. Fmoc-AEEA-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (30 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0060] 2. The mixture was reacted in an environment of 25 °C for 20 minutes. The resin was tested with ninhydrin and was colorless and transparent.

[0061] 3. The reaction solution was aspirated. The reaction column was washed 5 times with DMF, for 1 min each time, and drained until no liquid flowed out.

[0062] 2.2 Coupling of Fmoc-Glu-OtBu

[0063] 1. 20% piperidine/DMF (100 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0064] 2. Fmoc-Glu-OtBu (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (30 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0065] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0066] 4. The reaction solution was aspirated. The reaction column was washed with DMF (100 mL) 5 times, for 1 min each time, and drained until no liquid flowed out.

[0067] 2.3 Coupling of 20-(tert-butoxy)-20-oxoicosanoic acid

[0068] 1. 20% piperidine/DMF (100 mL) was added to the reaction column and the system

was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (100 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0069] 2. 20-(tert-butoxy)-20-oxoicosanoic acid (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (30 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0070] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0071] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times, for 1 min each time, and drained until no liquid flowed out.

[0072] 3. Cleavage and drying of crude peptide

[0073] 3.1. The cleavage solution was prepared according to the following volume:

[0074] HFIP/DCM=20/80.

[0075] 3.2. 100 mL of the prepared cleavage solution was poured into a reactor containing the dried peptide resin. The reactor was blown for 20 minutes. The mixture was filtered, and the filtrate was added into a flask. This operation was repeated twice. The cleavage solution collected twice was rotary evaporated to dryness to give 4.3 g of crude peptide.

[0076] Synthesis of WX001

[0077] 1. Attachment of resin

[0078] 1.1 1.43 g of 4-(2',4'-dimethoxyphenyl-Fmoc-aminomethyl)-phenoxyacetamidomethyldiphenylmethylaniline resin (Rink Amide MBHA Resin, degree of substitution Sub=0.28 mmol/g) was weighed and added to a reaction column. Then, DCM (50 mL) was added to the reaction column and the system was blown with nitrogen for 2 h. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 minute each time, and the reaction column was drained until no liquid flowed out.

[0079] 1.2 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column

was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0080] 2. Coupling of amino acids

[0081] 2.1 Coupling of Fmoc-Gly-OH

[0082] 1. Fmoc-Gly-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0083] 2. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0084] 3. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[0085] 2.2 Coupling of Fmoc-Ser(tBu)-OH

[0086] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0087] 2. Fmoc-Ser(tBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0088] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0089] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[0090] 2.3 Coupling of Fmoc-Ser(tBu)-OH

[0091] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0092] 2. Fmoc-Ser(tBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0

eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0093] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0094] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times, for 1 min each time, and drained until no liquid flowed out.

[0095] 2.4 Coupling of Fmoc-Pro-OH

[0096] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[0097] 2. Fmoc-Pro-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[0098] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[0099] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00100] 2.5 Coupling of Fmoc-Gly-OH

[00101] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with chloranil and showed blue.

[00102] 2. Fmoc-Gly-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00103] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested

with chloranil and was colorless and transparent.

[00104] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00105] 2.6 Coupling of Fmoc-Gly-OH

[00106] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00107] 2. Fmoc-Gly-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00108] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00109] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00110] 2.7 Coupling of Fmoc-Glu(OtBu)-OH

[00111] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00112] 2. Fmoc-Glu(OtBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00113] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00114] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00115] 2.8 Coupling of Fmoc-Leu-OH

[00116] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00117] 2. Fmoc-Leu-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00118] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00119] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00120] 2.9 Coupling of Fmoc-Leu-OH

[00121] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00122] 2. Fmoc-Leu-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00123] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00124] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00125] 2.10 Coupling of Fmoc-Trp(Boc)-OH

[00126] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00127] 2. Fmoc-Trp(Boc)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00128] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00129] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00130] 2.11 Coupling of Fmoc-Glu(OtBu)-OH

[00131] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00132] 2. Fmoc-Glu(OtBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00133] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00134] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00135] 2.12 Coupling of Fmoc-Val-OH

[00136] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00137] 2. Fmoc-Val-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00138] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00139] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00140] 2.13 Coupling of Fmoc-Phe-OH

[00141] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00142] 2. Fmoc-Phe-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00143] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00144] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00145] 2.14 Coupling of Fmoc-Glu(OtBu)-OH

[00146] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00147] 2. Fmoc-Glu(OtBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00148] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00149] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00150] 2.15 Coupling of Fmoc-Lys(Dde)-OH

[00151] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00152] 2. Fmoc-Lys(Dde)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00153] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00154] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00155] 2.16 Coupling of Fmoc-Ala-OH

[00156] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00157] 2. Fmoc-Ala-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00158] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00159] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00160] 2.17 Coupling of Fmoc-Lys(Boc)-OH

[00161] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column

was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00162] 2. Fmoc-Lys(Boc)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00163] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00164] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00165] 2.18 Coupling of Fmoc-Lys(Alloc)-OH

[00166] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00167] 2. Fmoc-Lys(Alloc)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00168] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00169] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00170] 2.19 Coupling of Fmoc-Glu(OtBu)-OH

[00171] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00172] 2. Fmoc-Glu(OtBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was

added. The nitrogen was adjusted so that the resin was evenly blown.

[00173] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00174] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00175] 2.20 Coupling of Fmoc-Asp(OtBu)-OH

[00176] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00177] 2. Fmoc-Asp(OtBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00178] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00179] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00180] 2.21 Coupling of Fmoc-Leu-OH

[00181] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00182] 2. Fmoc-Leu-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00183] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00184] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5

times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00185] 2.22 Coupling of Fmoc-Tyr(tBu)-OH

[00186] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00187] 2. Fmoc-Tyr(tBu)-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00188] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00189] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00190] 2.23 Coupling of Fmoc-Lys(Boc)-OH

[00191] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00192] 2. Fmoc-Lys(Boc)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00193] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00194] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00195] 2.24 Coupling of Fmoc-Ser(tBu)-OH

[00196] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid

flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00197] 2. Fmoc-Ser(tBu)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00198] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00199] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00200] 2.25 Coupling of Fmoc-Tyr(tBu)-OH

[00201] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00202] 2. Fmoc-Tyr(tBu)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00203] 3. The mixture was reacted in an environment of 25 °C for 2 h. The resin was tested with ninhydrin and showed blue.

[00204] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00205] 2.26 Coupling of Fmoc-Asp(OtBu)-OH

[00206] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00207] 2. Fmoc-Asp(OtBu)-OH (6.0 eq) was weighed and added to the above resin. HOAT (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction

column was blown with nitrogen. After the amino acid and HOAT were dissolved, DIC (6.0 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00208] 3. The mixture was reacted in an environment of 25 °C for 1 h. The resin was tested with ninhydrin and was colorless and transparent.

[00209] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00210] 2.27 Coupling of Fmoc-Ser(tBu)-OH

[00211] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00212] 2. Fmoc-Ser(tBu)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00213] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00214] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00215] 2.28 Coupling of Fmoc-Thr(tBu)-OH

[00216] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00217] 2. Fmoc-Thr(tBu)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00218] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00219] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00220] 2.29 Coupling of Fmoc-Phe-OH

[00221] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00222] 2. Fmoc-Phe-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00223] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00224] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00225] 2.30 Coupling of Fmoc-Thr(tBu)-OH

[00226] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00227] 2. Fmoc-Thr(tBu)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00228] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00229] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00230] 2.31 Coupling of Fmoc-Gly-OH

[00231] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system

was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00232] 2. Fmoc-Gly-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.00 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00233] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00234] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00235] 2.32 Coupling of Fmoc-Gln(Trt)-OH

[00236] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00237] 2. Fmoc-Gln(Trt)-OH (6.0 eq) was weighed and added to the above resin. HOBT (6.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid and HOBT were dissolved, DIC (6.0 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00238] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00239] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00240] 2.33 Coupling of Fmoc-Aib-OH

[00241] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00242] 2. Fmoc-Aib-OH (3.0 eq) was weighed and added to the above resin. DIEA (6.0 eq)

was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (2.85 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00243] 3. The mixture was reacted in an environment of 25 °C overnight. The resin was tested with ninhydrin and was colorless and transparent.

[00244] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00245] 2.34 Coupling of Boc-His(Trt)-OH

[00246] 1. 20% piperidine/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with chloranil and showed blue.

[00247] 2. Boc-His(Trt)-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HATU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00248] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with chloranil and was colorless and transparent.

[00249] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00250] 2.35 Removal of Alloc

[00251] 1. PhSiH₃ (10.0 eq) and DCM (10 mL) were added to the reaction column and the system was blown with nitrogen. Then, Pd(PPh₃)₄ (0.1 eq) was added and the system was blown with nitrogen for 20 minutes. The mixture was reacted twice, and the reaction column was drained until no liquid flowed out.

[00252] 2. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00253] 2.36 Coupling of Fmoc-Ida-OH

[00254] 1. Fmoc-Ida-OH (6.0 eq) was weighed and added to the above resin. DIEA (12.0 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column

was blown with nitrogen. After the amino acid was dissolved, HBTU (5.70 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00255] 2. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00256] 3. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00257] 2.37 Coupling of intermediate 1

[00258] 1. 10% DBU/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 20 minutes. The reaction column was drained until no liquid flowed out. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00259] 2. **Intermediate 1** (1.50 eq) was weighed and added to the above resin. DIEA (3.00 eq) was added, and DMF (10 mL) was further added to the reaction column. The reaction column was blown with nitrogen. After the amino acid was dissolved, HBTU (1.45 eq) was added. The nitrogen was adjusted so that the resin was evenly blown.

[00260] 3. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00261] 4. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00262] 2.38 Removal of Dde

[00263] 1. 3% hydrazine hydrate/DMF (50 mL) was added to the reaction column and the system was blown with nitrogen for 15 minutes. The reaction column was drained. DMF (50 mL) was added to wash 5 times, for 1 min each time. The reaction column was drained until no liquid flowed out. The resin was tested with ninhydrin and showed blue.

[00264] 2.39 Closing of amide ring

[00265] 1. DIEA (3.0 eq) was added to the DMF solution of the above resin. Then, HATU (1.5 eq) dissolved in DMF was slowly added dropwise to the reaction column and the system was blown with nitrogen. The nitrogen was adjusted so that the resin was evenly blown.

[00266] 2. The mixture was reacted in an environment of 25 °C for 0.5 h. The resin was tested with ninhydrin and was colorless and transparent.

[00267] 3. The reaction solution was aspirated. The reaction column was washed with DMF 5 times (50 mL each time), for 1 min each time, and drained until no liquid flowed out.

[00268] 4. MeOH (50 mL) was used to shrink the resin for 3 minutes each time. The reaction column was drained until no liquid flowed out. The resin was poured out and dried for later use.


[00269] 3. Cleavage and drying of crude peptide

[00270] 3.1. The cleavage solution was prepared according to the following volume:

[00271] TFA/triisopropylsilane/H₂O/3-mercaptopropionic acid=90/2.5/2.5/5.

[00272] 3.2 The dried peptide resin was added to the prepared cleavage solution. The mixture was shaken on a shaker for 2.5 hours and filtered. The filtrate was added to 10 times the volume of ice-cold isopropyl ether. The mixture was centrifuged, washed 5 times with isopropyl ether, and vacuum dried for 2 h to give a crude peptide. The crude peptide was separated and purified by preparative HPLC (purification step: mobile phase acetonitrile/water (40%/60%), 0.075% TFA; salt conversion step: mobile phase acetonitrile/water (20%/80%), 0.01% ammonium acetate) to give polypeptide WX001. The molecular weight of the polypeptide was confirmed by ESI-MS, with a calculated value of 4660.1 and a measured value of 4660.2.


Example 2



WX002

[00273] Polypeptide **WX002** was obtained by referring to the synthesis of **WX001**. The molecular weight of the polypeptide was confirmed by ESI-MS, with a calculated value of 4659.2 and a measured value of 4659.0.


Example 3



WX003

[00274] Polypeptide **WX003** was obtained by referring to the synthesis of **WX001**. The molecular weight of the polypeptide was confirmed by ESI-MS, with a calculated value of 4659.2 and a measured value of 4659.3.


Example 4



WX004

[00275] Polypeptide **WX004** was obtained by referring to the synthesis of **WX001**. The molecular weight of the polypeptide was confirmed by ESI-MS, with a calculated value of 4659.2 and a measured value of 4659.6.

Example 5



WX005

[00276] Polypeptide **WX005** was obtained by referring to the synthesis of **WX001**. The molecular weight of the polypeptide was confirmed by ESI-MS, with a calculated value of 4658.3 and a measured value of 4658.7.

Biological assay data

Assay example 1: Assay of agonistic activity on GLP-1R/GIPR/GCGR *in vitro*

[00277] **A: Main materials:**

[00278] **(I) Cell line**

[00279] The cell lines were constructed by Shanghai WuXi AppTec. See Table 1 for details.

Table 1: Information of cell lines

Target	Host cell	Cloning
GLP-1R	HEK293	N/A

GCGR	HEK293	N/A
GIPR	CHO	N/A

[00280] (II) Reagents and consumables

[00281] See Table 2 for details.

Table 2: Information of reagents and consumables

Name	Batch	Item number	Manufacturer
cAMP assay kit	29F	62AM4PEJ	Cisbio
1M HEPES	2120919	15630-106	Invitrogen
Hanks' Balanced Salt Solution (HBSS)	2185775	14025	Invitrogen
Human serum albumin (HSA)	SLCF7301	A1653-10G	Sigma
Casein	SLCC9458	C4765-10mL	Sigma
3-isobutyl-1-methylxanthine (IBMX)	STBF6061V	I5879-5G	Sigma
ECHO qualified 384-well plate	0006433672	PP-0200	Labcyte
OptiPlate-384	8210-19481	6007299	PerkinElmer

[00282] (III) Instrument

[00283] See Table 3 for details.

Table 3: Information of instrument

Name	Model	Manufacturer
EnVision	envision2014	PerkinElmer
Vi-cell counter	Vi-CELL™ XR Cell Viability Analyzer	Beckman
Bravo	Bravo V11	Agilent
ECHO	ECHO 555	Labcyte
Centrifuge	Allegra™ 25R Centrifuge	Beckman

[00284] B. Method

[00285] i) Assay materials

[00286] The assay buffers are shown in Table 4.

Table 4: Information of assay buffers

<u>Reagent</u>	<u>Storage concentration</u>	<u>Volume</u>	<u>Final concentration</u>
Hanks' Balanced Salt Solution	1x	48.7 mL/ 44.7mL	≈1x
HEPES buffer	1mol/L	250 µL	5mmol/L
5% casein solution (HEPES)/ 10% human serum albumin solution (HSA)	5%/ 10%	1000 µL/ 5000µL	0.10%/ 1%
3-isobutyl-1-methylxanthine (IBMX)	500mmol/L	50 µL	0.5mmol/L

[00287] Preparation of the assay reagent is shown in Table 5.

Table 5: Information of the assay reagent preparation

<u>Reagent</u>	<u>Storage concentration</u>	<u>Volume</u>	<u>Final concentration</u>
Cell lysates	1x	9.5 mL	≈1x
D2-cAMP solution	40x	250 µL	1x
cAMP-antibody solution	40x	250 µL	1x

[00288] ii) Assay method

[00289] a) Preparation of compound plate:

[00290] The compounds to be assayed were serially diluted 4-fold with Bravo to obtain 10 concentrations, with a starting concentration of 30 µM.

[00291] b) Transfer of compound:

[00292] 1) 100 nL of the compound was transferred to an OptiPlate-384 plate using Echo.

[00293] 2) The OptiPlate-384 plate was centrifuged at 1000 rpm for 5 seconds.

[00294] c) Preparation of cell suspension

[00295] 1) A GLP-1R/GIPR/GCGR cell cryopreservation tube was quickly placed in warm water at 37 °C to thaw.

[00296] 2) The cell suspension was transferred to a 15 mL centrifuge tube, and gently washed with 10 ml of HBSS.

[00297] 3) The centrifuge tube was centrifuged at 1000 rpm at room temperature for 1 minute.

[00298] 4) The supernatant was discarded.

[00299] 5) The cells at the bottom were gently dispersed and gently washed with 10 mL of HBSS. The cells were centrifuged and sedimented, and finally resuspended with the assay buffer.

[00300] 6) Vi-cell was used to measure the cell density and viability.

[00301] 7) The concentration of GLP-1R/GIPR/GCGR cells was diluted to 2.0×10^5 /mL with the assay buffer.

[00302] 8) 100 nL of the diluted cell suspension was transferred to the OptiPlate-384 plate.

[00303] 9) The plate was incubated at room temperature for 30 minutes.

[00304] d) Addition of assay reagent:

[00305] 1) 10 μ L of 800nM gradient diluted cAMP standard was added into an empty well of the OptiPlate-384 plate.

[00306] 2) 10 μ L of cAMP assay reagent was added.

[00307] 3) The OptiPlate-384 plate was covered with a TopSeal-A film and incubated at room temperature for 60 minutes.

[00308] The TopSeal-A was peeled off and the plate was read on the EnVision.

[00309] **C Assay results**

[00310] The assay results are shown in Table 6.

Table 6: Assay results of in vitro agonistic activity on GLP-1R/GCGR/GIPR

ID	Assay results of the activity								
	Agonistic activity on GLP-1R			Agonistic activity on GCGR			Agonistic activity on GIPR		
	EC ₅₀ (nM)	Max Dose (nM)	Max Activity (%)	EC ₅₀ (nM)	Max Dose (nM)	Max Activity (%)	EC ₅₀ (nM)	Max Dose (nM)	Max Activit y (%)
WX001	0.02271	20	122.64	0.02669	100	98.93	>20	20	2.26
WX002	0.01692	20	106.52	0.1948	100	144.14	>20	20	4.06
WX003	0.05822	20	111.18	0.6244	100	86.01	>20	20	1.57
WX004	0.04459	20	85.77	0.05312	100	68.62	>20	20	4.12

WX005	0.1122	20	104.24	2.993	100	140.57	>20	20	1.42
-------	--------	----	--------	-------	-----	--------	-----	----	------

[00311] Conclusion: The compounds of the present disclosure have very strong agonistic activity on GLP-1R/GCGR, but have no agonistic activity on GIPR.

Assay example 2: Drug efficacy study in DIO mice-in vivo drug efficacy evaluation

[00312] A. Purpose of the assay

[00313] The weight loss effect of the assay compound in DIO mice was studied.

[00314] B. Procedure of the assay

[00315] 1. After arriving at the facility, the DIO mice were raised in an animal breeding room with strictly controlled environmental conditions. The temperature in the breeding room was maintained at 20 to 24 °C, and the humidity was maintained at 30 to 70%. The temperature and humidity in the breeding room were monitored in real time using a thermo-hygrometer, and the temperature and humidity were recorded twice a day (once in the morning and once in the afternoon). The lighting in the animal breeding room was controlled by an electronic timed lighting system. The lights were on for 12 hours and off for 12 hours every day (turned on at 7:00 am and turned off at 19:00 pm). During the assay, the animals were raised in individual cages and toys were provided in each cage. During the assay, the animals had free access to food (growth/breeding feed for rats and mice) and water.

[00316] 2. The animals in each group were subcutaneously injected with a solvent or an assay compound (10 nmol/kg), respectively. The administration time was 9:30 in the morning. The administration frequency was once every three days, and the administration cycle was 21 days.

[00317] C. Assay results

[00318] The assay results are shown in Table 7.

Table 7: Drug efficacy of the assay compound in DIO mice

Compound number	Solvent	WX001	WX004
Δ Body weight % (comparison of the body weight after 21 days with the body weight on day 1)	3.28%	-28.91%	-31.89%

[00319] Conclusion: The compounds of the present disclosure exhibit excellent weight loss

efficacy in DIO mice.

Assay example 3. Assay of plasma protein binding (PPB)

[00320] A. Purpose of the assay

[00321] The binding degree of the assay compound to human/mouse plasma albumin was studied.

[00322] B. Procedure of the assay

[00323] 1. Matrix preparation: on the day of the assay, plasma was thawed in cold water and centrifuged at 3220 rpm for 5 min to remove all clots. The pH of the resulting plasma was measured and adjusted to 7.4 ± 0.1 using 1% phosphoric acid or 1N sodium hydroxide as needed.

[00324] 2. Dilution procedure for the assay compound: the assay compound was dissolved in dimethyl sulfoxide (DMSO) to prepare stock solutions with concentrations of 10 mM and 2 mM, respectively. A 40 μ M working solution was prepared by diluting 2 μ L of stock solution (2 mM) with 98 μ L of DMSO. A 400 μ M working solution of the control compound was prepared by diluting 10 μ L of stock solution with 240 μ L of DMSO. The working solution of the compound (5 μ L) was mixed well with a blank matrix (995 μ L) at a ratio of 1:200 to prepare a loading matrix.

[00325] 3. Analysis steps

[00326] 3.1 An equal volume of 30 μ L of loading matrix (n=2) was transferred to a sample collection plate to prepare a time 0 (T0) sample for residue determination. The sample was immediately matched with the corresponding blank buffer to a final volume of 60 μ L, and the volume ratio of plasma to buffer in each well was 1:1. Then, 60 μ L of 4% H₃PO₄ in H₂O and 480 μ L of stop solution containing the internal standard were added to the T0 sample of the assay compound. They were then stored with other samples at 2-8 °C for further processing.

[00327] 3.2 The remaining plasma samples were pre-incubated in a carbon dioxide incubator at 37 ± 1 °C for 30 min. Protein-free samples (F samples) and samples loaded with matrix (230 μ L) were all transferred into polycarbonate tubes (n = 2) and ultracentrifuged at 37 °C and $155,000 \times g$ (35,000 rpm) for 4 h.

[00328] 3.3 To prepare T samples (assay samples), an additional matrix-containing sample was

transferred to a separate 96-well plate (sample incubation plate) and incubated at 37 °C for 4 h.

[00329] 3.4 At the end of centrifugation, 30 µL of protein-free samples and 30 µL of T samples were transferred from the second layer of the supernatant (below the top layer) to a new sample collection plate. Each sample was mixed with the corresponding blank buffer or matrix to a final volume of 60 µL with a matrix:buffer volume ratio of 1:1. 60 µL of 4% H₃PO₄ aqueous solution and 480 µL of stop solution (with internal standard) were added to all samples. The mixture was centrifuged at 4000 rpm for 20 min and 100 µL of supernatant from each sample was analyzed by LC-MS/MS.

[00330] C. Assay results

[00331] The assay results are shown in Table 8:

Table 8: Assay results of PPB

Compound number	WX001	WX004
PPB% unbound (human/mouse)	NA/NA	NA/NA

[00332] Note: NA indicates that the plasma protein binding is too high, and no free drug is detected at a normal plasma protein concentration.

[00333] Conclusion: The compounds of the present disclosure have extremely high plasma protein binding.

Assay example 4: Plasma stability assay (PLS)

[00334] A. Purpose of the assay

[00335] Stability of the assay compound in normal mouse plasma was studied.

[00336] B. Procedure of the assay

[00337] 1. Before the assay, the coagulated frozen plasma was thawed in a 37 °C water bath. The plasma was centrifuged at 4000 rpm for 5 minutes. If there were blood clots, the blood clots were removed. The pH value was adjusted to 7.4±0.1.

[00338] 2. Preparation of assay compound solution: An assay compound was dissolved in DMSO to prepare a 100 µM solution.

[00339] 3. 98 µL of blank control plasma was added to 2 µL of the assay compound solution

(100 μ M), so that the final concentration of the mixed solution reached 2 μ M. The mixed solution was incubated in a 37 °C water bath.

[00340] 4. 100 μ L of H₃PO₄ solution and 800 μ L of stop solution (a solution of 200 ng/mL tolbutamide and 200 ng/mL labetalol in 100% methanol) were added at each time point (0, 10, 30, 60 and 120 min), respectively, to precipitate proteins, and mixed thoroughly.

[00341] 5. The sample was centrifuged at 4000 rpm for 20 min. 100 μ L of supernatant from each well was taken for LC-MS/MS analysis.

[00342] C. Assay results

[00343] The assay results are shown in Table 9.

Table 9: Assay results of PLS

Compound number	WX001
PLS (human/mouse) T _{1/2} (min)	>289/>289

[00344] Conclusion: The compound of the present disclosure has excellent plasma stability.

Assay example 5: Pharmacokinetic evaluation of the compound in mice

[00345] A. Purpose of the assay

[00346] Pharmacokinetics of the compound in C57BL/6 mice were assayed.

[00347] B. Procedure of the assay

[00348] The pharmacokinetic characteristics of the compound in rodents after a subcutaneous injection were assayed using the standard protocol. In the assay, the candidate compound was formulated into a clear solution and given to mice by a single subcutaneous injection (SC, 0.048 mpk). The solvent of the subcutaneous injection was citrate buffer (20 mM, pH=7). Whole blood was collected, and plasma was prepared. Drug concentration was analyzed by LC-MS/MS method, and pharmacokinetic parameters were calculated using Phoenix WinNonlin software.

[00349] C. Assay results

[00350] The assay results are shown in Table 10.

Table 10: Pharmacokinetic assay results

Compound	Maximum	T _{max} (h)	Half-life T _{1/2}	Integral of the SC
----------	---------	----------------------	----------------------------	--------------------

number	plasma concentration (nM)		(h)	concentration AUC_{0-72h}(nM.hr)
WX001	46	8	21	1809
WX004	36	12	22	1161

[00351] Conclusion: The compounds of the present disclosure have excellent pharmacokinetic properties in mice.

Assay example 6: Pharmacokinetic evaluation of the compound in cynomolgus monkeys

[00352] A. Purpose of the assay

[00353] Pharmacokinetics of the compound in cynomolgus monkeys were assayed.

[00354] B. Procedure of the assay

[00355] The pharmacokinetic characteristics of the compound in mammals after a subcutaneous injection were assayed using the standard protocol. In the assay, the candidate compound was formulated into a clear solution and given to cynomolgus monkeys by a single subcutaneous injection (SC, 0.02 mpk). The solvent of the subcutaneous injection was citrate buffer (20 mM, pH=7). Whole blood was collected, and plasma was prepared. Drug concentration was analyzed by LC-MS/MS method, and pharmacokinetic parameters were calculated using Phoenix WinNonlin software.

[00356] C. Assay results

[00357] The assay results are shown in Table 11.

Table 11: Pharmacokinetic assay results

Compound number	Maximum plasma concentration (nM)	T_{max} (h)	Half-life T_{1/2} (h)	Integral of the SC concentration AUC_{0- 240h}(nM.hr)
WX001	43	24	93	5510
WX004	48	32	67	5719

[00358] Conclusion: The compounds of the present disclosure have excellent pharmacokinetic

properties in monkeys.

Assay example 7: Verification of in vivo drug efficacy in STZ-NASH mouse model

[00359] A. Purpose of the assay

[00360] Drug efficacy of the assay substance in the NASH model induced by streptozotocin (STZ) + high-fat diet (HFD) in C57BL/6 mice was verified.

[00361] B. Procedure of the assay

[00362] Modeling method: Newborn mice were subcutaneously injected with STZ (200 µg/mouse) within 48 hours after birth. After 4 weeks of breastfeeding, animals with a fasting blood glucose level of >12 mmol/L were selected and subjected to a HFD feeding for 6 consecutive weeks to finally establish the NASH model. Another 8 animals were selected without STZ injection and HFD feeding (normal control group).

[00363] Administration scheme: Administration was started after one week of HFD feeding, and the first day of administration was set as Day 1. Then, the mice were subcutaneously injected every two days for 5 consecutive weeks.

[00364] Detection at the endpoint of the assay: Pathological hematoxylin-eosin staining and picro sirius red staining were performed.

[00365] C. Assay results

[00366] The assay results are shown in Table 13.

Table 13 Animal pathological indicators of the STZ-NASH model at endpoint






	Solvent control group of the model	Solvent control group of the model	WX001 (20nmol/kg)
Steatosis score	0	2.54	1.0
Inflammation score	0	1.67	1.63
Ballooning score	0	1.29	0
NAS score	0	5.5	2.63
Fibrosis (%)	0.71	1.11	1.21

[00367] Conclusion: The compound of the present disclosure can significantly improve the



NAS score in the STZ-NASH mouse model.

CLAIMS



1. A polypeptide represented by a sequence shown below,



wherein



¹Lys represents a modified lysine, and the modification is that the amino groups on the two





n is selected from 8, 9 and 10;


2. The polypeptide according to claim 1, wherein m is 2.

3. The polypeptide according to claim 1, wherein n is 9.


4. The polypeptide according to claim 1, wherein p is 1.

5. The polypeptide according to claim 1, wherein X₂ is


6. The polypeptide according to claim 1, wherein X₀ is




7. The polypeptide according to claim 1, wherein the structural unit $\begin{matrix} R \\ | \\ -CH-CH_2- \end{matrix}$ is



8. A polypeptide represented by the following formulas,





9. Use of the polypeptide compound according to any one of claims 1 to 8 in the manufacture of a medicament for the treatment of a GLP-1R/GCGR-related disease.
10. The use according to claim 9, wherein the GLP-1R/GCGR-related disease is obesity or nonalcoholic steatohepatitis.