

PATENT SPECIFICATION

(11) 1263 517

NO DRAWINGS

- (21) Application No. 43080/69 (22) Filed 29 Aug. 1969
 (31) Convention Application No. 756375 (32) Filed 30 Aug. 1968 in
 (33) United States of America (US)
 (45) Complete Specification published 9 Feb. 1972
 (51) International Classification C 07 c 65/04 69/84 C 11 b 9/00
 (52) Index at acceptance

C2C 3A7V2A2 3A7V2E2 3A7V2H 3A7V2K2 3A7V4A2
 3A7V4E2 3A7V4H 3A7V4K2

A5B 775

- (72) Inventors JAMES DOUGLAS GROSSMAN, ROBERT
 SANTORA DE SIMONE and LAMBERTUS
 GERKE HEERINGA



(54) SUBSTITUTED RESORCYLIC ACIDS AND ESTERS AND PROCESS FOR MAKING SAME


(71) We, INTERNATIONAL FLAVORS &
 FRAGRANCES INC., a Corporation organised
 and existing under the laws of the State of
 New York, United States of America, of 521
 West 57th Street, New York, New York,
 United States of America, do hereby declare
 the invention, for which we pray that a patent
 may be granted to us, and the method by
 which it is to be performed, to be particu-
 larly described in and by the following state-
 ment:—

Natural oakmoss is commercially important
 in producing high-grade fragrance composi-
 tions. Oakmoss constitutes an important and
 basic part of the fragrance impression of
 chypre and lavender. In view of the limited
 availability of natural oakmoss synthetic sub-
 stitutes are desirable and have been long
 sought.

It has been found that certain di - alkyl
 ring-substituted resorcylic acids and esters
 thereof simulate and resemble the fragrance
 impression of oakmoss. While a number of
 routes for the production of such resorcylic
 acids and esters thereof are available, they
 are rather tedious, complicated and generally
 uneconomic. For instance, one of these
 methods, as reported by A. Sonn in *Berichte*,
 62B, 3012-6 (1929), involves utilization of
 the rather expensive, and difficulty recoverable
 palladium catalyst for the aromaticization of
 mono- or dialkyl ring-substituted dihydro
 resorcyate. Another method shown by
 Robertson et al in *J. Chem. Soc.*, pp 313—
 20 (1930) involves a multi-step sequence
 difficult to perform and tedious to carry out.

The invention thus relates to di - alkyl ring-
 substituted resorcylic acids and esters there-
 of, their mode of preparation, and their use
 as fragrance materials.

The novel dialkyl ring-substituted resorcylic
 acids and esters thereof of this invention may
 be represented by the following formula:





wherein each of R_1 and R_2 is a lower alkyl
 radical containing from 1 to 5 carbon atoms
 and at least one of R_1 or R_2 contains two
 or more carbon atoms and wherein R_3 is
 hydrogen or a lower alkyl radical containing
 1 to 3 carbon atoms. Representative of the
 compounds included within said formula
 are:

methyl 3 - methyl - 6 - isopropyl - re -
 sorcyate
 methyl 3 - isopropyl - 6 - methyl - re -
 sorcyate
 methyl 3,6 - diethyl - resorcyate
 methyl 3 - methyl - 6 - n - butyl -
 resorcyate
 methyl 3,6 - di - n - propyl - resorcyate
 methyl 3,6 - di - isobutyl - resorcyate
 methyl 3,6 - di - n - butyl - resorcyate
 ethyl 3 - n - propyl - 6 - methyl - resorcy-
 ate
 3,6 - di - isopropyl resorcylic acid
 methyl 3 - methyl - 6 - ethyl - resorcyate
 methyl 3 - ethyl - 6 - methyl - resorcyate

The resorcylic acids and esters of this inven-
 tion effectively simulate and resemble oak-
 moss fragrance and are suitable for use by
 themselves as fragrance materials or in com-
 bination with other ingredients in a perfume
 composition such as a chypre or lavender
 perfume. The resorcylic esters are preferred
 as fragrance materials and the methyl esters
 are most suitable. Particularly preferred is
 methyl 3 - ethyl - 6 - methyl - resorcyate
 by reason of its pronounced and long lasting


mousse-like odor.

Also, in accordance with this invention it has been found that di-alkyl ring-substituted resorcylic acid and esters thereof, as represented by the formula




wherein R_3 is hydrogen or a lower alkyl radical containing from 1 to 5 carbon atoms and both R_1 and R_2 are such an alkyl radical, can be conveniently and more economically prepared by reacting a mono- or di-alkyl ring-substituted dihydroresorcylic acid with an oxidative chlorine material. This process has been found to result in high yield of the products desired and suppresses undesirable side reactions.

Suitable di-alkyl ring-substituted hydro-alkyl resorcylic acids and esters for this reaction may be represented by the following formulae:



wherein R_3 is hydrogen or a lower alkyl radical containing from 1 to 5 carbon atoms including methyl, ethyl, propyl, butyl, isopropyl and amyl, and both R_1 and R_2 are such a lower alkyl radical. Mixtures of such isomers can be used. The particular starting material selected will depend upon the end product desired and appropriate choices will be readily apparent. The dihydro-resorcylic acid or ester starting material is preferably substantially pure so as to avoid undesirable side reactions and facilitate isolation and recovery of the end product. Such dihydro-resorcylic acid or ester starting materials may be prepared by any of the known techniques. For example, such dihydro-resorcylic acids or esters can be readily made by reacting a beta-keto alkanolic acid or ester of the formula



with an alkyl alpha, beta-alkyl alkenoate of the formula

in the presence of an alkali metal alcoholate such as sodium methylate. In the above formulae R_1 , R_2 , R_3 have the meanings heretofore given and R_4 is a lower alkyl radical. A further description of the preparation of the intermediate dihydroresorcylic acids and esters may be found in A. Sonn, *Berichte*, 62B, 3012-6 (1929).

The oxidative chlorine material may be molecular chlorine or any material which, under the conditions of the reaction, yields or is a source of oxidative chlorine such as hypochlorous acid or nitrosyl chloride. Chlorine is preferred.

The yield obtained when practicing the process of this invention is affected by the degree of contact of the reactants. Thus, if the contact time of the oxidative chlorine material with the dihydro-resorcylic acid or ester is too low, the yields are lower and large amounts of unreacted dihydro-resorcylic acid or ester remains unconverted and must be separated from the reaction mixture. Intermixing of the reactants as by agitation or turbulent mixing will decrease the contact time required and increase the yield of desired product.

When hypochlorous acid is employed a two phase aqueous-non-aqueous reaction system results whereby the degree of contact between the two phases is an important if not controlling factor in determining the yield of product. The degree of contact is modified or regulated by nature of the agitation or intermixing.

It is generally desirable to carry out the reaction in an inert solvent for the dihydro-resorcylic acid or ester since at the reaction temperatures usually preferred, i.e. from about 20–30°C, the reactant is in solid form. Such a solvent medium, therefore, is useful to effect adequate contact between the reactants. Suitable inert reaction solvents include non-reactive halogenated hydrocarbons such as chloroform and tetrachloroethane.

The reaction conditions of temperature and pressure will depend upon a variety of factors including the time of the reaction, the particular reactants used, and the products desired. Generally, it is expedient to run the reaction at atmospheric pressure although elevated pressures may be employed. The reaction is conducted in the range of from about 15–50°C, and preferably in the range of from about 20–30°C, although variations from these temperatures are permissible. The time of reaction is generally determined by the temperature of the reaction and by the absolute pressure above the reaction mass.

Other factors such as the nature of the reactants, the extent and duration of contact of the reactants, the mixing system utilized, and the yield and conversions desired may also effect the reaction time. It is usually preferred to carry out the reaction in from about one to ten hours since shorter periods adversely affect yields while longer periods give no further advantage in the process.

The process is suitably carried out by dissolving or suspending the dihydro - resorcylic acid or ester in an inert organic solvent such as chloroform, whereby a two phase slurry of solid and saturated solution is formed. The oxidative chlorine material, as for example chlorine gas, is introduced into the reaction system, as by bubbling it through the reaction mass, while vigorously agitating the reaction system to achieve intimate intermixing. Introduction of the oxidative chlorine material is discontinued when chlorine is no longer taken up and the reaction mixture becomes clear. The organic solvent or vehicle is then removed from the reaction mass, the reactants separated and the reaction product may be recovered and isolated by any conventional technique such as distillation, selective extraction, or the like.

The dialkyl resorcylic acids and esters thereof obtained by practicing the processes, particularly the novel materials of this invention, are useful fragrance materials. In perfume compositions, i.e. a mixture of compounds including for example, natural oils, synthetic oils, aldehydes, ketones, esters, lactones, alcohols, and frequently hydrocarbons, the resorcylic acid or ester thereof contributes its particular olfactory characteristics. The overall effect of the perfume composition can be more than the effect of each ingredient. Thus, the individual resorcylic acid or ester may be used to alter the aroma characteristics of a perfume composition, for example, by high-lighting or moderating the olfactory contribution of another ingredient of the composition.

Perfume compositions containing the resorcylic acid or ester of this invention or the novel materials themselves can be used in conjunction with carriers, vehicles, solvents, dispersants, emulsifiers, surface-active agents, aerosol propellants, fixatives, and the like.

The amounts of the resorcylic acid or ester of this invention which will be effective in perfume compositions depends upon several factors, including the other ingredients of the composition, their amounts and the particular effects which are desired. It has been found that perfume compositions containing as little as 0.5% by weight of the compound of the invention, or even less, can be used to impart a basic oakmoss note to such materials as lavender, or chypre for use in soaps, cosmetics, lotions, handkerchief perfumes and similar products. The amount employed will

depend on considerations of cost, nature of the end product, the effect desired on the finished product and the particular fragrance sought.

The following examples serve to illustrate embodiments of the invention as it is now preferred to practice it. It will be understood that these examples are illustrative and the invention is to be considered restricted thereto only as indicated in the appended claims.

EXAMPLE I

Preparation of Methyl 3,6 Dimethyl - Resorcyate

Into a 5 liter reaction flask equipped with a stirrer, thermometer, reflux condenser, a gas bubbler inlet tube and outlet are placed 450 grams methyl 3,6 dimethyl - dihydro - resorcyate suspended in 2.5 liters of CHCl_3 . Chlorine gas at atmospheric pressure and room temperature is bubbled through the gas bubbler into the ingredients in the reaction vessel. The chlorine gas is added until no more chlorine gas is taken into solution and the solution becomes clear. The reaction vessel is stirred continuously during the bubbling of the chlorine gas. The stirring is continued after the cessation of the bubbling and until hydrogen chlorine evolution ceases. The chloroform solvent is stripped from the reaction mass yielding a solid product having a weight of 476.3 grams. This solid is mixed with diethyl ether and the resulting mixture is placed into a separatory funnel. The reaction mass is subjected to a bicarbonate extraction to recover starting material. The ether mixture is washed twice with 10% aqueous sodium hydroxide. The resulting aqueous phase is acidified with hydrochloric acid and cooled to 0°C . Additional hydrochloric acid is added until the pH of this phase is reduced to 2. The precipitate formed during the addition of acid is filtered and washed twice with water and dried. The dried product has a weight of 224.4 grams. Infrared, NMR and mass spectral analysis verifies that the product is methyl 3,6 - dimethyl - resorcyate. The product's melting range is 130.8°C — 136.6°C . The material has a mousse-like odor which can be imparted to soap at a level of 1/4%. The yield is 50.4% and the conversion is 75.0% taking into account unreacted reactant.

EXAMPLE II

Methyl 3 - Isopropyl - 6 - Methyl - Resorcyate

Into a 2 liter round bottom flask equipped with stirrer, thermometer, inlet gas bubbler and outlet, the following ingredients are placed:

211 grams of methyl 3 - isopropyl - 6 - methyl dihydro - resorcyate
1000 ml of chloroform.

Chlorine gas is bubbled through the reaction mixture at atmospheric pressure and room temperature until the chlorine no longer dissolves in the reaction mass. The reaction mass is concentrated in a flash evaporator to yield an oil that turns solid on standing. The solid material is dissolved in ether and the ether solution is washed with a saturated aqueous solution of sodium bicarbonate until the aqueous wash remains basic and evolution of CO₂ ceases. The ether solution is extracted with 10% sodium hydroxide at about 50°C and the extracts are filtered on vacuum to remove traces of ether. The ether-free caustic extract is neutralized with concentrated hydrochloric acid at 5°C to a pH of 2. A gummy mass forms and turns solid on standing at °C. The solid is dissolved in methanol, discolored with charcoal, and recrystallized from a methanol-water mixture. The solid is dried, yielding a product weighing 36.2 grams. This product is identified by NMR and mass spectral analysis as methyl 3 - isopropyl - 6 - methyl - resorcyate having a melting range of 125.6°C—135.8°C, and the following mass spectral data: 178, 192, 164, 224, and 226 (Parent peak).

EXAMPLE III

Methyl 3 - Ethyl - 6 - Methyl - Resorcyate
The process as described in Example I is carried out to obtain methyl 3 - ethyl - 6 - methyl resorcyate by substituting methyl 3 - ethyl - 6 - methyl - dihydroresorcyate for the methyl, 3,6 - dimethyl dihydroresorcyate used in Example I. The product has a fruity and caramel-like top note and its basic note is mousse-like. Its mass spectral data is: peaks at 163, 178, 210 (parent peak).

EXAMPLE IV

Methyl - 3 - n - Propyl - 6 - Methyl - Resorcyate

The process as described in Example I is carried out to obtain methyl 3 - n - propyl - 6 - methyl - resorcyate by substituting methyl 3 - n - propyl - 6 - methyl - dihydroresorcyate for the dihydro - resorcyate used in Example I. The product obtained has a melting range of 133.2°—134.6°C and a basic mousse olfactory note.

EXAMPLE V

Methyl 3 - Isobutyl - 6 - Methyl - Resorcyate
The process as described in Example I is carried out to obtain methyl 3 - isobutyl - 6 - methyl - resorcyate by substituting methyl 3 - isobutyl - 6 - methyl dihydro - resorcyate for the dihydro - resorcyate used in Example I. The product obtained has a basic mousse olfactory character and a melting range of 127.4°—128.6°C.

EXAMPLE VI

Methyl 3 - n - Butyl - 6 - Methyl - Resorcyate

The process as described in Example I is carried out to obtain methyl 3 - n - butyl - 6 - methyl - resorcyate by substituting methyl 3 - n - butyl - 6 - methyl - dihydro - resorcyate for the dihydro - resorcyate used in Example I. The product obtained has a mousse olfactory note.

EXAMPLE VII

Perfume Composition

The following mixture is prepared:

Ingredient	Grams	
Jasmin liquid, A	15	
Rose liquid	5	75
Solution orris	6	
Santal oil E.I.	6	
Bergamot	120	
Patchouli oil	6	
Musk ketone, 1/5 in B.B.	60	80
Vetivert oil	5	
Methyl 3 - ethyl - 6 - methyl - resorcyate.		
[Tincture 1/5]	200	
Coumarin	2	85
Vanillin	1.5	
Heliotropin	2	
Rose synthetic	25	
Rose otto, Bulgarian	10	
Pimento oil	5	90
Olibanum resinoid	10	
Bitter orange oil	4	
Ambrette seed oil	2	
Musk tincture, 3 per cent	250	
Alcohol	4000	95
	4734	

The foregoing perfume formulation is an important part of chypre essence. The methyl 3 - ethyl - 6 - methyl - resorcyate is used as a replacement for oakmoss. This perfume is incorporated into a handkerchief perfume at the 0.1% by weight level. The methyl 3 - ethyl - 6 - methyl - resorcyate gives to this fragrance a natural and distinctly oakmoss note.

EXAMPLE VIII

The following mixture is prepared:

Ingredient	Grams	
Benzyl acetate	50	
Rosewood	100	110
Cedarwood	150	
Linalyl acetate	100	
α - Ionone	80	
Ethyl cinnamate	20	
Amyl cinnamic aldehyde	50	115
Iso - eugenol	50	
Coumarin	100	
Methyl 3 - isopropyl - 6 - methyl resorcyate	50	
Musk xylene	50	120
Styrax resin	100	

The foregoing perfume formulation is an important part of chypre essence. The methyl 3 - isopropyl - 6 - methyl - resorcyate is used as a replacement for oakmoss. This perfume is incorporated into a soap perfume at the 0.1% level. The methyl 3 - isopropyl - 6 - methyl - resorcyate gives to this fragrance a natural and distinctly oakmoss note.

EXAMPLE IX

Perfume Composition


The following mixture is prepared:

	Ingredient	Grams
	Benzyl acetate	50
	Rosewood	100
15	Cedarwood	150
	Linalyl acetate	100
	α - Ionone	80
	Ethyl cinnamate	20
	Amyl cinnamic aldehyde	50
20	Iso - eugenol	50
	Coumarin	100
	Methyl 3 - n - butyl - 6 - methyl - resorcyate	200
	Musk xylene	50
25	Styrax resin	100


The foregoing perfume formulation is an important part of chypre essence. The methyl 3 - n - butyl - 6 - methyl - resorcyate is used as a replacement for oakmoss. This perfume is incorporated into a soap perfume at the 0.1% level. The methyl 3 - n - butyl - 6 - methyl - resorcyate gives to this fragrance a natural and distinctly oakmoss note.

WHAT WE CLAIM IS:—

1. A process for preparing an alkyl ring-substituted resorcylic acid or ester thereof having the formula



which comprises reacting the corresponding dihydro - resorcylic compound of the formulae



or mixtures thereof with an oxidative chlorine material, and wherein R_3 in the above formulae is hydrogen or a lower alkyl radical containing from 1 to 5 carbon atoms and both R_1 and R_2 are such an alkyl radical.

2. The process of claim 1 wherein the oxidative chlorine material is selected from the group consisting of chlorine, hypochlorous acid, and nitrosyl chloride.

3. The process of claim 1 wherein the oxidative chlorine source is molecular chlorine.


4. The process of claim 1 wherein each of R_1 , R_2 and R_3 is methyl.

5. The process of claim 4 wherein the oxidative chlorine source is molecular chlorine.

6. The process of claim 1 wherein R_1 is isopropyl and each of R_2 and R_3 is methyl.

7. The process of claim 1 wherein R_1 is ethyl and each of R_2 and R_3 is methyl.

8. A dialkyl resorcylic acid or ester thereof having the formula



wherein each of R_1 and R_2 is a lower alkyl radical containing from 1 to 5 carbon atoms and at least one of R_1 or R_2 contains two or more carbon atoms and wherein R_3 is hydrogen or a lower alkyl radical containing from 1 to 3 carbon atoms.

9. A compound as defined in claim 8 wherein R_1 is isopropyl and each of R_2 and R_3 is methyl.

10. A compound as defined in claim 8 wherein R_1 is ethyl and each of R_2 and R_3 is methyl.

11. A process according to Claim 1, for preparing an alkyl ring-substituted resorcylic acid ester substantially as hereinbefore described in any of Examples I to VI.

12. A compound when prepared by the process of any one of Claims 1 to 7 and 11.

13. A perfume composition comprising a compound according to any of Claims 8 to 10 and 12.

14. A perfume composition substantially as hereinbefore described in any of Examples VII to IX.

For the Applicants,
CARPMAELS & RANSFORD,
Chartered Patent Agents,
24 Southampton Buildings,
Chancery Lane,
London, W.C.2.

The foregoing perfume formulation is an important part of chypre essence. The methyl 3 - isopropyl - 6 - methyl - resorcyate is used as a replacement for oakmoss. This perfume is incorporated into a soap perfume at the 0.1% level. The methyl 3 - isopropyl - 6 - methyl - resorcyate gives to this fragrance a natural and distinctly oakmoss note.

EXAMPLE IX

Perfume Composition


The following mixture is prepared:

	Ingredient	Grams
	Benzyl acetate	50
	Rosewood	100
15	Cedarwood	150
	Linalyl acetate	100
	α - Ionone	80
	Ethyl cinnamate	20
	Amyl cinnamic aldehyde	50
20	Iso - eugenol	50
	Coumarin	100
	Methyl 3 - n - butyl - 6 - methyl - resorcyate	200
	Musk xylene	50
25	Styrax resin	100


The foregoing perfume formulation is an important part of chypre essence. The methyl 3 - n - butyl - 6 - methyl - resorcyate is used as a replacement for oakmoss. This perfume is incorporated into a soap perfume at the 0.1% level. The methyl 3 - n - butyl - 6 - methyl - resorcyate gives to this fragrance a natural and distinctly oakmoss note.

WHAT WE CLAIM IS:—

1. A process for preparing an alkyl ring-substituted resorcylic acid or ester thereof having the formula



which comprises reacting the corresponding dihydro - resorcylic compound of the formulae



or mixtures thereof with an oxidative chlorine material, and wherein R_3 in the above formulae is hydrogen or a lower alkyl radical containing from 1 to 5 carbon atoms and both R_1 and R_2 are such an alkyl radical.

2. The process of claim 1 wherein the oxidative chlorine material is selected from the group consisting of chlorine, hypochlorous acid, and nitrosyl chloride.

3. The process of claim 1 wherein the oxidative chlorine source is molecular chlorine.


4. The process of claim 1 wherein each of R_1 , R_2 and R_3 is methyl.

5. The process of claim 4 wherein the oxidative chlorine source is molecular chlorine.

6. The process of claim 1 wherein R_1 is isopropyl and each of R_2 and R_3 is methyl.

7. The process of claim 1 wherein R_1 is ethyl and each of R_2 and R_3 is methyl.

8. A dialkyl resorcylic acid or ester thereof having the formula



wherein each of R_1 and R_2 is a lower alkyl radical containing from 1 to 5 carbon atoms and at least one of R_1 or R_2 contains two or more carbon atoms and wherein R_3 is hydrogen or a lower alkyl radical containing from 1 to 3 carbon atoms.

9. A compound as defined in claim 8 wherein R_1 is isopropyl and each of R_2 and R_3 is methyl.

10. A compound as defined in claim 8 wherein R_1 is ethyl and each of R_2 and R_3 is methyl.

11. A process according to Claim 1, for preparing an alkyl ring-substituted resorcylic acid ester substantially as hereinbefore described in any of Examples I to VI.

12. A compound when prepared by the process of any one of Claims 1 to 7 and 11.

13. A perfume composition comprising a compound according to any of Claims 8 to 10 and 12.

14. A perfume composition substantially as hereinbefore described in any of Examples VII to IX.

For the Applicants,
CARPMAELS & RANSFORD,
Chartered Patent Agents,
24 Southampton Buildings,
Chancery Lane,
London, W.C.2.

