

ALKYNYL PYRROLOPYRIMIDINES AND RELATED ANALOGS AS HSP90-INHIBITORS

CROSS-REFERENCE

[001] This application claims the benefit of U.S. Provisional Application No. 60/666,899 filed March 30, 2005, which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

[002] The invention relates in general to alkynyl pyrrolopyrimidines and their broad-spectrum utility, *e.g.*, in inhibiting heat shock protein 90 (HSP90) to thereby treat or prevent HSP90-mediated diseases.

BACKGROUND

[003] HSP90s are ubiquitous chaperone proteins that are involved in folding, activation and assembly of a wide range of proteins, including key proteins involved in signal transduction, cell cycle control and transcriptional regulation. Researchers have reported that HSP90 chaperone proteins are associated with important signaling proteins, such as steroid hormone receptors and protein kinases, including, *e.g.*, Raf-1, EGFR, v-Src family kinases, Cdk4, and ErbB-2 (Buchner *J. TIBS* **1999**, *24*, 136-141; Stepanova, L. *et al. Genes Dev.* **1996**, *10*, 1491-502; Dai, K. *et al. J. Biol. Chem.* **1996**, *271*, 22030-4). Studies further indicate that certain co-chaperones, *e.g.*, HSP70, p60/Hop/Sti1, Hip, Bag1, HSP40/Hdj2/Hsj1, immunophilins, p23, and p50, may assist HSP90 in its function (see, *e.g.*, Caplan, A. *Trends in Cell Biol.* **1999**, *9*, 262-68).

[004] HSP90 possesses a binding pocket at its N-terminus. This pocket is highly conserved and has weak homology to the ATP-binding site of DNA gyrase (Stebbins, C. *et al., supra*; Grenert, J.P. *et al. J. Biol. Chem.* **1997**, *272*, 23843-50). Further, ATP and ADP have both been shown to bind this pocket with low affinity and to have weak ATPase activity (Proromou, C. *et al. Cell* **1997**, *90*, 65-75; Panaretou, B. *et al. EMBO J.* **1998**, *17*, 4829-36). *In vitro* and *in vivo* studies have demonstrated that occupancy of this N-terminal pocket by ansamycins and other HSP90 inhibitors alters HSP90 function and inhibits protein folding. At high concentrations, ansamycins and other HSP90 inhibitors have been shown to prevent binding of protein substrates to HSP90 (Scheibel, T.H. *et al. Proc. Natl. Acad. Sci. USA* **1999**, *96*, 1297-302; Schulte, T. W. *et al. J. Biol. Chem.* **1995**, *270*, 24585-8; Whitesell, L., *et al. Proc. Natl. Acad. Sci. USA* **1994**, *91*, 8324-8328). HSP90 inhibitors, *e.g.* ansamycins, have also been demonstrated to inhibit the ATP-dependent release of chaperone-associated protein substrates (Schneider, C.L. *et al. Proc. Natl. Acad. Sci., USA* **1996**, *93*, 14536-41; Sepp-Lorenzino *et al. J. Biol. Chem.* **1995**, *270*,

16580-16587). In either event, the substrates are degraded by an ubiquitin-dependent process in the proteasome (Schneider, C. L., *supra*; Sepp-Lorenzino, L., *et al. J. Biol. Chem.* **1995**, 270, 16580-16587; Whitesell, L. *et al. Proc. Natl. Acad. Sci. USA* **1994**, 91, 8324-8328).

[005] HSP90 substrate destabilization occurs in tumor and non-transformed cells alike and has been shown to be especially effective on a subset of signaling regulators, *e.g.*, Raf (Schulte, T. W. *et al. Biochem. Biophys. Res. Commun.* **1997**, 239, 655-9; Schulte, T. W., *et al. J. Biol. Chem.* **1995**, 270, 24585-8), nuclear steroid receptors (Segnitz, B.; U. Gehring *J. Biol. Chem.* **1997**, 272, 18694-18701; Smith, D. F. *et al. Mol. Cell. Biol.* **1995**, 15, 6804-12), v-Src (Whitesell, L., *et al. Proc. Natl. Acad. Sci. USA* **1994**, 91, 8324-8328) and certain

transmembrane tyrosine kinases (Sepp-Lorenzino, L. *et al. J. Biol. Chem.* **1995**, 270, 16580-16587) such as EGF receptor (EGFR) and Her2/Neu (Hartmann, F., *et al. Int. J. Cancer* **1997**, 70, 221-9; Miller, P. *et al. Cancer Res.* **1994**, 54, 2724-2730; Mimnaugh, E. G., *et al. J. Biol. Chem.* **1996**, 271, 22796-801; Schnur, R. *et al. J. Med. Chem.* **1995**, 38, 3806-3812), CDK4, and mutant p53. Erlichman *et al. Proc. AACR* **2001**, 42, abstract 4474. The HSP90 inhibitor-induced loss of these proteins leads to the selective disruption of certain regulatory pathways and results in growth arrest at specific phases of the cell cycle (Muisse-Heimericks, R. C. *et al. J. Biol. Chem.* **1998**, 273, 29864-72), and apoptosis, and/or differentiation of cells so treated (Vasilevskaya, A. *et al. Cancer Res.*, **1999**, 59, 3935-40). HSP90 inhibitors thus hold great promise for the treatment and/or prevention of many types of cancers and proliferative disorders, and also hold promise as traditional antibiotics.

[006] In addition to anti-cancer and antitumorigenic activity, HSP90 inhibitors have also been implicated in a wide variety of other utilities, including use as anti-inflammation agents, anti-infectious disease agents, agents for treating autoimmunity, agents for treating stroke, ischemia, multiple sclerosis, cardiac disorders, central nervous system related disorders and agents useful in promoting nerve regeneration (See, *e.g.*, Rosen *et al.* WO 02/09696 (PCT/US01/23640); Degranco *et al.* WO 99/51223 (PCT/US99/07242); Gold, U.S. Patent 6,210,974 B1; DeFranco *et al.*, US Patent 6,174,875. Overlapping somewhat with the above, there are reports in the literature that fibrogenetic disorders including but not limited to scleroderma, polymyositis, systemic lupus, rheumatoid arthritis, liver cirrhosis, keloid formation, interstitial nephritis, and pulmonary fibrosis also may be treatable with HSP90 inhibitors. Strehlow, WO 02/02123 (PCT/US01/20578). Still further HSP90 modulation, modulators and uses thereof are reported in Application Nos. PCT/US03/04283, PCT/US02/35938, PCT/US02/16287, PCT/US02/06518, PCT/US98/09805, PCT/US00/09512, PCT/US01/09512, PCT/US01/23640, PCT/US01/46303, PCT/US01/46304, PCT/US02/06518, PCT/US02/29715, PCT/US02/35069, PCT/US02/35938,


PCT/US02/39993, 60/293,246, 60/371,668, 60/335,391, 60/128,593, 60/337,919, 60/340,762, 60/359,484 and 60/331,893.

[007] Recently, purine derivatives, including pyrrolopyrimidines showing HSP90 inhibitory activity have been reported, *e.g.*, in PCT/US02/35069; PCT/US02/36075. U.S. Patent Application No. 10/945,851 and PCT/US04/31248. However, a need remains for additional novel and potent pyrrolopyrimidine HSP90 inhibitors that meet the demanding biological and pharmaceutical criteria required to proceed towards human clinical trials.

SUMMARY OF THE INVENTION

[008] The present invention is directed towards alkynyl pyrrolo[2,3-d]pyrimidines and related compounds that show broad utility, *e.g.*, by inhibiting HSP90 and treating diseases that are HSP90-dependent. These compounds differ from a parent pyrrolopyrimidine which was disclosed in prior patent applications in that they are substituted with an alkyne, *e.g.*, acetylene, on the ring carbon at the fifth position (C-5 position) and they exhibit improved HSP90 inhibitory activity over the parent compounds.

[009] In one aspect, the invention comprises alkynyl pyrrolo[2,3-d]pyrimidine compounds of Formula I:



wherein:

R⁰ is selected from the group consisting of hydrogen, halogen, lower alkyl, -CN, -SR⁸, -OR⁸, and -NHR⁸;

R¹ is selected from the group consisting of halogen, -OR¹¹, -SR¹¹ and lower alkyl;

R² is -NHR⁸;

R³ is selected from the group consisting of hydrogen, -CN, -C(O)OH, -OR¹¹, -SR¹¹, -C(O)R⁹, -NR⁸R¹⁰, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower alkylsilyl, aryl, heteroaryl, alicyclyl and heterocyclyl, all optionally substituted, wherein:

the aryl, heteroaryl, alicyclyl and heterocyclyl groups are mono-, bi- or tri- cyclic;

R⁸ and R¹⁰ taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

~~PCT/US2006/011846~~

the optional substituents on R^3 are selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R⁸, lower aryl, heteroaryl, alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, amino, alkylamino, dialkylamino, arylalkylamino, diarylamino, heteroarylamino, diheteroarylamino, arylheteroarylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidiny, pyridiny, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl, phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R^8 and R^8 taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R^4 is selected from the group consisting of optionally substituted lower alkylene, -C(R¹²)₂-, -C(O)-, -C(S)-, -S(O)- and -SO₂-;

R^5 is selected from the group consisting of aryl, heteroaryl, alicyclyl and heterocyclyl, wherein:

the aryl group is substituted with 2 to 5 substituents;

the heteroaryl group is substituted with 2 to 5 substituents;

the alicyclyl group is substituted with 3 to 5 substituents;

the heterocyclyl group is substituted with 3 to 5 substituents;

the substituents on R^5 are selected from the group consisting of halogen, lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R¹⁰, lower aryl, lower heteroaryl, lower alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, thioalkyl, amino, alkylamino, dialkylamino, arylalkylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidiny, pyridiny, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl, phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R^8 and R^{10} taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R^8 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower heteroaryl and -C(O)R⁹;

R^9 is selected from the group consisting of H, lower alkyl, lower alkenyl, lower alkynyl, lower aryl, lower heteroaryl, -NR¹⁰R¹⁰ and -OR¹¹, wherein

~~PCT/US2006/011846~~
R⁹ and R¹⁰ taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R¹⁰ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower heteroaryl and -C(O)R¹¹;

R¹¹ is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower aryl and lower heteroaryl; and

R¹² is selected from the group consisting of hydrogen and lower alkyl.

[010] Also included in the scope of the present invention are stereoisomeric forms, including the individual enantiomers and diastereomers, racemic mixtures, and diastereomeric mixtures, and combinations thereof, where appropriate, as well as polymorphs, specific racemates and stereoisomers, solvates, esters, tautomers, pharmaceutically acceptable salts and prodrugs of these compounds.

[011] In another aspect, the invention features pharmaceutical compositions comprising the compounds of the invention, in particular, the compounds of Formula I, or a polymorph, solvate, ester, tautomer, diastereoisomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, and one or more pharmaceutical excipients, for use in treatment or prevention of diseases and conditions that are HSP90-dependent.

[012] In yet another aspect, the invention is related to methods of preventing or treating HSP90-mediated disorders or conditions by administering a pharmaceutical composition that comprises a pharmaceutically effective amount of a compound of Formula I, or a polymorph, solvate, ester, tautomer, diastereomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof.

[013] In one embodiment, the invention provides a method for treating an individual having a disorder selected from the group of inflammatory diseases, infections, autoimmune disorders, stroke, ischemia, cardiac disorders, neurological disorders, fibrogenetic disorders, proliferative disorders, tumors, leukemias, chronic lymphocytic leukemia, acquired immuno-deficiency syndrome, neoplasms, cancers, carcinomas, metabolic diseases, and malignant diseases.

[014] In another embodiment, the invention provides a method for treating an individual having a fibrogenetic disorder, such as, for example, scleroderma, polymyositis, systemic lupus, rheumatoid arthritis, liver cirrhosis, keloid formation, interstitial nephritis or pulmonary fibrosis.

[015] In another embodiment, the invention provides a combination therapy comprising the administration of a pharmaceutically effective amount of a compound of Formula I, or a solvate, tautomer, diastereomer, enantiomer, pharmaceutically acceptable salt, polymorph, or prodrug

thereof according to any of the preceding aspects or embodiments, and at least one therapeutic agent selected from the group of cytotoxic agents, anti-angiogenesis agents and anti-neoplastic agents. The anti-neoplastic agent may be selected from the group of alkylating agents, anti-metabolites, epidophyllotoxins, antineoplastic enzymes, topoisomerase inhibitors, procarbazines, mitoxantrones, platinum coordination complexes, biological response modifiers and growth inhibitors, hormonal/anti-hormonal therapeutic agents, and haematopoietic growth factors.

[016] In a further aspect, the invention is related to the use of the compounds of Formula I in the manufacture of a medicament.

[017] In yet a further aspect, the invention is related to the use of the compounds of Formula I in the manufacture of a medicament for the therapeutical and/or prophylactic treatment of HSP90-dependent diseases and conditions.

~~PCT/US06/11846~~
INCORPORATION BY REFERENCE

[018] All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[019] The novel features of the invention are set forth with particularity in the appended claims.

A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[020] FIGURE 1 represents a plot of tumor volume (mm^3) against time (days), for animals administered compounds of the present invention (and controls) in a mouse N87 Gastric Carcinoma Xenograft model, as described in example 70.

[021] FIGURE 2 represents a plot of tumor volume (mm^3) against time (days), for animals administered compounds of the present invention (and controls) in a mouse NCI295 Adrenocortical Carcinoma Xenograft model, as described in example 71.

[022] FIGURE 3 represents a plot of tumor volume (mm^3) against time (days), for animals administered compounds of the present invention (and controls) in a mouse SK-MEL-28 Melanoma Xenograft model, as described in example 72.

[023] FIGURE 4 represents a plot of tumor volume (mm^3) against time (days), for animals administered compounds of the present invention (and controls) in a mouse a plot of tumor volume (mm^3) against time (days), for animals administered compounds of the present invention (and controls) in a mouse HT29 Colon Carcinoma Xenograft model, as described in example 73.

[024] FIGURE 5 represents Western Blot protein analysis of N87 gastric carcinoma cells after treatment with compounds 2 and 23 at varying concentrations and timepoints, as described in example 74.

DETAILED DESCRIPTION OF THE INVENTION

[025] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

PCT/US06/11846

[026] Definitions

A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester or other derivative of a compound of this invention, which, upon
5 administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or a pharmaceutically active metabolite or residue thereof. Particularly favored derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a patient (*e.g.*, by allowing orally administered compound to be more readily absorbed into blood) or which enhance delivery of
10 the parent compound to a biological compartment (*e.g.*, the brain or lymphatic system).

A "pharmaceutically acceptable salt" may be prepared for any compound of the invention having functionality capable of forming a salt, for example, an acid or base functionality.

Pharmaceutically acceptable salts may be derived from organic or inorganic acids and bases.

Compounds of the invention that contain one or more basic functional groups, (*e.g.*, amino,
15 alkylamino), are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable organic and inorganic acids. These salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed. Examples of suitable acid salts include acetate, adipate, alginate,
20 aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate,
25 pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts. See, *e.g.*, Berge *et al.* "Pharmaceutical Salts", *J.*
30 *Pharm. Sci.* **1977**, 66:1-19.

Compounds of the present invention that contain one or more acidic functional groups are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable bases. The term "pharmaceutically acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can
35 likewise be prepared *in situ* during the final isolation and purification of the compounds, or by

~~PCT/US2006/011846~~
separately reacting the purified compound in its free acid form with a suitable base, such as the

hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation, with ammonia, or with a pharmaceutically acceptable organic primary, secondary or tertiary amine.

Representative pharmaceutically acceptable cations include alkali or alkaline earth salts such as

5 lithium, sodium, potassium, calcium, magnesium, and aluminum salts and the like. Illustrative examples of some of the bases that can be used include sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, $N^+(C_{1-4} \text{ alkyl})_4$, and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. This invention also

10 envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

See, for example, Berge *et al.*, *supra*.

It should be understood that a reference to a salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates contain either stoichiometric or non-
15 stoichiometric amounts of a solvent, and are often formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol.


Polymorphs include the different crystal packing arrangements of the same elemental composition of a compound. Polymorphs usually have different X-ray diffraction patterns,
20 infrared spectra, melting points, density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature may cause a single crystal form to dominate.

Pharmaceutically acceptable prodrugs of the compounds of this invention include, but are not limited to, esters, carbonates, thiocarbonates, N-acyl derivatives, N-acyloxyalkyl derivatives,
25 quaternary derivatives of tertiary amines, N-Mannich bases, Schiff bases, aminoacid conjugates, phosphate esters, metal salts and sulfonate esters.

Suitable positions for derivatization of the compounds of the invention to create "prodrugs" include but are not limited, to, 2-amino substitution. Those of ordinary skill in the art have the knowledge and means to accomplish this without undue experimentation. Various forms of
30 prodrugs are well known in the art. See for example *Design of Prodrugs*, Bundgaard, A. Ed., Elsevier, 1985 and *Method in Enzymology*, Widder, K. *et al.*, Ed.; Academic, 1985, vol. 42, p. 309-396; Bundgaard, H. "Design and Application of Prodrugs" in *A Textbook of Drug Design and Development*, Krosgaard-Larsen and H. Bundgaard, Ed., 1991, Chapter 5, p. 113-191; and Bundgaard, H., *Advanced Drug Delivery Review*, 1992, 8, 1-38, each of which is incorporated
35 herein by reference.

The term "prodrugs" as employed herein includes, but is not limited to, the following groups and combinations of these groups, for example bis carbomates, or a carbomate and an acyloxyalkyl ester or a carbomate and an amide, etc.

[027] Amine prodrugs:



[028] Hydroxy prodrugs include, but are not limited to acyloxyalkyl esters, alkoxycarbonyloxyalkyl esters, alkyl esters, aryl esters and disulfide containing esters.

[029] The term "alkyl" as used herein, alone or in combination, refers to an optionally substituted straight-chain, or optionally substituted branched-chain saturated hydrocarbon monoradical having from one to about thirty carbons, more preferably one to twelve carbons. Examples of alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, tert-amyl, pentyl, hexyl, heptyl, octyl and the like.

The term "cycloalkyl" as used herein, alone or in combination, refers to cyclic alkyl

monoradicals which include monocyclic, bicyclic, tricyclic, and higher multicyclic alkyl radicals wherein each cyclic moiety has from three to about eight carbon atoms. Examples of cycloalkyl radicals include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and the like.

The term "lower alkyl" as used herein, alone or in combination, refers to an alkyl containing fewer carbon atoms, *e.g.*, one containing from one to about six carbon atoms.

The term "alkenyl" as used herein, alone or in combination, refers to an optionally substituted straight-chain, or optionally substituted branched-chain hydrocarbon radical having one or more carbon-carbon double-bonds and having from two to about thirty carbon atoms, more preferably two to about eighteen carbons. Examples of alkenyl radicals include ethenyl, propenyl, butenyl, 1,3-butadienyl and the like.

The term "cycloalkenyl" as used herein, alone or in combination, refers to cyclic alkenyl radicals which include monocyclic, bicyclic, tricyclic, and higher multicyclic alkenyl radicals wherein each cyclic moiety has from three to about eight carbon atoms.

~~For purposes of this application~~
The term "lower alkenyl" as used herein, alone or in combination, refers to an alkenyl having from two to about six carbons.

The term "alkynyl" as used herein, alone or in combination, refers to an optionally substituted straight-chain or optionally substituted branched-chain hydrocarbon radical having one or more carbon-carbon triple-bonds and having from two to about thirty carbon atoms, more preferably from two to about twelve carbon atoms, or from two to about six carbon atoms, as well as those having from two to about four carbon atoms. Examples of alkynyl radicals include ethynyl, 2-propynyl, 2-butynyl, 1,3-butadiynyl and the like.

The term "cycloalkynyl" as used herein, alone or in combination, refers to cyclic alkynyl radicals that include monocyclic, bicyclic, tricyclic, and higher multicyclic alkynyl radicals wherein each cyclic moiety has from three to about eight carbon atoms.

The term "lower alkynyl" as used herein, alone or in combination, refers to an alkynyl having from two to about six carbons.

The terms "heteroalkyl", "heteroalkenyl" and "heteroalkynyl" as used herein, alone or in combination, refer to include optionally substituted alkyl, alkenyl and alkynyl structures, as described above, and which have one or more skeletal chain atoms selected from an atom other than carbon, *e.g.*, oxygen, nitrogen, sulfur, phosphorous or combinations thereof.

The terms "lower heteroalkyl", "lower heteroalkenyl" and "lower heteroalkynyl" as used herein, alone or in combination, refer to, respectively, a heteroalkyl, heteroalkenyl and heteroalkynyl having from two to about six carbons.

The term "alkylene" as used herein, alone or in combination, refers to a divalent hydrocarbyl group; because it is divalent, it can link two other groups together. Typically it refers to $-(CH_2)_n-$ where n is 1-8 and preferably n is 1-4, though where specified, an alkylene can also be substituted by other groups, and can be of other lengths, and the open valences need not be at opposite ends of a chain. Thus $-CH(Me)-$ and $-C(Me)_2-$ may also be referred to as alkylenes, as can a cyclic group such as cyclopropan-1,1-diyl. Where an alkylene group is substituted, the substituents include those typically present on alkyl groups as described herein.

The term "lower alkylene" as used herein, alone or in combination, refers to an alkylene group containing fewer carbon atoms, *e.g.*, one containing from one to about six carbon atoms.

The term "carbon chain" as used herein, alone or in combination, refers to any alkyl, alkenyl, alkynyl, or heteroalkyl, heteroalkenyl, or heteroalkynyl group, which is linear, cyclic, or any combination thereof. If the chain is part of a linker and that linker comprises one or more rings as part of the core backbone, for purposes of calculating chain length, the "chain" only includes those carbon atoms that compose the bottom or top of a given ring and not both, and where the top and bottom of the ring(s) are not equivalent in length, the shorter distance shall be used in

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35
determining the chain length. If the chain contains heteroatoms as part of the backbone, those atoms are not calculated as part of the carbon chain length.

The term "membered ring" as used herein, alone or in combination, refers to any cyclic structure, including aromatic, heteroaromatic, alicyclic, heterocyclic, monocyclic, polycyclic, and fused rings. The term "membered" is meant to denote the number of skeletal (or ring) atoms that constitute the ring system. Thus, for example, pyrrole, pyrrolidine, succinimide, maleimide, tetrahydrofuran and thiophene are five-membered rings; pyridine, pyran, morpholine, piperazine, piperidine and pyrimidine are six-membered rings; and phthalimide, indole and indane are nine membered fused rings.

The term "aryl" as used herein, alone or in combination, refers to an optionally substituted aromatic hydrocarbon monoradical of six to about twenty ring atoms, and includes mono-aromatic rings and fused aromatic rings. A fused aromatic ring radical contains from two to four fused rings where the ring of attachment is an aromatic ring, and the other individual rings within the fused ring may be aromatic, heteroaromatic, alicyclic or heterocyclic. Further, the term aryl includes mono-aromatic rings and fused aromatic rings containing from six to about twelve carbon atoms, as well as those containing from six to about ten carbon atoms. Examples of aryl groups include, without limitation, phenyl, naphthyl, anthryl, chrysenyl, and benzopyrenyl ring systems.

The term "lower aryl" as used herein, alone or in combination, refers to an aryl having six to about ten skeletal ring carbons, *e.g.*, phenyl and naphthyl ring systems.

The term "heteroaryl" as used herein, alone or in combination, refers to optionally substituted aromatic radicals containing from about five to about twenty skeletal ring atoms and where one or more of the ring atoms is a heteroatom such as, for example, oxygen, nitrogen, sulfur, selenium or phosphorus. The term heteroaryl includes optionally substituted mono-heteroaryl radicals and fused heteroaryl radicals having at least one heteroatom (*e.g.*, quinoline, benzothiazole). A fused heteroaryl radical may contain from two to four fused rings where the ring of attachment is a heteroaromatic ring, the other individual rings within the fused ring system may be aromatic, heteroaromatic, alicyclic or heterocyclic. The term heteroaryl also includes mono-heteroaryls or fused heteroaryls having from five to about twelve skeletal ring atoms, as well as those having from five to about ten skeletal ring atoms. Examples of heteroaryls include, without limitation, furanyl, benzofuranyl, chromenyl, pyridyl, pyrrolyl, indolyl, quinolinyl, pyridyl-N-oxide, pyrimidyl, pyrazinyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, benzothiozolyl, benzimidazolyl, benzoxazolyl, benzothiadiazolyl, benzoxadiazolyl, benzotriazolyl, quinolinyl, isoquinolinyl, indolyl, purinyl, indoliziny, thienyl and the like and their oxides.

~~PCT/US2006/011846~~
The term "lower heteroaryl" as used herein, alone or in combination, refers to a heteroaryl having

five to about ten skeletal ring atoms, *e.g.*, pyridyl, thienyl, pyrimidyl, pyrazinyl, pyrrolyl, or furanyl.

The terms "alicyclic" and "alicycyl" as used herein, alone or in combination, refer to an

5 optionally substituted saturated or unsaturated nonaromatic hydrocarbon ring system containing from three to about twenty ring atoms. The term alicyclic includes mono-alicyclic and fused alicyclic radicals. A fused alicyclic may contain from two to four fused rings where the ring of attachment is an alicyclic ring, and the other individual rings within the fused-alicyclic radical may be aromatic, heteroaromatic, alicyclic and heterocyclic. The term alicyclic also includes
10 mono-alicyclic and fused alicyclic radicals containing from three to about twelve carbon atoms, as well as those containing from three to about ten carbon atoms. Examples of alicyclics include, without limitation, cyclopropyl, cyclopropenyl, cyclobutyl, cyclopentyl, cyclodecyl, cyclododecyl, cyclopentadienyl, indanyl, and cyclooctatetraenyl ring systems.

The terms "lower alicyclic" and "lower alicycyl" as used herein, alone or in combination, refer
15 to an alicyclic having three to about ten skeletal ring carbons, *e.g.*, cyclopropyl, cyclopropenyl, cyclobutyl, cyclopentyl, decalinyl, and cyclohexyl.

The terms "heterocyclic" and "heterocycyl" as used herein, alone or in combination, refer to optionally substituted saturated or unsaturated nonaromatic ring radicals containing from five to about twenty ring atoms where one or more of the ring atoms are heteroatoms such as, for
20 example, oxygen, nitrogen, sulfur, and phosphorus. The term heterocyclic includes mono-heterocyclic and fused heterocyclic ring radicals. A fused heterocyclic radical may contain from two to four fused rings where the attaching ring is a heterocyclic, and the other individual rings within the fused heterocyclic radical may be aromatic, heteroaromatic, alicyclic or heterocyclic. The term heterocyclic also includes mono-heterocyclic and fused alicyclic radicals having from
25 five to about twelve skeletal ring atoms, as well as those having from five to about ten skeletal ring atoms. Example of heterocyclics include without limitation, tetrahydrofuranyl, benzodiazepinyl, tetrahydroindazolyl, dihydroquinolinyl, and the like.

The terms "lower heterocyclic" and "lower heterocycyl" as used herein, alone or in combination, refer to a heterocyclic ring system having five to about ten skeletal ring atoms, *e.g.*,
30 dihydropyranyl, pyrrolidinyl, dioxolanyl, piperidinyl, piperazinyl, and the like.

The term "alkylaryl" as used herein, alone or in combination, refers to an aryl radical as defined above in which at least one H atom is replaced by an alkyl radical as defined above, such as, for example, tolyl, xylyl and the like.

The terms "arylalkyl" and "araalkyl" as used herein, alone or in combination, refer to an alkyl radical as defined above in which at least one H atom is replaced by an aryl radical as defined above, such as, for example, benzyl, 2-phenylethyl and the like.

The term "heteroarylalkyl" as used herein, alone or in combination, refers to an alkyl radical as defined above in which at least one H atom is replaced by a heteroaryl radical as defined above, each of which may be optionally substituted.

The term "alkoxy" as used herein, alone or in combination, refers to an alkyl ether radical, alkyl-O-, wherein the term alkyl is defined as above. Examples of alkoxy radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, tert-butoxy and the like.

The term "lower alkoxy" as used herein, alone or in combination, refers to an alkoxy group having one to about six carbon atoms.

The term "aryloxy" as used herein, alone or in combination, refers to an aryl ether radical wherein the term aryl is defined as above. Examples of aryloxy radicals include phenoxy, thienyloxy and the like.

The terms "alkylthio" and "thioalkyl" as used herein, alone or in combination, refer to an alkyl thio radical, alkyl-S-, wherein the term alkyl is as defined above.

The term "arylthio" as used herein, alone or in combination, refers to an aryl thio radical, aryl-S-, wherein the term aryl is as defined above.

The term "heteroarylthio" as used herein, alone or in combination, refers to the group heteroaryl-S-, wherein the term heteroaryl is as defined above.


The term "acyl" as used herein, alone or in combination, refers to a radical -C(O)R where R includes alkyl, alkenyl, alkynyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl or heteroarylalkyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl or heteroaryl alkyl groups may be optionally substituted.

The term "acyloxy" as used herein, alone or in combination, refers to the ester group -OC(O)R, where R is H, alkyl, alkenyl, alkynyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl, or heteroarylalkyl wherein the alkyl, alkenyl, alkynyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl or heteroarylalkyl may be optionally substituted.

The term "carboxy esters" as used herein, alone or in combination, refers to -C(O)OR where R is alkyl, aryl or arylalkyl, wherein the alkyl, aryl and arylalkyl groups may be optionally substituted.


The term "BOC" as used herein, alone or in combination, refers to -C(O)Otbutyl.

The term "carboxamido" as used herein, alone or in combination, refers to



PCT/US2006/011846
 where each of R and R' are independently selected from the group consisting of H, alkyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl and heteroarylalkyl, wherein the alkyl, aryl, heteroaryl, alicyclic, heterocyclic, or arylalkyl groups may be optionally substituted.

The terms "thioamide" and "thioamidyl" as used herein, alone or in combination, refer to



where each of R and R' are independently selected from the group consisting of H, alkyl, aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl and heteroarylalkyl, wherein the alkyl, aryl, heteroaryl, alicyclic, heterocyclic, or arylalkyl groups may be optionally substituted.

The term "oxo" as used herein, alone or in combination, refers to =O.

The term "halogen" as used herein, alone or in combination, refers to F, Cl, Br and I.

The terms "haloalkyl", "haloalkenyl", "haloalkynyl" and "haloalkoxy" as used herein, alone or in combination, refer to alkyl, alkenyl, alkynyl and alkoxy structures, as described above, that are substituted with one or more fluorines, chlorines, bromines or iodines, or with combinations thereof.

The terms "perhaloalkyl", "perhaloalkyloxy" and "perhaloacyl" as used herein, alone or in combination, refer to alkyl, alkyloxy and acyl radicals as described above, in which all the H atoms are replaced by fluorines, chlorines, bromines or iodines, or combinations thereof.

The terms "lower perhaloalkyl", "lower perhaloalkyloxy" and "lower perhaloacyl" as used herein, alone or in combination, refer to perhaloalkyl, perhaloalkyloxy and perhaloacyl radicals as described above, having from two to about six carbons.

The terms cycloalkyl, arylalkyl, aryl, heteroaryl, alicyclic, heterocyclic, alkyl, alkynyl, alkenyl, haloalkyl, and heteroalkyl include optionally substituted cycloalkyl, arylalkyl, aryl, heteroaryl, alicyclic, heterocyclic, alkyl, alkynyl, alkenyl, haloalkyl and heteroalkyl groups.

The term "alkylsilyl" as used herein, alone or in combination, refers to -NRR'R" where R, R' and R" are alkyls.

The term "lower alkylsilyl" as used herein, alone or in combination, refers to -NRR'R" where R, R' and R" are lower alkyls.

The term "amino" as used herein, alone or in combination, refers to -NH₂.

The term "alkylamino" as used herein, alone or in combination, refers to the group -NHR where R is alkyl.

The term "aminoalkyl" as used herein, alone or in combination, refers to the group -alkylene-NH₂, wherein alkylene is as defined herein.

The term "dialkylamino" as used herein, alone or in combination, refers to the group -NRR' where R and R' are alkyls.

The term "aryalkylamino" as used herein, alone or in combination, refers to the group -NRR' where R is alkyl, and R' is aryl.

The term "diarylamino" as used herein, alone or in combination, refers to the group -NRR' where R and R' are aryls.

- 5 The term "heteroarylamino" as used herein, alone or in combination, refers to the group -NHR where R is heteroaryl.

The term "diheteroarylamino" as used herein, alone or in combination, refers to the group -NRR' where R and R' are heteroaryls.

- 10 The term "arylheteroarylamino" as used herein, alone or in combination, refers to the group -NRR' where R is aryl, and R' is heteroaryl.

- The term "carbamyl" as used herein, alone or in combination, refers to the -NHC(O)OR and -OC(O)NHR groups, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. Non-limiting examples of sulfanyl groups include methylsulfanyl (-SCH₃) and iso-propylsulfanyl (-SCH(CH₃)₂) and the like.
- 15

- The term "uryl" as used herein, alone or in combination, refers to the -NHC(O)NHR group, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. Non-limiting examples of sulfanyl groups include methylsulfanyl (-SCH₃) and iso-propylsulfanyl (-SCH(CH₃)₂) and the like.
- 20

- The term "thiouryl" as used herein, alone or in combination, refers to the -NHC(S)NHR group, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. Non-limiting examples of sulfanyl groups include methylsulfanyl (-SCH₃) and iso-propylsulfanyl (-SCH(CH₃)₂) and the like.
- 25

- The term "guanidinyl" as used herein, alone or in combination, refers to the -NHC(NH)NHR group, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. Non-limiting examples of sulfanyl groups include methylsulfanyl (-SCH₃) and iso-propylsulfanyl (-SCH(CH₃)₂) and the like.
- 30

~~PCT/US2006/011846~~
The terms "sulfide" and "thioether" as used herein, alone or in combination, refer to a sulfur atom covalently linked to two atoms; the formal oxidation state of said sulfur is (II). These terms may be used interchangeably.

The term "sulfanyl" as used herein, alone or in combination, refers to the -S-R group, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. Non-limiting examples of sulfanyl groups include methylsulfanyl (-SCH₃) and iso-propylsulfanyl (-SCH(CH₃)₂) and the like.

The term "sulfoxide" as used herein, alone or in combination, refers to a sulfur atom covalently linked to three atoms, at least one of which is an oxygen atom; the formal oxidation state of said sulfur atom is (IV).

The term "sulfinyl" as used herein, alone or in combination, refers to the groups -S(O)-R, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted. A non-limiting example of a sulfinyl group includes methylsulfinyl (-S(O)CH₃) and the like.

The term "sulfone" as used herein, alone or in combination, refers to a sulfur atom covalently linked to four atoms, at least two of which are oxygen atoms; the formal oxidation state of said sulfur atom is (VI).

The term "sulfonyl" as used herein, alone or in combination, refers to the groups -S(O₂)-R, wherein R may be, but is not limited to alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl, wherein the alkyl, alkenyl, alkynyl, aryl, alicyclic, heterocyclic, aryl, heteroaryl, arylalkyl and heteroarylalkyl groups may be optionally substituted.

A non-limiting example of a sulfonyl group includes methylsulfonyl (-S(O₂)CH₃) and the like.

The term "phosphite" as used herein, alone or in combination, refers to a phosphorus atom covalently linked to three carbon atoms, wherein the formal oxidation state of said phosphorus is (III).

The term "phosphinyl" as used herein, alone or in combination, refers to the monoradical derived from a phosphite group, as defined above.

The term "phosphonate" as used herein, alone or in combination, refers to a phosphorus atom covalently linked to four atoms, three of which are oxygen and one of which is carbon wherein the formal oxidation state of said phosphorus is (V).

The term "phosphonyl" as used herein, alone or in combination, refers to the monoradical derived from a phosphonate group, as defined above.

The term "phosphate" as used herein, alone or in combination, refers to a phosphorus atom covalently linked to four oxygen atoms, wherein the formal oxidation state of said phosphorus is (V).

The term "phosphatidyl" as used herein, alone or in combination, refers to the monoradical derived from a phosphate group, as defined above.

The term "phosphoramidate" as used herein, alone or in combination, refers to a phosphorus atom covalently linked to four atoms, three of which are nitrogen and one of which is oxygen wherein the formal oxidation state of said phosphorus is (V).

The term "phosphoramidyl" as used herein, alone or in combination, refers to the monoradical derived from a phosphoramidate group, as defined above.

The terms "optional" or "optionally" mean that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "aryl optionally mono- or di-substituted with an alkyl" means that the alkyl may but need not be present, or either one alkyl or two may be present, and the description includes situations where the aryl is substituted with one or two alkyls and situations where the aryl is not substituted with an alkyl.

"Optionally substituted" groups may be substituted or unsubstituted. The substituents of an "optionally substituted" group may include, without limitation, one or more substituents independently selected from the following groups or designated subsets thereof: lower alkyl, lower alkenyl, lower alkynyl, lower aryl, heteroaryl, alicyclic, heterocyclic, arylalkyl, heteroarylalkyl, lower alkoxy, lower aryloxy, amino, alkylamino, dialkylamino, diarylalkylamino, alkylthio, arylthio, heteroarylthio, oxo, oxa, acyl (-C(O)R), (-C(O)), carboxyesters (-C(O)OR), carboxamido (-C(O)NH₂), carboxy, acyloxy, -H, halo, -CN, -NO₂, -N₃, -SH, -OH, -(O)CH₃, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidine, pyridinyl, thiophene, furanyl, indole, indazole, esters, amides, phosphonates, phosphonic acid, phosphates, phosphoramides, sulfonates, sulfones, sulfates, sulphonamides, carbamates, ureas, thioureas, thioamides and thioalkyls. An optionally substituted group may be unsubstituted (*e.g.*, -CH₂CH₃), fully substituted (*e.g.*, -CF₂CF₃), monosubstituted (*e.g.*, -CH₂CH₂F) or substituted at a level anywhere in-between fully substituted and monosubstituted (*e.g.*, -CH₂CF₃).

The term "pyridine-1-oxy" also means "pyridine-N-oxy".


The term "heteroaryl group is substituted with 2 to 5 substituents" encompasses 1-oxy-pyridyl or N-oxy-pyridyl having 1 to 4 substituents, i.e. the oxygen atom of the pyridine-N-oxide should be counted as a substituent.

[030] Some of the compounds of the present invention may contain one or more chiral centers and therefore may exist in enantiomeric and diastereomeric forms. The scope of the present

invention is intended to cover all isomers *per se*, as well as mixtures of *cis* and *trans* isomers, mixtures of diastereomers and racemic mixtures of enantiomers (optical isomers) as well.

Further, it is possible using well known techniques to separate the various forms, and some embodiments of the invention may feature purified or enriched species of a given enantiomer or diastereomer.

[031] Some of the compounds of the present invention may exist in tautomeric forms. The scope of the present invention is intended to cover all tautomers. As a non-limiting example, compounds of formula (I), wherein R^0 is $-SR^8$ or $-OR^8$, and R^8 is hydrogen, could exist as either of the tautomers below:



[032] A “pharmacological composition” refers to a mixture of one or more of the compounds described herein, or pharmaceutically acceptable salts thereof, with other chemical components, such as pharmaceutically acceptable carriers and/or excipients. The purpose of a pharmacological composition is to facilitate administration of a compound to an organism.

[033] The phrase “pharmaceutically acceptable carrier” as used herein means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject agent from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations. A physiologically acceptable carrier should not cause significant irritation to an

~~PCT/US2006/011846~~
organism and does not abrogate the biological activity and properties of the administered compound.

An "excipient" refers to an inert substance added to a pharmacological composition to further facilitate administration of a compound. Examples of excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

A "pharmaceutically effective amount" means an amount which is capable of providing a therapeutic and/or prophylactic effect. The specific dose of compound administered according to this invention to obtain therapeutic and/or prophylactic effect will, of course, be determined by the particular circumstances surrounding the case, including, for example, the specific compound administered, the route of administration, the condition being treated, and the individual being treated. A typical daily dose (administered in single or divided doses) will contain a dosage level of from about 0.01 mg/kg to about 50-100 mg/kg of body weight of an active compound of the invention. Preferred daily doses generally will be from about 0.05 mg/kg to about 20 mg/kg and ideally from about 0.1 mg/kg to about 10 mg/kg. Factors such as clearance rate, half-life and maximum tolerated dose (MTD) have yet to be determined but one of ordinary skill in the art can determine these using standard procedures.

[034] In some method embodiments, the preferred therapeutic effect is the inhibition, to some extent, of the growth of cells characteristic of a proliferative disorder, *e.g.*, breast cancer. A therapeutic effect will also normally, but need not, relieve to some extent one or more of the symptoms other than cell growth or size of cell mass. A therapeutic effect may include, for example, one or more of 1) a reduction in the number of cells; 2) a reduction in cell size; 3) inhibition (*i.e.*, slowing to some extent, preferably stopping) of cell infiltration into peripheral organs, *e.g.*, in the instance of cancer metastasis; 3) inhibition (*i.e.*, slowing to some extent, preferably stopping) of tumor metastasis; 4) inhibition, to some extent, of cell growth; and/or 5) relieving to some extent one or more of the symptoms associated with the disorder.

As used herein, the term IC₅₀ refers to an amount, concentration or dosage of a particular test compound that achieves a 50% inhibition of a maximal response in an assay that measures such response. In some method embodiments of the invention, the "IC₅₀" value of a compound of the invention can be greater for normal cells than for cells exhibiting a proliferative disorder, *e.g.*, breast cancer cells. The value depends on the assay used.

[035] By a "standard" is meant a positive or negative control. A negative control in the context of Her2 expression levels is, *e.g.*, a sample possessing an amount of Her2 protein that correlates with a normal cell. A negative control may also include a sample that contains no Her2 protein.


By contrast, a positive control does contain Her2 protein, preferably of an amount that correlates

PCT/US2006/011846
 with overexpression as found in proliferative disorders, *e.g.*, breast cancers. The controls may be from cell or tissue samples, or else contain purified ligand (or absent ligand), immobilized or otherwise. In some embodiments, one or more of the controls may be in the form of a diagnostic "dipstick."

- 5 [036] By "selectively targeting" is meant affecting one type of cell to a greater extent than another, *e.g.*, in the case of cells with high as opposed to relatively low or normal Her2 levels.

[037] Compounds of the Invention

Compounds of the invention are related to alkynyl pyrrolo[2,3-d]pyrimidines of Formula I



10

and their polymorphs, solvates, esters, tautomers, diastereomers, enantiomers, pharmaceutically acceptable salts or prodrugs thereof, which show utility for inhibiting HSP90 and treating and preventing diseases that are HSP90-dependent, wherein:

- 15 R^0 is selected from the group consisting of hydrogen, halogen, lower alkyl, -CN, -SR⁸, -OR⁸, and -NHR⁸;

R^1 is selected from the group consisting of halogen, -OR¹¹, -SR¹¹ and lower alkyl;

R^2 is -NHR⁸;

- 20 R^3 is selected from the group consisting of hydrogen, -CN, -C(O)OH, -OR¹¹, -SR¹¹, -C(O)R⁹, -NR⁸R¹⁰, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower alkylsilyl, aryl, heteroaryl, alicyclyl and heterocyclyl, all optionally substituted, wherein:

the aryl, heteroaryl, alicyclyl and heterocyclyl groups are mono-, bi- or tri- cyclic;

- 25 R^8 and R^{10} taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

- 30 the optional substituents on R^3 are selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R⁸, lower aryl, heteroaryl, alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, amino, alkylamino, dialkylamino, arylalkylamino, diarylamino, heteroarylamino, diheteroarylamino, arylheteroarylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidiny, pyridinyl, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl,

PCT/US2006/011846

phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R^8 and R^8 taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R^4 is selected from the group consisting of optionally substituted lower alkylene, - $C(R^{12})_2$ -, -C(O)-, -C(S)-, -S(O)- and -SO₂-;

R^5 is selected from the group consisting of aryl, heteroaryl, alicyclyl and heterocyclyl, wherein:

the aryl group is substituted with 2 to 5 substituents;

the heteroaryl group is substituted with 2 to 5 substituents;

the alicyclyl group is substituted with 3 to 5 substituents;

the heterocyclyl group is substituted with 3 to 5 substituents;

the substituents on R^5 are selected from the group consisting of halogen, lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R¹⁰, lower aryl, lower heteroaryl, lower alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, thioalkyl, amino, alkylamino, dialkylamino, arylalkylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidiny, pyridinyl, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl, phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R^8 and R^{10} taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R^8 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower heteroaryl and -C(O)R⁹;

R^9 is selected from the group consisting of H, lower alkyl, lower alkenyl, lower alkynyl, lower aryl, lower heteroaryl, -NR¹⁰R¹⁰ and -OR¹¹, wherein

R^{10} and R^{10} taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

~~PCT/US2006/011846~~
~~R¹⁰ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower~~
 alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower
 heteroaryl and -C(O)R¹¹;

R¹¹ is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl,
 5 lower aryl and lower heteroaryl; and

R¹² is selected from the group consisting of hydrogen and lower alkyl.

[038] In some embodiments, R⁰ is hydrogen, halogen or -CN. In other embodiments, R⁰ is
 hydrogen, lower alkyl, -SR⁸ or -OR⁸. In other embodiments, R⁰ is hydrogen, -SR⁸, -OR⁸ or
 -NHR⁸. In other embodiments, R⁰ is -SR⁸ or -OR⁸. In other embodiments, R⁰ is hydrogen.

10 [039] In some embodiments, R¹ is halogen or lower alkyl. In other embodiments, R¹ is -OR¹¹ or
 -SR¹¹. In some embodiments, R¹ is halogen. In some embodiments, R¹ is chloro or bromo. In
 some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo.

[040] In some embodiments, R² is -NH₂ or -NHC(O)R⁹. In other embodiments, R² is -NH-
 lower alkyl, -NH-lower alkenyl, -NH-lower alkynyl, -NH-lower aryl or -NH-lower heteroaryl.
 15 In other embodiments, R² is -NHC(O)R⁹ where R⁹ is lower alkyl, lower alkenyl, lower alkynyl,
 lower aryl or lower heteroaryl. In other embodiments, R² is -NH₂. In other embodiments, R² is -
 NH(O)tBu.

[041] In some embodiments, R³ is hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower
 perhaloalkyl, aryl, heteroaryl, alicyclyl, heterocyclyl, -CN or -C(O)R⁹, all optionally substituted.
 20 In other embodiments, R³ is hydrogen, lower alkyl, aryl, heteroaryl, alicyclyl, heterocyclyl or
 -C(O)R⁹, all optionally substituted. In other embodiments, R³ is hydrogen, lower alkyl, aryl,
 heteroaryl, or -C(O)R⁹, all optionally substituted. In other embodiments, R³ is substituted lower
 alkyl wherein the substituent on the lower alkyl is selected from the group consisting of lower
 alkyl, -OR⁸, -C(O)R⁹ and -NR⁸R⁸. In other embodiments, R³ is lower alkyl, aryl, heteroaryl,
 25 -CN or -C(O)R⁹, all optionally substituted. In other embodiments, R³ is hydrogen. In other
 embodiments, R³ is optionally substituted lower alkyl. In other embodiments, R³ is optionally
 substituted phenyl or pyridinyl. In other embodiments, R³ is -(CH₂)_nOH, where n=1-3. In other
 embodiments, R³ is -(CH₂)_mC(R¹²)₂(CH₂)_nOH, where m=0-2 and n=1-2. In other embodiments,
 R³ is -(CH₂)_nNR⁸R⁸, wherein n=1-3, and each R⁸ is independently hydrogen or lower alkyl, or -
 30 NR⁸R⁸ are taken together forming an optionally substituted phthalimide or morpholin. In other
 embodiments, R³ is -(CH₂)_nC(O)NR¹⁰R¹⁰, wherein n=1-3, and each R¹⁰ is independently
 hydrogen or -C(O)R¹¹, or -NR¹⁰R¹⁰ are taken together forming an optionally substituted
 piperazine. In other embodiments, R³ is optionally substituted lower alkylsilyl. In other
 embodiments, R³ is -CO₂Et. In other embodiments, R³ is -(CH₂)_nC(O)NR¹⁰R¹⁰ where n=1-3, and

each R^0 is independently hydrogen, or $-C(O)R^{11}$. In other embodiments, R^3 is $-(CH_2)_nC(O)NHC(O)OtBu$.

[042] In some embodiments, R^4 is optionally substituted lower alkylene, $-C(O)-$, $-S(O)-$ or $-SO_2-$. In other embodiments, R^4 is $-CH_2-$, $-S(O)-$ or $-SO_2$. In other embodiments, R^4 is $-CHR^{12}-$. In other embodiments, R^4 is $-CH_2-$.

[043] In some embodiments, the aryl, heteroaryl, alicyclyl or heterocyclyl group of R^5 is monocyclic or bicyclic. In other embodiments, R^5 is substituted aryl or heteroaryl and the substituents on said aryl or heteroaryl are selected from the group consisting of halogen, lower alkoxy, lower alkyl, thioalkyl, amino, alkylamino, dialkylamino. In other embodiments, R^5 is substituted aryl or heteroaryl and the substituents on the aryl or heteroaryl are selected from the group consisting of halogen, lower alkoxy and lower alkyl.

[044] In some embodiments, R^0 is hydrogen, halogen, $-SH$, $-OH$, or $-CN$; R^1 is halogen; and R^2 is $-NH_2$ or $-NH-C(O)R^9$.

[045] In some embodiments, R^1 is chloro or bromo; R^2 is $-NH_2$ or $-NH-C(O)R^9$; and R^3 is lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower aryl, or lower heteroaryl, all optionally substituted with $-OR^8$, $-NR^8R^8$ or $-C(O)R^9$.

[046] In some embodiments, R^0 is hydrogen, halogen or $-CN$; R^2 is $-NH_2$ or $-NH-C(O)R^9$; and R^4 is $-CH_2-$.

[047] In other embodiments, R^0 is hydrogen, halogen, $-SH$, $-OH$ or $-CN$; R^1 is halogen; R^2 is $-NH_2$; R^3 is hydrogen, $-OR^{11}$, $-SR^{11}$, $-NR^8R^8$, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower aryl, or lower heteroaryl, wherein the R^8 in R^3 is hydrogen, lower alkyl, lower heteroalkyl, lower aryl, or $-C(O)R^9$; R^4 is $-CH_2-$; and R^5 is aryl or heteroaryl, substituted with 2 to 5 substituents.

[048] In some embodiments, R^0 is selected from hydrogen, halogen and $-CN$; R^1 is halogen; R^2 is $-NHR^8$; R^3 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, aryl, heteroaryl, alicyclyl and heterocyclyl, all optionally substituted; R^4 is $-CHR^{12}-$; and R^5 optionally substituted is aryl or heteroaryl.

[049] In other embodiments, R^0 is selected from hydrogen, halogen and $-CN$; R^1 is halogen; R^2 is $-NH_2$; R^3 is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, all optionally substituted with $-OR^8$, $-NR^8R^8$ or $-C(O)R^9$; R^4 is $-CH_2-$; and R^5 is aryl or heteroaryl.

[050] In other embodiments, R^1 is chloro or bromo; R^2 is $-NH_2$; and R^5 is a phenyl having at least three substituents.

[051] In other embodiments, R^1 is chloro or bromo; R^2 is $-NH_2$; and R^5 is a pyridyl having at least two substituents.

[052] In other embodiments, R^1 is chloro or bromo; R^2 is $-NH_2$; and R^5 is 1-oxy-pyridyl (N-oxy-pyridyl) having at least two substituents.


[053] In other selected embodiments, R^0 is hydrogen; R^1 is chloro or bromo; R^2 is $-NH_2$ or $-NHC(O)tBu$; R^3 is selected from the group consisting of hydrogen, lower alkyl, aryl, heteroaryl, $-CN$, optionally substituted lower alkylsilyl, $-CO_2Et$, $-(CH_2)_mC(CH_3)_nOH$ where $m=0-2$ and $n=1-2$, $-(CH_2)_nOH$ where $n=1-3$, and $-(CH_2)_nNR^8R^{10}$ wherein $n=1-3$, R^8 and R^{10} are independently hydrogen, lower alkyl, or are taken together forming a piperazine, a phthalimide or a morpholine; R^4 is $-CH_2-$; R^5 is substituted aryl or heteroaryl, and the substituents on the aryl or heteroaryl are selected from the group consisting of halogen, lower alkoxy, lower alkyl, thioalkyl, amino, alkylamino and dialkylamino.


[054] In some selected embodiments, R^0 is hydrogen; R^1 is chloro or bromo; R^2 is $-NH_2$; R^3 is selected from the group consisting of hydrogen, lower alkyl, $-(CH_2)_nOH$, where $n=1-3$, $-(CH_2)_mC(CH_3)_nOH$, where $m=0-2$ and $n=1-2$, $-(CH_2)_2C(O)NH_2$, $-(CH_2)OC(O)CH_2N(CH_3)_2$, $-(CH_2)_nNHC(O)OtBu$ and $-(CH_2)_nNR^8R^8$, wherein $n=1-3$, R^8 and R^8 are independently hydrogen, lower alkyl, or are taken together forming an optionally substituted phthalimide or morpholine; R^4 is $-CH_2-$; R^5 is substituted heteroaryl, and the substituents on said heteroaryl are independently selected from the group consisting of lower alkoxy and lower alkyl.


[055] In some selected embodiments, R^0 is hydrogen; R^1 is chloro or bromo; R^2 is $-NH_2$; R^3 is selected from the group consisting of hydrogen, lower alkyl, $-(CH_2)_nOH$, where $n=1-3$, $-(CH_2)_mC(CH_3)_nOH$, where $m=0-2$ and $n=1-2$, $-(CH_2)_nC(O)NR^{10}R^{10}$ where $n=1-3$, and , R^{10} and R^{10} are independently hydrogen or $-C(O)R^{11}$, or are taken together forming an optionally substituted piperazine; R^4 is $-CH_2-$; R^5 is substituted heteroaryl, and the substituents on said heteroaryl are independently selected from the group consisting of lower alkoxy and lower alkyl.


[056] In some selected embodiments R^3 is substituted lower alkyl, and in some further selected embodiments R^3 is substituted lower alkyl, and the substituent on said lower alkyl is phosphonyl or phosphatidyl.


[057] In other preferred embodiments, are the following compounds





















[058] It should be understood that any of the above described embodiments of the invention can be combined in anyway where practical; those of ordinary skill in the art will appreciate the ways the various embodiments may be combined usefully within the spirit of the invention. A non-limiting list of compounds based on Formula I of the invention is exemplified in TABLE 1.

[059] TABLE 1: Exemplary Compounds based on Formula I



wherein R⁰ is H, and R⁴ is -CH₂-.

I

No.	Ex	R ¹	R ²	R ³	R ⁵
1		Cl	NH ₂	H	3,4,5-Trimethoxyphenyl
2		Cl	NH ₂	H	2-Chloro-3,4,5-trimethoxyphenyl
3		Cl	NH ₂	H	2-Bromo-3,4,5-trimethoxyphenyl
4		Cl	NH ₂	H	2-Iodo-3,4,5-trimethoxyphenyl
5		Cl	NH ₂	H	2-Fluoro-3,4,5-trimethoxyphenyl
6		Cl	NH ₂	H	3,4,5-Trimethylphenyl
7		Cl	NH ₂	H	2-Chloro-3,4,5-trimethylphenyl
8		Cl	NH ₂	H	2-Bromo-3,4,5-trimethylphenyl
9		Cl	NH ₂	H	2-Iodo-3,4,5-trimethylphenyl
10		Cl	NH ₂	H	2-Fluoro-3,4,5-trimethylphenyl
11		Cl	NH ₂	H	3,5-Dimethoxy-4-methylphenyl
12		Cl	NH ₂	H	2-Chloro-3,5-dimethoxy-4-methylphenyl
13		Cl	NH ₂	H	2-Bromo-3,5-dimethoxy-4-methylphenyl
14		Cl	NH ₂	H	2-Iodo-3,5-dimethoxy-4-methylphenyl
15		Cl	NH ₂	H	2-Fluoro-3,5-dimethoxy-4-methylphenyl
16		Cl	NH ₂	2-Py	3,4,5-Trimethoxyphenyl
17		Cl	NH ₂	2-Py	2-Chloro-3,4,5-trimethoxyphenyl
18		Cl	NH ₂	2-Py	2-Bromo-3,4,5-trimethoxyphenyl
19		Cl	NH ₂	2-Py	2-Iodo-3,4,5-trimethoxyphenyl
20		Cl	NH ₂	2-Py	2-Fluoro-3,4,5-trimethoxyphenyl
21		Cl	NH ₂	2-Py	3,4,5-Trimethylphenyl
22		Cl	NH ₂	2-Py	2-Chloro-3,4,5-trimethylphenyl
23		Cl	NH ₂	2-Py	2-Bromo-3,4,5-trimethylphenyl
24		Cl	NH ₂	2-Py	2-Iodo-3,4,5-trimethylphenyl
25		Cl	NH ₂	2-Py	2-Fluoro-3,4,5-trimethylphenyl
26		Cl	NH ₂	2-Py	3,5-Dimethoxy-4-methylphenyl
27		Cl	NH ₂	2-Py	2-Chloro-3,5-dimethoxy-4-methylphenyl
28		Cl	NH ₂	2-Py	2-Bromo-3,5-dimethoxy-4-methylphenyl
29		Cl	NH ₂	2-Py	2-Iodo-3,5-dimethoxy-4-methylphenyl
30		Cl	NH ₂	2-Py	2-Fluoro-3,5-dimethoxy-4-methylphenyl
31		Cl	NH ₂	2-Py	3,4,5-Trimethoxyphenyl
32		Cl	NH ₂	2-Py	2-Chloro-3,4,5-trimethoxyphenyl
33		Cl	NH ₂	2-Py	2-Bromo-3,4,5-trimethoxyphenyl
34		Cl	NH ₂	2-Py	2-Iodo-3,4,5-trimethoxyphenyl
35		Cl	NH ₂	2-Py	2-Fluoro-3,4,5-trimethoxyphenyl
36		Cl	NH ₂	2-Py	3,4,5-Trimethylphenyl
37		Cl	NH ₂	2-Py	2-Chloro-3,4,5-trimethylphenyl
38		Cl	NH ₂	2-Py	2-Bromo-3,4,5-trimethylphenyl
39		Cl	NH ₂	2-Py	2-Iodo-3,4,5-trimethylphenyl
40		Cl	NH ₂	2-Py	2-Fluoro-3,4,5-trimethylphenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
41		Cl	NH ₂	Et	3,5-Dimethoxy-4-methylphenyl
42		Cl	NH ₂	Et	2-Chloro-3,5-dimethoxy-4-methylphenyl
43		Cl	NH ₂	Et	2-Bromo-3,5-dimethoxy-4-methylphenyl
44		Cl	NH ₂	Et	2-Iodo-3,5-dimethoxy-4-methylphenyl
45		Cl	NH ₂	Et	2-Fluoro-3,5-dimethoxy-4-methylphenyl
46		Cl	NH ₂	Me	3,4,5-Trimethoxyphenyl
47		Cl	NH ₂	Me	2-Chloro-3,4,5-trimethoxyphenyl
48		Cl	NH ₂	Me	2-Bromo-3,4,5-trimethoxyphenyl
49		Cl	NH ₂	Me	2-Iodo-3,4,5-trimethoxyphenyl
50		Cl	NH ₂	Me	2-Fluoro-3,4,5-trimethoxyphenyl
51		Cl	NH ₂	Me	3,4,5-Trimethylphenyl
52		Cl	NH ₂	Me	2-Chloro-3,4,5-trimethylphenyl
53		Cl	NH ₂	Me	2-Bromo-3,4,5-trimethylphenyl
54		Cl	NH ₂	Me	2-Iodo-3,4,5-trimethylphenyl
55		Cl	NH ₂	Me	2-Fluoro-3,4,5-trimethylphenyl
56		Cl	NH ₂	Me	3,5-Dimethoxy-4-methylphenyl
57		Cl	NH ₂	Me	2-Chloro-3,5-dimethoxy-4-methylphenyl
58		Cl	NH ₂	Me	2-Bromo-3,5-dimethoxy-4-methylphenyl
59		Cl	NH ₂	Me	2-Iodo-3,5-dimethoxy-4-methylphenyl
60		Cl	NH ₂	Me	2-Fluoro-3,5-dimethoxy-4-methylphenyl
61		Cl	NH ₂	Ph	3,4,5-Trimethoxyphenyl
62		Cl	NH ₂	Ph	2-Chloro-3,4,5-trimethoxyphenyl
63		Cl	NH ₂	Ph	2-Bromo-3,4,5-trimethoxyphenyl
64		Cl	NH ₂	Ph	2-Iodo-3,4,5-trimethoxyphenyl
65		Cl	NH ₂	Ph	2-Fluoro-3,4,5-trimethoxyphenyl
66		Cl	NH ₂	Ph	3,4,5-Trimethylphenyl
67		Cl	NH ₂	Ph	2-Chloro-3,4,5-trimethylphenyl
68		Cl	NH ₂	Ph	2-Bromo-3,4,5-trimethylphenyl
69		Cl	NH ₂	Ph	2-Iodo-3,4,5-trimethylphenyl
70		Cl	NH ₂	Ph	2-Fluoro-3,4,5-trimethylphenyl
71		Cl	NH ₂	Ph	3,5-Dimethoxy-4-methylphenyl
72		Cl	NH ₂	Ph	2-Chloro-3,5-dimethoxy-4-methylphenyl
73		Cl	NH ₂	Ph	2-Bromo-3,5-dimethoxy-4-methylphenyl
74		Cl	NH ₂	Ph	2-Iodo-3,5-dimethoxy-4-methylphenyl
75		Cl	NH ₂	Ph	2-Fluoro-3,5-dimethoxy-4-methylphenyl
76		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethoxyphenyl
77		Cl	NH ₂	(CH ₂) ₂ OH	2-Chloro-3,4,5-trimethoxyphenyl
78		Cl	NH ₂	(CH ₂) ₂ OH	2-Bromo-3,4,5-trimethoxyphenyl
79		Cl	NH ₂	(CH ₂) ₂ OH	2-Iodo-3,4,5-trimethoxyphenyl
80		Cl	NH ₂	(CH ₂) ₂ OH	2-Fluoro-3,4,5-trimethoxyphenyl
81		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethylphenyl
82		Cl	NH ₂	(CH ₂) ₂ OH	2-Chloro-3,4,5-trimethylphenyl
83		Cl	NH ₂	(CH ₂) ₂ OH	2-Bromo-3,4,5-trimethylphenyl
84		Cl	NH ₂	(CH ₂) ₂ OH	2-Iodo-3,4,5-trimethylphenyl
85		Cl	NH ₂	(CH ₂) ₂ OH	2-Fluoro-3,4,5-trimethylphenyl
86		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethoxy-4-methylphenyl
87		Cl	NH ₂	(CH ₂) ₂ OH	2-Chloro-3,5-dimethoxy-4-methylphenyl
88		Cl	NH ₂	(CH ₂) ₂ OH	2-Bromo-3,5-dimethoxy-4-methylphenyl
89		Cl	NH ₂	(CH ₂) ₂ OH	2-Iodo-3,5-dimethoxy-4-methylphenyl
90		Cl	NH ₂	(CH ₂) ₂ OH	2-Fluoro-3,5-dimethoxy-4-methylphenyl
91		Cl	NH ₂	4-Py	3,4,5-Trimethoxyphenyl
92		Cl	NH ₂	4-Py	2-Chloro-3,4,5-trimethoxyphenyl
93		Cl	NH ₂	4-Py	2-Bromo-3,4,5-trimethoxyphenyl
94		Cl	NH ₂	4-Py	2-Iodo-3,4,5-trimethoxyphenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
95		Cl	NH ₂	4-Py	2-Fluoro-3,4,5-trimethoxyphenyl
96		Cl	NH ₂	Ph	3,4,5-Trimethylphenyl
97		Cl	NH ₂	Ph	2-Chloro-3,4,5-trimethylphenyl
98		Cl	NH ₂	Ph	2-Bromo-3,4,5-trimethylphenyl
99		Cl	NH ₂	Ph	2-Iodo-3,4,5-trimethylphenyl
100		Cl	NH ₂	Ph	2-Fluoro-3,4,5-trimethylphenyl
101		Cl	NH ₂	Ph	3,5-Dimethoxy-4-methylphenyl
102		Cl	NH ₂	Ph	2-Chloro-3,5-dimethoxy-4-methylphenyl
103		Cl	NH ₂	Ph	2-Bromo-3,5-dimethoxy-4-methylphenyl
104		Cl	NH ₂	Ph	2-Iodo-3,5-dimethoxy-4-methylphenyl
105		Cl	NH ₂	Pr	2-Fluoro-3,5-dimethoxy-4-methylphenyl
106		Cl	NH ₂	Pr	3,5-Dimethoxy-4-methylphenyl
107		Cl	NH ₂	Pr	2-Chloro-3,5-dimethoxy-4-methylphenyl
108		Cl	NH ₂	Pr	2-Bromo-3,5-dimethoxy-4-methylphenyl
109		Cl	NH ₂	Pr	2-Iodo-3,5-dimethoxy-4-methylphenyl
110		Cl	NH ₂	Pr	2-Fluoro-3,5-dimethoxy-4-methylphenyl
111		Cl	NH ₂	Pr	3,4,5-Trimethoxyphenyl
112		Cl	NH ₂	Pr	2-Chloro-3,4,5-trimethoxyphenyl
113		Cl	NH ₂	Pr	2-Bromo-3,4,5-trimethoxyphenyl
114		Cl	NH ₂	Pr	2-Iodo-3,4,5-trimethoxyphenyl
115		Cl	NH ₂	Pr	2-Fluoro-3,4,5-trimethoxyphenyl
116		Cl	NH ₂	Pr	3,4,5-Trimethylphenyl
117		Cl	NH ₂	Pr	2-Chloro-3,4,5-trimethylphenyl
118		Cl	NH ₂	Pr	2-Bromo-3,4,5-trimethylphenyl
119		Cl	NH ₂	Pr	2-Iodo-3,4,5-trimethylphenyl
120		Cl	NH ₂	Pr	2-Fluoro-3,4,5-trimethylphenyl
121		Cl	NH ₂	Pr	3,5-Dimethoxy-4-methylphenyl
122		Cl	NH ₂	Pr	2-Chloro-3,5-dimethoxy-4-methylphenyl
123		Cl	NH ₂	Pr	2-Bromo-3,5-dimethoxy-4-methylphenyl
124		Cl	NH ₂	Pr	2-Iodo-3,5-dimethoxy-4-methylphenyl
125		Cl	NH ₂	Pr	2-Fluoro-3,5-dimethoxy-4-methylphenyl
126		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethoxyphenyl
127		Cl	NH ₂	(CH ₂) ₃ OH	2-Chloro-3,4,5-trimethoxyphenyl
128		Cl	NH ₂	(CH ₂) ₃ OH	2-Bromo-3,4,5-trimethoxyphenyl
129		Cl	NH ₂	(CH ₂) ₃ OH	2-Iodo-3,4,5-trimethoxyphenyl
130		Cl	NH ₂	(CH ₂) ₃ OH	2-Fluoro-3,4,5-trimethoxyphenyl
131		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethylphenyl
132		Cl	NH ₂	(CH ₂) ₃ OH	2-Chloro-3,4,5-trimethylphenyl
133		Cl	NH ₂	(CH ₂) ₃ OH	2-Bromo-3,4,5-trimethylphenyl
134		Cl	NH ₂	(CH ₂) ₃ OH	2-Iodo-3,4,5-trimethylphenyl
135		Cl	NH ₂	(CH ₂) ₃ OH	2-Fluoro-3,4,5-trimethylphenyl
136		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethoxy-4-methylphenyl
137		Cl	NH ₂	(CH ₂) ₃ OH	2-Chloro-3,5-dimethoxy-4-methylphenyl
138		Cl	NH ₂	(CH ₂) ₃ OH	2-Bromo-3,5-dimethoxy-4-methylphenyl
139		Cl	NH ₂	(CH ₂) ₃ OH	2-Iodo-3,5-dimethoxy-4-methylphenyl
140		Cl	NH ₂	(CH ₂) ₃ OH	2-Fluoro-3,5-dimethoxy-4-methylphenyl
141		Cl	NH ₂	i-Bu	3,4,5-Trimethoxyphenyl
142		Cl	NH ₂	i-Bu	2-Chloro-3,4,5-trimethoxyphenyl
143		Cl	NH ₂	i-Bu	2-Bromo-3,4,5-trimethoxyphenyl
144		Cl	NH ₂	i-Bu	2-Iodo-3,4,5-trimethoxyphenyl
145		Cl	NH ₂	i-Bu	2-Fluoro-3,4,5-trimethoxyphenyl
146		Cl	NH ₂	i-Bu	3,4,5-Trimethylphenyl
147		Cl	NH ₂	i-Bu	2-Chloro-3,4,5-trimethylphenyl
148		Cl	NH ₂	i-Bu	2-Bromo-3,4,5-trimethylphenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
149		Cl	NH ₂	i-Bu	2-Iodo-3,4,5-trimethylphenyl
150		Cl	NH ₂	i-Bu	2-Fluoro-3,4,5-trimethylphenyl
151		Cl	NH ₂	i-Bu	3,5-Dimethoxy-4-methylphenyl
152		Cl	NH ₂	i-Bu	2-Chloro-3,5-dimethoxy-4-methylphenyl
153		Cl	NH ₂	i-Bu	2-Bromo-3,5-dimethoxy-4-methylphenyl
154		Cl	NH ₂	i-Bu	2-Iodo-3,5-dimethoxy-4-methylphenyl
155		Cl	NH ₂	i-Bu	2-Fluoro-3,5-dimethoxy-4-methylphenyl
156		Cl	NH ₂	CN	3,4,5-Trimethoxyphenyl
157		Cl	NH ₂	CN	2-Chloro-3,4,5-trimethoxyphenyl
158		Cl	NH ₂	CN	2-Bromo-3,4,5-trimethoxyphenyl
159		Cl	NH ₂	CN	2-Iodo-3,4,5-trimethoxyphenyl
160		Cl	NH ₂	CN	3,4,5-Trimethoxyphenyl
161		Cl	NH ₂	CN	2-Chloro-3,4,5-trimethoxyphenyl
162		Cl	NH ₂	CN	2-Bromo-3,4,5-trimethoxyphenyl
163		Cl	NH ₂	CN	2-Iodo-3,4,5-trimethoxyphenyl
164		Cl	NH ₂	CN	2-Fluoro-3,4,5-trimethoxyphenyl
165		Cl	NH ₂	CN	3,4,5-Trimethylphenyl
166		Cl	NH ₂	CN	2-Chloro-3,4,5-trimethylphenyl
167		Cl	NH ₂	CN	2-Bromo-3,4,5-trimethylphenyl
168		Cl	NH ₂	CN	2-Iodo-3,4,5-trimethylphenyl
169		Cl	NH ₂	CN	2-Fluoro-3,4,5-trimethylphenyl
170		Cl	NH ₂	CN	3,5-Dimethoxy-4-methylphenyl
171		Cl	NH ₂	CN	2-Chloro-3,5-dimethoxy-4-methylphenyl
172		Cl	NH ₂	CN	2-Bromo-3,5-dimethoxy-4-methylphenyl
173		Cl	NH ₂	CN	2-Iodo-3,5-dimethoxy-4-methylphenyl
174		Cl	NH ₂	CN	2-Fluoro-3,5-dimethoxy-4-methylphenyl
175		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethoxyphenyl
176		Cl	NH ₂	CH ₂ OH	2-Chloro-3,4,5-trimethoxyphenyl
177		Cl	NH ₂	CH ₂ OH	2-Bromo-3,4,5-trimethoxyphenyl
178		Cl	NH ₂	CH ₂ OH	2-Iodo-3,4,5-trimethoxyphenyl
179		Cl	NH ₂	CH ₂ OH	2-Fluoro-3,4,5-trimethoxyphenyl
180		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethylphenyl
181		Cl	NH ₂	CH ₂ OH	2-Chloro-3,4,5-trimethylphenyl
182		Cl	NH ₂	CH ₂ OH	2-Bromo-3,4,5-trimethylphenyl
183		Cl	NH ₂	CH ₂ OH	2-Iodo-3,4,5-trimethylphenyl
184		Cl	NH ₂	CH ₂ OH	2-Fluoro-3,4,5-trimethylphenyl
185		Cl	NH ₂	CH ₂ OH	3,5-Dimethoxy-4-methylphenyl
186		Cl	NH ₂	CH ₂ OH	2-Chloro-3,5-dimethoxy-4-methylphenyl
187		Cl	NH ₂	CH ₂ OH	2-Bromo-3,5-dimethoxy-4-methylphenyl
188		Cl	NH ₂	CH ₂ OH	2-Iodo-3,5-dimethoxy-4-methylphenyl
189		Cl	NH ₂	CH ₂ OH	2-Fluoro-3,5-dimethoxy-4-methylphenyl
190		Cl	NH ₂	i-Pr	3,4,5-Trimethoxyphenyl
191		Cl	NH ₂	i-Pr	2-Chloro-3,4,5-trimethoxyphenyl
192		Cl	NH ₂	i-Pr	2-Bromo-3,4,5-trimethoxyphenyl
193		Cl	NH ₂	i-Pr	2-Iodo-3,4,5-trimethoxyphenyl
194		Cl	NH ₂	i-Pr	2-Fluoro-3,4,5-trimethoxyphenyl
195		Cl	NH ₂	i-Pr	3,4,5-Trimethylphenyl
196		Cl	NH ₂	i-Pr	2-Chloro-3,4,5-trimethylphenyl
197		Cl	NH ₂	i-Pr	2-Bromo-3,4,5-trimethylphenyl
198		Cl	NH ₂	i-Pr	2-Iodo-3,4,5-trimethylphenyl
199		Cl	NH ₂	i-Pr	2-Fluoro-3,4,5-trimethylphenyl
200		Cl	NH ₂	i-Pr	3,5-Dimethoxy-4-methylphenyl
201		Cl	NH ₂	i-Pr	2-Chloro-3,5-dimethoxy-4-methylphenyl
202		Cl	NH ₂	i-Pr	2-Bromo-3,5-dimethoxy-4-methylphenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
203		Cl	NH ₂	i-Pr	2-Iodo-3,5-dimethoxy-4-methylphenyl
204		Cl	NH ₂	i-Pr	2-Fluoro-3,5-dimethoxy-4-methylphenyl
205		Cl	NH ₂	CO ₂ Et	3,4,5-Trimethoxyphenyl
206		Cl	NH ₂	CO ₂ Et	2-Chloro-3,4,5-trimethoxyphenyl
207		Cl	NH ₂	CO ₂ Et	2-Bromo-3,4,5-trimethoxyphenyl
208		Cl	NH ₂	CO ₂ Et	2-Iodo-3,4,5-trimethoxyphenyl
209		Cl	NH ₂	CO ₂ Et	2-Fluoro-3,4,5-trimethoxyphenyl
210		Cl	NH ₂	CO ₂ Et	3,4,5-Trimethylphenyl
211		Cl	NH ₂	CO ₂ Et	2-Chloro-3,4,5-trimethylphenyl
212		Cl	NH ₂	CO ₂ Et	2-Bromo-3,4,5-trimethylphenyl
213		Cl	NH ₂	CO ₂ Et	2-Iodo-3,4,5-trimethylphenyl
214		Cl	NH ₂	CO ₂ Et	2-Fluoro-3,4,5-trimethylphenyl
215		Cl	NH ₂	CO ₂ Et	3,5-Dimethoxy-4-methylphenyl
216		Cl	NH ₂	CO ₂ Et	2-Chloro-3,5-dimethoxy-4-methylphenyl
217		Cl	NH ₂	CO ₂ Et	2-Bromo-3,5-dimethoxy-4-methylphenyl
218		Cl	NH ₂	CO ₂ Et	2-Iodo-3,5-dimethoxy-4-methylphenyl
219		Cl	NH ₂	CO ₂ Et	2-Fluoro-3,5-dimethoxy-4-methylphenyl
220		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethoxyphenyl
221		Cl	NH ₂	CH ₂ -NMe ₂	2-Chloro-3,4,5-trimethoxyphenyl
222		Cl	NH ₂	CH ₂ -NMe ₂	2-Bromo-3,4,5-trimethoxyphenyl
223		Cl	NH ₂	CH ₂ -NMe ₂	2-Iodo-3,4,5-trimethoxyphenyl
224		Cl	NH ₂	CH ₂ -NMe ₂	2-Fluoro-3,4,5-trimethoxyphenyl
225		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethylphenyl
226		Cl	NH ₂	CH ₂ -NMe ₂	2-Chloro-3,4,5-trimethylphenyl
227		Cl	NH ₂	CH ₂ -NMe ₂	2-Bromo-3,4,5-trimethylphenyl
228		Cl	NH ₂	CH ₂ -NMe ₂	2-Iodo-3,4,5-trimethylphenyl
229		Cl	NH ₂	CH ₂ -NMe ₂	2-Fluoro-3,4,5-trimethylphenyl
230		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethoxy-4-methylphenyl
231		Cl	NH ₂	CH ₂ -NMe ₂	2-Chloro-3,5-dimethoxy-4-methylphenyl
232		Cl	NH ₂	CH ₂ -NMe ₂	2-Bromo-3,5-dimethoxy-4-methylphenyl
233		Cl	NH ₂	CH ₂ -NMe ₂	2-Iodo-3,5-dimethoxy-4-methylphenyl
234		Cl	NH ₂	CH ₂ -NMe ₂	2-Fluoro-3,5-dimethoxy-4-methylphenyl
235		Cl	NH ₂	3-Py	3,4,5-Trimethoxyphenyl
236		Cl	NH ₂	3-Py	2-Chloro-3,4,5-trimethoxyphenyl
237		Cl	NH ₂	3-Py	2-Bromo-3,4,5-trimethoxyphenyl
238		Cl	NH ₂	3-Py	2-Iodo-3,4,5-trimethoxyphenyl
239		Cl	NH ₂	3-Py	2-Fluoro-3,4,5-trimethoxyphenyl
240	2	Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
241		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
242		Cl	NH ₂	(CH ₂) ₂ OH	6-Bromo-3,5-dimethyl-4-methoxypyridin-2-yl
243		Cl	NH ₂	(CH ₂) ₂ OH	6-Chloro-3,5-dimethyl-4-methoxypyridin-2-yl
244		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-bromopyridin-2-yl
245		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
246		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-chloropyridin-2-yl
247		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
248		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
249		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
250		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
251		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
252		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethyl-pyridin-2-yl
253		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
254		Cl	NH ₂	(CH ₂) ₂ OH	4,5,6-Trimethoxypyridin-2-yl
255		Cl	NH ₂	(CH ₂) ₂ OH	4,5,6-Trimethoxy-1-oxypyridin-2-yl
256		Cl	NH ₂	(CH ₂) ₂ OH	3-Bromo-4,5,6-trimethoxypyridin-2-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
257		Cl	NH ₂	(CH ₂) ₂ OH	3-Chloro-4,5,6-trimethoxypyridin-2-yl
258		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethoxy-pyridin-2-yl
259		Cl	NH ₂	(CH ₂) ₂ OH	3,4,5-Trimethoxy-1-oxypyridin-2-yl
260		Cl	NH ₂	(CH ₂) ₂ OH	3-Bromo-3,4,5-trimethoxy-pyridin-2-yl
261		Cl	NH ₂	(CH ₂) ₂ OH	3-Chloro-3,4,5-trimethoxy-pyridin-2-yl
262		Cl	NH ₂	(CH ₂) ₂ OH	4,5,6-Trimethyl-pyridin-2-yl
263		Cl	NH ₂	(CH ₂) ₂ OH	4,5,6-Trimethyl-1-oxypyridin-2-yl
264		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-methoxy-pyridin-2-yl
265		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
266		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
267		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-bromopyridin-3-yl
268		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-chloropyridin-3-yl
269		Cl	NH ₂	(CH ₂) ₂ OH	5,6-Dimethyl-4-bromopyridin-3-yl
270		Cl	NH ₂	(CH ₂) ₂ OH	5,6-Dimethyl-4-chloropyridin-3-yl
271		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-methoxypyridin-4-yl
272		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-pyridin-4-yl
273		Cl	NH ₂	(CH ₂) ₂ OH	2,3,6-Trimethyl-pyridin-4-yl
274		Cl	NH ₂	(CH ₂) ₂ OH	2,3,6-Trimethoxy-pyridin-4-yl
275		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-bromopyridin-4-yl
276		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-chloropyridin-4-yl
277		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-methoxy-1-oxy-pyridin-4-yl
278		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-1-oxy-pyridin-4-yl
279		Cl	NH ₂	(CH ₂) ₂ OH	2,3,6-Trimethyl-1-oxy-pyridin-4-yl
280		Cl	NH ₂	(CH ₂) ₂ OH	2,3,6-Trimethoxy-1-oxy-pyridin-4-yl
281		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-bromo1-oxy-pyridin-4-yl
282		Cl	NH ₂	(CH ₂) ₂ OH	2,6-Dimethyl-3-chloro1-oxy-pyridin-4-yl
283		Cl	NH ₂	(CH ₂) ₂ OH	4,6-Dimethyl-5-iodopyridin-3-yl
284		Cl	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-aminopyridin-2-yl
285		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-methoxypyridin-2-yl
286		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
287		Cl	NH ₂	i-Pr	6-Bromo-3,5-dimethyl-4-methoxypyridin-2-yl
288		Cl	NH ₂	i-Pr	6-Chloro-3,5-dimethyl-4-methoxypyridin-2-yl
289		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-bromopyridin-2-yl
290		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
291		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-chloropyridin-2-yl
292		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
293		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-iodopyridin-2-yl
294		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
295		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
296		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
297		Cl	NH ₂	i-Pr	3,4,5-Trimethyl-pyridin-2-yl
298		Cl	NH ₂	i-Pr	3,4,5-Trimethyl-1-oxypyridin-2-yl
299		Cl	NH ₂	i-Pr	4,5,6-Trimethoxypyridin-2-yl
300		Cl	NH ₂	i-Pr	4,5,6-Trimethoxy-1-oxypyridin-2-yl
301		Cl	NH ₂	i-Pr	3-Bromo-4,5,6-trimethoxypyridin-2-yl
302		Cl	NH ₂	i-Pr	3-Chloro-4,5,6-trimethoxypyridin-2-yl
303		Cl	NH ₂	i-Pr	3,4,5-Trimethoxy-pyridin-2-yl
304		Cl	NH ₂	i-Pr	3,4,5-Trimethoxy-1-oxypyridin-2-yl
305		Cl	NH ₂	i-Pr	3-Bromo-3,4,5-trimethoxy-pyridin-2-yl
306		Cl	NH ₂	i-Pr	3-Chloro-3,4,5-trimethoxy-pyridin-2-yl
307		Cl	NH ₂	i-Pr	4,5,6-Trimethyl-pyridin-2-yl
308		Cl	NH ₂	i-Pr	4,5,6-Trimethyl-1-oxypyridin-2-yl
309		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-methoxy-pyridin-2-yl
310		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-methoxypyridin-3-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
311		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
312		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-bromopyridin-3-yl
313		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-chloropyridin-3-yl
314		Cl	NH ₂	i-Pr	5,6-Dimethyl-4-bromopyridin-3-yl
315		Cl	NH ₂	i-Pr	5,6-Dimethyl-4-chloropyridin-3-yl
316		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-methoxypyridin-4-yl
317		Cl	NH ₂	i-Pr	2,6-Dimethyl-pyridin-4-yl
318		Cl	NH ₂	i-Pr	2,3,6-Trimethyl-pyridin-4-yl
319		Cl	NH ₂	i-Pr	2,3,6-Trimethoxy-pyridin-4-yl
320		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-bromopyridin-4-yl
321		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-chloropyridin-4-yl
322		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-methoxy-1-oxy-pyridin-4-yl
323		Cl	NH ₂	i-Pr	2,6-Dimethyl-1-oxy-pyridin-4-yl
324		Cl	NH ₂	i-Pr	2,3,6-Trimethyl-1-oxy-pyridin-4-yl
325		Cl	NH ₂	i-Pr	2,3,6-Trimethoxy-1-oxy-pyridin-4-yl
326		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-bromo1-oxy-pyridin-4-yl
327		Cl	NH ₂	i-Pr	2,6-Dimethyl-3-chloro1-oxy-pyridin-4-yl
328		Cl	NH ₂	i-Pr	4,6-Dimethyl-5-iodopyridin-3-yl
329		Cl	NH ₂	i-Pr	3,5-Dimethyl-4-aminopyridin-2-yl
330	3	Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
331		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
332		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-bromopyridin-2-yl
333		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
334		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-chloropyridin-2-yl
335		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
336		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-iodopyridin-2-yl
337		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
338		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
339		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
340		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethyl-pyridin-2-yl
341		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
342		Cl	NH ₂	(CH ₂) ₃ OH	4,5,6-Trimethoxypyridin-2-yl
343		Cl	NH ₂	(CH ₂) ₃ OH	4,5,6-Trimethoxy-1-oxypyridin-2-yl
344		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethoxy-pyridin-2-yl
345		Cl	NH ₂	(CH ₂) ₃ OH	3,4,5-Trimethoxy-1-oxypyridin-2-yl
346		Cl	NH ₂	(CH ₂) ₃ OH	4,5,6-Trimethyl-pyridin-2-yl
347		Cl	NH ₂	(CH ₂) ₃ OH	4,5,6-Trimethyl-1-oxypyridin-2-yl
348		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-methoxy-pyridin-2-yl
349		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
350		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
351		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-bromopyridin-3-yl
352		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-chloropyridin-3-yl
353		Cl	NH ₂	(CH ₂) ₃ OH	5,6-Dimethyl-4-bromopyridin-3-yl
354		Cl	NH ₂	(CH ₂) ₃ OH	5,6-Dimethyl-4-chloropyridin-3-yl
355		Cl	NH ₂	(CH ₂) ₃ OH	2,6-Dimethyl-3-methoxypyridin-4-yl
356		Cl	NH ₂	(CH ₂) ₃ OH	2,6-Dimethyl-pyridin-4-yl
357		Cl	NH ₂	(CH ₂) ₃ OH	2,3,6-Trimethyl-pyridin-4-yl
358		Cl	NH ₂	(CH ₂) ₃ OH	2,3,6-Trimethoxy-pyridin-4-yl
359		Cl	NH ₂	(CH ₂) ₃ OH	2,6-Dimethyl-3-bromopyridin-4-yl
360		Cl	NH ₂	(CH ₂) ₃ OH	2,6-Dimethyl-3-chloropyridin-4-yl
361		Cl	NH ₂	(CH ₂) ₃ OH	4,6-Dimethyl-5-iodopyridin-3-yl
362		Cl	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-aminopyridin-2-yl
363	1	Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
364		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl















No.	Ex	R ¹	R ²	R ³	R ⁵
365		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-bromopyridin-2-yl
366		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
367		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-chloropyridin-2-yl
368		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
369		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
370		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
371		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
372		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
373		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethyl-pyridin-2-yl
374		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
375		Cl	NH ₂	CH ₂ OH	4,5,6-Trimethoxypyridin-2-yl
376		Cl	NH ₂	CH ₂ OH	4,5,6-Trimethoxy-1-oxypyridin-2-yl
377		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethoxy-pyridin-2-yl
378		Cl	NH ₂	CH ₂ OH	3,4,5-Trimethoxy-1-oxypyridin-2-yl
379		Cl	NH ₂	CH ₂ OH	4,5,6-Trimethyl-pyridin-2-yl
380		Cl	NH ₂	CH ₂ OH	4,5,6-Trimethyl-1-oxypyridin-2-yl
381		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-methoxy-pyridin-2-yl
382		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
383		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
384		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-bromopyridin-3-yl
385		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-chloropyridin-3-yl
386		Cl	NH ₂	CH ₂ OH	5,6-Dimethyl-4-bromopyridin-3-yl
387		Cl	NH ₂	CH ₂ OH	5,6-Dimethyl-4-chloropyridin-3-yl
388		Cl	NH ₂	CH ₂ OH	2,6-Dimethyl-3-methoxypyridin-4-yl
389		Cl	NH ₂	CH ₂ OH	2,6-Dimethyl-pyridin-4-yl
390		Cl	NH ₂	CH ₂ OH	2,3,6-Trimethyl-pyridin-4-yl
391		Cl	NH ₂	CH ₂ OH	2,3,6-Trimethoxy-pyridin-4-yl
392		Cl	NH ₂	CH ₂ OH	2,6-Dimethyl-3-bromopyridin-4-yl
393		Cl	NH ₂	CH ₂ OH	2,6-Dimethyl-3-chloropyridin-4-yl
394		Cl	NH ₂	CH ₂ OH	4,6-Dimethyl-5-iodopyridin-3-yl
395		Cl	NH ₂	CH ₂ OH	3,5-Dimethyl-4-aminopyridin-2-yl
396	6	Cl	NH ₂	2-Py	3,5-Dimethyl-4-methoxypyridin-2-yl
397		Cl	NH ₂	2-Py	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
398		Cl	NH ₂	2-Py	3,5-Dimethyl-4-bromopyridin-2-yl
399		Cl	NH ₂	2-Py	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
400		Cl	NH ₂	2-Py	3,5-Dimethyl-4-chloropyridin-2-yl
401		Cl	NH ₂	2-Py	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
402		Cl	NH ₂	2-Py	3,5-Dimethyl-4-iodopyridin-2-yl
403		Cl	NH ₂	2-Py	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
404		Cl	NH ₂	2-Py	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
405		Cl	NH ₂	2-Py	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
406		Cl	NH ₂	2-Py	3,4,5-Trimethyl-pyridin-2-yl
407		Cl	NH ₂	2-Py	3,4,5-Trimethyl-1-oxypyridin-2-yl
408		Cl	NH ₂	2-Py	4,5,6-Trimethoxypyridin-2-yl
409		Cl	NH ₂	2-Py	4,5,6-Trimethoxy-1-oxypyridin-2-yl
410		Cl	NH ₂	2-Py	3,4,5-Trimethoxy-pyridin-2-yl
411		Cl	NH ₂	2-Py	3,4,5-Trimethoxy-1-oxypyridin-2-yl
412		Cl	NH ₂	2-Py	4,5,6-Trimethyl-pyridin-2-yl
413		Cl	NH ₂	2-Py	4,5,6-Trimethyl-1-oxypyridin-2-yl
414		Cl	NH ₂	2-Py	4,6-Dimethyl-5-methoxy-pyridin-2-yl
415		Cl	NH ₂	2-Py	4,6-Dimethyl-5-methoxypyridin-3-yl
416		Cl	NH ₂	2-Py	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
417		Cl	NH ₂	2-Py	4,6-Dimethyl-5-bromopyridin-3-yl
418		Cl	NH ₂	2-Py	4,6-Dimethyl-5-chloropyridin-3-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
419		Cl	NH ₂	2-Py	5,6-Dimethyl-4-bromopyridin-3-yl
420		Cl	NH ₂	2-Py	5,6-Dimethyl-4-chloropyridin-3-yl
421		Cl	NH ₂	2-Py	2,6-Dimethyl-3-methoxypyridin-4-yl
422		Cl	NH ₂	2-Py	2,6-Dimethyl-pyridin-4-yl
423		Cl	NH ₂	2-Py	2,3,6-Trimethyl-pyridin-4-yl
424		Cl	NH ₂	2-Py	2,3,6-Trimethoxy-pyridin-4-yl
425		Cl	NH ₂	2-Py	2,6-Dimethyl-3-bromopyridin-4-yl
426		Cl	NH ₂	2-Py	2,6-Dimethyl-3-chloropyridin-4-yl
427		Cl	NH ₂	2-Py	4,6-Dimethyl-5-iodopyridin-3-yl
428		Cl	NH ₂	2-Py	3,5-Dimethyl-4-aminopyridin-2-yl
429		Cl	NH ₂	Ph	3,5-Dimethyl-4-methoxypyridin-2-yl
430		Cl	NH ₂	Ph	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
431		Cl	NH ₂	Ph	3,5-Dimethyl-4-bromopyridin-2-yl
432		Cl	NH ₂	Ph	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
433		Cl	NH ₂	Ph	3,5-Dimethyl-4-chloropyridin-2-yl
434		Cl	NH ₂	Ph	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
435		Cl	NH ₂	Ph	3,5-Dimethyl-4-iodopyridin-2-yl
436		Cl	NH ₂	Ph	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
437		Cl	NH ₂	Ph	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
438		Cl	NH ₂	Ph	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
439		Cl	NH ₂	Ph	3,4,5-Trimethyl-pyridin-2-yl
440		Cl	NH ₂	Ph	3,4,5-Trimethyl-1-oxypyridin-2-yl
441		Cl	NH ₂	Ph	4,5,6-Trimethoxypyridin-2-yl
442		Cl	NH ₂	Ph	4,5,6-Trimethoxy-1-oxypyridin-2-yl
443		Cl	NH ₂	Ph	3,4,5-Trimethoxy-pyridin-2-yl
444		Cl	NH ₂	Ph	3,4,5-Trimethoxy-1-oxypyridin-2-yl
445		Cl	NH ₂	Ph	4,5,6-Trimethyl-pyridin-2-yl
446		Cl	NH ₂	Ph	4,5,6-Trimethyl-1-oxypyridin-2-yl
447		Cl	NH ₂	Ph	4,6-Dimethyl-5-methoxy-pyridin-2-yl
448		Cl	NH ₂	Ph	4,6-Dimethyl-5-methoxypyridin-3-yl
449		Cl	NH ₂	Ph	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
450		Cl	NH ₂	Ph	4,6-Dimethyl-5-bromopyridin-3-yl
451		Cl	NH ₂	Ph	4,6-Dimethyl-5-chloropyridin-3-yl
452		Cl	NH ₂	Ph	5,6-Dimethyl-4-bromopyridin-3-yl
453		Cl	NH ₂	Ph	5,6-Dimethyl-4-chloropyridin-3-yl
454		Cl	NH ₂	Ph	2,6-Dimethyl-3-methoxypyridin-4-yl
455		Cl	NH ₂	Ph	2,6-Dimethyl-pyridin-4-yl
456		Cl	NH ₂	Ph	2,3,6-Trimethyl-pyridin-4-yl
457		Cl	NH ₂	Ph	2,3,6-Trimethoxy-pyridin-4-yl
458		Cl	NH ₂	Ph	2,6-Dimethyl-3-bromopyridin-4-yl
459		Cl	NH ₂	Ph	2,6-Dimethyl-3-chloropyridin-4-yl
460		Cl	NH ₂	Ph	4,6-Dimethyl-5-iodopyridin-3-yl
461		Cl	NH ₂	Ph	3,5-Dimethyl-4-aminopyridin-2-yl
462		Cl	NH ₂	3-Py	3,5-Dimethyl-4-methoxypyridin-2-yl
463		Cl	NH ₂	3-Py	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
464		Cl	NH ₂	3-Py	3,5-Dimethyl-4-bromopyridin-2-yl
465		Cl	NH ₂	3-Py	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
466		Cl	NH ₂	3-Py	3,5-Dimethyl-4-chloropyridin-2-yl
467		Cl	NH ₂	3-Py	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
468		Cl	NH ₂	3-Py	3,5-Dimethyl-4-iodopyridin-2-yl
469		Cl	NH ₂	3-Py	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
470		Cl	NH ₂	3-Py	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
471		Cl	NH ₂	3-Py	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
472		Cl	NH ₂	3-Py	3,4,5-Trimethyl-pyridin-2-yl





















No.	Ex	R ¹	R ²	R ³	R ⁵
473		Cl	NH ₂	3-Py	3,4,5-Trimethyl-1-oxypyridin-2-yl
474		Cl	NH ₂	3-Py	4,5,6-Trimethoxypyridin-2-yl
475		Cl	NH ₂	3-Py	4,5,6-Trimethoxy-1-oxypyridin-2-yl
476		Cl	NH ₂	3-Py	3,4,5-Trimethoxy-pyridin-2-yl
477		Cl	NH ₂	3-Py	3,4,5-Trimethoxy-1-oxypyridin-2-yl
478		Cl	NH ₂	3-Py	4,5,6-Trimethyl-pyridin-2-yl
479		Cl	NH ₂	3-Py	4,5,6-Trimethyl-1-oxypyridin-2-yl
480		Cl	NH ₂	3-Py	4,6-Dimethyl-5-methoxy-pyridin-2-yl
481		Cl	NH ₂	3-Py	4,6-Dimethyl-5-methoxypyridin-3-yl
482		Cl	NH ₂	3-Py	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
483		Cl	NH ₂	3-Py	4,6-Dimethyl-5-bromopyridin-3-yl
484		Cl	NH ₂	3-Py	4,6-Dimethyl-5-chloropyridin-3-yl
485		Cl	NH ₂	3-Py	5,6-Dimethyl-4-bromopyridin-3-yl
486		Cl	NH ₂	3-Py	5,6-Dimethyl-4-chloropyridin-3-yl
487		Cl	NH ₂	3-Py	2,6-Dimethyl-3-methoxypyridin-4-yl
488		Cl	NH ₂	3-Py	2,6-Dimethyl-pyridin-4-yl
489		Cl	NH ₂	3-Py	2,3,6-Trimethyl-pyridin-4-yl
490		Cl	NH ₂	3-Py	2,3,6-Trimethoxy-pyridin-4-yl
491		Cl	NH ₂	3-Py	2,6-Dimethyl-3-bromopyridin-4-yl
492		Cl	NH ₂	3-Py	2,6-Dimethyl-3-chloropyridin-4-yl
493		Cl	NH ₂	3-Py	4,6-Dimethyl-5-iodopyridin-3-yl
494		Cl	NH ₂	3-Py	3,5-Dimethyl-4-aminopyridin-2-yl
495		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
496		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
497		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-bromopyridin-2-yl
498		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
499		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-chloropyridin-2-yl
500		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
501		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-iodopyridin-2-yl
502		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
503		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
504		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
505		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethyl-pyridin-2-yl
506		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethyl-1-oxypyridin-2-yl
507		Cl	NH ₂	CH ₂ -NMe ₂	4,5,6-Trimethoxypyridin-2-yl
508		Cl	NH ₂	CH ₂ -NMe ₂	4,5,6-Trimethoxy-1-oxypyridin-2-yl
509		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethoxy-pyridin-2-yl
510		Cl	NH ₂	CH ₂ -NMe ₂	3,4,5-Trimethoxy-1-oxypyridin-2-yl
511		Cl	NH ₂	CH ₂ -NMe ₂	4,5,6-Trimethyl-pyridin-2-yl
512		Cl	NH ₂	CH ₂ -NMe ₂	4,5,6-Trimethyl-1-oxypyridin-2-yl
513		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-methoxy-pyridin-2-yl
514		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-methoxypyridin-3-yl
515		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
516		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-bromopyridin-3-yl
517		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-chloropyridin-3-yl
518		Cl	NH ₂	CH ₂ -NMe ₂	5,6-Dimethyl-4-bromopyridin-3-yl
519		Cl	NH ₂	CH ₂ -NMe ₂	5,6-Dimethyl-4-chloropyridin-3-yl
520		Cl	NH ₂	CH ₂ -NMe ₂	2,6-Dimethyl-3-methoxypyridin-4-yl
521		Cl	NH ₂	CH ₂ -NMe ₂	2,6-Dimethyl-pyridin-4-yl
522		Cl	NH ₂	CH ₂ -NMe ₂	2,3,6-Trimethyl-pyridin-4-yl
523		Cl	NH ₂	CH ₂ -NMe ₂	2,3,6-Trimethoxy-pyridin-4-yl
524		Cl	NH ₂	CH ₂ -NMe ₂	2,6-Dimethyl-3-bromopyridin-4-yl
525		Cl	NH ₂	CH ₂ -NMe ₂	2,6-Dimethyl-3-chloropyridin-4-yl
526		Cl	NH ₂	CH ₂ -NMe ₂	4,6-Dimethyl-5-iodopyridin-3-yl



No.	Ex	R ¹	R ²	R ³	R ⁵
527		Cl	NH ₂	CH ₂ -NMe ₂	3,5-Dimethyl-4-aminopyridin-2-yl
528		Cl	NH ₂	H	3,5-Dimethyl-4-methoxypyridin-2-yl
529		Cl	NH ₂	H	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
530		Cl	NH ₂	H	3,5-Dimethyl-4-bromopyridin-2-yl
531		Cl	NH ₂	H	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
532		Cl	NH ₂	H	3,5-Dimethyl-4-chloropyridin-2-yl
533		Cl	NH ₂	H	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
534		Cl	NH ₂	H	3,5-Dimethyl-4-iodopyridin-2-yl
535		Cl	NH ₂	H	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
536		Cl	NH ₂	H	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
537		Cl	NH ₂	H	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
538		Cl	NH ₂	H	3,4,5-Trimethyl-pyridin-2-yl
539		Cl	NH ₂	H	3,4,5-Trimethyl-1-oxypyridin-2-yl
540		Cl	NH ₂	H	4,5,6-Trimethoxypyridin-2-yl
541		Cl	NH ₂	H	4,5,6-Trimethoxy-1-oxypyridin-2-yl
542		Cl	NH ₂	H	3,4,5-Trimethoxy-pyridin-2-yl
543		Cl	NH ₂	H	3,4,5-Trimethoxy-1-oxypyridin-2-yl
544		Cl	NH ₂	H	4,5,6-Trimethyl-pyridin-2-yl
545		Cl	NH ₂	H	4,5,6-Trimethyl-1-oxypyridin-2-yl
546		Cl	NH ₂	H	4,6-Dimethyl-5-methoxy-pyridin-2-yl
547		Cl	NH ₂	H	4,6-Dimethyl-5-methoxypyridin-3-yl
548		Cl	NH ₂	H	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
549		Cl	NH ₂	H	4,6-Dimethyl-5-bromopyridin-3-yl
550		Cl	NH ₂	H	4,6-Dimethyl-5-chloropyridin-3-yl
551		Cl	NH ₂	H	5,6-Dimethyl-4-bromopyridin-3-yl
552		Cl	NH ₂	H	5,6-Dimethyl-4-chloropyridin-3-yl
553		Cl	NH ₂	H	2,6-Dimethyl-3-methoxypyridin-4-yl
554		Cl	NH ₂	H	2,6-Dimethyl-pyridin-4-yl
555		Cl	NH ₂	H	2,3,6-Trimethyl-pyridin-4-yl
556		Cl	NH ₂	H	2,3,6-Trimethoxy-pyridin-4-yl
557		Cl	NH ₂	H	2,6-Dimethyl-3-bromopyridin-4-yl
558		Cl	NH ₂	H	2,6-Dimethyl-3-chloropyridin-4-yl
559		Cl	NH ₂	H	3,5-Dimethyl-4-aminopyridin-2-yl
560		Cl	NH ₂	Me	3,5-Dimethyl-4-methoxypyridin-2-yl
561		Cl	NH ₂	Me	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
562		Cl	NH ₂	Me	3,5-Dimethyl-4-bromopyridin-2-yl
563		Cl	NH ₂	Me	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
564		Cl	NH ₂	Me	3,5-Dimethyl-4-chloropyridin-2-yl
565		Cl	NH ₂	Me	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
566		Cl	NH ₂	Me	3,5-Dimethyl-4-iodopyridin-2-yl
567		Cl	NH ₂	Me	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
568		Cl	NH ₂	Me	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
569		Cl	NH ₂	Me	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
570		Cl	NH ₂	Me	3,4,5-Trimethyl-pyridin-2-yl
571		Cl	NH ₂	Me	3,4,5-Trimethyl-1-oxypyridin-2-yl
572		Cl	NH ₂	Me	4,6-Dimethyl-5-methoxypyridin-3-yl
573		Cl	NH ₂	Me	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
574		Cl	NH ₂	Me	3,5-Dimethyl-4-aminopyridin-2-yl
575		Cl	NH ₂	Et	3,5-Dimethyl-4-methoxypyridin-2-yl
576		Cl	NH ₂	Et	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
577		Cl	NH ₂	Et	3,5-Dimethyl-4-bromopyridin-2-yl
578		Cl	NH ₂	Et	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
579		Cl	NH ₂	Et	3,5-Dimethyl-4-chloropyridin-2-yl
580		Cl	NH ₂	Et	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl













No.	Ex	R ¹	R ²	R ³	R ⁵
581		Cl	NH ₂	Et	3,5-Dimethyl-4-iodopyridin-2-yl
582		Cl	NH ₂	Et	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
583		Cl	NH ₂	Et	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
584		Cl	NH ₂	Et	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
585		Cl	NH ₂	Et	3,4,5-Trimethyl-pyridin-2-yl
586		Cl	NH ₂	Et	3,4,5-Trimethyl-1-oxypyridin-2-yl
587		Cl	NH ₂	Et	4,6-Dimethyl-5-methoxypyridin-3-yl
588		Cl	NH ₂	Et	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
589		Cl	NH ₂	Et	3,5-Dimethyl-4-aminopyridin-2-yl
590		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-methoxypyridin-2-yl
591		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
592		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-bromopyridin-2-yl
593		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
594		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-chloropyridin-2-yl
595		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
596		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-iodopyridin-2-yl
597		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
598		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
599		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
600		Cl	NH ₂	CO ₂ Et	3,4,5-Trimethyl-pyridin-2-yl
601		Cl	NH ₂	CO ₂ Et	3,4,5-Trimethyl-1-oxypyridin-2-yl
602		Cl	NH ₂	CO ₂ Et	4,6-Dimethyl-5-methoxypyridin-3-yl
603		Cl	NH ₂	CO ₂ Et	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
604		Cl	NH ₂	CO ₂ Et	3,5-Dimethyl-4-aminopyridin-2-yl
605		Cl	NH ₂	CN	3,5-Dimethyl-4-methoxypyridin-2-yl
606		Cl	NH ₂	CN	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
607		Cl	NH ₂	CN	3,5-Dimethyl-4-bromopyridin-2-yl
608		Cl	NH ₂	CN	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
609		Cl	NH ₂	CN	3,5-Dimethyl-4-chloropyridin-2-yl
610		Cl	NH ₂	CN	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
611		Cl	NH ₂	CN	3,5-Dimethyl-4-iodopyridin-2-yl
612		Cl	NH ₂	CN	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
613		Cl	NH ₂	CN	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
614		Cl	NH ₂	CN	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
615		Cl	NH ₂	CN	3,4,5-Trimethyl-pyridin-2-yl
616		Cl	NH ₂	CN	3,4,5-Trimethyl-1-oxypyridin-2-yl
617		Cl	NH ₂	CN	4,6-Dimethyl-5-methoxypyridin-3-yl
618		Cl	NH ₂	CN	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
619		Cl	NH ₂	CN	3,5-Dimethyl-4-aminopyridin-2-yl
620		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
621		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
622		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-bromopyridin-2-yl
623		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
624		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-chloropyridin-2-yl
625		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
626		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
627		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
628		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
629		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
630		Cl	NH ₂	CMe ₂ OH	3,4,5-Trimethyl-pyridin-2-yl
631		Cl	NH ₂	CMe ₂ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
632		Cl	NH ₂	CMe ₂ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
633		Cl	NH ₂	CMe ₂ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
634		Cl	NH ₂	CMe ₂ OH	3,5-Dimethyl-4-aminopyridin-2-yl















No.	Ex	R ¹	R ²	R ³	R ⁵
635		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
636		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
637		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-bromopyridin-2-yl
638		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
639		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-chloropyridin-2-yl
640		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
641		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
642		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
643		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
644		Cl	NH ₂	CH ₂ CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
645		Cl	NH ₂	CH ₂ CMe ₂ OH	3,4,5-Trimethyl-pyridin-2-yl
646		Cl	NH ₂	CH ₂ CMe ₂ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
647		Cl	NH ₂	CH ₂ CMe ₂ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
648		Cl	NH ₂	CH ₂ CMe ₂ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
649		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
650		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
651		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-bromopyridin-2-yl
652		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
653		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-chloropyridin-2-yl
654		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
655		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
656		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
657		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
658		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
659		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,4,5-Trimethyl-pyridin-2-yl
660		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	3,4,5-Trimethyl-1-oxypyridin-2-yl
661		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	4,6-Dimethyl-5-methoxypyridin-3-yl
662		Cl	NH ₂	CH ₂ CH ₂ CMe ₂ OH	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
663		Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
664		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
665		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
666		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
667		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
668		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
669		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
670		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
671		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
672		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
673		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
674		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl
675		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
676		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl















No.	Ex	R ¹	R ²	R ³	R ⁵
677		Cl	NH ₂		3,5-Dimethyl-4-aminopyridin-2-yl
678	18	Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
679		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
680		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
681		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
682		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
683		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
684		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
685		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
686		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
687		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
688		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
689		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl
690		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
691		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
692	19	Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
693		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
694		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
695		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
696		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
697		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
698		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
699		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
700		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
701		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
702		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
703		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl
704		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
705		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
706		Cl	NH ₂		3,4,5-Trimethoxyphenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
707		Cl	NH ₂		2-Chloro-3,4,5-trimethoxyphenyl
708		Cl	NH ₂		2-Bromo-3,4,5-trimethoxyphenyl
709		Cl	NH ₂		3,5-Dimethyl-4-methoxyphenyl
710		Cl	NH ₂		2-Chloro-3,5-Dimethyl-4-methoxyphenyl
711		Cl	NH ₂		2-Bromo-3,5-Dimethyl-4-methoxyphenyl
712		Cl	NH ₂		3,4,5-Trimethoxyphenyl
713		Cl	NH ₂		2-Chloro-3,4,5-trimethoxyphenyl
714		Cl	NH ₂		2-Bromo-3,4,5-trimethoxyphenyl
715		Cl	NH ₂		3,5-Dimethyl-4-methoxyphenyl
716		Cl	NH ₂		2-Chloro-3,5-Dimethyl-4-methoxyphenyl
717		Cl	NH ₂		2-Bromo-3,5-Dimethyl-4-methoxyphenyl
718		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
719		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
720		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-bromopyridin-2-yl
721		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
722		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-chloropyridin-2-yl
723		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
724		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-iodopyridin-2-yl
725		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
726		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
727		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
728		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,4,5-Trimethyl-pyridin-2-yl
729		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	3,4,5-Trimethyl-1-oxypyridin-2-yl
730		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	4,6-Dimethyl-5-methoxypyridin-3-yl
731		Cl	NH ₂	(CH ₂) ₂ C(O)NMe ₂	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
732	11	Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
733		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
734		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
735		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
736		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
737		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
738		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
739		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
740		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
741		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
742		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
743		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl


















No.	Ex	R ¹	R ²	R ³	R ⁵
744		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
745		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
746	12	Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
747		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
748		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-bromopyridin-2-yl
749		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
750		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-chloropyridin-2-yl
751		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
752		Cl	NH ₂	(CH ₂) ₂ CO NH ₂	3,5-Dimethyl-4-iodopyridin-2-yl
753		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-methoxypyridin-2-yl
754		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
755		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	6-Bromo-3,5-dimethyl-4-methoxypyridin-2-yl
756		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	6-Chloro-3,5-dimethyl-4-methoxypyridin-2-yl
757		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-bromopyridin-2-yl
758		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
759		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-chloropyridin-2-yl
760		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
761		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-iodopyridin-2-yl
762		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
763		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
764		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
765		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,4,5-Trimethyl-pyridin-2-yl
766		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,4,5-Trimethyl-1-oxypyridin-2-yl
767		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,5,6-Trimethoxypyridin-2-yl
768		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,5,6-Trimethoxy-1-oxypyridin-2-yl
769		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3-Bromo-4,5,6-trimethoxypyridin-2-yl
770		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3-Chloro-4,5,6-trimethoxypyridin-2-yl
771		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,4,5-Trimethoxy-pyridin-2-yl
772		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3,4,5-Trimethoxy-1-oxypyridin-2-yl
773		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3-Bromo-3,4,5-trimethoxy-pyridin-2-yl
774		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	3-Chloro-3,4,5-trimethoxy-pyridin-2-yl
775		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,5,6-Trimethyl-pyridin-2-yl
776		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,5,6-Trimethyl-1-oxypyridin-2-yl
777		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-methoxy-pyridin-2-yl
778		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-methoxypyridin-3-yl
779		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
780		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-bromopyridin-3-yl
781		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-chloropyridin-3-yl
782		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	5,6-Dimethyl-4-bromopyridin-3-yl
783		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	5,6-Dimethyl-4-chloropyridin-3-yl
784		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-methoxypyridin-4-yl
785		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-pyridin-4-yl
786		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,3,6-Trimethyl-pyridin-4-yl
787		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,3,6-Trimethoxy-pyridin-4-yl
788		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-bromopyridin-4-yl
789		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-chloropyridin-4-yl
790		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-methoxy-1-oxy-pyridin-4-yl
791		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-1-oxy-pyridin-4-yl
792		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,3,6-Trimethyl-1-oxy-pyridin-4-yl
793		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,3,6-Trimethoxy-1-oxy-pyridin-4-yl
794		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-bromo1-oxy-pyridin-4-yl
795		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	2,6-Dimethyl-3-chloro1-oxy-pyridin-4-yl
796		Cl	NH ₂	(CH ₂) ₂ CO NHBOC	4,6-Dimethyl-5-iodopyridin-3-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
797		Cl	NH ₂	(CH ₂) ₂ CONHBOC	3,5-Dimethyl-4-aminopyridin-2-yl
798		Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
799		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
800		Cl	NH ₂		6-Bromo-3,5-dimethyl-4-methoxypyridin-2-yl
801		Cl	NH ₂		6-Chloro-3,5-dimethyl-4-methoxypyridin-2-yl
802		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
803		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
804		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
805		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
806		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
807		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
808		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
809		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
810		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
811		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl


No.	Ex	R ¹	R ²	R ³	R ⁵
812		Cl	NH ₂		4,5,6-Trimethoxypyridin-2-yl
813		Cl	NH ₂		4,5,6-Trimethoxy-1-oxypyridin-2-yl
814		Cl	NH ₂		3-Bromo-4,5,6-trimethoxypyridin-2-yl
815		Cl	NH ₂		3-Chloro-4,5,6-trimethoxypyridin-2-yl
816		Cl	NH ₂		3,4,5-Trimethoxy-pyridin-2-yl
817		Cl	NH ₂		3,4,5-Trimethoxy-1-oxypyridin-2-yl
818		Cl	NH ₂		3-Bromo-3,4,5-trimethoxy-pyridin-2-yl
819		Cl	NH ₂		3-Chloro-3,4,5-trimethoxy-pyridin-2-yl
820		Cl	NH ₂		4,5,6-Trimethyl-pyridin-2-yl
821		Cl	NH ₂		4,5,6-Trimethyl-1-oxypyridin-2-yl
822		Cl	NH ₂		4,6-Dimethyl-5-methoxy-pyridin-2-yl
823		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
824		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
825		Cl	NH ₂		4,6-Dimethyl-5-bromopyridin-3-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
826		Cl	NH ₂		4,6-Dimethyl-5-chloropyridin-3-yl
827		Cl	NH ₂		5,6-Dimethyl-4-bromopyridin-3-yl
828		Cl	NH ₂		5,6-Dimethyl-4-chloropyridin-3-yl
829		Cl	NH ₂		2,6-Dimethyl-3-methoxypyridin-4-yl
830		Cl	NH ₂		2,6-Dimethyl-pyridin-4-yl
831		Cl	NH ₂		2,3,6-Trimethyl-pyridin-4-yl
832		Cl	NH ₂		2,3,6-Trimethoxy-pyridin-4-yl
833		Cl	NH ₂		2,6-Dimethyl-3-bromopyridin-4-yl
834		Cl	NH ₂		2,6-Dimethyl-3-chloropyridin-4-yl
835		Cl	NH ₂		2,6-Dimethyl-3-methoxy-1-oxypyridin-4-yl
836		Cl	NH ₂		2,6-Dimethyl-1-oxy-pyridin-4-yl
837		Cl	NH ₂		2,3,6-Trimethyl-1-oxypyridin-4-yl
838		Cl	NH ₂		2,3,6-Trimethoxy-1-oxypyridin-4-yl
839		Cl	NH ₂		2,6-Dimethyl-3-bromo 1-oxypyridin-4-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
840		Cl	NH ₂		2,6-Dimethyl-3-chloro-1-oxypyridin-4-yl
841		Cl	NH ₂		4,6-Dimethyl-5-iodopyridin-3-yl
842		Cl	NH ₂		3,5-Dimethyl-4-aminopyridin-2-yl
843		Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
844		Cl	NH ₂		3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
845		Cl	NH ₂		6-Bromo-3,5-dimethyl-4-methoxypyridin-2-yl
846		Cl	NH ₂		6-Chloro-3,5-dimethyl-4-methoxypyridin-2-yl
847		Cl	NH ₂		3,5-Dimethyl-4-bromopyridin-2-yl
848		Cl	NH ₂		3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
849		Cl	NH ₂		3,5-Dimethyl-4-chloropyridin-2-yl
850		Cl	NH ₂		3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
851		Cl	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
852		Cl	NH ₂		3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
853		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
854		Cl	NH ₂		3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl

No.	Ex	R ¹	R ²	R ³	R ⁵
855		Cl	NH ₂		3,4,5-Trimethyl-pyridin-2-yl
856		Cl	NH ₂		3,4,5-Trimethyl-1-oxypyridin-2-yl
857		Cl	NH ₂		4,5,6-Trimethoxypyridin-2-yl
858		Cl	NH ₂		4,5,6-Trimethoxy-1-oxypyridin-2-yl
859		Cl	NH ₂		3-Bromo-4,5,6-trimethoxypyridin-2-yl
860		Cl	NH ₂		3-Chloro-4,5,6-trimethoxypyridin-2-yl
861		Cl	NH ₂		4,6-Dimethyl-5-methoxypyridin-3-yl
862		Cl	NH ₂		4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
863		Br	NH ₂	(CH ₂)OH	3,5-Dimethyl-4-methoxypyridin-2-yl
864		Br	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
865		Br	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
866		Br	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
867		Br	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
868		Br	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
869		Br	NH ₂	(CH ₂)OH	3,5-Dimethyl-4-iodopyridin-2-yl
870		Br	NH ₂	(CH ₂) ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
871		Br	NH ₂	(CH ₂) ₃ OH	3,5-Dimethyl-4-iodopyridin-2-yl
872		Br	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
873		Br	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
874		Br	NH ₂		3,5-Dimethyl-4-iodopyridin-2-yl
875		Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
876		Cl	NH ₂		2-Chloro-4,5-dimethoxylphenyl
877		Cl	NH ₂		2-Nitro-4,5-dimethoxylphenyl
878		Cl	NH ₂		3,4-Dichlorophenyl

No.	Ex	R ¹	R ²	R ³	R ⁵
879		Cl	NH ₂	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethoxyphenyl
880		Cl	NH ₂	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	2,5-Dimethoxyphenyl
881		Cl	NH ₂	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethoxyphenyl
882		Cl	NH ₂	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3-Methoxyphenyl
883		Cl	NH ₂	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	4-Methoxyphenyl
884		Cl	<chem>CC(C)C(=O)N</chem>	(CH ₂) ₂ OH	3,5-Dimethyl-4-iodopyridin-2-yl
885		Cl	<chem>CC(C)C(=O)N</chem>	(CH ₂) ₃ OH	3,5-Dimethyl-4-iodopyridin-2-yl
886		Cl	<chem>CC(C)C(=O)N</chem>	-CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethyl-4-iodopyridin-2-yl
887		Cl	<chem>CC(C)C(=O)N</chem>	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethyl-4-iodopyridin-2-yl
888		Cl	<chem>CC(C)C(=O)N</chem>	(CH ₂)OH	3,5-Dimethyl-4-methoxypyridin-2-yl
889		Cl	<chem>CC(C)C(=O)N</chem>	(CH ₂) ₂ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
890		Cl	<chem>CC(C)C(=O)N</chem>	(CH ₂) ₃ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
891		Cl	<chem>CC(C)C(=O)N</chem>	-CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethyl-4-methoxypyridin-2-yl
892		Cl	<chem>CC(C)C(=O)N</chem>	-CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethyl-4-methoxypyridin-2-yl
893		Cl	<chem>CC(C)C(=O)N</chem>	-CH ₂ CH ₂ CH ₂ -N <chem>C1CCOCC1</chem>	3,5-Dimethyl-4-methoxypyridin-2-yl
894	4	Cl	NH ₂	CH ₂ N(i-Bu) ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
895	5	Cl	NH ₂	CH ₂ N(i-Bu) ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
896	7	Cl	NH ₂	(CH ₂) ₄ OH	3,5-Dimethyl-4-methoxypyridin-2-yl
897	8	Cl	NH ₂	Si(CH ₃) ₃	3,5-Dimethyl-4-methoxypyridin-2-yl
898	9	Cl	NH ₂	(CH ₂) ₂ CO ₂ H	3,5-Dimethyl-4-methoxypyridin-2-yl
899	10	Cl	NH ₂	(CH ₂) ₂ CON(Et) ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
900	13	Cl	NH ₂	CH ₂ NHCOOtBu	3,5-Dimethyl-4-methoxypyridin-2-yl
901	14	Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
902		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-methoxy-1-oxypyridin-2-yl
903		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-bromopyridin-2-yl
904		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-bromo-1-oxypyridin-2-yl
905		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-chloropyridin-2-yl
906		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-chloro-1-oxypyridin-2-yl
907		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-iodopyridin-2-yl
908		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-iodo-1-oxypyridin-2-yl
909		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-thiomethyl-pyridin-2-yl
910		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-thiomethyl-1-oxypyridin-2-yl
911		Cl	NH ₂	CH ₂ NH ₂	3,4,5-Trimethyl-pyridin-2-yl
912		Cl	NH ₂	CH ₂ NH ₂	3,4,5-Trimethyl-1-oxypyridin-2-yl
913		Cl	NH ₂	CH ₂ NH ₂	4,5,6-Trimethoxypyridin-2-yl
914		Cl	NH ₂	CH ₂ NH ₂	4,5,6-Trimethoxy-1-oxypyridin-2-yl
915		Cl	NH ₂	CH ₂ NH ₂	3,4,5-Trimethoxy-pyridin-2-yl
916		Cl	NH ₂	CH ₂ NH ₂	3,4,5-Trimethoxy-1-oxypyridin-2-yl
917		Cl	NH ₂	CH ₂ NH ₂	4,5,6-Trimethyl-pyridin-2-yl
918		Cl	NH ₂	CH ₂ NH ₂	4,5,6-Trimethyl-1-oxypyridin-2-yl
919		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-methoxy-pyridin-2-yl
920		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-methoxypyridin-3-yl
921		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-methoxy-1-oxypyridin-3-yl
922		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-bromopyridin-3-yl


No.	Ex	R ¹	R ²	R ³	R ⁵
923		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-chloropyridin-3-yl
924		Cl	NH ₂	CH ₂ NH ₂	5,6-Dimethyl-4-bromopyridin-3-yl
925		Cl	NH ₂	CH ₂ NH ₂	5,6-Dimethyl-4-chloropyridin-3-yl
926		Cl	NH ₂	CH ₂ NH ₂	2,6-Dimethyl-3-methoxypyridin-4-yl
927		Cl	NH ₂	CH ₂ NH ₂	2,6-Dimethyl-pyridin-4-yl
928		Cl	NH ₂	CH ₂ NH ₂	2,3,6-Trimethyl-pyridin-4-yl
929		Cl	NH ₂	CH ₂ NH ₂	2,3,6-Trimethoxy-pyridin-4-yl
930		Cl	NH ₂	CH ₂ NH ₂	2,6-Dimethyl-3-bromopyridin-4-yl
931		Cl	NH ₂	CH ₂ NH ₂	2,6-Dimethyl-3-chloropyridin-4-yl
932		Cl	NH ₂	CH ₂ NH ₂	4,6-Dimethyl-5-iodopyridin-3-yl
933		Cl	NH ₂	CH ₂ NH ₂	3,5-Dimethyl-4-aminopyridin-2-yl
934	15	Cl	NH ₂	(CH ₂) ₃ NH COOtBu	3,5-Dimethyl-4-methoxypyridin-2-yl
935	16	Cl	NH ₂	(CH ₂) ₃ NH ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
936	17	Cl	NH ₂		3,5-Dimethyl-4-methoxypyridin-2-yl
937	20	Cl	NH ₂	(CH ₂) ₂ NHtBu	3,5-Dimethyl-4-methoxypyridin-2-yl
938	21	Cl	NH ₂	(CH ₂) ₂ NHtBu	3,5-Dimethyl-4-methoxypyridin-2-yl
939	22	Cl	NH ₂	CH ₂ OCO CH ₂ NMe ₂	3,5-Dimethyl-4-methoxypyridin-2-yl
940		Cl	NH ₂	CH ₂ CH(OH)CH ₃	3,5-Dimethyl-4-methoxypyridin-2-yl
941		Cl	NH ₂	CH ₂ C(O)CH ₃	3,5-Dimethyl-4-methoxypyridin-2-yl

wherein each abbreviation has its usual meaning, which would be known to one of skill in the chemical arts, i.e. Me = methyl; Et = ethyl; Pr = propyl; i-Pr = *iso*-propyl; Bu = butyl; i-Bu = *iso*-butyl; tBu = *tert*-butyl; Ph = phenyl; BOC = *tert*-butoxycarbonyl; 2-Py = 2-pyridyl; 3-Py = 3-pyridyl; 4-Py = 4-pyridyl.


- 5 [060] Selected compounds in TABLE 1 are compounds 2, 3, 4, 7, 8, 12, 13, 17, 18, 22, 23, 27, 28, 32, 33, 38, 42, 43, 47, 48, 52, 53, 57, 58, 62, 63, 67, 68, 72, 73, 77, 78, 82, 83, 87, 88, 92, 93, 102, 103, 107, 108, 112, 113, 117, 118, 122, 123, 127, 128, 132, 133, 137, 138, 147, 148, 152, 153, 157, 158, 161, 162, 166, 167, 171, 172, 176, 177, 181, 182, 186, 187, 191, 192, 196, 197, 201, 202, 206, 207, 211, 212, 216, 217, 221, 222, 226, 227, 231, 232, 236, 237, 240, 241, 244,
- 10 246, 248, 265, 266, 285, 289, 291, 293, 310, 311, 330, 331, 332, 334, 349, 350, 363, 364, 365, 367, 369, 382, 383, 396, 397, 398, 400, 402, 415, 416, 429, 430, 431, 433, 435, 448, 449, 462, 463, 464, 466, 468, 481, 482, 495, 496, 497, 499, 501, 514, 515, 528, 529, 530, 532, 534, 547, 548, 560, 561, 652, 566, 587, 588, 590, 591, 592, 594, 596, 602, 603, 620, 621, 622, 624, 626, 632, 633, 635, 636, 637, 639, 641, 661, 662, 663, 664, 665, 667, 669, 675, 676, 678, 679, 680,
- 15 682, 684, 690, 691, 692, 693, 694, 696, 698, 704, 705, 707, 708, 710, 711, 713, 714, 716, 717, 718, 719, 720, 722, 730, 731, 732, 733, 734, 736, 738, 744, 745, 746, 747, 748, 750, 752, 753, 754, 757, 759, 761, 778, 779, 782, 783, 784, 798, 799, 802, 804, 806, 823, 824, 843, 844, 847, 849, 851, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 875, 894, 895, 896, 899, 900, 901, 903, 905, 907, 920, 921, 934, 935, 936, 937, 938, 939, 940, and 941. Further selected compounds
- 20 are 77, 78, 82, 83, 87, 88, 92, 93, 240, 241, 248, 265, 285, 293, 310, 330, 349, 382, 396, 415, 429, 448, 462, 481, 495, 514, 528, 560, 587, 602, 620, 632, 635, 661, 663, 675, 678, 690, 692,

704, 718, 730, 732, 744, 746, 753, 778, 798, 823, 863, 864, 865, 866, 867, 868, 869, 875, 894, 895, 896, 899, 900, 901, 903, 905, 920, 934, 935, 936, 940, and 941.





[061] Compounds of the invention exhibit improved HSP90 inhibiting activities over some pyrrolo[2,3-d]pyrimidine compounds of Formula II:







wherein R^{3'} is not alkynyl, which were disclosed in U.S. Patent Application No. 10/945,851 and PCT Application US04/31248. The pyrrolopyrimidine compounds specified in the EXAMPLE sections of the above patent applications carry either no substituent or a substituted alkyl group. Compound 0, one of the embodiments of the compounds of Formula II, carries no substituents on the C-5 position (ie R^{3'} = H) and it has an IC₅₀ = 98 nM as measured by a Her-2 degradation assay (described in EXAMPLE section below). A simple alkyl or a substituted alkyl at C-5 (ie R^{3'} = alkyl or substituted alkyl) can only bring two-fold additional activity. Conversely, if C-5 is alkynyl substituted, the potency increases approximately 10-20 fold. (See, Series A: compounds A, B, C and D below).



Series A: Non-Alkynes



Series B: Alkynes



PCT/US2006/011846
[062]

For example, the alkynyl substituents at the C-5 position (see Series B: Cpd's E, F, G and H), showed an potency range of 6-17 nM an improvement of at least tenfold over the parent Compound 0. Further, when two compounds having exactly the same structure, except for the presence of a triple vs. single bond, *e.g.* comparing Compound D to Compound E, Compound E which has a propargylic alcohol substituted on C-5 has an IC_{50} = 17 nM, while Compound D which has a propanol substituent on C-5 has an IC_{50} = 38 nM, a two-fold improvement. The difference in potency index between alkynes and non-alkynes measured by a secondary assay is also striking. The secondary assay measures the efficiency of killing tumor cells. For a given compound, the the potency index is defined as:

$$\text{Index} = EC_{50}(\text{compound})/EC_{50}(\text{control})$$

where the control is 17-AAG (17-allyl-17-desmethoxy-geldanamycin). EC_{50} is defined as the amount of compound added to effect a 50% reduction in viable cell number. Selected compounds were tested on cells of MCF7 and BT474 breast tumor cell lines. The assay shows that Compound 0 is over 10 times less active than 17-AAG. The best non-alkyne substituted analogs are 6 to >50 times less potent than 17-AAG. In contrast, the alkyne substituted analogs can be more potent than 17-AAG (see Cpd. H) – a record in the field of HSP90 inhibitors, or nearly as active, with an index between 0.7–6. TABLE 2 summarized the result of the assay.


[063] TABLE 2: Potency Index against Tumor Cells from MCF 7 and BT474 Cell Lines

Compound	Substituent @ C-5	MCF7 Index (μ M)	BT474 Index (μ M)
0	none	0.4	0.3
A	non-alkyne	0.2	1.0
B	non-alkyne	0.08	0.3
C	non-alkyne	0.6	0.3
D	non-alkyne	0.23	1.0
E	alkyne	0.13	0.02
F	alkyne	0.006	0.13
G	alkyne	0.03	0.02
H	alkyne	0.01	0.1

[064] Synthesis of the Compounds of the Invention

The compounds of the present invention may be prepared from 5-halo-pyrrolo[2,3-d]pyrimidines according to Scheme A below. The preparation of the starting material, N-(4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-2-yl)pivalamide, wherein PG is the pivaloyl protecting group and X = I, has been reported in Seela, F. *Synthesis* **2004**, 8, 1203 and references therein.

PCT/US06/11846




Scheme A

[065] Treatment of N-(4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-2-yl)pivalamide with the appropriate alkylating agent in a polar solvent (*e.g.* DMF, DMSO, NMP), in the presence of a base (*e.g.* K₂CO₃, C₂CO₃, NaH, NaOH, *t*-BuOH) affords the N(7)-alkylated adduct. The alkylation is typically run at 20–60 °C for 0.5–24 h.

Cleavage of the NH₂ protecting group gives the 2-amino derivative. If the protecting group is a pivaloyl group, it can be cleaved with ZnCl₂. The depivaloylation is typically run in EtOH, with 1–10 vol% of water, at 50–85 °C, for 2–24 h. Zinc (II) chloride can be substituted with other Lewis acids, such as ZnI₂ or CuCl which, however, may not give yields as high.

Sonogashira coupling of 4-chloro-7-alkyl-5-halo-7H-pyrrolo[2,3-d]pyrimidin-2-amine with an alkyne of Formula HC≡C-R¹ gives the desired alkyne. Typical reaction conditions require Pd(PPh₃)₄/CuI as catalytic system, Et₃N as base, and DCM or DMF as solvent. The reactions are typically run at 20–50 °C for 0.5 to 24 h. However, a variety of alternative catalytic systems/base/conditons can be used (see Liang, B. et. al., *J. Org. Chem.* **2005**, *70*, 391 and references therein).

[066] The R group can be further manipulated if necessary, as illustrated below:



[067] For illustration purposes, the propargylic alcohol can be converted to the mesylate or bromide using methods well known in the art. The mesyl or bromide group can then be displaced by nucleophiles such as amines. Care must be taken to avoid the unwanted nucleophilic substitution of the 4-Cl atom.

[068] The sequence of these steps can be performed in a different order than the one indicated on the Scheme, as for instance (i) alkylation, (ii), Sonogashira coupling, and (iii) deprotection.

PCT/US06/11846

[069] Pharmaceutical Compositions, Medicaments, Dosaging and Modes of Administration

The present invention is also directed to the use of alkynyl pyrrolo[2,3-d]pyrimidine compounds of Formula I and their related analogs, and their polymorphs, solvates, esters, tautomers,

5 diastereomers, enantiomers, pharmaceutically acceptable salts and prodrugs thereof. In some embodiments, the compounds are used for the treatment or prevention of diseases that are HSP90-dependent. In some embodiments, the compounds are used in the manufacture of a medicament. In other embodiments, the compounds are used in the manufacture of a medicament for the therapeutic and/or prophylactic treatment of diseases and conditions that are
10 HSP90-dependent. Examples of such diseases and conditions include disorders such as inflammatory diseases, infections, autoimmune disorders, stroke, ischemia, cardiac disorder, neurological disorders, fibrogenetic disorders, proliferative disorders, tumors, leukemias, chronic lymphocytic leukemia, acquired immunodeficiency syndrome, neoplasms, cancers, carcinomas, metabolic diseases, and malignant disease. The fibrogenetic disorders include but are not limited
15 to scleroderma, polymyositis, systemic lupus, rheumatoid arthritis, liver cirrhosis, keloid formation, interstitial nephritis and pulmonary fibrosis.

[070] The present invention features pharmaceutical compositions and medicaments comprising the compound of Formula I, or a polymorph, solvate, ester, tautomer, enantiomer, diastereomer, pharmaceutically acceptable salt thereof, or prodrug thereof, of any of the
20 preceding aspects and embodiments and one or more pharmaceutical excipients. Those of ordinary skill in the art are familiar with formulation and administration techniques that can be employed with the compounds and methods of the invention, *e.g.*, as discussed in Goodman and Gilman, *The Pharmacological Basis of Therapeutics* (10th edition); Pergamon; and *Remington's, Pharmaceutical Sciences* (20th edition), Mack Publishing Co., Easton, Pa.

[071] The compounds utilized in the methods of the instant invention may be administered either alone or in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according to standard pharmaceutical practice. The
25 compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration. For example, the therapeutic or pharmaceutical compositions of the invention can be administered locally to the area in need of treatment. This may be achieved by, for example, but not limited to, local infusion during surgery, topical application, *e.g.*, cream, ointment, injection, catheter, or implant, said implant made, *e.g.*, out of a porous, non-porous, or gelatinous material, including
30 membranes, such as sialastic membranes, or fibers. The administration can also be by direct injection at the site (or former site) of a tumor or neoplastic or pre-neoplastic tissue. Still further,

~~PCT/US2006/011846~~
the compounds or compositions of the invention can be delivered in a vesicle, *e.g.*, a liposome (see, for example, Langer, *Science* **1990**, 249,1527-1533; Treat *et al.*, *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Bernstein and Fidler, Ed., Liss, N.Y., pp. 353-365, 1989).

5 [072] The compounds and pharmaceutical compositions used in the methods of the present invention can also be delivered in a controlled release system. In one embodiment, a pump may be used (see, Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:201; Buchwald *et al.* *Surgery*, **1980** 88, 507; Saudek *et al.* *N. Engl. J. Med.* **1989**, 321, (574). Additionally, a controlled release system can be placed in proximity of the therapeutic target. (See, Goodson, *Medical*
10 *Applications of Controlled Release*, 1984, Vol. 2, pp. 115-138).

[073] The pharmaceutical compositions used in the methods of the instant invention can also contain the active ingredient in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to
15 any method known in the art for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the
20 manufacture of tablets. These excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, such as microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be un-
25 coated or coated by known techniques to mask the taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropylmethyl-cellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, or cellulose acetate butyrate may be employed as appropriate.

30 [074] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

PCT/US2006/011846
[075]

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

[076] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

[077] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

[078] The compounds and pharmaceutical compositions used in the methods of the instant invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin, or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soybean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening agents, flavoring agents, preservatives and antioxidants.

PCT/US2006/011846
[079]

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

[080] The pharmaceutical compositions may be in the form of a sterile injectable aqueous solution. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

[081] The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution may then be introduced into a water and glycerol mixture and processed to form a microemulsion.

[082] The injectable solutions or microemulsions may be introduced into a patient's blood-stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

[083] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[084] The compounds of the present invention used in the methods of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the inhibitors with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol. For topical use, creams, ointments, jellies, solutions or suspensions, *etc.*, containing a compound or composition of the

~~PCT/US2006/011846~~
invention can be used. As used herein, topical application can include mouth washes and gargles.

[085] The compounds used in the methods of the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

[086] The methods, compounds and compositions of the instant invention may also be used in conjunction with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. For example, the instant compounds may be useful in combination with known anti-cancer and cytotoxic agents. Further, the instant methods and compounds may also be useful in combination with other inhibitors of parts of the signaling pathway that links cell surface growth factor receptors to nuclear signals initiating cellular proliferation.

[087] The methods of the present invention may also be useful with other agents that inhibit angiogenesis and thereby inhibit the growth and invasiveness of tumor cells, including, but not *limited* to VEGF receptor inhibitors, including ribozymes and antisense targeted to VEGF receptors, angiostatin and endostatin. Examples of antineoplastic agents that can be used in combination with the compounds and methods of the present invention include, in general, and as appropriate, alkylating agents, anti-metabolites, epidophyllotoxins, an antineoplastic enzyme, a topoisomerase inhibitor, procarbazine, mitoxantrone, platinum coordination complexes, biological response modifiers and growth inhibitors, hormonal/anti-hormonal therapeutic agents and haematopoietic growth factors. Exemplary classes of antineoplastic include the anthracyclines, vinca drugs, mitomycins, bleomycins, cytotoxic nucleosides, epothilones, discodermolide, pteridines, diynenes and podophyllotoxins. Particularly useful members of those classes include, for example, carminomycin, daunorubicin, aminopterin, methotrexate, methopterin, dichloromethotrexate, mitomycin C, porfiromycin, 5-fluorouracil, 6-mercaptopurine, gemcitabine, cytosine arabinoside, podophyllotoxin or podo-phyllotoxin derivatives such as etoposide, etoposide phosphate or teniposide, melphalan, vinblastine, vincristine, leurosine, vindesine, leurosine, paclitaxel and the like. Other useful antineoplastic agents include estramustine, carboplatin, cyclophosphamide, bleomycin, gemcitabine, ifosamide, melphalan, hexamethyl melamine, thiotepa, cytarabin, idatrexate, trimetrexate, dacarbazine, L-asparaginase, camptothecin, CPT-11, topotecan, ara-C, bicalutamide, flutamide, leuprolide, pyridobenzoindole derivatives, interferons and interleukins.

[088] ~~PCT/US2006/011846~~

When a compound or composition of the invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms. In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer, for example, breast cancer. Administration typically occurs in an amount of between about 0.01 mg/kg of body weight to about 100 mg/kg of body weight per day (administered in single or divided doses), more preferably at least about 0.1 mg/kg of body weight per day. A particular therapeutic dosage can include, *e.g.*, from about 0.01 mg to about 1000 mg of compound, and preferably includes, *e.g.*, from about 1 mg to about 1000 mg. The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.1 mg to 1000 mg, preferably from about 1 mg to 300 mg, more preferably 10 mg to 200 mg, according to the particular application. The amount administered will vary depending on the particular IC_{50} value of the compound used and the judgment of the attending clinician taking into consideration factors such as health, weight, and age. In combinational applications in which the compound is not the sole active ingredient, it may be possible to administer lesser amounts of compound and still have therapeutic or prophylactic effect.

[089] Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component, *e.g.*, an effective amount to achieve the desired purpose. The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small amounts until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. The amount and frequency of administration of the compounds and compositions of the present invention used in the methods of the present invention, and if applicable other chemotherapeutic agents and/or radiation therapy, will be regulated according to the judgment of the attending clinician (physician) considering such factors as age, condition and size of the patient as well as severity of the disease being treated.

[090] The chemotherapeutic agent and/or radiation therapy can be administered according to therapeutic protocols well known in the art. It will be apparent to those skilled in the art that the administration of the chemotherapeutic agent and/or radiation therapy can be varied depending on the disease being treated and the known effects of the chemotherapeutic agent and/or

~~PCT/US2006/011846~~
radiation therapy on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic protocols (*e.g.*, dosage amounts and times of administration) can be varied in view of the observed effects of the administered therapeutic agents (*i.e.*, antineoplastic agent or radiation) on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

[091] Also, in general, the compounds of the invention need not be administered in the same pharmaceutical composition as a chemotherapeutic agent, and may, because of different physical and chemical characteristics, be administered by a different route. For example, the compounds/compositions may be administered orally to generate and maintain good blood levels thereof, while the chemotherapeutic agent may be administered intravenously. The determination of the mode of administration and the advisability of administration, where possible, in the same pharmaceutical composition, is well within the knowledge of the skilled clinician. The initial administration can be made according to established protocols known in the art, and then, based upon the observed effects, the dosage, modes of administration and times of administration can be modified by the skilled clinician.

[092] The particular choice of compound (and where appropriate, chemotherapeutic agent and/or radiation) will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

[093] The compounds/compositions of the invention (and where appropriate chemotherapeutic agent and/or radiation) may be administered concurrently (*e.g.*, simultaneously, essentially simultaneously or within the same treatment protocol) or sequentially, depending upon the nature of the proliferative disease, the condition of the patient, and the actual choice of chemotherapeutic agent and/or radiation to be administered in conjunction (*i.e.*, within a single treatment protocol) with the compound/composition. In combinational applications and uses, the compound/composition and the chemotherapeutic agent and/or radiation need not be administered simultaneously or essentially simultaneously, and the initial order of administration of the compound/composition, and the chemotherapeutic agent and/or radiation, may not be important. Thus, the compounds/compositions of the invention may be administered first followed by the administration of the chemotherapeutic agent and/or radiation; or the chemotherapeutic agent and/or radiation may be administered first followed by the administration of the compounds/compositions of the invention. This alternate administration may be repeated during a single treatment protocol. The determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is well within the knowledge of the skilled physician after evaluation of the disease being treated and the condition of the patient. For example, the chemotherapeutic agent

~~PCT/US2006/011846~~
and/or radiation may be administered first, especially if it is a cytotoxic agent, and then the treatment continued with the administration of the compounds/compositions of the invention followed, where determined advantageous, by the administration of the chemotherapeutic agent and/or radiation, and so on until the treatment protocol is complete.

5 [094] Thus, in accordance with experience and knowledge, the practicing physician can modify each protocol for the administration of a compound/composition for treatment according to the individual patient's needs, as the treatment proceeds. The attending clinician, in judging whether treatment is effective at the dosage administered, will consider the general well-being of the patient as well as more definite signs such as relief of disease-related symptoms, inhibition of
10 tumor growth, actual shrinkage of the tumor, or inhibition of metastasis. Size of the tumor can be measured by standard methods such as radiological studies, *e.g.*, CAT or MRI scan, and successive measurements can be used to judge whether or not growth of the tumor has been retarded or even reversed. Relief of disease-related symptoms such as pain, and improvement in overall condition can also be used to help judge effectiveness of treatment.

15 [095] Assays for Determining HSP90 Binding and Downstream Effect
A variety of *in vitro* and *in vivo* assays are available to test the effect of the alkynyl pyrrolo[2,3-
d]pyrimidine compounds of the invention on HSP90. HSP90 competitive binding assays and functional assays can be performed as known in the art by substituting in the compounds of the
20 invention. Chiosis *et al. Chemistry & Biology* **2001**, 8, 289-299, describe some of the known ways in which this can be done. For example, competition binding assays using, *e.g.*, geldanamycin or 17-AAG as a competitive binding inhibitor of HSP90 can be used to determine relative HSP90 affinity of the compounds of the invention by immobilizing the compound of
25 interest or other competitive inhibitor on a gel or solid matrix, preincubating HSP90 with the other inhibitor, passing the preincubated mix over the gel or matrix, and then measuring the amount of HSP90 that retains or does not retain on the gel or matrix.

[096] Downstream effects can also be evaluated based on the known effect of HSP90 inhibition on function and stability of various steroid receptors and signaling proteins including, *e.g.*, Raf1 and Her2. Compounds of the present invention induce dose-dependent degradation of these
30 molecules, which can be measured using standard techniques. Inhibition of HSP90 also results in up-regulation of HSP90 and related chaperone proteins that can similarly be measured. Antiproliferative activity on various cancer cell lines can also be measured, as can morphological and functional differentiation related to HSP90 inhibition.

[097] Many different types of methods are known in the art for determining protein
35 concentrations and measuring or predicting the level of proteins within cells and in fluid

5 samples. Indirect techniques include nucleic acid hybridization and amplification using, *e.g.*, polymerase chain reaction (PCR). These techniques are known to the person of skill and are discussed, *e.g.*, in Sambrook, Fritsch & Maniatis *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989; Ausubel, *et al.* *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, 1994, and, as specifically applied to the quantification, detection, and relative activity of HER2/Neu in patient samples, *e.g.*, in U.S. Patents 4,699,877, 4,918,162, 4,968,603, and 5,846,749. A brief discussion of two generic techniques that can be used follows.

10 [098] The determination of whether cells overexpress or contain elevated levels of Her2 can be determined using well known antibody techniques such as immunoblotting, radioimmunoassays, western blotting, immunoprecipitation, enzyme-linked immunosorbant assays (ELISA), and derivative techniques that make use of antibodies directed against Her2. As an example, Her2 expression in breast cancer cells can be determined with the use of an immunohistochemical assay, such as the Dako Hercep™ test (Dako Corp., Carpinteria, CA). The Hercep™ test is an antibody staining assay designed to detect Her2 overexpression in tumor tissue specimens. This particular assay grades Her2 expression into four levels: 0, 1, 2, and 3, with level 3 representing the highest level of Her2 expression. Accurate quantitation can be enhanced by employing an Automated Cellular Imaging System (ACIS) as described, *e.g.*, by Press, M. *et al.* *Modern Pathology* **2000**, 13, 225A.

20 [099] Antibodies, polyclonal or monoclonal, can be purchased from a variety of commercial suppliers, or may be manufactured using well-known methods, *e.g.*, as described in Harlow *et al.* *Antibodies: A Laboratory Manual*, 2nd ed; Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1988.

25 [0100] Her2 overexpression can also be determined at the nucleic acid level since there is a reported high correlation between overexpression of the Her2 protein and amplification of the gene that codes for it. One way to test this is by using RT-PCR. The genomic and cDNA sequences for Her2 are known. Specific DNA primers can be generated using standard, well-known techniques, and can then be used to amplify template already present in the cell. An example of this is described in Kurokawa, H. *et al.* *Cancer Res.* **2000**, 60, 5887-5894. PCR can be standardized such that quantitative differences are observed as between normal and abnormal cells, *e.g.*, cancerous and noncancerous cells. Well known methods employing, *e.g.*, densitometry, can be used to quantitate and/or compare nucleic acid levels amplified using PCR.

30 [0101] Similarly, fluorescent *in situ* hybridization (FISH) assays and other assays can be used, *e.g.*, Northern and/or Southern blotting. These rely on nucleic acid hybridization between the Her2 gene or mRNA and a corresponding nucleic acid probe that can be designed in the same or

a similar way as for PCR primers, above. See, *e.g.*, Mitchell MS, and Press M.F. *Oncol., Suppl.* **1999**, *12*, 108-116. For FISH, this nucleic acid probe can be conjugated to a fluorescent molecule, *e.g.*, fluorescein and/or rhodamine, that preferably does not interfere with hybridization, and which fluorescence can later be measured following hybridization. See, *e.g.*, Kurokawa, H *et al.*, *Cancer Res.* **2000**, *60*, 5887-5894 (describing a specific nucleic acid probe having sequence 5'-FAM-NucleicAcid-TAMRA-p-3' sequence). ACIS-based approaches as described above can be employed to make the assay more quantitative (de la Torre-Bueno, J., *et al. Modern Pathology* **2000**, *13*, 221A).

[0102] Immuno and nucleic acid detection can also be directed against proteins other than HSP90 and HER2, which proteins are nevertheless affected in response to HSP90 inhibition. The following examples are offered by way of illustration only and are not intended to be limiting of the full scope and spirit of the invention.

EXAMPLES

[0103] Materials and Methods

The chemical reagents used to create the novel products of the invention below are all available commercially, *e.g.*, from Aldrich Chemical Co., Milwaukee, WI, USA. Otherwise their preparation is facile and known to one of ordinary skill in the art, or it is referenced or described herein.


The final compounds were usually purified by preparative TLC (silica gel 60 Å, Whatman Partisil PK6F) or flash chromatography (silica gel 60 Å, EMD Chemicals) using EtOAc/hexane or MeOH/CH₂Cl₂ as eluents. R_f's were measured using silica gel TLC plates (silica gel 60 Å, EMD Chemicals). Analytical HPLC chromatograms were obtained using a C18 column (Agilent Zorbax 300SB-C18; 5 microns; 4.6 mm x 150 mm). A gradient was applied between solvent A (0.1% TFA in H₂O) and solvent B (0.5% TFA in CH₃CN) increasing the proportion of A linearly from 5% (t=0) to 100% (t=7.00 min), with a constant flow rate of 1 mL/min. The samples were diluted to typically 0.1-1 mg/mL in MeOH or CH₃CN and the injection volumes were typically 10 µL. The column was not heated, and UV detection was effected at 254 nm. ¹H-NMR spectra were recorded on a Bruker Avance 400 MHz spectrometer.

The chemical names were generated using the Beilstein Autonom 2.1 software.

[0104] General Procedures


General Procedure A: Sonogashira coupling.

PCT/US06/11846



- 5 A mixture of the appropriate 5-iodo-pyrrolo[2,3-d]pyrimidine, alkyne (2–5 equiv.), Et_3N (2–5 equiv.), $Pd(PPh_3)_4$ (0.01–0.05 equiv.), and CuI (0.05–0.30 equiv.) in DCM (5 mL/mmol of starting iodide) was heated to reflux for 0.5–3 h. The reaction mixture was washed with sat. sq. $NaHCO_3$ and brine, dried (Na_2SO_4), concentrated. Flash chromatography ($EtOAc/DCM/Et_3N$ 15:74:1, gradually adding MeOH (0–7 vol %)) gave the desired 5-alkynyl-pyrrolo[2,3-d]pyrimidine in typically 20–70% yield.


General procedure B: $ZnCl_2$ -mediated deprotection.



- 10 A suspension of 2-(pivaloylamino)-pyrrolo[2,3-d]pyrimidine and $ZnCl_2$ (3–20 equiv.) in wet $EtOH$ (5 vol% water) was heated to 80 °C, and monitored by HPLC. When the reaction reached completion, DCM was added, and the organic layer was washed with sat. aq. $NaHCO_3$ and brine. Drying (Na_2SO_4) and concentration afforded the desired 2-amino-pyrrolo[2,3-d]pyrimidine.

15 **[0105] Example 1**

3-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-yn-1-ol



See Shih, C. *et al. Heterocycles*, **1993**, 35, 825 and US Patent 5,196,424.

Step 1. 2-Amino-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one

A mixture of 2,4-diamino-6-hydroxypyrimidine (300 g, 2.37 mol), chloroacetaldehyde (50% aq. solution, 382 g, 2.43 mol, 303 mL, 1.02 eq.), sodium acetate (195 g, 2.37 mol), DMF (2.5 L), and water (360 mL) was stirred mechanically at rt for 2 days. The resulting solid was collected by filtration, and washed with water (50 mL × 3). The mother liquor was concentrated to give additional material which was washed with water (50 mL × 3). The combined solid materials were recrystallized from MeOH to give the title compound as a white powder (186 g, 52% yield, HPLC purity: 100%). t_R : 2.21 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 11.00 (br. s, 1H), 10.33 (br. s, 1H), 6.63 (q, 1H), 6.21 (q, 1H), 6.12 (br. s, 2H).

Step 2. 2,2-Dimethyl-N-(4-oxo-4,7-dihydro-3H-pyrrolo[2,3-d]pyrimidin-2-yl)-propionamide

A solution of 2-amino-3,7-dihydro-pyrrolo[2,3-d]pyrimidin-4-one (186 g, 1.23 mol) in pyridine (2 L) was treated with trimethylacetyl chloride (475 g, 3.94 mol, 485 mL, 3.2 eq) at 90 °C for 2h, to give a mixture of N(2)-monoacylated and N(2), N(7)-bisacylated material. The solvent was evaporated and the residue was taken up in aqueous ammonia (37% NH_3 , 310 mL) and MeOH (2L), and stirred at rt for 30 min, to selectively cleave the N(7)-pivaloyl group. The solid was collected by filtration, washed with water (500 mL × 5), and dried on high vacuum to give the title compound as a solid (193 g). Concentration of the mother liquor provided additional solid material, which was collected, washed with water (50 mL × 5), and dried (77 g). The combined

yield was 93% (HPLC purity 98.6%). t_R : 4.57 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 12.00, (br. s, 1H), 7.40 (br. s, 1H), 6.96 (q, 1H), 6.40 (q, 1H), 1.24 (s, 9H).

Step 3. N-(4-Chloro-7H-pyrrolo[2,3-d]pyrimidin-2-yl)-2,2-dimethyl-propionamide

A mixture of 2,2-dimethyl-N-(4-oxo-4,7-dihydro-3H-pyrrolo[2,3-d]pyrimidin-2-yl)-

- 5 propionamide (210 g, 0.90 mol), POCl_3 (828 g, 5.40 mol, 503 mL, 6.0 eq), benzyltriethylammonium chloride (411 g, 1.80 mol), N,N-dimethylaniline (220 g, 231 mL, 1.80 mol), and acetonitrile (2.0 L) was heated to reflux for 40–60 min, monitoring with HPLC. The solvent was evaporated on a rotary evaporator, and the residue was carefully (**caution: exothermic and corrosive**) and slowly added to ice water (16 L). The pH was adjusted to 7
- 10 with solid NaOH, and the resulting precipitate was collected by filtration. Drying afforded the title compound (159 g, 70% yield 70%, HPLC purity 100%). t_R : 5.37 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 12.35 (br. s, 1H), 10.06 (br. s, 1H), 7.54 (q, 1H), 6.52 (q, 1H), 1.25 (s, 9H).

Step 4. N-(4-Chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-2-yl)-2,2-dimethyl-propionamide

- A solution of N-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-2-yl)-2,2-dimethyl-propionamide (101 g, 0.40 mol) in anhydrous THF (2 L) was treated with N-iodosuccinimide (98.9g, 0.44mol, 1.1 eq.)
- 15 under N_2 atmosphere at rt for 40 min. The solvent was evaporated, and the residue was taken up in CH_2Cl_2 (1.5 L), and washed with Na_2SO_3 (500 mL $\times$ 3) and brine (300 mL $\times$ 3). Evaporated and recrystallization from MeOH gave the title product as a white powder (122 g, 81% yield, HPLC purity: 98.2%). t_R : 6.19 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 12.65 (br.s, 1H), 10.11 (br. s, 1H), 7.76 (d,
- 20 1H), 1.24 (s, 9H).

Step 5. N-[4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide

- A mixture of N-(4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-2-yl)-2,2-dimethyl-propionamide (37.8 g, 0.1 mol), 2-chloromethyl-4-methoxy-3,5-dimethyl-pyridine hydrochloride (23.1 g, 0.104
- 25 mol), finely powdered K_2CO_3 , (41.5 g, 0.3 mol, 3.0 eq.), and anhydrous DMF (200 mL) was stirred at rt overnight. The solvent was evaporated, and the residue taken up in CH_2Cl_2 (500mL), washed with brine (200 mL $\times$ 3), evaporated, and recrystallized from MeOH to give the title product as a white powder (42.0g, 80% yield; HPLC purity 98%). t_R : 6.49 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 10.16 (br. s, 1H), 8.04 (s, 1H), 7.72 (s, 1H), 5.46 (s, 2H), 3.73 (s, 3H), 2.33 (s, 3H),
- 30 2.15 (s, 3H), 1.21 (s, 9H).

Step 6. 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine

- A mixture of N-[4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide (1.0g, 1.89mmol), ZnCl_2 (1.29 g,
- 35 9.47 mol, 5.0 eq.), and EtOH/ H_2O solution (25mL, volumetric ratio 100:5) was stirred at 80°C


overnight. The reaction mixture poured into water, and the solid was collected by filtration, washed with water (10 mL×3), and recrystallized from MeOH to give the title product (0.67 g, 85% yield; HPLC purity 98%). t_R : 5.39 min. $^1\text{H-NMR}$ (DMSO- d_6) δ 8.07 (s, 1H), 7.28 (s, 1H), 6.75 (br. s, 2H), 5.29 (s, 2H), 3.73 (s, 3H), 2.26 (s, 3H), 2.17 (s, 3H).

5 **Step 7. 3-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-yn-1-ol**

The title compound was prepared by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see example 1) with propargylic alcohol according to the General Procedure A. t_R : 4.42min. $^1\text{H-NMR}$ (DMSO- d_6) δ 8.04 (s, 1H), 7.32 (s, 1H), 6.71 (br. s, 2H), 5.27 (s, 2H), 5.22 (t, 1H), 4.26 (d, 1H), 3.71 (s, 3H), 2.24 (s, 3H), 2.14 (s, 3H).

[0106] Example 2


4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol



The title compound was prepared by Sonogashira coupling of 4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 3-butyn-1-ol according to the General Procedure A. t_R : 4.54min. $^1\text{H-NMR}$ (DMSO- d_6) δ 8.04 (s, 1H), 7.22 (s, 1H), 6.69 (br.s., 2H), 5.25 (s, 2H), 4.82 (d, 1H), 3.70 (s, 3H), 3.55 (q, 2H), 2.49 (t, 2H), 2.23 (s, 3H), 2.14 (s, 3H).

[0107] Example 3

5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-yn-1-ol




The title compound was prepared by Sonogashira coupling of 4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 4-

pentyn-1-ol according to the General Procedure A. t_R : 4.71 min. $^1\text{H-NMR}$ (DMSO-d_6) δ 8.04 (s, 1H), 7.22 (s, 1H), 6.69 (br. s, 2H), 5.25 (s, 2H), 4.48 (d, 1H), 3.71 (s, 3H), 3.50 (q, 2H), 2.42 (t, 2H), 2.23 (s, 3H), 2.14 (s, 3H), 1.65 (s, 2H).

[0108] Example 4

5 **4-Chloro-5-(3-diisobutylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine**



Step1. *N*-[4-Chloro-5-(3-hydroxy-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide

- 10 The title compound was prepared by Sonogashira coupling of *N*-[4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide (see Example 1) with propargylic alcohol according to the General Procedure A. t_R : 5.58 min. $^1\text{H-NMR}$ (DMSO-d_6): δ 10.15 (br. s, 1H), 8.04 (s, 1H), 7.80 (s, 1H), 5.48 (s, 2H), 5.31 (t, 1H), 4.33 (d, 2H), 3.74 (s, 3H), 2.33 (s, 3H), 2.15 (s, 3H), 1.21 (s, 9H).

- 15 Step2. *N*-[4-Chloro-5-(3-diisobutylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide

- A solution of *N*-[4-chloro-5-(3-hydroxy-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide (9.0g, 19.7 mmol) and Et_3N (2 mL) in DCM (50 mL) was treated with MsCl (15.3 mL, 19.7 mmol) at 0 °C for 20 min and evaporated to give a yellow solid, containing mostly the desired mesylate (10.7 g). An aliquot of this solid (106.8 mg, 0.2 mmol) was treated with diisobutylamine (25.9 mg, 24.8 μL) in DCM (2 mL) at 0°C for 10h, and then at 25 °C for 3 days. Evaporation and preparative plate chromatography gave the title compound.

- 20 Step 3. 4-Chloro-5-(3-diisobutylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine


The title compound was prepared by cleaving the pivaloyl protecting group of *N*-[4-chloro-5-(3-diisobutylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide with ZnCl_2 according to the General Procedure B.

t_R : 5.23 min. 1H -NMR ($CDCl_3$) δ 8.23 (s, 1H), 7.08 (s, 1H), 5.31 (s, 2H), 4.99 (s, 1H), 3.76 (s, 3H), 3.56 (s, 2H), 2.29 (s, 3H), 2.27 (d, 4H), 2.21 (s, 3H), 1.74 (m, 2H), 0.90 (d, 12H).

[0109] Example 5

4-Chloro-5-(3-diisopropylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-

5-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1. N-[4-Chloro-5-(3-diisopropylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide

A solution of N-[4-chloro-5-(3-hydroxy-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide (9.0 g, 19.7 mmol) and Et₃N (2 mL) in DCM (50 mL) was treated with MsCl (15.3 mL, 19.7 mmol) at 0 °C for 20 min and evaporated to give a yellow solid, containing mostly the desired mesylate (10.7 g). An aliquot of this solid (106.8 mg, 0.2 mmol) was treated with diisopropylamine (20.3 mg, 28.1 μ L) in DCM (2 mL) at 0 °C for 10 h, and then at 25 °C for 3 days. Evaporation and preparative plate chromatography gave the title compound.

Step 2. 4-Chloro-5-(3-diisopropylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine


The title compound was prepared by cleaving the pivaloyl protecting group of N-[4-chloro-5-(3-diisopropylamino-prop-1-ynyl)-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide with ZnCl₂ according to the General Procedure B. t_R : 4.71 min. 1H -NMR ($CDCl_3$) δ 8.24 (s, 1H), 7.05 (s, 1H), 5.30 (s, 2H), 4.96 (s, 1H), 3.76 (s, 3H), 3.68 (s, 2H), 3.27 (s, 2H), 2.27 (s, 2H), 2.20 (s, 2H), 1.15 (d, 12H).

[0110] Example 6

4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-pyridin-2-ylethynyl-7H-

pyrrolo[2,3-d]pyrimidin-2-ylamine


PCT/US06/11846



The title compound was prepared by Sonogashira coupling of 4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 2-ethynylpyridine according to the General Procedure A. t_R : 4.65 min. 1H -NMR (DMSO- d_6) δ 8.59 (s, 1H), 8.07 (s, 1H), 7.82 (td, 1H), 7.62 (s, 1H), 7.56 (d, 1H), 7.36 (td, 1H), 6.84 (br. s, 2H), 5.36 (s, 2H), 3.74 (s, 3H), 2.28 (s, 3H), 2.17 (s, 3H).

[0111] Example 7


6-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-hex-5-yn-1-ol



The title compound was prepared by Sonogashira coupling of 4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 5-hexyn-1-ol according to the General Procedure A. t_R : 4.92 min. 1H -NMR (DMSO- d_6) δ 8.04 (s, 1H), 7.22 (s, 1H), 6.69 (br. s, 2H), 5.25 (s, 2H), 4.39 (t, 1H), 3.70 (s, 3H), 3.40 (q, 2H), 2.39 (t, 2H), 2.23 (s, 3H), 2.14 (s, 3H), 1.55 (m, 4H).

[0112] Example 8

4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-trimethylsilanylethynyl-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1: N-[4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-trimethylsilanylethynyl-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide


The title compound was prepared by Sonogashira coupling of N-[4-chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide (see Example 1) with ethynyl-trimethylsilane according to the General Procedure A. $t_R = 7.48$ min. ^1H NMR (CDCl_3) δ 8.18 (s, 2H), 7.38 (s, 1H), 5.49 (s, 2H), 3.74 (s, 3H), 2.24 (s, 6H), 1.35 (s, 9H), 0.24 (s, 9H).

Step 2: 4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-trimethylsilanylethynyl-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine

The title compound was prepared by cleaving the pivaloyl protecting group of N-[4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-trimethylsilanylethynyl-7H-pyrrolo[2,3-d]pyrimidin-2-yl]-2,2-dimethyl-propionamide with ZnCl_2 according to the General Procedure B. $t_R = 6.56$ min. ^1H NMR (CDCl_3) δ 8.22 (s, 1H), 7.11 (s, 1H), 5.30 (s, 2H), 5.09 (s, 2H), 3.75 (s, 3H), 2.26 (s, 3H), 2.18 (s, 3H), 0.24 (s, 9H).

[0113] Example 9

5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynoic acid




The title compound was prepared by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 4-pentynoic acid according to the General Procedure A. $t_R = 4.73$ min. ^1H NMR (DMSO-d_6) δ 12.2 (s, 1H), 8.15 (s, 1H), 7.28 (s, 1H), 6.80 (s, 2H), 5.33 (s, 2H), 3.73 (s, 3H), 3.31 (br. s, 2H), 2.362 (br. t, 2H), 2.27 (s, 3H), 2.17 (s, 3H).

[0114] Example 10

5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynoic acid diethylamide


PCT/US06/11846



A mixture of 5-[2-amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynoic acid (see previous example, 25 mg), Et₂NH (100 μ L), EDCI (87 mg), HOBt (50 mg), and DMF (1 mL) was stirred at rt for 16 h. Preparative plate chromatography (EtOAc/hexane 2:1) gave the title compound. t_R = 5.33 min. ¹H NMR (CDCl₃) δ 8.23 (s, 1H), 7.01 (s, 1H), 5.32 (s, 2H), 4.98 (s, 2H), 3.76 (s, 3H), 3.42 (q, 2H), 3.33 (q, 2H), 2.78 (dd, 2H), 2.64 (dd, 2H), 2.27 (s, 3H), 2.18 (s, 3H), 1.20 (t, 3H), 1.13 (t, 3H).

[0115] Example 11

5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-1-(4-methyl-piperazin-1-yl)-pent-4-yn-1-one



Step 1: 1-(4-Methyl-piperazin-1-yl)-pent-4-yn-1-one

A solution of 4-pentynoic acid (519 mg, 5.29 mmol) and Et₃N (737 μ L, 5.29 mmol) in DCM (10 mL) was treated with ethyl chloroformate (504 μ L, 5.29 mmol) at rt for 15 min. Then N-methyl-piperazine (588 μ L, 5.29 mmol) was added, and stirring was prolonged for 45 min. Work-up (sat. aq. NaHCO₃) and concentration gave the title compound as a colorless oil. ¹H NMR (CDCl₃) δ 3.63 (t, 2H), 3.48 (t, 2H), 2.55 (m, 4H), 2.38 (quint., 4H), 2.30 (s, 3H).

Step 2: 5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-1-(4-methyl-piperazin-1-yl)-pent-4-yn-1-one


The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 1-(4-Methyl-piperazin-1-yl)-pent-4-yn-1-one according to the General Procedure A. t_R = 4.16 min.

¹H NMR (CDCl₃) δ 8.22 (s, 1H), 7.00 (s, 1H), 5.30 (s, 2H), 5.14 (s, 2H), 3.74 (s, 3H), 3.64 (t, 2H), 3.50 (t, 2H), 2.76 (dd, 2H), 2.64 (dd, 2H), 2.29 (s, 3H), 2.24 (s, 3H), 2.18 (s, 3H).

[0116] Example 12

5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-

5-d]pyrimidin-5-yl]-pent-4-ynoic acid amide



Step 1: Pent-4-ynoic acid amide


A solution of 4-pentynoic acid (517 mg, 5.29 mmol) and Et₃N (737 μL, 5.29 mmol) in DCM (10 mL) was treated with ethyl chloroformate (504 μL, 5.29 mmol) at rt for 15 min. Then NH₃ (7M in MeOH, 1mL, 7 mmol) was added, and stirring was prolonged for 5 min. Work-up (sat. aq. NaHCO₃) and concentration gave the title compound as a colorless solid. ¹H NMR (CDCl₃) δ 6.03 (s, 1H), 5.86 (s, 1H), 2.55 (m, 2H), 2.46 (m, 2H), 2.02 (t, 1H).

Step 2: 5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynoic acid amide

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with pent-4-ynoic acid amide according to the general procedure A. *t*_R = 4.35 min. ¹H NMR (DMSO-d₆) δ 8.07 (s, 1H), 7.36 (s, 1H), 7.23 (s, 1H), 6.87 (s, 1H), 6.71 (s, 2H), 5.28 (s, 2H), 3.72 (s, 3H), 2.59 (t, 2H), 2.34 (t, 2H), 2.25 (s, 3H), 2.16 (s, 3H).

[0117] Example 13

{3-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-ynyl}-carbamic acid tert-butyl ester



~~PCT/US2006/011846~~
 Step 1: *Prop-2-ynyl-carbamic acid tert-butyl ester*

A solution of propargylamine (1g, 18 mmol) in DCM (10 mL) was treated with (BOC)₂O (4.0 g, 18 mmol) at rt for 3h. Evaporation gave the title compound as an oil. ¹H NMR (CDCl₃) δ 4.80 (s, 1H), 3.93 (s, 2H), 2.23 (t, 1H), 1.47 (s, 9H).


5 *Step 2: {3-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-ynyl}-carbamic acid tert-butyl ester*

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with prop-2-ynyl-carbamic acid tert-butyl ester according to the General Procedure A. *t*_R = 5.91 min.

10 ¹H NMR (CDCl₃) δ 8.20 (s, 1H), 7.05 (s, 1H), 5.36 (s, 2H), 5.31 (s, 2H), 4.82 (s, 1H), 4.16 (s, 2H), 3.76 (s, 3H), 2.26 (s, 3H), 2.24 (s, 3H), 1.47 (s, 9H).

[0118] Example 14

5-(3-Amino-prop-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine




15

A solution of {3-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-ynyl}-carbamic acid tert-butyl ester (see previous example, 22 mg) in DCM (3 mL) was treated with TFA (0.6 mL) at rt for 15 min. Evaporation, work-up (DCM/sat. aq. NaHCO₃), drying (Na₂SO₄), and evaporation gave the title compound as an oil.

20 *t*_R = 3.85 min. ¹H NMR (CDCl₃) δ 8.24 (s, 1H), 7.06 (s, 1H), 5.32 (s, 2H), 5.02 (s, 2H), 3.75 (s, 3H), 2.27 (s, 3H), 2.20 (s, 3H).

[0119] Example 15

{5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynyl}-carbamic acid tert-butyl ester



Step 1: Pent-4-ynyl-carbamic acid tert-butyl ester


A solution of pent-4-ynylamine (480 mg; Li, Y. *et al.* J. Am. Chem. Soc. **1996**, *118*, 9295) in DCM (5 mL) was treated with (BOC)₂O at rt for 15 min. Evaporation gave the title compound as a colorless oil. ¹H NMR (CDCl₃) δ 4.68 (s, 1H), 3.24 (q, 2H), 2.25 (td, 2H), 1.97 (t, 1H), 1.70 (quint., 2H), 1.45 (s, 9H).

Step 2: {5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynyl}-carbamic acid tert-butyl ester.

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with pent-4-ynyl-carbamic acid tert-butyl ester according to the General Procedure A. *t*_R = 5.96 min. ¹H NMR (CDCl₃) δ 8.23 (s, 1H), 7.02 (s, 1H), 5.30 (s, 2H), 5.07 (s, 2H), 4.80 (br. t, 1H), 3.75 (s, 3H), 3.30 (q, 2H), 2.48 (t, 2H), 2.26 (s, 3H), 2.19 (s, 3H), 1.80 (quint., 2H), 1.43 (s, 9H).

[0120] Example 16

5-(5-Amino-pent-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine




A solution of {5-[2-amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-ynyl}-carbamic acid tert-butyl ester (see previous example, 33 mg) in DCM (1.0 mL) was treated with TFA (0.2 mL) at rt for 10 min. The reaction mixture was concentrated, taken in water (1.5 mL), washed with EtOAc (1.5 mL), made alkaline with sat. aq. NH₄OH (0.5 mL), and back-extracted into EtOAc (25 mL). Washing (NH₄OH 1M), drying (Na₂SO₄) and concentration afforded the title compound. *t*_R = 4.37 min. ¹H NMR (CDCl₃) δ 8.15

(s, 1H), 6.97 (s, 1H), 5.25 (s, 2H), 3.72 (s, 3H), 2.94 (m, 2H), 2.48 (t, 2H), 2.23 (s, 3H), 2.16 (s, 3H), 1.80 (quint., 2H).

[0121] Example 17

2-{5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo

5 [2,3-d]pyrimidin-5-yl]-pent-4-ynyl}-isoindole-1,3-dione




The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 2-pent-4-ynyl-isoindole-1,3-dione (Li, Y. *et al.* J. Am. Chem. Soc. **1996**, *118*, 9295) according to the General Procedure A. $t_R = 6.08$ min. 1H NMR ($CDCl_3$) δ 8.17 (s, 1H), 7.75 (dd, 2H), 7.58 (dd, 2H), 6.85 (s, 1H), 5.22 (s, 2H), 3.81 (t, 2H), 3.72 (s, 3H), 2.46 (t, 2H), 2.23 (s, 3H), 2.14 (s, 3H), 1.95 (quint., 2H).

[0122] Example 18

4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-(4-morpholin-4-yl-but-1-ynyl)-

7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1: Methanesulfonic acid but-3-ynyl ester

A solution of 3-but-1-ynol (5.62 g, 80.2 mmol) and Et_3N (14.5 mL, 104 mmol) in DCM (40 mL) was treated with $MsCl$ (7.48 mL, 96.2 mmol) at $0^\circ C$ for 10 min. Work-up (water; sat. aq. $NaHCO_3$), drying (Na_2SO_4), and concentration gave the title compound as a pale orange oil. 1H NMR ($CDCl_3$) δ 4.32 (t, 2H), 3.07 (s, 3H), 2.67 (td, 2H), 2.09 (t, 2H).

Step 2: 4-But-3-ynyl-morpholine

A mixture of methanesulfonic acid but-3-ynyl ester (10.1 g, 68 mmol) and morpholine (12.5 mL, 143 mmol) was heated to $100^\circ C$ for 1h. The mixture was diluted with Et_2O (50 mL) and filtered.

~~PCT/US2006/011846~~


The solid residue was discarded, and the mother liquor extracted with 3N HCl. The aqueous layer was basified with NaOH, and back-extracted into EtOAc. Drying (Na₂SO₄), concentration, and distillation (short-path, b.p.= 130 °C at ~5 mm Hg) gave the title compound as a colorless oil. ¹H NMR (CDCl₃) δ 3.72 (t, 4H), 2.60 (t, 2H), 2.48 (t, 4H), 2.39 (td, 2H), 2.00 (t, 1H).

5 **Step 3: 4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-(4-morpholin-4-yl-but-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine**

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 4-but-3-ynyl-morpholine according to the General Procedure A. *t*_R = 4.12 min. ¹H NMR (CDCl₃) δ 8.33 (s, 1H), 6.96 (s, 1H), 5.26 (s, 2H), 5.12 (s, 2H), 3.76 (s, 3H), 3.72 (t, 4H), 2.67 (m, 2H), 2.60 (m, 2H), 2.52 (t, 4H), 2.25 (s, 3H), 2.20 (s, 3H).

[0123] Example 19

4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-(5-morpholin-4-yl-pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1: 5-Iodo-pent-1-yne

A mixture of 5-chloro-pent-1-yne (5.0 mL, 47 mmol), NaI (9.7 g, 65 mmol) and acetone (20 mL) was heated to reflux for 15 h. Additional NaI (9.7 g, 65 mmol) was added, and the refluxed was prolonged for 24 h. After filtration and concentration, the residue was dissolved in hexane and washed with water and aq. Na₂S₂O₃. Drying (Na₂SO₄) and concentration afforded the title compound. ¹H NMR (CDCl₃) δ 3.33 (t, 2H), 2.36 (td, 2H), 2.03 (quint., 2H), 2.01 (t, 1H).

Step 2: 4-Pent-4-ynyl-morpholine


A mixture of 5-iodo-pent-1-yne (2.40 g, 12.3 mmol) and morpholine (2.70 g, 30.9 mmol) was heated to 80 °C for 15 min. The mixture was diluted with Et₂O and filtered. The solid residue was discarded, and the mother liquor was extracted with 3N HCl. The aqueous layer was made basic with NaOH, and back-extracted into EtOAc. Drying (Na₂SO₄), and concentration gave the title compound as a pale orange oil. ¹H NMR (CDCl₃) δ 3.67 (t, 4H), 2.39 (m, 6H), 2.21 (td, 2H), 1.93 (t, 1H), 1.67 (quint., 2H).

Step 3: 4-Chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-5-(5-morpholin-4-yl-pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with 4-pent-4-ynyl-morpholine according to the general procedure A. $t_R = 4.26$ min. ^1H NMR (CDCl_3) δ 8.21 (s, 1H), 7.00 (s, 1H), 5.29 (s, 2H), 5.13 (s, 2H), 3.74 (s, 3H), 3.71 (t, 4H), 2.51–2.45 (m, 8H), 2.25 (s, 3H), 2.18 (s, 3H), 1.78 (quint., 2H).

[0124] Example 20

5-(4-tert-Butylamino-but-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1: *tert*-Butyl-but-3-ynyl-amine


A solution of toluene-4-sulfonic acid but-3-ynyl ester (1.57 g, 7.1 mmol) in *tert*-BuNH₂ (3.54 g, 48 mmol) was heated to reflux for 18 h. The reaction mixture was diluted with Et₂O (20 mL), filtered, and concentrated to afford the title product. ^1H NMR (CDCl_3) δ 2.74 (t, 2H), 2.38 (td, 2H), 2.00 (t, 1H), 1.12 (s, 9H).

Step 2: **5-(4-tert-Butylamino-but-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine**

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with *tert*-butyl-but-3-ynyl-amine according to the General Procedure A. $t_R = 4.38$ min. ^1H NMR (CDCl_3) δ 8.21 (s, 1H), 7.03 (s, 1H), 5.29 (s, 2H), 5.10 (s, 2H), 3.74 (s, 3H), 2.81 (t, 2H), 2.62 (t, 2H), 2.25 (s, 3H), 2.18 (s, 3H), 1.13 (s, 9H).

[0125] Example 21

5-(5-tert-Butylamino-pent-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine



Step 1: tert-Butyl-pent-4-ynyl-amine


A solution of 5-iodo-pent-1-yne (1.0 g, 5.2 mmol) in *tert*-BuNH₂ (2.0 g, 27 mmol) was heated to reflux for 4 h, evaporated, diluted with Et₂O, filtered, and concentrated to afford the title compound. ¹H NMR (CDCl₃) δ 2.67 (t, 2H), 2.27 (td, 2H), 1.96 (t, 1H), 1.70 (quint., 2H), 1.11 (s, 9H).

Step 2: 5-(5-tert-Butylamino-pent-1-ynyl)-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine

The title compound was obtained by Sonogashira coupling of 4-Chloro-5-iodo-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylamine (see Example 1) with *tert*-butyl-pent-4-ynyl-amine according to the General Procedure A. *t*_R = 4.51 min. ¹H NMR (CDCl₃) δ 8.23 (s, 1H), 7.02 (s, 1H), 5.30 (s, 2H), 5.00 (s, 2H), 3.75 (s, 3H), 2.74 (t, 2H), 2.50 (t, 2H), 2.27 (s, 3H), 2.19 (s, 3H), 1.76 (quint., 2H), 1.12 (s, 9H).

[0126] Example 22

Dimethylamino-acetic acid 3-[2-amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-ynyl ester




A solution of 3-[2-amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-prop-2-yn-1-ol (see example 1, 177 mg, 0.48 mmol) in anhydrous pyridine (3 mL) was treated with N,N-dimethylamino-acetyl chloride hydrochloride (113 mg, 0.72 mmol) at rt for 30 min. Addition of toluene (10 mL) caused the formation of a sticky pellet. The toluene solution was discarded, and the pellet was partitioned between water and DCM. The DCM layer was concentrated, and purified by flash chromatography (EtOAc/DCM/Et₃N 33:66:1,

gradually adding MeOH (0–2%)) to afford the title compound. $t_R = 4.51$ min. ^1H NMR (CDCl_3) δ 8.23 (s, 1H), 7.14 (s, 1H), 5.31 (s, 2H), 5.00 (s, 2H), 4.98 (s, 2H), 3.76 (s, 3H), 3.25 (s, 2H), 2.38 (s, 6H), 2.28 (s, 3H), 2.20 (s, 3H).

[0127] Example 23


5 **5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol**



Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 2-methylpent-4-yn-2-ol (H. Zhang et al., *Tetrahedron Lett.* **1999**, 40, 7851) according to the general procedure A gave the title compound, as a solid. Mp = 163–165 °C. HPLC Rt = 4.98 min.

[0128] Example 24

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(5-(4-methylpiperazin-1-yl)pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Step 1: 1-methyl-4-(pent-4-ynyl)piperazine


A mixture of pent-4-ynyl methanesulfonate (887 mg) and N-methyl piperazine (607 μL) was heated to 80 °C for 4.5 h, diluted with 1,2-dichloroethane (5 mL) and heated to 70 °C for 18h. Saturated aq. NaHCO_3 was added (10 mL) and the mixture was extracted with DCM (3 x 40 mL), dried over Na_2SO_4 , and concentrated to afford 1-methyl-4-(pent-4-ynyl)piperazine (0.23 g).

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-methyl-4-(pent-4-ynyl)piperazine, according to the general procedure A gave the title compound, as a solid. Mp = 160.1–162.3 °C. HPLC Rt = 4.08 min.

~~PCT/US2006/011846~~
 [0129] Example 25

4-chloro-5-(5-(4-ethylpiperazin-1-yl)pent-1-ynyl)-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



5 Step 1: 1-ethyl-4-(pent-4-ynyl)piperazine


A mixture of pent-4-ynyl methanesulfonate (849 mg) and N-methyl piperazine (665 μ L) was heated to 80 °C for 3 h, diluted with 1,2-dichloroethane (5 mL) and heated to 70 °C for 18h. Saturated aq. NaHCO₃ was added (10 mL) and the mixture was extracted with DCM (1 x 40 mL), dried over Na₂SO₄, and concentrated to afford 0.56 g of 1-ethyl-4-(pent-4-ynyl)piperazine.

10 Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-ethyl-4-(pent-4-ynyl)piperazine, according to the general procedure A gave the title compound, as a solid. Mp = 142.3-144.1 °C. HPLC Rt = 4.11 min.

15 [0130] Example 26

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(4-(4-methylpiperazin-1-yl)but-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Step 1: 1-(but-3-ynyl)-4-methylpiperazine


20 A mixture of but-3-ynyl 4-methylbenzenesulfonate (972 mg) and N-methyl piperazine (482 μ L) was heated to 80°C for 4.5 h, diluted with 1,2-dichloroethane (5 mL) and heated to 70 °C for 18h. Saturated aq. NaHCO₃ was added (10 mL) and the mixture was extracted with DCM (3 x 40 mL), and concentrated to afford 0.70 g of 1-(but-3-ynyl)-4-methylpiperazine as a 3:2 mixture of tosylate salt and free base.

25 Step 2

~~PCT/US2006/011846~~
 Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(but-3-ynyl)-4-methylpiperazine according to the general procedure A gave the title compound, as an oil. HPLC Rt = 4.05 min.

[0131] Example 27

- 5 **4-(5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-ynyl)-N-methylpiperazine-1-carboxamide**



Step 1: 1-(pent-4-ynyl)piperazine

- A solution of pent-4-ynyl methanesulfonate (19.55 g) and piperazine (41.5 g) in EtOH (80 mL) was heated to reflux for 1h, and evaporated. The residue was taken in NaOH 2M (70 mL) and extracted with Et₂O (70 mL). Evaporation gave 1-(pent-4-ynyl)piperazine (4.4 g).

Step 2: 4-nitrophenyl 4-(pent-4-ynyl)piperazine-1-carboxylate

- A solution of 1-(pent-4-ynyl)piperazine (2.2 g) and Et₃N (22 mL) in 1,2-dichloroethane (22 mL) was treated with p-nitrophenyl chloroformate (2.9 g) at rt for 10 min. Work-up and silica gel chromatography (DCM:EtOAc 1:3 → 0:3) gave 4-nitrophenyl 4-(pent-4-ynyl)piperazine-1-carboxylate as a yellow oil (2.96 g).

Step 3: N-methyl-4-(pent-4-ynyl)piperazine-1-carboxamide

- A solution of 4-nitrophenyl 4-(pent-4-ynyl)piperazine-1-carboxylate (1.37 g) in THF (10 mL) was treated with 40% aq. MeNH₂ (10 mL) at 70 °C for 2 h. The mixture was concentrated and diluted with conc. NH₄OH (5 mL) and water (15 mL). Extraction (DCM, 80 mL) and concentration gave N-methyl-4-(pent-4-ynyl)piperazine-1-carboxamide (0.659 g).


Step 4

- Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with N-methyl-4-(pent-4-ynyl)piperazine-1-carboxamide according to the general procedure A gave the title compound, as a solid. Mp = 191.2-193.2 °C. HPLC Rt = 4.24 min.

[0132] Example 28

4-(5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-ynyl)-N-ethylpiperazine-1-carboxamide

PCT/US06/11846



Step 1: N-ethyl-4-(pent-4-ynyl)piperazine-1-carboxamide


A solution of 1-(pent-4-ynyl)piperazine (1.08) in DCM (5 mL) was treated with ethyl isocyanate (0.56 mL) at rt for 2h, and evaporated to yield N-ethyl-4-(pent-4-ynyl)piperazine-1-carboxamide.

5 *Step 2*

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with N-ethyl-4-(pent-4-ynyl)piperazine-1-carboxamide according to the general procedure A gave the title compound, as a solid. Mp = 200.5-203.3 °C. HPLC Rt = 4.35 min.

10 [0133] Example 29

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(4-(4-phenylpiperazin-1-yl)but-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Step 1: 1-(but-3-ynyl)-4-phenylpiperazine


15 A solution of but-3-ynyl 4-methylbenzenesulfonate (1.72 g), N-phenyl piperazine (1.17 mL) and diisopropylethylamine (1.45 mL) in 1,2-dichloroethane (5 mL) was heated to reflux overnight. The mixture was concentrated, diluted with sat. aq. NaHCO₃ (5 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (3% Et₃N in EtOAc) afforded 1-(but-3-ynyl)-4-phenylpiperazine (0.76 g)

20 *Step 2*

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(but-3-ynyl)-4-phenylpiperazine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.97 min.

PCT/US2006/011846
[0134] Example 30

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(4-(4-(pyridin-2-yl)piperazin-1-yl)but-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



5 *Step 1: 1-(but-3-ynyl)-4-(pyridin-2-yl)piperazine*


A solution of but-3-ynyl 4-methylbenzenesulfonate (1.27 g), 1-(pyridin-2-yl)piperazine (1.17 g) and diisopropylethylamine (1.37 mL) in 1,2-dichloroethane (7 mL) was heated to reflux overnight. The mixture was concentrated, diluted with sat. aq. NaHCO₃ (5 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (EtOAc/hexane
10 1:1 → 1:0) afforded 1-(but-3-ynyl)-4-(pyridin-2-yl)piperazine (0.59 g)

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(but-3-ynyl)-4-(pyridin-2-yl)piperazine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.09 min.

15 [0135] Example 31

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(4-(4-(pyrimidin-2-yl)piperazin-1-yl)but-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Step 1: 2-(4-(but-3-ynyl)piperazin-1-yl)pyrimidine

20 A solution of but-3-ynyl 4-methylbenzenesulfonate (1.14 g), 2-(piperazin-1-yl)pyrimidine (1.06 g) and diisopropylethylamine (1.24 mL) in 1,2-dichloroethane (6 mL) was heated to reflux overnight. The mixture was concentrated, diluted with sat. aq. NaHCO₃ (5 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (EtOAc/hexane 1:1 → 1:0) afforded 2-(4-(but-3-ynyl)piperazin-1-yl)pyrimidine (0.60 g).


25

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 2-(4-(but-3-ynyl)piperazin-1-yl)pyrimidine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.50 min.

5 [0136] Example 32

4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(5-(4-phenylpiperazin-1-yl)pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Step 1: 1-(pent-4-ynyl)-4-phenylpiperazine


- 10 A solution of pent-4-ynyl methanesulfonate (1.05 g), N-phenyl piperazine (1.02 g) and diisopropylethylamine (1.2 mL) in THF (5 mL) was heated to reflux overnight. The mixture was concentrated, diluted with aq. NaOH 2M (10 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (EtOAc/hexane 2:1 → 2:0) afforded 1-(pent-4-ynyl)-4-phenylpiperazine (1.03 g).

15 Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(pent-4-ynyl)-4-phenylpiperazine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 5.08 min.

[0137] Example 33

- 20 **4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(5-(4-(pyridin-2-yl)piperazin-1-yl)pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine**



Step 1: 1-(pent-4-ynyl)-4-(pyridin-2-yl)piperazine

- 25 A solution of pent-4-ynyl methanesulfonate (1.05 g), 1-(pyridin-2-yl)piperazine (1.02 g) and diisopropylethylamine (1.2 mL) in THF (5 mL) was heated to reflux overnight. The mixture was


~~PCT/US2006/011846~~
concentrated, diluted with aq. NaOH 2M (10 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (EtOAc/hexane 2:1 → 2:0) afforded 1-(pent-4-ynyl)-4-(pyridin-2-yl)piperazine (1.02 g).

Step 2

- 5 Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(pent-4-ynyl)-4-(pyridin-2-yl)piperazine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.23 min.

[0138] Example 34

- 10 **4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(5-(4-(pyrimidin-2-yl)piperazin-1-yl)pent-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine**



Step 1: 2-(4-(pent-4-ynyl)piperazin-1-yl)pyrimidine


- 15 A solution of pent-4-ynyl methanesulfonate (1.05 g), 2-(piperazin-1-yl)pyrimidine (1.02 g) and diisopropylethylamine (1.2 mL) in THF (5 mL) was heated to reflux overnight. The mixture was concentrated, diluted with aq. NaOH 2M (10 mL) and extracted with DCM (2 x 50 mL). Drying (Na₂SO₄) and silica gel flash chromatography (EtOAc/hexane 2:1 → 2:0) afforded 2-(4-(pent-4-ynyl)piperazin-1-yl)pyrimidine (0.95 g).

Step 2

- 20 Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 2-(4-(pent-4-ynyl)piperazin-1-yl)pyrimidine according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.61 min.

[0139] Example 35

5-(5-(1H-imidazol-1-yl)pent-1-ynyl)-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



~~PCT/US2006/011846~~
 Step 1: 1-(pent-4-ynyl)-1H-imidazole


A mixture of pent-4-ynyl methanesulfonate (1.06 g), imidazole (534 mg) and K₂CO₃ (3.66 g) in 2-butanone (11 mL) was heated to reflux overnight. Work-up (DCM, H₂O), drying (Na₂SO₄) and concentration gave 1-(pent-4-ynyl)-1H-imidazole (472 mg).

5 *Step 2*

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(pent-4-ynyl)-1H-imidazole according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.40 min.

[0140] Example 36


10 **4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl ethyl carbonate**



15 A solution of 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol (see example 2) (1.53 g) and NaH (95%, 0.38 g) in anhydrous DMA (25 mL) was treated with ethyl chloroformate (1.5 mL) at 0-23 °C for 0.5 h. Work-up and preparative HPLC gave the title compound. HPLC Rt = 5.65 min.

[0141] Example 37

Ethyl 4-chloro-5-(4-(ethoxycarbonyloxy)but-1-ynyl)-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-ylcarbamate




20

A suspension of 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol (see example 2) (0.73 g) in anhydrous pyridine (5 mL) and DCM (25 mL) was treated with ethyl chloroformate (1 mL) at rt for 30 min. The mixture was cooled to 0 °C, treated with addition EtOCOC₂H₅ (1 mL), and the temperature was
 25 allowed to slowly reach rt overnight. The reaction mixture was diluted with DCM (30 mL) and

~~PCT/US2006/011846~~
 washed sequentially with H₂O, NH₄OH 1M, and brine. The DCM was evaporated, but not the residual pyridine, Addition of MeOH (20 mL) induced crystallization of the product, as white needles. HPLC Rt = 6.28 min.

[0142] Example 38

5 **4-(4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl)-N-methylpiperazine-1-carboxamide**



Step 1: 1-(but-3-ynyl)piperazine

A solution of but-3-ynyl 4-methylbenzenesulfonate (2.0 mL) and piperazine (2.0 g) in EtOH (6 mL) was heated to reflux for 30 min. The mixture was concentrated, diluted with NaOH 2 M (8 mL) and extracted with Et₂O (50 mL). Evaporation of the organic layer gave a 2:1 mixture of mono and bis-alkylated piperazine (450 mg) which was discarded. The aqueous layer was further extracted with DCM (100 mL) to give of 1-(but-3-ynyl)piperazine (640 mg).

Step 2: 4-(but-3-ynyl)-N-methylpiperazine-1-carboxamide

15 A solution of 1-(but-3-ynyl)piperazine (445 mg) in THF (4 mL) was treated with 4-nitrophenyl carbonochloridate (649 mg) at rt for 5 min. A precipitate formed immediately, and the suspension was treated with Et₃N (0.45 mL) to ensure complete reaction. The suspension was diluted with H₂O (2 mL), MeOH (2 mL) and 40% aq. MeNH₂ (4.0 mL) and stirred at rt for 3 days. Extraction with DCM (50 mL), washing (NaOH 2M), drying (Na₂SO₄) and concentration
 20 gave 4-(but-3-ynyl)-N-methylpiperazine-1-carboxamide (370mg).


Step 3

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 4-(but-3-ynyl)-N-methylpiperazine-1-carboxamide according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.10 min

25 [0143] Example 39

5-(4-(1H-imidazol-1-yl)but-1-ynyl)-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine

PCT/US2006/11846

*Step 1: 1-(but-3-ynyl)-1H-imidazole*


A mixture of but-3-ynyl 4-methylbenzenesulfonate (1.04 g), imidazole (573 mg) and K₂CO₃ (3.88 g) in 2-butanone (10 mL) was heated to reflux overnight. Work-up (DCM, H₂O), drying (Na₂SO₄) and concentration gave 1-(but-3-ynyl)-1H-imidazole (501 mg).

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 1-(but-3-ynyl)-1H-imidazole according to the general procedure A gave the title compound., as a solid. HPLC Rt = 4.21 min.

[0144] Example 40

(S)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl 2-aminopropanoate




A solution of 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol (see example 2) (51.7 mg), N-Boc alanine (53.3 mg), DMAP (33.7 mg), and EDCI (54.9 mg) in anhydrous DMA (4 mL) was stirred at rt overnight. Work-up (EtOAc; H₂O) gave the crude Boc-protected intermediate. A solution of this intermediate (40 mg) in DCM (1.0 mL) was treated with TFA (0.2 mL) at rt for 5 min, and evaporated. The material was dissolved in MeOH (1 mL) and water (10 mL), and washed with Et₂O (10 mL). The organic layer was discarded. The aqueous layer was brought to pH >7 with sat. aq. NaHCO₃ and back-extracted with DCM. The title compound was isolated as a solid. HPLC Rt = 4.37 min.

[0145] Example 41

(R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl 2-amino-3-methylbutanoate


PCT/US2006/011846



This compound was obtained by the same procedure as used for example 40, using N-Boc valine in place of N-Boc alanine, to give the title compound as a solid. HPLC Rt = 4.58 min.

[0146] Example 42


- 5 **(R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl 2-amino-4-methylpentanoate**



This compound was obtained by the same procedure as used for example 40, using N-Boc isoleucine in place of N-Boc valine, to give the title compound as a solid. HPLC Rt = 4.76 min.

10 [0147] Example 43


- 4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl di-tert-butyl phosphate**



- 15 A solution of 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol (see example 2) (4.0 g) and 1-H-tetrazole (0.45 M in acetonitrile, 70 mL) in THF (160 mL) was treated with di-*tert*-butyl diisopropylphosphoramidite (16.4 mL) at rt for 2h. The solution was cooled to 0 °C and treated with 30% aqueous H₂O₂ (11 mL) for 30 min. Work-up (EtOAc, aq. NaHCO₃) and reverse-phase preparative HPLC gave the title compound, as a solid. HPLC Rt = 6.13 min.

PCT/US2006/11846
[0148] Example 44


4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl dihydrogen phosphate



- 5 A solution of 4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl di-tert-butyl phosphate (see example 43) in DCM was treated with TFA at rt for h. Evaporation gave the title compound as a TFA salt (5 g). The crude was dissolved in a mixture of Et₃N (0.7 mL and MeOH (60 mL), loaded on Dowex 50Wx2-400 (20 g, pre-washed with MeOH), and the resin was washed with MeOH (200 mL) to
- 10 remove the excess TFA. The desired phosphate was released from the resin with Et₃N:MeOH 1:10 (200 mL) The solution was concentrated to give the phosphate as an oily triethylamine salt (2.75 g). Crystallization was induced with EtOH (60 mL). The mixture was left at rt for 1.5 h and at -20 °C overnight to give a first crop of crystals (0.64 g). The mixture was concentrated and the crystallization procedure was repeated to give a second crop (0.74 g). The combined
- 15 triethylamine salts (1274 mg) were dissolved in MeOH (50 mL) with the help of Et₃N (326 μL) and treated with a 1.0 M solution of NaOH in MeOH (4.68 mL). Evaporation and drying on high vacuum overnight gave the title compound as a solid sodium salt. HPLC Rt = 4.21 min.

[0149] Example 45

- 2-((2-amino-4-chloro-5-(4-hydroxybut-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide**
- 20



Step 1: 2-((4-chloro-5-iodo-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide

- A mixture of N-(4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-2-yl)pivalamide (235 mg), 2-(chloromethyl)-4-methoxy-3,5-dimethylpyridine 1-oxide (129 mg), and K₂CO₃ (413 mg) in DMF (6.0 mL) was stirred at rt for 3 days, and diluted with water. The precipitate was collected
- 25

by filtration and washed with IPA to give 2-((4-chloro-5-iodo-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide (124 mg).

Solid. HPLC Rt = 6.73 min

Step 2: 2-((4-chloro-5-(4-hydroxybut-1-ynyl)-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide

Sonogashira coupling of 2-((4-chloro-5-iodo-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide (335 mg) with but-3-yn-1-ol (200 μ L) according to the general procedure A gave crude 2-((4-chloro-5-(4-hydroxybut-1-ynyl)-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide (291 mg).


Solid. HPLC Rt = 5.82 min.

Step 3

Crude 2-((4-chloro-5-(4-hydroxybut-1-ynyl)-2-pivalamido-7H-pyrrolo[2,3-d]pyrimidin-7-yl)methyl)-4-methoxy-3,5-dimethylpyridine 1-oxide (291 mg) was dissolved in 10 mL EtOH. An aliquot of this solution (2 mL) was further diluted with EtOH (5 mL) and treated with ZnCl₂ (103 mg) at 120 °C for 20 min in a microwave oven. Reverse-phase preparative HPLC gave the title compound (2.2 mg), as a solid. HPLC Rt = 4.88 min.

[0150] Example 46

5-(but-3-en-1-ynyl)-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine




A mixture of 4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl 4-methylbenzenesulfonate (516 mg) and CsF (2.0 g) in DMF (7.0 mL) was heated to 70 °C for 1 h. Work-up and flash chromatography (1 $\rightarrow$ 3% MeOH in DCM) gave the title compound (280 mg), as a pale yellow oil. HPLC Rt = 5.90 min.

[0151] Example 47

5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-ynyl methanesulfonate


PCT/US06/11846



Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with pent-4-ynyl methanesulfonate according to the general procedure A gave the title compound, as a solid. HPLC Rt = 5.31 min.

5 [0152] Example 48


4-chloro-5-(5-fluoropent-1-ynyl)-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



10 A solution of KF (77.8 mg), Kryptofix 222 (Aldrich, 839 mg) and water (10 mL) was evaporated to dryness, and dried first azeotropically with anh. CH₃CN (3 x 20 mL), then on high vacuum at 50 °C overnight. The solid was redissolved in anhydrous acetonitrile (13 mL) to give a 0.1 M fluoride solution. Treatment of 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-ynyl methanesulfonate (see example 47) (42 mg) with this solution (3 mL) at 60-70 °C for 2h, work-up and reverse-phase preparative HPLC
15 gave the title compound, as an oil. HPLC Rt = 5.88 min.

[0153] Example 49

(S)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-yn-1,2-diol




20 Treatment of 5-(but-3-en-1-ynyl)-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine (see example 46) (35 mg) with AD-mix- α (Aldrich, 540 mg) in

~~PCT/US2006/011846~~
 t-BuOH:THF:water 1:1:1 (6mL) at rt overnight, followed by work-up and reverse-phase
 preparative HPLC gave the title compound (10 mg), as a solid. HPLC Rt = 4.13 min

[0154] Example 50


(R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-yne-1,2-diol



Treatment of 5-(but-3-en-1-ynyl)-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine (see example 46) (35 mg) with AD-mix- β (Aldrich, 540 mg) in t-BuOH:THF:water 1:1:1 (6mL) at rt overnight, followed by work-up and reverse-phase
 preparative HPLC gave the title compound (10 mg), as a solid. HPLC Rt = 4.13 min

[0155] Example 51


4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(3-methylbut-3-en-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine



Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 2-methylbut-1-en-3-yne according to the general procedure A gave the title compound, as a solid. HPLC Rt = 6.00 min.

[0156] Example 52


(S)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylbut-3-yne-1,2-diol



~~PCT/US2006/011846~~
 Treatment of 4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(3-methylbut-3-en-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine (see example 51) (263 mg) with AD-mix- α (Aldrich, 1.41 g) in t-BuOH:water 1:1: (7 mL) at rt for 3 days, followed by quench (NaHSO₃, 1.6 g), work-up, and preparative HPLC gave the title compound (69 mg), as a solid. HPLC Rt = 4.36 min

[0157] Example 53


(R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylbut-3-yn-1,2-diol



- 10 Treatment of 4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-5-(3-methylbut-3-en-1-ynyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine (see example 51) (357 mg) with AD-mix- β (Aldrich, 1.57 g) in t-BuOH:water 1:1: (10 mL) at rt for 18 h, followed by quench (NaHSO₃, 1.2 g), work-up, and silica gel flash chromatography (1 $\rightarrow$ 10 % MeOH in DCM) gave the title compound (61 mg) and recovered starting material (263 mg).
- 15 Solid. HPLC Rt = 4.36 min.

[0158] Example 54


(R)-((S)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-hydroxy-2-methylbut-3-ynyl) 3,3,3-trifluoro-2-methoxy-2-phenylpropanoate



- 20 A mixture of (S)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylbut-3-yn-1,2-diol (see example 52) (18 mg), (S)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (45 μ L), and Et₃N (50 μ L) in THF (3.0 mL) was heated to 70 $^{\circ}$ C for 2 h, and then to 40 $^{\circ}$ C overnight. Preparative TLC (EtOAc/DCM 25 25:65) gave the title compound, as a solid. HPLC Rt = 6.45 min.

PCT/US2006/011846
[0159] Example 55

(R)-((R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-hydroxy-2-methylbut-3-ynyl) 3,3,3-trifluoro-2-methoxy-2-phenylpropanoate




5

A mixture of (R)-4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylbut-3-yne-1,2-diol (see example 53) (22 mg), (S)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (45 μ L), and Et₃N (50 μ L) in THF (3.0 mL) was heated to 40 °C overnight. Preparative TLC (EtOAc/DCM 25:65) gave the title compound, as a solid. HPLC Rt = 6.44 min.

10

[0160] Example 56

2-((2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)ethynyl)-2-hydroxypropane-1,3-diyl diacetate



15 *Step 1: 2-ethynyl-2-hydroxypropane-1,3-diyl diacetate*

An ethynylmagnesium bromide solution (0.5 M in THF) was cooled to -78 °C under nitrogen. A solution of 1,3-diacetoxyacetone (0.5 g) in THF (3.0 mL) was added, and the resulting mixture was stirred for 30 min at -78 °C, then at 0 °C for 5 min, quenched with 1.0 N aq. HCl (3 mL) and extracted in EtOAc. Evaporation gave crude 2-ethynyl-2-hydroxypropane-1,3-diyl diacetate, contaminated with about 50 mol% of 1,3-diacetoxyacetone. The crude material was used without further purification.

20


Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with crude 2-ethynyl-2-hydroxypropane-1,3-diyl diacetate according to the general procedure A gave the title compound, as an oil. HPLC Rt = 4.89 min.

25

PCT/US2006/11846
[0161] Example 57


2-((2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)ethynyl)propane-1,2,3-triol



- 5 A solution of 2-((2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)ethynyl)-2-hydroxypropane-1,3-diyl diacetate (see example 56) (60 mg) and Et₃N (1 mL) in THF (1 mL) and MeOH (6 mL) was heated in a microwave oven to 150 °C for 20 min. Concentration and preparative HPLC gave the title compound, as a solid. HPLC Rt = 3.90 min.

10 **[0162] Example 58**


4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylbut-3-yn-2-ol



- 15 Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with 2-methylbut-3-yn-2-ol according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.81 min.

[0163] Example 59


5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-yn-2-ol



~~PCT/US2006/011846~~
 Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with pent-4-yn-2-ol according to the general procedure A gave the title compound, as a solid. HPLC Rt = 4.29 min

[0164] Example 60

- 5 **4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl acetate**



Step 1: But-3-ynyl acetate


- 10 A solution of but-3-yn-1-ol (3.0 mL) and Et₃N (6.6 mL) in 1,2-dichloroethane (10 mL) was treated with AcCl (3.1 mL) with caution (very exothermic). Work-up gave but-3-ynyl acetate.

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with but-3-ynyl acetate according to the general procedure A gave the title compound, as a solid. HPLC Rt = 5.31 min.

- 15 [0165] Example 61


4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl sulfamate



- 20 A mixture of formic acid (0.463 g) and N,N-dimethylacetamide (1 drop) was added to a solution of chlorosulfonyl isocyanate (1.31 g) in DCM (9 mL), and heated to reflux for 3 h. The resulting solution was added to a solution of 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol (see example 2) (200 mg) in DMA (10 mL) and stirred at rt for 30 min. Work-up and reverse-phase preparative HPLC gave the title compound, as a solid. HPLC Rt = 4.96 min.

PCT/US06/11846
[0166] Example 62


5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)pent-4-ynyl sulfamate



- 5 The title compound was obtained by the same procedure as in example 61, carrying out the reaction on 5-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethylpyridin-2-yl)methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-pent-4-yn-1-ol (see example 2) .Solid. HPLC Rt = 4.77 min.

[0167] Example 63

- 10 **4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl hydrogen sulfate**



Step 1: But-3-ynyl hydrogen sulfate, pyridine salt

- 15 But-3-yn-1-ol (2.1 g) in DCM (20 mL) was treated with sulfur trioxide-pyridine complex (4.77 g) overnight. The solid was filtered off and the reaction mixture was concentrated and used without further purification.

Step 2


Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with the pyridine salt of but-3-ynyl hydrogen sulfate according to the general procedure A gave the title compound.

- 20 Solid. HPLC Rt = 4.24 min.

[0168] Example 64

4-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)but-3-ynyl bis(2,2,2-trichloroethyl) phosphate

PCT/US2006/011846



Step 1: But-3-ynyl bis(2,2,2-trichloroethyl) phosphate

A solution of but-3-yn-1-ol (0.413 g) in DCE (20 mL) was treated with Et₃N (2 mL) and bis(2,2,2-trichloroethyl)phosphorochloride (2.27 g) at 50 °C overnight. Work-up gave crude but-3-ynyl bis(2,2,2-trichloroethyl) phosphate as an oil that was used without further purification.

Step 2

Sonogashira coupling of 4-chloro-5-iodo-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine with but-3-ynyl bis(2,2,2-trichloroethyl) phosphate according to the general procedure I gave the title compound.

Solid. HPLC Rt = 6.69 min.

[0169] Example 65

rHSP90 Competitive Binding Assay

Five microgram of purified rHSP90 protein (Stressgen, BC, Canada, #SPP-770) in phosphate buffered saline (PBS) was coated on 96 well plates by incubating overnight at 4°C. Unbound protein was removed and the coated wells were washed twice with 200 µL PBS. DMSO controls (considered as untreated samples) or test compounds were then added at 100-30-10-3-1-0.3 µM dilutions (in PBS), the plates mixed for 30 seconds on the plate shaker, and then incubated for 60 min. at 37 °C. The wells were washed twice with 200 µL PBS, and 10 µM biotinylated-geldanamycin (biotin-GM) was added and incubated for 60 min. at 37 °C. The wells were washed again twice with 200 µL PBS, before the addition of 20 µg/mL streptavidin-phycoerythrin (streptavidin-PE) (Molecular Probes, Eugene, OR) and incubation for 60 min. at 37°C. The wells were washed again twice with 200 µL PBS. Relative fluorescence units (RFU) was measured using a SpectraMax Gemini XS Spectrofluorometer (Molecular Devices, Sunnyvale, CA) with an excitation at 485 nm and emission at 580 nm; data was acquired using SOFTmax[®] PRO software (Molecular Devices Corporation, Sunnyvale, CA). The background was defined as the RFU generated from wells that were not coated with HSP90 but were treated with the biotin-GM and streptavidin-PE. The background measurements were subtracted from each sample treated with biotin-GM and streptavidin-PE measurements before other

~~PCT/US2006/011846~~
computation. Percent inhibition of binding for each sample was calculated from the background subtracted values as follows:

$$\% \text{ binding inhibition} = [(\text{RFU untreated} - \text{RFU treated}) / \text{RFU untreated}] \times 100.$$

[0170] Example 66

5 Cell Lysate Binding Assay

MCF7 breast carcinoma cell lysates were prepared by douncing in lysing buffer (20 mM HEPES, pH 7.3, 1 mM EDTA, 5 mM MgCl₂, 100 mM KCl), and then incubated with or without test compound for 30 mins at 4 °C, followed by incubation with biotin-GM linked to BioMagTM streptavidin magnetic beads (Qiagen) for 1 hr at 4 °C. The tubes were placed on a magnetic rack, and the unbound supernatant removed. The magnetic beads were washed three times in lysis
10 buffer and boiled for 5 mins at 95 °C in SDS-PAGE sample buffer. Samples were analyzed on SDS protein gels, and Western Blots were done for rHSP90. Bands in the Western Blots were quantitated using the Bio-rad Fluor-S MultiImager, and the % inhibition of binding of rHSP90 to the biotin-GM was calculated. The lysate binding ability of selected compounds of the invention
15 based on the above assay is summarized in Table 3. The IC₅₀ reported is the concentration of test compound needed to achieve 50% inhibition of the biotin-GM binding to rHSP90 in the MCF7 cell lysates.

[0171] Example 67

Her2 Degradation Assay

20 Many cancers are associated with the over expression Her2 protein. It has been shown that compounds able to diminish Her2 levels show good promise as anti-cancer agents. Thus a good in vitro assay for identifying compounds of the present invention, which are likely to demonstrate anti-cancer activity, is the Her2 degradation assay, as described below:
MCF7 breast carcinoma cells (ATCC) were grown in Dulbecco's modified Eagle's medium
25 (DMEM) containing 10% fetal bovine serum (FBS) and 10 mM HEPES, and plated in 24 well plates (50% confluent). Twenty-four hrs later (cells are 65-70% confluent), test compounds were added and incubated overnight for 16 h. For the less potent compounds, the amounts added were 100 μM, 30 μM, 10 μM and 1 μM, and for more potent compounds, the amounts added were 1 μM, 0.3 μM, 0.1 μM, 0.03 μM, 0.01 μM and 0.003 μM. The wells were washed with 1
30 mL phosphate buffered saline (PBS), and 200 μL trypsin was added to each well. After trypsinization was complete, 50 μL of FBS was added to each well. Then 200 μL cells were transferred to 96 well plates. The cells were pipetted up and down to obtain a single cell suspension. The plates were centrifuged at 2,500 rpm for 1 min using a Sorvall Legend RTTM tabletop centrifuge (Kendro Laboratory Products, Asheville, NC). The cells were then washed
35 once in PBS containing 0.2% BSA and 0.2% sodium azide (BA buffer). Phycoerythrin (PE)

PCT/US2006/011846
conjugated anti HER2/Neu antibody (Becton Dickinson, #340552), or PE conjugated anti-keyhole limpet hemocyanin [KLH] (Becton Dickinson, #340761) control antibody was added at a dilution of 1:20 and 1:40 respectively (final concentration was 1 µg/mL) and the cells were pipeted up and down to form a single cell suspension, and incubated for 15 mins. The cells were washed twice with 200 µL BA buffer, and resuspended in 200 µL BA buffer, and transferred to FACSCAN tubes with an additional 250 µL BA buffer. Samples were analyzed using a FACSCalibur™ flow cytometer (Becton Dickinson, San Jose, CA) equipped with Argon-ion laser that emits 15 mW of 488 nm light for excitation of the PE fluorochrome. 10,000 events were collected per sample. A fluorescence histogram was generated and the mean fluorescence intensity (MFI) of each sample was determined using Cellquest software. The background was defined as the MFI generated from cells incubated with control IgG-PE, and was subtracted from each sample stained with the HER2/Neu antibody. Cells incubated with DMSO were used as untreated controls since the compounds were resuspended in DMSO. Percent degradation of Her2 was calculated as follows:

$$\% \text{ Her2 degraded} = [(MFI \text{ untreated cells} - MFI \text{ treated cells}) / MFI \text{ untreated cell}] \times 100.$$

The Her2 degradation ability of selected compounds of the invention based on this assay is summarized in Table 3. IC₅₀ is defined as the concentration at which there was 50% degradation of the HER2/Neu protein.

[0172] Example 68

MTS Assay


3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium assays measure the cytotoxicity of geldanamycin derivatives. MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) is a tetrazolium dye that is converted to a formazan product by dehydrogenase enzymes of metabolically active cells (Corey, A. *et al.* "Use of an aqueous soluble tetrazolium/formazan assay for cell growth assays in culture," *Cancer Commun.* **1991**, 3, 207-212). Cells were seeded in 96 well plates at 2000 cells/well and allowed to adhere overnight in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum. The final culture volume was 100 µL. Viable cell number was determined by using the Celltiter 96 Aqueous Non-radioactive Cell Proliferation Assay (Promega, Madison WI). The MTS /PMS (phenazine methosulfate) solution was mixed at a ratio of 20:1, and 20 µL was added per well to 100 µL of culture medium. After 2-4 hours, the formation of the formazan product was measured at 490 nm absorbance using a multiwell plate spectrophotometer. Background was determined by measuring the Abs 490 nm of cell culture medium and MTS-PMS in the absence of cells and was subtracted from all values. Percent viable cells was calculated as follows:

~~PCT/US2006/011846~~

$$\% \text{ viable cells} = (\text{Abs at 490 nm treated cells} / \text{Abs at 490 nm untreated cells}) \times 100$$

The effect of selected compounds of the invention on MCF7 breast carcinoma cells according to the MTS assay is summarized in table 3. IC₅₀ was defined as the concentration of the compound which gave rise to 50% reduction in viable cell number.

5 [0173] TABLE 3 Biological Activities of Selected Compounds of the Invention



Sample. No	Example	R ³	IC ₅₀ (μM)			
			Her2	Lysate binding	MTS BT474	MTS MCF7
1	1	-CH ₂ OH	0.012	ND	0.02	0.13
2	10	-CH ₂ CH ₂ CON(Et) ₂	0.03	0.018	0.05	0.7
3	2	-CH ₂ CH ₂ OH	0.006	0.003	0.006	0.013
4	3	-CH ₂ CH ₂ CH ₂ OH	0.006	0.003	0.008	0.013
5	5	-CH ₂ N(iPr) ₂	0.055	ND	0.08	0.1
6	11	-CH ₂ CH ₂ C(O)-N-Me	0.018	ND	0.02	0.1
7	12	-CH ₂ CH ₂ CONH ₂	0.016	0.012	0.02	0.03
8	16	-CH ₂ CH ₂ CH ₂ NH ₂	0.028	ND	>10	0.09
9	15	-CH ₂ CH ₂ CH ₂ NHBOC	0.009	ND	1.0	0.06
10	17	-CH ₂ CH ₂ CH ₂ -N>	0.007	ND	0.1	0.01
11	18	-CH ₂ CH ₂ -N>	0.01	ND	1.0	0.05
12	19	-CH ₂ CH ₂ CH ₂ -N>	0.015	ND	1.0	0.1
13	20	-CH ₂ CH ₂ NHtBu	0.012	ND	23	130
14	21	-CH ₂ CH ₂ CH ₂ NHt-Bu	0.015	ND	30	230
15	7	-CH ₂ CH ₂ CH ₂ CH ₂ OH	0.009	ND	20	200
16	22	-CH ₂ O(CO)CH ₂ NMe ₂	0.013	ND	20	200

ND, not determined.

[0174] Example 69

10 In Vivo Mouse Tumor Studies

Six to 8 week old Balb/C and nu/nu athymic female mice were obtained from Harlan Sprague Dawley, (Indianapolis, IN). The mice were maintained in sterilized filter topped cages or ventilated caging in a room with a 12 hour light/dark cycle. Irradiated pelleted food (Harlan Teklad #7912) and autoclaved deionized water were provided ad libitum. Animals were

~~PCT/US2006/011846~~
 identified by the use of individually numbered ear tags. Experiments were carried out under institutional guidelines for the proper and human use of animals in research established by the Institute for Laboratory Animal Research (ILAR).

Tumor fragments (approximately 2mm³) or 5 x 10⁶ tumor cells were inoculated subcutaneously in the right or left flank of the animal. Mice with established tumors (50-200mm³) were selected for study (n=7-10/treatment group). Tumor dimensions were measured using calipers and tumor volumes were calculated using the equation for an ellipsoid sphere $(l \times w^2)/2 = \text{mm}^3$, where l and w refer to the larger and smaller dimensions collected at each measurement.

Mice were followed until tumor volumes in the control group reached approximately 1000mm³ and were sacrificed by CO₂ euthanasia. The mean tumor volumes of each group were calculated. The change in mean treated tumor volume was divided by the change in mean control tumor volume, multiplied by 100 and subtracted from 100% to give the tumor growth inhibition for each group. Statistical analysis was performed using the standard T-test and using GraphPad Prism© Software.

[0175] Example 70

N87 Gastric Carcinoma Xenograft

Tumors were established in mice, by inoculation of human N87 stomach cancer cells, according to example 69. Four treatment groups (n=7-10/treatment group) were established for study:

I. 4-[2-Amino-4-chloro-7-(4-methoxy-3,5-dimethyl-pyridin-2-ylmethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-but-3-yn-1-ol, prepared as described in example 2 above, was administered at 8mg/kg po

II. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 8mg/kg po

III. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 16mg/kg po

IV. Vehicle alone was administered to the control group.

The results are shown in figure 1, a plot of tumor volume (mm³) against time (days).

[0176] Example 71

NCI295 Adrenocortical Carcinoma Xenograft

Tumors were established in mice, by inoculation of NCI295 adrenocortical carcinoma cells, according to example 69. Two treatment groups (n=7-10/treatment group) were established for study:

PCT/US2006/011846

I. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 8mg/kg po

II. Vehicle alone was administered to the control group.

5 The results are shown in figure 2, a plot of tumor volume (mm³) against time (days).

[0177] Example 72

SK-MEL-28 Melanoma Xenograft

Tumors were established in mice, by inoculation of SK-MEL-28 melanoma cells, according to example 69. Two treatment groups (n=7-10/treatment group) were established for study:

10 I. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 8mg/kg po

II. Vehicle alone was administered to the control group.

The results are shown in figure 3, a plot of tumor volume (mm³) against time (days).

15 [0178] Example 73

HT29 Colon Carcinoma Xenograft

Tumors were established in mice, by inoculation of HT29 Colon Carcinoma cells, according to example 69. Three treatment groups (n=7-10/treatment group) were established for study:

20 I. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 4mg/kg po

II. 5-(2-amino-4-chloro-7-((4-methoxy-3,5-dimethylpyridin-2-yl)methyl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)-2-methylpent-4-yn-2-ol, prepared as described in example 23 above, was administered at 8mg/kg po

25 III. Vehicle alone was administered to the control group.

The results are shown in figure 4, a plot of tumor volume (mm³) against time (days).

[0179] Example 74

Pharmacodynamic effects

Preparation of buffers and reagents

30 The following reagents and buffers are prepared ahead of the experiments.

Western Lysis Buffer (WLS) contains 10mM HEPES, 42mM KCl, 5mM MgCl₂, 0.1mM EDTA, 0.1mM EGTA, 1mM DTT, 1mM PMSF, 1ug/ml Pepstatin A, 1ug/ml Leupeptin, 5ug/ml Aprotinin, and 1% Triton X-100. Aliquot and store at -20°C.

~~PCT/US2006/011846~~

5X Western Sample Buffer (5X WSB) contains 20ml glycerol, 4ml β -mercaptoethanol, 5g SDS, 12.5ml 1M Tris pH 6.8, and 50mg bromophenol blue. Add water to a final volume of 50 ml. Aliquot and store at -20°C .

Western Transfer Buffer contains 23.3g Tris base, 116g glycine and 1.6L methanol. Add water to a final volume of 8L and store at 4°C .

TBST (1X) contains 10mM Tris pH8.0, 150mM NaCl and 0.1% Tweene 20.

Blocking solution contains 5% nonfat dry milk in 1X TBST. Keep at 4°C .

Processing of tumor samples:

Snap-frozen N87 gastric carcinoma tumors are transferred from liquid nitrogen into -80°C freezer. WLB is supplemented with protease inhibitor cocktail (stock at 100X) on ice. Each tumor is thawed on ice and transferred onto the lid of a Petri dish. It is covered with 50ul of WLB and is dissected into smaller pieces with disposable scalpels. Any residual skin attached to the tumor is removed. The tumor pieces are then chopped further down, and transferred into 300-500ul of ice-cold WLB. The minced sample is sonicated at setting 3 on Fisher Scientific's Sonic Dismembrator 550 until no more solid pieces can be broken down. The suspension is then centrifuged at 15,000g, 4°C for 5 minutes. The supernatant is collected into a clean Eppendorf tube on ice as lysate. 2ul of the lysate is used for total protein quantification by following directions in the BCA Protein Assay kit. The rest of the lysate is snap-frozen in liquid nitrogen while the BCA assay is in process. The total protein concentration in each lysate is calculated upon completion of BCA assay. The lysates are thawed in a water bath, and their total protein concentrations adjusted to 4-10mg/ml using 5X WSB (to a final of 20% of the total volume) and WLB (if necessary). The adjusted lysates are boiled at 95°C for 5 minutes and cooled to room temperature. At this stage, they are frozen at -20°C for future Western blotting analysis.

Western blotting analysis:

The quantified and adjusted lysates from tumor or spleen samples are thawed in a water bath, and loaded at equal total protein amount onto 4-12% Tris-glycine precast gels together with biotinylated protein marker. Electrophoresis is carried out at 140V for 1.5 hours. The separated proteins in the gels are transferred onto PVDF membranes at 100V for about 1 hour in Western transfer buffer. The blots are incubated in blocking buffer at room temperature for 1 hour or at 4°C overnight with gentle rocking. Primary antibodies against various HSP90 client proteins of interest are applied at room temperature for 1 hour with gentle rocking. Excess antibodies are washed off with six 5- minute washes in TBST. The blots are then incubated in HRP-conjugated secondary antibodies and streptavidin-HRP conjugate at room temperature for 1 hour with gentle rocking. Excess secondary antibodies and conjugate are washed off by six 5 minute washes in

~~PCT/US06/31346~~
TBST. The blots are then developed using Pierce's SuperSignal West Femto chemilluminescent substrate by mixing freshly prepared equal volumes of luminal enhancer and peroxide buffer and adding the mixture onto the blots one at a time. The protein bands can be visualized on Bio-Rad's fluor-S Max2 MultiImager using Bio-Rad's Quantity One software.

5 The results of the Western Blott analysis are shown in figure 5.

[0180] The foregoing examples are not limiting and are merely illustrative of various aspects and embodiments of the present invention. All documents cited herein are indicative of the levels of skill in the art to which the invention pertains and are incorporated by reference herein in their entireties. None, however, is admitted to be prior art.

10 One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The methods and compositions described illustrate preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Certain modifications and other uses will occur to those skilled in the art, and are encompassed within the spirit of the invention,
15 as defined by the scope of the claims.

The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations which is not specifically disclosed herein. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any
20 equivalents of the features shown and described, or portions thereof. It is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments, optional features, modifications and variations of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are
25 considered to be within the scope of this invention as defined by the description and the appended claims.


In addition, where features or aspects of the invention are described in terms of Markush groups or other grouping of alternatives, *e.g.*, genuses, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members
30 of the Markush group or subgenus, and exclusions of individual members as appropriate, *e.g.*, by proviso.

Other embodiments are within the following claims.

PCT/US2006/11846

[0181] What is claimed is:

1. A compound represented by Formula I, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof:



I

wherein:

R^0 is selected from the group consisting of hydrogen, halogen, lower alkyl, -CN, -SR⁸, -OR⁸, and -NHR⁸;

R^1 is selected from the group consisting of halogen, -OR¹¹, -SR¹¹ and lower alkyl;

R^2 is -NHR⁸;

R^3 is selected from the group consisting of hydrogen, -CN, -C(O)OH, -OR¹¹, -SR¹¹, -C(O)R⁹, -NR⁸R¹⁰, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower alkylsilyl, aryl, heteroaryl, alicyclyl and heterocyclyl, all optionally substituted, wherein:

the aryl, heteroaryl, alicyclyl and heterocyclyl groups are mono-, bi- or tri- cyclic;

R^8 and R^{10} taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

the optional substituents on R^3 are selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R⁸, lower aryl, heteroaryl, alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, amino, alkylamino, dialkylamino, arylalkylamino, diarylamino, heteroarylamino, diheteroarylamino, arylheteroarylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidiny, pyridiny, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl, phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R^8 and R^8 taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

PCT/US2006/011846

R⁴ is selected from the group consisting of optionally substituted lower alkylene, -

C(R¹²)₂-,

-C(O)-, -C(S)-, -S(O)- and -SO₂-;

R⁵ is selected from the group consisting of aryl, heteroaryl, alicyclyl and heterocyclyl,

5 wherein:

the aryl group is substituted with 2 to 5 substituents;

the heteroaryl group is substituted with 2 to 5 substituents;

the alicyclyl group is substituted with 3 to 5 substituents;

the heterocyclyl group is substituted with 3 to 5 substituents;

10 the substituents on R⁵ are selected from the group consisting of halogen, lower alkyl, lower alkenyl, lower alkynyl, -CN, -C(O)OH, -NO₂, -SR⁸, -OR⁸, -C(O)R⁹, -NR⁸R¹⁰, lower aryl, lower heteroaryl, lower alicyclyl, lower heterocyclyl, arylalkyl, heteroarylalkyl, thioalkyl, amino, alkylamino, dialkylamino, arylalkylamino, oxo, perhaloalkyl, perhaloalkoxy, perhaloacyl, guanidinyl, pyridinyl, thiophenyl, furanyl, indolyl, indazolyl, phosphonyl, phosphatidyl, 15 phosphoramidyl, sulfanyl, sulfinyl, sulfonyl, sulphonamidyl, carbamyl, uryl, thiouryl and thioamidyl, wherein

R⁸ and R¹⁰ taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

20 R⁸ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower heteroaryl and -C(O)R⁹;

R⁹ is selected from the group consisting of H, lower alkyl, lower alkenyl, lower alkynyl, lower aryl, lower heteroaryl, -NR¹⁰R¹⁰ and -OR¹¹, wherein

25 R¹⁰ and R¹⁰ taken together with the N atom to which they are attached optionally form an optionally substituted ring comprising 3-7 ring atoms, wherein, in addition to said N atom, 0-3 of the ring atoms are heteroatoms selected from the group consisting of O, S and N;

R¹⁰ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower heteroalkyl, lower heteroalkenyl, lower heteroalkynyl, lower aryl, lower 30 heteroaryl and -C(O)R¹¹;

R¹¹ is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower aryl and lower heteroaryl; and

R¹² is selected from the group consisting of hydrogen and lower alkyl.

2. The compound of claim 1, or a polymorph, solvate, ester, tautomer, enantiomer, 35 pharmaceutically acceptable salt or prodrug thereof, wherein

~~PCT/US2006/011846~~
~~R¹ is hydrogen, halogen or -CN.~~

3. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R^1 is halogen.

5 4. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R^2 is $-NH_2$ or $-NH-C(O)R^9$.

5. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

10 R^3 is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, aryl, heteroaryl, alicyclyl, heterocyclyl, $-CN$ and $-C(O)R^9$, all optionally substituted.

6. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

15 R^3 is selected from the group consisting of lower alkyl, aryl, heteroaryl, $-CN$ and $-C(O)R^9$, all optionally substituted.

7. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is hydrogen.

20 8. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is optionally substituted phenyl or pyridinyl.

9. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

25 R^3 is $-(CH_2)_nOH$, where $n=1-3$.

10. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is $-(CH_2)_mC(R^{12})_2(CH_2)_nOH$, wherein

$m=0-2$;

$n=1-2$; and

each R^{12} is independently hydrogen or lower alkyl.

11. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein


R^3 is $-(CH_2)_nNR^8R^8$, wherein

$n=1-3$;

PCT/US2006/011846

each R^8 is independently hydrogen or lower alkyl, or

NR^8R^8 is an optionally substituted



12. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
5 pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is $-(CH_2)_nC(O)NR^{10}R^{10}$, wherein

$n=1-3$;

each R^{10} is independently hydrogen or $-C(O)R^{11}$, or

$NR^{10}R^{10}$ is an optionally substituted .

- 10 13. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is $-(CH_2)_nC(O)NR^{10}R^{10}$, where

$n=1-3$; and

each R^{10} is independently hydrogen or $-C(O)R^{11}$.

- 15 14. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is $-(CH_2)_nC(O)R^9$, wherein $n=1-3$.

15. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
pharmaceutically acceptable salt or prodrug thereof, wherein

20 R^3 is optionally substituted lower alkyl and the substituent on said lower alkyl is selected
from the group consisting of $-SR^8$, $-OR^8$, $-C(O)R^9$ and $-NR^8R^{10}$.

16. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is substituted lower alkyl.

- 25 17. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
pharmaceutically acceptable salt or prodrug thereof, wherein

R^3 is substituted lower alkyl, and the substituent on said lower alkyl is phosphonyl or
phosphatidyl.

18. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer,
30 pharmaceutically acceptable salt or prodrug thereof, wherein

R^4 is $-CH_2-$.

~~PCT/US2006/11846~~

19. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein each of said aryl, heteroaryl, alicyclyl or heterocyclyl group of R⁵ is monocyclic or bicyclic.
20. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein R⁵ is substituted aryl or heteroaryl and the substituents on said aryl or heteroaryl are selected from the group consisting of halogen, lower alkoxy, lower alkyl, thioalkyl, amino and dialkylamino.
21. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein R⁵ is substituted aryl or heteroaryl and the substituents on said aryl or heteroaryl are selected from the group consisting of halogen, lower alkyl or lower alkoxy.
22. The compound of claim 1 or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein:
R⁰ is hydrogen, halogen, -SH, -OH, or -CN;
R¹ is halogen; and
R² is -NH₂ or -NH-C(O)R⁹.
23. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein:
R¹ is chloro or bromo;
R² is -NH₂ or -NH-C(O)R⁹; and
R³ is lower alkyl, lower alkenyl, or lower alkynyl, lower perhaloalkyl, lower aryl, or lower heteroaryl, all optionally substituted, wherein the optional substituents on R³ are selected from the group consisting of hydrogen, -OR⁸, -NR⁸R⁸ and -C(O)R⁹.
24. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein:
R⁰ is hydrogen, halogen or -CN;
R² is -NH₂ or -NH-C(O)R⁹; and
R⁴ is -CH₂-.
25. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein:
R⁰ is hydrogen, halogen, -SH, -OH or -CN;
R¹ is halogen;

PCT/US2006/011846

R³ is hydrogen, -OR¹, -SR¹, -NR⁸R¹⁰, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, lower aryl or lower heteroaryl;

R⁴ is -CH₂-; and

R⁵ is optionally substituted aryl or heteroaryl, wherein each of said aryl and heteroaryl is monocyclic or bicyclic.

26. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R⁰ is hydrogen, halogen or -CN;

R¹ is halogen;

10 R³ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, lower alkynyl, lower perhaloalkyl, aryl, heteroaryl, alicyclyl and heterocyclyl, all optionally substituted;

R⁴ is -CHR¹²-; and

R⁵ is optionally substituted aryl or heteroaryl.

15 27. The compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R⁰ is hydrogen;

R¹ is halogen;

R² is -NH₂;

20 R³ is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, all optionally substituted;

R⁴ is -CH₂-; and

R⁵ is optionally substituted aryl or heteroaryl.

25 28. The compound of claim 25, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R¹ is chloro or bromo;

R² is -NH₂; and

R⁵ is a phenyl having at least three substituents.

30 29. The compound of claim 25, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein

R¹ is chloro or bromo;

R² is -NH₂; and

R⁵ is a pyridyl having at least two substituents.

35 30. The compound of claim 25, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, wherein


PCT/US2006/011846
 R¹ is chloro or bromo;

R² is -NH₂; and


R⁵ is 1-oxy-pyridyl (N-oxy-pyridyl) having at least two substituents.

31. A compound, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically


5 acceptable salt or prodrug thereof, selected from the group:




PCT/US2006/011846




PCT/US2006/011846



PCT/US2006/011846



PCT/US2006/011846







5

~~PCT/US2006/011846~~

33. A method of treating an individual having an HSP90 mediated disorder comprising administering to said individual a pharmaceutical composition comprising a pharmaceutically effective amount of the compound, polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug of claim 1.
34. The method of claim 33, wherein the HSP90 mediated disorder is selected from the group consisting of inflammatory diseases, infections, autoimmune disorders, stroke, ischemia, cardiac disorders, neurological disorders, fibrogenetic disorders, proliferative disorders, tumors, leukemias, neoplasms, cancers, carcinomas, metabolic diseases, and malignant diseases.
35. The method of claim 34, wherein the fibrogenetic disorder is further selected from the group consisting of scleroderma, polymyositis, systemic lupus, rheumatoid arthritis, liver cirrhosis, keloid formation, interstitial nephritis and pulmonary fibrosis.
36. The method of claim 32, further comprising administering at least one therapeutic agent selected from the group consisting of cytotoxic agents, anti-angiogenesis agents and anti-neoplastic agents.
37. The method of claim 36, wherein the at least one anti-neoplastic agent is selected from the group consisting of alkylating agents, anti-metabolites, epidophyllotoxins, antineoplastic enzymes, topoisomerase inhibitors, procarbazines, mitoxantrones, platinum coordination complexes, biological response modifiers and growth inhibitors, hormonal/anti-hormonal therapeutic agents, and haematopoietic growth factors.
38. The use of the compound, polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug of claim 1 in the manufacture of a medicament.
39. The use of the compound, polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug of claim 1 in the manufacture of a medicament for the therapeutic and prophylactic treatment of HSP90-dependent diseases and conditions.
40. A compound, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, selected from the group:



41. ~~PCT/US2006/011846~~
A compound of claim 1, or a polymorph, solvate, tautomer, enantiomer, pharmaceutically acceptable salt or prodrug thereof, selected from the group:



PCT/US2006/11846

