

USE OF MILK EXOSOME IN PREPARATION OF DRUG CARRIER

FIELD

5 [0001] The present disclosure relates to the field of pharmaceutical preparations, and in particular to use of a milk exosome in the preparation of a drug carrier.

BACKGROUND

10 [0002] Exosomes refer to small membranous vesicles (30-150 nm) containing complex RNA and proteins, now specifically to discoidal vesicles with a diameter of 40-100 nm. Exosomes were first discovered in sheep reticulocytes in 1983 and named "exosome" by Johnstone in 1987. A variety of cells can secrete exosomes under normal and pathological conditions. Exosomes are mainly derived from multivesicular bodies that are formed by the invagination of intracellular lysosomal particulates and released into the extracellular matrix after the outer membrane of the multivesicular bodies fuses with the cell membrane.

15 [0003] Currently, existing purification techniques for exosomes mainly include the following methods:

[0004] 1. Ultracentrifugation (differential centrifugation)

20 [0005] Ultracentrifugation is the most commonly used means of exosome purification and is considered the gold standard for exosome extraction. However, it is small in scale, time-consuming, and has unstable recovery rates (which may be related to the type of rotor), and the purity is also questionable. In addition, repeated centrifugation may damage the vesicles, resulting in reduced quality.

[0006] 2. Density gradient centrifugation

25 [0007] This is a zonal separation method, wherein the sucrose solution is subjected to ultracentrifugal force to form density gradient layers that are continuously distributed from low to high. The problem existing in density gradient centrifugation is that the steps are cumbersome and time-consuming.

[0008] 3. Ultrafiltration centrifugation

[0009] Extraction of exosomes by ultrafiltration centrifugation may cause the blockage of filter pores, resulting in a shorter membrane life and lower separation efficiency.

[0010] 4. Immunomagnetic bead method

5 **[0011]** This method has low efficiency in extracting exosomes, and the biological activity of exosomes is easily affected by pH and salt concentration, which is not conducive to downstream experiments, making it difficult to be widely popularized.

[0012] 5. PEG-based precipitation

10 **[0013]** Precipitation of exosomes using polyethylene glycol (PEG) has many problems such as low purity and recovery rate, many impurity proteins (false positives), non-uniform particle size, production of polymers that are difficult to remove, and destruction of exosomes by mechanical force or chemical additives such as Tween-20.

[0014] 6. Extraction Kit

15 **[0015]** The method of extracting exosomes using commercial exosome extraction kits is small in scale and will produce a lot of impurity proteins, leading to low purity and recovery rate of exosomes.

[0016] Therefore, the present disclosure aims to provide a method for efficiently extracting milk exosomes on a large scale without relying on centrifugation.

20 **[0017]** More importantly, drug loading by exosomes is difficult, and the present disclosure achieves and improves drug loading of protein drugs by exosomes.

[0018] Currently, protein drugs are often administered by injection, which is complicated and has safety risks.

SUMMARY

25 **[0019]** In view of this, the present disclosure uses milk exosomes as a drug carrier. The exosomes are derived from milk with safety and reliability, and has the functional properties of milk exosomes. Using the exosomes as a drug carrier can achieve oral administration of protein

drugs. Using the milk exosomes to load liraglutide can achieve oral administration of liraglutide and treat type 2 diabetes.

[0020] In order to achieve the above object, the present disclosure provides the following technical solutions.

5 **[0021]** The present disclosure provides use of a milk exosome in the preparation of a drug carrier. In some specific embodiments of the present disclosure, the drug includes a protein drug. In some specific embodiments of the present disclosure, the drug includes liraglutide.

[0022] The present disclosure further provides use of a milk exosome as a drug carrier in the preparation of a medicament for the prevention and/or treatment of a disease. In some specific
10 embodiments of the present disclosure, the drug includes a protein drug. In some specific embodiments of the present disclosure, the drug includes liraglutide. In some specific embodiments of the present disclosure, the drug is administered orally. In some specific embodiments of the present disclosure, the disease includes type 2 diabetes.

[0023] More importantly, the present disclosure further provides an oral medicament,
15 comprising an active ingredient and a milk exosome as a drug carrier.

[0024] In the use of the oral medicament of the present disclosure, the milk exosome is prepared by:

[0025] step 1, pre-treatment;

[0026] step 2, preliminary purification and concentration;

20 **[0027]** step 3, refined purification;

[0028] wherein the pre-treatment comprises fat removal and/or protein removal;

[0029] the preliminary purification and concentration are performed by tangential flow ultrafiltration, and

[0030] the refined purification is performed by a method selected from the group consisting of
25 CIMmultus QA column purification, Capto core 700 column purification, S400 molecular sieve chromatography purification, and a combination thereof.

[0031] In some specific embodiments of the present disclosure, the fat removal in step 1

comprises:

[0032] (1) removing fat by standing for static natural sedimentation and filtration; and/or

[0033] (2) removing fat by centrifugation;

[0034] wherein the filtration comprises depth filtration and/or capsule filter filtration.

5 **[0035]** In some specific embodiments of the present disclosure, the protein removal in step 1 is carried out by a protein dissolution method and/or a protein precipitation method;

[0036] wherein the protein dissolution method includes a sodium citrate dissolution method; and

10 **[0037]** the protein precipitation method is selected from the group consisting of EDTA precipitation method, sodium phosphate precipitation method, ammonium sulfate precipitation method, and a combination thereof.

[0038] In some specific embodiments of the present disclosure, the sodium citrate dissolution method comprises the following steps: mixing an equal volume of milk and a sodium citrate solution with a mass concentration of 2%, shaking in an ice bath, and filtering through a
15 membrane filter.

[0039] The sodium citrate dissolution method specifically comprises: pouring an equal volume of milk into a sodium citrate solution with a mass concentration of 2%, shaking in an ice bath on a shaker for 60 min to clarify the milk sample, and filtering the clarified milk sample through a 0.2 μm membrane filter.

20 **[0040]** In some specific embodiments of the present disclosure, the EDTA precipitation method comprises the following steps: mixing an equal volume of milk and an EDTA solution, standing at room temperature, centrifuging, and filtering through a membrane filter; wherein the EDTA solution has a concentration of 0.35 M and a pH of 7.0.

[0041] In some specific embodiments of the present disclosure, the sodium citrate dissolution
25 method specifically comprises: pouring an equal volume of milk into an EDTA solution, standing at room temperature for 15 min to precipitate, centrifuging the milk sample at 12,000 rpm for 40 min after precipitation, and then filtering through 0.45 μm and 0.2 μm membrane filters.

[0042] In some specific embodiments of the present disclosure, the sodium phosphate solution has a concentration of 1 M and a pH of 6.3.

[0043] In some specific embodiments of the present disclosure, the sodium phosphate precipitation method is as follows.

5 **[0044]** Depending on the type of milk, the sodium phosphate precipitation method comprises different specific steps:

[0045] (1) Whole milk

[0046] centrifuging the milk at 16,000 rpm for 30 min to remove fat, mixing an equal volume of milk with a sodium phosphate solution, stirring, precipitating, and freezing at -80°C;

10 **[0047]** (2) Skimmed milk

[0048] centrifuging the milk at 16,000 rpm for 30 min to remove fat, mixing an equal volume of milk and a sodium phosphate solution, stirring, precipitating, standing at room temperature, centrifuging at 3500 rpm for 10 min, collecting the supernatant, and filtering it through a membrane filter.

15 **[0049]** In some specific embodiments of the present disclosure, the sodium phosphate precipitation method specifically comprises the following steps: centrifuging milk at 16,000 rpm for 30 min to remove fat, pouring an equal volume of milk into the sodium phosphate solution, stirring on a magnetic stirrer at 350 rpm for 15 min, and centrifuging the milk sample at 3500 rpm for 10 min after precipitation, and then sequentially filtering the supernatant through 0.8 μm,
20 0.45 μm and 0.2 μm membrane filters.

[0050] In some specific embodiments of the present disclosure, the ammonium sulfate precipitation method comprises the following steps: mixing an equal volume of milk and an ammonium sulfate solution, standing at room temperature, centrifuging, and filtering through a membrane filter; wherein the concentration of the ammonium sulfate solution is 3.0 M.

25 **[0051]** In some specific embodiments of the present disclosure, the ammonium sulfate precipitation method specifically comprises the following steps: pouring an equal volume of milk into an ammonium sulfate solution while gentle stirring using a glass rod, standing at room temperature for 1.5 h to precipitate, centrifuging the milk sample at 6000 rpm for 20 min after

precipitation, and then sequentially filtering the supernatant through 0.8 μm , 0.45 μm and 0.2 μm membrane filter.

[0052] The present disclosure further provides a method for treating a disease, comprising administering to a subject a drug loaded in a milk exosome, wherein the milk exosome is used as a drug carrier.

[0053] In some specific embodiments of the present disclosure, the drug includes a protein drug.

[0054] In some specific embodiments of the present disclosure, the drug includes liraglutide.

[0055] In some specific embodiments of the present disclosure, the drug is administered orally.

[0056] In some specific embodiments of the present disclosure, the disease includes type 2 diabetes.

[0057] The present disclosure uses milk exosomes as a drug carrier. The exosomes are derived from milk with safety and reliability, and has the functional properties of milk exosomes. Using the exosomes as a drug carrier can achieve oral administration of protein drugs. Using the milk exosomes to load liraglutide can achieve oral administration of liraglutide and treat type 2 diabetes.

BRIEF DESCRIPTION OF DRAWINGS

[0058] For more clearly illustrating embodiments of the present disclosure or the technical solutions in the prior art, drawings for describing the embodiments or the prior art will be briefly described hereinafter.

[0059] FIG. 1 shows the process route for extraction of milk exosomes by TFF+BE-SEC and UC.

[0060] FIG. 2 shows the morphology of exosomes extracted by three methods under TEM (Bar=500 nm) (A: TFF; B: TFF + BE-SEC; C: UC).

[0061] FIG. 3 shows the particle size distribution of exosomes extracted by three methods (A: TFF; B: TFF + BE-SEC; C: UC).

[0062] FIG. 4 shows the SEC-HPLC chromatogram of milk exosomes extracted by three methods (A: TFF; B: TFF + BE-SEC; C: UC).

[0063] FIG. 5 shows a liraglutide solution after ultrasonication (left: a colloidal solution with more precipitates after ultrasonication) and a mixture of milk exosomes and liraglutide (right).

5 **[0064]** FIG. 6 shows the result obtained from Core700 column purification of milk exosomes loaded with liraglutide by ultrasonication.

[0065] FIG. 7 shows the result obtained from Core700 column purification of ultrasonicated liraglutide.

10 **[0066]** FIG. 8 shows the result obtained from QA column purification of milk exosomes and liraglutide after ultrasonication.

[0067] FIG. 9 shows the result obtained from QA column purification of ultrasonicated liraglutide.

15 **[0068]** FIG. 10 shows the effects of the liraglutide solution and the exosome-liraglutide mixture after ultrasonication and Core700 column purification on luciferase expression (all positive controls are treatment groups of 0.375 μ g/ml liraglutide without ultrasonication).

20 **[0069]** FIG. 11 shows the conditions for exploring the use of electroporation to load PKH67 in exosomes; wherein, FIG. 11A shows the FITC positive rate and the particle concentration of the samples after electroporation of the PKH67-exosome mixture under different voltages (100 μ F capacitance); and FIG. 11B shows the FITC positive rate and the particle concentration of the samples after electroporation of the PKH67-exosome mixture under different capacitance (500V voltage).

[0070] FIG. 12 shows the result obtained from Core700 column purification of milk exosomes loaded with liraglutide by electroporation.

25 **[0071]** FIG. 13 shows the result obtained from Core700 column purification of electroporated liraglutide.

[0072] FIG. 14 shows the effects of the liraglutide solution and the exosome-liraglutide mixture after electroporation and Core700 column purification on luciferase expression (all positive controls are treatment groups of 0.375 μ g/ml liraglutide without electroporation).

[0073] FIG. 15 shows the conditions for exploring the use of incubation to load PKH67 in exosomes; wherein, FIG. 15A shows the FITC positive rate and the particle concentration of the samples after incubation of the PKH67-exosome mixture at different temperatures (0.5 h); and FIG. 15B shows the FITC positive rate and the particle concentration of the samples after incubation at 50°C for different periods of time.

[0074] FIG. 16 shows the result obtained from Core700 column purification of electroporated liraglutide.

[0075] FIG. 17 shows the effects of the liraglutide solution and the exosome-liraglutide mixture after co-incubation and core700 column purification on luciferase expression (all positive controls are treatment groups of 0.375 µg/ml liraglutide without co-incubation).

[0076] FIG. 18 shows the cellular efficacy of exosomes loaded with different drugs.

[0077] FIG. 19 shows the particle number of positive exosomes in serum of mice after administration by gavage at different time points as detected by nanoflow cytometry.

[0078] FIG. 20 shows the particle number of positive exosomes in serum of mice after sublingual administration at different time points as detected by nanoflow cytometry.

[0079] FIG. 21 shows changes over time in blood glucose of mice sublingually administered with exosomes loaded with different drugs and given glucose solution by gavage after 1 h.

[0080] FIG. 22 shows the changes over time in random blood glucose of mice sublingually administered with different concentrations of exosome-liraglutide.

[0081] FIG. 23 shows the changes over time in fasting blood glucose of mice sublingually administered with different concentrations of exosome-liraglutide after 6 h of fasting.

DETAILED DESCRIPTION

[0082] The present disclosure discloses use of a milk exosome in the preparation of a drug carrier. Those skilled in the art can learn from the contents of the disclosure and appropriately adapt the process parameters to achieve the present disclosure. Particularly, it should be noted that all such alternatives and modifications are obvious to those skilled in the art and are

considered to be included within the present disclosure. The method and the application of the present disclosure have been described through the preferred embodiments, and it is obvious that the method and application described herein may be changed or appropriately modified and combined to realize and apply the technology of the present disclosure by those skilled in the art without departing from the content, spirit and scope of the present disclosure.

[0083] All materials and reagents used in the use of a milk exosome in the preparation of a drug carrier provided by the present disclosure are commercially available.

[0084] The present disclosure is further illustrated below in combination with the embodiments.

Example 1 Industrially scalable purification process for milk exosomes

[0085] 1) Pre-treatment of milk samples by different protein removal methods

[0086] a. EDTA precipitation method: Skimmed milk was mixed with EDTA of the same volume, different concentrations and different pHs, and incubated at room temperature for 15 min to precipitate casein. The EDTA used had a concentration of 0.15-0.35 mol·L⁻¹ and a pH of 6.0-8.0. Centrifugation was carried out at 12,000 r·min⁻¹ for 40 min, and the supernatant was filtered through 0.80, 0.45 and 0.20 μm membrane filters sequentially.

[0087] b. Sodium phosphate precipitation method: An equal amount of skimmed milk was added to a sodium phosphate solution. Several pre-tests were carried out on the concentration and pH range of sodium phosphate, such as using 0.5 mol·L⁻¹ (pH 5.3, pH 6.5 and pH 7.6), 1.0 mol·L⁻¹ (pH 5.2, pH 6.3 and pH 7.6) and 2 mol·L⁻¹ (pH 6.0, pH 7.0 and pH 8.0) sodium phosphate. This mixture was placed at 4°C overnight to precipitate protein, and the supernatant after precipitation was sequentially filtered through 0.80, 0.45, and 0.20 μm membrane filters.

[0088] c. Sodium citrate dissolution method: skimmed milk was mixed with equal amounts of 2%, 4%, 6%, 8% and 16% sodium citrate, incubated on ice for 2 h to dissolve casein, and filtered through a 0.20 μm membrane filter.

[0089] 2) Preliminary purification and concentration of milk exosomes by TFF method

[0090] The milk supernatant after pre-treatment was processed using a hollow fiber column

with a relative molecular mass of 750,000. The system parameters were set as follows: TMP maximum pressure at 2 atmospheres, alarm pressure at 2 atmospheres; 100 mL minimum volume, and 100 mL alarm volume. The sample was poured into a reusable beaker, and exchanged with 1×PBS until the concentration of permeate proteins was 0.00 g·L⁻¹. Then the liquid in the beaker was further concentrated to 1/20 of the original solution, and the sample was collected.

[0091] 3) Refined purification of milk exosomes by BE-SEC and UC

[0092] The ÄKTA Pure 25 purification system was rinsed with ultrapure water, connected to a HiScreen CaptoCore 700 column, and then rinsed with 3 column volumes of ultrapure water. After rinsing, the equilibration was carried out with 3 column volumes of 1×PBS, and the preliminarily purified sample was purified using a HiScreen CaptoCore 700 column at a flow rate of 2 mL·min⁻¹. The sample was collected at the position of the flow-through peak, i.e. milk exosomes.

[0093] Experimental results

[0094] As shown in FIG. 1, the protein precipitation method was used to obtain a pretreated sample, and the purification method TFF in combination with BE-SEC was used to obtain milk exosomes. It was demonstrated that the sample obtained by this process has a higher yield and purity compared with UC, a traditional extraction method of exosomes.

[0095] As shown in FIG. 2, the membrane structures of exosomes extracted by TFF, BE-SEC and UC can all be observed under TEM, showing a saucer-like or disc-like shape. However, the exosomes extracted by UC under TEM had a cluttered background and more protein aggregates. In comparison, exosomes purified by TFF combined with BE-SEC had a cleaner background.

[0096] The particle size, concentration and purity of exosomes detected by NanoFCM are shown in FIG. 3. The exosomes obtained by TFF had the average particle size of (66.92±16.39) nm, a yield of 3.6×10¹³ particles·L⁻¹ and a purity of 58%. The milk exosomes prepared by TFF+BE-SEC had the diameter concentrated in a range of 30-150 nm, the average particle of (66.41±15.81) nm, a yield of 2.5×10¹³ particles·L⁻¹ and a purity of 90%. The exosomes obtained by UC had the average particle size of (68.87±18.04) nm, a yield of 2.4×10¹¹ particles·L⁻¹ and a purity of 25%. It was demonstrated that the exosomes prepared by TFF+BE-SEC had a higher yield and purity.

[0097] As shown in FIG. 4, the amount of free proteins (particle size smaller than that of exosomes) in the exosome sample was detected by HPLC. The retention time of milk exosomes extracted by the three methods on a TSKgel G6000 PWWL column was all 8 min. However, the proportion of impurity peaks (RT>8 min) of the exosome samples extracted by TFF and BE-SEC was smaller than that of the samples obtained by UC, and the final product obtained by TFF+BE-SEC showed more of a single peak.

[0098] The use of protein precipitation pre-treatment in combination with purification by TFF and BE-SEC of milk can obtain exosome samples with a higher yield and purity. More importantly, TFF could achieve the preliminary purification and concentration of samples from a few liters to hundreds of liters, and the large-scale preparation of milk exosomes could finally be achieved by combining TFF with BE-SEC.

Example 2 Drug loading process for milk exosomes - ultrasonication method

[0099] Principle: Ultrasonication can disrupt the arrangement of lipid chains in the membrane and create temporary pores. After exosomes are subjected to low-frequency continuous ultrasonication, the sound wave causes the encapsulated air to expand and the pressure inside the vesicle to rise to the elastic limit of the bilayer, resulting in pores in the phospholipid layer. Due to the high drug concentration outside the membrane, the difference in osmotic pressure causes the drug to diffuse into the liposome through the pores.

[00100] Operation steps

[00101] The optimal conditions for loading by ultrasonication were explored using PKH67. Specifically, after co-incubation of PKH67 with milk exosomes at room temperature for 30 min, ultrasonication was carried out using a cell crusher at the power of 150w, 300w, 450w, 600w, 800w, 1000w and 1200w for 0 s, 50 s, 250 s, 500 s, 750 s, 20 min, 30 min, 1 h, 2 h, 4 h and 6 h. The particle number and PKH67 positive rate of the ultrasonicated samples were determined by using NanoFCM. Based on the optimal conditions for loading by ultrasonication, the volume for loading by ultrasonication was explored. The particle number, particle size and PKH67 positive rate of the ultrasonicated samples were determined by using NanoFCM.

[00102] Experimental results

Table 1 Effects of power and duration of ultrasonication on positive rate of exosome loading

Time Power	0s	50s	250s	500s	750s	20min	30min	1h	2h	4h	6h
0w	5.4%	6.1%	7.2%	8.3%	8.9%	10.3%	10.0%	15.2%	15.4%	27.5%	31.0%
150w	5.4%	8.6%	23.5%	31.4%	31.9%	40.3%	47.6%	45.5%	48.3%	50.1%	50.6%
300w	5.4%	9.7%	12.1%	19.3%	35.0%	33.6%	50.6%	45.1%	48.3%	58.0%	52.7%
450w	5.4%	15.8%	31.2%	60.2%	54.6%	44.5%	63.3%	74.3%	76.1%	75.9%	79.1%
600w	5.4%	15.4%	67.6%	77.6%	76.1%	78.1%	76.5%	78.6%	85.6%	74.1%	78.7%
800w	5.4%	30.0%	65.5%	78.1%	78.4%	75.4%	80.4%	79.3%	78.0%	80.1%	80.0%
1000w	5.4%	42.2%	64.8%	77.3%	77.2%	80.4%	87.8%	80.0%	85.7%	81.2%	88.1%
1200w	5.4%	41.0%	66.3%	77.5%	78.9%	75.5%	75.4%	88.7%	88.3%	85.0%	87.5%

Table 2 Effects of power and duration of ultrasonication on particle number (10^{10} particles/mL) of exosomes

Time Power	0s	50s	250s	500s	750s	20min	30min	1h	2h	4h	6h
0w	7.39	7.90	7.68	7.21	6.56	6.43	5.88	6.31	5.21	5.10	4.52
150w	7.39	6.77	7.33	6.88	6.21	6.07	6.22	5.77	5.48	5.09	4.71
300w	7.39	6.53	7.08	7.01	6.38	6.37	6.11	6.41	5.88	4.87	5.00
450w	7.39	7.01	6.81	6.55	4.56	5.21	5.43	6.01	4.78	4.45	4.03
600w	7.39	5.91	5.99	5.76	3.80	3.90	4.01	3.54	3.48	3.61	3.67
800w	7.39	5.90	6.42	5.87	4.22	5.11	4.63	4.37	3.71	3.31	3.44
1000w	7.39	6.44	6.53	6.11	5.89	5.33	5.41	3.51	2.69	2.24	2.30
1200w	7.39	6.54	7.11	6.38	6.67	7.31	6.06	2.67	3.08	2.87	2.01

Table 3 Effects of different volumes for ultrasonication (600w, 500s) on exosome loading

Volume (ml)	Positive rate (%)	Particle concentration (10^{10} particles/mL)
3	77.8	5.23
10	75.3	5.60
50	80.1	4.85
100	78.1	5.08
200	75.6	4.66
500	74.7	4.97

[00103] There was a large difference in exosome loading efficiency after ultrasonication at different power, and the positive rate after ultrasound was between 5% and 90%. For ultrasonication at 450W-1200W power for more than 20 min, the difference in loading efficiency was not obvious and was basically at about 80%. With the prolongation of time of ultrasonication, the particle number decreased. The loading scale of 3 ml-500 ml system could be achieved, and the particle number and loading positive rate after loading were relatively stable.

[00104] Loading liraglutide by ultrasonication:

[00105] After mixing according to the ratio of 1×10^{11} exosome particles to 5 mg liraglutide, ultrasonication was performed followed by purification using Core700 or QA columns. There was an obvious change in the state of the solution before and after ultrasonication. FIG. 5 shows a liraglutide solution after ultrasonication (left: a colloidal solution with more precipitates after ultrasonication) and a mixture of milk exosomes and liraglutide (right).

[00106] 25 mg of liraglutide, and a mixture of 25 mg liraglutide and 5×10^{11} exosome particles were ultrasonicated and then passed through a Core700 column to collect flow-through samples. The results are shown in FIGs. 6-7. On a 5 mL Core700 chromatography column, 2-4 mL of flow-through sample was collected after the injection started. This fraction was the exosome particles loaded with liraglutide, and the free liraglutide was adsorbed on the column and then removed by elution.

[00107] 250 mg liraglutide, and a mixture of 250 mg liraglutide and 5×10^{12} exosome particles

were ultrasonicated and then passed through a QA column for purification, and flow-through and eluted samples were collected. The results are shown in FIGss 8-9. The liraglutide-loaded exosomes and free liraglutide were eluted separately under different salt concentrations based on the difference in electric charge quantity.

- 5 **[00108]** GLP-1R-CHO cells were seeded into a 96-well plate and cultured for 24 h, and then the medium was replaced with blank milk exosomes, milk exosome-liraglutide (10^5 /mL, 10^6 /mL, 10^7 /mL, 10^8 /mL, 10^9 /mL and 10^{10} /mL) or liraglutide solutions with the corresponding particle concentration for 5 h of culture. The samples were collected according to the instructions of luciferase kit, and 100 μ l of cell lysate was added to each well to lyse cells for 5 min. The cell
- 10 lysate was collected and centrifuged at 1200 rpm for 5 min. Then the substrate was added and incubated for 5 min, and the fluorescence was detected using an enzyme reader. The results are shown in FIG. 10 and Table 4.

Table 4

	Exosome-liraglutide_Core700			Liraglutide (ultrasonication)_Core700		
Blank	20617	19685	22845	19513	20557	21774
Positive control	160126	161819	164485	166442	168105	182139
Blank exosomes	21244	24053	24830	22927	25572	26806
1×10^{10}	117775	129093	137324	23620	23463	23987
1×10^9	49581	52138	56060	22659	23782	25679
1×10^8	24063	25156	24889	24413	23676	23367
1×10^7	20066	20339	22143	23953	23706	24012
1×10^6	17023	18083	20867	22732	22735	22400
1×10^5	18999	17390	18551	17634	17430	21039

- 15 **[00109]** Experimental conclusion: From the comparison of the liquid appearance of liraglutide with that of the milk exosome-liraglutide mixture after ultrasonication, it can be seen that liraglutide did not precipitate when containing exosomes, but was loaded on the exosomes. After ultrasonication and QA purification of the milk exosome-liraglutide mixture, the chromatogram

showed that the A260/A280 absorption ratio of the samples after loading by ultrasonication in high salt elution (conductivity of 70.44 Ms/cm) was increased to 1.88, which represented the absorption peak of nucleic acid contents released from exosomes of which the bilayer membrane was damaged by ultrasonication. The particle number of exosomes decreased after prolonged
5 ultrasound time. After purification through core700 or QA columns, the drug-treated cells were proportionally diluted, and it was shown that the expression of luciferase was dose-dependent. The treatment of 1×10^{10} particles of milk exosome-exosome-liraglutide on cells significantly increased luciferase expression, indicating that the loading of liraglutide in milk exosomes was achieved with biological activity.

10

Example 3 Drug loading process for milk exosomes - electroporation method

[00110] Principle: Electroporation is a microbiological technique, where an electric field is applied to a cell to increase the permeability of the cell membrane, thereby allowing chemicals, drugs or DNA to be introduced into the cell.

15 **[00111]** Operation steps:

[00112] The optimal conditions for drug loading by electroporation were explored using PKH67. Specifically, after co-incubation of PKH67 with milk exosomes at room temperature for 30 min, electroporation was carried out using an electroporator at a voltage of 100-500 v and a capacitance of 100-500 μ F. The particle number, particle size and PKH67 positive rate of the
20 electroporated samples were determined by using NanoFCM.

[00113] 25 mg of liraglutide, and a mixture of 25 mg liraglutide and 5×10^{11} exosome particles were electroporated and then passed through a core700 column to remove free liraglutide. The flow-through samples were collected.

[00114] GLP-1R-CHO cells were seeded into a 96-well plate and cultured for 24 h, and then the
25 medium was replaced with blank milk exosomes, milk exosome-liraglutide (10^5 /mL, 10^6 /mL, 10^7 /mL, 10^8 /mL, 10^9 /mL and 10^{10} /mL) or liraglutide solutions with the corresponding particle concentration for 5 h of culture. The samples were collected according to the instructions of luciferase kit, and 100 μ l of cell lysate was added to each well to lyse cells for 5 min. The cell lysate was collected and centrifuged at 1200 rpm for 5 min. Then the substrate was added and

incubated for 5 min, and the fluorescence was detected using an enzyme reader.

[00115] The results are shown in FIGs. 11-14 and Tables 5-7.

Table 5 Effect of voltage on exosomes loading

Voltage V	Positive rate %	Concentration particles $\times 10^{10}$ /mL
0	7.12	8.55
100	8.2	2.8
200	28.5	2.52
300	39	2.26
400	62.9	1.56
500	59.5	2.14

5

Table 6 Effect of capacitance on exosomes loading

Capacitance μ F	Positive rate %	Concentration particles $\times 10^{10}$ /mL
0	6.35	8.55
100	46.8	3.2
200	51.8	3.78
300	71.2	2.78
400	69.8	2.96
500	57.1	2.11

Table 7

	Exosome-liraglutide_Core700			Liraglutide (electroporation)_Core700		
Blank	16617	15685	21845	17513	18557	19774
Positive control	170126	131819	144485	196442	128105	122139
Blank exosomes	18244	20053	24830	22927	18572	20806
1×10^{10}	97775	92093	87324	20620	20463	20987

1×10^9	50581	42138	46060	21659	22782	18679
1×10^8	22063	22156	22889	19413	18676	20367
1×10^7	20066	20339	22143	23953	23706	24012
1×10^6	17023	18083	20867	22732	22735	22400
1×10^5	18999	17390	18551	17634	17430	21039

Example 4 Drug loading process for milk exosomes - co-incubation method

[00116] Principle: The drug will be adsorbed on the drug carrier by the interaction between substances.

5 **[00117]** Operation steps

[00118] The optimal conditions for loading by co-incubation were explored using PKH67. Specifically, after co-incubation of PKH67 with milk exosomes at room temperature for 30 min, the samples were incubated at 4°C, 37°C and 50°C with shaking at 500 rpm for 30 min, 1 h, and 2 h. The particle number, particle size and PKH67 positive rate of the co-incubated samples were determined by using NanoFCM.

[00119] 25 mg of liraglutide, and a mixture of 25 mg liraglutide and 5×10^{11} exosome particles were co-incubated and then passed through a core700 column to remove free liraglutide. The flow-through samples were collected.

[00120] GLP-1R-CHO cells were seeded into a 96-well plate and cultured for 24 h, and then the medium was replaced with blank milk exosomes, milk exosome-liraglutide ($10^5/\text{mL}$, $10^6/\text{mL}$, $10^7/\text{mL}$, $10^8/\text{mL}$, $10^9/\text{mL}$ and $10^{10}/\text{mL}$) or liraglutide solutions with the corresponding particle concentration for 5 h of culture. The samples were collected according to the instructions of luciferase kit, and 100 μl of cell lysate was added to each well to lyse cells for 5 min. The cell lysate was collected and centrifuged at 1200 rpm for 5 min. Then the substrate was added and incubated for 5 min, and the fluorescence was detected using an enzyme reader.

[00121] The results are shown in FIGs. 14-17 and Tables 8-10.

Table 8 Effect of incubation temperature on exosome loading

Temperature	Positive rate %	Concentration particles $\times 10^{10}$ /mL
4°C	7.12	8.55
37°C	8.2	3.18
42°C	28.5	3.32
50°C	49	3.26
60°C	22.9	1.56

Table 9 Effect of incubation time on exosome loading

Time	Positive rate %	Concentration particles $\times 10^{10}$ /mL
0.5h	32.1	3.6
1h	46.8	3.78
2h	29.8	3.78
4h	21.2	2.78
8h	9.8	1.96

Table 10 Cellular efficacy by co-incubation method

	Exosome-liraglutide_Core700			Liraglutide (incubation)_Core700		
Blank	16617	15685	21845	17513	18557	19774
Positive control	190126	131819	144485	156442	128105	122139
Blank exosomes	18244	20053	24830	22927	18572	20806
1×10^{10}	77775	72093	87324	14620	15463	13987
1×10^9	40581	32138	46060	11659	12782	11679
1×10^8	12063	12156	12889	9413	8676	80367
1×10^7	10066	10339	12143	7953	7306	74012
1×10^6	7023	8083	7867	5732	6735	6400
1×10^5	6999	7390	6551	4634	5430	4039

[00122] Experimental conclusion: After purification through core700, the drug-treated cells were proportionally diluted, and it was shown that the expression of luciferase was dose-dependent. The treatment of 1×10^{10} particles of milk exosome-exosome-liraglutide on cells significant increased luciferase expression, indicating that the loading of liraglutide in milk exosomes was achieved with biological activity.

Example 5 Loading different T2DM drugs in milk exosomes

[00123] At present, there are many drugs for treating type 2 diabetes mellitus (T2DM) on the market, such as insulin degludec, insulin detemir, liraglutide, semaglutide, and tirzepatide which has been listed abroad. The present disclosure realized loading different T2DM drugs in milk exosomes, so that these drugs can be administered orally instead of by injection, solving the dilemma of no oral diabetes drugs on the market.

[00124] Operation

[00125] 1) According to the drug loading method of milk exosomes introduced in 4.2, the electroporation method was selected. The milk exosomes were loaded with insulin degludec, insulin detemir, liraglutide, semaglutide and tirzepatide separately, and purified through a core700 column.

[00126] 2) GLP-1R-CHO cells were seeded into a 96-well plate and cultured for 24 h, and then the medium was replaced with blank milk exosomes, or milk exosome loaded with different drugs ($10^5/\text{mL}$, $10^6/\text{mL}$, $10^7/\text{mL}$, $10^8/\text{mL}$, $10^9/\text{mL}$ and $10^{10}/\text{mL}$) for 5 h of culture. The samples were collected according to the instructions of luciferase kit, and 100 μl of cell lysate was added to each well to lyse cells for 5 min. The cell lysate was collected and centrifuged at 1200 rpm for 5 min. Then the substrate was added and incubated for 5 min, and the fluorescence was detected using an enzyme reader.

[00127] The results are shown in FIG. 18 and Table 11.

Table 11 Cellular efficacy of exosomes loaded with different drugs

	Exosmome-liraglutide			Exosome-tirzepatide			Exosome-semaglutide		
Blank	16617	15685	21845	17513	18557	19774	20617	19685	22845
Positive control	170126	171819	144485	206442	208105	212139	160126	161819	164485
Blank exosomes	23244	24053	24830	22927	26572	20806	21244	24053	24830
1×10^{10}	97775	92093	87324	110620	110463	120987	97775	99093	87324
1×10^9	50581	42138	46060	51659	52782	58679	49581	52138	56060
1×10^8	22063	22156	22889	19413	18676	20367	24063	25156	24889
1×10^7	20066	20339	22143	23953	23706	24012	20066	20339	22143
1×10^6	17023	18083	20867	22732	22735	22400	17023	18083	20867
1×10^5	18999	17390	18551	17634	17430	21039	18999	17390	18551

[00128] Experimental conclusion:

[00129] From the results, milk exosomes loaded with different drugs enable cells to release GLP in a dose-dependent manner, indicating that the present disclosure can realize the loading of different types of protein and polypeptide drugs.

Example 6 Sublingual administration of milk exosomes

[00130] Principle: Utilizing the characteristics of the sublingual mucosa that is not keratinized, has abundant capillaries, and has fast blood flow speed, and aiming to solve problems that milk exosomes are easily destroyed by gastric acid after passing through the gastrointestinal tract, resulting in low bioavailability of the loaded drug, the present disclosure prepares liraglutide-loaded milk exosomes into sublingual tablets to achieve oral administration of liraglutide, which can be absorbed through the sublingual mucosa and directly enter the blood circulation through the jugular vein and superior vena cava, with a rapid onset of action. In addition, the drug can be taken without water by placing it under the tongue to dissolve, which is convenient to take and has a good taste.

[00131] Experimental operation:

[00132] A. PKH67-labelled milk exosomes were administered separately by gavage, sublingually, and were prepared into enteric capsules for administration.

[00133] B. After administration, blood was collected from mice at different time points, and the number of positive particles in the blood was detected by nanoflow to determine the absorption effect of exosomes into the blood.

[00134] The experimental results are shown in FIGs. 19-20 and Tables 12-14.

Table 12 Data from administration by gavage

Mode of administration	Positive particles					
PBS	8.70×10^7	1.74×10^7	6.46×10^7	3.57×10^7	4.29×10^7	6.83×10^7
PKH67-15min	3.30×10^8	2.44×10^8	2.01×10^8	2.09×10^8	3.04×10^8	9.87×10^7
Tail vein	1.68×10^{11}	2.34×10^{11}	2.12×10^{11}	1.98×10^{11}	2.01×10^{11}	2.05×10^{11}
5min	1.85×10^9	2.32×10^9	1.94×10^9	2.05×10^9	1.77×10^9	2.11×10^9
15min	7.48×10^8	3.66×10^8	6.96×10^8	4.87×10^8	6.12×10^8	5.34×10^8
30min	4.00×10^8	2.62×10^8	1.57×10^8	3.56×10^8	1.22×10^8	3.19×10^8
45min	3.30×10^8	2.26×10^8	1.22×10^8	2.98×10^8	3.09×10^8	1.78×10^8
1h	2.44×10^8	2.08×10^8	1.74×10^8	1.35×10^8	9.98×10^7	2.04×10^8
2h	1.57×10^8	2.78×10^8	1.39×10^8	1.34×10^8	1.24×10^8	8.99×10^7
4h	1.39×10^8	1.04×10^8	1.57×10^8	1.02×10^8	1.11×10^8	8.34×10^7

Table 13 Data from sublingual administration

Mode of administration	Positive particles					
PBS	6.60×10^7	6.60×10^7	5.89×10^7	6.01×10^7	5.46×10^7	5.28×10^7
Tail vein	2.60×10^{11}	2.30×10^{11}	2.45×10^{11}	1.98×10^{11}	2.45×10^{11}	2.19×10^{11}
30min	7.88×10^9	5.78×10^9	6.56×10^9	2.89×10^{10}	4.02×10^{10}	3.27×10^{10}
1h	3.50×10^9	5.08×10^9	1.04×10^9	7.54×10^8	4.78×10^9	1.67×10^9
2h	1.11×10^9	5.36×10^8	8.97×10^8	7.29×10^8	1.02×10^9	6.92×10^8

4h	4.26×10^8	4.72×10^8	4.72×10^8	6.30×10^8	2.78×10^9	3.10×10^9
----	--------------------	--------------------	--------------------	--------------------	--------------------	--------------------

Table 14 Data of enteric capsules

	mean	SD
0min	4.58×10^7	2.67×10^7
30min	6.10×10^7	1.89×10^7
1h	1.53×10^7	9.87×10^6
2h	1.53×10^7	8.22×10^6
3h	3.05×10^7	1.24×10^7
4h	6.86×10^7	2.78×10^7
5h	1.30×10^8	9.18×10^6
8h	1.53×10^7	7.35×10^6
22h	9.92×10^7	3.22×10^7

[00135] For PKH67-labelled milk exosomes administered to mice by gavage, the blood was collected at different time points to detect positive particles and 4×10^{11} particles were administered per mouse. After 5 min, the highest 2×10^9 positive particles were detected in the blood of the mice, corresponding to 0.5% exosomes entering the blood. For PKH67-labelled milk exosomes administered sublingually, after 5 min, the highest 2×10^{10} positive particles were detected, corresponding to 5% exosomes entering the blood. However, for administration by enteric capsules, the number of positive particles in the blood of mice did not increase significantly. Therefore, the bioavailability by sublingual administration is significantly higher compared to administration by gavage.

Example 7 Sublingual tablets of milk exosomes

[00136] Experimental operation: Lyophilized powder of milk exosomes loaded with liraglutide was mixed with a disintegrant, filler, binder, lubricant and flavoring agent to prepare into

sublingual tablets. In each 1,000 tablets of sublingual tablets of liraglutide milk exosomes, the main drug has a content of 10^5 - 10^{17} particles, the disintegrant 1-50 g, the filler 20-400g, the flavoring agent 0-50 g, the binder 5-100 g, and the lubricant 1 g-10 g;

[00137] The disintegrant used was selected from the group consisting of microcrystalline cellulose, cross-linked sodium carboxymethyl cellulose, cross-linked polyvinylpyrrolidone, sodium carboxymethyl starch, low-substituted hydroxypropyl cellulose, starch, Tween, sodium dodecyl sulfate, and a mixture thereof.

[00138] The filler used was selected from the group consisting of microcrystalline cellulose, microcrystalline cellulose-mannitol, microcrystalline cellulose-micronized silica gel, lactose, starch, modified starch, mannitol, sorbitol, xylitol, erythrose, trehalose, pregelatinized starch, icing sugar, glucose, dextrin, calcium sulfate, and a mixture thereof.

[00139] The flavoring agent used was selected from the group consisting of stevioside, icing sugar, glycyrrhizin, aspartame, sucralose, cyclamate, thaumatin, saccharin, and a mixture thereof.

[00140] The binder used was selected from the group consisting of purified water, starch slurry, hydroxypropyl methylcellulose, polyvinylpyrrolidone, carbomer, dextrin, gelatin slurry, gum arabic slurry, sodium alginate, syrup, and a mixture thereof.

[00141] The lubricant used was selected from the group consisting of stearic acid, magnesium stearate, calcium stearate, micronized silica gel, talc, hydrogenated vegetable oil, polyethylene glycol 4000, polyethylene glycol 6000, sodium dodecyl sulfate, magnesium dodecyl sulfate, sodium stearyl fumarate, and a mixture thereof.

[00142] The optimal formula is as follows: 1×10^{15} particles of exosome-liraglutide, 4 g cross-linked sodium carboxymethyl cellulose, 80 g mannitol, 25 g purified water and 3.5 g magnesium stearate. The preparation was performed according to the following preparation steps.

[00143] Preparation steps:

[00144] Step 1: The main drug was lyophilized and passed through 80-120 mesh sieves.

[00145] Step 2: The filler, disintegrant, flavoring agent and lubricant were dried, crushed and passed through 80-120 mesh sieves for pre-treatment.

[00146] Step 3: The pre-treated main drug, filler, flavoring agent and disintegrant were mixed

well.

[00147] Step 4: The binder was added to the above well-mixed materials to prepare soft materials, which were then granulated and dried.

[00148] Step 5: The lubricant was added to the dry granules, mixed well and then tableted to obtain the final product.

Example 8

[00149] 1×10^{13} particles of the main drug, liraglutide-milk exosomes, was added to 0.6 g mannitol, lyophilized, dried with 55 g microcrystalline cellulose the filler, 1 g sodium carboxymethyl cellulose the disintegrant, and 1 g magnesium stearate the lubricant, crushed, sieved, mixed, prepared into a soft material, granulated with a 20 mesh sieve, air dried to moisture $\leq 3\%$, then added with lubricant for final blending, and finally tableted to obtain sublingual tablets of liraglutide-milk exosomes.

Example 9

[00150] Experimental operation

[00151] By electroporation method, insulin degludec, insulin detemir, liraglutide, semaglutide, and tirzepatide were loaded into milk exosomes and purified to obtain drug-loaded exosomes, which were prepared into sublingual tablets according to the above preparation formula.

[00152] Each mouse was sublingually administered with 1×10^{11} particles of sublingual tablets and injected with glucose to detect the glucose tolerance in mice.

[00153] The results are shown in FIG. 21 and Table 15.

Table 15

Time (min)	-60	0	15	30	60	120
Vehicle	80	81	431	398	298	199
	83	85	452	382	301	203

	87	85	403	401	265	217
Exosome-liraglutide	87	88	135	121	110	100
	82	86	146	135	109	102
	85	81	132	129	101	90
Exosome-tirzepatide	91	93	101	99	95	93
	87	81	110	97	94	90
	89	84	109	98	92	88
Exosome-semaglutide	92	89	129	112	109	98
	91	93	137	130	110	95
	86	83	140	129	103	100
Exosome-insulin detemir	86	88	139	120	106	90
	83	89	125	111	103	92
	82	80	131	109	101	95
Exosome-insulin degludec	88	81	135	121	107	95
	89	84	126	114	102	93
	81	89	139	117	109	90
Exosome-insulin	83	89	155	141	120	116
	81	88	189	139	126	120
	89	85	167	129	118	111

[00154] Result analysis

[00155] The use of milk exosomes can achieve loading of multiple hypoglycaemic drugs. The drug-loaded exosomes can achieve significant hypoglycaemic effect within 15 min by oral administration. And it can maintain the stable blood glucose and improve the glucose tolerance of mice.

Example 10

[00156] By electroporation method, liraglutide (not limited to liraglutide, but also including insulin degludec, insulin detemir, semaglutide and tirzepatide) was loaded into milk exosomes and purified to obtain drug-loaded exosomes, which were prepared into sublingual tablets according to the optimal preparation formula. Here, the results of milk exosomes loaded with
5 liraglutide are illustrated as an example.

[00157] Diabetic mice were administered sublingually with different concentrations of milk exosomes loaded with liraglutide, and random blood glucose of the mice was detected.

[00158] The results are shown in FIG. 22 and Table 16.

Table 16

min	Blank									
0	277	245	288	266	329	312	289	287	290	321
30	302	282	340	299	340	340	323	319	307	285
60	215	234	265	221	298	284	290	290	279	320
120	194	194	238	213	259	251	283	283	261	266
150	210	219	253	220	223	225	275	274	281	272
180	225	211	327	340	298	291	284	284	273	297
210	235	267	262	273	320	322	296	291	286	279
240	306	259	243	249	299	316	314	316	307	309
300	169	150	296	255	320	303	340	333	320	286
min	1.5×10 ⁸ particles per mouse									
0	212	213	251	252	277	277	281	284	332	330
30	303	340	238	269	209	192	239	233	302	282
60	245	263	282	252	232	217	250	224	290	279
120	208	229	259	257	210	225	251	307	259	274
150	167	160	276	276	215	198	236	225	261	270
180	190	220	291	289	225	258	183	233	269	269
210	214	192	287	301	212	201	182	174	274	284

240	246	204	269	300	276	249	166	155	279	290
300	239	235	279	286	212	273	193	197	291	286
min	1.5×10 ¹⁰ particles per mouse									
0	298	254	271	253	292	281	299	274	310	302
30	340	333	166	156	258	272	198	176	192	192
60	207	205	75	78	197	169	89	82	179	189
120	209	199	95	91	212	210	187	285	192	196
150	188	190	166	186	220	240	219	211	227	219
180	191	194	151	165	218	232	229	221	231	228
210	237	219	162	156	221	229	235	230	239	239
240	226	175	211	199	229	231	239	231	228	240
300	221	202	209	206	237	230	238	229	242	241
min	1.5×10 ¹² particles per mouse									
0	284	271	283	294	277	245	238	226	231	233
30	115	118	151	158	246	282	105	85	94	83
60	125	113	116	126	215	234	85	104	90	71
120	124	138	137	138	194	194	115	118	56	58
150	133	130	149	121	210	219	129	135	39	44
180	151	161	181	143	155	151	176	169	162	156
210	216	218	194	197	195	207	194	194	176	197
240	222	226	174	154	176	199	194	196	188	197
300	229	207	209	204	179	210	195	193	202	204

[00159] Conclusion: From the results of random blood glucose, it could be seen that the exosomes loaded with liraglutide can significantly reduce blood glucose.

5 Example 11

[00160] By electroporation, liraglutide (not limited to liraglutide, but also including insulin degludec, insulin detemir, semaglutide and tirzepatide) was loaded into milk exosomes and purified to obtain drug-loaded exosomes, which were prepared into sublingual tablets. Here, the results of milk exosomes loaded with liraglutide are illustrated as an example.

- 5 **[00161]** Diabetic mice were fasted for 6 h, and then orally administered with different concentrations of milk exosomes loaded with liraglutide. The fasting blood glucose of the mice was detected.

[00162] The results are shown in FIG. 23 and Table 17.

Table 17

h	blank									
0	448.2	390.6	342	284.4	433.8	435.6	462.6	441	342	351
0.25	457.2	439.2	423	473.4	468	522	525.6	570.6	466.2	487.8
0.5	460.8	392.4	399.6	361.8	459	451.8	599.4	585	473.4	495
1	612	612	374.4	421.2	493.2	486	612	612	486	495
2	612	612	365.4	342	480.6	473.4	428.4	403.2	451.8	466.2
3	430.2	376.2	448.2	478.8	612	586.8	493.2	451.8	388.8	361.8
4	504	444.6	376.2	381.6	522	568.8	363.6	401.4	322.2	313.2
5	338.4	329.4	493.2	459	531	495	376.2	340.2	259.2	246.6
6	612	545.4	329.4	309.6	388.8	322.2	379.8	354.6	327.6	298.8
h	i.v. 1.5×10 ¹⁰ particles per mouse									
0	594	525.6	331.2	342	448.2	466.2	518.4	502.2	421.2	392.4
0.25	450	424.8	241.2	282.6	385.2	424.8	480.6	466.2	277.2	277.2
0.5	361.8	419.4	259.2	194.4	338.4	334.8	428.4	446.4	160.2	190.8
1	196.2	235.8	111.6	109.8	230.4	264.6	450	457.2	127.8	117
2	122.4	108	82.8	82.8	253.8	257.4	417.6	415.8	142.2	126

3	64.8	73.8	66.6	55.8	88.2	90	419.4	417.6	72	86.4
4	55.8	37.8	50.4	45	45	48.6	399.6	370.8	52.2	52.2
5	50.4	45	50.4	55.8	36	36	279	271.8	54	55.8
6	54	43.2	59.4	57.6	34.2	32.4	212.4	230.4	41.4	36
h	Oral 1.5×10^{12} particles per mouse									
0	502.2	428.4	352.8	351	414	451.8	435.6	471.6	351	423
0.25	516.6	599.4	558	504	532.8	612	612	576	612	612
0.5	511.2	487.8	379.8	437.4	496.8	453.6	457.2	466.2	428.4	365.4
1	612	594	180	216	329.4	300.6	257.4	244.8	228.6	223.2
2	457.2	462.6	151.2	135	210.6	271.8	262.8	210.6	257.4	226.8
3	417.6	498.6	120.6	115.2	241.2	235.8	196.2	153	185.4	199.8
4	462.6	493.2	95.4	88.2	196.2	136.8	192.6	169.2	163.8	169.2
5	336.6	347.4	73.8	66.6	118.8	108	95.4	106.2	154.8	160.2
6	286.2	316.8	77.4	64.8	70.2	73.8	100.8	90	169.2	156.6
h	Oral 1.5×10^{10} particles per mouse									
0	462.6	428.4	469.8	428.4	415.8	369	426.6	482.4	324	273.6
0.25	612	612	450	451.8	612	576	540	532.8	318.6	356.4
0.5	318.6	284.4	417.6	426.6	612	612	500.4	477	315	396
1	552.6	592.2	379.8	372.6	426.6	469.8	612	581.4	388.8	385.2
2	570.6	581.4	354.6	336.6	612	590.4	552.6	500.4	266.4	250.2
3	487.8	415.8	475.2	480.6	592.2	554.4	367.2	383.4	158.4	145.8
4	577.8	583.2	221.4	266.4	612	612	273.6	288	154.8	149.4
5	455.4	432	358.2	295.2	532.8	538.2	304.2	279	135	133.2
6	594	612	612	612	349.2	410.4	468	482.4	153	151.2

[00163] Conclusion: From the results of the fasting blood glucose, it could be seen that the exosomes loaded with liraglutide can significantly reduce blood glucose.

5 **[00164]** The use of a milk exosome in the preparation of a drug carrier provided by the present disclosure is described in detail above. The principle and implementation of the present disclosure are illustrated by using specific embodiments herein. The above descriptions of the embodiments are only used to facilitate understanding of the method and the core idea of the present disclosure. It should be noted that, several improvements and modifications may be made
10 by those skilled in the art to the present disclosure without departing from the principle of the present disclosure, and these improvements and modifications also fall within the protection scope of the claims of the present disclosure.

CLAIMS

1. Use of a milk exosome in the preparation of a drug carrier.
2. The use according to claim 1, wherein the drug includes a protein drug.
- 5 3. The use according to claim 1 or 2, wherein the drug includes liraglutide.
4. Use of a milk exosome as a drug carrier in the preparation of a medicament for the prevention and/or treatment of a disease.
5. The use according to claim 4, wherein the drug includes a protein drug.
6. The use according to claim 4 or 5, wherein the drug includes liraglutide.
- 10 7. The use according to any one of claims 4 to 6, wherein the drug is administered orally.
8. The use according to any one of claims 4 to 7, wherein the disease includes type 2 diabetes.
9. An oral medicament, comprising an active ingredient and a milk exosome as a drug carrier.
- 15 10. The use according to any one of claims 1 to 8 or the oral medicament according to claim 9, wherein the milk exosome is prepared by
 - step 1, pre-treatment;
 - step 2, preliminary purification and concentration;
 - step 3, refined purification;- 20 wherein the pre-treatment comprises fat removal and/or protein removal,
the preliminarily purification and concentration is performed by tangential flow ultrafiltration, and
the refined purification are performed by a method selected from the group consisting of CIMmultus QA column purification, Capto core 700 column purification, S400 molecular sieve
25 chromatography purification, and a combination thereof.
- 11. A method for treating a disease, comprising administering to a subject a drug loaded in a

milk exosome, wherein the milk exosome is used as a drug carrier.

12. The method according to claim 11, wherein the drug includes a protein drug.

13. The method according to claim 12, wherein the drug includes liraglutide.

14. The method according to any one of claims 11 to 13, wherein the drug is administered
5 orally.

15. The method according to any one of claims 11 to 14, wherein the disease includes type 2
diabetes.