

This Sheet Has Been Replaced



3-(SUBSTITUTED KETONE AND ALCOHOL)-2-(2-IMIDAZOLIN-2-YL) PYRIDINE AND QUINOLINE HERBICIDAL AGENTS

BACKGROUND OF THE INVENTION

Weeds cause tremendous global economic losses by reducing crop yields and lowering crop values. In the United States, crops must compete with 1,800 different weeds. Accordingly, there is an ongoing search in the art to create more effective and/or more selective herbicidal agents.


U.S. Patent 4,798,619 discloses a variety of effective herbicidal agents including 3-(benzoyl and hydroxyalkyl)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds. While these compounds and other known herbicides may be effective herbicidal agents, there are increasing concerns with the environment and therefore increasing efforts have been undertaken to find more potent herbicides, i.e. those herbicides which can be applied at a lower rate, but still give the same or better control of a broad spectrum of undesirable plant species.

It is an object of the present invention to provide compounds that are highly effective for controlling undesirable plant species.


SUMMARY OF THE INVENTION

The present invention describes 3-(substituted ketone and alcohol)-2-(2-imidazolin-2-yl)-pyridine and quinoline compounds that are highly effective herbicidal agents useful for the control of undesirable plant species and methods for their preparation.

The 3-(substituted ketone)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds of the present invention are illustrated as formula I and the 3-(substituted alcohol)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds of the present invention are illustrated as formula II:



(I)




(II)

wherein

R is C₁-C₄ alkyl;

R₁ is C₁-C₄ alkyl or C₃-C₆ cycloalkyl; and when R and R₁ are taken together with the carbon to which they are attached they may represent C₃-C₆ cycloalkyl optionally substituted with methyl;

Q is



R_2 , R_3 and R_4 are each independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy or phenyl optionally substituted with one or two halogen atoms, C_1 - C_4 alkyl or haloalkyl groups or C_1 - C_4 alkoxy or haloalkoxy groups.

m is an integer of 1 or 2;

A is O, S, or NR_5 ; provided that when m is 2, A is O;

R_5 is C_1 - C_4 alkyl;

B is hydrogen, COR_6 or SO_2R_7 ;

R_6 is C_1 - C_{11} alkyl, chloromethyl or phenyl optionally substituted with one chloro, one nitro or one methoxy group;

R_7 is C_1 - C_4 alkyl or phenyl optionally substituted with one methyl group;

W is O or S;

X is hydrogen, halogen or methyl;

Y and Z are each independently hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_4 alkoxymethyl, C_1 - C_6 alkoxy, C_1 - C_4 alkylthio, phenoxy, C_1 - C_4 haloalkyl, nitro, cyano, C_1 - C_4 alkylsulfonyl or phenyl optionally substituted with one C_1 - C_4 alkyl, C_1 - C_4 alkoxy or halogen; and, when taken together, Y and Z may form a ring in which YZ are represented by the structure: $-(CH_2)_n-$, where n is an integer selected from 3 and 4; or




where R_8 , R_9 , R_{10} and R_{11} are each independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, nitro, cyano, phenyl, phenoxy, amino, chlorophenyl, methylphenyl, or phenoxy substituted with one chloro, trifluoromethyl, nitro or methyl group, with the proviso that only one of R_8 , R_9 , R_{10} or R_{11} , may represent a substituent other than hydrogen, halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy; and when R and R_1 are not the same, the optical isomers thereof.

While 3-(benzoyl and hydroxyalkyl)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds have been known, surprisingly, the 3-(substituted ketone and alcohol)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds of the present invention give better control of undesirable plant species at low rates.

DETAILED DESCRIPTION OF THE INVENTION

Surprisingly, it has been found that the formula I and formula II compounds of this invention are more potent herbicidal agents at lower rates, especially postemergence, than the 3-(benzoyl and hydroxyalkyl)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds known in the art. Advantageously, compounds of the present invention also control a wider variety of undesirable plant species at those lower rates than the 3-(benzoyl and hydroxyalkyl)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds. Naturally, a low rate is desirable because it means increased safety for the farmer and the environment.

Preferred 3-(substituted ketone and alcohol)-2-(2-imidazolin-2-yl)pyridine and quinoline compounds of the present invention that are especially effective in the control of undesirable plant species are illustrated as formulas III and IV:



wherein R_2 , R_3 , R_4 and m are as described above; A is O or S ;

Y and Z are each independently hydrogen, halogen, C_1-C_6

alkyl, C₁-C₄ alkoxyethyl or C₁-C₆ alkoxy, and when taken together, Y and Z may form a ring in which YZ are represented by the structure:



5 Especially preferred compounds of the present invention are formula III compounds wherein R₃ is hydrogen and R₂ and R₄ are independently hydrogen, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy or C₁-C₄ haloalkoxy.

10 Exemplary of halogen hereinabove are fluorine, chlorine, bromine and iodine. The term "haloalkyl" is defined as a compound of the following formula C_pH_qM_r where M is halogen and p, q and r are integers such that q+r = 2p+1. The term "haloalkoxy" is defined as a compound of the following formula

15 C_pH_qM_rO where M is halogen and p, q and r are integers such that q+r = 2p+1.


 Compounds of formula I and formula II, wherein m is 2, may be prepared as shown in Flow Diagram I.

20

25

30

FLOW DIAGRAM I




The appropriately substituted formula V 2-(2-imidazolin-2-yl)pyridine or quinoline acid is reacted with about one equivalent of a dehydrating agent such as diisopropylcarbodiimide or dicyclohexylcarbodiimide (DCC) in the presence of an aprotic solvent such as methylene chloride or tetrahydrofuran preferably at a temperature between about 20°C to 32°C to form the formula VI compound. Said formula VI compound is then reacted with one or more equivalents of the appropriately substituted formula VII lithium agent in the presence of an organic solvent such as

tetrahydrofuran preferably at a temperature between about -78°C to 30°C to obtain the desired formula I and formula II compounds.

Compounds of formula II wherein m is 1 may be prepared as shown in Flow Diagram II.


FLOW DIAGRAM II



The appropriately substituted formula VIII imidazolinone compound is reacted with at least about 2.0 molar equivalents of a $\text{C}_1\text{-C}_4$ alkyl lithium in the presence of an inert solvent such as tetrahydrofuran preferably at a temperature between about -80°C to 0°C . The thus-formed salt is then treated with at least about 1.0 molar equivalents of the appropriately substituted formula IX aldehyde compound in an inert solvent such as tetrahydrofuran preferably at a temperature between about -80°C to 40°C to obtain the desired formula II compound.

Advantageously, the thus-formed formula II compound wherein m is 1 is reacted with an oxidizing agent such as manganese(IV) oxide in the presence of an inert organic solvent such as chloroform preferably at a temperature between about 20°C to 80°C to give a formula I compound. The reaction scheme is illustrated in Flow Diagram III.

FLOW DIAGRAM III



Compounds of formulas V, VI and VIII and methods of preparation thereof are described in U.S. Patent No. 4,798,619.

Preparation of formula I and formula II N-substituted derivatives, wherein B is COR₆ or SO₂R₇; m is 2; and R, R₁, Q, R₂, R₃, R₄, A, R₅, W, X, Y and Z are as described above, can be achieved by reaction of the appropriately substituted formula I or formula II compound wherein B is hydrogen with an excess of acyl halide, acyl anhydride or sulfonyl halide, alone or in a solvent such as pyridine or toluene at a temperature between about 50°C to 125°C.

Formula II compounds wherein m is 1 and B is COR₆ or SO₂R₇ may be prepared by reducing formula I compounds wherein B is COR₆ or SO₂R₇ with a reducing agent such as sodium borohydride.

The formula I 3-(substituted ketone)-2-(2-imidazolin-2-yl)pyridines and quinolines and the formula II 3-(substituted alcohol)-2-(2-imidazolin-2-yl)pyridines and quinolines of the present invention are effective herbicidal agents useful for the control of a wide variety of herbaceous and woody, annual and perennial, monocotyledonous and dicotyledonous plants. These compounds are effective for controlling weeds

native to both dry land and wet land areas. The compounds are effective in controlling the above-said plants when applied to the foliage thereof or to soil or water containing seeds or other propagating organs of said plants such as stolons, tubers or rhizomes, at rates of from about 0.016 to 4.0 kg/ha and preferably from about 0.125 to 4.0 kg/ha.

Advantageously, it has been found that the compounds of this invention are best suited for use as broad spectrum herbicides, especially when applied postemergence to the locus in which weed control is desired. This is not to say, however, that all of the compounds of this invention are non-selective. In fact, some of the compounds of this invention are selective in leguminous crops, such as soybeans.


The formula I and formula II compounds of this invention may be applied to crops in the form of a solid or liquid herbicidal composition, comprising a herbicidally effective amount of the formula I or formula II compound dispersed or dissolved in an inert solid or liquid carrier. The formulations may be applied as preemergence or postemergence treatments.

Advantageously, the formula I and formula II compounds can be formulated as emulsifiable concentrates, wettable powders, granular formulations, flowable concentrates and the like.

In order to facilitate a further understanding of the invention, the following examples are presented to illustrate more specific details thereof. The invention is not to be limited thereby except as defined in the claims.

EXAMPLE 1


Preparation of 4-Isopropyl-4-methyl-2-[5-methyl-3-(2-thienoyl)-2-pyridyl]-2-imidazolin-5-one




A solution of thiophene (3.3 mL) in anhydrous tetrahydrofuran (150 mL) is cooled to -78°C and treated with a 2.5 molar solution of n-butyllithium in hexanes (16.8 mL) for 30 minutes. 2-Isopropyl-2,7-dimethyl-5H-imidazo[1',2':1,2]pyrrolo[3,4-b]pyridine-3(2H),5-dione (10.19 g, 0.0396 mol) is added and the reaction mixture is stirred at room temperature for three days, diluted with water and extracted with ethyl acetate. The combined organic extracts are washed with brine, dried (Na_2SO_4) and concentrated in vacuo to obtain a colorless oil. Chromatography of the oil using silica gel and a 30% hexane in diethyl ether mixture gives 2-[3-(hydroxydi-2-thienylmethyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one as a white solid (1.0 g, mp $190-191^\circ\text{C}$) and 4-isopropyl-4-methyl-2-[5-

methyl-3-(2-thenoyl)-2-pyridyl]-2-imidazolin-5-one as a white solid (0.45 g, mp 168-170°C).

Using essentially the same procedure and employing the appropriately substituted 5H-imidazo-[1',2':1,2]pyrrolo[3,4-b]pyridine-3(2H),5-dione and the appropriate thiophene or furan, the following compounds are obtained.




A	Y	Z	mp°
O	H	H	145 - 146
S	H	H	144 - 145
O	CH ₃	H	159 - 160
O	C ₂ H ₅	H	108 - 110
S	C ₂ H ₅	H	153 - 154
O	CH ₂ OCH ₃	H	123 - 125
S	CH ₂ OCH ₃	H	144 - 147
S	-CH=CH-CH=CH-		235 - 238



A	Y	Z	mp ^o
O	H	H	138 - 139
O	CH ₃	H	133 - 134
O	CH ₂ OCH ₃	H	138 - 140

EXAMPLE 2


Preparation of 2-[3-(α -Hydroxy-5-methyl-2-thenyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one



A stirred solution of 4-isopropyl-4-methyl-2-(5-methyl-2-pyridyl)-2-imidazolin-5-one (10.0 g, 0.043 mol) in anhydrous tetrahydrofuran (200 mL) is

cooled to -78°C and treated with a 1.5 molar solution of methyl lithium in diethyl ether (60.6 mL). The resultant suspension is stirred at -78°C for 30 minutes and at 0°C for 30 minutes. The reaction mixture is recooled to -78°C and 5-methyl-2-thiophenecarboxaldehyde (15.35 g, 0.043 mol) is added dropwise. After 30 minutes, the cooling bath is removed and the reaction mixture is stirred at room temperature overnight then diluted with water. The aqueous mixture is acidified to pH 6 with acetic acid and extracted with diethyl ether. The combined organic extracts are washed with water, dried (Na_2SO_4) and concentrated in vacuo to obtain an orange oil. Chromatography of the oil using silica gel and a 20% diethyl ether in methylene chloride mixture gives a yellow oil which is triturated with diethyl ether to obtain the title product as a white solid (2.0 g, mp $147-148^{\circ}\text{C}$).

Using essentially the same procedure and employing the appropriately substituted 2-(2-pyridyl)-2-imidazolin-5-one and the appropriate 2-thiophenecarboxaldehyde or 2-furaldehyde, the following compounds are obtained:



5


10

A	Y	Z	R ₂	R ₃	R ₄	mp ^o C
O	H	H	H	H	H	151 - 158
O	H	H	CH ₃	H	H	glass
S	H	H	H	H	CH ₃	glass
O	CH ₃	H	H	H	H	163 - 175
O	CH ₃	H	CH ₃	H	H	130 - 135
O	CH ₃	H	C ₂ H ₅	H	H	oil
S	CH ₃	H	H	H	CH ₃	123 - 127
O	C ₂ H ₅	H	H	H	H	115 - 121
S	C ₂ H ₅	H	H	H	H	140 - 155
O	C ₂ H ₅	H	CH ₃	H	H	liquid
S	C ₂ H ₅	H	CH ₃	H	H	120 - 122
O	C ₂ H ₅	H	C ₂ H ₅	H	H	118 - 121
S	C ₂ H ₅	H	H	H	CH ₃	121 - 123
S	C ₂ H ₅	H	Cl	H	H	131 - 133
S	-CH=CH-CH=CH-	H	H	H	H	168 - 170
S	-CH=CH-CH=CH-	CH ₃	H	H	H	179 - 185

30


EXAMPLE 3

Preparation of 4-Isopropyl-4-methyl-2-[5-methyl-3-(5-methyl-2-thenoyl)-2-pyridyl]-2-imidazolin-5-one



A solution of 2-[(3-(α-hydroxy-5-methyl-2-thenyl)-5-methyl-2-pyridyl)-4-isopropyl-4-methyl-2-imidazolin-5-one (1.35 g, 0.0038 mol) in chloroform (75 mL) is treated with manganese dioxide (10 g, 0.12 mol) and heated to 60°C. After 10 minutes, additional manganese dioxide (10 g) is added and the reaction mixture is stirred at 60°C for 2 hours, cooled to room temperature and filtered through diatomaceous earth. The filtrate is concentrated in vacuo to obtain a residue which is crystallized from diethyl ether to give the title product as a white solid (1.2 g, mp 173-175°C).

Using essentially the same procedure and employing the appropriately substituted 2-{3-[α-hydroxy-2-(thenyl or furfuryl)]-2-pyridyl}-2-imidazolin-5-one, the following compounds are obtained:



A	Y	R ₂	R ₃	R ₄	mp ^o C
S	H	H	H	CH ₃	150 - 152
S	CH ₃	H	H	CH ₃	173 - 178
O	C ₂ H ₅	CH ₃	H	H	106 - 108
S	C ₂ H ₅	CH ₃	H	H	140 - 142
S	C ₂ H ₅	H	H	CH ₃	glass
S	C ₂ H ₅	Cl	H	H	glass

EXAMPLE 4

Preemergence herbicidal evaluation of test compounds

The preemergence herbicidal activity of the compounds of the present invention is exemplified by the following tests in which the seeds of a variety of monocotyledonous and dicotyledonous plants are separately mixed with potting soil and planted on top of approximately one inch of soil in separate pint cups. After planting, the cups are sprayed with the selected aqueous acetone solution containing test compound in sufficient quantity to provide the equivalent of about 0.125 to 4.0 kg per hectare of test compound per cup. The treated cups are then placed on greenhouse benches, watered and cared for in accordance with conventional greenhouse procedures. From four to five weeks after

treatment, the tests are terminated and each cup is examined and rated according to the rating system set forth below. Data obtained are reported in Table I below. Where more than one test is involved for a given compound, the data are averaged.

Plant species employed in these evaluations are reported by header abbreviation, common name and scientific name.

Compounds employed in this preemergence herbicidal evaluation and in the post-emergence evaluation in the following example are given a compound number and identified by name. Data in Table I are reported by compound number.

Herbicide Rating Scale

Results of herbicide evaluation are expressed on a rating scale (0-9). The scale is based upon a visual observation of plant stand, vigor, malformation, size, chlorosis and overall plant appearance as compared with a control.

	<u>Rating</u>	<u>Meaning</u>	<u>% Control Compared To Check</u>
	9	Complete Kill	100
	8	Approaching Complete kill	91 - 99
	7	Good Herbicidal Effect	80 - 90
	6	Herbicidal Effect	65 - 79
	5	Definite Injury	45 - 64
	4	Injury	30 - 44
	3	Moderate Effect	16 - 29
	2	Slight Effect	6 - 15
	1	Trace Effect	1 - 5
	0	No Effect	0

PLANT SPECIES EMPLOYED IN HERBICIDAL EVALUATIONS

	<u>Header Abb.</u>	<u>Common Name</u>	<u>Scientific Name</u>
	Barnydgr	Barnyardgrass	ECHINOCHLOA CRUS-GALLI, (L) BEAU
5	Foxtail Sp	Foxtail Spp.	SETARIA SPP.
	Wild Oats	Oat, Wild	AVENA FATUA, L.
	Quackgrass	Quackgrass	AGROPYRON REPENS, (L) BEAUV
	Fld Bindwd	Bindweed, Field	CONVOLVULUS ARVENSIS, (RHIZOME) L.
10	Mrnglry Sp	Morningglory Spp.	IPOMOEA SPP.
	Wild Mustd	Mustard, Wild	BRASSICA KABER, (DC) L. C. WHEELR
	Velvetleaf	Velvetleaf	ABUTILON THEOPHRASTI, MEDIC.
15	Wht Fenman	Wheat, Fenman	TRITICUM AESTIVUM, FENMAN
	Corn Field,	Corn, Field	ZEA MAYS, L.
	Cotton	Cotton	GOSSYPIUM HIRSUTUM, L.
20	Soybean	Soybean	GLYCINE MAX

COMPOUNDS EVALUATED AS HERBICIDAL AGENTS

Compound

No.

- | | | |
|----|----|---|
| 5 | 1 | 2-[3-(2-Furoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one |
| | 2 | 4-Isopropyl-4-methyl-2-[3-(3-methyl-2-thenoyl)-2-pyridyl]-2-imidazolin-5-one |
| 10 | 3 | 4-Isopropyl-4-methyl-2-[3-(2-thenoyl)-2-pyridyl]-2-imidazolin-5-one |
| | 4 | 2-[3-(2-Furoyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one |
| 15 | 5 | 4-Isopropyl-4-methyl-2-[5-methyl-3-(2-thenoyl)-2-pyridyl]-2-imidazolin-5-one |
| | 6 | 4-Isopropyl-4-methyl-2-[5-methyl-3-(5-methyl-2-thenoyl)-2-pyridyl]-2-imidazolin-5-one |
| 20 | 7 | 4-Isopropyl-4-methyl-2-[5-methyl-3-(3-methyl-2-thenoyl)-2-pyridyl]-2-imidazolin-5-one |
| | 8 | 2-[5-Ethyl-3-(2-furoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one |
| 25 | 9 | 2-[5-Ethyl-3-(2-thenoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one |
| | 10 | 2-[5-Ethyl-3-(5-methyl-2-furoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one |
| 30 | | |
| | | |
| 35 | | |

Compound		
No.		
11	2-[5-Ethyl-3-(5-methyl-2-thenoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one	
5	12	2-[5-Ethyl-3-(3-methyl-2-thenoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
10	13	2-[3-(5-Chloro-2-thenoyl)-5-ethyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	14	2-[3-(2-Furoyl)-5-(methoxymethyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
15	15	4-Isopropyl-2-[5-(methoxymethyl)-3-(2-thenoyl)-2-pyridyl]-4-methyl-2-imidazolin-5-one
	16	4-Isopropyl-4-methyl-2-[3-(2-thenoyl)-2-quinolyl]-2-imidazolin-5-one
20	17	2-[3-(α -Hydroxyfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
25	18	2-[3-(α -Hydroxy-5-methylfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	19	2-[3-(α -Hydroxy-3-methyl-2-thenyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
30	20	2-[3-(α -Hydroxyfurfuryl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one

Compound	<u>No.</u>	
5	21	2-[3-(α -Hydroxy-5-methylfurfuryl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	22	2-[3-(α -Hydroxy-5-methyl-2-thenyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
10	23	2-[3-(5-Ethyl- α -Hydroxyfurfuryl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
15	24	2-[3-(α -Hydroxy-3-methyl-2-thenyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
20	25	2-[5-Ethyl-3-(α -hydroxyfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	26	2-[5-Ethyl-3-(α -hydroxy-2-thenyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
25	27	2-[5-Ethyl-3-(α -hydroxy-5-methylfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
30	28	2-[5-Ethyl-3-(α -hydroxy-5-methyl-2-thenyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one

Compound	No.	
5	29	2-[5-Ethyl-3-(5-ethyl- α -hydroxyfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	30	2-[5-Ethyl-3-(α -hydroxy-3-methyl-2-thenyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
10	31	2-[3-(5-Chloro- α -hydroxy-2-thenyl)-5-ethyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
15	32	2-[3-(α -Hydroxy-2-thenyl)-2-quinolyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
	33	2-[3-(α -Hydroxy-5-methyl-2-thenyl)-2-quinolyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
20	34	2-{3-[(di-2-Furyl)hydroxymethyl]-2-pyridyl}-4-isopropyl-4-methyl-2-imidazolin-5-one
25	35	2-[3-(di-2-Furylhydroxymethyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one
30	36	2-[3-(di-2-Furylhydroxymethyl)-5-(methoxymethyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one

TABLE I

Preemergence Herbicidal Evaluation of Test Compounds

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
1	4.000	8.0	-	7.0	9.0	9.0	8.0	9.0	8.0	-	-	-	-
	.500	7.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	6.0	9.0	9.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	7.0
2	4.000	9.0	-	8.0	9.0	9.0	8.0	9.0	8.0	-	-	-	-
	.500	7.0	9.0	9.0	7.0	9.0	7.0	9.0	8.0	9.0	9.0	9.0	8.0
	.125	1.4	4.0	0.0	2.0	4.0	1.7	8.0	2.0	8.0	7.0	4.0	0.0
3	4.000	8.0	-	7.0	9.0	9.0	7.0	9.0	8.0	-	-	-	-
	.500	7.0	9.0	9.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	7.0
	.125	5.0	5.0	7.0	9.0	9.0	5.0	9.0	8.0	7.0	4.0	7.0	4.0
4	.500	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	6.0	7.0	9.0	0.0	9.0	6.0	9.0	8.0	6.0	8.0	8.0	8.0
5	.500	8.0	8.0	8.0	0.0	8.0	8.0	9.0	8.0	5.0	8.0	6.0	7.0
	.125	0.0	2.0	0.0	0.0	0.0	0.0	8.0	2.0	0.0	0.0	0.0	0.0

TABLE I (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
6	4.000	8.0	-	8.0	9.0	9.0	8.0	9.0	9.0	-	-	-	-
	.500	5.0	4.0	1.5	1.5	4.5	4.7	4.5	4.0	1.5	2.0	3.0	2.0
	.125	1.4	1.5	0.0	0.0	4.5	0.7	4.0	3.5	0.0	0.0	1.0	0.5
7	.500	8.0	9.0	3.0	2.0	9.0	8.0	9.0	9.0	0.0	2.0	8.0	2.0
	.125	0.0	0.0	0.0	0.0	4.0	0.0	7.0	2.0	0.0	0.0	0.0	0.0
8	.500	8.0	8.0	8.0	9.0	9.0	7.0	9.0	8.0	6.0	7.0	7.0	0.0
	.125	0.0	0.0	4.0	0.0	9.0	0.0	9.0	7.0	0.0	2.0	0.0	0.0
9	.500	6.0	2.0	4.0	9.0	9.0	0.0	9.0	5.0	1.0	2.0	3.0	0.0
	.125	0.0	0.0	0.0	0.0	9.0	0.0	6.0	0.0	0.0	0.0	0.0	0.0
10	4.000	8.0	-	9.0	9.0	9.0	8.0	9.0	9.0	-	-	-	-
	.500	8.0	9.0	9.0	9.0	9.0	8.0	9.0	9.0	8.0	8.0	9.0	2.0
	.125	3.0	3.0	1.0	3.0	9.0	4.0	9.0	8.0	1.0	2.0	4.0	0.0

TABLE I (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
11	4.000	8.0	-	8.0	8.0	9.0	7.0	9.0	8.0	-	-	-	-
	.500	1.0	2.0	1.0	1.5	4.5	1.0	5.0	3.5	1.0	1.0	2.0	0.0
	.125	0.0	0.0	0.0	0.0	0.0	0.0	2.0	0.5	0.0	0.0	0.0	0.0
12	.500	1.0	4.0	0.0	4.0	9.0	0.0	9.0	8.0	0.0	0.0	2.0	0.0
	.125	0.0	0.0	0.0	0.0	6.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0
13	.500	4.0	4.0	4.0	5.0	9.0	6.0	8.0	7.0	2.0	3.0	2.0	0.0
	.125	0.0	4.0	0.0	0.0	2.0	0.0	8.0	4.0	0.0	0.0	-	0.0
14	.500	7.0	8.0	9.0	9.0	9.0	7.0	9.0	8.0	9.0	9.0	9.0	2.0
	.125	0.0	2.0	7.0	6.0	4.0	0.0	7.0	4.0	8.0	7.0	0.0	0.0
15	.500	4.0	8.0	9.0	7.0	9.0	4.0	9.0	7.0	8.0	7.0	7.0	2.0
	.125	0.0	7.0	2.0	4.0	4.0	0.0	5.0	5.0	3.0	3.0	0.0	0.0

TABLE I (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
16	4.000	2.0	-	4.0	6.0	2.0	1.0	9.0	4.0	-	-	-	-
17	.500	8.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	6.0	9.0	9.0	9.0	9.0	8.0	9.0	8.0	9.0	9.0	8.0	7.0
18	4.000	9.0	-	8.0	9.0	9.0	7.0	9.0	8.0	-	-	-	-
	.500	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	2.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0	9.0	9.0	9.0	7.0
19	4.000	8.0	-	8.0	8.0	9.0	8.0	9.0	8.0	-	-	-	-
	.500	2.2	7.0	8.0	7.0	6.0	5.5	9.0	8.0	9.0	7.0	5.0	5.0
	.125	0.4	0.0	0.0	0.0	0.0	1.7	0.0	0.0	0.0	0.0	0.0	0.0
20	4.000	9.0	-	8.0	9.0	9.0	8.0	9.0	9.0	-	-	-	-
	.500	8.5	9.0	9.0	9.0	8.5	9.0	8.5	9.0	9.0	8.0	7.5	7.0
	.125	9.0	9.0	7.0	7.0	9.0	8.0	9.0	8.0	5.0	7.0	9.0	6.0

TABLE I (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
21	4.000	9.0	-	8.0	9.0	9.0	9.0	9.0	9.0	-	-	-	-
	.500	8.5	9.0	9.0	9.0	9.0	8.3	9.0	9.0	8.5	9.0	9.0	7.5
	.125	5.5	8.5	8.0	3.0	9.0	6.0	8.5	8.0	4.0	5.5	8.5	4.0
22	4.000	8.0	-	8.0	9.0	9.0	8.0	9.0	8.0	-	-	-	-
	.500	3.3	8.0	8.0	5.0	9.0	4.0	9.0	8.0	2.0	3.0	8.0	1.0
	.125	0.0	0.0	0.0	0.0	2.0	0.0	7.0	5.0	0.0	0.0	2.0	0.0
23	4.000	9.0	-	8.0	9.0	9.0	8.0	9.0	9.0	-	-	-	-
	.500	8.0	9.0	9.0	9.0	9.0	7.7	9.0	8.5	8.5	9.0	9.0	7.5
	.125	4.0	7.5	7.5	4.5	9.0	5.3	9.0	8.5	3.5	7.0	8.5	5.0
24	.500	0.0	8.0	3.0	2.0	4.0	2.0	7.0	7.0	1.0	1.0	6.0	3.0
25	.500	9.0	9.0	8.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	8.0	1.0
	.125	8.0	8.0	7.0	8.0	9.0	7.0	9.0	9.0	2.0	5.0	7.0	1.0

TABLE I (Continued)

<u>Compound</u> <u>No.</u>	<u>Rate</u> <u>(kg/ha)</u>	<u>Barny</u> <u>ardgr</u>	<u>Foxta</u> <u>il Sp</u>	<u>Wild</u> <u>oats</u>	<u>Quack</u> <u>grass</u>	<u>Fld B</u> <u>indwd</u>	<u>Mrngl</u> <u>ry Sp</u>	<u>Wild</u> <u>mustd</u>	<u>Velve</u> <u>tleaf</u>	<u>Wht F</u> <u>enman</u>	<u>Corn</u> <u>field</u>	<u>Cot</u> <u>ton</u>	<u>Soy</u> <u>bean</u>
26	4.000	7.0	-	8.0	9.0	9.0	6.0	9.0	7.0	-	-	-	-
	.500	4.0	6.0	0.0	0.0	4.0	0.0	8.0	3.0	0.0	0.0	0.0	0.0
27	.500	9.0	9.0	6.0	9.0	9.0	8.0	9.0	8.0	7.0	9.0	8.0	2.0
	.125	7.0	9.0	2.0	4.0	9.0	1.0	8.0	7.0	3.0	3.0	6.0	0.0
28	4.000	8.0	-	6.0	6.0	9.0	6.0	8.0	7.0	-	-	-	-
	.500	7.0	7.0	0.0	0.0	9.0	0.0	8.0	7.0	0.0	1.0	2.0	0.0
	.125	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0
29	.500	9.0	9.0	7.0	9.0	9.0	7.0	9.0	8.0	8.0	7.0	8.0	2.0
	.125	7.0	8.0	2.0	5.0	9.0	2.0	8.0	7.0	3.0	3.0	8.0	0.0
30	.500	2.0	6.0	2.0	2.0	8.0	2.0	7.0	7.0	1.0	4.0	3.0	3.0

TABLE I (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
31	.500	0.0	4.0	0.0	0.0	4.0	2.0	9.0	8.0	0.0	0.0	0.0	0.0
	.125	0.0	0.0	0.0	0.0	0.0	0.0	4.0	0.0	0.0	0.0	0.0	0.0
32	4.000	9.0	-	9.0	8.0	9.0	6.0	9.0	6.0	-	-	-	-
33	4.000	8.0	-	7.0	9.0	9.0	8.0	9.0	6.0	-	-	-	-
	.500	0.0	0.0	1.0	6.0	1.0	2.0	8.0	5.0	1.0	7.0	4.0	0.0
34	4.000	8.0	-	7.0	9.0	9.0	6.0	9.0	7.0	-	-	-	-
	.500	8.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0	9.0	7.0
	.125	6.0	9.0	9.0	8.0	8.0	7.0	9.0	8.0	8.0	8.0	8.0	3.0
35	.500	7.0	8.0	9.0	6.0	9.0	8.0	9.0	9.0	6.0	9.0	7.0	7.0
	.125	0.0	7.0	3.0	2.0	9.0	4.0	8.0	7.0	0.0	2.0	2.0	1.0

TABLE I (Continued)

<u>Compound</u> <u>No.</u>	<u>Rate</u> <u>(kg/ha)</u>	<u>Barny</u> <u>ardgr</u>	<u>Foxta</u> <u>il Sp</u>	<u>Wild</u> <u>oats</u>	<u>Quack</u> <u>grass</u>	<u>Fld B</u> <u>indwd</u>	<u>Mrngl</u> <u>ry Sp</u>	<u>Wild</u> <u>mustd</u>	<u>Velve</u> <u>tleaf</u>	<u>Wht F</u> <u>enman</u>	<u>Corn</u> <u>field</u>	<u>Cot</u> <u>ton</u>	<u>Soy</u> <u>bean</u>
36	.500	4.0	9.0	8.0	9.0	9.0	7.0	9.0	8.0	9.0	9.0	7.0	2.0
	.125	0.0	4.0	4.0	4.0	2.0	0.0	5.0	3.0	6.0	2.0	4.0	0.0

EXAMPLE 5

Postemergence herbicidal evaluation of test compounds

The postemergence herbicidal activity of the compounds of the present invention is demonstrated by the following tests, wherein a variety of monocotyledonous and dicotyledonous plants are treated with test compounds, dispersed in aqueous acetone mixtures. In the tests, seedling plants are grown in jiffy flats for about two weeks. The test compounds are dispersed in 50/50 acetone/water mixtures containing 0.5% TWEEN® 20, a polyoxyethylene sorbitan monolaurate surfactant of Atlas Chemical Industries, in sufficient quantities to provide the equivalent of about 0.125 kg to 4.0 kg per hectare of active compound when applied to the plants through a spray nozzle operating at 40 psi for a predetermined time. After spraying, the plants are placed on greenhouse benches and are cared for in the usual manner, commensurate with conventional greenhouse practices. From four to five weeks after treatment, the seedling plants are examined and rated according to the rating system provided in Example 4 above. The data obtained are recorded in Table II below. The compounds evaluated are reported by compound number given in Example 4.

TABLE II

Postemergence Herbicidal Evaluation of Test Compounds

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
1	4.000	9.0	-	9.0	9.0	9.0	9.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0
	.125	8.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0	7.0
2	4.000	9.0	-	8.0	0.0	6.0	6.0	9.0	8.0	-	-	-	-
	.500	7.4	9.0	9.0	6.0	7.0	7.7	9.0	7.0	9.0	9.0	8.0	7.0
	.125	4.4	8.0	9.0	4.0	6.0	6.3	9.0	8.0	8.0	9.0	8.0	7.0
3	4.000	9.0	-	9.0	9.0	9.0	9.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	7.0	9.0	9.0	8.0	8.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
4	.500	9.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	9.0	9.0	8.0	3.0	8.0	9.0	9.0	9.0	9.0	8.0	7.0	3.0
5	.500	8.0	9.0	9.0	4.0	9.0	9.0	9.0	9.0	9.0	9.0	7.0	7.0
	.125	8.0	9.0	6.0	2.0	9.0	8.0	9.0	9.0	7.0	9.0	7.0	7.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
6	4.000	9.0	-	9.0	1.0	8.0	6.0	9.0	8.0	-	-	-	-
	.500	8.3	9.0	9.0	3.5	9.0	8.7	9.0	9.0	8.5	8.5	9.0	7.0
	.125	6.5	8.0	8.0	1.0	7.5	7.3	7.0	8.0	7.0	6.5	8.5	4.5
7	.500	8.0	9.0	8.0	0.0	8.0	7.0	8.0	9.0	9.0	8.0	8.0	7.0
	.125	8.0	8.0	8.0	0.0	2.0	7.0	7.0	8.0	6.0	5.0	8.0	0.0
8	.500	7.0	8.0	3.0	9.0	9.0	5.0	9.0	9.0	9.0	9.0	4.0	2.0
	.125	4.0	2.0	2.0	4.0	9.0	4.0	9.0	3.0	8.0	8.0	3.0	0.0
9	.500	4.0	7.0	4.0	7.0	9.0	4.0	9.0	6.0	2.0	3.0	0.0	0.0
	.125	2.0	2.0	2.0	0.0	4.0	0.0	8.0	0.0	0.0	0.0	0.0	0.0
10	4.000	9.0	-	9.0	7.0	9.0	9.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	9.0	4.0	8.0	8.0	9.0	9.0	9.0	9.0	9.0	3.0
	.125	8.0	9.0	8.0	2.0	8.0	6.0	9.0	8.0	6.0	7.0	9.0	1.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
11	4.000	9.0	-	9.0	4.0	9.0	8.0	9.0	9.0	-	-	-	-
	.500	8.0	8.0	6.5	3.0	7.0	6.0	9.0	7.5	6.0	7.5	7.0	0.5
	.125	6.5	7.0	4.0	1.0	5.5	2.0	8.0	6.0	3.0	2.5	5.5	0.5
12	.500	8.0	8.0	6.0	0.0	8.0	4.0	8.0	7.0	2.0	2.0	7.0	0.0
	.125	4.0	2.0	4.0	0.0	2.0	0.0	4.0	0.0	1.0	2.0	2.0	0.0
13	.500	8.0	9.0	7.0	4.0	9.0	6.0	9.0	8.0	7.0	6.0	4.0	1.0
	.125	4.0	6.0	4.0	2.0	4.0	0.0	7.0	2.0	2.0	3.0	-	0.0
14	.500	9.0	9.0	9.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	3.0
	.125	7.0	9.0	5.0	8.0	7.0	8.0	9.0	9.0	8.0	9.0	9.0	2.0
15	.500	2.0	9.0	9.0	8.0	9.0	7.0	9.0	9.0	9.0	9.0	8.0	0.0
	.125	2.0	9.0	2.0	8.0	6.0	0.0	2.0	8.0	8.0	8.0	6.0	0.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barney ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
16	4.000	6.0	-	7.0	8.0	8.0	0.0	9.0	1.0	-	-	-	-
17	.500	9.0	9.0	9.0	7.0	8.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	5.0	8.0	5.0	6.0	9.0	9.0	9.0	9.0	8.0	9.0	8.0	7.0
18	4.000	9.0	-	9.0	8.0	9.0	9.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	9.0	9.0	7.0	9.0	9.0	9.0	9.0	9.0	8.0	7.0
	.125	7.0	8.0	7.0	7.0	5.0	6.0	9.0	9.0	9.0	9.0	7.0	5.0
19	4.000	9.0	-	4.0	0.0	2.0	6.0	8.0	8.0	-	-	-	-
	.500	4.8	9.0	8.0	6.0	7.0	5.0	8.0	8.0	8.0	7.0	7.0	3.0
	.125	2.2	0.0	5.0	4.0	2.0	0.7	8.0	7.0	2.0	2.0	7.0	1.0
20	4.000	6.0	-	6.0	0.0	4.0	4.0	8.0	5.0	-	-	-	-
	.500	9.0	9.0	8.0	7.0	7.5	6.5	9.0	9.0	8.0	9.0	7.0	7.0
	.125	9.0	9.0	7.0	4.0	5.0	8.0	9.0	9.0	7.0	8.0	7.0	4.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
21	4.000	9.0	-	9.0	2.0	7.0	6.0	9.0	8.0	-	-	-	-
	.500	9.0	9.0	9.0	2.0	9.0	8.5	9.0	9.0	9.0	9.0	9.0	7.0
	.125	8.3	9.0	9.0	2.0	7.0	7.3	9.0	8.0	6.5	4.5	8.5	3.5
22	4.000	9.0	-	9.0	2.0	7.0	6.0	8.0	9.0	-	-	-	-
	.125	4.0	2.0	6.0	0.0	6.0	8.0	7.0	0.0	2.0	2.0	8.0	8.0
23	4.000	9.0	-	9.0	2.0	7.0	6.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	8.5	6.5	9.0	8.7	9.0	9.0	8.5	8.5	8.0	6.5
	.125	7.5	8.0	8.0	3.0	5.0	6.0	7.5	7.5	8.0	3.0	7.5	1.5
24	.500	8.0	8.0	2.0	0.0	2.0	6.0	6.0	7.0	2.0	2.0	6.0	0.0
	.125	2.0	6.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
25	.500	9.0	9.0	7.0	8.0	7.0	7.0	9.0	8.0	9.0	8.0	7.0	0.0
	.125	8.0	9.0	2.0	4.0	4.0	4.0	9.0	7.0	3.0	3.0	3.0	0.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardqr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
26	4.000	9.0	-	4.0	6.0	9.0	1.0	9.0	4.0	-	-	-	-
	.500	6.0	8.0	4.0	2.0	8.0	3.0	9.0	4.0	0.0	0.0	4.0	0.0
	.125	0.0	0.0	0.0	0.0	0.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0
27	.500	9.0	8.0	6.0	6.0	7.0	7.0	8.0	8.0	6.0	6.0	7.0	0.0
	.125	8.0	7.0	2.0	4.0	4.0	0.0	6.0	6.0	1.0	1.0	2.0	0.0
28	4.000	7.0	-	5.0	4.0	4.0	4.0	9.0	5.0	-	-	-	-
	.500	8.0	7.0	2.0	1.0	7.0	2.0	7.0	4.0	3.0	2.0	6.0	0.0
	.125	2.0	5.0	0.0	0.0	2.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0
29	.500	9.0	9.0	8.0	8.0	9.0	4.0	9.0	7.0	8.0	9.0	7.0	1.0
	.125	7.0	7.0	4.0	0.0	2.0	0.0	4.0	2.0	2.0	3.0	4.0	0.0
30	.500	2.0	4.0	0.0	0.0	4.0	0.0	8.0	0.0	0.0	0.0	0.0	0.0

TABLE II (Continued)

<u>Compound No.</u>	<u>Rate (kg/ha)</u>	<u>Barny ardgr</u>	<u>Foxta il Sp</u>	<u>Wild oats</u>	<u>Quack grass</u>	<u>Fld B indwd</u>	<u>Mrngl ry Sp</u>	<u>Wild mustd</u>	<u>Velve tleaf</u>	<u>Wht F enman</u>	<u>Corn field</u>	<u>Cot ton</u>	<u>Soy bean</u>
31	.500	2.0	4.0	0.0	2.0	4.0	0.0	8.0	7.0	1.0	2.0	3.0	0.0
	.125	0.0	2.0	0.0	0.0	0.0	0.0	6.0	2.0	0.0	0.0	0.0	0.0
32	4.000	8.0	-	9.0	9.0	9.0	2.0	9.0	3.0	-	-	-	-
33	4.000	9.0	-	8.0	0.0	4.0	6.0	8.0	5.0	-	-	-	-
	.500	2.3	7.0	4.0	2.0	0.0	0.0	8.0	2.0	2.0	7.0	7.0	0.0
	.125	0.0	0.0	2.0	0.0	0.0	0.0	8.0	2.0	0.0	6.0	6.0	0.0
34	4.000	9.0	-	9.0	9.0	9.0	7.0	9.0	9.0	-	-	-	-
	.500	9.0	9.0	9.0	8.0	9.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0
	.125	4.0	8.0	8.0	8.0	8.0	9.0	9.0	9.0	-	-	8.0	6.0
35	.500	9.0	9.0	8.0	7.0	9.0	9.0	9.0	9.0	9.0	9.0	8.0	8.0
	.125	7.0	8.0	7.0	2.0	7.0	2.0	9.0	9.0	6.0	8.0	6.0	4.0


TABLE II (Continued)

<u>Compound</u> <u>No.</u>	<u>Rate</u> <u>(kg/ha)</u>	<u>Barny</u> <u>ardgr</u>	<u>Foxta</u> <u>il Sp</u>	<u>Wild</u> <u>oats</u>	<u>Quack</u> <u>grass</u>	<u>Fld B</u> <u>indwd</u>	<u>Mrngl</u> <u>ry Sp</u>	<u>Wild</u> <u>mustd</u>	<u>Velve</u> <u>tleaf</u>	<u>Wht F</u> <u>enman</u>	<u>Corn</u> <u>field</u>	<u>Cot</u> <u>ton</u>	<u>Soy</u> <u>bean</u>
36	.500	7.0	9.0	4.0	8.0	4.0	6.0	9.0	8.0	9.0	9.0	7.0	3.0
	.125	4.0	8.0	2.0	7.0	4.0	2.0	7.0	7.0	7.0	8.0	7.0	1.0

WE CLAIM:



1. A compound having the structure




wherein

R is C₁-C₄ alkyl;

R_1 is C_1-C_4 alkyl or C_3-C_6 cycloalkyl; and when R and R_1 are taken together with the carbon to which they are attached they may represent C_3-C_6 cycloalkyl optionally substituted with methyl;

Q is




R_2 , R_3 and R_4 are each independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy or phenyl optionally substituted with one or two halogen atoms, C_1 - C_4 alkyl or haloalkyl groups or C_1 - C_4 alkoxy or haloalkoxy groups;

m is an integer of 1 or 2;

A is O, S, or NR₅; provided that when m is 2, A is O;




- R_5 is C_1-C_4 alkyl;
 B is hydrogen, COR_6 or SO_2R_7 ;
 R_6 is C_1-C_{11} alkyl, chloromethyl or phenyl optionally substituted with one chloro, one nitro or one methoxy group;
 R_7 is C_1-C_4 alkyl or phenyl optionally substituted with one methyl group;
 W is O or S;
 X is hydrogen, halogen or methyl;
 Y and Z are each independently hydrogen, halogen, C_1-C_6 alkyl, C_1-C_4 alkoxymethyl, C_1-C_6 alkoxy, C_1-C_4 alkylthio, phenoxy, C_1-C_4 haloalkyl, nitro, cyano, C_1-C_4 alkylsulfonyl or phenyl optionally substituted with one C_1-C_4 alkyl, C_1-C_4 alkoxy or halogen; and, when taken together, Y and Z may form a ring in which YZ are represented by the structure: $-(CH_2)_n-$, where n is an integer selected from 3 and 4; or



where R_8 , R_9 , R_{10} and R_{11} are each independently hydrogen, halogen, C_1-C_4 alkyl, C_1-C_4 alkoxy, C_1-C_4 alkylthio, C_1-C_4 alkylsulfonyl, C_1-C_4 haloalkyl, nitro, cyano, phenyl, phenoxy, amino, chlorophenyl, methylphenyl, or phenoxy substituted with one chloro, trifluoromethyl, nitro or methyl group, with the proviso that only one of R_8 , R_9 , R_{10} or R_{11} , may represent a substituent other than hydrogen, halogen, C_1-C_4 alkyl or C_1-C_4 alkoxy; and when R and R_1 are not the same, the optical isomers thereof.



2. The compound according to claim 1 having the structure




wherein

A is O or S; and

Y and Z are each independently hydrogen, halogen, C_1-C_6 alkyl, C_1-C_4 alkoxyethyl or C_1-C_6 alkoxy, and when taken together, Y and Z may form a ring in which YZ are represented by the structure:



3. The compound according to claim 2 having the structure






wherein R_2 and R_4 are independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, or C_1 - C_4 haloalkoxy.

4. The compound according to claim 2
2-[3-(2-furoyl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one; 2-[3-(2-furoyl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one; 2-[3-(α -hydroxyfurfuryl)-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one; 2-[3-(α -hydroxyfurfuryl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one; 2-[3-(α -hydroxy-5-methylfurfuryl)-5-methyl-2-pyridyl]-4-isopropyl-4-methyl-2-imidazolin-5-one; or 2-{3-[(di-2-furyl)-hydroxymethyl]-2-pyridyl}-4-isopropyl-4-methyl-2-imidazolin-5-one.

5. A method for controlling undesirable plant species which comprises applying to the foliage of said plants or to the soil or water containing seeds or other propagating organs thereof, a herbicidally effective amount of a compound as described in claim 1.

6. The method according to claim 5 wherein the compound has the structure



wherein



A is O or S; and


Y and Z are each independently hydrogen, halogen, C_1-C_6 alkyl, C_1-C_4 alkoxyethyl or C_1-C_6 alkoxy, and when taken together, Y and Z may form a ring in which YZ are represented by the structure:



7. The method according to claim 5 which comprises applying said compound at a rate of about 0.016 kg/ha to 4.0 kg/ha.

8. A herbicidal composition which comprises an inert solid or liquid diluent and a herbicidally effective amount of a compound as described in claim 1.

9. A method for the preparation of a 3-(substituted ketone)-2-(2-imidazolin-2-yl)pyridine or quinoline compound having the structure




wherein

R is C_1-C_4 alkyl;

R_1 is C_1-C_4 alkyl or C_3-C_6 cycloalkyl; and when R and R_1 are taken together with the carbon to which they are attached they may represent C_3-C_6 cycloalkyl optionally substituted with methyl;



Q is




R_2 , R_3 and R_4 are each independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy or phenyl optionally substituted with one or two halogen atoms, C_1 - C_4 alkyl or haloalkyl groups or C_1 - C_4 alkoxy or haloalkoxy groups.

A is O, S or NR_5 ;

R_5 is C_1 - C_4 alkyl;

X is hydrogen, halogen or methyl; and


Y and Z are each independently hydrogen, halogen, C_1 - C_6 alkyl, C_1 - C_4 alkoxymethyl, C_1 - C_6 alkoxy, C_1 - C_4 alkylthio, phenoxy, C_1 - C_4 haloalkyl, nitro, cyano, C_1 - C_4 alkylsulfonyl or phenyl optionally substituted with one C_1 - C_4 alkyl, C_1 - C_4 alkoxy or halogen; and, when taken together, Y and Z may form a ring in which YZ are represented by the structure: $-(CH_2)_n-$, where n is an integer selected from 3 and 4; or




where R_8 , R_9 , R_{10} and R_{11} are each independently hydrogen, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, nitro, cyano, phenyl, phenoxy, amino, chlorophenyl, methylphenyl, or phenoxy substituted




with one chloro, trifluoromethyl, nitro or methyl group, with the proviso that only one of R_8 , R_9 , R_{10} or R_{11} , may represent a substituent other than hydrogen, halogen, C_1-C_4 alkyl or C_1-C_4 alkoxy; and when R and R_1 are not the same, the optical isomers thereof; which comprises reacting an imidazolinone compound having the structure



wherein X, Y, Z, R and R_1 are as described above with at least about 2.0 molar equivalents of a C_1-C_4 alkyl lithium in the presence of an inert solvent to form a salt and reacting said salt with at least 1.0 molar equivalent of an aldehyde compound having the structure



wherein Q is as described above in the presence of an inert solvent to form a 3-(substituted alcohol)-2-(2-imidazolin-2-yl)pyridine or quinoline compound having the structure





wherein R, R₁, Q, X, Y and Z are as described above and reacting said 3-(substituted alcohol)-2-(2-imidazolin-2-yl)pyridine or quinoline compound with an oxidizing agent in the presence of an inert organic solvent to form said 3-(substituted ketone)-2-(2-imidazolin-2-yl)pyridine or quinoline compound.

10. The method according to claim 9 wherein the oxidizing agent is manganese(IV) oxide, the C₁-C₄ alkyl lithium is selected from the group consisting of n-butyl lithium and methyl lithium, and Q is

