

CLAIMS

What is claimed is:

1. A humanized anti-CD19 antibody or antigen binding fragment thereof comprising a light chain variable (VL) region and a heavy chain variable (VH) region, wherein the VL region comprises a VL complementarity determining region (CDR) 1 (VL CDR1), a VL CDR2, and a VL CDR3 and the VH region comprises a VH CDR1, a VL CDR2, and a VL CDR3,

wherein the VL region has an amino acid sequence at least 95% identical to SEQ ID NO: 14, SEQ ID NO: 20, SEQ ID NO: 11, or SEQ ID NO: 17; and/or

wherein the VH region has an amino acid sequence at least 95% identical to SEQ ID NO: 15, SEQ ID NO: 21, SEQ ID NO: 12, or SEQ ID NO: 18;

wherein the VL CDR1 is SEQ ID NO: 27, the VL CDR2 is SEQ ID NO: 28, and the VL CDR3 is SEQ ID NO: 29; and

wherein the VH CDR1 is SEQ ID NO: 30 or 33, the VH CDR2 is SEQ ID NO: 31 or 34, and the VH CDR3 is SEQ ID NO: 32.

2. The humanized anti-CD19 antibody or antigen binding fragment thereof of claim 1, wherein the antibody or the antigen binding fragment thereof is selected from the group consisting of an IgG, an Fab, an Fab', an F(ab')₂, an Fv, and an scFv.

3. A polypeptide comprising the humanized anti-CD19 antibody or antigen binding fragment thereof of claim 1.

4. A chimeric antigen receptor (CAR) or a T cell receptor (TCR), comprising: (i) an antigen binding domain, (ii) a costimulatory domain, and (iii) an activating domain,

wherein the costimulatory domain comprises an extracellular domain, a transmembrane domain, and an intracellular domain, and

wherein the antigen binding domain comprises at least the polypeptide of claim 3.

5. The CAR or TCR of claim 4, wherein the costimulatory domain is from or is derived from CD2, CD3 delta, CD3 epsilon, CD3 gamma, CD4, CD7, CD8 α , CD8 β , CD11a (ITGAL), CD11b (ITGAM), CD11c (ITGAX), CD11d (ITGAD), CD18 (ITGB2), CD19 (B4), CD27 (TNFRSF7), CD28, CD29 (ITGB1), CD30 (TNFRSF8), CD40 (TNFRSF5),

CD48 (SLAMF2), CD49a (ITGA1), CD49d (ITGA4), CD49f (ITGA6), CD66a (CEACAM1), CD66b (CEACAM8), CD66c (CEACAM6), CD66d (CEACAM3), CD66e (CEACAM5), CD69 (CLEC2), CD79A (B-cell antigen receptor complex-associated alpha chain), CD79B (B-cell antigen receptor complex-associated beta chain), CD84 (SLAMF5), CD96 (Tactile), CD100 (SEMA4D), CD103 (ITGAE), CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD158A (KIR2DL1), CD158B1 (KIR2DL2), CD158B2 (KIR2DL3), CD158C (KIR3DP1), CD158D (KIRDL4), CD158F1 (KIR2DL5A), CD158F2 (KIR2DL5B), CD158K (KIR3DL2), CD160 (BY55), CD162 (SELPLG), CD226 (DNAM1), CD229 (SLAMF3), CD244 (SLAMF4), CD247 (CD3-zeta), CD258 (LIGHT), CD268 (BAFFR), CD270 (TNFSF14), CD272 (BTLA), CD276 (B7-H3), CD279 (PD-1), CD314 (NKG2D), CD319 (SLAMF7), CD335 (NK-p46), CD336 (NK-p44), CD337 (NK-p30), CD352 (SLAMF6), CD353 (SLAMF8), CD355 (CRTAM), CD357 (TNFRSF18), inducible T cell co-stimulator (ICOS), LFA-1 (CD11a/CD18), NKG2C, DAP-10, ICAM-1, NKp80 (KLRP1), IL-2R beta, IL-2R gamma, IL-7R alpha, LFA-1, SLAMF9, LAT, GADS (GrpL), SLP-76 (LCP2), PAG1/CBP, a CD83 ligand, Fc gamma receptor, MHC class 1 molecule, MHC class 2 molecule, a TNF receptor protein, an immunoglobulin protein, a cytokine receptor, an integrin, activating NK cell receptors, a Toll ligand receptor, and fragments or combinations thereof.

6. The CAR or TCR of claim 4, wherein the transmembrane domain is from or is derived from 4-1BB/CD137, an alpha chain of a T cell receptor, a beta chain of a T cell receptor, CD2, CD3 delta, CD3 epsilon, CD3 gamma, CD4, CD7, CD8 α , CD8 β , CD11a (ITGAL), CD11b (ITGAM), CD11c (ITGAX), CD11d (ITGAD), CD18 (ITGB2), CD19 (B4), CD27 (TNFRSF7), CD28, CD29 (ITGB1), CD30 (TNFRSF8), CD40 (TNFRSF5), CD48 (SLAMF2), CD49a (ITGA1), CD49d (ITGA4), CD49f (ITGA6), CD66a (CEACAM1), CD66b (CEACAM8), CD66c (CEACAM6), CD66d (CEACAM3), CD66e (CEACAM5), CD69 (CLEC2), CD79A (B-cell antigen receptor complex-associated alpha chain), CD79B (B-cell antigen receptor complex-associated beta chain), CD84 (SLAMF5), CD96 (Tactile), CD100 (SEMA4D), CD103 (ITGAE), CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD158A (KIR2DL1), CD158B1 (KIR2DL2), CD158B2 (KIR2DL3), CD158C (KIR3DP1), CD158D (KIRDL4), CD158F1 (KIR2DL5A), CD158F2 (KIR2DL5B), CD158K (KIR3DL2), CD160 (BY55), CD162 (SELPLG), CD226 (DNAM1), CD229 (SLAMF3), CD244 (SLAMF4), CD247 (CD3-zeta), CD258 (LIGHT), CD268 (BAFFR), CD270 (TNFSF14), CD272 (BTLA), CD276 (B7-H3), CD279 (PD-1), CD314 (NKG2D),

CD319 (SLAMF7), CD335 (NK-p46), CD336 (NK-p44), CD337 (NK-p30), CD352 (SLAMF6), CD353 (SLAMF8), CD355 (CRTAM), CD357 (TNFRSF18), inducible T cell co-stimulator (ICOS), LFA-1 (CD11a/CD18), NKG2C, DAP-10, ICAM-1, NKp80 (KLRF1), IL-2R beta, IL-2R gamma, IL-7R alpha, LFA-1, SLAMF9, LAT, GADS (GrpL), SLP-76 (LCP2), PAG1/CBP, a CD83 ligand, Fc gamma receptor, MHC class 1 molecule, MHC class 2 molecule, a TNF receptor protein, an immunoglobulin protein, a cytokine receptor, an integrin, activating NK cell receptors, a Toll ligand receptor, and combinations thereof.

7. The CAR or TCR of claim 4, wherein the intracellular domain is from or is derived from 4-1BB/CD137, activating NK cell receptors, B7-H3, BAFFR, BLAME (SLAMF8), BTLA, CD100 (SEMA4D), CD103, CD160 (BY55), CD18, CD19, CD19a, CD2, CD247, CD27, CD276 (B7-H3), CD28, CD29, CD3 delta, CD3 epsilon, CD3 gamma, CD30, CD4, CD40, CD49a, CD49D, CD49f, CD69, CD7, CD84, CD8alpha, CD8beta, CD96 (Tactile), CD11a, CD11b, CD11c, CD11d, CDS, CEACAM1, CRT AM, cytokine receptors, DAP-10, DNAM1 (CD226), Fc gamma receptor, GADS, GITR, HVEM (LIGHTR), IA4, ICAM-1, ICAM-1, Ig alpha (CD79a), IL2R beta, IL2R gamma, IL7R alpha, Immunoglobulin-like proteins, inducible T cell costimulator (ICOS), integrins, ITGA4, ITGA4, ITGA6, ITGAD, ITGAE, ITGAL, ITGAM, ITGAX, ITGB2, ITGB7, ITGB1, KIRDS2, LAT, LFA-1, LFA-1, a ligand that specifically binds with CD83, LIGHT, LIGHT (tumor necrosis factor superfamily member 14; TNFSF14), LTBR, Ly9 (CD229), lymphocyte function-associated antigen-1 (LFA-1 (CD11a/CD18), MHC class I molecule, NKG2C, NKG2D, NKp30, NKp44, NKp46, NKp80 (KLRF1), OX-40, PAG/Cbp, programmed death-1 (PD-1), PSGL1, SELPLG (CD162), signaling lymphocytic activation molecules (SLAM proteins), SLAM (SLAMF1; CD150; IPO-3), SLAMF4 (CD244; 2B4), SLAMF6 (NTB-A; Lyl08), SLAMF7, SLP-76, TNF receptor proteins, TNFR2, a Toll ligand receptor, TRANCE/RANKL, VLA1, or VLA-6, or a combination thereof.

8. The CAR or TCR of claim 4, wherein the extracellular domain is from or is derived from CD2, CD3 delta, CD3 epsilon, CD3 gamma, CD4, CD7, CD8 α , CD8 β , CD11a (ITGAL), CD11b (ITGAM), CD11c (ITGAX), CD11d (ITGAD), CD18 (ITGB2), CD19 (B4), CD27 (TNFRSF7), CD28, CD29 (ITGB1), CD30 (TNFRSF8), CD40 (TNFRSF5), CD48 (SLAMF2), CD49a (ITGA1), CD49d (ITGA4), CD49f (ITGA6), CD66a (CEACAM1), CD66b (CEACAM8), CD66c (CEACAM6), CD66d (CEACAM3), CD66e (CEACAM5), CD69 (CLEC2), CD79A (B-cell antigen receptor complex-associated alpha

chain), CD79B (B-cell antigen receptor complex-associated beta chain), CD84 (SLAMF5), CD96 (Tactile), CD100 (SEMA4D), CD103 (ITGAE), CD134 (OX40), CD137 (4-1BB), CD150 (SLAMF1), CD158A (KIR2DL1), CD158B1 (KIR2DL2), CD158B2 (KIR2DL3), CD158C (KIR3DP1), CD158D (KIRDL4), CD158F1 (KIR2DL5A), CD158F2 (KIR2DL5B), CD158K (KIR3DL2), CD160 (BY55), CD162 (SELPLG), CD226 (DNAM1), CD229 (SLAMF3), CD244 (SLAMF4), CD247 (CD3-zeta), CD258 (LIGHT), CD268 (BAFFR), CD270 (TNFSF14), CD272 (BTLA), CD276 (B7-H3), CD279 (PD-1), CD314 (NKG2D), CD319 (SLAMF7), CD335 (NK-p46), CD336 (NK-p44), CD337 (NK-p30), CD352 (SLAMF6), CD353 (SLAMF8), CD355 (CRTAM), CD357 (TNFRSF18), inducible T cell co-stimulator (ICOS), LFA-1 (CD11a/CD18), NKG2C, DAP-10, ICAM-1, NKp80 (KLRP1), IL-2R beta, IL-2R gamma, IL-7R alpha, LFA-1, SLAMF9, LAT, GADS (GrpL), SLP-76 (LCP2), PAG1/CBP, a CD83 ligand, Fc gamma receptor, MHC class 1 molecule, MHC class 2 molecule, a TNF receptor protein, an immunoglobulin protein, a cytokine receptor, an integrin, activating NK cell receptors, a Toll ligand receptor, and fragments or combinations thereof.

9. The CAR or TCR of claim 4, wherein the activating domain is from or is derived from CD3-zeta or CD3-epsilon.
10. A vector comprising the humanized anti-CD19 antibody or antigen binding fragment thereof of claim 4.
11. A cell comprising a chimeric antigen receptor (CAR) or T cell receptor (TCR) of claim 4.
12. A composition comprising a plurality of cells of claim 11.
13. A method for manufacturing a cell expressing a chimeric antigen receptor (CAR) or a T cell receptor (TCR), comprising a step of transducing a cell with a vector of claim 10.