


[0011] A further example provides a compound of Formula (III):




wherein

R₁₁ is H or methyl;


R₁₀ is H or, when both R₁₄ and R₉ are H, is methyl-1,4'-bipiperidinyl;

R₉ is -H or is -CH₂- substituted by 1,4'-bipiperidinyl, oxo-, hydroxyl, methylpyridinyl, or piperidinyl optionally substituted with hydroxyl, -N(CH₃)₂, or piperidinyl.

[0011a] In one aspect, the present disclosure refers to a method of preparing a compound of formula




which method comprises the following steps:-



wherein the oxidation conditions comprise reacting the starting compound with 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO) and iodobenzene diacetate (PhI(OAc)₂), and wherein R is C₁-C₆ alkyl that is optionally substituted with: -OH, methoxy, ethoxy, -OCH(CH₃)₂, -O(CH₂)₂CH₃, phenyl, furanyl, -O(CH₂)₂OH, phenoxy, methylthio, -F, -N(CH₃)₂, cyano, pyridinyloxy, fluorophenoxy, isochromanyl, phenol, benzylamino, -NHCH₃, oxo-, amino, carboxyl, 7-member spiroaminyl, a three to six member cycloalkyl, saturated or unsaturated and optionally including one or more heteroatoms selected from O and N, and optionally substituted at one or more C or N atoms by methyl, cyano, fluoro, methylamino, or trifluoromethyl; or C₃-C₇ cycloalkane, saturated or unsaturated, optionally bridged, optionally including one or more heteroatoms selected from

O, S, and N, and optionally substituted at one or more C or N atoms by methyl, ethyl, pyridinyl, azetidiny, acetamidyl, carboxamidyl, cyano, fluoro, methylamino, or trifluoromethyl.

[0011b] In another aspect, the present disclosure refers to a compound of formula




[0012] In a further example the compound is selected from *rel*-(2*R*,6*R*)-4-(8-cyanoquinolin-5-yl)-N-((3*R*,4*S*)-4-fluoropyrrolidin-3-yl)-6-methylmorpholine-2-carboxamide hydrochloride, (2*R*,6*R*)-4-(8-cyanoquinolin-5-yl)-6-methyl-N-(1-methylpiperidin-4-yl)morpholine-2-carboxamide, 5-((2*S*,6*R*)-2-([1,4'-bipiperidin]-1'-ylmethyl)-6-methylmorpholino)quinoline-8-carbonitrile, and 5-((2*R*,7*R*)-2-(hydroxymethyl)-7-methyl-1,4-oxazepan-4-yl)quinoline-8-carbonitrile.


[0013] In a further example the compound or pharmaceutically effective salt thereof of the preceding paragraph has an IC₅₀ less than or equal to 20 nM against human TLR7 receptors expressed in a HEK-293 cell line. In a further example the compound or pharmaceutically effective salt thereof of the preceding paragraph of this disclosure has an IC₅₀ less than or equal to 100 nM against human TLR7 receptors expressed in a HEK-293 cell line. In a further example the IC₅₀ against human TLR7 receptors expressed in a HEK-293 cell line is measured by (1) plating cells of the HEK-293 cell line stably expressing TLR7 in Dulbecco's modified Eagle's medium containing 10 % fetal bovine serum at a density of 2.22X10⁵ cells/ml into a 384-well plate and incubating for 2 days at 37 °C, 5 % CO₂; (2) adding the compound or pharmaceutically acceptable salt thereof and incubating the cells for 30 minutes; (3) adding CL097 (InvivoGen) at 3ug/ml and incubating the cells for approximately 20 hours; and (4) quantifying NF-kappaB dependent reporter activation by measuring luminescence.

We claim:

1. A method of preparing a compound of formula



which method comprises the following steps:-



wherein the oxidation conditions comprise reacting the starting compound with 2,2,6,6-tetramethylpiperidine 1-oxyl (TEMPO) and iodobenzene diacetate (PhI(OAc)₂), and wherein R is

C₁-C₆ alkyl that is optionally substituted with:

-OH, methoxy, ethoxy, -OCH(CH₃)₂, -O(CH₂)₂CH₃, phenyl, furanyl, -O(CH₂)₂OH, phenoxy, methylthio, -F, -N(CH₃)₂, cyano, pyridinyloxy, fluorophenoxy, isochromanyl, phenol, benzylamino, -NHCH₃, oxo-, amino, carboxyl, 7-member spiroaminyl, a three to six member cycloalkyl, saturated or unsaturated and optionally including one or more heteroatoms selected from O and N, and optionally substituted at one or more C or N atoms by methyl, cyano, fluoro, methylamino, or trifluoromethyl; or

C₃-C₇ cycloalkane, saturated or unsaturated, optionally bridged, optionally including one or more heteroatoms selected from O, S, and N, and optionally substituted at one or more C or N atoms by methyl, ethyl, pyridinyl, azetidyl, acetamidyl, carboxamidyl, cyano, fluoro, methylamino, or trifluoromethyl.


2. A method according to claim 1, which further comprises one or two additional steps to modify the R group to a different group R', wherein R' is

C₁-C₆ alkyl that is optionally substituted with:

-OH, methoxy, ethoxy, -OCH(CH₃)₂, -O(CH₂)₂CH₃, phenyl, furanyl, -O(CH₂)₂OH, phenoxy, methylthio, -F, -N(CH₃)₂, cyano, pyridinyloxy, fluorophenoxy, isochromanyl, phenol, benzylamino, -NHCH₃, oxo-, amino, carboxyl, 7-member spiroaminyl, a three to six member cycloalkyl, saturated or unsaturated and optionally including one or more heteroatoms selected from O and N, and optionally substituted at one or more C or N atoms by methyl, cyano, fluoro, methylamino, or trifluoromethyl; or

C₃-C₇ cycloalkane, saturated or unsaturated, optionally bridged, optionally including one or more heteroatoms selected from O, S, and N, and optionally substituted at one or more C or N atoms by methyl, ethyl, pyridinyl, azetidiny, acetamidyl, carboxamidyl, cyano, fluoro, methylamino, or trifluoromethyl.

3. A compound of formula



wherein R is

C₁-C₆ alkyl that is optionally substituted with:

-OH, methoxy, ethoxy, -OCH(CH₃)₂, -O(CH₂)₂CH₃, phenyl, furanyl, -O(CH₂)₂OH, phenoxy, methylthio, -F, -N(CH₃)₂, cyano, pyridinyloxy, fluorophenoxy, isochromanyl, phenol, benzylamino, -NHCH₃, oxo-, amino, carboxyl, 7-member spiroaminyl, a three to six member cycloalkyl, saturated or unsaturated and optionally including one or

more heteroatoms selected from O and N, and optionally substituted at one or more C or N atoms by methyl, cyano, fluoro, methylamino, or trifluoromethyl; or

C₃-C₇ cycloalkane, saturated or unsaturated, optionally bridged, optionally including one or more heteroatoms selected from O, S, and N, and optionally substituted at one or more C or N atoms by methyl, ethyl, pyridinyl, azetidiny, acetamidyl, carboxamidyl, cyano, fluoro, methylamino, or trifluoromethyl;

or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 3, or a pharmaceutically acceptable salt thereof, wherein R is selected from

ethyl,
isopropyl,
3,4-difluorobenzyl,
cyclopropyl,
2-hydroxyethyl,
2-methoxyethyl,
(*R*)-2-hydroxypropyl,
(*S*)-1-hydroxypropan-2-yl,
(*R*)-1-hydroxypropan-2-yl,
(*S*)-1-hydroxybutan-2-yl,
(*S*)-1-hydroxy-3-methylbutan-2-yl,
(*S*)-2-hydroxy-1-phenylethyl,
(*R*)-2-hydroxy-1-phenylethyl,
2-hydroxybutyl,
2-ethoxyethyl,
(*R*)-1-hydroxybutan-2-yl,
1,3-dihydroxypropan-2-yl,
2,3-dihydroxypropyl,
((*R*)-tetrahydrofuran-2-yl)methyl,

(tetrahydrofuran-2-yl)methyl,
 2-propoxyethyl,
 (*R*)-1-hydroxypentan-2-yl,
 2-isopropoxyethyl,
 1-methoxybutan-2-yl,
 2-(2-fluorophenyl)-2-hydroxyethyl,
 (*R*)-1-hydroxy-3-methylbutan-2-yl,
 2,2-dimethoxyethyl,
 2-(2-hydroxyethoxy)ethyl,
 1-hydroxyhexan-2-yl,
 (*S*)-1-hydroxy-3,3-dimethylbutan-2-yl;
 (*S*)-1-hydroxyhexan-2-yl,
 (2*S*,3*S*)-1-hydroxy-3-methylpentan-2-yl,
 (*S*)-1-hydroxy-4-methylpentan-2-yl,
 (*R*)-1-hydroxy-4-methylpentan-2-yl,
 ((4-methylmorpholin-2-yl)methyl,
 (*S*)-1-hydroxy-4-(methylthio)butan-2-yl,
 2-phenoxyethyl,
 (*S*)-1-hydroxy-3-phenylpropan-2-yl,
 2-phenoxypropyl,
 2-hydroxy-3-phenylpropyl,
 (2-(pyridin-3-yloxy)propyl,
 2-(4-fluorophenoxy)ethyl,
 2-(3-fluorophenyl)-2-hydroxyethyl,
 (*S*)-1-cyclohexyl-3-hydroxypropan-2-yl,
 isochroman-1-ylmethyl,
 2-hydroxy-3-phenoxypropyl,
 (1*S*)-1,3-dihydroxy-1-phenylpropan-2-yl,
 2-hydroxy-3-(piperazin-1-yl)propyl,
 azetidin-3-yl,
 (*S*)-pyrrolidin-3-yl,
 (*R*)-pyrrolidin-3-yl,

(S)-piperidin-3-yl,
 (R)-piperidin-3-yl,
 (S)-pyrrolidin-2-ylmethyl,
 2-(benzylamino)ethyl,
 pyridin-2-yl,
 6-methylpyridin-2-yl,
 5-methylisoxazol-3-yl,
 2,2,2-trifluoroethyl,
 2,2-difluoroethyl,
 3,3,3-trifluoropropyl,
 (2-(dimethylamino)-2-methylpropyl),
 (S)-morpholin-2-ylmethyl,
 ((S)-4-methylmorpholin-2-yl)methyl,
 2-fluoroethyl,
 3-fluoropropyl,
 (S)-1,1,1-trifluoropropan-2-yl,
 (R)-1,1,1-trifluoropropan-2-yl,
 1,3-dimethyl-1*H*-pyrazol-5-yl,
 1-methyl-1*H*-pyrazol-5-yl,
 cyanomethyl, 1-cyanocyclopropyl,
 1,2,4-thiadiazol-5-yl,
 3-methyl-1,2,4-thiadiazol-5-yl,
 piperidin-4-yl,
 (4-fluoropiperidin-4-yl)methyl,
 azetidin-3-ylmethyl,
 3-carbamoyl-1-methyl-1*H*-pyrazol-4-yl,
 oxetan-3-ylmethyl,
 tetrahydro-2*H*-pyran-4-yl,
 tetrahydrofuran-3-yl,
 (3-methyloxetan-3-yl)methyl,
 oxetan-3-yl(3-(aminomethyl)oxetan-3-yl)methyl,
 oxetan-2-ylmethyl,

2-(methylamino)ethyl,
 (S)-3,3,3-trifluoro-2-hydroxypropyl,
 (R)-3,3,3-trifluoro-2-hydroxypropyl,
 2-(dimethylamino)ethyl,
 (S)-4,4,4-trifluoro-3-hydroxybutyl,
 (R)-4,4,4-trifluoro-3-hydroxybutyl,
 3-amino-4,4,4-trifluorobutyl,
 acetic acid,
 3-carbamoylpyridin-4-yl,
 (R)-morpholin-2-ylmethyl,
 pyridin-4-yl,
 pyridin-3-yl,
 1-(piperidin-4-yl)-1*H*-pyrazol-4-yl,
 1-(azetidin-3-yl)-1*H*-pyrazol-4-yl,
 (1*H*-pyrazol-5-yl)methyl,
 (1*H*-pyrazol-4-yl)methyl,
 (3-(trifluoromethyl)pyridin-2-yl)methyl,
 1-(pyridin-2-yl)ethyl,
 pyridin-2-ylmethyl,
 (6-methylpyridin-2-yl)methyl,
 (1-methylpiperidin-2-yl)methyl,
 (3-methylpyridin-2-yl)methyl,
 4-cyano-1*H*-pyrazol-3-yl,
 (3*S*,4*R*)-4-fluoro-1-methylpyrrolidin-3-yl,-
 (3*S*,4*R*)-4-fluoropiperidin-3-yl,
 ((2*R*,6*R*)-6-methylmorpholin-2-yl)methyl,
 (R)-pyrrolidin-2-ylmethyl,
 (S)-piperidin-2-ylmethyl,
 (1*R*,5*S*,6*S*)-3-azabicyclo[3.1.0]hexan-6-yl,
 methyl,
 ((2*S*,6*R*)-4,6-dimethylmorpholin-2-yl)methyl
 1-methylpiperidin-4-yl,

(3*S*,4*R*)-4-fluoropiperidin-3-yl,
1-(2-amino-2-oxoethyl)piperidin-4-yl,
2-carbamoylpyridin-4-yl,
(*S*)-piperidin-3-yl,
1-ethylpiperidin-4-yl,
1-ethylpiperidin-3-yl,
(*R*)-1-methylpiperidin-3-yl,
4-methylpiperidin-4-yl,
2-amino-2-methylpropyl,
(3*R*,4*S*)-4-fluoropyrrolidin-3-yl,
(3*S*,4*R*)-4-fluoropyrrolidin-3-yl,
azepan-3-yl;
1-methylazepan-3-yl,
1,4-dimethylpiperidin-4-yl and
4-fluoro-1-methylpiperidin-3-yl.