

MELANOCORTIN SUBTYPE-2 RECEPTOR (MC2R) ANTAGONISTS AND USES THEREOF

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 62/681,011 filed on June 5, 2018, and U.S. Provisional Patent Application No. 62/734,873 filed on September 21, 2018, each of which is herein incorporated by reference in its entirety.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with the support of the United States government under SBIR Grant No. 1R43DK115245-01 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

[0003] Described herein are compounds that modulate the activity of one or more melanocortin receptors, methods of making such compounds, pharmaceutical compositions and medicaments comprising such compounds, and methods of using such compounds in the treatment of conditions, diseases, or disorders that would benefit from modulating melanocortin subtype-2 receptor (MC2R) activity.

BACKGROUND OF THE INVENTION


[0004] The melanocortin receptors form a family of G protein-coupled receptor (GPCRs) (MC1R, MC2R, MC3R, MC4R, and MC5R) that are selectively activated by different melanocortin peptides adrenocorticotrophic hormone (ACTH), and the melanocortin peptides α -, β -, and γ -melanocyte-stimulating hormone (α -MSH, β -MSH, and γ -MSH) that are all derived proteolytically from proopiomelanocortin hormone, or POMC. ACTH is a 39 amino acid peptide that is the primary regulator of adrenal glucocorticoid synthesis and secretion and only has affinity for MC2R. As the central actor in this hypothalamic-pituitary-adrenal (HPA) axis, ACTH is secreted by the pituitary in response to stressful stimuli and acts at the adrenal gland to stimulate the synthesis and secretion of cortisol. Modulation of MC2R is attractive for the treatment of conditions, diseases, or disorders that would benefit from modulating melanocortin receptor activity.

SUMMARY OF THE INVENTION

[0005] Compounds described herein are melanocortin receptor modulator compounds. In some embodiments, compounds described herein modulate one or more of the subtype melanocortin receptor proteins. In some embodiments, compounds described herein modulate two or more of

the subtype melanocortin receptor proteins. In some embodiments, compounds described herein modulate MC2R.

[0006] In one aspect, described herein is a compound of Formula (I), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (I)

wherein:

- R^A is unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, wherein if R^A is substituted then R^A is substituted with R^a , R^b and R^c ;
 R^a , R^b and R^c are independently selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^a , R^b and R^c is substituted with one or more R^9 groups;
 L is absent, $-C(=O)-$, $-C(=O)NR^7-$, or $-SO_2-$;
 R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle, unsubstituted or substituted C_1 - C_7 alkyl, unsubstituted or substituted C_1 - C_7 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f ;
 or R^B and R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 7-membered heterocycle, wherein if the 3- to 7-membered heterocycle is substituted then the 3- to 7-membered heterocycle is substituted with R^d , R^e and R^f ;
 R^d , R^e and R^f are independently selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted

monocyclic heteroaryl, wherein any substituted group of R^d , R^e and R^f is substituted with one or more R^9 groups;

X^1 is CR^6 or N;

X^2 is CR^6 or N;

X^3 is CR^6 or N;

R^1 is an unsubstituted or substituted C_1 - C_6 alkyl, or unsubstituted or substituted C_3 - C_6 cycloalkyl, wherein if R^1 is substituted then R^1 is substituted with hydrogen, $-OR^8$, halogen, $-N(R^7)_2$, or $-CN$;

R^2 and R^3 are independently hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, or unsubstituted or substituted C_3 - C_6 cycloalkyl, wherein any substituted group of R^2 and R^3 is substituted with hydrogen, $-OR^8$, $-N(R^7)_2$, halogen, or $-CN$;

or R^2 and R^3 are taken together with the carbon atom to which they are attached to form $-C(=O)-$;

or R^2 and R^3 are taken together with the carbon atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic carbocycle;

R^4 and R^5 are independently selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -carbocycle, or unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -heterocycle, wherein any substituted group of R^4 and R^5 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$;

or R^4 and R^5 are taken together with the nitrogen atom to which they are attached to form a substituted or unsubstituted 3- to 6-membered heterocycle, wherein if the 3- to 6-membered heterocycle is substituted then the 3- to 6-membered heterocycle is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$;

or R^2 and R^4 are taken together with the intervening atoms to which they are attached to form a substituted or unsubstituted 5- to 6-membered N-containing heterocycle;

each R^6 is independently selected from the group consisting of hydrogen, halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl,

unsubstituted or substituted C₁-C₆fluoroalkyl, -CN, -OR⁸, -SR⁸, -CO₂R⁸, -C(=O)N(R⁷)₂, or -N(R⁷)₂;

each R⁷ is independently selected from the group consisting of hydrogen, substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl;

or two R⁷ are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R⁸ is independently selected from the group consisting of hydrogen, substituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R⁹ is independently hydrogen, halogen, unsubstituted or substituted C₁-C₄alkyl, unsubstituted or substituted C₁-C₄alkoxy, unsubstituted or substituted C₁-C₄fluoroalkyl, unsubstituted or substituted C₁-C₄fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂R⁸, -CH₂CO₂R⁸, -C(=O)N(R⁷)₂, -C(=O)N(R⁷)OR⁸, -CH₂C(=O)N(R⁷)₂, -N(R⁷)₂, -CH₂N(R⁷)₂, -C(R⁸)₂N(R⁷)₂, -NR⁷C(=O)R⁸, -CH₂NR⁷C(=O)R⁸, -NR⁷C(=O)N(R⁷)₂, -NR⁷C(=O)N(R⁷)₂, C(R⁸)=N(R⁷)-OR⁸, -SR⁸, -S(=O)R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂;

each R¹⁰ is independently selected from the group consisting substituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl;


n is 1 or 2.

[0007] Also described herein are pharmaceutical compositions comprising a compound described herein, or a pharmaceutically acceptable salt, or solvate thereof, and at least one pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is formulated for administration to a mammal by intravenous administration, subcutaneous administration, oral administration, inhalation, nasal administration, dermal administration, or ophthalmic administration. In some embodiments, the pharmaceutical composition is formulated for administration to a mammal by oral administration. In some embodiments, the pharmaceutical composition is in the form of a tablet, a pill, a capsule, a liquid, a suspension, a

gel, a dispersion, a solution, an emulsion, an ointment, or a lotion. In some embodiments, the pharmaceutical composition is in the form of a tablet, a pill, or a capsule.

[0008] In any of the aforementioned aspects are further embodiments in which the effective amount of the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is: (a) systemically administered to the mammal; and/or (b) administered orally to the mammal; and/or (c) intravenously administered to the mammal; and/or (d) administered by inhalation; and/or (e) administered by nasal administration; or and/or (f) administered by injection to the mammal; and/or (g) administered topically to the mammal; and/or (h) administered by ophthalmic administration; and/or (i) administered rectally to the mammal; and/or (j) administered non-systemically or locally to the mammal.

[0008a] In one aspect, described herein is a compound of Formula (IX), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (IX)

wherein:

V is CH or N;

R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, wherein any substituted group of R^a is substituted with one or more R⁹ groups;

R^b and R^c are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆

heteroalkyl, wherein any substituted group of R^b and R^c is substituted with one or more R^9 groups;

R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle, unsubstituted or substituted C_1 - C_7 alkyl, unsubstituted or substituted C_1 - C_7 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f ;

R^d is selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_2 - C_7 heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R^9 groups;

R^e and R^f are each independently selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein any substituted group of R^d is substituted with one or more R^9 groups;


X^1 is N or CR^6 ;

R^2 and R^3 are hydrogen;

or R^2 and R^3 are taken together with the carbon atom to which they are attached to form $-C(=O)-$;

R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -carbocycle, or unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -heterocycle, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$;

or R⁴ is H, -CH₃, HOCH₂CH₂-, HOCH(CH₃)CH₂-, -CH₂CH₂NH₂, -CH₂(CH₂)₂NH₂,



R⁶ is H or F;

each R⁷ is independently selected from the group consisting of hydrogen, substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl;

or two R⁷ are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R⁸ is independently selected from the group consisting of hydrogen, substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R⁹ is independently hydrogen, halogen, unsubstituted or substituted C₁-C₄alkyl, unsubstituted or substituted C₁-C₄alkoxy, unsubstituted or substituted C₁-C₄fluoroalkyl, unsubstituted or substituted C₁-C₄fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂R⁸, -CH₂CO₂R⁸, -C(=O)N(R⁷)₂, -C(=O)N(R⁷)OR⁸, -CH₂C(=O)N(R⁷)₂, -N(R⁷)₂, -CH₂N(R⁷)₂, -C(R⁸)₂N(R⁷)₂, -NR⁷C(=O)R⁸, -CH₂NR⁷C(=O)R⁸, -NR⁷C(=O)N(R⁷)₂, -NR⁷C(=O)N(R⁷)₂, C(R⁸)=N(R⁷)-OR⁸, -SR⁸, -S(=O)R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂; and

each R¹⁰ is independently selected from the group consisting of unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl.

[0008b] In one aspect, described herein is a Use of a compound of any one of claims 1-18 in the manufacture of a medicament for treating a disease or condition in a mammal that would benefit from the modulation of melanocortin subtype-2 receptor (MC2R) activity; wherein the disease or condition comprises growth of fat pads in the collarbone, back of neck, face and trunk, excessive sweating, dilation of capillaries, thinning of the skin, muscle weakness, hirsutism, depression/anxiety, hypertension, osteoporosis, insulin resistance, hyperglycemia, and heart disease; or wherein the disease or condition is Cushing's syndrome; or wherein the disease or condition is Cushing's disease; or wherein the disease or condition is ectopic ACTH syndrome; or wherein the disease or condition is congenital adrenal hyperplasia.

[0009] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in which the compound is administered once a day to the mammal or the compound is administered to the mammal multiple times over the span of one day. In some embodiments, the compound is administered on a continuous dosing schedule. In some embodiments, the compound is administered on a continuous daily dosing schedule.

[0010] In any of the embodiments disclosed herein, the mammal is a human.

[0011] In some embodiments, compounds provided herein are orally administered to a human.

[0012] Articles of manufacture, which include packaging material, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, within the packaging material, and a label that indicates that the compound or composition, or pharmaceutically acceptable salt, tautomers, pharmaceutically acceptable N-oxide, pharmaceutically active metabolite, pharmaceutically acceptable prodrug, or pharmaceutically acceptable solvate thereof, is used for modulating one or more subtype melanocortin receptor proteins, or for the treatment, prevention or amelioration of one or more symptoms of a disease or condition that would benefit from modulating one or more subtype melanocortin receptor proteins, are provided.

[0013] Other objects, features and advantages of the compounds, methods and compositions described herein will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments, are given by way of illustration only, since various changes and modifications within the spirit and scope of the instant disclosure will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE INVENTION

[0014] Adrenocorticotrophic hormone (ACTH) is a 39 amino acid peptide synthesized by anterior pituitary corticotrophic cells by proteolytic cleavage of the proopiomelanocortin hormone (POMC). ACTH is the primary regulator of adrenal glucocorticoid (GC; cortisol in humans and most other species; corticosterone in rodents) synthesis and secretion. As the central actor in this hypothalamic-pituitary-adrenal (HPA) axis, ACTH is secreted by the pituitary in response to stressful stimuli and acts at the adrenal gland to stimulate the synthesis and secretion of cortisol. This stimulation is mediated through a highly specific G protein-coupled receptor (GPCR) which is expressed almost uniquely in the adrenal cortex. The receptor is the melanocortin 2 receptor (MC2R), and, along with ACTH, is part of the larger melanocortin system.

[0015] The melanocortin system comprises a family of five GPCRs (MC1R, MC2R, MC3R, MC4R, and MC5R); their natural agonists, the melanocortin peptides α -, β -, and γ -melanocyte-stimulating hormone (α -MSH, β -MSH, and γ -MSH) and ACTH; and endogenous melanocortin antagonists agouti and agouti-related protein (AGRP). The melanocortin receptors (MCRs) have different selectivities for endogenous agonist and antagonist peptides and are expressed in diverse tissues where they serve varied and discreet physiological functions (Gantz, I. and T.M. Fong, *Am. J. Physiol. Endocrinol. Metab.*, 284: E468–E474, 2003).

[0016] It is possible to selectively modulate any one of the MCRs, or combinations thereof. In some embodiments, selectively modulating any one of the MCRs relative to the other MCRs, or combinations thereof, is useful in a variety of clinical applications. In some embodiments, selectively modulating any one of the MCRs relative to the other MCRs, or combinations thereof, reduces unwanted side effects in a variety of clinical applications. In one aspect, compounds described herein are antagonists of MC2R. In some embodiments, compounds described herein are selective antagonists for MC2R relative to other MCRs.

[0017] MC2R is a highly selective receptor for ACTH. Although ACTH can activate all five MCRs, at physiological levels, the sensitivity of the other receptors is not high enough to be activated, and ACTH selectively activates MC2R. Importantly, the other naturally occurring agonists α -MSH, β -MSH, and γ -MSH have no affinity for MC2R (Gantz, I. and T.M. Fong, *Am. J. Physiol. Endocrinol. Metab.*, 284: E468–E474, 2003). The major function of MC2R is to stimulate the fasciculata cells of the adrenal cortex to synthesize and secrete cortisol. MC2R requires the GPCR accessory protein MRAP (melanocortin 2 receptor protein) to be successfully secreted to the cell surface and as well as to function. MRAP is a small protein with a single transmembrane domain that forms an antiparallel homodimer in stable complex with MC2R and

is necessary for both cell surface expression of MC2R and its ability to bind ACTH. MRAP can bind to any of the MCRs and affect their activities, but is only essential for MC2R activity. Binding of ACTH to the MC2R/MRAP complex on adrenal cortical cells activates G_s to elevate intracellular cAMP levels which in turn stimulates cortisol synthesis and secretion by regulating multiple steps in the steroidogenic pathway.

[0018] Cushing's syndrome is a rare disorder characterized by chronic, excess glucocorticoid exposure. Clinical signs of Cushing's syndrome include growth of fat pads (collarbone, back of neck, face and trunk), excessive sweating, dilation of capillaries, thinning of the skin, muscle weakness, hirsutism, depression/anxiety, hypertension, osteoporosis, insulin resistance, hyperglycemia, heart disease, and a range of other metabolic disturbances resulting in high morbidity. If inadequately controlled in its severe forms, Cushing's syndrome is associated with high mortality. Although glucocorticoid excess can sometimes be ACTH independent, for example from excessive autonomous secretion of cortisol from a hyperfunctioning adrenal adenoma, carcinoma, or steroid abuse, about 60-80% of all cases are ACTH dependent Cushing's syndrome, known as Cushing's disease. Cushing's disease is caused by microadenomas of pituitary corticotrophic cells that secrete excess ACTH. Corticotroph adenomas are small, usually slow growing, benign tumors that normally come to clinical attention as a result of the effects of glucocorticoid excess, rather than because of the physical effects of an expanding tumor. First line treatments for Cushing's disease are surgical and involve removal of either the ACTH-secreting tumor in the pituitary or the adrenal glands themselves. As surgery is often unsuccessful, contraindicated, or delayed, medical therapy for these patients becomes necessary. Current treatment options include inhibitors of steroid synthesis enzymes that can prevent the production of cortisol and improve symptoms, but these treatments also induce a host of unwanted side effects due to the accumulation of other steroid products. In one aspect, an MC2R antagonist is used in the treatment of Cushing's syndrome. In some embodiments, an MC2R antagonist is used in the treatment of Cushing's disease. In some embodiments, glucocorticoid excess is ACTH independent. In some embodiments, glucocorticoid excess is ACTH dependent.

[0019] Ectopic ACTH syndrome, or ectopic Cushing's syndrome or disease, is essentially the same as Cushing's disease, except that the underlying tumor expressing ACTH is outside the pituitary gland. In some embodiments, the tumors are small carcinoid tumors that occur anywhere in the lungs or gastrointestinal tract. In some embodiments, an MC2R antagonist is used in the treatment of ectopic ACTH syndrome.

[0020] Congenital adrenal hyperplasia (CAH) is characterized by a reduction or loss of cortisol synthesis and excessive ACTH and corticotropin-releasing hormone. CAH can result from a variety of genetic defects in the adrenal steroidal biosynthesis pathway. In some embodiments, CAH is due to a mutation in 21 β -hydroxylase. The lack of cortisol removes the negative feedback to the pituitary which leads to excessive ACTH secretion. The resulting excessive adrenal stimulation causes overproduction of steroid precursors which also have negative consequences (e.g., hyperandrogenism). Administration of replacement glucocorticoids typically does not adequately suppress ACTH without also causing Cushing's-like symptoms. In some embodiments, an MC2R antagonist is used in the treatment of CAH.


[0021] In addition to Cushing's disease and CAH it has also been hypothesized that there might be a role for an MC2R antagonist in the treatment of depressive illness and septic shock. In some embodiments, an MC2R antagonist is used in the treatment of depressive illness. In some embodiments, an MC2R antagonist is used in the treatment of septic shock.

[0022] In some embodiments, compounds described herein are amenable to administration to a mammal in need of treatment with an MC2R antagonist.

Compounds

[0023] Compounds of Formula (I), including pharmaceutically acceptable salts, prodrugs, active metabolites and pharmaceutically acceptable solvates thereof, are melanocortin receptor modulators. In some embodiments, the compounds of Formula (I), including pharmaceutically acceptable salts, prodrugs, active metabolites and pharmaceutically acceptable solvates thereof, are MC2R modulators. In some embodiments, the MC2R modulators are MC2R antagonists.

[0024] Provided in one aspect is a compound of Formula (I), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (I)

wherein:

R^A is unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic heteroaryl, wherein if R^A is substituted then R^A is substituted with R^a, R^b and R^c;
 R^a, R^b and R^c are independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl,

unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^a, R^b and R^c is substituted with one or more R⁹ groups;

L is absent, -C(=O)-, -C(=O)NR⁷-, or -SO₂-;

R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle, unsubstituted or substituted C₁-C₇ alkyl, unsubstituted or substituted C₁-C₇ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein if R^B is substituted then R^B is substituted with R^d, R^e and R^f;

or R^B and R⁷ are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 7-membered heterocycle, wherein if the 3- to 7-membered heterocycle is substituted then the 3- to 7-membered heterocycle is substituted with R^d, R^e and R^f;

R^d, R^e and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^d, R^e and R^f is substituted with one or more R⁹ groups;

X¹ is CR⁶ or N;

X² is CR⁶ or N;

X³ is CR⁶ or N;

R¹ is an unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₃-C₆ cycloalkyl, wherein if R¹ is substituted then R¹ is substituted with hydrogen, -OR⁸, halogen, -N(R⁷)₂, or -CN;

R² and R³ are independently hydrogen, unsubstituted or substituted C₁-C₆ alkyl, or unsubstituted or substituted C₃-C₆ cycloalkyl, wherein any substituted group of R² and R³ is substituted with hydrogen, -OR⁸, -N(R⁷)₂, halogen, or -CN;

or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-;

or R² and R³ are taken together with the carbon atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic carbocycle;

R^4 and R^5 are independently selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -carbocycle, or unsubstituted or substituted $-(C_1-C_6 \text{ alkyl})$ -heterocycle, wherein any substituted group of R^4 and R^5 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$;

or R^4 and R^5 are taken together with the nitrogen atom to which they are attached to form a substituted or unsubstituted 3- to 6-membered heterocycle, wherein if the 3- to 6-membered heterocycle is substituted then the 3- to 6-membered heterocycle is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$;

or R^2 and R^4 are taken together with the intervening atoms to which they are attached to form a substituted or unsubstituted 5- to 6-membered N-containing heterocycle;

each R^6 is independently selected from the group consisting of hydrogen, halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, $-CN$, $-OR^8$, $-SR^8$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, or $-N(R^7)_2$;

each R^7 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl;

or two R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^8 is independently selected from the group consisting of hydrogen, substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R^9 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkyl, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or

substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂R⁸, -CH₂CO₂R⁸, -C(=O)N(R⁷)₂, -C(=O)N(R⁷)OR⁸, -CH₂C(=O)N(R⁷)₂, -N(R⁷)₂, -CH₂N(R⁷)₂, -C(R⁸)₂N(R⁷)₂, -NR⁷C(=O)R⁸, -CH₂NR⁷C(=O)R⁸, -NR⁷C(=O)N(R⁷)₂, -NR⁷C(=O)N(R⁷)₂, C(R⁸)=N(R⁷)-OR⁸, -SR⁸, -S(=O)R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂;

each R¹⁰ is independently selected from the group consisting substituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₁-C₆fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl;

n is 1 or 2.

[0025] In some embodiments, L is absent, -C(=O)-, -C(=O)NR⁷-, or -SO₂-. In some embodiments, L is absent, -C(=O)-, or -C(=O)NR⁷-. In some embodiments, L is -C(=O)- or -C(=O)NR⁷-. In some embodiments, L is -C(=O)-. In some embodiments, L is -C(=O)NR⁷-. In some embodiments, L is -C(=O)-.

[0026] In some embodiments, L is -C(=O)-; R² and R³ are independently hydrogen, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂OR⁸, or -CH₂NHR⁷; or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-. In some embodiments, R² and R³ are hydrogen; or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-. In some embodiments, R² and R³ are hydrogen. In some embodiments, R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-.

[0027] In some embodiments, L is -C(=O)-; R² and R³ are independently hydrogen or -CH₃; or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-; or R² and R³ are taken together with the carbon atom to which they are attached to form a cyclopropyl.






[0028] In some embodiments, R^A is unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted monocyclic 5-membered heteroaryl, wherein if R^A is substituted then R^A is substituted with R^a, R^b and R^c.



[0029] In some embodiments, R^A is unsubstituted or substituted phenyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, unsubstituted or substituted pyridazinyl, unsubstituted or substituted triazinyl, unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted pyrrolyl, unsubstituted or substituted oxazolyl, unsubstituted or substituted thiazolyl, unsubstituted or substituted imidazolyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted triazolyl, unsubstituted or substituted tetrazolyl, unsubstituted or substituted isoxazolyl, unsubstituted or substituted isothiazolyl, unsubstituted or substituted oxadiazolyl, or



unsubstituted or substituted thiadiazolyl; wherein if R^A is substituted then R^A is substituted with R^a , R^b and R^c .



[0030] In some embodiments, R^A is unsubstituted or substituted phenyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, or unsubstituted or substituted pyridazinyl; wherein if R^A is substituted then R^A is substituted with R^a , R^b and R^c .


[0031] In some embodiments, R^A is unsubstituted or substituted phenyl, or unsubstituted or substituted pyridinyl, wherein if R^A is substituted then R^A is substituted with R^a , R^b and R^c .



[0032] In some embodiments, R^A is , , , , or .

[0033] In some embodiments, R^A is , or . In some embodiments, R^A is

, or . In some embodiments, R^A is


. In some embodiments, R^A is , where V is CH or N.

[0035] In some embodiments, R^A is , where V is CH or N. In some embodiments,

R^A is , or . In some embodiments, R^A is

[0036] In some embodiments, n is 1. In some embodiments, n is 2.

[0037] In some embodiments, the compound has the structure of Formula (II), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (II)

wherein,

V is CH or N.

[0038] In some embodiments, R^B is an unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted bicyclic carbocycle, unsubstituted or substituted polycyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted bicyclic heterocycle, unsubstituted or substituted polycyclic heterocycle, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

[0039] In some embodiments, R^B is an unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted bridged carbocycle, unsubstituted or substituted spiro carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted bridged heterocycle, unsubstituted or substituted spiro heterocycle, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

[0040] In some embodiments, R^B is an unsubstituted or substituted phenyl, unsubstituted or substituted naphthyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted monocyclic 5-membered heteroaryl, unsubstituted or substituted bicyclic heteroaryl, monocyclic C_3 - C_8 cycloalkyl, unsubstituted or substituted bridged C_5 - C_{10} cycloalkyl, unsubstituted or substituted spiro C_5 - C_{10} cycloalkyl, unsubstituted or substituted monocyclic C_2 - C_8 heterocycloalkyl, unsubstituted or substituted bridged C_5 - C_{10} heterocycloalkyl, or unsubstituted or substituted spiro C_5 - C_{10} heterocycloalkyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

[0041] In some embodiments, R^B is an unsubstituted or substituted carbocycle that is an unsubstituted or substituted phenyl, unsubstituted or substituted naphthyl, unsubstituted or substituted indanyl, unsubstituted or substituted indenyl, unsubstituted or substituted tetrahydronaphthyl, unsubstituted or substituted cyclopropyl, unsubstituted or substituted cyclobutyl, unsubstituted or substituted cyclopentyl, unsubstituted or substituted cyclopentenyl, unsubstituted or substituted cyclohexyl, unsubstituted or substituted cyclohexenyl, unsubstituted or substituted cycloheptyl, unsubstituted or substituted cyclooctyl, unsubstituted or substituted spiro[2.2]pentyl, unsubstituted or substituted spiro[3.3]heptyl, unsubstituted or substituted

spiro[3.5]nonyl, unsubstituted or substituted spiro[4.4]nonyl, unsubstituted or substituted spiro[4.5]decyl, unsubstituted or substituted norbornyl, unsubstituted or substituted norbornenyl, unsubstituted or substituted bicyclo[1.1.1]pentyl, unsubstituted or substituted adamantyl, or unsubstituted or substituted decalanyl.

[0042] In some embodiments, R^B is an unsubstituted or substituted furanyl, unsubstituted or substituted thienyl, unsubstituted or substituted pyrrolyl, unsubstituted or substituted oxazolyl, unsubstituted or substituted thiazolyl, unsubstituted or substituted imidazolyl, unsubstituted or substituted pyrazolyl, unsubstituted or substituted triazolyl, unsubstituted or substituted tetrazolyl, unsubstituted or substituted isoxazolyl, unsubstituted or substituted isothiazolyl, unsubstituted or substituted oxadiazolyl, unsubstituted or substituted thiadiazolyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, unsubstituted or substituted pyridazinyl, unsubstituted or substituted triazinyl, unsubstituted or substituted quinolinyl, unsubstituted or substituted isoquinolinyl, unsubstituted or substituted cinnolinyl, unsubstituted or substituted phthalazinyl, unsubstituted or substituted quinazolinyl, unsubstituted or substituted quinoxalinyl, unsubstituted or substituted naphthyridinyl, unsubstituted or substituted pteridinyl, unsubstituted or substituted indolizinyl, unsubstituted or substituted azaindolizinyl, unsubstituted or substituted indolyl, unsubstituted or substituted azaindolyl, unsubstituted or substituted indazolyl, unsubstituted or substituted azaindazolyl, unsubstituted or substituted benzimidazolyl, unsubstituted or substituted azabenzimidazolyl, unsubstituted or substituted benzotriazolyl, unsubstituted or substituted azabenzotriazolyl, unsubstituted or substituted benzoxazolyl, unsubstituted or substituted azabenzoxazolyl, unsubstituted or substituted benzisoxazolyl, unsubstituted or substituted azabenzisoxazolyl, unsubstituted or substituted benzofuranyl, unsubstituted or substituted azabenzofuranyl, unsubstituted or substituted benzothienyl, unsubstituted or substituted azabenzothienyl, unsubstituted or substituted benzothiazolyl, unsubstituted or substituted azabenzothiazolyl, or unsubstituted or substituted purinyl. In some embodiments, R^B is an unsubstituted or substituted indolyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .


[0043] In some embodiments, R^B is an unsubstituted or substituted aziridinyl, unsubstituted or substituted azetidyl, unsubstituted or substituted oxetanyl, unsubstituted or substituted thietanyl, unsubstituted or substituted pyrrolidinyl, unsubstituted or substituted tetrahydrofuranyl, unsubstituted or substituted tetrahydrothienyl, unsubstituted or substituted oxazolidinonyl, unsubstituted or substituted tetrahydropyranyl, unsubstituted or substituted piperidinyl, unsubstituted or substituted morpholinyl, unsubstituted or substituted thiomorpholinyl,


unsubstituted or substituted piperaziny, unsubstituted or substituted homopiperidinyl, unsubstituted or substituted oxepanyl, unsubstituted or substituted thiepanyl, unsubstituted or substituted oxazepiny, unsubstituted or substituted diazepiny, unsubstituted or substituted thiazepiny, unsubstituted or substituted azaspiro[3.3]heptanyl, unsubstituted or substituted azaspiro[3.4]octanyl, unsubstituted or substituted azaspiro[3.4]octanyl, or unsubstituted or substituted azaspiro[4.4]nonyl.

[0044] In some embodiments, R^B is unsubstituted or substituted phenyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted pyrazinyl, or unsubstituted or substituted pyridazinyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .





[0045] In some embodiments, R^B is unsubstituted or substituted cyclopropyl, unsubstituted or substituted cyclobutyl, unsubstituted or substituted cyclopentyl, unsubstituted or substituted cyclopentenyl, or unsubstituted or substituted cyclohexyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f . In some embodiments, R^B is unsubstituted or substituted cyclobutyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .




[0046] In some embodiments, R^B is an unsubstituted or substituted aziridinyl, unsubstituted or substituted azetidiny, unsubstituted or substituted oxetanyl, unsubstituted or substituted thietanyl, unsubstituted or substituted pyrrolidinyl, unsubstituted or substituted tetrahydrofuranyl, unsubstituted or substituted tetrahydrothienyl, unsubstituted or substituted oxazolidinonyl, unsubstituted or substituted tetrahydropyranyl, unsubstituted or substituted piperidinyl, unsubstituted or substituted morpholinyl, unsubstituted or substituted thiomorpholinyl, or unsubstituted or substituted piperaziny, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f . In some embodiments, R^B is an unsubstituted or substituted aziridinyl, unsubstituted or substituted azetidiny, unsubstituted or substituted pyrrolidinyl, unsubstituted or substituted piperidinyl, unsubstituted or substituted morpholinyl, unsubstituted or substituted thiomorpholinyl, or unsubstituted or substituted piperaziny, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f . In some embodiments, R^B is an unsubstituted or substituted pyrrolidinyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .






[0047] In some embodiments, R^B is , where m is 0, 1, 2, or 3. In some emodiments, m is 2 or 3. In some embodiments, m is 0. In some embodiments, m is 1. In some embodiments, m is 2. In some embodiments, m is 3.


[0048] In some embodiments, R^B is unsubstituted or substituted phenyl, or unsubstituted or substituted pyridinyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

[0049] In some embodiments, R^B is , , , , or


. In some embodiments, R^B is  or . In some embodiments,

R^B is . In some embodiments, R^B is .

[0050] In some embodiments, R^B is , where Z is CH or N. In some embodiments,

R^B is , where Z is CH or N.

[0051] In some embodiments, the compound has the structure of Formula (III), or a pharmaceutically acceptable salt, or solvate thereof:




Formula (III)

wherein,

V is CH or N;


Z is CH or N.

[0052] In some embodiments, the compound has the structure of Formula (IV), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (IV).

[0053] In some embodiments, the compound has the structure of Formula (V), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (V).

[0054] In some embodiments, R^1 is $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH(CH_3)(CH_2CH_3)$, $-C(CH_3)_3$, $-CH_2OH$, $-CH_2CN$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CH_2CH_2OH$, $-CH_2CH_2CN$, $-CH_2CH_2F$, $-CH_2CHF_2$, $-CH_2CF_3$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, $-CH_2NH_2$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2NHCH_3$, $-CH_2CH_2N(CH_3)_2$, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

[0055] In some embodiments, R^1 is $-CH_3$, $-CH_2CH_3$, or $-CH_2CH_2CH_3$.


[0056] In some embodiments, R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted $-(C_1$ - C_6 alkyl)-carbocycle, or unsubstituted or substituted $-(C_1$ - C_6 alkyl)-heterocycle, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$; R^5 is selected from the group consisting of hydrogen, and C_1 - C_6 alkyl; or R^4 and R^5 are taken together with the nitrogen atom to which they are attached to form a substituted or unsubstituted 3- to 6-membered heterocycle, wherein if the 3- to 6-membered heterocycle is substituted then the 3- to 6-membered heterocycle is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic heterocycle, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, -

$C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$.

[0057] In some embodiments, R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$; R^5 is selected from the group consisting of hydrogen, $-CH_3$, $-CH_2CH_3$, and $-CH_2CH_2CH_3$.


[0058] In some embodiments, R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted monocyclic carbocycle, or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, C_1 - C_6 alkyl that is unsubstituted or substituted with a heterocycle, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$; R^5 is selected from the group consisting of hydrogen, $-CH_3$, $-CH_2CH_3$, and $-CH_2CH_2CH_3$. In some embodiments, R^4 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, C_1 - C_6 alkyl that is unsubstituted or substituted with a heterocycle, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$; R^5 is selected from the group consisting of hydrogen, $-CH_3$, $-CH_2CH_3$, and $-CH_2CH_2CH_3$. In some embodiments, R^4 is unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, C_1 - C_6 alkyl that is unsubstituted or substituted with a 1,3-dioxol-2-one group, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, $-N(R^7)_2$, $-OR^8$, $-CN$, $-CO_2R^8$, $-C(=O)N(R^7)_2$, $-SR^8$, $-S(=O)R^{10}$, $-S(=O)_2R^{10}$, $-NR^7C(=O)R^8$, $-NR^7SO_2R^{10}$, $-SO_2R^{10}$, or $-SO_2N(R^7)_2$; R^5 is selected from the group consisting of hydrogen, $-CH_3$, $-CH_2CH_3$, and $-CH_2CH_2CH_3$.

[0059] In some embodiments, the compound has the structure of Formula (VI), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (VI).

[0060] In some embodiments, the compound has the structure of Formula (VII), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (VII)

wherein:

R^A is unsubstituted or substituted phenyl, or unsubstituted or substituted pyridinyl,


wherein if R^A is substituted then R^A is substituted with R^a , R^b and R^c ;


L is absent, $-C(=O)-$, or $-C(=O)NR^7-$;


R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle,



wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f ;

or R^B and R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 7-membered heterocycle, wherein if the 3- to 7-membered heterocycle is substituted then the 3- to 7-membered heterocycle is substituted with R^d , R^e and R^f .



[0061] In some embodiments, R^A is  or .



[0062] In some embodiments, R^A is . In some embodiments, R^A is .

[0063] In some embodiments, X^1 is CR^6 ; X^2 is CR^6 ; X^3 is CR^6 .

[0064] In some embodiments, X^1 is N; X^2 is CR^6 ; X^3 is CR^6 .

[0065] In some embodiments, X^1 is CR^6 ; X^2 is N; X^3 is CR^6 .

[0066] In some embodiments, X^1 is CR^6 ; X^2 is CR^6 ; X^3 is N.

[0067] In some embodiments, each R^6 is independently selected from the group consisting of hydrogen, halogen, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, -CN, and -OR⁸. In some embodiments, each R^6 is independently selected from the group consisting of hydrogen, halogen, and unsubstituted or substituted C_1 - C_6 alkyl. In some embodiments, each R^6 is independently selected from the group consisting of hydrogen and halogen. In some embodiments, each R^6 is independently selected from the group consisting of hydrogen and fluorine.

[0068] In some embodiments, X^1 is CH, CF, or N; X^2 is CH, CF, or N; and X^3 is CH, CF, or N.

[0069] In some embodiments, R^4 is an unsubstituted or substituted C_1 - C_6 alkyl or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-2 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂.

[0070] In some embodiments, R^4 is an unsubstituted or substituted C_1 - C_6 alkyl, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂.

[0071] In some embodiments, R^4 is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-2 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂. In some embodiments, R^4 is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-2 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, -N(R⁷)₂, -OR⁸, or -CN. In some embodiments, R^4 is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1 atom, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C_1 - C_6 alkyl, -N(R⁷)₂, -OR⁸, or -CN.

[0072] In some embodiments, R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6

fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, wherein any substituted group of R^a is substituted with one or more R⁹ groups; R^b and R^c are independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^b and R^c is substituted with one or more R⁹ groups.


[0073] In some embodiments, R^a is hydrogen, Cl, Br, -CN, -OH, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂OH, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -CH₂CH₂OH, -CH₂CH₂CN, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂NH₂, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CH₂CH₂NH₂, -CH₂CH₂NHCH₃, -CH₂CH₂N(CH₃)₂, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; R^b and R^c are independently selected from the group consisting of hydrogen, Cl, Br, -CN, -OH, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂OH, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -CH₂CH₂OH, -CH₂CH₂CN, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂NH₂, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CH₂CH₂NH₂, -CH₂CH₂NHCH₃, or -CH₂CH₂N(CH₃)₂.

[0074] In some embodiments, R^d is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R⁹ groups; R^e and R^f are independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^e and R^f is substituted with one or more R⁹ groups. In some embodiments, R^d is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, and unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^d is substituted with one or more R⁹ groups; and R^e and R^f are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, and unsubstituted or substituted C₁-C₆ alkyl, wherein any substituted group of R^e and R^f is substituted with one or more R⁹ groups.

[0075] In some embodiments, R^d is selected from the group consisting of hydrogen, Cl, Br, -CN, -OH, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -

CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂OH, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -CH₂CH₂OH, -CH₂CH₂CN, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂NH₂, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CH₂CH₂NH₂, -CH₂CH₂NHCH₃, -CH₂CH₂N(CH₃)₂, unsubstituted or substituted cyclopropyl, unsubstituted or substituted cyclobutyl, unsubstituted or substituted cyclopentyl, unsubstituted or substituted cyclohexyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R⁹ groups; R^e and R^f are independently selected from the group consisting of F, Cl, Br, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂OH, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -CN, -OH, -OCH₃, and -OCH₂CH₃.


[0076] In some embodiments, the compound has the structure of Formula (VIII), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (VIII).

[0077] In some embodiments, R^A, R^B, R⁴, and X¹ are as described in Table 1, Table 2 or Table 3.

[0078] In some embodiments, the compound of Formula (I) or Formula (II) has the structure of Formula (IX), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (IX)

wherein,

V is CH or N.

[0079] In some embodiments, the compound has the structure of Formula (IX), or a pharmaceutically acceptable salt, or solvate thereof, wherein:

V is CH or N;

R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆

fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, wherein any substituted group of R^a is substituted with one or more R⁹ groups;

R^b and R^c are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^b and R^c is substituted with one or more R⁹ groups;

R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle, unsubstituted or substituted C₁-C₇ alkyl, unsubstituted or substituted C₁-C₇ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein if R^B is substituted then R^B is substituted with R^d, R^e and R^f,

R^d is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, unsubstituted or substituted C₂-C₇ heterocycloalkyl, unsubstituted or substituted phenyl or unsubstituted or substituted monocyclic heteroaryl, wherein any substituted group of R^d is substituted with one or more R⁹ groups;

R^e and R^f are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^e and R^f is substituted with one or more R⁹ groups;

X¹ is CR⁶ or N;

R² and R³ are hydrogen;

or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-;

R⁴ is selected from the group consisting of hydrogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted -(C₁-C₆ alkyl)-carbocycle, or unsubstituted or substituted -(C₁-C₆ alkyl)-heterocycle, wherein any substituted group of R⁴ is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂;

R^6 is H or F;

each R^7 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl;

or two R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^8 is independently selected from the group consisting of hydrogen, substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R^9 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂R⁸, -CH₂CO₂R⁸, -C(=O)N(R⁷)₂, -C(=O)N(R⁷)OR⁸, -CH₂C(=O)N(R⁷)₂, -N(R⁷)₂, -CH₂N(R⁷)₂, -C(R⁸)₂N(R⁷)₂, -NR⁷C(=O)R⁸, -CH₂NR⁷C(=O)R⁸, -NR⁷C(=O)N(R⁷)₂, -NR⁷C(=O)N(R⁷)₂, C(R⁸)=N(R⁷)-OR⁸, -SR⁸, -S(=O)R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂; and

each R^{10} is independently selected from the group consisting substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl.

[0080] In some embodiments, V is CH. In some embodiments, V is N.

[0081] In some embodiments, R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein any substituted group of R^a is substituted with one or more R^9 groups; and R^b and R^c are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -N(R⁷)₂, and unsubstituted or substituted C_1 - C_6 alkyl, wherein any substituted group of R^b and R^c is substituted with one or more R^9 groups. In some embodiments, R^a is hydrogen, Cl, Br, -CN, -OH, -OCH₃, -OCH₂CH₃, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH(CH₃)(CH₂CH₃), -C(CH₃)₃, -CH₂OH, -CH₂CN, -CH₂F, -CHF₂, -CF₃, -CH₂CH₂OH, -CH₂CH₂CN, -CH₂CH₂F, -CH₂CHF₂, -CH₂CF₃, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂NH₂, -CH₂NHCH₃, -CH₂N(CH₃)₂, -CH₂CH₂NH₂, -

CH₂CH₂NHCH₃, -CH₂CH₂N(CH₃)₂, cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; R^b and R^c are hydrogen. In some embodiments, R^a is -OR⁸; and R^b and R^c are hydrogen. In some embodiments, R^a is -OH, -OCH₃, or -OCH₂CH₃; and R^b and R^c are hydrogen.

[0082] In some embodiments, X¹ is CH, CF or N; V is CH or N; R² and R³ are hydrogen; or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-; R^a is -OCH₂CH₃; R^b and R^c are hydrogen.

[0083] In some embodiments, X¹ is CH, CF or N; V is CH or N; R² and R³ are hydrogen; or R² and R³ are taken together with the carbon atom to which they are attached to form -C(=O)-; R^a is -OCH₂CH₃; R^b and R^c are hydrogen; R^B is as described in Table 1 and/or Table 2. R⁴ is as described in Table 1 and/or Table 2..


[0084] In some embodiments, each R⁷ is independently selected from the group consisting of hydrogen, substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, and unsubstituted or substituted C₁-C₆fluoroalkyl; or two R⁷ are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle. In some embodiments, each R⁷ is independently selected from the group consisting of hydrogen, and substituted C₁-C₆ alkyl; or two R⁷ are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle.

[0085] In some embodiments, each R⁸ is independently selected from the group consisting of hydrogen, substituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, and unsubstituted or substituted C₁-C₆fluoroalkyl. In some embodiments, each R⁸ is independently selected from the group consisting of hydrogen, substituted or substituted C₁-C₆ alkyl, and unsubstituted or substituted C₃-C₆ cycloalkyl. In some embodiments, each R⁸ is independently selected from the group consisting of hydrogen and substituted or substituted C₁-C₆ alkyl.

[0086] In some embodiments, each R⁹ is independently hydrogen, halogen, unsubstituted or substituted C₁-C₄alkyl, unsubstituted or substituted C₁-C₄alkoxy, unsubstituted or substituted C₁-C₄fluoroalkyl, unsubstituted or substituted C₁-C₄fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂R⁸, -CH₂CO₂R⁸, -C(=O)N(R⁷)₂, -CH₂C(=O)N(R⁷)₂, -N(R⁷)₂, -CH₂N(R⁷)₂, -NR⁷C(=O)R⁸, or -CH₂NR⁷C(=O)R⁸. In some embodiments, each R⁹ is independently hydrogen, halogen, unsubstituted or substituted C₁-C₄alkyl, unsubstituted or substituted C₁-C₄alkoxy, unsubstituted or substituted C₁-C₄fluoroalkyl, -CN, -OH, -CO₂R⁸, -C(=O)N(R⁷)₂, -N(R⁷)₂, or -CH₂N(R⁷)₂.

[0087] In some embodiments, each R^{10} is independently selected from the group consisting substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted phenyl. In some embodiments, each R^{10} is independently selected from the group consisting substituted or substituted C_1 - C_6 alkyl and unsubstituted or substituted phenyl.

[0088] In some embodiments, the compound of Formula (IX) has the structure of Formula (IXa), or a pharmaceutically acceptable salt, or solvate thereof:




Formula (IXa)

wherein,

V is CH or N.

[0089] In some embodiments, R^B , R^4 , and X^1 are as described in Table 1 and/or Table 2.


[0090] In some embodiments, compounds described herein have the following structure:



[0091] In some embodiments, R^A , R^B , R^4 , and X^1 are as described herein.

[0092] In some embodiments, R^A , R^B , R^4 , and X^1 are as described in Table 1.


[0093] In some embodiments, compounds described herein have the following structure:



[0094] In some embodiments, R^A , R^B , R^4 , and X^1 are as described herein.

[0095] In some embodiments, R^A , R^B , R^4 , and X^1 are as described in Table 2.

[0096] In some embodiments, compounds described herein have the following structure:




[0097] In some embodiments, R^A , R^B , R^1 , R^4 , and X^1 , X^2 , X^3 are as described herein.

[0098] In some embodiments, R^A , R^B , R^1 , R^4 , and X^1 , X^2 , X^3 are as described in Table 3.



















[0099] Any combination of the groups described above for the various variables is contemplated herein. Throughout the specification, groups and substituents thereof are chosen by one skilled in the field to provide stable moieties and compounds.

[00100] Exemplary compounds of Formula (I) include the compounds described in the following Tables:



















Table 1:



Compound No.	R^A	R^B	X^1	R^4
1-1			CH	$-\text{CH}_2\text{CH}_2\text{NH}_2$
1-2			CH	$-\text{CH}_2\text{CH}_2\text{NH}_2$
1-3			CH	$-\text{CH}_2\text{CH}_2\text{NH}_2$
1-4			CH	$-\text{CH}_2\text{CH}_2\text{NH}_2$

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-5			CH	-CH ₂ CH ₂ NH ₂
1-6			CH	-CH ₂ CH ₂ NH ₂
1-7			CH	-CH ₂ CH ₂ NH ₂
1-8			CH	-CH ₂ CH ₂ NH ₂
1-9			CH	-CH ₂ CH ₂ NH ₂
1-10			CH	-CH ₂ CH ₂ NH ₂
1-11			CH	-CH ₂ CH ₂ NH ₂
1-12			CH	-CH ₂ CH ₂ NH ₂
1-13			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-14			CH	-CH ₂ CH ₂ NH ₂
1-15			CH	-CH ₂ CH ₂ NH ₂
1-16			CH	-CH ₂ CH ₂ NH ₂
1-17			CH	-CH ₂ CH ₂ NH ₂
1-18			CH	-CH ₂ CH ₂ NH ₂
1-19			CH	-CH ₂ CH ₂ NH ₂
1-20			CH	-CH ₂ CH ₂ NH ₂
1-21			CH	-CH ₂ CH ₂ NH ₂
1-22			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-23			CH	-CH ₂ (CH ₂) ₂ NH ₂
1-24			CH	-CH ₂ (CH ₂) ₂ NH ₂
1-25			CH	-CH ₂ CH ₂ NH ₂
1-26			CH	-CH ₂ CH ₂ NH ₂
1-27			CH	-CH ₂ CH ₂ NH ₂
1-28			CH	-CH ₂ CH ₂ NH ₂
1-29			CH	-CH ₂ CH ₂ NH ₂
1-30			CH	-CH ₂ CH ₂ NH ₂
1-31			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-32			CH	-CH ₂ CH ₂ NH ₂
1-33			CH	-CH ₂ CH ₂ NH ₂
1-34			CH	-CH ₂ CH ₂ NH ₂
1-35			CH	-CH ₂ CH ₂ NH ₂
1-36			CH	-CH ₂ CH ₂ NH ₂
1-37			CH	-CH ₂ CH ₂ NH ₂
1-38			CH	-CH ₂ CH ₂ NH ₂
1-39			CH	-CH ₂ CH ₂ NH ₂
1-40			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-41			CH	-CH ₂ CH ₂ NH ₂
1-42			CH	-CH ₂ CH ₂ NH ₂
1-43			CH	-CH ₂ CH ₂ NH ₂
1-44			CH	-CH ₂ CH ₂ NH ₂
1-45			CH	
1-46			CH	
1-47			CH	
1-48			CH	
1-49			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-50			CH	-CH ₂ CH ₂ NH ₂
1-51			CH	-CH ₂ CH ₂ NH ₂
1-52			CH	-CH ₂ CH ₂ NH ₂
1-53			CH	-CH ₂ CH ₂ NH ₂
1-54			CH	-CH ₂ CH ₂ NH ₂
1-55			CH	-CH ₂ CH ₂ NH ₂
1-56			CH	-CH ₂ CH ₂ NH ₂
1-57			CH	-CH ₂ CH ₂ NH ₂
1-58			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-59			CH	-CH ₂ CH ₂ NH ₂
1-60			CH	-CH ₂ CH ₂ NH ₂
1-61			CH	-CH ₂ CH ₂ NH ₂
1-62			CH	-CH ₂ CH ₂ NH ₂
1-63			CH	-CH ₂ CH ₂ NH ₂
1-64			CH	
1-65			CH	
1-66			CH	
1-67			CH	-CH ₂ CH ₂ NH ₂
























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-68			CH	-CH ₂ CH ₂ NH ₂
1-69			CF	-CH ₂ CH ₂ NH ₂
1-70			CH	
1-71			CH	-CH ₂ CH ₂ NH ₂
1-72			CF	
1-73			CF	
1-74			CH	-CH ₂ CH ₂ NH ₂
1-75			CH	-CH ₂ CH ₂ NH ₂
1-76			CH	-CH ₂ CH ₂ NH ₂

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-77			CH	-CH ₂ CH ₂ NH ₂
1-78			CH	-CH ₂ CH ₂ NH ₂
1-79			CH	-CH ₂ CH ₂ NH ₂
1-80			CH	-CH ₂ CH ₂ NH ₂
1-81			CH	-CH ₂ CH ₂ NH ₂
1-82			CH	
1-83			CH	
1-84			CH	
1-85			CH	-CH ₂ CH ₂ NH ₂




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-86			CH	-CH ₂ CH ₂ NH ₂
1-87			CH	-CH ₂ CH ₂ NH ₂
1-88			CH	-CH ₂ CH ₂ NH ₂
1-89			CH	-CH ₂ CH ₂ NH ₂
1-90			CH	
1-91			CH	
1-92			CH	
1-93			CH	
1-94			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-95			CF	
1-96			CF	
1-97			CH	
1-98			CH	
1-99			CH	
1-100			CH	
1-101			CH	
1-102			CH	
1-103			CH	




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-104			CH	-CH ₂ CH ₂ NH ₂
1-105			CH	-CH ₂ CH ₂ NH ₂
1-106			CH	-CH ₂ CH ₂ NH ₂
1-107			CH	-CH ₂ CH ₂ NH ₂
1-108			CH	
1-109			CH	
1-110			CF	
1-111			CF	
1-112			CF	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-113			CF	
1-114			CF	
1-115			CH	
1-116			CF	
1-117			CH	-CH ₂ CH ₂ NH ₂
1-118			CH	-CH ₂ CH ₂ NH ₂
1-119			CH	-CH ₂ CH ₂ NH ₂
1-120			CH	-CH ₂ CH ₂ NH ₂
1-121			CF	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-122			CF	
1-123			CF	
1-124			CF	
1-125			CH	
1-126			CH	-CH ₂ CH ₂ NH ₂
1-127			CH	-CH ₂ CH ₂ NH ₂
1-128			CH	
1-129			CH	
1-130			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-131			CH	
1-132			CH	
1-133			CH	
1-134			CH	
1-135			N	
1-136			N	
1-137			N	
1-138			CH	
1-139			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-140			CF	
1-141			CF	
1-142			CH	
1-143			CH	
1-144			CH	
1-145			CH	-CH ₂ CH ₂ NH ₂
1-146			CH	-CH ₂ CH ₂ NH ₂
1-147			CH	
1-148			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-149			N	
1-150			N	
1-151			N	
1-152			N	
1-153			CF	
1-154			CF	
1-155			CF	
1-156			CF	
1-157			CF	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-158			N	
1-159			CH	-CH ₂ CH ₂ NH ₂
1-160			CH	-CH ₂ CH ₂ NH ₂
1-161			CH	
1-162			CH	
1-163			CF	
1-164			CF	
1-165			CF	
1-166			CF	




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-167			N	
1-168			CF	
1-169			N	
1-170			CF	
1-171			N	
1-172			N	
1-173			CF	
1-174			CF	
1-175			CF	


























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-176			CF	
1-177			CF	
1-178			CF	
1-179			CF	
1-180			CF	
1-181			N	
1-182			CF	
1-183			CF	
1-184			CF	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-185			N	
1-186			CF	
1-187			N	
1-188			CF	
1-189			CF	
1-190			CF	
1-191			CH	
1-192			CH	
1-193			CH	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-194			N	
1-195			N	
1-196			N	
1-197			CF	
1-198			CF	
1-199			CF	
1-200			CF	
1-201			CF	
1-202			CF	
























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-203			CF	
1-204			CF	
1-205			N	
1-206			N	
1-207			N	
1-208			N	
1-209			N	
1-210			N	
1-211			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-212			N	
1-213			N	
1-214			N	
1-215			N	
1-216			CH	
1-217			CF	
1-218			N	
1-219			CF	
1-220			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-221			N	
1-222			N	
1-223			N	
1-224			N	
1-225			N	
1-226			N	
1-227			N	
1-228			CH	
1-229			CF	































Compound No.	R ^A	R ^B	X ¹	R ⁴
1-230			N	
1-231			N	
1-232			N	
1-233			N	
1-234			N	
1-235			N	
1-236			N	
1-237			N	
1-238			N	




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-239			N	
1-240			N	
1-241			N	
1-242			N	
1-243			N	
1-244			N	
1-245			N	
1-246			N	
1-247			N	



























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-248			N	
1-249			N	
1-250			N	
1-251		 cis	CF	
1-252		 trans	CF	
1-253			N	
1-254			N	
1-255			N	
1-256			N	































Compound No.	R ^A	R ^B	X ¹	R ⁴
1-257			N	
1-258			N	
1-259			N	
1-260			N	
1-261			N	
1-262			N	
1-263			N	
1-264			N	
1-265			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-266			N	
1-267			N	
1-268			N	
1-269			N	
1-270			CF	
1-271			CF	
1-272			N	
1-273			N	
1-274			N	


































Compound No.	R ^A	R ^B	X ¹	R ⁴
1-275			N	
1-276			N	
1-277			N	
1-278			N	
1-279			N	
1-280			N	
1-281			N	
1-282			N	
1-283			N	
1-284			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-285			N	
1-286		 cis	N	
1-287			N	
1-288			N	
1-289			N	
1-290			N	
1-291			N	
1-292			N	
1-293			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-294			N	
1-295			N	
1-296			N	
1-297			N	
1-298			N	
1-299		 cis	N	
1-300		 cis	N	
1-301		 trans	N	
1-302			N	































Compound No.	R ^A	R ^B	X ¹	R ⁴
1-303			N	
1-304			N	
1-305			N	
1-306			N	
1-307			N	
1-308		 racemic	N	
1-309			N	
1-310			N	
1-311			N	
1-312			N	




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-313			N	
1-314			N	
1-315			N	
1-316			N	
1-317			N	
1-318			N	
1-319			N	
1-320			N	
1-321			N	
1-322			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-323			N	
1-324			N	
1-325			N	
1-326			N	
1-327			N	
1-328			N	
1-329			N	
1-330			N	
1-331			N	
1-332			N	
1-333			N	































Compound No.	R ^A	R ^B	X ¹	R ⁴
1-334			N	
1-335			N	
1-336			N	
1-337			N	
1-338			N	
1-339			N	
1-340			N	
1-341			N	
1-342			N	
1-343			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-344			N	
1-345			N	
1-346			N	
1-347			N	
1-348			N	
1-349			N	
1-350			N	
1-351			N	
1-352			N	
1-353			N	




























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-354			N	
1-355			N	
1-356			N	
1-357			N	
1-358			N	
1-359			N	
1-360			N	
1-361			N	
1-362			N	
1-363			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-364			N	
1-365			N	
1-366			N	
1-367			N	
1-368			N	
1-369			N	
1-370			N	
1-371			N	
1-372			N	
1-373			N	


























Compound No.	R ^A	R ^B	X ¹	R ⁴
1-374			N	
1-375			N	
1-376			N	
1-377			N	
1-378			N	
1-379			N	
1-380			N	
1-381			N	
1-382			N	
1-383			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-384			N	
1-385			N	
1-386			N	
1-387			N	
1-388			N	
1-389			N	
1-390			N	
1-391			N	
1-392			N	
1-393			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-394			N	
1-395			N	
1-396			N	
1-397			N	
1-398			N	
1-399			N	
1-400			N	
1-401			N	
1-402			N	
1-403			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-404			N	
1-405			N	
1-406			N	
1-407			N	
1-408			N	
1-409			N	
1-410			N	
1-411			N	
1-412			N	
1-413			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-414			N	
1-415			N	
1-416			N	
1-417			N	
1-418			N	
1-419			N	
1-420			N	
1-421			N	
1-422			N	
1-423			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-424			N	
1-425			N	
1-426			N	
1-427			N	
1-428			N	
1-429			N	
1-430			N	
1-431			N	
1-432			N	

Compound No.	R ^A	R ^B	X ¹	R ⁴
1-433			N	
1-434			N	
1-435			N	
1-436			N	
1-437			N	
1-438			N	
1-439			N	

* absolute stereochemistry not determined

[00101] Compounds in Table 1 are named:

1-1: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide;

1-2: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methoxy-[1,1'-biphenyl]-3-carboxamide;

1-3: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

- 1-4:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-fluoro-[1,1'-biphenyl]-3-carboxamide;
- 1-5:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethyl-[1,1'-biphenyl]-3-carboxamide;
- 1-6:** N-(2-aminoethyl)-2'-chloro-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-carboxamide;
- 1-7:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyano-[1,1'-biphenyl]-3-carboxamide;
- 1-8:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methyl-[1,1'-biphenyl]-3-carboxamide;
- 1-9:** N-(2-aminoethyl)-2'-chloro-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3'-fluoro-[1,1'-biphenyl]-3-carboxamide;
- 1-10:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-11:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-12:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-(propan-2-yloxy)-[1,1'-biphenyl]-3-carboxamide;
- 1-13:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-hydroxy-[1,1'-biphenyl]-3-carboxamide;
- 1-14:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-15:** (R)-N-(2-aminoethyl)-4-(4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl)-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-16:** (R)-N-(2-aminoethyl)-4-(4-(2-chloro-4-(trifluoromethyl)benzoyl)-2-ethylpiperazin-1-yl)-2'-(methoxymethyl)-[1,1'-biphenyl]-3-carboxamide;
- 1-17:** (R)-N-(2-aminoethyl)-4-(4-(4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl)-2-ethylpiperazin-1-yl)-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-18:** (R)-N-(2-aminoethyl)-4-(4-(4'-(aminomethyl)-5-methyl-[1,1'-biphenyl]-2-carbonyl)-2-ethylpiperazin-1-yl)-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-19:** methyl (R)-3'-((2-aminoethyl)carbamoyl)-4'-(4-(2-chloro-4-(trifluoromethyl)benzoyl)-2-ethylpiperazin-1-yl)-[1,1'-biphenyl]-2-carboxylate;
- 1-20:** (R)-N³-(2-aminoethyl)-4'-(4-(2-chloro-4-(trifluoromethyl)benzoyl)-2-ethylpiperazin-1-yl)-N²-methyl-[1,1'-biphenyl]-2,3'-dicarboxamide;

- 1-21:** (R)-N3'-(2-aminoethyl)-4'-(4-(2-chloro-4-(trifluoromethyl)benzoyl)-2-ethylpiperazin-1-yl)-N2,N2-dimethyl-[1,1'-biphenyl]-2,3'-dicarboxamide;
- 1-22:** N-(2-aminoethyl)-2'-(aminomethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-carboxamide;
- 1-23:** N-(3-aminopropyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-24:** N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-25:** N-(2-aminoethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-26:** N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-3'-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-27:** N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-3-fluoro-5-methyl-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 1-28:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propyl-[1,1'-biphenyl]-3-carboxamide;
- 1-29:** N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-4'-propyl-[1,1'-biphenyl]-3-carboxamide;
- 1-30:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(pyridin-2-yl)benzamide;
- 1-31:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(pyridin-3-yl)benzamide;
- 1-32:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(3-ethoxypyridin-2-yl)benzamide;
- 1-33:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(3-ethoxypyridin-4-yl)benzamide;
- 1-34:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-35:** N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(4-ethoxypyridin-3-yl)benzamide;
- 1-36:** N-(2-aminoethyl)-2-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-37:** N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

- 1-38:** N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-39:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(pyridin-4-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-40:** N-(2-aminoethyl)-2-[(2R)-4-{2-[(3R)-3-(aminomethyl)pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-41:** N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-(aminomethyl)pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-42:** N-(2-aminoethyl)-2-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-43:** N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-44:** N-(2-aminoethyl)-2-[(2R)-4-[4'-(aminomethyl)-3-fluoro-5-methyl-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-45:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-46:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;
- 1-47:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-48:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;
- 1-49:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(pyrrolidin-1-yl)ethyl]benzamide;
- 1-50:** N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-(trifluoromethyl)benzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-51:** N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-fluorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-52:** N-(2-aminoethyl)-2-[(2R)-4-[2-(4-aminopiperidin-1-yl)-4-chlorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-53:** N-(2-aminoethyl)-2-[(2R)-4-(3,5-dichloropyridine-2-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-54:** N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-methoxybenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

- 1-55:** N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-56:** N-(2-aminoethyl)-2-[(2R)-4-{4-[3-(aminomethyl)pyrrolidin-1-yl]-2-fluorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-57:** N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-aminopyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-58:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;
- 1-59:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;
- 1-60:** N-(2-aminoethyl)-2-[(2R)-4-(2,4-dimethylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-61:** N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(pyrrolidin-1-yl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-62:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;
- 1-63:** N-(2-aminoethyl)-2-[(2R)-4-(2-cyclopropylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-64:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-2-yl)methyl]benzamide;
- 1-65:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-3-yl)methyl]benzamide;
- 1-66:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-4-yl)methyl]benzamide;
- 1-67:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-methyl-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]benzamide;
- 1-68:** N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-4-methylbenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-69:** N-(2-aminoethyl)-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-70:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-(pyridin-4-yl)benzamide;
- 1-71:** N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-methoxypyridin-3-yl)benzamide;

- 1-72:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-73:** 6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-74:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(piperidin-4-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-75:** N-(2-aminoethyl)-2-[(2R)-4-[2-(6-aminopyridin-3-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-76:** N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-77:** N-(2-aminoethyl)-2-[(2R)-4-(2,6-dimethylpyridine-3-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-78:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(4-fluoro-2-methylbenzoyl)piperazin-1-yl]benzamide;
- 1-79:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(4-fluoro-2-methoxybenzoyl)piperazin-1-yl]benzamide;
- 1-80:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(2-methoxy-4-methylbenzoyl)piperazin-1-yl]benzamide;
- 1-81:** N-(2-aminoethyl)-2-[(2R)-4-(4-tert-butylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-82:** 2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-83:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(1H-imidazol-5-yl)methyl]benzamide;
- 1-84:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-imidazol-5-yl)ethyl]benzamide;
- 1-85:** N-(2-aminoethyl)-2-[(2R)-4-[2-(2-aminopyridin-4-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-86:** N-(2-aminoethyl)-2-[(2R)-4-(4,6-dimethylpyridine-3-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-87:** N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-88:** N-(2-aminoethyl)-2-[(2R)-4-(4-cyclopropyl-2-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

- 1-89:** N-(2-aminoethyl)-2-[(2R)-4-(3-chloro-5-fluoropyridine-2-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-90:** 2-[(2R)-4-[4-chloro-2-(piperazin-1-yl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-91:** 2-[(2R)-4-[4-chloro-2-(dimethylamino)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-92:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-pyrrolidin-3-yl]benzamide;
- 1-93:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;
- 1-94:** 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-methyl-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-95:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-96:** 6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-97:** N-{[(2S)-azetidin-2-yl]methyl}-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-98:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2S)-pyrrolidin-2-yl]methyl}benzamide;
- 1-99:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-piperidin-3-yl]benzamide;
- 1-100:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-piperidin-3-yl]benzamide;
- 1-101:** 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-fluoro-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-102:** N-{[(2R)-azetidin-2-yl]methyl}-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-103:** 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{2-fluoro-4-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-104:** N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-methylbenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-105:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(morpholin-4-yl)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;

- 1-106:** N-(2-aminoethyl)-2-[(2R)-4-[3-(dimethylamino)-5-(trifluoromethyl)pyridine-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-107:** N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-108:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2R)-pyrrolidin-2-yl]methyl}benzamide;
- 1-109:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2S)-morpholin-2-yl]methyl}benzamide;
- 1-110:** 6-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-111:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-112:** N-{[(2S)-azetidin-2-yl]methyl}-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-113:** 6-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-114:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-115:** 2-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-116:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;
- 1-117:** N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-118:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[5-fluoro-3-(trifluoromethyl)pyridine-2-carbonyl]piperazin-1-yl]benzamide;
- 1-119:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(methylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;
- 1-120:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(pyrrolidin-1-yl)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;
- 1-121:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-122:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

- 1-123:** N-{[(2R)-azetidin-2-yl]methyl}-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-124:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-{[(2R)-pyrrolidin-2-yl]methyl}benzamide;
- 1-125:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-126:** 2-[(2R)-4-[2-amino-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-127:** N-(2-aminoethyl)-2-[(2R)-4-[6-(dimethylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-128:** 2-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;
- 1-129:** N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-imidazol-1-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-130:** N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-pyrrol-1-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-131:** N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-pyrazol-1-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-132:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(1H-imidazol-2-yl)methyl]benzamide;
- 1-133:** 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[5-(trifluoromethyl)pyridine-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-134:** 2-[(2R)-4-[3-chloro-5-(trifluoromethyl)pyridine-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-135:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-136:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-137:** 2'-ethoxy-5-[(2R)-4-[4-ethoxy-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-138:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-morpholin-3-yl]methyl}benzamide;
- 1-139:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]benzamide;

- 1-140:** 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;
- 1-141:** 6-[(2R)-4-(4-chloro-2-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;
- 1-142:** 2-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-143:** 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-144:** 2-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-145:** 2-[(2R)-4-[6-amino-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-146:** N-(2-aminoethyl)-2-[(2R)-4-[4-(dimethylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-147:** 2-[(2R)-4-[2-(dimethylamino)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-148:** 2-[(2R)-4-[2-(3-aminoazetidin-1-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-149:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-6-(2-ethoxyphenyl)pyridine-2-carboxamide;
- 1-150:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-151:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-152:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-153:** 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-154:** 6-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-155:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-156:** 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

- 1-157:** 6-[(2R)-4-[2-cyano-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-158:** 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-159:** N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-methoxy-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;
- 1-160:** N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-6-methylpyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-161:** 2-[(2R)-4-[2-(azetidin-1-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 1-162:** N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(pyrrolidin-1-yl)benzoyl]piperazin-1-yl]benzamide;
- 1-163:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-(methylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-164:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-165:** 6-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-166:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-methoxy-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-167:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-168:** N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;
- 1-169:** N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-170:** N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;
- 1-171:** 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-172:** 5-[(2R)-4-(2-carbamoyl-4-chlorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

- 1-173:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-174:** N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;
- 1-175:** 6-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-176:** 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-177:** 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-178:** 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-179:** 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-180:** 6-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-181:** 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-182:** N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;
- 1-183:** 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-184:** 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-185:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methyl-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-186:** 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-187:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-188:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-189:** 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

- 1-190:** 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-191:** N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{2-[ethyl(methyl)amino]-4-fluorobenzoyl}piperazin-1-yl]benzamide;
- 1-192:** 2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(4,6-dimethoxypyrimidin-5-yl)-N-[2-(methylamino)ethyl]benzamide;
- 1-193:** 2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(6-methoxy-4-oxo-1,4-dihydropyrimidin-5-yl)-N-[2-(methylamino)ethyl]benzamide;
- 1-194:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethyl-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-195:** 5-[(2R)-4-(2-chloro-4-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-196:** 5-[(2R)-4-(2-chloro-4-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-197:** 6-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-198:** N-(azetidin-3-yl)-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-199:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;
- 1-200:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3S)-pyrrolidin-3-yl]benzamide;
- 1-201:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3S)-1-methylpyrrolidin-3-yl]benzamide;
- 1-202:** N-(azetidin-3-yl)-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;
- 1-203:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-(1-methylazetidin-3-yl)benzamide;
- 1-204:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-(1-methylazetidin-3-yl)benzamide;
- 1-205:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethyl-[2,3'-bipyridine]-6-carboxamide;
- 1-206:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

- 1-207:** N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-208:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[2-(methylamino)ethyl]pyridine-2-carboxamide;
- 1-209:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-210:** 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-211:** N-(azetidin-3-yl)-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-212:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-213:** 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-214:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-215:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide.
- 1-216:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-(1-methylazetidin-3-yl)benzamide;
- 1-217:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(4-ethoxypyrimidin-5-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-218:** 5-[(2R)-4-(2-chloro-4-cyclopropoxybenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-219:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(4,5-dihydro-1H-imidazol-2-yl)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;
- 1-220:** N-(azetidin-3-yl)-3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridine-2-carboxamide;
- 1-221:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide.
- 1-222:** N-[(azetidin-3-yl)methyl]-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-223:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

- 1-224:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(1H-imidazol-2-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-225:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-(ethylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-226:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-227:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(1-methyl-4,5-dihydro-1H-imidazol-2-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-228:** 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(2R)-1-methylazetidin-2-yl]methyl}benzamide;
- 1-229:** 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(2S)-1-methylazetidin-2-yl]methyl}benzamide.
- 1-230:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-231:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(2R)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-232:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-233:** 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-234:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(1-methyl-1H-imidazol-2-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-235:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopropanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-236:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;
- 1-237:** 3-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;
- 1-238:** 5-[(2R)-4-(adamantane-2-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-239:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-240:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-241:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-242:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-243:** 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-244:** N-[[[(2S)-azetidin-2-yl]methyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-245:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[[[(2S)-1-methylazetidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-246:** 5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2'-methoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-247:** 5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-2'-methoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-248:** 5-[(2R)-4-[1-(2,2-difluoroethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-249:** 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-250:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-ethylcyclobutanecarbonyl)piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-251:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-252:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-*trans*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 1-253:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-254:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-255:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-256:** N-{{[(2S)-azetidin-2-yl]methyl}-2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-257:** 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-{{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-258:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-259:** 5-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-260:** 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-261:** 5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-262:** 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-263:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-264:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-ethylcyclopentanecarbonyl)piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-265:** 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-266:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-267:** 5-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-268:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(3-ethoxypyrazin-2-yl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-269:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(3-ethoxypyrazin-2-yl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;
- 1-270:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;
- 1-271:** 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;
- 1-272:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(3-ethoxypyrazin-2-yl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

- 1-273:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-274:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-275:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-276:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-277:** 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-278:** 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-279:** 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-280:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-281:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-282:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-283:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-284:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,5,7-trifluoroadamantane-1-carbonyl)piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-285:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,5,7-trifluoroadamantane-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-286:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-287:** 3-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-288:** 3-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-289:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-290:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-291:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3S,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-292:** 3-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-293:** 3-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-294:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]hept-5-ene-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-295:** 5-[(2R)-4-[1-(difluoromethyl)-3,3-difluorocyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-296:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3S,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-297:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]heptane-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-298:** 5-[(2R)-4-[3,3-difluoro-1-(trifluoromethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-299:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-300:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-3-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-301:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[*trans*-3-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-302:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,3,3-trifluoro-2,2-dimethylpropanoyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-303:** 5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-304:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

- 1-305:** 3-[(2R)-4-(2-tert-butylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-306:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-307:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-[(5-methyl-2-oxo-2H-1,3-dioxol-4-yl)methyl]pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-308:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1,2,2,3-tetramethylcyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-309:** 1,1,1-trifluoro-2-methylpropan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-310:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2R)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-311:** 5-[(2R)-4-(2-cyclopropylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-312:** 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-313:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-314:** 5-[(2R)-4-{7,7-dimethylbicyclo[2.2.1]heptane-1-carbonyl}-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-315:** 3-[(2R)-4-[(2S)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-316:** 3-[(2R)-4-[(2R)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-317:** 5-[(2R)-4-(2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-318:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3-fluoroazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-319:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3-fluoro-1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-320:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-321:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-322:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-323:** 5-[(2R)-4-(2,2-diethylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-324:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-325:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-326:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-methyl-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-327:** N-[[2,3'-bipyridine]-6-carboxamide];
- 1-328:** 5-[(2R)-4-[1-(2,2-difluoroethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-329:** N-[[2,3'-bipyridine]-6-carboxamide];
- 1-330:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-331:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-332:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-333:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-334:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2,3'-bipyridine]-6-carboxamide];
- 1-335:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-336:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

- 1-337:** N-[2-(azetidin-1-yl)ethyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-338:** 1-(trifluoromethyl)cyclobutyl (3R)-4-[6-(2-ethoxyphenyl)-2-[(3R)-pyrrolidin-3-yl]carbamoyl}pyridin-3-yl]-3-ethylpiperazine-1-carboxylate;
- 1-339:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]heptane-2-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-340:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]pyridine-2-carboxamide;
- 1-341:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-342:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-343:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-344:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-345:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-phenylpyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-346:** 5-[(2R)-4-(7-chloro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-347:** 5-[(2R)-4-(5-chloro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-348:** 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-[6-(2-ethoxyphenyl)-2-[(3R)-pyrrolidin-3-yl]carbamoyl}pyridin-3-yl]-3-ethylpiperazine-1-carboxylate;
- 1-349:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(2,2,2-trifluoroethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-350:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(7-fluoro-2,3-dihydro-1H-indole-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-351:** (1R)-2,2,2-trifluoro-1-phenylethyl (3R)-4-(2'-ethoxy-6-[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

- 1-352:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclopentanecarbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-353:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclobutanecarbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-354:** 5-[(2R)-4-[2,2-difluoro-7,7-dimethylbicyclo[2.2.1]heptane-1-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-355:** 3-[(2R)-4-[(2S)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-356:** 3-[(2R)-4-[(2R)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-357:** 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(3-fluoroazetidin-1-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-358:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclobutanecarbonyl)piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-359:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(2,2,2-trifluoroethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-360:** 2'-ethoxy-5-[(2R)-2-ethyl-4-(7-methyl-2,3-dihydro-1H-indole-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-361:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-362:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]pyridine-2-carboxamide;
- 1-363:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-364:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-365:** 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-366:** (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-367:** (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-368:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[5-(trifluoromethyl)thiophene-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-369:** 1,1,1,3,3,3-hexafluoropropan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-370:** 3-(trifluoromethyl)pentan-3-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-371:** 5-[(2R)-4-[1-(2,4-difluorophenyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-372:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-373:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-374:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-375:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-376:** 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-377:** (1S)-2,2,2-trifluoro-1-phenylethyl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-378:** 5-[(2R)-4-(2,5-dichlorothiophene-3-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-379:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[3-(trifluoromethyl)bicyclo[1.1.1]pentane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-380:** 5-[(2R)-4-(3,5-dichlorothiophene-2-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-381:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(3-fluorophenyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-382:** 5-[(2R)-4-(7-chloro-2,3-dihydro-1H-indene-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-383:** 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-384:** 1-phenylcyclobutyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-385:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

- 1-386:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2R)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-387:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}pyridine-2-carboxamide;
- 1-388:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-389:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-390:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-391:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;
- 1-392:** (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate; or (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-393:** 3-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-394:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-395:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-396:** (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(1-methylazetidin-3-yl)methyl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate; or (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(1-methylazetidin-3-yl)methyl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-397:** 1-ethylcyclopentyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-398:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;
- 1-399:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

- 1-400:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-401:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-402:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-403:** (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(6-{[(3S)-1-azabicyclo[2.2.2]octan-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate; or (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(6-{[(3S)-1-azabicyclo[2.2.2]octan-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-404:** 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-405:** N-{1-azabicyclo[2.2.1]heptan-4-yl}-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-406:** (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl(3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-407:** (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl(3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-408:** 2,2-dimethyl-1-phenylpropyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;
- 1-409:** 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-410:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxamide;
- 1-411:** N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-412:** 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-413:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-414:** 3-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

- 1-415:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methyl-4,5-dihydro-1H-imidazol-2-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;
- 1-416:** 5-[(2R)-4-[1-(difluoromethyl)cyclopentanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-417:** 3-[(2R)-4-[1-(difluoromethyl)cyclopentanecarbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-418:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-419:** 5-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-420:** 5-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-421:** 3-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-422:** 5-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;
- 1-423:** 5-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-424:** 3-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;
- 1-425:** 5-[(2R)-4-(7-chloro-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-426:** 3-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-427:** 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-428:** 5-[(2R)-4-(7-chloro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-429:** 5-[(2R)-4-(7-cyano-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;
- 1-430:** 3-[(2R)-4-(7-chloro-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;
- 1-431:** 3-[(2R)-4-(7-chloro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-432: 3-[(2R)-4-(7-cyano-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-433: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-434: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-435: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;


1-436: 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-437: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-438: 5-[(2R)-4-(7-cyano-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide; and

1-439: 3-[(2R)-4-(7-cyano-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide.

Table 2:



Compound No.	R^A	R^B	X^1	R^4
2-1			CH	
2-2			CH	
2-3			CH	H

Compound No.	R ^A	R ^B	X ¹	R ⁴
2-4			CH	H
2-5			CH	H
2-6			CH	H
2-7			CH	H
2-8			CH	H
2-9			CH	H
2-10			CH	H
2-11			CF	H
2-12			N	H

Compound No.	R ^A	R ^B	X ¹	R ⁴
2-13			N	H
2-14			N	H
2-15			N	H
2-16			N	-CH ₃
2-17			N	H
2-18			CF	H
2-19			N	H
2-20			N	H
2-21			CH	
2-22			CH	H


Compound No.	R ^A	R ^B	X ¹	R ⁴
2-23			CH	H
2-24			CH	H
2-25			N	H
2-26			CH	H
2-27			CH	H
2-28			CH	H
2-29			CH	H
2-30			N	H
2-31			N	H

[00102] Compounds in Table 2 are named:

2-1: 2-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methylamino]ethan-1-ol;

- 2-2:** (2S)-1-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl]amino]propan-2-ol;
- 2-3:** 1-{4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methoxy-[1,1'-biphenyl]-3-yl}methanamine;
- 2-4:** 1-{4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-(2-methoxyethoxy)-[1,1'-biphenyl]-3-yl}methanamine;
- 2-5:** 1-{4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propoxy-[1,1'-biphenyl]-3-yl}methanamine;
- 2-6:** 2-{[3'-(aminomethyl)-4'-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-2-yl]oxy}ethan-1-ol;
- 2-7:** 2-{[3'-(aminomethyl)-4'-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[1,1'-biphenyl]-2-yl]oxy}ethan-1-ol;
- 2-8:** [3'-(aminomethyl)-4'-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-2-yl]methanol;
- 2-9:** 1-{2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}methanamine;
- 2-10:** 1-{2-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}methanamine;
- 2-11:** 1-{6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}methanamine;
- 2-12:** 1-{3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridin-2-yl}methanamine;
- 2-13:** 1-{5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;
- 2-14:** 1-{5-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;
- 2-15:** 1-{5-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;
- 2-16:** ({5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl)methyl}(methyl)amine;
- 2-17:** 1-{5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;
- 2-18:** 1-{6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}methanamine;

- 2-19:** 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-(2,3,4-trifluorobenzoyl)piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;
- 2-20:** 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;
- 2-21:** {[5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]phenyl]methyl}[2-(methylamino)ethyl]amine;
- 2-22:** 13-[3-(aminomethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]phenyl]-N-methylpyridin-2-amine;
- 2-23:** 3-[3-(aminomethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]phenyl]-N,N-dimethylpyridin-2-amine;
- 2-24:** 1-{2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-[2-(pyrrolidin-1-yl)pyridin-3-yl]phenyl}methanamine;
- 2-25:** 1-{2'-ethoxy-5-[(2R)-4-[4-ethoxy-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;
- 2-26:** 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-3'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;
- 2-27:** 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[1,1'-biphenyl]-3-yl}methanamine;
- 2-28:** 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-4'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;
- 2-29:** 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-5'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;
- 2-30:** 1-{2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine; and
- 2-31:** 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine.

Table 3:

Compound No.	R ^A	-L-R ^B	X ¹	X ²	X ³	R ¹	R ⁴
3-1			CH	CH	CH	H	
3-2			CH	CH	CH	H	
3-3			CF	CH	CH	(2 <i>R</i>)-CH ₂ CH ₃	
3-4			CF	CH	CH	(2 <i>R</i>)-CH ₂ CH ₃	
3-5			CH	CH	CH	(3 <i>R</i>)-CH ₂ CH ₃	
3-6			CH	CH	CH	(3 <i>R</i>)-CH ₃	
3-7			N	CH	N	(2 <i>R</i>)-CH ₂ CH ₃	
3-8			N	CH	N	(2 <i>R</i>)-CH ₂ CH ₃	
3-9			CH	N	CH	(2 <i>R</i>)-CH ₂ CH ₃	

[00103] Compounds in Table 3 are named:

- 3-1:** N-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide;
- 3-2:** N-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;
- 3-3:** 6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;
- 3-4:** (3R)-N-(2,4-dichlorophenyl)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-{[2-(methylamino)ethyl]carbonyl}phenyl]-3-ethylpiperazine-1-carboxamide;
- 3-5:** N-(2-aminoethyl)-2-[(3R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-3-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;
- 3-6:** 2-[(3R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-3-methylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;
- 3-7:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxamide;
- 3-8:** 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]pyrazine-2-carboxamide; and
- 3-9:** N-(2-aminoethyl)-5-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2-(2-ethoxyphenyl)pyridine-4-carboxamide.

[00104] In one aspect, compounds described herein are in the form of pharmaceutically acceptable salts. As well, active metabolites of these compounds having the same type of activity are included in the scope of the present disclosure. In addition, the compounds described herein can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. The solvated forms of the compounds presented herein are also considered to be disclosed herein.

[00105] “Pharmaceutically acceptable,” as used herein, refers a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively nontoxic, i.e., the material is administered to an individual without causing undesirable biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

[00106] The term “pharmaceutically acceptable salt” refers to a form of a therapeutically active agent that consists of a cationic form of the therapeutically active agent in combination with a suitable anion, or in alternative embodiments, an anionic form of the therapeutically active agent in combination with a suitable cation. Handbook of Pharmaceutical Salts: Properties, Selection

and Use. International Union of Pure and Applied Chemistry, Wiley-VCH 2002. S.M. Berge, L.D. Bighley, D.C. Monkhouse, J. Pharm. Sci. 1977, 66, 1-19. P. H. Stahl and C. G. Wermuth, editors, *Handbook of Pharmaceutical Salts: Properties, Selection and Use*, Weinheim/Zürich:Wiley-VCH/VHCA, 2002. Pharmaceutical salts typically are more soluble and more rapidly soluble in stomach and intestinal juices than non-ionic species and so are useful in solid dosage forms. Furthermore, because their solubility often is a function of pH, selective dissolution in one or another part of the digestive tract is possible and this capability can be manipulated as one aspect of delayed and sustained release behaviours. Also, because the salt-forming molecule can be in equilibrium with a neutral form, passage through biological membranes can be adjusted.

[00107] In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I) with an acid. In some embodiments, the compound of Formula (I) (i.e. free base form) is basic and is reacted with an organic acid or an inorganic acid. Inorganic acids include, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, and metaphosphoric acid. Organic acids include, but are not limited to, 1-hydroxy-2-naphthoic acid; 2,2-dichloroacetic acid; 2-hydroxyethanesulfonic acid; 2-oxoglutaric acid; 4-acetamidobenzoic acid; 4-aminosalicylic acid; acetic acid; adipic acid; ascorbic acid (L); aspartic acid (L); benzenesulfonic acid; benzoic acid; camphoric acid (+); camphor-10-sulfonic acid (+); capric acid (decanoic acid); caproic acid (hexanoic acid); caprylic acid (octanoic acid); carbonic acid; cinnamic acid; citric acid; cyclamic acid; dodecylsulfuric acid; ethane-1,2-disulfonic acid; ethanesulfonic acid; formic acid; fumaric acid; galactaric acid; gentisic acid; glucoheptonic acid (D); gluconic acid (D); glucuronic acid (D); glutamic acid; glutaric acid; glycerophosphoric acid; glycolic acid; hippuric acid; isobutyric acid; lactic acid (DL); lactobionic acid; lauric acid; maleic acid; malic acid (- L); malonic acid; mandelic acid (DL); methanesulfonic acid; naphthalene-1,5-disulfonic acid; naphthalene-2-sulfonic acid; nicotinic acid; oleic acid; oxalic acid; palmitic acid; pamoic acid; phosphoric acid; propionic acid; pyroglutamic acid (- L); salicylic acid; sebacic acid; stearic acid; succinic acid; sulfuric acid; tartaric acid (+ L); thiocyanic acid; toluenesulfonic acid (*p*); and undecylenic acid.

[00108] In some embodiments, a compound of Formula (I) is prepared as a chloride salt, sulfate salt, bromide salt, mesylate salt, maleate salt, citrate salt or phosphate salt.

[00109] In some embodiments, pharmaceutically acceptable salts are obtained by reacting a compound of Formula (I) with a base. In some embodiments, the compound of Formula (I) is acidic and is reacted with a base. In such situations, an acidic proton of the compound of Formula (I) is replaced by a metal ion, e.g., lithium, sodium, potassium, magnesium, calcium, or

an aluminum ion. In some cases, compounds described herein coordinate with an organic base, such as, but not limited to, ethanolamine, diethanolamine, triethanolamine, tromethamine, meglumine, N-methylglucamine, dicyclohexylamine, tris(hydroxymethyl)methylamine. In other cases, compounds described herein form salts with amino acids such as, but not limited to, arginine, lysine, and the like. Acceptable inorganic bases used to form salts with compounds that include an acidic proton, include, but are not limited to, aluminum hydroxide, calcium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydroxide, lithium hydroxide, and the like. In some embodiments, the compounds provided herein are prepared as a sodium salt, calcium salt, potassium salt, magnesium salt, meglumine salt, N-methylglucamine salt or ammonium salt.

[00110] It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms. In some embodiments, solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and are formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described herein are conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein optionally exist in unsolvated as well as solvated forms.

[00111] The methods and formulations described herein include the use of *N*-oxides (if appropriate), or pharmaceutically acceptable salts of compounds having the structure of Formula (I), as well as active metabolites of these compounds having the same type of activity.

[00112] In some embodiments, sites on the organic radicals (e.g. alkyl groups, aromatic rings) of compounds of Formula (I) are susceptible to various metabolic reactions. Incorporation of appropriate substituents on the organic radicals will reduce, minimize or eliminate this metabolic pathway. In specific embodiments, the appropriate substituent to decrease or eliminate the susceptibility of the aromatic ring to metabolic reactions is, by way of example only, a halogen, deuterium, an alkyl group, a haloalkyl group, or a deuterioalkyl group.

[00113] In another embodiment, the compounds described herein are labeled isotopically (e.g. with a radioisotope) or by another other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

[00114] Compounds described herein include isotopically-labeled compounds, which are identical to those recited in the various formulae and structures presented herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be

incorporated into the present compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, sulfur, fluorine, chlorine, iodine, phosphorus, such as, for example, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , ^{124}I , ^{125}I , ^{131}I , ^{32}P and ^{33}P . In one aspect, isotopically-labeled compounds described herein, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. In one aspect, substitution with isotopes such as deuterium affords certain therapeutic advantages resulting from greater metabolic stability, such as, for example, increased *in vivo* half-life or reduced dosage requirements.

[00115] In some embodiments, the compounds of Formula (I) possess one or more stereocenters and each stereocenter exists independently in either the R or S configuration. In some embodiments, the compound of Formula (I) exists in the R configuration. In some embodiments, the compound of Formula (I) exists in the S configuration. The compounds presented herein include all diastereomeric, individual enantiomers, atropisomers, and epimeric forms as well as the appropriate mixtures thereof. The compounds and methods provided herein include all *cis*, *trans*, *syn*, *anti*, *entgegen* (E), and *zusammen* (Z) isomers as well as the appropriate mixtures thereof.

[00116] Individual stereoisomers are obtained, if desired, by methods such as, stereoselective synthesis and/or the separation of stereoisomers by chiral chromatographic columns or the separation of diastereomers by either non-chiral or chiral chromatographic columns or crystallization and recrystallization in a proper solvent or a mixture of solvents. In certain embodiments, compounds of Formula (I) are prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds/salts, separating the diastereomers and recovering the optically pure individual enantiomers. In some embodiments, resolution of individual enantiomers is carried out using covalent diastereomeric derivatives of the compounds described herein. In another embodiment, diastereomers are separated by separation/resolution techniques based upon differences in solubility. In other embodiments, separation of stereoisomers is performed by chromatography or by the forming diastereomeric salts and separation by recrystallization, or chromatography, or any combination thereof. Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981. In some embodiments, stereoisomers are obtained by stereoselective synthesis.

[00117] In some embodiments, compounds described herein are prepared as prodrugs. A "prodrug" refers to an agent that is converted into the parent drug *in vivo*. Prodrugs are often useful because, in some situations, they are easier to administer than the parent drug. They are,

for instance, bioavailable by oral administration whereas the parent is not. Further or alternatively, the prodrug also has improved solubility in pharmaceutical compositions over the parent drug. In some embodiments, the design of a prodrug increases the effective water solubility. An example, without limitation, of a prodrug is a compound described herein, which is administered as an ester (the “prodrug”) but then is metabolically hydrolyzed to provide the active entity. A further example of a prodrug is a short peptide (polyaminoacid) bonded to an acid group where the peptide is metabolized to reveal the active moiety. In certain embodiments, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically active form of the compound. In certain embodiments, a prodrug is enzymatically metabolized by one or more steps or processes to the biologically, pharmaceutically or therapeutically active form of the compound.

[00118] Prodrugs of the compounds described herein include, but are not limited to, esters, ethers, carbonates, thiocarbonates, N-acyl derivatives, N-acyloxyalkyl derivatives, N-alkyloxyacyl derivatives, quaternary derivatives of tertiary amines, N-Mannich bases, Schiff bases, amino acid conjugates, phosphate esters, and sulfonate esters. See for example Design of Prodrugs, Bundgaard, A. Ed., Elsevier, 1985 and Method in Enzymology, Widder, K. *et al.*, Ed.; Academic, 1985, vol. 42, p. 309-396; Bundgaard, H. “Design and Application of Prodrugs” in A Textbook of Drug Design and Development, Krosgaard-Larsen and H. Bundgaard, Ed., 1991, Chapter 5, p. 113-191; and Bundgaard, H., Advanced Drug Delivery Review, 1992, 8, 1-38, each of which is incorporated herein by reference. In some embodiments, a hydroxyl group in the compounds disclosed herein is used to form a prodrug, wherein the hydroxyl group is incorporated into an acyloxyalkyl ester, alkoxycarbonyloxyalkyl ester, alkyl ester, aryl ester, phosphate ester, sugar ester, ether, and the like. In some embodiments, a hydroxyl group in the compounds disclosed herein is a prodrug wherein the hydroxyl is then metabolized *in vivo* to provide a carboxylic acid group. In some embodiments, a carboxyl group is used to provide an ester or amide (i.e. the prodrug), which is then metabolized *in vivo* to provide a carboxylic acid group. In some embodiments, compounds described herein are prepared as alkyl ester prodrugs.

[00119] Prodrug forms of the herein described compounds, wherein the prodrug is metabolized *in vivo* to produce a compound of Formula (I) as set forth herein are included within the scope of the claims. In some cases, some of the herein-described compounds is a prodrug for another derivative or active compound.

[00120] In some embodiments, any one of the hydroxyl group(s), amino group(s) and/or carboxylic acid group(s) are functionalized in a suitable manner to provide a prodrug moiety. In some embodiments, the prodrug moiety is as described above.

[00121] In additional or further embodiments, the compounds described herein are metabolized upon administration to an organism in need to produce a metabolite that is then used to produce a desired effect, including a desired therapeutic effect.

[00122] A “metabolite” of a compound disclosed herein is a derivative of that compound that is formed when the compound is metabolized. The term “active metabolite” refers to a biologically active derivative of a compound that is formed when the compound is metabolized. The term “metabolized,” as used herein, refers to the sum of the processes (including, but not limited to, hydrolysis reactions and reactions catalyzed by enzymes) by which a particular substance is changed by an organism. Thus, enzymes may produce specific structural alterations to a compound. For example, cytochrome P450 catalyzes a variety of oxidative and reductive reactions while uridine diphosphate glucuronyltransferases catalyze the transfer of an activated glucuronic-acid molecule to aromatic alcohols, aliphatic alcohols, carboxylic acids, amines and free sulphydryl groups. Metabolites of the compounds disclosed herein are optionally identified either by administration of compounds to a host and analysis of tissue samples from the host, or by incubation of compounds with hepatic cells in vitro and analysis of the resulting compounds.

Synthesis of Compounds


[00123] Compounds of Formula (I) described herein are synthesized using standard synthetic techniques or using methods known in the art in combination with methods described herein.

[00124] Unless otherwise indicated, conventional methods of mass spectroscopy, NMR, HPLC, protein chemistry, biochemistry, recombinant DNA techniques and pharmacology are employed.

[00125] Compounds are prepared using standard organic chemistry techniques such as those described in, for example, March's Advanced Organic Chemistry, 6th Edition, John Wiley and Sons, Inc. Alternative reaction conditions for the synthetic transformations described herein may be employed such as variation of solvent, reaction temperature, reaction time, as well as different chemical reagents and other reaction conditions.

[00126] In some embodiments, compounds described herein are prepared as described in Scheme A.

Scheme A




a) DMSO, Et₃N, 100 °C; b) Fe, NH₄Cl, EtOH/H₂O, 70 °C; c) n-Butylnitrite, CuBr₂, ACN, 70 °C; d) R^AB(OH)₂, Pd₂(dba)₃HCl₃, P(t-Bu)₃HBF₄, K₃PO₄, toluene, 70 °C; e) LiOH.H₂O, H₂O/MeOH/THF; f) TFA, DCM.

[00127] Compound **I** is obtained by heating methyl 2-fluoro-5-nitrobenzoate or methyl 2,6-difluoro-3-nitrobenzoate with a mono-protected substituted piperazine in the presence of organic base such as *N,N*-diisopropylethylamine. An iron-catalyzed reduction of nitro group yields an aryl amine, which is subsequently transformed to an aryl bromide (**II**) *via* preparation of diazonim salt and subsequent displacement with Br anion under copper (I) catalysis. Compound **II** is subjected to organometallic coupling reaction such as Suzuki–Miyaura reaction with R^AB(OH)₂ or its corresponding ester and is followed by a standard hydrolysis to generate compound **III**. Compound **IV** is converted to the intermediate **V** *via* an iron-catalyzed nitro reduction and a subsequent Sandmeyer reaction of the aniline. Alternatively, Compound **V** can be also obtained from a standard deprotection of **II**.

[00128] In some embodiments, compounds described herein are prepared as described in Scheme B.

Scheme B




a) $R^B\text{-CO}_2\text{H}$ HATU, Et_3N , or $R^B\text{COCl}$, Et_3N ; b) $R^A\text{B(OH)}_2$, $\text{Pd}_2(\text{dba})_3\text{HCl}_3$, $\text{P}(\text{t-Bu})_3\text{HBF}_4$, K_3PO_4 , toluene, 70°C ;
 c) Aryl tributyltin, $\text{Pd}(\text{dppf})\text{Cl}_2$, CuO , DMF, 100°C ; d) IPA, $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$, 70°C ; e) Bis(pinacolato)diboron, $\text{Pd}(\text{DTBPF})\text{Cl}_2$, KOAc , dioxane, H_2O , 90°C ; f) $R^A\text{-Br}$, $\text{Pd}(\text{DTBPF})\text{Cl}_2$, K_2CO_3 , dioxane, H_2O , 80°C .

[00129] Compound **V** is converted to the intermediate **VI** by treating with an activated carboxylic acid using HATU and triethylamine or an acyl chloride in the presence of triethylamine or other bases. Compound **VII** is prepared from the aryl bromide (**VI**) by organometallic coupling reactions such as Suzuki-Miyaura reaction with boronic acid $R^A\text{B(OH)}_2$ or its ester or Stille coupling with $R^A\text{SnBu}_3$. The direct conversion of the carboxylic ester (**VII**) to the amide (**VIII**) is achieved by heating it with a diamine in a polar protic solvent. Alternatively, compound **VI** is converted into the corresponding boronic ester intermediate (**IX**) which is subsequently subject to coupling reaction such as Suzuki-Miyaura reaction with an aryl halide such as $R^A\text{-Br}$ and is followed by the conversion of the ester to the amide to yield **VIII**.

[00130] In some embodiments, compounds described herein are prepared as described in Scheme C.

Scheme C




a) R^4R^5NH , HATU, Et_3N ; b) TFA, DCM; c) R^B CO₂H, HATU, Et_3N , or R^B COCl, Et_3N ; d) deprotection (optically).

[00131] Compound **III** is reacted with an amine (R^4R^5NH) in the presence of HATU and triethylamine yielding intermediate **X**. Compound **X** is subjected to removal of Boc group with TFA or other acid and then formation of the amide with R^A CO₂H, or R^A COCl to give **XI**. If R^4 or R^5 contains a protecting group, then an additional deprotection step takes place to produce **XI**. Alternatively, intermediate **XII** is obtained from deprotection of **III** by TFA is converted to **XIII** by treating with pre-activated R^B CO₂H. Compound **XIII** is utilized to produce the corresponding amides (**XI**) similarly.

[00132] In some embodiments, compounds described herein are prepared as described in Scheme D.

Scheme D




a) R^A B(OH)₂, Pd₂(dba)₃HCl₃, P(t-Bu)₃HBF₄, K₃PO₄, toluene, 70 °C; b) NaBH₄, MeOH, 50 °C; c) IBX; d) R^4R^5NH , NaBH(OAc)₃; e) TFA, DCM; f) R^B CO₂H, HATU, Et_3N ; g) deprotection (if needed)

[00133] Compound **II** is subjected to a Suzuki coupling with boronic acid $R^B(OH)_2$ in the presence of a transition metal catalyst, yielding the intermediate (**XIV**). The carboxylic ester on **XIV** is reduced to an alcohol using $NaBH_4$ followed by oxidation with 2-iodoxybenzoic acid to afford the benzaldehyde **XV**, which is subjected to reductive amination reaction with an appropriate amine (R^4R^5NH) and a reducing agent such as $NaBH(OAc)_3$ to afford **XVI**. A subsequent Boc deprotection and then amide formation with $R^B(OH)_2$ yields compound **XVII**. If R^4 contains a protecting group, then an additional deprotection step takes place to produce **XVII**.

[00134] In some embodiments, compounds described herein are prepared as described in Scheme E.

Scheme E




a) $R^A-B(OH)_2$, $PdCl_2(t-Bu_2PPhMe_2)_2$, K_2CO_3 , dioxane/ H_2O ; b) DMSO, DIEA, 140 °C; c) TFA or HCl
 d) R^B-CO_2H , HATU, Et_3N or R^B-COCl , Et_3N ; e) KOH, $EtOH/H_2O$, 100 °C; f) R^4R^5NH , HATU, Et_3N ; g) $NiCl_2$, $NaBH_4$; h) Cbz-Cl, DIPEA DCM or CbzOSu, DCM; i) TFA, 70°C.

[00135] Compound **XVIII** is prepared by Suzuki-Miyaura coupling reaction with a boronic acid $R^A-B(OH)_2$ or Stille coupling with R^A-SnBu_3 , which reacts with a mono-protected substituted piperazine in the presence of an organic base such as DIEA to afford **XIX**. Subsequently, Boc group is removed by an acid and the corresponding amide is formed via standard amide formation reaction with $R^B(OH)_2$ to afford **XX**. Nickel (II) catalyzed reduction of the carbonitrile produces benzyl amine analogues (**XXII-B**). Hydrolysis of $-CN$ on **XX** with a strong inorganic base such as KOH affords the carboxylic acid intermediate (**XXI**), which is converted to compound **XXII-A** by coupling reaction with R^4R^5NH . If R^4 contains a protecting group, then an additional deprotection step takes place to produce **XXII**. Alternatively, **XXII-A**

is prepared via 4-step synthesis starting from **XIX** by first hydrolysis of cyano group to the carboxylic acid, followed by amide formation with R^4R^5NH to form **XXIII**, then Boc group is removed and $R^B\text{CO}_2\text{H}$ or $R^B\text{COCl}$ is used to obtain **XXII-A**. If R^4 contains a protecting group, then appropriate deprotection step to remove the protecting group is conducted to produce **XXII**. In some embodiments, compound **XXII-B** is prepared via an alternative pathway from **XIX** as follows: cyano group is reduced by NaBH_4 in the presence of NiCl_2 . The newly formed amine is then protected by a Cbz group. Subsequently Boc group is removed, followed by amide formation with $R^B\text{CO}_2\text{H}$. Finally, Cbz group is removed by stirring in hot TFA to afford **XXII-B**.

[00136] In some embodiments, compounds described herein are prepared as described in Scheme F.

Scheme F




a) DMSO, DIEA, 120 °C; b) $R^AB(OH)_2$, $\text{PdCl}_2(\text{t-Bu}_2\text{PPhNMe}_2)_2$, K_2CO_3 , dioxane/ H_2O , 100 °C; c) $\text{LiOH} \cdot \text{H}_2\text{O}$, $\text{MeOH}/\text{H}_2\text{O}/\text{THF}$; d) TFA, DCM, e) $R^B\text{CO}_2\text{H}$, HATU, Et_3N ; f) HATU, R^4R^5NH ; g) deprotection (if necessary)

[00137] Heating of methyl 6-bromo-3-chloropyrazine-2-carboxylate with a mono-protected substituted piperazine yields compound **XXIV**, which is subsequently subjected to Suzuki–Miyaura coupling reaction with $R^AB(OH)_2$ and is followed by hydrolysis of ester to acid and a standard de-Boc procedure to yield **XXV**. **XXVI** is obtained from a coupling reaction with a pre-activated $R^B\text{CO}_2\text{H}$ and further reaction with R^4R^5NH in the presence of HATU affords **XXVII**. If R^4 or R^5 contains a protecting group, then an additional deprotection step takes place to produce **XXII**.

[00138] In some embodiments, compounds described herein are prepared as described in Scheme G.

Scheme G




a) TFA, DCM; b) $R^B\text{-Cl}$ (or F), DIEA, DMSO; c) $\text{LiOH}\cdot\text{H}_2\text{O}$, $\text{H}_2\text{O}/\text{MeOH}/\text{THF}$;
 d) $R^4R^5\text{NH}$, HATU, Et_3N ; e) $R^AB(\text{OH})_2$, Pd (II), K_2CO_3 , dioxane/ H_2O ; f) deprotection

[00139] The Boc group on **II** is removed to yield **XXVIII**, which is treated with $R^B\text{-Cl}$ or $R^B\text{-F}$ and is followed by saponification of ester to form **XXIX**. Coupling reaction of **XXIX** with $R^4R^5\text{NH}$ yields **XXX**, which undergoes Suzuki-Miyaura reaction with the boronic acid $R^AB(\text{OH})_2$ to afford **XXXI**. An additional removal of protecting group takes place if R^4 or R^5 contains a protecting group on **XXXI**.

[00140] In some embodiments, compounds described herein are prepared as described in Scheme H.

Scheme H



a) TFA; b) $R^B\text{SO}_2\text{Cl}$, Et_3N ;
 c) $R^B\text{NCO}$ or $R^B R^7\text{NCO}_2\text{Ph}$, Et_3N or triphosgene, Et_3N , then $R^B R^7\text{NH}$, Et_3N .
 d) CDI, $R^B\text{OH}$, DIEA, Δ

[00141] The Boc group of on **X** is removed to form **XXXII**, which is reacted with aryl sulfonyl chloride in the presence of an organic base to afford **XXXIII**. In a similar manner, urea analogues (**XXXIV**) are prepared by either reaction with an isocyanates ($R^B NCO$) or reaction $R^B R^7 CO_2 Ph$ or reacting **XXXII** with triphosgene and then $R^B R^7 NH$. In addition, an efficient synthesis of carbamate derivatives (**XXXV**) was achieved by reacting **XXXII** with the CDI activated alcohols. If R^4 contains a protecting group, then it is removed subsequently with an appropriate condition on **XXXIII**, **XXXIV**, or **XXXV**.

[00142] In some embodiments, compounds described herein are synthesized as outlined in the Examples.

Certain Terminology

[00143] Unless otherwise stated, the following terms used in this application have the definitions given below. The use of the term “including” as well as other forms, such as “include”, “includes,” and “included,” is not limiting. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

[00144] As used herein, C_1-C_x includes C_1-C_2 , C_1-C_3 . . . C_1-C_x . By way of example only, a group designated as " C_1-C_6 " indicates that there are one to six carbon atoms in the moiety, i.e. groups containing 1 carbon atom, 2 carbon atoms, 3 carbon atoms or 4 carbon atoms. Thus, by way of example only, " C_1-C_4 alkyl" indicates that there are one to four carbon atoms in the alkyl group, i.e., the alkyl group is selected from among methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, and *t*-butyl.

[00145] An “alkyl” group refers to an aliphatic hydrocarbon group. The alkyl group is branched or straight chain. In some embodiments, the “alkyl” group has 1 to 10 carbon atoms, i.e. a C_1-C_{10} alkyl. Whenever it appears herein, a numerical range such as “1 to 10” refers to each integer in the given range; e.g., “1 to 10 carbon atoms” means that the alkyl group consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 10 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated. In some embodiments, an alkyl is a C_1-C_6 alkyl. In one aspect the alkyl is methyl, ethyl, propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, or *t*-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, tertiary butyl, pentyl, neopentyl, or hexyl.

[00146] An “alkylene” group refers refers to a divalent alkyl radical. Any of the above mentioned monovalent alkyl groups may be an alkylene by abstraction of a second hydrogen atom from the alkyl. In some embodiments, an alkylene is a C_1-C_6 alkylene. In other embodiments, an alkylene is a C_1-C_4 alkylene. Typical alkylene groups include, but are not

limited to, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, and the like. In some embodiments, an alkylene is $-\text{CH}_2-$.

[00147] An “alkoxy” group refers to a (alkyl)O- group, where alkyl is as defined herein.

[00148] The term “alkylamine” refers to the $-\text{N}(\text{alkyl})_x\text{H}_y$ group, where x is 0 and y is 2, or where x is 1 and y is 1, or where x is 2 and y is 0.

[00149] An “hydroxyalkyl” refers to an alkyl in which one hydrogen atom is replaced by a hydroxyl. In some embodiments, a hydroxyalkyl is a C_1 - C_4 hydroxyalkyl. Typical hydroxyalkyl groups include, but are not limited to, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$, and the like.

[00150] An “aminoalkyl” refers to an alkyl in which one hydrogen atom is replaced by an amino. In some embodiments, aminoalkyl is a C_1 - C_4 aminoalkyl. Typical aminoalkyl groups include, but are not limited to, $-\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$, and the like.

[00151] The term “alkenyl” refers to a type of alkyl group in which at least one carbon-carbon double bond is present. In one embodiment, an alkenyl group has the formula $-\text{C}(\text{R})=\text{CR}_2$, wherein R refers to the remaining portions of the alkenyl group, which may be the same or different. In some embodiments, R is H or an alkyl. In some embodiments, an alkenyl is selected from ethenyl (*i.e.*, vinyl), propenyl (*i.e.*, allyl), butenyl, pentenyl, pentadienyl, and the like. Non-limiting examples of an alkenyl group include $-\text{CH}=\text{CH}_2$, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{CH}=\text{CHCH}_3$, $-\text{C}(\text{CH}_3)=\text{CHCH}_3$, and $-\text{CH}_2\text{CH}=\text{CH}_2$.

[00152] The term “alkynyl” refers to a type of alkyl group in which at least one carbon-carbon triple bond is present. In one embodiment, an alkenyl group has the formula $-\text{C}\equiv\text{C}-\text{R}$, wherein R refers to the remaining portions of the alkynyl group. In some embodiments, R is H or an alkyl. In some embodiments, an alkynyl is selected from ethynyl, propynyl, butynyl, pentynyl, hexynyl, and the like. Non-limiting examples of an alkynyl group include $-\text{C}\equiv\text{CH}$, $-\text{C}\equiv\text{CCH}_3$, $-\text{C}\equiv\text{CCH}_2\text{CH}_3$, $-\text{CH}_2\text{C}\equiv\text{CH}$.

[00153] The term “heteroalkyl” refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, *e.g.*, oxygen, nitrogen (*e.g.* $-\text{NH}-$, $-\text{N}(\text{alkyl})-$, sulfur, or combinations thereof. A heteroalkyl is attached to the rest of the molecule at a carbon atom of the heteroalkyl. In one aspect, a heteroalkyl is a C_1 - C_6 heteroalkyl.

[00154] The term “aromatic” refers to a planar ring having a delocalized π -electron system containing $4n+2$ π electrons, where n is an integer. The term “aromatic” includes both carbocyclic aryl (“aryl”, *e.g.*, phenyl) and heterocyclic aryl (or “heteroaryl” or “heteroaromatic”)

groups (*e.g.*, pyridine). The term includes monocyclic or fused-ring polycyclic (*i.e.*, rings which share adjacent pairs of carbon atoms) groups.

[00155] The term “carbocyclic” or “carbocycle” refers to a ring or ring system where the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from “heterocyclic” rings or “heterocycles” in which the ring backbone contains at least one atom which is different from carbon. In some embodiments, a carbocycle is a monocyclic carbocycle or a bicyclic carbocycle. In some embodiments, a carbocycle is a monocyclic carbocycle. Carbocycles are non-aromatic or aromatic. Non-aromatic carbocycles are saturated or partially unsaturated. In some embodiments, a carbocycle is a bicyclic carbocycle. In some embodiments, at least one of the two rings of a bicyclic carbocycle is aromatic. In some embodiments, both rings of a bicyclic carbocycle are aromatic. Carbocycles include aryls and cycloalkyls.

[00156] As used herein, the term “aryl” refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. In one aspect, aryl is phenyl or a naphthyl. In some embodiments, an aryl is a phenyl. In some embodiments, an aryl is a phenyl, naphthyl, indanyl, indenyl, or tetrahydronaphthyl. In some embodiments, an aryl is a C₆-C₁₀aryl. Depending on the structure, an aryl group is a monoradical or a diradical (*i.e.*, an arylene group).

[00157] The term “cycloalkyl” refers to a monocyclic, bicyclic or polycyclic aliphatic, non-aromatic radical, wherein each of the atoms forming the ring (*i.e.* skeletal atoms) is a carbon atom. In some embodiments, cycloalkyls are spirocyclic or bridged compounds. In some embodiments, cycloalkyls are optionally fused with an aromatic ring, and the point of attachment is at a carbon that is not an aromatic ring carbon atom. Cycloalkyl groups include groups having from 3 to 10 ring atoms. In some embodiments, cycloalkyl groups are selected from among cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cyclooctyl, spiro[2.2]pentyl, norbornyl, norbornenyl, bicycle[1.1.1]pentyl, adamantyl, norbornyl, norbornenyl, decalinyl, or 7,7-dimethyl-bicyclo[2.2.1]heptanyl. In some embodiments, a cycloalkyl is a C₃-C₆cycloalkyl. In some embodiments, a cycloalkyl is a monocyclic cycloalkyl. Monocyclic cycloalkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyls include, for example, adamantyl, norbornyl (*i.e.*, bicyclo[2.2.1]heptanyl), norbornenyl, decalinyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like.

[00158] The term “halo” or, alternatively, “halogen” or “halide” means fluoro, chloro, bromo or iodo. In some embodiments, halo is fluoro, chloro, or bromo.

[00159] The term “fluoroalkyl” refers to an alkyl in which one or more hydrogen atoms are replaced by a fluorine atom. In one aspect, a fluoralkyl is a C₁-C₆fluoroalkyl.

[00160] The term “heterocycle” or “heterocyclic” refers to heteroaromatic rings (also known as heteroaryls) and heterocycloalkyl rings containing one to four heteroatoms in the ring(s), where each heteroatom in the ring(s) is selected from O, S and N, wherein each heterocyclic group has from 3 to 10 atoms in its ring system, and with the proviso that any ring does not contain two adjacent O or S atoms. Non-aromatic heterocyclic groups (also known as heterocycloalkyls) include rings having 3 to 10 atoms in its ring system and aromatic heterocyclic groups include rings having 5 to 10 atoms in its ring system. The heterocyclic groups include benzo-fused ring systems. Examples of non-aromatic heterocyclic groups are pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, oxazolidinonyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, thioxanyl, piperazinyl, aziridinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepinyl, diazepinyl, thiazepinyl, 1,2,3,6-tetrahydropyridinyl, pyrrolin-2-yl, pyrrolin-3-yl, indolinyl, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazolinyl, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl, indolin-2-onyl, isoindolin-1-onyl, isoindoline-1,3-dionyl, 3,4-dihydroisoquinolin-1(2H)-onyl, 3,4-dihydroquinolin-2(1H)-onyl, isoindoline-1,3-dithionyl, benzo[d]oxazol-2(3H)-onyl, 1H-benzo[d]imidazol-2(3H)-onyl, benzo[d]thiazol-2(3H)-onyl, and quinolizinyl. Examples of aromatic heterocyclic groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indolizinyl, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, and furopyridinyl. The foregoing groups are either C-attached (or C-linked) or N-attached where such is possible. For instance, a group derived from pyrrole includes both pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole includes imidazol-1-yl or imidazol-3-yl (both N-attached) or imidazol-2-yl, imidazol-4-yl or imidazol-5-yl (all C-attached). The heterocyclic groups include benzo-fused ring systems. Non-aromatic heterocycles are optionally substituted with one or two oxo (=O) moieties, such as pyrrolidin-2-one. In some embodiments, at least one of the two rings of a bicyclic heterocycle is aromatic. In some embodiments, both rings of a bicyclic heterocycle are aromatic.

[00161] The terms “heteroaryl” or, alternatively, “heteroaromatic” refers to an aryl group that includes one or more ring heteroatoms selected from nitrogen, oxygen and sulfur. Illustrative examples of heteroaryl groups include monocyclic heteroaryls and bicyclic heteroaryls. Monocyclic heteroaryls include pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, thiadiazolyl, and furazanyl. Monocyclic heteroaryls include indolizine, indole, benzofuran, benzothiophene, indazole, benzimidazole, purine, quinolizine, quinoline, isoquinoline, cinnoline, phthalazine, quinazoline, quinoxaline, 1,8-naphthyridine, and pteridine. In some embodiments, a heteroaryl contains 0-4 N atoms in the ring. In some embodiments, a heteroaryl contains 1-4 N atoms in the ring. In some embodiments, a heteroaryl contains 0-4 N atoms, 0-1 O atoms, and 0-1 S atoms in the ring. In some embodiments, a heteroaryl contains 1-4 N atoms, 0-1 O atoms, and 0-1 S atoms in the ring. In some embodiments, heteroaryl is a C₁-C₉heteroaryl. In some embodiments, monocyclic heteroaryl is a C₁-C₅heteroaryl. In some embodiments, monocyclic heteroaryl is a 5-membered or 6-membered heteroaryl. In some embodiments, bicyclic heteroaryl is a C₆-C₉heteroaryl.

[00162] A “heterocycloalkyl” group refers to a cycloalkyl group that includes at least one heteroatom selected from nitrogen, oxygen and sulfur. In some embodiments, a heterocycloalkyl is fused with an aryl or heteroaryl. In some embodiments, the heterocycloalkyl is oxazolidinonyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, morpholinyl, thiomorpholinyl, piperazinyl, piperidin-2-onyl, pyrrolidine-2,5-dithionyl, pyrrolidine-2,5-dionyl, pyrrolidinonyl, imidazolidinyl, imidazolidin-2-onyl, or thiazolidin-2-onyl. The term heterocycloalkyl also includes all ring forms of the carbohydrates, including but not limited to the monosaccharides, the disaccharides and the oligosaccharides. In one aspect, a heterocycloalkyl is a C₂-C₁₀heterocycloalkyl. In another aspect, a heterocycloalkyl is a C₄-C₁₀heterocycloalkyl. In some embodiments, a heterocycloalkyl contains 0-2 N atoms in the ring. In some embodiments, a heterocycloalkyl contains 0-2 N atoms, 0-2 O atoms and 0-1 S atoms in the ring.

[00163] The term “bond” or “single bond” refers to a chemical bond between two atoms, or two moieties when the atoms joined by the bond are considered to be part of larger substructure. In one aspect, when a group described herein is a bond, the referenced group is absent thereby allowing a bond to be formed between the remaining identified groups.

[00164] The term “moiety” refers to a specific segment or functional group of a molecule. Chemical moieties are often recognized chemical entities embedded in or appended to a molecule.

[00165] The term “optionally substituted” or “substituted” means that the referenced group is optionally substituted with one or more additional group(s) individually and independently selected from halogen, -CN, -NH₂, -NH(alkyl), -N(alkyl)₂, -OH, -CO₂H, -CO₂alkyl, -C(=O)NH₂, -C(=O)NH(alkyl), -C(=O)N(alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(alkyl), -S(=O)₂N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, fluoroalkoxy, heterocycloalkyl, aryl, heteroaryl, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, and arylsulfone. In some other embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(alkyl), -N(alkyl)₂, -OH, -CO₂H, -CO₂alkyl, -C(=O)NH₂, -C(=O)NH(alkyl), -C(=O)N(alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(alkyl), -S(=O)₂N(alkyl)₂, alkyl, cycloalkyl, fluoroalkyl, heteroalkyl, alkoxy, fluoroalkoxy, heterocycloalkyl, aryl, heteroaryl, aryloxy, alkylthio, arylthio, alkylsulfoxide, arylsulfoxide, alkylsulfone, and arylsulfone. In some other embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂(C₁-C₄alkyl), -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(C₁-C₄alkyl), -S(=O)₂N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, C₁-C₄fluoroalkoxy, -SC₁-C₄alkyl, -S(=O)C₁-C₄alkyl, and -S(=O)₂C₁-C₄alkyl. In some other embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -NH(CH₃), -N(CH₃)₂, -OH, -CO₂H, -CO₂(C₁-C₄alkyl), -C(=O)NH₂, -C(=O)NH(C₁-C₄alkyl), -C(=O)N(C₁-C₄alkyl)₂, -S(=O)₂NH₂, -S(=O)₂NH(C₁-C₄alkyl), -S(=O)₂N(C₁-C₄alkyl)₂, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₂-C₆heterocycloalkyl, C₁-C₄fluoroalkyl, C₁-C₄heteroalkyl, C₁-C₄alkoxy, C₁-C₄fluoroalkoxy, -SC₁-C₄alkyl, -S(=O)C₁-C₄alkyl, and -S(=O)₂C₁-C₄alkyl. In some embodiments, optional substituents are independently selected from halogen, -CN, -NH₂, -OH, -NH(CH₃), -N(CH₃)₂, -CH₃, -CH₂CH₃, -CF₃, -OCH₃, and -OCF₃. In some embodiments, substituted groups are substituted with one or two of the preceding groups. In some embodiments, an optional substituent on an aliphatic carbon atom (acyclic or cyclic) includes oxo (=O).

[00166] The term “acceptable” with respect to a formulation, composition or ingredient, as used herein, means having no persistent detrimental effect on the general health of the subject being treated.

[00167] The term “modulate” as used herein, means to interact with a target either directly or indirectly so as to alter the activity of the target, including, by way of example only, to enhance the activity of the target, to inhibit the activity of the target, to limit the activity of the target, or to extend the activity of the target.

[00168] The term “modulator” as used herein, refers to a molecule that interacts with a target either directly or indirectly. The interactions include, but are not limited to, the interactions of an

agonist, partial agonist, an inverse agonist, antagonist, degrader, or combinations thereof. In some embodiments, a modulator is an agonist.

[00169] The terms "administer," "administering," "administration," and the like, as used herein, refer to the methods that may be used to enable delivery of compounds or compositions to the desired site of biological action. These methods include, but are not limited to oral routes, intraduodenal routes, parenteral injection (including intravenous, subcutaneous, intraperitoneal, intramuscular, intravascular or infusion), topical and rectal administration. Those of skill in the art are familiar with administration techniques that can be employed with the compounds and methods described herein. In some embodiments, the compounds and compositions described herein are administered orally.

[00170] The terms "co-administration" or the like, as used herein, are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are administered by the same or different route of administration or at the same or different time.

[00171] The terms "effective amount" or "therapeutically effective amount," as used herein, refer to a sufficient amount of an agent or a compound being administered, which will relieve to some extent one or more of the symptoms of the disease or condition being treated. The result includes reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an "effective amount" for therapeutic uses is the amount of the composition comprising a compound as disclosed herein required to provide a clinically significant decrease in disease symptoms. An appropriate "effective" amount in any individual case is optionally determined using techniques, such as a dose escalation study.

[00172] The terms "enhance" or "enhancing," as used herein, means to increase or prolong either in potency or duration a desired effect. Thus, in regard to enhancing the effect of therapeutic agents, the term "enhancing" refers to the ability to increase or prolong, either in potency or duration, the effect of other therapeutic agents on a system. An "enhancing-effective amount," as used herein, refers to an amount adequate to enhance the effect of another therapeutic agent in a desired system.

[00173] The term "pharmaceutical combination" as used herein, means a product that results from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term "fixed combination" means that the active ingredients, e.g. a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term "non-fixed combination" means that the active ingredients, e.g. a compound of

Formula (I), or a pharmaceutically acceptable salt thereof, and a co-agent, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[00174] The terms “article of manufacture” and “kit” are used as synonyms.

[00175] The term “subject” or “patient” encompasses mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. In one aspect, the mammal is a human.

[00176] The terms “treat,” “treating” or “treatment,” as used herein, include alleviating, abating or ameliorating at least one symptom of a disease or condition, preventing additional symptoms, inhibiting the disease or condition, e.g., arresting the development of the disease or condition, relieving the disease or condition, causing regression of the disease or condition, relieving a condition caused by the disease or condition, or stopping the symptoms of the disease or condition either prophylactically and/or therapeutically.

Pharmaceutical compositions

[00177] In some embodiments, the compounds described herein are formulated into pharmaceutical compositions. Pharmaceutical compositions are formulated in a conventional manner using one or more pharmaceutically acceptable inactive ingredients that facilitate processing of the active compounds into preparations that are used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), herein incorporated by reference for such disclosure.

[00178] In some embodiments, the compounds described herein are administered either alone or in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition. Administration of the compounds and compositions described herein can be effected by any method that enables delivery of the compounds to the site of action. These methods include, though are not limited to delivery via enteral routes (including oral,

gastric or duodenal feeding tube, rectal suppository and rectal enema), parenteral routes (injection or infusion, including intraarterial, intracardiac, intradermal, intraduodenal, intramedullary, intramuscular, intraosseous, intraperitoneal, intrathecal, intravascular, intravenous, intravitreal, epidural and subcutaneous), inhalational, transdermal, transmucosal, sublingual, buccal and topical (including epicutaneous, dermal, enema, eye drops, ear drops, intranasal, vaginal) administration, although the most suitable route may depend upon for example the condition and disorder of the recipient. By way of example only, compounds described herein can be administered locally to the area in need of treatment, by for example, local infusion during surgery, topical application such as creams or ointments, injection, catheter, or implant. The administration can also be by direct injection at the site of a diseased tissue or organ.

[00179] In some embodiments, pharmaceutical compositions suitable for oral administration are presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. In some embodiments, the active ingredient is presented as a bolus, electuary or paste.

[00180] Pharmaceutical compositions which can be used orally include tablets, push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. Tablets may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with binders, inert diluents, or lubricating, surface active or dispersing agents. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. In some embodiments, the tablets are coated or scored and are formulated so as to provide slow or controlled release of the active ingredient therein. All formulations for oral administration should be in dosages suitable for such administration. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In some embodiments, stabilizers are added. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer

solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or Dragee coatings for identification or to characterize different combinations of active compound doses.

[00181] In some embodiments, pharmaceutical compositions are formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. The compositions may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in powder form or in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, saline or sterile pyrogen-free water, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

[00182] Pharmaceutical compositions for parenteral administration include aqueous and non-aqueous (oily) sterile injection solutions of the active compounds which may contain antioxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[00183] Pharmaceutical compositions may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[00184] For buccal or sublingual administration, the compositions may take the form of tablets, lozenges, pastilles, or gels formulated in conventional manner. Such compositions may comprise the active ingredient in a flavored basis such as sucrose and acacia or tragacanth.

[00185] Pharmaceutical compositions may be administered topically, that is by non-systemic administration. This includes the application of a compound of the present invention externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

[00186] Pharmaceutical compositions suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as gels, liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the formulation.

[00187] Pharmaceutical compositions for administration by inhalation are conveniently delivered from an insufflator, nebulizer pressurized packs or other convenient means of delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Alternatively, for administration by inhalation or insufflation, pharmaceutical preparations may take the form of a dry powder composition, for example a powder mix of the compound and a suitable powder base such as lactose or starch. The powder composition may be presented in unit dosage form, in for example, capsules, cartridges, gelatin or blister packs from which the powder may be administered with the aid of an inhalator or insufflator.

[00188] It should be understood that in addition to the ingredients particularly mentioned above, the compounds and compositions described herein may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

Methods of Dosing and Treatment Regimens

[00189] In one embodiment, the compounds of Formula (I), or a pharmaceutically acceptable salt thereof, are used in the preparation of medicaments for the treatment of diseases or conditions in a mammal that would benefit from modulation of melanocortin receptor activity. Methods for treating any of the diseases or conditions described herein in a mammal in need of such treatment, involves administration of pharmaceutical compositions that include at least one compound of Formula (I) or a pharmaceutically acceptable salt, active metabolite, prodrug, or pharmaceutically acceptable solvate thereof, in therapeutically effective amounts to said mammal.

[00190] In certain embodiments, the compositions containing the compound(s) described herein are administered for prophylactic and/or therapeutic treatments. In certain therapeutic applications, the compositions are administered to a patient already suffering from a disease or condition, in an amount sufficient to cure or at least partially arrest at least one of the symptoms of the disease or condition. Amounts effective for this use depend on the severity and course of the disease or condition, previous therapy, the patient's health status, weight, and response to the drugs, and the judgment of the treating physician. Therapeutically effective amounts are optionally determined by methods including, but not limited to, a dose escalation and/or dose ranging clinical trial.

[00191] In prophylactic applications, compositions containing the compounds described herein are administered to a patient susceptible to or otherwise at risk of a particular disease, disorder or condition. Such an amount is defined to be a "prophylactically effective amount or dose." In this use, the precise amounts also depend on the patient's state of health, weight, and the like. When used in patients, effective amounts for this use will depend on the severity and course of the disease, disorder or condition, previous therapy, the patient's health status and response to the drugs, and the judgment of the treating physician. In one aspect, prophylactic treatments include administering to a mammal, who previously experienced at least one symptom of the disease being treated and is currently in remission, a pharmaceutical composition comprising a compound of Formula (I), or a pharmaceutically acceptable salt thereof, in order to prevent a return of the symptoms of the disease or condition.

[00192] In certain embodiments wherein the patient's condition does not improve, upon the doctor's discretion the administration of the compounds are administered chronically, that is, for an extended period of time, including throughout the duration of the patient's life in order to ameliorate or otherwise control or limit the symptoms of the patient's disease or condition.

[00193] Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, in specific embodiments, the dosage or the frequency of administration, or both, is reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In certain embodiments, however, the patient requires intermittent treatment on a long-term basis upon any recurrence of symptoms.

[00194] The amount of a given agent that corresponds to such an amount varies depending upon factors such as the particular compound, disease condition and its severity, the identity (*e.g.*, weight, sex) of the subject or host in need of treatment, but nevertheless is determined according to the particular circumstances surrounding the case, including, *e.g.*, the specific agent being

administered, the route of administration, the condition being treated, and the subject or host being treated.

[00195] In general, however, doses employed for adult human treatment are typically in the range of 0.01 mg-2000 mg per day. In one embodiment, the desired dose is conveniently presented in a single dose or in divided doses administered simultaneously or at appropriate intervals, for example as two, three, four or more sub-doses per day.

[00196] In one embodiment, the daily dosages appropriate for the compound of Formula (I), or a pharmaceutically acceptable salt thereof, described herein are from about 0.01 to about 50 mg/kg per body weight. In some embodiments, the daily dosage or the amount of active in the dosage form are lower or higher than the ranges indicated herein, based on a number of variables in regard to an individual treatment regime. In various embodiments, the daily and unit dosages are altered depending on a number of variables including, but not limited to, the activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[00197] Toxicity and therapeutic efficacy of such therapeutic regimens are determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD₅₀ and the ED₅₀. The dose ratio between the toxic and therapeutic effects is the therapeutic index and it is expressed as the ratio between LD₅₀ and ED₅₀. In certain embodiments, the data obtained from cell culture assays and animal studies are used in formulating the therapeutically effective daily dosage range and/or the therapeutically effective unit dosage amount for use in mammals, including humans. In some embodiments, the daily dosage amount of the compounds described herein lies within a range of circulating concentrations that include the ED₅₀ with minimal toxicity. In certain embodiments, the daily dosage range and/or the unit dosage amount varies within this range depending upon the dosage form employed and the route of administration utilized.

[00198] In any of the aforementioned aspects are further embodiments in which the effective amount of the compound of Formula (I), or a pharmaceutically acceptable salt thereof, is: (a) systemically administered to the mammal; and/or (b) administered orally to the mammal; and/or (c) intravenously administered to the mammal; and/or (d) administered by injection to the mammal; and/or (e) administered topically to the mammal; and/or (f) administered non-systemically or locally to the mammal.

[00199] In any of the aforementioned aspects are further embodiments comprising single administrations of the effective amount of the compound, including further embodiments in

which (i) the compound is administered once a day; or (ii) the compound is administered to the mammal multiple times over the span of one day.

[00200] In any of the aforementioned aspects are further embodiments comprising multiple administrations of the effective amount of the compound, including further embodiments in which (i) the compound is administered continuously or intermittently: as in a single dose; (ii) the time between multiple administrations is every 6 hours; (iii) the compound is administered to the mammal every 8 hours; (iv) the compound is administered to the mammal every 12 hours; (v) the compound is administered to the mammal every 24 hours. In further or alternative embodiments, the method comprises a drug holiday, wherein the administration of the compound is temporarily suspended or the dose of the compound being administered is temporarily reduced; at the end of the drug holiday, dosing of the compound is resumed. In one embodiment, the length of the drug holiday varies from 2 days to 1 year.

Combination Treatments

[00201] In certain instances, it is appropriate to administer at least one compound of Formula (I), or a pharmaceutically acceptable salt thereof, in combination with one or more other therapeutic agents.

[00202] In one embodiment, the therapeutic effectiveness of one of the compounds described herein is enhanced by administration of an adjuvant (*i.e.*, by itself the adjuvant has minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced). Or, in some embodiments, the benefit experienced by a patient is increased by administering one of the compounds described herein with another agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

[00203] In one specific embodiment, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is co-administered with a second therapeutic agent, wherein the compound of Formula (I), or a pharmaceutically acceptable salt thereof, and the second therapeutic agent modulate different aspects of the disease, disorder or condition being treated, thereby providing a greater overall benefit than administration of either therapeutic agent alone.

[00204] In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient is simply be additive of the two therapeutic agents or the patient experiences a synergistic benefit.

[00205] For combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated and so forth. In additional embodiments, when co-administered with

one or more other therapeutic agents, the compound provided herein is administered either simultaneously with the one or more other therapeutic agents, or sequentially.

[00206] In combination therapies, the multiple therapeutic agents (one of which is one of the compounds described herein) are administered in any order or even simultaneously. If administration is simultaneous, the multiple therapeutic agents are, by way of example only, provided in a single, unified form, or in multiple forms (e.g., as a single pill or as two separate pills).

[00207] The compounds of Formula (I), or a pharmaceutically acceptable salt thereof, as well as combination therapies, are administered before, during or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound varies. Thus, in one embodiment, the compounds described herein are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. In another embodiment, the compounds and compositions are administered to a subject during or as soon as possible after the onset of the symptoms. In specific embodiments, a compound described herein is administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease. In some embodiments, the length required for treatment varies, and the treatment length is adjusted to suit the specific needs of each subject.

[00208] **Abbreviations:**

DIEA: N,N-diisopropylethylamine;

DMSO: dimethyl sulfoxide;

Pd(PPh₃)₄: tetrakis(triphenylphosphine)palladium(0)

Pd(dppf)Cl₂: [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride;

CuI: copper(I) iodide;

TBAF: tetra-n-butylammonium fluoride;

P(t-Bu)₃: tri-tert-butylphosphine;

HBF₄: tetrafluoroboric acid;

DBU: 1,8-diazabicyclo[5.4.0]undec-7-ene;

Prep-HPLC: preparative high performance liquid chromatography;

TFA: trifluoroacetic acid;

CH₃CN, MeCN, or ACN: acetonitrile;

MeOD: deuterated methanol;

CDCl₃: deuterated chloroform


DME: 1,2-dimethoxyethane;
DMF: N,N-dimethylformamide
DCM: dichloromethane
H₂O: water;
KOAc: potassium acetate;
NaOAc: sodium acetate;
Cs₂CO₃: cesium carbonate
P-TsOH: p-toluenesulfonic acid;
NaNO₂: sodium nitrite;
THF: tetrahydrofuran;
NBS: N-bromosuccinimide;
Br₂: bromine;
AgF: silver fluoride;
LiAlH₄: lithium aluminium hydride;
IBX: 2-iodoxybenzoic acid;
TEA: trimethylamine;
DMAP: N,N-dimethylpyridine-4-amine
HOBT: hydroxybenzotriazole;
EDCI: 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide;
Pd(PPh₃)₂Cl₂: bis(triphenylphosphine)palladium(II) dichloride;
PdAMphos or Pd (amphos)Cl₂ or : bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II);
Pd(DTBPF)Cl₂: [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II);
cc: column chromatography
rt: room temperature;
h: hour or hours.

EXAMPLES

[00209] The following examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein.

Synthesis of Compounds

Example 1: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide (compound 1-1)



[00210] Step 1-1, preparation of tert-butyl (3R)-3-ethyl-4-[2-(methoxycarbonyl)-4-nitrophenyl]piperazine-1-carboxylate: into a 40-mL round bottom flask, was placed methyl 2-fluoro-5-nitrobenzoate (2.2 g, 0.01 mmol, 1.2 eq.), tert-butyl (3R)-3-ethylpiperazine-1-carboxylate (2.0 g, 9.33 mmol), TEA (2.8 g, 0.03 mmol, 3.0 eq.) and DMSO (20 mL). The resulting solution was stirred at 100 °C for 2 h and cooled to rt. The reaction was diluted with water (50 mL) and extracted with ethyl acetate (3x50 mL). The combined organic layers were washed with brine (2x50 mL), dried over anhydrous sodium sulfate, and concentrated under vacuum. The residue was purified by a silica gel column eluting with ethyl acetate/petroleum ether (1:3). 3.3 g (90 %) of the title compound as a brown oil. LCMS (M+H)⁺: 394.2.

[00211] Step 1-2, preparation of methyl 2-[(2R)-2-ethylpiperazin-1-yl]-5-nitrobenzoate: a solution of tert-butyl (3R)-3-ethyl-4-[2-(methoxycarbonyl)-4-nitrophenyl]piperazine-1-carboxylate (3.3 g, 8.39 mmol) in TFA (3 mL)/DCM (6 mL) was stirred at 30 °C for 2 hrs. The resulting mixture was concentrated under vacuum. This resulted in 3.3 g (97%) of the title compound as a brown oil. LCMS (M+H)⁺: 294.1.

[00212] Step 1-3, preparation of tert-butyl (3R)-4-[4-bromo-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate: 2-chloro-4-(trifluoromethyl)benzoic acid (860 mg, 3.83 mmol, 1.2 eq.) and HATU (1.46 g, 3.83 mmol, 1.2 eq.) were dissolved in DMF (20 mL). After stirring for 5 min at rt, the resulting solution was treated with methyl 2-[(2R)-2-ethylpiperazin-1-yl]-5-nitrobenzoate trifluoroacetate (1.3 g, 3.19 mmol) and DIEA (1.23 g, 9.57 mmol, 3.0 eq.). The reaction was stirred for 1 h at room temperature, diluted with water (50 mL), and extracted with ethyl acetate (3x50 mL). The combined organic layers were washed with brine (2x50 mL), dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified by a silica gel column eluting with ethyl acetate/petroleum ether (1:1). 1.8 g (83%) of the title compound was obtained as a yellow oil. LCMS (M+H)⁺: 500.1.

[00213] Step 1-4, preparation of 5-amino-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]benzoate: into a 40-mL round-bottom flask, was placed methyl 2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-nitrobenzoate (1.8 g, 3.60 mmol), Fe

(1 g, 18.00 mmol, 5.0 eq.), NH₄Cl (963 mg, 18.00 mmol, 5.0 eq.), and EtOH (20 mL)/H₂O (4 mL). The resulting solution was stirred at 70 °C for 1 h and cooled to rt. The reaction was diluted with water (50 mL) and extracted with ethyl acetate (3x60 mL). The combined organic layers were dried and concentrated under vacuum. The residue was purified by a silica gel column eluting with ethyl acetate/petroleum ether (1:1). 1.1 g (65 %) of the title compound was isolated as a yellow oil. LCMS (M+H)⁺: 470.1.

[00214] Step 1-5, preparation of methyl 5-bromo-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]benzoate: into a 40-mL round-bottom flask, was placed CuBr₂ (1.31 g, 5.85 mmol, 2.5 eq.), n-butyl nitrite (482 mg, 4.68 mmol, 2.0 eq.), and ACN (20 mL). The resulting solution was stirred for 5 min at 0 °C and treated with a solution of methyl 5-amino-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]benzoate (1.1 g, 2.34 mmol) in ACN (5 mL) dropwise at 0 °C. The resulting solution was stirred at 50 °C for 1 h. The reaction mixture was cooled to rt and diluted with water (50 mL). The resulting solution was extracted with ethyl acetate (3x50 mL). The combined organic layers were dried and concentrated under vacuum. The residue was purified by a silica gel column eluting with ethyl acetate/petroleum ether (1:1). 520 mg (42 %) of the title compound was isolated as a light yellow oil. LCMS (M+H)⁺: 533.0.

[00215] Step 1-6, preparation of methyl 4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxylate: Into a 8-mL pressure tube vessel, was placed methyl 5-bromo-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]benzoate (30 mg, 0.06 mmol), 2,3-difluorophenylboronic acid (13.3 mg, 0.08 mmol), Pd₂(dba)₃CHCl₃ (5.8 mg, 0.01 mmol), P(t-Bu)₃.HBF₄ (1.6 mg, 0.01 mmol), K₃PO₄ (35.8 mg, 0.17 mmol), and toluene/H₂O (1 mL/0.1 mL). The resulting solution was degassed for 5 min with N₂, sealed, and stirred at 70 °C for 2 hrs in an oil bath. The resulting mixture was cooled to rt and concentrated under vacuum. The residue was applied onto a silica gel column eluting with ethyl acetate/petroleum ether (1:3), resulting in 20 mg (63%) of the title compound. LCMS (M+H)⁺: 567.1.


[00216] Step 1-7, preparation of N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide: a 8-mL pressure tube vessel, was placed methyl 4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxylate (20 mg, 0.04 mmol), ethane-1,2-diamine (0.5 mL), and IPA (0.5 mL). The resulting solution was sealed and stirred at 70 °C for 16 hrs. The resulting mixture was cooled to rt and concentrated under vacuum. The residue was purified by Prep-HPLC with the following conditions: SunFire

Prep C18 OBD column; mobile phase, phase A: water (0.05% TFA); phase B: ACN (30% up to 45% in 6 min); flow rate: 20 mL/min; detector, 220 & 254 nm, resulting in 7.4 mg (30%) of the title compound. LCMS (M+H)⁺: 595.2.

[00217] The following compounds were prepared similarly to **Example 1** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings.

Compound no.	MS (M+H) ⁺
1-2	589.1
1-3	603.1
1-4	577.2
1-5	593.2
1-6	593.2
1-7	584.2
1-8	573.2
1-9	611.1
1-10	603.2
1-11	617.2
1-12	617.3
1-13	575.2
1-16	603.2
1-22	588.2
1-23	617.3
1-28	601.2
1-29	601.2
1-30	560.2
1-31	560.2
1-34	604.3

Example 2: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxamide (compound 1-14)



[00218] Step 2-1, preparation of methyl 2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate: Into a 8-mL

pressure tube vessel purged and maintained with an inert atmosphere of nitrogen, was placed methyl 5-bromo-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]benzoate from **Example 1**, Step 1-5 (250 mg, 0.47 mmol), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (178 mg, 0.70 mmol), Pd(DTBPF)Cl₂ (25 mg, 0.04 mmol), KOAc (138 mg, 1.4 mmol), and dioxane (5 mL). The resulting solution was degassed for 10 min with N₂, sealed, and stirred at 90 °C for 3 h in an oil bath. The resulting mixture was cooled to rt and concentrated under vacuum. The residue was applied onto a silica gel column eluting with ethyl acetate/petroleum ether (1:3). This resulted in 120 mg (44%) of the title compound. LCMS (M+H)⁺: 581.2.

[00219] Step 2-2, preparation of methyl 4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxylate: to a 8-mL pressure tube vessel, was placed 2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (43 mg, 0.07 mmol), 1-bromo-2-cyclopropoxybenzene (28 mg, 0.13 mmol), Pd(DTBPF)Cl₂ (2 mg, 0.003 mmol), K₂CO₃ (20 mg, 0.14 mmol), and dioxane/H₂O (2 mL/0.2 mL). The resulting solution was degassed for 10 min with N₂, sealed, and stirred at 90 °C for 2 h in an oil bath. The reaction mixture was cooled to rt and concentrated under vacuum. The residue was purified by a prep-TLC with ethyl acetate/petroleum ether (1:3), resulting in 25 mg (57%) of the title compound. LCMS (M+H)⁺: 587.0.


[00220] Step 2-3, preparation of N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxamide: to a 8-mL pressure tube vessel, was placed methyl 4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxylate (25 mg, 0.04 mmol), ethane-1,2-diamine (2 mL), and IPA (2 mL). The resulting solution was sealed and stirred at 70 °C for 16 h. The resulting mixture was cooled to rt and concentrated under vacuum. The residue was purified by Prep-HPLC with the following conditions: SunFire Prep C18 OBD column; mobile phase, phase A: water (0.05% TFA); phase B: ACN (21% up to 38% in 6 min); flow rate: 20 mL/min; detector, 220 & 254 nm, yielding 10.7 mg (34%) of the title compound. LCMS (M+H)⁺: 615.2.

[00221] The following compounds were prepared similarly to **Example 2** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-32	604.2
1-33	604.2

Compound no.	MS (M+H) ⁺
1-35	604.2

Example 3: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide (compound 1-95)



[00222] Step 3-1, preparation of *tert*-butyl (3*R*)-3-ethyl-4-[3-fluoro-2-(methoxycarbonyl)-4-nitrophenyl]piperazine-1-carboxylate: to a solution of *tert*-butyl-(3*R*)-3-ethylpiperazine-1-carboxylate (555 mg, 2.59 mmol) and methyl 2,6-difluoro-3-nitrobenzoate (843 mg, 3.89 mmol) in DMSO (2 mL) was added DIEA (0.90 mL, 5.2 mmol). The mixture was heated at 130°C for 1 h. The mixture was purified by C18 reversed phase column chromatography to give the title compound (813 mg, 76 % yield) as a brownish yellow gum. LCMS (M+H)⁺: 412.3.

[00223] Step 3-2, preparation of *tert*-butyl (3*R*)-4-[4-amino-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate: to a solution of *tert*-butyl (3*R*)-3-ethyl-4-[3-fluoro-2-(methoxycarbonyl)-4-nitrophenyl]piperazine-1-carboxylate (813 mg, 1.98 mmol) in MeOH (30 mL) was added 10% Pd/C (213 mg, 0.0941 mmol) and NH₄Cl (1.188 g, 18.83 mmol). The mixture was heated at 80°C for 20 min. The mixture was filtered through Celite and the filtrate was concentrated and purified by C18 reversed phase column chromatography to give the title compound (612 mg, 81% yield) as a reddish brown gum. LCMS (M+H)⁺: 382.3.

[00224] Step 3-3, preparation of *tert*-butyl (3*R*)-4-[4-bromo-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate: to a solution of *tert*-butyl (3*R*)-4-[4-amino-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate (612 mg, 1.60 mmol) in MeCN (10 mL) under nitrogen was added CuBr₂ (197 mg, 0.882 mmol) and 90% *tert*-butyl nitrite (0.24 mL, 1.8 mmol). The mixture was heated at 70°C for 30 min. The mixture was quenched with ice water and extracted with DCM (3X). The combined organic layers were washed with saturated NaHCO₃ (aq), concentrated, and purified by C18 reversed phase column chromatography to give the title compound (267 mg, 0.600 mmol, 38% yield) as a brown gum. LCMS (M+H)⁺: 445.2.

[00225] Step 3-4, preparation of *tert*-butyl (3*R*)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate: to a mixture of *tert*-butyl (3*R*)-4-[4-bromo-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate (267 mg, 0.600 mmol), (2-ethoxypyridin-3-yl)boronic acid (150 mg, 0.900 mmol), Pd[*t*-Bu₂P(4-NMe₂C₆H₄)₂Cl₂] (42.5 mg, 0.0600 mmol), and K₂CO₃ (249 mg, 1.80 mmol) in a sealed tube was added dioxane (4 mL) and H₂O (0.4 mL). The resulting solution was degassed with N₂ (g) for 10 min, sealed, and stirred at 100°C for 30 min. The reaction was treated with additional (2-ethoxypyridin-3-yl)boronic acid (37.8 mg, 0.226 mmol), Pd[*t*-Bu₂P(4-NMe₂C₆H₄)₂Cl₂] (13.4 mg, 0.0189 mmol), and K₂CO₃ (78.3 mg, 0.567 mmol) and stirred at 100°C for additional 30 min. The mixture was concentrated and purified by C18 reversed phase column chromatography to give the title compound (255 mg, 87% yield) as a brown gum. LCMS (M+H)⁺: 488.4.

[00226] Step 3-5, preparation of 6-[(2*R*)-4-[(*tert*-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid: to a solution of *tert*-butyl (3*R*)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate (255 mg, 0.523 mmol) in THF/MeOH/H₂O (4.5/1.5/1.5 mL) was added LiOH·H₂O (220 mg, 5.24 mmol). The mixture was heated at 60°C overnight. About 50% of the reaction was completed. After addition of additional LiOH·H₂O (220 mg, 5.24 mmol), the mixture was continued to heat at 70°C for 2 days. After removal of volatile solvent, the aqueous residue was diluted with ice water and acidified with 1N HCl (aq) to pH 4. The solid was collected by vacuum filtration and dried to give the title compound (239 mg, 97% yield) as a brown solid. LCMS (M+H)⁺: 474.2.

[00227] Step 3-6, preparation of 3-(2-ethoxypyridin-3-yl)-6-[(2*R*)-2-ethylpiperazin-1-yl]-2-fluorobenzoic acid dihydrochloride: to a solution of 6-[(2*R*)-4-[(*tert*-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid (100 mg, 0.211 mmol) in DCM (0.3 mL) was added 4N HCl in dioxane (0.30 mL, 1.2 mmol). The mixture was stirred at rt for 1 hr. The solution was decanted and the residue was rinsed with DCM (3X) and dried under vacuum to give the title compound as a medium brown solid. LCMS (M+H)⁺: 374.3.

[00228] Step 3-7, preparation of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid: to a solution of 4-chloro-2-(trifluoromethyl)benzoic acid (94.9 mg, 0.423 mmol) and HATU (145 mg, 0.381 mmol) in DMF (0.5 mL) was added DIEA (0.073 mL, 0.42 mmol). After stirring at rt for 5 min, the resulting solution was added to a solution of 3-(2-ethoxypyridin-3-yl)-6-[(2*R*)-2-ethylpiperazin-1-yl]-2-fluorobenzoic acid dihydrochloride (0.211 mmol) and DIEA (0.11 mL, 0.42 mmol) in DMF (0.2 mL). The mixture was stirred at RT for 10 min. The reaction was directly purified by C18

reversed phase column chromatography to give the title compound (69.8 mg, 57% yield) as a white solid. LCMS (M+H)⁺: 580.3.

[00229] Step 3-8, preparation of *tert*-butyl *N*-[2-({6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}formamido)ethyl]-*N*-methylcarbamate: to a solution of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid (35.0 mg, 0.06 mmol) and HATU (34.4 mg, 0.091 mmol) in DMF (0.2 mL) was added DIEA (0.021 mL, 0.12 mmol). After stirring at rt for 5 min, the reaction was treated with *tert*-butyl *N*-(2-aminoethyl)-*N*-methylcarbamate (21.0 mg, 0.121 mmol) and stirred at rt for 5 min. The mixture was purified by C18 reversed phase column chromatography to give the title compound (35.6 mg, 80% yield) as an off-white solid. LCMS (M+H)⁺: 736.4.


[00230] Step 3-9, preparation of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-*N*-[2-(methylamino)ethyl]benzamide: To a solution of *tert*-butyl *N*-[2-({6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}formamido)ethyl]-*N*-methylcarbamate (35.6 mg, 0.0484 mmol) in DCM (0.1 mL) was added 4N HCl in dioxane (0.10 mL, 0.40 mmol). The mixture was stirred at rt for 10 min. The solution was decanted and the residue was rinsed with DCM (3X) and dried under vacuum. The HCl salt of product was triturated with saturated NaHCO₃ (aq) and dried to give the title compound (24.6 mg, 80% yield) as an off-white solid. LCMS (M+H)⁺: 636.3.

[00231] The following compounds were prepared similarly to **Example 3** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings. In some examples, the final deprotecting step may not be required:

Compound no.	MS (M+H) ⁺
1-15	569.2
1-19	617.2
1-20	616.2
1-21	630.2
1-25	603.3
1-45	630.4
1-46	618.3
1-48	644.5
1-49	658.5
1-69	622.3
1-92	630.5
1-97	630.5

Compound no.	MS (M+H) ⁺
1-98	644.5
1-99	644.5
1-100	644.4
1-102	630.6
1-108	644.4
1-109	660.2
1-112	648.3
1-123	648.4
1-124	662.3
1-138	660.3
1-198	634.3
1-199	662.3
1-200	648.4
1-201	662.3
1-202	630.4
1-203	644.5
1-204	648.5
3-5	604.2
3-6	590.3
1-216	630.5
1-217	659.2

Example 4: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide (compound 1-173)



[00232] Step 4-1, preparation of *tert*-butyl (3R)-4-(2-{[2-(dimethylamino)ethyl]carbamoyl}-4-(2-ethoxypyridin-3-yl)-3-fluorophenyl)-3-ethylpiperazine-1-carboxylate: to a solution of 6-[(2R)-4-[(*tert*-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid from **Example 3**, step 3-5 (1.000 g, 2.111 mmol) and HATU (963 mg, 2.53 mmol) in DMF (1 mL) was added DIEA (0.92 mL, 5.3 mmol). After stirring at RT for 5 min, the resulting solution was treated with (2-aminoethyl)dimethylamine (0.35 mL, 3.1 mmol) and stirred at rt for 10 min. The reaction was purified by C18 reversed phase column chromatography to give the title compound (973.7 mg, 85% yield) as a light yellow solid. LCMS (M+H)⁺: 544.4.

[00233] Step 4-2, preparation of *N*-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethylpiperazin-1-yl]-2-fluorobenzamide trihydrochloride: to a solution of *tert*-butyl (3R)-4-(2-{[2-(dimethylamino)ethyl]carbamoyl}-4-(2-ethoxypyridin-3-yl)-3-fluorophenyl)-3-

ethylpiperazine-1-carboxylate (973.7 mg, 1.791 mmol) in DCM (2 mL) was added 4N HCl in dioxane (2.3 mL, 9.2 mmol). The mixture was stirred at rt for 2 hrs. The reaction was treated with additional 4N HCl in dioxane (2.0 mL, 8.0 mmol) and continued to stir for 2 hrs. The solution was decanted and the residue was rinsed with DCM (3X) and dried under vacuum to give the title compound as a light yellow solid. LCMS (M+H)⁺: 444.3.

[00234] Step 4-3, preparation of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-*N*-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide: to a solution of 4-chloro-2-(trifluoromethyl)benzoic acid (40.6 mg, 0.181 mmol) and HATU (61.9 mg, 0.163 mmol) in DMF (0.2 mL) was added DIEA (0.047 mL, 0.27 mmol). After stirring at rt for 5 min, the resulting solution was added to a solution of *N*-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2*R*)-2-ethylpiperazin-1-yl]-2-fluorobenzamide trihydrochloride (40.0 mg, 0.0723 mmol) and DIEA (0.110 mL, 0.631 mmol) in DMF (0.2 mL). After stirring at rt for 10 min, the reaction was purified by C18 reversed phase column chromatography to give the title compound (35.3 mg, 75% yield) as an off-white solid. LCMS (M+H)⁺: 650.3.


[00235] The following compounds were prepared similarly to **Example 4** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings. In some examples, an additional deprotection is required in the final step:

Compound no.	MS (M+H) ⁺
1-37	604.4
1-38	554.2
1-39	597.3
1-42	570.2
1-43	550.2
1-47	632.4
1-53	571.2
1-54	566.3
1-55	561.2
1-58	588.3
1-59	584.0
1-60	530.3
1-61	605.3
1-62	570.4
1-63	542.4
1-71	590.3
1-72	648.3
1-73	598.2
1-74	603.4
1-76	614.4

Compound no.	MS (M+H) ⁺
1-77	531.5
1-78	534.3
1-79	550.3
1-80	546.3
1-81	558.4
1-82	580.3
1-86	531.4
1-87	550.3
1-88	556.4
1-89	555.5
1-91	605.4
1-96	586.2
1-105	656.4
1-106	614.4
1-107	613.4
1-110	594.3
1-111	632.3
1-113	582.2
1-114	620.4
1-117	561.3
1-118	589.4
1-119	600.2
1-120	640.4
1-121	602.5
1-122	614.4
1-126	586.3
1-127	614.3
1-129	614.3
1-130	613.3
1-131	614.3
1-133	597.4
1-134	631.5
1-142	587.3
1-143	626.5
1-144	621.5
1-145	586.3
1-146	614.3
1-147	591.4
1-153	593.2
1-154	627.2
1-155	632.4
1-156	602.4
1-157	627.4
1-159	601.3
1-160	560.4
1-161	603.3
1-162	617.4
1-163	632.3

Compound no.	MS (M+H) ⁺
1-164	633.3
1-165	647.3
1-166	633.3
1-168	646.4
1-174	634.4
1-175	641.4
1-179	616.3
1-186	607.1
1-191	605.3
1-192	585.2
1-193	571.2
1-228	644.5
1-229	662.2
1-270	659.4
1-271	645.5

Example 5: 6-[(2R)-4- [4-chloro- 2-(difluoromethyl)benzoyl]- 2-ethylpiperazin- 1-yl]-N-[2-(dimethylamino)ethyl]-3- (2-ethoxypyridin- 3-yl)- 2-fluorobenzamide (compound 1-184)



[00236] Step 5-1, preparation of 4-chloro-2-(difluoromethyl)benzoic acid: to a mixture of 4-chloro-2-methylbenzoic acid (0.460 g, 2.70 mmol), Selectfluor (2.697 g, 7.612 mmol) and Na₂S₂O₈ (0.365 g, 1.53 mmol) in a sealed tube was added MeCN (2 mL) and water (2 mL). N₂ (g) was bubbled and then the mixture was cooled to -78°C under nitrogen and AgNO₃ (51.8 mg, 0.305 mmol) was added. After addition, the mixture was warmed to rt under nitrogen and then heated at 80°C for 3 hrs, filtered through Celite and rinsed with EtOAc. The filtrate was washed with saturated NaHCO₃ (aq) (3 X 25 mL). The combined basic aqueous solution was acidified with 1N HCl (aq) to pH 2-3 and extracted with EtOAc (2X). The combined organic layers were concentrated and purified by C18 reversed phase column chromatography to give the title compound (128.8 mg, 23% yield) as an off-white solid.


[00237] Step 5-2, preparation of 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide: to a solution of 4-chloro-2-(difluoromethyl)benzoic acid (37.3 mg, 0.181 mmol) and HATU (61.9 mg, 0.163 mmol) in DMF (0.2 mL) was added DIEA (0.047 mL, 0.27 mmol). After stirring at rt for 5 min, the resulting solution was added to a solution of N-[2-(dimethylamino)ethyl]-3-(2-

ethoxypyridin-3-yl)-6-[(2*R*)-2-ethylpiperazin-1-yl]-2-fluorobenzamide trihydrochloride from **Example 4**, step 4-2 (43.0 mg, 0.0778 mmol) and DIEA (0.110 mL, 0.631 mmol) in DMF (0.2 mL). The mixture was stirred at rt for 10 min. The mixture was purified by C18 reversed phase column chromatography to give the title compound (20.6 mg, 0.0326 mmol, 41.9% yield) as an off-white solid. LCMS (M+H)⁺: 632.3.

[00238] The following compounds were prepared similarly to **Example 5** with appropriate substituting reagents and substrates at different steps. In some examples, an additional deprotection is required in the final step:

Compound no.	MS (M+H) ⁺
1-128	600.4
1-176	616.5
1-177	602.5
1-183	618.6
1-189	614.4
1-190	630.6
1-210	615.4
1-213	601.4

Example 6: 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1*H*-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide (compound 1-188)




[00239] Step 6-1, preparation of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1*H*-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide: to a solution of 6-[(2*R*)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzoic acid from **Example 3**, step 3-7 (48 mg, 0.082 mmol) in ACN (1 mL) was added HATU (31 mg, 0.082 mmol) and Et₃N (17 μ L, 0.12 mmol). After stirring for 5 min at rt, the resulting solution was treated with (4,5-dihydro-1*H*-imidazol-2-yl)methylamine dihydrochloride (14 mg, 0.081 mmol) which was pre-neutralized with Et₃N (34 μ L, 0.24 mmol) in ACN (1 mL). The reaction was stirred at rt for 20 min, diluted with DCM, and washed with H₂O and brine. The organic layer was dried with anhydrous Na₂SO₄ and concentrated. The residue was purified from a silica gel column chromatography eluting

with DCM/MeOH (100/0 to 85/15) to afford the title compound (21 mg, 39 %). LCMS (M+H)⁺: 661.2.

[00240] The following compounds were prepared similarly to **Example 6** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-64	652.4
1-65	652.4
1-66	652.4
1-70	638.2
1-83	641.4
1-84	655.6
1-93	655.1
1-116	673.4
1-125	643.4
1-132	641.3
1-139	656.6
1-140	630.6
1-141	649.5
1-170	657.5
1-178	627.3
1-182	645.4
1-219	675.2
1-224	640.3
1-226	644.5
1-227	658.4
1-234	670.4

Example 7: N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-aminopyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide (compound 1-57)



[00241] Step 7-1, preparation of 2-[(3S)-3-{[(tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoic acid: to a 15-mL pressure tube vessel, was placed a solution of 4-chloro-2-fluorobenzoic acid methyl ester (80 mg, 0.42 mmol), pyrrolidin-3-yl-carbamic acid *tert*-butyl ester (118 mg, 0.63 mmol), and DIEA (0.11 mL, 0.63 mmol) in DMSO (2 mL). The resulting solution was stirred at 140 °C for 2.5 hrs in an oil bath. The reaction progress was monitored by

LCMS. The resulting mixture was cooled to rt and purified from C18 reversed phase column chromatography with 0.1%TFA-ACN/0.1%TFA-H₂O to afford 2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoic acid methyl ester (150 mg, 100%). LCMS (M+H)⁺: 355.3.

To a 25-mL round-bottom flask, was placed a solution of 2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoic acid methyl ester (150 mg, 0.42 mmol) in H₂O/MeOH/THF (0.6/0.6/1.8 mL). The resulting solution was treated with LiOH monohydrate (170 mg, 4.2 mmol) and stirred overnight at 70 °C. The reaction was cooled to rt, concentrated, and then diluted with H₂O (~2mL). The resulting solution was acidified with 1N-HCl, and then extracted with EtOAc (2x). The combined organic layers were washed with brine, dried, and concentrated to afford 2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoic acid (110 mg, 82%). LCMS (M+H)⁺: 341.1.


[00242] Step 7-2, preparation of benzyl N-[2-({2-[(2R)-4-{2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}formamido)ethyl]carbamate: to a 10-mL round-bottom flask, was placed a solution of 2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoic acid (25 mg, 0.073 mmol) in ACN (1.0 mL). The solution was treated with HATU (30 mg, 0.079 mmol) and Et₃N (11 µL, 0.079 mmol). After stirring for 5 min, the resulting solution was treated with benzyl N-(2-{[5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethylpiperazin-1-yl]phenyl]formamido}ethyl) carbamate (40 mg, 0.073 mmol) and stirred at rt for 20 min. The resulting mixture was directly purified from C18 reversed phase column chromatography with 0.1%TFA-ACN/0.1%TFA-H₂O to afford benzyl N-[2-({2-[(2R)-4-{2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}formamido)ethyl]carbamate (30 mg, 48 %). LCMS (M+H)⁺: 854.5.

[00243] Step 7-3, preparation of N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-aminopyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide: A solution of benzyl N-[2-({2-[(2R)-4-{2-[(3S)-3-[[tert-butoxy)carbonyl]amino}pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}formamido)ethyl]carbamate (28 mg, 0.032 mmol) and catalytic amount of thioanisole (1~2 drops) in TFA (0.5 mL) was stirred at 60 °C for 30 min. The reaction was cooled to rt and concentrated. The residue was purified from C18 reversed phase column chromatography with 0.1%TFA-ACN/0.1%TFA-H₂O to afford N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-aminopyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide (13 mg, 65%). LCMS (M+H)⁺: 620.5.

[00244] The following compounds were prepared similarly to **Example 7** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings. In some examples, the acid-catalyzed deprotection is not required in the final step:

Compound no.	MS (M+H) ⁺
1-40	634.3
1-41	634.2
1-50	668.3
1-51	618.3
1-52	634.3
1-56	618.4
1-67	614.4
1-68	628.6
1-90	646.5
1-94	640.3
1-101	644.6
1-103	644.7
1-104	614.4
1-115	674.4
1-148	618.6
1-180	694.5
1-197	680.4

Example 8: N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide (compound 1-17)



[00245] Step 8-1, preparation of benzyl N-[2-({4-[(2R)-4-(2-bromo-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate: to a 10-mL round-bottom flask, was placed a solution of 2-bromo-4-fluorobenzoic acid (40 mg, 0.18 mmol) in ACN (2.0 mL). The solution was treated with HATU (76 mg, 0.2 mmol) and Et₃N (36 μ L, 0.27 mmol). After stirring for 5 min, the resulting solution was treated with benzyl N-[2-({2'-ethoxy-4-[(2R)-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate trifluoroacetate (120 mg, 0.18 mmol) and stirred at rt for 20 min. The resulting mixture was

directly purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford benzyl N-[2-({4-[(2R)-4-(2-bromo-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate (54 mg, 41%). LCMS (M+H)⁺: 731.5.

[00246] Step 8-2, preparation of benzyl N-[2-({4-[(2R)-4-[4'-({[(tert-butoxy)carbonyl]amino}methyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate: to a 15-mL pressure tube vessel, was placed benzyl N-[2-({4-[(2R)-4-(2-bromo-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate (50 mg, 0.068 mmol), [4-({[(tert-butoxy)carbonyl]amino}methyl)phenyl]boronic acid (35 mg, 0.14 mmol), PdCl₂(t-Bu₂PPhNMe₂)₂ (5 mg, 0.007 mmol), K₂CO₃ (19 mg, 0.14 mmol), and dioxane/H₂O (2 mL/0.2 mL). The resulting solution was degassed with N₂ for 5 min and then sealed with a cap. The reaction was stirred at 85 °C for 2 hrs in an oil bath. The resulting mixture was cooled to rt, diluted with EtOAc, and washed with 1N-HCl and brine. The organic layer was dried with anhydrous MgSO₄ and concentrated. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford benzyl N-[2-({4-[(2R)-4-[4'-({[(tert-butoxy)carbonyl]amino}methyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate (43mg, 74%). LCMS (M+H)⁺: 858.8.


[00247] Step 8-3, preparation of N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide: a solution of benzyl N-[2-({4-[(2R)-4-[4'-({[(tert-butoxy)carbonyl]amino}methyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}formamido)ethyl]carbamate (43 mg, 0.050 mmol) and catalytic amount of thioanisole (1~2 drops) in TFA (0.5 mL) was stirred at 60 °C for 30 min. The reaction was cooled to rt and concentrated. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide (22 mg, 71%). LCMS (M+H)⁺: 624.4.

[00248] The following compounds were prepared similarly to **Example 8** with appropriate substituting reagents and substrates at different steps. Some examples may require –CN reduction to introduce the benzyl amine in the final step:

Compound no.	MS (M+H) ⁺
1-18	620.5
1-24	606.3
1-26	624.4

Compound no.	MS (M+H) ⁺
1-27	638.5
1-36	625.6
1-44	639.4
1-75	612.3
1-85	612.4

Example 9. 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide (compound 1-135)



[00249] Step 9-1, preparation of 2'-ethoxy-5-fluoro-[2,3'-bipyridine]-6-carbonitrile: to a solution of the mixture of 6-bromo-3-fluoropyridine-2-carbonitrile and 6-chloro-3-fluoropyridine-2-carbonitrile (4:1 ratio, 1.0 equiv, 1.0 mmol, 200 mg) in dioxane (4 mL) was added Pd(Amphos)Cl₂ (0.025 equiv, 0.025 mmol, 18 mg), 2-ethoxypyridin-3-ylboronic acid (2.0 equiv, 2.0 mmol, 334 mg), K₂CO₃ (2.2 equiv, 2.2 mmol, 304 mg) and water (0.4 mL). The resulting solution was purged with nitrogen and heated at 100 °C for 1 h. LCMS showed complete consumption of starting material and formation of desired product. This reaction was repeated on 5.0 mmol scale under similar conditions. These two batches of reaction mixture were combined, diluted with ethyl acetate (50 mL), washed with water and brined, dried with anhydrous Na₂SO₄, filtered and concentrated. The remaining residue was purified by silica gel chromatography eluting with EtOAc/hexane (0-50%) to afford 1.7 g (>100%) of the desired product containing unidentified impurities. LCMS (M+H)⁺ = 244.1.

[00250] Step 9-2, preparation of tert-butyl (3R)-4-{6-cyano-2'-ethoxy-[2,3'-bipyridin]-5-yl}-3-ethylpiperazine-1-carboxylate: to a solution of 2'-ethoxy-5-fluoro-[2,3'-bipyridine]-6-carbonitrile (1.0 equiv, 6.0 mmol, 1.7 g) in DMSO (10 mL) was added tert-butyl (3R)-3-ethylpiperazine-1-carboxylate (1.5 equiv, 9.0 mmol, 1.9 g) and DIEA (2.0 equiv, 12 mmol, 2.0 mL). The resulting solution was purged with nitrogen and heated at 140 °C for 2 days. LCMS showed the desired product formed but Boc protecting group was partially lost. Thus, the reaction was cooled to rt and followed by addition of DIPEA (2.0 equiv, 12 mmol, 2.0 mL) and di-tert-butyl dicarbonate (2.0 equiv, 12 mmol, 2.75 mmol) and stirred for 1 h. Then reaction solution was diluted with ethyl acetate (50 mL), washed with water and brine, dried with anhydrous Na₂SO₄, filtered and

concentrated. The residue was purified by C18 reversed phase column chromatography eluting with MeCN/water (5-100%). Pure fractions were combined, neutralized with saturated NaHCO_3 , and extracted with ethyl acetate. The combined organic layers were dried with MgSO_4 , concentrated, and dried under high vacuum to give 1.20 g (46%) of the desired product. LCMS $(\text{M}+\text{H})^+ = 438.3$.

[00251] Step 9-3, preparation of 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carbonitrile trifluoroacetate: to a solution of tert-butyl (3R)-4-{6-cyano-2'-ethoxy-[2,3'-bipyridin]-5-yl}-3-ethylpiperazine-1-carboxylate (1.0 equiv, 0.73 mmol, 320 mg) in DCM (3.0 mL) was added TFA (1.0 mL). The resulting mixture was stirred at ambient temperature for 1 h. The reaction was concentrated under vacuum to afford crude product (560 mg) as TFA salt. This material was used for next step without further purification. LCMS $(\text{M}+\text{H})^+ = 338.4$.

[00252] Step 9-4, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carbonitrile: to a solution of 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carbonitrile trifluoroacetate (560 mg crude, 0.73 mmol,) in DMF (3.0 mL) was added DIEA (1.06 mL, 4.38 mmol, 6.0 eq.), HATU (570 mg, 1.5 mmol, 2.0 eq.) and 2-trifluoromethyl-4-chlorobenzoic acid (224 mg, 1.0 mmol, 1.35 eq.). The resulting mixture was stirred at rt for 10 min and LCMS showed complete consumption of starting material. The reaction solution was diluted with ethyl acetate (50 mL), washed with water and brine, with anhydrous Na_2SO_4 , filtered and concentrated. The residue was purified by silica gel chromatography eluting with EtOAc/hexane (0-50%) to afford 230 mg (58%) of the desired product. LCMS $(\text{M}+\text{H})^+ = 544.3$.

[00253] Step 9-5, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid: to a solution of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carbonitrile (27 mg, 0.05 mmol,) in EtOH (0.25 mL) was added KOH (85%, 28 mg, 0.50 mmol, 10 eq.) and water (0.25 mL). The resulting mixture was heated at 100 °C for overnight. The reaction mixture was cooled to rt, diluted with ethyl acetate (10 mL), washed with saturated NaHSO_4 (5.0 mL) and brine, dried and concentrated to afford 26 mg of crude title compound. This material was used for next step without further purification. LCMS $(\text{M}+\text{H})^+ = 563.3$.

[00254] Step 9-6, preparation of tert-butyl N-[2-({5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl})formamido]ethyl]-N-methylcarbamate: to a solution of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid (26 mg, 0.05 mmol) in DMF (0.5 mL) was added DIEA (0.10 mL, 0.55 mmol, 11 eq.), HATU

(38 mg, 0.10 mmol, 2.0 eq.) and 1-boc-1-methylethylenediamine (19 mg, 0.10 mmol, 2.0 eq.). The resulting mixture was stirred at ambient temperature for 0.5 h. The crude reaction solution was purified by a C18 reversed phase column chromatography eluting with MeCN/water (5-100%). Pure fractions were combined, neutralized with saturated NaHCO₃ and NaCl, extracted with ethyl acetate and dried with MgSO₄. The organic layer was concentrated and dried under high vacuum to give 15 mg (47%) of the desired product. LCMS (M+H)⁺ = 719.3.


[00255] Step 9-7, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide: to a solution of tert-butyl N-[2-({5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}formamido)ethyl]-N-methylcarbamate (15 mg, 0.021 mmol) in DCM (0.5 mL) was added TFA (0.2 mL). The resulting mixture was stirred at ambient temperature for 0.5 h. The reaction was concentrated and purified by a C18 reversed phase column chromatography eluting with MeCN/water (5-60%). Pure fractions were combined, neutralized with saturated NaHCO₃ and NaCl, extracted with ethyl acetate and dried with MgSO₄. The organic layer was concentrated and dried under high vacuum to give 9 mg (70%) of the desired product. LCMS (M+H)⁺ = 619.4.

[00256] The following compounds were prepared similarly to **Example 9** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings. In some examples, an additional deprotection is not required in the final step.

Compound no.	MS (M+H) ⁺
1-136	602.6
1-137	629.6
1-149	632.3
1-151	631.4
1-167	615.4
1-171	576.4
1-172	594.5
1-181	630.6
1-185	600.4
1-187	616.6
1-194	603.4
1-195	595.3
1-205	617.4
1-206	644.4
1-208	618.6
1-209	630.5
1-211	617.5
1-214	631.4
1-215	645.5

Compound no.	MS (M+H) ⁺
1-218	607.3
1-220	616.3
1-221	630.5
1-222	631.3
1-223	645.4
1-230	628.4
1-231	645.3
1-232	645.5
1-233	642.4
1-236	627.4
1-237	641.4
1-239	642.5
1-240	628.5
1-242	642.5
1-243	656.5
1-244	628.4
1-245	642.4
1-246	614.3
1-247	628.4
1-249	656.6
1-256	642.4
1-257	656.6
1-268	646.3
1-269	632.3
1-272	632.1
1-275	649.3
1-276	649.5
1-289	646.4
1-290	660.5
1-291	646.5
1-296	660.3
1-303	581.2

Example 10. 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide Acetate (compound 1-150)



[00257] Step 10-1, preparation of 5-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid: to a solution of tert-butyl (3R)-4-{6-cyano-2'-ethoxy-[2,3'-bipyridin]-5-yl}-3-ethylpiperazine-1-carboxylate from **Example 9**, step 9-2 (271 mg, 0.62 mmol) in EtOH (3 mL) was added KOH (85%, 409 mg, 6.2 mmol, 10 eq.) and water (3.0 mL). The resulting mixture was heated at 100 °C for overnight. The reaction mixture was cooled to rt, diluted with ethyl acetate (10 mL), washed with saturated NaHSO₄ (10 mL) and brine, dried and concentrated to afford 283 mg of crude title compound. This material was used for next step without further purification. LCMS (M+H)⁺ = 457.3.

[00258] Step 10-2, preparation of tert-butyl (3R)-4-(6-{[2-(dimethylamino)ethyl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate: to a solution of 5-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid (283 mg, 0.619 mmol) in DMF (5.0 mL) was added HATU (353 mg, 0.93 mmol, 1.5 eq.), DIEA (0.31 mL, 1.86 mmol, 3.0 eq.) and N,N-dimethylethylenediamine (84 mg, 0.93 mmol, 1.5 eq.). The resulting mixture was stirred at 50 °C for 0.5 h. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried with Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography eluting with MeOH/DCM (0-15%) to afford 396 mg of the desired product as brown oil. LCMS (M+H)⁺ = 527.4.

[00259] Step 10-3, preparation of N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of tert-butyl (3R)-4-(6-{[2-(dimethylamino)ethyl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (396 mg, 0.75 mmol) in DCM (4.0 mL) was added TFA (2.0 mL). The resulting mixture was stirred at ambient temperature for 0.5 h and concentrated under vacuum. The residue was neutralized with saturated NaHCO₃, extracted with ethyl acetate, dried with Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography eluting with MeOH/DCM (0-20%) to afford 201 mg (64%) of the desired product as brown oil. LCMS (M+H)⁺ = 427.4.


[00260] Step 10-4, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide acetate: to a solution of N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide (89 mg, 0.20 mmol) in DMF (2 mL) was added HATU (91 mg, 0.24 mmol, 1.2 eq.), DIEA (0.092 mL, 0.50 mmol, 2.5 eq) and 2-trifluoromethyl-4-chlorobenzoic acid (45 mg, 0.20 mmol, 1.0 eq.). The resulting mixture was heated at 50 °C for 0.5 h. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried with Na₂SO₄, filtered and concentrated. The residue was purified by silica gel chromatography eluting with

MeOH/DCM (0-9%). Pure fraction were combined and concentrated with HOAc (0.1 mL) to afford 90 mg (64%) of the desired product. LCMS (M+H)⁺ = 633.4.

[00261] The following compounds were prepared similarly to **Example 10** with appropriate substituting reagents and substrates at different steps. In some examples, an additional deprotection is required in the final step.

Compound no.	MS (M+H) ⁺
1-158	590.1
1-169	629.5
1-196	609.3
1-207	644.5
1-225	629.3
1-253	641.1
1-254	627.4
1-258	629.4
1-259	611.2
1-260	602.4
1-266	615.2
1-267	597.2
1-277	588.2

Example 11: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]pyrazine-2-carboxamide (compound 3-8)



[00262] Step 11-1, preparation of methyl 6-bromo-3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]pyrazine-2-carboxylate: to a 15-mL pressure tube vessel, was placed 6-bromo-3-chloropyrazine-2-carboxylic acid methyl ester (300 mg, 1.19 mmol), tert-butyl (3R)-3-ethylpiperazine-1-carboxylate (370 mg, 1.72 mmol, 1.4 equiv), DIEA (0.3 mL, 1.78 mmol, 1.5 equiv), and DMSO (6 mL). The resulting solution was stirred at 140 °C for 1 h, cooled to rt, and diluted with EtOAc (~15 mL). The organic layer was washed with 1N HCl (2x) and brine (1x), dried with anhydrous Na₂SO₄, and concentrated. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O. 140 mg (27%) of methyl 6-bromo-3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]pyrazine-2-carboxylate was isolated. LCMS (M+H)⁺: 429.3.

[00263] Step 11-2, preparation of methyl 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylate: to a 15-mL pressure tube vessel, was placed 66-bromo-3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]pyrazine-2-carboxylate (100 mg, 0.23 mmol), 2-ethoxypyridine-3-boronic acid (78 mg, 0.46 mmol), PdCl₂(t-Bu₂PPhNMe₂)₂ (24 mg, 0.033 mmol), K₂CO₃ (64 mg, 0.46 mmol), and dioxane/H₂O (2.5 mL/0.25 mL). The resulting solution was degassed with N₂ for 5 min and then sealed with a cap. The reaction was stirred at 100 °C for 1 h in an oil bath. The resulting mixture was cooled to rt, diluted with EtOAc, and washed with 1N-HCl and brine. The organic layer was dried with anhydrous Na₂SO₄ and concentrated. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford methyl 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylate (85 mg, 77%). LCMS (M+H)⁺: 472.3.

[00264] Step 11-3, preparation of 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid: to a solution of methyl 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylate (85 mg, 0.18 mmol) in MeOH/H₂O/THF (1/1/3 mL) was added lithium hydroxide hydrate (108 mg, 15 eq.). The resulting solution was stirred at rt for 3 hrs. After removal of the volatile solvent, the aqueous residue was diluted with H₂O (~3 mL), and then acidified with 1N HCl. The resulting aqueous layer was extracted with DCM (3x). The combined organic layers were washed with brine, dried with anhydrous Na₂SO₄, and concentrated under vacuum to afford 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid (75 mg, 91%). LCMS (M+H)⁺: 458.3.

[00265] Step 11-4, preparation of 6-(2-ethoxypyridin-3-yl)-3-[(2R)-2-ethylpiperazin-1-yl]pyrazine-2-carboxylic acid: a solution of 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid (75 mg, 0.16 mmol) in TFA (0.2 mL)/DCM (0.6 mL) was stirred at rt for 1.5 h. The resulting mixture was concentrated under vacuum. The residue (~100%) was diluted with ACN (1 mL) and neutralized with Et₃N, which was used in the next step without further purification. LCMS (M+H)⁺: 358.2.

[00266] Step 11-5, preparation of 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid: to a solution of 4-chloro-2-(trifluoromethyl)benzoic acid (45 mg, 0.2 mmol, 1.33 eq.) in ACN (1 mL) was added HATU (75 mg, 0.2 mmol) and followed by Et₃N (40 µL, 0.35 mmol). The resulting solution was stirred for 5 min at rt and treated with 6-(2-ethoxypyridin-3-yl)-3-[(2R)-2-ethylpiperazin-1-yl]pyrazine-2-carboxylic acid (~55 mg, 0.15 mmol, 1 eq.) prepared from the previous step. The

resulting solution was stirred for 20 min at room temperature and concentrated under vacuum. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid (55 mg, 64%). LCMS (M+H)⁺: 564.5.


[00267] Step 11-6, preparation of tert-butyl N-[2-({3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazin-2-yl}formamido)ethyl]-N-methylcarbamate: to a 10-mL vial, was placed 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxylic acid (30 mg, 0.053 mmol, 1.00 eq.), HATU (21 mg, 0.056 mmol), Et₃N (11 μ L, 0.08 mmol) and ACN (1 mL). The resulting solution was stirred for 5 min at rt and treated with tert-butyl N-(2-aminoethyl)-N-methylcarbamate (11 mg, 0.063 mmol, 1.2 eq.). The reaction was stirred for 20 min at room temperature and concentrated under vacuum. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford tert-butyl N-[2-({3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazin-2-yl}formamido)ethyl]-N-methylcarbamate (18 mg, 47%). LCMS (M+H)⁺: 720.3.

[00268] Step 11-7, preparation of 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]pyrazine-2-carboxamide: a solution of tert-butyl N-[2-({3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)pyrazin-2-yl}formamido)ethyl]-N-methylcarbamate (18 mg, 0.025 mmol) in TFA (0.2 mL)/DCM (0.6 mL) was stirred at rt for 1.5 h. The resulting mixture was concentrated under vacuum. The residue was purified from C18 reversed phase column chromatography eluting with 0.1%TFA-ACN/0.1%TFA-H₂O to afford 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]pyrazine-2-carboxamide (10 mg, 67 %). LCMS (M+H)⁺: 620.4.

[00269] The following compounds were prepared similarly to **Example 11** with appropriate substituting reagents and substrates at different steps. In some examples, an additional deprotection is not required in the final step.

Compound no.	MS (M+H) ⁺
3-7	634.2

Example 12: 1-{5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine (compound 2-13)




[00270] Step 12-1, preparation of 1-{5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine: to a solution of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carbonitrile from **Example 9**, step 9-4 (65 mg, 0.12 mmol) in MeOH (1.0 mL) was added NiCl₂ (8.0 mg, 0.06 mmol, 0.5 eq.) and NaBH₄ (36 mg, 0.96 mmol, 8.0 eq.) at 0 °C. The resulting mixture was stirred at the same temperature for 15 min and LCMS analysis showed about 40% conversion to the desired product. At 0 °C, an additional NiCl₂ (0.5 equiv, 0.06 mmol, 8.0 mg) and NaBH₄ (8.0 equiv, 0.96 mmol, 36 mg) were added and the resulting mixture was stirred for another 15 min. The reaction mixture was diluted with ethyl acetate and washed with saturated NH₄Cl. The organic layer was dried and concentrated. The residue was purified by a C18 reversed-phase column chromatography eluting with MeCN/water (5-60%). Pure fractions were combined, neutralized with saturated NaHCO₃ and NaCl, extracted with ethyl acetate and dried with MgSO₄. The organic layer was concentrated and dried under high vacuum to give 8 mg (12%) of the desired product. LCMS (M+H)⁺ = 548.3.

[00271] The following compounds were prepared similarly to **Example 12** with appropriate substituting reagents and substrates at different steps. Some examples may require additional functional group transformations to introduce an appropriate substituent on aryl rings.

Compound no.	MS (M+H) ⁺
2-3	532.2
2-4	576.4
2-5	560.2
2-6	562.4
2-7	528.2
2-8	532.2
2-9	547.3
2-10	568.5
2-11	515.3
2-12	547.3
2-14	494.3
2-15	548.2

Compound no.	MS (M+H) ⁺
2-16	562.3
2-17	498.2
2-18	515.3
2-19	500.3
2-20	531.9
2-22	532.1
2-23	546.3
2-24	572.3
2-25	558.1
2-26	544.3
2-27	526.3
2-28	544.3
2-29	544.4

Example 13: 2-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl]amino]ethan-1-ol (compound 2-1)



[00272] Step 13-1, preparation of tert-butyl (3R)-4-[2'-ethoxy-3-(methoxycarbonyl)-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate: into a 8-mL pressure tube vessel, was placed tert-butyl (3R)-4-[4-bromo-2-(methoxycarbonyl)phenyl]-3-ethylpiperazine-1-carboxylate from **Example 1**, step 1.3 (80 mg, 0.19 mmol), (2-ethoxyphenyl)boronic acid (46.6 mg, 0.28 mmol), Pd₂(dba)₃CHCl₃ (19.4 mg, 0.02 mmol), P(t-Bu)₃.HBF₄ (5.4 mg, 0.02 mmol), K₃PO₄ (119.2 mg, 0.56 mmol), and toluene/H₂O (1 mL/0.1 mL). The resulting solution was degassed for 5 min with N₂, sealed, and stirred at 70 °C for 2 hrs in an oil bath. The reaction was cooled to rt and concentrated under vacuum. The residue was applied onto a silica gel column eluting with ethyl acetate/petroleum ether (1:1). 80 mg (91%) of the desired product was isolated. LCMS (M+H)⁺ = 469.1.

[00273] Step 13-2, preparation of tert-butyl (3R)-4-[2'-ethoxy-3-(hydroxymethyl)-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate: to a heterogenous solution of tert-butyl (3R)-4-[2'-ethoxy-3-(hydroxymethyl)-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate (60 mg, 0.13 mmol) and NiCl₂ (16.6 mg, 0.13 mmol) in MeOH (1.0 mL) was added NaBH₄ (48.4 mg, 1.28 mmol, 10 eq.) at rt. The resulting mixture was stirred at 50 °C for 1 h. The reaction mixture was

cooled to rt, diluted with H₂O, and extracted with ethyl acetate (3x). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated to afford 60 mg (~100%) of the desired product. LCMS (M+H)⁺ = 441.3.

[00274] Step 13-3, preparation of tert-butyl (3R)-4-{2'-ethoxy-3-formyl-[1,1'-biphenyl]-4-yl}-3-ethylpiperazine-1-carboxylate: a solution of tert-butyl (3R)-4-[2'-ethoxy-3-(hydroxymethyl)-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate (90 mg, 0.2 mmol) and IBX (171.6 mg, 0.62 mmol) in ACN (2 mL) was stirred at rt for 1 h. After removal of the solids by filtration, the filtrate was concentrated under vacuum to afford the crude desired product (90 mg, ~100%), which was used in the next step without further purification. LCMS (M+H)⁺ = 439.3.

[00275] Step 13-4, preparation of tert-butyl (3R)-4-[3-({[2-(benzyloxy)ethyl]amino}methyl)-2'-ethoxy-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate: to a solution of tert-butyl (3R)-4-{2'-ethoxy-3-formyl-[1,1'-biphenyl]-4-yl}-3-ethylpiperazine-1-carboxylate (30 mg, 0.07 mmol) and 2-(benzyloxy)ethan-1-amine (21 mg, 0.14 mmol) in DCM (2.0 mL) was added NaBH(OAc)₃ (44 mg, 0.21 mmol, 3 eq.). The resulting mixture was stirred at rt for 30 min and diluted with H₂O (20 mL). The resulting solution was extracted with EtOAc (3x). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated to afford the desired product (25 mg, 64%). LCMS (M+H)⁺ = 574.4.

[00276] Step 13-5, preparation of [2-(benzyloxy)ethyl]({2'-ethoxy-4-[(2R)-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-yl}methyl)amine trifluoroacetate: to a solution of tert-butyl (3R)-4-[3-({[2-(benzyloxy)ethyl]amino}methyl)-2'-ethoxy-[1,1'-biphenyl]-4-yl]-3-ethylpiperazine-1-carboxylate (25 mg, 0.04 mmol) in DCM (2.0 mL) was added TFA (1 mL). The resulting mixture was stirred at rt for 1 h and concentrated under vacuum. This resulted in the desired product (25 mg, 98%) as TFA salt. LCMS (M+H)⁺ = 474.3.


[00277] Step 13-6, preparation of [2-(benzyloxy)ethyl]({4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl}methyl)amine: to a solution of 2-chloro-4-(trifluoromethyl)benzoic acid (2 mg, 0.01 mmol, 0.3 eq.), HATU (5.7 mg, 0.01 mmol, 0.3 eq.) and [2-(benzyloxy)ethyl]({2'-ethoxy-4-[(2R)-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-yl}methyl)amine trifluoroacetate (17.5 mg, 0.03 mmol) in DMF (2 mL) was added DIEA (11.6 mg, 0.09 mmol, 3 eq.). The resulting mixture was stirred at rt for 1 h and diluted with H₂O (20 mL). The resulting solution was extracted with EtOAc (3x). The combined organic layers were dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by a silica gel column chromatography eluting with DCM/MeOH (10/1) to afford the desired product (20 mg, 38%). LCMS (M+H)⁺ = 680.2.

[00278] Step 13-7, preparation of 2-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl]amino]ethan-1-ol trifluoroacetate: A solution of [2-(benzyloxy)ethyl]({4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl}amine (20 mg, 0.03 mmol) in TFA (1.0 mL) was stirred at 80 °C for 3 hrs. The resulting mixture was cooled to rt and concentrated under vacuum. The residue was purified by Prep-HPLC with the following conditions: SunFire Prep C18 OBD column; mobile phase, phase A: water (0.05% TFA); phase B: ACN (31% up to 45% in 6 min); flow rate: 20 mL/min; detector, 220 & 254 nm. 3.9 mg (19%) of the desired product was isolated as a TFA salt. LCMS (M+H)⁺ = 590.2.

[00279] The following compounds were prepared similarly to **Example 13** with appropriate substituting reagents and substrates at different steps. In some examples, an additional deprotection is not required in the final step.

Compound no.	MS (M+H) ⁺
2-2	604.3
2-21	588.4

Example 14: N-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide (compound 3-2)



[00280] Step 14-1, preparation of tert-butyl 4-[4-bromo-2-(methoxycarbonyl)phenyl]piperazine-1-carboxylate: to a solution of *tert*-butyl piperazine-1-carboxylate (1.600 g, 8.588 mmol) and methyl 4-bromo-2-fluorobenzoate (1.000 g, 4.292 mmol) in DMSO (5 mL) was added DIEA (2.20 mL, 12.6 mmol). The mixture was heated at 130°C for 1 day. The mixture was distributed between water and DCM and the organic layer was separated and concentrated. The residue was purified by C18 reversed phase column chromatography to give the title compound (0.5088 g, 30% yield) as a medium brown solid. MS (M+H)⁺: 399.2. The hydrolyzed acid (0.7537 g, 46% yield) was also isolated as a light brown solid. LCMS (M+H)⁺= 385.3.

[00281] Step 14-2, preparation of methyl 5-bromo-2-(piperazin-1-yl)benzoate hydrochloride: to a solution of *tert*-butyl 4-[4-bromo-2-(methoxycarbonyl)phenyl]piperazine-1-carboxylate (508.8 mg, 1.274 mmol) in DCM (2 mL) was added 4N HCl in dioxane (2.0 mL, 8.0 mmol). The

mixture was stirred at rt for 2 hrs. The mixture was concentrated to dryness to give the title compound as a light yellow solid. LCMS (M+H)⁺ = 299.2.

[00282] Step 14-3, preparation of methyl 5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}benzoate: To a solution of methyl 5-bromo-2-(piperazin-1-yl)benzoate hydrochloride (0.4538 g, 1.352 mmol) and 2-fluoro-5-(trifluoromethyl)benzonitrile (353.0 mg, 1.867 mmol) in DMSO (3 mL) was added DIEA (0.81 mL, 4.7 mmol). The mixture was heated at 130°C overnight. The mixture was purified by C18 reversed phase column chromatography to give the title compound (453 mg, 72% yield) as a yellow solid. LCMS (M+H)⁺ = 468.3.

[00283] Step 14-4, preparation of 5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}benzoic acid: To a solution of methyl 5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}benzoate (453 mg, 0.967 mmol) in a mixed solvent of THF:MeOH:H₂O (6:2:2 mL) was added LiOH·H₂O (406 mg, 9.67 mmol). The mixture was stirred at rt for 4 hrs. After removal of volatile solvent, the aqueous residue was diluted with water and acidified with 1N HCl (aq) to pH 2-3. The solid was collected by vacuum filtration and washed with water and dried to give title compound (421 mg, 96% yield) as a yellow solid. LCMS (M+H)⁺ = 454.1.

[00284] Step 14-5, preparation of *tert*-butyl *N*-{3-[(5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}phenyl)formamido]propyl}carbamate: To a solution of 5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}benzoic acid (201.9 mg, 0.4444 mmol) and HATU (220 mg, 0.579 mmol) in DMF (1 mL) was added DIEA (0.15 mL, 0.86 mmol). After stirring at RT for 5 min, the resulting solution was treated with *tert*-butyl *N*-(3-aminopropyl)carbamate (126.5 mg, 0.7262 mmol) and stirred at rt for 5 min. The reaction was purified by C18 reversed phase column chromatography to give the title compound (198.7 mg, 73% yield) as an off-white solid. LCMS (M+H)⁺ = 610.3.

[00285] Step 14-6, preparation of *tert*-butyl *N*-{3-[(4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-yl)formamido]propyl}carbamate: To a mixture of *tert*-butyl *N*-{3-[(5-bromo-2-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}phenyl)formamido]propyl}carbamate (20.0 mg, 0.0328 mmol), 2-ethoxyphenylboronic acid (10.9 mg, 0.0657 mmol), Pd[*t*-Bu₂P(4-NMe₂C₆H₄)]₂Cl₂ (9.2 mg, 0.013 mmol), and K₂CO₃ (27.2 mg, 0.197 mmol) in a sealed tube was added dioxane (2 mL) and H₂O (0.2 mL). The resulting mixture was degassed with N₂ for 10 min and stirred at 100°C for 30 min. The mixture was concentrated and purified by C18 reversed phase column


chromatography to give the title compound (18.5 mg, 87% yield) as a white solid. LCMS (M+H)⁺ = 652.5.

[00286] Step 14-7, preparation of *N*-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide: to a solution of *tert*-butyl *N*-{3-[(4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-yl)formamido]propyl} carbamate (18.5 mg, 0.0284 mmol) in DCM (0.1 mL) was added 4N HCl in dioxane (0.1 mL, 0.4 mmol). The mixture was stirred at rt for 30 min. The mixture was concentrated to dryness to give 1HCl salt of the title compound which was triturated with saturated NaHCO₃ (aq) to give the title compound (12.9 mg, 82% yield) as an off-white solid. LCMS (M+H)⁺ = 552.2.

[00287] The following compounds were prepared similarly to **Example 14** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
3-1	544.3

Example 15: 6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide (compound 3-3)




[00288] Step 15-1, preparation of benzyl N-[2-[(6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl]formamido)ethyl]-N-methylcarbamate: into a 8-mL vial, was placed benzyl N-(2-[[3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethylpiperazin-1-yl]-2-fluorophenyl]formamido]ethyl)-N-methylcarbamate trifluoroacetate (30 mg, 0.027 mmol), which was prepared by a similar manner described in **Example 4**, step 1 & 2, TEA (8 mg, 0.079 mmol, 2.98 eq.), 2,4-dichlorobenzene-1-sulfonyl chloride (8 mg, 0.033 mmol, 1.23 eq.), and DCM (2 mL). The resulting solution was stirred for 2 hr at 25 °C. The reaction was diluted with DCM (10 mL) and washed with H₂O (2x20 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 20 mg (crude) of the title compound as off-white oil. LCMS (M+H)⁺ = 772.2.

[00289] Step 15-2, preparation of 6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide: Into a 8-mL vial, was placed benzyl N-[2-[(6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-

ethoxypyridin-3-yl)-2-fluorophenyl]formamido)ethyl]-N-methylcarbamate (20 mg, 0.021 mmol) and trifluoroacetic acid (2 mL). The resulting solution was stirred for 3 hrs at 60 °C, cooled to rt, and concentrated under vacuum. The residue was dissolved in DMF (4 mL) and purified by reverse phase HPLC resulting in 7.9 mg (56 %) of the title compound as a formic acid salt.

LCMS (M+H)⁺ = 638.2

Example 16: (3R)-N-(2,4-dichlorophenyl)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-[[2-(methvlamino)ethyl]carbamoyl]phenyl]-3-ethylpiperazine-1-carboxamide (compound 3-4)




[00290] Step 16-1, preparation of phenyl N-(2,4-dichlorophenyl)carbamate: Into a 8-mL vial, was placed 2,4-dichloroaniline (200 mg, 1.23 mmol) and THF (5 mL). The resulting solution was treated with NaH (59 mg, 1.47 mmol, 1.19 eq, 60% purity) and stirred for 20 min at 0 °C. The reaction was treated with phenyl carbonochloridate (232 mg, 1.48 mmol, 1.2 eq.), stirred for an additional 3 hr at 0 °C, and concentrated under vacuum. The residue was purified from a silica gel column chromatography eluting with ethyl acetate/petroleum ether (1:5). This resulted in 120 mg (34%) of the title compound as yellow oil. LCMS (M+H)⁺ = 282.0.

[00291] Step 16-2, preparation of benzyl N-[2-([6-[(2R)-4-[(2,4-dichlorophenyl)carbamoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl]formamido)ethyl]-N-methylcarbamate: Into a 8-mL vial, was placed benzyl N-[2-([3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethylpiperazin-1-yl]-2-fluorophenyl]formamido)ethyl]-N-methylcarbamate trifluoroacetate (36 mg, 0.053 mmol), DCM (2 mL), TEA (16 mg, 0.158 mmol, 2.97 eq.), and phenyl N-(2,4-dichlorophenyl)carbamate (15 mg, 0.053 mmol). The resulting solution was stirred for 2 hrs at 25 °C, diluted with DCM (10 mL), and washed with water (2x20 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 30 mg (crude) of the title compound as yellow oil. LCMS (M+H)⁺ = 751.2.

[00292] Step 16-3, preparation of (3R)-N-(2,4-dichlorophenyl)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-[[2-(methvlamino)ethyl]carbamoyl]phenyl]-3-ethylpiperazine-1-carboxamide: Into a 8-mL vial, was placed benzyl N-[2-([6-[(2R)-4-[(2,4-dichlorophenyl)carbamoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl]formamido)ethyl]-N-methylcarbamate (30 mg, 0.04 mmol) and TFA (2 mL). The resulting solution was stirred for 2 hrs at 60 °C, cooled to rt,


and concentrated under vacuum. The residue was dissolved in DMF (4 mL) and purified by Prep-HPLC with the following conditions (Prep-HPLC-013): Column, Atlantis Prep T3 OBD Column, 19*150mm 5um; mobile phase, Water (0.1% FA) and ACN (24% PhaseB up to 53% in 6 min); 20 mL/min. Detector, UV 220, 254 nm. This resulted in 5.6 mg (21 %) of the title compound as a formic acid salt. LCMS (M+H)⁺ = 617.2.

Example 17: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-152)



[00293] Step 17-1, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid from **Example 9**, step 5 (40 mg, 0.071 mmol) in ACN (0.5 mL) was added HATU (30 mg, 0.078 mmol, 1.1 eq.) and Et₃N (19 µL, 2.0 eq.). The resulting solution was stirred at rt for 5 min and treated with (*R*)-1-methylpyrrolidin-3-ylamine (10.6 mg, 0.106 mmol, 1.5 eq.). The reaction was stirred at rt for 0.5 h. The crude reaction solution was purified by a C18 reversed phase column chromatography eluting with MeCN/water (5-100%). The combined pure fractions were neutralized with saturated NaHCO₃, removed volatile solvent, and extracted with DCM (3x). The combined organic layer was dried with anhydrous Na₂SO₄, concentrated and dried under high vacuum to give 25 mg (55%) of the title compound. LCMS (M+H)⁺ = 645.5.

Example 18: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide (compound 1-212)




[00294] Step 18-1, preparation of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide: the title compound was prepared by a similar manner described in **Example 17** starting from 40 mg (0.071 mmol) of 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid. 20 mg (45%) of the title compound was isolated. LCMS (M+H)⁺ = 631.4.

[00295] The following compounds were prepared similarly to **Example 18** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-359	663.3
1-398	671.4

Example 19: 1-{2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine (compound 2-30)



[00296] Step 19-1, preparation of 5-[(2R)-4-[6-chloro-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carbonitrile: to a DMF (3.0 mL) of 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carbonitrile (Step 9-3, 1.0 equiv, 0.26 mmol, 87 mg) was added DIPEA (2.5 equiv, 0.65 mmol, 0.1 mL), HATU (1.2 equiv, 0.39 mmol, 118 mg) and 6-chloro-2-(trifluoromethyl)pyridine-3-carboxylic acid (1.0 equiv, 0.26 mmol, 61 mg). The resulting mixture was stirred at rt for 0.5 h and LCMS showed complete consumption of starting material. The reaction solution was diluted with ethyl acetate (50 mL), washed with water and brine, dried with anhydrous Na₂SO₄, filtered and concentrated. The remaining residue was purified by silica gel chromatography eluting with EtOAc/hexane (0-50%) to afford 104 mg of the title compound. LCMS (M+H)⁺ = 545.3.

[00297] Step 19-2, preparation of 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carbonitrile: to an EtOH (0.5 mL) solution of 5-[(2R)-4-[6-chloro-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carbonitrile (1.0 equiv, 0.05 mmol, 28 mg) was added NaOEt (21% w/w in EtOH, 2.5 equiv, 0.13 mmol, 0.05 mL). The resulting mixture was heated at 70 °C for 0.5 h. The

reaction solution was diluted with ethyl acetate (20 mL), washed with saturated NH_4Cl and brine, dried with anhydrous Na_2SO_4 , filtered and concentrated to afford 25 mg of crude product. This material was used for next step without purification. LCMS $(\text{M}+\text{H})^+ = 555.2$.


[00298] Step 19-3, preparation of tert-butyl N-({2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methyl)carbamate: to a MeOH (1.0 mL) solution of 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carbonitrile (1.0 equiv, 0.05 mmol, 25 mg) was added NiCl_2 (0.5 equiv, 0.025 mmol, 3.3 mg) and di-tert-butyl dicarbonate (2.0 equiv, 0.10 mmol, 0.023 mL). At 0 °C, NaBH_4 (7.0 equiv, 0.35 mmol, 13.5 mg) was added portion-wise and the resulting mixture was stirred at the same temperature for 0.5 h. The reaction mixture was filtered and filtrate was concentrated. The remaining residue was purified by silica gel chromatography eluting with EtOAc/hexane (0-50%) to afford 24 mg of the title compound. LCMS $(\text{M}+\text{H})^+ = 659.4$.

[00299] Step 19-4, preparation of 1-{2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine: to a DCM (0.5 mL) of tert-butyl N-({2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methyl)carbamate (1.0 equiv, 0.036 mmol, 24 mg) was added TFA (0.10 mL). The resulting mixture was stirred at ambient temperature for 0.5 h. The reaction solution was concentrated and the residue obtained was purified by C18 reversed-phase chromatography eluting with MeCN/water (5-50%). Pure fractions were combined, neutralized with saturated NaHCO_3 and NaCl, extracted with ethyl acetate and dried with MgSO_4 . The organic layer was concentrated and dried under high vacuum to give 6 mg of the title compound. LCMS $(\text{M}+\text{H})^+ = 559.2$

[00300] The following compounds were prepared similarly to **Example 19** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS $(\text{M}+\text{H})^+$
2-31	545.4

Example 20: 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-279)



[00301] Step 20-1, preparation of 2-(difluoromethyl)-4-fluorobenzoic acid: to a mixture of 4-fluoro-2-methylbenzoic acid (1.000 g, 6.49 mmol), Selectfluor (6.895 g, 19.46 mmol) and sodium persulfate (0.772 g, 3.24 mmol) under nitrogen was added MeCN (3.2 mL) and water (3.2 mL). The mixture was cooled to -78°C under nitrogen and silver nitrate (110.2 mg, 0.649 mmol) was added. The frozen mixture was degassed under vacuum and charged with nitrogen (repeated 3X). The mixture was warmed to rt under nitrogen and then heated at 80°C for 7 hrs. The reaction was not complete and thus more Selectfluor (2.298 g, 6.49 mmol), sodium persulfate (0.257 g, 1.08 mmol) and silver nitrate (36.7 mg, 0.216 mmol) were added under nitrogen and the reaction was continued to heat at 80°C for 6 hrs. The mixture was filtered through Celite and rinsed with EtOAc. The filtrate was washed with saturated NaHCO₃ (aq) (5 X 25 mL). The combined basic aqueous solution was acidified with 1N HCl (aq) to pH 2-3 to form a solid. The solid was collected by vacuum filtration, washed with water and dried under vacuum to give the title compound (484.1 mg). The filtrate was extracted with EtOAc (1X) and the organic layer was concentrated and purified by C18 reversed phase column chromatography to give more product (314.4 mg). The reaction overall gave the title compound (798.5 mg, 4.20 mmol, 64.7% yield) as an off-white solid.

[00302] Step 20-2, preparation of *tert*-butyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbonyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate: to a solution of 5-[(2R)-4-[(*tert*-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid (400 mg, 0.876 mmol) from Step 10-1 in DMF (2.0 mL) was added HATU (500 mg, 1.31 mmol), DIEA (0.46 mL, 2.63 mmol). After stirring at rt for 5 min, (3R)-1-methylpyrrolidin-3-amine (176 mg, 1.75 mmol) was added. The resulting mixture was stirred at rt for 10 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (284.6 mg, 0.53 mmol, 60.3% yield) as a light brown solid. LCMS (M+H)⁺ = 539.5.

[00303] Step 20-3, preparation of 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide *tris*(2,2,2-trifluoroacetate): to a solution


of *tert*-butyl (3*R*)-4-(2'-ethoxy-6-{[(3*R*)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (268.8 mg, 0.50 mmol) in DCM (1.5 mL) was added TFA (0.76 mL). The resulting mixture was stirred at rt for 30 min and concentrated *in vacuo* to dryness to give the title compound (389.5 mg, 0.50 mmol, 100% yield) as a brown gum which was used in the next step without further purification. LCMS (M+H)⁺ = 439.1.

[00304] Step 20-4, preparation of 5-[(2*R*)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 2-(difluoromethyl)-4-fluorobenzoic acid (142.3 mg, 0.75 mmol) in DMF (0.5 mL) was added HATU (265.6 mg, 0.70 mmol) and DIEA (0.26 mL, 1.50 mmol). After stirring at rt for 5 min, this HATU-activated solution was added to a solution of 2'-ethoxy-5-[(2*R*)-2-ethylpiperazin-1-yl]-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide *tris*(2,2,2-trifluoroacetate) (389.5 mg, 0.50 mmol) and DIEA (0.52 mL, 3.00 mmol) in DMF (0.5 mL). The resulting mixture was stirred at rt for 10 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (245.0 mg, 0.40 mmol, 80.4% yield) as a light brown solid. LCMS (M+H)⁺ = 611.3.

[00305] The following compounds were prepared similarly to **Example 20** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-262	627.3
1-287	626.5
1-293	610.4

Example 21: 5-[(2*R*)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-*N*-[(3*R*)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-278)



[00306] Step 21-1, preparation of *tert*-butyl (3*R*)-4-(6-{[(3*R*)-1-(benzyloxy)carbonyl]pyrrolidin-3-yl}carbamoyl)-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate: to a solution of 5-[(2*R*)-4-[(*tert*-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxylic acid (300 mg, 0.657 mmol) from Step 10-1 in DMF (1.0 mL) was added HATU (325 mg, 0.854 mmol), DIEA (0.23 mL, 1.3 mmol). After stirring at rt for 5 min, benzyl (3*R*)-3-aminopyrrolidine-1-carboxylate (217 mg,

0.986 mmol) was added. The resulting mixture was stirred at rt for 10 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (270.8 mg, 0.411 mmol, 62.6% yield) as a brown solid. LCMS (M+H)⁺ = 659.5.

[00307] Step 21-2, preparation of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate bis(2,2,2-trifluoroacetate): to a solution of *tert*-butyl (3R)-4-(6-{[(3R)-1-[(benzyloxy)carbonyl]pyrrolidin-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (90.0 mg, 0.137 mmol) in DCM (0.4 mL) was added TFA (0.2 mL). The resulting mixture was stirred at rt for 30 min and concentrated *in vacuo* to dryness to give the title compound (107 mg, 0.137 mmol, 100% yield) as a brown gum which was used in the next step without further purification. LCMS (M+H)⁺ = 559.4.


[00308] Step 21-3, preparation of benzyl (3R)-3{5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate: to a solution of 2-(difluoromethyl)-4-fluorobenzoic acid (38.8 mg, 0.204 mmol) from Step 20-1 in DMF (0.2 mL) was added HATU (72.4 mg, 0.190 mmol) and DIEA (0.071 mL, 0.41 mmol). After stirring at rt for 5 min, this HATU-activated solution was added to a solution of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate *bis*(2,2,2-trifluoroacetate) (107 mg, 0.136 mmol) and DIEA (0.14 mL, 0.80 mmol) in DMF (0.2 mL). The resulting mixture was stirred at rt for 10 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (63.0 mg, 0.0862 mmol, 63.4% yield) as a light yellow solid. LCMS (M+H)⁺ = 731.3.

[00309] Step 21-4, preparation of 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-*N*-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to benzyl (3R)-3{5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate (63.0 mg, 0.0862 mmol) was added (methylsulfanyl)benzene (0.050 mL) and TFA (0.25 mL). The resulting mixture was heated at 60°C for 30 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (39.9 mg, 0.0669 mmol, 77.6% yield) as an off-white solid. LCMS (M+H)⁺ = 597.4.

[00310] The following compounds were prepared similarly to **Example 21** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-265	613.3
1-288	612.1
1-292	596.4

Example 22: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-263)



[00311] Step 22-1, preparation of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate: to a solution of *tert*-butyl (3R)-4-(6-{[(3R)-1-(benzyloxy)carbonyl]pyrrolidin-3-yl}carbonyl)-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate from **Example 21**, step 1 (27 mg, 0.041 mmol) in DCM (0.4 mL) was added TFA (0.1 mL). The resulting solution was stirred at rt for 1 h and concentrated under vacuum to remove the volatile solvent. The residue was dissolved in ACN (0.3 mL) and neutralized with TEA (16 μ L, $\sim$ 3eq.), which was used in the next step without further purification. LCMS (M+H)⁺ = 559.3.


[00312] Step 22-2, preparation of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate: to a solution of 1-(trifluoromethyl)cyclopentane-1-carboxylic acid (11 mg, 0.06 mmol) in ACN (0.5 mL) was added HATU (22 mg, 0.06 mmol) and TEA (11 μ L, 0.08 mmol). After stirring at rt for 5 min, the resulting solution was treated with benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate from step 1 (22 mg, 0.039 mmol). The reaction was stirred at rt for 30 min and directly purified by C18 reversed phase column chromatography to give the title compound (21 mg, 74% yield). LCMS (M+H)⁺ = 723.8.

[00313] Step 22-3, preparation of 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-amido}pyrrolidine-1-carboxylate (21 mg, 0.029 mmol) in TFA (0.5 mL) was added one drop of methyl(phenyl)sulfane. The resulting mixture was stirred at 60 °C for 1 hr and cooled to rt. After removal of the volatile solvent, the residue was purified by C18 reversed phase column chromatography to give the title compound (14 mg, 82 % yield) as a white powder. LCMS (M+H)⁺ = 589.4.

[00314] The following compounds were prepared similarly to **Example 22** with appropriate substituting reagents and substrates at different steps:

Compound no.	MS (M+H) ⁺
1-250	523.6
1-251	608.3
1-252	608.3
1-264	537.6
1-273	603.5
1-274	629.5
1-280	575.5
1-285	641.4
1-294	613.4
1-295	593.4
1-297	615.5
1-298	611.3
1-299	603.3
1-300	575.3
1-301	575.3
1-302	563.3
1-304	588.4
1-308	577.6
1-314	575.5
1-318	607.5
1-321	589.3
1-324	603.5
1-325	603.5
1-327	589.3
1-328	571.4
1-329	589.4
1-335	602.5
1-352	597.4
1-353	583.4
1-354	611.4
1-368	603.3

Example 23: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-255)




[00315] Step 23-1, preparation of 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 1-(trifluoromethyl)cyclopentane-1-carboxylic acid (13 mg, 0.071 mmol) in ACN (0.5 mL) was added HATU (27 mg, 0.071 mmol) and TEA (19 μ L, 0.14 mmol). After stirring at rt for 5 min, the resulting solution was treated with neutral 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide from **Example 20**, step 3 (20 mg, 0.045 mmol). The reaction was stirred at rt for 30 min and directly purified by C18 reversed phase column chromatography to give the title compound (19 mg, 70% yield). LCMS (M+H)⁺ = 603.6.

[00316] The following compounds were prepared similarly to **Example 23** with appropriate substituting reagents and substrates at different steps. Some examples may require a reductive amination to introduce the tert-amine moiety in the final step:

Compound no.	MS (M+H) ⁺
1-235	561.4
1-238	587.3
1-241	589.5
1-248	571.5
1-281	629.5
1-282	643.5
1-284	655.4
1-286	617.6
1-313	603.6
1-319	621.5
1-322	617.4
1-330	603.5
1-331	603.5
1-332	617.5
1-333	617.6
1-334	633.4
1-336	616.6
1-337	603.6
1-339	629.4
1-340	602.6
1-341	602.4
1-359	617.4
1-363	603.4
1-378	617.2
1-379	601.4
1-380	617.3
1-382	617.5
1-385	629.5
1-388	628.5
1-394	615.4
1-395	629.5

Compound no.	MS (M+H) ⁺
1-399	588.4
1-400	629.5
1-401	602.4
1-405	615.5
1-415	616.4
1-416	585.3
1-417	584.4

Example 24: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-283)




[00317] Step 24-1, preparation of 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 1-(trifluoromethyl)cyclobutane-1-carboxylic acid (632 mg, 3.76 mmol) in DMF (5.0 mL) was added HATU (1.43 g, 3.76 mmol) and TEA (0.67 mL, 5.02 mmol). After stirring at rt for 5 min, the resulting solution was treated with 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide from **Example 20**, step 3 (1.10 g, 2.51 mmol). The reaction was stirred at rt for 30 min and directly purified by C18 reversed phase column chromatography to give the title compound (0.85 g, 58% yield). LCMS (M+H)⁺ = 589.4.

[00318] The following compounds were prepared similarly to **Example 24** with appropriate substituting reagents and substrates at different steps. Some examples may require a reductive amination to introduce the tert-amine moiety in the final step:

Compound no.	MS (M+H) ⁺
1-344	588.4
1-349	602.5
1-358	597.2
1-364	589.4
1-371	633.4
1-390	589.4
1-411	615.4

Example 25: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide (compound 1-374)



[00319] Step 25-1, preparation of tert-butyl (3R)-4-(6-bromo-2-cyanopyridin-3-yl)-3-ethylpiperazine-1-carboxylate: Into a 1000-mL round-bottom flask, was placed 6-bromo-3-fluoropicolinonitrile (50.0 g, 249 mmol), tert-butyl (R)-3-ethylpiperazine-1-carboxylate (59 g, 275 mmol), DIEA (97 g, 750 mmol), and DMSO (500 mL). The resulting reaction mixture was stirred for 24 hrs at 90 °C. The reaction mixture was cooled to room temperature and quenched with water (1000 mL). The resulting solution was extracted with ethyl acetate (3x1000 mL). The organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in the title compound (100 g, 66%) as a yellow oil. LCMS (M+H)⁺ = 395.1.

[00320] Step 25-2, preparation of tert-butyl (3R)-4-[2-cyano-6-(2-ethoxyphenyl)pyridin-3-yl]-3-ethylpiperazine-1-carboxylate: Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl (3R)-4-(6-bromo-2-cyanopyridin-3-yl)-3-ethylpiperazine-1-carboxylate (5.0 g, 13 mmol), (2-ethoxyphenyl)boronic acid (4.0 g, 26 mmol), potassium carbonate (5.0 g, 36 mmol), Pd(DtBPF)Cl₂ (0.4 g, 0.62 mmol), and dioxane/water (50 mL/5 mL). The resulting solution was stirred at 80 °C for 1 hour and cooled to rt. The inorganic solid was filtered off, and the filtrate was concentrated. The residue was purified from a silica gel column eluting with ethyl acetate/petroleum ether (1:3). This resulted in the title compound (4.0 g, 67 %) as a yellowish oil. LCMS (M+H)⁺ = 437.2.

[00321] Step 25-3, preparation of 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridine-2-carboxylic acid: Into a 250-mL round-bottom flask purged, was placed tert-butyl (3R)-4-[2-cyano-6-(2-ethoxyphenyl)pyridin-3-yl]-3-ethylpiperazine-1-carboxylate (4.0 g, 9.2 mmol), EtOH (40 mL), and water (40 mL). This resulting solution was treated with KOH (5.3 g, 94 mmol) at rt and then stirred at 100 °C for 50 hrs. The reaction was cooled to rt and concentrated to remove EtOH. The pH of aqueous layer was adjusted to 6-7 with 3N-HCl. The resulting solution was extracted with ethyl acetate (3x50 mL). The combined organic layers were

washed with brine, dried over anhydrous sodium sulfate, and concentrated to afford the title compound (2.3 g, 55%) as a yellow solid. LCMS (M+H)⁺ = 456.2.

[00322] Step 25-4, preparation of ethyl 6-(2-ethoxyphenyl)-3-[(2R)-2-ethylpiperazin-1-yl]pyridine-2-carboxylate: Into a 50-mL round-bottom flask, was placed 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridine-2-carboxylic acid (1.0 g, 2.2 mmol), EtOH (10 mL), and sulfuric acid (0.5 mL). The resulting reaction mixture was stirred at 80 °C for 3 hours and cooled to rt. The reaction was diluted with water (50 ml) and then adjusted to pH 6~7 with sodium bicarbonate. The resulting solution was extracted with EtOAc (3x10 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, and concentrated under vacuum to afford the title compound (400 mg, 48 %) as a yellow oil. LCMS (M+H)⁺ = 384.2.

[00323] Step 25-5, preparation of ethyl 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]pyridine-2-carboxylate: Into a 8-mL vial, was placed (S)-2-(trifluoromethyl)pyrrolidine (75 mg, 0.54 mmol), DIEA (210 mg, 1.62 mmol), and DCM (1 mL). The reaction mixture was cooled to 0 °C and treated with bis(trichloromethyl) carbonate (64 mg, 0.22 mmol). The resulting mixture was stirred at 0 °C for 3 hours, quenched with water (20 mL), and extracted with DCM (3x20 mL). The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The activated pyrrolidine was used in the next step directly without further purification. Ethyl 6-(2-ethoxyphenyl)-3-[(2R)-2-ethylpiperazin-1-yl]pyridine-2-carboxylate (210 mg, 5.43 mmol), the activated pyrrolidine described above, and potassium carbonate (750 mg, 5.43 mmol), and ACN (2 mL) were placed into a 8-mL vial. The resulting reaction mixture was stirred at 30 °C for 2 hours and cooled to room temperature. The crude product was purified by a reverse phase C18 column chromatography to afford the title compound (200 mg, 68 %) as a yellow oil. LCMS (M+H)⁺ = 549.4.

[00324] Step 25-6, preparation of 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]pyridine-2-carboxylic acid: Into a 8-mL vial, was placed ethyl 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]pyridine-2-carboxylate (200 mg, 0.365 mmol), LiOH (88 mg, 3.7 mmol), EtOH (2 mL), and water (1 mL). The resulting reaction mixture was stirred at 60 °C for 1 h and cooled to rt. The reaction was diluted with water (10 ml), and then adjusted to pH 6~7 with 3N-HCl. The resulting solution was extracted with EtOAc (3x10 mL). The combined organic layers were washed with brine, dried over anhydrous sodium

sulfate, and concentrated to afford the title compound (170 mg, 90%) as a yellow oil. LCMS $(M+H)^+ = 521.6$.


[00325] Step 25-7, preparation of 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide formate: Into a 50-mL round-bottom flask, was placed 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]pyridine-2-carboxylic acid (80 mg, 0.15 mmol), HATU (58 mg, 0.15 mmol), DIEA (40 mg, 0.30 mmol), and DMF (1 mL). The resulting reaction mixture was stirred for 10 min at rt, and then treated with (R)-1-methylpyrrolidin-3-amine (20 mg, 0.20 mmol). The resulting reaction mixture was stirred at rt for 1 h. The crude product was purified by Prep-HPLC with the following conditions (Prep-HPLC-013): Column, Atlantis Prep T3 OBD Column, 19*150mm 5um; mobile phase, Water (0.1%FA) and ACN (30% Phase B up to 90% in 8min); Total flow 20 mL/min. Detector, UV 220 nm. This resulted in the title compound (62 mg, 62%) as a white solid. LCMS $(M+H)^+ = 603.4$.

[00326] The following compounds were prepared similarly to **Example 25** with appropriate substituting reagents and substrates at different steps. Some examples may require a chiral separation to get optically pure compounds, but their absolute stereochemistries were not determined. In some examples, an additional deprotection is required in the final step:

Compound no.	MS $(M+H)^+$
1-305	577.4
1-306	590.3
1-310	590.4
1-311	562.4
1-315	563.4
1-316	563.4
1-317	570.5
1-320	582.2
1-323	578.4
1-326	604.4
1-342	589.4
1-343	589.4
1-345	598.4
1-346	604.5
1-347	604.4
1-350	588.4
1-355	577.4
1-356	577.4
1-360	584.3
1-361	638.3
1-362	603.3
1-365	595.3

Compound no.	MS (M+H) ⁺
1-372	604.4
1-373	604.3
1-375	603.4
1-381	630.6
1-383	609.3
1-386	604.4
1-387	603.4
1-389	652.5
1-391	604.4
1-393	608.2
1-402	651.5
1-404	609.3
1-409	595.5
1-412	595.6
1-413	596.6
1-414	594.5
1-418	638.5
1-419	636.4
1-420	622.3
1-421	635.3
1-422	613.4
1-423	627.4
1-424	612.3
1-426	626.4
1-427	618.4
1-433	617.4
1-434	618.4
1-435	617.3


Example 26: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-[(5-methyl-2-oxo-2H-1,3-dioxol-4-yl)methyl]pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-307)



[00327] Step 26-1, preparation of 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-[(5-methyl-2-oxo-2H-1,3-dioxol-4-yl)methyl]pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: To a heterogeneous solution of **Example 22** (50 mg, 0.085 mmol) and potassium hydrogen carbonate (10 mg, 0.10 mmol) in DMF (1.5 mL) was slowly added 4-(bromomethyl)-5-methyl-1,3-dioxol-2-one (20 mg,

0.10 mmol) at rt. The resulting reaction was stirred at rt for 16 hours. The inorganic solid was filtered off and then the filtrate was purified from a reverse phase C18 column chromatography to afford the title compound (18 mg, 30%). LCMS (M+H)⁺ = 701.4.

Example 27: (2S*)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (compound 1-367) [*absolute stereochemistry not determined]



[00328] Step 27-1, preparation of (2S*)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate: to a solution of 1,1,1-trifluoro-3,3-dimethylbutan-2-ol (100 mg, 0.640 mmol) in THF (3.0 mL) was added 1-(1H-imidazole-1-carbonyl)-1H-imidazole (156 mg, 0.961 mmol). The mixture was stirred at rt for 2.5 days. The mixture was concentrated to dryness and the residue was purified by silica gel column chromatography to give the racemic mixture of title compound (145 mg, 0.578 mmol, 90.3% yield) as a white solid. LCMS (M+H)⁺ = 251.1. The racemic mixture was purified by following conditions to give the enantiomerically pure title compound (2nd enantiomer: tentative assignment).

Chiral Resolution Conditions:

Column: normal phase Daicel Chiralpak ID column, 20 mm ID X 250 mm L, 5 µm particle size

Gradient: isocratic 5% EtOH in Hexanes

Flow rate: 20 mL/min

Retention Times: 1st enantiomer: 1.22 min (absolute stereochemistry not determined; tentatively assigned as compound 1-366); 2nd enantiomer: 1.52 min (absolute stereochemistry not determined; tentatively assigned as compound 1-367).

[00329] Step 27-2, preparation of benzyl (3R)-3-{2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-amido}pyrrolidine-1-carboxylate: to a solution of *tert*-butyl (3R)-4-(6-{[(3R)-1-[(benzyloxy)carbonyl]pyrrolidin-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (181 mg, 0.274 mmol) from Step 21-1 in DCM (0.7 mL) was added TFA (0.38 mL, 4.9 mmol). The resulting mixture was stirred at rt for 30 min and concentrated *in vacuo* to dryness. The crude TFA salt was neutralized with saturated NaHCO₃

(aq) to give the title compound (153 mg, 0.274 mmol, 100% yield) as a brown solid. LCMS $(M+H)^+ = 559.3$.

[00330] Step 27-3, preparation of benzyl (2*S**)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3*R*)-4-(6-[[[(3*R*)-1-[(benzyloxy)carbonyl]pyrrolidin-3-yl]carbamoyl]-2'-ethoxy-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate: to a solution of benzyl (3*R*)-3-{2'-ethoxy-5-[(2*R*)-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-amido}pyrrolidine-1-carboxylate (110 mg, 0.197 mmol) in THF (5.0 mL) was added DIEA (0.270 mL, 1.55 mmol) and (2*S*)-1,1,1-trifluoro-3,3-dimethylbutane-2-yl 1*H*-imidazole-1-carboxylate (75 mg, 0.30 mmol) in a sealed tube. The mixture was heated at 100°C for 24 h. The mixture was concentrated to dryness and the residue was purified by C18 reversed phase column chromatography to give the title compound (120 mg, 0.162 mmol, 82.2% yield) as yellow oil. LCMS $(M+H)^+ = 741.4$.


[00331] Step 27-4, preparation of (2*S**)-1,1,1-trifluoro-3,3-dimethylbutane-2-yl (3*R*)-4-(2'-ethoxy-6-[[[(3*R*)-pyrrolidin-3-yl]carbamoyl]-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate formate: to benzyl (2*S**)-1,1,1-trifluoro-3,3-dimethylbutane-2-yl (3*R*)-4-(6-[[[(3*R*)-1-[(benzyloxy)carbonyl]pyrrolidin-3-yl]carbamoyl]-2'-ethoxy-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate (120 mg, 0.162 mmol) was added TFA (2.0 mL). The resulting mixture was heated at 60°C for 2 h. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (72.0 mg, 0.110 mmol, 67.9% yield) as a white solid. LCMS $(M+H)^+ = 597.4$.

[00332] The following compounds were prepared similarly to **Example 27** with appropriate substituting reagents and substrates at different steps. Some examples' absolute stereochemistries were not determined. In some examples, an additional deprotection is not required in the final step:

Compound no.	MS $(M+H)^+$
1-309	579.4
1-312	607.3
1-338	590.4
1-348	606.3
1-351	627.3
1-366*	607.4
1-369	619.3
1-377	627.2
1-396*	621.4
1-403*	647.4
1-407*	606.3

* absolute stereochemistry not determined

Example 28: (2*S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3*R*)-4-(2'-ethoxy-6-{[(3*R*)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate (compound 1-392) [*absolute stereochemistry not determined]**




[00333] Step 28-1, preparation of (2*S**)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3*R*)-4-(2'-ethoxy-6-{[(3*R*)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate: to a solution of (2*S*)-1,1,1-trifluoro-3,3-dimethylbutane-2-yl (3*R*)-4-(2'-ethoxy-6-{[(3*R*)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate formate (40.0 mg, 0.0613 mmol) in MeOH (3.0 mL) was added paraformaldehyde (38.5 mg) and sodium cyanoborohydride (38.5 mg, 0.613 mmol). The resulting mixture was stirred at rt for 1 h. The reaction mixture was quenched by TFA (0.1 mL) and re-basified with saturated NaHCO₃ (aq). The mixture was extracted with DCM and the organic layer was concentrated *in vacuo* to dryness. The residue was purified by C18 reversed phase column chromatography to give the title compound (26.1 mg, 0.0420 mmol, 68.5% yield) as an off-white solid. LCMS (M+H)⁺ = 621.5.

[00334] The following compounds were prepared similarly to **Example 28** with appropriate substituting reagents and substrates at different steps. Some examples' absolute stereochemistries were not determined:

Compound no.	MS (M+H) ⁺
1-370	621.4
1-376	621.5
1-384	613.3
1-397	579.4
1-406*	621.5
1-408	629.6

* absolute stereochemistry not determined

Example 29: 5-[(2*R*)-4-(7-cyano-5-fluoro-1*H*-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-438)



[00335] Step 29-1, preparation of 5-fluoro-1*H*-indole-7-carbonitrile: to a mixture of 7-bromo-5-fluoro-1*H*-indole (920 mg, 4.30 mmol), dicyanozinc (1.514 g, 12.89 mmol) and Pd(PPh₃)₄ (993 mg, 0.860 mmol) in a sealed tube was added DMF (10 mL). Nitrogen (g) was bubbled through the reaction mixture. The resulting mixture was heated at 120°C for 16 h. The mixture was quenched by water and extracted with DCM (3X). The combined organics were concentrated to dryness and the residue was purified by C18 reversed phase column chromatography to give the title compound (676 mg, 4.22 mmol, 98 % yield) as an off-white solid.


[00336] Step 29-2, preparation of 4-nitrophenyl 7-cyano-5-fluoro-1*H*-indole-1-carboxylate: to a suspension of 5-fluoro-1*H*-indole-7-carbonitrile (306 mg, 1.91 mmol) and 4-nitrophenyl carbonochloridate (463 mg, 2.30 mmol) in THF (20 mL) was added TEA (0.80 mL, 5.74 mmol) and DMAP (23.4 mg, 0.191 mmol). The resulting mixture was stirred at rt for 2 h but the reaction was not complete. More 4-nitrophenyl carbonochloridate (463 mg, 2.30 mmol) and triethylamine (0.80 mL, 5.73 mmol) were added and the reaction was continued for 2 h. The mixture was concentrated to dryness and the residue was purified by silica gel column chromatography to give the title compound (411 mg, 1.26 mmol, 66 % yield) as an off-white solid.

[00337] Step 29-3, preparation of 5-[(2*R*)-4-(7-cyano-5-fluoro-1*H*-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 2'-ethoxy-5-[(2*R*)-2-ethylpiperazin-1-yl]-*N*-[(3*R*)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide from **Example 20**, Step 3 (50.0 mg, 0.114 mmol) and 4-nitrophenyl 7-cyano-5-fluoro-1*H*-indole-1-carboxylate (44.5 mg, 0.137 mmol) in MeCN (1 mL) was added DMAP (34.8 mg, 0.285 mmol). The mixture was heated at 50°C for 1 h. The mixture was purified by C18 reversed phase column chromatography to give the title compound (14.5 mg, 0.023 mmol, 20 % yield) as a light yellow solid. LCMS (M+H)⁺ = 625.4.

[00338] The following compounds were prepared similarly to **Example 29** with appropriate substituting reagents and substrates at different steps.


Compound no.	MS (M+H) ⁺
1-425	634.1
1-428	616.2
1-429	607.4
1-430	633.3
1-431	615.2
1-432	606.3
1-436	650.5
1-437	649.4
1-439	624.3

Example 30: 5-[(2R)-4-[2-chloro-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide (compound 1-261)



[00339] Step 30-1, preparation of 5-[(2R)-4-[2-chloro-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide: to a solution of 2-chloro-4-fluorobenzoic acid (35.9 mg, 0.206 mmol) and HATU (73.0 mg, 0.192 mmol) in DMF (0.2 mL) was added DIEA (0.072 mL, 0.41 mmol). After stirring at rt for 5 min, this HATU-activated solution was added to a solution of 2'-ethoxy-5-[(2R)-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide *tris*(2,2,2-trifluoroacetate) from **Example 20**, Step 3 (107 mg, 0.137 mmol) and DIEA (0.072 mL, 0.41 mmol) in DMF (0.2 mL). The resulting mixture was stirred at rt for 10 min. The reaction mixture was purified by C18 reversed phase column chromatography to give the title compound (39.4 mg, 0.0645 mmol, 47% yield) as a light yellow solid. LCMS (M+H)⁺ = 595.5.

Example 31: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxamide (compound 1-410)



[00340] Step 31-1, preparation of 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxylic acid: to a solution of 3-[(2R)-4-[(tert-butoxy)carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridine-2-carboxylic acid (450 mg, 0.98 mmol) from **Example 25**, step 3 in DCM (5.0 mL) was added TFA (1.14 mL, 14.8 mmol) at rt. The resulting solution was stirred at rt for 1 h and concentrated under vacuum to afford 6-(2-ethoxyphenyl)-3-[(2R)-2-ethylpiperazin-1-yl]pyridine-2-carboxylic acid. This residue was dissolved in ACN (2 mL) and neutralized with Et₃N (~0.3 mL). The solution was used in the next HATU coupling step without further purification. To a solution of 1-(trifluoromethyl)cyclobutane-1-carboxylic acid (332 mg, 1.98 mmol) in ACN (5 mL) was added HATU (751 mg, 1.98 mmol) and followed by Et₃N (0.26 mL, 1.98 mmol) at rt. After stirring at rt for 5 min, this HATU-activated solution was treated with the solution of 6-(2-ethoxyphenyl)-3-[(2R)-2-ethylpiperazin-1-yl]pyridine-2-carboxylic acid described above. The resulting mixture was stirred at rt for 30 min and concentrated under vacuum. The residue was purified by C18 reversed phase column chromatography to give the title compound (290 mg, 58% yield). LCMS (M+H)⁺ = 506.3.

[00341] Step 31-2, preparation of N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxamide: to a solution of 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxylic acid (70 mg, 0.14 mmol) and HATU (58.0 mg, 0.15 mmol) in DMF (1.5 mL) was added Et₃N (0.074 mL, 0.55 mmol). After stirring at rt for 5 min, the resulting solution was treated with (S)-quinuclidin-3-amine dihydrochloride (33 mg, 0.17 mmol). The resulting mixture was stirred at rt for 1 hr and directly purified by C18 reversed phase column chromatography to give the title compound (40 mg, 47% yield). LCMS (M+H)⁺ = 614.3.

Example A-1: Parenteral Pharmaceutical Composition

[00342] To prepare a parenteral pharmaceutical composition suitable for administration by injection (subcutaneous, intravenous), 1-100 mg of a water-soluble salt of a compound Formula (I), or a pharmaceutically acceptable salt or solvate thereof, is dissolved in sterile water and then mixed with 10 mL of 0.9% sterile saline. A suitable buffer is optionally added as well as optional acid or base to adjust the pH. The mixture is incorporated into a dosage unit form suitable for administration by injection

Example A-2: Oral Solution

[00343] To prepare a pharmaceutical composition for oral delivery, a sufficient amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is added to water (with optional solubilizer(s), optional buffer(s) and taste masking excipients) to provide a 20 mg/mL solution.

Example A-3: Oral Tablet

[00344] A tablet is prepared by mixing 20-50% by weight of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, 20-50% by weight of microcrystalline cellulose, 1-10% by weight of low-substituted hydroxypropyl cellulose, and 1-10% by weight of magnesium stearate or other appropriate excipients. Tablets are prepared by direct compression. The total weight of the compressed tablets is maintained at 100 -500 mg.

Example A-4: Oral Capsule

[0001] To prepare a pharmaceutical composition for oral delivery, 10-500 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is mixed with starch or other suitable powder blend. The mixture is incorporated into an oral dosage unit such as a hard gelatin capsule, which is suitable for oral administration.

[00345] In another embodiment, 10-500 mg of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is placed into Size 4 capsule, or size 1 capsule (hypromellose or hard gelatin) and the capsule is closed.

Example A-5: Topical Gel Composition

[00346] To prepare a pharmaceutical topical gel composition, a compound of Formula (I), or a pharmaceutically acceptable salt thereof, is mixed with hydroxypropyl cellulose, propylene glycol, isopropyl myristate and purified alcohol USP. The resulting gel mixture is then incorporated into containers, such as tubes, which are suitable for topical administration.

Example B-1: MC2R assays**Membrane preparation:**

Crude membrane fractions are prepared from CRE-bla-CHO-K1 cells stably expressing hMC2 receptor and hMRAP accessory protein (Thermo Fisher). The cells are grown to 85 - 100% confluence on standard tissue culture dishes in GlutaMax DMEM growth media (Gibco) with following additives: 10% dialyzed FBS (Gemini), 0.1 mM NEAA (Gibco), 25 mM HEPES

(Gibco), 5 µg/mL blasticidin (Goldbio), 100 µg/mL zeocin (Invitrogen), 600 µg/mL hygromycin (Goldbio). To prepare membranes, cells are scraped and collected in 1X Dulbecco's phosphate buffered saline (Corning) and then pelleted at 1000 RPM's. The cell pellet is reconstituted in membrane preparation buffer (20 mM HEPES, 6 mM MgCl₂ and 1 mM EGTA, protease inhibitor tablets (Pierce) adjusted to pH 7.4), homogenized using a dounce homogenizer, and the resulting membrane fraction is pelleted by centrifugation at 12,000 RPM's. The membrane pellet is resuspended in membrane preparation buffer, snap frozen and stored at -80°C for later use.

Binding assay for hMC2 antagonists protocol:

The hMC2 membrane binding assay utilizes the following components: radiolabel [¹²⁵I]ACTH (1-39) Tyr23 (PerkinElmer), wheatgerm agglutinin coated PVT SPA beads (PerkinElmer), crude hMC2R membranes, and compounds. Briefly hMC2R membranes are incubated with SPA beads in binding assay buffer (50mM HEPES, 5mM MgCl₂, 1mM CaCl₂, 0.2%BSA, protease inhibitor tablets (Pierce) adjusted to pH7.4) prior to assay initiation. A dose response of compound (the final concentration of compound are typically 0-10,000nM), SPA membranes, and [¹²⁵I]ACTH (1-39) Tyr23 at a final concentration of 0.2 nM is plated in a 96-well assay plate and allowed to incubate 1.5 hours at room temperature. Assay plates are read using a Top Count NXT and K_i values for compounds are determined using a GraphPad Prism 6 non-linear regression analysis.

[00347] Illustrative binding affinities of selective compounds are described in Table A. The potencies are divided into four criteria: + means that K_i is between 1,000 nM and 10,000 nM; ++ means that K_i is between 100 nM and 1,000 nM; +++ means K_i is between 10 nM and 100 nM; ++++ means K_i is below 10 nM.

Table A

Compound No.	MC2R binding potency
1-1	+++
1-2	++++
1-3	++++
1-4	+++
1-5	+++
1-6	+++
1-7	+++
1-8	+++
1-9	+++
1-11	+++

Compound No.	MC2R binding potency
1-15	++++
1-16	+++
1-17	++++
1-18	++++
1-19	+++
1-20	+++
1-23	+++
1-24	++++
1-25	++++
1-26	++++
1-27	++++
1-32	+++
1-33	+++
1-34	++++
1-35	++++
1-36	++++
1-37	++++
1-38	++++
1-40	++++
1-41	++++
1-42	++++
1-43	++++
1-44	++++
1-45	++++
1-46	++++
1-47	++++
1-48	++++
1-49	++++
1-50	++++
1-51	++++
1-52	++++
1-54	++++
1-57	++++
1-58	++++
1-59	++++
1-60	++++
1-62	++++

Compound No.	MC2R binding potency
1-67	++++
1-68	++++
1-69	++++
1-71	++++
1-72	++++
1-73	++++
1-76	++++
1-82	++++
1-84	++++
1-87	++++
1-88	++++
1-90	++++
1-91	++++
1-92	++++
1-93	++++
1-94	++++
1-95	++++
1-96	++++
1-97	++++
1-98	++++
1-99	++++
1-100	++++
1-101	++++
1-102	++++
1-104	++++
1-107	++++
1-108	++++
1-109	++++
1-110	++++
1-111	++++
1-112	++++
1-113	++++
1-114	++++
1-115	++++
1-116	++++
1-117	++++
1-123	++++

Compound No.	MC2R binding potency
1-124	++++
1-125	++++
1-136	++++
1-137	++++
1-138	++++
1-142	++++
1-143	++++
1-148	++++
1-149	++++
1-150	++++
1-151	++++
1-152	++++
1-155	++++
1-156	++++
1-163	++++
1-164	++++
1-165	++++
1-167	++++
1-168	++++
1-169	++++
1-170	++++
1-171	++++
1-173	++++
1-174	++++
1-176	++++
1-177	++++
1-178	++++
1-181	++++
1-182	++++
1-183	++++
1-184	++++
1-187	++++
1-188	++++
1-189	++++
1-190	++++
1-198	++++
1-199	++++

Compound No.	MC2R binding potency
1-200	++++
1-201	++++
1-202	++++
1-203	++++
1-206	++++
1-207	++++
1-208	++++
1-209	++++
1-210	++++
1-211	++++
1-212	++++
1-213	++++
1-214	++++
1-215	++++
1-216	++++
1-219	++++
1-220	++++
1-221	++++
1-222	++++
1-223	++++
1-224	++++
1-225	++++
1-226	++++
1-227	++++
1-228	++++
1-229	++++
1-230	++++
1-231	++++
1-232	++++
1-233	++++
1-234	++++
1-236	++++
1-237	++++
1-238	++++
1-239	++++
1-240	++++
1-241	++++

Compound No.	MC2R binding potency
1-242	++++
1-243	++++
1-244	++++
1-245	++++
1-246	++++
1-247	++++
1-249	++++
1-253	++++
1-254	++++
1-255	++++
1-256	++++
1-257	++++
1-258	++++
1-259	++++
1-260	++++
1-261	++++
1-262	++++
1-263	++++
1-264	++++
1-265	++++
1-266	++++
1-267	++++
1-270	++++
1-271	++++
1-273	++++
1-274	++++
1-275	++++
1-276	++++
1-277	++++
1-278	++++
1-279	++++
1-280	++++
1-282	++++
1-283	++++
1-286	++++
1-287	++++
1-288	++++

Compound No.	MC2R binding potency
1-289	++++
1-290	++++
1-291	++++
1-292	++++
1-293	++++
1-296	++++
1-297	++++
1-298	++++
1-299	++++
1-304	++++
1-305	++++
1-306	++++
1-312	++++
1-313	++++
1-314	++++
1-315	++++
1-320	++++
1-321	++++
1-324	++++
1-325	++++
1-326	++++
1-327	++++
1-328	++++
1-329	++++
1-330	++++
1-331	++++
1-332	++++
1-333	++++
1-334	++++
1-335	++++
1-336	++++
1-337	++++
1-338	++++
1-339	++++
1-340	++++
1-341	++++
1-342	++++

Compound No.	MC2R binding potency
1-343	++++
1-344	++++
1-345	++++
1-346	++++
1-348	++++
1-349	++++
1-350	++++
1-351	++++
1-352	++++
1-353	++++
1-354	++++
1-356	++++
1-357	++++
1-358	++++
1-360	++++
1-361	++++
1-362	++++
1-363	++++
1-365	++++
1-367*	++++
1-368	++++
1-371	++++
1-372	++++
1-373	++++
1-374	++++
1-375	++++
1-376	++++
1-377	++++
1-378	++++
1-380	++++
1-381	++++
1-382	++++
1-383	++++
1-385	++++
1-386	+++
1-387	++++
1-388	++++

Compound No.	MC2R binding potency
1-389	++++
1-390	+++
1-391	++++
1-392*	++++
1-393	++++
1-394	++++
1-395	++++
1-396*	++++
1-397	++++
1-398	++++
1-399	++++
1-400	++++
1-401	++++
1-402	++++
1-403*	++++
1-404	++++
1-405	++++
1-406*	++++
1-407*	++++
1-408	++++
1-409	++++
1-410	++++
1-411	++++
1-412	++++
1-413	++++
1-414	++++
1-415	++++
1-416	++++
1-417	++++
1-418	++++
1-419	++++
1-420	++++
1-421	++++
1-422	++++
1-423	++++
1-424	++++
1-425	++++

Compound No.	MC2R binding potency
1-426	++++
1-427	++++
1-428	++++
1-429	++++
1-430	++++
1-431	++++
1-432	++++
1-433	++++
1-434	++++
1-435	++++
1-436	++++
1-437	++++
1-438	++++
1-439	++++
2-2	++++
2-3	+++
2-13	++++
2-16	++++
2-20	++++
2-21	++++
2-25	++++
2-30	++++
2-31	++++

* absolute stereochemistry not determined

[00348] The examples and embodiments described herein are for illustrative purposes only and various modifications or changes suggested to persons skilled in the art are to be included within the spirit and purview of this application and scope of the appended claims.

WHAT IS CLAIMED IS:

-

wherein:

R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, unsubstituted or substituted C₃-C₆ cycloalkyl, wherein any substituted group of R^a is substituted with one or more R⁹ groups;

R^b and R^c are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^b and R^c is substituted with one or more R⁹ groups;

R^B is unsubstituted or substituted carbocycle, unsubstituted or substituted heterocycle, unsubstituted or substituted C₁-C₇ alkyl, unsubstituted or substituted C₁-C₇ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein if R^B is substituted then R^B is substituted with R^d, R^e and R^f;

-198-


R^e and R^f are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R⁷)₂, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted C₁-C₆ fluoroalkyl, unsubstituted or substituted C₁-C₆ heteroalkyl, wherein any substituted group of R^d is substituted with one or more R^9 groups;


X^1 is N or CR⁶;

R^2 and R^3 are hydrogen;

or R^2 and R^3 are taken together with the carbon atom to which they are attached to form -C(=O)-;

R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted -(C₁-C₆ alkyl)-carbocycle, or unsubstituted or substituted -(C₁-C₆ alkyl)-heterocycle, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic heterocycle, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂;





R^6 is H or F;

each R^7 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, unsubstituted or substituted heteroaryl; or two R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^8 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted aryl, and unsubstituted or substituted heteroaryl;

each R^9 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkyl, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, -CN, -OH, -CO₂ R^8 , -CH₂CO₂ R^8 , -C(=O)N(R^7)₂, -C(=O)N(R^7)OR⁸, -CH₂C(=O)N(R^7)₂, -N(R^7)₂, -CH₂N(R^7)₂, -C(R^8)₂N(R^7)₂, -NR⁷C(=O) R^8 , -CH₂NR⁷C(=O) R^8 , -NR⁷C(=O)N(R^7)₂, -NR⁷C(=O)N(R^7)₂, C(R^8)=N(R^7)-OR⁸, -SR⁸, -S(=O) R^{10} , -SO₂ R^{10} , or -SO₂N(R^7)₂; and

each R^{10} is independently selected from the group consisting of unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, unsubstituted or substituted phenyl, and unsubstituted or substituted heteroaryl.

2. The compound of claim 1, or a pharmaceutically acceptable salt, or solvate thereof, wherein: R^a is selected from the group consisting of hydrogen, halogen, -OR⁸, -CN, -N(R^7)₂, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein any substituted group of R^a is substituted with one or more R^9 groups; and

R^b and R^c are each independently selected from the group consisting of hydrogen, halogen, -OR⁸, -N(R⁷)₂, and unsubstituted or substituted C₁-C₆ alkyl, wherein any substituted group of R^b and R^c is substituted with one or more R⁹ groups.

3. The compound of claim 1 or claim 2, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^a is -OR⁸; and

R^b and R^c are each hydrogen.


4. The compound of any one of claims 1-3, or a pharmaceutically acceptable salt, or solvate thereof, wherein:


R^4 is selected from the group consisting of hydrogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic carbocycle, or unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-4 N atoms and 0 or 1 O or S atoms, -N(R⁷)₂, -OR⁸, -CN, -CO₂R⁸, -C(=O)N(R⁷)₂, -SR⁸, -S(=O)R¹⁰, -S(=O)₂R¹⁰, -NR⁷C(=O)R⁸, -NR⁷SO₂R¹⁰, -SO₂R¹⁰, or -SO₂N(R⁷)₂.

5. The compound of any one of claims 1-4, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^4 is an unsubstituted or substituted monocyclic 4-, 5- or 6-membered heterocycle containing 1-2 N atoms and 0 or 1 O or S atoms, wherein any substituted group of R^4 is substituted with one or more halogen, unsubstituted or substituted C₁-C₆ alkyl, -N(R⁷)₂, -OR⁸, or -CN.

6. The compound of any one of claims 1-5, or a pharmaceutically acceptable salt, or solvate thereof, wherein:





7. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^B is an unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted bicyclic carbocycle, unsubstituted or substituted polycyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, unsubstituted or substituted bicyclic heterocycle, unsubstituted or substituted polycyclic heterocycle, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .


8. The compound of any one of claims 1-7, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^B is an unsubstituted or substituted phenyl, unsubstituted or substituted naphthyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted monocyclic 5-membered heteroaryl, unsubstituted or substituted bicyclic heteroaryl, monocyclic C_3 - C_8 cycloalkyl, unsubstituted or substituted bridged C_5 - C_{10} cycloalkyl, unsubstituted or substituted monocyclic C_2 - C_8 heterocycloalkyl, or unsubstituted or substituted bridged C_5 - C_{10} heterocycloalkyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

9. The compound of any one of claims 1-8, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^B is unsubstituted or substituted phenyl, unsubstituted or substituted monocyclic 6-membered heteroaryl, unsubstituted or substituted monocyclic 5-membered heteroaryl, unsubstituted or substituted bicyclic heteroaryl, monocyclic C_3 - C_8 cycloalkyl, or unsubstituted or substituted monocyclic C_2 - C_8 heterocycloalkyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .


10. The compound of any one of claims 1-9, or a pharmaceutically acceptable salt, or solvate thereof, wherein:



11. The compound of any one of claims 1-9, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^B is unsubstituted or substituted cyclopropyl, unsubstituted or substituted cyclobutyl, unsubstituted or substituted cyclopentyl, unsubstituted or substituted cyclopentenyl, or unsubstituted or substituted cyclohexyl, wherein if R^B is substituted then R^B is substituted with R^d , R^e and R^f .

12. The compound of any one of claims 1-9, or a pharmaceutically acceptable salt, or solvate thereof, wherein:



13. The compound of any one of claims 1-12, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

R^d is selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, unsubstituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted C_1 - C_6 heteroalkyl, wherein any substituted group of R^d is substituted with one or more R^9 groups;

R^e and R^f are independently selected from the group consisting of hydrogen, halogen, $-OR^8$, $-CN$, $-N(R^7)_2$, and unsubstituted or substituted C_1 - C_6 alkyl, wherein any substituted group of R^e and R^f is substituted with one or more R^9 groups.

14. The compound of any one of claims 1-13, or a pharmaceutically acceptable salt, or solvate thereof, wherein:

each R^7 is independently selected from the group consisting of hydrogen, substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl;


or two R^7 are taken together with the nitrogen atom to which they are attached to form an unsubstituted or substituted 3- to 6-membered monocyclic heterocycle;

each R^8 is independently selected from the group consisting of hydrogen, substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, and unsubstituted or substituted C_1 - C_6 fluoroalkyl;

each R^9 is independently hydrogen, halogen, unsubstituted or substituted C_1 - C_4 alkyl, unsubstituted or substituted C_1 - C_4 alkoxy, unsubstituted or substituted C_1 - C_4 fluoroalkyl, unsubstituted or substituted C_1 - C_4 fluoroalkoxy, unsubstituted or substituted monocyclic carbocycle, unsubstituted or substituted monocyclic heterocycle, $-CN$, $-OH$, $-CO_2R^8$, $-CH_2CO_2R^8$, $-C(=O)N(R^7)_2$, $-CH_2C(=O)N(R^7)_2$, $-N(R^7)_2$, $-CH_2N(R^7)_2$, $-NR^7C(=O)R^8$, or $-CH_2NR^7C(=O)R^8$; and


each R^{10} is independently selected from the group consisting substituted or substituted C_1 - C_6 alkyl, unsubstituted or substituted C_3 - C_6 cycloalkyl, unsubstituted or substituted C_1 - C_6 fluoroalkyl, and unsubstituted or substituted phenyl.


15. The compound of claim 1, or a pharmaceutically acceptable salt, or solvate thereof, wherein the compound of Formula (IX) has the structure of Formula (IXa), or a pharmaceutically acceptable salt, or solvate thereof:



Formula (IXa).

16. The compound of claim 15, or a pharmaceutically acceptable salt, or solvate thereof, wherein:





17. The compound of claim 1, wherein the compound is:

1-1: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide;

1-2: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methoxy-[1,1'-biphenyl]-3-carboxamide;

1-3: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-4: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-fluoro-[1,1'-biphenyl]-3-carboxamide;

1-5: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethyl-[1,1'-biphenyl]-3-carboxamide;

1-6: N-(2-aminoethyl)-2'-chloro-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-carboxamide;

1-7: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyano-[1,1'-biphenyl]-3-carboxamide;

1-8: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methyl-[1,1'-biphenyl]-3-carboxamide;

1-9: N-(2-aminoethyl)-2'-chloro-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3'-fluoro-[1,1'-biphenyl]-3-carboxamide;

1-10: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-11: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propoxy-[1,1'-biphenyl]-3-carboxamide;

1-12: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-(propan-2-yloxy)-[1,1'-biphenyl]-3-carboxamide;

1-13: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-hydroxy-[1,1'-biphenyl]-3-carboxamide;

1-14: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-cyclopropoxy-[1,1'-biphenyl]-3-carboxamide;

1-22: N-(2-aminoethyl)-2'-(aminomethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-3-carboxamide;

1-23: N-(3-aminopropyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-24: N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-25: N-(2-aminoethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-26: N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-3'-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-27: N-(2-aminoethyl)-4-[(2R)-4-[4'-(aminomethyl)-3-fluoro-5-methyl-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

1-28: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propyl-[1,1'-biphenyl]-3-carboxamide;

1-29: N-(2-aminoethyl)-4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-4'-propyl-[1,1'-biphenyl]-3-carboxamide;

1-31: N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(pyridin-3-yl)benzamide;

1-34: N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-35: N-(2-aminoethyl)-2-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(4-ethoxypyridin-3-yl)benzamide;

1-36: N-(2-aminoethyl)-2-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-37: N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-38: N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-39: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(pyridin-4-yl)benzoyl]piperazin-1-yl]benzamide;

1-40: N-(2-aminoethyl)-2-[(2R)-4-{2-[(3R)-3-(aminomethyl)pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-41: N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-(aminomethyl)pyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-42: N-(2-aminoethyl)-2-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-43: N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-44: N-(2-aminoethyl)-2-[(2R)-4-[4'-(aminomethyl)-3-fluoro-5-methyl-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-45: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-46: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;

1-47: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-48: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;

1-49: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(pyrrolidin-1-yl)ethyl]benzamide;

1-50: N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-(trifluoromethyl)benzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-51: N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-fluorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-52: N-(2-aminoethyl)-2-[(2R)-4-[2-(4-aminopiperidin-1-yl)-4-chlorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-53: N-(2-aminoethyl)-2-[(2R)-4-(3,5-dichloropyridine-2-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-54: N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-methoxybenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-55: N-(2-aminoethyl)-2-[(2R)-4-(2-chloro-4-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-56: N-(2-aminoethyl)-2-[(2R)-4-{4-[3-(aminomethyl)pyrrolidin-1-yl]-2-fluorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-57: N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-aminopyrrolidin-1-yl]-4-chlorobenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-58: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;

1-59: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;

1-60: N-(2-aminoethyl)-2-[(2R)-4-(2,4-dimethylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-61: N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(pyrrolidin-1-yl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-62: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]benzamide;

1-63: N-(2-aminoethyl)-2-[(2R)-4-(2-cyclopropylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-64: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-2-yl)methyl]benzamide;

1-65: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-3-yl)methyl]benzamide;

1-66: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(pyridin-4-yl)methyl]benzamide;

1-67: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-methyl-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]benzamide;

1-68: N-(2-aminoethyl)-2-[(2R)-4-{2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]-4-methylbenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-69: N-(2-aminoethyl)-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-70: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-(pyridin-4-yl)benzamide;

1-71: N-(2-aminoethyl)-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-methoxypyridin-3-yl)benzamide;

1-72: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-73: 6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-74: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(piperidin-4-yl)benzoyl]piperazin-1-yl]benzamide;

1-75: N-(2-aminoethyl)-2-[(2R)-4-[2-(6-aminopyridin-3-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-76: N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-77: N-(2-aminoethyl)-2-[(2R)-4-(2,6-dimethylpyridine-3-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-78: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(4-fluoro-2-methylbenzoyl)piperazin-1-yl]benzamide;

1-79: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(4-fluoro-2-methoxybenzoyl)piperazin-1-yl]benzamide;

1-80: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-(2-methoxy-4-methylbenzoyl)piperazin-1-yl]benzamide;

1-81: N-(2-aminoethyl)-2-[(2R)-4-(4-tert-butylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-82: 2-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-83: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(1H-imidazol-5-yl)methyl]benzamide;

1-84: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-imidazol-5-yl)ethyl]benzamide;

1-85: N-(2-aminoethyl)-2-[(2R)-4-[2-(2-aminopyridin-4-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-86: N-(2-aminoethyl)-2-[(2R)-4-(4,6-dimethylpyridine-3-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-87: N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-88: N-(2-aminoethyl)-2-[(2R)-4-(4-cyclopropyl-2-methylbenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-89: N-(2-aminoethyl)-2-[(2R)-4-(3-chloro-5-fluoropyridine-2-carbonyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-90: 2-[(2R)-4-[4-chloro-2-(piperazin-1-yl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-91: 2-[(2R)-4-[4-chloro-2-(dimethylamino)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-92: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-pyrrolidin-3-yl]benzamide;

1-93: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;

1-94: 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-methyl-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-95: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-96: 6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-97: N-{[(2S)-azetidin-2-yl]methyl}-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-98: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2S)-pyrrolidin-2-yl]methyl}benzamide;

1-99: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-piperidin-3-yl]benzamide;

1-100: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-piperidin-3-yl]benzamide;

1-101: 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{4-fluoro-2-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-102: N-{[(2R)-azetidin-2-yl]methyl}-2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-103: 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{2-fluoro-4-[(3S)-3-(methylamino)pyrrolidin-1-yl]benzoyl}piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-104: N-(2-aminoethyl)-2-[(2R)-4-{2-[3-(aminomethyl)pyrrolidin-1-yl]-4-methylbenzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-105: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(morpholin-4-yl)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;

1-106: N-(2-aminoethyl)-2-[(2R)-4-[3-(dimethylamino)-5-(trifluoromethyl)pyridine-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-107: N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-108: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2R)-pyrrolidin-2-yl]methyl}benzamide;

1-109: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2S)-morpholin-2-yl]methyl}benzamide;

1-110: 6-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-111: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-112: N-{[(2S)-azetidin-2-yl]methyl}-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-113: 6-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-114: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-115: 2-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-116: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;

1-117: N-(2-aminoethyl)-2-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-118: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[5-fluoro-3-(trifluoromethyl)pyridine-2-carbonyl]piperazin-1-yl]benzamide;

1-119: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(methylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;

1-120: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-(pyrrolidin-1-yl)-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;

1-121: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-122: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-123: N-[(2R)-azetidin-2-yl]methyl}-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-124: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(2R)-pyrrolidin-2-yl]methyl}benzamide;

1-125: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-126: 2-[(2R)-4-[2-amino-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)benzamide;

1-127: N-(2-aminoethyl)-2-[(2R)-4-[6-(dimethylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-128: 2-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;

1-129: N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-imidazol-1-yl)benzoyl]piperazin-1-yl]benzamide;

1-130: N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-pyrrol-1-yl)benzoyl]piperazin-1-yl]benzamide;

1-131: N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(1H-pyrazol-1-yl)benzoyl]piperazin-1-yl]benzamide;

1-132: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(1H-imidazol-2-yl)methyl]benzamide;

1-133: 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[5-(trifluoromethyl)pyridine-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-134: 2-[(2R)-4-[3-chloro-5-(trifluoromethyl)pyridine-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-135: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-136: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-137: 2'-ethoxy-5-[(2R)-4-[4-ethoxy-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-138: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3S)-morpholin-3-yl]methyl}benzamide;

1-139: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]benzamide;

1-140: 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;

1-141: 6-[(2R)-4-(4-chloro-2-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(1H-imidazol-2-yl)ethyl]benzamide;

1-142: 2-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-143: 5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-144: 2-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-145: 2-[(2R)-4-[6-amino-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)benzamide;

1-146: N-(2-aminoethyl)-2-[(2R)-4-[4-(dimethylamino)-6-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-147: 2-[(2R)-4-[2-(dimethylamino)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-148: 2-[(2R)-4-[2-(3-aminoazetidin-1-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-149: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-6-(2-ethoxyphenyl)pyridine-2-carboxamide;

1-150: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-151: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-152: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-153: 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-154: 6-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-155: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-156: 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-157: 6-[(2R)-4-[2-cyano-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-158: 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-159: N-(2-aminoethyl)-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[2-methoxy-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]benzamide;

1-160: N-(2-aminoethyl)-2-[(2R)-4-[2-(dimethylamino)-6-methylpyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-161: 2-[(2R)-4-[2-(azetidin-1-yl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)benzamide;

1-162: N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(pyrrolidin-1-yl)benzoyl]piperazin-1-yl]benzamide;

1-163: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-(methylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-164: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-165: 6-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-166: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[2-methoxy-6-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-167: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-168: N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;

1-169: N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-170: N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;

1-171: 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-173: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-174: N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;

1-175: 6-[(2R)-4-[4-cyano-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-176: 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-177: 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-178: 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-179: 6-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-180: 6-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-181: 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-182: N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;

1-183: 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-184: 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-185: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methyl-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-186: 6-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-187: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-188: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-189: 6-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-190: 6-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-191: N-[2-(dimethylamino)ethyl]-5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-{2-[ethyl(methyl)amino]-4-fluorobenzoyl}piperazin-1-yl]benzamide;

1-194: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethyl-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-195: 5-[(2R)-4-(2-chloro-4-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-196: 5-[(2R)-4-(2-chloro-4-ethoxybenzoyl)-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-197: 6-[(2R)-4-{4-chloro-2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]benzoyl}-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-198: N-(azetidin-3-yl)-6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-199: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;

1-200: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3S)-pyrrolidin-3-yl]benzamide;

1-201: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(3S)-1-methylpyrrolidin-3-yl]benzamide;

1-202: N-(azetidin-3-yl)-3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluorobenzamide;

1-203: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[4-methoxy-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-2-fluoro-N-(1-methylazetidin-3-yl)benzamide;

1-204: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-(1-methylazetidin-3-yl)benzamide;

1-205: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethyl-[2,3'-bipyridine]-6-carboxamide;

1-206: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-207: N-[2-(dimethylamino)ethyl]-2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-208: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[2-(methylamino)ethyl]pyridine-2-carboxamide;

1-209: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-210: 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-211: N-(azetidin-3-yl)-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-212: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-213: 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-214: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-215: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-216: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-(1-methylazetidin-3-yl)benzamide;

1-218: 5-[(2R)-4-(2-chloro-4-cyclopropoxybenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-219: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(4,5-dihydro-1H-imidazol-2-yl)ethyl]-3-(2-ethoxypyridin-3-yl)-2-fluorobenzamide;

1-220: N-(azetidin-3-yl)-3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridine-2-carboxamide;

1-221: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide.

1-222: N-[(azetidin-3-yl)methyl]-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-223: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-224: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[2-(1H-imidazol-2-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-225: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-(ethylamino)-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-226: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[(4,5-dihydro-1H-imidazol-2-yl)methyl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-227: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(1-methyl-4,5-dihydro-1H-imidazol-2-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-228: 2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-{[(2R)-1-methylazetidin-2-yl]methyl}benzamide;

1-229: 6-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[(2S)-1-methylazetidin-2-yl]methyl} benzamide.

1-230: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-231: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(2R)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-232: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-233: 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-234: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(1-methyl-1H-imidazol-2-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-235: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopropanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-236: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

1-237: 3-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

1-238: 5-[(2R)-4-(adamantane-2-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-239: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-240: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-241: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-242: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-243: 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-244: N-[(2S)-azetidin-2-yl]methyl}-2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-245: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-246: 5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2'-methoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-247: 5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-2'-methoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-248: 5-[(2R)-4-[1-(2,2-difluoroethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-249: 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-250: 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-ethylcyclobutanecarbonyl)piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-251: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-252: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[*trans*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

1-253: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-254: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-255: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-256: N-{[(2S)-azetidin-2-yl]methyl}-2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-257: 2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-258: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-259: 5-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-260: 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-261: 5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-262: 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-263: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-264: 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-ethylcyclopentanecarbonyl)piperazin-1-yl]-N-[2-(methylamino)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-265: 5-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-266: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-267: 5-[(2R)-4-(2,4-dichlorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-270: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[(3R)-1-methylpyrrolidin-3-yl]benzamide;

1-271: 3-(2-ethoxypyridin-3-yl)-6-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-2-fluoro-N-[(3R)-pyrrolidin-3-yl]benzamide;

1-273: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-274: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-275: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-276: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-277: 5-[(2R)-4-(4-chloro-2-cyanobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-278: 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-279: 5-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-280: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-281: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-282: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.2]octane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-283: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-284: 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,5,7-trifluoroadamantane-1-carbonyl)piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-285: 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,5,7-trifluoroadamantane-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-286: 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-287: 3-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-288: 3-[(2R)-4-[4-chloro-2-(difluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-289: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-290: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3R,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-291: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3S,4R)-4-fluoropyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-292: 3-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-293: 3-[(2R)-4-[2-(difluoromethyl)-4-fluorobenzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-294: 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]hept-5-ene-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-295: 5-[(2R)-4-[1-(difluoromethyl)-3,3-difluorocyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-296: 2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-N-[(3S,4R)-4-fluoro-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-297: 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]heptane-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-298: 5-[(2R)-4-[3,3-difluoro-1-(trifluoromethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-299: 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-300: 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-3-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-301: 2'-ethoxy-5-[(2R)-2-ethyl-4-[*trans*-3-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-302: 2'-ethoxy-5-[(2R)-2-ethyl-4-(3,3,3-trifluoro-2,2-dimethylpropanoyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-303: 5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-304: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-305: 3-[(2R)-4-(2-tert-butylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-306: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-307: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-[(5-methyl-2-oxo-2H-1,3-dioxol-4-yl)methyl]pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-309: 1,1,1-trifluoro-2-methylpropan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-310: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2R)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-311: 5-[(2R)-4-(2-cyclopropylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-312: 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-313: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-314: 5-[(2R)-4-{7,7-dimethylbicyclo[2.2.1]heptane-1-carbonyl}-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-315: 3-[(2R)-4-[(2S)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-316: 3-[(2R)-4-[(2R)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-methoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-317: 5-[(2R)-4-(2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-318: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3-fluoroazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-319: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3-fluoro-1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-320: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-321: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-322: 2'-ethoxy-5-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-323: 5-[(2R)-4-(2,2-diethylpyrrolidine-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-324: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-{[(2R)-pyrrolidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-325: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2S]-pyrrolidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxamide;

1-326: 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-methyl-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-327: N-[[2R)-azetidin-2-yl]methyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-328: 5-[(2R)-4-[1-(2,2-difluoroethyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-329: N-[[2S)-azetidin-2-yl]methyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-330: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2R)-1-methylazetidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxamide;

1-331: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2S)-1-methylazetidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxamide;

1-332: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2S)-1-methylpyrrolidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxamide;

1-333: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[[2R)-1-methylpyrrolidin-2-yl]methyl]-[2,3'-bipyridine]-6-carboxam;

1-334: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3-methoxy-1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-335: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-336: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[*cis*-4-(trifluoromethyl)cyclohexanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-337: N-[2-(azetidin-1-yl)ethyl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-338: 1-(trifluoromethyl)cyclobutyl (3R)-4-[6-(2-ethoxyphenyl)-2-[[3R)-pyrrolidin-3-yl]carbonyl]pyridin-3-yl]-3-ethylpiperazine-1-carboxylate;

1-339: 2'-ethoxy-5-[(2R)-2-ethyl-4-[2-(trifluoromethyl)bicyclo[2.2.1]heptane-2-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-340: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]pyridine-2-carboxamide;

1-341: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-342: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-343: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-344: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-345: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-phenylpyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-346: 5-[(2R)-4-(7-chloro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-347: 5-[(2R)-4-(5-chloro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-348: 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-[6-(2-ethoxyphenyl)-2-{[(3R)-pyrrolidin-3-yl]carbamoyl}pyridin-3-yl]-3-ethylpiperazine-1-carboxylate;

1-349: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(2,2,2-trifluoroethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-350: 2'-ethoxy-5-[(2R)-2-ethyl-4-(7-fluoro-2,3-dihydro-1H-indole-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-351: (1R)-2,2,2-trifluoro-1-phenylethyl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-352: 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclopentanecarbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-353: 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclobutanecarbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-354: 5-[(2R)-4-[2,2-difluoro-7,7-dimethylbicyclo[2.2.1]heptane-1-carbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide xylate;

1-355: 3-[(2R)-4-[(2S)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-356: 3-[(2R)-4-[(2R)-2-tert-butylpyrrolidine-1-carbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-pyrrolidin-3-yl]pyridine-2-carboxamide;

1-357: 5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[2-(3-fluoroazetidin-1-yl)ethyl]-[2,3'-bipyridine]-6-carboxamide;

1-358: 2'-ethoxy-5-[(2R)-2-ethyl-4-(1-phenylcyclobutanecarbonyl)piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-359: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(2,2,2-trifluoroethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-360: 2'-ethoxy-5-[(2R)-2-ethyl-4-(7-methyl-2,3-dihydro-1H-indole-1-carbonyl)piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-361: 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-362: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]pyridine-2-carboxamide;

1-363: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-364: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-365: 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-366: (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-[[[(3R)-pyrrolidin-3-yl]carbamoyl]-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate);

1-367: (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-[[[(3R)-pyrrolidin-3-yl]carbamoyl]-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate);

1-368: 2'-ethoxy-5-[(2R)-2-ethyl-4-[5-(trifluoromethyl)thiophene-2-carbonyl]piperazin-1-yl]-N-[(3R)-pyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-369: 1,1,1,3,3,3-hexafluoropropan-2-yl (3R)-4-(2'-ethoxy-6-[[[(3R)-pyrrolidin-3-yl]carbamoyl]-[2,3'-bipyridin]-5-yl]-3-ethylpiperazine-1-carboxylate);

1-370: 3-(trifluoromethyl)pentan-3-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-371: 5-[(2R)-4-[1-(2,4-difluorophenyl)cyclobutanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-372: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-373: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-374: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-375: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-376: 1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-377: (1S)-2,2,2-trifluoro-1-phenylethyl (3R)-4-(2'-ethoxy-6-{[(3R)-pyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-378: 5-[(2R)-4-(2,5-dichlorothiophene-3-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-379: 2'-ethoxy-5-[(2R)-2-ethyl-4-[3-(trifluoromethyl)bicyclo[1.1.1]pentane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-380: 5-[(2R)-4-(3,5-dichlorothiophene-2-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-381: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(3-fluorophenyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-382: 5-[(2R)-4-(7-chloro-2,3-dihydro-1H-indene-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-383: 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-384: 1-phenylcyclobutyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-385: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-386: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2R)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-387: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}pyridine-2-carboxamide;

1-388: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-389: 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-390: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-391: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-{[(2S)-1-methylazetidin-2-yl]methyl}-[2,3'-bipyridine]-6-carboxamide;

1-392: (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-392: (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-393: 3-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-394: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-395: 2'-ethoxy-5-[(2R)-2-ethyl-4-[4-(trifluoromethyl)bicyclo[2.2.1]heptane-1-carbonyl]piperazin-1-yl]-N-[(1-methylazetidin-3-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-396: (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(1-methylazetidin-3-yl)methyl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-396: (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(1-methylazetidin-3-yl)methyl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-397: 1-ethylcyclopentyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-398: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridine]-6-carboxamide;

1-399: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

1-400: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-401: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-402: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-403: (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(6-{[(3S)-1-azabicyclo[2.2.2]octan-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-403: (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(6-{[(3S)-1-azabicyclo[2.2.2]octan-3-yl]carbamoyl}-2'-ethoxy-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-404: 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-405: N-{1-azabicyclo[2.2.1]heptan-4-yl}-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-406: (2R)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-407: (2S)-1,1,1-trifluoro-3,3-dimethylbutan-2-yl (3R)-4-[6-(2-ethoxyphenyl)-2-{[(3R)-pyrrolidin-3-yl]carbamoyl}pyridin-3-yl]-3-ethylpiperazine-1-carboxylate;

1-408: 2,2-dimethyl-1-phenylpropyl (3R)-4-(2'-ethoxy-6-{[(3R)-1-methylpyrrolidin-3-yl]carbamoyl}-[2,3'-bipyridin]-5-yl)-3-ethylpiperazine-1-carboxylate;

1-409: 5-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-410: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]pyridine-2-carboxamide;

1-411: N-[(3S)-1-azabicyclo[2.2.2]octan-3-yl]-2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclobutanecarbonyl]piperazin-1-yl]-[2,3'-bipyridine]-6-carboxamide;

1-412: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-413: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(2-fluoropropan-2-yl)pyrrolidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-414: 3-[(2R)-4-(7-cyano-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

1-415: 2'-ethoxy-5-[(2R)-2-ethyl-4-[1-(trifluoromethyl)cyclopentanecarbonyl]piperazin-1-yl]-N-[(1-methyl-4,5-dihydro-1H-imidazol-2-yl)methyl]-[2,3'-bipyridine]-6-carboxamide;

1-416: 5-[(2R)-4-[1-(difluoromethyl)cyclopentanecarbonyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-417: 3-[(2R)-4-[1-(difluoromethyl)cyclopentanecarbonyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-418: 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-2,3-dihydro-1H-indole-1-carbonyl]piperazin-1-yl]-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-419: 5-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-420: 5-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-421: 3-[(2R)-4-(7-chloro-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-422: 5-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-(1-methylazetidin-3-yl)-[2,3'-bipyridine]-6-carboxamide;

1-423: 5-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-424: 3-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-(1-methylazetidin-3-yl)pyridine-2-carboxamide;

1-425: 5-[(2R)-4-(7-chloro-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-426: 3-[(2R)-4-(7-cyano-5-fluoro-2,3-dihydro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-427: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-428: 5-[(2R)-4-(7-chloro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-429: 5-[(2R)-4-(7-cyano-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-430: 3-[(2R)-4-(7-chloro-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-431: 3-[(2R)-4-(7-chloro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-432: 3-[(2R)-4-(7-cyano-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-433: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-434: 2'-ethoxy-5-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-435: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[(2S)-2-(trifluoromethyl)piperidine-1-carbonyl]piperazin-1-yl]-N-[(3S)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-436: 2'-ethoxy-5-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-437: 6-(2-ethoxyphenyl)-3-[(2R)-2-ethyl-4-[7-(trifluoromethyl)-1H-indole-1-carbonyl]piperazin-1-yl]-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

1-438: 5-[(2R)-4-(7-cyano-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-N-[(3R)-1-methylpyrrolidin-3-yl]-[2,3'-bipyridine]-6-carboxamide;

1-439: 3-[(2R)-4-(7-cyano-5-fluoro-1H-indole-1-carbonyl)-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)-N-[(3R)-1-methylpyrrolidin-3-yl]pyridine-2-carboxamide;

2-1: 2-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl]amino]ethan-1-ol;

2-2: (2S)-1-[(4-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[1,1'-biphenyl]-3-yl)methyl]amino]propan-2-ol;

2-3: 1-[4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-methoxy-[1,1'-biphenyl]-3-yl]methanamine;

2-4: 1-[4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-(2-methoxyethoxy)-[1,1'-biphenyl]-3-yl]methanamine;

2-5: 1-{4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-propoxy-[1,1'-biphenyl]-3-yl}methanamine;

2-6: 2-{[3'-(aminomethyl)-4'-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-2-yl]oxy}ethan-1-ol;

2-7: 2-{[3'-(aminomethyl)-4'-[(2R)-2-ethyl-4-[2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[1,1'-biphenyl]-2-yl]oxy}ethan-1-ol;

2-8: [3'-(aminomethyl)-4'-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[1,1'-biphenyl]-2-yl]methanol;

2-9: 1-{2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}methanamine;

2-10: 1-{2-[(2R)-4-[4'-(aminomethyl)-5-fluoro-[1,1'-biphenyl]-2-carbonyl]-2-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)phenyl}methanamine;

2-11: 1-{6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}methanamine;

2-12: 1-{3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxyphenyl)pyridin-2-yl}methanamine;

2-13: 1-{5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;

2-14: 1-{5-[(2R)-4-(2-chloro-4-methylbenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;

2-15: 1-{5-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;

2-16: ({5-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methyl)(methyl)amine;

2-17: 1-{5-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-2'-ethoxy-[2,3'-bipyridin]-6-yl}methanamine;

2-18: 1-{6-[(2R)-4-(2-chloro-4-fluorobenzoyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluorophenyl}methanamine;

2-19: 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-(2,3,4-trifluorobenzoyl)piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;

2-20: 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;

2-21: {[5-(2-ethoxypyridin-3-yl)-2-[(2R)-2-ethyl-4-[4-fluoro-2-(trifluoromethyl)benzoyl]piperazin-1-yl]phenyl]methyl}[2-(methylamino)ethyl]amine;

2-22: 13-[3-(aminomethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]phenyl]-N-methylpyridin-2-amine;

2-23: 3-[3-(aminomethyl)-4-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]phenyl]-N,N-dimethylpyridin-2-amine;

2-24: 1-{2-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-5-[2-(pyrrolidin-1-yl)pyridin-3-yl]phenyl}methanamine;

2-25: 1-{2'-ethoxy-5-[(2R)-4-[4-ethoxy-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;

2-26: 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-3'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;

2-27: 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-[1,1'-biphenyl]-3-yl}methanamine;

2-28: 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-4'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;

2-29: 1-{2'-ethoxy-4-[(2R)-2-ethyl-4-[4-methyl-2-(trifluoromethyl)benzoyl]piperazin-1-yl]-5'-fluoro-[1,1'-biphenyl]-3-yl}methanamine;

2-30: 1-{2'-ethoxy-5-[(2R)-4-[6-ethoxy-2-(trifluoromethyl)pyridine-3-carbonyl]-2-ethylpiperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;

2-31: 1-{2'-ethoxy-5-[(2R)-2-ethyl-4-[6-methoxy-2-(trifluoromethyl)pyridine-3-carbonyl]piperazin-1-yl]-[2,3'-bipyridin]-6-yl}methanamine;

3-1: N-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2',3'-difluoro-[1,1'-biphenyl]-3-carboxamide;

3-2: N-(3-aminopropyl)-4-{4-[2-cyano-4-(trifluoromethyl)phenyl]piperazin-1-yl}-2'-ethoxy-[1,1'-biphenyl]-3-carboxamide;

3-3: 6-[(2R)-4-(2,4-dichlorobenzenesulfonyl)-2-ethylpiperazin-1-yl]-3-(2-ethoxypyridin-3-yl)-2-fluoro-N-[2-(methylamino)ethyl]benzamide;

3-4: (3R)-N-(2,4-dichlorophenyl)-4-[4-(2-ethoxypyridin-3-yl)-3-fluoro-2-{[2-(methylamino)ethyl]carbonyl}phenyl]-3-ethylpiperazine-1-carboxamide;

3-5: N-(2-aminoethyl)-2-[(3R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-3-ethylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)benzamide;

3-6: 2-[(3R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-3-methylpiperazin-1-yl]-5-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]benzamide;


3-7: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-N-[2-(dimethylamino)ethyl]-6-(2-ethoxypyridin-3-yl)pyrazine-2-carboxamide;






3-8: 3-[(2R)-4-[4-chloro-2-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-6-(2-ethoxypyridin-3-yl)-N-[2-(methylamino)ethyl]pyrazine-2-carboxamide; and


3-9: N-(2-aminoethyl)-5-[(2R)-4-[2-chloro-4-(trifluoromethyl)benzoyl]-2-ethylpiperazin-1-yl]-2-(2-ethoxyphenyl)pyridine-4-carboxamide;

or a pharmaceutically acceptable salt, or solvate thereof.

18. The compound of claim 1, wherein the compound is:







pharmaceutically acceptable salt, or solvate thereof.

19. A pharmaceutical composition comprising a compound of any one of claims 1-18, or a pharmaceutically acceptable salt, or solvate thereof, and at least one pharmaceutically acceptable excipient.
20. Use of a compound of any one of claims 1-18 in the manufacture of a medicament for treating a disease or condition in a mammal that would benefit from the modulation of melanocortin subtype-2 receptor (MC2R) activity;
wherein the disease or condition comprises growth of fat pads in the collarbone, back of neck, face and trunk, excessive sweating, dilation of capillaries, thinning of the skin, muscle weakness, hirsutism, depression/anxiety, hypertension, osteoporosis, insulin resistance, hyperglycemia, and heart disease;

or wherein the disease or condition is Cushing's syndrome;
or wherein the disease or condition is Cushing's disease;
or wherein the disease or condition is ectopic ACTH syndrome;
or wherein the disease or condition is congenital adrenal hyperplasia.