

BICYCLIC HETEROCYCLIC DERIVATIVES AS MNK1 AND MNK2 MODULATORS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 The present application claims priority to UK Patent Application No. 1205669.3, filed March 30, 2012, which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

10 The human MAP Kinase-interacting kinases, also known as MAP Kinase signal-integrating kinases, (MNKs¹), are ubiquitously expressed protein-serine/threonine kinases that are directly activated by ERK or p38 MAP kinases.^{2,3} They comprise a group of four proteins derived from two genes (gene symbols: MKNK1 and MKNK2) by alternative splicing. MNK1a/b and MNK2a/b proteins differ at their C-termini, in each case the a-form possessing a longer C-terminal region than the b-form which lacks the MAP Kinase-binding region. The
15 N-termini of all forms contain a polybasic region which binds Importin α and the translation factor scaffold protein eukaryotic Initiation Factor (eIF4G). The catalytic domains of MNK1a/b and MNK2a/b share three unusual features, two short inserts and a DFD tripeptide feature where other kinases have DFG. Mnk isoforms differ markedly in their activity, regulation, and in subcellular localization. The best-characterized MNK substrate is eIF4E.
20 Although the cellular role of eIF4E phosphorylation remains unclear, it may promote export of a defined set of mRNAs from the nucleus. Other Mnk substrates bind to AU-rich elements that modulate the stability/translation of specific mRNAs.¹ MNK1 is highly expressed in hematological malignancies^{4,5} and both MNK1 and MNK2 are up-regulated in solid tumors such as gliomas and ovarian cancers.^{6,7}

25 The Eukaryotic Initiation Factor-4 E (eIF4E) regulates the expression of genes involved in proliferation and survival as a cap dependent mRNA translation and mRNA export factor. eIF4E is dysregulated in several human cancers, including breast,¹⁰ prostate¹¹, and some leukemias,¹² and elevated levels of eIF4E are a marker of poor prognosis.^{12,13} Moreover,
30 overexpression and dysregulation of eIF4E leads to an increased tumor number, invasion, and metastases in mouse models¹³ and transgenic expression of eIF4E leads to a variety of cancers.^{13,14} eIF4E overexpression results in a specific increase in the translation of these weakly competitive mRNAs, many of which encode products that stimulate cell growth and angiogenesis, *e.g.*, fibroblast growth factor 2 and vascular endothelial growth factor,¹⁵⁻¹⁷

cyclin D1,¹⁸ and ribonucleotide reductase.¹⁹ Several lines of evidence support the key role that eIF4E plays in cancer development and/or progression. First, overexpression of eIF4E can cause neoplastic transformation of cells or accentuate neoplastic features.^{20;21} Second, reducing eIF4E with antisense RNA or reducing its function by overexpression of the inhibitory 4E-BP proteins can suppress the oncogenic properties of many cell lines.^{10;22;23} Third, increased expression of eIF4E found in the most classes of solid tumors. These include bladder,²⁴ breast,^{25;26} cervical,²⁷ head and neck,²⁸⁻³⁰ and prostate tumors.³¹ Finally, both the expression and activity of eIF4E are regulated at multiple levels by growth factors and oncogenes,³² suggesting that the protein is a nexus of converging transformation signaling pathways.

eIF4E is phosphorylated by the MNK1/2 serine/threonine kinases in response to activation by mitogenic and stress signals downstream of ERK1/2 and p38 MAP kinase respectively.^{2;3} eIF4E phosphorylation at serine 209 by MNK1/2 is required to promote its transformation activity.^{33;34} Surprisingly, MNK1/2 double knock-out mice do not have any apparent phenotype,³⁵ calling into question whether phosphorylation by MNK has any impact on the functionality of the mammalian eIF4E.³⁶ Nevertheless, MNK phosphorylation of eIF4E on Ser-209 is believed to be critical to eIF4E oncogenic activity.³⁴

Thus, inhibitors of MNK1/2, by preventing the phosphorylation of eIF4E could provide a viable therapeutic approach in high-eIF4E dependent cancers.

Studies have shown that overexpression of eIF4E, as well as eIF4E phosphorylation, promote cancer cell survival, at least in part through the elevation of the anti-apoptotic protein Mcl-1.³⁴ Mcl-1 is a Bcl2 family member with a very short half-life, and Mcl-1 mRNA translation highly depends on eIF4E. Thus, it is possible that the inhibition of eIF4E phosphorylation by Mnk might induce the death tumor cells, as shown for Myc-induced lymphoma.³⁴

Blast crisis chronic myeloid leukemia (BC-CML) is characterized by an expansion of a population of granulocyte macrophage progenitor-like cells (GMPs) that have acquired self-renewal capacity,³⁷ a feature not seen in normal or chronic phase (CP) GMPs. The ability to self-renew is thought to be mediated by β -catenin activation, and may contribute to disease persistence, as well as activity as a reservoir for drug resistance. The mechanisms contributing to β -catenin activation remain obscure, and will need to be identified to improve

the control of BC-CML. The role of the translation machinery in mediating β -catenin-mediated self-renewal was investigated, since prior work had implicated aberrant mRNA translation in drug-resistance and BC pathophysiology.³⁸⁻⁴⁰ Using immunofluorescence (IF), it was confirmed that BC-GMPs have activated nuclear beta-catenin compared to GMPs isolated from normal cord blood, and that this was associated with increased eIF4E expression and phosphorylation at Ser209. Trough biochemical and genetic approaches in CML cell lines (K562 and KCL22), it was demonstrated that eIF4E overexpression was sufficient to increase beta-catenin activity (as measured by immunofluorescence for nuclear beta-catenin, beta-catenin reporter assays, and expression of beta-catenin-regulated genes). By expressing phospho-mutant forms of eIF4E (S209A, S209D), it was found that the increase in beta-catenin transcriptional activity is dependent on phosphorylation of at Ser209. In line with these observations, siRNA-mediated knockdown or inhibition of the MNK1/2 kinases (which mediate *in vivo* eIF4E phosphorylation) with small molecules prevented the increased beta-catenin activity induced by eIF4E overexpression. Mechanistically, eIF4E activates beta-catenin signaling via a two-step mechanism. First, eIF4E overexpression increased total cell beta-catenin and secondly, eIF4E phosphorylation facilitated beta-catenin nuclear translocation. The latter step was associated with increased beta-catenin phosphorylation at Ser552, a site known to be involved in nuclear translocation, and directly regulated by AKT. Consistent with this model, siRNA-mediated knockdown or small molecule inhibition of AKT (AKT inhibitor IV) prevented eIF4E-mediated increases in beta-catenin transcriptional activity. The importance of eIF4E phosphorylation on beta-catenin activation and the self-renewal capacity of primary BC GMPs cells was assessed. Treatment with CGP57380, but not imatinib or dasatinib, inhibited eIF4E phosphorylation, as well as preventes accumulation of active nuclear beta-catenin in BC GMPs. The effect of MNK1/2 inhibition was evaluated on the stem cell function of BC cells using both *in vitro* and *in vivo* assays. In an *in vitro* serial replating assay, it was shown that CGP57380 impaired the ability of CD34+ BC cells (including those carrying T315I mutation), but not normal CD34+ cells, to serially replate for more than 8 weeks in methylcellulose. Interestingly, treatment with either imatinib or dasatinib only partially impaired the ability of BC-CML to serially replate. *In vitro* treatment of BC CD34+ CML cells, but not normal cord blood CD34+ cells, with CGP57380 retarded their ability to engraft NSG mice. Finally, *in vivo* serial transplantation assay for assessing the leukemia stem cell (LSC) function of patient-derived BC-GMPs was developped. BC GMPs or BC CD34+ CML cells were injected intrafemorally into 8- to 10-week old sublethally irradiated NSG mice. Following engraftment, mice were treated with

vehicle, CGP53780 (40 mg/kg/d), or dasatinib (5mg/kg/d) for three consecutive weeks.

Following treatment, human CD34+ cells were isolated from the mice, and transplanted into a second recipient mouse. At 16 weeks, it was found that *in vivo* treatment with CGP57380, but not dasatinib, prevented BC cells from serially transplanting NSG mice. In summary, these results demonstrate that: 1. eIF4E is overexpressed and phosphorylated at Ser209 in BC, but not in normal GMPs; 2. eIF4E phosphorylation activates beta-catenin signalling in BC GMPs; 3. MNK inhibition prevents eIF4E phosphorylation and beta-catenin signalling in BC GMPs; and 4. MNK inhibition prevents BC GMPs from functioning as leukemia stem cells. These studies suggest that pharmacologic inhibition of the MNK1/2 kinases may be therapeutically useful in BC CML.

The level of expression of eIF4E and the degree of eIF4E phosphorylation is regulated by pathways that include the P38 kinase, MAPK kinase and AKT/mTOR pathways⁴¹ as shown in Figure 4. Inhibitors of mTOR such as rapamycin, decrease the level of phosphorylated eIF4E.⁴² mTOR inhibitors, as single agents, have proven efficacious in several cancer types such as transplant-associated lymphoma^{43,44} and Kaposi sarcoma,^{45,46} tuberous sclerosis-related astrocytoma,⁴⁷ and mantle cell and other non-Hodgkin lymphomas.⁴⁸ Two mTOR inhibitors are currently marketed for the treatment of the renal cell carcinoma.^{49,50}

Inhibition of mTOR by rapamycin also suppresses mTOR catalyzed phosphorylation of EBP1 leading to an increased level eIF4E-EBP1. Consequently, rapamycin inhibits translation initiation by decreasing the phosphorylation of eIF4E-binding proteins, thus decreasing eIF4E availability to the initiation complex.

The treatment with rapalogs typically leads to the clinically stable disease or partial remission rather than the tumor elimination.⁵¹ This suboptimal drug effect is likely due at least in part to the cytostatic rather than cytotoxic properties of the mTORC1 inhibitors. Therefore, there is a potential for a drug combination therapy that ideally would culminate in the complete remission of the cancer. However, most of the attempts to combine mTORC1 inhibitors with other drugs, typically the standard chemotherapeutic agents targeting DNA replication, have been disappointing, on occasion even leading to drug antagonism. Preclinical studies of mTOR combined with cis-platin⁵² or methotrexate⁵³ show the most promise.

Combination therapy with MNK1/2 and mTOR kinases inhibitors could be a viable strategy to treat certain types of cancer.⁵⁴ WO 2010/055072 discloses MNK and mTOR combination therapy with small molecules, antibodies and siRNA for the treatment of cancer,⁵⁵ and recent findings support that MNK and mTOR combination induces apoptosis in cutaneous T cell lymphoma cells.⁴²

Macrophages are major effectors of innate immunity, stimulated by a broad variety of bacterial products through specific TLRs on the cell surface to produce proinflammatory cytokines, such as TNF. *E. coli* LPS is a potent stimulus to macrophage gene expression, especially TNF, by engaging the TLR4 membrane signaling complex.⁵⁶ It was shown that TLR signaling pathways require Mnk expression through the use of a panel of commercial TLR agonist panel on macrophage. TNF production was increased as a response to Salmonella LPS (TLR4), ODN2006 (TLR9), HKLM (TLR2), FSL (TLR6/2) and imiquimod (TLR7) stimulation. In each case the production of TNF was inhibited by MNK kinases inhibitor CGP57380 in a dose dependant fashion⁵⁷ and the release of multiple innate proinflammatory cytokines were affected, supporting a central role for MNK in inflammation.⁵⁸

It is reported that heterogeneous nuclear ribonucleoprotein A1 (hnRNPA1) when phosphorylated by MNK1/2 accumulates in the cytoplasmic stress granules (SGs) under stress related conditions. hnRNPA1 exit the nucleus bound to poly(A) mRNA and this complex is required for hnRNPA1 phosphorylation by MNK1/2 and for its relocalization to the cytoplasmic SGs. Phosphorylation of hnRNPA1 by MNK1/2 reduces its binding affinity to 3'UTR mRNA and consequently Mnk inhibition enhances hnRNPA1 association with TNF mRNA. TNF gene transcript level is undetectable in unstimulated T Cell and greatly increased upon stimulation. MNK inhibition effect on TNF appears to be more at the translation level as MNK inhibition has no influence on the level of TNF mRNA.⁵⁹ Moreover, the formation of SG is reported to be prevented by MNK inhibition⁶⁰ thus removing the protection that was offered by the SGs where the phosphorylated hnRNPA1 bound mRNA could localize.

MNK inhibitors can regulate the innate immune response in macrophage. A compound with anti inflammatory properties will inhibit the release of pro-inflammatory cytokines.

It has been shown that CGP57380, a Mnk inhibitor, inhibits the release of TNF alpha by macrophage⁶¹ (and not eIF4E). According to WO2005/003785 MNK kinases are promising targets for anti-inflammatory therapy.

- 5 MNK1/2 were also reported to phosphorylate a number of different proteins in addition to eIF4E. Three of these are hnRNPA1,⁶⁰ cPLA2 and Sprouty2.^{62;63} Their role and function is still being investigated. Among these substrates, hnRNPA1 is overexpressed in colorectal cancer and it could contribute to maintenance of telomere repeats in cancer cells with enhanced cell proliferation.⁶⁴ It is also reported that the expression levels of hnRNPA/B is
 10 deregulated in non small cell lung cancer.⁶⁵


MNK inhibitors are potentially useful in the treatment of cancers including breast,⁶⁶ prostate,¹¹ hematological malignancies (e.g., CML, AML), head and neck, colon,⁶⁷ bladder, prostatic adenocarcinoma, lung, cervical, and lymphomas.^{68;68;69}

15

SUMMARY OF THE INVENTION

In one aspect, the present invention relates to compounds that act as kinase inhibitors, in particular as inhibitors of the MAP kinase interacting kinases 1 and 2 (MNK1 and MNK2).

In certain embodiments, a compound of the present invention is of Formula (I):



20

or a pharmaceutically acceptable form thereof, wherein R¹, R², R³, R⁵, R⁶, R⁷, R⁸, R⁹, X₁, X₂, X₃, X₄, X₅, X₆, X₇, and X₈ are as described herein.

- 25 In another aspect, the present invention relates to pharmaceutical compositions comprising these compounds and to their use for the preparation of a medicament for the prophylaxis and treatment diseases associated with a dysfunction linked to MNK1 and MNK2 pathway, where MNK1 and MNK2 play a role (MNK overexpression, eIF4E overexpression, P38 MAPK

kinase pathway), such as but not limited to, cancer (blood and solid tumors), inflammatory conditions, Alzheimer's disease, metabolic disorders (obesity, diabetes) as well as methods of treatment of these disorders using compounds described herein as single agents or in combination with one or more additional agents. In some embodiments, an additional agent

5 is a kinase inhibitor. In some embodiments, an additional agent is an mTOR inhibitor.

Exemplary mTOR inhibitors include sirolimus, temsirolimus, everolimus, and ridaforolimus.

In some embodiments, an additional agent is a PI3-kinase inhibitor. Exemplary PI3-kinase inhibitors include wortmannin, demethoxyviridin, LY294002, perifosine, CAL101, PX-886, BEZ235, SF1126, INK1117, INK1197, IPI-145, GDC-0941, BKM120, XL147, XL765,

10 palomid 529, GSK1059615, ZSTK474, PWT33597, IC87114, TG100-115, CAL263, PI-103, GNE-477, CUDC-907, GSK 2126458, GDC-0980, PF-46915032, CAL263, SF1126 and PX-886. In some embodiments, the PI3-kinase inhibitor inhibits PI3K- α , PI3K- β , PI3K- γ , and/or PI3K- δ .

15 In yet another aspect, the present invention describes methods for the synthesis and isolation of a compound of Formula (I), such as, for example, substituted 3,6-diphenylimidazo[1,2-a]pyridine; 3,6-diphenylimidazo[1,2-a]pyrazine; 3,6-diphenylimidazo[1,2-b]pyridazine; 3,5-diphenylpyrazolo[1,5-a]pyrimidine; 1,6-diphenyl-1H-imidazo[4,5-c]pyridine; 1,6-diphenyl-1H-benzo[d]imidazole.

20

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 demonstrates that Example 6 (ETC-10445-3), Example 1 (ETC-5336) and Example 2 (ETC-6740) prevent eIF4E phosphorylation in BC-CML cell lines and primary cells. (A) Western blot analysis of K562 cell lines treated with various concentrations of compounds for 24 h. (B-D) BC-CML cells were treated with various compounds for 48 h. 48 hours post drug treatment, cells were harvested for immunofluorescence analysis for nuclear active β -catenin and phosphor-eIF4E level.

30

Figure 2 shows the effects of Example 6 (ETC-0445-3), Example 1 (ETC-5336), and Example 2 (ETC-6740) on serial replating efficiency of BC-CML primary cells. Colony forming and serial replating assays were performed on normal and BC-CML primary samples. CD34+ cells from (A) cord blood and (B-D) BC CML cells were treated with compounds for 48 hr. Forty-eight hours post drug treatment, 1×10^4 cells were plated for

35

colony-forming cell (CFC) assay; colonies were enumerated and individually picked for serial replating. The serial replating efficiency was assessed by the ability of individual clones to replate to the third plating over 8 weeks and displayed as percentage relative to the first colony forming assay readout.

5

Figure 3 demonstrates that Example 176 (ETC-7114) and Example 62 (ETC-7168) prevent eIF4E phosphorylation in BC-CML cell line. (A) Western blot analysis of K562 cell line treated with various concentration of compounds for 24 h. Samples from BC-CML patients were treated with various compounds for 24 h. 24 hours post drug treatment, cells were harvested for immunofluorescence analysis for nuclear active beta-catenin and phosphor-eIF4E level.

10

Figure 4 shows the pathway connection between MNK and mTOR (*PNAS* 2010, 107 (32), 13975-13976).

15

DEFINITIONS

Chemical definitions

Definitions of specific functional groups and chemical terms are described in more detail below. The chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in Thomas Sorrell, *Organic Chemistry*, University Science Books, Sausalito, 1999; Smith and March, *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; and Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

20

25

30

Compounds described herein can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, e.g., enantiomers and/or diastereomers. For example, the compounds described herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. Isomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid

chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred isomers can be prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972). The invention additionally encompasses compounds described herein as individual isomers substantially free of other isomers, and alternatively, as mixtures of various isomers.

- 10 The term "aliphatic," as used herein, includes both saturated and unsaturated, nonaromatic, straight chain (*i.e.*, unbranched), branched, acyclic, and cyclic (*i.e.*, carbocyclic) hydrocarbons, and combinations thereof. In some embodiments, an aliphatic group is optionally substituted with one or more functional groups. As will be appreciated by one of ordinary skill in the art, "aliphatic" is intended herein to include, but is not limited to, alkyl, 15 alkenyl, alkynyl, cycloalkyl, and cycloalkenyl moieties, and combinations thereof.

When a range of values is listed, it is intended to encompass each value and sub-range within the range. For example "C₁₋₆ alkyl" is intended to encompass, C₁, C₂, C₃, C₄, C₅, C₆, C₁₋₆, C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₅, C₂₋₄, C₂₋₃, C₃₋₆, C₃₋₅, C₃₋₄, C₄₋₆, C₄₋₅, and C₅₋₆ alkyl.

20

- "Alkyl" refers to a radical of a straight-chain or branched saturated hydrocarbon group having from 1 to 20 carbon atoms ("C₁₋₂₀ alkyl"). In some embodiments, an alkyl group has 1 to 10 carbon atoms ("C₁₋₁₀ alkyl"). In some embodiments, an alkyl group has 1 to 9 carbon atoms ("C₁₋₉ alkyl"). In some embodiments, an alkyl group has 1 to 8 carbon atoms ("C₁₋₈ 25 alkyl"). In some embodiments, an alkyl group has 1 to 7 carbon atoms ("C₁₋₇ alkyl"). In some embodiments, an alkyl group has 1 to 6 carbon atoms ("C₁₋₆ alkyl"). In some embodiments, an alkyl group has 1 to 5 carbon atoms ("C₁₋₅ alkyl"). In some embodiments, an alkyl group has 1 to 4 carbon atoms ("C₁₋₄ alkyl"). In some embodiments, an alkyl group has 1 to 3 carbon atoms ("C₁₋₃ alkyl"). In some embodiments, an alkyl group has 1 to 2 30 carbon atoms ("C₁₋₂ alkyl"). In some embodiments, an alkyl group has 1 carbon atom ("C₁ alkyl"). In some embodiments, an alkyl group has 2 to 6 carbon atoms ("C₂₋₆ alkyl").

Examples of C₁₋₆ alkyl groups include methyl (C₁), ethyl (C₂), n-propyl (C₃), isopropyl (C₃), n-butyl (C₄), tert-butyl (C₄), sec-butyl (C₄), iso-butyl (C₄), n-pentyl (C₅), 3-pentanyl (C₅), amyl (C₅), neopentyl (C₅), 3-methyl-2-butanyl (C₅), tertiary amyl (C₅), and n-hexyl (C₆).

Additional examples of alkyl groups include n-heptyl (C₇), n-octyl (C₈) and the like. Unless otherwise specified, each instance of an alkyl group is independently optionally substituted, *i.e.*, unsubstituted (an “unsubstituted alkyl”) or substituted (a “substituted alkyl”) with one or more substituents. In certain embodiments, the alkyl group is unsubstituted C₁₋₁₀ alkyl (*e.g.*,
5 -CH₃). In certain embodiments, the alkyl group is substituted C₁₋₁₀ alkyl.

“Alkenyl” refers to a radical of a straight-chain or branched hydrocarbon group having from 2 to 20 carbon atoms, one or more carbon-carbon double bonds, and no triple bonds (“C₂₋₂₀ alkenyl”). In some embodiments, an alkenyl group has 2 to 10 carbon atoms (“C₂₋₁₀ alkenyl”). In some embodiments, an alkenyl group has 2 to 9 carbon atoms (“C₂₋₉ alkenyl”).
10 In some embodiments, an alkenyl group has 2 to 8 carbon atoms (“C₂₋₈ alkenyl”). In some embodiments, an alkenyl group has 2 to 7 carbon atoms (“C₂₋₇ alkenyl”). In some embodiments, an alkenyl group has 2 to 6 carbon atoms (“C₂₋₆ alkenyl”). In some embodiments, an alkenyl group has 2 to 5 carbon atoms (“C₂₋₅ alkenyl”). In some
15 embodiments, an alkenyl group has 2 to 4 carbon atoms (“C₂₋₄ alkenyl”). In some embodiments, an alkenyl group has 2 to 3 carbon atoms (“C₂₋₃ alkenyl”). In some embodiments, an alkenyl group has 2 carbon atoms (“C₂ alkenyl”). The one or more carbon-carbon double bonds can be internal (such as in 2-butenyl) or terminal (such as in 1-butenyl). Examples of C₂₋₄ alkenyl groups include ethenyl (C₂), 1-propenyl (C₃), 2-propenyl
20 (C₃), 1-butenyl (C₄), 2-butenyl (C₄), butadienyl (C₄), and the like. Examples of C₂₋₆ alkenyl groups include the aforementioned C₂₋₄ alkenyl groups as well as pentenyl (C₅), pentadienyl (C₅), hexenyl (C₆), and the like. Additional examples of alkenyl include heptenyl (C₇), octenyl (C₈), octatrienyl (C₈), and the like. Unless otherwise specified, each instance of an alkenyl group is independently optionally substituted, *i.e.*, unsubstituted (an “unsubstituted
25 alkenyl”) or substituted (a “substituted alkenyl”) with one or more substituents. In certain embodiments, the alkenyl group is unsubstituted C₂₋₁₀ alkenyl. In certain embodiments, the alkenyl group is substituted C₂₋₁₀ alkenyl.

“Alkynyl” refers to a radical of a straight-chain or branched hydrocarbon group having from
30 2 to 20 carbon atoms, one or more carbon-carbon triple bonds, and optionally one or more double bonds (“C₂₋₂₀ alkynyl”). In some embodiments, an alkynyl group has 2 to 10 carbon atoms (“C₂₋₁₀ alkynyl”). In some embodiments, an alkynyl group has 2 to 9 carbon atoms (“C₂₋₉ alkynyl”). In some embodiments, an alkynyl group has 2 to 8 carbon atoms (“C₂₋₈ alkynyl”). In some embodiments, an alkynyl group has 2 to 7 carbon atoms (“C₂₋₇ alkynyl”).

In some embodiments, an alkynyl group has 2 to 6 carbon atoms ("C₂₋₆ alkynyl"). In some embodiments, an alkynyl group has 2 to 5 carbon atoms ("C₂₋₅ alkynyl"). In some embodiments, an alkynyl group has 2 to 4 carbon atoms ("C₂₋₄ alkynyl"). In some embodiments, an alkynyl group has 2 to 3 carbon atoms ("C₂₋₃ alkynyl"). In some
 5 embodiments, an alkynyl group has 2 carbon atoms ("C₂ alkynyl"). The one or more carbon-carbon triple bonds can be internal (such as in 2-butyne) or terminal (such as in 1-butyne). Examples of C₂₋₄ alkynyl groups include, without limitation, ethynyl (C₂), 1-propynyl (C₃), 2-propynyl (C₃), 1-butyne (C₄), 2-butyne (C₄), and the like. Examples of C₂₋₆ alkynyl groups include the aforementioned C₂₋₄ alkynyl groups as well as pentynyl (C₅), hexynyl
 10 (C₆), and the like. Additional examples of alkynyl include heptynyl (C₇), octynyl (C₈), and the like. Unless otherwise specified, each instance of an alkynyl group is independently optionally substituted, *i.e.*, unsubstituted (an "unsubstituted alkynyl") or substituted (a "substituted alkynyl") with one or more substituents. In certain embodiments, the alkynyl group is unsubstituted C₂₋₁₀ alkynyl. In certain embodiments, the alkynyl group is substituted
 15 C₂₋₁₀ alkynyl.

"Carbocyclyl" or "carbocyclic" refers to a radical of a non-aromatic cyclic hydrocarbon group having from 3 to 10 ring carbon atoms ("C₃₋₁₀ carbocyclyl") and zero heteroatoms in the non-aromatic ring system. In some embodiments, a carbocyclyl group has 3 to 8 ring
 20 carbon atoms ("C₃₋₈ carbocyclyl"). In some embodiments, a carbocyclyl group has 3 to 6 ring carbon atoms ("C₃₋₆ carbocyclyl"). In some embodiments, a carbocyclyl group has 3 to 6 ring carbon atoms ("C₃₋₆ carbocyclyl"). In some embodiments, a carbocyclyl group has 5 to 10 ring carbon atoms ("C₅₋₁₀ carbocyclyl"). Exemplary C₃₋₆ carbocyclyl groups include, without limitation, cyclopropyl (C₃), cyclopropenyl (C₃), cyclobutyl (C₄), cyclobutenyl (C₄),
 25 cyclopentyl (C₅), cyclopentenyl (C₅), cyclohexyl (C₆), cyclohexenyl (C₆), cyclohexadienyl (C₆), and the like. Exemplary C₃₋₈ carbocyclyl groups include, without limitation, the aforementioned C₃₋₆ carbocyclyl groups as well as cycloheptyl (C₇), cycloheptenyl (C₇), cycloheptadienyl (C₇), cycloheptatrienyl (C₇), cyclooctyl (C₈), cyclooctenyl (C₈), bicyclo[2.2.1]heptanyl (C₇), bicyclo[2.2.2]octanyl (C₈), and the like. Exemplary C₃₋₁₀
 30 carbocyclyl groups include, without limitation, the aforementioned C₃₋₈ carbocyclyl groups as well as cyclononyl (C₉), cyclononenyl (C₉), cyclodecyl (C₁₀), cyclodecenyl (C₁₀), octahydro-1*H*-indenyl (C₉), decahydronaphthalenyl (C₁₀), spiro[4.5]decanyl (C₁₀), and the like. As the foregoing examples illustrate, in certain embodiments, the carbocyclyl group is either monocyclic ("monocyclic carbocyclyl") or contain a fused, bridged or spiro ring

system such as a bicyclic system ("bicyclic carbocyclyl") and can be saturated or can be partially unsaturated. "Carbocyclyl" also includes ring systems wherein the carbocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups wherein the point of attachment is on the carbocyclyl ring, and in such instances, the number of carbons continue to designate the number of carbons in the carbocyclic ring system. Unless otherwise specified, each instance of a carbocyclyl group is independently optionally substituted, *i.e.*, unsubstituted (an "unsubstituted carbocyclyl") or substituted (a "substituted carbocyclyl") with one or more substituents. In certain embodiments, the carbocyclyl group is unsubstituted C₃₋₁₀ carbocyclyl. In certain embodiments, the carbocyclyl group is a substituted C₃₋₁₀ carbocyclyl.

In some embodiments, "carbocyclyl" is a monocyclic, saturated carbocyclyl group having from 3 to 10 ring carbon atoms ("C₃₋₁₀ cycloalkyl"). In some embodiments, a cycloalkyl group has 3 to 8 ring carbon atoms ("C₃₋₈ cycloalkyl"). In some embodiments, a cycloalkyl group has 3 to 6 ring carbon atoms ("C₃₋₆ cycloalkyl"). In some embodiments, a cycloalkyl group has 5 to 6 ring carbon atoms ("C₅₋₆ cycloalkyl"). In some embodiments, a cycloalkyl group has 5 to 10 ring carbon atoms ("C₅₋₁₀ cycloalkyl"). Examples of C₅₋₆ cycloalkyl groups include cyclopentyl (C₅) and cyclohexyl (C₆). Examples of C₃₋₆ cycloalkyl groups include the aforementioned C₅₋₆ cycloalkyl groups as well as cyclopropyl (C₃) and cyclobutyl (C₄). Examples of C₃₋₈ cycloalkyl groups include the aforementioned C₃₋₆ cycloalkyl groups as well as cycloheptyl (C₇) and cyclooctyl (C₈). Unless otherwise specified, each instance of a cycloalkyl group is independently unsubstituted (an "unsubstituted cycloalkyl") or substituted (a "substituted cycloalkyl") with one or more substituents. In certain embodiments, the cycloalkyl group is unsubstituted C₃₋₁₀ cycloalkyl. In certain embodiments, the cycloalkyl group is substituted C₃₋₁₀ cycloalkyl.

"Heterocyclyl" or "heterocyclic" refers to a radical of a 3- to 10-membered non-aromatic ring system having ring carbon atoms and 1 to 4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("3-10 membered heterocyclyl"). In heterocyclyl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. A heterocyclyl group can either be monocyclic ("monocyclic heterocyclyl") or a fused, bridged or spiro ring system such as a bicyclic system ("bicyclic heterocyclyl"), and can be saturated or can be partially unsaturated. Heterocyclyl bicyclic ring systems can include one or more heteroatoms in one or both rings.

“Heterocyclyl” also includes ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more carbocyclyl groups wherein the point of attachment is either on the carbocyclyl or heterocyclyl ring, or ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups, wherein the point of attachment is on the heterocyclyl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heterocyclyl ring system. Unless otherwise specified, each instance of heterocyclyl is independently optionally substituted, *i.e.*, unsubstituted (an “unsubstituted heterocyclyl”) or substituted (a “substituted heterocyclyl”) with one or more substituents. In certain embodiments, the heterocyclyl group is unsubstituted 3–10 membered heterocyclyl. In certain embodiments, the heterocyclyl group is substituted 3–10 membered heterocyclyl.

In some embodiments, a heterocyclyl group is a 5–10 membered non-aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–10 membered heterocyclyl”). In some embodiments, a heterocyclyl group is a 5–8 membered non-aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–8 membered heterocyclyl”). In some embodiments, a heterocyclyl group is a 5–6 membered non-aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur (“5–6 membered heterocyclyl”). In some embodiments, the 5–6 membered heterocyclyl has 1–3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heterocyclyl has 1–2 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heterocyclyl has one ring heteroatom selected from nitrogen, oxygen, and sulfur.

Exemplary 3-membered heterocyclyl groups containing one heteroatom include, without limitation, azirdinyl, oxiranyl, thioenyl. Exemplary 4-membered heterocyclyl groups containing one heteroatom include, without limitation, azetidiny, oxetanyl and thietanyl. Exemplary 5-membered heterocyclyl groups containing one heteroatom include, without limitation, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothiophenyl, dihydrothiophenyl, pyrrolidinyl, dihydropyrrolyl and pyrrolyl-2,5-dione. Exemplary 5-membered heterocyclyl groups containing two heteroatoms include, without limitation, dioxolanyl, oxasulfuranyl, disulfuranyl, and oxazolidin-2-one. Exemplary 5-membered heterocyclyl groups containing

three heteroatoms include, without limitation, triazoliny, oxadiazoliny, and thiadiazoliny. Exemplary 6-membered heterocyclyl groups containing one heteroatom include, without limitation, piperidiny, tetrahydropyrany, dihydropyridiny, and thianyl. Exemplary 6-membered heterocyclyl groups containing two heteroatoms include, without limitation, 5 piperaziny, morpholiny, dithianyl, and dioxany. Exemplary 6-membered heterocyclyl groups containing two heteroatoms include, without limitation, triazinany. Exemplary 7-membered heterocyclyl groups containing one heteroatom include, without limitation, azepany, oxepany and thiepany. Exemplary 8-membered heterocyclyl groups containing one heteroatom include, without limitation, azocany, oxecany, and thiocany. Exemplary 5- 10 membered heterocyclyl groups fused to a C₆ aryl ring (also referred to herein as a 5,6-bicyclic heterocyclic ring) include, without limitation, indoliny, isoindoliny, dihydrobenzofurany, dihydrobenzothienyl, benzoxazolinony, and the like. Exemplary 6-membered heterocyclyl groups fused to an aryl ring (also referred to herein as a 6,6-bicyclic heterocyclic ring) include, without limitation, tetrahydroquinoliny, tetrahydroisoquinoliny, and the like.

15 “Aryl” refers to a radical of a monocyclic or polycyclic (*e.g.*, bicyclic or tricyclic) 4n+2 aromatic ring system (*e.g.*, having 6, 10, or 14 π electrons shared in a cyclic array) having 6–14 ring carbon atoms and zero heteroatoms provided in the aromatic ring system (“C_{6–14} aryl”). In some embodiments, an aryl group has six ring carbon atoms (“C₆ aryl”; *e.g.*, 20 phenyl). In some embodiments, an aryl group has ten ring carbon atoms (“C₁₀ aryl”; *e.g.*, naphthyl such as 1-naphthyl and 2-naphthyl). In some embodiments, an aryl group has fourteen ring carbon atoms (“C₁₄ aryl”; *e.g.*, anthracyl). “Aryl” also includes ring systems wherein the aryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the radical or point of attachment is on the aryl ring, and in such instances, 25 the number of carbon atoms continue to designate the number of carbon atoms in the aryl ring system. Unless otherwise specified, each instance of an aryl group is independently optionally substituted, *i.e.*, unsubstituted (an “unsubstituted aryl”) or substituted (a “substituted aryl”) with one or more substituents. In certain embodiments, the aryl group is unsubstituted C_{6–14} aryl. In certain embodiments, the aryl group is substituted C_{6–14} aryl.

30 “Heteroaryl” refers to a radical of a 5–10 membered monocyclic or bicyclic 4n+2 aromatic ring system (*e.g.*, having 6 or 10 π electrons shared in a cyclic array) having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each

heteroatom is independently selected from nitrogen, oxygen and sulfur ("5-10 membered heteroaryl"). In heteroaryl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. Heteroaryl bicyclic ring systems can include one or more heteroatoms in one or both rings. "Heteroaryl" includes
5 ring systems wherein the heteroaryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the point of attachment is on the heteroaryl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heteroaryl ring system. "Heteroaryl" also includes ring systems wherein the heteroaryl ring, as defined above, is fused with one or more aryl groups wherein the point of
10 attachment is either on the aryl or heteroaryl ring, and in such instances, the number of ring members designates the number of ring members in the fused (aryl/heteroaryl) ring system. Bicyclic heteroaryl groups wherein one ring does not contain a heteroatom (*e.g.*, indolyl, quinolinyl, carbazolyl, and the like) the point of attachment can be on either ring, *i.e.*, either the ring bearing a heteroatom (*e.g.*, 2-indolyl) or the ring that does not contain a heteroatom
15 (*e.g.*, 5-indolyl).

In some embodiments, a heteroaryl group is a 5-10 membered aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5-10
20 membered heteroaryl"). In some embodiments, a heteroaryl group is a 5-8 membered aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5-8 membered heteroaryl"). In some embodiments, a heteroaryl group is a 5-6 membered aromatic ring system having ring carbon atoms and 1-4 ring heteroatoms
25 provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5-6 membered heteroaryl"). In some embodiments, the 5-6 membered heteroaryl has 1-3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5-6 membered heteroaryl has 1-2 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5-6 membered
30 heteroaryl has 1 ring heteroatom selected from nitrogen, oxygen, and sulfur. Unless otherwise specified, each instance of a heteroaryl group is independently optionally substituted, *i.e.*, unsubstituted ("unsubstituted heteroaryl") or substituted ("substituted heteroaryl") with one or more substituents. In certain embodiments, the heteroaryl group is

unsubstituted 5–14 membered heteroaryl. In certain embodiments, the heteroaryl group is substituted 5–14 membered heteroaryl.

- Exemplary 5-membered heteroaryl groups containing one heteroatom include, without
5 limitation, pyrrolyl, furanyl and thiophenyl. Exemplary 5-membered heteroaryl groups containing two heteroatoms include, without limitation, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, and isothiazolyl. Exemplary 5-membered heteroaryl groups containing three heteroatoms include, without limitation, triazolyl, oxadiazolyl, and thiadiazolyl. Exemplary 5-membered heteroaryl groups containing four heteroatoms include, without
10 limitation, tetrazolyl. Exemplary 6-membered heteroaryl groups containing one heteroatom include, without limitation, pyridinyl. Exemplary 6-membered heteroaryl groups containing two heteroatoms include, without limitation, pyridazinyl, pyrimidinyl, and pyrazinyl. Exemplary 6-membered heteroaryl groups containing three or four heteroatoms include, without limitation, triazinyl and tetrazinyl, respectively. Exemplary 7-membered heteroaryl
15 groups containing one heteroatom include, without limitation, azepinyl, oxepinyl, and thiepinyl. Exemplary 5,6-bicyclic heteroaryl groups include, without limitation, indolyl, isoindolyl, indazolyl, benzotriazolyl, benzothiophenyl, isobenzothiophenyl, benzofuranyl, benzoisofuranyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzoxadiazolyl, benzthiazolyl, benzisothiazolyl, benzthiadiazolyl, indoliziny, and purinyl. Exemplary 6,6-
20 bicyclic heteroaryl groups include, without limitation, naphthyridinyl, pteridinyl, quinolinyl, isoquinolinyl, cinnolinyl, quinoxalinyl, phthalazinyl, and quinazolinyl.

- “Partially unsaturated” refers to a group that includes at least one double or triple bond. The term “partially unsaturated” is intended to encompass rings having multiple sites of
25 unsaturation, but is not intended to include aromatic groups (*e.g.*, aryl or heteroaryl groups) as herein defined. Likewise, “saturated” refers to a group that does not contain a double or triple bond, *i.e.*, contains all single bonds.

- Alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl groups, as defined
30 herein, are optionally substituted (*e.g.*, “substituted” or “unsubstituted” alkyl, “substituted” or “unsubstituted” alkenyl, “substituted” or “unsubstituted” alkynyl, “substituted” or “unsubstituted” carbocyclyl, “substituted” or “unsubstituted” heterocyclyl, “substituted” or “unsubstituted” aryl or “substituted” or “unsubstituted” heteroaryl group). In general, the term “substituted”, whether preceded by the term “optionally” or not, means that at least one

hydrogen present on a group (e.g., a carbon or nitrogen atom) is replaced with a permissible substituent, e.g., a substituent which upon substitution results in a stable compound, e.g., a compound which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction. Unless otherwise indicated, a “substituted” group has a substituent at one or more substitutable positions of the group, and when more than one position in any given structure is substituted, the substituent is either the same or different at each position. The term “substituted” is contemplated to include substitution with all permissible substituents of organic compounds, any of the substituents described herein that results in the formation of a stable compound. The present invention contemplates any and all such combinations in order to arrive at a stable compound. For purposes of this invention, heteroatoms such as nitrogen may have hydrogen substituents and/or any suitable substituent as described herein which satisfy the valencies of the heteroatoms and results in the formation of a stable moiety.

Exemplary carbon atom substituents include, but are not limited to, halogen, $-\text{CN}$, $-\text{NO}_2$, $-\text{N}_3$, $-\text{SO}_2\text{H}$, $-\text{SO}_3\text{H}$, $-\text{OH}$, $-\text{OR}^{\text{aa}}$, $-\text{ON}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_2$, $-\text{N}(\text{R}^{\text{bb}})_3^+\text{X}^-$, $-\text{N}(\text{OR}^{\text{cc}})\text{R}^{\text{bb}}$, $-\text{SH}$, $-\text{SR}^{\text{aa}}$, $-\text{SSR}^{\text{cc}}$, $-\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{CO}_2\text{H}$, $-\text{CHO}$, $-\text{C}(\text{OR}^{\text{cc}})_2$, $-\text{CO}_2\text{R}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OCO}_2\text{R}^{\text{aa}}$, $-\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{CO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{C}(=\text{O})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{R}^{\text{aa}}$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{OR}^{\text{aa}}$, $-\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OC}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{C}(=\text{NR}^{\text{bb}})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{NR}^{\text{bb}}\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{SO}_2\text{R}^{\text{aa}}$, $-\text{SO}_2\text{OR}^{\text{aa}}$, $-\text{OSO}_2\text{R}^{\text{aa}}$, $-\text{S}(=\text{O})\text{R}^{\text{aa}}$, $-\text{OS}(=\text{O})\text{R}^{\text{aa}}$, $-\text{Si}(\text{R}^{\text{aa}})_3$, $-\text{OSi}(\text{R}^{\text{aa}})_3$, $-\text{C}(=\text{S})\text{N}(\text{R}^{\text{bb}})_2$, $-\text{C}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{C}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{S})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{OC}(=\text{O})\text{SR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{OR}^{\text{aa}}$, $-\text{SC}(=\text{O})\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})_2\text{R}^{\text{aa}}$, $-\text{OP}(=\text{O})_2\text{R}^{\text{aa}}$, $-\text{P}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{OP}(=\text{O})(\text{R}^{\text{aa}})_2$, $-\text{OP}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{P}(=\text{O})_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{OP}(=\text{O})_2\text{N}(\text{R}^{\text{bb}})_2$, $-\text{P}(=\text{O})(\text{NR}^{\text{bb}})_2$, $-\text{OP}(=\text{O})(\text{NR}^{\text{bb}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{OR}^{\text{cc}})_2$, $-\text{NR}^{\text{bb}}\text{P}(=\text{O})(\text{NR}^{\text{bb}})_2$, $-\text{P}(\text{R}^{\text{cc}})_2$, $-\text{P}(\text{R}^{\text{cc}})_3$, $-\text{OP}(\text{R}^{\text{cc}})_2$, $-\text{OP}(\text{R}^{\text{cc}})_3$, $-\text{B}(\text{R}^{\text{aa}})_2$, $-\text{B}(\text{OR}^{\text{cc}})_2$, $-\text{BR}^{\text{aa}}(\text{OR}^{\text{cc}})$, C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

or two geminal hydrogens on a carbon atom are replaced with the group $=\text{O}$, $=\text{S}$, $=\text{NN}(\text{R}^{\text{bb}})_2$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{R}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{C}(=\text{O})\text{OR}^{\text{aa}}$, $=\text{NNR}^{\text{bb}}\text{S}(=\text{O})_2\text{R}^{\text{aa}}$, $=\text{NR}^{\text{bb}}$, or $=\text{NOR}^{\text{cc}}$; each instance of R^{aa} is, independently, selected from C_{1-10} alkyl, C_{1-10} perhaloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{3-10} carbocyclyl, 3–14 membered heterocyclyl, C_{6-14} aryl, and 5–14 membered heteroaryl, or two R^{aa} groups are joined to form a 3–14 membered

heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

- each instance of R^{bb} is, independently, selected from hydrogen, –OH, –OR^{aa}, –N(R^{cc})₂, –CN, –C(=O)R^{aa}, –C(=O)N(R^{cc})₂, –CO₂R^{aa}, –SO₂R^{aa}, –C(=NR^{cc})OR^{aa}, –C(=NR^{cc})N(R^{cc})₂, –SO₂N(R^{cc})₂, –SO₂R^{cc}, –SO₂OR^{cc}, –SOR^{aa}, –C(=S)N(R^{cc})₂, –C(=O)SR^{cc}, –C(=S)SR^{cc}, –P(=O)₂R^{aa}, –P(=O)(R^{aa})₂, –P(=O)₂N(R^{cc})₂, –P(=O)(NR^{cc})₂, C_{1–10} alkyl, C_{1–10} perhaloalkyl, C_{2–10} alkenyl, C_{2–10} alkynyl, C_{3–10} carbocyclyl, 3–14 membered heterocyclyl, C_{6–14} aryl, and 5–14 membered heteroaryl, or two R^{bb} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

- each instance of R^{cc} is, independently, selected from hydrogen, C_{1–10} alkyl, C_{1–10} perhaloalkyl, C_{2–10} alkenyl, C_{2–10} alkynyl, C_{3–10} carbocyclyl, 3–14 membered heterocyclyl, C_{6–14} aryl, and 5–14 membered heteroaryl, or two R^{cc} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups;

- each instance of R^{dd} is, independently, selected from halogen, –CN, –NO₂, –N₃, –SO₂H, –SO₃H, –OH, –OR^{ee}, –ON(R^{ff})₂, –N(R^{ff})₂, –N(R^{ff})₃⁺X[–], –N(OR^{ee})R^{ff}, –SH, –SR^{ee}, –SSR^{ee}, –C(=O)R^{ee}, –CO₂H, –CO₂R^{ee}, –OC(=O)R^{ee}, –OCO₂R^{ee}, –C(=O)N(R^{ff})₂, –OC(=O)N(R^{ff})₂, –NR^{ff}C(=O)R^{ee}, –NR^{ff}CO₂R^{ee}, –NR^{ff}C(=O)N(R^{ff})₂, –C(=NR^{ff})OR^{ee}, –OC(=NR^{ff})R^{ee}, –OC(=NR^{ff})OR^{ee}, –C(=NR^{ff})N(R^{ff})₂, –OC(=NR^{ff})N(R^{ff})₂, –NR^{ff}C(=NR^{ff})N(R^{ff})₂, –NR^{ff}SO₂R^{ee}, –SO₂N(R^{ff})₂, –SO₂R^{ee}, –SO₂OR^{ee}, –OSO₂R^{ee}, –S(=O)R^{ee}, –Si(R^{ee})₃, –OSi(R^{ee})₃, –C(=S)N(R^{ff})₂, –C(=O)SR^{ee}, –C(=S)SR^{ee}, –SC(=S)SR^{ee}, –P(=O)₂R^{ee}, –P(=O)(R^{ee})₂, –OP(=O)(R^{ee})₂, –OP(=O)(OR^{ee})₂, C_{1–6} alkyl, C_{1–6} perhaloalkyl, C_{2–6} alkenyl, C_{2–6} alkynyl, C_{3–10} carbocyclyl, 3–10 membered heterocyclyl, C_{6–10} aryl, 5–10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups, or two geminal R^{dd} substituents can be joined to form =O or =S;

each instance of R^{ee} is, independently, selected from C_{1–6} alkyl, C_{1–6} perhaloalkyl, C_{2–6} alkenyl, C_{2–6} alkynyl, C_{3–10} carbocyclyl, C_{6–10} aryl, 3–10 membered heterocyclyl, and 3–10 membered heteroaryl, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups;

each instance of R^{ff} is, independently, selected from hydrogen, C_{1-6} alkyl, C_{1-6} perhaloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, 3–10 membered heterocyclyl, C_{6-10} aryl and 5–10 membered heteroaryl, or two R^{ff} groups are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{gg} groups; and

each instance of R^{gg} is, independently, halogen, $-CN$, $-NO_2$, $-N_3$, $-SO_2H$, $-SO_3H$, $-OH$, $-OC_{1-6}$ alkyl, $-ON(C_{1-6}$ alkyl) $_2$, $-N(C_{1-6}$ alkyl) $_2$, $-N(C_{1-6}$ alkyl) $_3^+X^-$, $-NH(C_{1-6}$ alkyl) $_2^+X^-$, $-NH_2(C_{1-6}$ alkyl) $^+X^-$, $-NH_3^+X^-$, $-N(OC_{1-6}$ alkyl)(C_{1-6} alkyl), $-N(OH)(C_{1-6}$ alkyl), $-NH(OH)$, $-SH$, $-SC_{1-6}$ alkyl, $-SS(C_{1-6}$ alkyl), $-C(=O)(C_{1-6}$ alkyl), $-CO_2H$, $-CO_2(C_{1-6}$ alkyl), $-OC(=O)(C_{1-6}$ alkyl), $-OCO_2(C_{1-6}$ alkyl), $-C(=O)NH_2$, $-C(=O)N(C_{1-6}$ alkyl) $_2$, $-OC(=O)NH(C_{1-6}$ alkyl), $-NHC(=O)(C_{1-6}$ alkyl), $-N(C_{1-6}$ alkyl) $C(=O)(C_{1-6}$ alkyl), $-NHCO_2(C_{1-6}$ alkyl), $-NHC(=O)N(C_{1-6}$ alkyl) $_2$, $-NHC(=O)NH(C_{1-6}$ alkyl), $-NHC(=O)NH_2$, $-C(=NH)O(C_{1-6}$ alkyl), $-OC(=NH)(C_{1-6}$ alkyl), $-OC(=NH)OC_{1-6}$ alkyl, $-C(=NH)N(C_{1-6}$ alkyl) $_2$, $-C(=NH)NH(C_{1-6}$ alkyl), $-C(=NH)NH_2$, $-OC(=NH)N(C_{1-6}$ alkyl) $_2$, $-OC(NH)NH(C_{1-6}$ alkyl), $-OC(NH)NH_2$, $-NHC(NH)N(C_{1-6}$ alkyl) $_2$, $-NHC(=NH)NH_2$, $-NHOSO_2(C_{1-6}$ alkyl), $-SO_2N(C_{1-6}$ alkyl) $_2$, $-SO_2NH(C_{1-6}$ alkyl), $-SO_2NH_2$, $-SO_2C_{1-6}$ alkyl, $-SO_2OC_{1-6}$ alkyl, $-OSO_2C_{1-6}$ alkyl, $-SOC_{1-6}$ alkyl, $-Si(C_{1-6}$ alkyl) $_3$, $-OSi(C_{1-6}$ alkyl) $_3$, $-C(=S)N(C_{1-6}$ alkyl) $_2$, $-C(=S)NH(C_{1-6}$ alkyl), $-C(=S)NH_2$, $-C(=O)S(C_{1-6}$ alkyl), $-C(=S)SC_{1-6}$ alkyl, $-SC(=S)SC_{1-6}$ alkyl, $-P(=O)_2(C_{1-6}$ alkyl), $-P(=O)(C_{1-6}$ alkyl) $_2$, $-OP(=O)(C_{1-6}$ alkyl) $_2$, $-OP(=O)(OC_{1-6}$ alkyl) $_2$, C_{1-6} alkyl, C_{1-6} perhaloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} carbocyclyl, C_{6-10} aryl, 3–10 membered heterocyclyl, 5–10 membered heteroaryl; or two geminal R^{gg} substituents can be joined to form $=O$ or $=S$; wherein X^- is a counterion.

A “counterion” or “anionic counterion” is a negatively charged group associated with a cationic quaternary amino group in order to maintain electronic neutrality. Exemplary counterions include halide ions (*e.g.*, F^- , Cl^- , Br^- , I^-), NO_3^- , ClO_4^- , OH^- , $H_2PO_4^-$, HSO_4^- , sulfonate ions (*e.g.*, methanesulfonate, trifluoromethanesulfonate, *p*-toluenesulfonate, benzenesulfonate, 10-camphor sulfonate, naphthalene-2-sulfonate, naphthalene-1-sulfonic acid-5-sulfonate, ethan-1-sulfonic acid-2-sulfonate, and the like), and carboxylate ions (*e.g.*, acetate, ethanoate, propanoate, benzoate, glycerate, lactate, tartrate, glycolate, and the like).

"Halo" or "halogen" refers to fluorine (fluoro, -F), chlorine (chloro, -Cl), bromine (bromo, -Br), or iodine (iodo, -I).

Nitrogen atoms can be substituted or unsubstituted as valency permits, and include primary, secondary, tertiary, and quaternary nitrogen atoms. Exemplary nitrogen atom substituents include, but are not limited to, hydrogen, -OH, -OR^{aa}, -N(R^{cc})₂, -CN, -C(=O)R^{aa}, -C(=O)N(R^{cc})₂, -CO₂R^{aa}, -SO₂R^{aa}, -C(=NR^{bb})R^{aa}, -C(=NR^{cc})OR^{aa}, -C(=NR^{cc})N(R^{cc})₂, -SO₂N(R^{cc})₂, -SO₂R^{cc}, -SO₂OR^{cc}, -SOR^{aa}, -C(=S)N(R^{cc})₂, -C(=O)SR^{cc}, -C(=S)SR^{cc}, -P(=O)₂R^{aa}, -P(=O)(R^{aa})₂, -P(=O)₂N(R^{cc})₂, -P(=O)(NR^{cc})₂, C₁₋₁₀ alkyl, C₁₋₁₀ perhaloalkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₁₀ carbocyclyl, 3-14 membered heterocyclyl, C₆₋₁₄ aryl, and 5-14 membered heteroaryl, or two R^{cc} groups attached to a nitrogen atom are joined to form a 3-14 membered heterocyclyl or 5-14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa}, R^{bb}, R^{cc} and R^{dd} are as defined above.

15

In certain embodiments, the substituent present on a nitrogen atom is a nitrogen protecting group (also referred to as an amino protecting group). Nitrogen protecting groups include, but are not limited to, -OH, -OR^{aa}, -N(R^{cc})₂, -C(=O)R^{aa}, -C(=O)N(R^{cc})₂, -CO₂R^{aa}, -SO₂R^{aa}, -C(=NR^{cc})R^{aa}, -C(=NR^{cc})OR^{aa}, -C(=NR^{cc})N(R^{cc})₂, -SO₂N(R^{cc})₂, -SO₂R^{cc}, -SO₂OR^{cc}, -SOR^{aa}, -C(=S)N(R^{cc})₂, -C(=O)SR^{cc}, -C(=S)SR^{cc}, C₁₋₁₀ alkyl (e.g., aralkyl, heteroaralkyl), C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, C₃₋₁₀ carbocyclyl, 3-14 membered heterocyclyl, C₆₋₁₄ aryl, and 5-14 membered heteroaryl groups, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aralkyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa}, R^{bb}, R^{cc}, and R^{dd} are as defined herein. Nitrogen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

Amide nitrogen protecting groups (e.g., -C(=O)R^{aa}) include, but are not limited to, formamide, acetamide, chloroacetamide, trichloroacetamide, trifluoroacetamide, phenylacetamide, 3-phenylpropanamide, picolinamide, 3-pyridylcarboxamide, *N*-benzoylphenylalanyl derivative, benzamide, *p*-phenylbenzamide, *o*-nitrophenylacetamide, *o*-nitrophenoxyacetamide, acetoacetamide, (*N*'-dithiobenzylxyacylamino)acetamide, 3-(*p*-hydroxyphenyl)propanamide, 3-(*o*-nitrophenyl)propanamide, 2-methyl-2-(*o*-

nitrophenoxy)propanamide, 2-methyl-2-(*o*-phenylazophenoxy)propanamide, 4-chlorobutanamide, 3-methyl-3-nitrobutanamide, *o*-nitrocinnamide, *N*-acetylmethionine, *o*-nitrobenzamide, and *o*-(benzoyloxymethyl)benzamide.

- 5 Carbamate nitrogen protecting groups (*e.g.*, $\text{-C(=O)OR}^{\text{aa}}$) include, but are not limited to, methyl carbamate, ethyl carbamate, 9-fluorenylmethyl carbamate (Fmoc), 9-(2-sulfo)fluorenylmethyl carbamate, 9-(2,7-dibromo)fluorenylmethyl carbamate, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10-tetrahydrothioxanthyl)]methyl carbamate (DBD-Tmoc), 4-methoxyphenacyl carbamate (Phenoc), 2,2,2-trichloroethyl carbamate (Troc), 2-
- 10 trimethylsilylethyl carbamate (Teoc), 2-phenylethyl carbamate (hZ), 1-(1-adamantyl)-1-methylethyl carbamate (Adpoc), 1,1-dimethyl-2-haloethyl carbamate, 1,1-dimethyl-2,2-dibromoethyl carbamate (DB-*t*-BOC), 1,1-dimethyl-2,2,2-trichloroethyl carbamate (TCBOC), 1-methyl-1-(4-biphenyl)ethyl carbamate (Bpoc), 1-(3,5-di-*t*-butylphenyl)-1-methylethyl carbamate (*t*-Bumeoc), 2-(2'- and 4'-pyridyl)ethyl carbamate (Pyoc), 2-(*N,N*-
- 15 dicyclohexylcarboxamido)ethyl carbamate, *t*-butyl carbamate (BOC), 1-adamantyl carbamate (Adoc), vinyl carbamate (Voc), allyl carbamate (Alloc), 1-isopropylallyl carbamate (Ipaoc), cinnamyl carbamate (Coc), 4-nitrocinnamyl carbamate (Noc), 8-quinolyl carbamate, *N*-hydroxypiperidiny carbamate, alkylidithio carbamate, benzyl carbamate (Cbz), *p*-methoxybenzyl carbamate (Moz), *p*-nitrobenzyl carbamate, *p*-bromobenzyl carbamate, *p*-
- 20 chlorobenzyl carbamate, 2,4-dichlorobenzyl carbamate, 4-methylsulfinylbenzyl carbamate (MsZ), 9-anthrylmethyl carbamate, diphenylmethyl carbamate, 2-methylthioethyl carbamate, 2-methylsulfonylethyl carbamate, 2-(*p*-toluenesulfonyl)ethyl carbamate, [2-(1,3-dithianyl)]methyl carbamate (Dmoc), 4-methylthiophenyl carbamate (Mtpc), 2,4-dimethylthiophenyl carbamate (Bmpc), 2-phosphonioethyl carbamate (Peoc), 2-
- 25 triphenylphosphonioisopropyl carbamate (Ppoc), 1,1-dimethyl-2-cyanoethyl carbamate, *m*-chloro-*p*-acyloxybenzyl carbamate, *p*-(dihydroxyboryl)benzyl carbamate, 5-benzisoxazolylmethyl carbamate, 2-(trifluoromethyl)-6-chromonylmethyl carbamate (Troc), *m*-nitrophenyl carbamate, 3,5-dimethoxybenzyl carbamate, *o*-nitrobenzyl carbamate, 3,4-dimethoxy-6-nitrobenzyl carbamate, phenyl(*o*-nitrophenyl)methyl
- 30 carbamate, *t*-amyl carbamate, *S*-benzyl thiocarbamate, *p*-cyanobenzyl carbamate, cyclobutyl carbamate, cyclohexyl carbamate, cyclopentyl carbamate, cyclopropylmethyl carbamate, *p*-decyloxybenzyl carbamate, 2,2-dimethoxyacetylvinyl carbamate, *o*-(*N,N*-dimethylcarboxamido)benzyl carbamate, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl carbamate, 1,1-dimethylpropynyl carbamate, di(2-pyridyl)methyl carbamate, 2-

furanylmethyl carbamate, 2-iodoethyl carbamate, isoborynl carbamate, isobutyl carbamate, isonicotinyln carbamate, *p*-(*p*'-methoxyphenylazo)benzyl carbamate, 1-methylcyclobutyl carbamate, 1-methylcyclohexyl carbamate, 1-methyl-1-cyclopropylmethyl carbamate, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl carbamate, 1-methyl-1-(*p*-phenylazophenyl)ethyl carbamate, 1-methyl-1-phenylethyl carbamate, 1-methyl-1-(4-pyridyl)ethyl carbamate, phenyl carbamate, *p*-(phenylazo)benzyl carbamate, 2,4,6-tri-*t*-butylphenyl carbamate, 4-(trimethylammonium)benzyl carbamate, and 2,4,6-trimethylbenzyl carbamate.

Sulfonamide nitrogen protecting groups (*e.g.*, $-S(=O)_2R^{aa}$) include, but are not limited to, *p*-toluenesulfonamide (Ts), benzenesulfonamide, 2,3,6-trimethyl-4-methoxybenzenesulfonamide (Mtr), 2,4,6-trimethoxybenzenesulfonamide (Mtb), 2,6-dimethyl-4-methoxybenzenesulfonamide (Pme), 2,3,5,6-tetramethyl-4-methoxybenzenesulfonamide (Mte), 4-methoxybenzenesulfonamide (Mbs), 2,4,6-trimethylbenzenesulfonamide (Mts), 2,6-dimethoxy-4-methylbenzenesulfonamide (iMds), 2,2,5,7,8-pentamethylchroman-6-sulfonamide (Pmc), methanesulfonamide (Ms), β -trimethylsilylthanesulfonamide (SES), 9-anthracenesulfonamide, 4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonamide (DNMBS), benzylsulfonamide, trifluoromethylsulfonamide, and phenacysulfonamide.

Other nitrogen protecting groups include, but are not limited to, phenothiazinyl-(10)-acyl derivative, *N*'-*p*-toluenesulfonylaminoacyl derivative, *N*'-phenylaminothioacyl derivative, *N*-benzoylphenylalanyl derivative, *N*-acetylmethionine derivative, 4,5-diphenyl-3-oxazolin-2-one, *N*-phthalimide, *N*-dithiasuccinimide (Dts), *N*-2,3-diphenylmaleimide, *N*-2,5-dimethylpyrrole, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct (STABASE), 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridone, *N*-methylamine, *N*-allylamine, *N*-[2-(trimethylsilyl)ethoxy]methylamine (SEM), *N*-3-acetoxypyrrolamine, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrroline-3-yl)amine, quaternary ammonium salts, *N*-benzylamine, *N*-di(4-methoxyphenyl)methylamine, *N*-5-dibenzosuberylamine, *N*-triphenylmethylamine (Tr), *N*-[(4-methoxyphenyl)diphenylmethyl]amine (MMTr), *N*-9-phenylfluorenylamine (PhF), *N*-2,7-dichloro-9-fluorenylmethyleneamine, *N*-ferrocenylmethylamino (Fcm), *N*-2-picolylamino *N*'-oxide, *N*-1,1-dimethylthiomethyleneamine, *N*-benzylideneamine, *N*-*p*-methoxybenzylideneamine, *N*-diphenylmethyleneamine, *N*-[(2-pyridyl)mesityl]methyleneamine, *N*-(*N*',*N*'-

dimethylaminomethylene)amine, *N,N'*-isopropylidenediamine, *N-p*-nitrobenzylideneamine, *N*-salicylideneamine, *N*-5-chlorosalicylideneamine, *N*-(5-chloro-2-hydroxyphenyl)phenylmethyleamine, *N*-cyclohexylideneamine, *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)amine, *N*-borane derivative, *N*-diphenylborinic acid derivative, *N*-

5 [phenyl(pentaacylchromium- or tungsten)acyl]amine, *N*-copper chelate, *N*-zinc chelate, *N*-nitroamine, *N*-nitrosoamine, amine *N*-oxide, diphenylphosphinamide (Dpp), dimethylthiophosphinamide (Mpt), diphenylthiophosphinamide (Ppt), dialkyl phosphoramidates, dibenzyl phosphoramidate, diphenyl phosphoramidate, benzenesulfenamide, *o*-nitrobenzenesulfenamide (Nps), 2,4-dinitrobenzenesulfenamide,

10 pentachlorobenzenesulfenamide, 2-nitro-4-methoxybenzenesulfenamide, triphenylmethylsulfenamide, and 3-nitropyridinesulfenamide (Npys).

In certain embodiments, the substituent present on an oxygen atom is an oxygen protecting group (also referred to as a hydroxyl protecting group). Oxygen protecting groups include,

15 but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, and $-P(=O)(NR^{bb})_2$, wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein. Oxygen protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W.

20 Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

Exemplary oxygen protecting groups include, but are not limited to, methyl, methoxymethyl (MOM), methylthiomethyl (MTM), *t*-butylthiomethyl, (phenyldimethylsilyl)methoxymethyl

25 (SMOM), benzyloxymethyl (BOM), *p*-methoxybenzyloxymethyl (PMBM), (4-methoxyphenoxy)methyl (*p*-AOM), guaiacolmethyl (GUM), *t*-butoxymethyl, 4-pentenylloxymethyl (POM), siloxymethyl, 2-methoxyethoxymethyl (MEM), 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl (SEMOR), tetrahydropyranyl (THP), 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-

30 methoxycyclohexyl, 4-methoxytetrahydropyranyl (MTHP), 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl (CTMP), 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl, 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl, 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-

- benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, benzyl (Bn), *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p,p'*-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl, α -naphthylidiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxyphenyl)diphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-yl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodisulfuran-2-yl, benzisothiazolyl S,S-dioxido, trimethylsilyl (TMS), triethylsilyl (TES), triisopropylsilyl (TIPS), dimethylisopropylsilyl (IPDMS), diethylisopropylsilyl (DEIPS), dimethylthexylsilyl, *t*-butyldimethylsilyl (TBDMS), *t*-butyldiphenylsilyl (TBDPS), tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl (DPMS), *t*-butylmethoxyphenylsilyl (TBMPS), formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-(ethylenedithio)pentanoate (levulinoyldithioacetal), pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, 2,4,6-trimethylbenzoate (mesitoate), alkyl methyl carbonate, 9-fluorenylmethyl carbonate (Fmoc), alkyl ethyl carbonate, alkyl 2,2,2-trichloroethyl carbonate (Troc), 2-(trimethylsilyl)ethyl carbonate (TMSEC), 2-(phenylsulfonyl) ethyl carbonate (Psec), 2-(triphenylphosphonio) ethyl carbonate (Peoc), alkyl isobutyl carbonate, alkyl vinyl carbonate alkyl allyl carbonate, alkyl *p*-nitrophenyl carbonate, alkyl benzyl carbonate, alkyl *p*-methoxybenzyl carbonate, alkyl 3,4-dimethoxybenzyl carbonate, alkyl *o*-nitrobenzyl carbonate, alkyl *p*-nitrobenzyl carbonate, alkyl *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl carbonate, methyl dithiocarbonate, 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxy)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenate, *o*-(methoxyacyl)benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-

tetramethylphosphorodiamidate, alkyl *N*-phenylcarbamate, borate, dimethylphosphinothioyl, alkyl 2,4-dinitrophenylsulfenate, sulfate, methanesulfonate (mesylate), benzyisulfonate, and tosylate (Ts).

- 5 In certain embodiments, the substituent present on a sulfur atom is a sulfur protecting group (also referred to as a thiol protecting group). Sulfur protecting groups include, but are not limited to, $-R^{aa}$, $-N(R^{bb})_2$, $-C(=O)SR^{aa}$, $-C(=O)R^{aa}$, $-CO_2R^{aa}$, $-C(=O)N(R^{bb})_2$, $-C(=NR^{bb})R^{aa}$, $-C(=NR^{bb})OR^{aa}$, $-C(=NR^{bb})N(R^{bb})_2$, $-S(=O)R^{aa}$, $-SO_2R^{aa}$, $-Si(R^{aa})_3$, $-P(R^{cc})_2$, $-P(R^{cc})_3$, $-P(=O)_2R^{aa}$, $-P(=O)(R^{aa})_2$, $-P(=O)(OR^{cc})_2$, $-P(=O)_2N(R^{bb})_2$, and $-P(=O)(NR^{bb})_2$,
 10 wherein R^{aa} , R^{bb} , and R^{cc} are as defined herein. Sulfur protecting groups are well known in the art and include those described in detail in *Protecting Groups in Organic Synthesis*, T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, incorporated herein by reference.

- 15 These and other exemplary substituents are described in more detail in the Detailed Description, Examples, and claims. The present invention is not intended to be limited in any manner by the above exemplary listing of substituents.

Other definitions

- 20 The term "pharmaceutically acceptable form thereof" as used herein refers to pharmaceutically acceptable salts, solvates, hydrates, prodrugs, tautomers, isomers, enantiomers, diastereomers, and/or polymorphs of a compound of the present invention.

- In certain embodiments, the pharmaceutically acceptable form is a pharmaceutically acceptable salt. The term "pharmaceutically acceptable salt" as used herein refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, Berge *et al.*, describe
 25 pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric

acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, 5 camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, 10 persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and N+(C1-4alkyl)4 salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable 15 salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

20 In certain embodiments, the pharmaceutically acceptable form is a hydrate or solvate. The term "hydrate" as used herein refers to a compound non-covalently associated with one or more molecules of water. Likewise, the term "solvate" refers to a compound non-covalently associated with one or more molecules of an organic solvent.

In certain embodiments, the pharmaceutically acceptable form is a prodrug. The term 25 "prodrug" as used herein refers to a derivative of a parent compound that requires transformation within the body in order to release the parent compound. In certain cases, a prodrug has improved physical and/or delivery properties over the parent compound. Prodrugs are typically designed to enhance pharmaceutically and/or pharmacokinetically based properties associated with the parent compound. The advantage of a prodrug can lie in 30 its physical properties, such as enhanced water solubility for parenteral administration at physiological pH compared to the parent compound, or it enhances absorption from the digestive tract, or it may enhance drug stability for long-term storage. In recent years several types of bioreversible derivatives have been exploited for utilization in designing prodrugs. Using esters as a prodrug type for compounds containing a carboxyl or hydroxyl functionality

is known in the art as described, for example, in *The Organic Chemistry of Drug Design and Drug Interaction* by Richard Silverman, published by Academic Press (1992).

In certain embodiments, the pharmaceutically acceptable form is a tautomer. The term
5 “tautomer” as used herein includes two or more interconvertable compounds resulting from at least one formal migration of a hydrogen atom and at least one change in valency (e.g., a single bond to a double bond, a triple bond to a single bond, or vice versa). The exact ratio of the tautomers depends on several factors, including temperature, solvent, and pH. Tautomerizations (i.e., the reaction providing a tautomeric pair) may catalyzed by acid or
10 base. Exemplary tautomerizations include keto-to-enol; amide-to-imide; lactam-to-lactim; enamine-to-imine; and enamine-to-(a different) enamine tautomerizations.

In certain embodiments, the pharmaceutically acceptable form is an isomer. The term
“isomer” as used herein includes any and all geometric isomers and stereoisomers (e.g.,
15 enantiomers, diastereomers, etc.). For example, “isomer” include cis- and trans-isomers, E- and Z- isomers, R- and S-enantiomers, diastereomers, (D)-isomers, (L)-isomers, racemic mixtures thereof, and other mixtures thereof, as falling within the scope of the invention. For instance, an isomer/enantiomer may, in some embodiments, be provided substantially free of the corresponding enantiomer, and may also be referred to as “optically enriched.”
20 “Optically-enriched,” as used herein, means that the compound is made up of a significantly greater proportion of one enantiomer. In certain embodiments the compound of the present invention is made up of at least about 90% by weight of a preferred enantiomer. In other embodiments the compound is made up of at least about 95%, 98%, or 99% by weight of a preferred enantiomer. Preferred enantiomers may be isolated from racemic mixtures by any
25 method known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts or prepared by asymmetric syntheses. See, for example, Jacques, *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen, S.H., *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, E.L. *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); Wilen,
30 S.H. *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972).

In certain embodiments, the pharmaceutically acceptable form is a polymorph. The term
“polymorph” as used herein refers to a crystalline compound existing in more than one

crystalline form/structure. When polymorphism exists as a result of difference in crystal packing it is called packing polymorphism. Polymorphism can also result from the existence of different conformers of the same molecule in conformational polymorphism. In pseudopolymorphism the different crystal types are the result of hydration or solvation.

5

A "subject" to which administration is contemplated includes, but is not limited to, humans (i.e., a male or female of any age group, e.g., a pediatric subject (e.g, infant, child, adolescent) or adult subject (e.g., young adult, middle-aged adult or senior adult)) and/or other primates (e.g., cynomolgus monkeys, rhesus monkeys); mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, goats, cats, and/or dogs; and/or birds, including commercially relevant birds such as chickens, ducks, geese, and/or turkeys.

10

"Treat," "treating" and "treatment" encompasses an action that occurs while a subject is suffering from a condition (e.g., a "MNK1- or MNK2-related" disease, disorder, or condition, e.g., a disease, disorder, or condition in which MNK1 and/or MNK2 is known to play role) which reduces the severity of the condition or retards or slows the progression of the condition ("therapeutic treatment"). "Treat," "treating" and "treatment" also encompasses an action that occurs before a subject begins to suffer from the condition and which inhibits or reduces the severity of the condition ("prophylactic treatment").

15

20

As used herein "inhibition," "inhibiting," and "inhibit", refer to the ability of a compound to reduce, slow, halt or prevent activity of a particular biological process relative to vehicle. In certain embodiments, the biological process is *in vitro* (e.g., a biochemical or cellular assay). In certain embodiments, the biological process is *in vivo*.

25

An "effective amount" of a compound refers to an amount sufficient to elicit the desired biological response, e.g., treat the condition. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound described herein may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the condition being treated, the mode of administration, and the age and health of the subject. An effective amount encompasses therapeutic and prophylactic treatment.

30

A “therapeutically effective amount” of a compound is an amount sufficient to provide a therapeutic benefit in the treatment of a condition or to delay or minimize one or more symptoms associated with the condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment of the condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of the condition, or enhances the therapeutic efficacy of another therapeutic agent. In some embodiments, a therapeutically effective amount is an amount effective to inhibit cell growth or induce cell death.

A “prophylactically effective amount” of a compound is an amount sufficient to prevent a condition, or one or more symptoms associated with the condition or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the condition. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.


As used herein, the term “kinase” represent transferase class enzymes that are able to transfer a phosphate group from a donor molecule to an acceptor molecule, *e.g.*, an amino acid residue of a protein or a lipid molecule. Representative, non-limiting examples of kinases include Abl, ACK, Akt1/PKB α , Akt2/PKB β , Akt3/PKB γ , ALK1, ALK2, Alk4, AMPK α 1/ β 1/ γ 1, AMPK α 1/ β 1/ γ 2, AMPK α 1/ β 1/ γ 3, AMPK α 1/ β 2/ γ 1, AMPK α 2/ β 1/ γ 1, AMPK α 2/ β 2/ γ 2, Abl2, ARKS, Ask1, Aurora A, Aurora B, Aurora C, Axl, BARK1, Blk, Bmx, B-Raf, Brk, BrSK1, BrSK2, Btk, CaMK1 α , CaMK1 β , CaMK1 γ , CaMK1 δ , CAMK2 α , CaMK2 β , CAMK2 δ , CAMK2 γ , CAMK4, CAMKK1, CAMKK2, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK9, CDK1/cyclin B, CDK2/cyclin A, CDK2/cyclin E, CDK3/cyclin E, CDK5/p25, CDK5/p35, CDK6/cyclinD3, CDK7/cyclin H/MAT1, CDK9/cyclin T1, CHK1, CHK2, CK1(γ), CK1 δ , CK2 α 1, CK2 α 2, cKit, c-RAF, CLK1, CLK2, CLK3, COT, Csk, DAPK1, DAPK2, DAPK3, DCAMLK2, DDR2, DMPK, DRAK1, DYRK1A, DYRK2, DYRK3, eEF2K, EGFR, EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EphB1, EphB2, EphB3, EphB4, ErbB4, Erk1, Erk2, FAK, Fer, Fes, FGFR1, Flt2, Flt4, FLT3 D835Y, FGFR2, FGFR3, FGFR4, Fgr, Flt1, Flt3, Fms, FRK, FynA, GCK, GPRK5, GRK2, GRK4, GRK6, GRK7, GSK3 α , GSK3 β , Hck, HER2, HER4,

HIPK1, HIPK2, HIPK3, HIPK4, IGF1R, IKK β , IKK α , IKK ϵ , IR, InsR, IRR, IRAK1, IRAK2, IRAK4, Itk, JAK2, JAK3, JNK1, JNK2, JNK3, KDR, KHS1, Kit, Lck, LIMK1, LKB1, LOK, LRRK2, Lyn A, Lyn B, MAPK1, MAPK2, MAPK12, MAPKAP-K2, MAPKAP-K3, MAPKAPK2, MAPKAPK3, MAPKAPK5, MARK1, MARK2, MARK3, MARK4, MELK, MEK1, MEK2, MEKK2, MEKK3, Mer, Met, MET M1250T, MINK, MKK4, MKK6, MKK7 β , MLCK, MLK1, MLK3, MNK1, MNK2, MRCK α , MRCK β , MSK1, MSK2, MSSK1, STK23, STK4, STK3, STK24, MST1, MST2, MST3, MST4, MUSK, mTOR, MYO3 β , MYT1, NDR1, NEK11, NEK2, NEK3, NEK6, NEK7, NEK9, NLK, NUA2, p38 α , p38 β , p38 δ , p38 γ , p70S6K, S6K, SRK, PAK1/CDC42, PAK2, PAK3, PAK4, PAK5, PAK6, PAR-1B α , PASK, PBK, PDGFR α , PDGFR β , PDK1, PEK, PHKG2, PI3K α , PI3K β , PI3K γ , PI3K δ , Pim1, Pim2, PKA α , PKA β , PKA γ , PKA(b), PKA, PKB α , PKB β , PKB γ , PKC α , PKC β 1, PKC β 2, PKC β 11, PKC δ , PKC ϵ , PKC γ , PKC μ , PKC η , PKC ι , PKC θ , PKC ζ , PKD1, PKD2, PKD3, PKG1 α , PKG1B, PKN1, PKN2, PKR, PLK1, PLK2, PLK3, PLK4, Polo, PRAK, PRK2, PrKX, PTK5, PYK2, QIK, Raf1, Ret, RIPK2, RIPK5, ROCK1, ROCK2, RON, ROS, Rse, RSK1, RSK2, RSK3, RSK4, SAPK2a, SAPK2b, SAPK3, SAPK4, SGK1, SGK2, SGK3, SIK, MLCK, SLK, Snk, Src, SRPK1, SRPK2, STK33, SYK, TAK1-TAB1, TAK1, TBK1, TAO1, TAO2, TAO3, TBK1, TEC, TESK1, TGF β R1, TGF β R2, Tie2, TLK2, TrkA, TrkB, TrkC, TSSK1, TSSK2, TTK, TXK, TYK2, TYRO3, ULK1, ULK2, WEE1, WNK2, WNK3, Yes1, YSK1, ZAK, ZAP70, ZC3, and ZIPK.

20

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

In one aspect, the present invention provides a compound of Formula (I):



or a pharmaceutically acceptable form thereof, wherein

25 X_1 , X_2 , and X_3 are independently N or C;

X_4 , X_5 , X_6 are independently N or CR⁴;

wherein at least one of X_1 , X_2 , X_3 , X_4 , X_5 , and X_6 is N;

X_7 is N or CR¹⁰;

X_8 is N or CR¹¹;

---- is a single or double bond, as valency allows;

$R^1, R^2, R^3, R^5, R^6, R^7, R^8, R^9, R^{10}$, and R^{11} are independently hydrogen, halogen,
 5 optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl,
 optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂,
 -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -
 NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂;

each R is independently hydrogen, optionally substituted C₁₋₆ aliphatic, optionally
 10 substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or
 optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken
 together to form an optionally substituted heterocycle;

each R⁴ is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic;
 and

15 R^1 and R², R² and R³, R³ and R⁴, R¹ and R¹¹, R¹⁰ and R¹¹, R⁵ and R⁶, R⁶ and R⁷, R⁷
 and R⁸, or R⁸ and R⁹ may be optionally taken together to form an optionally substituted 5-6
 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

As described generally above, R¹ is hydrogen, halogen, optionally substituted aliphatic,
 20 optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted
 heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -
 CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -
 NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R¹ is hydrogen. In some embodiments,
 R¹ is halogen. In some embodiments, R¹ is -CN, -CH₂OH, -CH₂NH₂, -C(O)R_a, -CH₂R_b, -
 25 CH(OH)R_c, or -SO₂R_a, wherein R_a, R_b, and R_c are as described herein. In some
 embodiments, R¹ is -C(O)R, where R is optionally substituted heterocyclyl.

As described generally above, R² is hydrogen, halogen, optionally substituted aliphatic,
 optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted
 30 heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -
 CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -
 NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R² is hydrogen. In some embodiments,
 R² is halogen. In some embodiments, R² is -H, -F, -Cl, -CH₃, -CF₃, -CHF₂, -OH, -OC₁₋₆
 alkyl, -SR, -NH₂, -NHCH₃, -N(CH₃)₂, -NH₂Et, -N(Et)₂, -NH₂-Pr, -NC(O)R_c, SR_c, -S(O)R_e, -

S(O)₂R_e, -CN, -CH₂OH, -CH₂NH₂, -N(R)₂, -NO₂, -CONH₂, -CONHR, -CONRR, -C(O)R, C(O)R_e, -C(O)OR_e, -C(O)NR_fR_g, -C(OH)RR, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, C₄-C₇ cycloalkyl, mono or polycyclic aryl, or mono or polycyclic heteroaryl. In some
 5 embodiments, R² is optionally substituted from 1 to 3 times with substituents selected from the group consisting of halogen, -H, -OR_c, -C(O)R_b, -C(O)OR_c, -C(O)NR_fR_g, -NH₂, -NHR_d, -NR_dR_f, -SR_c, -S(O)R, -S(O)₂R, -CN, -NO₂, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, C₄-C₇ cycloalkylalkyl, and mono or polycyclic aryl. In some embodiments, R² is -H, -F, -Cl, -CN, -CH₃, -CF₃, -CHF₂, -C(O)NH₂, -OH, -OC₁₋₄alkyl, or -OCF₃.

10 In some embodiments, R¹ and R² are taken together to form an optionally substituted 5-membered ring heterocyclic or heteroaryl moiety.

As described generally above, R³ is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted
 15 heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R³ is hydrogen. In some embodiments, R³ is halogen. In some embodiments, R³ is -H, -F, -Cl, -CH₃, -CF₃, -CHF₂, -OH, or -OMe. In some embodiments, R³ is -H, -F, -Cl, -CN, -CH₃, -CF₃, -CHF₂, -C(O)NH₂, -OH, -OC₁₋₄alkyl, or -OCF₃.
 20

As described generally above, each R⁴ is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic. In some embodiments, R⁴ is hydrogen. In some embodiments, R⁴ is halogen. In some embodiments, R⁴ is fluoro or chloro. In some embodiments, R⁴ is
 25 optionally substituted C₁₋₆ aliphatic. In some embodiments, R⁴ is optionally substituted C₁₋₆ alkyl. In some embodiments, R⁴ is unsubstituted C₁₋₆ alkyl. In some embodiments, R⁴ is H or Me.


As described generally above, R⁵ is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted
 30 heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R⁵ is hydrogen. In some embodiments,

R^5 is halogen. In some embodiments, R^5 is -F, -Cl, Me, -CF₃, -OCF₃, -OH, -OMe, -CN, or -CONH₂.

As described generally above, R^6 is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R^6 is hydrogen. In some embodiments, R^6 is halogen. In some embodiments, R^6 is -F, -Cl, Me, -CF₃, -OCF₃, -OH, -OMe, -CN, or -CONH₂. In some embodiments, R^6 is -H, -F, -Cl, -CN, -CH₃, -CF₃, -CHF₂, -C(O)NH₂, -OH, -OC₁₋₄alkyl, or -OCF₃.

As described generally above, R^7 is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R^7 is hydrogen. In some embodiments, R^7 is halogen. In some embodiments, R^7 is -H, -F, -Cl, Me, -CF₂H, -CF₃, -OH, -NH₂, -NHMe, -N(Me)₂, -CO₂H, -CONH₂, -CN, -OMe, alkyne, -CH₂OH, -CH₂NH₂, -NHCOCH₃, -CONHOH, -SO₂NH₂, -SO₂Me, morpholine, 1-methylpiperazine, carbocyclyl, heterocyclyl, aryl, heteroaryl, or substituted heteroaryl. In some embodiments, R^7 is -CN. In some embodiments, R^7 is -H, -F, -Cl, -CN, -CH₃, -CF₃, -CHF₂, -C(O)NH₂, -OH, -OC₁₋₄alkyl, or -OCF₃.

In some embodiments, R^6 and R^7 are taken together to form an optionally substituted 5- or 6-membered ring heterocyclic or heteroaryl moiety. In certain embodiments, a heterocyclic or heteroaryl moiety formed by R^6 and R^7 together with the benzene ring fused thereto is one of the following formulae:



As described generally above, R^8 is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R^8 is hydrogen. In some embodiments, R^8 is halogen.

As described generally above, R^9 is hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂. In some embodiments, R^9 is hydrogen. In some embodiments, R^9 is halogen.

In some embodiments, each R_a is independently -OR_f, NHR_c, NR_cR_f, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted carbocyclyl, optionally substituted aryl, optionally substituted heteroaryl, or optionally substituted heterocyclyl. In some embodiments, each R_a is independently -OR_f, NHR_c, NR_cR_f, alkyl, cycloalkyl, aryl, or optionally substituted 3- to 7-membered heterocyclyl having 1-2 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

In some embodiments, each R_b is independently -OR_f, -NHR_c, -NR_cR_f, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted carbocyclyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -SR_c, -S(O)R_c, -S(O)₂R_c, C(O)R_c, -C(O)OR_c, or -C(O)NR_cR_f. In some embodiments, R_b is optionally substituted 3- to 7-membered heterocyclyl having 1-2 heteroatoms independently selected from nitrogen, oxygen, and sulfur. In some embodiments, R_b is C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, C₄-C₇ cycloalkylalkyl, mono or polycyclic aryl, or mono or polycyclic heteroaryl. In certain embodiments, R_b is optionally substituted from 1 to 3 times with substituents independently selected from the group consisting of halogen, -H, -OR_c, -C(O)R_b, -C(O)OR_c, -C(O)NR_cR_f, -NHR_d, -NR_dR_f, -SR_c, -S(O)R, -S(O)₂R, -NH₂, -CN, -NO₂, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, C₄-C₇ cycloalkylalkyl, monocyclic aryl, and polycyclic aryl.

In some embodiments, each R_c is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted carbocyclyl, optionally substituted heterocyclyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R_c is C_{1-6} alkyl (e.g., methyl, ethyl, isopropyl, cyclopropyl, t-butyl), cycloalkyl, 3- to 7-membered heterocyclyl, aryl, or heteroaryl. In some
 5 embodiments, R_c is optionally substituted phenyl. In some embodiments, R_c is an optionally substituted 5- to 10-membered heteroaryl having 1-4 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

- 10 In some embodiments, each R_d is independently optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, $-COR_c$, $S(O)R_c$, $S(O)_2R_c$, optionally substituted alkylaryl, or optionally substituted aryl. In some embodiments, R_d is alkyl, $-COR_c$, $S(O)R_c$, $S(O)_2R_c$, phenyl, benzyl, substituted benzyl, or aryl.
- 15 In some embodiments, each R_e is independently hydrogen, halo, $-OH$, $-OR_c$, $-NH_2$, $-NHR_c$, $-NHSO_2R_c$, $-NR_cSO_2R_c$, $-NR_cR_f$, $-CN$, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, $-COR_c$, or $-NHCOR_c$. In some embodiments, R_e is $-H$, $-F$, $-OH$, $-OR_c$, $-NH_2$, $-NHR_c$, $-NHSO_2R_c$, $-NR_cR_f$, $-CN$, aryl, heterocyclyl,
 20 trifluoroacetyl, acetyl, or NH-acetyl.

In some embodiments, each R_f is independently $-H$ or C_{1-6} alkyl.


- In some embodiments, each R_g is independently hydrogen, halo, $-OH$, $-OR_c$, $-NH_2$, $-NHR_c$, $-NHSO_2R_c$, $-NR_cSO_2R_c$, $-NR_cR_f$, $-CN$, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, $-COR_c$, or $-NHCOR_c$. In some embodiments, R_g is $-H$, $-F$, $-OH$, $-OR_c$, $-NH_2$, $-NHR_c$, $-NHSO_2R_c$, $-NR_cR_f$, $-CN$, aryl, heterocyclyl,
 25 trifluoroacetyl, acetyl, or NH-acetyl.

30

In some embodiments, R_d and R_e are each fluoro. In some embodiments, R_d and R_e are each methyl. In some embodiments, R_d and R_e are taken together to form $=O$. In some embodiments, R_d and R_e are taken together to form a 4-, 5-, or 6-membered spiro-fused heterocycle or carbocycle.


- In some embodiments, R_d and R_g are each fluoro. In some embodiments, R_d and R_g are each methyl. In some embodiments, R_d and R_g are taken together to form =O. In some embodiments, R_d and R_g are taken together to form a 4-, 5-, or 6-membered spiro-fused heterocycle or carbocycle.

In some embodiments, R^1 is -CN or -SONH₂. In some embodiments, R^1 is one of the following:




- wherein R_a , R_b , R_c , and R_d are described herein, and n is 0, 1, or 2.

In some embodiments, R^1 is -C(O)R, wherein R is optionally substituted heterocyclyl. In




some embodiments, R^1 is -C(O)R, wherein R is R_d or R_e . In some embodiments, R^1 is -C(O)R, wherein R is one of the following:




15

In some embodiments, R_a is one of the following:




In some embodiments, R_a is one of the following:




5


In some embodiments, R_b is one of the following:




10 In some embodiments, R_c is one of the following:



In some embodiments, R^1 and R^2 are taken together to form:




In some embodiments, R^2 is $-H$, $-CN$, $-NH_2$, $-SO_2NH_2$,



- 5 In some embodiments, R^2 , R^3 , R^4 , R^5 , R^6 , R^8 , and R^9 are each hydrogen.

In some embodiments, a provided compound is of Formula (Ia):



or a pharmaceutically acceptable form thereof, wherein

- 10 X_1 , X_2 , and X_3 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

wherein at least one of X_1 , X_2 , X_3 , X_4 , and X_5 is N;

X_7 is N or CR^{10} ;

X_8 is N or CR^{11} ;

- 15 ----- is a single or double bond, as valency allows;

R^1 , R^2 , R^3 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , and R^{11} are independently hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, $-CN$, $-OR$, $-SR$, $-N(R)_2$, $-NO_2$, $-C(O)R$, $-CO_2R$, $-C(O)N(R)_2$, $-C(O)NROR$, $-S(O)R$, $-SO_2R$, $-SO_2N(R)_2$, $-OC(O)R$, $-NRC(O)R$, $-NRSO_2R$, or $-NRC(O)N(R)_2$;

each R is independently hydrogen, optionally substituted C_{1-6} aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or


optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken together to form an optionally substituted heterocycle;

each R⁴ is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic;

and

- 5 R¹ and R², R² and R³, R³ and R⁴, R¹ and R¹¹, R¹⁰ and R¹¹, R⁵ and R⁶, R⁶ and R⁷, R⁷ and R⁸, or R⁸ and R⁹ may be optionally taken together to form an optionally substituted 5-6 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

In some embodiments, a provided compound is of Formula (Ia-i):



or a pharmaceutically acceptable form thereof, wherein

X₁, X₂, and X₃ are independently N or C;

X₄ and X₅ are independently N or CR⁴;

wherein at least one of X₁, X₂, X₃, X₄, and X₅ is N;

- 15 X₇ is N or CR¹⁰;

X₈ is N or CR¹¹;

----- is a single or double bond, as valency allows;


- 20 R¹, R², R³, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, and R¹¹ are independently hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂;

- 25 each R is independently hydrogen, optionally substituted C₁₋₆ aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken together to form an optionally substituted heterocycle;

each R^4 is independently hydrogen, halogen, or optionally substituted C_{1-6} aliphatic;
and

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^1 and R^{11} , R^{10} and R^{11} , R^5 and R^6 , R^6 and R^7 , R^7 and R^8 , or R^8 and R^9 may be optionally taken together to form an optionally substituted 5-6 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

In some embodiments, a provided compound is of Formula (Ib):



or a pharmaceutically acceptable form thereof, wherein

10 X_1 , X_2 , and X_3 are independently N or C;

X_4 , X_5 , X_6 are independently N or CR^4 ;

wherein at least one of X_1 , X_2 , X_3 , X_4 , X_5 , and X_6 is N;

X_7 is N or CH;

X_8 is N or CH;


15 $---$ is a single or double bond, as valency allows;

R^1 is $-C(O)R$, wherein R is optionally substituted heterocyclyl;

R^2 , R^3 , R^6 , and R^7 are independently $-H$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CF_3$, $-CHF_2$, $-C(O)NH_2$, $-OH$, $-OC_{1-4}alkyl$, or $-OCF_3$; or R^6 and R^7 are taken together to form a 5-6 membered fused heterocyclyl or heteroaryl moiety; and

20 each R^4 is independently hydrogen or C_{1-6} alkyl.

In some embodiments, a provided compound is of Formula (Ib-i):



or a pharmaceutically acceptable form thereof, wherein

X_1 , X_2 , and X_3 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

5 wherein at least one of X_1 , X_2 , X_3 , X_4 , and X_5 is N;

X_7 is N or CH;

X_8 is N or CH;

----- is a single or double bond, as valency allows;


R^1 is $-C(O)R$, wherein R is optionally substituted heterocyclyl;

10 R^2 , R^3 , R^6 , and R^7 are independently $-H$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CF_3$, $-CHF_2$, $-C(O)NH_2$, $-OH$, $-OC_{1-4}alkyl$, or $-OCF_3$; or R^6 and R^7 are taken together to form a 5-6 membered fused heterocyclyl or heteroaryl moiety; and

each R^4 is independently hydrogen or $C_{1-6}alkyl$.

15 In some embodiments, R^4 is hydrogen.

In some embodiments, a provided compound is of Formula **(Ib-ii)**:



or a pharmaceutically acceptable form thereof, wherein

X_1 , X_2 , and X_3 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

wherein at least one of X_1 , X_2 , X_3 , X_4 , and X_5 is N;

X_7 is N or CH;

5 X_8 is N or CH;

----- is a single or double bond, as valency allows;


R^1 is $-C(O)R$, wherein R is optionally substituted heterocyclyl;

R^2 , R^3 , R^6 , and R^7 are independently $-H$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CF_3$, $-CHF_2$, $-C(O)NH_2$, $-OH$, $-OC_{1-4}alkyl$, or $-OCF_3$; or R^6 and R^7 are taken together to form a 5-6 membered fused

10 heterocyclyl or heteroaryl moiety; and

each R^4 is independently hydrogen or C_{1-6} alkyl.

In some embodiments, a provided compound is of Formula (Ic):



15 or a pharmaceutically acceptable form thereof, wherein

X_1 , and X_2 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

X_7 is N or CR^{10} ;

X_8 is N or CR^{11} ;

20 R^1 , R^2 , R^3 , R^5 , R^6 , R^7 , R^8 , R^{10} , and R^{11} are independently hydrogen, halogen, optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, $-CN$, $-OR$, $-SR$, $-N(R)_2$, $-NO_2$, $-C(O)R$, $-CO_2R$, $-C(O)N(R)_2$, $-C(O)NROR$, $-S(O)R$, $-SO_2R$, $-SO_2N(R)_2$, $-OC(O)R$, $-NRC(O)R$, $-NRSO_2R$, or $-NRC(O)N(R)_2$;

25 each R is independently hydrogen, optionally substituted C_{1-6} aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or


optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken together to form an optionally substituted heterocycle;

each R⁴ is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic;

and

- 5 R¹ and R², R² and R³, R³ and R⁴, R¹ and R¹¹, R¹⁰ and R¹¹, R⁵ and R⁶, R⁶ and R⁷, R⁷ and R⁸, or R⁸ and R⁹ may be optionally taken together to form an optionally substituted 5-6 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

In some embodiments, a provided compound is of Formula (Ic-i):



or a pharmaceutically acceptable form thereof, wherein

X₁ and X₂ are independently N or C;

X₄ and X₅ are independently N or CR⁴;

X₇ is N or CH;


- 15 X₈ is N or CH;

R¹ is -C(O)R, wherein R is optionally substituted heterocyclyl;

R², R³, R⁶, and R⁷ are independently -H, -F, -Cl, -CN, -CH₃, -CF₃, -CHF₂, -C(O)NH₂, -OH, -OC₁₋₄alkyl, or -OCF₃; or R⁶ and R⁷ are taken together to form a 5-6 membered fused heterocyclyl or heteroaryl moiety; and

- 20 each R⁴ is independently hydrogen or C₁₋₆ alkyl.

In some embodiments, a provided compound is of Formula (Id):



or a pharmaceutically acceptable form thereof, wherein

X_1 , and X_2 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

5 X_7 is N or CR^{10} ;

X_8 is N or CR^{11} ;

one ----- is a single bond, and the other ----- is a double bond;

$R^1, R^2, R^3, R^5, R^6, R^7, R^8, R^{10}$, and R^{11} are independently hydrogen, halogen,

optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl,

10 optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂,

-NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂;

each R is independently hydrogen, optionally substituted C₁₋₆ aliphatic, optionally

substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or

15 optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken

together to form an optionally substituted heterocycle;


each R^4 is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic;

and

R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^1 and R^{11} , R^{10} and R^{11} , R^5 and R^6 , R^6 and R^7 , R^7

20 and R^8 , or R^8 and R^9 may be optionally taken together to form an optionally substituted 5-6 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

In some embodiments, a provided compound is of Formula (Id-i):



or a pharmaceutically acceptable form thereof, wherein

X_1 , and X_2 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

5 X_7 is N or CH;

X_8 is N or CH;

one --- is a single bond, and the other --- is a double bond;

R^1 is $-C(O)R$, wherein R is an optionally substituted heterocyclyl;







10 R^2 , R^3 , R^6 , and R^7 are independently $-H$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CF_3$, $-CHF_2$, $-C(O)NH_2$, $-OH$, $-OC_{1-4}alkyl$, or $-OCF_3$; or R^6 and R^7 are taken together to form a 5-6 membered fused heterocyclyl or heteroaryl; and

each R^4 is independently hydrogen or $C_{1-6} alkyl$.

In some embodiments, for any of the formulae described herein, X_7 and X_8 are each CH.







15

In some embodiments, a provided compound is of Formula (II), (III), (IV), (V), (VI), or (VII):


**Formula II****Formula III****Formula IV****Formula V****Formula VI****Formula VII**

In some embodiments, for Formula (II), (III), (IV), (V), (VI), or (VII), R¹ is -C(O)R as described herein.


- 5 In some embodiments, a provided compound is of Formula (IIa), (IIIa), (IVa), (Va), (VIa), or (VIIa):

**Formula IIa****Formula IIIa****Formula IVa****Formula Va****Formula VIa****Formula VIIa**


In some embodiments, a provided compound is of Formula (IIa-i), (IIIa-i), (IVa-i), (Va-i), (VIa-i), or (VIIa-i):




Formula IIa-i




Formula IIIa-i




Formula IVa-i



Formula Va-i




Formula VIa-i




Formula VIIa-i


- 5 In some embodiments, a provided compound is of Formula (IIb), (IIIb), (IVb), (Vb), (VIb), or (VIIb):




Formula IIb




Formula IIIb




Formula IVb



Formula Vb




Formula VIb




Formula VIIb


In some embodiments, a provided compound is of Formula (IIb-i), (IIIb-i), (IVb-i), (Vb-i), (VIb-i), or (VIIb-i):




Formula IIb-i




Formula IIIb-i




Formula IVb-i



Formula Vb-i




Formula VIb-i




Formula VIIb-i


5 In some embodiments, a provided compound is of Formula (IIc), (IIIc), (IVc), (Vc), (VIc), or (VIIc):




Formula IIc




Formula IIIc




Formula IVc



Formula Vc




Formula VIc



Formula VIIc

In some embodiments, R is optionally substituted heterocyclyl.

10 In some embodiments, a provided compound is one of the following:



The present invention also provides pharmaceutical compositions comprising an effective amount of a compound described herein, or a pharmaceutically acceptable form (e.g., pharmaceutically acceptable salt or prodrug) thereof, and, optionally, a pharmaceutically acceptable excipient.

Pharmaceutically acceptable excipients include any and all solvents, diluents or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. General considerations in formulation and/or manufacture of pharmaceutical compositions agents can be found, for example, in *Remington's Pharmaceutical Sciences*, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980), and *Remington: The Science and Practice of Pharmacy*, 21st Edition (Lippincott Williams & Wilkins, 2005).

Pharmaceutical compositions described herein can be prepared by any method known in the art of pharmacology. In general, such preparatory methods include the steps of bringing the compound of the present invention (the "active ingredient") into association with a carrier

and/or one or more other accessory ingredients, and then, if necessary and/or desirable, shaping and/or packaging the product into a desired single- or multi-dose unit.

Pharmaceutical compositions can be prepared, packaged, and/or sold in bulk, as a single unit
5 dose, and/or as a plurality of single unit doses. As used herein, a "unit dose" is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

10

Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition of the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of
15 example, the composition may comprise between 0.1% and 100% (w/w) active ingredient.

Pharmaceutically acceptable excipients used in the manufacture of provided pharmaceutical compositions include inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering
20 agents, lubricating agents, and/or oils. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and perfuming agents may also be present in the composition.

Exemplary diluents include calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate
25 lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, etc., and combinations thereof.

Exemplary granulating and/or dispersing agents include potato starch, corn starch, tapioca
30 starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch

(starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, etc., and combinations thereof.

- 5 Exemplary surface active agents and/or emulsifiers include natural emulsifiers (*e.g.* acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (*e.g.* bentonite (aluminum silicate) and Veegum (magnesium aluminum silicate)), long chain amino acid derivatives, high molecular weight alcohols (*e.g.* stearyl alcohol, cetyl alcohol,
- 10 oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (*e.g.* carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (*e.g.* carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose),
- 15 sorbitan fatty acid esters (*e.g.* polyoxyethylene sorbitan monolaurate (Tween 20), polyoxyethylene sorbitan (Tween 60), polyoxyethylene sorbitan monooleate (Tween 80), sorbitan monopalmitate (Span 40), sorbitan monostearate (Span 60), sorbitan tristearate (Span 65), glyceryl monooleate, sorbitan monooleate (Span 80)), polyoxyethylene esters (*e.g.* polyoxyethylene monostearate (Myrj 45), polyoxyethylene hydrogenated castor oil,
- 20 polyethoxylated castor oil, polyoxymethylene stearate, and Solutol), sucrose fatty acid esters, polyethylene glycol fatty acid esters (*e.g.* CREMOPHOR), polyoxyethylene ethers, (*e.g.* polyoxyethylene lauryl ether (Brij 30)), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic F-68, Poloxamer P188, cetrimonium bromide,
- 25 cetylpyridinium chloride, benzalkonium chloride, docusate sodium, etc. and/or combinations thereof.

- Exemplary binding agents include starch (*e.g.* cornstarch and starch paste), gelatin, sugars (*e.g.* sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol, etc.), natural
- 30 and synthetic gums (*e.g.* acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum), and larch arabogalactan), alginates, polyethylene oxide, polyethylene

glycol, inorganic calcium salts, silicic acid, polymethacrylates, waxes, water, alcohol, etc., and/or combinations thereof.

Exemplary preservatives include antioxidants, chelating agents, antimicrobial preservatives, 5 antifungal preservatives, alcohol preservatives, acidic preservatives, and other preservatives.

Exemplary antioxidants include alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and 10 sodium sulfite.

Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA) and salts and hydrates thereof (e.g., sodium edetate, disodium edetate, trisodium edetate, calcium disodium edetate, dipotassium edetate, and the like), citric acid and salts and hydrates thereof (e.g., 15 citric acid monohydrate), fumaric acid and salts and hydrates thereof, malic acid and salts and hydrates thereof, phosphoric acid and salts and hydrates thereof, and tartaric acid and salts and hydrates thereof. Exemplary antimicrobial preservatives include benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, 20 hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and thimerosal.

Exemplary antifungal preservatives include butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, 25 sodium benzoate, sodium propionate, and sorbic acid.

Exemplary alcohol preservatives include ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and phenylethyl alcohol.

30 Exemplary acidic preservatives include vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and phytic acid.

Other preservatives include tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium

lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, Glydant Plus, Phenonip, methylparaben, Germall 115, Germaben II, Neolone, Kathon, and Euxyl. In certain embodiments, the preservative is an anti-oxidant. In other embodiments, the preservative is a
 5 chelating agent.

Exemplary buffering agents include citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid,
 10 calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate,
 15 sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, etc., and combinations thereof.

Exemplary lubricating agents include magnesium stearate, calcium stearate, stearic acid,
 20 silica, talc, malt, glyceryl behenate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, etc., and combinations thereof.

Exemplary natural oils include almond, apricot kernel, avocado, babassu, bergamot, black
 25 current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach
 30 kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary synthetic oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric

triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and combinations thereof.

Liquid dosage forms for oral and parenteral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredients, the liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (*e.g.*, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents. In certain embodiments for parenteral administration, the conjugates of the invention are mixed with solubilizing agents such as CREMOPHOR, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and combinations thereof.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions can be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing the conjugates of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may comprise buffering agents.

Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only,

or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type can be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The active ingredients can be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active ingredient can be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may comprise buffering agents. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

Dosage forms for topical and/or transdermal administration of a compound of this invention may include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants and/or patches. Generally, the active ingredient is admixed under sterile conditions with a pharmaceutically acceptable carrier and/or any needed preservatives and/or buffers as can be required. Additionally, the present invention contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of an active ingredient to the body. Such dosage forms can be prepared, for example, by dissolving and/or dispensing the active ingredient in the proper medium. Alternatively or additionally, the rate can be controlled by either providing a rate controlling membrane and/or by dispersing the active ingredient in a polymer matrix and/or gel.

Suitable devices for use in delivering intradermal pharmaceutical compositions described herein include short needle devices such as those described in U.S. Patents 4,886,499; 5,190,521; 5,328,483; 5,527,288; 4,270,537; 5,015,235; 5,141,496; and 5,417,662. Intradermal compositions can be administered by devices which limit the effective

penetration length of a needle into the skin, such as those described in PCT publication WO 99/34850 and functional equivalents thereof. Jet injection devices which deliver liquids to the dermis via a liquid jet injector and/or via a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Jet injection devices are described, for example, in U.S. Patents 5,480,381; 5,599,302; 5,334,144; 5,993,412; 5,649,912; 5,569,189; 5,704,911; 5,383,851; 5,893,397; 5,466,220; 5,339,163; 5,312,335; 5,503,627; 5,064,413; 5,520,639; 4,596,556; 4,790,824; 4,941,880; 4,940,460; and PCT publications WO 97/37705 and WO 97/13537. Ballistic powder/particle delivery devices which use compressed gas to accelerate vaccine in powder form through the outer layers of the skin to the dermis are suitable. Alternatively or additionally, conventional syringes can be used in the classical mantoux method of intradermal administration.

Formulations suitable for topical administration include, but are not limited to, liquid and/or semi liquid preparations such as liniments, lotions, oil in water and/or water in oil emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions. Topically-administrable formulations may, for example, comprise from about 1% to about 10% (w/w) active ingredient, although the concentration of the active ingredient can be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration may further comprise one or more of the additional ingredients described herein.

A pharmaceutical composition of the invention can be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration via the buccal cavity. Such a formulation may comprise dry particles which comprise the active ingredient and which have a diameter in the range from about 0.5 to about 7 nanometers or from about 1 to about 6 nanometers. Such compositions are conveniently in the form of dry powders for administration using a device comprising a dry powder reservoir to which a stream of propellant can be directed to disperse the powder and/or using a self propelling solvent/powder dispensing container such as a device comprising the active ingredient dissolved and/or suspended in a low-boiling propellant in a sealed container. Such powders comprise particles wherein at least 98% of the particles by weight have a diameter greater than 0.5 nanometers and at least 95% of the particles by number have a diameter less than 7 nanometers. Alternatively, at least 95% of the particles by weight have a diameter greater than 1 nanometer and at least 90% of the particles by number have a diameter less than 6 nanometers. Dry powder compositions may

include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50 to 99.9% (w/w) of the composition, and the active ingredient may constitute 0.1 to 20% (w/w) of the composition. The propellant may further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

10

Pharmaceutical compositions of the invention formulated for pulmonary delivery may provide the active ingredient in the form of droplets of a solution and/or suspension. Such formulations can be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising the active ingredient, and may conveniently be administered using any nebulization and/or atomization device. Such formulations may further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. The droplets provided by this route of administration may have an average diameter in the range from about 0.1 to about 200 nanometers.

20

Formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutical composition of the invention. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 to 500 micrometers. Such a formulation is administered by rapid inhalation through the nasal passage from a container of the powder held close to the nares.

25

Formulations for nasal administration may, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of the active ingredient, and may comprise one or more of the additional ingredients described herein. A pharmaceutical composition of the invention can be prepared, packaged, and/or sold in a formulation for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may contain, for example, 0.1 to 20% (w/w) active ingredient, the

30

balance comprising an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising the active ingredient. Such powdered, aerosolized, and/or aerosolized
5 formulations, when dispersed, may have an average particle and/or droplet size in the range from about 0.1 to about 200 nanometers, and may further comprise one or more of the additional ingredients described herein.

A pharmaceutical composition of the invention can be prepared, packaged, and/or sold in a
10 formulation for ophthalmic administration. Such formulations may, for example, be in the form of eye drops including, for example, a 0.1/1.0% (w/w) solution and/or suspension of the active ingredient in an aqueous or oily liquid carrier. Such drops may further comprise buffering agents, salts, and/or one or more other of the additional ingredients described herein. Other ophthalmically-administrable formulations which are useful include those
15 which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this invention.

Although the descriptions of pharmaceutical compositions provided herein are principally
20 directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist
25 can design and/or perform such modification with ordinary experimentation.

Still further encompassed by the invention are kits (e.g., pharmaceutical packs). The kits provided may comprise an inventive pharmaceutical composition or compound and a container (e.g., a vial, ampule, bottle, syringe, and/or dispenser package, or other suitable
30 container). In some embodiments, provided kits may optionally further include a second container comprising a pharmaceutical excipient for dilution or suspension of an inventive pharmaceutical composition or compound. In some embodiments, the inventive pharmaceutical composition or compound provided in the container and the second container are combined to form one unit dosage form.

Optionally, a single container may comprise one or more compartments for containing an inventive pharmaceutical composition or compound, and/or a pharmaceutically acceptable excipient for suspension or dilution. In some embodiments, a single container can be appropriate for modification such that the container may receive a physical modification so as to allow combination of compartments and/or components of individual compartments. For example, a foil or plastic bag may comprise two or more compartments separated by a perforated seal which can be broken so as to allow combination of contents of two individual compartments once the signal to break the seal is generated. A kit may thus comprise such multi-compartment containers providing an inventive pharmaceutical composition or compound and one or more pharmaceutically acceptable excipients.

Optionally, instructions for use are additionally provided in such kits of the invention. Such instructions may provide, generally, for example, instructions for dosage and administration. In other embodiments, instructions may further provide additional detail relating to specialized instructions for particular containers and/or systems for administration. Still further, instructions may provide specialized instructions for use in conjunction and/or in combination with an additional therapeutic agent.

Compounds provided herein are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject or organism will depend upon a variety of factors including the disease, disorder, or condition being treated and the severity of the disorder; the activity of the specific active ingredient employed; the specific composition employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific active ingredient employed; the duration of the treatment; drugs used in combination or coincidental with the specific active ingredient employed; and like factors well known in the medical arts.

The compounds and compositions provided herein can be administered by any route, including enteral (e.g., oral), parenteral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal,

intravaginal, intraperitoneal, topical (as by powders, ointments, creams, and/or drops), mucosal, nasal, bucal, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; and/or as an oral spray, nasal spray, and/or aerosol. Specifically contemplated routes are oral administration, intravenous administration (e.g., systemic intravenous injection), regional administration via blood and/or lymph supply, and/or direct administration to an affected site. In general the most appropriate route of administration will depend upon a variety of factors including the nature of the agent (e.g., its stability in the environment of the gastrointestinal tract), and/or the condition of the subject (e.g., whether the subject is able to tolerate oral administration).

10

The exact amount of a compound required to achieve an effective amount will vary from subject to subject, depending, for example, on species, age, and general condition of a subject, severity of the side effects or disorder, identity of the particular compound(s), mode of administration, and the like. The desired dosage can be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage can be delivered using multiple administrations (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

15

20 In certain embodiments, an effective amount of a compound for administration one or more times a day to a 70 kg adult human may comprise about 0.0001 mg to about 3000 mg, about 0.0001 mg to about 2000 mg, about 0.0001 mg to about 1000 mg, about 0.001 mg to about 1000 mg, about 0.01 mg to about 1000 mg, about 0.1 mg to about 1000 mg, about 1 mg to about 1000 mg, about 1 mg to about 100 mg, about 10 mg to about 1000 mg, or about 100 mg to about 1000 mg, of a compound per unit dosage form.

25

In certain embodiments, the compounds of the invention may be at dosage levels sufficient to deliver from about 0.001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, preferably from about 0.1 mg/kg to about 40 mg/kg, preferably from about 0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, and more preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

30

It will be appreciated that dose ranges as described herein provide guidance for the administration of provided pharmaceutical compositions to an adult. The amount to be administered to, for example, a child or an adolescent can be determined by a medical practitioner or person skilled in the art and can be lower or the same as that administered to an adult.

It will be also appreciated that a compound or composition, as described herein, can be administered in combination with one or more additional therapeutically active agents. The compounds or compositions can be administered in combination with additional therapeutically active agents that improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. It will also be appreciated that the therapy employed may achieve a desired effect for the same disorder, and/or it may achieve different effects.

The compound or composition can be administered concurrently with, prior to, or subsequent to, one or more additional therapeutically active agents. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. It will further be appreciated that the additional therapeutically active agent utilized in this combination can be administered together in a single composition or administered separately in different compositions. The particular combination to employ in a regimen will take into account compatibility of the inventive compound with the additional therapeutically active agent and/or the desired therapeutic effect to be achieved. In general, it is expected that additional therapeutically active agents utilized in combination be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

Exemplary additional therapeutically active agents include, but are not limited to, small organic molecules such as drug compounds (*e.g.*, compounds approved by the U.S. Food and Drug Administration as provided in the Code of Federal Regulations (CFR)), peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoproteins, mucoproteins, lipoproteins, synthetic polypeptides or proteins, small molecules linked to proteins, glycoproteins, steroids, nucleic acids, DNAs, RNAs, nucleotides, nucleosides, oligonucleotides, antisense oligonucleotides, lipids, hormones, vitamins, and cells. In some embodiments, an additional therapeutically active agent is a kinase inhibitor.

Also encompassed by the present disclosure are kits (e.g., pharmaceutical packs). The kits provided may comprise a provided pharmaceutical composition or compound and a container (e.g., a vial, ampule, bottle, syringe, and/or dispenser package, or other suitable container).

- 5 In some embodiments, provided kits may optionally further include a second container comprising a pharmaceutical excipient for dilution or suspension of a provided pharmaceutical composition or compound. In some embodiments, a provided pharmaceutical composition or compound provided in the container and the second container are combined to form one unit dosage form. In some embodiments, a provided kits further includes
- 10 instructions for use.

- Compounds and compositions described herein are generally useful for the inhibition of one or more kinases. In certain embodiments, compounds and compositions described herein are generally useful for the inhibition of MNK1 and/or MNK2. In some embodiments, methods
- 15 of treating kinase-related disorder in a subject are provided which comprise administering an effective amount of a compound described herein (e.g., a compound of Formula (I)), or a pharmaceutically acceptable form thereof, to a subject in need of treatment. In some embodiments, methods of treating MNK1- and/or MNK2-related disorder in a subject are provided which comprise administering an effective amount of a compound described herein
- 20 (e.g., a compound of Formula (I)), or a pharmaceutically acceptable form thereof, to a subject in need of treatment. In certain embodiments, the effective amount is a therapeutically effective amount. In certain embodiments, the effective amount is a prophylactically effective amount. In certain embodiments, the subject is suffering from a MNK1-related disorder. In certain embodiments, the subject is susceptible to a MNK1-
- 25 mediated disorder. In certain embodiments, the subject is suffering from a MNK2-related disorder. In certain embodiments, the subject is susceptible to a MNK2-mediated disorder.

- As used herein, the term "kinase-related disorder" (e.g., "MNK1- and/or MNK2-related disorder") means any disease, disorder, or other pathological condition in which a kinase
- 30 (e.g., MNK1 and/or MNK2) is known to play a role. In some embodiments, the present disclosure relates to treating or lessening the severity of one or more diseases in which MNK1 and/or MNK2 is known to play a role.

In certain embodiments, the kinase-related condition (*e.g.*, MNK1- and/or MNK2-related condition) is selected from the group consisting of proliferative diseases, neurodegenerative diseases, autoimmune diseases, and inflammatory diseases.

- 5 In certain embodiments, a provided compound is useful for treating a proliferative disease, *e.g.*, cancer. Exemplary cancers include, but are not limited to, acoustic neuroma; adenocarcinoma; adrenal gland cancer; anal cancer; angiosarcoma (*e.g.*, lymphangiosarcoma, lymphangioendotheliosarcoma, hemangiosarcoma); appendix cancer; benign monoclonal gammopathy; biliary cancer (*e.g.*, cholangiocarcinoma); bladder cancer; breast cancer (*e.g.*,
10 adenocarcinoma of the breast, papillary carcinoma of the breast, mammary cancer, medullary carcinoma of the breast); brain cancer (*e.g.*, meningioma, glioblastomas, glioma (*e.g.*, astrocytoma, oligodendroglioma), medulloblastoma); bronchus cancer; carcinoid tumor; cervical cancer (*e.g.*, cervical adenocarcinoma); choriocarcinoma; chordoma; craniopharyngioma; colorectal cancer (*e.g.*, colon cancer, rectal cancer, colorectal
15 adenocarcinoma); connective tissue cancer; epithelial carcinoma; ependymoma; endotheliosarcoma (*e.g.*, Kaposi's sarcoma, multiple idiopathic hemorrhagic sarcoma); endometrial cancer (*e.g.*, uterine cancer, uterine sarcoma); esophageal cancer (*e.g.*, adenocarcinoma of the esophagus, Barrett's adenocarcinoma); Ewing sarcoma; eye cancer (*e.g.*, intraocular melanoma, retinoblastoma); familial hypereosinophilia; gall bladder cancer;
20 gastric cancer (*e.g.*, stomach adenocarcinoma); gastrointestinal stromal tumor (GIST); germ cell cancer; head and neck cancer (*e.g.*, head and neck squamous cell carcinoma, oral cancer (*e.g.*, oral squamous cell carcinoma), throat cancer (*e.g.*, laryngeal cancer, pharyngeal cancer, nasopharyngeal cancer, oropharyngeal cancer)); hematopoietic cancers (*e.g.*, leukemia such as acute lymphocytic leukemia (ALL) (*e.g.*, B-cell ALL, T-cell ALL), acute myelocytic
25 leukemia (AML) (*e.g.*, B-cell AML, T-cell AML), chronic myelocytic leukemia (CML) (*e.g.*, B-cell CML, T-cell CML), and chronic lymphocytic leukemia (CLL) (*e.g.*, B-cell CLL, T-cell CLL)); lymphoma such as Hodgkin lymphoma (HL) (*e.g.*, B-cell HL, T-cell HL) and non-Hodgkin lymphoma (NHL) (*e.g.*, B-cell NHL such as diffuse large cell lymphoma (DLCL) (*e.g.*, diffuse large B-cell lymphoma), follicular lymphoma, chronic lymphocytic
30 leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), marginal zone B-cell lymphomas (*e.g.*, mucosa-associated lymphoid tissue (MALT) lymphomas, nodal marginal zone B-cell lymphoma, splenic marginal zone B-cell lymphoma), primary mediastinal B-cell lymphoma, Burkitt lymphoma, lymphoplasmacytic lymphoma (*i.e.*, Waldenström's macroglobulinemia), hairy cell leukemia (HCL), immunoblastic large cell

- lymphoma, precursor B-lymphoblastic lymphoma and primary central nervous system (CNS) lymphoma; and T-cell NHL such as precursor T-lymphoblastic lymphoma/leukemia, peripheral T-cell lymphoma (PTCL) (*e.g.*, cutaneous T-cell lymphoma (CTCL) (*e.g.*, mycosis fungoides, Sezary syndrome), angioimmunoblastic T-cell lymphoma, extranodal natural killer T-cell lymphoma, enteropathy type T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, and anaplastic large cell lymphoma); a mixture of one or more leukemia/lymphoma as described above; and multiple myeloma (MM)), heavy chain disease (*e.g.*, alpha chain disease, gamma chain disease, mu chain disease); hemangioblastoma; hypopharynx cancer; inflammatory myofibroblastic tumors; immunocytic amyloidosis;
- 10 kidney cancer (*e.g.*, nephroblastoma *a.k.a.* Wilms' tumor, renal cell carcinoma); liver cancer (*e.g.*, hepatocellular cancer (HCC), malignant hepatoma); lung cancer (*e.g.*, bronchogenic carcinoma, small cell lung cancer (SCLC), non-small cell lung cancer (NSCLC), adenocarcinoma of the lung); leiomyosarcoma (LMS); mastocytosis (*e.g.*, systemic mastocytosis); muscle cancer; myelodysplastic syndrome (MDS); mesothelioma;
- 15 myeloproliferative disorder (MPD) *e.g.*, polycythemia Vera (PV), essential thrombocytosis (ET), agnogenic myeloid metaplasia (AMM) *a.k.a.* myelofibrosis (MF), chronic idiopathic myelofibrosis, chronic myelocytic leukemia (CML), chronic neutrophilic leukemia (CNL), hypereosinophilic syndrome (HES)); neuroblastoma; neurofibroma (*e.g.*, neurofibromatosis (NF) type 1 or type 2, schwannomatosis); neuroendocrine cancer (*e.g.*, gastroenteropancreatic
- 20 neuroendocrine tumor (GEP-NET), carcinoid tumor); osteosarcoma (*e.g.*, bone cancer); ovarian cancer (*e.g.*, cystadenocarcinoma, ovarian embryonal carcinoma, ovarian adenocarcinoma); papillary adenocarcinoma; pancreatic cancer (*e.g.*, pancreatic adenocarcinoma, intraductal papillary mucinous neoplasm (IPMN), Islet cell tumors); penile cancer (*e.g.*, Paget's disease of the penis and scrotum); pinealoma; primitive neuroectodermal
- 25 tumor (PNT); plasma cell neoplasia; paraneoplastic syndromes; intraepithelial neoplasms; prostate cancer (*e.g.*, prostate adenocarcinoma); rectal cancer; rhabdomyosarcoma; salivary gland cancer; skin cancer [*e.g.*, squamous cell carcinoma (SCC), keratoacanthoma (KA), melanoma, basal cell carcinoma (BCC)]; small bowel cancer (*e.g.*, appendix cancer); soft tissue sarcoma (*e.g.*, malignant fibrous histiocytoma (MFH), liposarcoma, malignant
- 30 peripheral nerve sheath tumor (MPNST), chondrosarcoma, fibrosarcoma, myxosarcoma); sebaceous gland carcinoma; small intestine cancer; sweat gland carcinoma; synovioma; testicular cancer (*e.g.*, seminoma, testicular embryonal carcinoma); thyroid cancer (*e.g.*, papillary carcinoma of the thyroid, papillary thyroid carcinoma (PTC), medullary thyroid

cancer); urethral cancer; vaginal cancer; and vulvar cancer (*e.g.*, Paget's disease of the vulva).

In certain embodiments, a provided compound is useful for treating a neurodegenerative disease. Exemplary neurodegenerative diseases include, but are not limited to, Alzheimer's

5 disease, Huntington's disease, progressive supranuclear palsy, corticobasal degeneration, frontotemporal lobar degeneration, Pick's disease, Parkinson's disease, Lewy body disease, and amyotrophic lateral sclerosis (ALS).

- 10 In certain embodiments, a provided compound is useful for treating an autoimmune disease. Exemplary autoimmune diseases include, but are not limited to, rheumatoid arthritis, spondyloarthropathies, gouty arthritis, degenerative joint diseases such as osteoarthritis, systemic lupus erythematosus, Sjogren's syndrome, ankylosing spondylitis, undifferentiated spondylitis, Behcet's disease, haemolytic autoimmune anaemias, multiple sclerosis,
- 15 amyotrophic lateral sclerosis, amylosis, acute painful shoulder, psoriatic arthritis, juvenile arthritis, asthma, atherosclerosis, osteoporosis, bronchitis, tendonitis, bursitis, skin condition (*e.g.*, psoriasis, eczema, burns, dermatitis, pruritus (itch)), enuresis, eosinophilic disease, gastrointestinal disorder (*e.g.*, selected from peptic ulcers, regional enteritis, diverticulitis, gastrointestinal bleeding, eosinophilic gastrointestinal disorders (*e.g.*, eosinophilic
- 20 esophagitis, eosinophilic gastritis, eosinophilic gastroenteritis, eosinophilic colitis), gastritis, diarrhea, gastroesophageal reflux disease (GERD), inflammatory bowel disease (IBD) (*e.g.*, Crohn's disease, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behcet's syndrome, indeterminate colitis) and inflammatory bowel syndrome (IBS)), Still's disease, myasthenia gravis, Hashimoto's thyroiditis, Ord's
- 25 thyroiditis, Grave's disease, acute disseminated encephalomyelitis, Addison's disease, opsoclonus-myoclonus syndrome, antiphospholipid antibody syndrome, autoimmune hepatitis, celiac disease, Goodpasture's syndrome, optic neuritis, scleroderma, primary biliary cirrhosis, Reiter's syndrome, warm autoimmune hemolytic anemia, alopecia universalis, chronic fatigue, dysautonomia, neuromyotonia, vulvodynia and disorders ameliorated by a
- 30 gastroprokinetic agent (*e.g.*, ileus, postoperative ileus and ileus during sepsis; gastroesophageal reflux disease (GERD); eosinophilic esophagitis, gastroparesis such as diabetic gastroparesis; food intolerances and food allergies and other functional bowel disorders, such as non-ulcerative dyspepsia (NUD) and non-cardiac chest pain (NCCP, including costo-chondritis)).

- In certain embodiments, a provided compound is useful for treating an inflammatory disease. The term "inflammatory disease" refers to those conditions that are characterized by signs of pain (dolor, from the generation of noxious substances and the stimulation of nerves), heat (calor, from vasodilatation), redness (rubor, from vasodilatation and increased blood flow), swelling (tumor, from excessive inflow or restricted outflow of fluid), and/or loss of function (functio laesa, which can be partial or complete, temporary or permanent. Inflammation takes on many forms and includes, but is not limited to, acute, adhesive, atrophic, catarrhal, chronic, cirrhotic, diffuse, disseminated, exudative, fibrinous, fibrosing, focal, granulomatous, hyperplastic, hypertrophic, interstitial, metastatic, necrotic, obliterative, parenchymatous, plastic, productive, proliferous, pseudomembranous, purulent, sclerosing, seroplastic, serous, simple, specific, subacute, suppurative, toxic, traumatic, and/or ulcerative inflammation.
- Exemplary inflammatory diseases include, but are not limited to inflammation associated with acne, asthma, arteritis (*e.g.*, polyarteritis, temporal arteritis, periarteritis nodosa, Takayasu's arteritis), arthritis (*e.g.*, crystalline arthritis, osteoarthritis, psoriatic arthritis, gouty arthritis, reactive arthritis, rheumatoid arthritis, and Reiter's arthritis), ankylosing spondylitis, amylosis, amyotrophic lateral sclerosis, autoimmune diseases, allergies or allergic reactions, atherosclerosis, bronchitis, bursitis, chronic prostatitis, conjunctivitis, Chagas disease, chronic obstructive pulmonary disease, dermatomyositis, dry eye syndrome, diverticulitis, diabetes (*e.g.*, type I diabetes mellitus, type 2 diabetes mellitus), a skin condition (*e.g.*, psoriasis, eczema, burns, dermatitis, pruritus (itch)), endometriosis, Guillain-Barre syndrome, infection, ischaemic heart disease, Kawasaki disease, glomerulonephritis, gingivitis, hypersensitivity, headaches (*e.g.*, migraine headaches, tension headaches), ileus (*e.g.*, postoperative ileus and ileus during sepsis), idiopathic thrombocytopenic purpura, interstitial cystitis (painful bladder syndrome), gastrointestinal disorder (*e.g.*, selected from peptic ulcers, regional enteritis, diverticulitis, gastrointestinal bleeding, eosinophilic gastrointestinal disorders (*e.g.*, eosinophilic esophagitis, eosinophilic gastritis, eosinophilic gastroenteritis, eosinophilic colitis), gastritis, diarrhea, gastroesophageal reflux disease (GERD), inflammatory bowel disease (IBD) (*e.g.*, Crohn's disease, ulcerative colitis, collagenous colitis, lymphocytic colitis, ischaemic colitis, diversion colitis, Behcet's syndrome, indeterminate colitis), inflammatory bowel syndrome (IBS), lupus, multiple sclerosis, morphea, myasthenia gravis, myocardial ischemia, nephrotic syndrome,

- pemphigus vulgaris, pernicious anaemia, peptic ulcers, polymyositis, primary biliary cirrhosis, neuroinflammation associated with brain disorders (*e.g.*, Parkinson's disease, Huntington's disease, and Alzheimer's disease), prostatitis, chronic inflammation associated with cranial radiation injury, pelvic inflammatory disease, reperfusion injury, regional enteritis, rheumatic fever, systemic lupus erythematosus, scleroderma, sclerodoma, sarcoidosis, spondyloarthropathies, Sjogren's syndrome, thyroiditis, transplantation rejection, tendonitis, trauma or injury (*e.g.*, frostbite, chemical irritants, toxins, scarring, burns, physical injury), vasculitis, vitiligo and Wegener's granulomatosis. In certain embodiments, the inflammatory disorder is selected from arthritis (*e.g.*, rheumatoid arthritis), inflammatory bowel disease, inflammatory bowel syndrome, asthma, psoriasis, endometriosis, interstitial cystitis, prostatitis, appendicitis, Blau syndrome, blepharitis, bronchiolitis, cervicitis, cholangitis, cholecystitis, chronic recurrent multifocal osteomyelitis (CRMO), cryopyrin associated periodic syndrome (CAPS), dacryoadenitis, dermatomyositis, dry eye syndrome, encephalitis, endocarditis, endometritis, enterocolitis, epicondylitis, epididymitis, familial cold-induced autoinflammatory syndrome, familial Mediterranean fever (FMF), fasciitis, fibrositis, hepatitis, hidradenitis suppurativa, laryngitis, mastitis, meningitis, mevalonate kinase deficiency (MKD), Muckle-Wells syndrome, myelitis myocarditis, myositis, nephritis, oophoritis, orchitis, osteitis, inflammatory osteolysis, otitis, pancreatitis, parotitis, pericarditis, peritonitis, pharyngitis, pleuritis, phlebitis, pneumonitis, pneumonia, proctitis, pulmonary fibrosis, pyelonephritis, pyoderma gangrenosum and acne syndrome (PAPA), pyogenic sterile arthritis, rhinitis, salpingitis, sinusitis, stomatitis, synovitis, TNF receptor associated periodic syndrome (TRAPS), tonsillitis, undifferentiated arthropathy, uveitis, vaginitis and vulvitis. In certain embodiments, the inflammatory condition is an acute inflammatory condition (*e.g.*, inflammation resulting from infection). In certain
- embodiments, the inflammatory condition is a chronic inflammatory condition (*e.g.*, conditions resulting from asthma, arthritis and inflammatory bowel disease). The compounds may also be useful in treating inflammation associated with trauma and non-inflammatory myalgia. The compounds may also be useful in treating inflammation associated with cancer.
- In certain embodiments, a provided compound is useful for treating or lessening the severity of arthropathies and osteopathological diseases including, but not limited to, rheumatoid arthritis, osteoarthritis, gout, polyarthritis, and psoriatic arthritis.

In certain embodiments, a provided compound is useful for treating or lessening the severity of acute and chronic inflammatory diseases including, but not limited to, ulcerative colitis, inflammatory bowel disease, Crohn's disease, dry eye syndrome, allergic rhinitis, allergic dermatitis, cystic fibrosis, chronic obstructive bronchitis, and asthma.

5

In certain embodiments, a provided compound is useful for treating or lessening the severity of hyperproliferative diseases including, but not limited to, psoriasis or smooth muscle cell proliferation including vascular proliferative disorders, atherosclerosis, and restenosis.

- 10 In certain embodiments, a provided compound is useful for treating or lessening the severity of endometriosis, uterine fibroids, endometrial hyperplasia, and benign prostate hyperplasia.

In some embodiments, a provided compound is useful for treating or lessening the severity of one or more diseases and conditions, wherein the disease or condition is selected from

- 15 immune-related conditions or diseases, which include, but are not limited to graft versus host disease, transplantation, transfusion, anaphylaxis, allergies (*e.g.*, allergies to plant pollens, latex, drugs, foods, insect poisons, animal hair, animal dander, dust mites, or cockroach calyx), type I hypersensitivity, allergic conjunctivitis, allergic rhinitis, and atopic dermatitis.

- 20 In some embodiments, a provided compound is useful for treating tumorogenesis.

In some embodiments, a provided compound is useful for treating a metabolic disorder (*e.g.*, obesity, diabetes).

- 25 In some embodiments, the present disclosure provides a method of inhibiting MNK1 comprising contacting MNK1 with an effective amount of a compound described herein (*e.g.*, a compound of Formula (I)), or a pharmaceutically acceptable form thereof. In some embodiments, the present disclosure provides a method of inhibiting MNK2 comprising contacting MNK2 with an effective amount of a compound described herein (*e.g.*, a
30 compound of Formula (I)), or a pharmaceutically acceptable form thereof. The MNK1 or MNK2 may be purified or crude, and may be present in a cell, tissue, or subject. Thus, such methods encompass both inhibition of *in vitro* and *in vivo* MNK1 or MNK2 activity. In certain embodiments, the method is an *in vitro* method, *e.g.*, such as an assay method. It will be understood by one of ordinary skill in the art that inhibition of MNK1 or MNK2 does not

necessarily require that all of the MNK1 or MNK2 be occupied by an inhibitor at once. Exemplary levels of inhibition of MNK1 or MNK2 include at least 10% inhibition, about 10% to about 25% inhibition, about 25% to about 50% inhibition, about 50% to about 75% inhibition, at least 50% inhibition, at least 75% inhibition, about 80% inhibition, about 90% inhibition, and greater than 90% inhibition.

In some embodiments, provided is a method of inhibiting kinase activity in a subject in need thereof (*e.g.*, a subject diagnosed as having a kinase-related disorder) comprising administering to the subject an effective amount of a compound described herein (*e.g.*, a compound of Formula (I)), or a pharmaceutically acceptable form thereof, or a pharmaceutical composition thereof. In some embodiments, provided is a method of inhibiting MNK1 and/or MNK2 activity in a subject in need thereof (*e.g.*, a subject diagnosed as having a MNK1- and/or MNK2-related disorder) comprising administering to the subject an effective amount of a compound described herein (*e.g.*, a compound of Formula (I)), or a pharmaceutically acceptable form thereof, or a pharmaceutical composition thereof.

EXAMPLES

Compounds of formula (I) were synthesized and their ability to inhibit MNK1/2 kinase was evaluated both in enzymatic and in cell-based assays. In parallel, their ability to inhibit other kinases was also assessed. Enzymatic assays have shown that compounds of formula (I) are inhibitors of MNK1 and MNK2 with IC_{50} values in the range of 1000 nM to lower than 10 nM. eIF4e phosphorylation inhibition in Hela cell line was found to have IC_{50} s in average 10 time higher than the enzymatic IC_{50} s. These cell-based IC_{50} s vary from as low as 40 nM to 3 μ M.

To further evaluate the therapeutic potential of these bicyclic molecules, their ability to inhibit growth of several tumotype cell lines was assessed. Below are the technical descriptions of these assays.

Enzymatic and cell-based eIF4E phosphorylation assays and Thermal Shift assay

Biological Methods

The compounds described in this invention bind to and inhibit the kinases MNK1 and MNK2. They were analyzed using both *in vitro* and *in vivo* assays that are known in the art.

Thermal Shift Assays

- 5 Thermal shift assays can be used to monitor the binding of small molecules to proteins by measuring the change in melting temperature that results from an increase in thermal stability of the protein on binding a compound. The thermally induced unfolding of the protein is monitored in the presence of a fluorescent dye, Sypro Orange using a real time PCR system. The innate fluorescence of Sypro Orange is quenched in an aqueous environment; however in
- 10 a hydrophobic non-polar environment such as the exposed hydrophobic residues of an unfolding protein, the compound fluoresces profusely. This enables the unfolding of proteins to be monitored by measuring an increase in fluorescence. Melting temperature is defined as the temperature at which half the protein is in an unfolded state.
- 15 The increase in melting temperature of MNK1 and MNK2 on binding small molecular weight compounds was determined using recombinant full length human enzymes with N-terminal GST tags (glutathione-S-transferase). The proteins were expressed in *E.coli* and purified on glutathione sepharose using the Profinia Protein Purification System (BioRad, Hercules, CA) and the GST tag was removed using PreScission protease. Both recombinant enzymes were
- 20 diluted in buffer A (10mM Tris/HCl pH7.5, 50mM NaCl, 1mM DTT) to a concentration of 62.5μM. The compounds to be analyzed were dissolved in 100% DMSO to a final concentration of 100μM. The reaction mixture consisted of 5μM enzyme (MNK1 or MNK2), 4μM compound, 17μl buffer A and 5μl of 5X Sypro Orange in a final volume of 25μl. The thermal analysis was performed on a BioRad CFX96 RTPCR instrument from 20°C to 90°C
- 25 in 0.5°C increments with a dwell time of 20s.

The data is analyzed by plotting the fluorescent intensity as a function of temperature. Readings that occur before or after the sigmoidal region of the melt curve are discarded. The fluorescent values are normalized for each compound, 100 for the maximum reading (UL)

30 and 0 for the minimum reading (LL). The curve is fitted to the Boltzmann equation using non-linear regression (GraphPad Prism, GraphPad Software Inc.). The inflection point of the transition curve, T_m (V50) is calculated using the Boltzmann equation:

$$Y = [LL + (UL - LL)] / [1 + \exp(T_m - t/\alpha)]$$

Where t = temperature, LL and UL are the minimum and maximum intensities respectively and α is the slope of the curve within T_m . The thermal shift is defined as the difference between the melting temperatures in the presence and absence of compound.

5 In Vitro MNK Kinase Assay

MNK1 and MNK2 inhibitor activity was determined using recombinant kinase domains expressed in *E.coli*. MNK1 and MNK2 were expressed as GST fusion proteins and the GST tag was removed using PreScission protease. After concentration to 10-15 mg/ml the proteins were flash frozen in liquid nitrogen and stored at -80°C. MNK1 and MNK2 were activated using recombinant ERK2 which was activated using a constitutively active mutant of MEK1, both ERK2 and MEK1 were expressed in *E.coli* as N-terminally his tagged proteins. Recombinant ERK2 was activated by incubating 11.3 μ M of the kinase with 1 μ M MEK1 and 100 μ M ATP. This reaction mixture was then used immediately for the activation of the MNKs. The activation of the MNK1 was performed by incubating 5.0 μ M of MNK1 with 0.3 μ M of activated ERK2 and 500 μ M ATP at 30°C for 6 hours. The activation of MNK2 was performed by incubating 50 μ M of MNK2 with 3.0 μ M of activated ERK2 and 500 μ M ATP at 30°C for 2 hours. The activated MNKs were stored at -20°C until required for assay.

Kinase assays were performed on the Caliper Life Sciences (Mountain View, CA) Microfluidics LabChip® Platform. Enzyme activity is analyzed by 'sipping' reactions from a microtitre plate into LabChip. The data signature is generated by the shift in mobility of non-phosphorylated peptide substrates and phosphorylated products by electrophoresis in the chip and detected by LED induced fluorescence. The magnitude of the fluorescent signal reveals the extent of the reaction. The data is analyzed by calculating the relative heights of the substrate and product peaks and the product / (product+substrate) peak ratio is reported.

The following buffers were used to assay kinase activity:

Reconstitution buffer: 10mM HEPES/NaOH pH7.5, 0.003% Brij® L23, 0.004% TWEEN® 20.

30 Substrate buffer: 245mM HEPES/NaOH pH7.5, 0.003% Brij® L23, 0.004% TWEEN® 20, 26mM MgCl₂.

Termination buffer: 100mM HEPES/NaOH pH7.3, 0.022% Brij® L23, 5.6% DMSO, 0.16% CR3, 11.2mM EDTA pH8.0.

Separation buffer: 100mM HEPES/NaOH pH7.3, 0.02% Brij® L23, 5% DMSO, 0.1% CR3, 1mM EDTA pH8.0.

Peptide substrate (JH3): 5-FAM-TATKSGSTTKNRFVV-CONH₂.

- 5 The MNK1 assay was performed by adding 65 nM of activated MNK1 and 1 µl of test compound to a microtitre plate in a volume of 15 µl of reconstitution buffer. The plate was incubated at 22 °C for 15 minutes before the addition of 3.9 µM of JH3 and 3.12 mM ATP in 10µl of substrate buffer and a further incubation period of 60 minutes at 28 °C. The reaction was stopped by the addition of 45 µl of termination buffer. The final concentration of MNK1, 10 JH3 peptide, ATP and compound in a 26 µl assay volume was 40 nM, 1.5 µM, 1.2 mM, and 1X respectively.

- The MNK2 assay was performed by adding 32.5 nM of activated MNK1 and 1 µl of test compound to a microtitre plate in a volume of 15 µl of reconstitution buffer. The plate was 15 incubated at 22 °C for 15 minutes before the addition of 3.9 µM of JH3 and 650 µM ATP in 10 µl of substrate buffer and a further incubation period of 60 minutes at 28 °C. The reaction was stopped by the addition of 45 µl of termination buffer. The final concentration of MNK2, JH3 peptide, ATP and compound in a 26 µl assay volume was 20 nM, 1.5 µM, 250 µM, and 1X respectively.

- 20 Inhibition constants (IC₅₀) were determined by plotting kinase activity versus log compound concentration and fitting with a non-linear regression algorithm using GraphPad Prism (GraphPad Software Inc.).

MNK Cell-Based Assay

- 25 It has been reported that Ser209 of eIF4E is solely phosphorylated by the MNK enzymes. The ability of compounds to inhibit this process in Hela cells was investigated using the AlphaScreen *SureFire*® assay platform from Perkin Elmer (Waltham, MA). eIF4E phosphorylated on Ser209 is recognized by two antibodies, the first which is fused to a streptavidin coated donor bead binds to an epitope away from Ser209, the second which is 30 fused to a protein A conjugated acceptor bead binds to phosphorylated Ser209. The phosphorylation of eIF4E on Ser209 brings the two antibodies into close proximity and when excited by a laser a singlet oxygen is released by the donor bead which excites the acceptor bead resulting in the emission of light. This enables the monitoring of eIF4E Ser209 phosphorylation and its inhibition in a cellular context.

Hela cells were seeded into microtitre plates (30,000 cells per well) in 100 μ l of culture medium and incubated at 37 °C for 24 hours. The media was then removed by aspiration and the cells resuspended in 50 μ l of serum free medium containing the test compound and
5 incubated at 37 °C for 2 hours. The culture medium was again removed by aspiration and the cells resuspended in lysis buffer (provided in Perkin Elmer SureFire® Assay Kit). After agitation at 350 rpm for 20 minutes at 22 °C, 4 μ l was transferred to a 384 well OptiPlate™ (Perkin Elmer, Waltham, MA). To each well was added 5 μ l of acceptor mix; the plate was sealed and agitated gently at 22 °C for 2 hours. Then, in subdued light, 2 μ l of donor mix was
10 added to each well, the plate was sealed, wrapped in aluminum foil and agitated gently at 22 °C for 2 hours. Emission was measured using the EnVision® plate reader (Perkin Elmer, Waltham, MA).

Inhibition constants (IC_{50}) were determined by plotting AlphaScreen signal versus log compound concentration and fitting with a non-linear regression algorithm using GraphPad
15 Prism (GraphPad Software Inc.).

Cell Cytotoxicity Assays

Methodology:

Cancer cell lines, MV-4-11 (leukemia), P116.c139 (acute T cell leukemia) and D1.1 (acute T
20 cell leukemia), were purchased from ATCC and cultured according to supplier's recommendations. K562 cells (myelogenous leukemia) over-expressing eIF4E, were also used for the cytotoxicity assay. For cells treated for 48 hours, 5000 cells were seeded in 70 μ l of growth medium in black, flat-bottom 96-well plate. For cells treated for six days, 1000 cells were seeded in 70 μ l of growth medium in black, flat-bottom 96-well plate. The
25 compounds, Cercosporamide, Example 176 (ETC-7114), Example 175 (ETC-7117), Example 133 (ETC-7165) and Example 62 (ETC-7168), were treated with doses ranging from 0.003 μ M to 50 μ M. 50 μ l of the diluted compounds was added to the cells and incubated at 37 °C in 5% CO₂. After 48 hours or six days treatment, cell viability was determined by CellTiter-Glo Luminescent Cell Viability Assay (Promega, Madison, WI). 120 μ l of the reagent was
30 added to the cells and luminescence was measured using Tecan Safire Reader. Data was analysed with Graphpad Prism software and the figures represented indicate the half maximal inhibitory concentration (IC_{50}). Error bars denote standard deviation (SD).

Results:**Table 1: IC₅₀ (μM) after 48 hours of treatment**

	K562	MV-4-11	D1.1	P116.c139
Cercosporamide	> 50	> 50	20.2	> 50
ETC-7114 Example 176	14.2	1.0	1.0	0.4
ETC-7117 Example 175	18.2	0.9	12.3	12.5
ETC-7165 Example 133	23.0	2.8	15.8	10.9
ETC-7168 Example 62	7.5	0.8	1.6	0.8

Table 2: IC₅₀ (μM) after 6 days of treatment

	K562	MV-4-11	D1.1	P116.c139
Cercosporamide	4.3	4.0	17.3	4.9
ETC-7114 Example 176	3.4	0.4	0.8	0.1
ETC-7117 Example 175	21.2	0.3	10.1	5.7
ETC-7165 Example 133	16.0	1.5	11.2	4.0
ETC-7168 Example 62	3.1	0.2	1.7	0.3

Discussion:

5 Generally, the IC₅₀ is lower in cells treated for 6 days than in the cells treated for 48 hours. Amongst the three test cell lines, the MV-4-11 cell line is the most sensitive to all test compounds.

Amongst the four compounds, Example 176 (ETC-7114) and Example 62 (ETC-7168) are the most cytotoxic. The IC₅₀ of example 176 (ETC-7114) and Example 62 (ETC-7168) is ≤ 1 μM and ≤ 2 μM, respectively, in the three test cell lines, regardless of the treatment duration. The IC₅₀ of example 176 (ETC-7114) and Example 62 (ETC-7168) in K562 cells over-expressing eIF4E were reduced by more than half when treatment duration was extended to 6 days. These data suggest that the cytotoxic effects of Example 176 (ETC-7114) and Example 62 (ETC-7168) are probably non-cell line specific.

15

The treatment duration affects the cytotoxic effects of compounds Example 175 (ETC-7117) and Example 133 (ETC-7165) in the test cell lines. The extended treatment duration from 48 hours to 6 days could decrease the IC₅₀ by more than half-fold in P116.c139 cells. However, the IC₅₀ of example 175 (ETC-7117) and example 133 (ETC-7165) in K562 cells over-expressing eIF4E and in D1.1 cells was not influenced by the treatment duration. These data

20

suggests that the cytotoxic effects of Example 175 (ETC-7117) and Example 133 (ETC-7165) are probably more cell line specific.

Immunofluorescence and Serial Replating Assays

Methods

- 5 Cord Blood (CB) samples were purchased from the Singapore Cord Blood bank. CML samples were the Singapore General Hospital after signed informed consent under local IRB-approved procedures. MNCs were obtained using Ficoll separation, and CD34⁺ cells selected by immunomagnetic beads (Miltenyi Biotech, Germany).

10 Cell Culture and Generation of Cell Lines

K562 cell line was obtained from the ATCC, and grown in RPMI supplemented with 10% FCS, L-glutamine, and penicillin/streptomycin.

Serial Replating Assay

- 15 CD34-enriched CB and BC cells were thawed and allowed to recover overnight in serum-free StemPro media (Invitrogen, Carlsbad, CA), supplemented with human growth factors and 1x nutrient supplement (Invitrogen). Cells were then subjected to drug treatment for 48 hr, harvested, washed, and seeded in methycellulose (H4434, STEMCELL Technologies, Canada). Colonies were enumerated after 2 weeks, individually picked, and replated in fresh
20 methylcellulose in a 96-well format, and counted at 2 weeks. Three rounds of serial replating (representing > 8 weeks in culture) were performed.

Immunofluorescence Analysis

- Cells (1×10^5) were cytospun onto glass slides, fixed with 4% paraformaldehyde, and stained
25 with mouse monoclonal antibodies against activated β -catenin (clone 8E7, Millipore, UK), or rabbit monoclonal antibodies against phospho-eIF4E S209 (EP2151Y, Abcam, UK). Slides were then stained with either PE-conjugated anti-mouse or FITC-conjugated anti-rabbit antibodies. Images were obtained with the use of a fluorescence microscope (Olympus IX71SIF3) at 40x magnification.

30

Western Blotting

Exponentially growing cells were plated at 2×10^5 cells/ml, and whole cell lysate processed for western blot analysis using antibodies recognizing eIF4E, phospho-eIF4E (Cell signaling Technology),

Discussion

As shown in figure 1A, treatment of K562 cells with increasing concentration of drugs causes a dose dependent decrease in eIF4E phosphorylation. Similar to what was observed in BC-CML cell lines, we showed that treatment of BC-CML primary cells with increasing dose of drugs also resulted in a dose dependent decrease in eIF4E phosphorylation as well as active beta-catenin (Figure 1B-D). The functional consequence of decrease beta-catenin as well as eIF4E phosphorylation on the self renewal capacity of BC leukemia stem cells (LSCs) was assessed next. We performed a serial replating assay as previously described (Jamieson et al., 2004). Importantly, the serial replating assay has been found to correlate well with beta-catenin-driven self-renewal in BC-GMPs, and also the *in vivo* serial-transplanting ability of a variety of fusion-gene driven LSCs (Huntly et al., 2004). Using normal CB CD34⁺ cells, we found that control treated cells were capable of serial replating up to three times (equivalent to > 8 weeks *in vitro*). We also found that treatment with Example 6 (ETC-0445) did not significantly alter the serial replating efficiency compared to DMSO (Figure 2A). In contrast, for CD34⁺ BC cells, treatment with increasing dose of Example 6 (ETC-0445) and Example 1 (ETC-5336) retards the serial replating efficiency as compared to DMSO (Figure 2B and 2C). Treatment of CD34⁺ BC cells with increasing dose of Example 2 (ETC-6740) not only retards serial replating efficiency, at 10.0 μ M and 20.0 μ M, it potently impaired the ability of BC cells to serial replate (Figure 2D).

References

- Huntly, B. J., Shigematsu, H., Deguchi, K., Lee, B. H., Mizuno, S., Duclos, N., Rowan, R., Amaral, S., Curley, D., Williams, I. R., *et al.* (2004). MOZ-TIF2, but not BCR-ABL, confers properties of leukemic stem cells to committed murine hematopoietic progenitors. *Cancer Cell* 6, 587-596.
- Jamieson, C. H., Ailles, L. E., Dylla, S. J., Muijtjens, M., Jones, C., Zehnder, J. L., Gotlib, J., Li, K., Manz, M. G., Keating, A., *et al.* (2004). Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. *N Engl J Med* 351, 657-667.

Treatment of K562 cells with of Example 176 (ETC-7114) or Example 62 (ETC-7168) cause a decrease in eIF4E phosphorylation (Figure 1A); in addition to inhibition of eIF4E phosphorylation, a decrease in BCR-ABL activity is observed, as demonstrated by the

decrease in phosphorylation of Crkl, an *in vivo* substrate of BCR-ABL. The decrease in eIF4E phosphorylation is also accompanied by the decrease in nuclear beta-catenin accumulation as demonstrated using immunofluorescence analysis (figure 1B and C).

5

Synthetic Examples

The following examples serve to illustrate the invention without limiting the scope thereof.

10

Abbreviations

CAN: acetonitrile

AcOEt: ethyl acetate

AcOH: acetic acid

AUC: area under the curve

15

Brine: saturated solution of NaCl in water

cat.: catalyst

d: day(s)

DCM: dichloromethane

DIEA: diisopropyl-ethyl-amine

20

DMF: *N,N*-dimethylformamide

DMSO: dimethylsulfoxide

DMSO-*d*₆: per-deuterated dimethylsulfoxide

dppf: 1,1'- Bis(diphenylphosphino) ferrocene

EDCI: 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide)

25

Ether: diethylether

EtOH: ethanol

h: hour(s)

HATU: 2-(1H-7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate

Methanaminium

30

HBTU: O-Benzotriazole-*N,N,N',N'*-tetramethyl-uronium-hexafluoro-phosphate

hOBt: *N*-Hydroxybenzotriazole

HPLC: high pressure liquid chromatography

L: litre(s)


LC-MS: Liquid chromatography-mass spectrometry


- Me: methyl
MeOH: methanol
min: minute(s)
m.p.: melting point
5 MS: mass spectrometry
NBS: *N*-Bromosuccinimide
NEt₃: triethylamine
NIS: *N*-iodosuccinimide
NMM: *N*-methylmorpholine
10 NMR: Nuclear Magnetic Resonance
Pd(dppf)Cl₂: [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II)
rt: room temperature
THF: tetrahydrofuran
TFA: trifluoroacetic acid
15 TLC: thin layer chromatography

Compounds according to the invention, for example, starting materials, intermediates or products, are prepared as described herein or by the application or adaptation of known methods, by which is meant methods used heretofore or described in the literature.

- 20 Compounds useful according to the invention may be prepared by the application or adaptation of known methods, by which is meant methods used heretofore or described in the literature, for example those described by Larock, R.C., *Comprehensive Organic Transformations*, VCH publishers, (1989), which is hereby incorporated by reference in its entirety.


25

**Formula 2**General procedure A




- To a solution of 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoic acid, *N*-methylmorpholine and HATU in DMF, was added the desired amine (RR'NH) and the resulting mixture was stirred for 10 h under nitrogen atmosphere. The reaction mixture was poured onto ice water and the precipitate was isolated by filtration and dried to afford the desired amide used as such or purified as indicated.

General procedure B




- To a solution of the halogenated bicyclic derivative in 1,4-dioxane was added the boronic acid or the boronate ester derivative, K₃PO₄ and water and Pd(PPh₃)₄. The reaction mixture was heated at 90 °C for 12 h under argon atmosphere then, was filtered through celite and concentrated under reduced pressure. The residue was purified by flash column chromatography to afford the desired product.

15



Intermediate 1: 4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)benzoic acid




Step 1: To a solution of 6-bromoimidazo [1, 2-a] pyrazine (30 g, 151 mmol) in toluene (400 mL) were sequentially added Cs_2CO_3 (123 g, 378 mmol), 4-(ethoxycarbonyl)phenylboronic acid (35.2 g, 181 mmol) and $\text{Pd}(\text{dppf})_2\text{Cl}_2$ (3.70 g, 4.50 mmol). The reaction mixture was stirred at 90 °C for 12 h under argon atmosphere then, was filtered through celite. The filtrate was concentrated and the residue was purified the by flash column chromatography (silica gel, eluent n-hexane/EtOAc 70:30) to afford ethyl 4-(imidazo [1, 2-a]pyrazin-6-yl)benzoate (16 g, 39%) as yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.35 (s, 1H), 9.17 (s, 1H), 8.15 (d, $J = 8.4$ Hz, 2H), 8.03 (d, $J = 8.4$ Hz, 2H), 7.88 (s, 1H), 7.79 (s, 1H), 4.32 (q, $J = 6.8$ Hz, 2H), 1.44 (t, $J = 7.2$ Hz 3H); MS (ESI) m/z 268 [$\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_2 + \text{H}$] $^+$.

Step 2: To a solution of ethyl 4-(imidazo [1, 2-a]pyrazin-6-yl)benzoate (16.0 g, 59.8 mmol) in $\text{CCl}_4/\text{CH}_3\text{OH}$ (200/ 50 mL) was added *N*-bromosuccinamide (12.7 g, 71.8 mmol). The reaction mixture was stirred at room temperature for 30 min then, was filtered through celite. The filtrate was concentrated, resuspended in water and extracted with CH_2Cl_2 . The organic phase was concentrated to afford ethyl 4-(3-bromoimidazo [1, 2-a]pyrazin-6-yl)benzoate (18.0 g, 87%) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.17 (s, 1H), 8.78 (s, 1H), 8.28 (d, $J = 8.4$ Hz, 2H), 8.09 (d, $J = 8.0$ Hz, 2H), 7.94 (s, 1H), 4.34 (q, $J = 7.2$ Hz, 2H), 1.35 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 345 [$\text{C}_{15}\text{H}_{12}\text{BrN}_3\text{O}_2$] $^+$.

Step 3: To a solution of ethyl 4-(3-bromoimidazo [1, 2-a]pyrazin-6-yl)benzoate (27.0 g, 77.9 mmol) in $\text{THF}/\text{CH}_3\text{OH}$ (300/100 mL) were added LiOH (26.0 g, 623 mmol) and water (100 mL). The reaction mixture was stirred for 5 h at room temperature then, was concentrated, diluted with water (200 mL) and acidified with aqueous HCl solution till pH 3. The precipitate was isolated by filtration and dried to afford 4-(3-bromoimidazo [1, 2-a]pyrazin-6-yl)benzoic acid (14.0 g, 56%) as white solid. ^1H NMR (400 MHz, CDCl_3) δ : 9.17 (s, 1H), 8.68 (s, 1H), 8.25 (d, $J = 8.4$ Hz 2H), 8.09 (d, $J = 8.0$ Hz, 2H), 7.99 (s, 1H); MS (ESI) m/z 316 [$\text{C}_{13}\text{H}_8\text{BrN}_3\text{O}_2$] $^+$.


Intermediates 2 and 3: (4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



Step 1: To a solution of 6-bromoimidazo[1,2-a]pyrimidine (2.4 g, 12.1 mmol) in DMF, were sequentially added (4-methylpiperazin-1-yl)(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (80% of 4.1 g, 12.1 mmol), a solution of NaHCO₃ (3.1 g, 36.3 mmol) in water (30 mL), water (3 mL) and (A-Phos)₂PdCl₂ (400 mg, 0.60 mmol). The reaction mixture was heated at 90 °C for 4 h, then was diluted with water (100 mL) and extracted with EtOAc (3×100 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (silica gel, eluent CHCl₃/MeOH 95:5 to 85:15) to afford of (4-(imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (1.5 g, 65%). ¹H NMR (400 MHz, CDCl₃) δ 8.60 (d, *J* = 2.0 Hz, 1 H), 8.10 (d, *J* = 4.0, 1 H), 7.80 (d, *J* = 8.0 Hz, 1 H), 7.58 (m, 4 H) 7.39 (d, *J* = 8.0, 1 H), 3.84 (br s, 2 H), 3.50 (br s, 2 H), 2.40 (br s, 4 H), 2.35 (s, 3 H); MS (ESI) *m/z* 322 [M+1]

Step 2: To a solution of ((4-(imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (1.4 g, 4.36 mmol) in ACN (15 mL) and DCM (5 mL) was added *N*-Bromosuccinimide (0.93 g, 5.23 mmol) at 0 °C. The reaction mixture was stirred for 90 min while warming to room temperature. The reaction mixture was diluted with DCM and washed with water. The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (silica gel, eluent CHCl₃/MeOH 95:5 to 90:10) to afford ((4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (1.0 g, 83%) as a brown solid. ¹H NMR (400 MHz, CDCl₃) δ 8.82 (d, *J* = 4.0 Hz, 1 H), 8.53 (d, *J* = 4.0 Hz, 1 H), 7.87 (s, 1 H), 7.50 (m, 4 H), 3.84 (bs, 2 H), 3.50 (br s, 2 H), 2.39 (bs, 4 H), 2.34 (s, 3 H); MS (ESI) *m/z* 400.

Intermediates 4 and 5: 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoic acid and (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



Step 1: To a solution of ethyl 4-(imidazo [1, 2-a]pyrazin-6-yl)benzoate (8.00 g, 29.9 mmol) in DMF (200 mL), was added *N*-Iodosuccinimide (8.10 g, 36.0 mmol) and the mixture was heated at 60 °C for 2 h and poured onto ice water. The precipitate was isolated by filtration and dried to afford ethyl 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoate (11 g, 94%) as yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.13 (s, 1H), 8.80 (s, 1H), 8.29 (d, *J* = 8.4 Hz, 2H), 8.07 (d, *J* = 8.4 Hz, 2H), 8.00 (s, 1H), 4.37–4.31 (m, 2H), 1.34 (t, *J* = 7.2 Hz, 3H); MS (ESI) *m/z* 393 [C₁₅H₁₂IN₃O₂]⁺.

Step 2: A solution of ethyl 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoate (11 g, 30.5 mmol) and LiOH·H₂O (5.00 g, 121 mmol) in THF/CH₃OH/H₂O (200/50/50 mL) was stirred for 12 h. The reaction mixture was concentrated, diluted with water (20 mL) and acidified with an aqueous solution of HCl till pH 2. The precipitate was isolated by filtration and dried to afford 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoic acid (8.00 g, 72%) as off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.12 (s, 1H), 8.78 (s, 1H), 8.26 (d, *J* = 8.4 Hz, 2H), 8.06 (d, *J* = 8.4 Hz, 2H), 8.05 (s, 1H); MS (ESI) *m/z* 366 [C₁₃H₈IN₃O₂ + H]⁺.

Step 3: *N*-methylmorpholine (6.90 mL, 54.64 mmol), HATU (15.58 g, 40.98 mmol) and *N*-methylpiperazine (4.15 mL, 40.98 mmol) were added sequentially to a solution of 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoic acid (10.0 g, 27.32 mmol) in DMF (50 mL). The resulting mixture was stirred at room temperature for 3 h under nitrogen atmosphere and, was diluted with water (100 mL). The precipitate was isolated by filtration and dried to afford (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino)methanone (10.5 g, 86%) as off-white solid. ¹H NMR (400 MHz, CDCl₃) δ: 9.15 (d, *J* = 1.2 Hz, 1H), 8.44 (d, *J* = 1.2 Hz, 1H), 8.05 (d, *J* = 8.0 Hz, 2H), 7.83 (s, 1H), 7.57 (d, *J* = 8.0 Hz, 2H), 3.85 (bs, 2H), 3.51 (bs, 2H), 2.53 (bs, 2H), 2.41 (bs, 2H), 2.35 (s, 3H); MS (ESI) *m/z* 447 [C₁₈H₁₈IN₅O]⁺.

Intermediate 6: Synthesis of (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl) (morpholino) methanone

N-methyl-morpholine (3.0 mL, 7.5 mmol), HATU (7.5 g, 27 mmol) and morpholinè (1.26 g, 14.85 mmol) were added sequentially to a solution of 4-(3-iodoimidazo[1,2-a]pyrazin-6-

yl)benzoic acid (5.0 g, 3.25 mmol) in DMF (10 mL) and the resulting mixture was stirred at room temperature for 3 h under nitrogen atmosphere. The reaction mixture was diluted with water (50 mL) and the precipitate that has formed was filtered off and dried to afford (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino)methanone (2.0 g, 65%) as off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.10 (s, 1H), 8.50 (s, 1H), 8.10 (d, 2H), 7.90 (s, 1H), 7.60 (d, 2H), 3.4-3.9 (m, 8H); MS (ESI) *m/z* 434 [C₁₇H₁₅IN₄O₂ + H]⁺.

Intermediate 7: (4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino)methanone

To a solution of 4-(3-bromoimidazo [1, 2-a]pyrazin-6-yl)benzoic acid (8.00 g, 25.14 mmol) in DMF (70 mL), were added sequentially *N*-methyl-morpholine (5.5 mL, 50.28 mmol), HATU (14.4 g, 37.71 mmol) and Morpholine (3.32 mL, 37.71 mmol). The reaction mixture was stirred at room temperature for 3 h under nitrogen atmosphere then, was diluted with water (50 mL). The precipitate was isolated by filtration and dried to afford compound (4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino) methanone (6.40 g, 66%) as off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.15 (d, *J* = 1.2 Hz, 1H), 8.44 (d, *J* = 1.2 Hz, 1H), 8.05 (d, *J* = 8.0 Hz, 2H), 7.83 (s, 1H), 7.57 (d, *J* = 8.0 Hz, 2H), 3.85 (bs, 2H), 3.51 (bs, 2H), 2.53 (bs, 2H), 2.41 (bs, 2H); MS (ESI) *m/z* 387 [C₁₇H₁₅BrN₄O₂ + H]⁺.

Intermediate 8: (6-bromoimidazo[1,2-a]pyrazin-3-yl)benzamide 5a and 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile

To a solution of 3-bromo-6-iodoimidazo [1, 2-a]pyrazine (500 mg, 1.50 mmol) in DMF (20.0 mL) was added 4-carbamoylphenylboronic acid (305 mg, 1.84 mmol), Na₂CO₃ (408 mg, 3.84 mmol), water (2.00 mL) and Pd(PPh₃)₄ (36 mg, 0.03 mmol). The mixture was heated at 90 °C for 2 h and diluted with water. The precipitate was isolated by filtration and dried under vacuum to afford 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzamide (300 mg, 61%) as an off white solid. ¹H NMR (400 MHz, CDCl₃) δ: 9.04 (s, 1H), 8.82 (s, 1H), 8.20 (s, 1H), 8.12 (s, 1H), 8.05 (d, *J* = 8.4 Hz, 2H), 7.86 (d, *J* = 8.4 Hz, 2H), 7.50 (s, 1H); MS (ESI) *m/z* 317 [C₁₃H₉BrN₄O]⁺.


Intermediate 9: 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile

To a solution of 6-bromo-3-iodoimidazo[1,2-a]pyrazine (3 g, 9.26 mmol) in DMF (50 mL), were added 4-cyanophenylboronic acid 3 (1.632 g, 11.1 mmol), K₃PO₄ (4.91 g, 23.15 mmol), Pd(PPh₃)₄ (0.534 mg, 0.46 mmol) and water (5 mL). The reaction mixture was heated at 90 °C for 1 h and water was added to the mixture to induce precipitation. The precipitate was

isolated by filtration and was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 1:1) to afford of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (2.5 g, 90%) as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 9.88 (s, 1H), 8.90 (s, 1H), 8.27 (s, 1H), 8.05-7.99 (m, 4H); MS (ESI) m/z 301.1 $[\text{C}_{13}\text{H}_7\text{BrN}_4+2]^+$.

5

Example 1: (4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




The title was prepared following General procedure B using (4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylcyclohexyl)methanone (intermediate 7) and 4-hydroxyphenylboronic acid as starting materials. The reaction crude product was purified by flash column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford (4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (930 mg, 40%, AUC HPLC 98.2%) as an off-white solid. ^1H NMR (400 MHz, CDCl_3) δ 9.21 (s, 1H), 8.54 (s, 1H), 7.96 (d, $J = 8.4$ Hz, 2H), 7.83 (s, 1H), 7.54 (d, $J = 8.4$ Hz, 2H), 7.43 (d, $J = 8.4$ Hz, 2H), 7.06 (d, $J = 8.8$ Hz, 2H), 3.86 (bs, 2H), 3.51 (bs, 2H), 2.53 (bs, 2H), 2.41 (bs, 2H), 2.35 (s, 3H); MM (ESI) m/z 414 $[\text{C}_{24}\text{H}_{23}\text{N}_5\text{O}_2+\text{H}]^+$.

15

Example 2: (4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino)methanone

20




The title compound was prepared following General procedure B using ((4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino) methanone (intermediate 7) and 4-hydroxyphenylboronic acid as starting materials. The reaction crude product was purified by flash column chromatography (silica gel, 100-200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford 4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(morpholino)methanone (150 mg, 32%, AUC HPLC 97.6%) as an off-white solid. ^1H NMR (400 MHz, CD_3OD) δ 9.88 (s, 1H),

25

9.21 (s, 1H), 8.83 (s, 1H), 8.11 (d, $J = 8.0$ Hz, 2H), 7.94 (s, 1H), 7.61 (d, $J = 8.4$ Hz, 2H), 7.51 (d, $J = 8.4$ Hz, 2H), 6.97 (d, $J = 8.4$ Hz, 2H), 3.60 (bs, 8H); MS (ESI) m/z 401 $[C_{23}H_{20}N_4O_3+H]^+$.

5 Example 3: 4-(6-(4-(4,4-dimethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzamide




Step 1: The title compound was prepared following General procedure A and the reaction crude product was used without further purification in the next step.

- 10 1H NMR (400 MHz, $CDCl_3$) δ : 9.08 (s, 1H), 8.41 (s, 1H), 8.04 (d, $J = 8.4$ Hz, 2H), 7.90 (s, 1H), 7.56 (d, $J = 8.4$ Hz, 2H), 3.75 (bs, 2H), 3.40 (bs, 2H), 1.51 (bs, 2H), 1.34 (s, 2H), 1.02 (s, 6H); MS (ESI) m/z 460 $[C_{20}H_{21}N_4O]^+$.

- Step 2: The title was prepared following General procedure B with (4,4-dimethylpiperidin-1-yl)(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)methanone and 4-carbamoylphenylboronic acid as starting materials. The reaction crude product was purified by column chromatography (silica gel, 100-200 eluent CH_2Cl_2/CH_3OH 90:10) to afford 6-(4-(4,4-dimethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-ylbenzamide (130 mg, 34%, AUC HPLC 95.4%) as an off-white solid. 1H NMR (400 MHz, $CDCl_3$) δ : 9.26 (s, 1H), 8.62 (s, 1H), 8.06 (d, $J = 8.4$ Hz, 2H), 7.96 (s, 1H), 7.95 (d, $J = 4.8$ Hz, 2H), 7.73 (d, $J = 8.4$ Hz, 2H), 7.52 (d, $J = 8.0$ Hz, 2H), 6.25 (bs, 1H), 5.74 (bs, 1H), 3.75 (bs, 2H), 3.39 (bs, 2H), 1.49 (bs, 2H), 1.32 (bs, 2H), 1.02 (s, 6H); MS (ESI) m/z 453 $[C_{27}H_{27}N_5O_2+H]^+$.
- 15
- 20


Intermediate 10: 4-(3-(4-carbamoylphenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid



Step 1: To a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzamide (2.00 g, 6.30 mmol) in a mixture of 1,4-dioxane (25 mL) and water (5 mL), were sequentially added 4-(ethoxycarbonyl)phenylboronic acid (1.30 g, 6.90 mmol), Na_2CO_3 (3.30 g, 6.90 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (363 mg, 0.31 mmol). The reaction mixture was heated at 90 °C for 16 h under argon atmosphere then, was filtered through a short pad of celite. The filtrate was concentrated under reduced pressure and the residue was purified by flash column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford ethyl 4-(3-(4-carbamoylphenyl)imidazo[1,2-a]pyrazin-6-yl)benzoate (1.50 g, 62%) as yellow solid. MS (ESI) m/z 387 [$\text{C}_{22}\text{H}_{18}\text{N}_4\text{O}_3 + \text{H}$] $^+$.

Step 2: To a solution of ethyl 4-(3-(4-carbamoylphenyl)imidazo[1,2-a]pyrazin-6-yl)benzoate (1.50 g, 12.7 mmol) in $\text{THF}/\text{CH}_3\text{OH}$ (20/10 mL) was added $\text{LiOH}\cdot\text{H}_2\text{O}$ (870 mg, 20.7 mmol). The mixture was stirred at rt for 5 h then was concentrated under reduced pressure to a smaller volume. The residue was diluted with water (20 mL) and acidified till pH 3 with an aqueous 2M HCl solution. The precipitate was isolated by filtration and dried to afford 4-(3-(4-carbamoylphenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (750 mg, 93%) as an off-white solid. ^1H NMR (400 MHz, CDCl_3) δ 9.30 (s, 1H), 9.06 (s, 1H), 8.17 (d, $J = 8.4$ Hz, 2H), 8.14 (s, 1H), 8.10 (d, $J = 8.4$ Hz, 2H), 8.03 (d, $J = 8.4$ Hz, 2H), 7.93 (d, $J = 8.4$ Hz, 3H), 7.49 (bs, 1H); MS (ESI) m/z 359 [$\text{C}_{20}\text{H}_{14}\text{N}_4\text{O}_3 + \text{H}$] $^+$.

Example 4: 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzamide hydrochloride




Step 1: *tert*-butyl 1-(4-(3-(4-carbamoylphenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-ylcarbamate was prepared following General procedure A using *tert*-butyl piperidin-4-ylcarbamate and *tert*-butyl piperidin-4-ylcarbamate as starting materials. The reaction crude product was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford the boc protected intermediate. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.28 (s, 1H), 9.00 (s, 1H), 8.16–8.13 (m, 3H), 8.08 (d, $J = 8$ Hz, 2H), 7.93 (d, $J = 8.4$ Hz, 2H), 7.49–7.45 (m, 3H),

6.89 (s, 1H), 4.30–4.20 (m, 1H), 3.55–3.46 (m, 2H), 3.15–3.10 (m, 4H), 1.79–1.71 (m, 2H), 1.23 (s, 9H); MS (ESI) m/z 541 $[\text{C}_{30}\text{H}_{32}\text{N}_6\text{O}_4+\text{H}]^+$.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-carbamoylphenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-ylcarbamate (500 mg, 0.98 mmol) in CH_2Cl_2 (5 mL) was added a 20% HCl solution in 1,4-dioxane (0.15 mL, 0.82 mmol). The reaction mixture was stirred at room temperature for 2 h then, was concentrated under reduced pressure to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl) benzamide hydrochloride salt (90 mg, 15%, AUC HPLC 96%) as yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.54 (s, 1H), 9.09 (s, 1H), 8.48 (s, 1H), 8.23–8.18 (m, 4H), 7.96 (d, $J = 8.0$ Hz, 2H), 7.60 (d, $J = 8.0$ Hz, 2H), 4.80 (bs, 1H), 3.89–3.85 (m, 2H), 3.10–3.00 (m, 2H), 2.10–2.00 (m, 2H), 1.59–1.50 (m, 2H); MS (ESI) m/z 477 $[\text{C}_{25}\text{H}_{24}\text{N}_6\text{O}_2+\text{H}]^+$.


Example 5: 4-(6-(4-(4-*tert*-butylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzamide



The title compound was prepared following General procedure A using *N*-*t*-butyl piperazine. The reaction crude product was purified by column chromatography (silica gel, 100–200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97:3) to afford 4-(6-(4-(4-*tert*-butylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzamide (158 mg, 70%, AUC HPLC 96%) as an off-white solid. ^1H NMR (400 MHz, CDCl_3) δ 9.28 (s, 1H), 9.00 (s, 1H), 8.16 (d, $J = 8.0$ Hz, 2H), 8.13 (s, 1H), 8.08 (d, $J = 8.0$ Hz, 2H), 7.94 (d, $J = 8.0$ Hz, 2H), 7.52 (d, $J = 8.0$ Hz, 3H), 7.45 (bs, 1H), 3.65 (bs, 1H), 2.82 (bs, 1H), 2.70–2.65 (m, 3H), 1.80 (bs, 3H), 1.54 (s, 9H); MS (ESI) m/z 483 $[\text{C}_{28}\text{H}_{30}\text{N}_6\text{O}_2+\text{H}]^+$.


25

Example 6: 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile




To a solution of (4-(3-bromoimidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (200 mg, 0.50 mmol) in toluene (4 mL) and Ethanol (2 mL) under inert atmosphere, was added K₂CO₃ (138 mg, 1.00 mmol), 4-cyanophenylboronic acid (110 mg, 0.75 mmol) and Pd(PPh₃)₄ (58 mg, 0.05 mmol). The resulting mixture was heated in a microwave oven at 140 °C for 15 min, and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (154.7 mg, 73%, AUC HPLC 99%) as a yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.31 (s, 1H), 9.04 (s, 1H), 8.25 (s, 1H), 8.16 (d, *J* = 8.1 Hz, 2H), 8.11-8.04 (m, 4H), 7.51 (d, *J* = 8.1 Hz, 2H), 3.63 (bs, 2H), 3.46-3.20 (m, 2H), 2.32 (bs, 4H), 2.20 (s, 3H); MS (ESI) *m/z* 423 [C₂₅H₂₁FN₆O + H]⁺.

Example 7: 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile




To a solution of (4-(3-iodoimidazo[1,2-*a*]pyrazin-6-yl)phenyl)(morpholino)methanone (200 mg, 0.52 mmol) in a mixture toluene (3 mL) and water (1.5 mL) under inert atmosphere was successively added K₂CO₃ (127 g, 0.92 mmol), 4-cyanophenylboronic acid (111 mg, 0.52 mmol) and Pd(PPh₃)₄ (53 mg, 0.05 mmol). The resulting mixture was microwaved at 140 °C for 15 min then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent DCM/MeOH 95:5) to afford 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (100 mg, 53%, AUC HPLC 95%), yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 9.17 (d, *J* = 0.7 Hz, 1H), 8.55 (d, *J* = 1.0 Hz, 1H), 7.95-7.86 (m, 3H), 7.80 (d, *J* = 8.3 Hz, 2H), 7.72 (d, *J* = 8.3 Hz, 2H), 7.43 (d, *J* = 8.2 Hz, 2H), 3.88-3.22 (m, 8H); MS (ESI) *m/z* 410 [C₂₄H₁₉N₅O₂ + H]⁺.

Example 8: 2-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile



To a solution of 4-(3-iodoimidazo[1,2-*a*]pyrazin-6-yl)phenyl(4-methylpiperazin-1-yl)methanone (400 mg, 0.89 mmol) in a mixture of toluene (8 mL) and ethanol (4 mL) under inert atmosphere, was added K_2CO_3 (247 mg, 1.79 mmol), 2-cyanophenylboronic acid (197 mg, 1.34 mmol) and $Pd(PPh_3)_4$ (103 mg, 0.089 mmol). The resulting mixture was heated in a microwave oven at 140 °C for 15 min, and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5 to 90:10) to afford 2-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (134 mg, 35%, AUC HPLC 99%) as a yellow solid. 1H NMR (400 MHz, CD_3OD) δ 9.13 (d, $J = 0.8$ Hz, 1H), 8.65 (d, $J = 0.8$ Hz, 1H), 8.06 (s, 1H), 8.03-7.98 (m, 3H), 7.91-7.84 (m, 2H), 7.73-7.68 (m, 1H), 7.46 (d, $J = 8.2$ Hz, 2H), 3.77 (bs, 2H), 3.48 (bs, 2H), 2.52-2.42 (m, 4H), 2.33 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.99, 143.89, 141.99, 140.85, 139.26, 137.46, 136.94, 135.46, 135.09, 132.04, 131.99, 131.26, 128.72, 127.97, 125.66, 55.96, 55.53, 46.01, 43.00; MS (ESI) m/z 423 [$C_{25}H_{22}N_6O + H$] $^+$.


Example 9: 4-(6-(4-(4-(methylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile



Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (10.0 mL), were added HATU (168 mg, 0.441 mmol), *N*-methylmorpholine (59 mg, 0.588 mmol) and *tert*-butyl methyl(piperidin-4-yl)carbamate (126 mg, 0.588 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, and then was diluted with water (20 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5 to 90:10) to afford the boc-protected derivative as a yellow solid. MS (ESI) m/z 537 [$C_{31}H_{32}N_6O_3 + H$] $^+$.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl(methyl)carbamate (0.294 mmol) in methanol (3 mL) and dioxane (3 mL), was added a solution of 4M HCl in dioxane (3 mL). The resulting mixture was stirred at room temperature for 18 h and then was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-(methylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (63.3 mg, 50%, AUC HPLC 96%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 9.22 (s, 1H), 8.97 (s, 1H), 8.17 (d, *J* = 8.2 Hz, 2H), 8.13 (s, 1H), 8.03-7.98 (m, 4H), 7.57 (d, *J* = 8.2 Hz, 2H), 4.75-4.72 (m, 1H), 3.91 (bs, 1H), 3.17-3.11 (m, 2H), 3.01-2.98 (m, 1H), 2.64 (s, 3H), 2.19-2.05 (m, 2H), 1.51 (bs, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 172.10, 144.07, 142.39, 141.09, 139.45, 137.02, 136.83, 134.47, 133.68, 129.78, 128.56, 128.03, 119.37, 115.42, 113.53, 57.32, 31.51; MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.

Example 10: 4-(6-(4-(4-(2-aminoethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoic acid (50 mg, 0.147 mmol) in DMF (5.0 mL) were added HATU (84 mg, 0.221 mmol), *N*-methyl morpholine (30 mg, 0.294 mmol) and *tert*-butyl 2-(piperidin-4-yl)ethylcarbamate (67 mg, 0.294 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h and then diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford the carbamate derivative as a yellow solid. MS (ESI) *m/z* 551 [C₃₂H₃₄N₆O₃+H]⁺.

Step 2: To a solution of *tert*-butyl 2-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl)ethylcarbamate (0.147 mmol) in DCM (1 mL) was added TFA (1 mL). The reaction mixture was stirred at room temperature for 18 h and then concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-(2-aminoethyl)piperidine-1-

carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (15.9 mg, 22%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 9.19 (d, *J* = 1.3 Hz, 1H), 8.93 (d, *J* = 1.3 Hz, 1H), 8.13-8.11 (m, 3H), 8.01-7.96 (m, 4H), 7.53 (d, *J* = 8.4 Hz, 2H), 4.69-4.66 (m, 1H), 3.82-3.80 (m, 1H), 3.18-3.15 (m, 1H), 3.04-3.00 (m, 2H), 2.91 (bs, 1H), 1.89 (bs, 1H), 1.75-1.64 (m, 4H), 1.30-1.27 (m, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 171.92, 167.39, 144.03, 142.35, 141.11, 139.14, 137.48, 136.83, 134.46, 133.65, 129.74, 128.50, 127.94, 119.39, 115.32, 113.48, 43.46, 38.38, 34.93, 34.58, 33.46, 32.58; MS (ESI) *m/z* 451 [C₂₇H₂₆N₆O + H]⁺.


10 Example 11: 4-(6-(4-(4-methyl-1,2,3,6-tetrahydropyridine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile



To a solution of 4-(3-(4-carbamoylphenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoic acid (100mg, 0.279 mmol) in DMF (0.56 mL) was added phosphorus oxychloride (51 μl, 0.558 mmol) dropwise. The reaction mixture was stirred at room temperature under inert atmosphere for 4 h, followed by the addition of 4-methylpiperidin-4-ol (43 mg, 0.279 mmol) and anhydrous pyridine (1 mL). The reaction mixture was cooled to 0°C, and phosphorus oxychloride (28 μl, 0.307 mmol) was added dropwise. The reaction mixture was stirred at 0°C for at least 30min, and allowed to warm to room temperature overnight. The reaction mixture was then diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford 4-(6-(4-(4-methyl-1,2,3,6-tetrahydropyridine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (36 mg, 31%, AUC HPLC 93%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 9.29 (s, 1H), 8.59 (s, 1H), 7.98-7.95 (m, 3H), 7.91-7.88 (m, 2H), 7.77-7.75 (m, 2H), 7.56-7.54 (m, 2H), 5.48-5.23 (br s, 1H), 4.17 (bs., 1H), 3.88 (bs., 2H), 3.50 (bs., 1H), 2.16-2.07 (m, 2H), 1.74 (s., 3H); ¹³C NMR (400 MHz, CDCl₃) δ 144.12, 141.20, 140.31, 137.37, 136.99, 136.25, 133.45, 132.49, 128.18, 126.49, 125.66, 118.13, 112.69, 112.38, 23.10; MS (ESI) *m/z* 420 [C₂₆H₂₁N₅O + H]⁺.


30

Example 12: 4-(6-(4-(4-ethylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



- Step 1: (4-ethylpiperazin-1-yl)(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)methanone was prepared following General procedure A and the reaction crude product was used without further purification in the next step. ^1H NMR (400 MHz, CDCl_3) δ 9.11 (s, 1H), 8.72 (s, 1H), 8.10 (s, 1H), 8.19 (d, $J = 8.0$ Hz, 2H), 7.51 (d, $J = 8.0$ Hz, 2H), 3.61 (bs, 2H), 3.51 (bs, 4H), 2.39 (bs, 2H), 2.34 (m, 2H), 1.00 (s, 3H); MS (ESI) m/z 461 [$\text{C}_{19}\text{H}_{20}\text{IN}_5\text{O}$] $^+$
- Step 2: The title compound was prepared following General procedure B with (4-ethylpiperazin-1-yl)(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)methanone and 4-cyanophenylboronic acid as starting materials. The reaction crude product was purified by column chromatography (silica gel, 100-200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford 4-(6-(4-(4-ethylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (95 mg, 20%, AUC HPLC 95.7%) as a light green solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.30 (s, 1H), 8.60 (s, 1H), 7.99 (s, 1H), 7.97 (d, $J = 8$ Hz, 2H), 7.91 (d, $J = 8.4$ Hz, 2H), 7.77 (d, $J = 8$ Hz, 2H), 7.55 (d, $J = 8$ Hz, 2H), 3.86–3.80 (m, 2H), 3.50–3.43 (m, 2H), 2.60–2.40 (m, 6H), 1.10 (t, $J = 5.4$ Hz, 3H); MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.

Example 13: 4-(6-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



- Step 1: (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-isopropylpiperazin-1-yl)methanone was prepared following General procedure A and the reaction crude product was used without further purification in the next step.

^1H NMR (400 MHz, CDCl_3) δ 9.15 (d, $J = 1.2$ Hz, 1H), 8.44 (d, $J = 1.2$ Hz, 1H), 8.05 (d, $J = 8.0$ Hz, 2H), 7.83 (s, 1H), 7.57 (d, $J = 8.0$ Hz, 2H), 3.85 (bs, 2H), 3.51 (bs, 2H), 2.53 (bs, 2H), 2.41 (bs, 2H), 2.05–2.00 (m, 1H), 1.35 (d, $J = 8.0$ Hz, 6H); MS (ESI) m/z 475 $[\text{C}_{20}\text{H}_{22}\text{N}_5\text{O}]^+$.


5

Step 2: 4-(6-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile was prepared following General procedure B with (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-isopropylpiperazin-1-yl)methanone and 4-cyanophenylboronic acid as starting materials. The reaction crude product was purified by column chromatography

10 (silica gel, 100–200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98:2) to afford 4-(6-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (60 mg, 23%, AUC HPLC >99%) as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.30 (s, 1H), 9.03 (s, 1H), 8.23 (s, 1H), 8.14 (d, $J = 8.0$ Hz, 2H), 8.03–8.09 (m, 4H), 7.50 (d, $J = 8.4$ Hz, 2H), 3.60 (bs, 4H), 2.66–2.72 (m, 1H), 2.49 (bs, 4H), 0.96 (d, $J = 6.4$ Hz, 6H); MS (ESI) m/z 451

15 $[\text{C}_{27}\text{H}_{26}\text{N}_6\text{O}+\text{H}]^+$.

Example 14: 4-(6-(4-(4-cyclopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



20 Step 1: (4-cyclopropylpiperazin-1-yl)(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl) methanone was prepared following General procedure A and the reaction crude product was used without further purification in the next step.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.21 (s, 1H), δ 8.89 (s, 1H), 7.96 (d, $J = 8.0$ Hz, 2H), 7.83 (s, 1H), 7.55 (d, $J = 8.0$ Hz, 2H), 3.75 (bs, 4H), 3.53 (bs, 4H), 2.84–2.90 (m, 1H), 1.10 (s, 2H), 0.77 (d, $J = 8.0$ Hz, 2H); MS (ESI) m/z 473 $[\text{C}_{20}\text{H}_{20}\text{N}_5\text{O}+\text{H}]^+$.


25

Step 2: The title compound was prepared following General procedure B with (4-cyclopropylpiperazin-1-yl)(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl) methanone and 4-cyanophenylboronic acid as starting materials. The reaction crude product was purified by

30 column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97:3) to afford 4-(6-(4-(4-

cyclopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (65 mg, 28%, AUC HPLC 96.9%) as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 9.30 (s, 1H), 9.03 (s, 1H), 8.23 (s, 1H), 8.15 (d, J = 8.0 Hz, 2H), 8.03–8.09 (m, 4H), 7.50 (d, J = 8.0 Hz, 2H), 3.58 (bs, 4H), 2.62 (bs, 4H), 1.64–1.67 (m, 1H), 0.32 (d, J = 8.0 Hz, 2H), 0.42 (d, J = 8.0 Hz, 2H); MS (ESI) m/z 449 [$\text{C}_{27}\text{H}_{24}\text{N}_6\text{O}+\text{H}$] $^+$.

Example 15: 4-(6-(4-(4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



Step 1: 1-(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-one was prepared following General procedure A and the reaction crude product was used without further purification in the next step.


^1H NMR (400 MHz, CDCl_3) δ 9.15 (d, J = 1.2 Hz, 1H), 8.44 (d, J = 1.2 Hz, 1H), 8.05 (d, J = 8.0 Hz, 2H), 7.83 (s, 1H), 7.57 (d, J = 8.0 Hz, 2H), 3.85 (bs, 2H), 3.51 (bs, 2H), 2.53 (bs, 2H), 2.41 (bs, 2H); MS (ESI) m/z 446 [$\text{C}_{18}\text{H}_{15}\text{IN}_4\text{O}_2+\text{H}$] $^+$.

Step 2: 4-(6-(4-(4-oxopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile was prepared following General procedure B with 1-(4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-one and 4-cyanophenylboronic acid as starting materials. The reaction crude product was purified by column chromatography (silica gel, 100-200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97:3) to afford 4-(6-(4-(4-oxopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (20 mg, 24%, AUC HPLC 96.9%) as an off-white solid; ^1H NMR (400 MHz, DMSO- d_6) δ 9.31 (s, 1H), 9.05 (s, 1H), 8.24 (s, 1H), 8.18 (d, J = 8.4 Hz, 2H), 8.04–8.09 (m, 4H), 7.61 (d, J = 8.4 Hz, 2H), 3.75 (bs, 8H); MS (ESI) m/z 422 [$\text{C}_{25}\text{H}_{19}\text{N}_5\text{O}_2+\text{H}$] $^+$.

Step 3: To a solution of 4-(6-(4-(4-oxopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (140 mg, 0.34 mmol) in methanol (5 mL) was added sodium borohydride (19 mg, 0.5 mmol). The reaction mixture was stirred at room temperature for 1 h and was concentrated under reduced pressure. The residue was purified by flash column

chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 90:10) to afford 4-(6-(4-(4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (65 mg, 50%, AUC HPLC >99%) as gray solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.30 (s, 1H), 9.03 (s, 1H), 8.23 (s, 2H), 8.03–8.09 (m, 5H), 7.48 (d, $J = 8.4$ Hz, 2H), 4.10 (d, $J = 8.0$ Hz, 1H), 1.22 (bs, 8H); MS (ESI) m/z 424 $[\text{C}_{25}\text{H}_{21}\text{N}_5\text{O}_2 + \text{H}]^+$.

Intermediate 11: ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoate




Intermediate 11

Step 1: To a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (5.00 g, 16.7 mmol) in DMF (100 mL), were added 4-(ethoxycarbonyl)phenylboronic acid (4.90 g, 25.2 mmol), Na_2CO_3 (4.40 g, 41.5 mmol), water (5.0 mL) and $\text{Pd}(\text{PPh}_3)_4$ (386 mg, 0.33 mmol). The reaction mixture was heated at 90°C for 2 h under argon atmosphere then, was diluted with water. The precipitate was isolated by filtration and dried under vacuum to afford ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoate (2.80 g, 46%) as off white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.28 (s, 1H), 9.07 (s, 1H), 8.25 (s, 1H), 8.22 (d, $J = 6.0$ Hz, 2H), 8.08–8.02 (m, 6H), 4.38–4.33 (m, 2H), 1.36 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 369 $[\text{C}_{22}\text{H}_{16}\text{N}_4\text{O}_2 + \text{H}]^+$.


Step 2: To a solution of ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoate (2.80 g, 7.61 mmol) in $\text{THF}/\text{CH}_3\text{OH}/\text{H}_2\text{O}$ (50/20/20 mL) was added $\text{LiOH}\cdot\text{H}_2\text{O}$ (1.25 g, 30.5 mmol). The reaction mixture was stirred at rt for 12 h and was concentrated to a smaller volume. The residue was diluted with water (20 mL) and acidified till pH 2. The precipitate was filtered off and dried to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (2.00 g, 80%) as off white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.30 (s, 1H), 9.08 (s, 1H), 8.22 (d, $J = 9.6$ Hz, 3H), 8.06–8.02 (m, 6H); MS (ESI) m/z 341 $[\text{C}_{20}\text{H}_{12}\text{N}_4\text{O}_2 + \text{H}]^+$.

Example 16: 4-(6-(4-(4-hydroxy-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methyl morpholine (130 μ l, 1.178 mmol). The reaction mixture was stirred at room temperature
 5 under inert atmosphere for 1 h, followed by the addition of 4-methylpiperidin-4-ol hydrochloride (89 mg, 0.587 mmol). The reaction mixture was left to stir for 18 h, then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford 4-
 10 (6-(4-(4-hydroxy-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (73 mg, 57%, AUC HPLC 97%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.28 (s, 1H), 8.59 (s, 1H), 7.98-7.94 (m, 3H), 7.90-7.88 (m, 2H), 7.77-7.75 (m, 2H), 7.53 (d, *J*=8.4 Hz, 2H), 4.35 (bs, 1H), 3.51-3.38 (m, 3H), 1.70-1.55 (m, 4H), 1.32 (s, 3H); ¹³C NMR (400 MHz, CDCl₃) δ 169.70, 144.12, 141.20, 140.29, 137.33, 136.86, 136.24, 133.45,
 15 132.47, 128.19, 127.70, 126.54, 125.68, 118.13, 112.70, 112.39, 68.13, 30.41; MS (ESI) *m/z* 438 [C₂₆H₂₃N₅O₂ + H]⁺.


Example 17: 4-(6-(4-(4-methyl-1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



20 The title compound was prepared according to genral procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and 1-methyl-homopiperazine as starting materials. The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (10 mL) and extracted with EtOAc (3x30
 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(4-methyl-1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-

- a]pyrazin-3-yl)benzonitrile (11 mg, 17%, AUC HPLC 99%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.30 (d, $J = 1.2$ Hz, 1H), 9.03 (d, $J = 1.2$ Hz, 1H), 8.24 (s, 1H), 8.15 (d, $J = 8.4$ Hz, 2H), 8.11–8.01 (m, 4H), 7.50 (d, $J = 7.2$ Hz, 2H), 3.70–3.60 (m, 4H), 2.70–2.64 (m, 1H), 2.61–2.55 (m, 1H), 2.60–2.50 (m, 2H), 2.29 (d, $J = 26.0$ Hz, 3H), 1.91–1.82 (m, 1H), 1.80–1.70 (m, 1H); MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.


Example 18: 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile Hydrochloride



- Step 1: *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl) benzoyl)piperidin-4-ylcarbamate was prepared following General procedure A and the reaction crude product was purified by column chromatography (silica gel, 100–200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 97:3). ^1H NMR (400 MHz, CDCl_3) δ 9.30 (s, 1H), 8.61 (s, 1H), 8.02 (bs, 1H), 8.00 (s, 1H), 7.98 (d, $J = 8.4$ Hz, 2H), 7.96 (d, $J = 8.0$ Hz, 2H), 7.79 (d, $J = 8.4$ Hz, 2H), 7.53 (d, $J = 8.0$ Hz, 2H), 4.30–4.25 (m, 1H), 3.74–3.70 (m, 4H), 2.10–2.06 (m, 4H), 1.45 (bs, 9H); MS (ESI) m/z 523 [$\text{C}_{30}\text{H}_{30}\text{N}_6\text{O}_3 + \text{H}$] $^+$.

- Step 2: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl) benzoyl)piperidin-4-ylcarbamate (100 mg, 0.90 mmol) in a mixture of CH_2Cl_2 (5 mL), and 20% HCl solution in 1,4-dioxan (0.15 mL, 0.82 mmol) was stirred at room temperature for 2 h. The reaction mixture was concentrated to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile Hydrochloride (52 mg, 59%, AUC HPLC 98.0%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 9.32 (s, 1H), 9.04 (s, 1H), 8.3 (bs, 1H), 8.26 (s, 1H), 8.18–8.16 (m, 3H), 8.07–8.04 (m, 4H), 7.48 (d, $J = 8.0$ Hz, 2H), 3.30–3.20 (m, 2H), 2.01–1.96 (m, 4H), 1.49–1.39 (m, 2H); MS (ESI) m/z 458 [$\text{C}_{25}\text{H}_{22}\text{N}_6\text{O} + \text{H}$] $^+$.


Example 19: 4-(6-(4-(4,4-dimethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



The title compound was prepared following General procedure A with 4,4-dimethylpiperidine as starting material. The reaction crude product was purified by flash column chromatography (silica gel, 100-200 eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95:5) to afford 4-(6-(4-(4,4-dimethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (35 mg, 14%, AUC HPLC 95%) as a light brown solid. ^1H NMR (400 MHz, CDCl_3) δ 9.29 (s, 1H), 8.59 (s, 1H), 7.99–7.89 (m, 5H), 7.77 (d, $J = 8.4$ Hz, 2H), 7.53 (d, $J = 8.4$ Hz, 2H), 3.75 (bs, 2H), 3.39 (bs, 2H), 1.49 (bs, 2H), 1.32 (bs, 2H), 1.01 (s, 6H); MS (ESI) m/z 436 $[\text{C}_{27}\text{H}_{25}\text{N}_5\text{O} + \text{H}]^+$.

10

Example 20: 4-(6-(4-(4-(aminomethyl)-4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



Step 1: To a solution of *t*-butyl-4-(aminomethyl)-4-hydroxypiperidine-1-carboxylate (2.0 g, 8.69 mmol) in THF, were added DMAP (110 mg, 0.86 mmol) and ethyl 2,2,2-trifluoroacetate (1.5 g, 10.43 mmol). The reaction mixture was heated at 70 °C and stirred for 6 h then was diluted with ethyl acetate. The organic layer was washed in turn with 1N HCl (2×10 mL) and brine then, dried over Na_2SO_4 , filtered and concentrated. The residue was purified by flash column chromatography (silica gel, eluent *n*-Hexane/EtOAc 70:30) to afford *tert*-butyl 4-hydroxy-4-((2,2,2-trifluoroacetamido)methyl)piperidine-1-carboxylate (2.00 g, 84%) as an oil. ^1H NMR (400 MHz, CDCl_3) δ 6.90 (bs, 1H), 3.79 (bs, 2H), 3.41 (bs, 2H), 3.25–3.18 (m, 4H), 1.57 (bs, 4H), 1.48 (s, 9H); MS (ESI) m/z 327 $[\text{C}_{13}\text{H}_{21}\text{F}_3\text{N}_2\text{O}_4 + \text{H}]^+$.


Step 2: A solution of *tert*-butyl 4-hydroxy-4-((2,2,2-trifluoroacetamido)methyl)piperidine-1-carboxylate (500 mg, 8.69 mmol) in a mixture of 1,4-dioxane and 30% HCl solution in 1,4-

dioxane (10 mL). The reaction mixture was stirred for 3 h at room temperature and was concentrated to afford 2,2,2-trifluoro-N-((4-hydroxypiperidin-4-yl)methyl)acetamide (400 mg, 90%) as an oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 5.08 (bs, 1H), 3.37 (bs, 2H), 3.20 (d, *J* = 6.4 Hz, 2H), 3.10–3.07 (m, 2H), 3.01–2.94 (m, 2H), 1.72–1.64 (m, 2H), 1.57–1.54 (m, 2H); MS (ESI) *m/z* 227 [C₈H₁₄ClF₃N₂O₂ + H]⁺.

Step 3: To a solution of 4-(3-(4-hydroxyphenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-indazol-5-yl)benzoic acid (300 mg, 0.88 mmol) in DMF (5 mL) were added *N*-methylmorpholine (270 mg, 3.57 mmol), HATU (503 mg, 1.32 mmol) and 2,2,2-trifluoro-N-((4-hydroxypiperidin-4-yl)methyl)acetamide (350 mg, 3.57 mmol). The reaction mixture was stirred for 10 h at room temperature and was quenched by adding water. The aqueous phase was extracted with CH₂Cl₂ and the organic layer was washed with brine solution, dried over Na₂SO₄, filtered and concentrated. The crude compound was purified by flash column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 95:5) to afford N-((1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)-4-hydroxypiperidin-4-yl)methyl)-2,2,2-trifluoroacetamide (70 mg, 20%) as an off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.29 (s, 2H), 9.03 (s, 1H), 8.23 (s, 1H), 8.14 (d, *J* = 8.4 Hz, 2H), 8.09–8.03 (m, 4H), 7.49 (d, *J* = 8.0 Hz, 2H), 4.78 (s, 1H), 4.20 (bs, 1H), 3.40 (bs, 1H), 3.28 (bs, 1H), 3.22 (d, *J* = 6.0 Hz, 2H), 3.15 (bs, 1H), 1.52–1.40 (m, 4H); MS (ESI) *m/z* 549 [C₂₈H₂₃F₃N₆O₃ + H]⁺.

Step 4: To a solution of N-((1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)-4-hydroxypiperidin-4-yl)methyl)-2,2,2-trifluoroacetamide (100 mg, 0.17 mmol) in methanol (5 mL) was added K₂CO₃ (97 mg, 0.44 mmol) was added. The reaction mixture was stirred for 5 h then was concentrated under reduced pressure. The residue was purified by flash column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 90:10) to afford 4-(6-(4-(4-(aminomethyl)-4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (50 mg, 40%, AUC HPLC 96.7%) as a light yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.20 (s, 1H), 8.95 (s, 1H), 8.15–8.11 (m, 3H), 8.02–7.96 (m, 4H), 7.55 (d, *J* = 8.4 Hz, 2H), 4.41 (bs, 1H), 3.65–3.49 (bs, 2H), 3.35 (s, 2H), 2.94 (s, 2H), 1.77–1.63 (m, 4H); MS (ESI) *m/z* 453 [C₂₆H₂₄N₆O₂ + H]⁺.


Example 21: 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile



The title compound was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and NN-dimethyl-4-aminopiperidine as starting materials. The reaction crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 96:4) to afford 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (65 mg, 22%, AUC HPLC 96%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.31 (s, 1H), 9.03 (s, 1H), 8.24 (s, 1H), 8.16 (d, *J* = 8.0 Hz, 2H), 8.04–8.09 (m, 4H), 7.52 (d, *J* = 8.0 Hz, 2H), 3.55 (bs, 4H), 3.08–3.16 (m, 1H), 2.85 (t, 6H), 2.25 (bs, 4H); MS (ESI) *m/z* 451 [C₂₇H₂₆N₆O+H]⁺.

10


Example 22: *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-yl)acetamide



The title compound was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and 4-*N*-acetamido-piperidine as starting materials. The reaction crude production was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 93:7) to afford *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-yl)acetamide (80 mg, 29%, AUC HPLC 96%) as an off-white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.21 (s, 1H), 8.96 (s, 1H), 8.14 (d, *J* = 8.4 Hz, 2H), 8.11 (s, 1H), 8.02–7.96 (m, 4H), 7.54 (d, *J* = 8.0 Hz, 2H), 3.96 (bs, 1H), 3.72 (s, 3H), 3.09 (bs, 1H), 2.51 (bs, 2H), 4.59 (bs, 1H), 2.02 (bs, 1H), 1.94 (s, 3H), 1.88 (bs, 1H); MS (ESI) *m/z* 465 [C₂₇H₂₄N₆O₂+H]⁺.

20


Example 23: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile hydrochloride salt



Step 1: *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and *tert*-butyl 4-methylpiperidin-4-ylcarbamate as starting materials. The reaction crude production was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 95:5). ¹H NMR (400 MHz, CDCl₃) δ 9.30 (s, 1H), 9.03 (s, 1H), 8.23 (s, 1H), 8.12 (d, *J* = 8.0 Hz, 2H), 8.09–8.03 (m, 4H), 7.49 (d, *J* = 8.0 Hz, 2H), 6.62 (bs, 1H), 3.39–3.35 (m, 1H), 3.21 (bs, 3H), 2.10–1.96 (m, 3H), 1.37–1.30 (m, 13H); MS (ESI) *m/z* 537 [C₃₁H₃₂N₆O₃ + H]⁺.


Step 2: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (150 mg, 0.27 mmol) in a mixture of CH₂Cl₂ (5 mL) and 20% HCl solution in 1,4-dioxane (0.15 mL, 0.82 mmol) was stirred at room temperature for 2 h. The reaction mixture was concentrated to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile hydrochloride salt (76 mg, 60%, AUC HPLC 97%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.32 (s, 1H), 9.04 (s, 1H), 8.26 (s, 1H), 8.24 (bs, 1H), 8.17 (d, *J* = 12.0 Hz, 2H), 8.09–8.04 (m, 4H), 7.49 (d, *J* = 8.0 Hz, 2H), 3.39–3.29 (m, 3H), 1.76–1.58 (m, 4H), 1.39 (bs, 3H); MS (ESI) *m/z* 473 [C₂₆H₂₄N₆O + H]⁺.

Example 24: 4-(6-(4-(2-oxa-7-azaspiro[3.5]nonane-7-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




The title compound was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and 2-oxa-7-azaspiro[3.5]nonane as starting materials. The reaction crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 95:5) to afford 4-(6-(4-(2-oxa-7-azaspiro[3.5]nonane-7-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (58 mg, 25%, AUC HPLC 96%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.30 (s, 1H), 9.04 (s, 1H), 8.24 (s, 1H), 8.14 (d, *J* = 8.0 Hz, 2H), 8.09–8.03 (m, 4H), 7.27 (d, *J* = 8.0 Hz, 2H), 4.33 (bs, 4H), 3.53 (bs, 2H), 3.28 (bs, 2H), 1.78–1.72 (m, 4H); MS (ESI) *m/z* 450 [C₂₇H₂₃N₅O₂ + H]⁺.

10 Example 25: 4-(6-(4-(2-oxa-6-azaspiro[3.3]heptane-6-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



The title compound was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and 2-oxa-6-azaspiro[3.3]heptane as starting material. The reaction crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 93:7) to afford 4-(6-(4-(2-oxa-6-azaspiro[3.3]heptane-6-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (40 mg, 33%, AUC HPLC 96%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.30 (s, 1H), 9.09 (s, 1H), 8.25 (s, 1H), 8.12 (d, *J* = 8.4 Hz, 2H), 8.01–8.10 (m, 4H), 7.27 (d, *J* = 8.0 Hz, 2H), 4.70 (bs, 4H), 4.50 (bs, 2H), 4.20 (bs, 2H); MS (ESI) *m/z* 422 [C₂₅H₁₉N₅O₂ + H]⁺.


20 Example 26: 4-(6-(4-(4-tert-butylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



The title compound was prepared following General procedure A using 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid and N-*t*-butyl piperazine as starting materials. The reaction crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 95:5) to afford 4-(6-(4-(4-*tert*-butylpiperazine-1-

- 5 carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (137 mg, 27%, AUC HPLC 98%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.29 (s, 1H), 9.02 (s, 1H), 8.21 (s, 1H), 8.14 (d, *J* = 8.0 Hz, 2H), 8.13–8.03 (m, 4H), 7.49 (d, *J* = 8.0 Hz, 2H), 3.59 (bs, 2H), 2.49–2.43 (m, 6H), 1.02 (s, 9H); MS (ESI) *m/z* 465 [C₂₈H₂₈N₆O + H]⁺.

10 Example 27: 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-methylphenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




- Step 1: To a solution of 6-bromo-3-iodoimidazo[1,2-a]pyrazine (5.00 g, 15.4 mmol) in DMF (60 mL) under inert atmosphere, was added Na₂CO₃ (4.08 g, 38.6 mmol), 4-cyanophenylboronic acid (2.66 g, 16.9 mmol) and Pd(PPh₃)₄ (356 mg, 0.31 mmol). The resulting mixture was stirred at 80 °C for 48 h, then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent hexane/EtOAc 3:2) to afford the crude 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (1.51 g, 33%) as a pink solid. ¹H NMR (400 MHz, CD₃OD) δ 9.07 (d, *J* = 1.2 Hz, 1H), 8.88 (d, *J* = 1.2 Hz, 1H), 8.27 (s, 1H), 8.02 (q, *J* = 7.8 Hz, 4H); MS (ESI) *m/z* 301 [C₁₃H₇BrN₄ + H]⁺.
- 20

- Step 2: 4-borono-2-methylbenzoic acid (100 mg, 0.55 mmol), K₂CO₃ (138 mg, 1.00 mmol) and Pd(dppf)Cl₂ (73.1 mg, 0.01 mmol) were added sequentially to a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (150 mg, 0.50 mmol) in DMF at room temperature under inert atmosphere. The reaction mixture was refluxed for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methylbenzoic acid (105 mg, 83 %) as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) 12.87 (bs, 1H), 9.31 (s, 1H), 9.07 (s, 1H), 8.24 (s, 1H), 8.10–8.04 (m, 5H), 8.02 (d, *J* = 8.2 Hz, 1H), 7.94 (d, *J* = 8.2 Hz, 1H), 2.62 (s, 3H); MS (ESI) *m/z* 355 [C₂₁H₁₄N₄O₂ + H]⁺.
- 25
- 30

Step 3: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methylbenzoic acid (50 mg, 0.14 mmol) in DMF (3 mL), were added HATU (80 mg, 0.21 mmol), *N*-methyl morpholine (57 mg, 0.56 mmol) and *tert*-butyl piperidin-4-ylcarbamate hydrochloride (56 mg, 0.30 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent EtOAc) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methylbenzoyl)piperidin-4-ylcarbamate (63 mg, 83%) as off-white powder. MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.

Step 4: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methylbenzoyl)piperidin-4-ylcarbamate in DCM (3 mL) and TFA (3 mL) was stirred 18 h at room temperature. The reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-methylphenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (38 mg, 94%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.16 (d, *J* = 1.4 Hz, 1H), 8.89 (d, *J* = 1.4 Hz, 1H), 8.08 (s, 1H), 8.00 – 7.88 (m, 6H), 7.33 (t, *J* = 8.6 Hz, 1H), 4.78 (s, 1H), 3.60 (bs, 1H), 3.43 (t, *J* = 11.3 Hz, 1H), 3.21 (t, *J* = 11.9 Hz, 1H), 2.97 (bs, 3H), 2.38 (d, *J* = 3.8 Hz, 3H), 2.17 (d, *J* = 12.3 Hz, 1H), 1.97 (bs, 1H), 1.71 – 1.42 (m, 2H); ¹³C NMR (400 MHz, CD₃OD) δ 171.84, 163.22, 163.88, 162.53, 143.96, 142.33, 141.26, 138.66, 137.22, 136.79, 134.45, 133.65, 129.74, 127.91, 127.57, 127.21, 125.59, 119.72, 119.38, 115.20, 113.47, 46.00, 40.81, 31.85, 31.36, 30.84, 19.05; MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.


Example 28: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-methylphenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methylbenzoic acid (55 mg, 0.15 mmol) in DMF (3 mL), were added HATU (86 mg, 0.30 mmol), *N*-methyl

- morpholine (31 mg, 0.30 mmol) and *tert*-butyl piperidin-4-ylcarbamate hydrochloride (65 mg, 0.30 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and the aqueous phase was extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent EtOAc) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)-2-methylbenzoyl)-4-methylpiperidin-4-ylcarbamate, **7** (36 mg, 83%) as an off-white powder. MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.
- Step 2: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)-2-methylbenzoyl)-4-methylpiperidin-4-ylcarbamate in DCM (3 mL) and TFA (3 mL) was stirred 18 h at room temperature. The reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-methylphenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (37 mg, 54%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, CD₃OD) 9.18 (d, *J* = 1.4 Hz, 1H), 8.91 (d, *J* = 1.4 Hz, 1H), 8.10 (s, 1H), 8.05–7.86 (m, 6H), 7.32 (d, *J* = 8.0 Hz, 1H), 4.37–3.90 (m, 2H), 3.78–3.51 (m, 1H), 3.61–3.32 (m, 1H), 2.98 (t, *J* = 8.6 Hz, 3H), 1.93–1.64 (m, 2H), 1.61 (bs, 2H), 1.36 (s, 3H); ¹³C NMR (400 MHz, CD₃OD) δ 170.32, 142.54, 140.94, 139.92, 137.19, 136.04, 135.36, 134.88, 132.27, 128.43, 126.51, 124.11, 117.95, 113.77, 112.07, 37.45, 36.95, 36.30, 17.72; MS (ESI) *m/z* 451 [C₂₇H₂₆N₆O + H]⁺.

Example 29: 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-fluorophenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile




- Step 1: 4-bromo-2-fluorobenzonitrile (102 mg, 0.55 mmol), K₂CO₃ (138 mg, 1.00 mmol) and Pd(dppf)Cl₂ (73.1 mg, 0.01 mmol) were added sequentially to a solution of 4-(6-bromoimidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (150 mg, 0.50 mmol) in DMF under inert atmosphere. The reaction mixture was refluxed for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude -(3-(4-

cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoic acid (161 mg, 67 %) as brown solid. MS (ESI) m/z 359 [$C_{20}H_{11}FN_4O_2 + H$] $^+$.

- Step 2: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoic acid (60 mg, 0.17 mmol) in DMF (3 mL), were successively added HATU (95 mg, 0.25 mmol), *N*-methylmorpholine (67 mg, 0.67 mmol) and *tert*-butyl piperidin-4-ylcarbamate hydrochloride (67 mg, 0.34 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and the aqueous phase was extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent EtOAc) to afford 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoyl)piperidin-4-ylcarbamate (70 mg, 76%) as an off-white powder. MS (ESI) m/z 441 [$C_{25}H_{21}FN_6O + H$] $^+$.
- Step 3: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoyl)piperidin-4-ylcarbamate (70 mg, 0.13 mmol) in DCM (3 mL) and TFA (3 mL) was stirred 18 h at room temperature. The reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /HCOOH 0.01%) to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-fluorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (39 mg, 53%, AUC HPLC 98%) as a white solid. 1H NMR (400 MHz, CD_3OD) δ 9.20 (d, $J=1.4$ Hz, 1H), 9.00 (d, $J=1.4$ Hz, 1H), 8.11 (s, 1H), 8.07-7.91 (m, 6H), 7.50 (t, $J=7.5$ Hz, 1H), 4.75 (d, $J=13.7$ Hz, 1H), 3.69 (d, $J=12.6$ Hz, 1H), 3.39-3.18 (m, 2H), 2.98 (t, $J=11.7$ Hz, 1H), 2.11 (d, $J=12.1$ Hz, 1H), 1.97 (d, $J=11.7$ Hz, 1H), 1.97-1.45 (m, 2H); ^{13}C NMR (400 MHz, CD_3OD) δ 170.23, 166.98, 161.24, 158.80, 144.16, 142.48, 141.84, 139.82, 136.98, 134.49, 133.61, 130.45, 129.86, 128.20, 125.04, 124.85, 124.06, 124.03, 119.40, 115.980, 115.24, 115.01, 113.61, 46.78, 41.45, 32.61, 31.88; MS (ESI) m/z 441 [$C_{25}H_{21}FN_6O + H$] $^+$.

Example 30: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-fluorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoic acid (90 mg, 0.25 mmol) in DMF (3 mL), was successively added HATU (95 mg, 0.25 mmol), *N*-methylmorpholine (55 mg, 0.50 mmol) and *tert*-butyl piperidin-4-ylcarbamate hydrochloride (106 mg, 0.50 mmol). The reaction mixture was stirred under inert atmosphere
 5 for 18 h at room temperature, then was diluted with water (10 mL) and the aqueous phase was extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent EtOAc) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoyl)-4-methylpiperidin-4-ylcarbamate
 10 as off-white powder. MS (ESI) *m/z* 455 [C₂₆H₂₃FN₆O + H]⁺.

Step 2: A solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoyl)-4-methylpiperidin-4-ylcarbamate in DCM (3 mL) and TFA (3 mL) was stirred 18 h at room temperature. The reaction mixture was concentrated under reduced
 15 pressure and the residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-fluorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (42 mg, 37% over 2 steps, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.20 (d, *J* = 1.4 Hz, 1H), 8.99 (d, *J* = 1.4 Hz, 1H), 8.11 (s, 1H), 8.02–7.94 (m, 6H), 7.50 (t, *J* = 5.0 Hz, 1H), 4.41 (d, *J* = 14.1 Hz,
 20 1H), 3.60 (d, *J* = 14.5 Hz, 1H), 3.49–3.34 (m, 2H), 2.05–1.76 (m, 4H), 1.50 (s, 3H); ¹³C NMR (400 MHz, CD₃OD) δ 165.53, 159.80, 157.35, 142.70, 140.99, 140.55, 140.47, 138.36, 138.33, 135.50, 133.03, 132.13, 129.03, 128.40, 126.75, 123.36, 123.18, 122.60, 117.94, 114.54, 113.81, 113.57, 11.16, 52.22, 42.86, 37.46, 35.17, 34.42, 20.74; MS (ESI) *m/z* 455 [C₂₆H₂₃FN₆O + H]⁺.

25

Example 31: 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-chlorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




Step 1: 4-borono-2-Chlorobenzoic acid (110 mg, 0.55 mmol), K₂CO₃ (138 mg, 1.00 mmol)
 30 and Pd(dppf)Cl₂ (73.1 mg, 0.01 mmol) were added sequentially to a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (150 mg, 0.50 mmol) in DMF under inert

atmosphere. The reaction mixture was stirred at 80 °C for 18 h then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-chlorobenzoic acid (127 mg, 68 %) as brown solid. MS (ESI) *m/z* 375 [C₂₀H₁₁ClN₄O₂ + H]⁺.

Step 2: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-chlorobenzoic acid (53 mg, 0.14 mmol) in DMF (3 mL) under inert atmosphere, was added HATU (80 mg, 0.21 mmol), *N*-methyl morpholine (57 mg, 0.56 mmol) and *tert*-butyl piperidin-4-ylcarbamate hydrochloride (56 mg, 0.28 mmol). The reaction mixture was stirred at room temperature for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent EtOAc) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-chlorobenzoyl)piperidin-4-ylcarbamate (70 mg, 89%) as off-white powder. ¹H NMR (400 MHz, CDCl₃) δ 9.26 (s, 1H), 8.61 (s, 1H), 8.23 (d, *J* = 2.7 Hz, 1H), 8.07–7.96 (m, 2H), 7.93–7.81 (m, 4H), 7.35 (m, *J* = 9.8 Hz, 1H), 4.64 (s, 2H), 3.72 (s, 1H), 3.44 (t, *J* = 12.5 Hz, 1H), 3.18 (t, *J* = 11.5 Hz, 1H), 3.00–2.94 (m, 3H), 2.08 (d, *J* = 10.6 Hz, 1H), 1.93 (t, *J* = 10.3 Hz, 1H), 1.45 (s, 9H); MS (ESI) *m/z* 457 [C₂₃H₂₁ClN₆O + H]⁺.


Step 3: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-fluorobenzoyl)piperidin-4-ylcarbamate (70 mg, 0.13 mmol) in DCM (5 mL) were added TFA (5 mL) and the resulting mixture stirred 18 h at room temperature. The reaction mixture was concentrated under reduced pressure and purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)-3-chlorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (15 mg, 26%, AUC HPLC 97%) as white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.20 (d, *J* = 1.4 Hz, 1H), 8.93 (d, *J* = 1.2 Hz, 1H), 8.23 (d, *J* = 2.7 Hz, 1H), 8.04–8.15 (m, 2H), 7.98 (m, *J* = 7.6 Hz, 4H), 7.45 (m, *J* = 9.7 Hz, 1H), 4.73 (t, *J* = 14.9 Hz, 1H), 3.53 (t, *J* = 12.5 Hz, 1H), 3.11–3.27 (m, 2H), 2.98 (m, *J* = 6.5 Hz, 1H), 2.09 (d, *J* = 14.2 Hz, 1H), 1.93 (t, *J* = 13.0 Hz, 1H), 1.70–1.45 (m, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 167.27, 167.15, 142.72, 140.99, 139.16, 139.04, 138.33, 135.51, 135.34, 135.21, 133.03, 132.15, 130.59, 130.50, 128.42, 127.91, 127.49, 127.44, 126.75, 125.32, 125.27, 114.50, 112.14, 45.44, 44.87, 39.84, 39.03, 31.72, 31.13, 30.85; MS (ESI) *m/z* 457 [C₂₃H₂₁ClN₆O + H]⁺.

Example 32: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-chlorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



- 5 Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-chlorobenzoic acid (75 mg, 0.20 mmol) in DMF (2 mL), were successively added HATU (114 mg, 0.30 mmol), *N*-methyl morpholine (61 mg, 0.40 mmol) and *tert*-butyl 4-methylpiperidin-4-ylcarbamate hydrochloride (86 mg, 0.40 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and the aqueous layer was extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent EtOAc) to afford *tert*-butyl 1-(2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate as an off-white powder. MS (ESI) *m/z* 471 [C₂₆H₂₃ClN₆O + H]⁺.
- 15 Step 2: A solution of *tert*-butyl 1-(2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate in DCM (3 mL) and TFA (3 mL) was stirred at room temperature for 18h. The reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)-3-chlorophenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (11 mg, 13% over 2 steps, AUC HPLC 99%), white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.20 (d, *J* = 1.3 Hz, 1H), 8.99 (d, *J* = 1.3 Hz, 1H), 8.24 (s, 1H), 8.21–7.88 (m, 2H), 7.98 (m, *J* = 5.7 Hz, 4H), 7.46 (d, *J* = 8.0 Hz, 1H), 4.32–4.00 (m, 1H), 3.71–3.51 (m, 1H), 3.50–3.30 (m, 2H), 1.93–1.51 (m, 4H), 1.37 (d, *J* = 4.0 Hz, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 170.32, 142.54, 140.94, 139.92, 137.19, 136.04, 135.36, 134.88, 132.27, 128.43, 126.51, 124.11, 117.95, 113.77, 112.07, 43.01, 37.45, 36.95, 36.3, 17.72; MS (ESI) *m/z* 471 [C₂₆H₂₃ClN₆O + H]⁺.
- 25

Example 33: 4-(6-(3-methoxy-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



Step 1: To a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (2.00 g, 6.91 mmol) in 1,4-dioxane, were added 3-methoxy-4-(methoxycarbonyl)phenylboronic acid (1.76 g, 8.7 mmol), an aqueous solution of Na_2CO_3 (1.46 g, 13.8 mmol) in water and $\text{Pd}(\text{PPh}_3)_4$ (498 mg, 0.43 mmol). The reaction mixture heated for 3 h at 90 °C under argon atmosphere and was filtered through celite. The filtrate was concentrated and the residue was purified by flash column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 98:2) to afford methyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methoxybenzoate (1.20 g, 44%) as yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.87 (s, 1H), 9.22 (s, 1H), 8.89 (s, 1H), 7.95 (s, 1H), 7.80 (s, 1H), 7.73 (s, 1H), 7.61 (d, $J=8.0$ Hz, 2H), 6.98 (d, $J=8.0$ Hz, 2H), 3.91 (s, 3H), 3.79 (s, 3H); MS (ESI), m/z 385 [$\text{C}_{22}\text{H}_{16}\text{N}_4\text{O}_3 + \text{H}$] $^+$.


Step 2: To a solution of methyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methoxybenzoate (1.20 g, 3.20 mmol) in $\text{THF}/\text{CH}_3\text{OH}$ (40/10 mL) was added a solution of LiOH (440 mg, 9.7 mmol) in water (10 mL). The reaction mixture was stirred at room temperature for 5 h and was concentrated *in vacuo*. The residue was first diluted with water (10 mL) and then was acidified till pH 3 using an aqueous solution of 2N HCl. The yellow precipitated was isolated by filtration and washed with water then was dried under vacuum to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methoxybenzoic acid (800 mg, 80%) as light yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.91 (s, 1H), 9.23 (s, 1H), 8.89 (s, 1H), 7.97 (s, 1H), 7.73 (d, $J=8.0$ Hz, 2H), 7.61 (d, $J=8.0$ Hz, 2H), 6.98 (d, $J=8.0$ Hz, 2H), 3.91 (s, 3H); MS (ESI), m/z 371 [$\text{C}_{21}\text{H}_{14}\text{N}_4\text{O}_3 + \text{H}$] $^+$.

Step 3: To a stirred solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-2-methoxybenzoic acid (300 mg, 0.81 mmol) in DMF (5 mL) were added *N*-methylmorpholine (250 mg, 2.43 mmol), HATU (465 mg, 1.22 mmol) and *tert*-butyl piperazine-1-carboxylate

(230 mg, 1.22 mmol). The reaction mixture was stirred for 10 h at room temperature, and was diluted with water followed by an extraction with CH₂Cl₂. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated. The residue was purified by flash column chromatography (silica gel, eluent CH₂Cl₂/CH₃OH 97:3) to afford *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)-2-methoxybenzoyl)piperazine-1-carboxylate (100 mg, 30%) as off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.29 (s, 1H), 9.02 (s, 1H), 8.17 (s, 1H), 8.08 (d, *J* = 8.4 Hz, 2H), 7.94 (d, *J* = 8.4 Hz, 2H), 7.75 (s, 1H), 7.69 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 7.6 Hz, 1H), 3.89 (s, 3H), 3.62–3.56 (m, 2H), 3.46–3.41 (m, 2H), 3.37–3.29 (m, 4H), 1.39 (s, 9H); MS (ESI) *m/z* 539 [C₃₀H₃₀N₆O₄ + H]⁺.

Step 4: A solution of *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)-2-methoxybenzoyl)piperazine-1-carboxylate (50 mg, 0.09 mmol) in a mixture of 1,4-dioxane and 30% HCl solution in 1,4-dioxane (1 mL) was stirred at room temperature for 3 h and was concentrated under reduced pressure. The residue was purified by flash column chromatography (silica gel, eluent 10% CH₂Cl₂/CH₃OH 90:10) to afford 4-(6-(3-methoxy-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (25 mg, 30%, AUC HPLC 88%) as a light yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.21 (bs, 1H), 8.97 (s, 1H), 8.12 (bs, 1H), 8.02–7.96 (m, 4H), 7.81 (s, 1H), 7.69 (d, *J* = 7.6 Hz, 1H), 7.39 (d, *J* = 8.0 Hz, 1H), 3.99 (s, 3H), 3.89–3.84 (m, 2H), 3.52–3.43 (m, 2H), 3.14 (bs, 2H), 3.03 (bs, 2H); MS (ESI) *m/z* 439 [C₂₅H₂₂N₆O₂ + H]⁺.

Example 34: 4-(6-(2-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrile




Step 1: To a solution of 4-(6-bromoimidazo[1,2-*a*]pyrazin-3-yl)benzonitrile (50 mg, 0.167 mmol) in a mixture of DMF (2 mL) and water (1.3 mL), was added Na₂CO₃ (35 mg, 0.334 mmol), 4-borono-3-fluorobenzoic acid (33 mg, 0.183 mmol) and Pd(PPh₃)₄ (19 mg, 0.016 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 0.5 h and was filtered through celite. The filtrate was concentrated under reduced pressure and the residue was purified by column chromatography to afford 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyrazin-6-yl)-3-fluorobenzoic acid (25.0 mg, 42.3%, AUC HPLC 99%) as a off white solid.

^1H NMR (600 MHz, DMSO- d_6) δ 9.33 (s, 1H), 8.97 (s, 1H), 8.26 (s, 1H), 8.10-8.01 (m, 5H), 7.84 (d, J = 8.4 Hz, 2H), 7.72 (d, J = 12.6 Hz, 2H); MS (ESI) m/z 359 [$\text{C}_{20}\text{H}_{11}\text{FN}_4\text{O}_2 + \text{H}$] $^+$.

Step 2: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)-3-fluorobenzoic acid (30 mg, 0.083 mmol) in a mixture of CH_2Cl_2 (2 mL) and DMF (1, 3 mL) was added sequentially DIPEA (60 μL , 0.351 mmol), HOBT (23 mg, 0.175 mmol), EDCI $\cdot\text{HCl}$ (33 mg, 0.175 mmol) and *N*-methyl piperazine (14 μL , 0.125 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, and then solvents were removed under vacuum. The crude residue was purified by column chromatography to afford 4-(6-(2-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (16 mg, 45%, AUC HPLC 99%) as a white solid. ^1H NMR (600 MHz, DMSO- d_6) δ 9.36 (s, 1H), 9.01 (s, 1H), 8.30 (s, 1H), 8.14 (t, J = 7.8 Hz, 1H), 8.15-8.03 (m, 4H), 7.44 (d, J = 11.4 Hz, 1H), 7.39 (d, J = 9.6 Hz, 1H), 3.64-3.62 (m, 2H), 2.40-2.26 (m, 4H), 2.20 (s, 3H); MS (ESI) m/z 441 [$\text{C}_{25}\text{H}_{21}\text{FN}_6\text{O} + \text{H}$] $^+$.


Example 35: 3-fluoro-4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



To a solution of 4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl(4-methylpiperazin-1-yl)methanone (200 mg, 0.45 mmol) in a mixture of toluene (4 mL) and ethanol (2 mL) under inert atmosphere, was added K_2CO_3 (124 mg, 0.89 mmol), 4-cyano-2-fluorophenylboronic acid (88 mg, 0.54 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (52 mg, 0.045 mmol). The resulting mixture was heated in a microwave reactor at 140 $^\circ\text{C}$ for 15 min, and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) to afford 3-fluoro-4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (18.1 mg, 9%, AUC HPLC 96%) as a yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 9.21 (s, 1H), 8.74 (d, J = 1.5 Hz, 1H), 8.12 (d, J = 8.2 Hz, 2H), 8.10 (s, 1H), 8.00-7.96 (m, 1H), 7.89-7.87 (m, 1H), 7.82 (d, J = 8.0 Hz, 1H), 7.54 (d, J = 8.2 Hz, 2H), 3.81 (bs, 2H), 3.53 (bs, 2H), 2.55-2.52 (m, 4H), 2.36 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.97, 160.63, 143.81, 142.39, 140.94, 139.19, 138.00, 136.99,

133.17, 130.43, 128.75, 127.91, 122.44, 122.00, 121.61, 118.25, 116.42, 115.57, 55.98, 55.58, 46.01; MS (ESI) m/z 441 $[\text{C}_{25}\text{H}_{21}\text{FN}_6\text{O} + \text{H}]^+$.

Example 36: 4-(6-(6-(4-methylpiperazine-1-carbonyl)pyridin-3-yl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



Step: 1 To a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (140 mg, 0.468 mmol) and methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)picolinate (246 mg, 0.936 mmol) in a mixture of DMF/H₂O (6:1, 3.5 mL) was added Na₂CO₃ (148 mg, 1.40 mmol).


- 10 The suspension was stirred vigorously whilst de-gassing with N₂ for 5 min before adding Pd(PPh₃)₄ (54 mg, 0.046 mmol). The mixture was heated to 100°C for 3 h and then, concentrated under reduced pressure. The crude residue was purified by column chromatography to afford methyl 5-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)picolinate (120 mg, 72%) as an off white solid. ¹H NMR (600 MHz, DMSO-*d*₆) δ 9.42 (s, 1H), 9.35 (s, 1H), 9.25 (s, 1H), 8.66 (d, J = 10.8 Hz, 1H), 8.28 (s, 1H), 8.18 (d, J = 8.4 Hz, 1H), 8.12-8.05 (m, 4H), 3.92 (s, 3H); MS (ESI) m/z 356 $[\text{C}_{20}\text{H}_{13}\text{N}_5\text{O}_2 + \text{H}]^+$.

- Step: 2 A solution of methyl 5-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)picolinate (60 mg, 0.169 mmol) and a 5N sodium hydroxide solution (135 μ L, 0.676 mmol) in methanol (3 ml) was stirred at room temperature for 1 hr. 5N hydrochloric acid (319 μ L) was added to the reaction solution at room temperature, followed by extraction with ethyl acetate. The extract was dried over Na₂SO₄, concentrated under reduced pressure. To a solution of the crude acid (50 mg, 0.146 mmol) in DMF (1.5 mL) was added sequentially *N*-methyl piperazine (41 μ L, 0.366 mmol), HBTU (139 mg, 0.366 mmol), HOBT (38 mg, 0.278 mmol) and DIPEA (127 μ L, 0.733 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 12 h, then it was diluted with water (3 mL) and extracted with CH₂Cl₂ (3 $\times$ 10 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography to afford 4-(6-(6-(4-methylpiperazine-1-carbonyl)pyridin-3-yl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (35

mg, 56%, AUC HPLC 96%), light brown solid. ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 9.33 (s, 1H), 9.28 (s, 1H), 9.27 (s, 1H), 8.57 (d, $J = 10.2$ Hz, 1 H), 8.26 (s, 1H), 8.13-8.02 (m, 4H), 7.70 (d, $J = 8.4$ Hz, 1H), 3.71-3.64 (m, 2H), 3.50-3.42 (m, 2H), 2.45-2.37 (m, 2H), 2.34-2.26 (m, 2H), 2.20 (s, 3H); MS (ESI) m/z 424 [$\text{C}_{24}\text{H}_{21}\text{N}_7\text{O} + \text{H}$] $^+$.

5

Example 37: 4-(6-(5-(4-methylpiperazine-1-carbonyl)pyridin-2-yl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




Step 1: A mixture of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (50 mg, 0.167 mmol), ethyl 6-bromonicotinate (39 mg, 0.167 mmol), $\text{PdCl}_2(\text{PPh}_3)_2$ (12 mg, 0.016 mmol), and hexamethylditin (35 μL , 0.167 mmol) in anhydrous DMF (0.5 mL) was heated at 100°C for 3 h, and then cooled to room temperature and solvents were removed under vacuum. The crude residue was purified by column chromatography to afford methyl 6-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)nicotinate (29.6 mg, 48%) as an off white solid. ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 9.42 (s, 1H), 9.36 (s, 1H), 9.16 (s, 1H), 8.53-8.46 (m, 2H), 8.29 (s, 1H), 8.14 (d, $J = 8.4$ Hz, 2H), 8.05 (d, $J = 8.4$ Hz, 2H), 4.38 (q, $J = 7.2$ Hz, 2H), 1.36 (t, $J = 7.2$ Hz, 2H); MS (ESI) m/z 370 [$\text{C}_{21}\text{H}_{15}\text{N}_5\text{O}_2 + \text{H}$] $^+$.

Step 2: A solution of methyl 6-(3-(4-cyanophenyl)imidazo[1,2-a]pyrazin-6-yl)nicotinate (20 mg, 0.054 mmol) and a 5N sodium hydroxide solution (43 μL , 0.216 mmol) in ethanol (1 mL) was stirred at room temperature for 1 hr. 5N hydrochloric acid (101 μL) was added to the reaction solution at room temperature, followed by extraction with ethyl acetate. The extract was dried over Na_2SO_4 , filtered and concentrated under reduced pressure and directly used in the next step without further purification.

To a solution of above acid (15 mg, 0.043 mmol) in DMF (0.5 mL) was added sequentially *N*-methylpiperazine (12 μL , 0.109 mmol), HBTU (42 mg, 0.109 mmol), HOBt (12 mg, 0.083 mmol) and DIPEA (13 μL , 0.073 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 12 h, then was diluted with water (1 mL) and extracted with EtOAc (3 $\times$ 5 mL). The combined organic layer was dried over Na_2SO_4 and was

concentrated under reduced pressure. The crude residue was purified by column chromatography to afford 4-(6-(5-(4-methylpiperazine-1-carbonyl)pyridin-2-yl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (12 mg, 66%, AUC HPLC 93.6%) as a light brown solid. ¹H NMR (600 MHz, DMSO-*d*₆) δ 9.34 (d, *J* = 12.0 Hz, 2H), 8.68 (s, 1H), 8.40 (d, *J* = 7.8 Hz, 1H), 8.26 (s, 1H), 8.12 (d, *J* = 8.4 Hz, 2H), 8.05-7.98 (m, 3H), 3.68-3.63 (m, 2H), 2.40-2.35 (m, 4H), 2.20 (s, 3H); MS (ESI) *m/z* 424 [C₂₄H₂₁N₇O + H]⁺.

Example 38: 4-(6-(2-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



10

Step 1: To a solution of 3-methoxy-4-bromobenzoic acid (500 mg, 2.16 mmol) in DMF (5 mL) was added NMM (0.44 g, 4.32 mmol) followed by addition of HATU (1.23 g, 3.24 mmol) at rt and stirred for 30 min. 1-methylpiperazine (0.24 g, 2.38 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain 350 mg (83%) of (4-bromo-3-methoxyphenyl)(4-methylpiperazin-1-yl)methanone (AUC LC-MS 83%) as a brown solid.

15

Step 2: To a solution of (4-bromo-3-methoxyphenyl)(4-methylpiperazin-1-yl)methanone (350 mg), 1,4-dioxane (10 mL), were added bis(pinacolato)diboron (0.30 g, 1.19 mmol), KOAc (330 mg, 3.36 mmol), PdCl₂dppf (24 mg, 0.03 mmol) and dppf (18 mg, 0.033 mmol). The reaction mixture was heated at 90 °C for 16 h, was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 350 mg of (3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone as a brown liquid which was used in next step without further purification.

20

25


Step 3: To a mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (232 mg), (3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (350 mg), K₃PO₄ (412 mg, 1.94 mmol) in 1,4-dioxane (5 mL) and water (5

30

mL) was added Pd(PPh₃)₄ (56 mg, 0.048 mmol). The reaction mixture was heated at 90 °C for 16 h under argon atmosphere, was diluted with water and extracted with EtOAc. The organic phase was washed in turn with water, brine solution, dried over anhydrous Na₂SO₄ filtered and concentrated under reduced pressure. The residue was purified by column chromatography to afford 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl) phenyl)imidazo [1,2-a]pyridin-3-yl)benzonitrile (80 mg, 30%, AUC HPLC 99.3%) as an off-white solid; m.p. 311-318°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.27 (s, 1 H), 9.14 (s, 1 H), 8.19 (d, *J* = 8.0 Hz, 1 H), 7.99 (s, 1 H), 7.88 (d, *J* = 8.0 Hz, 2 H), 7.79 (d, *J* = 8.0 Hz, 2 H), 7.14-7.12 (m, 2 H), 3.95 (s, 3 H), 3.84 (bs, 2 H), 3.54 (bs, 2 H), 2.54 (s, 2 H), 2.36 (bs, 2 H), 2.34 (s, 3 H); MS (ESI) *m/z* 453.32 [C₂₆H₂₄N₆O₂ + H]⁺.

Step 4: A solution of 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (150 mg) in anhydrous DCM (5 mL) was treated with BBr₃ at rt for 3 h. After the completion of reaction as indicated by TLC, the reaction mixture was washed with NaHCO₃. The organic layer was dried over Na₂SO₄ filtered and concentrated. The residue was purified by preparative HPLC to afford 4-(6-(2-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (50 mg, 34%, AUC HPLC 97%) as an off white solid; m.p. 237-241°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 11.15 (s, 1 H), 9.43 (s, 1 H), 9.31 (s, 1 H), 8.26 (s, 1 H), 8.16 (d, *J* = 12.0 Hz, 1 H), 8.09-8.01 (m, 4 H), 6.95 (s, 2 H), 3.59 (bs, 2 H), 3.34 (bs, 2 H), 2.49 (bs, 2 H), 2.27 (bs, 2 H), 2.19 (s, 3 H); MS (ESI) *m/z* 439.2 [C₂₅H₂₂N₆O₂ + H]⁺.

Example 39: 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile




Step: 1 To a solution of 4-bromo-2-methoxy benzoic acid (1 g, 4.34 mmol) in DMF (15 mL) was added NMM (0.876 mL, 8.68 mmol) followed by HATU (2.08 g, 6.51 mmol) at rt and stirred for 30 min. 1-Methylpiperazine (0.478 mg, 4.78 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc, washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced

pressure. The residue was purified by column chromatography (silica gel, Petroleum ether/EtOAc 100:0 to 50:50) to afford (4-bromo-2-methoxyphenyl)(4-methylpiperazin-1-yl)methanone (1.1 g, 81%) of as an off white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.33 (s, 1H), 7.25-7.19 (m, 2H), 3.84 (s, 3H), 3.6-2.9 (m, 8H), 2.83 (s, 3 H); MS (ESI) *m/z* 315.10 [C₁₃H₁₇BrN₂O₂+2]⁺.

Step 2: A mixture of (4-bromo-2-methoxyphenyl)(4-methylpiperazin-1-yl)methanone (1.1 g, 3.52 mmol), Bis(pinacolato)diboron (1.07 g, 4.23 mmol), KOAc (1.03 g, 10.56 mmol) in 1,4-dioxane (20 mL) was degassed with argon for 30 min. PdCl₂dppf (75 mg, 0.1 mmol), dppf (56 mg, 0.1 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 800 mg of (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone as a dark brown liquid which was used in next step without purification.


Step 3: To a mixture of 2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (678 mg, 1.88 mmol), 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (400 mg, 1.34 mmol), K₃PO₄ (568.1 mg, 2.68 mmol) in 1,4-dioxane (20 mL) and water (5 mL) was added Pd(PPh₃)₄ (77 mg, 0.05 mmol). The reaction mixture was heated at 90 °C for 1 h and was diluted with water and extracted with EtOAc. The organic phase was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by Preparative HPLC to afford 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (180 mg, 30%, AUC HPLC 96.7%) as a pale brown solid; m.p. 188-192°C. ¹H NMR (400 MHz, CD₃OD) δ (ppm): 9.19 (s, 1 H), 8.94 (s, 1 H), 8.89 (s, 1 H), 8.00-7.94 (m, 4 H), 7.78 (s, 1 H), 7.64 (d, *J* = 8.0 Hz, 1 H), 7.32 (d, *J* = 8.0 Hz, 1 H), 3.95 (s, 3 H), 3.80 (bs, 2 H), 3.34 (bs, 2 H), 2.54 (s, 2 H), 2.45 (bs, 1 H), 2.38 (bs, 1 H), 2.33 (s, 3 H); MS (ESI) *m/z* 453.30 [C₂₆H₂₄N₆O₂ + H]⁺.

Example 40: 4-(6-(4-(4-methylpiperazin-1-yl)sulfonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile



1-methyl-4-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)sulfonyl)piperazine (240 mg, 0.656 mmol), K_3PO_4 (212 mg, 1 mmol) and $Pd(PPh_3)_4$ (28 mg, 0.025 mmol) were added sequentially to a solution of 4-(6-bromoimidazo[1,2-a]pyrazin-3-yl)benzonitrile (150 mg, 0.5 mmol) in a mixture of 1,4-dioxane/ H_2O (10:1 mL) at room temperature under argon atmosphere. The reaction mixture was refluxed for 6 h and was diluted with water (10 mL) and extracted with EtOAc (3×50 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $CHCl_3/MeOH$ 95:5) to afford 4-(6-(4-(4-methylpiperazin-1-yl)sulfonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzonitrile (80 mg, 36%, AUC HPLC 99.1%) as an off-white solid; m.p. 220-228°C. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 9.30 (s, 1 H), 8.61 (s, 1 H), 8.08 (d, J = 8.8 Hz, 2 H), 8.00 (s, 1 H), 7.92-7.85 (m, 4 H), 7.77 (d, J = 8.4 Hz, 2 H), 3.08 (bs, 4 H), 2.50-2.48 (m, 4 H), 2.27 (s, 3 H); MS (ESI) m/z 459.27 [$C_{24}H_{22}N_6O_2S + H$] $^+$.


Example 41: (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



To a solution of (4-(3-iodoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (150 mg, 0.34 mmol) in a mixture of toluene (4 mL) and ethanol (2 mL) under inert atmosphere, was added K_2CO_3 (93 mg, 0.67 mmol), 4-chlorophenylboronic acid (79 mg, 0.50 mmol) and $Pd(PPh_3)_4$ (39 mg, 0.034 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 15 min, and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5 to 90:10) to afford (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-


methylpiperazin-1-yl)methanone (128.4 mg, 66%, AUC HPLC 99%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 9.24 (d, $J = 1.4$ Hz, 1H), 8.54 (d, $J = 1.4$ Hz, 1H), 7.95 (d, $J = 8.4$ Hz, 2H), 7.89 (s, 1H), 7.60-7.55 (m, 4H), 7.52 (d, $J = 8.4$ Hz, 2H), 3.82 (bs, 2H), 3.49 (s, 2H), 2.50-2.33 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.78, 143.84, 140.70, 139.63, 137.76, 136.14, 135.34, 135.33, 129.98, 129.36, 127.84, 126.45, 126.41, 126.24, 112.54, 55.27, 54.79, 46.03; MS (ESI) m/z 432 [$\text{C}_{24}\text{H}_{22}\text{ClN}_5\text{O} + \text{H}$] $^+$.

Example 42: 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzamide



To a solution of 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyrazine (200 mg, 0.65 mmol) in a mixture of toluene (3 mL) and EtOH (1.5 mL) under inert atmosphere, was consecutively added K_2CO_3 (180 mg, 1.30 mmol), 4-carbamoylphenylboronic acid (117 mg, 0.71 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (75 mg, 0.06 mmol). The resulting mixture was heated in a microwave reactor at 140°C for 30 min and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent hexane/EtOAc 3:2) to afford 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzamide (31 mg, 83%, AUC HPLC 97%) as white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.26 (s, 1H), 8.97 (s, 1H), 8.16 (d, $J = 8.2$ Hz, 2H), 8.13 – 8.02 (m, 2H), 7.99 (d, $J = 8.2$ Hz, 2H), 7.87 (d, $J = 8.2$ Hz, 2H), 7.66 (d, $J = 8.3$ Hz, 2H), 7.42 (m, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 167.43, 142.93, 140.29, 138.83, 187.92, 135.55, 133.99, 133.30, 129.74, 129.38, 127.87, 126.43, 126.21, 126.01, 114.30; MS (ESI) m/z 390 [$\text{C}_{19}\text{H}_{13}\text{ClN}_4\text{O} + \text{H}$] $^+$.

Example 43: 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-one




Step 1: To a solution of 6-bromo-3-iodoimidazo[1,2-a]pyrazine (3.00 g, 9.26 mmol) in DMF (20 mL), was added Na_2CO_3 (2.46 g, 23.2 mmol), 4-chlorophenylboronic acid (1.60 g, 10.2 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (214 mg, 0.18 mmol). The resulting mixture was stirred at 80°C for 18

h under inert atmosphere and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent Hex/EtOAc 3:2) to afford 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyrazine (930 mg, 32%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 9.03 (d, *J* = 1.2 Hz, 1H), 8.36 (d, *J* = 1.2 Hz, 1H), 7.91 (s, 1H), 7.54 (q, *J* = 12.3 Hz, 4H); MS (ESI) *m/z* 308 [C₁₃H₇BrN₄ + H]⁺.

Step 2: To a solution of 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyrazine (2.78 g, 9.03 mmol) in a mixture of DMF (24 mL) and water (4.8 mL) under inert atmosphere, was added K₂CO₃ (2.50 g, 18.1 mmol), 4-boronobenzoic acid (1.65 mg, 9.94 mmol) and PdCl₂dppf (1.32 g, 1.81 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 30 min then, was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 9:1) to afford 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid 5 (326 mg, 11%) as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.02 (bs, 1H), 9.28 (d, *J* = 1.3 Hz, 1H), 9.00 (d, *J* = 1.4 Hz, 1H), 8.22 (d, *J* = 8.5 Hz, 2H), 8.11 (s, 1H), 8.04 (d, *J* = 8.5 Hz, 2H), 7.88 (d, *J* = 8.5 Hz, 2H), 7.67 (d, *J* = 8.5 Hz, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.05, 142.99, 140.33, 140.30, 137.64, 135.62, 133.35, 130.56, 129.81, 129.66, 129.40, 126.42, 126.31, 114.72; MS (ESI) *m/z* 350 [C₁₉H₁₂ClN₃O₂ + H]⁺.


Step 3: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (45 mg, 0.13 mmol) in DMF (5 mL), were sequentially added HATU (73 mg, 0.19 mmol), *N*-methyl morpholine (52 mg, 0.51 mmol) and piperidin-4-one (45 mg, 0.40 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-one (41 mg, 75%, AUC HPLC 99%) as brown solid. ¹H NMR (400 MHz, CD₃OD) δ 9.11 (s, 1H), 8.74 – 8.79 (m, 1H), 8.00 – 8.08 (m, 2H), 7.96 (s, 1H), 7.71 (d, *J* = 8.5 Hz, 2H), 7.59 (d, *J* = 8.5 Hz, 2H), 7.48 (d, *J* = 8.4 Hz, 2H), 4.12–3.61 (m, 3H), 3.45 (bs, 1H), 2.70–2.36 (m, 1H), 1.91–1.62 (m, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 208.88, 172.48, 171.94, 164.41, 143.52, 143.47, 141.05, 140.96, 139.31, 138.96, 137.41, 137.35, 136.89, 136.40, 135.00, 134.92, 130.90, 128.70, 128.53, 127.89, 127.84, 127.22, 115.16, 115.09, 99.62, 96.49, 46.52, 40.93, 36.99, 36.14; MS (ESI) *m/z* 431 [C₂₄H₁₉ClN₄O₂ + H]⁺.

Example 44: (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-hydroxypiperidin-1-yl)methanone



- 5 To a solution of 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperidin-4-one (22 mg, 0.05 mmol) in MeOH (3 mL) was added NaBH₄ (5 mg, 0.13 mmol) and the reaction mixture was stirred at room temperature for 18 h. The reaction was quenched by adding water (10 mL) dropwise, and the aqueous solution was extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄ and were concentrated under reduced
- 10 pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-hydroxypiperidin-1-yl)methanone (10 mg, 45%, AUC HPLC >99%) as a white solid. ¹H
- 15 NMR (400 MHz, CD₃OD) δ 9.15 (d, *J* = 1.3 Hz, 1H), 8.82 (d, *J* = 1.3 Hz, 1H), 8.08 (d, *J* = 8.4 Hz, 2H), 7.98 (s, 1H), 7.75 (d, *J* = 8.6 Hz, 2H), 7.63 (d, *J* = 8.6 Hz, 2H), 7.51 (d, *J* = 8.4 Hz, 2H), 4.19 (bs, 1H), 3.91 (m, *J* = 4.0 Hz, 1H), 3.68 (bs, 1H), 3.46–3.19 (m, 2H), 2.04–1.74 (m, 2H), 1.67–1.39 (m, 2H); ¹³C NMR (400 MHz, CD₃OD) δ 172.00, 143.69, 141.66, 140.95, 139.15, 137.42, 136.35, 135.43, 130.89, 128.56, 128.46, 127.44, 115.12, 67.64, 46.48, 40.87, 35.45, 34.75; MS (ESI) *m/z* 433 [C₂₄H₂₁ClN₄O₂ + H]⁺.

- 20 Example 45: (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-ethylpiperazin-1-yl)methanone




- To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (50 mg, 0.14 mmol) in DCM (5 mL) at 0 °C, was added oxalyl chloride (36 mg, 0.29 mmol) and the mixture thus obtained was stirred to room temperature for 3h. The reaction mixture was
- 25 concentrated under reduced vacuum and to the residue, was added DMF (5 mL), 1-ethylpiperazine (92 mg, 0.23 mmol) and a solution of KOH (0.10 ml, 13 M in THF). The

resulting mixture was stirred at room temperature for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford 4-(3-(4-

5 chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(4-ethylpiperazin-1-yl)methanone (28 mg, 44%, AUC HPLC 99%) as brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.08 (d, *J* = 1.0 Hz, 1H), 8.76 (d, *J* = 1.1 Hz, 1H), 8.05 (d, *J* = 8.3 Hz, 2H), 7.93 (s, 1H), 7.71 (d, *J* = 8.5 Hz, 2H), 7.60 (d, *J* = 8.5 Hz, 2H), 7.52 (d, *J* = 8.3 Hz, 2H), 3.80 (bs, 4H), 3.10–2.79 (m, 6H), 1.25 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 171.92, 143.84, 141.82, 140.56, 139.58, 136.24, 136.22, 135.84, 130.88, 130.81, 128.89, 128.46, 127.86, 127.49, 115.11, 53.17, 52.79, 10.67; MS (ESI) *m/z* 446 [C₂₅H₂₄ClN₅O + H]⁺.

Example 46: (4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)phenyl)(piperazin-1-yl)methanone 2,2,2-trifluoroacetate salt




15

Step 1: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoic acid (50 mg, 0.14 mmol) in DMF (5 mL) were added HATU (82 mg, 0.22 mmol), *N*-methyl morpholine (29 mg, 0.29 mmol) and *tert*-butyl piperazine-1-carboxylate (54 mg, 0.29 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/HCOOH 0.01%) to afford *tert*-butyl 4-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyrazin-6-yl)benzoyl)piperazine-1-carboxylate (23 mg, 31%, AUC HPLC 100%) as a brown solid. ¹H NMR (400 MHz, CDCl₃) δ 9.33 (s, 1H), 8.54 (s, 1H), 7.96 (d, *J* = 8.0 Hz, 2H), 7.91 (s, 1H), 7.57 (d, *J* = 7.5 Hz, 4H), 7.51 (d, *J* = 8.0 Hz, 2H), 3.84–3.60 (bs, 2H), 3.60–3.28 (bs, 6H), 1.46 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 169.97, 162.65, 154.55, 143.07, 140.40, 139.28, 137.46, 136.05, 135.90, 133.08, 129.61, 127.89, 126.69, 125.36, 112.64, 80.47, 28.36; MS (ESI) *m/z* 518 [C₂₈H₂₈ClN₅O₃ + H]⁺.

30

Step 2: A solution of *tert*-butyl 4-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperazine-1-carboxylate (66 mg, 0.65 mmol) in a mixture of DCM (5 mL) and TFA (5 mL) was stirred at room temperature for 18 h, then was concentrated under reduced pressure to afford (4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)(piperazin-1-yl)methanone as a TFA salt (24 mg, 69 %, AUC HPLC 98%) as a brown solid. ¹H NMR (400 MHz, CDCl₃) δ 9.29 (d, *J* = 1.3 Hz, 1H), 8.96 (d, *J* = 1.3 Hz, 1H), 8.18 (d, *J* = 8.4 Hz, 2H), 8.12 (s, 1H), 7.88 (d, *J* = 8.5 Hz, 2H), 7.67 (d, *J* = 8.5 Hz, 2H), 7.58 (d, *J* = 8.4 Hz, 2H), 3.68 (s, 5H), 3.20 (s, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 170.01, 142.99, 140.34, 138.97, 138.12, 136.21, 134.45, 132.67, 130.23, 129.65, 127.99, 126.99, 126.88, 125.02, 112.75, 43.42, 29.69; MS (ESI) *m/z* 418 [C₂₃H₂₀ClN₅O + H]⁺.

Example 47: (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)methanone 2,2,2-trifluoroacetate salt




Step 1: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoic acid (70 mg, 0.20 mmol) in DMF (5 mL), was successively added HATU (114 mg, 0.30 mmol), *N*-methyl morpholine (41 mg, 0.40 mmol) and *tert*-butyl piperidin-4-ylcarbamate (80 mg, 0.40 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by preparative (C18, eluent ACN/H₂O/0.01% HCOOH acid) to afford *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-ylcarbamate (47 mg, 44 %, AUC HPLC 98%) as a brown solid. ¹H NMR (400 MHz, CDCl₃) δ 9.30 (s, 1H), 8.53 (s, 1H), 7.92 (t, *J* = 7.3 Hz, 3H), 7.56 (d, *J* = 1.2 Hz, 4H), 7.47 (d, *J* = 8.1 Hz, 2H), 4.58 (d, *J* = 7.1 Hz, 2H), 3.69 (bs, 2H), 3.21–2.83 (m, 2H), 1.97 (bs, 2H), 1.43 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 169.72, 162.92, 155.09, 143.03, 140.39, 139.36, 137.27, 136.51, 135.80, 133.20, 130.09, 129.59, 127.64, 126.66, 126.60, 125.44, 112.64, 79.67, 47.92, 46.55, 41.22, 33.10, 32.15, 28.39; MS (ESI) *m/z* 532 [C₂₉H₃₀ClN₅O₃ + H]⁺.

Step 2: A solution of *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-ylcarbamate (40 mg, 0.08 mmol) in a mixture of DCM (5 mL) and TFA (5 mL) was stirred at room temperature for 18 h, then was concentrated under reduced pressure to afford (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)methanone as a TFA salt (54 mg, >99%, AUC HPLC 96%) as a brown solid. ¹H NMR (400 MHz, CD₃OD) δ 9.22 (s, 1H), 8.83 (s, 1H), 8.07 (d, *J* = 7.8 Hz, 3H), 7.73 (d, *J* = 8.2 Hz, 2H), 7.62 (d, *J* = 8.2 Hz, 2H), 7.51 (d, *J* = 8.0 Hz, 2H), 4.71 (bs, 1H), 3.85 (bs, 1H), 3.45 (m, *J* = 3.4 Hz, 1H), 3.30–2.85 (m, 2H), 2.28–1.91 (m, 2H), 1.62 (bs, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 172.04, 143.03, 141.45, 140.50, 139.00, 137.041, 136.68, 133.66, 131.50, 131.04, 130.94, 130.58, 128.76, 128.60, 128.02, 126.78, 117.47, 115.32, 114.65, 47.09, 41.61, 32.74, 31.48, 30.76; MS (ESI) *m/z* 432 [C₂₃H₂₀ClN₅O + H]⁺.


Example 48: *N*-(1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl)-

2,2,2-trifluoroacetamide




To a solution of (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)methanone (16 mg, 0.04 mmol) in DMF (3 mL) were added pyridine (0.05 mL) and 2,2,2-trifluoroacetic anhydride (0.3 mL). The reaction mixture was stirred at room temperature for 18 h and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford *N*-(1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl)-2,2,2-trifluoroacetamide (13 mg, 63 %, AUC HPLC 99%) as yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 9.12 (d, *J* = 1.3 Hz, 1H), 8.80 (d, *J* = 1.3 Hz, 1H), 8.08 (d, *J* = 8.4 Hz, 2H), 7.96 (s, 1H), 7.74 (d, *J* = 8.6 Hz, 2H), 7.61 (d, *J* = 8.6 Hz, 2H), 7.51 (d, *J* = 8.4 Hz, 2H), 4.65 (bs, 1H), 4.05 (m, *J* = 5.7 Hz, 1H), 3.80 (bs, 1H), 3.25 (bs, 1H), 3.00 (bs, 1H), 2.00 (bs, 1H), 1.88 (bs, 1H), 1.58 (bs, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 170.63, 157.09, 156.35, 142.41, 140.45, 139.30, 137.92, 135.67, 134.86, 134.86, 134.38, 129.44, 127.09, 126.46, 126.11, 120.32, 117.47, 114.62, 113.69, 111.77, 46.44, 40.91, 31.11, 30.32; MS (ESI) *m/z* 528 [C₂₆H₂₁ClF₃N₅O + H]⁺.

Example 49: *N*-(1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl)-2,2,2-acetamide



To a solution of (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)methanone (16 mg, 0.04 mmol) in DMF (3 mL), were successively added pyridine (0.05 mL) and acetic anhydride (0.3 mL, 3.18 mmol). The reaction mixture was stirred at room temperature for 18 h and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford *N*-(1-(4-(3-(4-chlorophenyl)imidazo[1,2-*a*]pyrazin-6-yl)benzoyl)piperidin-4-yl)-2,2,2-acetamide (11 mg, 61 %, AUC HPLC >99%) as white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.13 (d, *J* = 1.3 Hz, 1H), 8.81 (d, *J* = 1.3 Hz, 1H), 8.07 (d, *J* = 8.4 Hz, 2H), 7.96 (s, 1H), 7.74 (d, *J* = 8.6 Hz, 2H), 7.62 (d, *J* = 8.6 Hz, 2H), 7.51 (d, *J* = 8.4 Hz, 2H), 4.55 (bs, 1H), 3.95 (d, *J* = 5.3 Hz, 1H), 3.74 (bs, 1H), 3.24 (bs, 1H), 3.07 (bs, 1H), 2.03–1.81 (m, 5H), 1.32–1.60 (m, 2H); ¹³C NMR (400 MHz, CD₃OD) δ 172.61, 172.03, 143.83, 141.86, 140.75, 139.39, 137.21, 136.27, 135.79, 130.87, 128.49, 127.87, 127.54, 115.10, 42.30, 42.30, 33.22, 32.31, 22.66; MS (ESI) *m/z* 474 [C₂₆H₂₄ClN₅O₂ + H]⁺.

Example 50: (4-(3-(4-(2H-tetrazol-5-yl)phenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




To a solution of 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzonitrilemethanone (100 mg, 0.16 mmol) in DMF (3 mL), was added NH₄Cl (30 mg, 0.56 mmol) and NaN₃ (30 mg, 0.46 mmol). The resulting mixture was heated at 100 °C for 18 h and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford (4-(3-(4-(2H-tetrazol-5-yl)phenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (25 mg, 33%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 9.15 (d, *J* = 1.2 Hz, 1H), 9.91 (d, *J* = 1.2 Hz, 1H), 8.29 (d, *J* = 8.4 Hz, 2H), 8.13 (t, *J* = 8.7 Hz, 3H), 8.06 (s, 1H),

7.91 (d, $J = 8.4$ Hz, 2H), 7.60 (d, $J = 8.3$ Hz, 2H), 3.86 (m, 4H), 3.19 (m, 4H), 2.84 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 172.09, 165.22, 160.86, 143.88, 141.92, 140.60, 139.91, 135.96, 135.77, 130.35, 129.80, 129.15, 127.98, 115.40, 54.60, 44.2; MS (ESI) m/z 366 $[\text{C}_{25}\text{H}_{23}\text{N}_9\text{O} + \text{H}]^+$.


5

Example 51: (4-methylpiperazin-1-yl)(4-(3-(4-(trifluoromethyl)phenyl)imidazo[1,2-
a]pyrazin-6-yl)phenyl)methanone



To a solution of (4-(3-iodoimidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (200 mg, 0.45 mmol) in a mixture of toluene (4 mL) and ethanol (2 mL) under inert atmosphere, was added K_2CO_3 (124 mg, 0.89 mmol), 4-(trifluoromethyl)phenylboronic acid (101 mg, 0.54 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (52 mg, 0.045 mmol). The resulting mixture was heated in a microwave reactor at 140°C for 15 min then, was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) to afford (4-methylpiperazin-1-yl)(4-(3-(4-(trifluoromethyl)phenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)methanone (150 mg, 72%, AUC HPLC 97%) as a yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 9.18 (s, 1H), 8.91 (s, 1H), 8.11 (d, $J = 8.2$ Hz, 2H), 8.08 (s, 1H), 8.00 (d, $J = 8.2$ Hz, 2H), 7.94 (d, $J = 8.2$ Hz, 2H), 7.54 (d, $J = 8.3$ Hz, 2H), 3.81 (bs, 2H), 3.54 (bs, 2H), 2.55-2.47 (m, 4H), 2.36 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 172.00, 143.96, 142.16, 140.93, 139.31, 136.90, 136.43, 132.89, 131.85, 129.77, 128.75, 128.19, 127.90, 127.57, 125.52, 115.24, 45.99; MS (ESI) m/z 466 $[\text{C}_{25}\text{H}_{22}\text{F}_3\text{N}_5\text{O} + \text{H}]^+$.

Example 52: *N*-hydroxy-4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-
a]pyrazin-3-yl)benzamide



Step 1: To a solution of (4-(3-bromoimidazo[1,2-a]pyrazin-6-yl)phenyl)(4-methylcyclohexyl)methanone (1.00 g, 2.5 mmol) in 1,4-dioxane (25 mL), were added 4-(ethoxycarbonyl)phenylboronic acid (970 mg, 5.0 mmol), K_3PO_4 (1.06 g, 5.0 mmol), water (5 mL) and $Pd(PPh_3)_4$ (580 mg, 0.5 mmol). The reaction mixture was heated at 90 °C for 12 h under argon atmosphere then, was cooled and filtered through a pad of celite. The filtrate was concentrated under reduce pressure and the residue was purified by flash column chromatography (eluent $CHCl_3/MeOH$ 94:6) to afford ethyl 3-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzoate (600 mg, 55%) as a off-white solid. 1H NMR (400 MHz, CD_3OD) δ 9.28–9.27 (d, J = 1.2 Hz, 1H), δ 8.64–8.64 (d, J = 1.2 Hz, 1H), 8.29–8.27 (d, J = 8.0 Hz, 2H), 7.99–7.98 (d, J = 2.0 Hz, 2H), 7.96(s, 1H), 7.74–7.72 (d, J = 8.4 Hz, 2H), 7.55–7.53 (d, J = 8.4 Hz, 2H), 4.48–4.43 (q, J = 7.2 Hz, 2H), 3.84 (bs, 2H), 3.50 (bs, 2H), 2.53 (bs, 2H), 2.39 (bs, 2H), 2.35 (s, 3H), 1.47–1.44 (t, J = 7.2 Hz, 3H); MS (ESI) m/z 470 $[C_{27}H_{27}N_5O_3 + H]^+$.


Step 2: A solution of ethyl 3-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzoate (600 mg, 1.28 mmol) and LiOH (107 mg, 2.56 mmol) in THF/MeOH/ H_2O (6/2.5/2 20 mL) was stirred at room temperature for 5 h. The reaction mixture was evaporated and diluted with H_2O (200 mL) and acidified with aqueous HCl till pH 3. The white precipitate was isolated by filtration and dried to afford the 3-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)benzoic acid as a white solid (450 mg, 80%). 1H NMR (400 MHz, $DMSO-d_6$) δ 9.28 (s, 1H), 9.01 (s, 1H), 8.17–8.11 (m, 5H), 7.94–7.92 (d, J = 8.0 Hz, 2H), 7.50–7.48 (d, J = 8.0 Hz, 2H), 3.61 (bs, 4H), 2.32 (bs, 4H), 2.19 (s, 3H); MS (ESI) m/z 440 $[C_{25}H_{23}N_5O_3 - H]^+$.

Step 3: To a solution of acid (50 mg, 0.113 mmol) in CH_2Cl_2 : DMF (3:1, 4 mL) was added sequentially DIPEA (83 μ L, 0.476 mmol), HOBt (32 mg, 0.238 mmol), EDCI·HCl (45 mg, 0.238 mmol) and NH_2 -OTHP (26 mg, 0.226 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 12 h, then it was diluted with water (3 mL) and extracted with CH_2Cl_2 (3×10 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyrazin-3-yl)-N-(tetrahydro-2H-pyran-2-yloxy)benzamide (44.6 mg, 73%, AUC HPLC 94.5%) as a white solid. 1H NMR (600 MHz, $DMSO-d_6$) δ 11.81 (bs, 1H), 9.30 (s, 1H), 9.00


(s, 1H), 8.19 (s, 1H), 8.16 (d, $J = 12$ Hz, 2H), 7.99 (dd, $J = 10.6, 8.3$ Hz, 4H), 7.51 (d, $J = 12$ Hz, 2H), 5.08-5.04 (m, 1H), 4.13-4.06 (m, 1H), 3.70-3.53 (m, 3H), 2.43-2.27 (m, 4H), 2.22 (s, 3H), 1.79-1.73 (m, 3H), 1.63-1.53 (m, 3H); MS (ESI) m/z 541 [$C_{30}H_{32}N_6O_4 + H$] $^+$.

- 5 Step 4: To a solution of protected hydroxamate (10 mg, 0.018 mmol) in a CH_3CN : MeOH (1:1, 0.6 mL) was added 1M aqueous HCl (0.111 mL) at room temperature. After stirring for 4 h, the solution was concentrated *in vacuo*. The solid was triturated in hexane dried to yield *N*-hydroxy-4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyrazin-3-yl)benzamide (5.3 mg, 63%, AUC HPLC 99%), light yellow solid. 1H NMR (600 MHz, DMSO- d_6) δ 11.39 (bs, 1H), 9.35 (s, 1H), 9.04 (s, 1H), 8.26 (s, 1H), 8.20 (d, $J = 8.4$ Hz, 2H), 8.00 (d, $J = 7.8$ Hz, 2H), 7.95 (d, $J = 8.4$ Hz, 2H), 7.59 (d, $J = 8.4$ Hz, 2H), 4.39-3.60 (m, 6H), 3.05-3.15 (m, 2H), 2.77 (s, 3H); MS (ESI) m/z 457 [$C_{25}H_{24}N_6O_3 + H$] $^+$.


15 Example 53: (4-(3-(4-(difluoromethyl)phenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



- To a mixture of 2-(4-(difluoromethyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (200 mg, 0.5 mmol), (4-(3-bromoimidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (317 mg, 1.25 mmol), K_3PO_4 (212 mg, 1 mmol) in 1,4-Dioxane (10 mL) and water (1 mL) was added $Pd(PPh_3)_4$ (30 mg) and the reaction mixture was heated at 90 °C for 1 h. Water was added to the reaction mixture and extracted with EtOAc. The organic layer was dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $CHCl_3/MeO$ 96.5:3.5%) and by preparative HPLC to give (4-(3-(4-(difluoromethyl)phenyl)imidazo[1,2-*a*]pyrazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (50 mg, 23%, AUC HPLC 99.59%) as an off-white solid; m.p. 73-86 °C. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 9.26 (s, 1 H), 8.60 (s, 1 H), 7.96-7.94 (m, 3 H), 7.77-7.71 (m, 4 H), 7.52 (d, $J = 8.0$ Hz, 2 H), 6.76 (t, $J = 5.6$ Hz, 1 H), 3.82 (bs, 2 H), 3.47 (bs, 2 H), 2.51 (bs, 2 H), 2.34 (bs, 2H), 2.33 (s, 3 H); MS (ESI) m/z 448.46 [$C_{25}H_{23}F_2N_5O + H$] $^+$.


**Formula 3**

Intermediate 12: 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid



- 5 Step 1: To a solution of 6-chloro-3-iodoimidazo[1,2-*b*]pyridazine (1.00 g, 3.58 mmol) in a mixture of DMF (20 mL) and water (1 mL) under inert atmosphere, were added Na₂CO₃ (759 mg, 7.16 mmol), 4-cyanophenylboronic acid (885 mg, 5.37 mmol) and Pd(PPh₃)₄ (414 mg, 0.358 mmol). The resulting mixture was stirred at 90 °C for 18 h and quenched with ice water. The precipitate was isolated by filtration and dried *in vacuo* to afford 4-(6-
- 10 chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (897 mg, 98%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.24-8.21 (m, 2H), 8.18 (s, 1H), 8.02 (d, *J* = 9.4 Hz, 1H), 7.81-7.79 (m, 2H), 7.45 (d, *J* = 9.4 Hz, 1H); MS (ESI) *m/z* 255 [C₁₃H₇CIN₄ + H]⁺.


- Step 2: To a solution of 4-(6-chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (484 mg, 1.9
- 15 mmol) in DMF (20 mL) and water (4 mL) under inert atmosphere were added Cs₂CO₃ (1.24 g, 3.8 mmol), 4-boronobenzoic acid (473 mg, 2.85 mmol) and Pd(dppf)₂Cl₂ (278 mg, 0.38 mmol). The resulting mixture was heated at 90 °C for 18 h, and then was diluted with water (20 mL) and extracted with DCM (3×40 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by
- 20 column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (310 mg, 48%) as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.54 (s, 1H), 8.51 (d, *J* = 8.6 Hz, 2H), 8.40 (d, *J* = 9.6 Hz, 1H), 8.26 (d, *J* = 8.4 Hz, 2H), 8.13 (d, *J* = 8.4 Hz, 2H), 8.03 (d, *J* = 8.6 Hz, 2H), 8.02 (d, *J* = 9.6 Hz, 1H); MS (ESI) *m/z* 341 [C₂₀H₁₂N₄O₂ + H]⁺.

Intermediate 13: 4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid

Step 1: 4-boronobenzoic acid (11.88 g, 71.61 mmol), K_3PO_4 (27.60 g, 13.02 mmol) and $Pd(PPh_3)_4$ (3.75 g, 3.25 mmol) were added sequentially to a solution of 6-chloroimidazo[1,2-b]pyridazine (10 g, 65.10 mmol) in a mixture of 1,4-dioxane/ H_2O (250:50 mL) at room temperature under argon atmosphere. The reaction mixture was refluxed for 6 h and was diluted with water (100 mL) and extracted with EtOAc (3x100 mL). The combined aqueous layer was acidified to pH 2 using citric acid. The precipitate was isolated by filtration and dried under reduced pressure to afford 4-(imidazo[1,2-b]pyridazin-6-yl)benzoic acid (4.5 g, 57%) as an off-white solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 13.01 (bs, 1H), 8.04 (s, 1H), 8.3-8.01 (m, 5H), 7.85 (d, $J = 7.2$ Hz, 2H); MS (ESI) m/z 240 $[M + H]^+$.

Step 2: To a solution of 4-(imidazo[1,2-b]pyridazin-6-yl)benzoic acid (4.5 g, 18.82 mmol) in DCM (25 mL) and ACN (55 mL) was added NBS (3.68 g, 20.71 mmol) and stirred at 0 °C for 3 h. The reaction mixture was filtered, washed with water and dried to afford 4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid (5.5 g, 93%) as a yellow solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 13.01 (bs, 1H), 8.3-8.01 (m, 5H), 7.85 (d, $J = 7.2$ Hz, 2H); MS (ESI) m/z 317 $[M]^+$.

Example 54: (4-(3-(4-chlorophenyl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




Step 1: To a solution of 6-chloro-3-iodoimidazo[1,2-b]pyridazine (1.00 g, 3.58 mmol) in a mixture of DMF (20 mL) and water (1 mL) under inert atmosphere, were added Na_2CO_3 (759 mg, 7.16 mmol), 4-cyanophenylboronic acid (885 mg, 5.37 mmol) and $Pd(PPh_3)_4$ (414 mg, 0.358 mmol). The resulting mixture was stirred at 90 °C for 18 h and quenched with ice water. The precipitate was isolated by filtration and dried *in vacuo* to afford 4-(6-chloroimidazo[1,2-b]pyridazin-3-yl)benzonitrile (897 mg, 98%) as a yellow solid. 1H NMR

(400 MHz, CDCl_3) δ 8.24-8.21 (m, 2H), 8.18 (s, 1H), 8.02 (d, $J = 9.4$ Hz, 1H), 7.81-7.79 (m, 2H), 7.45 (d, $J = 9.4$ Hz, 1H); MS (ESI) m/z 255 [$\text{C}_{13}\text{H}_7\text{ClN}_4 + \text{H}$] $^+$.

Step 2: To a solution of 4-(6-chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (200 mg, 0.785 mmol) in toluene (3 mL) and ethanol (1.5 mL) under inert atmosphere were added K_2CO_3 (347 mg, 2.51 mmol), 4-(4-methylpiperazine-1-carbonyl)phenylboronic acid hydrochloride (268 mg, 0.942 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (91 mg, 0.079 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 15 min and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (91.2 mg, 43%, AUC HPLC 99%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 8.30 (d, $J = 8.6$ Hz, 2H), 8.23-8.21 (m, 2H), 8.06 (d, $J = 8.4$ Hz, 2H), 7.81 (d, $J = 8.6$ Hz, 2H), 7.66-7.61 (m, 3H), 7.31 (bs, 1H), 3.97-3.74 (m, 4H), 2.84 (bs, 4H), 2.56 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 169.57, 165.35, 151.41, 140.15, 137.01, 136.67, 134.25, 132.82, 132.60, 128.12, 127.49, 127.05, 126.70, 126.54, 118.75, 116.71, 111.15, 53.78, 44.43; MS (ESI) m/z 423 [$\text{C}_{25}\text{H}_{22}\text{N}_6\text{O} + \text{H}$] $^+$.


Example 55: 4-(6-(4-(4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



To a solution (4-hydroxypiperidin-1-yl) (4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (94 mg, ~0.28 mmol) in a mixture of DMF (8 mL) and water (2 mL) blanketed with nitrogen, was successively added K_2CO_3 (217 mg, 1.58 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (114 mg, 0.08 mmol) and 4-(6-chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (200 mg, 0.78 mmol). The reaction mixture was stirred at 80 °C for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent $\text{ACN}/\text{H}_2\text{O}/0.01\% \text{HCOOH}$) to afford 4-(6-(4-(4-hydroxypiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (52 mg,

61%, AUC HPLC 99%) as yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.51–8.47 (m, 3H), 8.37 (d, $J = 8.0$ Hz, 1H) 8.19 (d, $J = 8.0$ Hz, 2H), 8.02–7.96 (m, 3H), 7.59 (d, $J = 8.0$ Hz, 2H), 4.82 (s, 1H), 4.03 (s, 1H), 3.77 (s, 1H), 3.54 (s, 1H), 3.37–3.20 (m, 2H), 1.82–1.74 (m, 2H), 1.40 (s, 2H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_4$) ppm 168.14, 150.88, 140.10, 138.02, 135.60, 135.37, 132.84, 132.64, 127.37, 127.17, 126.58, 126.22, 125.87, 118.79, 116.89, 109.47, 65.32, 44.49, 34.391, 33.63; MS (ESI) m/z 424 $[\text{C}_{25}\text{H}_{21}\text{N}_5\text{O}_2 + \text{H}]^+$.

Example 56: 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



10

Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (10 mL) were added HATU (168 mg, 0.441 mmol), *N*-methyl morpholine (59 mg, 0.588 mmol) and *tert*-butyl piperidin-4-ylcarbamate (126 mg, 0.588 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) to afford title compound as a yellow solid. MS (ESI) m/z 523 $[\text{C}_{30}\text{H}_{30}\text{N}_6\text{O}_3 + \text{H}]^+$.


20

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-ylcarbamate (0.294 mmol) in DCM (1 mL) was added TFA (1 mL). The resulting mixture was stirred at room temperature for 18 h and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /0.01% HCOOH) to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (20.5 mg, 15%, AUC HPLC 98%) as a yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 8.30 (d, $J = 8.4$ Hz, 2H), 8.21 (bs, 1H), 8.14–8.10 (m, 3H), 7.83 (d, $J = 9.4$ Hz, 1H), 7.76 (d, $J = 8.4$ Hz, 2H), 7.62 (d, $J = 8.2$ Hz, 2H), 4.78 (bs, 1H), 3.89 (bs, 1H), 3.51–3.45 (m, 1H), 3.32–3.22 (m, 1H), 3.07–2.99 (m, 1H), 2.18–2.05 (m, 2H), 1.67 (bs, 2H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.72, 167.66, 152.79, 141.68, 138.46, 138.03, 135.25, 134.11, 133.59, 128.88, 128.64, 128.17, 127.68,

30

127.07, 119.71, 118.50, 111.88, 47.10, 41.61, 31.59, 30.85; MS (ESI) m/z 423 [$C_{25}H_{22}N_6O + H$] $^+$.


Example 57: *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)acetamide



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (10 mL) were added HATU (224 mg, 0.588 mmol), *N*-methyl morpholine (89 mg, 0.882 mmol) and *N*-(piperidin-4-yl)acetamide (125 mg, 0.882 mmol).

- The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5 to 90:10) to afford *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)acetamide, (33 mg, 24%, AUC HPLC 98%) as a yellow solid. 1H NMR (400 MHz, DMSO- d_6) δ 8.53-8.49 (m, 3H), 8.39 (d, J = 9.6 Hz, 1H), 8.22 (d, J = 8.3 Hz, 2H), 8.03-7.98 (m, 3H), 7.86 (d, J = 7.6 Hz, 1H), 7.59 (d, J = 8.3 Hz, 2H), 4.34 (bs, 1H), 3.85-3.83 (m, 1H), 3.59 (bs, 1H), 3.19-3.03 (m, 2H), 1.89-1.70 (m, 5H), 1.35-1.34 (m, 2H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 168.31, 168.23, 150.88, 140.12, 137.83, 135.71, 135.38, 132.85, 132.67, 127.39, 127.23, 126.61, 126.27, 125.91, 118.79, 116.93, 109.49, 45.48, 22.63; MS (ESI) m/z 465 [$C_{27}H_{24}N_6O_2 + H$] $^+$.

Example 58: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (10 mL) were added HATU (168 mg, 0.441 mmol), *N*-methyl morpholine (59 mg, 0.588 mmol) and *tert*-butyl 4-methylpiperidin-4-ylcarbamate (126 mg,

0.588 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate as a yellow solid. MS (ESI) *m/z* 537 [C₃₁H₃₂N₆O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (0.294 mmol) in DCM (1 mL) was added TFA (1 mL). The resulting mixture was stirred at room temperature for 18 h and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (23.5 mg, 17%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.28 (d, *J* = 8.4 Hz, 2H), 8.18 (bs, 1H), 8.12-8.08 (m, 3H), 7.82 (d, *J* = 9.4 Hz, 1H), 7.74 (d, *J* = 8.4 Hz, 2H), 7.61 (d, *J* = 8.0 Hz, 2H), 4.36 (bs, 1H), 3.73 (bs, 1H), 3.50-3.47 (m, 2H), 1.94-1.85 (m, 4H), 1.53 (s, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 171.70, 152.75, 138.42, 137.99, 135.25, 134.08, 133.58, 128.89, 128.62, 128.10, 127.63, 127.06, 119.72, 118.48, 111.84, 53.50, 44.88, 39.31, 36.73, 36.01, 22.66; MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.


Example 59: 4-(6-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (55 mg, 0.162 mmol) in DMF (10 mL) were added HATU (123 mg, 0.324 mmol), *N*-methylmorpholine (49 mg, 0.486 mmol) and 1-isopropylpiperazine (62 mg, 0.486 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude

residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 4-(6-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (32 mg, 44%, AUC HPLC 98%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.53-8.49 (m, 3H), 8.39 (d, *J* = 9.5 Hz, 1H), 8.22 (d, *J* = 8.3 Hz, 2H), 8.03-8.01 (m, 3H), 7.64 (d, *J* = 8.2 Hz, 2H), 3.90-3.59 (m, 3H), 2.99-2.92 (m, 2H), 2.75 (bs, 4H), 1.08 (d, *J* = 6.3 Hz, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.14, 150.84, 137.18, 135.92, 135.42, 132.85, 132.66, 127.79, 127.18, 126.64, 126.28, 125.93, 118.79, 116.92, 109.50, 47.54, 17.43; MS (ESI) *m/z* 451 [C₂₇H₂₆N₆O + H]⁺.

10 Example 60: 4-(6-(4-(4-(methylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (55 mg, 0.162 mmol) in DMF (10 mL) were added HATU (123 mg, 0.324 mmol), *N*-methyl morpholine (49 mg, 0.486 mmol) and *tert*-butyl piperidin-4-ylcarbamate (104 mg, 0.486 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl(methyl)carbamate as a yellow solid. MS (ESI) *m/z* 537 [C₃₁H₃₂N₆O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl(methyl)carbamate (0.162 mmol) in DCM (1 mL) was added TFA (1 mL). The resulting mixture was stirred at room temperature for 18 h and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(4-(4-(methylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (29.8 mg, 42%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.29 (d, *J* = 8.4 Hz, 2H), 8.19 (bs, 1H), 8.13-8.09 (m, 3H), 7.82 (d, *J* = 9.4 Hz, 1H), 7.75 (d, *J* = 8.4 Hz, 2H), 7.62 (d, *J* = 8.0 Hz,

2H), 3.94 (bs, 1H), 3.40 (bs, 1H), 3.34-3.20 (m, 2H), 3.01(bs, 1H), 2.77 (s, 3H), 2.37-2.16 (m, 2H), 1.66 (bs, 2H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.69, 152.76, 138.39, 138.02, 135.24, 134.09, 133.59, 128.91, 128.62, 128.12, 127.65, 127.06, 119.72, 118.49, 111.84, 57.16, 47.09, 41.58, 30.66, 29.87, 29.16; MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.


5

Example 61: 4-(6-(4-(4-ethylpiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (120 mg, 0.353 mmol) in DMF (10 mL) were added HATU (402 mg, 1.06 mmol), *N*-methyl morpholine (143 mg, 1.41 mmol) and 1-ethylpiperazine (181 mg, 1.41 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h before diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 ~ 90:10) to afford 4-(6-(4-(4-(methylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (64.6 mg, 42%, AUC HPLC 99%) as yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.54 (s, 1H), 8.51 (d, $J = 8.7$ Hz, 2H), 8.42 (d, $J = 9.6$ Hz, 1H), 8.25 (d, $J = 8.4$ Hz, 2H), 8.03-8.00 (m, 3H), 7.69 (d, $J = 8.2$ Hz, 2H), 3.44-3.22 (m, 4H), 3.17-3.04 (m, 6H), 1.24-1.20 (m, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 168.45, 162.96, 150.86, 140.21, 136.40, 135.54, 132.93, 132.73, 128.01, 127.34, 126.77, 126.39, 126.02, 118.86, 117.00, 109.62, 51.01, 50.26, 9.12; MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.


Example 62: 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



To a solution of 4-(6-chloroimidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (150 mg, 0.59 mmol) in DMF (2.5 mL) and water (0.5 mL) under inert atmosphere were added Cs_2CO_3 (384 mg,

1.18 mmol), morpholino(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (279 mg, 0.88 mmol) and Pd(dppf)₂Cl₂ (86 mg, 0.118 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 30 min, and then was diluted with water (10 mL) and extracted with DCM (3×15 mL). The combined organic layer was dried over Na₂SO₄,
 5 filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (72.1 mg, 35%, AUC HPLC 96%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.31-8.25 (m, 4H), 8.06-8.05 (m, 2H), 7.82-7.81 (m, 2H), 7.70-7.61 (m, 3H), 3.78-3.53 (m, 8H); ¹³C NMR (100 MHz, CDCl₃) δ 169.39, 151.86, 137.43, 136.45, 133.24, 132.63, 132.52, 128.14, 128.08, 127.47,
 10 127.26, 126.86, 126.30, 118.66, 117.32, 111.39, 66.87, 48.21, 42.67; MS (ESI) *m/z* 410 [C₂₄H₁₉N₅O₂ + H]⁺.

Example 63: 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid (500 mg, 1.57 mmol) in DMF (5 mL) was added NMM (317 mg, 3.14 mmol) followed by HATU (754.7 mg, 2.35 mmol) and the solution was stirred for 30 min at rt. *N,N*-dimethylpiperidin-4-amine
 20 (221 mg, 1.72 mmol) was added to the reaction mixture and stirred for an additional 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-(dimethylamino)piperidin-1-yl)methanone (350 mg, 52%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.01 (bs, 1H), 8.3-8.01 (m, 5H), 7.85 (d, *J* = 7.2 Hz, 2H); MS (ESI) *m/z* 317 [M]⁺.

Step 2: To a mixture of (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-(dimethylamino)piperidin-1-yl)methanone (350 mg, 0.817 mmol), 4-cyanophenylboronic
 30


- acid (144 mg, 0.981 mmol), K_3PO_4 (346 mg, 1.63 mmol) in 1,4-dioxane (25 mL) and water (5 mL) was added $Pd(PPh_3)_4$ (47 mg, 0.04 mmol). The reaction mixture was heated at 90 °C for 6 h under argon atmosphere then, was diluted with water (200 mL). The precipitate was isolated by filtration and was purified by column chromatography (silica gel, eluent $CHCl_3/MeOH$ 95:5) to afford 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (150 mg, 41%, AUC HPLC 98.9%) as a yellow solid; 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.78 (d, $J = 8.77$ Hz, 1H), 8.52 (s, 1H), 8.30 (d, $J = 8.2$ Hz, 2H), 8.23 (d, $J = 8.2$ Hz, 2H), 7.76 (d, $J = 7.7$ Hz, 2H), 7.62 (d, $J = 7.6$ Hz, 2H), 7.41 (d, $J = 7.4$ Hz, 1H), 4.9 (bs, 1H), 3.9 (bs, 1H), 3.17 (bs, 1H), 2.91 (bs, 1H), 2.45 (bs, 1H), 2.38 (bs, 6H), 2.08 (bs, 1H), 1.95 (bs, 1H), 1.52 (b. s, 2H); MS (ESI) m/z 451.2 $[C_{27}H_{26}N_6O + H]^+$.

Example 64: 4-(6-(4-(1,4-oxazepane-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



- To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid (200 mg, 0.588 mmol) in DMF (5 mL) was added NMM (118.8 mg, 1.176 mmol) followed by HATU (376.3 mg, 1.17 mmol) and the solution was stirred at rt for 30 min. 1,4-oxazepane (65 mg, 0.646 mmol) was added to the reaction mixture which was stirred for an additional 16 h.
- The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) to afford 4-(6-(4-(1,4-oxazepane-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (120 mg, 50%, AUC HPLC 99.42%) as a pale-yellow solid; 1H NMR (400 MHz, $CDCl_3$) δ 8.33 (d, $J = 7.9$ Hz, 2H), 8.21 (s, 1H), 8.16 (d, $J = 9.7$ Hz, 1H), 8.05 (d, $J = 7.5$ Hz, 2H), 7.82 (d, $J = 8.3$ Hz, 2H), 7.63-7.59 (m, 3H), 3.9 (m, 5H), 3.7 (bs, 1H), 3.59 (bs, 2H), 2.17 (bs, 1H), 1.87 (bs, 1H); MS (ESI) m/z 424.31 $[C_{25}H_{21}N_5O_2 + H]^+$.


Example 65: 4-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile



Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (120 mg, 0.353 mmol) in DMF (10 mL) were added HATU (402 mg, 1.06 mmol), *N*-methyl morpholine (143 mg, 1.41 mmol) and *tert*-butyl piperazine-1-carboxylate (263 mg, 1.41 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate as a yellow solid. MS (ESI) *m/z* 509 [C₂₉H₂₈N₆O + H]⁺.

Step 2: To a solution of *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate (0.353 mmol) in DCM (1 mL) was added TFA (1 mL). The resulting mixture was stirred at room temperature for 18 h and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (15.4 mg, 11%, AUC HPLC 92%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.45 (d, *J* = 8.6 Hz, 2H), 8.34 (s, 1H), 8.26-8.22 (m, 3H), 7.95 (d, *J* = 9.6 Hz, 1H), 7.89 (d, *J* = 8.6 Hz, 2H), 7.71 (d, *J* = 8.4 Hz, 2H), 3.87 (bs, 4H), 3.27 (bs, 4H); ¹³C NMR (100 MHz, CD₃OD) δ 171.80, 152.88, 138.46, 137.54, 135.28, 134.19, 133.64, 129.23, 129.11, 128.72, 128.27, 127.85, 127.14, 119.70, 118.58, 112.00, 44.56; MS (ESI) *m/z* 409 [C₂₄H₂₀N₆O + H]⁺.

Example 66: N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-ethylpiperidin-4-yl)acetamide



Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (120 mg, 0.353 mmol) in DMF (10 mL) were added HATU (402 mg, 1.06 mmol), *N*-methyl morpholine (143 mg, 1.41 mmol) and *tert*-butyl 4-ethylpiperidin-4-ylcarbamate (322 mg, 1.41 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-ethylpiperidin-4-ylcarbamate as a yellow solid. MS (ESI) *m/z* 509 [C₂₉H₂₈N₆O + H]⁺.


Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-ethylpiperidin-4-ylcarbamate (0.353 mmol) in DCM (1 mL) was added TFA (1 mL). The reaction mixture was stirred at room temperature for 18 h and was concentrated under reduced pressure. The crude compound was carried forth to the next step without further purification. MS (ESI) *m/z* 409 [C₂₇H₂₆N₆O + H]⁺.

Step 3: To a solution of 4-(6-(4-(4-amino-4-ethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (0.353 mmol) in DCM (1 mL) were added acetic anhydride (0.1 mL, 1.06 mmol) and triethylamine (0.15 mL, 1.06 mmol). The resulting mixture was stirred at room temperature for 1 h and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) and preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-4-ethylpiperidin-4-yl)acetamide (44.2 mg, 25%, AUC HPLC 96%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.20 (d, *J* = 8.3 Hz, 2H), 8.02-7.98 (m, 3H), 7.72 (d, *J* = 9.5 Hz, 1H), 7.67 (d, *J* = 8.3 Hz, 2H), 7.55-7.53 (m, 3H), 4.39-4.36 (m, 1H), 3.58-3.55 (m, 1H), 3.37-3.34 (m, 1H), 3.24-3.18 (m, 1H), 2.31-

2.29 (m, 2H), 2.01(s, 3H), 1.86-1.77 (m, 2H), 1.55-1.50 (m, 2H), 0.86 (t, $J = 7.4$ Hz, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.50, 171.37, 152.67, 138.98, 137.55, 135.19, 133.98, 133.59, 133.54, 128.83, 128.45, 127.49, 126.93, 119.70, 118.35, 111.80, 56.18, 45.30, 39.55, 35.26, 34.54, 31.58, 23.64, 7.70; MS (ESI) m/z 493 [$\text{C}_{29}\text{H}_{28}\text{N}_6\text{O}_2 + \text{H}$] $^+$.

5

Example 67: *N*-((1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)methyl)acetamide



Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (120 mg, 0.353 mmol) in DMF (10 mL) were added HATU (402 mg, 1.06 mmol), *N*-methylmorpholine (143 mg, 1.41 mmol) and *tert*-butyl piperidin-4-ylmethylcarbamate (302 mg, 1.41 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) to afford *tert*-butyl (1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)methylcarbamate as a yellow solid. MS (ESI) m/z 537 [$\text{C}_{31}\text{H}_{32}\text{N}_6\text{O}_3 + \text{H}$] $^+$.

Step 2: A solution of *tert*-butyl (1-(4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)methylcarbamate (0.353 mmol) in DCM (1 mL) and TFA (1 mL) was stirred at room temperature for 18 h and concentrated under reduced pressure. The crude compound was used for next step without further purification. MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.


25

Step 3: To a solution of 4-(6-(4-(4-(aminomethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (0.353 mmol) in DCM (1 mL) were added acetic anhydride (0.1 mL, 1.06 mmol) and triethylamine (0.15 mL, 1.06 mmol). The resulting mixture was stirred at room temperature for 1 h and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5 to 90:10) and preparative HPLC (C18, eluent $\text{ACN}/\text{H}_2\text{O}/0.01\% \text{HCOOH}$) to afford *N*-((1-(4-(3-(4-

30


cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperidin-4-yl)methyl)acetamide (40.3 mg, 24%, AUC HPLC 96%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.42 (d, *J* = 8.6 Hz, 2H), 8.29 (bs, 1H), 8.18 (d, *J* = 8.6 Hz, 3H), 7.90 (d, *J* = 9.6 Hz, 1H), 7.86 (d, *J* = 8.6 Hz, 2H), 7.60 (d, *J* = 8.4 Hz, 2H), 4.68-4.65 (m, 1H), 3.79-3.76 (m, 1H), 3.13 (bs, 3H), 2.93-2.87 (m, 1H), 1.95 (s, 3H), 1.88-1.79 (m, 2H), 1.73-1.70 (m, 1H), 1.28-1.24 (m, 2H); ¹³C NMR (100 MHz, CD₃OD) δ 173.51, 171.60, 153.21, 139.20, 137.96, 135.24, 134.37, 133.75, 128.78, 128.69, 128.60, 128.07, 127.62, 127.17, 119.75, 118.77, 45.65, 43.39, 37.59, 31.55, 30.70, 29.07, 22.56; MS (ESI) *m/z* 479 [C₂₈H₂₆N₆O₂ + H]⁺.

10 Example 68: 4-(6-(4-(3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (50 mg, 0.147 mmol) in DMF (0.74 mL) were added HATU (84 mg, 0.221 mmol), *N*-methyl morpholine (65 μL, 0.588 mmol) and 2-oxopiperazine (18 mg, 0.177 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 92:8) to afford 4-(6-(4-(3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)benzonitrile (28 mg, 45%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.53-8.46 (m, 3H), 8.38 (d, *J* = 9.6 Hz, 1H), 8.22 (d, *J* = 8.4 Hz, 2H), 8.15 (s, 1H), 8.03-7.96 (m, 3H), 7.67 (d, *J* = 8.4 Hz, 2H), 4.20-4.05 (m, 1H), 4.05-3.90 (m, 1H), 3.90-3.70 (m, 1H), 3.60-3.50 (m, 1H), 3.30-3.20 (m, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.44, 150.95, 140.24, 137.06, 136.24, 135.51, 132.94, 132.79, 127.93, 127.36, 126.74, 126.39, 126.04, 118.92, 117.07, 109.62, 30.70; MS (ESI) *m/z* 423 [C₂₄H₁₈N₆O₂ + H]⁺.

Example 69: 4-(6-(4-(4-methyl-3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid (50 mg, 0.147 mmol) in DMF (0.74 mL) were added HATU (84 mg, 0.221 mmol), *N*-methyl morpholine (65 μ L, 0.588 mmol) and 1-methyl-2-oxopiperazine hydrochloride (27 mg, 0.177 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 92:8), followed by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(4-methyl-3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (16 mg, 24%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.52-8.47 (m, 3H), 8.39 (d, *J* = 9.6 Hz, 1H), 8.23 (d, *J* = 8.4 Hz, 2H), 8.04-7.97 (m, 3H), 7.67 (d, *J* = 8.4 Hz, 2H), 4.25-3.95 (m, 2H), 3.95-3.75 (m, 1H), 3.75-3.55 (m, 1H), 3.50-3.20 (m, 2H), 2.88 (s, 3H); MS (ESI) *m/z* 437 [C₂₅H₂₀N₆O₂ + H]⁺.

Example 70: 4-(6-(4-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid (50 mg, 0.147 mmol) in DMF (0.74 mL) were added HATU (84 mg, 0.221 mmol), *N*-methyl morpholine (65 μ L, 0.588 mmol) and 1-methyl-4-(piperidin-4-yl)piperazine (32 mg, 0.177 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 90:10) to

afford 4-(6-(4-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (17 mg, 23%, AUC HPLC 99%) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.53-8.47 (m, 3H), 8.38 (d, $J = 9.6$ Hz, 1H), 8.20 (d, $J = 8.4$ Hz, 2H), 8.04-7.96 (m, 3H), 7.60 (d, $J = 8.4$ Hz, 2H), 4.55-4.40 (m, 1H), 3.70-3.60 (m, 1H), 3.50-3.20 (m, 4H), 3.15-3.00 (m, 1H), 2.90-2.75 (m, 1H), 2.55-2.40 (m, 2H), 2.40-2.20 (m, 3H), 2.13 (s, 3H), 1.95-1.65 (m, 2H), 1.50-1.35 (m, 2H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 168.18, 151.03, 140.24, 138.08, 135.75, 135.47, 132.97, 132.80, 127.56, 127.29, 126.73, 126.41, 126.04, 118.92, 117.07, 109.61, 60.71, 55.16, 48.50, 45.75, 30.70; MS (ESI) m/z 506 $[\text{C}_{30}\text{H}_{31}\text{N}_7\text{O} + \text{H}]^+$.

10

Example 71: 4-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.47 mL) were added HATU (168 mg, 0.441 mmol), *tert*-butyl 1,9-diazaspiro[5.5]undecane-1-carboxylate hydrochloride (103 mg, 0.353 mmol) and *N*-methylmorpholine (162 μL , 1.47 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with H_2O (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CHCl}_3/\text{MeOH}$ 96:4) to afford *tert*-butyl 9-(4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)phenyl)carbonyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (70 mg, 41%) as a yellow solid. MS (ESI) m/z 577 $[\text{C}_{34}\text{H}_{36}\text{N}_6\text{O}_3 + \text{H}]^+$.

Step 2: To *tert*-butyl 9-(4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (50 mg, 0.131 mmol) was added 20% TFA in CH_2Cl_2 (5 mL). The reaction mixture was stirred at room temperature for 20 min then concentrated under reduced pressure. To the residue was added saturated sodium bicarbonate (10 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(1,9-diazaspiro[5.5]undecane-

9-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (37 mg, 89%, AUC HPLC 99%) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.54-8.48 (m, 3H), 8.38 (d, $J = 9.6$ Hz, 1H), 8.28 (s, 1H), 8.21 (d, $J = 8.0$ Hz, 2H), 8.04-7.97 (m, 3H), 7.59 (d, $J = 8.0$ Hz, 2H), 3.75-3.60 (m, 3H), 2.85-2.70 (m, 2H), 1.80-1.40 (m, 11H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 168.36, 164.63, 151.10, 140.29, 138.00, 135.87, 135.50, 133.01, 132.86, 127.61, 127.40, 126.79, 126.49, 126.11, 118.99, 117.14, 109.68, 30.76, 24.82, 18.89; MS (ESI) m/z 477 $[\text{C}_{29}\text{H}_{28}\text{N}_6\text{O} + \text{H}]^+$.

Example 72: 4-(6-(3-chloro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 4-(6-chloroimidazo[1,2-b]pyridazin-3-yl)benzonitrile (100 mg, 0.39 mmol) in a mixture of DMF (4 mL) and water (1 mL) under inert atmosphere, was successively added K_2CO_3 (108 mg, 0.78 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (57 mg, 0.08 mmol) and 4-borono-2-chlorobenzoic acid (78 mg, 0.39 mmol) were added at room temperature. The reaction mixture was stirred at 80 °C for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid as a black solid. MS (ESI) m/z 375 $[\text{C}_{20}\text{H}_{11}\text{ClN}_4\text{O}_2 + \text{H}]^+$.

Step 2: To a solution of 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)benzoic acid in DMF (5 mL) was successively added HATU (223 mg, 0.59 mmol), *N*-methyl morpholine (80 mg, 0.78 mmol) and 1-methylpiperazine (78 mg, 0.78 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and the aqueous phase extracted with DCM (3×10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/ H_2O / HCOOH 0.01%) to afford 4-(6-(3-chloro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (50 mg, 28%, AUC HPLC 95%) as yellow solid. ^1H NMR (400 MHz,

CD₃OD) δ 8.06–7.97 (m, 3H), 7.96–7.85 (m, 3H), 7.64 (d, J = 9.5 Hz, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.49 (d, J = 8.0 Hz, 1H), 3.97 (s, 2H), 3.48 (s, 2H), 2.93 (s, 2H), 2.85 (s, 2H), 2.60 (s, 3H); ¹³C NMR (100 MHz, CD₃OD) δ 168.03, 151.17, 141.47, 138.71, 137.81, 135.52, 133.65, 132.25, 130.16, 130.01, 129.61, 128.93, 128.01, 127.73, 127.23, 127.17, 127.11, 119.65, 117.95, 111.78, 55.23, 54.75, 48.42, 46.57, 45.08, 41.43, 30.75; MS (ESI) m/z 457 [C₂₅H₂₁ClN₆O + H]⁺.

Example 73: 4-(6-(3-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



10

Step 1: To a solution of 4-(6-chloroimidazo[1,2-b]pyridazin-3-yl)benzonitrile (100 mg, 0.39 mmol) in a mixture of DMF (4 mL) and water (1 mL) at room temperature, was successively added K₂CO₃ (108 mg, 0.78 mmol), Pd(dppf)Cl₂ (57 mg, 0.08 mmol) and 4-borono-2-fluorobenzoic acid (72 mg, 0.39 mmol) under inert atmosphere. The reaction mixture was stirred at 80 °C for 18 h and was diluted with water (10 mL). The aqueous phase was extracted with DCM (3×10 mL) and the combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-fluorobenzoic acid as black solid. MS (ESI) m/z 359 [C₂₀H₁₁FN₄O₂ + H]⁺.


20

Step 2: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-fluorobenzoic acid (~ 0.39 mmol) in DMF (5 mL) was added HATU (223 mg, 0.59 mmol), *N*-methylmorpholine (80 mg, 0.78 mmol) and 1-methylpiperazine (78 mg, 0.78 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄, filtered and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(3-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (35 mg, 20%, AUC HPLC 97%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.07–8.33 (m, 3H), 8.02 (d, J = 9.0 Hz, 1H), 7.87 (d, J = 7.6 Hz, 1H), 7.63–7.82 (m, 4H), 7.58 (t, J = 7.2

30

Hz, 1H), 3.99 (bs, 2H), 3.61 (bs, 2H), 3.03 (bs, 2H), 2.93 (bs, 2H), 2.67 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 167.15, 166.38, 161.13, 158.67, 151.44, 141.60, 140.24, 140.16, 135.51, 133.56, 135.51, 133.80, 133.56, 131.29, 131.19, 130.15, 127.97, 127.44, 127.15, 126.08, 125.90, 124.73, 124.70, 119.67, 118.14, 115.68, 115.45, 111.89, 55.11, 54.66, 46.59, 44.77, 41.44; MS (ESI) m/z 441 [$\text{C}_{25}\text{H}_{21}\text{FN}_6\text{O} + \text{H}$] $^+$.

Example 74: 4-(6-(3-methyl-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




- Step 1: To a solution of 4-(6-chloroimidazo[1,2-b]pyridazin-3-yl)benzonitrile (200 mg, 0.67 mmol) in a mixture of DMF (4 mL) and water (1 mL) blanketed with nitrogen, was successively added K_2CO_3 (185 mg, 1.34 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (98 mg, 0.13 mmol) and 4-borono-2-methylbenzoic acid (131 mg, 0.73 mmol). The reaction mixture was stirred at 80 °C for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-methylbenzoic acid (271 mg) as black solid which was used carried forth to the next step without further purification. MS (ESI) m/z 355 [$\text{C}_{21}\text{H}_{14}\text{FN}_4\text{O}_2 + \text{H}$] $^+$.
- Step 2: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-methylbenzoic acid (90 mg, crude) in DMF (5 mL), was added HATU (142 mg, 0.38 mmol), *N*-methylmorpholine (52 mg, 0.50 mmol) and 1-methylpiperazine (51 mg, 0.50 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /0.01% HCOOH) to afford 4-(6-(3-methyl-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (30 mg, 28%, AUC HPLC 99%) as a brown solid. ^1H NMR (400 MHz, CD_3OD) δ 8.12 (s, 1H), 8.02 (d, $J = 9.5$ Hz, 1H), 7.91–7.82 (m, 2H), 7.75 (d, $J = 9.5$ Hz, 3H), 7.70 (d, $J = 8.4$ Hz, 2H), 7.39 (d, $J = 7.9$ Hz, 1H), 4.09–3.85 (m, 2H), 3.48 (s, 2H), 2.95 (s, 2H), 2.79 (s, 2H),

2.60 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 169.86, 165.89, 151.43, 140.16, 136.98, 135.48, 133.71, 132.62, 132.10, 128.96, 126.58, 126.51, 126.08, 125.48, 124.59, 118.29, 117.07, 110.36, 53.90, 53.46, 45.20, 43.59, 39.84, 17.97; MS (ESI) m/z 437 $[\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}]^+$.

5


Example 75: 4-(6-(3-isopropoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



To a solution of 4-(6-chloroimidazo[1,2-a]pyridazin-3-yl)benzonitrile (100 mg, 0.39 mmol) in a mixture of DMF (4 mL) and water (1 mL) blanketed with nitrogen, was sequentially added K_2CO_3 (108 mg, 0.78 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (57 mg, 0.08 mmol) and (2-isopropoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (200 mg, 0.39 mmol). The reaction mixture was stirred at 80 °C for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /0.01% HCOOH) to afford 4-(6-(3-isopropoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (74 mg, 40%, AUC HPLC 99%) as yellow solid. ^1H NMR (400 MHz, CD_3OD) δ ppm 8.00–8.14 (m, 3H), 7.95 (d, $J = 8.16$ Hz, 1H), 7.66–7.74 (m, 1H), 7.51–7.60 (m, 4H), 7.40 (d, $J = 6.90$ Hz, 1H), 4.69–4.82 (m, 1H), 3.82–4.20 (m, 2H), 3.57 (s, 2H), 2.90–3.18 (m, 4H), 2.74 (s, 3H), 1.31–1.49 (m, 6H); ^{13}C NMR (100 MHz, CD_3OD) δ ppm 169.42, 155.57, 152.50, 138.95, 135.28, 133.94, 130.30, 129.05, 128.73, 127.23, 126.91, 120.88, 119.67, 118.36, 112.90, 111.70, 72.498, 55.09, 55.092, 54.63, 48.42, 46.06, 44.56, 40.80, 22.52, 22.36; MS (ESI) m/z 481 $[\text{C}_{28}\text{H}_{28}\text{N}_6\text{O}_2 + \text{H}]^+$.


25

Example 76: 4-(6-(3-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



Boron tribromide (195 mg, 0.78 mmol) was added slowly to a solution of 4-(6-(3-isopropoxy-4-(4-methylpiperazine-1-carbonyl) phenyl) imidazo[1,2-b]pyridazin-3-yl)benzonitrile (237 mg, 0.49 mmol) in DCM (10 ml) at -78 °C then the reaction mixture was stirred at room temperature for 6 h. The reaction mixture was quenched by adding MeOH and concentrated under vacuum. The crude residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(3-isopropoxy-4-(4-methylpiperazine-1-carbonyl)phenyl) imidazo[1,2-b]pyridazin-3-yl)benzonitrile (18 mg, 11%, AUC HPLC 95%) as yellow solid. ¹H NMR (400 MHz, CD₃OD) δ ppm 8.29 (d, *J* = 8.0 Hz, 3H), 8.19 (s, 1 H), 8.00–8.12 (m, 1H), 7.70–7.82 (m, 3H), 7.49–7.63 (m, 2H), 7.40 (d, *J* = 8.0 Hz, 1H), 3.48–4.11 (m, 4H), 3.01 (s, 4H), 2.64–2.74 (m, 3 H); ¹³C NMR (100 MHz, CD₃OD) δ ppm 169.82, 167.16, 155.65, 152.74, 141.68, 139.22, 135.16, 134.13, 133.71, 133.58, 130.52, 128.06, 127.66, 126.91, 126.42, 125.79, 119.83, 119.72, 118.45, 115.15, 111.86, 54.97, 44.70, 40.47; MS (ESI) *m/z* 439 [C₂₅H₂₂N₆O₂+H]⁺

Example 77: 4-(6-(3-hydroxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



20

- Step 1: To a solution of 4-bromo-2-methoxybenzoic acid (2 g, 8.66 mmol) in DMF (20 mL) were added HATU (4.94 g, 12.99 mmol), N-methyl morpholine (1.75 g, 17.32 mmol) and morpholine (0.83 g, 0.95 mmol). The reaction mixture was stirred at 0 °C to rt for 16 h, then was diluted with H₂O (15 mL) and extracted with EtOAc (3 × 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure to afford (4-bromo-2-methoxyphenyl)(morpholino)methanone (1.4 g, 54%). ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 7.07–7.04 (m, 2H), 6.99 (s, 1H), 3.78 (s, 3H), 3.74–3.71 (bs, 4H), 3.69–3.52 (bs, 2H), 3.19–3.66 (bs, 2H); MS (ESI) *m/z* 301 [C₁₂H₁₄BrNO₃ + H]⁺.
- 10 Step 2: To a mixture of (4-bromo-2-methoxyphenyl)(morpholino)methanone (1.19 g, 3.96 mmol), bis(pinacolato)diboron (1.07 g, 402 mmol), KOAc (1.16 g, 11.89 mmol) in 1,4-dioxane (18 mL) was added PdCl₂dppf (87.08 mg, 0.12 mmol), dppf (65.97 mg, 0.12 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous
- 15 Na₂SO₄ and concentrated under reduced pressure to afford (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone (1.23 g, 90%) as a brown liquid which was used in next step without purification. Direct mass indicated desired *m/z* 345
- 20 Step 3: To a mixture of (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone 2 (1.23 g, 3.54 mmol), 6-chloroimidazo[1,2-b]pyridazine (0.43 g, 2.84 mmol), K₃PO₄ (1.5 g, 7.08 mmol) in 1,4-dioxane (18 mL) and water (3 mL) was added Pd(PPh₃)₄ (204 mg, 0.17 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with water and extracted with EtOAc and
- 25 washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford (4-(imidazo[1,2-b]pyridazin-6-yl)-2-methoxyphenyl)(morpholino)methanone (0.7 g, 58.8 %, LC-MS 85%) as a light yellow solid. ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 8.04 (d, *J* = 6.8
- 30 Hz, 2H), 7.82 (s, 1H), 7.59 (s, 1H), 7.49–7.40 (m, 3H), 3.98 (s, 3H), 3.87–3.65 (m, 4H), 3.61–3.33 (m, 2H), 3.29 (bs, 2H); MS (ESI) *m/z* 339.14 [C₁₈H₁₈N₄O₃ + H]⁺.


Step 4: To a solution of (4-(imidazo[1,2-b]pyridazin-6-yl)-2-methoxyphenyl)(morpholino)methanone (0.66 g, 1.95 mmol) in ACN (10 mL) was added

NIS (0.52 g, 2.34 mmol) and stirred at 70 °C for 3 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-(3-iodoimidazo[1,2-b]pyridazin-6-yl)-2-methoxyphenyl(morpholino)methanone (0.88 g, 97%, LC-MS 65%) as a green solid. ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 7.99 (d, *J* = 9.2 Hz, 1H), 7.88 (s, 1H), 7.71 (s, 1H), 7.60 (d, *J* = 7.6 Hz, 1H), 7.54 (d, *J* = 9.2 Hz, 1H), 7.42 (d, *J* = 8.0 Hz, 1H), 3.99 (s, 3H), 3.83–3.80 (m, 4H), 3.61–3.59 (m, 2H), 3.39–3.33 (m, 2H); MS (ESI) *m/z* 465 [C₁₈H₁₇N₄O₃ + H]⁺.

Step 5: To a mixture of 4-(3-iodoimidazo[1,2-b]pyridazin-6-yl)-2-methoxyphenyl(morpholino)methanone (0.86 g, 1.85 mmol), 4-cyanophenylboronic acid (0.33 g, 2.22 mmol), K₃PO₄ (0.78 g, 3.7 mmol) in 1,4-dioxane (14 mL) and water (2 mL) was added Pd(PPh₃)₄ (106.8 mg, 0.09 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford 4-(6-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (0.35 g, 43.2 %, LC-MS 99%) as a light yellow solid.


Step 6: To a solution of 4-(6-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (0.25 g, 0.57 mmol) in CH₂Cl₂ (10 mL) cooled to 0 °C was added BBr₃ (2 mL). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of a saturated aqueous NaHCO₃ (10 mL) and extracted with CH₂Cl₂ (3 x 30 mL). The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent MeOH/CHCl₃ 3:97) followed by preparative TLC to afford 4-(6-(3-hydroxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (80 mg, 33%, AUC HPLC 98.2%) as a pale yellow solid, m.p: 149-153 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 9.86 (s, 1H), 8.39 (d, *J* = 8.4 Hz, 2 H), 8.14 (d, *J* = 9.2 Hz, 1H), 7.81 (d, *J* = 8.4 Hz, 2H), 7.61 (t, *J* = 10 Hz, 2H), 7.52 (d, *J* = 8.0, 1H), 7.42 (d, *J* = 8.2 Hz, 1H), 3.81 (bs, 8 H); MS (ESI) *m/z* 426 [C₂₄H₁₉N₅O₃ + H]⁺.

Example 78: 4-(6-(2-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile



To a solution of (4-bromo-3-methoxyphenyl)(4-methylpiperazin-1-yl)methanone (312 mg, 1.00 mmol) in DMF (10 mL) under inert atmosphere, were added 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi[1,3,2-dioxaborolane] (280 mg, 1.10 mmol), KOAc (295 mg, 3.00 mmol) and PdCl₂dppf (22 mg, 0.03 mmol). The resulting mixture was stirred at 100 °C for 2 h after which time the reaction mixture was diluted with DMF (8 mL) and water (2 mL) prior to the addition of 4-(6-chloroimidazo[1,2-a]pyrazin-3-yl)benzonitrile (100 mg, 0.39 mmol), K₂CO₃ (108 mg, 0.78 mmol) and additional PdCl₂dppf (57 mg, 0.08 mmol). The reaction mixture thus obtained was heated to reflux for 3 h and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent, ACN/H₂O/0.01% HCOOH) to afford 4-(6-(2-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (30 mg, AUC HPLC 97%) as yellow solid. ¹H NMR (400 MHz, CD₃OD) δ ppm 8.33 (s, 1H), 8.27 (d, *J* = 8.2 Hz, 2H), 8.18 (s, 1H), 8.01 (d, *J* = 9.5 Hz, 1H), 7.66–7.78 (m, 4H), 7.16 (d, *J* = 7.7 Hz, 1H), 3.65–4.00 (m, 7H), 2.83–3.01 (m, 4H), 2.57–2.67 (m, 3H); ¹³C NMR (100 MHz, CD₃OD) δ ppm 171.45, 167.58, 159.10, 152.89, 141.44, 139.44, 134.71, 134.21, 133.41, 132.29, 130.47, 129.04, 127.95, 127.76, 127.58, 125.35, 122.63, 120.58, 119.73, 111.91, 111.69, 56.66, 54.99, 44.92; MS (ESI) *m/z* 453 [C₂₆H₂₄N₆O₂+H]⁺.

Example 79: 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-nitrophenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile




Step 1: To a solution of 2-nitro-4-bromobenzoic acid (9.3 mmol) in DMF (6 mL) was added NMM (1.87 g, 18.6 mmol) followed by addition of HATU (5.3 g, 13.95 mmol) and the mixture was stirred at rt for 30 min. 1-Methylpiperazine (1.39 g, 13.95 mmol) was added and stirred at rt for an additional 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and

concentrated under reduced pressure to give ((4-bromo-2-methylphenyl)(4-methylpiperazin-1-yl)methanone (1.9 g, 70%, AUC HPLC 98%) as a brown solid. ^1H NMR (400 MHz, CDCl_3): δ 8.4 (s, 1 H), 8.01 (d, $J = 6.2$ Hz, 1 H), 7.4 (d, $J = 8.3$ Hz, 1 H) 3.8 (bs, 2 H), 2.3-2.5 (s, 7 H).


5

Step 2: To a solution of (4-bromo-2-nitrophenyl)(4-methylpiperazin-1-yl)methanone 6 (328 mg, 1.00 mmol) in DMF (10 mL) under inert atmosphere, was added 4,4,4',4',5,5,5'-octamethyl-2,2'-bi1,3,2-dioxaborolane (280 mg, 1.10 mmol), KOAc (295 mg, 3.00 mmol) and PdCl_2dppf (22 mg, 0.03 mmol). The resulting mixture was stirred at 100 °C for 2 h after which time, it was diluted with DMF (8 mL) and water (2 mL) prior to the addition of 4-(6-chloroimidazo[1,2-a]pyrazin-3-yl)benzonitrile 3 (200 mg, 0.78 mmol), K_2CO_3 (217 mg, 1.58 mmol) and additional PdCl_2dppf (114 mg, 0.08 mmol). The reaction mixture thus obtained was heat to reflux for 3 h and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /0.01% HCOOH) to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-nitrophenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (129 mg, 36% over 2 steps, AUC HPLC 97%) as yellow solid. ^1H NMR (400 MHz, CD_3OD) 8.94 (d, $J = 1.6$ Hz, 1H), 8.56 (dd, $J = 8.0, 1.63$ Hz, 1H), 8.43 (d, $J = 8.5$ Hz, 2H), 8.36 (s, 1H), 8.28 (d, $J = 9.5$ Hz, 1H), 8.02 (d, $J = 9.5$ Hz, 1H), 7.89 (d, $J = 8.5$ Hz, 2H), 7.72 (d, $J = 7.9$ Hz, 1H), 3.71-4.05 (m, 2H), 3.41 (s, 2H), 2.70 (t, $J = 5.0$ Hz, 2H), 2.53 (s, 2H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ ppm 168.22, 151.26, 147.63, 141.86, 139.15, 135.75, 134.46, 134.18, 133.78, 130.40, 128.68, 128.24, 127.63, 124.68, 119.67, 118.41, 112.39, 55.35, 54.99, 47.61, 45.88, 42.47; MS (ESI) m/z 468 $[\text{C}_{25}\text{H}_{21}\text{N}_7\text{O}_3 + \text{H}]^+$.

25 Example 80: N-(5-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide



Step 1: To a solution of 4-bromo-2-nitrobenzoic acid (2.5 g, 10.16 mmol) in DMF (15 mL) were added HATU (5.79 g, 15.24 mmol), N-methyl morpholine (2.05 g, 20.32 mmol) and morpholine (0.97 g, 11.18 mmol). The reaction mixture was stirred at 0 °C to room temperature for 16 h, then it was diluted with H₂O (30 mL) and extracted with EtOAc (3 × 100 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure to afford (4-bromo-2-nitrophenyl)(morpholino)methanone (4.2 g, 91.8%, LC-MS 69.8%). ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm) 8.40 (s, 1H), 8.08 (d, *J* = 8.4 Hz, 1H), 7.54 (d, *J* = 8.0 Hz, 1H), 3.67-3.51 (m, 4H), 3.51 (bs, 2H), 3.21 (bs, 2H); MS (ESI) *m/z* 316 [C₁₁H₁₁BrN₂O₄ + H]⁺.



Step 2: To a mixture of 6-chloroimidazo[1,2-*b*]pyridazine (1.5 g, 9.76 mmol), Bis(pinacolato)diboron (2.65 g, 10.45 mmol), KOAc (2.87 g, 29.29 mmol) in 1,4-dioxane (15 mL) was added PdCl₂dppf (214 mg, 0.29 mmol) and dppf (162.4 mg, 0.29 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-*b*]pyridazine (2.15 g, 90%) as a brown liquid which was used in next step without purification. MS (ESI) *m/z* 246 [C₁₂H₁₆BN₃O₂ + H]⁺.

Step 3: To a mixture of (4-bromo-2-nitrophenyl)(morpholino)methanone (2 g, 7.32 mmol), 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-*b*]pyridazine (2.15 g, 8.78 mmol), K₃PO₄ (3.1 g, 14.64 mmol) in 1,4-dioxane (30 mL) and water (4 mL) was added Pd(PPh₃)₄ (423 mg, 0.36 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford (4-(imidazo[1,2-*b*]pyridazin-6-yl)-2-nitrophenyl)(morpholino)methanone (1.25 g, 55.8 %, LC-MS 96.5%) as a light yellow solid. ¹H NMR: (400 MHz, DMSO-*d*₆) δ (ppm) 8.84 (s, 1H), 8.54 (d, *J* = 6.0 Hz, 1H), 8.46 (s, 1H), 8.32 (d, *J* = 9.6 Hz, 1H), 7.98 (d, *J* = 9.6 Hz, 1H), 7.89 (s, 1H), 7.77 (d, *J* = 8.0 Hz, 1H), 3.7 (bs, 4H), 3.55 (bs, 2H), 3.17 (bs, 2H); MS (ESI) *m/z* 354 [C₁₇H₁₅N₅O₄ + H]⁺.


- Step 4: To a solution of (4-(imidazo[1,2-b]pyridazin-6-yl)-2-nitrophenyl)(morpholino)methanone (1.24 g, 3.51 mmol) in ACN (18 mL) was added N-iodosuccinimide (0.95 g, 4.2 mmol). The reaction mixture was stirred at 80 °C for 2 h and was diluted with EtOAc and washed with water. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (4-(3-iodoimidazo[1,2-b]pyridazin-6-yl)-2-nitrophenyl)(morpholino)methanone (1.7 g, 94%, LC-MS 93.3%) as a green solid. ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 8.84 (s, 1H), 8.45 (d, *J* = 8.0 Hz, 1H), 8.08 (d, *J* = 9.6 Hz, 2H), 7.94 (s, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 3.85-3.66 (m, 6H), 3.29 (bs, 2H); MS (ESI) *m/z* 480 [C₁₇H₁₄N₅O₄ + H]⁺.
- Step 5: To a mixture of (4-(3-iodoimidazo[1,2-b]pyridazin-6-yl)-2-nitrophenyl)(morpholino)methanone (1.7 g, 3.54 mmol), 4-cyanophenylboronic acid (0.62 g, 4.25 mmol), K₃PO₄ (1.5 g, 7.08 mmol) in 1,4-dioxane (18 mL) and water (2.5 mL) was added Pd(PPh₃)₄ (204 mg, 0.17 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford 4-(6-(4-(morpholine-4-carbonyl)-3-nitrophenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (1.1 g, 68.7 %, LC-MS 91%) as a light yellow solid. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.80 (s, 1H), 8.67 (d, *J* = 8.0 Hz, 1H), 8.55-8.23 (m, 4H), 7.83 (d, *J* = 8.4 Hz, 2H), 7.67 (d, *J* = 9.6 Hz, 1H), 7.61 (d, *J* = 8.0, 1H), 3.86-3.67 (m, 4H), 3.66 (bs, 2H), 3.31 (t, *J* = 4.4 Hz, 2H); MS (ESI) *m/z* 455 [C₂₄H₁₈N₆O₄ + H]⁺.
- Step 6: To a solution of 4-(6-(4-(morpholine-4-carbonyl)-3-nitrophenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (0.6 g, 1.32 mmol) in ethanol (10 mL) was added SnCl₂·2H₂O (0.89 g, 3.96 mmol) and concentrated HCl (1 mL). The reaction mixture was stirred at 90 °C for 3 h. The solvent was evaporated under reduced pressure and basified with a saturated aq. NaHCO₃ and extracted with EtOAc (3 x 30 mL). The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford 4-(6-(3-amino-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (250 mg, 45%, LC-MS 77%) as a yellow solid.

Step 7: To a solution of 4-(6-(3-amino-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)benzonitrile (0.24 g, 0.56 mmol) in dichloromethane (10 mL) were added

TEA (0.15 mL, 1.13 mmol) and acetyl chloride (0.06 mL, 0.85 mmol). The reaction mixture was stirred for 2 h at room temperature. The reaction mixture was diluted with H₂O (20 mL), basified with saturated aqueous NaHCO₃ and extracted with CHCl₃ (3 × 50 mL). The combined organic layer was concentrated under reduced pressure to afford crude compound.

- 5 The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 97:3) to afford N-(5-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide (100 mg, 38%, AUC HPLC 98.09%) as a pale yellow solid, m.p: 226-229 °C. NMR: (400 MHz, CDCl₃) δ (ppm) 9.12 (d, J = 11.6 Hz, 2H), 8.39 (d, J = 8.4 Hz, 2H), 8.22 (s, 1H), 8.14 (d, J = 9.1 Hz, 1H), 7.84 (d, J = 8.0 Hz, 2H), 7.75 (d, J = 8.0 Hz, 1H), 7.66 (d, J = 9.2 Hz, 1H), 7.39 (d, J = 7.6 Hz, 1H), 3.76 (bs, 8H), 2.26 (s, 3H); MS (ESI) m/z 467 [C₂₆H₂₂N₆O₃ + H]⁺.

Example 81: N-(5-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)acetamide



- Step 1: To a solution of 2-amino-4-bromobenzoic acid (4 g, 18.51 mmol) in 1,4-dioxane (40 mL) was added bis(trichloromethyl)carbonate in 1,4-dioxane (1.81 g, 6.11 mmol) drop wise at 0 °C. The reaction mixture was stirred at 110 °C for 16 h. The solvent was evaporated under reduced pressure and washed with n-hexane to afford 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (4.3 g, 95.8% LC-MS 97%). ¹H NMR: (400 MHz, DMSO-*d*₆) δ (ppm) 11.8 (s, 1H), 7.80 (d, J = 8.4 Hz, 1H), 7.40 (d, J = 8.4 Hz, 2H); MS (ESI) m/z 243.02 [C₈H₄BrNO₃ + H]⁺.
- 20

Step 2: To a solution of 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (1 g, 4.13 mmol), N,N-dimethylpiperidin-4-amine (0.58 g, 4.54 mmol) in dry DMF (10 mL) was added DMAP (50.4 mg, 0.41 mmol) at rt. The reaction mixture was stirred for 16 h, then was diluted with H₂O (30 mL) and extracted with EtOAc (3 × 100 mL). The combined organic layer was dried over Na₂SO₄ and concentrated under reduced pressure to afford (2-amino-4-bromophenyl)(4-(dimethylamino)piperidin-1-yl)methanone (0.97 g, 72.4%, LC-MS 99.5%). ¹H NMR (400 MHz, CDCl₃) δ (ppm) 6.92 (d, *J* = 8.0 Hz, 1H), 6.88 (s, 1H), 6.83 (d, *J* = 6.8 Hz, 1H), 4.4 (s, 2H), 2.96 (bs, 2H), 2.39-2.37 (m, 1H), 2.36 (s, 6H), 1.89 (d, *J* = 11.6 Hz, 2H), 1.49-1.41 (m, 2H); MS (ESI) *m/z* 327 [C₁₄H₂₀BrN₃O + H]⁺.

10

Step 3: To a solution of (2-amino-4-bromophenyl)(4-(dimethylamino)piperidin-1-yl)methanone (0.5 g, 1.53 mmol) in dichloromethane (10 mL) were added triethylamine (0.63 mL, 4.59 mmol) and acetyl chloride (0.16 mL, 2.29 mmol). The reaction mixture was stirred for 3 h at rt. The reaction mixture was diluted with H₂O (20 mL), basified with saturated aqueous solution of NaHCO₃ and extracted with CHCl₃ (3 × 50 mL). The combined organic layer was concentrated under reduced pressure to afford N-(5-bromo-2-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)acetamide (0.6 g, 88.23%, LC-MS 83%). ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 9.01 (s, 1H), 8.53 (s, 1H), 7.42 (d, *J* = 8.4 Hz, 1H), 7.06 (d, *J* = 7.6 Hz, 1H), 4.6 (bs, 1H), 2.95 (bs, 3H), 2.41 (m, 1H), 2.32 (s, 6H), 2.08 (s, 3H), 1.95 (bs, 2H), 1.52 (bs, 2H); MS (ESI) *m/z* 369.07 [C₁₆H₂₂BrN₃O₂ + H]⁺.

20

Step 4: To a mixture of N-(5-bromo-2-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)acetamide (0.52 g, 1.42 mmol), 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-b]pyridazine (416.3 mg, 1.69 mmol), K₃PO₄ (602 mg, 2.84 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added Pd(PPh₃)₄ (82 mg, 0.07 mmol) and the reaction mixture was heated at 90 °C for 6 h. The reaction mixture was filtered through celite bed and washed with 10% MeOH in CHCl₃ and dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH/NH₃ 90:10:1) to afford N-(2-((4-(dimethylamino)piperidin-1-yl)carbonyl)-5-(imidazo(1,2-a)pyridazin-6-yl)phenyl)acetamide (0.5 g, 70 %, LC-MS 81%) as a light yellow solid. NMR: (400 MHz, CDCl₃) δ (ppm) 9.07 (s, 1H), 8.91 (s, 1H), 8.02 (d, *J* = 8.8 Hz, 2H), 7.76 (s, 1H), 7.75 (d, *J* = 6.4 Hz, 1H), 7.42 (d, *J* = 1.6 Hz, 1H), 4.75 (bs, 1H), 3.95 (bs, 1H), 3.15-2.95 (m, 2H), 2.41 (m, 1H), 2.25 (s, 6H), 2.19 (s, 3H), 1.98 (bs, 1H), 1.75 (bs, 2H), 1.25 (s, 3H); MS (ESI) *m/z* 406.16 [C₂₃H₂₇N₅O₂ + H]⁺.


30

Step 5: To a solution of N-(2-(4-(dimethylamino)piperidine-1-carbonyl)-5-(imidazo[1,2-a]pyridin-6-yl)phenyl)acetamide (0.49 g, 1.21 mmol) in ACN (10 mL) was added N-iodosuccinimide (0.32 g, 1.45 mmol) and stirred at 80 °C for 3 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford N-(2-((4-(dimethylamino)piperidin-1-yl)carbonyl)-5-(3-iodoimidazo[1,2-a]pyridazin-6-yl)phenyl)acetamide (0.3 g, 46.7%) as a green solid. The crude product was used in the next step without purification.

10

Step 6: To a mixture of N-(2-(4-(dimethylamino)piperidine-1-carbonyl)-5-(3-iodoimidazo[1,2-a]pyridin-6-yl)phenyl)acetamide (0.3 g, 0.56 mmol), 4-cyanophenylboronic acid (99.4 mg, 0.67 mmol), K₃PO₄ (237 mg, 101 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added Pd(PPh₃)₄ (32.3 mg, 0.03 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was filtered through celite and washed with 10% MeOH in CHCl₃ and dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by preparative HPLC to afford N-(5-(3-(4-cyanophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)acetamide (50 mg, 17.4 %, LC-MS 95%) as a yellow solid. m.p: 141-146 °C. ¹H NMR: (400 MHz, CDCl₃) δ (ppm) 9.13 (s, 1H), 9.08 (s, 1H), 8.39 (d, *J* = 8.4 Hz, 2H), 8.22 (s, 1H), 8.13 (d, *J* = 9.2 Hz, 1H), 7.85 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 7.6, 1H), 6.67 (d, *J* = 9.6 Hz, 1H), 7.40 (d, *J* = 8.4 Hz, 1H), 4.75 (bs, 1H), 3.95 (bs, 1H), 3.15-2.95 (m, 2H), 2.41 (m, 1H), 2.3 (s, 6H), 2.21 (s, 3H), 1.95 (bs, 2H), 1.65 (bs, 2H); MS (ESI) *m/z* 508.16 [C₂₉H₂₉N₇O₂ + H]⁺.

25 Example 82: (4-(3-(4-chlorophenyl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




Step 1: To a solution of 3-bromo-6-chloroimidazo[1,2-b]pyridazine (500 mg, 2.15 mmol) in 1,4-dioxane (4.5 mL) under inert atmosphere were added aqueous Na₂CO₃ solution (3.76 mL, 2 M), 4-chlorophenylboronic acid (336 mg, 2.15 mmol) and Pd(dppf)₂Cl₂ (124 mg, 0.108 mmol). The resulting mixture was heated to reflux for 18 h, and then diluted with water (20

30

mL) and extracted with DCM (3×30 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 6-chloro-3-(4-chlorophenyl)imidazo[1,2-*b*]pyridazine (243 mg, 43%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.06 (s, 1H), 8.01-7.96 (m, 3H), 7.50-7.47 (m, 2H), 7.11 (d, *J* = 12 Hz, 1H); MS (ESI) *m/z* 264 [C₁₂H₇Cl₂N₃ + H]⁺.

Step 2: To a solution of 6-chloro-3-(4-chlorophenyl)imidazo[1,2-*b*]pyridazine (200 mg, 0.757 mmol) in toluene (3 mL) and ethanol (1.5 mL) under inert atmosphere were added K₂CO₃ (335 mg, 2.4 mmol), 4-(4-methylpiperazine-1-carbonyl)phenylboronic acid hydrochloride (259 mg, 0.909 mmol) and Pd(PPh₃)₄ (88 mg, 0.076 mmol). The resulting mixture was heated in a microwave reactor at 140 °C for 15 min, and then was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford (4-(3-(4-chlorophenyl)imidazo[1,2-*b*]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (41 mg, 17%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.19 (d, *J* = 9.5 Hz, 1H), 8.11-8.06 (m, 5H), 7.62-7.58 (m, 3H), 7.54-7.51 (m, 2H), 3.94-3.67 (m, 4H), 2.72-2.66 (m, 4H), 2.49 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 169.59, 165.33, 150.94, 139.36, 137.12, 136.71, 133.88, 132.95, 129.00, 128.02, 127.98, 127.90, 127.36, 126.89, 126.24, 115.76, 54.12, 44.83; MS (ESI) *m/z* 432 [C₂₄H₂₂ClN₅O + H]⁺.

Example 83: (4-(3-(4-chlorophenyl)imidazo[1,2-*b*]pyridazin-6-yl)-2-methylphenyl)(4-methylpiperazin-1-yl)methanone




Step 1: To a solution of 6-chloro-3-(4-chlorophenyl)imidazo[1,2-*b*]pyridazine (308 mg, 1.00 mmol) in a mixture of DMF (4 mL) and water (1 mL), were successively added K₂CO₃ (138 mg, 1.00 mmol), Pd(dppf)Cl₂ (146 mg, 0.20 mmol) and 4-borono-2-methylbenzoic acid (198 mg, 1.10 mmol). The reaction mixture was stirred at 80 °C for 18 h under inert atmosphere, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under vacuum to afford the

crude 4-(3-(4-chlorophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-methylbenzoic acid (724 mg) as black solid. MS (ESI) m/z 364 [$C_{20}H_{14}ClN_3O_2 + H$] $^+$.


Step 2: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-methylbenzoic acid (363 mg, ~1.00 mmol) in DMF (5 mL), was successively added HATU (1.14 g, 3.00 mmol), *N*-methyl morpholine (400 mg, 4.00 mmol) and 1-methylpiperazine (200 mg, 2.00 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /0.01% $HCOOH$) to afford (4-(3-(4-chlorophenyl)imidazo[1,2-b]pyridazin-6-yl)-2-methylphenyl)(4-methylpiperazin-1-yl)methanone (24 mg, 5%, AUC HPLC 99%) as yellow solid. 1H NMR (400 MHz, CD_3OD) δ 8.05–7.93 (m, 4H), 7.89–7.82 (m, 2H), 7.67 (d, J = 9.5 Hz, 1H), 7.39 (d, J = 8.6 Hz, 3H), 4.18–3.83 (m, 2H), 3.50 (s, 2H), 3.08 (s, 2H), 2.91 (s, 2H), 2.70 (s, 3H), 2.39 (s, 3H); ^{13}C -NMR (100 MHz, CD_3OD) δ 171.40, 166.86, 152.68, 140.84, 138.16, 137.71, 136.82, 134.77, 133.56, 130.39, 129.87, 129.14, 128.97, 128.33, 127.93, 126.71, 126.03, 117.84, 55.26, 54.85, 46.54, 44.94, 41.19, 19.34; MS (ESI) m/z 446 [$C_{25}H_{24}ClN_6O + H$] $^+$.

Example 84: (4-(3-(1H-indol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




Step 1: To a solution of 4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid (100 mg, 0.314 mmol) in DMF (1.5 mL) was added HATU (179 mg, 0.471 mmol) and *N*-methyl morpholine (138 μ L, 1.257 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 1-methylpiperazine (70 μ L, 0.628 mmol). The reaction mixture was left to stir for 18 h, then it was diluted with H_2O (25 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) to afford the product as a yellow

solid (127 mg, 99%): ^1H NMR (400 MHz, CDCl_3) δ 8.09 (d, $J = 8.0$ Hz, 2H), 8.02 (d, $J = 9.2$ Hz, 1H), 7.81 (s, 1H), 7.59-7.54 (m, 3H), 3.84 (bs, 2H), 3.49 (bs, 3H), 2.50-2.34 (m, 6H); MS (ESI) m/z 400 [$\text{C}_{18}\text{H}_{18}\text{BrN}_5\text{O} + \text{H}$] $^+$.



- 5 Step 2: (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (100 mg, 0.250 mmol), 1H-indol-5-ylboronic acid (48 mg, 0.300 mmol), $\text{Pd}(\text{PPh}_3)_4$ (29 mg, 0.025 mmol) and Cs_2CO_3 (163 mg, 0.50 mmol) was dissolved in dioxane (2.0 mL) and water (0.5 mL). The resulting mixture was heated in a microwave reactor for 30 min at 110°C , then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic
- 10 layer was filtered through celite, dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford (4-(3-(1H-indol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (51 mg, 47%, AUC HPLC 97%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 8.45 (s, 1H), 8.32 (bs, 1H), 8.10-8.07 (m, 4H), 7.91 (d, $J = 8.4$ Hz, 1H),
- 15 7.57-7.50 (m, 4H), 7.29 (t, $J = 2.8$ Hz, 1H), 6.67 (s, 1H), 3.83 (bs, 2H), 3.49 (bs, 2H), 2.50-2.33 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3): 169.67, 150.44, 138.93, 137.35, 137.10, 135.61, 133.08, 130.37, 128.11, 127.80, 127.27, 126.09, 125.00, 121.77, 120.35, 119.87, 114.56, 111.32, 103.30, 46.06; MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}$] $^+$.


- 20 Example 85: 5-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)indolin-2-one



- (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (100 mg, 0.250 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (78 mg, 0.300 mmol), $\text{Pd}(\text{PPh}_3)_4$ (29 mg, 0.025 mmol) and Cs_2CO_3 (163 mg, 0.500 mmol) was
- 25


dissolved in dioxane (2.0 mL) and water (0.5 mL). The reaction mixture was heated in a microwave reactor for 30 min at 110°C, then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was filtered through celite, dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford 5-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)indolin-2-one (33 mg, 29%, AUC HPLC 98%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.12 (d, *J* = 9.2 Hz, 1H), 8.06-8.04 (m, 4H), 7.99 (s, 1H), 7.60 (d, *J* = 8.4 Hz, 3H), 7.55 (d, *J* = 9.6 Hz, 1H), 7.04 (d, *J* = 8.0 Hz, 1H), 3.85 (bs, 2H), 3.68 (s, 2H), 3.52 (bs, 2H), 2.53-2.36 (m, 7H); ¹³C NMR (100 MHz, CDCl₃) δ 176.79, 169.52, 150.82, 142.26, 139.19, 137.34, 137.08, 132.93, 128.78, 127.89, 127.25, 127.00, 126.29, 125.78, 123.54, 123.03, 115.13, 109.71, 46.04, 36.22; MS (ESI) *m/z* 453 [C₂₆H₂₄N₆O₂ + H]⁺.

Example 86: (4-(3-(1H-indazol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




(4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (100 mg, 0.250 mmol), 1H-indazol-5-ylboronic acid (49 mg, 0.300 mmol), Pd(PPh₃)₄ (29 mg, 0.025 mmol) and Cs₂CO₃ (163 mg, 0.500 mmol) was dissolved in dioxane (2.0 mL) and water (0.5 mL). The reaction mixture was heated in a microwave reactor for 30 min at 110 °C, then it was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was filtered through celite, dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford (4-(3-(1H-indazol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (21 mg, 19%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.62 (s, 1H), δ 8.20 (s, 1H), 8.13-8.05 (m, 5H), 7.66 (d, *J* = 8.8 Hz, 1H), 7.59-7.53 (m, 3H), 3.84 (bs, 2H), 3.50 (bs, 2H), 2.51-2.34 (m, 7H); MS (ESI) *m/z* 438 [C₂₅H₂₃N₇O + H]⁺.

Example 87: (4-(3-(1H-benzo[d]imidazol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



(4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (50
 5 mg, 0.125 mmol), 1H-benzo[d]imidazol-5-ylboronic acid (22 mg, 0.137 mmol) and
 Pd(PPh₃)₄ (22 mg, 0.019 mmol) was dissolved in dioxane (0.6 mL) and saturated NaHCO₃
 solution (0.6 mL) was added. The reaction mixture was heated at 150 °C in a microwave
 reactor for 10 min, then was diluted with water (25 mL) and extracted with EtOAc (3x25
 mL). The combined organic layer was filtered over celite, dried over Na₂SO₄ and was
 10 concentrated under reduced pressure. The crude residue was purified by preparative HPLC
 (C18, eluent ACN, water, formic acid 0.1%) to afford (4-(3-(1H-benzo[d]imidazol-5-
 yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone (15 mg, 27%,
 AUC HPLC 95%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.55 (s, 1H), 8.18-8.15
 (m, 2H), 8.12 (t, *J* = 4.8 Hz, 2H), 8.03 (d, *J* = 8.0 Hz, 2H), 7.95-7.92 (m, 1H), 7.82-7.80 (m,
 15 1H), 7.56-7.49 (m, 3H), 3.89 (bs, 2H), 3.55 (bs, 2H), 2.60-2.50 (m, 4H), 2.39 (s, 3H); MS
 (ESI) *m/z* 438 [C₂₅H₂₃N₇O + H]⁺.

Example 88 (4-(3-(1H-benzo[d][1,2,3]triazol-5-yl)imidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone




20 A mixture of (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-methylpiperazin-1-
 yl)methanone (53 mg, 0.132 mmol), 1H-benzo[d][1,2,3]triazol-5-ylboronic acid (24 mg,
 0.146 mmol) and Tetrakis (23 mg, 0.020 mmol) in 1,4-dioxane (0.65 mL) and saturated
 NaHCO₃ solution (0.65 mL), was heated in a microwave reactor for 10 min at 150 °C, then it
 25 was diluted with H₂O (25 mL) and washed with EtOAc (3x25 mL). The crude product

remained in the aqueous layer, hence it was concentrated under reduced pressure, dissolved in methanol and filtered over celite and the filtrate was concentrated under reduced pressure.

The residue was then purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford (4-(3-(1H-benzo[d][1,2,3]triazol-5-yl)imidazo[1,2-b]pyridazin-6-

- 5 yl)phenyl)(4-methylpiperazin-1-yl)methanone (13 mg, 23%, AUC HPLC 98%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.91 (s, 1H), 8.21-8.14 (m, 3H), 8.06-8.03 (m, 4H), 7.61 (d, *J* = 8.4 Hz, 2H), 7.55 (d, *J* = 8.8 Hz, 1H), 3.95 (bs, 2H), 3.63 (bs, 2H), 2.68-2.60 (m, 4H), 2.43 (s, 3H); MS (ESI) *m/z* 439 [C₂₄H₂₂N₈O + H]⁺.

10 Example 89: 5-(6-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)indolin-2-one



- Step 1: To a solution of 4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid (200 mg, 0.589 mmol) in DMF (3.0 mL) was added HATU (336 mg, 0.884 mmol) and *N*-methyl morpholine (259 μl, 2.357 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 1-methyl-4-(piperidin-4-yl)piperazine (162 mg, 0.884 mmol). The reaction mixture was stirred for 18 h then, was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue
- 20 was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH/NH₄OH 94:5:1) to afford a light yellow solid (128 mg, 45%). ¹H NMR (400 MHz, CD₃OD) δ 8.25 (d, *J* = 8.4 Hz, 2H), 8.14 (d, *J* = 9.6 Hz, 1H), 7.92 (d, *J* = 9.6 Hz, 1H), 7.86 (s, 1H), 7.63 (d, *J* = 8.4 Hz, 2H), 4.73 (bs, 1H), 4.58 (bs, 1H), 3.85 (bs, 1H), 3.15 (bs, 1H), 2.93 (bs, 1H), 2.68-2.57 (m, 8H), 2.32 (s, 3H), 2.09-1.92 (m, 2H), 1.51 (bs, 2H); MS (ESI) *m/z* 483 [C₂₃H₂₇BrN₆O + H]⁺.


25

- Step 2: (4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)phenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone (125 mg, 0.258 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (81 mg, 0.310 mmol), Pd(PPh₃)₄ (30 mg, 0.026 mmol) and Cs₂CO₃ (169 mg, 0.517 mmol) was dissolved in dioxane (2.0 mL) and water (0.5 mL). The
- 30 reaction mixture was heated in a microwave reactor for 30min at 110°C, diluted with water

(25 mL) and washed with EtOAc (3x25 mL). The aqueous layer was concentrated under reduced pressure. The crude residue was dissolved in dichloromethane and filtered through a pad of celite. The filtrate was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 5-(6-(4-(4-(4-methylpiperazin-1-yl)piperidine-1-

- 5 carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)indolin-2-one (15 mg, 11%, AUC HPLC 98%) as a yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 8.21 (d, J = 8.4 Hz, 2H), 8.16 (d, J = 9.6 Hz, 1H), 8.10-8.07 (m, 3H), 7.86 (d, J = 9.6 Hz, 1H), 7.63 (d, J = 8.4 Hz, 2H), 7.08 (d, J = 8.0 Hz, 1H), 4.69-4.58 (m, 2H), 3.86 (bs, 1H), 3.32-3.20 (m, 3H), 2.94-2.62 (m, 9H), 2.45 (s, 3H), 2.05-1.89 (m, 2H), 1.55 (bs, 2H); MS (ESI) m/z 536 [$\text{C}_{31}\text{H}_{33}\text{N}_7\text{O}_2 + \text{H}$] $^+$.

- 10 Example 90: 5-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-b]pyridazin-3-yl)indolin-2-one




- Step 1: *tert*-butyl 9-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-b]pyridazin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate was prepared following General procedure A using 4-
15 (3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoic acid and *tert*-butyl 1,9-diazaspiro[5.5]undecane-1-carboxylate as starting materials. Purification by column chromatography (silica gel, eluent EtOAc/Hexane 60:40) to afford a light yellow solid (330 mg). MS (ESI) m/z 554 [$\text{C}_{27}\text{H}_{32}\text{BrN}_5\text{O}_3 + \text{H}$] $^+$.

- 20 Step 2: *tert*-butyl 9-(4-(3-bromoimidazo[1,2-b]pyridazin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (340 mg, 0.521 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (162 mg, 0.625 mmol), $\text{Pd}(\text{PPh}_3)_4$ (60 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.042 mmol) was dissolved in dioxane (4.0 mL) and water (1.0 mL). The reaction mixture was heated in a microwave reactor for 30 min at 110 $^\circ\text{C}$, then it was diluted
25 with H_2O (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was filtered over celite, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford a yellow solid (144 mg). MS (ESI) m/z 607 [$\text{C}_{35}\text{H}_{38}\text{N}_6\text{O}_4 + \text{H}$] $^+$.

Step 3: To a solution of *tert*-butyl 9-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (142 mg, 0.234 mmol) in DCM (3.0 mL) was added trifluoroacetic acid (1.0 mL). The reaction mixture was stirred at room temperature under inert atmosphere for 30 min, then it was neutralized with NaHCO₃, and diluted with water (25 mL) and washed with EtOAc (3x25 mL). The crude product remained in the aqueous layer; hence it was concentrated under reduced pressure. The crude residue was dissolved in dichloromethane, filtered, and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 5-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)indolin-2-one (85 mg, 79%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.23 (d, *J* = 8.4 Hz, 2H), 8.17 (d, *J* = 9.6 Hz, 1H), 8.09-8.06 (m, 3H), 7.86 (d, *J* = 9.6 Hz, 1H), 7.64 (d, *J* = 8.4 Hz, 2H), 7.08 (d, *J* = 8.0 Hz, 1H), 4.57-4.38 (m, 2H), 3.73-3.44 (m, 4H), 3.23 (bs, 2H), 2.15-1.79 (m, 10H); ¹³C NMR (400 MHz, CD₃OD) δ 179.83, 171.80, 152.36, 144.81, 140.39, 138.61, 138.07, 132.82, 130.46, 128.81, 128.58, 128.01, 127.56, 126.70, 124.44, 123.70, 117.22, 110.84, 56.81, 40.76, 30.79, 23.56, 18.69; MS (ESI) *m/z* 507 [C₃₀H₃₀N₆O₂ + H]⁺.


Example 91: 5-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)isoindolin-1-one



Step 1: To a solution of 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (200 mg, 0.63 mmol) in DMF (5 mL) were added HATU (479 mg, 1.26 mmol), *N*-methyl morpholine (191 mg, 1.89 mmol) and *N,N*-dimethylpiperidin-4-amine (242 mg, 1.89 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3x20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford (4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)phenyl)(4-(dimethylamino)piperidin-1-yl)methanone as a yellow solid. MS (ESI) *m/z* 428 [C₂₀H₂₂BrN₅O + H]⁺.

Step 2: To a solution of *tert*-butyl 4-(4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate (0.63 mmol) in DMF (4 mL) and water (0.8 mL) under inert atmosphere were added Cs₂CO₃ (411 mg, 1.26 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (245 mg, 0.945 mmol) and Pd(dppf)₂Cl₂ (92 mg, 0.126 mmol). The resulting mixture was stirred and heated to 90 °C for 18 h, and then was diluted with water (20 mL) and extracted with DCM (3×40 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 5-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)isoindolin-1-one (66 mg, 22%, AUC HPLC 96%) as yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.56(s, 1H), 8.30-8.02 (m, 6H), 7.88-7.86 (m, 1H), 7.60-7.58 (m, 2H), 7.02-7.00 (m, 1H), 4.47 (bs, 1H), 3.62 (bs, 3H), 3.06 (bs, 1H), 2.83 (bs, 1H), 2.35 (bs, 1H), 2.18 (s, 6H), 1.84-1.71 (m, 2H), 1.37 (bs, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 176.28, 168.12, 150.21, 143.46, 138.61, 137.72, 135.92, 132.61, 128.02, 127.43, 126.97, 126.28, 126.24, 125.99, 122.76, 121.43, 115.21, 109.19, 61.09, 54.80, 41.32, 35.81; MS (ESI) *m/z* 481 [C₂₈H₂₈N₆O₂ + H]⁺.

Example 92: 5-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)indolin-2-one




Step 1: To a solution of 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (200 mg, 0.63 mmol) in DMF (5 mL), were added HATU (479 mg, 1.26 mmol), *N*-methyl morpholine (191 mg, 1.89 mmol) and *tert*-butyl piperazine-1-carboxylate (352 mg, 1.89 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 4-(4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate as a yellow solid. MS (ESI) *m/z* 486 [C₂₂H₂₄BrN₅O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 4-(4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate (0.63 mmol) in DMF (4 mL) and water (0.8 mL) under inert atmosphere were added Cs₂CO₃ (411 mg, 1.26 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (245 mg, 0.945 mmol) and Pd(dppf)₂Cl₂ (92 mg, 0.126 mmol). The resulting mixture was stirred and heated at 90 °C for 18 h, and then was diluted with water (20 mL) and extracted with DCM (3×40 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 4-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate. MS (ESI) *m/z* 539 [C₃₀H₃₀N₆O₄ + H]⁺

Step 3: To a solution of *tert*-butyl 4-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)piperazine-1-carboxylate (0.63 mmol) in DCM (1 mL) was added TFA (1 mL). The resulting mixture was stirred at room temperature for 18 h and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) and preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 5-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)indolin-2-one (71 mg, 26%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.56 (s, 1H), 8.29 (d, *J* = 9.6 Hz, 1H), 8.19-8.17 (m, 3H), 8.09 (d, *J* = 8.0 Hz, 1H), 8.02 (s, 1H), 7.87 (d, *J* = 9.5 Hz, 1H), 7.58 (d, *J* = 7.7 Hz, 2H), 7.01 (d, *J* = 8.0 Hz, 1H), 3.62-3.51 (m, 7H), 2.75-2.69 (m, 4H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 176.29, 168.24, 150.21, 143.46, 138.61, 137.56, 135.92, 132.60, 128.02, 127.60, 126.97, 126.28, 126.24, 126.00, 122.76, 121.43, 115.20, 109.20, 54.79, 45.71, 35.81; MS (ESI) *m/z* 439 [C₂₅H₂₂N₆O₂ + H]⁺.

Example 93: 5-(6-(4-(1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)isoindolin-1-one




Step 1: To a solution of 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (200 mg, 0.63 mmol) in DMF (5 mL) were added HATU (479 mg, 1.26 mmol), *N*-methyl morpholine (191

mg, 1.89 mmol) and *tert*-butyl 1,4-diazepane-1-carboxylate (379 mg, 1.89 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 4-(4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate as a yellow solid. MS (ESI) *m/z* 500 [C₂₃H₂₆BrN₅O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 4-(4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate (0.63 mmol) in DMF (4 mL) and water (0.8 mL) under inert atmosphere were added Cs₂CO₃ (411 mg, 1.26 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (245 mg, 0.945 mmol) and Pd(dppf)₂Cl₂ (92 mg, 0.126 mmol). The resulting mixture was stirred and heated to 90 °C for 18 h, and then was diluted with water (20 mL) and extracted with DCM (3×40 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford *tert*-butyl 4-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate 5 (310 mg, 48%) as a brown solid. MS (ESI) *m/z* 553 [C₃₁H₃₂N₆O₄ + H]⁺.



Step 3: A solution of *tert*-butyl 4-(4-(3-(2-oxoindolin-5-yl)imidazo[1,2-*b*]pyridazin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate (0.63 mmol) in a mixture of DCM (1 mL) and TFA (1 mL), was stirred at room temperature for 18 h and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) and preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 5-(6-(4-(1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)isoindolin-1-one (46 mg, 16%, AUC HPLC 98%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.55 (s, 1H), 8.29 (d, *J* = 9.5 Hz, 1H), 8.19-8.17 (m, 3H), 8.09 (d, *J* = 8.2 Hz, 1H), 8.03 (s, 1H), 7.87 (d, *J* = 9.4 Hz, 1H), 7.58 (d, *J* = 8.0 Hz, 2H), 7.01 (d, *J* = 8.2 Hz, 1H), 3.68-3.62 (m, 4H), 3.41-3.38 (m, 2H), 3.17 (d, *J* = 5.2 Hz, 1H), 2.91-2.85 (m, 1H), 2.81-2.74 (m, 3H), 1.78-1.72 (m, 1H), 1.63-1.56 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) 176.28, 169.60, 150.24, 143.46, 138.61, 135.56, 128.02, 127.27, 127.01, 126.92, 126.28, 126.24, 126.00, 122.76, 121.45, 115.20, 109.20, 48.49, 44.53, 35.80; MS (ESI) *m/z* 453 [C₂₆H₂₄N₆O₂ + H]⁺.

Example 94: 5-(6-(4-(4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)indolin-2-one



Step 1: To a solution of 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)benzoic acid (200 mg, 0.63 mmol) in DMF (5 mL) were added HATU (479 mg, 1.26 mmol), *N*-methyl morpholine (191 mg, 1.89 mmol) and 4-(piperidin-4-yl)morpholine (321 mg, 1.89 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, and then was diluted with water (10 mL) and extracted with DCM (3×20 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)phenyl(4-morpholinopiperidin-1-yl)methanone as a yellow solid. MS (ESI) *m/z* 470 [C₂₂H₂₄BrN₅O₂ + H]⁺.

Step 2: To a solution of 4-(3-bromoimidazo[1,2-*b*]pyridazin-6-yl)phenyl(4-morpholinopiperidin-1-yl)methanone (0.63 mmol) in DMF (4 mL) and water (0.8 mL) under inert atmosphere were added Cs₂CO₃ (411 mg, 1.26 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (245 mg, 0.945 mmol) and Pd(dppf)₂Cl₂ (92 mg, 0.126 mmol). The resulting mixture was heated to 90 °C for 18 h, and then was diluted with water (20 mL) and extracted with DCM (3×40 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to 90:10) to afford 5-(6-(4-(4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-*b*]pyridazin-3-yl)indolin-2-one (16.2 mg, 5%, AUC HPLC 98%) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.56 (s, 1H), 8.29 (d, *J* = 9.5 Hz, 1H), 8.19-8.17 (m, 3H), 8.09 (d, *J* = 8.0 Hz, 1H), 8.02 (s, 1H), 7.88 (d, *J* = 9.6 Hz, 1H), 7.59 (d, *J* = 8.2 Hz, 2H), 7.01 (d, *J* = 8.1 Hz, 1H), 4.48-4.46 (m, 1H), 3.62-3.56 (m, 7H), 3.08 (bs, 1H), 2.84 (bs, 1H), 2.44-2.41 (m, 5H), 1.91-1.76 (m, 2H), 1.05-1.03 (m, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 176.85, 168.69, 150.78, 144.04, 139.19, 138.27, 136.50, 133.18, 128.59, 128.00, 127.54, 126.85, 126.82, 126.57, 123.33, 122.01, 115.78, 109.76, 67.03, 61.41, 49.87, 36.38; MS (ESI) *m/z* 523 [C₃₀H₃₀N₆O₃ + H]⁺.


**Formula 4****Intermediate 14: Synthesis of 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyridine**

- 5 Step 1: To a solution of 6-bromo-3-iodoimidazo[1,2-a]pyridine (21 g, 65.4 mmol), 4-Chlorophenylboronic acid (11.22 g, 71.9 mmol), K_3PO_4 (27.7 g, 130.8 mmol) in a mixture of DMF (100 mL) and water (15 mL), was added $Pd(PPh_3)_4$ (3.77 g, 3.27 mmol). The reaction mixture was heated at 90 °C for 6 h under argon atmosphere prior to the addition of water (200 mL). The precipitate was isolated by filtration and was purified by column
- 10 chromatography (silica gel, eluent $CHCl_3/MeOH$ 95:5) to afford of 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyridine (10.6 g, 52%) as a yellowish green solid. 1H NMR (400 MHz, $CDCl_3$) δ 8.38 (s, 1H), 7.69 (s, 1H), 7.60 (d, $J = 8.0$ Hz, 2H), 7.82 (t, $J = 9.2$ Hz, 1H), 7.58 (q, $J = 5.2$ Hz, 4H), MS (ESI) m/z 309 [$C_{13}H_8BrN_2Cl + 2H$] $^+$.
- 15 Step 2: 4-(ethoxycarbonyl) phenylboronic acid (3.81 g, 19.67 mmol), K_3PO_4 (6.95 g, 32.78 mmol) and $Pd(PPh_3)_4$ (0.56 g, 0.49 mmol) were added sequentially to a solution of ethyl 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (5 g, 16.39 mmol) in a mixture of 1,4-dioxane (50 mL) and water (5 mL) at room temperature. The reaction mixture was refluxed for 6 h under argon atmosphere, and was diluted with water (100 mL) and extracted with
- 20 EtOAc (3x100 mL). The combined organic layer was dried over Na_2SO_4 , filtered and

concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 1:1) to afford ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (2.7 mg, 45%) as an off-white solid. ^1H NMR (400 MHz, CD_3OD) δ 9.21 (s, 1 H), 8.54 (s, 1 H), 8.16 (d, $J = 8.4$ Hz, 2 H), 7.85–7.44 (m, 9 H), 4.44 (q, $J = 7.2$ Hz, 2 H), 1.42 (t, $J = 7.2$ Hz, 3 H); MS (ESI) m/z 369 $[\text{M} + \text{H}]^+$.

Step 3: To a solution of ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (2.7 g, 7.18 mmol) in THF (40 mL) was added a solution of LiOH (0.9 g, 21.54 mmol) in water (10 mL) and MeOH (10 mL). The reaction mixture was stirred for 5 h at rt and was concentrated under reduced pressure to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (900 mg, 37%) of as an off white solid which was used in the next step without purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.88 (s, 1 H), 8.13–8.03 (m, 6 H), 7.92–7.90 (m, 4 H); MS (ESI) m/z 340 $[\text{C}_{21}\text{H}_{13}\text{N}_3\text{O}_2 + \text{H}]^+$.


Example 95: 4-(6-(4-(4-Methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (210 mg, 0.9 mmol), (4-methylpiperazin-1-yl)(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (388 mg, 1.17 mmol), NaHCO_3 (228 mg, 2.71 mmol) in a mixture of DMF (11 mL) and water (2 mL) was added $(\text{A-Phos})_2\text{PdCl}_2$ (31 mg, 0.04 mmol). The reaction mixture was heated at 90°C for 1 h under argon atmosphere then, was diluted with water and extracted with EtOAc. The organic phase was washed with water and brine solution, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by Preparative TLC to afford of 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (130 mg, 45%, AUC HPLC purity 98.4%) as a pale yellow solid; m.p. $106\text{--}116^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.50 (s, 1 H), 7.85–7.79 (m, 4 H), 7.74 (d, $J = 8.0$ Hz, 2 H), 7.60–7.52 (m, 5 H), 3.83 (bs, 2 H), 3.50 (bs, 2 H), 2.51 (bs, 2 H), 2.37 (bs, 2 H), 2.34 (s, 3 H). ^{13}C NMR (75 MHz, CDCl_3) δ (ppm): 169.5, 146.3, 138.5, 135.6,

134.8, 133.8, 133.2, 128.1, 127.8, 127.2, 127.1, 125.6, 124.5, 120.6, 118.7, 118.4, 111.5, 55.3, 47.7, 46.0; MS (ESI) m/z 422.30 [$C_{26}H_{23}N_3O + H$] $^+$.


Example 96: 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



5

To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (0.2 g, 0.58 mmol) in DMF (5 mL) were added HATU (0.33 g, 0.88 mmol), *N*-methyl morpholine (0.18 g, 1.76 mmol) and morpholine (60 mg, 0.70 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h then, was diluted with water (15 mL) and
 10 extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na_2SO_4 , filtered concentrated under reduced pressure. The crude residue was first purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) and by Prep-HPLC to afford 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (39 mg, 16.2%, AUC HPLC >99) as an off white solid; m.p. 117-130 °C. 1H NMR (400 MHz, $CDCl_3$) δ
 15 (ppm): 8.49 (s, 1 H), 7.84-7.71 (m, 5 H), 7.60-7.50 (m, 5 H), 3.72 (m, 8 H); MS (ESI) m/z 409.19 [$C_{25}H_{20}N_4O_2 + H$] $^+$.

Example 97: 4-(6-(4-(1,4-oxazepane-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




20

To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (0.2 g, 0.58 mmol) in DMF (5 mL) at 0 °C, were added HATU (0.33 g, 0.88 mmol), *N*-methyl morpholine (0.18 g, 1.76 mmol) and 1,4-oxazepane (64 mg, 0.638 mmol). The reaction mixture was allowed to warm to rt and stirred under inert atmosphere for 16 h, then was
 25 diluted with water (15 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) and


followed by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford 4-(6-(4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (130 mg, 54%, AUC HPLC 99.85%) as an off-white solid; m.p. 107-112 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.49 (s, 1H), 7.84-7.71 (m, 5H), 7.60-7.50 (m, 5H), 4.09-3.8 (m, 5H), 3.7 (bs, 1H), 3.67 (b. s, 2H), 2.1 (bs, 1H); 1.9 (bs, 1H); MS (ESI) *m/z* 423.32 [C₂₆H₂₂N₄O₂ + H]⁺.

Example 98: 4-(6-(4-(4-ethylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



- 10 To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid 1 (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methyl morpholine (130 μL, 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 1-ethylpiperazine (75 μL, 0.589 mmol). The reaction mixture was left to stir for 18 h, then was diluted with water (25 mL)
- 15 and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford 4-(6-(4-(4-ethylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (93 mg, 73%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.03-7.98 (m, 5H), 7.84-7.81 (m, 3H), 7.74-7.72 (m, 1H), 7.50 (d, *J* = 8.0 Hz, 2H), 3.62-3.38 (m, 4H), 2.37 (s, 6H), 1.01 (t, *J* = 6.8 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.49, 145.73, 137.74, 135.24, 135.08, 133.49, 133.15, 127.73, 127.66, 127.02, 125.73, 125.51, 124.40, 121.82, 118.78, 117.76, 109.77, 51.40, 11.69; MS (ESI) *m/z* 436 [C₂₇H₂₅N₅O + H]⁺.
- 20

Example 99: *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-yl)acetamide




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoic acid (1 g, 2.9 mmol) in DMF (15.0 mL) were added HATU (1.65 g, 4.38 mmol), *N*-methyl morpholine (0.641 mL, 5.8 mmol) and *tert*-butyl 4-methylpiperidin-4-ylcarbamate (691 mg, 3.24 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then was diluted with water (15 mL) and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (800 mg, 53%) as an off-white solid. ¹H NMR (400 MHz, CD₃OD) δ (ppm): 8.47 (s, 1H), 7.88 (d, *J* = 7.6 Hz, 2H), 7.83 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 2H), 7.72 (d, *J* = 9.2 Hz, 1H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.51 (d, *J* = 7.6 Hz, 2H), 4.42 (bs, 1H), 4.19 (bs, 1H), 3.55-3.32 (m, 3H), 2.07-1.95 (m, 4H), 1.44 (s, 9H), 1.40 (s, 3H); MS (ESI) *m/z* 536.2 [C₃₂H₃₃N₅O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (800 mg, 1.49 mmol) in DCM (10 mL) were added TFA (3 mL) in DCM (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL), NaHCO₃ (2 x 100 mL) and extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyridin-3-yl)benzonitrile (600 mg, 61%, AUC HPLC >99%) as an off white solid; m.p. 233-235°C. ¹H NMR (400 MHz, CD₃OD) δ (ppm): 8.75 (s, 1 H), 7.92 (m, 3 H), 7.87 (s, 1 H), 7.79-7.76 (m, 4 H), 7.52 (d, *J* = 11.2 Hz, 2 H), 3.8 (bs, 2 H), 3.62 (m, 2 H), 1.72-1.41 (m, 4 H), 1.20 (s, 3 H); MS (ESI) *m/z* 436.2 [C₂₇H₂₅N₅O + H]⁺.

Step 3: To a solution of 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (200 mg, 0.46 mmol) in DCM (10 mL) were added TEA (0.13 mL, 0.92 mmol) and acetyl chloride (0.05 mL, 0.596 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 4 h, then was diluted with cold water (10 mL) and extracted with EtOAc (3 x 25 mL). The combined organic layer was washed with NaHCO₃ (2 x 10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) and by preparative HPLC to afford *N*-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-yl)acetamide (100 mg, 47%, AUC HPLC 98.1%) as an off white solid; m.p. 188-210 °C. ¹H NMR (100 MHz, DMSO-*d*₆) δ (ppm): 8.79 (s, 1 H), 8.02-7.96 (m, 5 H), 7.81 (d, *J* = 8.0 Hz, 2 H), 7.72 (d, *J* = 12.0 Hz, 2 H), 7.50-7.45 (m, 3 H), 4.12 (bs, 1 H), 3.23 (bs, 2 H), 2.09 (bs, 2 H), 1.82 (s, 3 H), 1.44 (bs, 2 H), 1.29 (s, 3 H); MS (ESI) *m/z* 478.39 [C₂₉H₂₇N₅O₂ + H]⁺.


Example 100: 4-(6-(4-(4-(diethylamino)-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (200 mg, 0.46 mmol) in methanol (15.0 mL) were added acetaldehyde (40 mg, 0.92 mmol), catalytic amount of acetic acid (0.5 mL). The reaction mixture was stirred at 0 °C under inert atmosphere for 2 h, then was slowly added NaBH₃CN (97 mg, 1.38 mmol) then the reaction was slowly warmed to rt and stirred for 24 h. Reaction mixture was diluted with water and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to afford followed by preparative HPLC to afford 4-(6-(4-(4-(diethylamino)-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 46%, AUC HPLC 97.2%) as an off white solid; m.p. 204-208 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.49 (s, 1 H), 7.84-7.71 (m, 6 H), 7.60-7.50 (m, 5 H), 3.85-3.59 (m, 3 H), 3.33 (bs, 1 H), 2.57-2.49


(m, 4 H), 1.90 (bs, 1 H), 1.74 (bs, 1 H), 1.42-1.30 (m, 2 H), 1.06-1.01 (m, 9 H); MS (ESI) m/z 492.34 [$C_{31}H_{33}N_5O + H$] $^+$.

Example 101: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-methyl-N-(2-morpholino-2-oxoethyl)benzamide




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methylmorpholine (130 μ L, 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 2-(methylamino)-1-morpholinoethanone hydrochloride (115 mg, 0.589 mmol). The reaction mixture was stirred for 18 h, then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 97:3) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-*N*-methyl-*N*-(2-morpholino-2-oxoethyl)benzamide (87 mg, 61%, AUC HPLC 97%) as a white solid. 1H NMR (400 MHz, $CDCl_3$) δ 8.50 (s, 1H), 7.84-7.79 (m, 4H), 7.75-7.73 (m, 2H), 7.63-7.52 (m, 5H), 4.36 (s, 2H), 3.73-3.54 (m, 8H), 3.11 (s, 3H); ^{13}C NMR (100 MHz, $DMSO-d_6$) δ 170.16, 166.08, 145.71, 137.68, 135.54, 135.04, 133.49, 133.16, 127.75, 127.60, 126.95, 125.75, 124.42, 121.83, 118.78, 117.76, 109.80, 66.06, 52.21, 48.37, 44.53, 41.68, 34.39; MS (ESI) m/z 480 [$C_{28}H_{25}N_5O_3 + H$] $^+$.

Example 102: 4-(6-(4-(3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μ L, 0.824 mmol) and 2-oxopiperazine (25 mg, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 95:5) to afford 4-(6-(4-(3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (80 mg, 92%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (s, 1H), 8.13 (bs, 1H), 7.95-8.05 (m, 5H), 7.78-7.89 (m, 3H), 7.70-7.78 (m, 1H), 7.57 (d, *J* = 8.53 Hz, 2H), 4.09 (bs, 2H), 3.72 (bs, 1H), 3.55 (bs, 1H), 3.26 (bs, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.65, 145.74, 138.12, 135.06, 133.48, 133.17, 127.75, 128.00, 125.69, 125.55, 124.44, 121.93, 118.79, 117.76, 109.79; MS (ESI) *m/z* 422 [C₂₅H₁₉N₅O₂ + H]⁺.

Example 103: 4-(6-(4-(4-methyl-1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μ L, 0.824 mmol) and *N*-methylhomopiperazine (31 μ L, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 90:10) to afford 4-(6-(4-(4-methyl-1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (47 mg, 52%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.76 (s, 1H), 7.90-7.97 (m, 4H), 7.88 (s, 1H), 7.74-7.82 (m, 4H), 7.56 (d, *J* = 8.2 Hz, 2H), 3.80-3.83 (m, 1H), 3.74-3.80 (m, 1H), 3.56-3.62 (m, 1H), 3.52-3.56 (m, 1H), 2.95 (s, 1H), 2.86 (bs, 1H), 2.70-2.76 (m, 1H), 2.64-2.70 (m, 2H), 2.42 (d, *J* = 0.8 Hz, 3H), 1.86-1.94 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 169.94, 145.78, 135.10, 133.55, 133.22, 127.81, 127.47, 127.03, 125.80, 125.59, 124.47, 121.81, 118.84, 117.83, 109.84, 47.98; MS (ESI) *m/z* 436 [C₂₇H₂₅N₅O + H]⁺.

Example 104: 4-(6-(4-(4-(piperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




5 Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methyl morpholine (129 μ L, 1.18 mmol) and 1-Boc-4-(piperidin-4-yl)piperazine (95 mg, 0.353 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined
10 organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96:4) to afford *tert*-butyl 4-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)piperazine-1-carboxylate (57 mg, 33%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.76 (bs., 1H), 7.96-7.88 (m, 5H), 7.83-7.77 (m, 4H), 7.56 (d, *J* = 8.0 Hz, 2H), 4.00-3.85 (m, 1H), 3.65-3.55 (m, 4H), 3.25-3.10 (m, 2H), 3.10-3.00 (m, 4H), 3.00-2.70 (m, 2H), 2.25-1.95 (m, 2H), 1.75-1.55 (m, 2H), 1.47 (s, 9H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 171.93, 155.72, 140.09, 136.30, 134.67, 134.40, 134.25, 129.50, 129.96, 128.76, 128.55, 128.10, 122.92, 119.49, 118.30, 112.80, 82.11, 64.13, 28.53; MS (ESI) *m/z* 591 [C₃₅H₃₈N₆O₃ + H]⁺.
15
20

Step 2: A solution of *tert*-butyl 4-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)piperazine-1-carboxylate (56 mg, 0.095 mmol) was added 20% TFA in CH₂Cl₂ (5 mL). The reaction mixture was stirred at room temperature for 20 min then, was concentrated under reduced pressure. To the residue was added saturated sodium
25 bicarbonate (10 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(4-(piperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (47 mg, 99%, AUC HPLC 99%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 8.88 (s, 1H), 8.36-8.30 (m, 2H), 8.11 (d, *J* = 9.6 Hz, 1H), 8.05-7.98 (m, 4H), 7.84 (d, *J* = 8.4 Hz, 2H), 7.59 (d, *J* = 8.0 Hz, 2H), 4.75-4.65 (m, 1H), 3.90-3.75 (m, 1H), 3.35-3.25 (m, 1H), 3.25-3.15 (m,
30

1H), 3.10-3.00 (m, 4H), 3.00-2.85 (m, 2H), 2.15-1.85 (m, 2H), 1.70-1.50 (m, 2H), 1.40-1.20 (m, 2H), 1.00-0.85 (m, 1H) ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.47, 158.52, 158.19, 136.84, 135.75, 133.25, 131.81, 128.85, 127.55, 127.38, 125.16, 122.92, 118.59, 117.87, 115.44, 114.92, 111.09, 61.85, 45.27; MS (ESI) *m/z* 491 [C₃₀H₃₀N₆O + H]⁺.


5

Example 105: 4-(6-(4-(1'-methyl-4,4'-bipiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methylmorpholine (129 μL, 1.18 mmol) and, 1-methyl-4,4'-bipiperidine (64 mg, 0.353 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96:4), followed by cold CH₃OH/CH₃CN wash to afford 4-(6-(4-(1'-methyl-4,4'-bipiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (131 mg, 89%, AUC HPLC 97%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.04-7.98 (m, 5H), 7.87-7.79 (m, 3H), 7.77-7.70 (m, 1H), 7.50 (d, *J* = 8.0 Hz, 2H), 4.60-4.40 (m, 1H), 3.75-3.60 (m, 1H), 3.45-3.35 (m, 1H), 3.15-2.70 (m, 9H), 2.65-2.55 (m, 1H), 2.50-2.35 (m, 2H), 1.90-1.65 (m, 2H), 1.50-1.30 (m, 2H), 1.30-1.10 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.39, 145.66, 137.60, 135.02, 133.44, 133.09, 127.69, 127.41, 126.94, 125.67, 125.46, 124.34, 121.74, 118.71, 117.71, 109.72, 53.11, 45.67; MS (ESI) *m/z* 505 [C₃₂H₃₃N₅O + H]⁺.

Example 106: 4-(6-(4-(4-hydroxy-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (200 mg, 0.58 mmol) in DMF (15.0 mL) were added HATU (335 mg, 0.88 mmol), *N*-methyl morpholine (0.21 mL, 1.17 mmol) and 4-methylpiperidin-4-ol hydrochloride (115 mg, 0.76 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then it was diluted with water (25 mL) and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) followed by preparative HPLC to afford 4-(6-(4-(4-hydroxy-4-methylpiperidine-1-carbonyl) phenyl)imidazo [1,2-a]pyridin-3-yl)benzonitrile (100 mg, 40%, AUC HPLC >99%) as an off white solid; m.p. 149-154 °C. ¹H NMR (400 MHz, DMSO-d₆) δ (ppm): 8.80 (bs, 1 H), 8.03-7.97 (m, 5 H), 7.82-7.71 (m, 4 H), 7.48 (d, *J* = 8.0 Hz, 2 H), 4.45 (s, 1 H), 4.11 (bs, 1 H), 3.41 (bs, 1 H), 3.27 (bs, 2 H), 1.53-1.44 (m, 4 H), 1.16 (s, 3 H); MS (ESI) *m/z* 437.26 [C₂₇H₂₄N₄O₂ + H]⁺.

Example 107: 4-(6-(4-(1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl) imidazo[1,2-a]pyridin-6-yl) benzoic acid (100 mg, 0.29 mmol) in DMF (3 mL) were added HATU (165 mg, 0.44 mmol), *N*-methyl morpholine (117 mg, 1.16 mmol) and *tert*-butyl 1,4-diazepane-1-carboxylate (118 mg, 0.59 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h before being diluted with water (10 mL) and extracted with DCM (3×10 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent DCM/MeOH 7:3) to

afford *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate (120 mg, 79%, AUC HPLC 98%) as white solid. MS (ESI) m/z 522 [$C_{31}H_{31}N_5O_3 + H$]⁺.

- 5 Step 2: To a solution of *tert*-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)-1,4-diazepane-1-carboxylate in DCM (3 mL) was added TFA (3 mL). The reaction mixture was stirred at room temperature for 18 h then, was concentrated under reduced pressure and the residue was purified by preparative HPLC (C18, eluent ACN/H₂O/0.01% HCOOH) to afford 4-(6-(4-(1,4-diazepane-1-carbonyl)phenyl)imidazo[1,2-
10 a]pyridin-3-yl)benzonitrile (78 mg, 64%, AUC HPLC 98%) as white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.75 (s, 1H), 7.93 (s, 4H), 7.89 (s, 1H), 7.81 (d, J = 8.2 Hz, 2H), 7.77 (s, 2H), 7.61 (d, J = 8.2 Hz, 2H), 4.05–3.60 (m, 4H), 3.50–3.32 (m, 4H), 2.27–2.00 (m, 2H); ¹³C NMR (400 MHz, CD₃OD) δ 173.64, 166.40, 147.45, 140.31, 136.31, 134.85, 134.72, 134.43, 129.51, 129.00, 128.59, 127.84, 126.63, 122.92, 119.53, 118.53, 112.79, 46.73, 45.87, 43.46,
15 27.48; MS (ESI) m/z 422 [$C_{26}H_{23}N_5O + H$]⁺.

Example 108: 4-(6-(4-(4-(piperidin-4-yl)piperazine-1-carbonyl)phenyl)imidazo[1,2-
a]pyridin-3-yl)benzonitrile



- 20 Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl) benzoic acid (200 mg, 0.58 mmol) in DMF (3 mL), were sequentially added HATU (330 mg, 0.87 mmol), *N*-methyl morpholine (235 mg, 2.32 mmol) and *tert*-butyl 4-(piperazin-1-yl)piperidine-1-carboxylate (312 mg, 1.16 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was diluted with water (10 mL) and extracted with
25 DCM (3×15 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent DCM/MeOH 9:1) to afford *tert*-butyl 4-(4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)piperazin-1-yl)piperidine-1-carboxylate (239 mg, 69%, AUC HPLC 95%) as an off-white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.64
30 (s, 1H), 7.84 (s, 4H), 7.80 (s, 1H), 7.20–7.65 (m, 4H), 7.49 (d, J = 8.0 Hz, 2H), 4.72 (s, 1H),

4.00–3.69 (s, 1H), 3.55–3.36 (m, 4H), 3.29–2.75 (m, 2H), 2.65–2.49 (m, 5H), 2.19–1.59 (m, 2H), 1.59–1.38 (m, 11H); MS (ESI) m/z 591 [$C_{35}H_{38}N_6O_3 + H$] $^+$.

Step 2: To a solution of *tert*-butyl 4-(4-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-

5 yl)benzoyl)piperazin-1-yl)piperidine-1-carboxylate in DCM (3 mL), was added TFA (3 mL).

The reaction mixture was stirred at room temperature for 18 h then, was concentrated under reduced pressure and the residue was purified by column chromatography (silica gel, eluent DCM/MeOH 4:1) to afford 4-(6-(4-(4-(piperidin-4-yl)piperazine-1-

carbonyl)phenyl)imidazo[1,2-*a*]pyridin-3-yl)benzonitrile (141 mg, 48%, AUC HPLC 99%) as


10 pink solid. 1H NMR (400 MHz, CD_3OD) δ 8.79 (s, 1H), 8.09–8.03 (s, 1H), 8.02–7.98 (m, 1H), 7.96 (s, 4H), 7.94–7.88 (m, 1H), 7.80 (d, $J = 8.0$ Hz, 2H), 7.55 (d, $J = 8.0$ Hz, 2H), 4.66 (s, 1H), 3.82 (s, 1H), 3.28–3.06 (m, 5H), 3.00–2.83 (m, 5H), 2.76 (m, $J = 5.5$ Hz, 1H), 2.08 – 1.74 (m, 2H), 1.67–1.40 (m, 2H); ^{13}C NMR (100 MHz, CD_3OD) δ 171.78, 163.59, 163.24,

162.90, 162.56, 145.15, 139.22, 137.14, 135.65, 134.53, 133.45, 130.87, 130.28, 130.20,

15 129.37, 128.96, 128.86, 128.73, 127.23, 123.65, 119.75, 119.36, 116.94, 113.75, 62.66,

47.18, 45.13, 44.43, 42.58, 29.70, 28.94; MS (ESI) m/z 491 [$C_{35}H_{30}N_6O + H$] $^+$

Example 109: 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyridin-3-yl)benzonitrile




20 Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoic acid (300 mg, 0.88 mmol) in DMF (2 mL) was added NMM (134 mg, 1.33 mmol) followed by addition of HATU (504 mg, 1.33 mmol) at rt and stirred for 30 min. *tert*-butyl piperidin-4-ylcarbamate (195 mg, 0.97 mmol) was added and stirred at rt for 1 h. The reaction mixture

25 was diluted with water and filtered to obtain of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)piperidin-4-ylcarbamate (325 mg, 55%, LC-MS 78.5%) as a pale yellow solid. MS (ESI) m/z : 522.2(M+H).


Step 2: Trifluoroacetic acid (3.5 mL) in CH_2Cl_2 (15 mL) was added to a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-ylcarbamate (325 mg, 0.624 mmol) in CH_2Cl_2 at 0 °C and stirred at rt for 1 h. Reaction mixture was diluted with water and basified with sat. Na_2CO_3 solution. The aqueous layer was extracted with EtOAc, dried and concentrated under reduced pressure to give crude product which, was purified by preparative HPLC to afford 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (30 mg, 21%, AUC HPLC 97.3%) as an off-white solid; m.p. 129-138 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.80 (s, 1 H), 8.03-7.97 (m, 5 H), 7.82 (d, $J = 11.0$ Hz, 2 H), 7.72 (d, $J = 14.0$ Hz, 2 H), 7.46 (d, $J = 11.0$ Hz, 2 H), 4.29 (bs, 1 H), 3.57 (bs, 1 H), 3.07-2.84 (m, 3 H), 2.27-2.18 (m, 2 H), 1.76 (bs, 2 H), 1.23 (bs, 2 H); MS (ESI) m/z 422.24 [$\text{C}_{26}\text{H}_{23}\text{N}_5\text{O} + \text{H}$] $^+$.

Example 110: N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)-2,2,2-trifluoroacetamide




To a solution of 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (300 mg, 0.71 mmol) in dichloromethane (10 mL) were added trifluoroacetic anhydride (0.12 mL, 0.85 mmol) and triethyl amine (0.2 mL, 1.42 mmol) in dichloromethane. The reaction mixture was stirred for 2 h at rt. The reaction mixture was diluted with H_2O (100 mL), NaHCO_3 (100 mL) and extracted with CH_2Cl_2 (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. The crude product was purified by Prep-HPLC (C18, ACN/ H_2O / 10 mM NH_4HCO_3) to afford N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)-2,2,2-trifluoroacetamide (120 mg, 33%, AUC HPLC 97.7%) as a white solid; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 9.40 (d, $J = 9.3$ Hz, 1H), 8.80 (s, 1H), 8.03-7.97 (m, 5H), 7.86-7.82 (m, 4H), 7.50 (d, $J = 7.48$ Hz, 2H), 4.47 (s, 1H), 3.97 (q, $J = 3.2$ Hz, 1H), 3.69-3.36 (m, 1H), 3.31-3.28 (m, 1H), 2.97-2.93 (m, 1H), 2.07-1.77 (m, 2H), 1.51-1.49 (m, 2H); MS (ESI) m/z 518.31 [$\text{C}_{28}\text{H}_{22}\text{F}_3\text{N}_5\text{O}_2 + \text{H}$] $^+$.

Example 111: N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)methanesulfonamide



To a solution of 4-(6-(4-(4-aminopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (130 mg, 0.30 mmol) in dichloromethane (10 mL) were added TEA (0.12 mL, 0.91 mmol) and MsCl (0.42 mL, 0.37 mmol). The reaction mixture was stirred for 2 h at ambient temperature and diluted with H₂O (100 mL), NaHCO₃ (100 mL) and extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. The crude product was purified by preparative HPLC to afford N-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)methanesulfonamide (60 mg, 42%, AUC HPLC 99.02%) as an off-white solid; mp 211-215 °C, ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.50 (s, 1H), 7.84-7.80 (m, 4H), 7.73 (d, *J* = 8.4 Hz, 2H), 7.59 (d, *J* = 8.0 Hz, 2H), 7.53-7.50 (m, 3H), 4.60 (bs, 1H), 4.29 (d, *J* = 7.6 Hz, 1H), 3.85 (bs, 1H), 3.63-3.60 (m, 1H), 3.15 (bs, 2H), 3.09 (s, 3H), 2.03 (bs, 2H), 1.45 (bs, 2H); MS (ESI) *m/z* 500 [C₂₇H₂₅N₅O₃S + H]⁺.


Example 112: 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (0.2 g, 0.58 mmol) in DMF (5 mL) were added HATU (0.33 g, 0.88 mmol), N-methyl morpholine (0.18 g, 1.76 mmol) and N,N-dimethylpiperidin-4-amine (81 mg, 0.70 mmol). The reaction mixture was stirred at 0 °C to room temperature under inert atmosphere for 16 h, then it was diluted with water (15 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was

dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford 4-(6-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (140 mg, 57%, AUC HPLC >99%) as an off-white solid; m.p. 128-132 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.49 (s, 1H), 7.84-7.71 (m, 5H), 7.60-7.50 (m, 5H), 7.80 (m, 3H), 7.71 (m, 1H), 7.45 (q, 2H), 4.45 (bs, 1H), 3.65 (bs, 1H), 3.15-2.8 (m, 2H), 2.39 (m, 1H), 2.22 (bs, 6H), 1.9-1.6 (m, 2H), 1.35 (m, 2H); MS (ESI) m/z 450.3 [$\text{C}_{28}\text{H}_{27}\text{N}_5\text{O} + \text{H}$] $^+$.

10 Example 113: 4-(6-(4-(4-amino-4-ethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methyl morpholine (130 μL , 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of *tert*-butyl 4-ethylpiperidin-4-ylcarbamate (134 mg, 0.589 mmol). The reaction mixture was stirred for 18 h, then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford a white solid (45 mg, 28%, AUC HPLC 96%); MS (ESI) m/z 550 [$\text{C}_{33}\text{H}_{35}\text{N}_5\text{O}_3 + \text{H}$] $^+$.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-ethylpiperidin-4-ylcarbamate (45 mg, 0.082 mmol) in DCM (1.0 mL) was added trifluoroacetic acid (1 mL). The reaction mixture was stirred at room temperature under inert atmosphere for 30min, then was neutralized with NaHCO_3 , diluted with water (25 mL) and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 90:9:1) to afford 4-(6-(4-(4-amino-4-ethylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (6

mg, 16%, AUC HPLC 98%) as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 8.77 (s, 1H), 7.97-7.90 (m, 5H), 7.82-7.79 (m, 4H), 7.55 (d, $J=8.4$ Hz, 2H), 3.90- 3.54 (m, 4H), 1.75-1.51 (m, 6H), 0.98 (t, $J=7.6$ Hz, 3H); MS (ESI) m/z 451 [$\text{C}_{28}\text{H}_{27}\text{N}_5\text{O} + \text{H}$] $^+$.

5 Example 114: 4-(6-(4-(2,7-diazaspiro[3.5]nonane-7-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of 4-(3-iodoimidazo [1,2-a]pyridin-6-yl)benzoic acid (500 mg, 1.37 mmol) in DMF (15.0 mL) were added HATU (782.9 mg, 2.05 mmol), *N*-methyl morpholine (0.03 mL, 2.74 mmol) and 1-methylpiperazine (340 mg, 1.5 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then it was diluted with water (25 mL) and extracted with EtOAc (3 $\times$ 25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford *tert*-butyl 7-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-2,7-diazaspiro[3.5]nonane-2-carboxylate (440 mg, 56%) as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.45 (s, 1H), 7.94-7.67 (m, 5H), 4.08 (s, 1H), 3.98 (s, 1H), 3.79 (s, 1H), 3.29 (m, 4H), 2.89 (s, 3H), 1.77-1.70 (m, 4H), 1.38 (s, 9H); MS (ESI) m/z 573.1 [$\text{C}_{26}\text{H}_{29}\text{N}_4\text{O}_3 + \text{H}$] $^+$.


Step 2: 4-cyanophenylboronic acid (135 mg, 0.92 mmol), K_3PO_4 (326 mg, 1.53 mmol), and $\text{Pd}(\text{PPh}_3)_4$ (66 mg, 0.04 mmol) were added sequentially to a solution of *tert*-butyl 7-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-2,7-diazaspiro[3.5]nonane-2-carboxylate (440 mg, 0.769 mmol) in a mixture of 1,4-Dioxane/ H_2O (30:10 mL) at room temperature under argon atmosphere. The reaction mixture was refluxed for 6 h and was diluted with water (50 mL) and extracted with EtOAc (3 $\times$ 50 mL). The combined organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CHCl}_3/\text{MeOH}$ 95:5) to afford 4-(6-(4-(2,7-diazaspiro[3.5]nonane-7-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (250 mg, 59%). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.80 (s, 1H), 8.01 (s, 4H), 7.85-7.73 (m, 6H),

4.06 (s, 1H), 3.78 (s, 1H), 3.34 (m, 4H), 1.66 (bs, 4H), 1.38 (s, 9H); MS (ESI) m/z 548 $[C_{33}H_{33}N_5O_3 + H]^+$

Step 3: To a solution of 4-(6-(4-(2,7-diazaspiro[3.5]nonane-7-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (150 mg, 0.274 mmol) in dichloromethane (10 mL) were added TFA (3 mL) in dichloromethane (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL), $NaHCO_3$ (100 mL) and extracted with CH_2Cl_2 (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. The crude product was purified by preparative HPLC (C18, ACN/ H_2O /10 mM NH_4HCO_3) to afford (4-amino-4-methylpiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)methanone (80 mg, 16%, AUC HPLC 97.23%) as a brown solid; 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.81 (s, 1H), 8.01 (m, 5H), 7.85-7.82 (d, $J = 7.83$ Hz, 3H), 7.76-7.73 (d, $J = 7.74$ Hz, 3H), 4.01 (bs, 2H), 3.73 (bs, 2H), 2.50 (m, 4H), 1.60 (m, 4H); MS (ESI) m/z 448.26 $[C_{28}H_{25}N_5O + H]^+$.

15

Example 115: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(2-morpholino-2-oxoethyl)benzamide




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methylmorpholine (130 μ L, 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 2-amino-1-morpholinoethanone (106 mg, 0.589 mmol). The reaction mixture was left to stir for 18 h, then it was diluted with water (25 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 97:3) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(2-morpholino-2-oxoethyl)benzamide (84 mg, 61%, AUC HPLC 98%) as a white solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 8.84 (s, 1H), 8.65 (t, $J = 5.6$ Hz, 1H), 8.04-7.98 (m, 7H), 7.89-7.87 (m, 2H), 7.84-7.75 (m, 2H), 4.16 (d, $J = 5.6$

25

Hz, 2H), 3.61-3.57 (m, 4H), 3.51-3.46 (m, 4H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 167.20, 165.82, 145.76, 139.39, 135.11, 133.49, 133.28, 133.17, 127.92, 127.75, 126.88, 125.57, 125.52, 124.46, 121.97, 118.79, 117.76, 109.79, 66.04, 44.61, 41.78, 40.81; MS (ESI) m/z 466 $[\text{C}_{27}\text{H}_{23}\text{N}_3\text{O}_3 + \text{H}]^+$.

5


Example 116: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(2-morpholinoethyl)benzamide



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methyl morpholine (130 μL , 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 2-morpholinoethanamine (77 μL , 0.589 mmol). The reaction mixture was left to stir for 18 h, then it was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-*N*-(2-morpholinoethyl)benzamide (90 mg, 67%, AUC HPLC 98%) as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 8.52 (s, 1H), 7.90-7.88 (m, 2H), 7.84-7.80 (m, 4H), 7.75-7.73 (m, 2H), 7.64-7.61 (m, 2H), 7.56-7.53 (m, 1H), 3.74 (t, $J = 4.4$ Hz, 4H), 3.61-3.57 (m, 2H), 2.64 (t, $J = 6.0$ Hz, 2H), 2.53 (s, 4H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 165.59, 145.69, 139.17, 135.05, 133.54, 133.42, 133.09, 127.75, 127.65, 126.77, 125.51, 125.44, 124.37, 121.85, 118.71, 117.69, 109.72, 65.92, 57.09, 53.04; MS (ESI) m/z 452 $[\text{C}_{27}\text{H}_{25}\text{N}_3\text{O}_2 + \text{H}]^+$.

20

Example 117: 4-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (120 mg, 0.353 mmol) in DMF (1.7 mL) was added HATU (201 mg, 0.530 mmol) and *N*-methylmorpholine (155 μ L, 1.41 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of *tert*-butyl 1,9-diazaspiro[5.5]undecane-1-carboxylate hydrochloride (205 mg, 0.707 mmol). The reaction mixture was left to stir for 18 h, then it was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford *tert*-butyl 9-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (192 mg, 94%, AUC HPLC 96%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (s, 1H), 8.03-7.98 (m, 5H), 7.84-7.81 (m, 3H), 7.75-7.72 (m, 1H), 7.49-7.47 (m, 2H), 3.42-3.31 (m, 5H), 1.70-1.51 (m, 9H), 1.40 (bs, 11H); MS (ESI) *m/z* 576 [C₃₅H₃₇N₅O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 9-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-1,9-diazaspiro[5.5]undecane-1-carboxylate (183 mg, 0.318 mmol) in DCM (1.0 mL) was added trifluoroacetic acid (1.0 mL). The reaction mixture was stirred at room temperature under inert atmosphere for 30 min, then was neutralized with solid NaHCO₃, then diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 92:8) to afford 4-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (96 mg, 64%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.03-7.98 (m, 5H), 7.86-7.82 (m, 3H), 7.75-7.72 (m, 1H), 7.49-7.47 (m, 2H), 3.50-3.31 (m, 4H), 2.98 (bs, 2H), 1.69-1.58 (m, 10H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 168.64, 145.73, 137.80, 135.26, 135.08, 133.49, 133.16, 127.75, 127.44, 127.09,

125.71, 125.51, 124.42, 121.81, 118.78, 117.78, 115.76, 109.79, 53.66, 22.92, 17.80; MS (ESI) m/z 476 [$C_{30}H_{29}N_5O + H$] $^+$.


5 Example 118: 4-(6-(4-(1-acetyl-1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (30 mg, 0.063 mmol) in DCM (0.5 mL) at 0°C was added triethylamine (26 μ L, 0.189 mmol), followed by acetyl chloride (10 μ L, 0.126 mmol). The reaction mixture was stirred at 0°C under inert atmosphere and allowed to warm to room temperature over 30 minutes. The reaction mixture was then diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH/NH_4OH$ 90:9:1) to afford 4-(6-(4-(1-acetyl-1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (14 mg, 43%, AUC HPLC 99%) as a white solid. 1H NMR (400 MHz, $CDCl_3$) δ 8.49 (s, 1H), 7.85-7.82 (m, 4H), 7.81-7.78 (m, 2H), 7.75-7.73 (m, 5H), 3.96-3.40 (m, 6H), 3.09-2.96 (m, 2H), 2.13 (s, 3H), 1.76-1.69 (m, 6H), 1.42 (bs, 2H); MS (ESI) m/z 518 [$C_{32}H_{31}N_5O_2 + H$] $^+$.


20

Example 119: 4-(6-(4-(1-methyl-1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



A mixture of 4-(6-(4-(1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (104 mg, 0.219 mmol) and paraformaldehyde (70 mg) in acetonitrile (3 mL) and acetic acid (1.7 mL) was treated with sodium borohydride (35 mg). The reaction mixture was left to stir for 24 h, then it was diluted with saturated NaHCO₃ (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH/NH₄OH 90:9:1) to afford 4-(6-(4-(1-methyl-1,9-diazaspiro[5.5]undecane-9-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (49 mg, 45%, AUC HPLC 93%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.52 (s, 1H), 7.86-7.82 (m, 4H), 7.77-7.74 (m, 2H), 7.61-7.59 (m, 2H), 7.55-7.53 (m, 3H), 4.45 (bs, 1H), 3.77 (bs, 1H), 3.24 (bs, 2H), 2.94 (bs, 2H), 2.56 (bs, 3H), 2.01-1.52 (m, 10H); ¹³C NMR (100 MHz, CDCl₃) δ 169.72, 146.34, 138.62, 134.78, 133.79, 133.25, 127.95, 127.84, 127.23, 127.19, 125.68, 124.52, 120.61, 118.73, 118.43, 111.60, 50.03, 35.99, 22.12, 21.78, 18.94; MS (ESI) *m/z* 490 [C₃₁H₃₁N₅O + H]⁺.


Example 120: 4-(6-(4-(4-hydroxy-4-phenylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methylmorpholine (130 µl, 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 4-phenylpiperidin-4-ol (105 mg, 0.589 mmol). The reaction mixture was left to stir for 18 h, then was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(4-hydroxy-4-phenylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (52 mg, 35%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.03-7.98


(m, 5H), 7.84-7.81 (m, 3H), 7.75-7.72 (m, 1H), 7.58-7.53 (m, 4H), 7.35-7.31 (m, 2H), 7.24-7.21 (m, 1H), 5.18 (s, 1H), 4.46 (bs, 1H), 3.53 (br s, 2H), 3.31 (bs, 1H), 1.96 (bs, 2H), 1.72-1.58 (m, 2H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 168.44, 149.11, 145.66, 137.42, 135.80, 134.99, 133.44, 133.08, 127.79, 127.66, 127.47, 126.91, 126.32, 125.74, 125.49, 124.69, 124.33, 121.71, 118.71, 117.67, 109.69, 70.07; MS (ESI) m/z 499 [$\text{C}_{32}\text{H}_{26}\text{N}_4\text{O}_2 + \text{H}$] $^+$.

Example 121: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(2-(1-methylpiperidin-4-yl)ethyl)benzamide




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methylmorpholine (130 μL , 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of 2-(1-methylpiperidin-4-yl)ethanamine (84 mg, 0.589 mmol). The reaction mixture was left to stir for 18 h, then it was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was washed with MeOH (3x25 mL) and dried under vacuum to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(2-(1-methylpiperidin-4-yl)ethyl)benzamide (36 mg, 26%, AUC HPLC 97%) as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.82 (s, 1H), 8.49 (br s, 1H), 8.03-7.99 (m, 5H), 7.96-7.94 (m, 2H), 7.88-7.82 (m, 3H), 7.76-7.74 (m, 1H), 3.41 (b. s, 3H), 3.02-2.56 (m., 11H), 2.32 (m, 2H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 145.68, 139.25, 135.05, 133.42, 133.09, 127.76, 127.67, 126.78, 125.49, 125.44, 124.37, 121.84, 118.71, 117.71, 109.75, 55.77, 52.67, 49.48, 20.65; MS (ESI) m/z 465 [$\text{C}_{29}\text{H}_{29}\text{N}_5\text{O} + \text{H}$] $^+$.

Example 122: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-methyl-N-(pyridin-4-yl)benzamide



- To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.295 mmol) in DMF (1.5 mL) was added HATU (168 mg, 0.442 mmol) and *N*-methyl morpholine (130 μ L, 1.178 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 1 h, followed by the addition of *N*-methylpyridin-4-amine (64 mg, 0.589 mmol). The reaction mixture was stirred for 18 h, then it was diluted with water (25 mL) and extracted with EtOAc (3 x 25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by preparative HPLC (C18, eluent ACN/water/formic acid 0.1%) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-*N*-methyl-*N*-(pyridin-4-yl)benzamide (20 mg, 16%, AUC HPLC 97%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.77 (s, 1H), 8.45-8.43 (m, 2H), 7.99-7.98 (m, 5H), 7.79-7.68 (m, 4H), 7.45-7.43 (m, 2H), 7.24-7.23 (m, 2H), 3.43 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ 169.33, 151.39, 150.40, 145.72, 138.18, 135.09, 134.83, 133.45, 133.15, 129.10, 127.76, 126.53, 125.32, 125.22, 124.69, 121.88, 120.50, 118.78, 117.72, 109.79, 37.04; MS (ESI) m/z 430 $[\text{C}_{27}\text{H}_{19}\text{N}_5\text{O} + \text{H}]^+$.

Example 123: 4-(6-(4-(4-(aminomethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




- Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methyl morpholine (129 μ L, 1.18 mmol) and 4-(*boc*-aminomethyl)piperidine (95 mg, 0.441 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (10 mL) and extracted with EtOAc (3x20 mL). The combined organic

layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $\text{CHCl}_3/\text{MeOH}$ 96:4) to afford *tert*-butyl (1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)piperidin-4-yl)methylcarbamate (157 mg, 99%, AUC HPLC 99%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.80 (s, 1H), 8.03-7.98 (m, 5H), 7.84-7.80 (m, 3H), 7.75-7.70 (m, 1H), 7.48 (d, $J = 8.4$ Hz, 2H), 6.90-6.85 (m, 1H), 4.55-4.35 (m, 1H), 3.70-3.55 (m, 1H), 3.10-2.95 (m, 1H), 2.90-2.70 (m, 4H), 1.75-1.55 (m, 2H), 1.37 (s, 9H), 1.15-1.00 (m, 2H); MS (ESI) m/z 536 [$\text{C}_{32}\text{H}_{33}\text{N}_5\text{O}_3 + \text{H}$] $^+$.

Step 2: To *tert*-butyl (1-(4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoyl)piperidin-4-yl)methylcarbamate (70 mg, 0.131 mmol) was added 20% TFA in CHCl_3 (3 mL). The reaction mixture was stirred at room temperature for 20 min then concentrated under reduced pressure. To the residue was added saturated sodium bicarbonate (10 mL) and extracted with EtOAc (3 x 30 mL). The combined organic was dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/ H_2O /formic acid 0.1%) to afford 4-(6-(4-(4-(aminomethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyridin-3-yl)benzonitrile (26 mg, 46%, AUC HPLC 99%) as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 8.76 (s, 1H), 8.55 (s, 1H), 7.95-7.90 (m, 4H), 7.89 (s, 1H), 7.81-7.76 (m, 3H), 7.54 (d, $J = 8.4$ Hz, 2H), 4.80-4.65 (m, 1H), 3.90-3.75 (m, 1H), 3.30-3.15 (m, 1H), 3.00-2.85 (m, 1H), 2.86 (d, $J = 6.8$ Hz, 2H), 2.00-1.85 (m, 2H), 1.85-1.70 (m, 1H), 1.40-1.20 (m, 2H); ^{13}C NMR (100 MHz, CD_3OD): δ 171.96, 147.44, 140.03, 136.66, 134.84, 134.72, 134.39, 129.44, 128.84, 128.61, 128.50, 127.80, 126.56, 122.83, 119.50, 118.48, 112.72, 45.70, 36.32; MS (ESI) m/z 436 [$\text{C}_{27}\text{H}_{25}\text{N}_5\text{O} + \text{H}$] $^+$.

Example 124: 4-(6-(4-(4-(2-aminoethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-*a*]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-*a*]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methyl morpholine (129 μL , 1.18 mmol) and *tert*-butyl 2-(piperidin-4-yl)ethylcarbamate (101 mg,

0.441 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (10 mL) and extracted with EtOAc (3×20 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 97:3) to afford *tert*-butyl 2-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)ethylcarbamate (79 mg, 49%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.70 (s, 1 H), 8.05-7.97 (m, 4 H), 7.85-7.78 (m, 3 H), 7.75-7.70 (m, 1 H), 7.48 (d, *J* = 8.0 Hz, 2 H), 6.80-6.73 (m, 1 H), 4.55-4.35 (bs, 1 H), 3.70-3.50 (bs, 1 H), 3.10-2.90 (m, 3 H), 2.85-2.70 (bs, 1 H), 1.85-1.50 (m, 3 H), 1.37 (s, 9 H), 1.35-1.30 (m, 2 H), 1.20-1.00 (m, 3 H); MS (ESI) *m/z* 550 [C₃₃H₃₅N₅O₃ + H]⁺.


Step 2: To *tert*-butyl 2-(1-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl)ethylcarbamate (70 mg, 0.127 mmol) was added 20% TFA in CHCl₃ (3 mL). The reaction mixture was stirred at room temperature for 20 min then it was concentrated under reduced pressure. To the residue was added saturated sodium bicarbonate (10 mL) and extracted with EtOAc (3 x 30 mL). The combined organic was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN/water/formic acid 0.1%) to afford 4-(6-(4-(4-(2-aminoethyl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (16 mg, 29%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.75 (s, 1H), 8.55 (s, 1H), 7.95-7.90 (m, 4H), 7.89 (s, 1H), 7.81-7.76 (m, 3H), 7.53 (d, *J* = 8.4 Hz, 2H), 4.75-4.60 (m, 1H), 3.90-3.75 (m, 1H), 3.25-3.10 (m, 1H), 3.00-2.80 (m, 3H), 1.95-1.80 (m, 1H), 1.80-1.65 (m, 2H), 1.65-1.60 (m, 2H), 1.40-1.15 (m, 2H); ¹³C NMR (400 MHz, CD₃OD) δ 171.89, 147.44, 139.93, 136.82, 134.84, 134.71, 134.39, 129.43, 128.81, 128.64, 128.48, 127.81, 126.55, 122.81, 119.50, 118.47, 112.71, 38.53, 35.50, 34.61; MS (ESI) *m/z* 450 [C₂₈H₂₇N₅O + H]⁺.

Example 125: 4-(6-(4-(4-(pyridin-4-yl)piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methylmorpholine (129 μ L, 1.18 mmol) and 1-(4-pyridinyl)-piperazine (58 mg, 0.441 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 92:8) to afford 4-(6-(4-(4-(pyridin-4-yl)piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (34 mg, 24%, AUC HPLC 96%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.77 (s, 1H), 8.17 (d, *J* = 7.2 Hz, 2H), 7.96-7.90 (m, 4H), 7.90 (s, 1H), 7.83 (d, *J* = 8.0 Hz, 2H), 7.79 (s, 2H), 7.63 (d, *J* = 8.0 Hz, 2H), 7.08 (d, *J* = 7.2 Hz, 2H) 3.95-3.68 (m, 8H); ¹³C NMR (400 MHz, DMSO-*d*₆): δ 168.98, 156.53, 145.76, 140.38, 138.13, 135.11, 134.76, 133.51, 133.19, 127.91, 127.79, 127.10, 125.70, 125.55, 124.46, 121.92, 118.81, 117.81, 109.83, 107.56; MS (ESI) *m/z* 485 [C₃₀H₂₄N₆O + H]⁺.


Example 126: 4-(6-(4-(4-benzoylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (100 mg, 0.294 mmol) in DMF (1.5 mL) were added HATU (168 mg, 0.441 mmol), *N*-methylmorpholine (129 μ L, 1.18 mmol) and 1-benzoylpiperazine (67 mg, 0.441 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was


diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 92:8) to afford 4-(6-(4-(4-benzoylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (75 mg, 50%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.75 (s, 1H), 7.95-7.89 (m, 4H), 7.88 (s, 1H), 7.82-7.55 (m, 4H), 7.59 (d, *J* = 8.0 Hz, 2H), 7.63-7.43 (m, 5H), 4.00-3.40 (m, 8H); ¹³C NMR (100 MHz, CD₃OD): δ 172.90, 172.34, 147.49, 140.29, 136.50, 136.11, 134.87, 134.76, 134.43, 131.43, 129.87, 129.48, 129.22, 128.63, 128.59, 128.21, 127.85, 126.61, 122.92, 119.55, 118.51, 112.76; MS (ESI) *m/z* 512 [C₃₂H₂₅N₅O + H]⁺.

Example 127: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(4-(4-methylpiperidin-1-yl)phenyl)benzamide




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μL, 0.824 mmol) and 4-(4-methylpiperidin-1-yl)aniline (47 mg, 0.309 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was washed with cold CHCl₃/MeOH mixture and the precipitates obtained were further washed with cold CH₃CN to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(4-(4-methylpiperidin-1-yl)phenyl)benzamide (17 mg, 17%, AUC HPLC 96%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.80 (s, 1H), 8.07 (d, *J* = 8.4 Hz, 2H), 7.98-7.91 (m, 4H), 7.90 (s, 1H), 7.85 (d, *J* = 8.4 Hz, 2H), 7.80 (s, 2H), 7.69 (d, *J* = 8.4 Hz, 2H), 7.30 (d, *J* = 8.4 Hz, 2H), 3.60-3.45 (m, 2H), 3.13 (bs, 1H), 3.10-2.95 (m, 2H), 2.85 (s, 3H), 2.20-2.05 (m, 2H), 2.03-1.85 (m, 2H); ¹³C NMR (400 MHz, DMSO-*d*₆): δ 164.93, 145.82, 139.73, 139.29, 137.63, 135.20, 134.03, 133.53, 133.21, 128.37, 127.78, 126.95, 126.70, 125.55, 124.51, 122.08, 120.67, 118.82, 117.85, 109.86, 37.72; MS (ESI) *m/z* 512 [C₃₃H₂₉N₅O + H]⁺.

Example 128: 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-N-(4-((4-methylpiperazin-1-yl)methyl)phenyl)benzamide



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μ L, 0.824 mmol) and 4-((4-methylpiperazin-1-yl)methyl)aniline (51 mg, 0.309 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-*N*-(4-((4-methylpiperazin-1-yl)methyl)phenyl)benzamide (40 mg, 37%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.78 (s, 1H), 8.05 (d, *J* = 8.4 Hz, 2H), 7.96-7.90 (m, 4H), 7.84 (s, 1H), 7.83 (d, *J* = 8.4 Hz, 2H), 7.80-7.76 (m, 2H), 7.68 (d, *J* = 8.8 Hz, 2H), 7.34 (d, *J* = 8.4 Hz, 2H), 3.56 (s, 2H), 2.70-2.40 (m, 8H), 2.36 (s, 3H); ¹³C NMR (100 MHz, CD₃OD): δ 168.12, 147.47, 141.62, 139.17, 135.76, 134.82, 134.74, 134.53, 134.40, 131.08, 129.56, 129.44, 128.49, 128.30, 127.79, 126.61, 122.95, 122.18, 119.51, 118.48, 112.72, 63.14, 55.55, 53.15, 45.65; MS (ESI) *m/z* 527 [C₃₃H₃₀N₆O + H]⁺.


Example 129: 4-((4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazin-1-yl)methyl)benzamide



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (113 μ L, 0.824 mmol) and 4-(1-piperazinylmethyl)benzamide dihydrochloride (72 mg, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18


h, then it was diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-((4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazin-1-yl)methyl)benzamide (7 mg, 7%, AUC HPLC 99%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.80 (s, 1H), 8.03-7.97 (m, 5H), 7.95 (bs, 1H), 7.90-7.80 (m, 5H), 7.96-7.71 (m, 1H), 7.51 (d, *J* = 7.6 Hz, 2H), 7.43 (bs, 2H), 7.34 (bs, 1H), 3.80-3.40 (m, 8H), 2.69 (s, 2H); MS (ESI) *m/z* 541 [C₃₃H₂₈N₆O₂ + H]⁺.

10 Example 130: 4-(6-(4-(4-methyl-3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μL, 0.824 mmol) and 1-methylpiperazin-2-one (37 mg, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (15 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96:4) to afford 4-(6-(4-(4-methyl-3-oxopiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (64 mg, 57%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 8.76 (bs, 1H), 7.97-7.89 (m, 4H), 7.88 (s, 1H), 7.84-7.78 (m, 2H), 7.77 (d, *J* = 1.2 Hz, 2H), 7.64-7.57 (m, 2H), 4.40-4.10 (m, 2H), 4.10-3.60 (m, 2H), 3.57-3.42 (m, 2H), 3.01 (s, 3H); ¹³C NMR (100 MHz, CD₃OD): δ 171.72, 147.44, 140.53, 135.51, 134.81, 134.72, 134.38, 129.42, 129.27, 128.56, 128.52, 127.79, 126.58, 122.93, 119.50, 118.47, 112.70, 34.68; MS (ESI) *m/z* 436 [C₂₆H₂₁N₅O₂ + H]⁺.

Example 131: 4-(6-(4-(4-phenylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μ L, 0.824 mmol) and 4-phenylpiperazine (38 μ L, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent EtOAc) to afford 4-(6-(4-(4-phenylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (83 mg, 84%, AUC HPLC 98%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (s, 1H), 8.04-7.98 (m, 5H), 7.88-7.80 (m, 3H), 7.66-7.62 (m, 1H), 7.56 (d, *J* = 8.0 Hz, 2H), 7.37-7.20 (m, 2H), 6.96 (d, *J* = 8.0 Hz, 2H), 6.83-6.79 (m, 1H), 3.90-3.45 (m, 4H), 3.30-3.10 (m, 4H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 168.66, 150.74, 145.77, 137.86, 135.18, 135.10, 133.52, 133.19, 128.99, 127.79, 127.77, 127.09, 125.77, 125.57, 124.45, 121.88, 119.40, 118.81, 117.79, 115.94, 109.81; MS (ESI) *m/z* 484 [C₃₁H₂₅N₅O + H]⁺.


Example 132: 4-(6-(4-(4-benzylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μ L, 0.824 mmol) and 4-benzylpiperazine (43 μ L, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h, then it was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The residue was purified by column


chromatography (silica gel, eluent $\text{CHCl}_3/\text{MeOH}$ 96:4) to afford 4-(6-(4-(4-benzylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (48 mg, 47%, AUC HPLC 97%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.79 (bs, 1H), 8.03-7.96 (m, 5H), 7.84-7.78 (m, 3H), 7.74-7.69 (m, 1H), 7.49 (d, $J = 8.4$ Hz, 2H), 7.36-7.28 (m, 4H), 7.28-7.22 (m, 1H), 3.75-3.55 (m, 2H), 3.51 (s, 2H), 3.50-3.33 (m, 2H), 2.49-2.30 (m, 4H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 168.54, 145.75, 137.77, 137.74, 135.28, 135.09, 133.51, 133.18, 128.89, 128.21, 127.76, 127.70, 127.04, 125.75, 125.54, 124.43, 121.83, 118.81, 117.79, 109.80, 61.83; MS (ESI) m/z 498 [$\text{C}_{32}\text{H}_{27}\text{N}_5\text{O} + \text{H}$] $^+$.

10 Example 133: 4-(6-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (70 mg, 0.206 mmol) in DMF (1.0 mL) were added HATU (117 mg, 0.309 mmol), *N*-methyl morpholine (90 μL , 0.824 mmol) and 1-methyl-4-(piperidin-4-yl)piperazine (45 mg, 0.247 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 18 h then, was diluted with water (15 mL) and extracted with EtOAc (3 $\times$ 30 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The residue was purified by preparative HPLC (C18, eluent ACN, water, formic acid 0.1%) to afford 4-(6-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (23 mg, 22%, AUC HPLC 99%) as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 8.76 (br s, 1H), 7.97-7.90 (m, 4H), 7.89 (s, 1H), 7.81-7.76 (m, 4H), 7.54 (d, $J = 8.4$ Hz, 2H), 4.75-4.60 (m, 1H), 3.90-3.75 (m, 1H), 3.25-3.10 (m, 1H), 3.00-2.85 (m, 1H), 2.80-2.55 (m, 9H), 2.40 (s, 3H), 2.10-1.95 (m, 1H), 1.95-1.80 (m, 1H), 1.65-1.40 (m, 2H); ^{13}C NMR (100 MHz, CD_3OD): δ 171.87, 140.01, 136.75, 134.89, 134.75, 134.44, 129.48, 128.89, 128.71, 128.53, 127.89, 122.89, 119.56, 118.51, 112.75, 62.65, 55.83, 45.55; MS (ESI) m/z 505 [$\text{C}_{31}\text{H}_{32}\text{N}_6\text{O} + \text{H}$] $^+$.

Example 134: 4-(6-(4-(4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (0.2 g, 0.58 mmol) in DMF (5 mL) were added HATU (0.33 g, 0.88 mmol), *N*-methyl morpholine (0.18 g, 1.76 mmol) and 4-(piperidin-4-yl)morpholine (108.4 mg, 0.70 mmol). The reaction mixture was stirred at 0 °C to room temperature under inert atmosphere for 16 h, then it was diluted with water (15 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) and followed by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford 4-(6-(4-(4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (140 mg, 50%, AUC HPLC 98.7%) as a white solid; m.p. 131-141°C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.6 (s, 1H), 8.05-7.91 (m, 5H), 7.80 (m, 2H), 7.48 (m, 1H), 4.45 (bs, 1H), 3.65-3.56 (m, 5H), 3.01-2.82 (m, 2H), 2.46-2.39 (m, 5H), 1.85-1.75 (m, 2H), 1.38 (bs, 2H); MS (ESI) *m/z* 492.23 [C₃₀H₂₉N₅O₂ + H]⁺.


Example 135: 4-(6-(4-(4-methyl-4-(5-methyl-1,3,4-oxadiazol-2-yl)piperidin-4-yl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (150 mg, 0.44 mmol) and 2-methyl-5-(4-methylpiperidin-4-yl)-1,3,4-oxadiazole B (105 mg, 0.48 mmol) in DMF (5 mL) was added HATU (250 mg, 0.66 mmol) followed by NMM (0.09 mL, 0.88

mmol) and the reaction mixture was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain 4-(6-(4-(4-methyl-4-(5-methyl-1,3,4-oxadiazol-2-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (60 mg, 27%, AUC HPLC 95.3%) as a white solid, m.p : 150-160 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.80 (s, 1H), 8.03-7.98 (m, 5H), 7.82 (d, *J* = 7.9 Hz, 4H), 7.51 (d, *J* = 8.3 Hz, 2H), 4.08 (bs, 1H), 3.54 (bs, 1H), 3.31 (bs, 2H), 2.48 (s, 3H), 2.07 (bs, 2H), 1.70 (bs, 2H), 1.35 (s, 3H); MS (ESI) *m/z* 503.34 [C₃₀H₂₆N₆O₂ + H]⁺.


10 Example 136: 4-(6-(4-(4-methyl-4-(1H-pyrazol-3-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (150 mg, 0.442 mmol) in DMF (5.0 mL) were added HATU (251 mg, 0.663 mmol), *N*-methyl morpholine (0.17 mL, 1.547 mmol) and tert-butyl 4-methyl-4-(1H-pyrazol-3-yl)piperidine hydrochloride (126 mg, 0.53 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then was diluted with H₂O (15 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford 4-(6-(4-(4-methyl-4-(1H-pyrazol-3-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (80 mg, 38%, AUC HPLC 97.2%) as a white solid; mp. 223-226 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.50 (s, 1H), 7.83-7.73 (m, 6H), 7.57-7.50 (m, 6H), 6.18 (s, 1H), 4.21 (s, 1H), 3.58 (s, 1H), 3.38 (s, 2H), 2.22-2.16 (m, 2H), 1.85-1.77 (m, 2H), 1.32 (s, 3H); MS (ESI) *m/z* 487.5 [C₃₀H₂₆N₆O + H]⁺.

25

Example 137: 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)-7-methylimidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: A solution of 5-bromo-4-methylpyridin-2-amine (8 g, 42.78 mmol) and
 5 chloroacetaldehyde (approx. 50 %wt) solution in water (27.52 ml, 214 mmol) in ethanol (50 ml) was heated at 100 °C under N₂ for 16 h. The solvent was removed and the residue was dissolved in EtOAc (100 ml). The organic layer was washed with sat. NaHCO₃ (2x50 ml), water (2x50 ml), brine (2x100 ml), dried over Na₂SO₄ and the solvent was removed completely under vacuum to afford 6-bromo-7-methylimidazo [1,2-a]pyridine (8.5 g, 94 %)
 10 as a brown solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.93 (s, 1H), 7.85 (s, 1H), 7.57 (d, *J* = 7.56 Hz, 2H), 2.38 (s, 3H); LC-MS (95%) *m/z* 211 [C₈H₇BrN₂ + H]⁺.

Step 2: To a solution of 6-bromo-7-methylimidazo(1,2-a)pyridine (9 g, 43.0 mmol) and anhydrous sodium acetate (9.52 g, 116.1 mmol) in MeOH (100 mL) at 0 °C was added iodine
 15 (12.0 g, 47.3 mmol). The reaction mixture was stirred at rt for 20 h. The precipitate was collected by filtration and washed with MeOH to afford 6-bromo-3-iodo-7-methylimidazo[1,2-a]pyridine (6 g, 41%) as a light grey solid. ¹H NMR (400 MHz, CDCl₃) δ 8.30 (s, 1H), 7.64 (s, 1H) 7.49 (s, 1H) 2.50 (s, 3H); MS (ESI) *m/z* 336.7 [M+H]⁺.

Step 3: To mixture of 6-bromo-3-iodo-7-methylimidazo[1,2-a]pyridine (5 g, 14.9 mmol), 4-cyanophenylboronic acid (2.41 g, 16.4 mmol) and K₃PO₄ (6.32 g, 29.8 mmol) in 1,4-dioxane (200 mL) and water (50 mL) was added Pd(PPh₃)₄ (860 mg, 0.74 mmol). The reaction mixture was heated at 90 °C for 1 h. The reaction mixture was diluted with water and extracted with EtOAc (2x100 ml) and washed with water (2x50 ml) and brine solution (2x50
 25 ml). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 95:5) to give 4-(6-bromo-7-methylimidazo[1,2-a]pyridin-3-yl)benzonitrile (2.72 g, 59%). ¹H NMR (400 MHz, CDCl₃) δ

8.52 (s, 1H), 7.83 (d, $J = 7.82$ Hz, 2H), 7.74 (s, 1H), 7.68 (d, $J = 7.6$ Hz, 2H), 2.49 (s, 3H); MS (ESI) m/z 312 $[M + H]^+$.

Step 4: To a mixture of 4-(6-bromo-7-methylimidazo[1,2-a]pyridin-3-yl)benzonitrile (2.7 g, 8.68 mmol), 4-(ethoxycarbonyl) phenylboronic acid (1.85 g, 9.54 mmol), K_3PO_4 (3.68 g, 17.36 mmol) in 1,4-dioxane (200 mL) and water (50 mL), was added $Pd(PPh_3)_4$ (501 mg, 0.43 mmol). The reaction mixture was heated at 90 °C for 1 h and was diluted with water, extracted with EtOAc (2x100 mL) and washed with water (2x50 mL) and brine solution (2 x 50 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent $CHCl_3/MeOH$ 95:5) to 1.6 g (50 %) of ethyl 4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoate. 1H NMR (400 MHz, CD_3OD) δ 9.21 (s, 1H), 8.95 (s, 1H), 8.21 (s, 1H), 8.17–8.12 (m, 5H), 8.08 (s, 1H), 7.91 (d, $J = 8.0$ Hz, 1H), 7.82 (t, $J = 8.4$ Hz, 1H), 4.03 (q, $J = 7.2$ Hz, 2H), 2.39 (s, 3H) 2.1–2.23 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 353 $[M + H]^+$.

15

Step 5: To a solution of 4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoate (1.6 g, 3.26 mmol) in THF (20 mL) was added LiOH (0.411 g, 9.79 mmol) in water (15 mL) and MeOH (15 mL) at rt and stirred for 16 h. TLC indicated absence of SM and formation of a polar spot. The reaction mixture was concentrated under reduced pressure to afford 4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoic acid (900 mg, 64%) as an off white solid which was used in the next step with out purification. 1H NMR (400 MHz, CD_3OD) δ 9.21 (s, 1H), 8.95 (s, 1H), 8.21 (s, 1H), 8.17–8.12 (m, 5H), 8.08 (s, 1H), 7.91 (d, $J = 8.0$ Hz, 1H), 7.82 (t, $J = 8.4$ Hz, 1H), 4.03 (q, $J = 7.2$ Hz, 2H), 2.39 (s, 3H) 2.1–2.23 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 353 $[M + H]^+$.

25

Step 6: To a solution of 4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoic acid (250 mg, 0.7 mmol) in DMF (15.0 mL) were added HATU (400 mg, 1.05 mmol), *N*-methyl morpholine (0.155 mL, 1.4 mmol) and *tert*-butyl 4-methylpiperidin-4-ylcarbamate (165 mg, 0.77 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h then, was diluted with water (25 mL) and extracted with EtOAc (3x25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (300 mg, 77%,


30

LC-MS 92%) as an off-white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.41 (s, 1 H), 7.95-7.94 (m, 5 H), 7.67 (s, 1 H), 7.54 (d, $J = 12.0$ Hz, 2 H), 7.44 (d, $J = 12.0$ Hz, 2 H), 3.79 (bs, 2 H), 3.45 (bs, 3 H), 2.29 (s, 3 H), 1.58-1.24 (m, 6 H), 1.09 (s, 9 H); MS (ESI) m/z 510 $[\text{M} + \text{H}]^+$.

5

Step 7: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (300 mg, 0.546 mmol) in dichloromethane (10 mL) was added TFA (3 mL) in dichloromethane (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL) and NaHCO_3 (100 mL) and
 10 extracted with CH_2Cl_2 (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford 4-(6-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)-7-methylimidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 46%, AUC HPLC 95.5%) as an off-white solid; m.p. 124-132 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.41 (s, 1 H), 7.95-7.94 (m, 5 H), 7.67 (s, 1 H), 7.54 (d, $J = 12.0$ Hz, 2 H), 7.44 (d, $J = 12.0$ Hz, 2 H), 3.79 (bs, 2
 15 H), 3.45 (bs, 3 H), 2.29 (s, 3 H), 1.58-1.24 (m, 6 H), 1.09 (s, 3 H); MS (ESI) m/z 450.31 $[\text{C}_{28}\text{H}_{27}\text{N}_5\text{O} + \text{H}]^+$.

Example 138: 4-(7-methyl-6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




20

To a solution of 4-(3-(4-cyanophenyl)-7-methylimidazo[1,2-a]pyridin-6-yl)benzoic acid (200 mg, 0.566 mmol) in DMF (10.0 mL) were added HATU (322 mg, 0.849 mmol), *N*-methylmorpholine (0.158 mL, 1.132 mmol) and 1-methylpiperazin (0.068 mL, 0.623 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then it was
 25 diluted with water (25 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) and followed by preparative HPLC to afford 4-(7-methyl-6-(4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 50%, AUC HPLC 97.6%)
 30 as an off-white solid; m.p. 162-164 °C. ^1H NMR (400 MHz, CD_3OD) δ (ppm): 8.47 (s, 1 H),

7.85-7.81 (m, 5 H), 7.57-7.53 (m, 5 H), 3.83 (bs, 2 H), 3.53 (bs, 2 H), 2.54-2.34 (bm, 4 H), 2.34 (s, 3 H), 2.33 (s, 3 H); MS (ESI) m/z 436.27 [$C_{27}H_{25}N_5O + H$]⁺.

Example 139: 4-(6-(2-methyl-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 2-methyl-4-bromobenzoic acid (2 g, 9.30 mmol) in DMF (6 mL) was added NMM (1.87g, 18.604 mmol) followed by addition of HATU (5.3 g, 13.95 mmol) at rt and stirred for 30 min. 1-methylpiperazine (1.39 g, 13.95 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain (4-bromo-2-methylphenyl)(4-methylpiperazin-1-yl)methanone (1.9 g, 70%, AUC-LC-MS 98%) as a brown solid.

Step 2: A mixture of (4-bromo-2-methylphenyl)(4-methylpiperazin-1-yl)methanone (1.9 g, 6.39 mmol), Bis(pinacolato)diboron (1.74 g, 6.84 mmol), KOAc (41.88 g, 19.19 mmol) in 1,4-dioxane (30 mL) was degassed with argon for 30 min. $PdCl_2dppf$ (140.5 mg, 0.19 mmol) and $dppf$ (106.4 mg, 0.19 mmol) were added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude (2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (~1.3 g) as a brown liquid which was used in next step without further purification. MS (ESI) m/z 345.

Step 3: To a mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (3.95 g, 13.33 mmol), (2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (1.3 g, 3.78 mmol), K_3PO_4 (1.6 g, 7.56 mmol) in 1,4-dioxane (30 mL) and water (6mL) was added $Pd(PPh_3)_4$ (218 mg, 0.188 mmol). The reaction mixture was heated at 90 °C for 1 h and was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated

under reduced pressure to obtain the crude product. The crude product was purified by column chromatography followed by preparative HPLC to afford 4-(6-(2-methyl-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (45 mg, AUC HPLC >99%); White solid; m.p. 169-173 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.46 (s, 1 H), 7.85-7.80 (m, 3 H), 7.77-7.72 (m, 2 H), 7.52-7.48 (m, 1 H), 7.39-7.37 (m, 2 H), 7.28 (s, 1 H), 3.86 (bs, 2 H), 3.31 (bs, 2 H), 2.51 (bs, 2 H), 2.39 (s, 3 H), 2.33 (s, 5 H); MS (ESI) *m/z* 436 [C₂₇H₂₅N₅O + H]⁺.

10 Example 140: 4-(6-(2-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 2-fluoro-4-bromobenzoic acid (1 g, 4.56 mmol) in DMF (3 mL) was added NMM (922 mg, 9.13 mmol) followed by addition of HATU (2.6 g, 6.85 mmol) at rt and stirred for 30 min. 1-Methylpiperazine (684 mg, 6.85 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain (4-bromo-2-fluorophenyl)(4-methylpiperazin-1-yl)methanone (1.6 g, 100%, LC-MS 87%) as a brown solid. MS (ESI) *m/z* 302.

20 Step 2: A mixture of (4-bromo-3-fluorophenyl)(4-methylpiperazin-1-yl)methanone (1.6 g, 5.31 mmol), Bis(pinacolato)diboron (1.45 g, 5.68 mmol), KOAc (1.56 g, 15.95 mmol) in 1,4-dioxane (30 mL) was degassed with argon for 30 min. PdCl₂dppf (116 mg, 0.159 mmol) and dppf (88.4mg, 0.159 mmol) were added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 2-fluoro-4-(4-methylpiperazine-1-carbonyl)phenylboronic acid (900 mg, LC-MS 55%) as a brown liquid which was used in next step without purification.

30

Step 3: To a solution of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (400 mg, 1.342 mmol), 2-fluoro-4-(4-methylpiperazine-1-carbonyl)phenylboronic acid (900 mg, 3.30 mmol), K_3PO_4 (711 mg, 3.35 mmol) in a mixture of 1,4-dioxane (12 mL) and water (3 mL) was added $Pd(PPh_3)_4$ (77.5 mg, 0.067 mmol). The reaction mixture was heated at 90 °C for 1 h, was diluted with water, extracted with EtOAc and the organic phase was washed with water and brine solution, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced. The residue was purified by preparative HPLC to give 4-(6-(2-methyl-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (35 mg, AUC HPLC 99.56%); m.p. 170-174 °C. 1H NMR (400 MHz, CD_3OD) δ (ppm): 8.77 (s, 1 H), 7.91 (m, 5 H), 7.79-7.66 (m, 3 H), 7.38-7.36 (m, 2 H), 3.80 (bs, 2 H), 3.54 (bs, 2 H), 2.61 (bs, 4 H), 2.40 (s, 3 H); MS (ESI) m/z 440.40 [$C_{26}H_{22}FN_5O + H$] $^+$.

Example 141: 4-(6-(3-chloro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (436 mg, 1.46 mmol), 3-chloro-4-(ethoxycarbonyl)phenylboronic acid (500 mg, 1.61 mmol), K_3PO_4 (775 mg, 3.65 mmol) in 1,4-dioxane (14 mL) and water (3.5 mL), was added $Pd(PPh_3)_4$ (85 mg, 0.0736 mmol). The reaction mixture was heated at 90 °C for 1 h, then was diluted with water and extracted with EtOAc. The organic layer was washed in turn with water and brine then was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain ethyl 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (1 g, 90%). The product was used in the next step without further purification.

Step 2: To a solution of ethyl 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (1 g, 2.49 mmol) in THF (10 mL) and MeOH (10 mL) was added LiOH (261 mg, 6.23 mmol) in water (4 mL). The reaction mixture was stirred for 16 h at rt and was concentrated under reduced pressure then diluted with water and acidified with potassium bisulphate. The aqueous layer was extracted with EtOAc, which was dried and concentrated

under reduced pressure to give 300 mg (38%) of 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (LC-MS 68%) as a yellow solid.

Step 3: To a solution of 2-chloro-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (300 mg, 0.80 mmol) in DMF (2 mL) was added NMM (162 mg, 1.60 mmol) followed by addition of HATU (458 mg, 1.20 mmol) at rt and stirred for 30 min. 1-Methylpiperazine (120 mg, 1.21 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with water and filtered to obtain a solid which was further purified by preparative HPLC to afford 4-(6-(3-chloro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (60 mg, 24%) as an off white solid; m.p. 143-157 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.46 (s, 1 H), 7.85-7.79 (m, 4 H), 7.73-7.71 (m, 2 H), 7.56 (s, 1 H), 7.49-7.38 (m, 3 H), 3.88-3.83 (m, 2 H), 3.34 (bs, 2 H), 2.51-2.41 (m, 4 H), 2.33 (s, 3 H); MS (ESI) *m/z* 456.32 [C₂₆H₂₂ClN₅O + H]⁺.

Example 142: 4-(6-(3-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of 4-(ethoxycarbonyl)-3-fluorophenylboronic acid (1.29 g, 6.091 mmol), K₃PO₄ (3.22 g, 15.22 mmol) in 1,4-dioxane (20 mL), and water (3 mL) was added 6-bromoimidazo[1,2-a]pyridine (1 g, 5.07 mmol) and degassed with argon for 30 min. (A-Phos)₂PdCl₂ (293 mg, 0.25 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for overnight. The reaction mixture was washed with water. The organic layer was concentrated and the crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96.5: 3.5) to afford 2-fluoro-4-(imidazo[1,2-a]pyridin-6-yl)benzoate (1.16 g, 81%, AUC LC-MS 84%). ¹H NMR (300 MHz, DMSO-*d*₆) ppm δ 9.1 (s, 1 H), 7.9 (m, 2 H), 7.6 (m, 5 H), 4.3 (q, *J* = 6.9 Hz, 2 H), 1.3 (t, *J* = 6.9 Hz, 3 H); MS (ESI) *m/z* 285 (M+1).

Step 2: To a solution of 2-fluoro-4-(imidazo[1,2-a]pyridin-6-yl)benzoate (1.16 g, 4.10 mmol) in DMF (20 mL) and added to NIS (1.10 g, 4.92 mmol). The reaction mixture stirred for 3 h at 100 °C. Water was added to the reaction mixture to induce precipitation of the product
5 which was filtered and dried under reduced pressure to give ethyl 2-fluoro-4-(3-iodoimidazo[1,2-a]pyridin-6-yl)benzoate (1.2 g, 75%, AUC LC-MS 92.5%). ¹H NMR (300 MHz, DMSO-*d*₆) δ 8.5 (s, 1 H), 7.8 (m, 6 H), 4.3 (q, *J* = 6.9 Hz, 2 H), 1.3 (t, *J* = 6.9 Hz, 3 H); MS (ESI) *m/z* 410 (M+1).


10 Step 3: A solution of ethyl 2-fluoro-4-(3-iodoimidazo[1,2-a]pyridin-6-yl)benzoate (900 mg, 2.23 mmol) and 4-cyanophenylboronic acid (394 mg, 2.68 mmol), K₃PO₄ (949 mg, 4.47 mmol) in 1,4-dioxane (10 mL), and water (1 mL) was degassed with argon for 30 min. (A-Phos)₂PdCl₂ (129 mg, 0.111 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for overnight. TLC indicated absence of SM. The
15 reaction mixture was washed with water. The organic layer was concentrated and purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96.5:3.5) to afford ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoate (600 mg, 71%). ¹H NMR (300 MHz, DMSO-*d*₆) ppm δ 8.9 (s, 1 H), 7.8 (m, 6 H), 4.3 (q, *J* = 6.9 Hz, 2 H), 1.3 (t, *J* = 6.9 Hz, 3 H); MS (ESI) *m/z* 386 (M+1).

20 Step 4: To a solution of ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoate (600 mg, 1.55 mmol) dissolved in THF (5 mL), water (5 mL), EtOH (5 mL) was added LiOH (130 mg, 3.11 mmol). The reaction mixture was stirred for 1 h. After completion of the reaction as indicated by TLC, the reaction mixture was added to citric acid
25 and washed with water and extracted with EtOAc. The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by recrystallization to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoic acid (450 mg, 81%, AUC LC-MS 83.2%). ¹H NMR (300 MHz, DMSO-*d*₆) ppm δ 8.07 (s, 1 H), 7.6 (d, *J* = 8.0 Hz, 2 H), 7.5 (m,
30 2 H), 7.4 (m, 4 H), 7.3 (d, *J* = 11.2 Hz, 2 H); MS (ESI) *m/z* 358 (M+1).

Step 5: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoic acid (450 mg, 1.26 mmol) in DMF (5 mL) was added NMM (0.28 mL, 2.52 mmol) followed by addition of HATU (957 mg, 2.52 mmol) at rt and stirred for 20 min. 1-Methylpiperazine

(0.138 mL, 1.38 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain 4-(6-(3-fluoro-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (150 mg, 27.1%, AUC HPLC 97.4%) as an off white solid; m.p. 176-185 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.86 (s, 1 H), 8.01 (m, 5 H), 7.84-7.67 (m, 4 H), 7.49 (t, $J = 8.0$ Hz, 1 H), 3.65 (bs, 2 H), 3.26 (bs, 4 H), 2.37 (bs, 2 H), 2.27 (bs, 4 H), 2.20 (s, 3 H); MS (ESI) m/z 440 $[\text{C}_{26}\text{H}_{22}\text{FN}_5\text{O} + \text{H}]^+$.

10 Example 143: 4-(6-(2-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




- Step 1: To a mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (3.95 g, 13.33 mmol), (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (4.9 g, 13.61 mmol), K_3PO_4 (5.8 g, 27.22 mmol) in 1,4-dioxane (80 mL) and water (20 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (785.9 mg, 0.68 mmol). The reaction mixture was heated at 90 °C for 1 h. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product.
- 20 The crude product was purified by column chromatography and by preparative HPLC of 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (54 mg, AUC HPLC 98.1%) as an off-white solid; m.p. 121-134 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.46 (s, 1 H), 7.84-7.71 (m, 4 H), 7.49 (d, $J = 15.0$ Hz, 1 H), 7.34 (d, $J = 10.4$ Hz, 1 H), 7.13 (d, $J = 10.4$ Hz, 1 H), 7.01 (s, 1 H), 3.90 (s, 3 H), 3.85 (bs, 2 H), 3.32 (bs, 2 H), 2.61-2.40 (m, 4 H), 2.32 (s, 3 H); MS (ESI) m/z 452 $[\text{C}_{27}\text{H}_{25}\text{N}_5\text{O}_2 + \text{H}]^+$.
- 25

Step 2: A solution of 4-(6-(3-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (600 mg, 1.33 mmol) in CH_2Cl_2 (10 mL) was cooled to 0 °C followed by the addition of BBr_3 (738 mg, 2.2 mmol). The reaction mixture was stirred at rt for 2 h. The reaction mixture was cooled to 0 °C and quenched by

30

dropwise addition of a saturated aqueous solution of NaHCO_3 (100 mL) followed by extraction with EtOAc (3 x 100 mL). The organic layer was washed in turn with water and brine then, was dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent $\text{CHCl}_3/\text{MeOH}$ 99/1 to 97/3) and by purification by preparative HPLC to afford 4-(6-(3-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (80 mg, 20%, AUC HPLC 98.7%) as an off-white solid; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.69 (s, 1 H), 8.28 (s, 1 H), 8.00 (m, 4 H), 7.82-7.79 (m, 1 H), 7.64-7.61 (m, 1 H), 7.21-7.15 (m, 3 H), 3.39 (bs, 4 H), 2.29 (bs, 2 H), 2.18 (s, 3 H); MS (ESI) m/z 438.2 $[\text{C}_{26}\text{H}_{23}\text{N}_5\text{O}_2 + \text{H}]^+$.

Example 144: 4-(6-(3-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: A mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (2.38 g, 8 mmol), (3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (2.88 g, 8 mmol), K_3PO_4 (3.4 g, 2 mmol) in 1,4-dioxane (30 mL) and water (6 mL) was degassed with argon for 30 min. $\text{Pd}(\text{PPh}_3)_4$ (462 mg, 0.4 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 1 h. The reaction mixture was diluted with water and extracted with EtOAc, washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography to obtain 4-(6-(2-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (1 g, 28%, AUC 99.3%) as an off-white solid; m.p. 123-178 °C. ^1H NMR (400 MHz, CD_3OD) δ (ppm): 8.72 (s, 1 H), 7.92-7.86 (m, 5 H), 7.70-7.52 (m, 3 H), 7.17 (s, 1 H), 7.10 (d, $J = 8.0$ Hz, 1 H), 3.90 (s, 3 H), 3.81-3.56 (br. m, 4 H), 2.60 (bs, 4 H), 2.41 (s, 3 H); MS (ESI) m/z 452.36 $[\text{C}_{27}\text{H}_{25}\text{N}_5\text{O}_2 + \text{H}]^+$.

Step 2: A solution of 4-(6-(2-methoxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (500 mg, 1.1086 mmol) in CH_2Cl_2

(10 mL) was cooled to 0 °C followed by addition of BBr₃ (1.72 g, 6.87 mmol). The reaction mixture was stirred at rt for 2h and heated to 40 °C for 30 min. After completion, the reaction mixture was cooled to 0 °C and quenched by drop wise addition of a saturated aqueous solution of NaHCO₃ (100 mL) followed by extraction with EtOAc (3 x 100 mL). The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 99:1 to 97:3) to afford 4-(6-(2-hydroxy-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 20%, AUC HPLC 96.8) as an off white solid; m.p. 263-272°C. ¹H NMR (400 MHz, CD₃OD) δ (ppm): 10.20 (s, 1 H), 8.81 (s, 1 H), 8.00-7.93 (m, 5 H), 7.76-7.73 (m, 1 H), 7.63-7.60 (m, 1 H), 7.53-7.50 (m, 1 H), 6.96-6.89 (m, 2 H), 3.70-3.35 (m, 4 H), 2.34 (bs, 4 H), 2.21 (s, 3 H); MS (ESI) *m/z* 438 [C₂₆H₂₃N₅O₂+H]⁺.

Example 145: 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-(trifluoromethyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 2-trifluoromethyl-4-bromobenzoic acid (1 g, 3.74 mmol) in DMF (15 mL) was added NMM (755 mg, 7.41 mmol) followed by addition of HATU (2.14 g, 5.61 mmol) at rt and stirred for 30 min. 1-Methylpiperazine (411 mg, 7.41 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 100:0 to 60:40) to afford (4-bromo-2-(trifluoromethyl)phenyl)(4-methylpiperazin-1-yl)methanone (1.2 g, 91%) as a red liquid. ¹H NMR (400 MHz, CDCl₃) δ 7.56 (d, *J* = 8.1 Hz, 2 H), 7.39 (s, 1 H), 3.79 (bs, 2 H), 3.58 (bs, 2 H), 2.45 (bs, 4 H), 2.32 (s, 3 H); MS (ESI) *m/z* 352 [M + H]⁺.

Step 2: To a solution of 4-bromo-2-(trifluoromethyl)phenyl)(4-methylpiperazin-1-yl)methanone (1.2 g, 3.42 mmol), bis(pinacolato)diboron (1.04 g, 4.11 mmol), KOAc (1 g,

10.26 mmol) in 1,4-dioxane (20 mL), was added PdCl₂dppf (75 mg, 0.1 mmol), dppf (56 mg, 0.1 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (4-methylpiperazin-1-yl)(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-(trifluoromethyl)phenyl)methanone (600 mg) as a brown liquid which was used in next step without purification.

Step 3: To a solution of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (390 mg, 1.3 mmol), (4-methylpiperazin-1-yl)(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-(trifluoromethyl)phenyl)methanone (400 mg, 1 mmol), K₃PO₄ (636 mg, 3 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was degassed with argon for 30 min. Pd(PPh₃)₄ (34 mg, 0.03 mmol), was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 1 h. TLC indicated absence of SM. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by preparative HPLC to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-(trifluoromethyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (40 mg, 8%, AUC HPLC >99%) as a pale yellow solid; m.p. 126-131 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.49 (s, 1 H), 7.86-7.83 (m, 5 H), 7.76-7.72 (m, 3 H), 7.51-7.44 (m, 2 H), 3.89 (bs, 1 H), 3.81 (bs, 1 H), 3.26 (bs, 2 H), 2.50 (bs, 2 H), 2.33 (s, 5 H); MS (ESI) *m/z* 490 [C₂₇H₂₂F₃N₅O + H]⁺.

Example 146: 4-(6-(3-amino-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: A solution of (4-bromo-2-nitrophenyl)(4-methylpiperazin-1-yl)methanone (750 mg, 2.29 mmol), bis(pinacolato)diboron (623 mg, 2.45 mmol), KOAc (675 mg, 6.87 mmol) in 1,4-dioxane (15 mL) was degassed with argon for 30 min. PdCl₂dppf (50 mg, 0.068 mmol), dppf (38 mg, 0.0687 mmol) was added and again degassed with argon for 30 min and the

reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (4-methylpiperazin-1-yl)(2-nitro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (400 mg) of as a brown liquid which direct-mass indicated the desired m/z. The crude product was used in the next step without purification.

Step 2: To a solution of (4-methylpiperazin-1-yl)(2-nitro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (400 mg, 1.06 mmol), K₃PO₄ (451 mg, 2.13 mmol) in 1,4-dioxane (5 mL) and water (1 mL) was degassed with argon for 30 min. (A-Phos)₂PdCl₂ (61 mg, 0.053 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 1 h. Water was added to the reaction mixture to induce precipitation which was filtered to give crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 96.5:3.5) to afford 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-nitrophenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (340 mg, 68.5%, LC-MS 89%). ¹H NMR (400 MHz, CDCl₃): δ 8.5 (s, 1 H), 8.3 (s, 1 H), 7.8 (s, 4 H), 7.7 (d, *J* = 8.3 Hz, 2 H), 7.5 (d, *J* = 7.9 Hz, 2 H), 3.9 (bs, 1 H), 3.7 (bs, 1 H), 3.3 (s, 2 H), 2.5 (bs, 2 H), 2.3 (s, 5 H).

Step 3: A solution of 4-(6-(4-(4-methylpiperazine-1-carbonyl)-3-nitrophenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (340 mg, 0.73 mmol) in EtOH (10 mL) was added HCl (1 mL) and SnCl₂ (492 mg, 2.18 mmol). The reaction mixture was heated at 90 °C for 2 h. Work-up and purification by column chromatography gave 4-(6-(3-amino-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (120 mg, 99.2%, AUC HPLC >99%) as an off-white solid; m.p. 218-222 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.67 (s, 1 H), 8.02-7.99 (m, 5 H), 7.80 (d, *J* = 8.0 Hz, 1 H), 7.61 (d, *J* = 8.0 Hz, 1 H), 7.10 (d, *J* = 8.0 Hz, 1 H), 7.02 (s, 1 H), 6.93 (d, *J* = 8.0 Hz, 1 H), 5.32 (s, 2 H), 3.47 (bs, 4 H), 2.31 (bs, 4 H), 2.19 (s, 3 H); MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.


Example 147: 4-(6-(3-(diethylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a solution of 4-(6-(3-amino-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (200 mg) in methanol (20 mL) were added formaldehyde (6 mL), and acetic acid (2 mL) and the reaction mixture was stirred for 2 h at rt. NaCNBH₃ (4 eq) was added at 0 °C and the reaction mixture was stirred at rt for 6 h. The reaction mixture was basified with NaHCO₃ solution and extracted with EtOAc. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography to obtain (150 mg, 75%) which further purified by preparative HPLC to afford 4-(6-(3-(dimethylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (90 mg, 45%, AUC HPLC 97.52%) as an off-white solid; m.p. 161-166 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.47 (s, 1H), 7.84-7.74 (m, 6H), 7.51 (d, *J* = 9.2 Hz, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 7.05 (d, *J* = 7.6 Hz, 1H), 3.85 (s, 2H), 3.40-3.27 (m, 2H), 2.87 (s, 6H), 2.51 (bs, 2H), 2.34 (bs, 6H); MS (ESI) *m/z* 465 [C₂₈H₂₈N₆O+H]⁺.

15

Example 148: 4-(6-(3-(diethylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




To a solution of 4-(6-(3-amino-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (200 mg) in methanol (20 ml) were added acetaldehyde (4 mL) and acetic acid (2 mL) and the reaction mixture was stirred for 2 h at rt. NaCNBH₃ (4 eq) was added at 0 °C and the reaction mixture was stirred at rt for 6 h. The reaction mixture was basified with NaHSO₄ solution and extracted with EtOAc. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford the crude product which was purified by column chromatography to obtain the desired product (90 mg, 90%)

25


which was purified further by preparative HPLC to give 4-(6-(3-(diethylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (70 mg, 31%, AUC HPLC 99.15%) as an off-white solid; m.p. 118-122 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.48 (s, 1H), 7.83-7.73 (m, 6H), 7.50 (d, *J* = 9.2 Hz, 1H), 7.33 (d, *J* = 8.0 Hz, 1H), 7.11 (d, *J* = 7.6 Hz, 1H), 7.05 (s, 1H), 4.00 (bs, 1H), 3.71 (bs, 1H), 3.39 (bs, 1H), 3.26-3.15 (m, 5H), 3.58 (bs, 1H), 2.42 (bs, 2H), 2.32 (s, 2H), 2.22 (bs, 2H), 1.11 (t, *J* = 6.0 Hz, 6H); MS (ESI) *m/z* 493 [C₃₀H₃₂N₆O+H]⁺.

Example 149: N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(4-methylpiperazine-1-carbonyl)phenyl)acetamide



To a solution of 4-(6-(3-amino-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (200 mg, 0.45 mmol) in DCM (20 mL) were added acetic anhydride (70 mg, 0.68 mmol) and pyridine (72 mg, 0.91 mmol) at 0 °C and the reaction mixture was stirred at 0 °C to rt for 16 h. The reaction mixture was acidified with aq NaHSO₄ and extracted with EtOAc. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure and purified by preparative HPLC to afford N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(4-methylpiperazine-1-carbonyl)phenyl)acetamide (50 mg, 22%, AUC HPLC 96.7%) as an off-white solid; m.p. 156-172 °C. ¹H NMR (400 MHz, DMSO *d*₆) δ 9.68 (s, 1H), 8.73 (s, 1H), 7.99 (s, 5H), 7.81-7.80 (m, 2H), 7.65 (d, *J* = 11.2 Hz, 1H), 7.58 (d, *J* = 12.0 Hz, 1H), 7.34 (d, *J* = 10.8 Hz, 1H), 3.6 (bs, 2H), 3.25 (bs, 2H), 2.39 (bs, 2H), 2.25 (bs, 2H), 2.20 (s, 3H), 2.00 (s, 3H); MS (ESI) *m/z* 479 [C₂₈H₂₆N₆O₂+H]⁺.

Example 150: 4-(6-(3-(methylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of methylamine (2 M in THF, 128 mL, 255 mmol) in THF (23 mL) at 0 °C was added n-butyl lithium (2.5 M in hexanes, 80 mL, 201 mmol) slowly. The mixture was stirred for one hour at 0 °C and then it was transferred via cannula to a solution of 4-bromo-2-fluorobenzoic acid (5 g, 22.8 mmol) in THF (5 mL) at -78 °C. The reaction was stirred for 30 minutes before it was quenched at -78 °C with 270 mL of 1N HCl. The aqueous layer was extracted ethyl acetate (4x200 mL) and the combined organic extracts were dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by flash chromatography (silica, eluent 100% CH₂Cl₂ to CH₂Cl₂/MeOH 93:7) to give 4-bromo-2-(methyamino)benzoic acid (1.83 g, 35%, LC-MS 94%) as a light orange solid. ¹H NMR (400 MHz, CDCl₃) δ 7.82 (d, *J* = 8.8 Hz, 1H), 6.86 (d, *J* = 1.2 Hz, 1H), 6.76 (dd, *J* = 1.6, 8.4 Hz, 1H), 2.94 (s, 3H); LC-MS *m/z* 230 [M]⁺.

Step 2: To a solution of 4-bromo-2-(methyamino)benzoic acid (1 g, 4.38 mmol) in DMF (15 mL) was added NMM (664 mg, 6.57 mmol) followed by addition of HATU (2.49 g, 6.57 mmol) at rt and stirred for 30 min. 1-methylpiperazine (486 mg, 4.81 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent petroleum ether to petroleum ether/EtOAc 60/40) to afford (4-bromo-2-(trifluoromethyl)phenyl)(4-methylpiperazin-1-yl)methanone (0.8 g, 60%, LC-MS 68%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.01 (s, 1H), 6.91 (d, *J* = 7.9 Hz, 2H), 6.79 (d, *J* = 4.8 Hz, 1H), 5.3 (bs, 1H), 3.61 (bs, 4H), 2.54 (bs, 4H). LC-MS *m/z* 312 [M]⁺.

Step 3: To a solution of (4-bromo-2-(methyamino)phenyl)(4-methylpiperazin-1-yl)methanone (0.8 g, 2.564 mmol) 1,4-dioxane (20 mL), were sequentially added bis(pinacolato)diboron (716 mg, 2.82 mmol), KOAc (544 mg, 5.56 mmol), PdCl₂dppf (56 mg, 0.076 mmol) and dppf (42 mg, 0.076 mmol). The reaction mixture was heated at 90 °C


for 16 h, then was diluted with EtOAc and washed with water. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford (2-(methylamino)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone as a brown liquid (700 mg) which was used in the next step without purification.

5

Step 4: A solution of (2-(methylamino)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(4-methylpiperazin-1-yl)methanone (300 mg, 1.01 mmol), (2-(methylamino)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone (400 mg, 1.1 mmol), K_3PO_4 (428 mg, 2.02 mmol) in 1,4-dioxane (20 mL) and water (4 mL) was degassed with argon for 30 min prior to the addition of $\text{Pd}(\text{PPh}_3)_4$ (34 mg, 0.03 mmol). The reaction mixture was heated at 90 °C for 1 h under argon atmosphere, then was diluted with water and extracted with EtOAc. The organic phase was washed with water and brine, was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) followed by Preparative HPLC (C18, $\text{ACN}/\text{H}_2\text{O}/10\text{ mM NH}_4\text{HCO}_3$) to afford 4-(6-(3-(methylamino)-4-(4-methylpiperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (145 mg, 26%, AUC HPLC 99.2%) as a light brown solid; m.p. 128-133 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.5 (s, 1H), 7.84-7.54 (m, 6H), 7.54-7.52 (d, $J = 7.5$ Hz, 1H), 7.15-7.13 (d, $J = 7.1$ Hz, 1H), 6.80-6.76 (m, 2H), 5.04 (s, 1H), 3.67 (bs, 4H), 2.88 (d, $J = 0.08$ Hz, 3H), 2.44 (bs, 4H), 2.33 (s, 3H); MS (ESI) m/z 451.1 [$\text{C}_{27}\text{H}_{26}\text{N}_6\text{O} + \text{H}$] $^+$.

20

Example 151: 4-(6-(3-(methylamino)-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of methylamine (2M in THF, 128 mL, 255 mmol) in THF (23 mL) at 0 °C was slowly added n-butyl lithium (2.5M in hexanes, 80.3 mL, 201 mmol). The mixture was stirred for one hour at 0 °C and then it was transferred via cannula to a solution of 4-bromo-2-fluorobenzoic acid (5 g, 22.8 mmol) in THF (5 mL) at -78 °C. The reaction was stirred for 30 minutes before it was quenched at -78 °C with 270 mL of 1N HCl. The aqueous layer was extracted with ethyl acetate (4x200 mL). The combined organic extracts were dried

30

over Na₂SO₄, filtered and concentrated. The crude product was purified by flash column chromatography (silica gel, eluent 100% CH₂Cl₂ to CH₂Cl₂/MeOH 93:7) to give 4-bromo-2-(methylamino)benzoic acid (1.83 g, 35%, LC-MS 94%) as a light orange solid. ¹H NMR (400 MHz, CDCl₃) δ 7.82 (d, *J* = 8.8 Hz, 1H), 6.86 (d, *J* = 1.2 Hz, 1H), 6.76 (dd, *J* = 1.6, 8.4 Hz, 1H), 2.94 (s, 3H); LC-MS *m/z* 230.8 [M]⁺.


Step 2: To a solution of 4-bromo-2-(methylamino)benzoic acid (1 g, 4.38 mmol) in DMF (15 mL) was added NMM (664 mg, 6.57 mmol) followed HATU (2.49 g, 6.57 mmol) and the resulting mixture stirred at rt for 30 min. Morpholine (419 mg, 4.81 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent 100% Petroleum ether to petroleum ether/EtOAc 40:60) to afford (4-bromo-2-(trifluoromethyl)phenyl)(4-methylpiperazin-1-yl)methanone (1.2 g, 92%, AUC LC-MS 88%) as a solid. ¹H NMR (400 MHz, CDCl₃) δ 7.82 (d, *J* = 8.8 Hz, 1H), 6.86 (d, *J* = 1.2 Hz, 1H), 6.76 (dd, *J* = 8.4 Hz, 1H), 5.41 (bs, 1H), 3.84-3.61 (m, 8H), 2.94 (d, *J* = 4.4 Hz, 3H). LC-MS *m/z* 299 [M]⁺.

Step 3: To a solution of (4-bromo-2-(methylamino)phenyl)(morpholino)methanone (1.2 g, 4.02 mmol), Bis(pinacolato)diboron (1.12 g, 4.42 mmol), KOAc (1.18 g, 12.06 mmol) in 1,4-dioxane (20 mL) was added PdCl₂(dppf) (87 mg, 0.12 mmol) and dppf (66 mg, 0.12 mmol) were added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford (2-(methylamino)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone (800 mg) as a brown liquid which was used in the next step without purification.

Step 4: To a solution of 4-(6-bromoimidazo[1,2-*a*]pyridin-3-yl)benzonitrile (300 mg, 1.013 mmol), (2-(methylamino)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone (385 mg, 1.1 mmol), K₃PO₄ (428 mg, 2.02 mmol) in 1,4-dioxane (10 mL) and water (2 mL), was added Pd(PPh₃)₄ (58 mg, 0.05 mmol). The reaction mixture was heated at 90 °C for 1 h then, was diluted with water and extracted with EtOAc. The organic layer was washed with water and brine, dried over anhydrous Na₂SO₄ and

concentrated under reduced pressure to obtain crude product. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) followed by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford 4-(6-(3-(methylamino)-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (4.5 mg, 10.2%, AUC HPLC 94.67%) as an off-White solid; m.p. 302-307 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.5 (s, 1H), 7.84-7.54 (m, 6H), 7.54-7.52 (d, *J* = 7.5 Hz, 1H), 7.15-7.13 (d, *J* = 7.1 Hz, 1H), 6.80-6.76 (m, 2H), 5.04 (s, 1H), 3.70-3.79 (m, 8H), 2.88 (d, *J* = 2.8 Hz, 3H); MS (ESI) *m/z* 436.1 [C₂₆H₂₃N₅O₂ + H]⁺.

10. Example 152: 4-(6-(4-(2-(4-methylpiperazin-1-yl)acetyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile







Step 1: To a solution of 2-bromo-1-(4-bromophenyl)ethanone (2.5 g, 8.99 mmol) in ACN (30 mL) was added K₂CO₃ (2.5 g, 173 mmol) followed by 1-methylpiperazine (1.17 g, 11.69 mmol) at rt and stirred for 16 h. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure to obtain crude product. The crude product was passed through a plug of silica gel using 10% MeOH in CHCl₃ as eluent and concentrated to afford 1-(4-bromophenyl)-2-(4-methylpiperazin-1-yl)ethanone 1 (500 mg, 40%, LC-MS-84%) as a yellow solid.


Step 2: To a mixture of 1-(4-bromophenyl)-2-(4-methylpiperazin-1-yl)ethanone (250 mg, 0.84 mmol), bis(pinacolato)diboron (278 mg, 1.09 mmol), KOAc (323 mg, 3.29 mmol) in 1,4-dioxane (10 mL) were added PdCl₂(dppf) (18.5 mg, 0.02 mmol) and dppf (14 mg, 0.02 mmol) and the reaction mixture was heated at 90 °C for 16 h. The direct mass indicated desired *m/z*. The reaction mixture was used in the next step without isolation.

Step 3: To a mixture of (4-methylpiperazin-1-yl)-1-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)ethanone (1.2 g, 3.48 mmol), 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (1.24 g, 4.18 mmol), K_3PO_4 (2.21 g, 10.44 mmol) in 1,4-dioxane (20 mL) and water (3 mL) was added $Pd(PPh_3)_4$ (120 mg, 0.1 mmol) and the reaction mixture was heated at 90 °C for 4 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude product. The crude product was purified by preparative HPLC to afford 4-(6-(4-(2-(4-methylpiperazin-1-yl)acetyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (150 mg, 15%, AUC HPLC 95.07%) as a yellow solid, m.p: 93-98 °C. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.53 (s, 1H), 8.12 (d, J = 10.8 Hz, 2H), 7.85-7.77 (m, 6H), 7.62 (d, J = 10.8 Hz, 2H), 7.54 (d, J = 12.8 Hz, 1H), 3.82 (s, 2H), 2.67-2.62 (m, 8H), 2.32 (s, 3H); MS (ESI) m/z 436 $[C_{27}H_{25}N_5O+H]^+$.

The following compounds in the table below were prepared following a procedure similar to that used in step 3 of example 152

Ex. #	Boronate ester	Compound	Characterization
153			(200 mg, 22.9%, AUC HPLC 99.03%) as an off-white solid, m.p: 215-218 °C. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ (ppm): 8.67 (s, 1H), 7.96-8.03 (m, 5H), 7.79 (d, <i>J</i> = 9.4 Hz, 1H), 7.78-7.61 (m, 2H), 7.56-7.40 (m, 2H), 3.65 (bs, 2H), 3.31 (bs, 2H), 2.32 (bs, 4H), 2.20 (s, 3H); MS (ESI) <i>m/z</i> 456 [C ₂₆ H ₂₂ ClN ₅ O+H] ⁺ .
154			(150 mg, 16.9%, AUC HPLC 98.91%) as a off-white solid, m.p: 188-191 °C. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ (ppm): 8.51 (s, 1H), 8.01 (s, 1H), 7.95 (s, 4H), 7.76 (d, <i>J</i> = 9.2 Hz, 1H), 7.42 (d, <i>J</i> = 8.2 Hz, 2H), 7.34 (bs, 2H), 3.31 (bs, 4H), 2.31 (s, 7H), 2.20 (s, 3H); MS (ESI) <i>m/z</i> 436 [C ₂₇ H ₂₅ N ₅ O + H] ⁺
155			(100 mg, 23.5%, AUC HPLC 99.85%) as an off-white solid, m.p: 195-197 °C. ¹ H NMR (400 MHz, CDCl ₃) δ (ppm): 8.31 (s, 1H), 7.86 (s, 3H), 7.80-7.75 (m, 2H), 7.69-7.63 (m, 3H), 7.44 (d, <i>J</i> = 7.9 Hz, 1H), 7.26 (s, 1H), 3.86 (bs, 2H), 3.54 (bs, 2H), 2.60-2.34 (m, 4H), 2.39 (s, 3H); MS (ESI) <i>m/z</i> 490 [C ₂₇ H ₂₂ F ₃ N ₅ O+ H] ⁺
156			(120 mg, 37.7%, AUC HPLC 98.77%) as an off-white solid, m.p: 140 °C. ¹ H NMR (400 MHz, CDCl ₃) δ (ppm) 8.46 (s, 1H), 7.84-7.72 (m, 6H), 7.51 (dd, <i>J</i> = 1.6 Hz, 1H), 7.38 (bs, 2H), 7.29 (s, 1H), 3.84 (d, <i>J</i> = 4.81 Hz, 4H), 3.60 (s, 2H), 3.30 (s, 2H), 2.40 (s, 3H); MS (ESI) <i>m/z</i> 423 [C ₂₆ H ₂₂ N ₄ O ₂ + H] ⁺


157			(45 mg, 50%, AUC HPLC 96.9%) as a white solid; m.p. 109-113 °C. ¹ H NMR (400 MHz, CDCl ₃) δ 8.47 (s, 1H), 7.85-7.71 (m, 6H), 7.50-7.43 (m, 4H), 4.92-4.75 (m, 2H), 3.35 (bs, 2H), 2.50 (bs, 2H), 2.39 (bs, 2H), 2.30 (s, 3H); MS (ESI) <i>m/z</i> 506 [C ₂₇ H ₂₂ F ₃ N ₅ O ₂ +H] ⁺
158			(120 mg, 20%, AUC HPLC 95.96%) as an off-white solid; m.p. 169-174 °C. ¹ H NMR (400 MHz, CDCl ₃) δ (ppm): 8.46 (s, 1H), 7.84-7.71 (m, 6H), 7.49 (d, <i>J</i> = 15.0 Hz, 1H), 7.34 (d, <i>J</i> = 10.4 Hz, 1H), 7.13 (d, <i>J</i> = 10.4 Hz, 1H), 7.01 (s, 1H), 3.90 (s, 3H), 3.85 (br. s, 4H), 3.42 (br. s, 2H), 3.35 (m, 2H); MS (ESI) <i>m/z</i> 437 [C ₂₆ H ₂₂ N ₄ O ₃ - H] ⁺



Intermediate 15: 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile

A solution of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (450 g, 1.51 mmol), Bis(pinacolato)diboron (410 mg, 1.61 mmol), KOAc (444 mg, 4.53 mmol) in 1,4-dioxane (15 mL) was degassed with argon for 30 min. PdCl₂dppf (33 mg, 0.045 mmol), dppf (25 mg, 0.045 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water, the organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile as a brown liquid which was used in next step without purification. MS (ESI) *m/z* 346 [M+H].

Example 159: 4-(6-(3-hydroxy-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of 2-methoxy-4-bromobenzoic acid (1 g, 4.32 mmol) in DMF (5 mL) was added NMM (0.97 mL, 8.65 mmol) followed by HATU (2.46 g, 6.49 mmol) at rt and stirred for 30 min. N-Boc-piperazine (886 mg, 4.76 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain tert-butyl 4-(4-bromo-2-methoxybenzoyl)piperazine-1-carboxylate (1.7 g, 95%, LC-MS 92%) as a solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 7.29 (s, 1H), 7.17 (q, *J* = 11.6 Hz, 2H), 3.82 (s, 3H), 3.5 (bs, 2H), 3.31–3.26 (m, 2H), 3.25–3.09 (m, 4H), 1.40 (s, 9H); MS (ESI) *m/z* 398/401 [C₁₇H₂₃BrN₂O₄]⁺.


Step 2: A mixture of tert-butyl 4-(4-bromo-2-methoxybenzoyl)piperazine-1-carboxylate (649 mg, 1.62 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (468 mg, 1.35 mmol), K₃PO₄ (862 mg, 4.06 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added Pd(PPh₃)₄ (78 mg, 0.06 mmol) and the reaction mixture was heated at 90 °C for 1 h. The reaction mixture was diluted with water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography to obtain tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-methoxybenzoyl)piperazine-1-carboxylate (500 mg, 50%, LC-MS 87%) as a solid.

Step 3: To a solution of tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-methoxybenzoyl)piperazine-1-carboxylate (100 mg, 0.18 mmol) in CH₂Cl₂ (5 mL) cooled to 0 °C, was added BBr₃ (2 mL). The reaction mixture was stirred at rt for 2 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated aq. NaHCO₃ (50

mL) and extracted with EtOAc (3 x 50 mL). The organic layer was washed with water, brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent MeOH/CHCl₃ 5:95) followed by preparative HPLC to give 4-(6-(3-hydroxy-4-(piperazine-1-

- 5 carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (50 mg, 63%, AUC HPLC 96.65%) as a white solid, m.p: 170–175 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 10.0 (bs, 1H), 8.69 (s, 1H), 7.99 (s, 5H), 7.80 (d, *J* = 9.3 Hz, 1H), 7.62 (d, *J* = 7.9 Hz, 1H), 7.19 (s, 2H), 7.12 (s, 1H), 3.51 (bs, 2H), 3.21 (bs, 2H), 2.63 (s, 4H); MS (ESI) *m/z* 422.0 [C₂₅H₂₁N₅O₂ - H].

10 Example 160: 4-(6-[3-(morpholin-4-yl)-4-(morpholin-4-ylcarbonyl)phenyl]imidazo[1,2-a]pyridin-3-yl)benzonitrile




- Step 1: To a solution of 2-fluoro-4-bromobenzoic acid (1 g, 4.56 mmol) in DMF (6 mL) was added K₂CO₃ (1.26 g, 9.13 mmol) followed by morpholine (0.47 g, 5.47 mmol) at rt and reaction was heated to reflux and stirred for 20 h at the same temperature. The reaction mixture was cooled to rt, and water (20 mL) was added and acidified with 1N HCl (5 mL). The reaction mixture was extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain 4-bromo-2-(morpholin-4-yl)benzoic acid (700 mg, 90%, LC-MS 95%).
- 20 Step 2: To a solution of 4-bromo-2-morpholinobenzoic acid (700 mg, 2.46 mmol) in DMF (6 mL) was added NMM (0.49 g, 4.91 mmol) followed by HATU (1.39 g, 3.68 mmol) at rt and stirred for 30 min. Morpholine (0.25 g, 2.94 mmol) was added to the reaction mixture and stirring was continued at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain (4-bromo-2-
- 25 morpholinophenyl)(morpholino)methanone (600 mg, LC-MS 86%).

- Step 3: To a mixture of (4-bromo-2-morpholinophenyl)(morpholino)methanone (400 mg, 1.12 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (470 mg, 1.35 mmol), K₃PO₄ (478 mg, 2.25 mmol) in 1,4-dioxane (5 mL),
- 30

and water (1 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (50 mg) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was concentrated to give the crude product. The crude product was purified by preparative HPLC to obtain 4-(6-[3-(morpholin-4-yl)-4-(morpholin-4-ylcarbonyl)phenyl]imidazo[1,2-a]pyridin-3-yl)benzonitrile (50 mg, 20%, AUC HPLC 99.20%) as an off-white solid; m.p. 237-240 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.46 (s, 1H), 7.84–7.79 (m, 4H), 7.73 (d, J = 8.4 Hz, 2H), 7.51 (t, J = 5.6 Hz, 1H), 7.37 (d, J = 8.0 Hz, 1H), 7.24 (s, 1H), 7.11 (s, 1H), 3.99–3.95 (m, 1H), 3.85–3.66 (m, 8H), 3.55–3.40 (m, 1H), 3.42–3.32 (m, 3H), 3.21–3.18 (m, 1H), 2.90–2.89 (m, 2H); MS (ESI) m/z 494 $[\text{C}_{29}\text{H}_{27}\text{N}_5\text{O}_3 + \text{H}]$.

10

Examples 161 a and b: 4-(6-(3-hydroxy-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: To a solution of (4-bromophenyl)(piperidin-4-yl)methanone (600 mg, 2.23 mmol) in MeOH (5 mL) was added $(\text{Boc})_2\text{O}$ (561 mg, 2.57 mmol). The reaction mixture was stirred for 2 h at rt and concentrated. The residues was recrystallized from methanol to give tert-butyl 4-(4-bromobenzoyl)piperidine-1-carboxylate (650 mg, 74.5%, LC-MS 98%) as a solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 7.94 (d, J = 8.4 Hz, 2H), 7.78 (d, J = 8.8 Hz, 2H), 3.82–3.59 (m, 3H), 3.02 (bs, 2H), 1.92 (bs, 2H), 1.75 (bm, 2H), 1.46 (bs, 9H); MS (ESI) m/z 370 $[\text{C}_{17}\text{H}_{22}\text{BrNO}_3 + 2]^+$.

Step 2: A mixture of tert-butyl 4-(4-bromobenzoyl)piperidine-1-carboxylate (599 mg, 1.62 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (468 mg, 1.35 mmol), K_3PO_4 (862 mg, 4.06 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (78 mg, 0.06 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with water and brine solution then dried over

anhydrous Na_2SO_4 and concentrated under reduced pressure to give the crude product, which was purified by column chromatography to afford tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidine-1-carboxylate (250 mg, 28%, LC-MS 39.6 %) as a solid.


5

Step 3: tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidine-1-carboxylate (250 mg) was treated with HCl in 1,4-dioxane (5 mL) at 0 °C for 2 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of a saturated aqueous solution of NaHCO_3 (50 mL) and the mixture was extracted with EtOAc (3 x 50 mL). The organic layer was washed with water and brine, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent MeOH/ CHCl_3 5:95) followed by preparative HPLC to furnish 4-(6-(4-(piperidine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (50 mg, 25%, AUC HPLC 98.84%) as a white solid, m.p: 129–133 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 8.87 (s, 1H), 8.59 (bs, 1H), 8.30 (bs, 1H), 8.11 (d, J = 8.4 Hz, 1H), 8.07 (s, 1H), 8.04–7.97 (m, 4H), 7.95–7.87 (m, 4H), 3.80 (bs, 1H), 3.34 (d, J = 12.7 Hz, 2H), 3.10 (t, J = 11.2 Hz, 2H), 1.97 (d, J = 13.2 Hz, 2H), 1.76 (q, J = 10.8 Hz, 2H); MS (ESI) m/z 407 [$\text{C}_{26}\text{H}_{22}\text{N}_4\text{O} + \text{H}$].

20 Step 4: To a solution of 4-(6-(4-(piperidine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl) (20 mg, 0.04 mmol) in MeOH (5 mL) cooled to 0 °C was added NaBH_4 (37 mg, 0.09 mmol) and the reaction mixture was stirred at rt for 2 h. The reaction mixture was extracted with EtOAc and the organic layer was washed with water and brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent MeOH/ CHCl_3 5:95) followed by preparative HPLC to give 4-(6-(4-(hydroxy(piperidin-4-yl)methyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (50 mg, 25%, AUC HPLC 97%) as a white solid. The racemic compound was resolved by chiral HPLC to afford both enantiomers as white solids. First enantiomer (example 161a) (10 mg, 30%, AUC HPLC 98%, chiral HPLC 99.6%) and second enantiomer (example 161b) (15 mg, 30%, AUC HPLC 98%, chiral HPLC 95.3%). ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.48 (s, 1H), 7.83–7.71 (m, 6H), 7.53–7.50 (m, 3H), 7.41 (d, J = 7.6 Hz, 2H), 4.48 (d, J = 6.8 Hz, 1H), 3.20 (dd, J = 12.48 Hz, 2H), 2.60 (q, J = 9.6 Hz, 3H), 2.10 (d, J = 13.2 Hz, 2H), 1.80 (bs, 2H), 1.40 (d, J = 12.8 Hz, 2H); MS (ESI) m/z 409 [$\text{C}_{26}\text{H}_{24}\text{N}_4\text{O} + \text{H}$] $^+$.

30

Example 162: 4-(6-(3-fluoro-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a mixture of 4-(ethoxycarbonyl)-3-fluorophenylboronic acid (400 mg, 1.34 mmol),
 5 4-6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (313 mg, 1.47 mmol), K_3PO_4 (853 mg, 4.02 mmol) in 1,4-dioxane (5 mL) and water was added (A-Phos) $_2PdCl_2$ (77 mg, 0.06 mmol) and the reaction mixture was heated at 100 °C for overnight under argon. The reaction mixture was washed with water and extracted with ethylacetate and concentrated. The crude product was purified by column chromatography (silica gel, eluent MeOH/ $CHCl_3$ 10:90) to afford
 10 ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoate (350 mg, 67.8%, LC-MS 98.5%). 1H NMR; (400 MHz, $DMSO-d_6$) δ (ppm): 8.91 (s, 1H), 7.99–8.02 (m, 6H), 7.83–7.79 (m, 4H), 4.3 (q, $J = 7$ Hz, 2H), 1.3 (t, $J = 7.4$ Hz, 3H); MS (ESI) m/z 386 $[C_{23}H_{16}FN_3O_2 + H]^+$.

Step 2: To a solution of ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-
 15 fluorobenzoate (350 mg, 0.90 mmol) in 15 mL of THF/ H_2O /EtOH 1:1:1 was added LiOH (76 mg, 1.81 mmol) and the reaction mixture was stirred at rt for 3 h. The reaction mixture was quenched with citric acid, washed with water and extracted with EtOAc (3 x 25 mL). The organic layer was washed with water and brine, dried over anhydrous Na_2SO_4 , filtered and
 20 concentrated under reduced pressure. The crude product was purified by recrystallization to afford 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoic acid (200 mg, 61.7%, LC-MS 95.7%).

Step 3: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-fluorobenzoic
 25 acid (200 mg, 0.56 mmol) in DMF (5 mL) was added NMM (0.12 mL, 1.12 mmol) followed by HATU (319 g, 0.84 mmol) and stirred for 20 min. To the reaction mixture, morpholine (0.05 g, 0.61 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain 4-(6-(3-fluoro-4-
 30 (morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 42%, AUC HPLC 99.33%) as a white solid, m.p: 115-135 °C. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.48


(s, 1H), 7.85-7.80 (m, 4H), 7.72 (d, $J = 8.3$ Hz, 2H), 7.54-7.45 (m, 2H), 7.47 (dd, $J = 3.0$ Hz, 1H), 7.29 (d, $J = 1.2$ Hz, 1H), 3.83 (d, $J = 4.3$ Hz, 4H), 3.79 (s, 2H), 3.39 (s, 2H); MS (ESI) m/z 427.35 [$C_{25}H_{19}FN_4O_2 + H$] $^+$.

5 Example 163: 4-(6-(4-(4-methyl-4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



A mixture of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (250 mg, 0.73 mmol), 4-(4-methylpiperidin-4-yl)morpholine (149 mg, 0.81 mmol) in DMF (10 mL) was added HATU (41 mg, 1.10 mmol) and NMM (0.16 mL, 1.47 mmol) and the reaction mixture was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford of 4-(6-(4-(4-methyl-4-morpholinopiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (60 mg, 16%, AUC HPLC 97.67%) as an off-white solid, m.p: 238-242 °C. 1H NMR (400 MHz, $DMSO-d_6$) δ 8.79 (s, 1H), 8.00 (s, 5H), 7.82-7.72 (m, 4H), 7.49 (d, $J = 7.4$ Hz, 2H), 3.92 (bs, 1H), 3.58 (s, 4H), 3.41 (s, 2H), 3.31 (s, 1H), 2.44 (s, 4H), 1.82 (bs, 1H), 1.75 (bs, 1H), 1.39 (bs, 2H), 0.90 (s, 3H); MS (ESI) m/z 506.46 [$C_{31}H_{31}N_5O_2 + H$].


20 Example 164: 4-(6-(4-(4-methyl-4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



To a mixture of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (250 mg, 0.73 mmol), 1-methyl-4-(4-methylpiperidin-4-yl)piperazine (159 mg, 0.81 mmol) in DMF (5 mL)

was added HATU (419 mg, 1.10 mmol), followed by NMM (0.16 mL, 1.47 mmol) and the reaction mixture was stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford 4-(6-(4-(4-methyl-4-(5-methyl-1,3,4-oxadiazol-2-yl)piperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 26%, AUC HPLC 98.77%) as a white solid, m.p: 262-267 °C. ¹H NMR (400 MHz, CDCl₃): δ (ppm): 8.49 (s, 1H), 7.84-7.72 (m, 6H), 7.57-7.50 (m, 5H), 4.01 (bs, 1H), 3.55 (bs, 2H), 3.37 (bs, 1H), 2.55-2.45 (m, 8H), 2.28 (s, 3H), 1.93 (bs, 1H), 1.77 (bs, 1H), 1.51 (bs, 1H), 1.42 (bs, 1H), 0.97 (s, 3H); MS (ESI) *m/z* 519 [C₃₂H₃₄N₆O + H].

Example 165: 4-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (0.25 g, 0.74 mmol) in DMF (5 mL) were added HATU (0.419 g, 1.1 mmol), N-methyl morpholine (0.15 g, 1.47 mmol) and tert-butyl piperazine-4-carboxylate A (0.15 g, 0.81 mmol). The reaction mixture was stirred at 0 °C to rt under inert atmosphere for 16 h, then was diluted with H₂O (15 mL) and extracted with EtOAc (3 × 25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazine-1-carboxylate (0.23 g, LC-MS 92%). ¹H NMR (400 MHz, CDCl₃): δ 8.48 (s, 1H), 7.87-7.83 (m, 4H), 7.72 (d, *J* = 8 Hz, 2H), 7.59-7.51 (m, 5H), 3.7-3.46 (m, 8H), 1.43 (s, 9H); MS (ESI) *m/z* 508 [C₃₀H₂₉N₅O₃ + H]⁺.

Step 2: A solution of hydrochloric acid in Ether was added drop wise to tert-butyl 4-(4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl) piperazine-1-carboxylate (0.23 g, 0.45 mmol) at 0 °C. Reaction mixture was stirred at rt for 2 h. Ether was evaporated under reduced pressure and basified with saturated NaHCO₃ solution. The solid thus formed was filtered to afford 4-(6-(4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (80 mg, 43.3%, AUC HPLC 95.22%) as an off-white solid, m.p: 88-98 °C. ¹H NMR: (400 MHz,

CDCl_3 δ (ppm) 8.50 (s, 1H), 7.84-7.73 (m, 6H), 7.59-7.51 (m, 5H), 3.84 (bs, 2H), 3.25 (bs, 2H), 3.05-2.8 (m, 4H), 1.62 (s, 1H); MS (ESI) m/z 408.2 $[\text{C}_{25}\text{H}_{21}\text{N}_5\text{O} + \text{H}]^+$

Example 166: N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide



Step 1: To a solution of methyl 2-acetamido-4-bromobenzoate (1 g, 4.37 mmol) in DCM (15 mL) was added TEA (0.873 g, 8.6 mmol) followed by acetyl chloride (0.406 g, 5.21 mmol) at 0 °C and stirred for 4 h. The reaction mixture was diluted with DCM and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain methyl 2-acetamido-4-bromobenzoate (600 mg, 51%) as a white solid.

Step 2: To a solution of methyl 2-acetamido-4-bromobenzoate (0.6 g, 2.21 mmol), bis(pinacolato)diboron (0.61 g, 2.435 mmol), KOAc (0.65g, 6.64 mmol) in 1,4-dioxane (15 mL) were added $\text{Pd}_2(\text{dba})_3$ (60 mg, 0.066 mmol), $\text{P}(\text{Cy})_3$ (60 mg, 0.066 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with ethyl acetate and washed with water. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford methyl 2-acetamido-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (600 mg, LC-MS 70%) as a green solid which was used in the next step without purification.


Step 3: To a mixture of methyl 2-acetamido-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.6 g, 1.87 mmol), 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (0.4 g, 1.34 mmol), K_3PO_4 (0.57 g, mmol) in 1,4-dioxane (5 mL), and water (1 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (40 mg) and the reaction mixture was heated at 90 °C for overnight and concentrated to give crude product. The crude product was purified by column

chromatography (silica gel, eluent MeOH/CHCl₃ 3.5: 96.5) to afford methyl 2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (350 mg, 63%, LC-MS 85%)

Step 4: To a solution of methyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-formamidobenzoate (350 mg, 0.85 mmol) in 10 mL of THF: H₂O: EtOH (1:1:1) was added LiOH (72 mg, 1.723 mmol) at 0 °C and stirred at rt for 2 h. The reaction mixture was concentrated under reduced pressure then water (10 mL) was added to the reaction mixture, the resulting solid was filtered to give 2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (200 mg, LC-MS 92%) as a white solid.

Step 5: To a solution of 2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (200 mg) in DMF (6 mL) was added NMM (0.178 g, 1.76 mmol) followed by HATU (0.59 g, 1.56 mmol) at rt and stirred for 30 min. Morpholine (92 mg, 1.06 mmol) was added and the stirring was continued at rt for overnight. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude compound which was purified by preparative HPLC to give N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide (151 mg, AUC HPLC 96.2%) of as a pale yellow solid: mp 150-153 °C. ¹H NMR (400 MHz, CDCl₃) δ 9.10 (s, 1H), 8.57 (s, 1H), 8.53 (s, 1H), 7.84-7.73 (m, 6H), 7.53 (d, *J* = 9.6 Hz, 1H), 7.31 (d, *J* = 8.0 Hz, 2H), 3.73 (bs, 8H), 2.21 (s, 3H); MS (ESI) *m/z* 466 [M+1]

Example 167: 4-(6-(3-hydroxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a mixture of 4-(6-bromoimidazo[1,2-a]pyridin-3-yl)benzonitrile (400 mg, 1.35 mmol), (2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)(morpholino)methanone (562 mg, 1.62 mmol), K₃PO₄ (572 mg, 2.7 mmol) in 1,4-dioxane (80 mL) and water (20 mL) was added Pd(PPh₃)₄ (77 mg, 0.068 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with water

and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by preparative HPLC to give 4-(6-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (120 mg, 20%, AUC HPLC 95.96%) as an off-white solid; m.p. 169-174 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.46 (s, 1H), 7.84-7.71 (m, 6H), 7.49 (d, $J = 15.0$ Hz, 1H), 7.34 (d, $J = 10.4$ Hz, 1H), 7.13 (d, $J = 10.4$ Hz, 1H), 7.01 (s, 1H), 3.90 (s, 3H), 3.85 (bs, 4H), 3.42 (b s, 2H), 3.35 (m, 2H); MS (ESI) m/z 437 [$\text{C}_{26}\text{H}_{22}\text{N}_4\text{O}_3 - \text{H}^+$] $^+$.

- 10 Step 2: To a solution of 4-(6-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (120 mg, 0.273 mmol) in CH_2Cl_2 (10 mL) cooled to 0 °C was added BBR_3 (136 mg, 0.547 mmol). The reaction mixture was stirred at rt for 2 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated aq. NaHCO_3 (100 mL) and extracted with EtOAc (3x100 mL). The organic layer was washed
- 15 with water, brine, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The crude compound was purified by preparative HPLC to give 4-(6-(3-hydroxy-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (80 mg, 72%, AUC HPLC 98.42%) as an off-white solid. mp 104-106 °C. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm): 10.06 (s, 1H), 8.69 (s, 1H), 7.99 (m, 5H), 7.80 (d, $J = 9.2$ Hz, 1H), 7.62 (d, $J = 9.2$ Hz,
- 20 1H), 7.27-7.54 (m, 2H), 7.13 (s, 1H), 3.59-3.26 (m, 8H); MS (ESI) m/z 425 [$\text{C}_{25}\text{H}_{20}\text{N}_4\text{O}_3 + \text{H}^+$] $^+$.

Example 168: 4-(6-(3-((1,3-dioxoisindolin-2-yl)methyl)-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



Step 1: A solution of 5-bromoisobenzofuran-1(3H)-one (5 g, 23.47 mmol) and KOH (1.58 g, 28.17 mmol) in MeOH (100 mL) was refluxed for 4 h. MeOH was removed and the residue was dissolved in DMF (30 mL) and treated with imidazole (4.79 g, 70.41 mmol) and tert-butyl dimethyl silyl chloride (7.07 g, 46.94 mmol) at 0 °C and reaction was stirred for 16 h at rt. The reaction mixture was poured into water and extracted with Et₂O (3 x 100 mL). The combined organic layer was washed with brine solution, dried over anhydrous Na₂SO₄ and concentrated in vacuo. The crude residue was dissolved in MeOH (15 mL) and THF (15 mL) and then treated with aqueous solution of K₂CO₃ (6.47 g, 46.94 mmol). The mixture was stirred for 16 h at rt and diluted with water (60 mL). The mixture was cooled to 0 °C and pH was adjusted to 5-6 with saturated solution of citric acid. The two phases were separated and the aqueous layer was extracted with Et₂O (3 x 50 mL). The combined organic layer was washed with brine solution, dried over anhydrous Na₂SO₄ and concentrated in vacuum to afford 4-bromo-2-((tert-butyldimethylsilyloxy)methyl)benzoic acid (4 g, 49%, LC-MS 90.5%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.21 (s, 1H), 7.88-7.81 (m, 2H), 7.62 (d, *J* = 8 Hz, 1H), 5.08 (s, 2H), 0.97 (s, 9H), 0.48 (s, 6H); MS (ESI) *m/z* 346 [C₁₄H₂₁BrSiO₃ + H]⁺.

Step 2: To a solution of 4-bromo-2-((tert-butyldimethylsilyloxy)methyl)benzoic acid (3 g, 8.69 mmol) in DMF (20 mL) were added HATU (4.95 g, 13.03 mmol), N-methylmorpholine (1.75 g, 17.38 mmol) and morpholine (0.9 g, 10.43 mmol). The reaction mixture was stirred at 0 °C to rt for 16 h, then was diluted with H₂O (50 mL) and extracted with

EtOAc (3x150 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent hexane/EtOAc 70:30) to afford (4-bromo-2-((tert-butyl)dimethylsilyloxy)methyl)phenyl(morpholino) methanone (1 g, 28%, LC-MS 98.56%).

- 5 ¹H NMR (400 MHz, CDCl₃) δ 7.70 (s, 1H), 7.43 (d, *J* = 1.6 Hz, 1H), 7.02 (d, *J* = 8 Hz, 1H), 4.69 (bs, 2H), 3.77 (bs, 4H), 3.58 (bs, 2H), 3.26 (bs, 2H), 0.94 (s, 9H), 0.11 (s, 6H); MS (ESI) *m/z* 415 [C₁₈H₂₂BrNSiO₃+H]⁺.


- Step 3: To a solution of (4-bromo-2-((tert-butyl)dimethylsilyloxy)methyl)phenyl(morpholino) methanone (1 g, 2.41 mmol) in THF (10 mL) was added tetrabutylammonium fluoride in THF (1M solution in THF, 4.83 mL, 4.83 mmol) drop wise at 0 °C. The reaction mixture was stirred at rt for 1 h and was diluted with H₂O (20 mL) and extracted with EtOAc (3 × 50 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent hexane/EtOAc 30:70) to afford (4-bromo-2-(hydroxymethyl)phenyl(morpholino)methanone (570 mg, 79%, LC-MS 90%). ¹H NMR (400 MHz, CDCl₃) δ 7.62 (s, 1H), 7.47 (d, *J* = 8.0 Hz, 1H), 7.09 (d, *J* = 8 Hz, 1H), 4.55 (bs, 2H), 3.78 (bs, 4H), 3.60 (bs, 2H), 3.35 (bs, 2H); MS (ESI) *m/z* 301.15 [C₁₂H₁₄BrNO₃+H]⁺.

- 20 Step 4: To a solution of (4-bromo-2-(hydroxymethyl)phenyl(morpholino)methanone (2.8 g, 9.33 mmol), phthalimide (1.5 g, 10.26 mmol) and triphenyl phosphine (3.66 g, 13.99 mmol) in THF (30 mL) was added DEAD (2.43 g, 13.99 mmol) drop wise at 0 °C. The reaction mixture was stirred at rt for 2 h then was diluted with H₂O (50 mL) and extracted with EtOAc (3x150 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent hexane/EtOAc 70:30) to afford 2-(5-bromo-2-(morpholine-4-carbonyl)benzyl)isoindoline-1,3-dione (2 g, LC-MS 39%) which was used in the next step without purification.

- 30 Step 5: To a mixture of 2-(5-bromo-2-(morpholine-4-carbonyl)benzyl)isoindoline-1,3-dione (540 mg, 1.26 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (521 mg, 1.51 mmol), K₃PO₄ (534 mg, 2.52 mmol) in 1,4-dioxane (15 mL) and water (3 mL) was added Pd(PPh₃)₄ (72.7 mg, 0.06 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with

water and extracted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 97:3) to afford 4-(6-(3-((1,3-dioxoisindolin-2-yl)methyl)-4-(morpholine-4-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (250 mg, 35%, AUC HPLC 98%) as an light yellow solid; ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.44 (s, 1H), 7.86-7.64 (m, 10H), 7.46 (d, $J = 8.4$ Hz, 2H), 7.31 (d, $J = 8.2$ Hz, 2H), 5.03-4.97 (m, 2H), 3.85-3.66 (m, 6H), 3.40 (s, 2H); MS (ESI) m/z 568 [$\text{C}_{34}\text{H}_{25}\text{N}_5\text{O}_4 + \text{H}$] $^+$;

10 Example 169: N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(4-(methylamino)piperidine-1-carbonyl)phenyl)acetamide



Step: 1 To a solution of 2-amino-4-bromobenzoic acid (2.5 g, 11.57 mmol) in 1,4-dioxane (10 mL) was added bis(trichloromethyl) carbonate at 0 °C and the reaction mixture was heated at 110 °C for 16 h. The reaction mixture was concentrated, filtered and washed with hexane to afford 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (2.3 g, 82%, LC-MS 99%) as an off white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.8 (s, 1H), 7.83 (d, $J = 8.4$ Hz, 1H), 7.43 (d, $J = 8.4$ Hz, 2H); MS (ESI) m/z 242 [$\text{C}_8\text{H}_4\text{BrNO}_3 + \text{H}$] $^+$.

20 Step 2: To a solution of 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (400 mg, 1.65 mmol), tert-butyl methyl(piperidin-4-yl)carbamate (389 mg, 1.81 mmol) in DMF (5 mL) was added DMAP (20 mg, 0.16 mmol) at rt and stirred at same temperature for 16 h. The reaction

mixture was added ice cooled water and extracted with EtOAc (3×30 mL). The combined organic layer was washed with water, brine solution and dried over Na₂SO₄ and was concentrated under reduced pressure to obtain the crude product. The crude product was washed with n-pentane to afford tert-butyl 1-(2-amino-4-bromobenzoyl)piperidin-4-yl(methyl)carbamate (670 mg, 98%, LC-MS 94%). ¹H NMR (400 MHz, CDCl₃) δ 6.93 (d, *J* = 8.4 Hz, 1H), 6.88 (s, 1H), 6.83 (d, *J* = 6.8 Hz, 1H), 4.42 (s, 3H), 4.21 (br s, 2H), 2.95 (s, 2H), 2.88 (s, 3H), 1.72-1.62 (m, 4H), 1.46 (s, 9H); MS (ESI) *m/z* 412/414 [C₁₈H₂₆BrN₃O₃]⁺.


Step 3: To a solution of tert-butyl 1-(2-amino-4-bromobenzoyl)piperidin-4-yl(methyl)carbamate (0.73 g, 1.78 mmol) in DCM (10 mL) was added triethylamine (0.74 mL, 5.34 mmol). Acetyl chloride (0.19 mL, 2.67 mmol) was added to the reaction mixture at 0 °C and stirred at rt for 2 h. The reaction mixture was added water and extracted with EtOAc (3 × 30 mL). The combined organic layer was washed with water, brine solution and dried over Na₂SO₄ and was concentrated under reduced pressure to obtain the crude product. The crude product was washed with n-pentane to afford tert-butyl 1-(2-acetamido-4-bromobenzoyl)piperidin-4-yl(methyl)carbamate (650 mg, 80.5%, LC-MS 92.5%). ¹H NMR (400 MHz, CDCl₃) δ 9.01 (s, 1H), 8.55 (s, 1H), 7.25 (d, *J* = 8.4 Hz, 1H), 7.01 (d, *J* = 6.8 Hz, 1H), 4.18 (s, 2H), 2.95 (s, 3H), 2.74 (s, 4H), 2.18 (s, 3H), 1.82-1.65 (m, 3H), 1.46 (s, 9H); MS (ESI) *m/z* 554/556 [C₂₀H₂₈BrN₃O₄]⁺.

Step 4: To a mixture of tert-butyl 1-(2-acetamido-4-bromobenzoyl)piperidin-4-yl(methyl)carbamate (215 mg, 0.47 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (196.8 mg, 0.57 mmol), K₃PO₄ (201 mg, 0.95 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added Pd(PPh₃)₄ (27.4 mg, 0.02 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was filtered through celite bed and washed with 10% MeOH in CHCl₃ and the filtrate was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 97:3) to afford tert-butyl 1-(2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl(methyl)carbamate (210 mg, 75%, LC-MS 61%) as a light yellow solid.

Step 5: To a solution of tert-butyl 1-(2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperidin-4-yl(methyl)carbamate (200 mg, 0.33 mmol) in DCM (10

mL) was added TFA (1.2 mL) at 0 °C. The reaction mixture was stirred at rt for 3 h, concentrated under reduced pressure and basified with saturated NaHCO₃ solution. The precipitated was isolated by filtration and was dried. The crude product was purified by preparative HPLC to afford N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(4-(methylamino)piperidine-1-carbonyl)phenyl)acetamide (30 mg, 18%, AUC HPLC 95.9%) as a pale yellow solid. m.p. 147-151 °C. ¹H NMR (400 MHz, CDCl₃) δ 9.07 (s, 1H), 8.53 (s, 2H), 7.84-7.73 (m, 6H), 7.76 (d, *J* = 7.6 Hz, 1H), 7.73 (s, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 4.60 (s, 1H), 3.95 (s, 1H), 3.15 (s, 2H), 2.77 (s, 1H), 2.49 (s, 3H), 2.19 (s, 3H), 2.05 (s, 3H), 1.44 (s, 2H); MS (ESI) *m/z* 493.05 [C₂₉H₂₈N₆O₂+H]⁺.

Example 170: N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(piperazine-1-carbonyl)phenyl)acetamide



Step: 1 To a solution of 2-amino-4-bromobenzoic acid (2.5 g, 11.57 mmol) in 1,4-dioxane (10 mL) was added bis(trichloromethyl) carbonate at 0 °C and the reaction mixture was heated at 110 °C for 16 h. The reaction mixture was concentrated, filtered and washed with hexane to afford 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (2.3 g, 82%, LC-MS 99%) as an off white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.8 (s, 1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.43 (d, *J* = 8.4 Hz, 2H); MS (ESI) *m/z* 242 [C₈H₄BrNO₃+H]⁺.

Step 2: To a solution of 7-bromo-1H-benzo[d][1,3]oxazine-2,4-dione (400 mg, 1.65 mmol), tert-butyl methyl(piperidin-4-yl)carbamate (389 mg, 1.81 mmol) in DMF (5 mL) was added DMAP (20 mg, 0.16 mmol) at rt and the mixture was stirred for 16 h. To the reaction mixture

was added ice cooled water and the solid thus precipitated was filtered and washed in turn with water, n-pentane and dried to afford tert-butyl 4-(2-amino-4-bromobenzoyl)piperazine-1-carboxylate (910 mg, 57%, LC-MS 64.6%). MS (ESI) m/z 384/386 [$C_{16}H_{22}BrN_3O_3+H$]⁺.


- 5 Step 3: To a solution of tert-butyl 4-(2-amino-4-bromobenzoyl)piperazine-1-carboxylate (0.9 g, 2.35 mmol) in dichloromethane (10 mL) was added TEA (0.98 mL, 7.05 mmol). Acetyl chloride (0.25 mL, 3.53 mmol) was added to the reaction mixture at 0 °C and stirred at rt for 2 h. To the reaction mixture was added water and the mixture was extracted with EtOAc (3 × 50 mL). The combined organic layer was washed in turn with water and brine. The organic
10 solution and dried over Na₂SO₄, filtered and was concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 97:3) to afford tert-butyl 4-(2-acetamido-4-bromobenzoyl)piperazine-1-carboxylate (600 mg, 60%, LC-MS 96.3%) as a light yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.96 (s, 1H), 8.54 (s, 1H), 7.24 (d, J = 10 Hz, 1H), 7.04
15 (d, J = 8.0 Hz, 1H), 3.58-3.41 (m, 8H), 2.15 (s, 3H), 1.47 (s, 9H); MS (ESI) m/z 426/428 [$C_{18}H_{24}BrN_3O_4$]⁺.

- Step 4: To a mixture of tert-butyl 4-(2-acetamido-4-bromobenzoyl)piperazine-1-carboxylate (373 mg, 0.87 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-
20 a]pyridin-3-yl)benzonitrile C (364 mg, 1.05 mmol), K₃PO₄ (372.7 mg, 1.76 mmol) in 1,4-dioxane (10 mL) and water (2 mL) was added Pd(PPh₃)₄ (50.7 mg, 0.04 mmol) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was filtered through celite bed and washed with 10% MeOH in CHCl₃ and the filtrate was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The
25 crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 97:3) to afford tert-butyl 4-(2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazine-1-carboxylate (310 mg, 62%, LC-MS 76%) as a light yellow solid. MS (ESI) m/z 565 [$C_{32}H_{32}N_6O_4+H$]⁺.

- 30 Step 5: To a solution of tert-butyl 4-(2-acetamido-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazine-1-carboxylate (250 mg, 0.44 mmol) in dichloromethane (10 mL) was added TFA (1.2 mL) at 0 °C. The reaction mixture was stirred at rt for 3 h. Reaction mixture was diluted with water and basified with saturated NaHCO₃ solution and extracted with 10% MeOH in CHCl₃. The combined organic layer was washed with water,

brine solution and dried over Na_2SO_4 and was concentrated under reduced pressure to obtain the crude product. The compound was further purified by preparative HPLC to afford N-(5-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)-2-(piperazine-1-carbonyl)phenyl)acetamide (30 mg, 14.6%, AUC HPLC 98.3%) as an off-white solid. mp 160-165 °C. ^1H NMR (400 MHz, CDCl_3) δ 9.08 (s, 1H), 8.56-8.53 (m, 2H), 7.84-7.77 (m, 6H), 7.75-7.73 (m, 1H), 7.55-7.30 (m, 3H), 3.62 (s, 4H), 2.93 (s, 4H), 2.20 (s, 3H); MS (ESI) m/z 465 [$\text{C}_{27}\text{H}_{24}\text{N}_6\text{O}_2 + \text{H}$] $^+$.

Example 171: 4-(6-(3-(piperazin-1-yl)-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile



10

Step 1: To a solution of 4-bromo-2-(4-(tert-butoxycarbonyl)piperazin-1-yl)benzoic acid (0.8 g, 2.08 mmol) in DMF (6 mL) was added NMM (0.41 g, 4.16 mmol) followed by HATU (1.18 g, 3.12 mmol) at rt and the reaction mixture was stirred for 30 min. tert-butyl piperazine-1-carboxylate (0.46 g, 2.49 mmol) was added and stirred was continued at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water, brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain tert-butyl 4-(4-bromo-2-(4-(tert-butoxycarbonyl)piperazin-1-yl)benzoyl)piperazine-1-carboxylate (600 mg, 52%, LC-MS-80%) as an off white solid.

15


Step 2: To a mixture of tert-butyl 4-(4-bromo-2-(4-(tert-butoxycarbonyl)piperazin-1-yl)benzoyl)piperazine-1-carboxylate (612 mg, 1.10 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (470 mg, 1.00 mmol), K_3PO_4 (427 mg, 2.01 mmol) in 1,4-dioxane (15 mL) and water (1 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (40 mg, 0.01 mmol) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent methanol/dichloromethane 10:90) to afford tert-butyl 4-(2-(4-(tert-butoxycarbonyl)piperazin-

25

1-yl)-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazine-1-carboxylate (200 mg, 29%, LC-MS 52%) as a light brown solid.

Step 3: To a solution of tert-butyl-4-(2-(4-(tert-butoxycarbonyl)piperazin-1-yl)-4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)piperazine-1-carboxylate (200 mg) in DCM (5 mL) was added a HCl solution in ether (5 mL) at 0 °C and stirred at rt for 2 h. The reaction mixture was concentrated basified with NaHCO₃, diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent methanol/dichloromethane 10:90) to afford 4-(6-(3-(piperazin-1-yl)-4-(piperazine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (100 mg, 70%, AUC HPLC 94.35%) as an off-white solid; mp 170-174 °C. ¹H NMR (400 MHz, DMSO *d*₆) δ 8.76 (s, 1H), 8.02 (s, 5H), 7.80 (d, *J* = 9.2 Hz, 1H), 7.71 (d, *J* = 8.0 Hz, 1H), 7.39 (d, *J* = 8.0 Hz, 1H), 7.28 (s, 1H), 7.25 (d, *J* = 8.0 Hz, 2H), 3.67-3.64 (m, 1H), 3.51-3.47 (m, 1H), 3.16-3.13 (m, 3H), 3.01 (s, 1H), 2.97-2.67 (m, 9H), 2.5 (s, 1H); MS (ESI) *m/z* 492.31 [C₂₉H₂₉N₇O + H]⁺.

Example 172: Synthesis of 4-(6-(4-(4-methoxy-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile




Step 1: To a solution of 4-bromobenzoic acid (500 mg, 4.97 mmol) in DMF (10 mL) were added HATU (1.41 g, 7.46 mmol), N-methyl morpholine (0.69 mL, 12.44 mmol) and 4-methylpiperidin-4-ol hydrochloride (452 mg, 5.97 mmol). The reaction mixture was stirred at 0 °C to room temperature under inert atmosphere for 16 h, then was diluted with H₂O (30 mL) and extracted with EtOAc (3×30 mL). The combined organic layer was washed with water, brine and dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 97:3) to afford (4-bromophenyl)(4-hydroxy-4-methylpiperidin-1-yl)methanone (435 mg, 58%, LC-MS 92.2%) as a pale yellow solid. MS (ESI) *m/z* 298/300 [C₁₃H₁₆BrNO₂ + H]⁺.

25

Step 2: A solution of (4-bromophenyl)(4-hydroxy-4-methylpiperidin-1-yl)methanone (375 mg, 1.26 mmol) in anhydrous DMF (8 mL) was added to sodium hydride (120.8 mg, 50%, 5.03 mmol) that was prewashed with n-hexane, cooled to 0 °C. Methyl iodide (2.5 mL, 40.14 mmol) was added to the reaction mixture and stirred at rt for 16 h. The reaction mixture was quenched with ice-cold water and extracted with EtOAc (3×30 mL). The combined organic layer was washed with water, brine solution and dried over Na₂SO₄, filtered and was concentrated under reduced pressure to afford (4-bromophenyl)(4-methoxy-4-methylpiperidin-1-yl)methanone (250 mg) as a green solid. MS (ESI) *m/z* 312/314 [C₁₄H₁₈BrNO₂+H]⁺.

Step 3: To a mixture of (4-bromophenyl)(4-methoxy-4-methylpiperidin-1-yl)methanone (240 mg, 0.77 mmol), 4-(6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (318.4 mg, 0.92 mmol), K₃PO₄ (326 mg, 1.54 mmol) in 1,4-dioxane (15 mL) and water (2 mL) was added Pd(PPh₃)₄ (44 mg, 0.04 mmol) and the reaction mixture was heated at 100 °C for 16 h. The reaction mixture was diluted with EtOAc and washed in turn with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced. The residue was purified by column chromatography (silica gel, eluent MeOH/DCM/MeOH 97:3) to afford 4-(6-(4-(4-methoxy-4-methylpiperidine-1-carbonyl)phenyl)imidazo[1,2-a]pyridin-3-yl)benzonitrile (70 mg, 29%, AUC HPLC 96.73%) as an off white solid. mp 159-163 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.50 (s, 1H), 7.84-7.72 (m, 6H), 7.58-7.50 (m, 5H), 4.40 (s, 1H), 3.55-3.40 (m, 2H), 3.33-3.12 (m, 4H), 1.92 (s, 1H), 1.75 (s, 1H), 1.45 (s, 2H), 1.20 (s, 3H); MS (ESI) *m/z* 451 [C₂₈H₂₆N₄O₂+H]⁺.

Example 173: (4-amino-4-methylpiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)methanone



- Step: 1 To a solution of 6-bromoimidazo [1,2-a]pyridine (20 g, 101.5 mmol) in ACN (300 mL) was added *N*-iodosuccinimide (22.8 g, 101.5 mmol) at rt and stirred for 5 h. The reaction mixture was filtered and washed with hot acetonitrile to afford 6-bromo-3-iodoimidazo[1,2-a]pyridine (22 g, 67%) as a pale yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.29 (s, 1H), 7.58 (d, *J* = 9.2 Hz, 1H), 7.35 (d, *J* = 7.6 Hz, 1H), 7.27 (s, 1H).
- Step 2: A mixture of 6-bromo-3-iodoimidazo[1,2-a]pyridine (21 g, 65.4 mmol), 4-chlorophenylboronic acid (11.22 g, 71.9 mmol), K₃PO₄ (27.7g, 130.8 mmol) in DMF (100 mL), and water (15 mL) was degassed with argon for 30 min. Pd(PPh₃)₄ (3.77 g, 3.27 mmol) was added and again degassed with argon for 30 min and the reaction mixture was heated at 90 °C for 6 h. TLC indicated absence of SM and formation of two polar spots. Water (2x100 ml) was added to the reaction mixture to induce precipitation which was filtered to give crude product. The crude product was purified by column chromatography (silica gel, eluent (CHCl₃/MeOH 95:5) to afford 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyridine (10.6 g, 52%) as a yellowish green solid. ¹H NMR (400 MHz, CDCl₃) δ 8.38 (s, 1H), 7.69 (s, 1H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.82 (t, *J* = 9.2 Hz, 1H), 7.58 (q, *J* = 5.2 Hz, 4H), MS (ESI) *m/z* 309 [C₁₃H₈BrN₂Cl + 2H]⁺

- Step 3: 4-(ethoxycarbonyl) phenylboronic acid (3.81 g, 19.67 mmol), K₃PO₄ (6.95 g, 32.78 mmol), and Pd(PPh₃)₄ (0.56 g, 0.49 mmol) were added sequentially to a solution of ethyl 6-bromo-3-(4-chlorophenyl)imidazo[1,2-a]pyridine (5 g, 16.39 mmol) in a mixture of 1,4-dioxane/H₂O (50:5 mL) at room temperature under argon atmosphere. The reaction mixture was refluxed for 6 h and was diluted with water (100 mL) and extracted with EtOAc (3x100 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under

reduced pressure. The residue was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 1:1) to afford ethyl 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (2.7 mg, 58%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 9.21 (s, 1H), 8.95 (s, 1H), 8.21 (s, 1H), 8.17–8.12 (m, 5H), 8.08 (s, 1H), 7.91 (d, *J* = 8.0 Hz, 2H), 7.82 (t, *J* = 8.4 Hz, 1H), 4.03 (q, *J* = 7.2 Hz, 2H), 1.42 (t, *J* = 7.2 Hz, 3H); MS (ESI) *m/z* 378 [C₂₂H₁₇ClN₂O₂ + H]⁺


Step 4: To a solution of ethyl 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (2.7 g, 7.18 mmol) and LiOH (0.9 g, 21.54 mmol) in a mixture of THF (40 mL), water (10 mL) and MeOH (10 mL) was stirred for 5 h at rt. The reaction mixture was concentrated under reduced pressure to afford 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (900 mg, 37%) as an off white solid which was used in the next step without purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.87 (s, 1 H), 8.14 (s, 1 H), 8.12–8.02 (m, 5 H), 7.95–7.87 (m, 5 H); MS (ESI) *m/z* 349 [C₂₀H₁₃ClN₂O₂ + H]⁺.

Step 5: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (500 mg, 1.43 mmol) in DMF (15.0 mL) were added HATU (817 mg, 2.14 mmol), *N*-methylmorpholine (0.317 mL, 2.86 mmol) and *tert*-butyl 4-methylpiperidin-4-ylcarbamate (336 mg, 1.58 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then was diluted with water (15 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na₂SO₄, filtered and was concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) to afford *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (410 mg, 52%, LC-MS 92%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.68 (s, 1 H), 7.85–7.77 (m, 6 H), 7.70–7.66 (m, 1 H), 7.61 (d, *J* = 12.0 Hz, 2 H), 7.46 (d, *J* = 12.0 Hz, 2 H), 3.79 (bs, 2 H), 3.45 (bs, 3 H), 1.45 (bs, 5 H), 1.38 (s, 9 H), 1.10 (s, 3 H); MS (ESI) *m/z* 545 [C₃₁H₃₃ClN₄O₃ + H]⁺.

Step 6: To a solution of *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (400 mg, 0.73 mmol) in dichloromethane (10 mL) was added TFA (3 mL) in dichloromethane (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL), NaHCO₃ (100 mL) and extracted with CH₂Cl₂ (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. Crude compound was purified by prep-

HPLC to afford (4-amino-4-methylpiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)methanone (80 mg, 16%, AUC HPLC 97.59%) as a yellow solid; m.p. 109-124°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.68 (s, 1 H), 7.85-7.77 (m, 6 H), 7.70-7.66 (m, 1 H), 7.61 (d, *J* = 12.0 Hz, 2 H), 7.46 (d, *J* = 12.0 Hz, 2 H), 3.79 (bs, 2 H), 3.45 (bs, 3 H), 1.45 (bs, 5 H), 1.10 (s, 3 H); MS (ESI) *m/z* 445 [C₂₆H₂₅ClN₄O + H]⁺.

Example 174: (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)methanone




Step 1: To a solution of 4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoic acid (400 mg, 1.14 mmol) in DMF (15.0 mL) were added HATU (651 mg, 1.71 mmol), *N*-methylmorpholine (0.253 mL, 2.28 mmol) and *tert*-butyl piperidin-4-ylcarbamate (251 mg, 1.26 mmol). The reaction mixture was stirred at room temperature under inert atmosphere for 16 h, then it was diluted with water (15 mL) and extracted with EtOAc (3×25 mL). The combined organic layer was dried over Na₂SO₄ and was concentrated under reduced pressure. The crude residue was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5 to afford *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (310 mg, 50%, LC-MS 91%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.42 (s, 1H), 8.0 (s, 1H), 7.79 (d, *J* = 9.2 Hz, 2H), 7.73 (s, 1H), 7.58-7.45 (m, 7H), 4.60 (bs, 1H), 4.45 (bs, 1H), 3.78 (bs, 2H), 3.18 (bs, 2H), 3.00 (bs, 2H), 2.18-1.89 (m, 2H), 1.44 (s, 9H); MS (ESI) *m/z* 531 [C₃₀H₃₁ClN₄O₃ + H]⁺.


Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (310 g, 0.73 mmol) in DCM (15 mL) was added TFA (3 mL) and the reaction mixture was stirred at rt for 3 h. The reaction mixture was concentrated and poured into ice cold water, extracted with EtOAc (3 x 50 mL). The organic layer was washed with NaHCO₃ solution (2x20 mL), water (2x10 mL) and brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude

compound was purified by preparative HPLC to afford (4-aminopiperidin-1-yl)(4-(3-(4-chlorophenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)methanone (80 mg, 21%, AUC HPLC 97.7%) as a white solid; m.p. 85-89 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.69 (s, 1H), 7.85 (s, 1H), 7.82-7.78 (m, 5H), 7.70-7.61 (m, 3H), 7.46 (d, *J* = 8.0 Hz, 2H), 4.28 (bs, 1H), 3.58 (s, 1H), 3.07 (bs, 1H), 2.94 (bs, 1H), 2.82 (m, 1H), 1.75-1.67 (m, 4H), 1.23 (s, 1H), 1.18 (bs, 1H); MS (ESI) *m/z* 431.44 [C₂₅H₂₃ClN₄O + H]⁺.


Example 175: (4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)(4-methylpiperazin-1-yl)methanone



4-hydroxyphenylboronic acid (136 mg, 0.98 mmol), K₃PO₄ (380 mg, 1.79 mmol) and Pd(PPh₃)₄ (51 mg, 0.04 mmol) were added sequentially to a solution of 4-(3-iodoimidazo[1,2-a]pyridin-6-yl)phenyl (4-methylpiperazin-1-yl)methanone (400 mg, 0.896 mmol) in a mixture of 1,4-dioxane/H₂O (15:1.5 mL) at room temperature under argon atmosphere. The reaction mixture was refluxed for 6 h and was diluted with water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 95:5) and by preparative HPLC to afford of (4-(3-(4-hydroxyphenyl)imidazo[1,2-a]pyridin-6-yl)phenyl)(4-methylpiperazin-1-yl) (100 mg, 23%, AUC HPLC 98.3%) as an off-white solid; m.p. 151-217 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.79 (s, 1H), 8.55 (s, 1H), 7.78-7.72 (m, 3H), 7.67-7.46 (m, 6H), 6.94 (d, *J* = 12.0 Hz, 2H), 3.62 (bs, 2H), 3.35 (bs, 2H), 2.31 (m, 4H), 2.20 (s, 3H); MS (ESI) *m/z* 413 [C₂₅H₂₄N₄O + H]⁺.

**Formula 5**

Example 176: 4-(5-(4-(4-methylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



5
Step 1: To a solution of 5-bromopyrazolo[1,5-a]pyrimidine (300 mg, 1.5 mmol) in 1,4-dioxane (10 mL) and water (2 mL), were sequentially added 4-ethoxycarbonyl phenyl boronic acid (380 mg, 1.95 mmol), K_3PO_4 (955 mg, 4.5 mmol) and $Pd(PPh_3)_4$ (52 mg, 0.04 mmol). The reaction mixture was heated at 90 °C for 4 h under argon atmosphere then, was
10 diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 9:1 to 1:1) to afford ethyl 4-(pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (200 mg, 49%) of as a yellow solid. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.77 (d, $J = 8.0$, 1H), 8.18-8.15 (m, 4H), 7.33 (d, $J = 7.5$ Hz, 2H), 7.12 (d, $J = 4.4$ Hz, 1H), 6.77 (d, $J = 2.2$ Hz, 1H), 4.45 (q, $J = 6.8$ Hz, 2H),
15 1.44 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 268 [$C_{15}H_{13}N_3O_2 + H$] $^+$.

Step 2: To a solution of ethyl 4-(pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (2.3 g, 9.36 mmol) in DCM (15 mL) and ACN (35 mL) was added NBS (2.02 g, 11.23 mmol) at 0 °C and stirred
20 at same temperature for 1 h. The reaction mixture was filtered and washed with water and dried to afford ethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)benzoate (2.0 g, 67%) as a yellow solid. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.77 (d, $J = 8.0$ Hz, 1H), 8.18-8.15 (m,

4H), 8.14 (s, 1H), 7.38 (d, $J = 4.4$ Hz, 1H), 4.40 (q, $J = 6.8$ Hz, 2H), 1.44 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 346 [$C_{15}H_{12}BrN_3O_2$] $^+$.


Step 3: A solution of ethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)benzoate (120 mg, 0.34 mmol), 4-cyanophenylboronic acid (66 mg, 0.44 mmol), K_3PO_4 (220 mg, 1.03 mmol) in 1,4-dioxane (3 mL) and water (1 mL) was degassed with argon for 30 min prior to the addition of $Pd(PPh_3)_4$ (12 mg, 0.01 mmol). The reaction mixture was heated at 90 °C for 3 h under argon then, was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 90/10 to 50/50) to afford of ethyl 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (80 mg, 61%) as a yellow solid. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.80 (d, $J = 7.5$ Hz, 1H), 8.58 (s, 1H), 8.30-8.20 (m, 6H), 7.79 (d, $J = 12.0$ Hz, 2H), 7.45 (d, $J = 12.0$ Hz, 1H), 4.40 (q, $J = 6.8$ Hz, 2H), 1.44 (t, $J = 7.2$ Hz, 3H); MS (ESI) m/z 369 [$C_{22}H_{16}N_4O_2 + H$] $^+$.

Step 4: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (500 mg, 1.35 mmol) in THF (20 mL) was added LiOH (228 mg, 5.43 mmol) in water (4 mL) at rt and stirred for 16 h. TLC indicated absence of SM and formation of a polar spot. The reaction mixture was concentrated under reduced pressure to afford 700 mg of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid as a yellow solid which was used in the next step without purification. 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 1H NMR (400 MHz, DMSO- d_6) δ (ppm): 12.2 (bs, 1H), 9.35 (d, $J = 6.8$ Hz, 1H), 8.97 (s, 1H), 8.48-8.44 (m, 2H), 8.34 (d, $J = 8.0$ Hz, 1H), 8.14 (d, $J = 8.4$ Hz, 2H), 8.09 (d, $J = 8.0$ Hz, 2H), 7.93 (d, $J = 8.0$ Hz, 1H), 7.86 (d, $J = 8.0$ Hz, 1H); MS (ESI) m/z 340 [$C_{20}H_{12}N_4O_2 + H$] $^+$.

Step 5: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (460 mg, 1.35 mmol) in DMF (5 mL) was added NMM (348 mg, 2.7 mmol) followed by addition of HATU (775 mg, 2.03 mmol) at rt and stirred for 30 min. 1-Methylpiperazine (270 mg, 2.7 mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent $CHCl_3/MeOH$ 90:10) and followed by preparative HPLC to afford 4-(5-(4-(4-methylpiperazine-1-


carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile, (50 mg, 9%, AUC HPLC 97.6%) as a yellow solid; m.p. 111-124 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.78 (d, *J* = 8.0 Hz, 1H), 8.52 (s, 1H), 8.29 (d, *J* = 16.0 Hz, 2H), 8.23 (d, *J* = 16.0 Hz, 2H), 7.75 (d, *J* = 16.0 Hz, 2H), 7.61 (d, *J* = 16.0 Hz, 1H), 7.41 (d, *J* = 12.0 Hz, 1H), 3.84 (bs, 2H), 3.49 (bs, 2H), 2.53 (bs, 2H), 2.39 (bs, 2H), 2.34 (s, 3H); MS (ESI) *m/z* 423 [C₂₅H₂₂N₆O + H]⁺.

Example 177: 4-(5-(4-(4-isopropylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic (400 mg, 1.17 mmol) in DMF (5 mL) was added NMM (303 mg, 2.3 mmol) followed by HATU (670 mg, 1.76 mmol) at rt and the reaction mixture was stirred for 30 min. 1-isopropylpiperazine (196 mg, 1.52 mmol) was added to the reaction mixture and stirred was continued at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by Prep-HPLC to afford 4-(5-(4-(4-methylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (45 mg, 8%, AUC HPLC >99%) as a yellow solid; m.p. 106-116 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.77 (d, *J* = 8.0 Hz, 1H), 8.51 (m, 1H), 8.30 (d, *J* = 12 Hz, 2H), 8.22 (d, *J* = 12 Hz, 2H), 7.75 (d, *J* = 8.0 Hz, 2H), 7.61 (d, *J* = 8.0 Hz, 2H), 7.40 (d, *J* = 8.0 Hz, 1H), 3.83 (s, 2H), 3.47 (bs, 2H), 2.75 (m, 1H), 2.63 (bs, 2H), 2.49 (bs, 2H), 1.06 (d, *J* = 8.0 Hz, 6H); MS (ESI) *m/z* 451 [C₂₇H₂₆N₆O + H]⁺.

Example 178: 4-(5-(4-(4-ethylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



- To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (400 mg, 1.17 mmol) in DMF (5 mL) was added NMM (303 mg, 2.35 mmol) followed by HATU (670 mg, 1.76 mmol) at rt and the solution was stirred for 30 min. 1-methylpiperazine (270 mg, 2.7 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by preparative HPLC to afford 4-(5-(4-(4-ethylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (15 mg, AUC HPLC 97.4%) as a yellow solid; m.p. 200-202 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.77 (d, $J = 8.0$ Hz, 1 H), 8.51 (s, 1 H), 8.30-8.20 (m, 4H), 7.75 (d, $J = 12.0$ Hz, 2H), 7.61 (d, $J = 12.0$ Hz, 2H), 7.40 (d, $J = 8.0$ Hz, 1H), 3.85 (bs, 2H), 3.50 (bs, 2H), 2.56 (bs, 4H), 2.47 (q, $J = 8.0$ Hz, 2H), 1.11 (t, $J = 8.0$ Hz, 3H); MS (ESI) m/z 437 $[\text{C}_{26}\text{H}_{24}\text{N}_6\text{O} + \text{H}]^+$.


Example 179: 4-(5-(4-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



- To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (200 mg, 0.588 mmol) in DMF (5 mL), was added NMM (117 mg, 1.17 mmol) followed by HATU (667 mg, 0.88 mmol) and the solution was stirred at room temperature for 30 min. 1-methyl-

4-(piperidin-4-yl)piperazine (118 mg, 0.646 mmol) was added to the reaction mixture and stirring was maintained for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by Preparative HPLC (C18, ACN/H₂O/0.075% NH₄HCO₃) to afford 4-(5-(4-(4-(4-methylpiperazin-1-yl)piperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (30 mg, 10%, AUC HPLC 98.6%) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.78 (d, *J* = 8.7 Hz, 1H), 8.52 (s, 1H), 8.30 (d, *J* = 8.2 Hz, 2H), 8.23 (d, *J* = 8.2 Hz, 2H), 7.768 (d, *J* = 7.7 Hz, 2H), 7.62 (d, *J* = 7.6 Hz, 2H), 7.41 (d, *J* = 7.4 Hz, 1H), 5.6 (bs, 1H), 3.8 (bs, 1H), 3.19 (bs, 1H), 2.6 (bs, 4H), 2.4 (bs, 4H), 2.3 (bs, 3H), 2.06-1.82 (m, 2H), 1.45 (bs, 3H); MS (ESI) *m/z* 506 [C₃₀H₃₁N₇O + H]⁺.


Example 180: 4-(5-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (200 mg, 0.588 mmol) in DMF (5 mL) was added NMM (117 mg, 1.17 mmol) followed by HATU (667 mg, 0.88 mmol) and the solution was stirred at rt for 30 min. *N,N*-dimethylpiperidin-4-amine (82 mg, 0.646 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent CH₂Cl₂/MeOH 95:5) followed by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford 4-(5-(4-(4-(dimethylamino)piperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (50 mg, 19%, AUC HPLC 99.8%) as a yellow solid; ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.78 (d, *J* = 8.7 Hz, 1H), 8.52 (s, 1H), 8.30 (d, *J* = 8.2 Hz, 2H), 8.23 (d, *J* = 8.2 Hz, 2H), 7.76 (d, *J* = 7.7 Hz, 2H), 7.62 (d, *J* = 7.6 Hz, 2H), 7.41 (d, *J* = 7.4 Hz, 1H), 4.9 (bs, 1H), 3.90 (bs, 1H), 3.17 (bs, 1H), 2.91

(bs, 1H), 2.45 (bs, 1H), 2.38 (bs, 6H), 2.08 (bs, 1H), 1.95 (bs, 1H), 1.52 (bs, 2H); MS (ESI) m/z 451 [$C_{27}H_{26}N_6O + H$] $^{+}$.


5 Example 181: 4-(5-(4-(1,4-oxazepane-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (200 mg, 0.58 mmol) in DMF (5 mL) was added NMM (117 mg, 1.17 mmol) followed by HATU (667 mg, 0.88 mmol) at rt and the solution was stirred for 30 min. 1,4-oxazepane (65 mg, 0.646 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) followed by Preparative HPLC (C18, $ACN/H_2O/0.075\% NH_4HCO_3$) to afford 4-(5-(4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (20 mg, 8%, AUC HPLC 99%) as a yellow solid; 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.78 (d, $J = 8.7$ Hz, 1H), 8.52 (s, 1H), 8.30 (d, $J = 8.3$ Hz, 2H), 8.23 (d, $J = 8.2$ Hz, 2H), 7.768 (d, $J = 7.7$ Hz, 2H), 7.62 (d, $J = 7.6$ Hz, 2H), 7.41 (d, $J = 7.4$ Hz, 1H) 4.09-3.80 (m, 5H), 3.7 (bs, 1H), 3.67 (bs, 2H), 2.1 (bs, 1H); 1.9 (bs, 1H); MS (ESI) m/z 424 [$C_{25}H_{21}N_5O_2 + H$] $^{+}$.

20

Example 182: N-(1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)-4-methylpiperidin-4-yl)acetamide



Step 1: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (600 mg, 1.76 mmol) in DMF (5 mL) was added NMM (266 mg, 2.64 mmol) followed by

25


HATU (1.0 g, 2.64 mmol) at rt and the solution was stirred for 30 min. *tert*-butyl 4-methylpiperidin-4-ylcarbamate (414.3 mg, 1.93 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to obtain crude product to afford *tert*-butyl 1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate as a yellow solid (400 mg, 42.5%). ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.77 (d, *J* = 7.5 Hz, 1H), 8.52 (s, 1H), 8.30 (d, *J* = 8.3 Hz, 2H), 8.22 (d, *J* = 7.9 Hz, 2H), 7.75 (d, *J* = 8.4 Hz, 2H), 7.59 (d, *J* = 8.0 Hz, 2H), 7.41 (d, *J* = 8.8 Hz, 1H), 4.39 (bs, 1H), 4.1 (s, 1H), 3.41-3.32 (m, 3H), 2.3 (m, 2H), 1.67-1.44 (m, 2H), 1.44 (s, 3H), 1.25 (s, 9H); MS (ESI) *m/z* 537 [C₃₁H₃₂N₆O₃ + H]⁺.

Step 2: To a solution of *tert*-butyl 1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)-4-methylpiperidin-4-ylcarbamate (400 mg, 0.73 mmol) in dichloromethane (10 mL) was added TFA (3 mL) in dichloromethane (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL), NaHCO₃ (100 mL) and extracted with CH₂Cl₂ (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford 4-(5-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile as a crude product (300 mg, 93%) as a yellow solid; ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 9.33 (d, *J* = 7.2 Hz, 1H), 8.96 (s, 1H), 8.46 (d, *J* = 8.4 Hz, 2H), 8.41 (d, *J* = 8.4 Hz, 2H), 7.93 (d, *J* = 8.0 Hz, 2H), 7.84 (d, *J* = 7.2 Hz, 2H), 7.60 (d, *J* = 8.4 Hz, 1H), 5.20 (bs, 2H), 3.44-3.35 (m, 4H), 1.90-1.60 (m, 4H), 1.38 (s, 3H); MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺; MS (ESI) *m/z* 437 [C₂₆H₂₄N₆O + H]⁺.

Step 3: To a solution of 4-(5-(4-(4-amino-4-methylpiperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile 7 (300 mg, 0.73 mmol) in dichloromethane (10 mL) were added TEA (3 mL) and acetyl chloride in dichloromethane (63 mg, 0.825 mmol). The reaction mixture was stirred for 2 h at rt. The reaction mixture was diluted with water (100 mL), NaHCO₃ (100 mL) and extracted with CH₂Cl₂ (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. The crude residue was purified by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford N-(1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)-4-methylpiperidin-4-yl)acetamide (180 mg, 56%, AUC HPLC 99.48%) as a yellow solid; ¹H

- NMR (400 MHz, CDCl₃) δ (ppm): 8.78 (d, J = 7.6 Hz, 1H), 8.52 (s, 1H), 8.30 (d, J = 8.4 Hz, 2H), 8.23 (d, J = 8.4 Hz, 2H), 7.76 (d, J = 8.4 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.41 (d, J = 7.6 Hz, 1H), 5.19 (s, 1H), 4.1 (bs, 1H), 3.54-3.25 (m, 3H), 2.3-2.10 (m, 2H), 2.0 (s, 3H), 1.6-1.70 (m, 2H), 1.42 (s, 3H); MS (ESI) m/z 479.35 [C₂₈H₂₆N₆O₂ + H]⁺; MS (ESI) m/z 479
- 5 [C₂₈H₂₆N₆O₂ + H]⁺.

Example 183: 4-(5-(4-(piperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile




- 10 Step 1: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (400 mg, 1.17 mmol) in DMF (5 mL) was added NMM (234 mg, 2.35 mmol) followed by HATU (667 mg, 1.76 mmol) at rt and the solution was stirred for 30 min. *tert*-butyl piperazine-1-carboxylate (239 mg, 1.28 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water
- 15 and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product to afford *tert*-butyl 4-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)piperazine-1-carboxylate (250 mg, 42%); ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.78 (d, J = 7.6 Hz, 1H), 8.52 (s, 1H), 8.30 (d, J = 8.0 Hz, 2H), 8.23 (d, J = 8.4 Hz, 2H), 7.75 (d, J = 8.4 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H),
- 20 7.41 (d, J = 7.6 Hz, 1H) 3.90-3.40 (m, 8H), 1.48 (s, 9H); MS (ESI) m/z 509 [C₂₉H₂₈N₆O₃ + H]⁺; MS (ESI) m/z 509 [C₂₉H₂₈N₆O₃ + H]⁺.

- Step 2: To a solution of *tert*-butyl 4-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)piperazine-1-carboxylate (250 mg, 0.492 mmol) in dichloromethane (10 mL)
- 25 were added TFA (3 mL) in dichloromethane (5 mL). The reaction mixture was stirred for 4 h at rt. The reaction mixture was diluted with water (100 mL), NaHCO₃ (100 mL) and extracted with CH₂Cl₂ (2×50 mL). The combined organic layer was concentrated to dryness under reduced pressure to afford crude compound. The crude product was purified by preparative HPLC (C18, ACN/H₂O/ 10 mM NH₄HCO₃) to afford 4-(5-(4-(piperazine-1-
- 30 carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (120 mg, 38% , AUC HPLC

97.14%) as a yellow solid; 8.78 (d, $J = 7.6$ Hz, 1H), 8.52 (s, 1H), 8.30 (d, $J = 8.4$ Hz, 2H), 8.23 (d, $J = 8.4$ Hz, 2H), 7.76 (d, $J = 8.4$ Hz, 2H), 7.6 (d, $J = 8.0$ Hz, 2H), 7.42 (d, $J = 7.6$ Hz, 1H), 3.8 (bs, 2H), 3.43 (bs, 2H), 3.00-2.80 (m, 4H); MS (ESI) m/z 409 [$C_{24}H_{20}N_6O + H$] $^+$.

5 Example 184: 4-(5-(4-(4-methylpiperazine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile




To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid 6 (400 mg, 1.17 mmol) in DMF (5 mL) was added NMM (234 mg, 2.35 mmol) followed by HATU (667 mg, 1.76 mmol) at rt and the solution was stirred for 30 min. 1-(piperazin-1-yl)ethanone (164.7 mg, 1.28 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude residue was purified by column chromatography (silica gel, eluent $CH_2Cl_2/MeOH$ 95:5) to afford 4-(5-(4-(4-acetyl piperazine-1-

10 carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (220 mg, 42%, AUC HPLC 98.4%) as a yellow solid; 1H NMR (400 MHz, $CDCl_3$) δ (ppm): 8.78 (d, $J = 8.7$ Hz, 1H), 8.52 (s, 1H), 8.30 (d, $J = 8.2$ Hz, 2H), 8.23 (d, $J = 8.2$ Hz, 2H), 7.76 (d, $J = 7.7$ Hz, 2H), 7.62 (d, $J = 7.6$ Hz, 2H), 7.41 (d, $J = 7.4$ Hz, 1H) 3.9-3.4 (m, 8H), 2.16 (bs, 3H); MS (ESI) m/z 451.1

15 [C₂₆H₂₂N₆O₂ + H] $^+$.

20


Example 185: 4-(5-(4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



25 To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoic acid (200 mg, 0.58 mmol) in DMF (5 mL) was added NMM (117 mg, 1.17 mmol) followed by HATU (667

mg, 0.88 mmol) at rt and the solution was stirred for 30 min. Morpholine (56 mg, 0.646 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 95:5) to afford 4-(5-(4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (120 mg, 75%, AUC HPLC 97.6%) as a yellow solid; ^1H NMR (400 MHz, CDCl_3) δ (ppm): 8.78 (d, $J = 8.7$ Hz, 1H), 8.52 (s, 1H), 8.30 (d, $J = 8.2$ Hz, 2H), 8.23 (d, $J = 8.2$ Hz, 2H), 7.76 (d, $J = 7.7$ Hz, 2H), 7.62 (d, $J = 7.6$ Hz, 2H), 7.41 (d, $J = 7.4$ Hz, 1H) 3.9-3.4 (m, 8H); MS (ESI) m/z 410 [$\text{C}_{24}\text{H}_{19}\text{N}_5\text{O}_2 + \text{H}$] $^+$.

Example 186: N-(1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)piperidin-4-yl)acetamide




Step 1: To a solution of tert-butyl 1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)piperidin-4-ylcarbamate (450 mg, 0.865 mmol) in DCM (10 mL) was added HCl in 1,4 dioxane (7 mL) at 0 °C and stirred at 0 °C to rt for 4 h. The reaction mixture was basified with an aqueous solution of NaHCO_3 and then extracted with DCM (60 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford of 4-(5-(4-(4-aminopiperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (300 mg, 80%, LC-MS 91%) as a yellow solid which was used in the next step without purification.

Step 2: To a solution of 4-(5-(4-(4-aminopiperidine-1-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (300 mg, 0.71 mmol) in DCM (5 mL) were added TEA (0.25 mL, 2.13 mmol) and acetyl chloride (0.07 g, 0.855 mmol). The reaction mixture was stirred at 0 °C to room temperature under inert atmosphere for 1 h, and was diluted with ice-cold water (10 mL) and extracted with DCM (25 mL). The organic layer was dried over Na_2SO_4 and was concentrated under reduced pressure. The crude product was purified by column

chromatography (silica gel, eluent MeOH/DCM 5:95) to afford N-(1-(4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoyl)piperidin-4-yl)acetamide (61 mg, 18%, AUC HPLC 96.26%) as a yellow solid; mp. 295-297 °C; ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.77 (d, *J* = 7.6 Hz, 1H), 8.52 (s, 1H), 8.29 (d, *J* = 8.0 Hz, 2H), 8.22 (d, *J* = 8.4 Hz, 2H), 7.75 (d, *J* = 8.4 Hz, 2H), 7.60 (d, *J* = 8.4 Hz, 2H), 7.40 (d, *J* = 7.2 Hz, 1H), 5.37 (d, *J* = 8.0 Hz, 1H), 4.71 (bs, 1H), 4.09-4.06 (m, 1H), 3.78 (bs, 1H), 3.19 (bs, 1H), 2.98 (bs, 1H), 2.10 (bs, 2H), 2.00 (s, 3H), 1.37 (bs, 2H); MS (ESI) *m/z* 465 [C₂₇H₂₄N₆O₂ + H]⁺.

10 Example 187: 4-(5-(3-hydroxy-4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile



Step 1: To a mixture of methyl 4-bromo-2-methoxybenzoate (3 g, 12.24 mmol), bis(pinacolato)diboron (3.42 g, 13.46 mmol), KOAc (3.59 g, 36.72 mmol) in 1,4-dioxane (15 mL) was added PdCl₂(PPh₃)₂ (300 mg), and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was used in next step without isolation.

Step 2: To a mixture methyl 2-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3.2 g, 11.04 mmol), 5-chloropyrazolo[1,5-a]pyrimidine (1.3 g, 8.49 mmol), K₃PO₄ (3.6 g, 16.99 mmol) in 1,4-dioxane (20 mL) and water (2 mL) was added Pd(PPh₃)₄ (150 mg) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain Preparation of methyl 2-methoxy-4-(pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (850 mg, LC-MS 83%) as a crude product which was used in the next step without purification.

25 Step 3: To a solution of methyl 2-methoxy-4-(pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (850 mg, 3.00 mmol) in ACN (30 mL) was added N-iodosuccinimide (810 mg, 3.60 mmol) at rt and stirred for 2 h. Water was added to the reaction mixture and the resulting solid was

filtered and washed with water to afford methyl 4-((4-iodo-1-methyl-1H-pyrazol-5-ylimino)methyl)-2-methoxybenzoate (1.1 g, 90%, LC-MS 70%).

Step 4: To a mixture of (E)-methyl 4-((4-iodo-1-methyl-1H-pyrazol-5-ylimino)methyl)-2-methoxybenzoate (1.1 g, 2.68 mmol), 4-cyanophenylboronic acid (0.51 g, 3.49 mmol), K_3PO_4 (1.14 g, 5.37 mmol) in 1,4-dioxane (20 mL) and water (3 mL) was added $Pd(PPh_3)_4$ (150 mg) and the reaction mixture was heated at 90 °C for 16 h under argon. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent $CHCl_3/MeOH$ 95:5) to afford methyl 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-methoxybenzoate (1 g, 90%, LC-MS 44%) as a yellow solid.

Step 5: To a solution of ethyl 4-(3-(4-cyanophenyl)imidazo[1,2-a]pyridin-6-yl)benzoate (2.7 g, 7.18 mmol) in THF (40 mL) was added LiOH (0.9 g, 21.54 mmol) in water (10 mL) and MeOH (10 mL) at rt and the reaction mixture was stirred for 5 h. The reaction mixture was concentrated under reduced pressure to give crude compound which was washed with EtOAc and water was added to the reaction mixture then diluted with EtOAc. The aqueous layer was acidified with sat. $KHSO_4$ solution then solid thus formed was filtered and dried to afford 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-methoxybenzoic acid (600 mg, 64%, LC-MS 51%) which was used in the next step without purification.

Step 6: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-methoxybenzoic acid (300 mg, 0.81 mmol) in DMF (6 mL) was added NMM (0.163 g, 1.62 mmol) followed by HATU (0.42 g, 1.21 mmol) at rt and stirred for 30 min. Morpholine (84 mg, 0.97 mmol) was added and stirring was continued at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude product which was purified by column chromatography (silica gel, eluent MeOH/DCM 10:90) to afford 4-(5-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (300 mg, 84%, LC-MS 27%) as a yellow solid.

Step 7: To a solution of 4-(5-(3-methoxy-4-(morpholine-4-carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (300 mg) in DCM was added BBr_3 (3 mL) at 0 °C and stirred

for 2 h at rt. The reaction mixture was basified with aq NaHSO₄ and extracted with DCM.


The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product which purified by column chromatography (silica gel, eluent MeOH-

DCM 10:90) followed by preparative HPLC to afford 4-(5-(3-hydroxy-4-(morpholine-4-

- 5 carbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-3-yl)benzonitrile (25 mg, 8%, AUC HPLC 98.06%) as a yellow solid; mp. 290-295 °C; ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.81 (s, 1H), 8.77 (d, *J* = 7.2 Hz, 1H), 8.52 (s, 1H), 8.29 (d, *J* = 8.0 Hz, 2H), 7.78-7.72 (m, 4H), 7.44 (d, *J* = 7.6 Hz, 1H), 7.38 (d, *J* = 7.6 Hz, 1H), 3.81-3.79 (m, 8H); MS (ESI) *m/z* 426 [C₂₄H₁₉N₅O₃+H]⁺.

-10

Example 188: N-(5-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide



- Step 1: To a solution of methyl-2-acetamido-4-bromobenzoate (2 g, 7.38 mmol), bis(pinacolato)diboron (2.06 g, 8.11 mmol), KOAc (2.16 g, 22.14 mmol) in 1,4-dioxane (15 mL) was added Pd₂(dba)₃ (200 mg) and P(Cy)₃ (200 mg) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was used in next step without isolation.

- Step 2: To a mixture of methyl 2-acetamido-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (2.3 g, 7.18 mmol), 5-chloropyrazolo[1,5-a]pyrimidine (1 g, 6.53 mmol), K₃PO₄ (2.77 g, 13.07 mmol) in 1,4-dioxane (20 mL) and water (2 mL) was added Pd(PPh₃)₄ (150 mg) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent EtOAc/ petroleum ether 50:50) to afford methyl 2-acetamido-4-(pyrazolo[1,5-a]pyrimidin-5-yl)benzoate (600

mg, 30%, LC-MS 98%) as a pale yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 11.14 (s, 1H), 9.43 (s, 1H), 8.74 (d, *J* = 7.6 Hz, 1H), 8.17 (d, *J* = 8.1 Hz, 1H), 7.94 (d, *J* = 1.6 Hz, 1H), 7.93-7.92 (m, 1H), 7.38 (d, *J* = 7.6 Hz, 1H), 6.76 (s, 1H), 3.97 (s, 3H), 2.29 (s, 3H); MS (ESI) *m/z* 311 [M+H]⁺.

5

Step 3: To a solution of methyl 2-acetamido-4-(pyrazolo[1,5-*a*]pyrimidin-5-yl)benzoate (600 mg, 1.93 mmol) in ACN (30 mL) was added *N*-iodosuccinimide (522 mg, 2.32 mmol) at rt and stirred for 2 h. Water was added to the reaction mixture and the resulting solid was filtered and washed with water to afford methyl 2-acetamido-4-(3-iodopyrazolo[1,5-

10 *a*]pyrimidin-5-yl)benzoate (0.7 g, 83%, LC-MS 87%) as a light brown solid. ¹H NMR (400 MHz, CDCl₃) δ 11.15 (s, 1H), 9.46 (s, 1H), 8.70 (d, *J* = 7.6 Hz, 1H), 8.18 (d, *J* = 8.8 Hz, 2H), 8.09 (d, *J* = 8.4 Hz, 1H), 7.43 (d, *J* = 7.6 Hz, 1H), 3.98 (s, 3H), 2.29 (s, 3H); MS (ESI) *m/z* 437 [C₁₆H₁₃N₄O₃+H].

15 Step 4: To a mixture of methyl 2-acetamido-4-(3-iodopyrazolo[1,5-*a*]pyrimidin-5-yl)benzoate (0.7 g, 1.60 mmol), 4-cyanophenylboronic acid (0.3 g, 2.08 mmol), K₃PO₄ (0.68 g, 3.21 mmol) in 1,4-dioxane (20 mL), and water (3 mL) was added Pd(PPh₃)₄ (100 mg) and the reaction mixture was heated at 90 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous

20 Na₂SO₄ and concentrated under reduced pressure to obtain the crude product. The crude product was purified by column chromatography (silica gel, eluent CHCl₃/MeOH 95:5) to afford methyl 2-acetamido-4-((*E*)-1-(4-cyanostyrylimino)allyl)benzoate (550 mg, 85%, LC-MS 30%) as a yellow solid. The intermediate was used in the next step without purification.


25 Step 5: To a solution of methyl 2-acetamido-4-((*E*)-1-(4-cyanostyrylimino)allyl)benzoate (0.55 g, 1.33 mmol) in THF (6 mL) was added LiOH (0.11 g, 2.67 mmol) in water (2 mL) and MeOH (4 mL) at rt and stirred for 5 h. The reaction mixture was concentrated under reduced pressure to 2-acetamido-4-(3-(4-cyanophenyl)pyrazolo[1,5-*a*]pyrimidin-5-yl)benzoic acid (350 mg, lithium salt, LC-MS 30%) as a yellow solid which was used in the

30 next step without purification.


Step 6: To a solution of 2-acetamido-4-(3-(4-cyanophenyl)pyrazolo[1,5-*a*]pyrimidin-5-yl)benzoic acid 5 (350 mg, 0.88 mmol) in DMF (6 mL) was added NMM (0.17 g, 1.76 mmol) followed by HATU (0.44 g, 1.16 mmol) at rt and stirred for 30 min. Morpholine (92 mg, 1.05

mmol) was added and stirred at rt for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain crude compound which was purified by column chromatography (silica gel, eluent MeOH/DCM 10:90) followed by preparative HPLC to afford *N*-(5-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-(morpholine-4-carbonyl)phenyl)acetamide (28 mg, 7%, AUC HPLC 95.24%) as a yellow solid. m.p 151-155 °C. ^1H NMR (400 MHz, CDCl_3) δ 9.16 (s, 1H), 9.10 (s, 1H), 8.76 (d, $J = 7.6$ Hz, 1H), 8.52 (s, 1H), 8.31 (d, $J = 8.0$ Hz, 1H), 8.04 (d, $J = 1.2$ Hz, 1H), 7.76 (d, $J = 8.4$ Hz, 2H), 7.46-7.41 (m, 2 H), 3.76 (bs, 8H), 2.25 (s, 2H); MS (ESI) m/z 465 $[\text{M}-1]$.

10

**Formula 6**

Example 189: 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)-1H-imidazo[4,5-c]pyridin-1-yl)benzonitrile



15

Step: 1 To a solution of anhydrous NH_3 (150 mL) in THF (50 mL) at -78°C was Potassium *t*-butoxide (19.1 g, 170 mmol) and the reaction mixture was allowed to warm to -

35 °C. A solution of 4-Methoxy-3-nitropyridine (10.5 g, 68.2 mmol) in 100 mL of THF was cooled to 0 °C and 15 mL of *t*-BuOOH (5 M in decane, 75 mmol) was added slowly over 5 min. The resulting solution was then added dropwise to the *t*-BuOK solution prepared above over 1 h, then stirred for 2 h at -35 °C and then carefully quenched with 50 mL of sat. aqueous solution of NH₄Cl. The reaction mixture was allowed to vent and warm to rt overnight, then the organics were concentrated and the residue was acidified with NH₄Cl solution and filtered. The resulting solid was washed with cold water and dried under vacuum to give the title compound as a brown solid (10.6 g, 91%). ¹H NMR (400 MHz, CDCl₃) δ 8.48 (s, 1H); 5.58 (s, 1H); 3.84 (s, 3H); MS (ESI) *m/z* 171 [M + H]⁺.


Step 2: Phosphorous oxybromide (5.73 g, 20 mmol) was added to a suspension of 4-methoxy-5-nitro-1H-pyridin-2-one (1.70 g, 10 mmol) in acetonitrile (20 mL) at rt and heated at reflux for 3 h. The reaction mixture was cooled and carefully poured on to icy saturated aqueous solution of K₂CO₃ then extracted with EtOAc. The organic extracts were combined, washed with water and brine, dried over MgSO₄, filtered and concentrated to give 2,4-dibromo-5-nitropyridine (2.1 g, 75 %) as a black solid. ¹H NMR (400 MHz, CDCl₃) δ 8.87 (s, 1H), 7.94 (s, 1H), MS (ESI) *m/z* 285 [M + H]⁺.

Step 3: A solution of 2,4-dibromo-5-nitropyridine (3 g, 7.16 mmol) in ethanol (25 mL) was cooled to 0 °C followed by addition of 4-cyanoaniline (0.845 g, 7.16 mmol) and triethylamine (1.52 mL, 10.75 mmol). The reaction mixture was stirred at rt for 24 h. After completion, the reaction mixture was cooled to 0 °C, filtered and dried with vacuum to afford 4-(2-bromo-5-nitropyridin-4-ylamino)benzonitrile (1.7 g, 38.8%) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 9.73 (bs, 1H), 9.09 (s, 1H), 7.82 (d, *J* = 7.80 Hz, 2H), 7.43 (d, *J* = 7.42 Hz, 2H); MS (ESI) *m/z* 317 [M]⁺.

Step 4: A solution of 4-(2-bromo-5-nitropyridin-4-ylamino)benzonitrile (1 g, 3.15 mmol) in ethanol (25 mL) was cooled to 0 °C followed by addition of SnCl₂·H₂O (2.13 g, 9.46 mmol). The reaction mixture was stirred at 80 °C for 4 h. After completion, the reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated NaHCO₃ (100 mL) followed by extraction with EtOAc (3x100 mL). The organic layer was washed with water, brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford 4-(5-amino-2-bromopyridin-4-ylamino)benzonitrile (0.8 g). MS (ESI) *m/z* 288 [M]⁺. Crude product was used without purification in the next step.


Step 5: A solution of 4-(5-amino-2-bromopyridin-4-ylamino)benzonitrile (0.8 g, 2.77 mmol) in triethylorthoformate (15 mL) was stirred at 100 °C for 16 h. After completion, the reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated aqueous solution of NaHCO₃ (100 mL) followed by extraction with EtOAc (3 x 100 mL). The organic layer was washed with water, brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography (silica gel, eluent petroleum ether/EtOAc 70:30) to afford 4-(6-bromo-1H-imidazo[4,5-c]pyridin-1-yl)benzonitrile (0.4 g, 48%); ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.9 (d, *J*=8.9 Hz, 2H), 8.15 (d, *J*=8.1 Hz, 2H), 7.9 (t, *J*=7.9 Hz, 3H), MS (ESI) *m/z* 297 [M]⁺.

Step 6: To a mixture of 4-(6-bromo-1H-imidazo[4,5-c]pyridin-1-yl)benzonitrile (300 mg, 1.0 mmol), (4-methylpiperazin-1-yl)(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)methanone (366 mg, 1.11 mmol), NaHCO₃ (169 mg, 2.02 mmol) in DMF (15 mL) and water (5 mL), was added (A-Phos)₂PdCl₂ (35 mg, 0.05 mmol) and the reaction mixture was heated at 90 °C for 1 h. The reaction mixture was diluted with water and extracted with EtOAc, the organic phase was washed with water and brine, dried over Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by Preparative TLC to afford of 4-(6-(4-(4-methylpiperazine-1-carbonyl)phenyl)-1H-imidazo[4,5-c]pyridin-1-yl)benzonitrile (60 mg, 15%, AUC HPLC 99.6%) as an off white solid. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 9.32 (s, 1 H), 8.23 (s, 1 H), 8.05 (d, *J*=8.0 Hz, 2 H), 7.97 (d, *J*=8.8 Hz, 2 H), 7.86 (s, 1 H), 7.72 (d, *J*=8.8 Hz, 2 H), 7.53 (d, *J*=8.4 Hz, 2 H), 3.83 (bs, 2 H), 3.48 (bs, 2 H), 2.51 (bs, 2 H), 2.34 (bs, 2 H), 2.33 (s, 3 H); MS (ESI) *m/z* 423.24 [C₂₅H₂₂N₆O +H]⁺.



Formula 7

Example 190: Synthesis of 4-(6-(4-(morpholine-4-carbonyl)phenyl)-1H-benzo[d]imidazol-1-yl)benzonitrile



Step 1: To an ice-bath cooled solution of 1,3-dibromobenzene (8.2 mL, 34.7 mmol) in concentrated sulfuric acid (80 mL) was slowly added KNO_3 (6.8 g, 34.7 mmol) so as to maintain the internal reaction temperature below 10 °C. The reaction mixture was stirred for an additional 1 h and then poured into crushed ice (500 mL). The yellow precipitate isolated by filtration, washed with water and dried under reduced pressure to afford 2,4-dibromo-1-nitrobenzene (8.4 g, 90%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 7.92 (s, 1H), 7.75 (d, J = 8.4 Hz, 1H), 7.61 (d, J = 1.6 Hz, 1H); MS (ESI) m/z 280.9 $[\text{C}_6\text{H}_3\text{Br}_2\text{NO}_2]^+$.

Step 2: To a solution of 2,4-dibromonitrobenzene (8 g, 28.47 mmol) and 4-aminobenzonitrile (3.69 g, 28.79 mmol) in DMF (100 mL) was slowly added $t\text{-BuOK}$ (6.38 g, 56.94 mmol) while maintaining temperature between 0 °C to 5 °C. The reaction mixture was stirred at rt for 1 h then, was poured into ice-water and pH was adjusted to 6 with 5% aqueous HCl. The solid was collected, washed with water, and purified by recrystallization from EtOH to afford 4-(5-bromo-2-nitrophenylamino)benzonitrile (7.0 g, 77%) as a pale yellow solid. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 9.54 (s, 1H), 8.19 (d, J = 9.2, 1H), 7.71 (d, J = 8.4 Hz, 2H), 7.35 (d, J = 8.8 Hz, 2H), 6.91 (d, J = 2.0 Hz, 1H); MS (ESI) m/z 318 $[\text{C}_{13}\text{H}_8\text{BrN}_3\text{O}_2]^+$.

Step 3: To a solution of 4-(5-bromo-2-nitrophenylamino)benzonitrile (7 g, 22 mmol) in ethanol (70 mL) cooled at 0 °C was added $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ (14.9 g, 66.0 mmol). The reaction mixture was stirred at 80 °C for 4 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated aqueous solution of NaHCO_3 (100 mL) and extracted with EtOAc (3x100 mL). The organic layer was washed with water, brine, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to afford 4-(2-amino-5-bromophenylamino)benzonitrile (5.0 g, 79%, LC-MS 60%). The crude product was used in the next step without purification. MS (ESI) m/z 288 $[\text{C}_{13}\text{H}_{10}\text{BrN}_3]^+$.

Step 4: A solution of 4-(2-amino-5-bromophenylamino)benzonitrile (5 g, 17.36 mmol) in triethylorthoformate (25 mL) was stirred at 100 °C for 16 h. The reaction mixture was cooled to 0 °C and quenched by drop wise addition of saturated aqueous solution of NaHCO₃ (100 mL) then, extracted with EtOAc (3 x 100 mL). The organic layer was washed with water, brine, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (silica gel, eluent EtOAc/petroleum ether 30:70) to afford 4-(6-bromo-1H-benzo[d]imidazol-1-yl)benzonitrile (3.5 g, 68%, LC-MS 80%); ¹H NMR (400 MHz, CDCl₃) δ 8.11 (s, 1H), 7.92 (d, *J* = 4.0 Hz, 2H), 7.76 (d, *J* = 8.8 Hz, 1H), 7.71 (s, 1H), 7.66 (d, *J* = 8.0 Hz, 2H), 7.51 (d, *J* = 1.6 Hz, 1H); MS (ESI) *m/z* 298 [C₁₄H₈BrN₃]⁺

Step 5: A mixture of 4-(6-bromo-1H-benzo[d]imidazol-1-yl)benzonitrile (3.5 g, 11.74 mmol), 4-ethoxycarbonyl phenylboronic acid (2.5 g, 12.94 mmol), K₃PO₄ (4.97 g, 23.48 mmol), Pd(PPh₃)₄ (0.67 g, 0.587 mmol) in 1,4-dioxane (100 mL) and water (20 mL) was heated at 90 °C for 4 h under argon. The reaction mixture was diluted with EtOAc (2 x 100 mL) and washed with water (2x50 mL), brine solution (2x50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by column chromatography (silica gel, eluent EtOAc/petroleum ether 10:90 to 50:50) to afford ethyl 4-(1-(4-cyanophenyl)-1H-benzo[d]imidazol-6-yl)benzoate (2.8 g, 65%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.19 (s, 1H), 8.12 (d, *J* = 8.4 Hz, 2H), 7.99-7.92 (m, 3H), 7.75-7.65 (m, 6H), 4.41 (d, *J* = 7.2 Hz, 2H), 1.44 (t, *J* = 7.6 Hz, 3H); MS (ESI) *m/z* 368 [C₂₃H₁₇N₃O₂ + H]⁺.

Step 6: To a solution of 4-(3-(4-cyanophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)benzoate 5 (2.5 g, 6.8 mmol) in THF (20 mL) was added LiOH (571 mg, 13.6 mmol) in water (4 mL) and stirred for 16 h at rt. The reaction mixture was worked up and concentrated under reduced pressure to afford 4-(1-(4-cyanophenyl)-1H-benzo[d]imidazol-6-yl)benzoic acid (2 g, 86%, LC-MS 65%) as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm): 8.77 (s, 1H), 8.14 (d, *J* = 8.0 Hz, 2H), 8.10-7.95 (m, 5H), 7.95-7.83 (m, 3H), 7.72 (d, *J* = 8.4 Hz, 1H); MS (ESI) *m/z* 340.13 [C₂₁H₁₃N₃O₂ + H]⁺.

Step 7: To a solution of 4-(1-(4-cyanophenyl)-1H-benzo[d]imidazol-6-yl)benzoic acid 6 (300 mg, 0.88 mmol) in DMF (5 mL) was added NMM (0.178 mL, 1.76 mmol) followed by

HATU (501 mg, 1.32 mmol) at rt and the solution was stirred for 30 min. Morpholine (0.84 mL, 0.968 mmol) was added to the reaction mixture and stirring was continued for 16 h. The reaction mixture was diluted with EtOAc and washed with water and brine solution. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to obtain crude product. The crude product was purified by preparative HPLC to afford 4-(6-(4-(morpholine-4-carbonyl)phenyl)-1H-benzo[d]imidazol-1-yl)benzonitrile (100 mg, 27%, AUC HPLC 95.9%) as an off-white solid; m.p. 207-210 °C; ¹H NMR (400 MHz, CDCl₃) δ (ppm): 8.18 (s, 1H), 7.98-7.92 (m, 3H), 7.72 (d, *J* = 6.6 Hz, 3H), 7.66-7.61 (m, 3H), 7.51 (d, *J* = 7.9 Hz, 2H), 3.90-3.40 (m, 8H); MS (ESI) *m/z* 409.15 [C₂₅H₂₀N₄O₂ + H]⁺.

10

ANNEXURE

Enzymatic and Cellular IC₅₀ table:

Examples	IC ₅₀ (μM)		
	Mnk1	Mnk2	Hela
1	<0.1	<0.1	<1.0
2	<1.0	<1.0	<1.0
3	<1.0	<0.1	<1.0
4	<0.1	<0.1	<1.0
5	<0.1	<0.1	<1.0
6	<1.0	<1.0	<1.0
7	<1.0	<1.0	<1.0
9	<1.0	<1.0	<1.0
14	<1.0	<1.0	<1.0
15	<1.0	<1.0	<1.0
17	<1.0	<1.0	<1.0
21	<1.0	<1.0	<1.0
23	<1.0	<0.1	<1.0
24	<1.0	<1.0	<1.0
26	<1.0	<1.0	<1.0
27	<0.1	<0.1	<1.0
28	<0.1	<0.1	<1.0
29	<1.0	<1.0	<1.0
30	<1.0	<1.0	<1.0
31	<1.0	<1.0	<1.0
32	<0.1	<1.0	<1.0
38	<1.0	<1.0	<1.0

Examples	IC50 (μ M)		
	Mnk1	Mnk2	Hela
39	<1.0	<1.0	<1.0
41	<1.0	<1.0	<1.0
42	<1.0	<1.0	<1.0
43	<1.0	<1.0	<1.0
45	<1.0	<1.0	<1.0
46	<1.0	<1.0	<1.0
47	<1.0	<0.1	<1.0
48	<1.0	<1.0	<1.0
49	<1.0	<1.0	<1.0
54	<0.1	<0.1	<1.0
55	<0.1	<0.1	<1.0
56	<0.1	<0.1	<1.0
57	<0.1	<0.1	<1.0
58	<0.1	<0.1	<1.0
59	<0.1	<0.1	<1.0
60	<0.1	<0.1	<1.0
61	<0.1	<0.1	<1.0
62	<0.1	<0.1	<1.0
63	<0.1	<0.1	<1.0
64	<0.1	<0.1	<1.0
65	<0.1	<0.1	<1.0
66	<0.1	<0.1	<1.0

Examples	IC50 (μ M)		
	Mnk1	Mnk2	HeLa
68	<0.1	<0.1	<1.0
69	<0.1	<0.1	<1.0
70	<0.1	<0.1	<1.0
71	<0.1	<0.1	<1.0
72	<0.1	<0.1	<1.0
73	<0.1	<0.1	<1.0
74	<0.1	<0.1	<1.0
78	<0.1	<0.1	<1.0
79	<0.1	<0.1	<1.0
80	<0.1	<0.1	<1.0
81	<0.1	<0.1	<1.0
85	<0.1	<0.1	<1.0
86	<0.1	<0.1	<1.0
87	<0.1	<0.1	<1.0
89	<0.1	<0.1	<1.0
94	<0.1	<0.1	<1.0
95	<0.1	<0.1	<1.0
96	<0.1	<0.1	<1.0
97	<0.1	<0.1	<1.0
98	<0.1	<0.1	<1.0
99	<0.1	<0.1	<1.0
100	<0.1	<0.1	<1.0

Examples	IC50 (μ M)		
	Mnk1	Mnk2	Hela
101	<0.1	<0.1	<1.0
102	<0.1	<0.1	<1.0
103	<0.1	<0.1	<1.0
105	<0.1	<0.1	<1.0
106	<0.1	<0.1	<1.0
107	<0.1	<0.1	<1.0
109	<0.1	<0.1	<1.0
111	<0.1	<0.1	<1.0
112	<0.1	<0.1	<1.0
113	<0.1	<0.1	<1.0
115	<1.0	<1.0	<1.0
116	<1.0	<1.0	<1.0
117	<0.1	<0.1	<1.0
118	<0.1	<0.1	<1.0
119	<0.1	<0.1	<1.0
122	<1.0	<1.0	<1.0
123	<0.1	<0.1	<1.0
124	<0.1	<0.1	<1.0
125	<0.1	<0.1	<1.0
126	<0.1	<0.1	<1.0
129	<1.0	<0.1	<1.0
130	<0.1	<0.1	<1.0

Examples	IC50 (μ M)		
	Mnk1	Mnk2	Hela
133	<0.1	<0.1	<1.0
134	<0.1	<0.1	<1.0
135	<0.1	<0.1	<1.0
137	<1.0	<0.1	<1.0
138	<1.0	<1.0	<1.0
139	<0.1	<0.1	<1.0
140	<0.1	<0.1	<1.0
141	<0.1	<0.1	<1.0
142	<1.0	<0.1	<1.0
143	<0.1	<0.1	<1.0
144	<0.1	<0.1	<1.0
145	<0.1	<1.0	<1.0
146	<0.1	<0.1	<1.0
147	<0.1	<0.1	<1.0
149	<1.0	<1.0	<1.0
150	<0.1	<0.1	<1.0
151	<1.0	<0.1	<1.0
153	<1.0	<1.0	<1.0
156	<1.0	<1.0	<1.0
157	<1.0	<1.0	<1.0
158	<1.0	<0.1	<1.0
159	<0.1	<0.1	<1.0

Examples	IC50 (μ M)		
	Mnk1	Mnk2	Hela
160	<0.1	<0.1	<1.0
162	<1.0	<1.0	<1.0
163	<0.1	<0.1	<1.0
164	<0.1	<0.1	<1.0
165	<0.1	<0.1	<1.0
166	<0.1	<0.1	<1.0
167	<0.1	<0.1	<1.0
169	<0.1	<0.1	<1.0
170	<0.1	<0.1	<1.0
172	<0.1	<0.1	<1.0
173	<0.1	<0.1	<1.0
174	<0.1	<0.1	<1.0
175	<1.0	<0.1	<1.0
176	<0.1	<0.1	<1.0
179	<0.1	<0.1	<1.0
181	<0.1	<0.1	<1.0
183	<0.1	<0.1	<1.0
184	<0.1	<0.1	<1.0
185	<0.1	<0.1	<1.0
186	<0.1	<0.1	<1.0
187	<0.1	<0.1	<1.0
188	<0.1	<0.1	<1.0
190	<1.0	<1.0	<1.0

Reference List

- 5 (1) Buxade, M.; Parra-Palau, J. L.; Proud, C. G. *Front Biosci.* **2008**, *13*, 5359-5373.
- (2) Fukunaga, R.; Hunter, T. *EMBO J.* **1997**, *16*, 1921-1933.
- (3) Waskiewicz, A. J.; Flynn, A.; Proud, C. G.; Cooper, J. A. *EMBO J.* **1997**, *16*, 1909-1920.

- (4) Worch, J.; Tickenbrock, L.; Schwable, J.; Steffen, B.; Cauvet, T.; Mlody, B.; Buerger, H.; Koeffler, H. P.; Berdel, W. E.; Serve, H.; Muller-Tidow, C. *Oncogene* **2004**, *23*, 9162-9172.
- (5) Pellagatti, A.; Esoof, N.; Watkins, F.; Langford, C. F.; Vetrie, D.; Campbell, L. J.; Fidler, C.; Cavenagh, J. D.; Eagleton, H.; Gordon, P.; Woodcock, B.; Pushkaran, B.; Kwan, M.; Wainscoat, J. S.; Boultonwood, J. *Br.J.Haematol.* **2004**, *125*, 576-583.
- (6) Bredel, M.; Bredel, C.; Juric, D.; Harsh, G. R.; Vogel, H.; Recht, L. D.; Sikic, B. I. *Cancer Res.* **2005**, *65*, 4088-4096.
- (7) Hendrix, N. D.; Wu, R.; Kuick, R.; Schwartz, D. R.; Fearon, E. R.; Cho, K. R. *Cancer Res.* **2006**, *66*, 1354-1362.
- (8) Culjkovic, B.; Topisirovic, I.; Borden, K. L. *Cell Cycle* **2007**, *6*, 65-69.
- (9) Culjkovic, B.; Tan, K.; Orolicki, S.; Amri, A.; Meloche, S.; Borden, K. L. *J.Cell Biol.* **2008**, *181*, 51-63.
- (10) Nathan, C. O.; Carter, P.; Liu, L.; Li, B. D.; Abreo, F.; Tudor, A.; Zimmer, S. G.; De Benedetti, A. *Oncogene* **1997**, *15*, 1087-1094.
- (11) Bianchini, A.; Loiarro, M.; Bielli, P.; Busa, R.; Paronetto, M. P.; Loreni, F.; Geremia, R.; Sette, C. *Carcinogenesis* **2008**, *29*, 2279-2288.
- (12) Topisirovic, I.; Guzman, M. L.; McConnell, M. J.; Licht, J. D.; Culjkovic, B.; Neering, S. J.; Jordan, C. T.; Borden, K. L. *Mol.Cell Biol.* **2003**, *23*, 8992-9002.
- (13) Graff, J. R.; Zimmer, S. G. *Clin.Exp.Metastasis* **2003**, *20*, 265-273.
- (14) Ruggero, D.; Montanaro, L.; Ma, L.; Xu, W.; Londei, P.; Cordon-Cardo, C.; Pandolfi, P. P. *Nat.Med.* **2004**, *10*, 484-486.
- (15) Kevil, C.; Carter, P.; Hu, B.; DeBenedetti, A. *Oncogene* **1995**, *11*, 2339-2348.
- (16) Kevil, C. G.; De Benedetti, A.; Payne, D. K.; Coe, L. L.; Laroux, F. S.; Alexander, J. S. *Int.J.Cancer* **1996**, *65*, 785-790.
- (17) Scott, P. A.; Smith, K.; Poulsom, R.; De Benedetti, A.; Bicknell, R.; Harris, A. L. *Br.J.Cancer* **1998**, *77*, 2120-2128.
- (18) Rosenwald, I. B.; Lazaris-Karatzas, A.; Sonenberg, N.; Schmidt, E. V. *Mol.Cell Biol.* **1993**, *13*, 7358-7363.
- (19) Abid, M. R.; Li, Y.; Anthony, C.; De Benedetti, A. *J.Biol.Chem.* **1999**, *274*, 35991-35998.
- (20) De Benedetti, A.; Rhoads, R. E. *Proc.Natl.Acad.Sci.U.S.A* **1990**, *87*, 8212-8216.

- (21) Lazaris-Karatzas, A.; Montine, K. S.; Sonenberg, N. *Nature* **1990**, *345*, 544-547.
- (22) Rinker-Schaeffer, C. W.; Graff, J. R.; De Benedetti, A.; Zimmer, S. G.; Rhoads, R. E. *Int.J.Cancer* **1993**, *55*, 841-847.
- 5 (23) Rousseau, D.; Gingras, A. C.; Pause, A.; Sonenberg, N. *Oncogene* **1996**, *13*, 2415-2420.
- (24) Crew, J. P.; Fuggle, S.; Bicknell, R.; Cranston, D. W.; De Benedetti, A.; Harris, A. L. *Br.J.Cancer* **2000**, *82*, 161-166.
- (25) DeFatta, R. J.; Turbat-Herrera, E. A.; Li, B. D.; Anderson, W.; De Benedetti, A. *Int.J.Cancer* **1999**, *80*, 516-522.
- 10 (26) Kerekatte, V.; Smiley, K.; Hu, B.; Smith, A.; Gelder, F.; De Benedetti, A. *Int.J.Cancer* **1995**, *64*, 27-31.
- (27) Turbat-Herrera, E. A. T. A. B. D. L. S. d. A. B. M. a. M.-G. *Southwest Assoc.Clin.Microbiol* **1999**, *8*.
- 15 (28) Franklin, S.; Pho, T.; Abreo, F. W.; Nassar, R.; De Benedetti, A.; Stucker, F. J.; Nathan, C. O. *Arch.Otolaryngol.Head Neck Surg.* **1999**, *125*, 177-182.
- (29) Nathan, C. O.; Liu, L.; Li, B. D.; Abreo, F. W.; Nandy, I.; De Benedetti, A. *Oncogene* **1997**, *15*, 579-584.
- (30) Villar, H. O.; Yan, J.; Hansen, M. R. *Curr.Opin.Chem.Biol.* **2004**, *8*, 387-391.
- 20 (31) Williams, B. J.; Eastham, J. A.; Venables, D.; DeBenedetti, A.; Acree, D. T. *AACR Special Conference on Angiogenesis and Cancer* **1998**, B69-B70.
- (32) Raught, B.; Gingras, A. C. *Int.J.Biochem.Cell Biol.* **1999**, *31*, 43-57.
- (33) Topisirovic, I.; Ruiz-Gutierrez, M.; Borden, K. L. *Cancer Res.* **2004**, *64*, 8639-8642.
- 25 (34) Wendel, H. G.; Silva, R. L.; Malina, A.; Mills, J. R.; Zhu, H.; Ueda, T.; Watanabe-Fukunaga, R.; Fukunaga, R.; Teruya-Feldstein, J.; Pelletier, J.; Lowe, S. W. *Genes Dev.* **2007**, *21*, 3232-3237.
- (35) Ueda, T.; Watanabe-Fukunaga, R.; Fukuyama, H.; Nagata, S.; Fukunaga, R. *Mol.Cell Biol.* **2004**, *24*, 6539-6549.
- 30 (36) Scheper, G. C.; Proud, C. G. *Eur.J.Biochem.* **2002**, *269*, 5350-5359.
- (37) Sharon Lim, Tzuen Yih Saw, Sandy Chang, Min Zhang, Matthew Robert Janes, David A Fruman, David A.Rizzieri, Soo-Yong Tan, Charles Chuah, and S.Tiong Ong. Targeting of a Novel MNK-eIF4E-b-Catenin Axis in Blast Crisis Chronic Myelogenous Leukemia Inhibits Leukemia Stem-Cell Function. 2011: 53st ASH Annual

Meeting and Exposition. Oral Session: Chronic Myeloid Leukemia - Biology and Pathophysiology, excluding Therapy: Extrinsic and Intrinsic Survival Mechanisms. 12-10-2011. Ref Type: Conference Proceeding

- (38) Ly, C.; Arechiga, A. F.; Melo, J. V.; Walsh, C. M.; Ong, S. T. *Cancer Res.* **2003**, *63*, 5716-5722.
- (39) Prabhu, S.; Saadat, D.; Zhang, M.; Halbur, L.; Fruehauf, J. P.; Ong, S. T. *Oncogene* **2007**, *26*, 1188-1200.
- (40) Zhang, M.; Fu, W.; Prabhu, S.; Moore, J. C.; Ko, J.; Kim, J. W.; Druker, B. J.; Trapp, V.; Fruehauf, J.; Gram, H.; Fan, H. Y.; Ong, S. T. *Mol. Cell Biol.* **2008**, *28*, 6496-6509.
- (41) Hay, N. *Proc. Natl. Acad. Sci. U.S.A* **2010**, *107*, 13975-13976.
- (42) Marzec, M.; Liu, X.; Wysocka, M.; Rook, A. H.; Odum, N.; Wasik, M. A. *PLoS. One.* **2011**, *6*, e24849.
- (43) Jimenez-Rivera, C.; Avitzur, Y.; Fecteau, A. H.; Jones, N.; Grant, D.; Ng, V. L. *Pediatr. Transplant.* **2004**, *8*, 243-248.
- (44) Zaltzman, J. S.; Prasad, R.; Chun, K.; Jothy, S. *Nephrol. Dial. Transplant.* **2005**, *20*, 1748-1751.
- (45) Campistol, J. M.; Gutierrez-Dalmau, A.; Torregrosa, J. V. *Transplantation* **2004**, *77*, 760-762.
- (46) Stallone, G.; Schena, A.; Infante, B.; Di Paolo, S.; Loverre, A.; Maggio, G.; Ranieri, E.; Gesualdo, L.; Schena, F. P.; Grandaliano, G. *N. Engl. J. Med.* **2005**, *352*, 1317-1323.
- (47) Franz, D. N.; Leonard, J.; Tudor, C.; Chuck, G.; Care, M.; Sethuraman, G.; Dinopoulos, A.; Thomas, G.; Crone, K. R. *Ann. Neurol.* **2006**, *59*, 490-498.
- (48) Hess, G.; Smith, S. M.; Berkenblit, A.; Coiffier, B. *Semin. Oncol.* **2009**, *36 Suppl 3*, S37-S45.
- (49) Chan, H. Y.; Grossman, A. B.; Bukowski, R. M. *Adv. Ther.* **2010**, *27*, 495-511.
- (50) Hudes, G.; Carducci, M.; Tomczak, P.; Dutcher, J.; Figlin, R.; Kapoor, A.; Staroslawska, E.; Sosman, J.; McDermott, D.; Bodrogi, I.; Kovacevic, Z.; Lesovoy, V.; Schmidt-Wolf, I. G.; Barbarash, O.; Gokmen, E.; O'Toole, T.; Lustgarten, S.; Moore, L.; Motzer, R. J. *N. Engl. J. Med.* **2007**, *356*, 2271-2281.
- (51) Gibbons, J. J.; Abraham, R. T.; Yu, K. *Semin. Oncol.* **2009**, *36 Suppl 3*, S3-S17.

- (52) Beuvink, I.; Boulay, A.; Fumagalli, S.; Zilbermann, F.; Ruetz, S.; O'Reilly, T.; Natt, F.; Hall, J.; Lane, H. A.; Thomas, G. *Cell* **2005**, *120*, 747-759.
- (53) Teachey, D. T.; Sheen, C.; Hall, J.; Ryan, T.; Brown, V. I.; Fish, J.; Reid, G. S.; Seif, A. E.; Norris, R.; Chang, Y. J.; Carroll, M.; Grupp, S. A. *Blood* **2008**, *112*, 2020-2023.
- (54) Wang, X.; Yue, P.; Chan, C. B.; Ye, K.; Ueda, T.; Watanabe-Fukunaga, R.; Fukunaga, R.; Fu, H.; Khuri, F. R.; Sun, S. Y. *Mol. Cell Biol.* **2007**, *27*, 7405-7413.
- (55) HEMMING, Brian, GRZMIL, Michal, MORIN, Pier, and MERLO, Adrian. Treating Cancer by Modulating a Mnk. (WO2010055072A2). 2010. Ref Type: Patent
- (56) Hou, L.; Sasaki, H.; Stashenko, P. *Infect. Immun.* **2000**, *68*, 4681-4687.
- (57) Andersson, K.; Sundler, R. *Cytokine* **2006**, *33*, 52-57.
- (58) Rowlett, R. M.; Chrestensen, C. A.; Nyce, M.; Harp, M. G.; Pelo, J. W.; Cominelli, F.; Ernst, P. B.; Pizarro, T. T.; Sturgill, T. W.; Worthington, M. T. *Am.J.Physiol Gastrointest.Liver Physiol* **2008**, *294*, G452-G459.
- (59) Buxade, M.; Parra, J. L.; Rousseau, S.; Shpiro, N.; Marquez, R.; Morrice, N.; Bain, J.; Espel, E.; Proud, C. G. *Immunity*. **2005**, *23*, 177-189.
- (60) Guil, S.; Long, J. C.; Caceres, J. F. *Mol. Cell Biol.* **2006**, *26*, 5744-5758.
- (61) Buxade, M.; Morrice, N.; Krebs, D. L.; Proud, C. G. *J.Biol.Chem.* **2008**, *283*, 57-65.
- (62) Hefner, Y.; Borsch-Haubold, A. G.; Murakami, M.; Wilde, J. I.; Pasquet, S.; Schieltz, D.; Ghomashchi, F.; Yates, J. R., III; Armstrong, C. G.; Paterson, A.; Cohen, P.; Fukunaga, R.; Hunter, T.; Kudo, I.; Watson, S. P.; Gelb, M. H. *J.Biol.Chem.* **2000**, *275*, 37542-37551.
- (63) DaSilva, J.; Xu, L.; Kim, H. J.; Miller, W. T.; Bar-Sagi, D. *Mol. Cell Biol.* **2006**, *26*, 1898-1907.
- (64) Ushigome, M.; Ubagai, T.; Fukuda, H.; Tsuchiya, N.; Sugimura, T.; Takatsuka, J.; Nakagama, H. *Int.J.Oncol.* **2005**, *26*, 635-640.
- (65) Boukakis, G.; Patrinoou-Georgoula, M.; Lekarakou, M.; Valavanis, C.; Guialis, A. *BMC.Cancer* **2010**, *10*, 434.
- (66) Soni, A.; Akcakanat, A.; Singh, G.; Luyimbazi, D.; Zheng, Y.; Kim, D.; Gonzalez-Angulo, A.; Meric-Bernstam, F. *Mol. Cancer Ther.* **2008**, *7*, 1782-1788.
- (67) Berkel, H. J.; Turbat-Herrera, E. A.; Shi, R.; De Benedetti, A. *Cancer Epidemiol.Biomarkers Prev.* **2001**, *10*, 663-666.


(68) Wendel, H. G.; De Stanchina, E.; Fridman, J. S.; Malina, A.; Ray, S.; Kogan, S.; Cordon-Cardo, C.; Pelletier, J.; Lowe, S. W. *Nature* **2004**, *428*, 332-337.

(69) De Benedetti, A.; Graff, J. R. *Oncogene* **2004**, *23*, 3189-3199.

CLAIMS

What is claimed is:

1. A compound of Formula (Ia):



or a pharmaceutically acceptable form thereof, wherein

X_1 , X_2 , and X_3 are independently N or C;

X_4 and X_5 are independently N or CR⁴;

- wherein at least one of X_1 , X_2 , X_3 , X_4 , and X_5 is N;

X_7 is N or CR¹⁰;

X_8 is N or CR¹¹;

----- is a single or double bond, as valency allows;

R^1 , R^2 , R^3 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} , and R^{11} are independently hydrogen, halogen,

- optionally substituted aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, optionally substituted heterocyclyl, -CN, -OR, -SR, -N(R)₂, -NO₂, -C(O)R, -CO₂R, -C(O)N(R)₂, -C(O)NROR, -S(O)R, -SO₂R, -SO₂N(R)₂, -OC(O)R, -NRC(O)R, -NRSO₂R, or -NRC(O)N(R)₂;


- each R is independently hydrogen, optionally substituted C₁₋₆ aliphatic, optionally substituted aryl, optionally substituted carbocyclyl, optionally substituted heteroaryl, or optionally substituted heterocyclyl; or two R groups on the same nitrogen may be taken together to form an optionally substituted heterocycle;

each R⁴ is independently hydrogen, halogen, or optionally substituted C₁₋₆ aliphatic;

and


R^1 and R^2 , R^2 and R^3 , R^3 and R^4 , R^1 and R^{11} , R^{10} and R^{11} , R^5 and R^6 , R^6 and R^7 , R^7 and R^8 , or R^8 and R^9 may be optionally taken together to form an optionally substituted 5-6 membered carbocyclic, aryl, heterocyclic, or heteroaryl ring.

- 5 2. The compound of claim 1, wherein the compound is of Formula (Ia-i):



or a pharmaceutically acceptable form thereof.

3. The compound of claim 1, wherein the compound is of Formula (Ib-i):



or a pharmaceutically acceptable form thereof, wherein

X_1 , X_2 , and X_3 are independently N or C;

X_4 and X_5 are independently N or CR^4 ;

wherein at least one of X_1 , X_2 , X_3 , X_4 , and X_5 is N;

15 X_7 is N or CH;

X_8 is N or CH;


----- is a single or double bond, as valency allows;

R^1 is $-C(O)R$, wherein R is an optionally substituted heterocyclyl;

R^2 , R^3 , R^6 , and R^7 are independently $-H$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CF_3$, $-CHF_2$, $-C(O)NH_2$, $-OH$, $-OC_{1-4}alkyl$, or $-OCF_3$; or R^6 and R^7 are taken together to form a 5-6 membered fused heterocyclyl or heteroaryl; and
each R^4 is independently hydrogen or C_{1-6} alkyl.

5


4. The compound of claim 3, wherein the compound is of Formula (Ib-ii):



or a pharmaceutically acceptable form thereof.


10 5. The compound of any one of claims 1-4, wherein R^7 is $-CN$.

6. The compound of any one of claims 1-5, wherein R^1 is $-C(O)R$, wherein R is one of the following:




15


7. The compound of any one of claims 1, 2, 5, and 6, wherein the compound is of Formula (II), (III), (IV), (V), (VI), or (VII):




Formula II




Formula III




Formula IV



Formula V




Formula VI



Formula VII

8. The compound of claim 1, wherein the compound is selected from the group consisting of:



and pharmaceutically acceptable forms thereof.

9. A compound of any one of claims 1-8, wherein the compound is an MNK1 and/or MNK2 inhibitor.
- 5 10. A composition comprising a compound of any one of claims 1-9, or a pharmaceutically acceptable form thereof, and a pharmaceutically acceptable excipient.
11. The composition of claim 10, wherein the composition comprises an additional therapeutic agent.
- 10 12. The composition of claim 11, wherein the additional therapeutic agent is a kinase inhibitor.
13. The composition of claim 12, wherein the kinase inhibitor is a PI3K inhibitor.
- 15 14. The composition of claim 12, wherein the kinase inhibitor is an mTOR inhibitor.
15. A method of inhibiting MNK1 and/or MNK2 in a cell comprising administering to a cell a compound of any one of claims 1-9, or a pharmaceutically acceptable form thereof.
- 20 16. The method of claim 15, wherein the cell is *in vitro*.
17. The method of claim 15, wherein the cell is in a subject.
- 25 18. A method of treating a MNK1- or MNK2-related disorder comprising administering to a subject in need of treatment a compound of any one of claims 1-9, or a pharmaceutically acceptable form thereof, or a composition of any one of claims 10-14.
19. The method of claim 18, wherein the disorder is cancer (*e.g.*, blood and solid tumors), an inflammatory conditions, Alzheimer's disease, or a metabolic disorder (*e.g.*, obesity, diabetes).
- 30 20. The method of claim 18, wherein the compound is administered in combination with an additional therapeutic agent.

21. The method of claim 20, wherein the additional therapeutic agent is a kinase inhibitor.
22. The method of claim 21, wherein the kinase inhibitor is a PI3K inhibitor.
- 5 23. The method of claim 21, wherein the kinase inhibitor is an mTOR inhibitor.
24. The method of claim 19, wherein the disorder is cancer.
- 10 25. The method of any one of claims 18-24, wherein the compound or composition is administered at a dosage level sufficient to deliver from about 0.001 mg/kg to about 100 mg/kg to a subject in need thereof.
26. Use of a compound of any one of claims 1-9, or a pharmaceutically acceptable form
15 thereof, or a composition of any one of claims 10-14 for treatment of a MNK1- or MNK2-related disorder.
27. The use of claim 26, wherein the disorder is cancer (*e.g.*, blood and solid tumors), an inflammatory conditions, Alzheimer's disease, or a metabolic disorder (*e.g.*, obesity,
20 diabetes).
28. The use of claim 26, wherein the compound is used in combination with an additional therapeutic agent.
- 25 29. The use of claim 28, wherein the additional therapeutic agent is a kinase inhibitor.
30. The use of claim 29, wherein the kinase inhibitor is a PI3K inhibitor.
31. The use of claim 29, wherein the kinase inhibitor is an mTOR inhibitor.
- 30 32. The use of claim 26, wherein the disorder is cancer.

33. The use of any one of claims 26-32, wherein the compound or composition is administered at a dosage level sufficient to deliver from about 0.001 mg/kg to about 100 mg/kg to a subject in need thereof.