

PROCESS FOR MAKING AMINO DIARYL ETHERS AND AMINO DIARYL
ETHERS HYDROCHLORIDE SALTS

FIELD OF THE INVENTION

The present invention relates to processes useful in the preparation of an amino diaryl ether of formula (I), in particular 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)).

5

The present invention further relates to a hydrochloride salt of the compound of formula (I) in particular a monohydrochloride salt of the compound of formula (I) and to processes useful in the preparation of said monohydrochloride salt of the compound of formula (I) in particular 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)).

10 BACKGROUND OF THE INVENTION

Compounds of Formula (I) are modulators of the Notch signalling pathway, useful in the treatment and/or prevention of cancer, in particular useful in the treatment and/or prevention of Notch-dependent cancer and may be prepared using the process disclosed in WO2013/093885. The process disclosed in WO2013/093885 suffers i.a. from the disadvantages that (i) the compounds of formula (I) obtained by said process are contaminated with the catalyst used and (ii) side reactions take place, leading to further contamination of the compounds of formula (I) with side products. These disadvantages affect e.g. physical appearance and solubility of the compounds of formula (I) and onward manufacturing steps like milling. In conclusion, the process described in WO2013/093885 is not suitable for large-scale production of pharmaceutical grade compounds of formula (I). It would thus be beneficial to develop alternative or improved processes for the production of compounds of Formula (I) that do not suffer from some or all of these disadvantages.

15
20

SUMMARY OF THE INVENTION

It has now surprisingly been found that contamination of compounds of formula (I) can be omitted by making them via the process comprising catalytic hydrogenation using a palladium catalyst in a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol. The inventive process has been found to be reliable and applicable on an industrial scale and the compounds of formula (I) achieved by the inventive process consistently have an unprecedented high level of purity (99.9% pure as determined by HPLC, Pd-content <1 ppm as determined by elemental analysis).


25
30

The inventors have surprisingly found that, when the compounds of formula (I) are transformed to their hydrochloride salts, stable salts were obtained whereas all attempts to form other acidic addition salts failed. The inventors have further surprisingly found that, when the compounds of formula (I) are transformed to their monohydrochloride salts, the level of purity is further increased in that said monohydrochloride salts are typically obtained as colourless, crystalline solids, whereas the colour of the respective free base varies from batch to batch (range from colourless to reddish), independent of the analytical level of purity.

Compounds disclosed in WO2013/093885, like 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)), have a quite low melting point of e.g. 92.0 °C (Fig. 2) which constitutes a certain risk regarding physical stability of the compound during manufacture, milling, storage and processing to pharmaceutical formulations. However, it has surprisingly been found that the monohydrochloride salts of the compounds of formula (I), e.g. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride (formula (Ia)), have a significantly higher melting point than the free base (e.g. 173 °C, Fig. 1). Thus, monohydrochloride salts of the compounds of formula (I) surprisingly have improved properties with respect to the compounds of formula (I) in their free base form, such as higher melting points, improved solubilities in aqueous media and a discrete polymorph according to XRPD.

Taking these surprising findings into account, i.e. the high level of purity of the compounds of formula (I) achieved by the process of the invention and the favorable properties of the hydrochloride salts of the compound of formula (I), in particular of the monohydrochloride salts of the compounds of formula (I), the inventors herewith provide the present invention in its following aspects.

In a first aspect, the present invention provides a process for making a compound of formula (I)



Formula (I)

- 3 -


wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

5 wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl comprising:

reducing a compound of formula (II) with hydrogen in the presence of a palladium catalyst and a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol



Formula (II)

wherein X is selected from O and NH; and


wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹¹ and R²² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³³ and R⁴⁴ are each independently selected from H, halogen and C1-C10 alkyl;

to form said compound of formula (I).

20 In a second aspect, the present invention provides a process for making a hydrochloride salt of the compound of formula (I)



Formula (I)

wherein X is selected from O and NH; and


wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl

comprising reacting said compound of formula (I) with hydrochloric acid.

In a third aspect, the present invention provides a process for making a monohydrochloride salt of the compound of formula (I)



Formula (I)

wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl comprising reacting said compound of formula (I) with less than 1.5 molar equivalents of hydrochloric acid.

In a fourth aspect, the present invention provides a hydrochloride salt of the compound of formula (I) as defined herein.

In a fifth aspect, the present invention provides a monohydrochloride salt of the compound of formula (I) as defined herein.

In a sixth aspect, the present invention provides a hydrochloride salt of the compound of formula (I) as defined herein for use as a medicament.

In a seventh aspect, the present invention provides a monohydrochloride salt of the compound of formula (I) as defined herein for use as a medicament.

In an eighth aspect, the present invention provides a hydrochloride salt of the compound of formula (I) as defined herein for use in a method of treatment and/or prevention of cancer.

In a ninth aspect, the present invention provides a monohydrochloride salt of the compound of formula (I) as defined herein for use in a method of treatment and/or prevention of cancer.

BRIEF DESCRIPTION OF THE FIGURES

In the following the present invention is described in detail with reference to accompanying figures in which:

5 Figure 1 shows a differential scanning calorimetry thermogram (DSC) of the monohydrochloride salt of the compound of formula (Ia), i.e. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride. The DSC thermogram indicates that 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride has a melting point with an onset of 170.9 °C and peaking at 173.4 °C.

10

Figure 2 shows a DSC thermogram of the compound of formula (Ia), i.e. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine. The DSC thermogram indicates a melting point of the compound of formula (Ia) with an onset of 90.8 °C and peaking at 92.0 °C.

15 Figure 3 shows an X-ray powder diffraction (XRPD) diffractogram of the monohydrochloride salt of the compound of formula (Ia), i.e. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride.

Figure 4 shows an HPLC chromatogram of the compound of formula (Ia), i.e. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (purity: 99.9 %), obtained by the process of the invention (see
20 Example 3A).

Figure 5 shows a reference HPLC chromatogram of the compound of formula (Ia), i.e. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine, obtained by the process described in WO2013/093885
25 (Example 2).

Figure 6 shows a DSC thermogram of a substance that was obtained when reacting 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)) with more than 1.5 molar equivalents of hydrochloric acid (Example 5). The DSC thermogram indicates that the substance has no
30 sharp melting point and suggests that at least two species are present.

DETAILED DESCRIPTION OF THE INVENTION

Hereinafter, the present invention is described in further detail and is exemplified.

Definitions

The term "about" refers to $\pm 10\%$ of a given measurement.

The term "polar, aprotic solvent" refers to solvents which do not contain acidic hydrogen (Morrison and Boyd, Organic Chemistry 3rd. Edition, 31 (1974)) and which have a polarity according to the $E_T(30)$ -Scale of between about 36 kcal/mol and about 49 kcal/mol (C. Reichardt, *Chem. Rev.* **1994**, 94, 2319 – 2358; C. Reichardt, G. Schäfer, *Liebigs Ann.* **1995**, 1579 – 1582; R. Eberhardt, S. Löbbecke, B. Neidhart, C. Reichardt, *Liebigs Ann. /Recueil* **1997**, 1195 – 1199; C. Reichardt, *Green Chem.* **2005**, 7, 339 – 351; V. G. Machado, R. I. Stock, C. Reichardt, *Chem. Rev.* **2014**, 114, 10429-10475.). The proton donating or proton accepting interaction is usually greatest when the atom attached to the proton is nitrogen or oxygen. This behavior has been attributed to hydrogen bonding. In general, the hydrogen bond strength increases with increasing acidity of the proton-donating group, and increasing basicity of the proton-accepting group. Polar, aprotic solvents suitable for use in this invention will be those solvents that do not contain acidic or basic functional groups and do not degrade into acids or bases, including, but not limited to, ketones, nitriles, substituted aromatics such as halogenated aromatics, amides, sulfoxides, alkyl carbonates, chlorinated aliphatics, aromatic aldehydes, sulfones, esters, and the like, or mixtures thereof. The preferred polar, aprotic solvents for use in this invention include, but are not limited to, acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform, trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl acetate, amyl acetate, dimethyl sulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate, methyl ethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone, and mixtures thereof. Various other solvents having the above polar and aprotic characteristics are known to those skilled in the art.

The term "C3-C10 alcohol" refers to a unsubstituted linear or branched alkanol group comprising 3-10 carbon atoms, preferably a unsubstituted linear or branched alkanol group comprising 3-5 carbon atoms e.g. 1-propanol, 2-propanol, 2-butanol, 2-Methyl-1-propanol, or 2-Methylpropan-2-ol, more preferably 2-butanol or 2-propanol, most preferably 2-propanol.

The term "non-polar solvent" refers to a solvent having a polarity according to the $E_T(30)$ -Scale of less than about 35 kcal/mol. The preferred non-polar solvents for use in this

invention include, but are not limited to, pentane, hexane, heptane, cyclohexane and toluene, in particular heptane. When the non-polar solvent is e.g. a C5-C8-alkane, such as pentane, hexane or heptane, said C5-C8-alkane may be isomerically pure (e.g. n-pentane, n-hexane or n-heptane) or a random mixture of constitutional isomers.

5

The term "protic solvent" refers to a solvent comprising a proton-donating group. Non-limiting examples of protic solvents are e.g. water.

10 The expression "Tout" refers to the outer temperature measured outside a reactor and the expression "Ti" refers to the internal temperature measured inside a reactor. When a reaction temperature is indicated without specifying "Tout" or "Ti", it is understood that the temperature is measured outside the reactor.

15 The term "alkyl" refers to an unsubstituted linear or branched alkyl group, preferably an unsubstituted linear or branched alkyl group comprising 1-10 carbon atoms ("C1-C10 alkyl"), more preferably an unsubstituted linear or branched alkyl group comprising 3-8 carbon atoms ("C3-C8 alkyl"). Non-limiting examples of alkyl groups are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl and 1,1-dimethylbutyl. A particularly preferred alkyl group is tert-butyl.

20

The term "halogen" refers to fluorine, chlorine, bromine or iodine.

25 The term "cycloalkyl" refers to unsubstituted monocyclic, bicyclic, tricyclic or tetracyclic hydrocarbon groups of 3 to 12 carbon atoms ("C3-C12 cycloalkyl"), preferably of 5 to 10 carbon atoms ("C5-C10 cycloalkyl"). Non-limiting examples of cycloalkyl groups are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and adamantyl. Particularly preferred cycloalkyl groups are cyclohexyl and adamantyl.

30 The term "activated charcoal" refers to a form of carbon processed to have small, low-volume pores that increase the surface area available for adsorption or chemical reactions. Activated charcoal has a surface area in excess of 500 m²/g. Typically, the surface area of activated charcoal is about 1000 m²/g, but it may be up to 3000 m²/g. A non-limiting example of activated charcoal is activated charcoal having an iodine absorption (0.05 mol I₂/l) of >70

mL/g and/or a methylene blue-absorption (0.15% sol.) of >12 mL/0.1 g or having an iodine absorption of 1200mg/g and/or a methylene blue-absorption of 255mg/g.

The term "base" refers to a chemical compound that deprotonates another compound when reacted with it. Suitable bases for use in accordance with this disclosure include but are not limited to, e.g., tertiary amines, organolithium compounds and basic alkaline metal salts and hydrides. Examples of tertiary amines include triethylamine, N-methylmorpholine and diisopropylethylamine. Examples of organolithium compounds include methyl lithium (CH_3Li , MeLi), n-butyllithium (n-BuLi), sec-butyllithium (sec-BuLi) and tert-butyllithium (tert-BuLi). Examples of basic alkaline metal hydrides and salts include, e.g., sodium hydride (NaH), potassium hydride (KH), sodium carbonate (Na_2CO_3), potassium carbonate (K_2CO_3), sodium bicarbonate (NaHCO_3), sodium and potassium alkoxides including, but not limited to, sodium and potassium t-butoxide, propoxide, i-propoxide, ethoxide, methoxide, and the like, sodium amide (NaNH_2), potassium amide (KNH_2), and the like. Preferred bases are MeLi , K_2CO_3 and Na_2CO_3 .

The term "hydrochloride salt" refers to the compound of formula (I) or (Ia), respectively, wherein said compound of formula (I) or (Ia) is protonated and the counterion is chloride (Cl⁻) and comprises mono- and/or dichloride salts, in particular a mixture of mono- and dichloride salts.

The term "monohydrochloride salt" refers to the compound of formula (I) or (Ia), respectively, wherein said compound of formula (I) or (Ia) is protonated once and the counterion is chloride (Cl^-). Thus, the stoichiometry of mono-protonated compound of formula (I) or (Ia) to chloride is 1:1.


The term "purity" is expressed in percent (%) and is calculated from an HPLC chromatogram recorded at 220 nm according to the following formula:

30 purity of compound of formula (I) = (A_i/A_x) $\times$ 100,

wherein A_i is the area of the peak corresponding to the compound of formula (I), e.g. the compound of formula (Ia) and A_x is the sum of the areas of all other peaks observed in the HPLC chromatogram.

Process for making compounds of formula (I)

In a first aspect, the present invention provides a process for making a compound of formula (I)



Formula (I)


wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl, comprising:

reducing a compound of formula (II) with hydrogen in the presence of a palladium catalyst and a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol



Formula (II)

wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹¹ and R²² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³³ and R⁴⁴ are each independently selected from H, halogen and C1-C10 alkyl;

to form said compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention wherein the palladium catalyst is palladium on activated charcoal.

In a preferred embodiment, there is provided a process according to the first aspect of the invention wherein said palladium catalyst is palladium on activated charcoal selected from palladium on activated charcoal comprising about 10% wt/wt palladium and palladium on activated charcoal comprising about 5% wt/wt palladium, in particular palladium on activated charcoal comprising about 5% wt/wt palladium.

In one embodiment, there is provided a process according to the first aspect of the invention wherein said reduction of the compound of formula (II) to afford the compound of formula (I) comprises:

- 10 (a) providing a solution of the compound of formula (II) in a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol; followed by
- (b) adding the palladium catalyst to the solution of step (a); and
- (c) contacting the mixture obtained in step (b) with hydrogen;

wherein said palladium catalyst added in step (b) is added as a dry powder or as a suspension in the solvent used in step (a), preferably as a suspension in the solvent used in step (a).

In a further embodiment, there is provided a process according to the first aspect of the invention wherein the amount of palladium relative to the compound of formula (II) is less than 7 mol%, preferably less than about 6 mol%, more preferably less than about 5 mol%, more preferably less than about 4 mol%, more preferably less than about 3 mol%, more preferably less than about 2 mol%, in particular about 1.25 mol%.

In a further embodiment, there is provided a process according to the first aspect of the invention wherein the amount of palladium relative to the compound of formula (II) is less than 7 mol% and equal or higher than 1 mol%, preferably less than about 6 mol% and equal or higher than 1 mol%, more preferably less than about 5 mol% and equal or higher than 1 mol%, more preferably less than about 4 mol% and equal or higher than 1 mol%, more preferably less than about 3 mol% and equal or higher than 1 mol%, more preferably less than about 2 mol% and equal or higher than 1 mol%, in particular about 1.25 mol%.

In a preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is selected from 2-propanol, acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform,

trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl acetate, amyl acetate, dimethylsulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate, methyl ethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone and mixtures thereof.

5

In a further preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is selected from 2-propanol, acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform, trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl
10 acetate, amyl acetate, dimethylsulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate and mixtures thereof.

In a more preferred embodiment, there is provided a process according to the first aspect of
15 the invention wherein the solvent is a polar, aprotic solvent.

In an even more preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is selected from acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform,
20 trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl acetate, amyl acetate, dimethylsulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate, methyl ethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone and mixtures thereof.

In a further even more preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is selected from acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform, trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl
25 acetate, amyl acetate, dimethylsulfoxide (DMSO), dimethylformamide (DMF),
30 dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate and mixtures thereof.

In a particular preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is ethyl acetate or 2-propanol.

In a more particular preferred embodiment, there is provided a process according to the first aspect of the invention wherein the solvent is ethyl acetate.

5 The inventors have surprisingly found that the process according to the invention allows for a shorter reaction time than the prior art process described in WO2013/093885. Accordingly, in one embodiment, there is provided a process according to the first aspect of the invention, wherein the reduction of said compound of formula (II) to said compound of formula (I) is complete after 1-3 h, preferably after 1.5-3 h, more preferably after 2-3 h, e.g. after 2.5 h, preferably when performed using 1.2 kg of the compound of formula (II).

10 In a further embodiment, there is provided a process according to the first aspect of the invention, wherein the reduction of said compound of formula (II) to said compound of formula (I) is complete after 0.5-1.5 h, preferably after 40 min, preferably when performed using 300 g of the compound of formula (II).

15 In a further embodiment, there is provided a process according to the first aspect of the invention, further comprising contacting the compound of formula (I) with activated charcoal.

In a further embodiment, there is provided a process according to the first aspect of the invention, further comprising crystallizing the compound of formula (I).

20 In a preferred embodiment, there is provided a process according to the first aspect of the invention, further comprising contacting the compound of formula (I) with activated charcoal; and crystallizing the compound of formula (I).


25 In a further embodiment, there is provided a process according to the first aspect of the invention, further comprising crystallizing the compound of formula (I), wherein said crystallizing comprises precipitating the compound of formula (I) in crystalline form from a solvent mixture comprising a solvent as described herein, preferably a polar, aprotic solvent used in the reduction of the compound of formula (II), and a non-polar solvent.

30 In a preferred embodiment, there is provided a process according to the first aspect of the invention, further comprising crystallizing the compound of formula (I), wherein said

- 13 -


crystallizing comprises precipitating the compound of formula (I) in crystalline form from a solvent mixture comprising ethyl acetate and heptane.

- In a further embodiment, there is provided a process according to the first aspect of the invention, wherein the starting material of formula (II) is obtained by a process comprising reacting a compound of formula (III)



Formula (III)

- wherein X¹ is selected from OH and NH₂; and
 10 wherein R¹¹ and R²² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl
 with a compound of formula (IV)



Formula (IV)

- 15 wherein LG is a leaving group; and
 wherein Y¹, Y² and Y³ are each independently selected from N and CH; and
 wherein R³³ and R⁴⁴ are each independently selected from H, halogen and C1-C10 alkyl;
 in the presence of a base to form said starting material of formula (II) wherein X is O or NH,
 20 wherein R¹¹ and R²² are as defined for the compound of formula (III); and wherein R³³, R⁴⁴, Y¹, Y² and Y³ are as defined for the compound of formula (IV); and optionally further comprising crystallizing said starting material of formula (II).

- In one embodiment the leaving group LG is selected from the group consisting of F, Cl, Br, mesylate, tosylate and triflate. Preferably, the leaving group LG is selected from the group consisting of F, Cl, and Br.


In a preferred embodiment, there is provided a process according to the first aspect of the

invention, wherein the starting material of formula (II) is obtained by a process comprising reacting a compound of formula (III) as defined above, with a compound of formula (IV) as defined above in the presence of a base in a polar, aprotic solvent, preferably in a polar, aprotic solvent selected from dimethyl sulfoxide (DMSO), dimethyl formamide (DMF) or a mixture thereof, most preferably in DMSO, to form said starting material of formula (II).

In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein the starting material of formula (II) is obtained by a process comprising reacting a compound of formula (III) as defined above, with a compound of formula (IV) as defined above in the presence of a base to form said starting material of formula (II), and optionally further comprising crystallizing said starting material of formula (II), wherein said base is selected from the group consisting of potassium carbonate (K_2CO_3), sodium carbonate (Na_2CO_3), cesium carbonate (Cs_2CO_3), NEt_3 , Hünig's base and 1,8-Diazabicyclo[5.4.0]undec-7-en (DBU), preferably selected from the group consisting of potassium carbonate (K_2CO_3), sodium carbonate (Na_2CO_3) and cesium carbonate (Cs_2CO_3). More preferably said base is potassium carbonate (K_2CO_3).

In a further preferred embodiment, there is provided a process according to the first aspect of the invention, wherein the starting material of formula (II) is obtained by a process comprising reacting a compound of formula (III) as defined above, with a compound of formula (IV) as defined above in the presence of a base in a polar, aprotic solvent, preferably in a polar, aprotic solvent selected from dimethyl sulfoxide (DMSO), dimethyl formamide (DMF) or a mixture thereof, most preferably in DMSO, to form said starting material of formula (II); and optionally further comprising crystallizing said starting material of formula (II), wherein said crystallizing comprises precipitating the starting material of formula (II) in crystalline form from a solvent mixture comprising a polar, aprotic solvent and a non-polar solvent, preferably from a solvent mixture comprising ethyl acetate and heptane.


In a particularly preferred embodiment, there is provided a process according to the first aspect of the invention, wherein said compound of formula (I) is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia));



- 15 -

Formula (Ia)


and wherein said compound of formula (II) is 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa));



5


Formula (IIa)

and optionally wherein the starting material of formula (IIa) is obtained by a process comprising reacting 4-(*tert*-butyl)-phenol (formula (IIIa))



Formula (IIIa)

10 with 2-chloro-5-nitro-pyridine (formula (IVa))




Formula (IVa)

in the presence of a base to form 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)), and optionally further comprising crystallizing said 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)).

15

Thus the present invention provides a process for making a compound of formula (Ia)




Formula (Ia)

comprising:

20 reducing a compound of formula (IIa) with hydrogen in the presence of a palladium catalyst and a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol

- 16 -



Formula (IIa)

to form said compound of formula (Ia).

- 5 In a further particularly preferred embodiment, there is provided a process according to the first aspect of the invention,

wherein said compound of formula (I) is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)); and

- 10 wherein the starting material of formula (II) is 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)); and

wherein the starting material of formula (IIa) is obtained by a process comprising reacting 4-(*tert*-butyl)-phenol (formula (IIIa)) with 2-chloro-5-nitro-pyridine (formula (IVa)) in the presence of a base to form 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)); and

- 15 further comprising crystallizing said 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (formula (IIa)), wherein 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) is precipitated in crystalline form from a solvent mixture comprising ethyl acetate and heptane.

- In a further particularly preferred embodiment of the first aspect of the invention, the present invention provides a process for making 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)), comprising reducing 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) with hydrogen in the presence of palladium on activated charcoal and ethyl acetate to form 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)).

- 25 In a further particularly preferred embodiment of the first aspect of the invention, the present invention provides a process for making 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)) comprising reducing 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) with hydrogen in the presence of palladium on activated charcoal and ethyl acetate to form 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)); wherein the amount of palladium relative to the compound of formula (IIa) is less than 7 mol%, preferably less than about 6 mol%, more preferably less than about 5 mol%, more preferably less than about 4 mol%, more preferably less than about 3 mol%, more preferably less than about 2 mol%, in particular about 1.25 mol%.

In a further particularly preferred embodiment of the first aspect of the invention, the present invention provides a process for making 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)) comprising reducing 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) with hydrogen in the presence of palladium on activated charcoal and ethyl acetate to form 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)); wherein the amount of palladium relative to the compound of formula (IIa) is less than 7 mol% and equal or higher than 1 mol%, preferably less than about 6 mol% and equal or higher than 1 mol%, more preferably less than about 5 mol% and equal or higher than 1 mol%, more preferably less than about 4 mol% and equal or higher than 1 mol%, more preferably less than about 3 mol% and equal or higher than 1 mol%, more preferably less than about 2 mol% and equal or higher than 1 mol%, in particular about 1.25 mol%.

In a further particularly preferred embodiment of the first aspect of the invention, the present invention provides a process for making 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)) comprising reducing 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) with hydrogen in the presence of palladium on activated charcoal and ethyl acetate to form 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)); wherein said reduction of 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) with hydrogen comprises:

- (a) providing a solution of 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)) in ethyl acetate; followed by
 - (b) adding palladium on charcoal to the solution of step (a); and
 - (c) contacting the suspension obtained in step (b) with hydrogen;
- wherein the palladium on activated charcoal added in step (b) is added as a dry powder or as a suspension in ethyl acetate, preferably as a suspension in ethyl acetate.

Compounds of formula (I)

Preferably the group NH₂ is in position 4 (para) with respect to group X of the ring having groups Y¹, Y² and Y³. Preferably the groups R¹ or R² are in position 3 (meta) or 4 (para) with respect to group X of the ring without groups Y¹, Y² and Y³, more preferably R¹ is in position 4 (para) with respect to group X of the ring without groups Y¹, Y² and Y³, whereas R² is H, even more preferably R¹ is H, whereas R² is in position 4 (para) with respect to group X of the ring without groups Y¹, Y² and Y³. Preferably the group R³ or R⁴ are in position 2 (ortho) (provided Y² and/or Y¹ is C) with respect to group X of the ring having groups Y¹, Y² and Y³. More

preferably R^3 is in position 2 (ortho) (provided Y^2 and/or Y^1 is C) with respect to group X of the ring having groups Y^1 , Y^2 and Y^3 , whereas R^4 is H, even more preferably R^3 is H, whereas R^4 is in position 2 (ortho) (provided Y^2 and/or Y^1 is C) with respect to group X of the ring having groups Y^1 , Y^2 and Y^3 . Preferably the groups R^{11} or R^{22} are in position 3 (meta) or 4 (para) with respect to group X of the ring without groups Y^1 , Y^2 and Y^3 , more preferably R^{11} is in position 4 (para) with respect to group X of the ring without groups Y^1 , Y^2 and Y^3 , whereas R^{22} is H, even more preferably R^{11} is H, whereas R^{22} is in position 4 (para) with respect to group X of the ring without groups Y^1 , Y^2 and Y^3 . Preferably the group R^{33} or R^{43} are in position 2 (ortho) (provided Y^2 and/or Y^1 is C) with respect to group X of the ring having groups Y^1 , Y^2 and Y^3 , more preferably R^{33} is in position 2 (ortho) (provided Y^2 and/or Y^1 is C) with respect to group X of the ring having groups Y^1 , Y^2 and Y^3 , whereas R^{44} is H, even more preferably R^{33} is H, whereas R^{43} is in position 2 (ortho) (provided Y^2 and/or Y^1 is C) with respect to group X of the ring having groups Y^1 , Y^2 and Y^3 .

15 In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) X is O; and wherein for said compound of formula (II) X is O.

20 In a further embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 , Y^2 and Y^3 is CH; and

wherein for said compound of formula (II) Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 , Y^2 and Y^3 is CH.

25 In a further embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 and Y^3 is CH; and

30 wherein for said compound of formula (II) Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 and Y^3 is CH.

In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

Y^1 is selected from N and CH; and

Y^2 and Y^3 are each CH; and

wherein for said compound of formula (II)

Y^1 is selected from N and CH; and Y^2 and Y^3 are each CH.

In a more preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

5 Y^1 is N; and

Y^2 and Y^3 are each CH; and

wherein for said compound of formula (II)

Y^1 is N; and Y^2 and Y^3 are each CH.

10 In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^1 and R^2 are each independently selected from H and C1-C10 alkyl; and wherein for said compound of formula (II) R^{11} and R^{22} are each independently selected from H and C1-C10 alkyl.

In a further embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^1 and R^2 are each independently
15 selected from H and C3-C12 cycloalkyl; and wherein for said compound of formula (II) R^{11} and R^{22} are each independently selected from H and C3-C12 cycloalkyl.

In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is selected from H, C1-
20 C10 alkyl and C3-C12 cycloalkyl; and wherein for said compound of formula (II) R^{11} is H and R^{22} is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl.

In a further preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-
25 dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and wherein for said compound of formula (II) R^{11} is H and R^{22} is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl.

In a particularly preferred embodiment, there is provided a process according to the first
30 aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is tert-butyl; and wherein for said compound of formula (II) R^{11} is H and R^{22} is tert-butyl.

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are each independently selected from H,

and halogen; and wherein for said compound of formula (II) R^{33} and R^{44} are each independently selected from H, and halogen.

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are both halogen, preferably fluorine; and
5 wherein for said compound of formula (II) R^{33} and R^{44} are both halogen, preferably fluorine.

In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^3 is H and R^4 is selected from H and halogen, preferably from H and fluorine; and wherein for said compound of formula (II) R^{33} is H and R^{44} is selected from H, and halogen, preferably from H and fluorine.

10 In a particularly preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are both H; and wherein for said compound of formula (II) R^{33} and R^{44} are both H.

In one embodiment, there is provided a process according to the first aspect of the invention,
15 wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 , Y^2 and Y^3 is CH; and

20 R^1 and R^2 are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R^3 and R^4 are each independently selected from H and halogen; and

wherein for said compound of formula (II)

X, Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and

25 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

30 Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 and Y^3 is CH; and

R^1 and R^2 are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R^3 and R^4 are each independently selected from H and halogen; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
compound of formula (I).

5

In one embodiment, there is provided a process according to the first aspect of the invention,
wherein for said compound of formula (I)

X is selected from O and NH; and

Y¹ is selected from N and CH; and

10 Y² and Y³ are each CH; and

R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12
cycloalkyl; and

R³ and R⁴ are each independently selected from H and halogen; and

wherein for said compound of formula (II)

15 X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention,

20 wherein for said compound of formula (I)

X is selected from O and NH; and

Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

25 R³ and R⁴ are each independently selected from H and halogen; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
compound of formula (I).

30

In one embodiment, there is provided a process according to the first aspect of the invention,
wherein for said compound of formula (I)

X is selected from O and NH; and

Y¹ is selected from N and CH; and

Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R^3 is H and R^4 is selected from H and halogen; and

wherein for said compound of formula (II)

- 5 X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and
 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
 compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention,

- 10 wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 is selected from N and CH; and

Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H, C3-C8 alkyl and C5-C10 cycloalkyl; and

- 15 R^3 is H and R^4 is selected from H and halogen; and

 wherein for said compound of formula (II)

X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and

R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
 compound of formula (I).

20

In one embodiment, there is provided a process according to the first aspect of the invention,

 wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 is selected from N and CH; and

- 25 Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H and C3-C8 alkyl; and

R^3 is H and R^4 is selected from H and halogen; and

 wherein for said compound of formula (II)

X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and

- 30 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
 compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention,

 wherein for said compound of formula (I)

- 23 -

X is selected from O and NH; and

Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H and C5-C10 cycloalkyl; and

5 R³ is H and R⁴ is selected from H and halogen; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

10

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is O; and

Y¹ is selected from N and CH; and

15 Y² and Y³ are each CH; and

R¹ is H and R² is selected from H and C3-C8 alkyl; and

R³ is H and R⁴ is selected from H and halogen; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

20 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

25 X is O; and

Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H and C5-C10 cycloalkyl; and

R³ is H and R⁴ is selected from H and halogen; and

30 wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is NH; and

Y¹ is selected from N and CH; and

5 Y² and Y³ are each CH; and

R¹ is H and R² is selected from H and C3-C8 alkyl; and

R³ is H and R⁴ is selected from H and halogen; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

10 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

15 X is NH; and

Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H and C5-C10 cycloalkyl; and

R³ is H and R⁴ is selected from H and halogen; and

20 wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

25 In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

30 R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ and R⁴ are each independently selected from H and F; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

- 5 In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)
- X is selected from O and NH; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
- 10 R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
R³ and R⁴ are each independently selected from H and F; and
- wherein for said compound of formula (II)
- 15 X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

- In one embodiment, there is provided a process according to the first aspect of the invention,
- 20 wherein for said compound of formula (I)
- X is selected from O and NH; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
- 25 R³ is H and R⁴ is selected from H and F; and
- wherein for said compound of formula (II)
- X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
- 30 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

- 26 -

X is O; and

Y¹ is N; and

Y² and Y³ are each CH; and

5 R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

10 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

15 X is O; and

Y¹, Y² and Y³ are each CH; and

R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

20 R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

25

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is NH; and

Y¹ is N; and

30 Y² and Y³ are each CH; and

R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

5

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is NH; and

Y¹, Y² and Y³ are each CH; and

10 R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

15 X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention,

20 wherein for said compound of formula (I)

X is O; and

Y¹ is N; and

Y² and Y³ are each CH; and

25 R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

30 X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In a preferred embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

- 28 -

X is O; and

Y¹ is N; and

Y² and Y³ are each CH; and

5 R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is H; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

10 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

15 In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is O; and

Y¹, Y² and Y³ are each CH; and

20 R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and

25 R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention, wherein for said compound of formula (I)

X is NH; and

30 Y¹ is N; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R^3 is H and R^4 is selected from H and F; and
wherein for said compound of formula (II)

X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and
 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
5 compound of formula (I).

In one embodiment, there is provided a process according to the first aspect of the invention,
wherein for said compound of formula (I)

X is NH; and
10 Y^1 , Y^2 and Y^3 are each CH; and
 R^1 is H and R^2 is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl,
tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl,
cyclohexyl and adamantyl; and
 R^3 is H and R^4 is selected from H and F; and

15 wherein for said compound of formula (II)
 X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and
 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
compound of formula (I).

20 In a preferred embodiment, there is provided a process according to the first aspect of the
invention, wherein for said compound of formula (I)

X is selected from O and NH; and
 Y^1 , Y^2 and Y^3 are each selected from N and CH; and
 R^1 is H and R^2 is tert-butyl; and
25 R^3 and R^4 are both H; and

wherein for said compound of formula (II)

X , Y^1 , Y^2 and Y^3 are as defined for the compound of formula (I); and
 R^{11} is equal to R^1 , R^{22} is equal to R^2 , R^{33} is equal to R^3 and R^{44} is equal to R^4 of the
compound of formula (I).

30

In a preferred embodiment, there is provided a process according to the first aspect of the
invention, wherein for said compound of formula (I)

X is selected from O and NH; and
 Y^1 is selected from N and CH; and

Y² and Y³ are each CH; and
R¹ is H and R² is tert-butyl; and
R³ and R⁴ are both H; and

wherein for said compound of formula (II)

- 5 X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
compound of formula (I).

- 10 In a further preferred embodiment, there is provided a process according to the first aspect of
the invention, wherein for said compound of formula (I)

X is selected from O and NH; and
Y¹ is N; and
Y² and Y³ are each CH; and
R¹ is H and R² is tert-butyl; and
15 R³ and R⁴ are both H; and

wherein for said compound of formula (II)

X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
compound of formula (I).

20


In a further preferred embodiment, there is provided a process according to the first aspect of
the invention, wherein for said compound of formula (I)

X is selected from O and NH; and
Y¹, Y² and Y³ are each CH; and
25 R¹ is H and R² is tert-butyl; and
R³ and R⁴ are both H; and

wherein for said compound of formula (II)


X, Y¹, Y² and Y³ are as defined for the compound of formula (I); and
R¹¹ is equal to R¹, R²² is equal to R², R³³ is equal to R³ and R⁴⁴ is equal to R⁴ of the
30 compound of formula (I).

In a particularly preferred embodiment, there is provided a process according to the first
aspect of the invention, wherein said compound of formula (I) is selected from




Formula V

4-(4-(tert-pentyl)phenoxy)aniline,



Formula VI


4-(4-cyclohexylphenoxy)aniline,



Formula VII


6-(4-((3r,5r,7r)-adamantan-1-yl)phenoxy)pyridin-3-amine,

5



Formula VIII

6-(3-(tert-butyl)phenoxy)pyridin-3-amine,




Formula IX


4-(4-(tert-butyl)phenoxy)-3-fluoroaniline,

Formula X

- 32 -




6-(4-(tert-Pentyl)phenoxy)pyridin-3-amine,




Formula XI

6-(4-Butylphenoxy)pyridin-3-amine,




Formula XII

4-(4-Cyclohexylphenoxy)-3-fluoroaniline,



Formula XIII


3-Fluoro-4-(4-(tert-pentyl)phenoxy)aniline,




Formula XIV

6-(4-(2-Methylpentan-2-yl)phenoxy)pyridin-3-amine,

Formula XV




4-((3r,5r,7r)-Adamantan-1-yl)phenoxyaniline,




4-((3r,5r,7r)-Adamantan-1-yl)phenoxy-3-fluoroaniline,

Formula XVI




6-(4-cyclohexylphenoxy)pyridin-3-amine,

Formula XVII



4-(4-(*tert*-butyl)phenoxy)aniline,


Formula XVIII



4-(4-isopropylphenoxy)aniline, and

Formula XIX

Formula XX



6-(4-(2,4,4-trimethylpentan-2-yl)phenoxy)pyridin-3-amine;

and

wherein said compound of formula (II) is selected from the nitroaryl-analogues of the compounds of formulas (V)-(XX) as shown above.

5

In a further particularly preferred embodiment, there is provided a process according to the first aspect of the invention,

wherein said compound of formula (I) is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (formula (Ia)); and

10

wherein said compound of formula (II) is 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa)).

Hydrochloride and monohydrochloride salts of the compounds of formula (I)

The inventors have surprisingly found that, when the compounds of formula (I) are transformed to their hydrochloride salts, stable salts were obtained whereas all attempts to form other acidic addition salts failed. The inventors have further surprisingly found that, when the compounds of formula (I) are transformed to their monohydrochloride salts, the level of purity is further increased in that said monohydrochloride salts are typically obtained as colourless, crystalline solids (see e.g., Example 4), whereas the colour of the respective free base varies from batch to batch (range from colourless to reddish), independent of the analytical level of purity, which is consistently and reproducibly high (99.9% pure as determined by HPLC, Pd-content <1 ppm as determined by elemental analysis). When providing active pharmaceutical ingredients (API's) for the pharmaceutical industry, consistent properties of the API's, including their colour, physical appearance, purity, solubility and reproducibility of manufacture are of utmost importance. Accordingly, the high purity, colourless and crystalline monohydrochloride salts of the compounds of formula (I) are particularly useful as API's, which have been found highly consistent in repeated multi-kg large scale manufacture.

Further surprisingly, attempts at forming other acid addition salts or the respective dihydrochloride salts failed (see Example 5). For example, when more than 1.5 molar equivalents of hydrochloric acid were used to form a hydrochloride salt of the compound of formula (I), a mixture of mono- and dichloride salts with unsharp melting points was obtained (see Example 5 and Fig. 6).

Thus, in one embodiment, there is provided a process according to the first aspect of the invention, further comprising reacting the compound of formula (I) or (Ia) with hydrochloric acid, usually with equal or more than about 1.5 molar equivalents of hydrochloric acid, preferably with equal or more than about 1.5 molar equivalents and less than about 5 molar equivalents of hydrochloric acid, more preferably with equal or more than about 2 molar equivalents and less than about 3 molar equivalents of hydrochloric acid, even more preferably with about 2.5 molar equivalents of hydrochloric acid to form a hydrochloride salt of said compound of formula (I) or (Ia).

Thus, in one embodiment, there is provided a process according to the first aspect of the invention, further comprising reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia).

In a preferred embodiment, there is provided a process according to the first aspect of the invention, further comprising reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia); wherein said reacting the compound of formula (I) or (Ia) with hydrochloric acid is performed in a protic solvent, preferably in 2-propanol.

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia), followed by crystallizing said monohydrochloride salt.

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia), followed by crystallizing said monohydrochloride salt, wherein said crystallizing comprises precipitating said monohydrochloride salt in crystalline form from a solvent mixture comprising a protic solvent and a non-polar solvent, most preferably from a solvent mixture comprising 2-propanol and heptane.

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising crystallizing the compound of formula (I) or (Ia) and reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia).

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising contacting the compound of formula (I) or (Ia) with activated charcoal; crystallizing the compound of formula (I) or (Ia); and reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than


about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia).

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising contacting the compound of formula (I) or (Ia) with activated charcoal; crystallizing the compound of formula (I) or (Ia); and reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia); wherein said crystallizing comprises precipitating said compound of formula (I) or (Ia) in crystalline form from a solvent mixture comprising a protic solvent and a non-polar solvent, most preferably from a solvent mixture comprising 2-propanol and heptane.

In one embodiment, there is provided a process according to the first aspect of the invention, further comprising crystallizing the compound of formula (I) or (Ia) and reacting the compound of formula (I) or (Ia) with less than about 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about 1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of said compound of formula (I) or (Ia), followed by crystallizing said monohydrochloride salt.

In a second aspect, the present invention provides a process for making a hydrochloride salt of the compound of formula (I)

- 38 -



Formula (I)


wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

5 wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl comprising reacting the compound of formula (I) with hydrochloric acid, usually with equal or more than about 1.5 molar equivalents of hydrochloric acid, preferably with equal or more than about 1.5 molar equivalents and less than about 5 molar equivalents of hydrochloric acid, more preferably with equal or more than about 2 molar equivalents and less than about 3 molar equivalents of hydrochloric acid, even more preferably with about 2.5 molar equivalents of hydrochloric acid to form a hydrochloride salt of said compound of formula (I).

15 In a third aspect, the present invention provides a process for making a monohydrochloride salt of the compound of formula (I)



Formula (I)

wherein X is selected from O and NH; and

20 wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl comprising reacting said compound of formula (I) with less than 1.5 molar equivalents of hydrochloric acid, usually with less than about 1.5 and equal or more than about 1 molar equivalents of hydrochloric acid, preferably about 1.4 molar equivalents, more preferably about 1.3 molar equivalents, yet more preferably about 1.2 molar equivalents, even more preferably about 1.1 molar equivalents, most preferably about 1 molar equivalent, e.g., about

1.01 molar equivalents of hydrochloric acid, to form the monohydrochloride salt of the compound of formula (I).

5 In one embodiment, there is provided a process according to the third aspect of the invention, further comprising crystallizing said monohydrochloride salt of the compound of formula (I).

10 In a preferred embodiment, there is provided a process according to the third aspect of the invention, further comprising crystallizing said monohydrochloride salt of the compound of formula (I), wherein said crystallizing comprises precipitating said monohydrochloride salt in crystalline form from a solvent mixture comprising a protic solvent and a non-polar solvent, most preferably from a solvent mixture comprising 2-propanol and heptane.

15 In one embodiment, there is provided a process according to the third aspect of the invention, wherein the compound of formula (I) is crystallized prior to reacting it with hydrochloric acid.

20 In one embodiment, there is provided a process according to the third aspect of the invention, wherein the compound of formula (I) is contacted with activated charcoal and crystallized prior to reacting it with hydrochloric acid.

25 In one embodiment, there is provided a process according to the third aspect of the invention, wherein the compound of formula (I) is contacted with activated charcoal and crystallized by precipitating it in crystalline form from a solvent mixture comprising a protic solvent and a non-polar solvent, most preferably from a solvent mixture comprising 2-propanol and heptane, prior to reacting it with hydrochloric acid.

30 In one embodiment, there is provided a process according to the third aspect of the invention, further comprising crystallizing said monohydrochloride salt of the compound of formula (VI); wherein the compound of formula (I) is crystallized prior to reacting it with hydrochloric acid.

In one embodiment, there is provided a process according to the third aspect of the invention, further comprising crystallizing said monohydrochloride salt of the compound of


- 40 -

formula (I); wherein the compound of formula (I) is contacted with carbon on activated charcoal and crystallized prior to reacting it with hydrochloric acid.

In a preferred embodiment, there is provided a process according to the third aspect of the invention, wherein said reacting the compound of formula (I) with hydrochloric acid is performed in a protic solvent, preferably in 2-propanol.

It has further surprisingly been found that monohydrochloride salts of the compounds of formula (I) have improved properties, such as higher melting points and improved solubilities in aqueous media, as compared to the compounds of formula (I) in their free base form. For example, 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride has a melting point of 173 °C (measured by DSC; see Fig. 1), whereas the respective free base of formula (Ia) has a melting point of only 92 °C (measured by DSC; see Fig. 2). Surprisingly, all attempts at forming other acid addition salts failed (Example 5). In addition, attempts at forming the respective dihydrochloride salt of the compound of formula (Ia) surprisingly resulted in the formation of a substance (Example 5) having two melting points (Fig. 6), suggesting that at least two species were present.

In a fourth aspect, the present invention provides a hydrochloride salt of the compound of formula (I)



Formula (I)

wherein X is selected from O and NH; and


wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl.

In a fifth aspect, the present invention provides a monohydrochloride salt of the compound of formula (I)

- 41 -



Formula (I)

wherein X is selected from O and NH; and

wherein Y¹, Y² and Y³ are each independently selected from N and CH; and

- 5 wherein R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl and

wherein R³ and R⁴ are each independently selected from H, halogen and C1-C10 alkyl.

- 10 In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) X is selected from O and NH.

- 15 In a particularly preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) X is O.

- 20 In a further embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y¹, Y² and Y³ are each selected from N and CH, wherein at least one of Y¹, Y² and Y³ is CH.

- 25 In a further embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y¹, Y² and Y³ are each selected from N and CH, wherein at least one of Y¹ and Y³ is CH.

- In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y¹ is selected from N and CH; and Y² and Y³ are each CH.

- 30 In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the

fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y^1 is N; and Y^2 and Y^3 are each CH.

In a further preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) each of Y^1 , Y^2 and Y^3 are CH.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y^1 and Y^2 are both CH; and Y^3 is N.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) Y^1 and Y^2 are both N; and Y^3 is CH.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^1 and R^2 are each independently selected from H and C1-C10 alkyl.

In a further embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^1 and R^2 are each independently selected from H and C3-C12 cycloalkyl.

In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl.

In a further preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl.

In a particularly preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^1 is H and R^2 is tert-butyl.

5

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are each independently selected from H, and halogen.

10

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are both halogen, preferably fluorine.

15 In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^3 is H and R^4 is selected from H, and halogen, preferably from H and fluorine.

20 In a particularly preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I) R^3 and R^4 are both H.

25 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 , Y^2 and Y^3 are each selected from N and CH, wherein at least one of Y^1 , Y^2 and Y^3 is CH; and

30 R^1 and R^2 are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R^3 and R^4 are each independently selected from H and halogen.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

5 Y¹, Y² and Y³ are each selected from N and CH, wherein at least one of Y¹ and Y³ is CH; and

R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R³ and R⁴ are each independently selected from H and halogen.

10

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

15 Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ and R² are each independently selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R³ and R⁴ are each independently selected from H and halogen.

20

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

25 Y¹ is selected from N and CH; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R³ and R⁴ are each independently selected from H and halogen.

30

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y¹ is selected from N and CH; and

- 45 -

Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H, C1-C10 alkyl and C3-C12 cycloalkyl; and

R^3 is H and R^4 is selected from H and halogen.

5 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 is selected from N and CH; and

10 Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H, C3-C8 alkyl and C5-C10 cycloalkyl; and

R^3 is H and R^4 is selected from H and halogen.

15 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 is selected from N and CH; and

Y^2 and Y^3 are each CH; and

20 R^1 is H and R^2 is selected from H and C3-C8 alkyl; and

R^3 is H and R^4 is selected from H and halogen.

25 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is selected from O and NH; and

Y^1 is selected from N and CH; and

Y^2 and Y^3 are each CH; and

R^1 is H and R^2 is selected from H and C5-C10 cycloalkyl; and

30 R^3 is H and R^4 is selected from H and halogen.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 46 -

X is O; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H and C3-C8 alkyl; and
5 R³ is H and R⁴ is selected from H and halogen.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

10 X is O; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H and C5-C10 cycloalkyl; and
R³ is H and R⁴ is selected from H and halogen.

15 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is NH; and
20 Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H and C3-C8 alkyl; and
R³ is H and R⁴ is selected from H and halogen.

25 In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is NH; and
Y¹ is selected from N and CH; and
30 Y² and Y³ are each CH; and
R¹ is H and R² is selected from H and C5-C10 cycloalkyl; and
R³ is H and R⁴ is selected from H and halogen.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- X is selected from O and NH; and
5 Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
10 R¹² and R¹³ are each independently selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 15 X is selected from O and NH; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
20 R³ and R⁴ are each independently selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 25 X is selected from O and NH; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
30 R³ is H and R⁴ is selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- X is O; and
5 Y¹ is N; and
Y² and Y³ are each CH; and
R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
10 R³ is H and R⁴ is selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 15 X is O; and
Y¹, Y² and Y³ are each CH; and
R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
20 R³ is H and R⁴ is selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 25 X is NH; and
Y¹ is N; and
Y² and Y³ are each CH; and
R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and
30 R³ is H and R⁴ is selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is NH; and

5 Y¹, Y² and Y³ are each CH; and

R¹ and R² are each independently selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F.

10

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is O; and

15 Y¹ is N; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

20 R³ is H and R⁴ is selected from H and F.

In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

25 X is O; and

Y¹ is N; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

30

R³ is H and R⁴ is H.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is O; and

5 Y¹, Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

R³ is H and R⁴ is selected from H and F.

10

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

X is NH; and

15 Y¹ is N; and

Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

20 R³ is H and R⁴ is selected from H and F.

In one embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

25 X is NH; and

Y¹, Y² and Y³ are each CH; and

R¹ is H and R² is selected from H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, 1,1-dimethylpropyl, 1,1-dimethyl-3,3-dimethylbutyl, 1,1-dimethylbutyl, cyclohexyl and adamantyl; and

30 R³ is H and R⁴ is selected from H and F.

In a preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 5 X is selected from O and NH; and
Y¹, Y² and Y³ are each selected from N and CH; and
R¹ is H and R² is tert-butyl; and
R³ and R⁴ are both H.

- 10 In a further preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- X is selected from O and NH; and
Y¹ is selected from N and CH; and
Y² and Y³ are each CH; and
15 R¹ is H and R² is tert-butyl; and
R³ and R⁴ are both H.

- In a further preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)


- 20 X is selected from O and NH; and
Y¹ is N; and
Y² and Y³ are each CH; and
R¹ is H and R² is tert-butyl; and
25 R³ and R⁴ are both H.

In a further preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein for said compound of formula (I)

- 30 X is selected from O and NH; and
Y¹, Y² and Y³ are each CH; and
R¹ is H and R² is tert-butyl; and
R³ and R⁴ are both H.


In a particularly preferred embodiment, there is provided a process according to the second and third aspect of the invention or a hydrochloride salt and a monohydrochloride salt according to the fourth and fifth aspect of the invention, wherein said compound of formula (I) is selected from

5




Formula V

4-(4-(tert-pentyl)phenoxy)aniline,



Formula VI


4-(4-cyclohexylphenoxy)aniline,



Formula VII

6-(4-((3r,5r,7r)-adamantan-1-yl)phenoxy)pyridin-3-amine,


10




Formula VIII

6-(3-(tert-butyl)phenoxy)pyridin-3-amine,

Formula IX




4-(4-(tert-butyl)phenoxy)-3-fluoroaniline,




6-(4-(tert-Pentyl)phenoxy)pyridin-3-amine,

Formula X




6-(4-Butylphenoxy)pyridin-3-amine,

Formula XI



4-(4-Cyclohexylphenoxy)-3-fluoroaniline,


Formula XII



3-Fluoro-4-(4-(tert-pentyl)phenoxy)aniline,


Formula XIII

- 54 -




Formula XIV

6-(4-(2-Methylpentan-2-yl)phenoxy)pyridin-3-amine,




Formula XV

4-((3r,5r,7r)-Adamantan-1-yl)phenoxy)aniline,



Formula XVI


3-(4-((3r,5r,7r)-Adamantan-1-yl)phenoxy)-3-fluoroaniline,



Formula XVII


6-(4-cyclohexylphenoxy)pyridin-3-amine,

5




Formula XVIII

4-(4-(*tert*-butyl)phenoxy)aniline,



Formula XIX

4-(4-isopropylphenoxy)aniline, and



Formula XX

6-(4-(2,4,4-trimethylpentan-2-yl)phenoxy)pyridin-3-amine.

In a particularly preferred embodiment, there is provided a process according to the second aspect of the invention or a hydrochloride salt according to the fourth aspect of the invention, wherein said compound of formula (I) is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine, i.e. the hydrochloride salt obtained by the process according to the second aspect of the invention or
5 the hydrochloride salt according to the fourth aspect of the invention is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine hydrochloride.

In a further particularly preferred embodiment, there is provided a process according to the third aspect of the invention or a monohydrochloride salt according to the fifth aspect of the invention, wherein said compound of formula (I) is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine,
10 i.e. the monohydrochloride salt obtained by the process according to the third aspect of the invention or the monohydrochloride salt according to the fifth aspect of the invention is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride.

15 In a particularly preferred embodiment, there is provided a monohydrochloride salt according to the fifth aspect of the invention, wherein said monohydrochloride salt is 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride and has a melting point of about 168 °C to about 178 °C, preferably about 171 °C to about 175 °C, most preferably about 173 °C (measured using differential scanning calorimetry (DSC)).

In a further particularly preferred embodiment, there is provided a monohydrochloride salt according to the fifth aspect of the invention, wherein said monohydrochloride salt is 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride and has an X-ray powder diffraction pattern comprising 2 θ values as shown in Figure 3 measured using CuK α radiation.

In a preferred embodiment, there is provided a monohydrochloride salt according to the fifth aspect of the invention, wherein said monohydrochloride salt is 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride and has a purity of 100% ($\geq$ 95% area measured by HPLC-UV) after storage for 24 months at 25°C, 60% rh.

In a further preferred embodiment, there is provided a monohydrochloride salt according to the fifth aspect of the invention, wherein said monohydrochloride salt is 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride and has a purity of 98.9% ($\geq$ 95% area measured by HPLC-UV) after storage for 6 months at 40°C, 75% rh.

Use of the hydrochloride and the monohydrochloride salts of the invention

In a sixth aspect, the present invention provides a hydrochloride salt of the compound of formula (I) according to the fourth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine hydrochloride, for use as a medicament.

In a seventh aspect, the present invention provides a monohydrochloride salt of the compound of formula (I) according to the fifth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride, for use as a medicament.

In an eighth aspect, the present invention provides a hydrochloride salt of the compound of formula (I) according to the fourth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine hydrochloride, for use in a method of treatment and/or prevention of cancer.

Also provided is the use of a hydrochloride salt of the compound of formula (I) according to the fourth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine hydrochloride for the manufacture of a medicament for the treatment and/or prevention of cancer.

Also provided is the use of a hydrochloride salt of the compound of formula (I) according to the fourth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine hydrochloride for the treatment and/or prevention of cancer.

- 5 In a ninth aspect, the present invention provides a monohydrochloride salt of the compound of formula (I) according to the fifth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride, for use in a method of treatment and/or prevention of cancer.

- Also provided is the use of a monohydrochloride salt of the compound of formula (I)
10 according to the fifth aspect of the invention and its embodiments described above, preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride for the manufacture of a medicament for the treatment and/or prevention of cancer.

- Also provided is the use of a monohydrochloride salt of the compound of formula (I)
according to the fifth aspect of the invention and its embodiments described above,
15 preferably 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride for the treatment and/or prevention of cancer.

- In one embodiment of the fourth and fifth aspect of the invention, said cancer is a Notch dependent cancer, preferably a Notch dependent cancer selected from the group consisting
20 of T cell-Acute lymphoblastic leukemia (T-ALL), chronic myeloid leukemia (CML), chronic lymphocytic leukemia (CLL), Mantle cell lymphoma (MCL), breast cancer, pancreatic cancer, prostate cancer, melanoma, brain tumors, tumor angiogenesis, and colorectal cancer.

EXAMPLES

Hereinafter, the present invention is described in more details and specifically with reference to the examples, which however are not intended to limit the present invention.

5 **Materials:**

4-(*tert*-butyl)-phenol (formula (IIIa)) and 2-chloro-5-nitro-pyridine (formula (IVa)) used for the exemplified embodiments of this invention are commercially available from various sources, e.g. may be purchased from Sigma-Aldrich. Generally, building blocks of formula (III) and (IV) are known and commercially available.

10

Compounds of formula (II) may be obtained according to general procedure A.

Compounds of formula (I) may be obtained according to general procedure B.

The monohydrochloride salts of the compounds of formula (I) described herein may be obtained according to general procedure C.

15

Instrumentation:HPLC:

Device: Agilent 1100 series with DAD

Software: CDS Chromeleon 6.8

20 Column: Advanced Materials Technology Halo C18, 2.7 μ m, 100 mm x 4.6 mm

Column temperature: 20°C

Mobile Phase A: Water

Mobile Phase B: Acetonitrile

25 Gradient:

Time [min]	%A	%B
0	50	50
5	0	100
5	50	50
7.5	50	50

 (Equilibration)

30 Flow rate: 1.2 ml/min

Injection Volume: 2.0 μ L

Sample solvent: Acetonitrile for the compound of formula (Ia)

Acetonitrile / water 3:1 v/v for the monohydrochloride salt of the compound of formula (Ia)

Sample Concentration: 0.8 mg/mL

Autosampler temperature: 20°C

5 Detection: UV at 220 nm

XRPD:

The X-ray powder diffraction studies were performed using a Bruker D8 Advance using a Cu anode at 40kV, 40 mA. The software used for data collection was Diffrac. plus Part 11,
10 Version 3. No background correction or smoothing was applied to the patterns.

General Procedure A: Coupling

A compound of formula (III) (1 equiv) and a compound of formula (IV) (1 equiv) are dissolved in DMF or in DMSO (5 vol) at room temperature (rt), followed by addition of K₂CO₃ (about 1.5
15 equiv). The mixture is stirred at rt until one or preferably both of the compounds of formula (III) and (IV) is fully converted, whereafter ethyl acetate (8 vol) and water (10 vol) are added. The layers are separated, the organic layer is washed with aqueous NaHCO₃ (4 vol) and brine (4 vol) and dried over anhydrous Na₂SO₄ (0.5-1 % wt/wt). The suspension is filtered and the filter cake (Na₂SO₄) is washed with ethyl acetate (3 vol).

20 *Optional Crystallization:*

The combined filtrates are concentrated to about half their volume (i.e. ca. 6 vol ethyl acetate are distilled off). Subsequently, heptane (10 vol) is added at T_{out} = 50 °C and part of the solvent (about 10 vol) is distilled off. The resulting concentrated solution is cooled to T_{out} = -20 °C to afford a suspension. The suspension is filtered and the filter cake is washed with
25 cold heptane (4 vol) and dried on a rotary evaporator to afford the respective nitro-diaryl compound of formula (II).

General Procedure B: Hydrogenation

In an autoclave, a compound of formula (II) as obtainable according to general procedure A
30 is dissolved in ethyl acetate (5 vol). Then, a suspension of palladium on activated charcoal (preferably having a Pd-content of 5% wt/wt; preferably about 1.25 mol% Pd) in ethyl acetate (1 vol) is added to the solution at rt via a funnel and the funnel is rinsed with ethyl acetate (3 vol). The autoclave is purged with hydrogen, then the suspension is stirred under a hydrogen

pressure of 0.5 bar at rt until full conversion. Afterwards, the suspension is filtered and the filter cake is rinsed with ethyl acetate (3 vol).

Optional workup comprising crystallization:

The filtrate is concentrated under reduced pressure to about half its volume. Then, activated
5 charcoal is added at rt and the resulting suspension is stirred at rt. Subsequently, the suspension is filtered and the filter cake is washed with ethyl acetate (2×2 vol). The combined filtrates are concentrated under reduced pressure to about 1/3 of the initial volume and heptane (7 vol) is added at $T_{out} = 40\text{ }^{\circ}\text{C}$, before distilling off about half of the solvent. The concentrated solution is cooled to $T_{out} = -20\text{ }^{\circ}\text{C}$ to afford a suspension which is further
10 cooled to $T_{out} = -24\text{ }^{\circ}\text{C}$. Afterwards, the suspension is filtered and the filter cake is washed with cold heptane (3 vol) and dried on the filter overnight to afford the respective amino-diaryl compound of formula (I).

General Procedure C: Salt formation

15 A compound of formula (I) is dissolved in 2-propanol (4 vol). The solution is heated to $T_{out} = 40\text{ }^{\circ}\text{C}$ and 5-6 N HCl in 2-propanol (about 1.0 equiv) is added dropwise at $T_{out} = 40\text{ }^{\circ}\text{C}$ ($T_{i_{max}} = 44.9\text{ }^{\circ}\text{C}$). Optionally, seeds of the monohydrochloride salt of the compound of formula (I) (0.1% wt/wt) are then added. The (seeded) mixture is stirred for about 0.5 h at $T_i = 40\text{ }^{\circ}\text{C}$ to give a suspension. Heptane (4 vol) is added at $T_i = 40\text{ }^{\circ}\text{C}$ and the suspension is further
20 warmed to $T_i = 45\text{ }^{\circ}\text{C}$. A second portion of heptane (8 vol) is added. Then the mixture is warmed to $T_{out} = 55\text{ }^{\circ}\text{C}$ and kept stirring at this temperature for about 1 h.

Workup:

The suspension is cooled to $T_i = 13\text{ }^{\circ}\text{C}$ over a period of about 16 h, followed by further cooling to $T_i = 0\text{ }^{\circ}\text{C}$ over a period of about 30 min and stirring at $T_i = 0\text{ }^{\circ}\text{C}$ for 1 h.
25 Subsequently, the suspension is filtered and the filter cake is washed with a 10:1 mixture of heptane and 2-propanol (3 vol). The filter cake is dried on a rotary evaporator ($T_{out} = 50\text{ }^{\circ}\text{C}$, <10 mbar) for 3 d to afford the monohydrochloride salt of the compound of formula (I), typically as a colourless, crystalline solid.

30 **Example 1A: 2-(4-(tert-butyl)phenoxy)-5-nitropyridine (Formula (IIa))**

According to general procedure A: 4-(tert-butyl)-phenol (formula (IIIa), 3.98 kg) and 2-chloro-5-nitro-pyridine (formula (IVa), 4.13 kg) were dissolved in DMSO (21 L) at room temperature (rt), followed by addition of K_2CO_3 (5.41 kg) over 17 min at rt. The mixture was stirred at rt for 17.25 h, whereafter ethyl acetate (33 L) and water (20 L) were added. The resulting biphasic

mixture was stirred for 35 min and allowed to stand for another 35 min before the layers were separated. The organic layer was washed with aqueous NaHCO_3 (8% wt/wt, 16.5 L) and brine (16.5 L) and dried over anhydrous Na_2SO_4 (2.5 kg). The suspension was filtered and the filter cake (Na_2SO_4) was washed with ethyl acetate (13 L). Then, the combined filtrates
5 were concentrated under reduced pressure (25 L ethyl acetate were distilled off). Subsequently, heptane (isomeric mixture, 41 L) was added at $T_{\text{out}} = 50\text{ }^\circ\text{C}$ over 7 min and 42 L solvent were distilled off under reduced pressure. The resulting yellow solution was cooled to $T_{\text{out}} = -20\text{ }^\circ\text{C}$ over 20 h and stirred at that temperature for 65 min to afford a suspension. The suspension was filtered and the filter cake was washed with cold heptane
10 (isomeric mixture, 16.5 L) and dried on a rotary evaporator overnight to afford 2-(4-(tert-butyl)phenoxy)-5-nitropyridine (Formula (IIa), 6.2 kg, 87%) as a white solid.

Example 1B: 2-(4-(tert-butyl)phenoxy)-5-nitropyridine (Formula (IIa))

According to general procedure A: 4-(tert-butyl)-phenol (formula (III a), 7.94 kg) and 2-chloro-
15 5-nitro-pyridine (formula (Va), 8.36 kg) were dissolved in DMSO (42 L) at room temperature (rt), followed by addition of K_2CO_3 (10.96 kg) over 12 min at rt. The mixture was stirred at rt for 21.5 h, whereafter ethyl acetate (66 L) and water (42 L) were added. The resulting biphasic mixture was stirred for 68 min and allowed to stand for another 2 h before the layers were separated. The organic layer was washed with aqueous NaHCO_3 (8% wt/wt, 33 L) and
20 brine (33 L) and dried over anhydrous Na_2SO_4 (5 kg). The suspension was filtered and the filter cake (Na_2SO_4) was washed with ethyl acetate (25 L). Then, the combined filtrates were concentrated under reduced pressure (48 L ethyl acetate were distilled off). Subsequently, heptane (isomeric mixture, 84 L) was added at $T_{\text{out}} = 50\text{ }^\circ\text{C}$ over 67 min and 79 L solvent were distilled off under reduced pressure. The resulting yellow solution was cooled to $T_{\text{out}} =$
25 $-20\text{ }^\circ\text{C}$ over 20 h and stirred at that temperature for 61 min to afford a suspension. The suspension was filtered and the filter cake was washed with cold heptane (isomeric mixture, 33 L) and dried on a rotary evaporator overnight to afford 2-(4-(tert-butyl)phenoxy)-5-nitropyridine (Formula (IIa), in two portions 4.96 and 5.06 kg, 70%) as a white solid. However, material coated in the vessel, dissolved with 20 L ethyl acetate combined with
30 mother liquor was re-transferred into solvent switch from ethyl acetate to heptane according to general process leading to third portion of 3.59 kg (23%) as a white solid. Total overall yield increased to 93% in stage 1.

Example 2 (comparative example): 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia)) obtained by the process described in WO2013/093885

- 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa), 8.6 g) and palladium on activated charcoal (2.35 g, 10% wt/wt palladium) in MeOH (86 mL) were stirred under a H₂-atmosphere (0.5 bar) at room temperature for 2 h to afford 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia)). However, *N*-methylated by-product was observed by HPLC and a Pd content of 6.9 ppm was detected by elemental analysis.

Example 3A: 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia))

- 10 According to general procedure B: In an autoclave, 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa), 1.2 kg) was dissolved in ethyl acetate (6 L). Then, a suspension of palladium on activated charcoal (119 g, 5% wt/wt palladium) in ethyl acetate (1.2 L) was added to the solution at rt via a funnel and the funnel was rinsed with ethyl acetate (3.6 L). The autoclave was purged with hydrogen (5 bar), then the suspension was stirred (832 rpm) under a
15 hydrogen pressure of 0.5 bar at rt until full conversion (2 h 20 min). Afterwards, the suspension was filtered and the filter cake was rinsed with ethyl acetate (3.6 L).

Workup: In this campaign five shots of hydrogenation runs were combined and treated according to general process:

- The filtrates (from five runs) were combined concentrated under reduced pressure (74 L ethyl acetate were distilled off). Then, activated charcoal (605g) was added at Tout = 25 °C and the resulting suspension was stirred at Tout = 25 °C for 1 h 10 min. Subsequently, the suspension was filtered and the filter cake was washed with ethyl acetate (2x 12 L). The combined filtrates were concentrated under reduced pressure (42 L ethyl acetate were distilled off) and heptane (isomeric mixture, 42 L) was added at Tout = 40 °C over 12 min,
25 before distilling off 36 L of solvent under reduced pressure. The concentrated solution was cooled to Tout = -20 °C over 6 h to afford a suspension which was further stirred for 1 h at Tout = -24 °C. Afterwards, the suspension was filtered and the filter cake was washed with cold heptane (isomeric mixture, 19 L) and dried on the filter overnight to afford 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia), 4.95 kg, 91%) as a reddish solid.

30

Example 3B: 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia))

According to general procedure B: In an autoclave, 2-(4-(*tert*-butyl)phenoxy)-5-nitropyridine (Formula (IIa), 50 g) was dissolved in 2-propanol (0.4 L). Then, a suspension of palladium on

activated charcoal (5 g, 5% wt/wt palladium) in 2-propanol (25 mL) was added to the solution at rt via a funnel and the funnel was rinsed with 2-propanol (25 mL). The autoclave was purged with hydrogen (5 bar), then the suspension was stirred (832 rpm) under a hydrogen pressure of 0.1 bar at rt until full conversion (7 h 30 min). Afterwards, the suspension was filtered and the filter cake was rinsed with 2-propanol (0.2 L).

Workup: The filtrate was concentrated under reduced pressure (0.65 L 2-propanol were distilled off) and 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine was isolated as a solid.

Example 4: 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride

- According to general procedure C: 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine (Formula (Ia), 4.95 kg), were dissolved in 2-propanol (20 L). The solution was heated to $T_{out} = 40\text{ }^{\circ}\text{C}$ and 5-6 N HCl in 2-propanol (3.7 L, 1.01 equiv) was added dropwise over 20 min at $T_{out} = 40\text{ }^{\circ}\text{C}$ ($T_{i_{max}} = 44.9\text{ }^{\circ}\text{C}$). Seeds of 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride (5 g) were added and the seeded mixture was stirred for 0.5 h at $T_i = 40\text{ }^{\circ}\text{C}$ to give a suspension.
- Heptane (isomeric mixture, 20 L) was added over a period of 33 min at $T_i = 40\text{ }^{\circ}\text{C}$ and the suspension was further warmed to $T_i = 45\text{ }^{\circ}\text{C}$. A second portion of heptane was added (isomeric mixture, 40 L) was added dropwise over 54 min. Then the mixture was warmed to $T_{out} = 55\text{ }^{\circ}\text{C}$ and kept stirring at this temperature for 1 h.

Workup:

- The suspension was cooled to $T_i = 13\text{ }^{\circ}\text{C}$ over a period of 16 h, followed by further cooling to $T_i = 0\text{ }^{\circ}\text{C}$ over a period of 30 min and stirring at $T_i = 0\text{ }^{\circ}\text{C}$ for 1 h. Subsequently, the suspension was filtered and the filter cake was washed with a 10:1 mixture of heptane (isomeric mixture) and 2-propanol (2×16.5 L). The filter cake was dried on a rotary evaporator ($T_{out} = 50\text{ }^{\circ}\text{C}$, <10 mbar) for 3 d to afford the title compound as a colourless, crystalline solid (5.02 kg, 88%).

Example 5: Salt formation with 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine

Formation of different salts were tested according to the procedure indicated in Table 1 below:

Table 1: Salt formation with 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine

Acid	Eq. acid used	procedure	comments
HCl as 5-6N HCl in 2-PrOH	1 eq (later using 1.01 eq.)	According to general procedure C	Established process

- 64 -

HCl as 1.25M HCl in 2-PrOH	2.5 eq	According to 1 eq. process, however, product turned out to be mixed HCl	DSC revealed unspecified mixture of Mono and Di-HCl salt
maleic acid	1 eq.	4.13 g dissolved in 10 vol EtOH, at 50°C added 1 eq. maleic acid to give solution; cooled down to 0°C, however, still solution obtained	Reaction discarded
fumaric acid	1 eq.	4.13 g dissolved in 10 vol EtOH, at 50°C added 1 eq. fumaric acid to give solution; cooled down to 0°C, however, still solution obtained	Reaction discarded
benzoic acid	1 eq.	4.13 g dissolved in 8 vol EtOH suspended, at 50°C added 1 eq. benzoic acid to give solution; cooled down to 0°C, however, still solution obtained	Reaction discarded
tartaric acid	0.5 eq.	4.13 g dissolved in 8 vol EtOH suspended, at 50°C added 0.5 eq. tartaric acid to give very fine white solid; which did not give a thicker suspension, even at rt upon cooling	Reaction discarded
tartaric acid	1 eq.	4.13 g dissolved in 8 vol EtOH suspended, at 50°C added 1 eq. tartaric acid to give solution, which did not give a suspension even at rt upon cooling	Reaction discarded

Example 6: *Forced degradation of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride*

5

Instrumentation:

Light stability chamber: Solarbox 1500e

Oven: Salvis Vacuum drying oven VC-20

Heating unit: Ikamag RET-GS with IKA ETS SD and stainless steel heating block

Vials: Supelco V-Vials and screw caps with PTFE septa

Gas tight vials: Agilent 10 mL HS-GC clear glass vials and crimp caps with Not applicable PTFE septa

5 Quartz glass cuvette: Hellma precision cuvette Suprasil, Type 117.100-QS, 10 mm path length, screw capped with silicone septa

HPLC: As described above

10 Analytical Methods:

The HPLC test method described above was used to determine the assay and purity of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride before the forced degradation experiments and the area % content of individual impurities and degradation products in the 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride samples obtained from the

15 degradation experiments.

Identity of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride was determined by comparison of the retention time against a reference.

The assay and the purity of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride were determined using HPLC with UV detection at 220 nm utilizing the area normalization methodology for purity determination and external standard calibration for assay

20 determination. During all experiments the UV spectra from 200 nm to 950 nm have been recorded and also the peak purity of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride was assessed.

Only single determinations for each test condition had been performed.

25 For each test condition described below, one sample was prepared and analyzed in accordance with the HPLC method.

1. Solid state with thermal stress

Approximately 20.4 mg and 50.1 mg of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride were weighed into a Supelco V-Vial. The vial was closed with the screw cap and stored in the heating block of the heating unit operated at 100°C ± 5°C for 20 h (20

30 mg sample) and 68 h (50 mg sample), respectively.

After storage at 100°C for the intended time each sample was cooled down to room temperature and the content of the vial was dissolved in about 4 mL of the sample solvent

(acetonitrile / water 3:1 v/v). The obtained solution was quantitatively transferred into a 25 mL volumetric flask (20 mg sample), or to a 50 mL volumetric flask (50 mg sample), respectively. Then the vial was rinsed three times with about 4 mL sample solvent, whereupon each time the content of the vial was also transferred into the 25 mL volumetric flask. After the flask
5 was filled up to volume with the sample solvent and was shaken well, an aliquot of this solution was transferred into a HPLC vial for analysis.

Results: 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride is stable in solid state at 100°C. No significant changes were observed in the purity and the purity profile after 68 h storage at 100°C. The observed decrease in the assay was within the accepted variability of
10 the method ($\pm 2\%$).

2. Solid state stressed by UV/VIS radiation

Two samples of 20.23 mg and 20.17 mg of 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride were weighed each into a quartz glass cuvette. Both cuvettes were
15 closed with the screw cap. One of the cuvettes was protected by wrapping it in aluminium foil. This sample served as dark control. Both cuvettes were placed alongside each other in the light stability chamber and were then exposed in accordance to guideline [4] to UV/VIS radiation applying energy of 500 W/m² until a total illumination of 1200 kLux•h was reached. After exposure to the UV/VIS radiation for an illumination of 1200 kLux•h the sample was
20 brought to room temperature and the content of the cuvette was dissolved in about 2 mL of the sample solvent (acetonitrile / water 3:1 v/v). The obtained solution was quantitatively transferred into a 25 mL volumetric flask. Then the cuvette was rinsed five times with about 2 mL sample solvent, whereupon each time the content of the cuvette was also transferred into the 25 mL volumetric flask. After the flask was filled up to volume with the sample solvent
25 and was shaken well, an aliquot of this solution was transferred into a HPLC vial for analysis. Result: In solid state 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride is stable against UV/VIS radiation. After an illumination of 1200 kLux•h, applying an energy of 500 W/m², the purity decreased by 0.6 area % which was mainly related to the occurrence of two unknown impurities. The observed decrease in the assay of 2.1% w/w was not related to the
30 UV/VIS radiation, as the assay of the dark control had a similar value (decrease by 2.4% w/w).

3. Alkaline and thermal stress in aqueous solution

Approximately 20 mg (4×) of 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride were weighed into a 10 mL GC-HS vial and 2.0 mL of 0.1 M aqueous NaOH were added. The four vials were crimped and shaken. Two of the vials were stored in an oven operated at 60°C ± 5°C and two in an oven operated at 90°C ± 5°C. In addition, a blank sample was prepared by transferring a volume of 2.0 mL of 0.1 M aqueous NaOH into a 10 mL GC-HS vial. After crimping, the vial containing the blank was stored in the oven operated at 90°C ± 5°C. The test conditions are summarized in Table 2:

Table 2: Test conditions of alkaline and thermal stress in aqueous solution

Sample weight	Concentration	Test condition	Storage time
0	0 mg/mL	0.1 M NaOH / 90 °C	66 h
20.37 mg	10.2 mg/mL	0.1 M NaOH / 60 °C	18 h
20.63 mg	10.3 mg/mL	0.1 M NaOH / 60 °C	66 h
20.32 mg	10.2 mg/mL	0.1 M NaOH / 90 °C	18 h
20.78 mg	10.4 mg/mL	0.1 M NaOH / 90 °C	66 h

10

After storage at the defined conditions for the intended time each sample was cooled down to room temperature. Then, the samples were neutralized by adding 2.0 mL of an aqueous 0.1 M HCl solution. For each sample the vial content was quantitatively transferred into a 25 mL volumetric flask. Then the vial was rinsed three times with about 5 mL sample solvent (acetonitrile / water 3:1 v/v), whereupon each time the content of the vial was also transferred into the 25 mL volumetric flask. After the flask was filled up to volume with the sample solvent and was shaken well, an aliquot of this solution was transferred into a HPLC vial for analysis.

15

All recorded chromatograms were evaluated for identity, assay and purity according to the analytical method [1]. In addition, peak purity was assessed in the chromatograms of the stability samples. For each analytical sequence a SST was performed as specified in the analytical method [1]. All SST criteria were fulfilled for each of the analytical sequences.

20

Example 7: Solubility of 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine monohydrochloride vs. 6-(4-(*tert*-butyl)phenoxy)pyridin-3-amine

25

The solubilities in water of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine (formula (Ia)) and the monohydrochloride salt thereof, i.e. 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride, were assessed at various pH-values according to Tables 3 and 4. The solubility in water of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine (formula (Ia)) according to

5 Table 3 was assayed as follows: Free base 500mg were charged into 14 mL of acidic media at pH=2, 4, at neutral pH=7 and basic pH=10 to afford suspensions within calculated concentration of appr. 35mg/mL. The solubility in water of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride according to Table 4 was assayed as follows: HCl salt 100mg

10 each were charged into acidic media at pH=2 and 4 and neutral media pH=7 to furnish solutions. Most important observation was to that at a neutral solution at pH=7 drops down to pH=1 by charging HCl salt 100mg.

It was found that 6-(4-(tert-butyl)phenoxy)pyridin-3-amine (formula (Ia)) is very poorly soluble in water, irrespective of the pH. The solubility limit of the compound of formula (Ia) in 0.01% HCl was determined to be 100 mg in 190 mL (0.53 mg / mL). In contrast, the

15 monohydrochloride salt of the compound of formula (Ia) was found to be well-soluble in water.

Table 3: Solubility of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine at different pH-values

solvent	pH solvent	mass in mg	volume in mL	conc. in mg/mL	soluble	pH solution
15% acetic acid soln.	2	500	14ml	33,33	no	2
Fluka 33643 (pH=4)	4	500	14ml	33,33	no	4
Fluka 33646 (pH=7)	7	500	14ml	33,33	no	7
Fluka 33649 (pH=10)	10	500	14ml	33,33	no	10

20

Table 4: HCl salt of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine solubilities at different pH-values

solvent	pH solvent	mass in mg	volume in mL	conc. in mg/mL	soluble	pH solution
---------	------------	------------	--------------	----------------	---------	-------------

15% acetic acid soln.	2	100	0.6ml	166,66	yes	0
Fluka 33643 (pH=4)	4	100	1.4ml	71,43	yes	1
Fluka 33646 (pH=7)	7	100	1.0ml	100,00	yes	1

Example 8: Stability studies of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride:

- 5 For 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride two different types of stability studies were performed including short-term and long-term stability studies, respectively, at 25°C (RH, 60%), -5°C and -20°C for 36 months and an accelerated study at 40°C (RH, 75%) over 6 months. The results of these studies are shown in tables 7-14 below. In terms of analytical determinations six tests were involved including appearance, assay
- 10 free base mean and purity mean, water, IR and XRPD. Appearance was determined by visual inspection. The product should be a white or off-white powder. Water content was determined according to USP <921> coulometric method. Identification of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride by IR spectroscopy was done according to USP <197>, comparing the spectrum obtained to that of the reference standard. X-Ray
- 15 powder diffraction (XRPD) was performed according to USP <941> against the reference XRPD spectrum. Purity and assay of free base was done by HPLC-UV. Two methods were used. Mobile phases in the first method were water and acetonitrile. Mobile phases in the second method were water (0.5 mL of trifluoroacetic acid was added to 1000 mL of water) and acetonitrile (0.4 mL of trifluoroacetic acid was added to 1000 mL of acetonitrile).
- 20 Equipment and chromatographic conditions for identification and determination of purity and assay of the 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride by HPLC-UV according to the first method and the second method, respectively, are shown in Tables 5 and 6.
- 25 **Table 5:** Equipment and chromatographic conditions for identification and determination of purity and assay of 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride by HPLC-UV according to the first method

HPLC model	Agilent 1100/1200 series
------------	--------------------------

Detector	Diode Array Detector (DAD)
Column	Halo C18, 2.7 µm, 100 mm x 4.6 mm column
Gradient	Time : 0 min. 5.0 min. 10.0 min. 10.0 min. 12.5 min. Mobile phase A: 50% 0% 0% 50% 50% Mobile phase B: 50% 100% 100% 50% 50%
Flow rate	1.2 mL/min
Autosampler temperature	20°C
Column temperature	20°C
Injection volume	2 µl
Detection	220 nm / UV
Slit width	4 nm
Bandwidth	4 nm
Reference wavelength	Off
Peak width	0.10 min
Step	0.01 sec
Acquisition time	11 min
Run time	15 min

Table 6: Equipment and chromatographic conditions for identification and determination of purity and assay of the 6-(4-(tert-butyl)phenoxy)pyridin-3-amine monohydrochloride by HPLC-UV according to the second method

HPLC model	Agilent 1100/1200 series
Detector	Diode Array Detector (DAD)
Column	Halo C18, 2.7 µm, 100 mm x 4.6 mm column
Gradient	Time : 0 min. 1.0 min. 5.0 min. 8.0 min. 11.0 min. 11.1 min.

- 71 -

	15.0 min. Mobile phase A: 60% 60% 30% 10% 10% 60% 60% Mobile phase B: 40% 40% 70% 90% 90% 40% 40%
Flow rate	0.8 mL/min
Autosampler temperature	20°C
Column temperature	30°C
Injection volume	3 µl
Detection	220 nm / UV
Slit width	2 nm
Bandwidth	4 nm
Reference wavelength	Off
Peak width	0.05 min
Step	0.05 sec
Acquisition time	11 min
Run time	15 min

Table 7: 6 months stability studies with storage conditions: 25°C, 60% rh

		Release data	3 mths	6 mths
Acceptance Criteria		Result	Result	Result
White to off-white powder		White powder	White powder	White powder
80.9-92.9% m/m (anhydrous, solvent free)		87.0	88.8	87.2
>=	95% area	99.2	99.5	98.8
Report results [%m/m]		0.02	0.01	0.01
Conforms to reference		complies	complies	complies
Conforms to reference		according to diffractogram	not tested	not tested

Table 8: 36 months stability studies with storage conditions: 25°C, 60% rh

	Release data	3 mths	6 mths	9 mths	12 mths	18 mths	24 mths	36 mths
Acceptance Criteria	Result	Result	Result	Result	Result	Result	Result	Result
White to off-white powder	White powder	White powder	White powder	White powder	White powder	White powder	White powder	Pending
80.9-92.9% m/m (anhydrous, solvent free)	not tested	not tested	88.6	88.0	86.9	87.7	87.6	Pending
>= 95% area		not tested	100.0	100.0	100.0	100.0	100.0	Pending

Report results [%m/m]	0.02	0.01	0.01	0.02	0.02	0.02	0.02	Pending
Conforms to reference	Complies	Complies	Complies	Complies	Complies	Complies	Complies	Pending
Conforms to reference	according to diffractogram	not tested	not tested	not tested	Complies	not tested	according to diffractogram	Pending

Table 9: 6 months stability studies with storage conditions: 5°C

	Release data	3 mths	6 mths
Acceptance Criteria	Result	Result	Result
White to off-white powder	White powder	White powder	White powder
80.9-92.9% m/m (anhydrous, solvent free)	89.4	87.2	87.6
>= 95% area	100.0	99	99
Report results [%m/m]	0.01	0.02	0.02
Conforms to reference	complies	complies	complies
Conforms to reference	according to diffract-togram	not tested	not tested

Table 10: 36 months stability studies with storage conditions: 5°C

[illegible]

off-white powder	powder	powder	powder	powder	powder	powder	powder	ding
80.9-92.9% m/m (anhydrous, solvent free)	not tested	not tested	87.5	89.6	87.3	87.0	88.3	Pending
>= 95% area	not tested	not tested	100.0	100.0	100.0	100.0	100.0	Pending
Report results [%m/m]	0.01	0.02	0.02	0.03	0.02	0.01	0.02	Pending
Conforms to reference	complies	Complies	Complies	Complies	Complies	Complies	Complies	Pending
Conforms to reference	according to diffractogram	not tested	not tested	not tested	see diffractogram	not tested	according to diffractogram	Pending

Table 11: 6 months stability studies with storage conditions: -20°C

	Release data	3 mths	6 mths
Acceptance Criteria	Result	Result	Result
White to off-white powder	White powder	White powder	White powder
80.9-92.9% m/m (anhydrous, solvent free)	89.4	86.1	87.3
>= 95% area	100	99	99
Report results [%m/m]	0.01	0.02	0.02
Conforms to reference	complies	complies	complies
Conforms to reference	according to	not tested	not tested

	diffractogram		
--	---------------	--	--

Table 12: 36 months stability studies with storage conditions: -20°C

	Release data	3 mths	6 mths	9 mths	12 mths	18 mths	24 mths	36 mths
Acceptance Criteria	Result	Result	Result	Result	Result	Result	Result	Result
White to off-white powder	White powder	White powder	White powder	White powder	White powder	White powder	pending	Pending
80.9-92.9% m/m (anhydrous, solvent free)	not tested	not tested	87.4	88.6	87.5	87.4	87.4	Pending
>= 95% area	not tested	not tested	100.0	100.0	100.0	100.0	100.0	Pending
Report results [%m/m]	0.01	0.02	0.02	0.03	0.02	0.01	0.02	Pending
Conforms to reference	Complies	Complies	Complies	Complies	Complies	Complies	Complies	Pending
Conforms to reference	according to diffractogram	not tested	not tested	not tested	see diffractogram	not tested	according to diffractogram	Pending

Table 13: 6 months stability studies with storage conditions: 40°C, 75% rh

	Release data	3 mths	6 mths
Acceptance Criteria	Result	Result	Result
White to off-white powder	White powder	White powder	White powder
80.9-92.9% m/m (anhydrous, solvent free)	87.0	89.2	87.4
>= 95% area	99.2	99.6	98.9
Report results [%m/m]	0.02	0.01	0.02
Conforms to reference	complies	complies	complies
Conforms to reference	accor-ding to diffract-togram	not tested	complies

Table 14: Summary of stability studies performed


[illegible]

- 78 -

18 months	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C
24 months	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C
36 months (data pen- ding)	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C	25°C, 60% rh +5°C -20°C

CLAIMS


1. A crystalline form of a monohydrochloride salt of the compound of formula (Ia)



Formula (Ia)

- 5 wherein the crystalline form is characterized by having an X-ray powder diffractogram which has one or more peaks chosen from peaks ($^{\circ} 2\theta$) at $7.3^{\circ}\pm 0.2^{\circ}$, $11.1^{\circ}\pm 0.2^{\circ}$, $14.6^{\circ}\pm 0.2^{\circ}$, $18.7^{\circ}\pm 0.2^{\circ}$, $21.4^{\circ}\pm 0.2^{\circ}$, $23.3^{\circ}\pm 0.2^{\circ}$, $23.8^{\circ}\pm 0.2^{\circ}$, $25.9^{\circ}\pm 0.2^{\circ}$, $27.4^{\circ}\pm 0.2^{\circ}$, and $30.7^{\circ}\pm 0.2^{\circ}$.
- 10 2. Use of the monohydrochloride salt of claim 1 in the manufacture of a medicament for the treatment of cancer in a patient in need thereof, wherein the monohydrochloride salt is to be administered to a patient in need thereof.
3. The monohydrochloride salt of claim 1, wherein the salt is anhydrous.
4. The monohydrochloride salt of claim 1, characterized by having a melting point with an onset of 170.9°C and peaking at 173.4°C .
- 15 5. The monohydrochloride salt of claim 1, characterized by having a melting point of 173°C .
6. The monohydrochloride salt of claim 1, characterized by differential scanning calorimetry (DSC) pattern substantially as set forth in FIG. 1.
- 20 7. The monohydrochloride salt of claim 1, having a purity of 99.9% pure as determined by HPLC.
8. The monohydrochloride salt of claim 1, wherein the salt is in a crystalline form.


9. The monohydrochloride salt of claim 8, characterized by an X-ray powder diffraction (XRPD) pattern substantially as set forth in FIG. 3.
10. The monohydrochloride salt of claim 8, characterized by having an X-ray powder diffractogram which has one or more peaks chosen from peaks ($^{\circ} 2\theta$) as set forth in FIG. 3, wherein the one or more peaks have a Cps intensity greater than 100.
11. The monohydrochloride salt of claim 8, wherein the one or more peaks ($^{\circ} 2\theta$) have a Cps intensity greater than 100 are selected from about 7.3° , about 14.6° , about 21.4° , about 27.4° , about 25.9° , about 30.7° , about 22.0° , about 33.8° , about 23.3° , about 23.8° and about 28.7° .
12. A pharmaceutical formulation comprising the monohydrochloride salt of claim 1 and an excipient.
13. Use of the monohydrochloride salt of claim 1 or a pharmaceutical formulation thereof in the manufacture of a medicament for the treatment of a notch dependent cancer, wherein the monohydrochloride salt or pharmaceutical formulation is to be administered.
14. A monohydrochloride salt of claim 1 or a pharmaceutical formulation thereof for use as a medicament.
15. A process for making a crystalline form of monohydrochloride salt of the compound of formula (Ia) according to claim 1



Formula (Ia)

,
the process comprising:

reducing a compound of formula (IIa) with hydrogen in the presence of a palladium catalyst and a solvent wherein the solvent is a polar, aprotic solvent or a C3-C10 alcohol




Formula (IIa)

to form said compound of formula (Ia), further comprising reacting the compound of formula (Ia) with less than about 1.5 molar equivalents of hydrochloric acid to form the monohydrochloride salt of said compound of formula (Ia).


- 5
- 10 16. The process according to claim 15, wherein said palladium catalyst is palladium on activated charcoal.
17. The process according to claim 15, wherein said palladium catalyst is present in an amount of less than 7 mol%.
- 15 18. The process according claim 15, wherein the solvent is selected from 2-propanol, acetone, 2-butanone, 3-methyl-2-butanone, cyclohexanone, acetonitrile, chlorobenzene, methylene chloride, chloroform, trichloroethane, ethylene chloride, benzaldehyde, sulfolane, ethyl acetate, propyl acetate, amyl acetate, dimethylsulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide, diethylcarbonate, propylene carbonate, ethylene carbonate, methyl ethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone and mixtures thereof.
- 20 19. The process according to claim 15, wherein said solvent is ethyl acetate.
20. The process according to claim 15, further comprising contacting the compound of formula (Ia) with activated charcoal.
- 25 21. The process according to claim 15, further comprising crystallizing said compound of formula (Ia).

22. The process according to claim 15, further comprising crystallizing said monohydrochloride salt.
23. The process according to claim 15, wherein said compound of formula (II) is obtained by a process comprising reacting a compound of formula (IIIa)



Formula (IIIa)

with a compound of formula (IVa)



Formula (IVa)

wherein Cl is a leaving group; and
optionally
further comprising the step of crystallizing said starting material of formula (IIa).