

FIGURE 1

Serum uric acid (mg/dL) levels 0, 3, 7 and 14 days after administering 4-(2-(5-bromo-4-(4-cyclopropyl-naphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetamido)-3-chlorobenzoic acid, potassium salt in humans at doses of 300mg, 400mg or 500mg b.i.d.




FIGURE 2

Serum uric acid ($\mu\text{mol/L}$) levels 0, 3, 7 and 14 days after administering 4-(2-(5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetamido)-3-chlorobenzoic acid, potassium salt in humans at doses of 300mg, 400mg or 500mg b.i.d.




FIGURE 3

Change in serum uric acid (mg/dL) levels 3, 7 and 14 days after administering 4-(2-(5-bromo-4-(4-cyclopropyl-naphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetamido)-3-chlorobenzoic acid, potassium salt in humans at doses of 300mg, 400mg or 500mg b.i.d.




FIGURE 4

Change in serum uric acid ($\mu\text{mol/L}$) levels 3, 7 and 14 days after administering 4-(2-(5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetamido)-3-chlorobenzoic acid, potassium salt in humans at doses of 300mg, 400mg or 500mg b.i.d.




FIGURE 5

Change in serum uric acid ($\mu\text{mol/dL}$) levels by treatment day after administering 4-(2-(5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetamido)-3-chlorobenzoic acid, potassium salt in humans at doses of 300mg, 400mg or 500mg b.i.d.




FIGURE 6

Increase in Daily Uric Acid Output Following Oral Administration of 2-(5-bromo-4-(4-cyclopropylnaphthalen-1-yl)-4H-1,2,4-triazol-3-ylthio)acetic acid solution

