

**ISTAROXIME-CONTAINING INTRAVENOUS FORMULATION FOR THE TREATMENT  
OF ACUTE HEART FAILURE (AHF)**

**Cross Reference To Related Applications**

[1] This claims benefit of U.S. Provisional Application No. 62/814,149, filed March 5, 2019, the entire contents of which are incorporated by reference herein.

**Field of the Invention**

[2] The present invention relates to the field of pharmaceuticals, in particular to an istaroxime-containing intravenous formulation for use for the treatment of acute heart failure.

**Background of the invention**

[3] The prevalence of heart failure (HF) is age-dependent, ranging from less than 2% of people younger than 60 years to more than 10% of people older than 75 years (Metra M, Teerlink JR, Lancet 2017; 390:1981-1995). Most patients with HF have a history of hypertension, coronary artery disease, cardiomyopathies, or valve disease, or a combination of these disorders (Metra M, Teerlink JR, Lancet 2017; 390:1981-1995). The calculated lifetime risk of developing HF is expected to increase and those with hypertension are at higher risk (Lloyd-Jones DM et al., Circulation 2002;106:3068-3072). Patients with HF have a poor prognosis with high rates of hospital admission and mortality.

[4] Clinical symptoms in HF are caused by a cardiac double pathological feature that consists in an inotropic abnormality, resulting in diminished systolic emptying (systolic dysfunction) and a compliance abnormality in which the ability of the ventricles to suck blood from the venous system is impaired (diastolic dysfunction), thus reducing the amount of blood available for systolic contraction, which is an impairment of left ventricle (LV) filling. Whatever the initial triggering mechanism of HF, an abnormal distribution of intracellular  $\text{Ca}^{2+}$  resulting from reduced  $\text{Ca}^{2+}$  uptake by the sarcoplasmic reticulum (SR), which is the main intracellular  $\text{Ca}^{2+}$  store (Schwinger RH et al., J Mol Cell Cardiol. 1999;31(3):479-91; Bers D et al., Ann N.Y. Acad Sci 2006; 1080:165-177), underlies the impaired contractility and relaxation. This abnormal  $\text{Ca}^{2+}$  distribution involves the  $\text{Ca}^{2+}$ -ATPase of the SR membrane (SERCA2a), an ATP dependent  $\text{Ca}^{2+}$  transport pump. SERCA2a activity is physiologically limited by its interaction with phospholamban (PLN) (Asahi M et

al., J Biol Chem 1999; 274: 32855–32862.; Toyoshima C et al., Proc Natl Acad Sci U S A 2003;100: 467–47; Bers DM., Annu Rev Physiol 2008;70:23-49.; MacLennan DH, Kranias EG., Nat Rev Mol Cell Biol 2003; 4(7): 566-577), SERCA2a restraint is normally relieved by PLN phosphorylation by protein kinase A (PKA), a signalling pathway severely depressed as a consequence of HF remodelling (Karim CB et al., J Mol Biol 2006; 358: 1032–1040; Lohse M et al., Circ Res 2003; 93:896-906; Bers DM, Physiology 2006;21: 380–387; Mann DL, Bristow MR, Circulation 2005; 111:2837–2849). A deficiency in cardiac SERCA2a activity is widely recognized as the main cause of the reduced  $\text{Ca}^{2+}$  uptake in the SR of the failing myocardium (Bers D et al., Ann N.Y. Acad Sci 2006; 1080:165-177; Bers DM, Physiology 2006;21: 380–387; Minamisawa S et al., Cell 1999; 99: 313–322.).

[5] In addition to its consequences on myocyte contractility and relaxation, abnormal  $\text{Ca}^{2+}$  distribution also facilitates cardiac arrhythmias (Zaza A & Rocchetti M, Curr Pharm Des 2015; 21:1053-1061) and, on the long term, it accelerates myocytes loss by apoptosis (Nakayama H et al., J Clin Invest 2007; 117:2431-44). Reduced SERCA2a function also increases the energy cost of contraction, because it requires a compensatory increase in  $\text{Ca}^{2+}$  extrusion through the Na-Ca exchanger (NCX), which is less energy efficient (Lipskaya L et al., Expert Opin Biol Ther 2010; 10:29-41). Substantial evidence indicates that normalization of SERCA2a function restores intracellular  $\text{Ca}^{2+}$  homeostasis and improves contractility and relaxation of cardiomyocytes and of the heart *in situ* (Byrne MJ et al., Gene Therapy 2008;15:1550-1557; Sato et al., J Biol Chem 2001;276:9392-99). To summarize, recovery of SERCA2a function in HF may improve cardiac relaxation and contractility while minimizing arrhythmias, myocardial oxygen consumption and myocyte death (Lipskaia L et al., Expert Opin Biol Ther. 2010; 10:29-41). In parallel to SERCA2a activation, inhibition of the Na,K-pump can further increase intracellular  $\text{Ca}^{2+}$  content without inducing excessive cytosolic  $\text{Ca}^{2+}$  accumulation (Shattock MJ et al., J Physiol. 2015; 15;593(6):1361-82). Therefore, the combination of Na,K-ATPase inhibition and SERCA2a stimulation may afford further positive inotropy at a reduced risk of arrhythmogenic  $\text{Ca}^{2+}$  triggering events.

[6] Current long-term therapy of HF is centred on prevention of “myocardial remodelling” and neuro-hormonal storm (  $\beta$ -blockers, ACE inhibitors, aldosterone antagonists), which is a chronic maladaptive response to reduced contractility, amplifies the initial damage and underlies disease evolution (Heineke J & Molkentin D, Nat Rev 2006;7:589-600). While this approach has indisputable merit, it does not target impaired heart “contractility” and “relaxation”, which are the functional derangements defining HF and responsible for its symptoms. Indeed, particularly in the advanced disease stages, such as in patients with acute heart failure (AHF), drugs that increase myocardial

contractility/relaxation ("inotropic/lusitropic agents") are still widely used and crucial for the management of patients with AHF (Metra M, Teerlink JR, Lancet 2017;390:1981-1995). These include sympathomimetic amines (dobutamine) and levosimendan, a  $\text{Ca}^{2+}$ -sensitizer with a strong vasodilator effect. Unfortunately, these agents act by mechanisms with potentially harmful components, such as facilitation of life-threatening arrhythmias, increased myocardial oxygen consumption and, in some patients, impairment of an already insufficient coronary blood flow due to the fall in blood pressure caused by vasodilatation (Ashkar H & Makaryus AN, Dobutamine [updated 2018 Oct 27], In StatPearls [Internet], Treasure Island (FL): StatPearls Publishing, 2018-Jan-2017 (available at <https://www.ncbi.nlm.nih.gov/books/NBK470431/>); Gong B. et al., J Cardiothorac Vasc Anesth 2015;29:1415-25; EDITORIAL Patel PA et al., Circ Heart Failure 2014;7:918-925). This limits the use of these agents for relieving the symptoms of the AHF, as clearly stated in both the US and EU guidelines that assign to them and evidence grade C, which is the lowest level of evidence based on the results of the available clinical trials (Rigopoulos AG et al., Herz 2017 Sep 22; Butler J et al., Eur J Heart Fail. 2018;20(5):839-841; Georgiade M et al., J Am Coll Cardiol. 2008;51:2276-85). Furthermore, these agents do not improve patient's prognosis and survival, and their therapeutic use must be carefully monitored (Ashkar H & Makaryus AN, Dobutamine [updated 2018 Oct 27], In StatPearls [Internet], Treasure Island (FL): StatPearls Publishing, 2018 Jan-2017 (available at <https://www.ncbi.nlm.nih.gov/books/NBK470431/>); Gong B. et al., J Cardiothorac Vasc Anesth 2015 29:1415-25).

[7] Among positive inotropes, the cardiac glycoside Digoxin, which is an inhibitor of the Na,K-ATPase enzymatic activity, has been one of the most commonly prescribed medications in the past. However, its use has been decreasing over the last decades because of the difficulty in maintaining digoxin serum concentration ranges at which digoxin displays its beneficial effects (0.5-0.7 ng/ml) without reaching the threshold level of 0.9 ng/ml, above which is observed an increased risk of death due mainly to arrhythmias (Packer M, Journal of Cardiac Failure 2016; 22:726-730; Packer M, Eur J Heart Failure 2018;20:851-852). OMECAMTIV MECARBIL, a cardiac myosin activator that increases cardiac contraction without improving the impaired relaxation, is under clinical development, but its cardiac effects are also associated with an increase of high sensitive troponin plasma levels that indicates some degree of cardiomyocytes injury/damages (Teerlink JR et al., J Am Coll Cardiol. 2016;67(12):1444-1455).

[8] Intensive research is also in progress for the development of HF drugs with mechanisms of action other than positive inotropy. The agents most investigated and under clinical development are: SERELAXIN - recombinant relaxin 2 mediator; ULARITIDE -

recombinant natriuretic peptide; BMS986231 - NO donor; ADRECIZUMAB - Adrenomedullin inhibitor; ANX-042 - spliced variant of NP; TD1439 - Nephylisin (NEP) inhibitor. However, when evaluated in clinical phase 2-3 trials, none of these new agents has met the primary end-point.

[9] The clinical course and prognosis of a patient with chronic heart failure (CHF) is much worse after an episode of AHF (Solomon SD et al., *Circulation* 2007;116:1482-87; Teneggi V et al., *Heart Failure Rev* 2018;23:667-691). AHF can be defined as the new onset or recurrence of symptoms and signs of heart failure, requiring urgent evaluation and treatment and resulting in unscheduled care or hospital admission (Teneggi V et al., *Heart Failure Rev* 2018;23:667-691; Packer M, *Eur J Heart Failure* 2018;20:851-852). Half of the patients with AHF have reduced systolic function (HFrEF), representing a target for potential therapies (Braunwald E., *Lancet* 2015; 385:812-24). Therapies for AHF in patients with reduced ejection fraction (HFrEF) have focused on alleviating congestion with vasodilators, diuretics, or ultrafiltration or increasing cardiac output with positive inotropes. Although this therapeutic strategy has reduced the risk of sudden cardiac death, the post-discharge event rate remains unacceptably high in patients hospitalized for AHF. Many unwanted cardiovascular side effects can be caused by the available therapy, such as myocardial ischemia, cardiac injury and arrhythmias consequent to the inotrope therapy, particularly in patients with coronary artery disease (CAD) (Abraham WT et al., *J Am Coll Cardiol* 2005; 46:57-64; Flaherty JD et al., *J Am Coll Cardiol*. 2009; 53(3):254-63), hypotension and low perfusion of the peripheral organs (kidney) caused by vasodilators particularly in HF patients with low blood pressure. Accordingly, the main goal during hospitalization is to improve cardiac output without causing cardiac and/or kidney injury.

[10] Moreover, there has been little focus on examining or treating an impaired left ventricular (LV) diastolic relaxation that, in the remaining 50% of patients with HF but preserved (50) ejection fraction (HFpEF) or mid-range (40-49) reduction ejection fraction (HFmrEF or HFmEF), is responsible for the symptoms of HF (Butler J et al., *Eur J Heart Fail*. 2018; 20, 839-841; Bonsu KO et al., *Heart Failure Reviews* 2018; 23:147-156). In addition, patients with AHF who have reduced EF also exhibit an impairment of ventricular relaxation that contributes to the overall failure of cardiac function. A variety of echocardiographic indexes has been developed to measure the cardiac relaxation capacity both in animal models and patients with HF (e.g., decreased early mitral annular tissue velocity [e'] and decreased early mitral inflow [E] deceleration time [DT]), along with echocardiographic parameters of increased LV filling pressure (e.g., E/e' ratio). Even though the correspondence of the single index changes is not perfectly superimposable in some animal models and patients, their overall changes in animal models of ventricular

relaxation impairment are certainly translatable to the human condition and used to study the drug effect in AHF (Shah SA et al., *Am Heart J* 2009;157:1035-41).

[11] Various therapeutic approaches that increase SERCA2a function have been previously investigated. These include SERCA2a overexpression by gene transfer (Byrne et al., *Gene Therapy* 2008;15:1550-1557) or PLN inactivation by expression of mutants with negative dominance (Hoshijima M et al., *Nat. Med.* 2002;8:864-871; Iwanaga Y et al., *J Clin Invest* 2004;113, 727-736), AdV-shRNA (Suckau L et al., *Circulation* 2009;119:1241-1252), microRNA (Gröbl T et al., *PLoS One* 2014;9:e92188) or antibodies (Kaye DM et al., *J. Am. Coll. Cardiol.* 2007;50:253-260). As highlighted by the negative results of the largest phase 2b clinical trial applying SERCA2a gene delivery in HF (CUPID 2), these approaches suffer from major problems in construct delivery (viral vectors etc.) and dose adjustment that are far from being solved (Hulot JS, *Eur Heart J* 2016;19:1534-1541). A small-molecule (pyridone derivative) attenuating the inhibitory effect of phospholamban on SERCA2a activity, which is structurally different from istaroxime, has been recently described (Kaneko M et al., *Eur J Pharmacol* 2017;814:1-7), but no data on patients are available.

[12] From the overall picture of the state of the art, and in spite of more than 30 years of trials and related publications, the treatment of patients admitted to hospital because of AHF symptoms is still largely "opinion based" rather than being "evidence based" (Rigopoulos AG et al., *Herz* 2017 Sep 22; Butler J et al., *Eur J Heart Fail.* 2018;20(5):839-841; Georgiade M et al., *J Am Coll Cardiol.* 2008;51:2276-85). Many of the available drugs were designed with rescue and symptom relief in mind and not necessarily to target and correct any specific underlying pathophysiology/biochemical mechanism that may be responsible for the symptoms of AHF.

[13] As a general paradigm, drugs are molecules that produce their wanted or unwanted effect by interacting with the molecules/proteins of patients. The therapeutic benefits of these drugs depend upon their selectivity in correcting the abnormalities of the protein underlying the disease symptoms over other possible effects on proteins with misappropriated or even counterbalancing activities.

[14] The deficiency in cardiac SERCA2a activity is widely recognized as one of the most important causes of the decreased relaxation of cardiomyocytes and increased susceptibility to arrhythmias in patients with cardiac failure (Bers D et al., *Ann N.Y. Acad Sci* 2006; 1080:165-177; Bers DM, *Physiology* 2006;21:380-387; Minamisawa S et al., *Cell* 1999; 99:313-322; Fernandez-Tenorio M & Niggli E., *J Mol Cell Cardiol.* 2018 Jun;119:87-95). To this end, the potential energy starved failing heart status may further

potentiate the consequences of SERCA2a deficiency (Ventura-Clapier R, Garnier A, Veksler V, Joubert F., *Biochim Biophys Acta*. 2011 Jul;1813(7):1360-72; Pinz I et al., *J Biol Chem* 2011; 286(12):10163–10168). In turn, these two causes, if acknowledged, may be adequately addressed. Moreover, for more than 200 years (first evidence in literature: No Author listed, *An Account of the Effects of the Digitalis Purpurea in Dropsy*, *Lond Med J*. 1785;6:55-60), Digitalis, which was subsequently recognized to act throughout the inhibition of the Na-K pump, has been used to increase cardiac pumping activity in spite of some unwanted side effects (*e.g.*, arrhythmias or long term cardiomyocytes damage) (Hougen TJ, Friedman WF., *Am J Physiol*. 1982 Oct;243(4):H517-22; Whitbeck MG et al., *Eur Heart J*. 2013 May;34(20):1481-8). The latter effects are very likely due to the increased cardiomyocytes cytoplasmic  $Ca^{2+}$  that, on one hand is useful for stimulate contraction but, on the other, may favor the above mentioned side effect that are further enhanced by the deficiency of SERCA2a activity (Zaza A & Rocchetti M, *Curr Parm Des* 2015; 21:1053-1061). Consequently, drugs with a combined "selective" effect on these two molecular targets may be beneficial to patients or, at least, may prove or disprove the clinical impact of these two molecular mechanisms.

[15] Notwithstanding the differences in the therapeutic response and outcome of subsets of patients having different degrees of deficiency in relaxation or contraction (Butler J, *Eur J Heart Fail*. 2018;20,839–841; Bonsu KO et al., *Heart Failure Reviews* 2018; 23:147–156) (considering the parameters HFrEF HF, HFmEF or HFpEF Heart Failure reduced Ejection Fraction ( $= < 40$ ), Heart Failure moderate reduction (m or mr) Ejection Fraction (between 40 and 50) Heart Failure preserved Ejection Fraction ( $> 50$ )), it is mandatory to develop therapeutic strategies aimed at assessing the proper combination of the two activities on SERCA2a activation and Na-K pump inhibition for the three subsets of patients.

[16] In particular, there is a strong and, to date, unmet need to improve the therapy of acute heart failure in HFpEF patients (Bonsu KO et al., *Heart Failure Reviews* 2018; 23:147–156) for whom an improvement of diastolic function by correcting the underlying molecular mechanism has not been yet achieved.

[17] Istaroxime (PST 2744) is disclosed in EP0825197 and in De Munari S. et al., *J. Med. Chem.* 2003, 64, 3644-3654 and is the compound (3Z,5a)-3-[(2-aminoethoxy)imino]androstane-6,17-dione. Istaroxime is a new small-molecule drug under clinical development for the treatment of AHFS that is endowed of the double mechanism of action of inhibiting the  $Na^+/K^+$  pump (Micheletti R et al., *J Pharmacol Exp Ther* 2002; 303:592-600) while activating SERCA2a (Rocchetti M et al., *J Pharmacol Exp Ther* 2005;313:207-15).

[18] At the same level of inotropy, the proarrhythmic effect of istaroxime is considerably lower than that of digoxin, a pure Na-K pump inhibitor (Rocchetti M et al., J Pharmacol Exp Ther. 2005;313:207-15). This suggests that, by improving  $\text{Ca}^{2+}$  clearance from the cytosol (Alemanni, J Mol Cell Cardiol 2011;50:910-8), SERCA2a stimulation may also minimize the proarrhythmic effect of Na-K pump blockade (Rocchetti M et al., J Pharmacol Exp Ther 2005;313:207-15; Zaza A & Rocchetti, M Curr Pharm Des 2015;21:1053-1061) while preserving its inotropic effect. The reduction of the proarrhythmic effect by istaroxime has been confirmed in clinical studies (Georghiade M et al., J Am Coll Cardiol 2008;51:2276-85), wherein istaroxime was administered as a continuous 6-hour infusion.

[19] In HF patients, istaroxime infusion improved both systolic and diastolic functions. Amelioration of systolic function was detected as increases in contraction tissue velocity ( $s'$ ) and in the slope of end-systolic elastance (ESPVR slope); increased diastolic compliance was revealed by an increment in the early relaxation tissue velocity ( $e'$ ) and decreased end-diastolic elastance (EDPVR slope) (Shah SA et al., Am Heart J 2009;157:1035-41).

[20] According to the results described in the Horizon study by Gheorghiade (Gheorghiade M et al., J Am Coll Cardiol 2008;51:2276-85), where istaroxime has been infused for 6 hours, the plateau effect on the improvement of diastolic relaxation, continuously measured as a decrease in PCWP (pulmonary capillary wedge pressure), occurs after 3 hours of infusion, after which the level of PCWP remains constant up to 6 hours. As it may be expected from the parallel dual targets, SERCA2a and the Na-K pump, there is no clear separation between the effects on the echocardiographic indexes of relaxation and those of contraction when increasing the infusions doses of istaroxime. Therefore, the potential beneficial effect due to the SERCA2a activation cannot be separated from the potential detrimental effect due to the Na-K pump inhibition when Istaroxime is infused up to 6 hours. Even though both the clinical (Gheorghiade M et al., J Am Coll Cardiol. 2008;51:2276-85; Shah SA et al., Am Heart J 2009;157:1035-41) and experimental studies in dog (Mattera GG et al., Am J Cardiol 2007;99[suppl]:33A-40A) have demonstrated that the presence of the SERCA2a-stimulating activity of istaroxime considerably reduces the pro-arrhythmic activity associated to the Na-K pump inhibition, studies are still not satisfactory as to the clinical outcome, in particular to ensure a properly improved diastolic function and a safer discharge of the patient from hospital. As a matter of fact, the Gheorghiade and Shah clinical trials (Gheorghiade M et al., J Am Coll Cardiol 2008; 51:2276-85; Shah SA et al., Am Heart J 2009;157:1035-41) found that, at the end of the 6 hours infusion, traditional parameters of LV systolic performance, such as stroke volume index (SVI) and Ejection fraction (EF), did not change dramatically with

istaroxime compared to placebo and the duration of the effect on the diastolic relaxation was not properly discussed.

[21] The overall HORIZON study where istaroxime has been infused for 6 hours (see Gheorghiade M et al., J Am Coll Cardiol. 2008; 51:2276-85; Shah SA et al., Am Heart J 2009;157:1035-41) showed a greater improvement of systolic contraction than of diastolic relaxation within a dose range of 0.5 µg/Kg/min and 1.5 µg/Kg/min.

[22] An improved diastolic function is expected to be achieved by a "pure" SERCA2a activator. However, notwithstanding the intense research on discovering small molecules or gene therapy aimed at selectively activating SERCA2a, no promising clinical outcomes have been reached so far.

[23] Accordingly, there is a long-felt need for an advance in the treatment of acute heart failure, in particular for improving diastolic function. The present invention satisfies the above needs and overcomes the problem of prior art.


#### **Summary of the invention**

[24] It has surprisingly been found that the intravenous infusion of istaroxime for a time longer than 6 hours and up to 48 hours or more provides unexpected improvements to the cardiac diastolic relaxation echocardiographic indexes with respect to the same infusion for 6 hours or less, while the echocardiographic indexes of systolic contraction are almost unchanged from 6 hours to 24 hours of infusion time.

[25] Described herein are pharmaceutical compositions containing istaroxime formulated for intravenous infusion in a human subject for use in a treatment method for acute heart failure. In particular, the administration by infusion is for a duration longer than 6 hours, whereby the diastolic relaxation is improved as compared to administration of istaroxime by intravenous infusion for a duration of 6 hours or less (e.g., from 3 to 6 hours). In some embodiments, the diastolic relaxation improvement is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure. In some embodiments, the infusion duration is up to about 24 hours. In others, it is up to about 36 or 48 hours. In such embodiments, the pharmaceutical composition containing istaroxime is administered at a dose between 0.2 µg/kg/min and 1.5 µg/kg/min; preferably it is administered at a dose between 0.25 µg/kg/min and 1.0 µg/kg/min.


[26] In some embodiments, the pharmaceutical composition containing istaroxime is administered for a duration sufficient to produce a plasma level of an istaroxime metabolite that is greater than about 5 ng/ml for at least about 6 hours. In particular, the istaroxime

metabolites may comprise a formula (II) or a formula (III) compound, such as:



**PST 2915**

Formula (II)




**PST 3093**

Formula (III)


[27] In some embodiments, the human subject suffers from heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF) and/or undergoing a therapeutic treatment for heart failure with one or more further therapeutically active ingredients.

[28] Another aspect of the invention features a compound having a formula (II) or formula (III):



**PST 2915**

Formula (II)

**PST 3093**

Formula (III)

[29] In some embodiments, the compounds having the formula (II) or (III) are used in the treatment of a disease requiring the activation of SERCA2a, such as a cardiovascular disease. In particular, the compounds may be used to treat acute heart failure. In some aspects, these compounds are included in an admixture with at least one pharmaceutically acceptable vehicle and/or excipient.

[30] Also disclosed herein are methods for treating an individual having heart failure that include the steps of (1) providing an individual having heart failure; (2) administering a therapeutically effective amount of a pharmaceutical composition containing istaroxime to the individual for an infusion duration of longer than 6 hours; and (3) measuring one or more parameters of heart function, such as diastolic relaxation. In such methods, the administering of the pharmaceutical composition results in an improvement in diastolic relaxation as compared to istaroxime administered by intravenous infusion for 6 hours or less, thereby treating the individual having acute heart failure. In preferred embodiments, the individual is human. In particular embodiments, the infusion duration is up to about 24 hours. In other embodiments, the infusion duration is up to about 36 hours. In still others, the infusion duration is up to about 48 hours. In such methods, the istaroxime may be administered at a dose of about 0.2 µg/kg/min to about 1.5 µg/kg/min. Preferably, it is administered at a dose of about 0.25 µg/kg/min to about 1.0 µg/kg/min.

[31] In some embodiments, the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmrEF). In other embodiments, the individual is undergoing a therapeutic treatment for heart failure with one or more further therapeutically active ingredients.

[32] Also featured herein are pharmaceutical compositions containing istaroxime for intravenous infusion in an individual for the treatment of acute heart failure. In such aspects, the administration is for an infusion duration sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite that is greater than about 5 ng/ml for an accumulation period of at least about 6 hours, whereby the diastolic relaxation is improved as compared to administration of istaroxime prior to the accumulation period. Preferably, the istaroxime metabolite comprises formula (II) or (III).

[33] In some embodiments, the infusion duration is sufficient to produce a plasma concentration level of istaroxime metabolite that is greater than about 10 ng/ml for an accumulation period of at least about 6 hours. In other embodiments, the plasma concentration level of the istaroxime metabolite is greater than about 15 ng/ml for an accumulation period of at least about 6 hours. In still others, it is greater than about 20 ng/ml. In some embodiments, the compositions are administered at a dose of about 0.2 µg/kg/min to about 1.5 µg/kg/min; preferably from about 0.25 µg/kg/min to about 1.0 µg/kg/min).

[34] Other aspects of the invention feature methods for treating an individual having heart failure, including the steps of: (1) providing an individual having heart failure; (2) administering a therapeutically effective amount of a pharmaceutical composition containing istaroxime to the individual for an infusion duration sufficient to produce a plasma concentration level of an istaroxime metabolite that is greater than about 5 ng/ml for an accumulation period of at least about 6 hours; and (3) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation.

[35] In some embodiments, the plasma concentration level of the istaroxime metabolite is greater than about 10 ng/ml for an accumulation period of at least about 6 hours. In other embodiments, it is greater than about 15 ng/ml for an accumulation period of at least about 6 hours. In still others, it is greater than about 20 ng/ml for an accumulation period of at least about 6 hours. The therapeutically effective dose of the composition is between about 0.2 µg/kg/min and about 1.5 µg/kg/min; preferably, between 0.25 µg/kg/min and 1.0 µg/kg/min.

[36] In yet another aspect of the invention, a method for treating an individual having acute heart failure is featured that includes the steps of (1) providing an individual having acute heart failure; (2) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation as measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure; and (3)

administering a therapeutically effective amount of a pharmaceutical composition containing istaroxime to the individual at a dose of at least about 1.0  $\mu\text{g/kg/min}$  for a duration sufficient to improve diastolic relaxation in the individual. Then, after improvement of diastolic relaxation in the individual, a therapeutically effective amount of a second pharmaceutical composition containing istaroxime is administered to the individual at a dose of between about 0.25  $\mu\text{g/kg/min}$  to about 1.0  $\mu\text{g/kg/min}$  for an infusion duration sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite that is greater than about 5 ng/ml for an accumulation period of at least about 6 hours. In such aspects, the istaroxime metabolite comprises formula (II) or (III).

[37] Still other aspects of the invention feature methods for treating an individual having heart failure that include (1) providing an individual having heart failure; (2) administering a therapeutically effective amount of a pharmaceutical composition containing an istaroxime metabolite to the individual for an infusion duration of longer than 6 hours; and (3) measuring one or more parameters of heart function, such as diastolic relaxation. In some aspects, the infusion duration is between 6 hours and 48 hours at a dose of between about 0.2  $\mu\text{g/kg/min}$  to about 1.5  $\mu\text{g/kg/min}$ ; preferably between about 0.25  $\mu\text{g/kg/min}$  to about 1.0  $\mu\text{g/kg/min}$ . In a particular embodiment, the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).

#### **Brief description of the drawings**

[38] These and other features and advantages of the present disclosure will become better understood with regard to the following description, appended claims, and accompanying drawings wherein:

[39] **Figure 1** shows a time course of changes in pulmonary capillary wedge pressure (PCWP) in patients infused with placebo (white diamond) as compared to istaroxime infusion for 6 hours at a dose of 0.5  $\mu\text{g/kg/min}$  (dark diamond), 1.0  $\mu\text{g/kg/min}$  (square), and 1.5  $\mu\text{g/kg/min}$  (triangle). The X-axis represents the average PCWP (mmHg), and the Y-axis represents time (hours).

[40] **Figure 2A** shows the plasma levels of istaroxime (square) and its metabolites in Caucasian patients intravenously infused with 0.5  $\mu\text{g/kg/min}$  istaroxime for 24 hours. The metabolites are PST 2915 (diamond), PST 2922 (circle), and PST 3093 (triangle). The X-axis represents the plasma concentration (ng/ml), and the Y-axis represents time (hours).

[41] **Figure 2B** shows the plasma levels of istaroxime (square) and its metabolites in Caucasian patients (left panel) and Chinese patients (right panel) intravenously infused with 0.5 µg/kg/min istaroxime for 24 hours. The metabolites are PST 2915 (diamond), PST 2922 (circle), and PST 3093 (triangle). The X-axis represents the plasma concentration (ng/ml), and the Y-axis represents time (hours).

[42] **Figure 2C** shows the plasma levels of istaroxime (square) and its metabolites in Chinese patients intravenously infused with 1.0 µg/kg/min istaroxime for 24 hours. The metabolites are PST 2915 (diamond), PST 2922 (circle), and PST 3093 (triangle). The X-axis represents the plasma concentration (ng/ml), and the Y-axis represents time (hours).

#### **Detailed description of the invention**

[43] The compositions and methods disclosed herein confer to individuals suffering from heart failure unexpected benefits. Provided herein are compositions comprising istaroxime or a metabolite thereof. Further, as disclosed herein, infusion with istaroxime or its metabolites for more than 6 hours improves selectively cardiac relaxation over cardiac contraction. Moreover, istaroxime infusion time can be extended to allow for the accumulation of one or more of its metabolites, at least one of which exhibits single-function SERCA2a activation (*i.e.*, behaves as a "pure" SERCA2a activator). The compositions and methods disclosed herein will be described in more detail below.

#### Definitions

[44] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as those commonly understood by one of ordinary skill in the art to which this invention belongs. Standard techniques are used unless otherwise specified. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. The materials, methods and examples are illustrative only, and are not intended to be limiting. All publications, patents and other documents mentioned herein are incorporated by reference in their entirety.

[45] As used herein, the singular forms "a," "an," and "the" include the plural referents unless the context clearly indicates otherwise.

[46] The term "about" refers to the variation in the numerical value of a measurement, *e.g.* used to obtain that measure. In one embodiment, the term "about" means within 5% of

the reported numerical value, preferably, the term "about" means within 3% of the reported numerical value.

[47] The term "heart failure" refers to a clinical syndrome characterized by typical symptoms (*e.g.*, breathlessness, ankle swelling and fatigue) that may be accompanied by signs (*e.g.*, elevated jugular venous pressure, pulmonary crackles and peripheral edema) caused by a structural and/or functional cardiac abnormality, resulting in a reduced cardiac output and/or elevated intracardiac pressures at rest or during stress.

[48] The terms "acute heart failure" or "AHF" are used interchangeably herein and refer generally to a rapid onset or worsening of symptoms and/or signs of HF requiring immediate treatment and hospitalization. The current definition of "acute heart failure" is rather nonspecific and may include a broad spectrum of conditions with several phenotypes characterized by different clinical presentation, etiology, precipitating factors, therapeutic approach, and prognosis. In addition, a large proportion of patients have a subacute course of the disease with a progressive worsening of signs and symptoms of HF which could develop days before hospital admission.

[49] The terms "chronic heart failure" or "CHF" are used interchangeably herein and refer to the current clinical classification of chronic HF based on the presence of signs and symptoms of HF and left ventricular ejection fraction (LVEF), recognizing three categories: "heart failure with reduced ejection fraction" or "HFrEF," which is characterized by an LVEF of less than about 40%; "heart failure with mid-range ejection fraction" or "HFmrEF" or "HFmrEF," which is characterized by an LVEF from about 40% to about 49%; and "heart failure with preserved ejection fraction" or "HFpEF," which is characterized by an LVEF of equal to or greater than about 50%. The terms "HFmrEF" and "HFpEF" include two additional criteria, namely increased natriuretic peptides levels (BNP >35 pg/ml and/or NT-proBNP >125 pg/mL) associated with the evidence of structural and/or functional heart disease (left ventricular hypertrophy and/or left atrium enlargement and/or evidence of diastolic dysfunction). The efficacy of HF evidence-based medications have been confirmed only in patients with "HFrEF," whereas no treatment demonstrated a significant improvement of outcomes in patients with "HFpEF".

[50] The term "treating" refers to any indicia of success in the treatment or amelioration of the disease or condition. Treating can include, for example, reducing or alleviating the severity of one or more symptoms of the disease or condition, or it can include reducing the frequency with which symptoms of a disease, defect, disorder, or adverse condition, and the like, are experienced by an individual, such as a human patient.

[51] The term "preventing" refers to the prevention of the disease or condition, *e.g.*, acute heart failure, in an individual, such as a human patient. For example, if an individual at risk of developing heart failure is treated with the methods of the present invention and does not later develop heart failure, then the disease has been prevented in that individual.

[52] The term "treat or prevent" is sometimes used herein to refer to a method that results in some level of treatment or amelioration of the disease or condition, and contemplates a range of results directed to that end, including but not restricted to prevention of the condition entirely.

[53] As used herein, the term "pharmaceutically acceptable carrier" means a chemical composition with which an istaroxime compound or a metabolite of istaroxime may be combined and which, following the combination, can be used to administer the compound to a mammal.

[54] As used herein, the term "pharmaceutically acceptable" salt, solvate, hydrate, or ester means a salt, solvate, hydrate, or ester form of the active ingredient which is compatible with any other ingredients of the pharmaceutical composition, which is not deleterious to the subject to which the composition is to be administered.

[55] The term "intravenous infusion" refers to the administration or delivery of liquid substances directly into a vein of a mammal. Typical "infusions" use only the pressure supplied by gravity.

[56] The term "parameter" as used herein to refer to measuring heart function means any heart function that is observable or measurable using suitable measuring techniques available in the art. A non-limiting list of exemplary "parameters" of heart function include heart rate, blood pressure, diastolic relaxation, systolic contraction, LVEF, diastolic blood pressure, systolic blood pressure, cardiac output, stroke volume, deceleration slope, cardiac index, mitral inflow velocity, and the like. As one having ordinary skill in the art will appreciate, measuring one or more "parameters" of heart function can be used to detect heart dysfunction as compared to the average normal parameters and can also be used to determine whether heart function has improved following or during treatment.

[57] The terms "therapeutically active" or "active" ingredient or compound refer to a substance that provides a beneficial effect to the individual to whom the substance is administered. A "therapeutically effective amount" or "therapeutically effective dose" is the amount of a composition or active ingredient sufficient to provide a beneficial effect to the individual to whom the composition or active ingredient is administered.

### Description

[58] The present invention is based on the unexpected discovery that istaroxime infusion for more than 6 hours provides a prevailing lusitropic effect or improvement of the reduced cardiac relaxation, as shown by the clear changes of the echo indexes of cardiac relaxation (E/A DTs,  $e'$ ,  $E/e'$  and left atrial area or volume) while those of contraction ( $S_a$  and  $s$ ) remained unchanged. In some embodiments, the istaroxime infusion is between about 6 hours and up to 48 hours or more, *e.g.*, 6 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19 h, 20 h, 21 h, 22 h, 23 h, 24 h, 25 h, 26 h, 27 h, 28 h, 29 h, 30 h, 31 h, 32 h, 33 h, 34 h, 35 h, 36 h, 37 h, 38 h, 39 h, 40 h, 41 h, 42 h, 43 h, 44 h, 45 h, 46 h, 47 h, 48 h, or more. For instance, the infusion can be for up to 24 hours, or up to 36 hours, or up to 48 hours, or more. In other embodiments, the duration of the infusion is greater than 6 hours and equal to or less than about 48 hours, or greater than 6 hours and equal to or less than about 36 hours, or greater than 6 hours and equal to or less than about 24 hours, or greater than 6 hours and equal to or less than about 12 hours. It being understood that the infusion of istaroxime or a metabolite thereof provides a prevailing lusitropic effect or improvement of the reduced cardiac relaxation as compared to infusion with istaroxime or a metabolite thereof for a period of 6 hours or less, *e.g.*, 0, 1, 2, 3, 5, or 6 hours.


[59] Importantly, an analysis of four independent groups of data confirms that istaroxime infusion for more than 6 hours provides a prevailing lusitropic effect and improvement of the cardiac relaxation. First, **Figure 1** shows a PCWP time course of patients intravenously infused with istaroxime for up to 6 hours. As previously described in the HORIZON study of patients intravenously infused with istaroxime for 6 hours, time course changes of PCWP, which the skilled artisan will appreciate as a valid marker of diastolic relaxation, shows an improvement of diastolic relaxation during the initial 3 hours, but then the average PCWP plateaus and remains unchanged for the following 3 hours (see **Figure 1**). In other words, patients infused for 6 hours with istaroxime showed no improvement in diastolic relaxation between 3 and 6 hours of the infusion. In contrast, patients administered intravenous infusion of istaroxime for 24 hours exhibit a clear increase in echocardiographic indexes of diastolic relaxation from 6 to 24 hours, while the indexes of systolic contraction remain unchanged (see **Tables 1A-1C**). Moreover, as shown in **Figures 2A-2C**, there is a progressive and remarkable increase in the plasma concentration of the istaroxime metabolite PST 3093 at the 6-hour and 24-hour time points, while the plasma concentration of istaroxime remains constant throughout this time interval. Finally, the synthesis of PST 3093 and the subsequent biochemical and pharmacological studies demonstrate that this compound is endowed of a selective

SERCA2a stimulatory activity (see, e.g., Example 2 and **Table 3**) at concentrations well below the that of the intravenous infusion studies shown in **Figure 2**. Moreover, its infusion in a rat model of diabetic cardiomyopathy is associated with an improvement in diastolic relaxation (see **Table 5**). According to Munafò *et al.* (Nature 2018, 553(7689):399-401), simply repeating single experiments is not sufficient, but rather many lines of evidence is needed. Thus, it is the consistency among these four independent findings that, *per se*, confers scientific robustness to the assertion that istaroxime infusion for more than 6 hours provides a prevailing lusitropic effect and improvement of the cardiac relaxation as compared to shorter infusion times.

[60] These findings are entirely unexpected, since the plasmatic level of istaroxime remains constant (*i.e.*, within about 10 ng/ml at 3 hour time point to about 8ng/ml) during the following time points for the whole duration of the infusion. The present inventors have discovered that in humans this prolonged infusion of istaroxime generates an increasing concentrations of istaroxime metabolites PST 3093 and PST 2915, which behave as selective or “pure” SERCA2a activator. Further, this selectively is even greater for PST 3093, thus explaining the unexpected effect in improving diastolic function over systolic function.

[61] Consequently, patients with HFpEF or HFmEF may benefit from a more selective correction of the impaired diastolic relaxation by increasing the plasma level of the metabolite over that of istaroxime. Thus, minimizing the Na<sup>+</sup>/K<sup>+</sup> pump inhibition, notwithstanding the constant plasma level of istaroxime, with its associated unwanted effects in term of arrhythmias or cardiomyocytes damage. Advantageously, the clinical outcome is a safer patient discharge after treatment of acute heart failure.


[62] Istaroxime is an inotropic compound having the following structural formula (I):




Formula (I)

[63] Upon administering to a mammal, such as a human, istaroxime is metabolized into several metabolites that are capable of activating SERCA2a.


[64] Istaroxime metabolic pathway is illustrated below:



[65] As such, disclosed herein are metabolites of istaroxime having SERCA2a activity that have the following structural formulas (II) and (III):



Formula (II)

**PST 3093**

Formula (III)

[66] In preferred embodiments, the metabolite of istaroxime (here also named PST 3093) endowed with selective or “pure” SERCA2a activity is the compound of formula (III).

[67] The present inventors have isolated and characterized PST 2915 and PST 3093. In rats with diabetic cardiomyopathy, administration of PST 3093 showed improved diastolic relaxation and overall cardiac function as measured as an increase of stroke volume SV.

[68] Therefore, it is also an object of the present invention to utilize the SERCA2a-activation properties of the compound of formula (II) or the compound of formula (III), or their respective pharmaceutically acceptable salts or esters, as well as their different hydrates, solvates, polymorphic forms. Another object of the present invention is the compound of formula (II) or formula (III) for use as medicament, in particular for treating diseases requiring the activation of SERCA2a, more preferably for the treatment of heart failure or acute heart failure.

[69] Also disclosed herein is a pharmaceutical composition comprising the compound of formula (II) or formula (III), in an admixture with at least one pharmaceutically acceptable vehicle and/or excipient. In preferred embodiments, the pharmaceutical composition is formulated for administering to an individual by infusion, preferably, it is by intravenous infusion.

[70] The surprising effect of the present invention on the improved cardiac diastolic relaxation can be better appreciated from the comparison of cardiac parameters between the 6-hour infusion of the previous study (HORIZON clinical trial) and the 24-hour infusion according to the present invention. The outline of the study is described below and the data at 6 hours of infusion, 24 hours of infusion, and 48 hours of infusion are summarized in **Tables 1A, 1B, and 1C**, respectively.

[71] The synopsis of the clinical trial is described herein:

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Title                      | The clinical study of the safety and efficacy of Istaroxime in Treatment of Acute Decompensated Heart Failure - A multicenter, randomized, double-blind, placebo controlled, parallel group clinical study.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Indication                 | Acute Decompensated Heart Failure (ADHF)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Objective                  | To Assess the safety, tolerability and efficacy of two different doses of istaroxime (0.5 and 1.0 µg/kg/min), a new agent with lusitropic and inotropic activities that improves the cardiac contraction-relaxation cycle. The 2 doses of istaroxime (0.5 and 1.0 µg/kg/min) will be infused i.v. for 24 hours in comparison with placebo, in treatment of Chinese and Caucasian patients with Acute Decompensated Heart Failure. In all the Caucasian patients and in a subset of Chinese patients pharmacokinetics and metabolism of istaroxime shall also be studied.                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Study Design               | A multicenter, randomized, double-blind, placebo-controlled, parallel group study.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Study Period               | This study includes a screening period (Days -1), a treatment period (Day 1), a post-treatment period (Days 2-4), and a follow-up period (which includes one patient visit on Day 30).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Subject Selection Criteria | <p><b>Inclusion criteria</b></p> <p>Patients who fulfill the following inclusion criteria at screening will be considered for the study:</p> <ol style="list-style-type: none"> <li>1. Signed informed consent;</li> <li>2. Male or female patients 18-85 years (inclusive);</li> <li>3. Admission for a recurrent ADHF episode with dyspnea at rest or minimal exertion and need of intravenous diuretic therapy (<math>\geq 40</math> mg iv. furosemide);</li> <li>4. Systolic blood pressure between 90 and 125 mmHg (limits included) without signs or symptoms of hypoperfusion including cardiogenic shock, cold extremities and peripheral vasoconstriction, oliguria/anuria, signs of cerebral hypo perfusion such as confusion;</li> <li>5. Left ventricular (LV) Ejection fraction (EF) <math>\leq 40</math> % measured by 2D-Echocardiography</li> <li>6. E/Ea ratio <math>&gt;10</math></li> <li>7. BNP <math>\geq 350</math>pg/mL or NT-pro-BNP <math>\geq 1400</math> pg/mL</li> </ol> |

8. Adequate echocardiography window (defined as visualization of at least 13/16 segment of the left ventricle);

#### **Exclusion Criteria**

Any of the following criteria established at screening would render a patient ineligible for the study:

1. Pregnant or breast-feeding women (women of child bearing potential must have the results of a negative pregnancy test recorded prior to study drug administration)
2. Current (within 12 hours prior to screening) or planned (through the completion of study drug infusion) treatment with any iv. therapies, including vasodilators (including nitrates or nesiritide), positive inotropic agents and vasopressors
3. Current or need of mechanical support (intra-aortic balloon pump, endotracheal intubation, mechanical ventilation, or any ventricular assist device),
4. Ongoing treatment with oral digoxin. Patient treated with digoxin cannot be randomized. However, if digoxin treatment has been stopped during the last week before randomization and the digoxin plasma level is < 0.5 ng / ml, patient may be randomized;
5. History of hypersensitivity to the study medication or any related medication
6. Diagnosis of cardiogenic shock within the past month;
7. Acute coronary syndrome or stroke within the past 3 months;
8. Coronary artery bypass graft or percutaneous coronary intervention within the past month or planned in the next month;
9. Primary hypertrophic or restrictive cardiomyopathy or systemic illness known to be associated with infiltrative heart disease;
10. Cor pulmonale or other causes of right-sided HF not related to left ventricular dysfunction;
11. Pericardial constriction or active pericarditis;
12. Atrial fibrillation with marked irregularities of heart rhythm;
13. Life threatening ventricular arrhythmia or ICD (implantable cardioverter defibrillator) shock within the past month;
14. CRT (cardiac resynchronization therapy), ICD or pacemaker implantation within the past month;
15. Valvular disease as primary cause of HF;
16. Heart rate >120 bpm or < 50 bpm
17. Acute respiratory distress syndrome or ongoing sepsis;
18. Fever >38°
19. History of bronchial asthma or porphyria;
20. Donation or loss of blood equal to or exceeding 500 mL, during the 8 weeks before administration of study medication;
21. Positive testing for Hepatitis B and/or Hepatitis C with abnormal liver functions;
22. Participation in another interventional study within the past 30 days;
23. The following laboratory exclusion criteria, verified based on results obtained within the last 24 hours of hospitalization:
  - a. Serum creatinine > 3.0 mg/dl (> 265 µmol/L);

|                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                     | <p>b. Aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) &gt; 3 x upper limit of normal,</p> <p>c. Hemoglobin (Hb) &lt; 10 g/dL,</p> <p>d. Platelet count &lt; 100,000/<math>\mu</math>L,</p> <p>e. Serum potassium &gt; 5.3 mmol/L or &lt; 3.8 mmol/L,</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Study Drugs         | <b>Test drug: Istaroxime (10 mg per vial)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Mode administration | <b>Intravenous infusion via a syringe pump.</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Treatment duration  | <b>Treatment by i.v. infusion will last 24 hours.</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Dosing scheme       | <b>Istaroxime 0.5 – 1.0 <math>\mu</math>g/kg/min since the beginning.</b> A continuous i.v. infusion for 24 hours not exceeding 144 mg for 24 hours of istaroxime for patients with body weight > 100 kg shall be carried out.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Sample Size         | 120 total patients (96 Chinese patients and 24 Caucasian patients)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Study Procedures    | <p><b>Screening period (between Hours -24 to -1)</b></p> <p>Within a maximum of 24 hours before administration of study medication (istaroxime), a medical screening will be performed on all prospective patients to assess suitability for the study. Prior to conducting any study specific procedures, the investigator or his/her designee will explain the study fully to the patient and provide him/her with a copy of the Patient Information Sheet and Informed Consent Document. If the patient is willing to participate in the study, s/he and the investigator or his/her designee will both sign the Informed Consent Document and a copy of the signed document will be kept by the patient.</p> <p><b>Treatment period (Day 1)</b></p> <ol style="list-style-type: none"> <li>1) Confirm eligibility;</li> <li>2) Randomization of patients (after eligibility has been confirmed)</li> <li>3) Insertion of multiple lumen intravenous catheter</li> <li>4) Start istaroxime or placebo infusion (date and time of infusion start must be recorded in the CRF)</li> <li>5) cTnT (at pre-dose: two samples, then at 3 and 6, 12, 24, 48 and 72 hours after start of infusion)</li> <li>6) NT pro-BNP at baseline and at the end of 24 hours infusion</li> <li>7) Blood samples collection for metabolites and PK (at pre-dose, 0.5-3-6-12-24 after start of infusion and at 0.25, 0.5, 1, 4 12, 24 hours after the end of infusion) in all the Italian patients and in a subset of Chinese patients pharmacokinetics and metabolism of istaroxime shall also be studied.</li> <li>8) Vital signs (including body temperature and dyspnoea at pre-dose, 3, 6,</li> </ol> |

|  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|--|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | <p>12 and 24 after the start of infusion)</p> <p>9) 12-lead ECG profile</p> <p>10) Stop Day -1 Holter</p> <p>11) Start 24-hour Holter ECG (Day 1 recording; to be started immediately before initiation of the study drug infusion)</p> <p>12) Echocardiography at baseline and 6 and 24 hours after infusion start</p> <p>13) 24-hours urine collections for measurement of istaroxime and its metabolites and urinary creatinine for the calculation of the creatinine clearance;</p> <p>14) Blood collection for K<sup>+</sup> and eGFR between 23 hours and 30 minutes and 23 hours and 55 minutes since infusion start;</p> <p>15) Concomitant medication monitoring (including chronic medication; dose, date and time must be recorded on CRF)</p> <p>16) Adverse events monitoring</p> <p><b>Post-treatment period (Day 2 to Day 4)</b></p> <p><b>Evaluations at 24 hours (day 2) from randomization include:</b></p> <p>1) Vital signs (including body temperature and dyspnea);</p> <p>2) 12-lead ECG (single ECGs);</p> <p>3) Stop 24-hour Holter ECG;</p> <p>4) Start 24-hour Holter ECG (Day 2 recording);</p> <p>5) Stop istaroxime infusion (date and time of infusion end must be recorded in the CRF);</p> <p>6) Serum potassium level and 24-hour urine collection for measurement of istaroxime metabolites and urinary creatinine for calculation of the creatinine clearance;</p> <p>7) Serum creatinine clearance and calculation of eGFR;</p> <p>8) cTnT (50 % or 20 % relative increase over the basal cTnT levels, respectively for patients with cTnT basal levels &lt; or &gt; of the 99 % URL (upper reference levels, as defined for the Roche hs test, in patients with normal renal function, eGFR ≥ 85 ml/min); in patients with eGFR below this value, the renal function variations must be considered in evaluating the significance of the cTnT changes);</p> <p>9) NT pro-BNP;</p> <p>10) Metabolites;</p> <p>11) Echocardiography;</p> <p>12) Concomitant medication monitoring (including chronic medication must be recorded in the CRF);</p> <p>13) Adverse Events monitoring.</p> <p><b>Evaluations at 48 hours (day 3) include:</b></p> <p>1) Vital signs (including body temperature and dyspnea);</p> |
|--|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

- 2) 12-Lead ECG (single ECGs);
- 3) Stop 24-hour Holter ECG;
- 4) Standard hematology;
- 5) Standard blood chemistry;
- 6) Serum potassium level;
- 7) 24-h urine collection for measurement of istaroxime metabolites and urinary creatinine for the calculation of creatinine clearance ;
- 8) Calculation of eGFR;
- 9) NT-proBNP;
- 10) cTnT;
- 11) Blood samples for istaroxime metabolites;
- 12) Echocardiography
- 13) Adverse events monitoring;
- 14) Concomitant medication monitoring (including chronic medication must be recorded in the CRF);

**Evaluations at 72 hours (day 4) include:**

- 1) cTnT and NTproBNP (at 72 hours after start of infusion)
- 2) Vital signs (including body temperature and dyspnoea)
- 3) Physical examination (HF signs included)
- 4) 12-lead ECG
- 5) Adverse events monitoring
- 6) Concomitant medication monitoring (including chronic medication)
- 7) Istaroxime metabolites
- 8) Serum potassium and creatinine levels for calculation of eGFR
- 9) Creatinine clearance

**Follow-up period and visit (Day 5 to Day 30)**

During the follow-up period the Investigator/designee will make every effort to establish patient outcomes.

**Evaluations on Day 30 (follow-up visit) include:**

- 1) Vital signs (including body temperature and dyspnoea);
- 2) 12-lead ECG in triplicate;
- 3) Calculation of eGFR;
- 4) Standard hematology;
- 5) Standard blood chemistry;
- 6) NT-proBNP
- 7) cTnT;
- 8) Urine pregnancy test ( $\beta$ -HCG) for females of childbearing potential
- 9) Urinalysis;
- 10) Physical examination (HF signs included);
- 11) Adverse events monitoring;
- 12) Concomitant medication monitoring (including chronic medication)

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Efficacy<br>Parameters | <p><b>Efficacy endpoints</b></p> <p><b>1. Primary efficacy end-point:</b><br/>Change from baseline to 24 hours after infusion start (treatment period Day 1) in the E/Ea ratio assessed by tissue Doppler.</p> <p><b>2. Secondary efficacy end-points:</b><br/>Change from baseline to 24 hours in the treatment period Day1 (addressing the differences between the changes at 6 and 24 hours from baseline) of the following Echo-Doppler parameters:</p> <ul style="list-style-type: none"> <li>- LV Ejection fraction (EF)</li> <li>- LV end systolic and end diastolic volumes</li> <li>- Stroke volume index (SVI)</li> <li>- E, A and E/A ratio</li> </ul> <p>Difference between the changes at 6 and 24 hours from baseline of the Tissue Doppler parameter E/Ea</p> <p>Others Tissue Doppler parameters such as Sa, Da and Aa</p> <p>Changes in dyspnoea assessed at 3, 6, 12, 24, 48 hours after infusion start by Visual Analog Scale (VAS) (including only patients presenting dyspnoea at baseline);</p> <p>Area under the curve (AUC) on changes in dyspnoea assessed at 3, 6, 12, 24, 48 hours after infusion start by VAS (including only patients presenting dyspnea at baseline);</p> <p>Changes in BNP from baseline at 24 hours;</p> <p>Proportion of patients with hospital readmissions or emergency visits for cardiovascular reasons by Day 30;</p> <p>Proportion of patients with episodes of worsening HF defined by the need to increase the dose or reinitiate i.v. therapy with diuretics and/ or other inotropic agents during the hospitalization;</p> <p>Length of the hospitalization;</p> |
| Safety<br>Parameters   | <p><b>Safety endpoints:</b></p> <p>The following safety endpoints will be assessed during treatment and the post-treatment/follow-up periods:</p> <ul style="list-style-type: none"> <li>Incidence of adverse events;</li> <li>Change in vital signs (including body temperature and dyspnoea);</li> <li>Change in 12-lead ECG parameters;</li> <li>Incidence of clinically or hemodynamically significant episodes of supraventricular or ventricular arrhythmias detected by continuous ECG dynamic monitoring;</li> <li>Change in laboratory parameters (hematology, blood chemistry and urinalysis);</li> <li>Change in renal function;</li> <li>Change in in cTnT;</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |

|                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                      | <p>Incidence of cTnT elevation (&gt;50% or &gt; 20% relative increase over the basal cTnT levels at baseline, for patients with cTnT levels at baseline &lt; or ≥ of the 99% URL (upper reference levels, as defined for the Roche hs test, in patients with normal renal function, eGFR ≥85 ml/min); in patients with eGFR below this value, the renal function variations must be considered in evaluating the significance of the cTnT changes);</p> <p>Mortality at Day 30;</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| PK parameters        | <p><b>Full plasma and urine PK profile:</b></p> <p>The following PK metrics will be computed for E and Z isomers (when applicable) of istaroxime plasma concentrations using non-compartmental analysis: C<sub>max</sub>, t<sub>max</sub>, AUC<sub>0-t</sub>, AUC<sub>0-∞</sub>, I<sub>z</sub>, t<sub>1/2</sub>, C<sub>IT</sub>, MRT, V<sub>ss</sub>, V<sub>z</sub>;</p> <p>the following PK metrics will be computed for E and Z isomers (when applicable) of istaroxime urine concentrations: Ae, Ae%, C<sub>IR</sub>;</p> <p>In addition, the following PK metrics will be computed as above for plasma and urine concentrations of the E and Z isomers (when applicable) of istaroxime metabolites 2915, 2922, and 3093: C<sub>max</sub>, t<sub>max</sub>, AUC<sub>0-t</sub>, AUC<sub>0-∞</sub>, I<sub>z</sub>, t<sub>1/2</sub> and, if possible, Ae and Ae%;</p>                                                                                                                                                                                                                                                                                                                                 |
| Statistical Analysis | <p><u>Primary efficacy endpoint:</u></p> <p>The primary efficacy endpoint (change from baseline in E/Ea ratio) will be analyzed using a linear mixed model for repeated measures including treatment, centre, timepoint, gender, baseline cTnT (normal &lt;URL, abnormal ≥URL), atrial fibrillation (Yes/No) and treatment*timepoint interaction as fixed effects and baseline and baseline*timepoint interaction as covariates.</p> <p>The primary comparison will be 0.5 µg/kg/min dose of istaroxime versus placebo at 24 hours. Highest dose of istaroxime (1.0 µg/kg/min) versus placebo will be tested as a secondary comparison.</p> <p>Additional analyses separated by cohort will be implemented for sensitivity purpose.</p> <p><u>Secondary efficacy endpoints</u></p> <p>The following secondary endpoints:</p> <p>Change from baseline to 24 hours (addressing the differences between the changes at 6 and 24 hours from baseline) of the following Echo-Doppler parameters:</p> <ul style="list-style-type: none"> <li>- LV Ejection fraction (EF)</li> <li>- LV end systolic and end diastolic volumes</li> <li>- Stroke volume index (SVI)</li> <li>- E, A and E/A ratio</li> </ul> |

|  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|--|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|  | <p>Change from baseline to 24 hours in the E/Ea ratio assessed by tissue Doppler (difference between the changes at 6 and 24 hours from baseline)</p> <p>Others Tissue Doppler parameters such as Sa, Da and Aa</p> <p>Changes in dyspnoea using VAS score will be analysed using a mixed model for repeated measures similar to the one used for the primary efficacy endpoint.</p> <p>AUC on changes in dyspnoea by VAS and changes in BNP will be analyzed using an ANCOVA model with treatment, centre, gender, baseline cTnT (normal &lt;URL, abnormal ≥URL) and atrial fibrillation (Yes/No) as fixed effects and baseline dyspnea as covariate.</p> <p>Number and proportion of patients with:</p> <p>hospital readmissions or emergency visits for cardiovascular reasons within Day 30</p> <p>episodes of worsening HF defined by the need to increase the dose or reinstitute i.v. therapy with diuretics and/ or other inotropic agents during the hospitalization</p> <p>will be summarized by treatment groups using descriptive statistics.</p> <p>Length of hospitalization will be summarized by treatment group using descriptive statistics.</p> <p><u>Safety endpoints</u></p> <p>The number and the percentage of patients experiencing adverse events, adverse drug reactions, serious adverse events and adverse events leading to study withdrawal will be summarized by treatment group. Adverse events will also be summarized by treatment group by means of System Organ Class and Preferred Term using the MedDRA dictionary.</p> <p>Vital signs (including body temperature and dyspnoea), 12-lead ECG parameters, incidence of clinically or hemodynamically significant episodes of supraventricular or ventricular arrhythmias, laboratory parameters, renal function, cTnT, increase of cTnT and mortality will be summarized by treatment group using descriptive statistics.</p> |
|--|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

[72] Echocardiography was performed on patients according to international standards (see, for example, Lang RM et al., J Am Soc Echocardiogr 2005;18(12):1440-63; Nagueh SF et al., Eur J Echocardiogr 2009;10(2):165-93; Evangelista A et al., Eur J Echocardiogr 2008;9(4):438-48). Echocardiography was performed by expert physicians or sonographers at the sites. Echocardiography was done at screening, baseline, 6 hours after

infusion start, 24 hours after infusion start (just before the end of infusion), and 48 hours after infusion start.

[73] The following parameters were recorded for each patient at each timepoint and centrally measured by the CoreLab:

1. Cardiac dimension measures:

- a. Left ventricle end diastolic diameter (EDD): measured with M-mode echocardiography at the level of mitral valve (MV) leaflets from parasternal long axis view (PLAX) (normal range [NR]: 42-59 mm males and 39-53 mm females);
- b. Left ventricle end systolic diameter (ESD): measured with M-mode echocardiography at the level of mitral valve (MV) leaflets from PLAX (NR: 25-35 mm);
- c. Left ventricle end diastolic volume (EDV): measured with M-mode echocardiography at the level of mitral valve (MV) leaflets from PLAX (NR: 67-155 mL males and 56-104 mL females);
- d. Left ventricle end systolic volume (ESV): measured with M-mode echocardiography at the level of mitral valve (MV) leaflets from PLAX (NR: 22-58 mL males and 19-49 mL females);
- e. Left atrium diameter (LAD): measured at end-ventricular systole with M-mode echocardiography from PLAX. (NR: 30-40 mm males and 27-38 mm females)
- f. Left atrium area (LAA): measured from apical four chamber view (NR:  $\leq 20$  cm<sup>2</sup>); and
- g. Left atrium volume (LAV): derived from area-length measured from apical four chamber view (NR: 18-58 mL males and 22-52 mL females).

2. Left ventricle diastolic function parameters:

- a. E wave: measured from mitral valve pulsed wave Doppler, is the peak velocity of early filling. Normal range for all the diastolic parameters significantly changes with age;
- b. A wave: measured from mitral valve pulsed wave Doppler is the peak velocity of late atrial filling. Not evaluable in patients with AF;
- c. E wave deceleration time (EDT): measured from mitral valve pulsed wave Doppler represent the slope of the descending part of E wave;
- d. E/A ratio: determines the type of diastolic filling pattern (normal E/A= 1-2 and EDT= 150-200 ms, abnormal relaxation E/A<1 and EDT $\geq$ 240 ms,

- pseudonormal  $E/A=0.8-1.5$ ; restrictive  $E/A \geq 2$  and  $EDT < 160$  ms). Not evaluable in patients with AF;
- e.  $E_a$ : measured with tissue Doppler method at the lateral and septal side of the mitral annulus from apical four chamber view is the early diastolic velocity. The value has been calculated as the average between  $E_a$  lateral and  $E_a$  septal. (NR  $\geq 10$  cm/s) (see Nagueh SF et al., Eur J Echocardiogr. 2009;10(2):165-93);
  - f.  $A_a$ : measured with tissue Doppler method at the lateral and septal side of the mitral annulus from apical four chamber view is the late atrial diastolic velocity. The value has been calculated as the average between  $A_a$  lateral and  $A_a$  septal. Not evaluable in patients with AF; and
  - g.  $E/E_a$  ratio: this is a derived measure from  $E$  and  $E_a$  value. This is highly correlated with left ventricle filling pressure and with prognosis in patients with HF. (NR:  $< 13$ ) (see Nagueh SF et al., *supra*).
3. Left ventricle systolic function parameters:
- a. Left ventricle ejection fraction (LVEF): measured with Simpson biplane method according to international recommendations from apical four chamber view and apical two chamber view. (NR  $\geq 55\%$ ) (see Lang RM et al., J Am Soc Echocardiogr. 2005;18(12):1440-63); and
  - b.  $S_a$ : measured with tissue Doppler method at the lateral and septal side of the mitral annulus from apical four chamber view. The value has been calculated as the average between  $S_a$  lateral and  $S_a$  septal. Validation studies demonstrated that  $S_a$  correlates with LVEF (NR  $\geq 6$  cm/s) (see Gulati VK et al., Am J Cardiol. 1996;77(11):979-84).
4. Overall cardiac contraction parameters:
- a. Stroke volume (SV): is a derived measure obtained with the application of Bernoulli's formula using the dimension of left ventricle outflow tract (LVOT) as diameter and LVOT time velocity integral as velocity. (NR  $> 60$  mL/beat);
  - b. Cardiac output (CO): is derived by the multiplication of SV x heart rate (HR) (NR:  $> 4$  L/min);
  - c. Stroke volume index (SVI): is a derived parameter obtained by the adjustment of SV by body surface area (BSA) (NR: 33-47 mL/beat/m<sup>2</sup>); and
  - d. Cardiac index (CI): is a derived parameter obtained by the adjustment of CO by body surface area (BSA) (NR: 2.5-4 L/min/m<sup>2</sup>).
5. Right ventricle function parameters:
- a. Pulmonary arterial systolic pressure (PASP): estimated by the sum of the peak velocity at tricuspidal continuous wave Doppler and a fixed value

- derived from inferior vena cava diameter and respiratory change. (NR<35 mmHg);
- b. Tricuspid annular plane systolic excursion (TAPSE): measured from M-mode echocardiography from apical four chamber view. TAPSE correlates with right ventricle ejection fraction and its reduction associated with worse prognosis in HF. (NR>16 mm) (see Ghio S et al., J Am Coll Cardiol 2001;37(1):183-8); and
  - c. Right ventricle Sa: measured with tissue Doppler method at right ventricle free wall from apical four chamber view. Sa is a derived parameter of systolic function and correlated with right ventricle ejection fraction. (NR >10 cm/s) (see Voelkel NF et al., Circulation 2006;114(17):1883-91; Haddad F et al., Circulation 2008;117(13):1717-31).
6. Other parameters:
- a. Mitral regurgitation (MR): evaluated with a visual qualitative assessment and graded in four categories: none, mild, moderate, and severe (see Lancellotti P et al., Eur J Echocardiogr 2010;11(4):307-32); and
  - b. Inferior vena cava diameter (IVC): measured with M-mode echocardiography from subcostal view at 1-2 cm from the junction with right atrium. This parameter has been used to estimate systolic pulmonary artery pressure. It correlated with right atrium pressure indicating the grade of congestion. Increased IVC diameter is associated with prognosis in patients with HF (NR: ≤1.5 cm) (see Pellicori P et al., JACC Cardiovasc Imaging 2013;6(1):16-28; Voelkel NF et al., Circulation 2006;114(17):1883-91).

**Table 1A. Cardiac changes at 6 hours of infusion.**

| Parameter                   | Ista 0.5<br>µg/kg/min | Ista 1.0<br>µg/kg/min | Placebo      | p Ist 0.5 | p Ist 1.0    |
|-----------------------------|-----------------------|-----------------------|--------------|-----------|--------------|
| <b>LAA (cm<sup>2</sup>)</b> | -0.33 1.885           | -0.84 2.421           | -0.52 1.840  | 0.663     | 0.521        |
| <b>LAV (ml)</b>             | -1.16 11.265          | -4.92 14.390          | -3.24 10.047 | 0.399     | 0.558        |
| <b>Diastolic function</b>   |                       |                       |              |           |              |
| <b>E wave (cm/s)</b>        | -3.33 14.764          | -9.13 17.464          | -1.10 10.990 | 0.451     | <b>0.018</b> |
| <b>A wave (cm/s)</b>        | 1.76 13.572           | 5.56 17.280           | 1.55 10.368  | 0.955     | 0.348        |
| <b>EDT (ms)</b>             | 6.83 47.921           | 20.37 50.849          | -0.18 38.939 | 0.491     | 0.052        |

|                          |        |        |        |       |        |       |              |              |
|--------------------------|--------|--------|--------|-------|--------|-------|--------------|--------------|
| <b>E/A ratio</b>         | -0.286 | 0.866  | -0.317 | 0.898 | -0.124 | 0.866 | 0.566        | 0.458        |
| <b>e' (cm/s)</b>         | 0.61   | 1.010  | 0.01   | 1.154 | 0.25   | 1.167 | 0.146        | 0.376        |
| <b>E/e' ratio</b>        | -3.183 | 5.628  | -2.028 | 3.652 | -0.740 | 3.994 | <b>0.032</b> | 0.150        |
| <b>Systolic function</b> |        |        |        |       |        |       |              |              |
| <b>Sa (cm/s) Left V</b>  | 0.613  | 1.035  | 0.908  | 0.936 | 0.197  | 0.919 | 0.065        | <b>0.001</b> |
| <b>S (cm/s) Right V</b>  | 1.25   | 2.185  | 2.00   | 1.907 | 0.43   | 1.441 | 0.125        | <b>0.003</b> |
| <b>Cardiac function</b>  |        |        |        |       |        |       |              |              |
| <b>CO (l/min)</b>        | 0.385  | 0.843  | 0.228  | 0.760 | 0.083  | 0.705 | 0.094        | 0.390        |
| <b>CI (l/min/m2)</b>     | 0.209  | 0.445  | 0.140  | 0.434 | 0.042  | 0.400 | 0.090        | 0.309        |
| <b>SV (ml/beat)</b>      | 7.724  | 11.752 | 7.269  | 8.134 | 2.405  | 7.244 | <b>0.020</b> | <b>0.007</b> |
| <b>SVI(ml/beat/m2)</b>   | 4.198  | 6.218  | 4.187  | 4.641 | 1.317  | 4.077 | <b>0.019</b> | <b>0.005</b> |

**Table 1B. Cardiac changes at 24 hours of infusion.**

| Parameter          | Ista 0.5<br>µg/kg/min |        | Ista 1.0<br>µg/kg/min |        | Placebo | p Ista 0.5 | p Ista 1.0   |                  |
|--------------------|-----------------------|--------|-----------------------|--------|---------|------------|--------------|------------------|
| LAA (cm²)          | -1.70                 | 2.463  | -2.56                 | 2.972  | -0.31   | 1.886      | <b>0.008</b> | <b>&lt;0.001</b> |
| LAV (ml)           | -7.94                 | 13.269 | -13.81                | 17.198 | -2.95   | 10.624     | 0.079        | <b>0.002</b>     |
| Diastolic function |                       |        |                       |        |         |            |              |                  |
| E wave (cm/s)      | -8.14                 | 17.640 | -14.24                | 24.416 | -4.16   | 12.502     | 0.267        | <b>0.031</b>     |
| A wave (cm/s)      | 5.13                  | 10.990 | 11.10                 | 16.498 | -1.14   | 7.580      | <b>0.045</b> | <b>0.003</b>     |
| EDT (ms)           | 12.88                 | 54.954 | 9.79                  | 52.212 | 4.58    | 35.759     | 0.456        | 0.628            |
| E/A ratio          | -0.647                | 0.812  | -0.722                | 1.068  | -0.164  | 0.833      | <b>0.005</b> | <b>0.004</b>     |
| e' (cm/s)          | 0.94                  | 1.089  | 0.91                  | 1.792  | 0.26    | 1.23       | <b>0.013</b> | 0.635            |
| E/e' ratio         | -4.548                | 4.754  | -3.191                | 2.623  | -1.285  | 3.351      | <b>0.001</b> | <b>0.011</b>     |
| Systolic function  |                       |        |                       |        |         |            |              |                  |
| Sa (cm/s) Left V   | 0.679                 | 0.907  | 0.803                 | 1.023  | 0.171   | 1.029      | <b>0.024</b> | <b>0.012</b>     |

|                                   |              |             |             |       |              |
|-----------------------------------|--------------|-------------|-------------|-------|--------------|
| <b>S (cm/s) Right V</b>           | 1.10 1.780   | 1.55 2.012  | 0.15 1.424  | 0.052 | <b>0.015</b> |
| <b>Cardiac function</b>           |              |             |             |       |              |
| <b>CO (l/min)</b>                 | 0.486 0.696  | 0.239 0.786 | 0.195 0.651 | 0.073 | <b>0.797</b> |
| <b>CI (l/min/m<sup>2</sup>)</b>   | 0.264 0.369  | 0.140 0.443 | 0.116 0.373 | 0.097 | <b>0.808</b> |
| <b>SV (ml/beat)</b>               | 9.725 12.192 | 9.339 8.875 | 4.039 6.259 | 0.015 | <b>0.005</b> |
| <b>SVI(ml/beat/m<sup>2</sup>)</b> | 5.333 6.664  | 5.318 4.955 | 2.336 3.516 | 0.019 | <b>0.005</b> |

**Table 1C. Cardiac changes a 48 hours of infusion.**

| <b>Parameter</b>                   | Ista 0.5<br>µg/kg/min | Ista 1.0<br>µg/kg/min | Placebo      | p Ist 0.5    | p Ist 1.0    |
|------------------------------------|-----------------------|-----------------------|--------------|--------------|--------------|
| <b>Diastolic function</b>          |                       |                       |              |              |              |
| <b>E/A ratio</b>                   | -0.278 1.043          | -0.356 0.967          | -0.172 0.741 | 0.729        | 0.509        |
| <b>E/e' ratio</b>                  | -2.570 3.654          | -2.218 3.193          | -1.800 4.013 | 0.388        | 0.629        |
| <b>Systolic function</b>           |                       |                       |              |              |              |
| <b>LVEF (%)</b>                    | 1.28 3.693            | 2.06 4.975            | 1.24 3.539   | 0.968        | 0.427        |
| <b>Sa (cm/s)</b>                   | 0.224 0.811           | 0.257 0.771           | 0.132 0.898  | 0.640        | 0.525        |
| <b>Cardiac function</b>            |                       |                       |              |              |              |
| <b>CO (l/min)</b>                  | 0.350 0.788           | 0.368 0.735           | -0.017 0.691 | <b>0.041</b> | <b>0.026</b> |
| <b>CI (l/min/m<sup>2</sup>)</b>    | 0.191 0.426           | 0.205 0.407           | -0.007 0.384 | <b>0.044</b> | <b>0.027</b> |
| <b>SV (ml/beat)</b>                | 4.815 9.832           | 5.203 8.078           | 1.787 7.108  | 0.139        | 0.062        |
| <b>SVI (ml/beat/m<sup>2</sup>)</b> | 2.762 5.603           | 2.913 4.572           | 1.039 4.026  | 0.139        | 0.071        |

[74] The quantitative determination of PST 2744 and its metabolite PST 2915 in human plasma was determined by the HPLC-MS/MS method that included a mobile phase of 70:30

acetonitrile/water, 1mL/L 1M formic acid, and 1mL/L 5M ammonium acetate. The flow rate was 1mL/min, and the chromatographic separation was by reversed phase HPLC (Column: SYNERGI 4 $\mu$  POLAR-RP80A 150 x 4.6mm equipped with a Security-guard Phenomenex Polar-RP 4x3mm). Detection was performed by MS/MS, and the acquisition mode was Multiple Reaction Monitoring (MRM).

[75] The quantitative determination of istaroxime metabolites PST 2922 and PST 3093 in human plasma was also measured by the HPLC-MS/MS method. In this case, the mobile phase was 50:50 H<sub>2</sub>O/CH<sub>3</sub>CN (v/v) and 500 $\mu$ L/L 98-100% HCOOH. The flow rate was 1mL/min and chromatographic separation was done by reversed-phase HPLC (Column: Phenomenex Phenyl hexyl, 150x4.6mm, equipped with a Phenomenex phenyl propyl guard-cartridge) under isocratic conditions. Detection was performed by MS/MS (376.0 $\rightarrow$  282.0amu for PST2922, 378.0 $\rightarrow$  284.0 amu for PST 3093 and 362.0 $\rightarrow$  268.0 amu for PST 3418, IS).

[76] The data reported in **Table 1A** clearly indicates that infusion at either 0.5  $\mu$ g/kg/min or 1  $\mu$ g/kg/min istaroxime does not significantly improve most of the altered echo parameters of diastolic function (E, A waves, E/A ratio), with significant reduction occurring only with E/e' at 0.5  $\mu$ g/kg/min. The systolic function (Sa and S wave) is improved at 1  $\mu$ g/kg/min.

[77] Surprisingly, when istaroxime was infused for 24 hours, a clear and statistically significant improvement is observed for most of the diastolic function parameters (E and A waves, E/A and E/e' ratios), while the positive effect of systolic function (Sa and S wave) is maintained but does not continue to increase (see **Table 1B**). At 48 hours, both the CO and CI changes are still significantly increased over placebo (see **Table 1C**). The shift from increased changes of SV and SVI at 6 and 24 hours to changes in CO and CI at 48 hours is also favoured by the normalization of the decreased HR at 6 and 24 hours, which was demonstrated in the previous Horizon study and confirmed in the present study. The changes of the other indexes of cardiac relaxation and contraction are still present, but do not achieve the statistical significance.

[78] **Figures 2A-2C** show the plasma concentrations of istaroxime and its metabolites in Caucasian and Chinese patients during and subsequent to infusion with 0.5 or 1  $\mu$ g/kg/min of istaroxime for 24 hours. During the infusion period, both istaroxime (2744) and istaroxime metabolite PST 2922 remain relatively constant and are rapidly cleared after infusion is stopped. On the other hand, istaroxime metabolites PST 2915 and PST 3093 continue to accumulate in the plasma throughout the infusion period with the average concentration of PST 3093 exceeding 60 ng/mL by the end of the infusion period and with

average concentration levels exceeding 10 ng/mL even at 70 hours, or 46 hours post-infusion.

[79] Notably, at 48 hours of infusion, the plasma levels of istaroxime are not detectable after 20 hours, while those of metabolite PST 3093 average about 25.02 ng/ml in Caucasian patients after 0.5 µg/kg/ml istaroxime infusion, about 12.5 ng/ml in Chinese patients after 0.5 µg/kg/ml istaroxime infusion, and about 21.2 ng/ml with 1 µg/kg/ml istaroxime infusion (see **Figures 2A-2C**). These concentrations are much higher than the concentrations of 3093 exhibiting SERCA2a-stimulatory activity in SR vesicles from normal canine heart as shown in **Table 3**. Moreover, according to Ferrandi M et al. (BJP 2013; 169:1849-1861), istaroxime exerts its maximum SERCA2a activation in SR vesicles from failing canine hearts at concentrations that are much lower (about 10 times) than those able to stimulate this activity in SR vesicles from healthy canine heart. Therefore, it is likely that this remarkable difference between normal and failing heart SR vesicles may also occur for PST 3093, which may continue to maintain its SERCA2a stimulatory activity even at the lower concentrations detected at 72 hours (see **Figures 2A-2C**), where no echocardiographic data are available.

[80] While not intending to be bound by theory, the above-discussed observation is consistent with the hypothesis that a pure SERCA2a activator may improve cardiac pump function. Finally this change in efficacy of cardiac pump function is not associated with any significant changes in plasma level of Hs TnT, which is considered by cardiologists as the most reliable biomarker of myocardial damage. This lack of Hs TnT change is likely due the activation of SERCA2a that, by reducing the cardiomyocytes plasma  $Ca^{2+}$  concentration, also minimizes the cardiomyocytes damage. At present, the stimulation of the cardiac pumping ability by the only available inotropic agent under development (omecamtiv) is associated with an increase in the plasma levels of Hs TnT, and different developmental strategies are under study to detect the dose that minimize these changes in plasma HsTnT (Teerlink J R et al., 2016 Lancet 388, 2895-903).

[81] The present invention may provide the basis for planning appropriate trials aimed at assessing whether different ratios between plasma levels of PST 3093 and istaroxime, achievable by varying the dose and duration of istaroxime infusion, may furnish greater therapeutic benefits to patients with HFpEF or HFmEF than to patients with HFReF, or in patients with or without Echocardiographic indexes revealing a status of diastolic impairment; thus increasing the precision of the therapeutic approach to AHF.

Pharmaceutical compositions

[82] Pharmaceutical compositions and formulations for intravenous infusion comprising istaroxime or a metabolite thereof in admixture with at least one conventional pharmaceutically acceptable carrier and/or vehicle and/or excipient are commonly known in the art.

[83] The pharmaceutical compositions and formulations for intravenous infusion can be formulated in any way and can be administered in a variety of unit dosage forms depending upon the condition or disease and the degree of illness, the general medical condition of each patient, the resulting preferred method of administration and the like. Details on techniques for formulation and administration are well described in the scientific and patent literature, see, *e.g.*, the latest edition of Remington's Pharmaceutical Sciences, Mack Publishing Co, Easton PA ("Remington's").

[84] The formulations may conveniently be presented in unit dosage form and may be prepared by any method known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier or vehicle material to produce a single dosage form will vary depending upon the subject being treated and the particular mode of administration. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be the amount of the compound which produces a therapeutic effect.

[85] Pharmaceutical formulations as provided herein can be prepared according to any method known to the art for the manufacture of pharmaceuticals. Such formulations can contain additional agents, such as preserving or stabilizing agents. A formulation can be admixed with nontoxic pharmaceutically acceptable carriers or excipients which are suitable for manufacture. Formulations may comprise one or more diluents, emulsifiers, preservatives, buffers, excipients, etc. and may be provided in such forms as liquids, powders, emulsions, lyophilized powders, etc.

[86] Aqueous suspensions can contain an active agent (*e.g.*, a composition used to practice the uses and methods as provided herein) in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl-methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally occurring phosphatide (*e.g.*, lecithin), a condensation product of an alkylene oxide with a fatty acid (*e.g.*, polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (*e.g.*,

heptadecaethylene oxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol (e.g., polyoxyethylene sorbitol mono-oleate), or a condensation product of ethylene oxide with a partial ester derived from fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan mono-oleate). The aqueous suspension can also contain one or more preservatives such as ethyl or n-propyl p-hydroxybenzoate. Formulations can be adjusted for osmolarity.

[87] According to the present invention, istaroxime is given by intravenous (IV) administration. These formulations can comprise a solution of active agent dissolved in a pharmaceutically acceptable carrier. Acceptable vehicles and solvents that can be employed are water, dextrose in water, and Ringer's solution, an isotonic sodium chloride. These solutions are sterile and generally free of undesirable matter. These formulations may be sterilized by conventional, well known sterilization techniques. The formulations may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents, e.g., sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate and the like. The concentration of active agent in these formulations can vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight, and the like, in accordance with the particular mode of administration selected and the patient's needs. The administration is by bolus or continuous infusion (e.g., substantially uninterrupted introduction into a blood vessel for a specified period of time).

[88] Istaroxime as provided herein can be lyophilized. Provided herein is a stable lyophilized formulation comprising a composition as provided herein, which can be made by lyophilizing a solution comprising a pharmaceutical as provided herein and a bulking agent, e.g., mannitol, trehalose, raffinose, and sucrose or mixtures thereof. There are many other conventional lyophilizing agents. Among the sugars, lactose is the most common. Also used are citric acid, sodium carbonate, EDTA, Benzyl alcohol, glycine, sodium chloride, etc. (see, for example, Journal of Excipients and Food Chemistry Vol. 1, Issue 1 (2010) pp 41-54; U.S. patent app. no. 20040028670). In a preferred embodiment, istaroxime can be prepared as powder for injection according to the teaching of CN103315968.

[89] According to the present invention, istaroxime as provided herein can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, compositions are administered to a subject already suffering from a condition, or disease in an amount sufficient to treat, prevent, cure, alleviate or partially arrest the clinical manifestations of the condition, or disease and its complications (i.e., a "therapeutically

effective amount"). For example, in alternative embodiments, pharmaceutical compositions as provided herein are administered in an amount sufficient to treat, prevent or ameliorate in an individual in need thereof. The amount of pharmaceutical composition adequate to accomplish this is defined as a "therapeutically effective dose." The dosage schedule and amounts effective for this use, *i.e.*, the "dosing regimen," will depend upon a variety of factors, including the stage of the disease or condition, the severity of the disease or condition, the general state of the patient's health, the patient's physical status, age and the like. In calculating the dosage regimen for a patient, the mode of administration also is taken into consideration.

[90] The dosage regimen also takes into consideration pharmacokinetics parameters well known in the art, *i.e.*, the active agents' bioavailability, metabolism, clearance, and the like (see, e.g., Hidalgo-Aragones J., *Steroid Biochem. Mol. Biol.* 1996;58:611-617; Groning, *Pharmazie* 1996;51:337-341; Fotherby *Contraception* 1996;54:59-69; Johnson, *J. Pharm. Sci.* 1995;84:1144-1146; Rohatagi, *Pharmazie* 1995;50:610-613; Brophy, *Eur. J. Clin. Pharmacol.* 1983;24:103-108; the latest Remington's, *supra*). The state of the art allows the clinician to determine the dosage regimen for each individual patient, active agent and disease or condition treated. Guidelines provided for similar compositions used as pharmaceuticals can be used as guidance to determine the dosage regimen, *i.e.*, dose schedule and dosage levels, administered practicing the methods as provided herein are correct and appropriate.

[91] Single or multiple administrations of formulations can be given depending on the dosage and frequency as required by the AHF clinical symptoms of patient. The formulations should provide a sufficient quantity of active agent to effectively treat or prevent or ameliorate a conditions, diseases or symptoms as described herein. A correct treatment of AHF, by selectively normalizing a depressed biochemical activity underlying the symptoms of subset of patients (HFpEF or HFmEF), may be expected to selectively improve the symptoms and to reduce the incidence of unwanted side effects produced by the available drugs either during hospital staying or after discharge. The term prevention is applicable when the continuous monitoring of the pulmonary pressure is possible with the appropriate chronic implantable devices that furnish and estimation of PCWP. In this condition a significant increase in the PCWP may precede the appearance of symptoms of AHF, thus providing the rational to infuse the Istaroxime at the right dose to prevent the symptoms and the consequent hospitalization

[92] In one embodiment, an effective amount of istaroxime or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof, administered to an individual

in need thereof comprises use of various dosing schedules, e.g.: from about 0.1 µg/kg/min to about 3.0 µg/kg/min, e.g., 0.1 µg/kg/min, 0.15 µg/kg/min, 0.2 µg/kg/min, 0.25 µg/kg/min, 0.3 µg/kg/min, 0.35 µg/kg/min, 0.4 µg/kg/min, 0.5 µg/kg/min, 0.6 µg/kg/min, 0.7 µg/kg/min, 0.8 µg/kg/min, 0.9 µg/kg/min, 1.0 µg/kg/min, 1.1 µg/kg/min, 1.2 µg/kg/min, 1.3 µg/kg/min, 1.4 µg/kg/min, 1.5 µg/kg/min, 1.6 µg/kg/min, 1.7 µg/kg/min, 1.8 µg/kg/min, 1.9 µg/kg/min, 2.0 µg/kg/min, 2.1 µg/kg/min, 2.2 µg/kg/min, 2.3 µg/kg/min, 2.4 µg/kg/min, 2.5 µg/kg/min, 2.6 µg/kg/min, 2.7 µg/kg/min, 2.8 µg/kg/min, 2.9 µg/kg/min, or 3.0 µg/kg/min. For instance, in some embodiments, istaroxime or its metabolite (e.g., PST 3093), is administered by infusion at an effective dose from about 0.2 µg/kg/min to about 2.0 µg/kg/min, or from about 0.2 µg/kg/min to about 1.5 µg/kg/min, or from about 0.25 µg/kg/min to about 1.0 µg/kg/min, or from about 0.5 µg/kg/min to about 1.0 µg/kg/min.

[93] In alternative embodiments, an effective amount of istaroxime or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof, administered to an individual in need thereof is individualized based on basal levels and subsequent changes of certain heart function parameters, such as echo indexes or Pulmonary Capillary Wedge Pressure (PCWP), dyspnea, peripheral and pulmonary venous congestion, urinary volume, serum biomarkers such as NT-proBNP and high sensitive cardiac Troponin (hs-cTnT).

[94] In alternative embodiments, an effective amount is demonstrated by reduction of PCWP, orthopnea, paroxysmal nocturnal dyspnea, reduction of peripheral and pulmonary venous congestion, such as pulmonary crepitations or rales, reduction of ankle swelling, reduction of biomarkers urinary output such as NT-proBNP and high sensitive cardiac Troponin (hs-cTnT).

[95] In alternative embodiments, an effective amount of istaroxime or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof, administered to an individual in need thereof is individualized based on basal levels and subsequent changes of certain heart function parameters, such as echo indexes or Pulmonary Capillary Wedge Pressure (PCWP), dyspnea, peripheral and pulmonary venous congestion, urinary volume, serum biomarkers such as NT-proBNP and high sensitive cardiac Troponin (hs-cTnT).

[96] In alternative embodiments, an effective amount is demonstrated by reduction of PCWP, orthopnea, paroxysmal nocturnal dyspnea, reduction of peripheral and pulmonary venous congestion, such as pulmonary crepitations or rales, reduction of ankle swelling, reduction of biomarkers urinary output such as NT-proBNP and high sensitive cardiac Troponin (hs-cTnT).

Methods of treatment

[97] Also provided herein are methods of treating an individual with heart failure. In preferred embodiments, the individual exhibits symptoms of, or has been diagnosed with, acute heart failure. While the individual can be a non-human animal, in a preferred embodiment, the individual is a human patient, such as a human patient suffering from heart failure.

[98] In general, the compositions described herein can be used to treat the individual having heart failure or acute heart failure. In an embodiment, the method of therapy includes providing or presenting the individual having heart failure or acute heart failure. In some cases, a measuring step is first carried out to determine the baseline heart function of the individual. For instance, an individual with heart failure may exhibit impaired or decreased diastolic relaxation function. The measuring step may include measuring one or more parameters of heart failure, such as, but not limited to, decreased heart rate, decreased heart pressure, decreased systolic and/or diastolic blood pressure, reduced left ventricular end-diastolic/systolic volume and function (LVEF), or increased E/Ea or E/A ratios reduced Ea ratio decreased stroke volume. As one having ordinary skill in the art will appreciate, any suitable measuring technique available in the art at the time of the measuring step is suitable for use herein, and it is well within the purview of such skilled artisan to select an appropriate measuring technique corresponding to the parameter of interest. A non-limiting list of suitable measuring equipment/techniques includes echocardiogram, cardiac catheterization, nuclear stress test, CAT scan, radionuclide ventriculography scan, stethoscope, sphygmomanometer, and the like. For instance, the diastolic relaxation can be measured by echocardiography or PCWP.

[99] The methods disclosed herein also include administering to the individual a therapeutically effective amount of istaroxime or a metabolite thereof, such as PST 3093. In preferred embodiments, the istaroxime or istaroxime metabolite is in a pharmaceutical composition, such as any one of the combinations discussed above. The istaroxime or istaroxime metabolite is administered in an therapeutically effective dose as disclosed elsewhere herein, e.g., between about 0.25 µg/kg/min to about 1.0 µg/kg/min. In a more preferred embodiment, the route of administration is infusion, such as intravenous fusion. The measuring step can be performed before, during, or after the administering step. For instance, it may be desired to continually monitor one or more of the parameters of heart function during treatment and for a period of time thereafter.

[100] As discussed above, it has been surprisingly discovered that administering istaroxime (or its metabolites) by infusion for an infusion duration of greater than 6 hours,


e.g., 6.1 h, 6.2 h, 6.3 h, 6.4 h, 6.5 h, 6.6 h, 6.7 h, 6.8 h, 6.9 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19 h, 20 h, 21 h, 22 h, 23 h, 24 h, 25 h, 26 h, 27 h, 28 h, 29 h, 30 h, 31 h, 32 h, 33 h, 34 h, 35 h, 36 h, 37 h, 38 h, 39 h, 40 h, 41 h, 42 h, 43 h, 44 h, 45 h, 46 h, 47 h, or 48 h or more, results in increased SERCA2a activity and improving diastolic relaxation without causing arrhythmogenic effects due to, for example,  $\text{Na}^+/\text{K}^+$  pump inhibition. In this manner, istaroxime infusion for greater than 6 hours exerts a lusitropic-SERCA2a activity that is prevailing on the inotropic activity and results in improved diastolic relaxation as compared to istaroxime infusion for less than 6 hours.

[101] While not intending to be bound by theory, it is believed that this later-arising “pure” SERCA2a activation is due to an accumulation of istaroxime metabolites in the plasma of the individual. As such, in some embodiments, istaroxime is administered via intravenous infusion for a period of time sufficient to enable the accumulation of istaroxime metabolites in the plasma of the individual. In preferred embodiments, the infusion duration is sufficient to allow for the accumulation of one or more istaroxime metabolites; preferably, the metabolite is PST 2915 having the structural formula (II) or PST 3093 having the structural formula (III); more preferably, the metabolite is PST 3093. In some embodiments, the accumulation of istaroxime metabolite in the plasma is at a concentration of least about 3 ng/mL, e.g., 3 ng/mL, 4 ng/mL, 5 ng/mL, 6 ng/mL, 7 ng/mL, 8 ng/mL, 9 ng/mL, 10 ng/mL, 11 ng/mL, 12 ng/mL, 13 ng/mL, 14 ng/mL, 15 ng/mL, 16 ng/mL, 17 ng/mL, 18 ng/mL, 19 ng/mL, 20 ng/mL, 21 ng/mL, 22 ng/mL, 23 ng/mL, 24 ng/mL, 25 ng/mL, 26 ng/mL, 27 ng/mL, 28 ng/mL, 29 ng/mL, 30 ng/mL, 35 ng/mL, 40 ng/mL, 45 ng/mL, 50 ng/mL or more for a period of time of at least about 3 hours, e.g., 3 h, 4 h, 5 h, 6 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19 h, 20 h, 21 h, 22 h, 23 h, 24 h or more. In one embodiment, the istaroxime metabolite accumulates in the plasma to the desired concentration within 6 hours; preferably, within 3 hours or within 2 hours or within 1 hour of istaroxime infusion initiation and is maintained at or above that concentration for at least about 3 additional hours, e.g., 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or more additional hours; preferably, for at least 6 additional hours; more preferably for at least about 12 additional hours. In some embodiments, the desired plasma concentration of metabolite is at least about 5 ng/mL. In another embodiment, the istaroxime metabolite accumulates in the plasma to a concentration of at least about 10 ng/mL and is maintained at or above that concentration for at least about 6 additional hours; preferably for at least about 12 additional hours. In yet another embodiment, the istaroxime metabolite accumulates in the plasma to a concentration of at least about 15 ng/mL and is maintained at or above that concentration for at least about 6 additional hours; preferably for at least about 12 additional hours. In some embodiments, the plasma concentration of the metabolite

remains above 5 ng/mL for at least about 6 hours following the completion of the istaroxime administration. In some embodiments, the plasma concentration of the metabolite remains above 10 ng/mL for at least about 6 hours following the completion of the istaroxime administration. In yet other embodiments, the plasma concentration of the metabolite remains above 20 ng/mL for at least about 6 hours following the completion of the istaroxime administration. In others, the concentration remains at least about 30 ng/mL, 40 ng/mL, or 50 ng/mL for an additional 6 hours or more, *e.g.*, 6 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19, 20 h, 21 h, 22 h, 23 h, 24 h or more following istaroxime infusion. For instance, the metabolite accumulation may remain at a concentration level of at least about 10 ng/mL for an additional 12 hours or more, *e.g.*, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19, 20 h, 21 h, 22 h, 23 h, 24 h, 25 h, 26 h, 27 h, 28 h, 29 h, 30 h, 31 h, 32 h, 33 h, 34 h, 35 h, 36 h, 37 h, 38 h, 39 h, 40 h, or more following the istaroxime infusion.


[102] As one having ordinary skill in the art would appreciate, the plasma concentration of istaroxime or istaroxime metabolite can be measured by conventional means, such as by HPLC-MS/MS.

[103] In some embodiments, the istaroxime metabolites have the structural formulas (II) or (III):



**PST 2915**

Formula (II)

**PST 3093**

Formula (III)

[104] In preferred embodiments, the metabolite of istaroxime is the compound PST 3093.

[105] In a further example, an individual is diagnosed with heart failure or acute heart failure, and is administered about 0.25 µg/kg/min to about 1.0 µg/kg/min of istaroxime for a period of time that is greater than 6 hours. As the istaroxime is metabolized by the individual, istaroxime metabolites, such as PST 3093, begin to accumulate in the plasma of the individual. For instance, istaroxime metabolite PST 3093 may accumulate to a plasma concentration of at least about 5 ng/mL within 3 hours of istaroxime infusion and is maintained at a plasma level of at least about 5ng/mL for the duration of the istaroxime infusion and for an additional 6 to about 36 hours. The presence of the PST 3093 acting as a "pure" SERCA2a activator confers to the individual improved diastolic relaxation.

[106] In some embodiments, the method of treatment may include administration of the istaroxime metabolite by infusion in combination with or instead of istaroxime. For instance, a method of treating an individual with heart failure is disclosed that includes administering to the individual a therapeutically effective amount of a pharmaceutical composition that comprises a pharmaceutically acceptable carrier and an istaroxime metabolite having the formula (II) or (III). In preferred embodiments, PST 3093 or PST 2915 is the metabolite; more preferably, it is PST 3093. Such methods may include a measuring step, wherein one or more parameters of heart function are measured using measuring techniques available in the art. The measuring step may be performed prior to, during, or subsequent to the administration of the pharmaceutical composition. In such methods, the therapeutically effective dose of PST 3093 will be between about 0.2 µg/kg/min to about 2.0 µg/kg/min; preferably between about 0.3 µg/kg/min and about 1.5 µg/kg/min; more preferably between about 0.5 µg/kg/min and about 1.0 µg/kg/min. In such methods, the duration of infusion will be greater than 6 hours, e.g., 6.1 h, 6.2 h,

6.3 h, 6.4 h, 6.5 h, 6.6 h, 6.7 h, 6.8 h, 6.9 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19 h, 20 h, 21 h, 22 h, 23 h, 24 h, 36 h, 48 h, or more.

[107] In some embodiments, it is desired to treat an individual with acute heart failure with two or more istaroxime dose regimens to initially treat the acute heart failure symptoms and subsequently normalize the heart functional defects associated with a more chronic heart failure condition. In such embodiments, the dosing regimen is manipulated to control the istaroxime to metabolite ratio by starting with the administration by infusion of a higher dose of istaroxime to treat acute heart failure symptoms followed by administering a lower istaroxime dose by infusion for a longer duration to allow for the accumulation of istaroxime metabolites. In this manner, the initial infusion will allow for rapid SERCA2a stimulation and Na,K-ATPase inhibition resulting in a rapid positive inotropy to treat the acute heart failure symptoms, while the second infusion will allow for accumulation of the selective SERCA2a-activating istaroxime metabolite PST 3093 and extended positive inotropy at a reduced risk of arrhythmogenic  $\text{Ca}^{2+}$  triggering events.

[108] Therefore, provided herein is a method of treating an individual having acute heart failure, wherein it is administered to the individual by infusion a first pharmaceutical composition that includes a pharmaceutically acceptable carrier and istaroxime at a therapeutic dose of at least about 1.5  $\mu\text{g/kg/min}$ . The heart function of the individual can be measured and monitored before, during, and/or after beginning infusion of the first pharmaceutical composition using any of the techniques discussed herein. For instance, in some embodiments, one or more parameters of heart function are measured prior to administering istaroxime in order to determine, *e.g.*, whether the individual is presenting with HFpEF or HFmEF. In other embodiments, initial measurements may include echocardiogram or PCWP values to measure diastolic relaxation dysfunction. These measurements may also be initiated concurrently with administering the istaroxime by infusion and/or may be continued throughout the duration of the infusion. In some embodiments, the step of measuring one or more parameters of heart function can be performed after administering the first pharmaceutical composition.

[109] In the dosing manipulation methods, once an improvement in the parameters of heart function are measured, a pharmaceutical composition that comprises a pharmaceutically acceptable carrier and istaroxime at a lower therapeutic dose is administered by infusion. For instance, an improvement in diastolic relaxation in the individual as measured by echocardiogram or PCWP as compared to the same measurements taken prior to and/or at the start of the initial infusion would indicate treatment of the acute heart failure symptoms. Also, the reduction of acute heart failure

symptoms such as breathlessness, ankle swelling, elevated jugular venous pressure, pulmonary crackles and peripheral edema may justify the change in the infusion rate. In some embodiments, the second therapeutic dose of istaroxime is less than about 1.5  $\mu\text{g/kg/min}$ ; preferably between about 0.3  $\mu\text{g/kg/min}$  and about 1.5  $\mu\text{g/kg/min}$ ; more preferably between about 0.5  $\mu\text{g/kg/min}$  and about 1.0  $\mu\text{g/kg/min}$ . For instance, in one particular embodiment, the second therapeutic dose of istaroxime is about 0.5  $\mu\text{g/kg/min}$ . Administering by infusion of the second therapeutic dose of istaroxime may then be continued for a duration of greater than 6 hours, *e.g.*, 6.1 h, 6.2 h, 6.3 h, 6.4 h, 6.5 h, 6.6 h, 6.7 h, 6.8 h, 6.9 h, 7 h, 8 h, 9 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 17 h, 18 h, 19 h, 20 h, 21 h, 22 h, 23 h, 24 h, 36 h, 48 h, or more.

[110] In some embodiments, the second, lower dose of istaroxime may be infused for a duration sufficient to produce an accumulated plasma concentration of an istaroxime metabolite, such as PST 3093. In such embodiments, infusion of the lower dose of istaroxime is continued until the plasma concentration of PST 3093 is at least about 20 ng/mL. In other embodiments, infusion of the lower dose of istaroxime is continued until the plasma concentration of PST 3093 is at least about 30 ng/mL. In still other embodiments, infusion is continued until the plasma concentration of PST 3093 is at least about 40 ng/mL or at least about 50 ng/mL or at least about 60 ng/mL. Once infusion of the lower dose of istaroxime is stopped, the istaroxime compound is cleared from the individual while the istaroxime metabolite exhibiting selective or "pure" SERCA2a activation (*i.e.*, 3093) remains in the individual's bloodstream for extended periods of time to confer to the individual improved heart function with a much lower risk of arrhythmogenic triggering events.

[111] In alternative embodiments, in evaluating the efficacy of a treatment, a treatment regimen or a particular dosage, or to determine if a treatment versus a maintenance dosage should be given, individuals, *e.g.*, patients affected by acute or chronic heart failure, are subject to regular periodic screening for the presence and extent of organ and tissue involvement or damage, *e.g.*, heart (ventricle dilatation, third heart sound cardiac hypertrophy), fatigue, tiredness, reduced exercise tolerance, increased time to recover after exercise, kidney (renal insufficiency, oliguria), lung (orthopnea, paroxysmal nocturnal dyspnea, tachypnea), ankle swelling, elevated jugular venous pressure. A thorough physical examination should be done at a time interval chosen by those experts in the treatment of a cardiovascular disease, in particular acute or chronic heart failure which would concentrate on cardiac, pulmonary and peripheral circulation functions. Accordingly, in alternative embodiments, therapy with istaroxime or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof as disclosed herein, is instituted as early as

possible, preferably in emergency, to prevent the rapid evolution of symptoms and continued after patient's discharge for years, preferably during the whole life of the patient or at least a period consistent with the way other drugs are used in heart failure. As the result of monitoring of patient's conditions by the medical doctor, istaroxime long infusion, longer than 6 hours may be given the patient up to once/twice a month in order to prevent occurrence of acute episodes of heart failure, thus avoiding emergency rescue of the patient and lessening the probability of life-threatening episodes.

[112] According to the present invention, uses and methods as provided herein can further comprise co-administration with other drugs or pharmaceuticals. In fact, the present invention selectively corrects a depressed cardiac biochemical function (namely the SERCA2a activity). This certainly contributes to relieving the existing HF clinical symptoms, with less unwanted side effects than those of the available therapies (just because the selectivity mentioned above). However, as CHF and AHF are complex clinical syndromes the present invention is potentially associable to existing and future drug classes and/or specific drugs such as: a) drug classes such as, ACE inhibitors, AIRBs, diuretics, Ca channel blockers, blockers, digitalis, NO donors, vasodilators, SERCA2a stimulators, neprilysin (NEP) inhibitors, myosin filament activators, recombinant relaxin-2 mediators, recombinant NP protein, activators of the soluble Guanylate Cyclase (sGC), beta-arrestin ligand of Angiotensin II receptor; b) specific drugs: hydrochlorothiazide, furosemide, verapamil, diltiazem, carvedilol, metoprolol, hydralazine, eplerenone, spironolactone, lisinopril, ramipril, nitroglycerin, nitrates, digoxin, valsartan, olmesartan, telmisartan, candesartan, losartan, entresto, omecamtiv, sacubitril, serelaxin, ularitide, levosimendan, cinaciguat. Subjects suffering from heart failure treated with the above drugs and undergoing regular clinical monitoring, for example having their pulmonary blood pressure continuously monitored with implanted probes, can be guarded in order to predict episode of AHF that may be prevented by the infusion of Istaroxime according to the present invention.

[113] Istaroxime as disclosed in the present invention, as used a therapeutic agent for treating acute heart failure, can be combined with other therapeutic agents used in the treatment of the same disease. Exemplary other therapeutic agents are diuretics, for example furosemide, bumetanide, and torasemide. Metolazone, an aldosterone antagonist, such as spironolactone or eplerenone; thiazide diuretics, such as Hydrochlorothiazide, metolazone, and chlorthalidone. Other agents are ACE inhibitors, for example Lisinopril and Ramipril. Also Angiotensin II receptor blockers (ARBs), such as valsartan, candesartan and losartan can be taken into consideration. Angiotensin receptor/neprilysin inhibitor (ARNI), sacubitril for example, are comprised. Other agents can be selected from Beta-

blockers, such as carvedilol and metoprolol for example, or Vasodilators, for example Hydralazine, optionally combined with isosorbide dinitrate, Nitrates, as nitroglycerin, amlodipine and felodipine; non-dihydropyridines such as diltiazem or verapamil. The compounds of the present invention can also be combined with Digoxin, if needed. Other drugs, as Ivabradine and other Anticoagulant may be considered.

[114] The compounds of the present invention can be combined with other therapeutic agents, in particular agents useful for treating cardiovascular diseases, more in particular in the combination therapy of heart failure. The combined active ingredients can be administered according to different protocols, decided by the medical doctor. According to an embodiment of the present invention, combination therapy can be carried out by administering istaroxime both at the same time or at different time of the further therapeutically active ingredient or ingredients. In case of concomitant administration, the compound of the present invention and the further active ingredient or ingredients can be each formulated in a respective pharmaceutical composition or in the same unitary dosage form. In the former case, the present invention provides a kit, in particular for the treatment of heart failure, comprising separate pharmaceutical compositions containing the compound of the present invention and the further active ingredient or ingredients, respectively. In another embodiment, the present invention provides a pharmaceutical unit dosage form kit, in particular for the treatment of acute heart failure, comprising compound of the present invention and the further active ingredient or ingredients in the same unit dosage form. Combination therapy according to the present invention provides advantageous treatment of heart failure due to the inotropic-lusitropic effect of istaroxime herein disclosed in addition to or synergically combined with the well-known therapeutic effect of the additional active agents herein disclosed.

[115] Also provided are nanoparticles, nanolipoparticles, vesicles and liposomal membranes comprising compounds used to practice the uses and methods as provided herein, *e.g.*, to deliver pharmaceutically active compounds and compositions as provided herein: istaroxime or a compound of formula (II) or formula (III) or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof, optionally combined with a further therapeutically active agent as disclosed above to a subject in need thereof. In alternative embodiments, these compositions are designed to target specific molecules, including biologic molecules, such as polypeptides, including cell surface polypeptides, *e.g.*, for targeting a desired cell type, *e.g.*, a myocyte or heart cell, an endothelial cell, and the like. A slow release of Istaroxime may provide a sufficient compound to selectively increase the plasma levels of the metabolite leaving the plasma levels of Istaroxime within very low ranges.

[116] Provided are multilayered liposomes comprising compounds used to practice methods as provided herein, *e.g.*, as described in Park, et al., U.S. application No. 20070082042. The multilayered liposomes can be prepared using a mixture of oil-phase components comprising squalane, sterols, ceramides, neutral lipids or oils, fatty acids and lecithins, to about 200 to 5000 nm in particle size, to entrap a composition used to practice uses and methods as provided herein.

[117] Liposomes can be made using any method, *e.g.*, as described in U.S. Patent No. 4,534,899; or Park, et al., U.S. application No. 20070042031, including method of producing a liposome by encapsulating an active agent according to the present invention (or a combination of active agents), the method comprising providing an aqueous solution in a first reservoir; providing an organic lipid solution in a second reservoir, and then mixing the aqueous solution with the organic lipid solution in a first mixing region to produce a liposome solution, where the organic lipid solution mixes with the aqueous solution to substantially instantaneously produce a liposome encapsulating the active agent; and immediately then mixing the liposome solution with a buffer solution to produce a diluted liposome solution.

[118] In one embodiment, liposome compositions used to practice uses and methods as provided herein comprise a substituted ammonium and/or polyanions, *e.g.*, for targeting delivery of istaroxime or an equivalent of a pharmaceutically acceptable salt, solvate or hydrate thereof used to practice methods as provided herein to a desired cell type, as described, *e.g.*, in U.S. application No. 20070110798.

[119] Provided are nanoparticles comprising compounds according to the present invention used to practice uses and methods as provided herein in the form of active agent-containing nanoparticles (*e.g.*, a secondary nanoparticle), as described, *e.g.*, in U.S. application No. 20070077286. In one embodiment, provided are nanoparticles comprising a fat-soluble active agent used to practice a use and method as provided herein or a fat-solubilized water-soluble active agent to act with a bivalent or trivalent metal salt.

[120] In one embodiment, solid lipid suspensions can be used to formulate and to deliver compositions used to practice uses and methods as provided herein to mammalian cells *in vivo*, *in vitro* or *ex vivo*, as described, *e.g.*, in U.S. application No. 20050136121.

[121] The compositions and formulations used to practice the uses and methods as provided herein can be delivered by the use of liposomes or nanoliposomes. By using liposomes, particularly where the liposome surface carries ligands specific for target cells, or are otherwise preferentially directed to a specific organ, one can focus the delivery of

the active agent into target cells in vivo. See, e.g., U.S. Patents Nos. 6,063,400; 6,007,839; Al-Muhammed, J. Microencapsul. 1996; 13:293-306; Chonn Curr. Opin. Biotechnol. 1995; 6:698-708; Ostro, Am. J. Hosp. Pharm. 1989; 46:1576-1587. A liposome formulation of istaroxime as disclosed in Eur J Pharm Biopharm. 2011;79(2):285-93 is also provided in the present invention.

#### Delivery vehicles


[122] In alternative embodiments, any delivery vehicle can be used to practice the uses and methods as provided herein, e.g., to deliver the compounds provided herein to a subject in need thereof. For example, delivery vehicles comprising polycations, cationic polymers and/or cationic peptides, such as polyethyleneimine derivatives, can be used e.g. as described, e.g., in U.S. application No. 20060083737.

[123] In one embodiment, a dried polypeptide-surfactant complex is used to formulate a composition used to practice a use and method as provided herein, e.g., as described in U.S. application No. 20040151766.

[124] In one embodiment, a composition used to practice uses and methods as provided herein can be applied to cells using vehicles with cell membrane-permeant peptide conjugates, e.g., as described in U.S. Patents No. 7,306,783; 6,589,503. In one aspect, the composition to be delivered is conjugated to a cell membrane-permeant peptide. In one embodiment, the composition to be delivered and/or the delivery vehicle are conjugated to a transport-mediating peptide, e.g., as described in U.S. Patent No. 5,846,743, describing transport-mediating peptides that are highly basic and bind to polyphosphoinositides.


[125] In one embodiment, electro-permeabilization is used as a primary or adjunctive means to deliver the composition to a cell, e.g., using any electroporation system as described e.g. in U.S. Patents No. 7,109,034; 6,261,815; 5,874,268.

[126] The following examples further illustrate the present invention.

**Example 1. Preparation of the compounds of formula (II) and (III)**Synthesis of PST 2915*Step 1: Hydroboration*


[127] To a solution of dehydroepiandrosterone **I** (30.0 g) in 450 mL of THF maintained under a nitrogen atmosphere and at a temperature of  $-10^{\circ}\text{C}$ . was added the complex BH.THF 1M in THF (260 mL). On completing the addition, the temperature was allowed to rise once again to ambient temperature; after 3 h 500 mL of  $\text{H}_2\text{O}$  were added and then  $\text{NaBO}_3 \cdot 4\text{H}_2\text{O}$  (31.4 g). The reaction was left to stir for one night. The precipitate formed was filtered, washed with THF and eliminated. The aqueous and organic phases were Separated, NaCl was added to the aqueous phase and this was re-extracted with THF (3x200 mL). The combined organic phases were anhydrified with NaCl and  $\text{Na}_2\text{SO}_4$  and evaporated under reduced pressure to obtain the crude product, which was crystallised by AcOEt/MeOH and then filtered and washed with AcOEt. Approximately 21 g of androstane  $3\beta$ ,  $6\alpha$ ,  $17\beta$ -triol **II** were obtained (known product: Nicholson, S. H., Turner, A. B. J. Chem. Soc., Perkin Trans. 1, 1976, 1357 and US 6,384,250 B2).

[128] The analytical results are in agreement with those reported in the literature.

*Step 2: Selective Oxidation.*


[129] To a solution of Androstane 3 $\beta$ , 6 $\alpha$ , 17 $\beta$ -triol **II** (30 grams) in a mixture composed by Dioxane (825 mL), Water (150 mL) and Pyridine (16.5 mL), N-Bromosuccinimide (52 grams) was added portion wise within 10 minutes protecting the vessel from light. The mixture was stirred for 16 hours at room temperature, diluted with 900 ml of water then Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (15.5 grams) were added portion wise within 15 minutes. The solution was concentrated (around 1500 mL were removed) and the suspension was filtered and the solid dried under vacuum giving 28.1 grams of 6 $\alpha$ -hydroxyandrostane-3,17-dione **III** (95% yield).

*Step 3: Ketone Protection.*



[130] A suspension of 6 $\alpha$ -hydroxyandrostane-3,17-dione **III** (18.85 grams) in 360 mL of glycol and P-Toluenesulfonic Acid (554 mg) was heated at 100° C and distilled under vacuum to remove the azeotropic mixture glycol/water (around 5 mL). The mixture was cooled and treated with 250 mg of KOH dissolved in 25 ml of Methanol. 15 mL of water were added and, after stirring for 2 hours, the suspension was filtered giving intermediate **IV** as white solid (20.2 grams, 83% yield). The product was used without further purification.


*Step 4: Oxidation.*



[131] A solution constituted by 3 mL of Sodium Hypochlorite (6%) and 28 mL of Ethyl Acetate were stirred at room temperature and 27 mg of RuO<sub>2</sub> hydrate were added. When


all the catalyst Ruthenium was solubilized the product **IV** (1 gram) was added portion wise waiting for the disappearance of black suspension. After 1 hour additional 3 ml of Sodium Hypochlorite (6%) were added and the clear solution stirred for 3 hours at room temperature. When the reaction is completed the mixture was filtered on a Celite pad and the aqueous phase was extracted with AcOEt. The combined organic phases were washed with a solution of  $\text{NaHCO}_3$  (5% in water) and with NaCl (10% in water). The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness giving intermediate **V** (950 mg, 94% yield).

*Step 5: Reduction.*



[132] A suspension of product **V** (5.76 grams) in Methanol (72 mL) was stirred at  $0^\circ\text{C}$  and  $\text{NaBH}_4$  (730 mg) was added. After 2 hours the reaction was completed, and the solvent was removed under reduced pressure. The crude product was suspended in 30 mL of water and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness. The crude solid was purified by flash chromatography ( $\text{SiO}_2$ , Cyclohexane/AcOEt 7/3 as eluent) giving the product **VI** (5.16 grams, 89% yield).


*Step 6: Ketone Deprotection.*



[133] To a stirred solution of product **VI** (2.85 grams) in 350 mL of distilled Acetone, P-toluenesulfonic acid (7.14 grams) was added. After 3 hours at room temperature a 5% solution of  $\text{NaHCO}_3$  was added and the solvent was removed under reduced pressure. The

product was extracted with  $\text{CH}_2\text{Cl}_2$ . The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and evaporated to dryness, giving the intermediate **VII** (2.13 grams, 95% yield).

*Step 7: Synthesis of PST 2915.*




[134] To a stirred solution of 6 $\beta$ -hydroxyandrostane-3,17-dione **VII** (3.5 grams) in THF (100 mL), a solution of 2-aminoethoxyamine dihydrochloride (1.728 grams) in  $\text{H}_2\text{O}$  (34 mL) was rapidly added dropwise. After 1.5 h at room temperature under vigorous stirring,  $\text{NaCl}$  (4 grams) was added and the mixture stirred for 15 min. The phases were separated, and the aqueous phase was extracted twice with THF (2x50 mL). The combined organic extracts were dried over  $\text{Na}_2\text{SO}_4$ , filtered and evaporated to give a white solid (4.52 grams).

[135] The crude product was suspended and slurried in 45 mL of  $\text{AcOEt/EtOH}$  97/3 for 1.5 hour then filtered and dried under reduce pressure at  $35^\circ\text{C}$  for 48 hours, giving (E,Z)-3-(2-Aminoethoxyimino)-6 $\beta$ -hydroxyandrostane-17-one hydrochloride, **PST2915** (4.069 grams, 89% yield).

#### Synthesis of PST 3093

*Step 1: Hydroboration.*




[136] To a solution of dehydroepiandrosterone **I** (30.0 g) in 450 mL of anhydrous THF maintained under a nitrogen atmosphere and at a temperature of  $-10^\circ\text{C}$ . was added the

complex BH.THF 1M in THF (260 mL). On completing the addition, the temperature was allowed to rise once again to room temperature; after 3 hours, 500 mL of H<sub>2</sub>O were added and then NaBO<sub>3</sub>·4H<sub>2</sub>O (31.4 g). The reaction was left to stir for one night. The precipitate formed was filtered, washed with THF and eliminated. The aqueous and organic phases were separated, NaCl was added to the aqueous phase and this was re-extracted with THF (3x200 mL). The combined organic phases were anhydridified with NaCl and Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure to obtain the crude product, which was crystallised by AcOEt/MeOH and then filtered and washed with AcOEt. Approximately 21 g of androstane 3 $\beta$ , 6 $\alpha$ , 17 $\beta$ -triol **II** were obtained (known product: Nicholson, S. H., Turner, A. B. J. Chem. Soc., Perkin Trans. 1, 1976, 1357 and US 6,384,250 B2).


[137] The analytical results are in agreement with those reported in the literature.

*Step 2: Selective Oxidation*




[138] To a solution of Androstane 3 $\beta$ , 6 $\alpha$ , 17 $\beta$ -triol **II** (30 grams) in a mixture composed by Dioxane (825 mL), Water (150 mL) and Pyridine (16.5 mL), N-Bromosuccinimide (52 grams) was added portion wise within 10 minutes protecting the vessel from light. The mixture was stirred for 16 hours at room temperature, diluted with 900 ml of water then Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (15.5 grams) were added portion wise within 15 minutes. The solution was concentrated (around 1500 mL were removed) and the suspension was filtered and the solid dried under vacuum giving 28.1 grams of 6 $\alpha$ -hydroxyandrostane-3,17-dione **III** (95% yield).

*Step 3: Ketone Protection.*




[139] A suspension of 6 $\alpha$ -hydroxyandrostane-3,17-dione **III** (18.85 grams) in 360 mL of glycol and P-Toluenesulfonic Acid (554 mg) was heated at 100° C and distilled under vacuum to remove the azeotropic mixture glycol/water (around 5 mL). The mixture was cooled and treated with 250 mg of KOH dissolved in 25 ml of Methanol. 15 mL of water were added and, after stirring for 2 hours, the suspension was filtered giving intermediate **IV** as white solid (20.2 grams, 83% yield). The product was used without further purification.

*Step 4: Oxidation.*




[140] A solution constituted by 3 mL of Sodium Hypochlorite (6%) and 28 mL of Ethyl Acetate were stirred and 27 mg of RuO<sub>2</sub> hydrate were added. When all the catalyst Ruthenium was solubilized the product **IV** (1 gram) was added portion wise waiting for the disappearance of black suspension. After 1 hour additional 3 ml of Sodium Hypochlorite (6%) were added and the clear solution stirred for 3 hours at room temperature. When the reaction is completed the mixture was filtered on a Celite pad and the aqueous phase was extracted with AcOEt. The combined organic phases were washed with a solution of NaHCO<sub>3</sub> (5% in water) and with NaCl (10% in water). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness giving intermediate **V** (950 mg, 94% yield).

*Step 5: Reduction.*




[141] A suspension of product **V** (5.76 grams) in Methanol (72 mL) was stirred at 0°C and NaBH<sub>4</sub> (730 mg) was added. After 2 hours the reaction was completed, and the solvent was removed under reduced pressure. The crude product was suspended in 30 mL of water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated and dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The crude solid was purified by flash chromatography (SiO<sub>2</sub>, Cyclohexane/AcOEt 7/3 as eluent) giving the product **VI** (5.16 grams, 89% yield).

*Step 6: Ketone Deprotection.*



[142] To a stirred solution of product **VI** (2.85 grams) in 350 mL of distilled Acetone, P-toluenesulfonic acid (7.14 grams) was added. After 3 hours at room temperature a 5% solution of NaHCO<sub>3</sub> was added and the solvent was removed under reduced pressure. The product was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness, giving the intermediate **VII** (2.13 grams, 95% yield).

*Step 7: Synthesis of PST 3093.*



[143] To a stirred solution of 6β-hydroxyandrostane-3,17-dione **VII** (4.5 grams) in THF (113 mL), a solution of O-(Carboxymethyl)hydroxylamine dihydrochloride (1.56 gram) in H<sub>2</sub>O (5 mL) was rapidly added dropwise. After 1.5 h at room temperature under vigorous stirring, NaCl (6.4 grams) was added and the mixture stirred for 15 min. The phases were separated, and the aqueous phase was extracted three times with THF (50 mL). The

combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated to dryness giving 5.95 grams of crude product.

[144] The crude product was purified by flash chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>/MeOH/Acetic acid, 92.5/7/0.5) giving (E,Z)-[(6-beta-hydroxy-17-oxoandrostan-3-ylidene)amino]oxyacetic acid, **PST3093** (3.7 grams, 69% yield).

## Example 2. Biological activity of Istaroxime metabolites

### Procedures

#### *Animal care*

[145] The investigation adheres to the Guide of the Care and Use of Laboratory Animals published by the National Institute of Health (NIH publication No. 85-23, revised 1996) and to the guidelines for animal care endorsed by the participating institutions.

#### *Purification of dog renal Na,K-ATPase and Na,K-ATPase activity assay*

[146] Purification of renal Na,K-ATPase was performed according to the method of Jørgensen (Methods Enzymol. 1988;156:29-43). Kidneys were excised from 1-3 year-old male beagle dogs (WuXi AppTec, Suzhou Co., Ltd. 1318 Wuzhong Ave., Wuzhong District Suzhou, 215104 P.R. China) under pentobarbital anesthesia (Import Authorization from Italian Health Ministry 0009171-09/04/2015-DGSAF-COD\_UO-P, 2015). Kidneys were sliced and the outer medulla was dissected, pooled and suspended (1g/10 ml) in a sucrose-histidine solution, containing 250 mM sucrose, 30 mM histidine and 5 mM EDTA, pH 7.2 and homogenized. The homogenate was centrifuged at 6.000 g for 15 min, the supernatant was decanted and centrifuged at 48.000 g for 30 min. The pellet was suspended in the sucrose-histidine buffer and incubated for 20 min with a sodium-dodecyl-sulphate (SDS) solution dissolved in a gradient buffer, containing 25 mM imidazole and 1 mM EDTA, pH 7.5. The sample was layered on the top of a sucrose discontinuous gradient (10, 15 and 29.4%) and centrifuged at 60.000 g for 115 min. The pellet was suspended in the gradient buffer.

[147] Na,K-ATPase activity was assayed *in vitro* by measuring the release of <sup>32</sup>P-ATP, as described previously (see Ferrandi M. et al., Hypertension 1996;28(6):1018-25). Increasing concentrations of the standard ouabain, or tested compound, were incubated with 0.3 g of purified dog kidney enzyme for 10 min at 37°C in 120 l final volume of a medium containing 140 mM NaCl, 3 mM MgCl<sub>2</sub>, 50 mM Hepes-Tris, 3 mM ATP at a pH 7.5. Then, 10 l of incubation solution containing 10 mM KCl and 20 nCi of <sup>32</sup>P-ATP (3-10

Ci/mmol, Perkin Elmer) was added, and the reaction was continued for 15 min at 37°C. The reaction was then stopped by acidification with 20% v/v ice-cold perchloric acid.  $^{32}\text{P}$  was separated by centrifugation with activated Charcoal (Norit A, Serva) and the radioactivity was measured. The inhibitory activity was expressed as percent of the control samples carried out in the absence of ouabain or tested compound. The concentration of compound causing 50% inhibition of the Na,K-ATPase activity ( $\text{IC}_{50}$ ) was calculated by using a multiple parameter non-linear regression best fitting program (Kaleidagraph<sup>TM</sup>, Sinergy Software).

*SERCA2a activity measurement in heart sarcoplasmic reticulum (SR) microsomes*

[148] Male beagle dogs were used for obtaining cardiac tissues for SERCA2a-enriched sarcoplasmic reticulum preparations. Healthy dogs were utilized for obtaining the data in **Table 3**. Chronic heart failure was induced in dogs by multiple intracoronary microembolizations with polystyrene latex microspheres (45-90  $\mu\text{m}$ , Polysciences, Warrington, PA, USA) as described previously (see Sabbah HN et al., Am J Physiol. 1991;260:H1379-84). The experiments were conducted in the General Pharmacology Department of Sigma-Tau, Rome, Italy.

[149] Left ventricle tissues were dissected, homogenized in 4 volumes of 10 mM  $\text{NaHCO}_3$  (pH 7), 1 mM PMSF, 10  $\mu\text{g}/\text{ml}$  Aprotinin and Leupeptin and centrifuged at 12.000 g for 15 minutes, as described in Nediani C. et al. (J Biol Chem. 1996;271:19066-73). Supernatants were filtered and centrifuged at 100.000 g for 30 min. Contractile proteins were extracted by suspending the pellets with 0.6 M KCl, 30 mM Histidine, pH 7 and further centrifugation at 100.000 g for 30 min. Final pellets were reconstituted with 0.3 M Sucrose, 30 mM Histidine, pH 7.

[150] SERCA2a activity was measured *in vitro* as  $^{32}\text{P}$ -ATP hydrolysis at different  $\text{Ca}^{2+}$  concentrations (100 to 3000 nM) in the absence and presence of the tested compounds as described previously (see Micheletti R. et al., Am J Card 2007;99:24A-32A). Increasing concentrations of each compound (from 0.05 to 300 nM) were pre-incubated with 2  $\mu\text{g}$  of microsomes for 5 min at 4° C in 80  $\mu\text{l}$  of a solution containing 100 mM KCl, 5 mM  $\text{MgCl}_2$ , 1  $\mu\text{M}$  A23187, 20 mM Tris, pH 7.5. Then, 20  $\mu\text{l}$  of 5 mM Tris-ATP containing 50 nCi of  $^{32}\text{P}$ -ATP (3-10 Ci/mmol, Perkin Elmer) was added. The ATP hydrolysis was continued for 15 min at 37°C and was stopped by acidification with 100  $\mu\text{l}$  of 20% v/v ice-cold perchloric acid.  $^{32}\text{P}$  was separated by centrifugation with activated charcoal (Norit A, SERVA) and the radioactivity was measured. SERCA2a-dependent activity was identified as the portion of total hydrolytic activity inhibited by 10  $\mu\text{M}$  cyclopiazonic acid (see Seidler NW. et al., J Biol Chem. 1989; 264:17816-23).

[151] Dose-response curves were fitted by using a sigmoidal curve fitting software and the maximal velocity ( $V_{max}$ ) activity and the  $K_d$   $Ca^{2+}$  were calculated (Synergy Software KaleidaGraph 3.6).

*Drug toxicity studies in mice*

[152] Acute toxicity has been determined in the mouse (Albino Swiss CD-1, body weight 30 g). Mice were orally treated, or intravenously injected, with single administration of increasing doses of the test substance to identify the dose causing 50% mortality. Mortality occurred within 30 min after the administration and survival after 24h. The acute toxicity ( $LD_{50}$ ) was then assessed.

*Haemodynamics in streptozotocin diabetic rat (echocardiography 2M-Doppler-Tissue Doppler)*

[153] Sprague Dawley male rats (150-175 g) were made diabetic by a single injection into the tail vein of a solution of streptozotocin (STZ, 50 mg/kg, Sigma-Aldrich), freshly prepared in 0.1 M sodium citrate buffer, pH 4.5. Control rats received citrate buffer. Fasting glycaemia was measured after 1 week and rats with values greater than 400 mg/dl were considered diabetic.

[154] Eight to nine weeks after STZ injection, rats were submitted to transthoracic echocardiographic and Doppler evaluation performed under pentobarbital anesthesia. Two-dimensionally guided M-mode recordings were used to obtain short-axis measurements of left ventricular end-diastolic diameter (LVEDD), left ventricular end-systolic diameter (LVESD), posterior (PW) and septal (SW) diastolic wall thickness according to the American Society of Echocardiography guidelines (Lang RM et al., Eur J Echocardiography 2006; 7:79-108). Fractional shortening was calculated as  $FS = (LVEDD - LVESD) / LVEDD$ . Relative wall thickness was calculated as  $PWTd + IVSTd / LVEDD$ .

[155] Mitral inflow was measured by pulsed Doppler at the tips of mitral leaflets from an apical 4-chamber view to obtain early and late filling velocities (E, A) and deceleration time of early filling velocity (DT). The deceleration slope was calculated as E/DT ratio. The mitral deceleration index was calculated as DT/E ratio.

[156] Tissue Doppler Imaging (TDI) was evaluated from the apical 4-chamber view to record septal mitral annular movements, i.e., peak myocardial systolic ( $s'$ ) and early and late diastolic velocity ( $e'$  and  $a'$ ).

[157] The compound PST 3093 was iv administered to STZ injected rats at the dose of 0.22 mg/kg and echocardiographic parameters were measured after 15 and 30 min from the beginning of iv infusion and 10 min after interruption of infusion.

#### *Statistical analysis*

[158] Data are reported as mean  $\pm$  SD, as indicated. Statistical analysis was performed by Student's t-test (pair t test for STZ rats).  $P < 0.05$  was regarded as statistically significant.

#### Biological results

##### *In vitro screening*

##### *Inhibition of dog renal Na,K-ATPase activity*

[159] **Table 2** shows the inhibitory effect of the tested compounds on the enzymatic activity of the purified dog renal Na,K-ATPase. The corresponding  $IC_{50}$  are expressed in  $\mu M$  concentration. Istaroxime inhibited the Na,K-ATPase activity with an  $IC_{50}$  of 0.14  $\mu M$ , similar to that of Digoxin, while PST 2915 inhibited the Na,K-ATPase activity with an  $IC_{50}$  of 2.1  $\mu M$ . Further, PST 3093 did not significantly inhibit Na,K-ATPase activity at all ( $IC_{50} > 100 \mu M$ ).

**Table 2 Inhibition of dog renal Na,K-ATPase.**

| Compound |  |
|----------|--|
|          |  |
|          |  |
|          |  |
|          |  |
|          |  |

*SERCA2a ATPase activity in heart-derived SR microsomes from normal dog*

[160] The compounds disclosed herein were tested on SERCA2a ATPase activity prepared from normal and failing dogs in a range of concentrations from 0.1 and 500 nM. The effect has been expressed as % increase of the Vmax activity of a control sample run in the absence of compound. Data are mean  $\pm$  SD, where n indicates the number of experiments.

[161] In the normal dog SR vesicles, istaroxime, PST 3093 and PST 2915 significantly stimulated SERCA2a activity at concentrations of 0.1 nM and 10 nM (see **Table 3**). SERCA2a activation by istaroxime, PST 3093 and PST 2915 was also tested in failing dog preparations. This effect was particularly evident in the failing preparations (data not shown), where SERCA2a activity is known to be depressed compared to a normal heart (Bers DM, Physiology 2006;21:380–387), and therefore implies that the compounds istaroxime and PST 3093 may correct SERCA2a alteration in the failing heart.

[162] In contrast, previous studies showed that Digoxin failed to stimulate SERCA2a activity (Rocchetti M et al., J Pharmacol Exp Ther 2005;313:207-215; Ferrandi M et al., Br J Pharmacol 2013;169:1849-61).

**Table 3 SERCA2a ATPase activity in heart-derived SR microsomes from normal dog.**

Data are expressed as % increase vs control and are mean  $\pm$  SD

| Compound   | Concentration nM<br>(ng/ml) | Vmax<br>( $\mu$ mol/min/mg prot)<br>mean $\pm$ SD | % increase vs<br>control |
|------------|-----------------------------|---------------------------------------------------|--------------------------|
| Istaroxime |                             | 1.252 $\pm$ 0.083 (n=5)                           |                          |
|            | 0.1 nM (0.039 ng/ml)        | 1.423 $\pm$ 0.123 (n=5)                           |                          |
|            | 10 nM (3.9ng/ml)            | 1.505 $\pm$ 0.111 (n=5)                           |                          |
|            |                             | 1.312 $\pm$ 0.050 (n=5)                           |                          |
|            | 0.1 nM (0.037 ng/ml)        | 1.641 $\pm$ 0.194 (n=4)                           |                          |
|            | 10 nM (3.7 ng/ml)           | 1.556 $\pm$ 0.106 (n=4)                           |                          |

| Compound | Concentration nM<br>(ng/ml) | V <sub>max</sub><br>( $\mu$ mol/min/mg prot)<br>mean $\pm$ SD | % increase vs<br>control |
|----------|-----------------------------|---------------------------------------------------------------|--------------------------|
|          |                             | 1.312 $\pm$ 0.050 (n=5)                                       |                          |
|          | 0.1 nM (0.041ng/ml)         | 1.464 $\pm$ 0.184 (n=5)                                       |                          |
|          | 10 nM (4.1 ng/ml)           | 1.526 $\pm$ 0.038 (n=5)                                       |                          |
|          |                             | 1.312 $\pm$ 0.050 (n=5)                                       |                          |
|          | 0.1 nM (0.037 ng/ml)        | 1.466 $\pm$ 0.112 (n=5)                                       |                          |
|          | 10 nM (3.7 ng/ml)           | 1.541 $\pm$ 0.170 (n=5)                                       |                          |

#### *In vivo studies*

##### *Acute toxicity in mouse*

[163] The acute toxicity of the tested compound PST 3093 was determined in the mouse (Albino Swiss CD-1, body weight 30 g). Compound PST 3093 was orally administered or intravenously injected at increasing doses to identify the dose causing 50% mortality. Mortality occurred within 30 min after the administration and survival after 24h.

[164] The results for PST 3093 acute toxicity are reported in **Table 4** and indicated that the compound had an LD<sub>50</sub> >250 and 200 mg/kg after iv or oral administration, respectively. For comparison, the acute toxicity for the reference compound istaroxime has been also included in **Table 4**.

**Table 4 Acute toxicity (LD<sub>50</sub>) of Istaroxime and PS3093 in mouse.**

| Compound        | mg/kg |
|-----------------|-------|
| Istaroxime i.v. |       |
| Istaroxime os   |       |
| 3093 i.v.       |       |
| 3093 os         |       |

i.v., intravenous

os, oral

*Haemodynamics in streptozotocin (STZ) diabetic rats (echocardiography 2M-Doppler-Tissue Doppler)*

[165] **Table 5** shows the echocardiographic parameters in STZ diabetic rats before and after 15 and 30 min from iv infusion of PST 3093 at 0.22 mg/kg, and 10 min after interruption of infusion. Data is presented as mean  $\pm$  SD, and values with an asterisk are statistically significant with at least  $p < 0.05$ .

[166] The data indicated that in an animal model characterized by a diastolic dysfunction, such as the STZ diabetic rats, PST 3093 administration ameliorated diastolic function. In particular, E wave (which represents the early filling velocity of transmitral inflow during the rapid filling phase and constitutes the energy dependent phase of LV relaxation, mainly mediated by SERCA2a activity) was significantly increased in STZ rats at 15 and 30 min after PST 3093 infusion (**Table 5**). This effect was consistent with a stimulation of SERCA2a activity by the compound, as shown in the *in vitro* assay (**Table 3**), suggesting the ability of PST 3093 to restore SERCA2a function activity, which is depressed in STZ rats as shown by Choi *et al.* (AJP 2002; H1398-H1408).

[167] However, it should be considered that the peak E velocity is influenced by the preload and is directly correlated with heart rate (HR) (Mihm MJ *et al.*, Life Sci. 2001; 22;69(5):527-42; do Carmo JM *et al.*, AJP 2008; 295:H1974-1981). Conversely, the deceleration time of E wave (DT), and the related changes in mitral deceleration index (DT/E) and deceleration slope of E wave (E/DT), are not affected by HR changes and are

considered robust indicators of diastolic function and early signs of diastolic dysfunction (Mihm MJ et al., Life Sci. 2001; 22;69(5):527-42). In particular, the behaviour of some echocardiographic diastolic parameters, such as the DT of E wave, can be affected even in opposite direction according to the various grades of diastolic dysfunction. As clearly indicated in previous publications (see Mitter SS et al., JACC 2017;69(11):1451-1464), the DT of E wave is usually prolonged when diastolic dysfunction is of grade 1 and becomes very short in patients with grade 3 diastolic dysfunction, being dynamically affected also by the state of pulmonary congestion of the patient.

[168] In this respect, the variation of the DT of E wave in animal models of diastolic dysfunction, as compared to the healthy controls, may vary in opposite directions. For example, in rats with diabetic cardiomyopathy induced by STZ injection, the DT results are equal to (see Thackeray JT et al., Cardiovasc Diabetol. 2011;10:75; Carillion A et al., PloS One 2017; e0180103) or even longer than in control rats (Joffe II et al., JACC Vol. 34, No. 7, 1999; 2111-2119; Guido MC et al., Oxid Med Cell Longev. 2017;5343972); while in dogs with heart failure induced by microembolization of the coronary arteries, the DT results are reduced as compared with control healthy dogs (Sabbah H et al., Am J Cardiol. 2007;99(2A):41A-46A). Furthermore, unlike transmitral Doppler flow, tissue-Doppler (TDI) parameters were relatively unaffected by load and a decrease in the early relaxation velocity ( $e'$ ) would be an unequivocal indicator of diastolic dysfunction.

[169] The data evidenced marked effects of PST 3093 on these parameters. The E wave is significantly prolonged by PST 3093 treatment, while the DT is reduced. A significant reduction of DT and DT/E with increased of E/DT and  $e'$  are shown in **Table 5**. The increase of  $e'$  appeared to be associated with a significant increase of CO and SV, while no significant change of heart rate was observed. Of note, the direction of the effects of PST 3093 on DT and E/ $e'$  are the same as those obtained when this STZ rat model was treated with Istaroxime.

[170] In contrast, in the HF dog model of coronary microembolization, where the DT was reduced as compared to control dogs (Sabbah H et al., Am J Cardiol. 2007;99(2A):41A-46A), the effect of istaroxime was to prolong the DT. In other words, independently from the variation of the DT of E wave in the HF animal models as compared with their respective controls, istaroxime reverses such parameters toward the levels present in the corresponding control animal and improves diastolic function.

[171] These effects were more evident after 30 min from the beginning of PST 3093 infusion and tended to disappear after 10 min from the interruption of infusion. The effects of PST 3093 on the impaired cardiac function of STZ rats were consistent with the SERCA2a

stimulatory activity of the compound that, by correcting the depressed cardiac relaxation, increased the amount of blood available for contraction and resulted in an increase of volume of blood pumped from the ventricle (SV).

[172] To evaluate the relevance of the above results to the human condition, the obvious pathophysiological differences between the patients with AHF and the STZ rats should be considered. In the latter, marked changes in the body fluids, sympathetic nervous system and heart rate, may, *per se*, affect echocardiographic parameters, independently from the changes in cellular  $\text{Ca}^{2+}$  handling and decrease of SERCA2a activity (Mihm MJ et al., Life Sci. 2001; 22;69(5):527-42; do Carmo JM et al., AJP 2008; 295:H1974-1981). Therefore, the similarities between human and rats in DT/E, E/DT and  $e'$  changes may be considered as having the same underlying mechanism – a stimulation of the SERCA2a activity by PST 3093.

**Table 5 Haemodynamic parameters after 3093 iv infusion in STZ diabetic rats**

\* P<0.05 compared to basal values


| Function              | Echo Parameter | before<br>(n=10) | 0.22 mg/kg iv<br>after 15 min<br>(n=10) | 0.22 mg/kg iv<br>after 30 min<br>(n=10) | after 10 min from<br>STOP infusion<br>(n=8) |
|-----------------------|----------------|------------------|-----------------------------------------|-----------------------------------------|---------------------------------------------|
|                       |                |                  |                                         |                                         |                                             |
| Diastolic<br>function |                |                  |                                         |                                         |                                             |
|                       |                |                  |                                         |                                         |                                             |
|                       |                |                  |                                         |                                         |                                             |
|                       | E/e'           |                  |                                         |                                         |                                             |
|                       | e'             |                  |                                         |                                         |                                             |

|         |  |  |  |  |  |
|---------|--|--|--|--|--|
| overall |  |  |  |  |  |
|         |  |  |  |  |  |
|         |  |  |  |  |  |


|                   |                                                 |
|-------------------|-------------------------------------------------|
| E,                | E wave, early filling velocity of mitral inflow |
| DT (ms),          | deceleration time of E wave                     |
| DT/E ( $s^2/m$ ), | mitral deceleration index                       |
| E/DT ( $m/s^2$ ), | deceleration slope                              |
| E/e',             | index of LV filling pressure                    |
| e' (cm/s) TDI,    | early relaxation velocity                       |
| CO (ml/min),      | cardiac output                                  |
| HR (beat/min),    | heart rate                                      |
| SV (ml/beat),     | stroke volume                                   |

**CLAIMS**

1. A pharmaceutical composition for intravenous infusion in a human subject, the pharmaceutical composition comprising istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof for use in a method for the treatment of acute heart failure characterized in that the administration is for an infusion duration longer than 6 hours, whereby diastolic relaxation is improved as compared to administration of istaroxime by intravenous infusion for a duration from 3 to 6 hours.
2. The pharmaceutical composition for use according to claim 1, wherein the diastolic relaxation improvement is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure.
3. The pharmaceutical composition for use according to claim 1 or 2, wherein the infusion duration is up to about 24 hours.
4. The pharmaceutical composition for use according to any one of claims 1, 2, or 3, wherein the infusion duration is up to about 36 hours.
5. The pharmaceutical composition for use according to any one of claims 1 to 4, wherein the infusion duration is up to about 48 hours.
6. The pharmaceutical composition for use according to any one of claims 1 to 5, wherein peripheral vascular resistance does not decrease prior to 24 hours after infusion.
7. The pharmaceutical composition for use according to any one of claims 1 to 6, wherein istaroxime is administered at a dose between 0.2 µg/kg/min and 1.5 µg/kg/min.
8. The pharmaceutical composition for use according to any one of claims 1 to 7, wherein istaroxime is administered at a dose between 0.25 µg/kg/min and 1.0 µg/kg/min.
9. The pharmaceutical composition for use according to any one of claims 1 to 8, wherein istaroxime is administered by infusion for a duration sufficient to produce a plasma level of an istaroxime metabolite, wherein the plasma level is greater than about 5 ng/ml for an accumulation period of at least about 6 hours, and wherein the istaroxime metabolite comprises a formula (II) or a formula (III) compound

**PST 2915**

Formula (II)

**PST 3093**

Formula (III)

or a pharmaceutically acceptable salt, an ester a solvate, a hydrate, a polymorph thereof.

10. The pharmaceutical composition for use according to any one of claims 1 to 9, wherein the human subject suffers from heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).

11. The pharmaceutical composition for use according to any one of claims 1-10 wherein the human subject is undergoing a therapeutic treatment for heart failure with one or more further therapeutically active ingredients.

12. The pharmaceutical composition for use according to claim 11, wherein the further therapeutically active ingredients are selected among ACE inhibitors, AIRBs, diuretics, Ca channel blockers, beta-blockers, digitalis, NO donors, vasodilators, SERCA2a stimulators, neprilisin (NEP) inhibitors, myosin filament activators, recombinant relaxin-2 mediators,

recombinant NP protein, activators of the soluble Guanylate Cyclase (sGC), beta-arrestin ligand of Angiotensin II receptor.

13. The pharmaceutical composition for use according to claim 12, wherein the diuretic, is selected among furosemide, bumetanide, torasemide, metolazone, an aldosterone antagonist, in particular spironolactone and eplerenone; thiazide diuretics, in particular hydrochlorothiazide, metolazone, chlorthalidone.

14. The pharmaceutical composition for use according to claim 12, wherein the ACE inhibitor is selected among Lisinopril and Ramipril.

15. The pharmaceutical composition for use according to claim 12, wherein the Angiotensin II receptor blocker is selected among valsartan, candesartan, olmesartan, telmisartan and losartan.

16. The pharmaceutical composition for use according to claim 12, wherein the Angiotensin receptor/neprilysin inhibitor is sacubitril.


17. The pharmaceutical composition for use according to claim 12, wherein the beta-blocker is selected among carvedilol and metoprolol.

18. The pharmaceutical composition for use according to claim 12, wherein the Vasodilator is selected among Hydralazine, optionally combined with isosorbide dinitrate, nitrates, in particular nitroglycerin or isosorbide nitrate, amlodipine, felodipine, non-dihydropyridines, in particular diltiazem or verapamil.


19. The pharmaceutical composition for use according to claim 12, wherein the agent for the treatment of heart failure is selected from Digoxin, entresto, omecantiv, serelaxin, ularitide, levosimendan.

20. The pharmaceutical composition for use according to any one of claims 11 to 19, wherein the therapeutic treatment for heart failure is preceding or concomitant or subsequent to istaroxime administration.

21. A compound having a formula (II) or formula (III)

**PST 2915**

Formula (II)

**PST 3093**

Formula (III)

or a pharmaceutically acceptable salt, an ester a solvate, a hydrate, a polymorph thereof.

22. The compound of claim 21 having the formula (III), or a pharmaceutically acceptable salt, an ester, a solvate, a hydrate, a polymorph thereof.

23. The compound of claim 21 or claim 22 for use in the treatment of a disease requiring the activation of SERCA2a.

24. The compound of claim 23 for use in the treatment of a cardiovascular disease.

25. The compound for use according to claim 24, wherein the disease is heart failure, in particular acute heart failure.

26. A pharmaceutical composition comprising the compound of claim 21 or claim 22, or a pharmaceutically acceptable salt, a solvate, a hydrate, a polymorph thereof in admixture with at least one pharmaceutically acceptable vehicle and/or excipient.

27. The pharmaceutical composition according to claim 26, further comprising at least one additional therapeutic ingredient as disclosed in any one of claims 12 to 19.

28. The pharmaceutical composition according to claim 27, wherein the at least one additional therapeutic ingredient is in the same or in a separate unit dosage form of the compound of claim 21 or claim 22, or a pharmaceutically acceptable salt, a solvate, a hydrate, a polymorph thereof.

29. A method for treating an individual having heart failure, the method comprising the steps of:

(1) providing an individual having heart failure;

(2) administering to the individual a therapeutically effective amount of a pharmaceutical composition comprising; (i) a pharmaceutically acceptable carrier; and (ii) istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof; wherein said administering comprises intravenous fusion for an infusion duration of longer than 6 hours;

(3) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation;

wherein the administering of the pharmaceutical composition results in an improvement in diastolic relaxation as compared to istaroxime administered by intravenous infusion for less than 6 hours, thereby treating the individual having acute heart failure.

30. The method of claim 29, wherein the individual is human.

31. The method of claim 29 or claim 30, wherein the diastolic relaxation improvement is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure.

32. The method of any one of claims 29 to 31, wherein the infusion duration is up to about 24 hours.

33. The method of claim 32, wherein the infusion duration is up to about 36 hours.

34. The method of claim 33, wherein the infusion duration is up to about 48 hours.

35. The method of any one of claims 29 to 34, wherein the istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof is administered at a dose of about 0.2 µg/kg/min to about 1.5 µg/kg/min.

36. The method of claim 35, wherein the istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof is administered at a dose of about 0.25 µg/kg/min to about 1.0 µg/kg/min.

37. The method of any one of claims 29 to 36, wherein the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).

38. The method of any one of claims 29 to 37, wherein the individual is undergoing a therapeutic treatment for heart failure with one or more further therapeutically active ingredients.

39. The method of claim 38, wherein the further therapeutically active ingredients are selected among ACE inhibitors, AIRBs, diuretics, Ca channel blockers, beta-blockers, digitalis, NO donors, vasodilators, SERCA2a stimulators, neprilisin (NEP) inhibitors, myosin filament activators, recombinant relaxin-2 mediators, recombinant NP protein, activators of the soluble Guanylate Cyclase (sGC), beta-arrestin ligand of Angiotensin II receptor.

40. The method of claim 39, wherein the diuretic, is selected among furosemide, bumetanide, torasemide, metolazone, an aldosterone antagonist, in particular spironolactone and eplerenone; thiazide diuretics, in particular hydrochlorothiazide, metolazone, chlorthalidone.

41. The method of claim 39, wherein the ACE inhibitor is selected among Lisinopril and Ramipril.

42. The method of claim 39, wherein the Angiotensin II receptor blocker is selected among valsartan, candesartan, olmesartan, telmisartan and losartan.

43. The method of claim 39, wherein the Angiotensin receptor/neprilysin inhibitor is sacubitril.

44. The method of claim 39, wherein the beta-blocker is selected among carvedilol and metoprolol.

45. The method of claim 39, wherein the Vasodilator is selected among Hydralazine, optionally combined with isosorbide dinitrate, nitrates, in particular nitroglycerin or isosorbide nitrate, amlodipine, felodipine, non-dihydropyridines, in particular diltiazem or verapamil.


46. The method of claim 39, wherein the agent for the treatment of heart failure is selected from Digoxin, entresto, omecantiv, serelaxin, ularitide, levosimendan.

47. The method of any one of claims 39 to 46, wherein the therapeutic treatment for heart failure is preceding or concomitant or subsequent to istaroxime administration.

48. A pharmaceutical composition for intravenous infusion in an individual comprising istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof, the pharmaceutical composition for use in a method for the treatment of acute heart failure characterized in that the administration is for an infusion duration sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite, wherein:


(a) the plasma concentration level is greater than about 5 ng/ml for an accumulation period of at least about 6 hours; and

(b) the istaroxime metabolite comprises formula (II) or (III):



**PST 2915**

Formula (II)



**PST 3093**

Formula (III)

or a pharmaceutically acceptable salt, an ester a solvate, a hydrate, a polymorph thereof; whereby the diastolic relaxation is improved as compared to administration of istaroxime prior to the accumulation period.

49. The pharmaceutical composition for use according to claim 48, wherein the diastolic relaxation improvement is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure.

50. The pharmaceutical composition for use according any to claims 48 or 49, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 10 ng/ml for an accumulation period of at least about 6 hours.

51. The pharmaceutical composition for use according to claim 50, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 15 ng/ml for an accumulation period of at least about 6 hours.

52. The pharmaceutical composition for use according to claim 51, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 20 ng/ml for an accumulation period of at least about 6 hours.

53. The pharmaceutical composition for use according to any one of claims 48 to 52, wherein istaroxime is administered at a dose comprised between 0.2 µg/kg/min and 1.5 µg/kg/min.

54. The pharmaceutical composition for use according to claim 53, wherein istaroxime is administered at a dose comprised between 0.25 µg/kg/min and 1.0 µg/kg/min.


55. A method for treating an individual having heart failure, the method comprising the steps of:

(1) providing an individual having heart failure;

(2) administering to the individual a therapeutically effective amount of a pharmaceutical composition comprising; (i) a pharmaceutically acceptable carrier; and (ii) istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof; wherein said administering comprises intravenous fusion for an infusion duration sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite, wherein:


(i) the plasma concentration level is greater than about 5 ng/ml for an accumulation period of at least about 6 hours; and

(ii) the istaroxime metabolite comprises formula (II) or formula (III):



**PST 2915**

Formula (II)



**PST 3093**

Formula (III)

or a pharmaceutically acceptable salt, an ester, a solvate, a hydrate, a polymorph thereof;

(3) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation;

wherein the administering of the pharmaceutical composition results in an improvement in diastolic relaxation as compared to istaroxime administered by intravenous infusion prior to the accumulation period, thereby treating the individual having acute heart failure.

56. The method of claim 55, wherein the individual is human.

57. The method of claim 55 or claim 56, wherein the improvement in diastolic relaxation is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure.
58. The method of any one of claims 55 to 57, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 10 ng/ml for an accumulation period of at least about 6 hours.
59. The method of claim 58, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 15 ng/ml for an accumulation period of at least about 6 hours.
60. The method of claim 59, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 20 ng/ml for an accumulation period of at least about 6 hours.
61. The method of any one of claims 55 to 59, wherein istaroxime is administered at a dose comprised between 0.2 µg/kg/min and 1.5 µg/kg/min.
62. The method of claim 61, wherein istaroxime is administered at a dose comprised between 0.25 µg/kg/min and 1.0 µg/kg/min.
63. The method of any one of claims 55 to 62, wherein the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).
64. The method of any one of claims 55 to 63, wherein the individual is undergoing a therapeutic treatment for heart failure with one or more further therapeutically active ingredients.
65. The method of claim 64, wherein the further therapeutically active ingredients are selected among ACE inhibitors, AIRBs, diuretics, Ca channel blockers, beta-blockers, digitalis, NO donors, vasodilators, SERCA2a stimulators, neprilisin (NEP) inhibitors, myosin filament activators, recombinant relaxin-2 mediators, recombinant NP protein, activators of the soluble Guanylate Cyclase (sGC), beta-arrestin ligand of Angiotensin II receptor.
66. The method of claim 65, wherein the diuretic, is selected among furosemide, bumetanide, torasemide, metolazone, an aldosterone antagonist, in particular spironolactone and eplerenone; thiazide diuretics, in particular hydrochlorothiazide, metolazone, chlorthalidone.

67. The method of claim 65, wherein the ACE inhibitor is selected among Lisinopril and Ramipril.

68. The method of claim 65, wherein the Angiotensin II receptor blocker is selected among valsartan, candesartan, olmesartan, telmisartan and losartan.

69. The method of claim 65, wherein the Angiotensin receptor/neprilysin inhibitor is sacubitril.

70. The method of claim 65, wherein the beta-blocker is selected among carvedilol and metoprolol.

71. The method of claim 65, wherein the Vasodilator is selected among Hydralazine, optionally combined with isosorbide dinitrate, nitrates, in particular nitroglycerin or isosorbide nitrate, amlodipine, felodipine, non-dihydropyridines, in particular diltiazem or verapamil.

72. The method of claim 65, wherein the agent for the treatment of heart failure is selected from Digoxin, entresto, omecantiv, serelaxin, ularitide, levosimendan.

73. The method of any one of claims 65 to 72, wherein the therapeutic treatment for heart failure is preceding or concomitant or subsequent to istaroxime administration.

74. A method for treating an individual having acute heart failure, the method comprising the steps of:

(1) providing an individual having acute heart failure;

(2) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation as measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure;


(3) administering to the individual a therapeutically effective amount of a first pharmaceutical composition comprising; (i) a first pharmaceutically acceptable carrier; and (ii) istaroxime or a pharmaceutically acceptable salt, solvate, or hydrate thereof administered at a first dose of at least about 1.0 µg/kg/min for a first infusion duration sufficient to improve diastolic relaxation in the individual;

(4) administering, upon improvement of diastolic relaxation in the individual, a therapeutically effective amount of a second pharmaceutical composition comprising; (i) a second pharmaceutically acceptable carrier; and (ii) istaroxime or a pharmaceutically

acceptable salt, solvate, or hydrate thereof administered at a second dose of between about 0.25 µg/kg/min to about 1.0 µg/kg/min for a second infusion duration sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite, wherein:


(a) the plasma concentration level is greater than about 5 ng/ml for an accumulation period of at least about 6 hours; and

(b) the istaroxime metabolite comprises formula (II) or (III):



**PST 2915**

Formula (II)



**PST 3093**

Formula (III)

or a pharmaceutically acceptable salt, an ester a solvate, a hydrate, a polymorph thereof;

wherein the administering of the first pharmaceutical composition, second pharmaceutical composition, or both, results in an improvement in diastolic relaxation, thereby treating the individual having acute heart failure.

75. The method of claim 74, wherein the individual is human.

76. The method of any one of claims 74 to 75, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 10 ng/ml for an accumulation period of at least about 6 hours.

77. The method of claim 76, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 15 ng/ml for an accumulation period of at least about 6 hours.

78. The method of claim 77, wherein the infusion duration is sufficient to produce in the individual a plasma concentration level of an istaroxime metabolite greater than about 20 ng/ml for an accumulation period of at least about 6 hours.

79. The method of any one of claims 74 to 78, wherein istaroxime is administered at a first dose of at least about 1.5 µg/kg/min.

80. The method of any one of claims 74 to 79, wherein istaroxime is administered at a second dose comprised about 0.5 µg/kg/min.

81. The method of any one of claims 74 to 80, wherein the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).

82. A method for treating an individual having heart failure, the method comprising the steps of:

(1) providing an individual having heart failure;

(2) administering to the individual a therapeutically effective amount of a pharmaceutical composition comprising; (i) a pharmaceutically acceptable carrier; and (ii) a compound as in claim 22; wherein said administering comprises intravenous infusion for an infusion duration of longer than 6 hours;

(3) measuring one or more parameters of heart function, wherein the one or more parameters of heart function comprises diastolic relaxation;

wherein the administering of the pharmaceutical composition results in an improvement in diastolic relaxation as compared to an equivalent compound administered by intravenous infusion for less than 6 hours, thereby treating the individual having acute heart failure.

83. The method of claim 82, wherein the individual is human.

84. The method of claim 82 or claim 83, wherein the diastolic relaxation improvement is measured by echocardiographic parameter E/A, E/e' or by pulmonary capillary wedge pressure.

85. The method of any one of claims 82 to 84, wherein the infusion duration is up to about 24 hours.

86. The method of claim 85, wherein the infusion duration is up to about 36 hours.

87. The method of claim 86, wherein the infusion duration is up to about 48 hours.

88. The method of any one of claims 82 to 87, wherein the compound is administered at a dose of about 0.2 µg/kg/min to about 1.5 µg/kg/min.

89. The method of claim 88, wherein the compound is administered at a dose of about 0.25 µg/kg/min to about 1.0 µg/kg/min.

90. The method of any one of claims 82 to 89, wherein the individual is diagnosed with heart failure with preserved ejection fraction (HFpEF) or mid-range reduction ejection fraction (HFmEF).