

BACKGROUND OF THE INVENTION**1. Field of the Invention**


This invention relates to novel analogs of rebeccamycin which posses antineoplastic properties.

2. Background Art

U.S. Pat. Nos. 4,487,925 and 4,552,842 disclose the anti-tumor agent designated rebeccamycin, and the 5'-methyl and 5',2',3'',6''-tetraacetate derivatives thereof, and a process for producing the same agent by cultivating a rebeccamycin-producing strain of Nocardia aerocolonigenes, preferably Nocardia aerocolonigenes ATCC 39243, or a rebeccamycin-producing mutant thereof in an aqueous nutrient medium containing assimilable sources of carbon and nitrogen under submerged aerobic conditions until a substantial amount of rebeccamycin is produced. Recently, Nocardia aerocolonigenes, ATCC 39243 was reclassified as Saccharothrix aerocolonigenes, ATCC 39243 (Bush et al, J. Antibiotics, 40:663-678, 1987).


SUMMARY OF THE INVENTION

The present invention provides new analogs of the antitumor agent designated rebeccamycin (Formula I).




Formula I

More specifically, there are provided rebeccamycin analogs of the Formulas II and III below,



Formula II



Formula III

wherein X_1 and X_2 are independently fluorine or hydrogen, provided that X_1 and X_2 are not simultaneously hydrogen; as well as pharmaceutically acceptable acid addition salts of such analogs.

The compounds of the Formulas II and III are produced by supplementing the cultures of a rebeccamycin producing strain of Saccharothrix aerocolonigenes with the appropriate tryptophan analog.

DESCRIPTION OF THE DRAWINGS

Figure 1 shows the IR spectrum for the compound of Formula IV.

Figure 2 shows the ^1H -NMR for the compound of Formula IV.

Figure 3 shows the ^{13}C -NMR for the compound of Formula IV.

Figure 4 shows the IR spectrum for the compound of Formula V.

Figure 5 shows the ^1H -NMR for the compound of Formula V.

Figure 6 shows the ^{13}C -NMR for the compound of Formula V.

Figure 7 shows the IR spectrum for the compound of Formula VIII.

Figure 8 shows the ^1H -NMR for the compound of Formula VIII.

Figure 9 shows the ^{13}C -NMR for the compound of Formula VIII.

Figure 10 shows the mass spectrum of the compound of Formula IV.

Figure 11 shows the UV spectrum of the compound of Formula IV.

Figure 12 shows the mass spectrum of the compound of Formula V.


Figure 13 shows the UV spectrum of the compound of Formula V.

Figure 14 shows the mass spectrum of the compound of Formula VIII.

Figure 15 shows the UV spectrum of the compound of Formula VIII.

DETAILED DESCRIPTION OF THE INVENTION

U.S. Pat. Nos. 4,487,925 and 4,552,842 disclose the production and isolation of the antitumor agent designated rebeccamycin (Formula I)




Formula I


The above-mentioned rebeccamycin compound is the principal component of the fermentation of the rebeccamycin producing strain of Saccharothrix aerocolonigenes.

It has now been found according to the present invention that the fermentation procedure disclosed in U.S. Pat. Nos. 4,487,925 and 4,552,842 can be carried out in the presence of certain

tryptophan analogs to produce new analogs of rebeccamycin having valuable antitumor properties. The rebeccamycin analogs of the present invention have the Formulas II and III below.



Formula II




Formula III

wherein X_1 and X_2 are fluorine or hydrogen, provided that X_1 and X_2 are not simultaneously hydrogen; as well as pharmaceutically acceptable acid addition salts thereof.


A preferred example of the compounds of the present invention

is the compound having the structural formula IV below:




Formula IV

Another preferred example of the compounds of the present invention is the compound having the structural formula V below:




Formula V

Another preferred example of the compounds of the present invention is the compound having the structural formula VI below:



Formula VI


Still another preferred example of the compounds of the present invention is the compound having the structural formula VII below:



Formula VII


Still another preferred example of the compounds of the present invention is the compound having the structural formula VIII below:

9




Formula VIII

Another preferred example of the compounds of the present invention is the compound having the structural formula IX below:




Formula IX

Another preferred example of the compounds of the present invention is the compound having the structural formula X below:



Formula X

Yet another preferred example of the compounds of the present invention is the compound having the structural formula XI below:



Formula XI

In the present process a tryptophan analog is added to the rebeccamycin fermentation medium and is incorporated during fermentation into the rebeccamycin structure creating a corresponding rebeccamycin analog. A more extensive description of the process is given below and in the illustrative examples which follow.

Preparation of the Antibiotics ^{1.1}

The compounds Formulas IV - XI are produced by cultivating a rebeccamycin producing strain of Saccharothrix aerocolonigenes, with DL-4-fluorotryptophan, DL-5-fluorotryptophan, DL-6-fluorotryptophan, or DL-7-fluorotryptophan. The preferred producing organism is a novel strain of Saccharothrix aerocolonigenes, previously designated as Nocardia aerocolonigenes strain C38,383-RK2 (ATCC 39243) in United States Patent 4,487,925. Recently, this strain was reclassified as Saccharothrix aerocolonigenes (Bush et al., J. Antibiotics 40:668-678, 1987) and is designated herein as Saccharothrix aerocolonigenes strain C38,383-RK2 (ATCC 39243). This strain was isolated from a soil sample collected in Panama. A biologically pure culture of strain C38,383-RK2 has been deposited with the American Type Culture Collection, Rockville, Maryland, and added to their permanent collection of microorganisms as ATCC 39243. This culture, designated as C38,383-RK2, is also maintained as a dormant culture in lyophile tubes and cryogenic vials in the Bristol-Myers Squibb Co. Pharmaceutical Research and Development Division Culture Collection, 5 Research Parkway, Wallingford, Connecticut 06492.

The taxonomic studies on strain C38,383-RK2 (ATCC 39243) have been described in detail in United States Patent 4,487,925 and in J. Antibiotics 40:668-678, 1987. The strain has been classified as a novel strain of Saccharothrix aerocolonigenes.

It is to be understood that the present invention is not limited to use of the particular preferred strain ATCC 39243 or to organisms fully answering its description. It is especially intended to include other rebeccamycin producing strains or mutants

of the described organism which can be produced by conventional means such as x-radiation, ultraviolet radiation, treatment with nitrogen mustard, phage exposure and the like.

In practicing the present process, a rebeccamycin-producing strain of Saccharothrix aerocolonigenes, having the identifying characteristics of strain C38,383-RK2 (ATCC 39243), or a mutant or variant thereof, is cultivated in a conventional aqueous nutrient medium supplemented with the appropriate tryptophan analog. For optimal production of the compound of Formulas VI and X, the medium should be supplemented with DL-4-fluorotryptophan. For optimal production of the compounds of Formulas IV and IX, the medium should be supplemented with DL-5-fluorotryptophan. For optimal production of the compounds of Formula V and VIII, the medium should be supplemented with DL-6-fluorotryptophan. For optimal production of the compounds of Formulas VII and XI, the medium should be supplemented with DL-7-fluorotryptophan. The organism is grown in a nutrient medium containing known nutritional sources for actinomycetes. Thus, the organism is grown in a nutrient medium containing an assimilable carbon source such as sucrose, lactose, glucose, rhamnose, fructose, glycerol or soluble starch. The medium should also contain an assimilable nitrogen source such as fishmeal, peptone, peanut meal, cottonseed meal, corn steep liquor, amino acids or ammonium salts. Nutrient inorganic salts can also be incorporated in the medium so as to provide sodium, potassium, ammonium, calcium, phosphate, sulfate, nitrate, carbonate and like ions. Trace elements such as copper, manganese, iron zinc, etc. are added to the medium if desired, or they may be present as impurities of other constituents of the media. Submerged aerobic

conditions are preferably employed for the production of large quantities of antibiotic, although for production of limited amounts, surface cultures and bottles may also be used. The general procedures used for the cultivation of other actinomycetes are applicable to the present invention.

Production of the antibiotics of the present invention can be effected by any temperature conducive to satisfactory growth of the producing organism, e.g. 18° to 39° C and is conveniently carried out at a temperature of about 28° C. The fermentation may be carried out in flasks or in laboratory or industrial fermentors of various capacity.

When tank fermentation is to be used it is desirable to produce a vegetative inoculum in a nutrient broth by inoculating a small volume of the culture medium with a slant, a cryopreservative culture or a lyophilized culture of the producing organism. After obtaining a viable and active inoculum in this manner, it is transferred aseptically to the fermentation tank charged with production medium for large scale production of the antibiotic of the present invention. The medium in which the vegetative inoculum is grown can be the same as, or different from, that utilized in the tank as long as it is such that a good growth of the producing organism is obtained and supplemented with the appropriate fluorotryptophan. Further agitation may be provided by a mechanical impeller. Antifoam agents such as lard oil or silicone oil may also be added if needed. Antibiotic production is monitored by high performance liquid chromatography assay or by conventional biological assay. In general, optimum production of the antibiotics of the present invention is achieved after

incubation of about 6 days.

Isolation and purification of the so-obtained derivatives may be carried out by conventional chromatographic procedures.

Physical and Chemical Properties:

The compounds of Formulas IV, V and VIII have the following Physical and Chemical Properties:

Compound of Formula IV

Description: Bright yellow amorphous solid

Molecular Formula: $C_{26}H_{19}F_2N_3O_7$

Molecular weight: 523.454

Mass Spectrum: Kratos MS 25 Mass Spectrometer. FABMS 524 ($M + H$)⁺, 361 ($M-162$, loss of glucose).

Ultraviolet Spectrum: Hewlett Packard 845A Diode Array Spectrophotometer. Concentration 1.0 mg/100 ml MeOH. Neutral λ_{max} nm ($E 1\%/1cm$): 405, 322(457), 288, 277, 259, 230(348).

Infrared Spectrum: Perkin-Elmer 1800 FTIR Spectrometer. KBr Pellet (cm^{-1}): 3340, 2915, 1752, 1708, 1628, 1591, 1484, 1462, 1392, 1331, 1292, 1247, 1191, 1112, 1081, 1052, 947, 904 795, 762, 752, 752, 511.

360 MHz 1H -NMR: Bruker Model AM-3000 Spectrometer. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical shifts (ppm): 11.76 (s, 1H), 11.23 (s, 1H), 8.85 (dd, 1H), 8.77 (dd, 1H), 8.01 (dd, 1H), 7.68 (dd, 1H), 7.46 (m, 2H), 6.29 (d, 1H), 6.12 (t, 1H), 5.45 (d, 1H), 5.17 (d, 1H), 4.95 (d, 1H), 4.07 (d, 1H), 3.95 (m, 2H), 3.81 (d, 1H), 3.59 (m, 2H).

90 MHz ^{13}C -NMR: Bruker Model AM-3000 Spectrometer. Proton

decoupled spectrum. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical shifts (ppm): 170.9, 170.8, 157.0(d), 157.0(d), 138.6, 137.2, 130.7, 129.2, 121.7(d), 121.4(d), 121.3, 119.6, 116.6, 115.1, 115.0(d), 114.6(d), 113.3(d), 113.2(d), 109.1(d), 109.1(d), 84.8, 78.7, 76.5, 73.2, 67.6, 58.3.

Solubility: Soluble in DMSO, DMF, THF, acetone, EtOAc, MeOH.

Thin Layer Chromatography (R_f values): Normal phase (silica gel 60); EtOAc: 0.10. EtOAc-MeOH (9:1 v/v): 0.41. Reversed phase (C_{18}); 0.1M NH_4OAc -MeOH- CH_3CN (2:4:4 v/v): 0.67.

Compound of Formula V

Description: Bright yellow amorphous solid

Molecular Formula: $C_{26}H_{19}F_2N_3O_7$

Molecular Weight: 523.454

Mass Spectrum: Kratos MS 25 Mass Spectrometer. FABMS 524 ($M + H$)⁺, 361 ($M-162$, loss of glucose).

Ultraviolet Spectrum: Hewlett Packard 8452A Diode Array Spectrophotometer. Concentration 1.0 mg/100 ml MeOH. Neutral, λ_{max} nm (E 1%/1cm): 398(60), 316(640), 280(259), 256(313), 226(525), 204(497).

Infrared Spectrum: Perkin-Elmer 1800 FTIR Spectrometer. KBr Pellet (cm^{-1}): 3324, 2927, 1745, 1701, 1623, 1580, 1491, 1471, 1452, 1412, 1384, 1330, 1233, 1172, 1116, 1075, 1048, 1016, 963, 916, 829, 764, 745, 646, 635, 617, 498, 490.

360 MHz 1H -NMR: Bruker Model AM-3000 Spectrometer. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical shifts (ppm): 11.77 (s, 1H), 11.18 (s, 1H), 9.12 (dd, 1H), 9.05 (dd, 1H),

7.85(dd,1H), 7.43(dd,1H), 7.23(t,2H), 6.26(d,1H), 6.24(s,1H), 5.31(d,1H), 5.04(d,1H), 3.98(m,2H), 3.87(m,1H), 3.66(s,2H).

90 MHz ^{13}C -NMR: Bruker Model AM-3000 Spectrometer. Proton decoupled spectrum. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical shifts (ppm): 170.8, 170.7, 161.8(d), 161.8(d), 143.2(d), 141.5(d), 130.1, 128.7, 126.0(d), 125.9(d), 120.9, 119.3, 118.2, 118.1, 117.7, 116.5, 108.8(d), 108.6(d), 98.9(d), 98.3(d), 84.7, 78.6, 76.5, 73.1, 67.5, 58.3.

Solubility: Soluble in DMSO, DMF, THF, MeOH, acetone, EtOAc.

Thin Layer Chromatography (R_f values): Normal phase (silica gel 60); EtOAc: 0.40. EtOAc-MeOH (9:1 v/v): 0.63. Reversed phase (C_{18}); 0.1M NH_4OAc -MeOH- CH_3CN (2:4:4: v/v): 0.60.

Compound of Formula VIII

Description: Bright yellow amorphous solid

Molecular Formula: $\text{C}_{27}\text{H}_{21}\text{F}_2\text{N}_3\text{O}_7$

Molecular Weight: 537.481

Mass Spectrum: Kratos MS 25 Mass Spectrometer. FABMS 538 ($\text{M} + \text{H}$)⁺, 361 ($\text{M}-176$, loss of 4-O-methylglucose).

Ultraviolet Spectrum: Hewlett Packard 8452A Diode Array Spectrophotometer. Concentration 1.2 mg/100 ml MeOH. Neutral λ_{max} nm ($E 1\%/1\text{cm}$): 398(103), 316(1050), 280(387), 256(454), 228(730).

Infrared Spectrum: Perkin-Elmer 1800 FTIR Spectrometer. KBr Pellet (cm^{-1}): 3324, 2938, 1747, 1703, 1623, 1580, 1491, 1471, 1452, 1412, 1384, 1330, 1233, 1172, 1140, 1116, 1087, 1054, 963, 918, 828, 764, 649, 635, 618, 498.

360 MHz ^1H -NMR: Bruker Model AM-3000 Spectrometer. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical

shifts (ppm): 11.77 (s, 1H), 11.18 (s, 1H), 9.12 (dd, 1H), 9.05 (dd, 1H), 7.88 (dd, 1H), 7.41 (dd, 1H), 7.23 (m, 2H), 6.25 (d, 1H), 6.24 (s, 1H), 5.36 (dd, 1H), 5.31 (d, 1H), 5.04 (d, 1H), 3.97 (m, 2H), 3.60-3.95 (m, 4H).

90 MHz ^{13}C -NMR: Bruker Model AM-3000 Spectrometer. Proton decoupled spectrum. Dual carbon-proton probe, 5mm. Solvent d_6 -DMSO. Observed chemical shifts (ppm): 170.8, 170.7, 161.8 (d), 161.7 (d), 143.1 (d), 141.5 (d), 130.1, 128.7, 126.0 (d), 125.9 (d), 120.9, 119.4, 118.2, 118.1, 117.7, 116.5, 108.8 (d), 108.6 (d), 98.7 (d), 98.3 (d), 84.4, 77.2, 77.2, 76.2, 73.2, 59.9, 58.5 ppm.

Solubility: Soluble in DMSO, DMF, THF, acetone, EtOAc, MeOH.

Thin Layer Chromatography (R_f values): Normal phase (silica gel 60); EtOAc: 0.72. EtOAc-MeOH (9:1 v/v): 0.89. Reversed phase (C_{18}); 0.1M NH_4OAc -MeOH- CH_3CN (2:4:4 v/v): 0.59.

Biological Properties:

Representative compounds of the present invention were tested against the transplanted mouse leukemia P388 to determine in vivo antitumor activity (Tables 1-3). CDF_1 mice were implanted intraperitoneally ("ip") with 10^6 P388 leukemia cells obtained from DBA/2 donor mice bearing this transplantable murine leukemia. The CDF_1 leukemic mice were treated ip with either saline (control mice) or doses of the compound of Formulas IV, V and VIII once at Day 1 post-tumor inoculation. These animals were observed daily and their deaths recorded. Average body weight changes (from the day of leukemia implant to the day of last treatment) were determined for all groups as a means of reflecting drug toxicity. The incidence of mice alive in each group on Day 5 post-tumor implant was recorded as an additional means of assessing drug

toxicity. No therapeutic result was considered as meaningful if more than 1 mouse per treatment group had died by Day 5. Each treatment group consisted of 4 to 6 mice; control groups contained 10 mice. The number of mice, if any, surviving to Day 30 (the 1st day of the experiment) was also recorded.

Therapeutic efficacy was evaluated by determining the median survival time ("MST") of mice treated with the compound of Formula IV, V and VIII and comparing it to the MST of parallel control mice. This comparison was made by dividing the MST of the former by the latter and multiplying by 100 to derive a parameter called the percent T/C value. A percent T/C of $\geq 125\%$ was considered to represent a meaningful increase in lifespan and, hence, an active result.

As shown in Table 1, the compound of Formula IV is active against P388 leukemia at dose levels ranging from 10 to 90 mg/kg. The best effect was achieved at a dosage of 30 mg/kg and consisted of a percent T/C of 175%. Toxicity was not observed even at the highest dose (90 mg/kg) tested.

As shown in Table 2, the compound of Formula V is active against P388 leukemia at dose levels ranging from 0.8 to 102.4 mg/kg. The best effect was achieved at a dosage of 102.4 mg/kg and consisted of a percent T/C of 178%. Toxicity was not observed even at the highest dose (102.4 mg/kg) tested.

As shown in Table 3, the compound of Formula VIII is active against P388 leukemia at dose levels ranging from 0.8 to 102.4 mg/kg. The best effect was achieved at a dosage of 51.2 mg/kg and consisted of a percent T/C of 206%. Toxicity was not observed at the highest dose (102.4 mg/kg) tested.

Table 1. Effect of Compound of Formula IV on P388 Leukemia^a
(Day 1 Treatment)

Dose, ip. (mg/kg/inj)	Median Survival Times (Days)	± T/C	Average Weight Change (g)	No. of Mice Alive on	
				Day 5	Day 30
90	16.5	165	-0.9	4/4	0/4
30	17.5	175	0.6	4/4	0/4
10	14.5	145	0.7	4/4	0/4
Control	10.0	100		10/10	0/10

^a Mice were implanted with 10^6 P388 leukemia cells, and treatments with the compound of Formula IV were begun 1 day later. Control mice were given saline injections.

Table 2. Effect of Compound of Formula V on P388 Leukemia^a
(Day 1 Treatment)

Dose, ip. (mg/kg/inj)	Median Survival Times (Days)	± T/C	Average Weight Change (g)	No. of Mice Alive on	
				Day 5	Day 30
102.4	16.0	178	-0.1	6/6	0/6
51.2	15.0	167	-0.2	6/6	0/6
25.6	14.5	161	-0.2	6/6	0/6
12.8	15.0	167	0.8	6/6	0/6
6.4	15.0	167	-1.1	6/6	0/6
3.2	14.0	156	0.0	6/6	0/6
1.6	14.0	156	-0.7	6/6	0/6
0.8	13.0	144	0.1	6/6	0/6
Control	9.0	100	1.3	10/10	0/10

^a Mice were implanted with 10^6 P388 leukemia cells and treatments with the compound of Formula V were begun 1 day later. Control mice were given saline injections.

Table 3. Effect of Compound of Formula VIII on P388 Leukemia^a
(Day 1 Treatment)

Dose, ip. (mg/kg/inj)	Median Survival Times (Days)	± T/C	Average Weight Change (g)	No. of Mice Alive on	
				Day 5	Day 30
102.4	14.5	161	0.2	6/6	0/6
51.2	18.5	206	0.3	6/6	0/6
25.6	14.0	156	0.1	6/6	0/6
12.8	14.5	161	-0.1	6/6	0/6
6.4	16.0	178	0.1	6/6	0/6
3.2	16.5	183	-0.3	6/6	0/6
1.6	14.0	156	-0.1	6/6	0/6
0.8	14.0	156	-0.4	6/6	0/6
Control	9.0	100	1.3	10/10	0/10

^a Mice were implanted with 10⁶ P388 leukemia cells and treatments with the compound of Formula VIII were begun 1 day later. Control mice were given saline injections.

The present invention includes within its scope pharmaceutical compositions which comprise an effective or tumor-inhibiting amount of a rebeccamycin analog of the present invention, or a pharmaceutically acceptable acid addition salt thereof, in combination with an inert pharmaceutically acceptable carrier or diluent.

According to another aspect of the invention, a method is provided for therapeutically treating an animal (preferably mammalian) host effected by a malignant tumor which comprises administering to such host an effective tumor-inhibiting dose of the antibiotic of the present invention or a pharmaceutically acceptable acid addition salt thereof.

Examples of suitable compositions include solid compositions

for oral administration such as tablets, capsules, pills, powders and granules, liquid compositions for oral administration such as solutions, suspensions, syrups and elixirs and preparations for parenteral administration such as sterile solutions, suspensions or emulsions. They may also be manufactured in the form of sterile solid compositions which can be dissolved in sterile water, physiological saline or some other sterile injectable medium immediately before use.

It will be appreciated that the actual preferred dosages of the rebeccamycin analogs of the present invention will vary according to the particular compound being used, the particular composition formulated, the mode of application and the particular situs, host and disease being treated. Many factors that modify the action of the drug will be taken into account by those skilled in the art, e.g. age, body weight, sex, diet, time of administration, rate of excretion, condition of the host, drug combinations, reaction sensitivities and severity of the disease. Administration can be carried out continuously or periodically within the maximum tolerated dose. Optimal application rates for a given set of conditions can be readily ascertained by those skilled in the art using conventional dosage determination tests.

The present invention is illustrated by the following examples which are not intended to be construed as limiting the scope of the invention.

Example 1. Preparation of cryopreservative culture of Saccharothrix aerocolonigenes strain C38,383-RK2 (ATCC 19243).

Saccharothrix aerocolonigenes strain C38,383-RK2 was

maintained as a cryopreservative culture stored at -80°C in a Revco ultralow temperature freezer. To prepare a cryopreservative culture, strain C38,383-RK2 was transferred in test tubes on slants of yeast-malt extract agar supplemented with CaCO_3 which consisted of

dextrose	4.0g
yeast extract	4.0g
malt extract	10g
CaCO_3	1.5g
agar	15g
deionized water	q.s. to 1 liter

The agar slant was incubated at 28°C for 7-10 days. Vegetative culture was prepared by transferring the surface growth from the slant culture to a 500 ml Erlenmeyer flask containing 100 ml of a sterile vegetative medium consisting of

Cerelose (Corn Products)	30g
Pharmamedia (Traders Oil Mill Co.)	10g
Nutrisoy (Archer Daniels Midland Co.)	10g
CaCO_3	3g
deionized water	q.s. to 1 liter

This vegetative culture was incubated at 28°C for 48 hours on a rotary shaker set at 250 rev/min. The vegetative culture was mixed with equal volume of cryoprotective solution consisting of

Sucrose	100g
glycerol	200g
deionized water	q.s. to 1 liter

Four ml portions of this mixture were transferred to sterile cryogenic tube (5 ml capacity, Corning) and were frozen in a dry ice-acetone bath. The frozen vegetative cultures were then stored at -80°C in a Revco ultralow temperature freezer.

Example 2. Preparation of vegetative culture of Saccharothrix aerocolonigenes strain C38,383-RK2 (ATCC 39243)

Vegetative culture was prepared by transferring 4 ml of the cryopreservative culture to a 500 ml Erlenmeyer flask containing 100 ml of a sterile vegetative medium having the same composition as the vegetative medium described in Example 1. The vegetative culture was incubated at 28°C for 48 hours on a rotary shaker set at 250 rev/min.

Example 3. Preparation of the compound of Formula IV

Three ml of the vegetative culture of Example 2 was inoculated into 500 ml Erlenmeyer flasks each containing 100 ml of production medium consisting of

Staclipse J-UB starch (A.E. Staley)	10g
KH_2PO_4	2g
MgSO_4	1g
L-threonine	2.5g
CaCO_3	2g
deionized water	q.s. to 1 liter

The production culture was incubated at 28°C on a rotary shaker set at 250 rev/min. After 48 hours of fermentation, DL-5-fluorotryptophan was added to the culture at a final concentration

of 1 mg/ml. The culture was allowed to incubate at 28°C and shaken at 250 rev/min for additional 4 days. Production of the compound of Formula IV was monitored by HPLC. Optimal production of the compound of Formula IV at a concentration of 23-32 mg/ml was generally obtained at 6 days of fermentation (i.e. 4 days after the addition of DL-5-fluorotryptophan).

Example 4. Preparation of the compound of Formula V and VIII.

Three ml of vegetative culture of Example 2 was inoculated into 500 ml Erlenmeyer flasks each containing 100 ml of production medium having the same composition as described in Example 3. The production culture was incubated at 28°C on a rotary shaker set at 250 rev/min. After 48 hours of fermentation, DL-6-fluorotryptophan was added to the culture at a final concentration of 1 mg/ml. The culture was allowed to incubate at 28°C and shaken at 250 rev/min for additional 4 days. Production of the compounds of Formula V and VIII were monitored by HPLC. Optimal production of the compound of Formula V and VIII were generally obtained at 6 days of fermentation (i.e. 4 days after the addition of DL-6-fluorotryptophan) at a concentration of 36-58 µg/ml and 31-42 µg/ml respectively.

Example 5. Isolation and purification of the compounds of Formulas IV, V and VIII.

a) General Method

Solvents were not redistilled before use. Methanol, acetone, ethyl acetate, isopropyl ether, chloroform, tetrahydrofuran, ethyl ether and hexanes were ACS reagent grade. Water for HPLC refers to

in-house deionized water from a Barnstead Nanopure II system. Tetrahydrofuran, methanol and acetonitrile for HPLC use were B & J Brand HPLC grade solvents. Ammonium acetate was Fisher HPLC grade.

Normal phase thin layer chromatograph (tlc) was carried out on Silica gel 60, F-254 plates (EM Reagents, Cat. #5765, 5 x 10 cm, by 0.25 mm thick). Reversed phase tlc was accomplished with Whatman MKC₁₈ plates (Cat. #4803-110, 0.2 mm thick). Plates were developed in Whatman cylindrical jars with caps and 10 ml of eluant. Rebeccamycin analogs were visible as yellow zones in normal lighting or as yellow fluorescing zones with 254 nm or 366 nm ultraviolet light.

To whole broths was added Dicalite (speed plus) filter aid. After brief stirring the broths were filtered on large Buchner funnels or on a Tolhurst Centerslung Centrifugal Filter Unit (Model 1B15, Ametek, Inc.). Filtrates were discarded. Mycelial mats were stirred in THF or THF-acetone mixtures for one hour, filtered, and the Dicalite further rinsed with acetone until it no longer fluoresced yellow under UV light. The combined filtrates were concentrated under reduced pressure to yield crude extracts.

A vacuum liquid chromatography (VLC) apparatus consists of a Buchner funnel (Kontes, Art. #K-954100) containing a sealed-in sintered glass disc (M porosity), a side hose connection for vacuum and a lower 24/40 joint for attachment of receiving flasks. Initially, the least polar eluting solvents pulled through under vacuum to form tightly packed 5 cm adsorbent bed heights. Samples were preadsorbed onto adsorbent and applied to funnels as slurries, or applied in a solution of the least polar eluting solvent. Step gradients were carried out where predetermined volumes of

increasingly polar eluant constituted the fractions. The funnel was sucked dry after each volume of eluant. Fractions were concentrated and combined on the basis of tlc analysis..

Apparatus for size exclusion chromatography consisted of the following: A Glenco column (2.5 I.D. x 100 cm) equipped with solvent resistant teflon and plates; Fluid Metering, Inc. FMI lab pump (Model RP-G150); Glenco glass reservoir (500 ml); Isco Model 328 fraction collector. Columns were slurry packed with Sephadex LH-20 (Pharmacia) preswollen in the eluting solvent. Solvent was delivered in a downward manner through the column at a rate controlled by the lab pump.

The following components were used to construct a semi-preparative HPLC system: Waters Associates Model 590 Solvent Delivery System pump; Knauer model 87 Variable Wavelength Detector. Waters Associates model SR-204 Strip Chart Recorder; Whatman Partisil 10 ODS-3 column (10 mm x 50 cm); 316 stainless steel tubing (0.23 mm i.d.).

b.) Isolation and purification of the compound of Formula IV

Whole broth (5 liters) was filtered with Dicalite and the mycelial mat extracted by stirring in THF (2 liters) for 1 hour. After further filtering, and an additional THF rinse (1 liter), the combined filtrates were concentrated under reduced pressure to yield 4.5 g crude extract. The extract was adsorbed onto 6.5g Lichroprep Si 60 Silica gel (EM Science, Art. 9336, 15-25 microns) and applied to a 60 ml VLC funnel containing an additional 24.5 g silica gel. A hexane-ethyl acetate step gradient was carried out (200 ml volumes), followed by a 200 ml volume THF wash. The THF

wash (156 mg) was dissolved in 4 ml THF and applied to a column containing 160 g Sephadex LH-20 equilibrated with THF (bed height 90 cm, bed volume 430 ml). Flow rate 3.75 ml/min. As could be determined visually, a major yellow band (62 mg) eluted in the first one-fourth of the second bed volume. Final purification was accomplished by reversed phase (C_{18}) HPLC chromatography with a flow rate of 4 ml/min (0.1 M NH_4OAc -THF (60-40)). Detection was at 320 nm. Eluting at 39 minutes was the major analog (23 mg) designated as the compound of the Formula IV.

c.) Isolation and purification of the compounds of Formula V and VIII

Whole broth (2 liters) was filtered using Dicalite filter aid. The mycelial mat, after stirring in THF (2 liters) for 45 minutes, was filtered, and rinsed with an additional volume of THF (1.5 liters). The filtrate was concentrated under reduced pressure to yield 3.39 g crude extract. The crude extract was triturated with five 50 ml volumes of THF, which were combined and concentrated to yield 0.74 g of THF soluble residue. This mass contained the bulk of the yellow fluorescing materials. The THF soluble material was adsorbed onto 2 g Silica Gel H (Merck, 10-40 microns) and applied to a 30 ml VLC funnel containing an additional 11 g Silica Gel H. A hexane-ethyl acetate step gradient was carried out using 100 ml volumes of eluant. The two major rebeccamycin analogs were separated cleanly in this manner. The less polar yellow zone (141 mg) eluted with hexane-ethyl acetate (1:1) and the more polar analog (105 mg) with hexane-ethyl acetate (1:3).

The more polar analog (105 mg) was dissolved in 2 ml THF and

applied to a column containing 160 g Sephadex LH-20 (bed height 90 cm, bed volume 430 ml) preswollen in THF. The flow rate was 4 ml/min. The main yellow band eluted at 1.25 bed volumes as could be determined visually to yield the compound of Formula V (77 mg).


The less polar analog from VLC (141 mg) was dissolved in 2 ml MeOH and applied to a column containing 110 g Sephadex LH-20 preswollen in MeOH (bed height 80cm, bed volume 400 ml). The flow rate 3.5 ml/min. The main yellow zone eluted at the end of the fourth bed volume to yield the compound of formula VIII (70 mg).



29


What is claimed is:

1. A rebeccamycin analog having the formula:



wherein X_1 and X_2 are independently fluorine or hydrogen, provided that X_1 and X_2 are not simultaneously hydrogen, as well as pharmaceutically acceptable acid addition salts thereof.

2. A rebeccamycin analog having the formula:





wherein X_1 and X_2 are independently fluorine or hydrogen, provided that X_1 and X_2 are not simultaneously hydrogen, as well as



30



1 pharmaceutically acceptable acid addition salts thereof.

3. The compounds of Claim 1 having the formulae







31







32

4. The compounds of Claim 2 having the formulae:





5. A process for producing the rebeccamycin analogs of claims 1 to 4,
or a pharmaceutically acceptable acid addition salt thereof,
which comprises cultivating a rebeccamycin-producing strain of
Saccharothrix aerocolonigenes in an aqueous nutrient medium in the
presence of a tryptophan analog, until a substantial amount of the
desired rebeccamycin analog is produced by said organism in said



1 culture medium and recovering the desired rebeccamycin derivative
from the culture medium in a substantially pure form.

5

6... The process of Claim 5, wherein the rebeccamycin-producing
strain is Saccharothrix aerocolonigenes ATCC 39243.

10

7. The process of Claim 11 wherein the tryptophan analog is DL-4-
fluorotryptophan;

DL-5-flurotryptophan;

15

DL-6-fluorotryptophan; or

DL-7-fluorotryptophan.

20

8. A pharmaceutical composition comprising an
effective tumor-inhibiting amount of at least one compound
of anyone of Claims 1 to 4, in association with a
pharmaceutical acceptable substantially nontoxic carrier
or excipient.

25

9. A process for preparing the pharmaceutical composition
of Claim 8 which comprises incorporating at least one
compound of anyone of Claims 1 to 4 into a pharmaceutically
acceptable substantially non-toxic carrier or excipient.

30

10. The use of at least one compound of anyone of
Claims 1 to 4 for preparing a pharmaceutical composition
for therapeutically treating a mammalian host affected
by a tumor.

35